

ON CERTAIN PROPERTIES OF THE CLASS $\mathcal{U}(\lambda)$

NAJLA M. ALARIFI, MILUTIN. OBRADOVIĆ, AND NIKOLA. TUNESKI

ABSTRACT. Let \mathcal{A} be the class of functions analytic in the unit disk $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ and normalized such that $f(z) = z + a_2z^2 + a_3z^3 + \dots$. In this paper we study the class $\mathcal{U}(\lambda)$, $0 < \lambda \leq 1$, consisting of functions f from \mathcal{A} satisfying

$$\left| \left(\frac{z}{f(z)} \right)^2 f'(z) - 1 \right| < \lambda \quad (z \in \mathbb{D})$$

and give results regarding the Zalcman Conjecture, the generalised Zalcman conjecture, the Krushkal inequality and the second and third order Hankel determinant.

1. INTRODUCTION AND PRELIMINARIES

Let \mathcal{A} be the class of functions that are analytic in the open unit disk $\mathbb{D} = \{z : |z| < 1\}$ and normalised such that $f(0) = f'(0) - 1 = 0$, i.e., have expansion $f(z) = z + a_2z^2 + a_3z^3 + \dots$.

Further, let

$$\mathcal{U}(\lambda) = \{f \in \mathcal{A} : |U_f(z) - 1| < \lambda, z \in \mathbb{D}\},$$

where $0 < \lambda \leq 1$ and

$$U_f(z) := \left(\frac{z}{f(z)} \right)^2 f'(z).$$

The functions from $\mathcal{U}(\lambda)$ are univalent, its special case when $\lambda = 1$ first studied in [5] and more details on them can be found in [6, 7, 12]).

An intriguing fact about $\mathcal{U}(\lambda)$ is that in spite the class \mathcal{S}^* of starlike functions is very large and contains most classes of univalent functions, it doesn't contain the class $\mathcal{U} \equiv \mathcal{U}(1)$, i.e., \mathcal{U} is not in \mathcal{S}^* , nor vice versa. Namely, the function $-\ln(1-z)$ is convex, thus starlike, but not in \mathcal{U} because $U_f(0.99) = 3.621\dots > 1$, while the function f defined by $\frac{z}{f(z)} = 1 - \frac{3}{2}z + \frac{1}{2}z^3 = (1-z)^2 \left(1 + \frac{z}{2}\right)$ is in \mathcal{U} and such that $\frac{zf'(z)}{f(z)} = -\frac{2(z^2+z+1)}{z^2+z-2} = -\frac{1}{5} + \frac{3i}{5}$ for $z = i$. This rear property is the main reason why the class \mathcal{U} (and $\mathcal{U}(\lambda)$) attracts huge attention in the past decades.

In this paper we study class $\mathcal{U}(\lambda)$ regarding the Zalcman Conjecture, the generalised Zalcman conjecture, the Krushkal inequality and the second and third order Hankel determinant which will be defined in corresponding sections further in the paper.

For the study we will need the following result proven in [8] as a part of the proof of Theorem 1.

2010 *Mathematics Subject Classification.* Primary: 30C45, 30C50; Secondary: 30C80.

Key words and phrases. analytic, class $\mathcal{U}(\lambda)$, Zalcman conjecture, generalised Zalcman conjecture, Krushkal inequality, Hankel determinant.

Lemma 1.1. *For each function f in $\mathcal{U}(\lambda)$, $0 < \lambda \leq 1$, there exists function ω_1 , analytic in \mathbb{D} , such that $|\omega_1(z)| \leq |z| < 1$, and $|\omega_1'(z)| \leq 1$, for all $z \in \mathbb{D}$, with*

$$(1.1) \quad \frac{z}{f(z)} = 1 - a_2z - \lambda z\omega_1(z).$$

Additionally, for $\omega_1(z) = c_1z + c_2z^2 + \dots$,

$$(1.2) \quad |c_1| \leq 1, \quad |c_2| \leq \frac{1}{2}(1 - |c_1|^2) \quad \text{and} \quad |c_3| \leq \frac{1}{3} \left[1 - |c_1|^2 - \frac{4|c_2|^2}{1 + |c_1|} \right].$$

Using (1.1), we have

$$z = [1 - a_2z - \lambda z\omega_1(z)]f(z),$$

and after equating the coefficients,

$$(1.3) \quad \begin{aligned} a_3 &= \lambda c_1 + a_2^2, \\ a_4 &= \lambda c_2 + 2\lambda a_2 c_1 + a_2^3, \\ a_5 &= \lambda c_3 + 2\lambda a_2 c_2 + \lambda^2 c_1^2 + 3\lambda a_2^2 c_1 + a_2^4, \end{aligned}$$

that we will use later on.

We will also use the next result ([10]).

Lemma 1.2. *Let $f \in \mathcal{U}(\lambda)$ for $0 < \lambda \leq 1$, and be given by $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$. Then*

$$(1.4) \quad |a_2| \leq 1 + \lambda.$$

If $|a_2| = 1 + \lambda$, then f must be of the form

$$(1.5) \quad f(z) = \frac{z}{1 - (1 + \lambda)e^{i\phi}z + \lambda e^{2i\phi}z^2}$$

for some $\phi \in [0, 2\pi]$.

In the same paper ([10]) it was conjectured that for functions in $\mathcal{U}(\lambda)$, $|a_n| \leq \sum_{i=0}^{n-1} \lambda^i$ holds sharply, and was claimed to be proven in the case $n = 3$:

$$(1.6) \quad |a_3| \leq 1 + \lambda + \lambda^2.$$

The proof rely on another claim, that for all functions f from $\mathcal{U}(\lambda)$,

$$(1.7) \quad \frac{f(z)}{z} \prec \frac{1}{(1+z)(1+\lambda z)}.$$

Recently, in [2], the second claim, and consequently the first one also, was proven to be wrong by giving a counterexample. Still, the subset of $\mathcal{U}(\lambda)$ when the inequality (1.6) and subordination (1.7) hold is nonempty, as the function

$$(1.8) \quad f_\lambda(z) = \frac{z}{(1-z)(1-\lambda z)} = \sum_{n=1}^{\infty} \frac{1-\lambda^n}{1-\lambda} z^n = z + (1+\lambda)z^2 + (1+\lambda+\lambda^2)z^3 + \dots$$

shows. Here $\left. \frac{1-\lambda^n}{1-\lambda} \right|_{\lambda=1} = n$ for all $n = 1, 2, 3, \dots$

Let note that from (1.3) and (1.6),

$$|a_3| = |\lambda c_1 + a_2^2| \leq 1 + \lambda + \lambda^2,$$

i.e.,

$$(1.9) \quad |\lambda c_1 + a_2^2| \leq 1 + \lambda + \lambda^2,$$

which we will use further in the proofs.

2. ZALCMAN AND GENERALISED ZALCMAN CONJECTURE FOR THE CLASS $\mathcal{U}(\lambda)$

In 1960 Zalcman posed the conjecture:

$$|a_n^2 - a_{2n-1}| \leq (n-1)^2 \quad (n \in \mathbb{N}, n \geq 2),$$

proven in 2014 by Krushkal ([1]) for the whole class \mathcal{S} by using complex geometry of the universal Teichmüller space. In 1999, Ma ([3]) proposed a generalized Zalcman conjecture,

$$|a_m a_n - a_{m+n-1}| \leq (m-1)(n-1) \quad (m, n \in \mathbb{N}, m \geq 2, n \geq 2),$$

which is still an open problem, closed by Ma for the class of starlike functions and for the class of univalent functions with real coefficients. Ravichandran and Verma in [11] closed it for the classes of starlike and convex functions of given order and for the class of functions with bounded turning.

In [9], the authors treated some particulars cases of those problems and obtained sharp results. Since $\mathcal{U}(\lambda) \subseteq \mathcal{U}$ for $0 < \lambda \leq 1$, the same results are valid for the class $\mathcal{U}(\lambda)$, but those results are not sharp for the class $\mathcal{U}(\lambda)$. In this part of the paper we find better and sharp results for the functions f from $\mathcal{U}(\lambda)$ satisfiung subordination (1.7).

Theorem 2.1. *Let f in $\mathcal{U}(\lambda)$ be of the form $f(z) = z + a_2 z^2 + a_3 z^3 + \dots$. Then*

- (i) $|a_2^2 - a_3| \leq \lambda$, $0 < \lambda \leq 1$;
- (ii) $|a_3^2 - a_5| \leq \lambda(1 + \lambda)^2$, $\frac{1}{2} \left(\frac{\sqrt{23}}{3} - 1 \right) \leq \lambda \leq 1$, if the third coefficient of f satisfies inequality (1.6).

Both results are sharp as the function f_λ given by (1.8) shows.

Proof.

(i) From $a_3 = \lambda c_1 + a_2^2$, we have $|a_2^2 - a_3| = |-\lambda c_1| \leq \lambda$.

(ii) Using the relations (1.2), (1.4) and (1.9), we have

$$\begin{aligned} |a_3^2 - a_5| &= |\lambda c_3 + 2\lambda a_2 c_2 + \lambda a_2^2 c_1| \\ &= \lambda |c_3 + 2a_2 c_2 - \lambda c_1^2 + c_1(\lambda c_1 + a_2^2)| \\ &\leq \lambda (|c_3| + 2|a_2||c_2| + \lambda|c_1|^2 + |c_1||\lambda c_1 + a_2^2|) \\ &\leq \lambda \left[\frac{1}{3} \left(1 - |c_1|^2 - \frac{4|c_2|^2}{1 + |c_1|} \right) + 2(1 + \lambda)|c_2| + \lambda|c_1|^2 + (1 + \lambda + \lambda^2)|c_1| \right] \\ (2.1) \quad &= \lambda g_1(|c_1|, |c_2|), \end{aligned}$$

where

$$g_1(x, y) = \frac{1}{3} \left(1 - x^2 - \frac{4y^2}{1+x} \right) + 2(1 + \lambda)y + \lambda x^2 + (1 + \lambda + \lambda^2)x,$$

$$0 \leq x = |c_1| \leq 1, \quad 0 \leq y = |c_2| \leq \frac{1}{2}(1 - x^2).$$

Since $\frac{\partial g_1(x,y)}{\partial x} > 0$ for all $(x, y) \in E := \{(x, y) : 0 \leq x \leq 1, 0 \leq y \leq \frac{1}{2}(1-x^2)\}$, then g_1 has no singular points in the interior of E and g_1 attains its maximum on the boundary of E .

For $x = 0$, we have $0 \leq y \leq \frac{1}{2}$ and

$$\begin{aligned} g_1(0, y) &= \frac{1}{3}(1-4y^2) + 2(1+\lambda)y \\ &= \frac{1}{3}[-4y^2 + 6(1+\lambda)y + 1] \\ &\leq 1 + \lambda. \end{aligned}$$

Similarly, for $y = 0$, we have $0 \leq x \leq 1$ and

$$\begin{aligned} g_1(x, 0) &= \frac{1}{3}(1-x^2) + \lambda x^2 + (1+\lambda+\lambda^2)x \\ &= \left(\lambda - \frac{1}{3}\right)x^2 + (1+\lambda+\lambda^2)x + \frac{1}{3} \\ &\leq (1+\lambda)^2. \end{aligned}$$

Finally, for $0 \leq x \leq 1$ and $y = \frac{1}{2}(1-x^2)$, we have

$$g_1\left(x, \frac{1}{2}(1-x^2)\right) = 1 + \lambda + \left(\frac{4}{3} + \lambda + \lambda^2\right)x - x^2 - \frac{1}{3}x^3 = \varphi_1(x).$$

Since $\varphi_1'(x) = \frac{4}{3} + \lambda + \lambda^2 - 2x - x^2 \geq \lambda^2 + \lambda - \frac{5}{3} \geq 0$ for $\frac{1}{2}\left(\frac{\sqrt{23}}{3} - 1\right) \leq \lambda \leq 1$, then φ_1 is an increasing function for $0 \leq x \leq 1$, and $\varphi_1(x) \leq (1+\lambda)^2$.

Using (2.1) and all these facts, we have the statement of Theorem 3.1(ii). \square

Remark 2.1. For $0 < \lambda < \frac{1}{2}\left(\frac{\sqrt{23}}{3} - 1\right) = 0.8844\dots$ we have that the function φ_1 attains its maximum for $x_0 = \sqrt{\lambda^2 + \lambda + \frac{7}{3}} - 1$ and $\varphi_1(x_0) = \frac{2}{3}\left(\lambda^2 + \lambda + \frac{7}{3}\right)^{\frac{3}{2}} - 1 - \lambda^2$, and so $|a_3^2 - a_5| \leq \lambda\left(\frac{2}{3}\left(\lambda^2 + \lambda + \frac{7}{3}\right)^{\frac{3}{2}} - 1 - \lambda^2\right)$.

Next we consider the Generalized Zalcman conjecture.

Theorem 2.2. Let f in $\mathcal{U}(\lambda)$ be of the form $f(z) = z + a_2z^2 + a_3z^3 + \dots$. Then

- (i) $|a_2a_3 - a_4| \leq \lambda(\lambda + 1)$, $0 < \lambda \leq 1$;
- (ii) $|a_2a_4 - a_5| \leq \lambda(1 + \lambda + \lambda^2)$, $\sqrt{\frac{2}{3}} \leq \lambda \leq 1$, if the third coefficient of f satisfies inequality (1.6).

Both results are sharp as the function f_λ given by (1.8) shows.

Proof.

(i) Using (1.2), (1.3) and (1.4) we obtain

$$\begin{aligned} |a_2a_3 - a_4| &= |\lambda c_2 + \lambda a_2 c_1| \\ &\leq \lambda(|c_2| + |a_2||c_1|) \\ &\leq \lambda\left[\frac{1}{2}(1 - |c_1|^2) + (1 + \lambda)|c_1|\right] \\ &\leq \lambda(1 + \lambda). \end{aligned}$$

(ii) Similarly,

$$\begin{aligned} |a_2a_4 - a_5| &= \lambda|c_3 + a_2c_2 + a_2^2c_1 + \lambda c_1^2| \\ &\leq \lambda(|c_3| + |a_2||c_2| + |c_1||a_2^2 + \lambda c_1|) \\ &\leq \lambda \left[\frac{1}{3} \left(1 - |c_1|^2 - \frac{4|c_2|^2}{1 + |c_1|} \right) + (1 + \lambda)|c_2| + (1 + \lambda + \lambda^2)|c_1| \right] \\ &= \lambda g_2(|c_1|, |c_2|), \end{aligned}$$

where

$$g_2(x, y) = \frac{1}{3} \left(1 - x^2 - \frac{4y^2}{1 + x} \right) + (1 + \lambda)y + (1 + \lambda + \lambda^2)x,$$

$$0 \leq x = |c_1| \leq 1, 0 \leq y = |c_2| \leq \frac{1}{2}(1 - x^2).$$

Since $\frac{\partial g_2(x, y)}{\partial x} > 0$ for all $(x, y) \in E$, then the function g_2 attains its maximum on the boundary of E . In that sense, let consider all cases

For $x = 0$, we have $0 \leq y \leq \frac{1}{2}$ and

$$\begin{aligned} g_2(0, y) &= \frac{1}{3}(1 - 4y^2) + (1 + \lambda)y \\ &= \frac{1}{3}[-4y^2 + 3(1 + \lambda)y + 1] \\ &\leq \frac{1}{2}(1 + \lambda), \end{aligned}$$

since $\lambda \geq \sqrt{\frac{2}{3}} > \frac{1}{3}$.

For $0 \leq x \leq 1$ and $y = 0$ we have

$$\begin{aligned} g_2(x, 0) &= \frac{1}{3}(1 - x^2) + (1 + \lambda + \lambda^2)x \\ &= \frac{1}{3}[-x^2 + 3(1 + \lambda + \lambda^2)x + 1] \\ &\leq 1 + \lambda + \lambda^2. \end{aligned}$$

Finally, for $0 \leq x \leq 1$, $y = \frac{1}{2}(1 - x^2)$ we have

$$g_2 \left(x, \frac{1}{2}(1 - x^2) \right) = \frac{1}{2}(1 + \lambda) + \left(\lambda^2 + \lambda + \frac{4}{3} \right) x - \frac{1}{2}(1 + \lambda)x^2 - \frac{1}{3}x^3 \equiv \varphi_2(x).$$

Since for $\lambda^2 \geq \frac{2}{3}$,

$$\varphi_2'(x) = \lambda^2 + \lambda + \frac{4}{3} - (1 + \lambda)x - x^2 \geq \lambda^2 - \frac{2}{3} \geq 0,$$

we realize that φ_2 is an increasing function and

$$\varphi_2(x) \leq \varphi_2(1) = 1 + \lambda + \lambda^2.$$

From all the previous facts, we have the statement of the theorem. \square

3. KRUSHKAL INEQUALITY FOR THE CLASS $\mathcal{U}(\lambda)$

The inequality

$$|a_n^p - a_2^{p(n-1)}| \leq 2^{p(n-1)} - n^p$$

was introduced by Krushkal who claims to have proven its sharpness for the whole class of univalent functions in [1]. This claim is not confirmed by the mathematical community.

In this section we give direct proof over the class $\mathcal{U}(\lambda)$ for the cases $n = 4$, $p = 1$ and $n = 5$, $p = 1$.

Theorem 3.1. *Let f in $\mathcal{U}(\lambda)$ be of the form $f(z) = z + a_2z^2 + a_3z^3 + \dots$. Then*

- (i) $|a_4 - a_2^3| \leq 2\lambda(\lambda + 1)$;
- (ii) $|a_5 - a_2^4| \leq \lambda(3 + 5\lambda + 3\lambda^2)$, if the third coefficient of f satisfies inequality (1.6).

Both results are sharp as the function f_λ given by (1.8) shows.

Proof.

(i) Similarly as in the proofs of two previous theorem, we have

$$\begin{aligned} |a_4 - a_2^3| &= \lambda|c_2 + 2a_2c_1| \\ &\leq \lambda(|c_2| + 2|a_2||c_1|) \\ &\leq \lambda \left[\frac{1}{2}(1 - |c_1|^2) + 2(1 + \lambda)|c_1| \right] \\ &\leq 2\lambda(1 + \lambda). \end{aligned}$$

(ii) We can easy verify that

$$\begin{aligned} |a_5 - a_2^4| &= \lambda|c_3 + 2a_2c_2 + \lambda c_1^2 + 3a_2^2c_1| \\ &= \lambda|c_3 + 2a_2c_2 - 2\lambda c_1^2 + 3c_1(\lambda c_1 + a_2^2)| \\ &\leq \lambda(|c_3| + 2|a_2||c_2| + 2\lambda|c_1|^2 + 3|c_1||\lambda c_1 + a_2^2|) \\ &\leq \lambda \left[\frac{1}{3} \left(1 - |c_1|^2 - \frac{4|c_2|^2}{1 + |c_1|} \right) + 2(1 + \lambda)|c_2| + 2\lambda|c_1|^2 + 3(1 + \lambda + \lambda^2)|c_1| \right] \\ &= \lambda g_3(|c_1|, |c_2|), \end{aligned}$$

where

$$g_3(x, y) = \frac{1}{3} \left(1 - x^2 - \frac{4y^2}{1 + x} \right) + 2(1 + \lambda)y + 2\lambda x^2 + 3(1 + \lambda + \lambda^2)x,$$

$$0 \leq x = |c_1| \leq 1, 0 \leq y = |c_2| \leq \frac{1}{2}(1 - x^2).$$

Since $\frac{\partial g_3(x, y)}{\partial x} > 0$, $(x, y) \in E$, then we have that g_3 attains its maximum on the boundary of E .

For $x = 0$, $0 \leq y \leq \frac{1}{2}$ we have

$$g_3(0, y) = \frac{1}{3}(1 - 4y^2) + 2(1 + \lambda)y \leq (1 + \lambda).$$

For $0 \leq x \leq 1$, $y = 0$ we have

$$\begin{aligned} g_3(x, 0) &= \left(2\lambda - \frac{1}{3}\right)x^2 + 3(1 + \lambda + \lambda^2)x + \frac{1}{3} \\ &\leq 2\lambda + 3(1 + \lambda + \lambda^2) \\ &= 3 + 5\lambda + 3\lambda^2. \end{aligned}$$

Finally, for $0 \leq x \leq 1$, $y = \frac{1}{2}(1 - x^2)$,

$$g_3\left(x, \frac{1}{2}(1 - x^2)\right) = 1 + \lambda + \left(\frac{1}{3} + 3(1 + \lambda + \lambda^2)\right)x - (1 - \lambda)x^2 - \frac{1}{3}x^3 \equiv \varphi_3(x).$$

Since

$$\begin{aligned} \varphi_3'(x) &= \frac{1}{3} + 3(1 + \lambda + \lambda^2) - 2(1 - \lambda)x - x^2 \\ &\geq \frac{1}{3} + 5\lambda + 3\lambda^2 > 0, \end{aligned}$$

φ_3 is an increasing function and

$$\varphi_3(x) \leq \varphi_3(1) = 3 + 5\lambda + 3\lambda^2.$$

All those facts imply the conclusion of theorem. \square

4. HANKEL DETERMINANT OF SECOND AND THIRD ORDER

Let $f \in \mathcal{A}$. Then the q th Hankel determinant of f is defined for $q \geq 1$, and $n \geq 1$ by

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \cdots & a_{n+q} \\ \vdots & \vdots & & \vdots \\ a_{n+q-1} & a_{n+q} & \cdots & a_{n+2q-2} \end{vmatrix}.$$

Thus, the second and the third Hankel determinants are, respectively,

$$\begin{aligned} H_2(2) &= a_2a_4 - a_3^2, \\ H_3(1) &= a_3(a_2a_4 - a_3^2) - a_4(a_4 - a_2a_3) + a_5(a_3 - a_2^2). \end{aligned}$$

In [8] the authors showed that for the class \mathcal{U} , the following estimates are sharp:

$$|H_2(2)| \leq 1 \quad \text{and} \quad |H_3(1)| \leq \frac{1}{4}.$$

Here we generalize this result for the classes $\mathcal{U}(\lambda)$.

Theorem 4.1. *Let $f \in \mathcal{U}(\lambda)$, $0 < \lambda \leq 1$, and $f(z) = z + a_2z^2 + a_3z^3 + \cdots$. Then*

$$|H_2(2)| \leq \frac{\lambda(\lambda+1)}{2} \quad \text{and} \quad |H_3(1)| \leq \frac{\lambda^2}{4}.$$

The second result is sharp due to the function $f(z) = \frac{z}{1-\lambda/2z^3} = z + \frac{\lambda z^4}{2} + \frac{\lambda^2 z^7}{4} + \cdots$.

Proof. Using the relation (1.3), after some calculations we have:

$$\begin{aligned}
|H_2(2)| &= |a_2 a_4 - a_3^2| = \lambda |a_2 c_2 - \lambda c_1^2| \\
&\leq \lambda (|a_2| |c_2| + \lambda |c_1|^2) \\
&\leq \lambda \left[(\lambda + 1) \left(\frac{1 - |c_1|^2}{2} \right) + \lambda |c_1|^2 \right] \\
&\leq \lambda \left[\frac{\lambda + 1}{2} + \left(\frac{\lambda - 1}{2} \right) |c_1|^2 \right] \\
&\leq \frac{\lambda(\lambda + 1)}{2},
\end{aligned}$$

when $0 \leq |c_1|^2 \leq 1$, because $\lambda - 1 \leq 0$.

In a similar way, using (1.2) and (1.3), after some calculations, for the third order Hankel determinant we have

$$\begin{aligned}
|H_3(1)| &= \lambda^2 |c_1 c_3 - c_2^2| \leq \lambda^2 (|c_1| |c_3| + |c_2|^2) \\
&\leq \lambda^2 \left[\frac{|c_1|}{3} \left(1 - |c_1|^2 - \frac{4|c_2|^2}{1 + |c_1|} \right) + |c_2|^2 \right] \\
&= \frac{\lambda^2}{3} \left[|c_1| - |c_1|^3 + \frac{3 - |c_1|}{1 + |c_1|} \cdot |c_2|^2 \right] \\
&\leq \frac{\lambda^2}{3} \left[|c_1| - |c_1|^3 + \frac{3 - |c_1|}{1 + |c_1|} \cdot \frac{(1 - |c_1|^2)^2}{4} \right] \\
&= \frac{\lambda^2}{12} (3 - 2|c_1|^2 - |c_1|^4) \leq \frac{3\lambda^2}{12} = \frac{\lambda^2}{4}.
\end{aligned}$$

Equality is attained for the function $f(z) = \frac{z}{1 - (\lambda/2)z^3} = z + \frac{\lambda z^4}{2} + \frac{\lambda^2 z^7}{4} + \dots$. \square

REFERENCES

- [1] S.L. Krushkal, A short geometric proof of the Zalcman and Bieberbach conjectures, preprint, arXiv:1408.1948.
- [2] Li L., Ponnusamy S., Wirths K.J., Relations of the class $\mathcal{U}(\lambda)$ to other families of functions, arXiv:2104.05346.
- [3] W. Ma, Generalized Zalcman conjecture for starlike and typically real functions, *J. Math. Anal. Appl.*, 234 (1999), pp. 328–339.
- [4] Obradović M., Starlikeness and certain class of rational functions, *Math. Nachr.* 175 (1995), 263–268.
- [5] Obradović, M.; Pascu, N. N.; Radomir, I. A class of univalent functions, *Math. Japon.*, 44 (1996), no. 3, 565–568.
- [6] Obradović M., Ponnusamy S., New criteria and distortion theorems for univalent functions, *Complex Variables Theory Appl.*, 44 (3) (2001), 173–191.
- [7] Obradović M., Ponnusamy S., On the class \mathcal{U} , *Proc. 21st Annual Conference of the Jammu Math. Soc. and a National Seminar on Analysis and its Application*, 11–26, 2011.
- [8] M. Obradović and N. Tuneski, Some properties of the class \mathcal{U} , *Ann. Univ. Mariae Curie-Skłodowska Sect. A* **73** (2019), no. 1, 49–56.
- [9] M. Obradović and N. Tuneski, Zalcman and generalized Zalcman conjecture for the class \mathcal{U} , arXiv:2005.14301.
- [10] M. Obradović, S. Ponnusamy and K.-J. Wirths, Geometric studies on the class $\mathcal{U}(\lambda)$, *Bull. Malays. Math. Sci. Soc.* **39** (2016), no. 3, 1259–1284.
- [11] V. Ravichandran, S. Verma, Generalized Zalcman conjecture for some classes of analytic functions. *J. Math. Anal. Appl.* 450 (2017), no. 1, 592–605.

- [12] Thomas D.K., Tuneski N., Vasudevarao A., *Univalent functions. A primer*, De Gruyter Studies in Mathematics, 69. De Gruyter, Berlin, 2018.

DEPARTMENT OF MATHEMATICS, IMAM ABDULRAHMAN BIN FAISAL UNIVERSITY, DAMMAM 31113,
KINGDOM OF SAUDI ARABIA

Email address: najarifi@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF CIVIL ENGINEERING, UNIVERSITY OF BELGRADE,
BULEVAR KRALJA ALEKSANDRA 73, 11000 BELGRADE, SERBIA

Email address: obrad@grf.bg.ac.rs

DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTY OF MECHANICAL ENGINEERING,
SS. CYRIL AND METHODIUS UNIVERSITY IN SKOPJE, KARPOŠ II B.B., 1000 SKOPJE, REPUBLIC OF
NORTH MACEDONIA

Email address: nikola.tuneski@mf.edu.mk