THE TRUNCATED HAMBURGER MOMENT PROBLEMS WITH GAPS IN THE INDEX SET

ALJAŽ ZALAR

ABSTRACT. In this article we solve four special cases of the truncated Hamburger moment problem (THMP) of degree 2k with one or two missing moments in the sequence. As corollaries we obtain, by using appropriate substitutions, the solutions to bivariate truncated moment problems of degree 2k for special curves. Namely, for the curves $y = x^3$ (first solved by Fialkow [Fia11]), $y^2 = x^3$, $y = x^4$ where a certain moment of degree 2k + 1 is known and $y^3 = x^4$ with a certain moment given. The main technique is the completion of the partial positive semidefinite matrix (ppsd) such that the conditions of Curto and Fialkow's solution of the THMP are satisfied. The main tools are the use of the properties of positive semidefinite Hankel matrices and a result on all completions of a ppsd matrix with one unknown entry, proved by the use of the Schur complements for 2×2 and 3×3 block matrices.

1. INTRODUCTION

For $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ and $i = (i_1, \ldots, i_d) \in \mathbb{Z}^d_+$, we set $|i| = i_1 + \ldots + i_d$ and $x^i = x_1^{i_1} \cdots x_d^{i_d}$. Given a real *d*-dimensional multisequence $\beta = \beta^{(2k)} = \{\beta_i\}_{i \in \mathbb{Z}^d_+, |i| \leq 2k}$ of degree 2k and a closed subset K of \mathbb{R}^d , the **truncated moment problem (TMP)** supported on K for β asks to characterize the existence of a positive Borel measure μ on \mathbb{R} with support in K, such that

(1.1)
$$\beta_i = \int_K x^i d\mu(x) \quad \text{for} \quad i \in \mathbb{Z}^d_+, |i| \le 2k$$

If such measure exists, we say that β has a representing measure supported on K and μ is its K-representing measure.

We denote by $M(k) = M(k)(\beta) = (\beta_{i,j})_{i,j=0}^k$ the moment matrix associated with β , where the rows and columns are indexed by X^i , $|i| \leq k$, in degree-lexicographic order. Let $\mathbb{R}[x]_k := \{p \in \mathbb{R}[x]: \deg p \leq k\}$ stand for the set of polynomials in d variables of degree at most k. To every $p := \sum_{i \in \mathbb{Z}_+^d, |i| \leq k} a_i x^i \in \mathbb{R}[x]_k$, we denote by $p(X) = \sum_{i \in \mathbb{Z}_+^d, |i| \leq k} a_i X^i$ the vector from the column space $\mathcal{C}(M(k))$ of the matrix M(k). Recall from [CF96], that β has a representing measure μ with the support supp μ being a subset of $\mathcal{Z}_p := \{x \in \mathbb{R}^d : p(x) = 0\}$ if and only if p(X) = 0. We say that the matrix M(k) is **recursively generated (rg)** if for $p, q, pq \in \mathbb{R}[x]_k$ such that p(X) = 0, it follows that (pq)(X) = 0.

The full moment problem (MP), where β_i is given for every $i \in \mathbb{Z}_+^d$, being the classical question in analysis and also due to its relation with real algebraic geometry via the duality with positive

FACULTY OF COMPUTER AND INFORMATION SCIENCE, UNIVERSITY OF LJUBLJANA, VEČNA POT 113, 1000 LJUBLJANA, SLOVENIA

E-mail address: aljaz.zalar@fri.uni-lj.si.

Date: July 21, 2020.

²⁰¹⁰ Mathematics Subject Classification. Primary 47A57, 47A20, 44A60; Secondary 15A04, 47N40.

Key words and phrases. Hamburger moment problem, truncated moment problems, representing measure, moment matrix.

Supported by the Slovenian Research Agency grants J1-2453, P1-0288.

ALJAŽ ZALAR

polynomials given by Haviland's theorem [Hav35], has been widely studied, see e.g., [Akh65, AhK62, KN77, Las09, Lau05, Lau09, Mar08, PS06, PS08, Put93, PV99, Sch91, Sch03, Sch17]. The TMP, which is more general than the full MP [Sto01], has been intensively studied in a series of papers by Curto and Fialkow [CF91, CF96, CF98a, CF98b, CF02, CF04, CF05, CF08] with the celeberated flat extension theorem they established as a core tool in the field. There are also various generalizations of the TMP (e.g., [?, Bol96, ?, DU18], to matrix moments, [BK10, ?] to tracial moments, [IKLS17] to infinitely many variables). Recently, Fialkow's core variety [Fia17] approach led to many new results on the TMP; see also [BF20, DS18]. A **concrete solution** to the TMP is a set of necessary and sufficient conditions for the existence of a K-representing measure. Among necessary conditions, M(k) must be psd and rg [CF91, CF98b], which also suffice in some cases. Concrete solutions to the TMP are known in the following cases:

- (1) (Truncated Hamburger moment problem (THMP)) d = 1 and $K = \mathbb{R}$. See [AhK62, Theorem I.3] or [Ioh82, Theorem A.II.1] for the special case of even k with an invertible moment matrix and [CF91, Section 3] for the general case.
- (2) (Truncated Hausdorff moment problem) d = 1 and $K = [0, \infty)$. See [KN77, p. 175] for the special case of an invertible moment matrix and [CF91, Section 5] for the general case.
- (3) (Truncated Stieltjes moment problem) d = 1 and K = [a, b], a < b. See [KN77, Theorems III.2.4 and II.2.3] and [CF91, Section 4] for the general case.
- (4) d = 2 and K is a curve p(x, y) = 0 with deg $p \le 2$. See [CF02, CF04, CF05, FN10, Fia14, CS16].
- (5) d = 2 and K is a curve $y = x^3$. See [Fia11].
- (6) d = 2 and the moment matrix has a special feature called *recursive determinateness*. See [CF13] for details.
- (7) (Extremal case) The rank of the moment matrix is the same as the cardinality of the corresponding variety; see [CFM08].
- (8) Some special cases are solved in [CS15, Fia17, Ble15, BF20].

In (5), β must satisfy certain numerical conditions, which are equivalent to the conditions from Corollary 3.3 below. The proof is by separating the nonsingular case from the singular one. In the nonsingular case the existence of a flat extension is established by a detailed and technically demanding analysis, while the singular case is done by the use of additional features of the moment matrix such as recursive determinateness and known results for such matrices.

In this article we present concrete solutions to the four cases of the THMP of degree 2k with some unknown moments $\beta_{i_1}, \ldots, \beta_{i_j}, 1 \leq i_1 \leq \cdots \leq i_j \leq 2k - 1$, in the sequence, which we call the **THMP with gaps** $(\beta_{i_1}, \ldots, \beta_{i_j})$. Namely, we solve the THMP with gaps (β_{2k-1}) , $(\beta_{2k-2}, \beta_{2k-1}), (\beta_1)$ and (β_1, β_2) . The motivation to solve this cases of the THMP with gaps is to obtain the solutions to the special cases of the 2-dimensional TMP. Namely, the solution of the THMP with gaps:

- (1) (β_{2k-1}) gives an alternative solution to the TMP with d = 2 and K being the curve $y = x^3$ (see (5) above). The advantage of our approach is that the proof is short and we also do not need to separate three subcases, i.e., k = 1, k = 2 and $k \ge 3$.
- (2) $(\beta_{2k-2}, \beta_{2k-1})$ solves the TMP with d = 2, K being the curve $y = x^4$ and in addition the moment $\beta_{3,2k-2}$ of degree 2k + 1 is known. To solve the TMP for the curve $y = x^4$ without this additional moment, one needs to solve the THMP with gaps $(\beta_{2k-5}, \beta_{2k-2}, \beta_{2k-1})$ which is a possible topic of future research.
- (3) (β_1) solves the TMP with d = 2 and K being the curve $y^2 = x^3$.

(4) (β_1, β_2) solves the TMP with d = 2, K being the curve $y^3 = x^4$ and known $\beta_{\frac{5}{3},0}$. By $\beta_{\frac{5}{3},0}$ we mean the moment of $x_1^{\frac{5}{3}}$, i.e., $\int_K x_1^{\frac{5}{3}} d\mu$. To solve the TMP for the curve $y^3 = x^4$ without this additional information, one needs to solve the THMP with gaps $(\beta_1, \beta_2, \beta_5)$, which is another open question for future research.

1.1. **Readers Guide.** The paper is organized as follows. In Section 2 we present the tools used in the proofs of our main results:

- Generalized Schur complements and verification of positive semidefiniteness of block matrices (Subsection 2.1).
- Properties of psd Hankel matrices (Subsection 2.2).
- The solution to the THMP (Subsection 2.3).
- A result about psd completions of partial psd matrices with one unknown entry (Subsetion 2.4).
- An extension principle for psd matrices (Subsection 2.5).
- A result about subsequences of moment sequences (Subsection 2.6).

In Section 3 we solve the THMP of degree 2k with gaps (β_{2k-1}) (see Theorem 3.1) and $(\beta_{2k-2}, \beta_{2k-1})$ (see Theorem 3.5). Corollary 3.3, being a special case of the (β_{2k-1}) -case, is the solution to the TMP with d = 2 and the curve $y = x^3$ as K, while Corollary 3.6, being a special case of the $(\beta_{2k-2}, \beta_{2k-1})$ -case, is the solution to the TMP with d = 2, the curve $y = x^4$ as K and an additional moment $\beta_{3,2k-2}$ known.

In Section 4 we solve the THMP of degree 2k with gaps (β_1) (see Theorem 4.1) and (β_1, β_2) (see Theorem 4.5). Corollary 4.4, being a special case of the (β_1) -case, is the solution to the TMP with d = 2 and the curve $y^2 = x^3$ as K, while Corollary 4.7, being a special case of the (β_1, β_2) -case, is the solution to the TMP with d = 2, the curve $y^3 = x^4$ as K and an additional moment $\beta_{\frac{5}{2},0}$ known.

Acknowledgement. I would like to thank Jaka Cimprič and Abhishek Bhardwaj for useful suggestions on the preliminary versions of this article.

2. PRELIMINARIES

In this section we present some tools which will be needed in the proofs of our main results in Sections 3 and 4.

We write $M_{n,m}$ (resp. M_n) for the set of $n \times m$ (resp. $n \times n$) real matrices. For a matrix M we denote by $\mathcal{C}(M)$ its column space. The set of real symmetric matrices of size n will be denoted by S_n . For a matrix $A \in S_n$ the notation $A \succ 0$ (resp. $A \succeq 0$) means A is positive definite (pd) (resp. positive semidefinite (psd)).

2.1. Generalized Schur complements. Let

(2.1)
$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in S_{n+n}$$

be a real matrix where $A \in M_n$, $B \in M_{n,m}$, $C \in M_{m,n}$ and $D \in M_m$. The generalized Schur complement [Zha05] of A (resp. D) in M is defined by

$$M/A = D - CA^+B$$
 (resp. $M/D = A - BD^+C$),

where A^+ (resp. D^+) stands for the Moore-Penrose inverse of A (resp. D).

- *Remark* 2.1. (1) If A (resp. D) is invertible, then M/A (resp. M/D) is the usual Schur complement of A (resp. D) in M.
 - (2) Note that $M/A = \begin{pmatrix} D & C \\ B & A \end{pmatrix} /A$.

The following theorem gives conditions for verifying positive semidefiniteness of a block matrix of size 2.

Theorem 2.2. [Alb69] Let

(2.2)
$$M = \begin{pmatrix} A & B \\ B^T & C \end{pmatrix} \in S_{n+m}$$

be a real symmetric matrix where $A \in S_n$, $B \in M_{n,m}$ and $C \in S_m$. Then the following conditions are equivalent:

- (1) $M \succeq 0$.
- (2) $C \succeq 0, C(B^T) \subseteq C(C) \text{ and } M/C \succeq 0.$
- (3) $A \succeq 0, C(B) \subseteq C(A)$ and $M/A \succeq 0$.

If m = 1 in (2.2), then rank $M \in \{ \operatorname{rank} A, \operatorname{rank} A+1 \}$. The following proposition characterizes w.r.t. the value of M/A when each of the possibilities occurs in the case M is psd.

Proposition 2.3. Let

$$M = \left(\begin{array}{cc} A & b \\ b^T & c \end{array}\right) \in S_{n+1}$$

be a real symmetric matrix where $A \in S_n$, $b \in \mathbb{R}^n$ and $c \in \mathbb{R}$. Then rank $M = \operatorname{rank} A$ if and only if M/A = 0. Otherwise rank $M = \operatorname{rank} A + 1$.

Proof. By Theorem 2.2, the psd assumption implies that $b \in C(A)$. By the properties of the Moore-Penrose inverse $\{A^+b + w : w \in \ker A\}$ is the set of solutions z of the system Az = b. Therefore,

(2.3)
$$\mathcal{C}(M) = \mathcal{C}\left(\begin{pmatrix} A & 0\\ b^T & c - b^T(A^+b + w) \end{pmatrix}\right) = \mathcal{C}\left(\begin{pmatrix} A & 0\\ b^T & M/A \end{pmatrix}\right),$$

where the second equality follows from the fact that A is symmetric, $b \in C(A)$ and $w \in \ker A$. Now, the statement of the proposition follows from (2.3).

The following proposition gives an explicit formula, called the *quotient formula* [CH69], for expressing the Schur complement of a 2×2 upper left-hand or a 2×2 lower right-hand block in a 3×3 block matrix using 2×2 block submatrices.

Proposition 2.4. Let

$$K = \begin{pmatrix} A & B & D \\ B^T & C & E \\ D^T & E^T & F \end{pmatrix} = \begin{pmatrix} M & D \\ E \\ \hline D^T & E^T & F \end{pmatrix} = \begin{pmatrix} A & B & D \\ B^T & N \\ D^T & N \end{pmatrix} \in S_{n_1+n_2+n_3}$$

be a 3×3 block real matrix, where $A \in S_{n_1}, C \in S_{n_2}, F \in S_{n_3}$ are real symmetric matrices and $B \in M_{n_1,n_2}, D_{n_1,n_3}, E_{n_2,n_3}$ are rectangular matrices. If M and A are nonsingular, then

(2.4)
$$K/M = \begin{pmatrix} A & D \\ D^T & F \end{pmatrix} / A - \left[\begin{pmatrix} A & B \\ D^T & E^T \end{pmatrix} / A \right] (M/A)^{-1} \left[\begin{pmatrix} A & D \\ B^T & E \end{pmatrix} / A \right]$$

If N and C are nonsingular, then

(2.5)
$$K/N = \begin{pmatrix} C & B^T \\ B & A \end{pmatrix} / C - \left[\begin{pmatrix} C & E \\ B & D \end{pmatrix} / C \right] (N/C)^{-1} \left[\begin{pmatrix} C & B^T \\ E^T & D^T \end{pmatrix} / C \right]$$

Proof. By an easy calculation we have that

$$K/A = \begin{pmatrix} M/A & \begin{pmatrix} A & D \\ B^T & E \end{pmatrix} / A \\ \begin{pmatrix} A & B \\ D^T & E^T \end{pmatrix} / A & \begin{pmatrix} A & D \\ B^T & E \end{pmatrix} / A \end{pmatrix}.$$

Now the quotient formula [CH69] K/M = (K/A)/(M/A) yields (2.4).

By Remark 2.1 (2), it is true that K/N = L/N where

$$L = \begin{pmatrix} N & B^T \\ D^T \\ \hline B & D & A \end{pmatrix}$$

Now (2.5) follows from (2.4).

2.2. Hankel matrices. Let $k \in \mathbb{N}$. For

$$\beta = (\beta_0, \dots, \beta_{2k}) \in \mathbb{R}^{2k+1}$$

we denote by

$$A_{\beta} := (\beta_{i+j})_{i,j=0}^{k} = \begin{pmatrix} \beta_{0} & \beta_{1} & \beta_{2} & \cdots & \beta_{k} \\ \beta_{1} & \beta_{2} & \ddots & \ddots & \beta_{k+1} \\ \beta_{2} & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \beta_{2k-1} \\ \beta_{k} & \beta_{k+1} & \cdots & \beta_{2k-1} & \beta_{2k} \end{pmatrix} \in S_{k+1}$$

the corresponding Hankel matrix. We denote by $\mathbf{v}_{\mathbf{j}} := (\beta_{j+\ell})_{\ell=0}^k$ the (j+1)-th column of A_{β} , $0 \le j \le k$, i.e.,

$$A_{\beta} = \left(\begin{array}{ccc} \mathbf{v_0} & \cdots & \mathbf{v_k} \end{array} \right).$$

As in [CF91], the rank of β , denoted by rank β , is defined by

$$\operatorname{rank} \beta = \begin{cases} k+1, & \text{if } A_{\beta} \text{ is nonsingular,} \\ \min \left\{ i \colon \mathbf{v_i} \in \operatorname{span} \{ \mathbf{v_0}, \dots, \mathbf{v_{i-1}} \} \right\}, & \text{if } A_{\beta} \text{ is singular.} \end{cases}$$

We denote the upper left-hand corner of A_{β} of size m + 1 by

$$A_{\beta}(m) = (\beta_{i+j})_{i,j=0}^m \in S_{m+1}.$$

The following proposition is the alternative description of rank β if A_{β} is singular.

Proposition 2.5. [CF91, Proposition 2.2] Let $k \in \mathbb{N}$, $\beta = (\beta_0, \ldots, \beta_{2k})$, and assume that A is positive semidefinite and singular. Then

rank
$$\beta = \min\{j : 0 \le j \le k \text{ such that } A_{\beta}(j) \text{ is singular}\}.$$

Important property of psd Hankel matrices is the following rank principle.

Theorem 2.6. [CF91, Corollary 2.5] Let $k \in \mathbb{N}$, $\beta = (\beta_0, \ldots, \beta_{2k})$, $\tilde{\beta} = (\beta_0, \ldots, \beta_{2k-2})$, $A_\beta \succeq 0$ and $r = \operatorname{rank} \tilde{\beta}$. Then:

(1) rank $A_{\tilde{\beta}} = r$. (2) $r \leq \operatorname{rank} A_{\beta} \leq r + 1$. (3) rank $A_{\beta} = r + 1$ if and only if

$$\beta_{2k} > \varphi_0 \beta_{2k-r} + \ldots + \varphi_{r-1} \beta_{2k-1},$$

where $(\varphi_0, \dots, \varphi_{r-1}) := A_{\beta}(r-1)^{-1}(\beta_r, \dots, \beta_{2r-1})^T$

We will use the following corollary of Proposition 2.5 and Theorem 2.6 in the sequel.

Corollary 2.7. In the notation of Theorem 2.6, under the assumptions $A_{\beta} \succeq 0$, A_{β} is singular, and $r = \operatorname{rank} \tilde{\beta}$, then

$$r = \operatorname{rank} \beta = \operatorname{rank} A_{\beta}(r-1) = \operatorname{rank} A_{\beta}(r) = \ldots = \operatorname{rank} A_{\beta}(k-1) = \operatorname{rank} A_{\widetilde{\beta}}$$

We denote the lower right-hand corner of A_{β} of size m + 1 by

$$A_{\beta}[m] = (\beta_{i+j})_{i,j=m-k}^{k} = \begin{pmatrix} \beta_{2(k-m)} & \beta_{2(k-m)+1} & \beta_{2(k-m+1)} & \cdots & \beta_{2k-m} \\ \beta_{2(k-m)+1} & \beta_{2(k-m+1)} & \ddots & \ddots & \beta_{2k-m+1} \\ \beta_{2(k-m+1)} & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \beta_{2k-1} \\ \beta_{2k-m} & \beta_{2.k-m+1} & \cdots & \beta_{2k-1} & \beta_{2k} \end{pmatrix} \in S_{m+1}$$

Let

$$\beta^{(\text{rev})} := (\beta_{2k}, \beta_{2k-1}, \dots, \beta_0)$$

be the sequence obtained from β by reversing the order of numbers. Using Corollary 2.7 for a reversed sequence implies the following corollary.

Corollary 2.8. In the notation of Theorem 2.6, under the assumption $A_{\beta} \succeq 0$, A_{β} is singular and $r = \operatorname{rank} \widetilde{\beta}^{(rev)}$, where $\widetilde{\beta}^{(rev)} := (\beta_{2k}, \ldots, \beta_2)$, it holds that

$$r = \operatorname{rank} \beta^{(rev)} = \operatorname{rank} A_{\beta}[r-1] = \operatorname{rank} A_{\beta}[r] = \ldots = \operatorname{rank} A_{\beta}[k-1] = \operatorname{rank} A_{\widetilde{\beta}^{(rev)}}.$$

Proof. Corollary 2.7 used for $\beta^{(rev)}$ implies that

(2.6)
$$r = \operatorname{rank} \beta^{(\operatorname{rev})} = \operatorname{rank} A_{\beta^{(\operatorname{rev})}}(r-1) = \operatorname{rank} A_{\beta^{(\operatorname{rev})}}(r) = \ldots = A_{\beta^{(\operatorname{rev})}}(k-1) = \operatorname{rank} A_{\widetilde{\beta}^{(\operatorname{rev})}}.$$

For $\ell = 0, ..., k$ define the permutation matrices $P_{\ell} : \mathbb{R}^{\ell+1} \to \mathbb{R}^{\ell+1}$ by $e_i^{(\ell)} \mapsto e_{\ell+2-i}^{(\ell)}$, $i = 1, ..., \ell+1$, where $e_1^{(\ell)}, ..., e_{\ell+1}^{(\ell)}$ is the standard basis for $\mathbb{R}^{\ell+1}$. Note that $A_{\beta^{(rev)}}(\ell) = P_{\ell}^T A_{\beta}[\ell] P_{\ell}$ and hence rank $A_{\beta^{(rev)}}(\ell) = \operatorname{rank} A_{\beta}[\ell]$, which together with (2.6) implies the statement of the corollary.

A sequence $\beta = (\beta_0, \dots, \beta_{2k})$ with $r := \operatorname{rank} \beta$ is **positively recursively generated** if $A_\beta(r-1) \succ 0$ and denoting $(\varphi_0, \dots, \varphi_{r-1}) := A_\beta(r-1)^{-1}(\beta_r, \dots, \beta_{2r-1})^T$, it is true that

(2.7)
$$\beta_j = \varphi_0 \beta_{j-r} + \dots + \varphi_{r-1} \beta_{j-1} \quad \text{for} \quad j = r, \dots, 2k.$$

Note that (2.7) is equivalent to

(2.8)
$$\mathbf{v_j} = \varphi_0 \mathbf{v_{j-r}} + \dots + \varphi_{r-1} \mathbf{v_{j-1}} \quad \text{for} \quad j = r, \dots, k.$$

2.3. Solution of the truncated Hamburger moment problem.

Theorem 2.9. [CF91, Theorem 3.9] For $k \in \mathbb{N}$ and $\beta = (\beta_0, \ldots, \beta_{2k})$ with $\beta_0 > 0$, the following statements are equivalent:

- (1) There exists a representing measure for β supported on $K = \mathbb{R}$.
- (2) There exists a $(\operatorname{rank} \beta)$ -atomic representing measure for β .
- (3) β is positively recursively generated.
- (4) $A_{\beta} \succeq 0$ and rank $A_{\beta} = \operatorname{rank} \beta$.

A straightforward corollary of Theorem 2.9 and Corollary 2.7 is the following.

Corollary 2.10. Let $k \in \mathbb{N}$ and $\beta = (\beta_0, \dots, \beta_{2k})$ with $\beta_0 > 0$. Suppose that A_β is singular. The following statements are equivalent:

- (1) There exists a representing measure for β supported on $K = \mathbb{R}$.
- (2) There exists a $(\operatorname{rank} \beta)$ -atomic representing measure for β .
- (3) β is positively recursively generated.
- (4) $A_{\beta} \succeq 0$ and rank $A_{\beta} = \operatorname{rank} A_{\beta}(k-1)$.

2.4. Partially positive semidefinite matrices and their completions. A partial matrix $A = (a_{ij})_{i,j=1}^n$ is a matrix of real numbers $a_{ij} \in \mathbb{R}$, where some of the entries are not specified.

A partial symmetric matrix $A = (a_{ij})_{i,j=1}^n$ is **partially positive semidefinite (ppsd)** (resp. **partially positive definite (ppd)**) if the following two conditions hold:

- (1) a_{ij} is specified if and only if a_{ji} is specified and $a_{ij} = a_{ji}$.
- (2) All fully specified principal minors of A are psd (resp. pd).

It is well-known that a ppsd matrix A(x) of the form as in Lemma 2.11 below admits a psd completion. (This follows from the fact that the corresponding graph is chordal, see e.g. [GJSW84, Dan92, BW11].) In the notation of Lemma 2.11, if $A(x_0)$, $x_0 \in \mathbb{R}$, is a psd Hankel matrix, then Corollary 2.7 implies that (2.9) below holds. Since we will need an additional information about the rank of the completion $A(x_0)$ and the explicit interval of all possible x_0 for our results, we give a proof of Lemma 2.11 based on the use of generalized Schur complements assuming (2.9) holds.

Lemma 2.11. Let

$$A(x) := \begin{pmatrix} A_1 & a & b \\ a^T & \alpha & x \\ b^T & x & \beta \end{pmatrix} \in S_n$$

be a partially positive semidefinite symmetric matrix, where $A_1 \in S_{n-2}$, $a, b \in \mathbb{R}^{n-2}$, $\alpha, \beta \in \mathbb{R}$ and x is a variable. Let

$$A_2 := \begin{pmatrix} A_1 & a \\ a^T & \alpha \end{pmatrix} \in S_{n-1}, \qquad A_3 := \begin{pmatrix} A_1 & b \\ b^T & \beta \end{pmatrix} \in S_{n-1},$$

and

$$x_{\pm} := b^T A_1^+ a \pm \sqrt{(A_2/A_1)(A_3/A_1)} \in \mathbb{R}.$$

Suppose the following holds:

(2.9)
$$A_1 \text{ is invertible } or \operatorname{rank} A_1 = \operatorname{rank} A_2.$$

Then:

(1) $A(x_0)$ is positive semidefinite if and only if $x_0 \in [x_-, x_+]$.

(2)

$$\operatorname{rank} A(x_0) = \begin{cases} \max \{ \operatorname{rank} A_2, \operatorname{rank} A_3 \}, & \text{for } x_0 \in \{ x_-, x_+ \}, \\ \max \{ \operatorname{rank} A_2, \operatorname{rank} A_3 \} + 1, & \text{for } x_0 \in (x_-, x_+). \end{cases}$$

(3) If A(x) is partially positive definite, then A(x') is positive definite for $x' \in (x_-, x_+)$.

Proof. By Theorem 2.2, $A(x) \succeq 0$ if and only if

(2.10)
$$A_2 \succeq 0, \qquad \begin{pmatrix} b \\ x \end{pmatrix} \in \mathcal{C}(A_2) \quad \text{and} \quad f(x) := A(x)/A_2 \ge 0,$$

The first condition of (2.10) is true by the ppsd assumption.

Since $A_2 \succeq 0$, it follows by Theorem 2.2 that $a \in C(A_1)$ and hence by the properties of the Moore-Penrose inverse we have that $A_1(A_1^+a) = a$. Thus,

(2.11)
$$\mathcal{C}(A_2) = \mathcal{C}\left(\begin{pmatrix} A_1 & 0\\ a^T & \alpha - a^T A_1^+ a \end{pmatrix}\right) = \mathcal{C}\left(\begin{pmatrix} A_1 & 0\\ a^T & A_2/A_1 \end{pmatrix}\right).$$

Now we separate two cases according to A_2/A_1 .

Case 1: $A_2/A_1 > 0$.

(2.11) and the assumption of Case 1 imply that $C(A_2) = C(A_1 \oplus 1)$. Since $A_3 \succeq 0$, it follows by Theorem 2.2 that $b \in C(A_1)$. Therefore $\begin{pmatrix} b & x \end{pmatrix}^T \in C(A_1 \oplus 1)$ for every $x \in \mathbb{R}$. Thus the second condition of (2.10) is true for every $x \in \mathbb{R}$.

Note that the assumption of Case 1 and Proposition 2.3 imply that rank $A_2 > \operatorname{rank} A_1$ and hence the assumption (2.9) implies invertibility of A_1 and A_2 . By Proposition 2.4, used for A(x) as K, A_2 as M and A_1 as A, we have that

(2.12)
$$f(x) = A_3/A_1 - (A_2/A_1)^{-1}(x - b^T A_1^+ a)^2$$

Therefore $f(x_0) \ge 0$ if and only if $x_0 \in [x_-, x_+]$, which is the third condition of (2.10). Now by Proposition 2.3 we know that rank $A(x) > \operatorname{rank} A_2$ if and only if $f(x_0) > 0$, which establishes (1),(2) in the case $A_2/A_1 > 0$.

Case 2: $A_2/A_1 = 0$.

(2.11) and the assumption of Case 2 imply that

(2.13)
$$\mathcal{C}(A_2) = \mathcal{C}\left(\begin{pmatrix} A_1\\ a^T \end{pmatrix}\right).$$

Therefore, using (2.13), it is true that

(2.14)
$$\begin{pmatrix} b \\ x \end{pmatrix} \in \mathcal{C}(A_2) \Leftrightarrow \begin{pmatrix} b \\ x \end{pmatrix} = \begin{pmatrix} A_1 \\ a^T \end{pmatrix} z = \begin{pmatrix} A_1z \\ a^Tz \end{pmatrix}$$
 for some $z \in \mathbb{R}^{n-2}$.

Since $A_3 \succeq 0$, it follows by Theorem 2.2 that $b \in C(A_1)$ and hence by the properties of the Moore-Penrose inverse $\{A_1^+b + w : w \in \ker A_1\}$ is the set of all solutions z of the system $A_1z = b$. Therefore, using (2.14), it follows that

$$\begin{pmatrix} b \\ x \end{pmatrix} \in \mathcal{C}(A_2) \iff x \in \{a^T A_1^+ b + a^T w \colon w \in \ker A_1\} = \{a^T A_1^+ b\},\$$

where we used the fact that A_1 is symmetric, $a \in C(A_1)$ and $w \in \ker A_1$ for the last equality. So only $x_0 = a^T A_1^+ b$ satisfies the second condition of (2.10).

Now by definition of the generalized Schur complement, we have

$$f(x) = \beta - \begin{pmatrix} b^T & x \end{pmatrix} A_2^+ \begin{pmatrix} b \\ x \end{pmatrix}.$$

By the properties of the Moore-Penrose inverse

$$A_2^+\begin{pmatrix}b\\x_0\end{pmatrix}=\begin{pmatrix}A_1^+b\\0\end{pmatrix}+v$$
 for some $v\in \ker A_2$.

Hence,

$$f(x_0) = \beta - (b^T \ x_0) \left(\begin{pmatrix} A_1^+ b \\ 0 \end{pmatrix} + v \right) = \beta - b^T A_1^+ b = A_3 / A_1 \ge 0,$$

where the second equality follows from the fact that A_2 is symmetric, $\begin{pmatrix} b^T & x_0 \end{pmatrix}^T \in C(A_2)$ and $v \in \ker A_2$, and the last inequality follows by the ppsd assumption. Note that $x_0 = x_+ = x_-$ and by Proposition 2.3, rank $A(x_0) = \operatorname{rank} A_2$ if and only if $A_3/A_1 = 0$, in which case also rank $A_3 = \operatorname{rank} A_2$. Otherwise we have $f(x_0) = A_3/A_1 > 0$, which implies by Proposition 2.3 that rank $A(x_0) = \operatorname{rank} A_1 + 1$. Thus (1),(2) are true in the case $A_2/A_1 = 0$.

(3) follows from (2) by noticing that $A_2/A_1 > 0$, $A_3/A_1 > 0$ and rank $A_2 = \operatorname{rank} A_3 = n-1$.

2.5. Extension principle. The extension principle for psd matrices is the following.

Lemma 2.12. Let $A \in S_n$ be a positive semidefinite matrix, $Q \subseteq \{1, ..., n\}$ a subset and A_Q be the restriction of A to rows and columns from the set Q. If $v \in \ker A_Q$ is a nonzero vector from the kernel of A_Q , then the vector \hat{v} with the only nonzero entries in rows from Q and such that the restriction $\hat{v}|_Q$ to the rows from Q equals to v, belongs to ker A.

Proof. By permuting rows and columns we may assume that A is of the form $A = \begin{pmatrix} A_Q & B \\ B^T & C \end{pmatrix}$. We have to prove that

Since A is psd, for every $w := \begin{pmatrix} v^T & u^T \end{pmatrix} \in \mathbb{R}^n$ we have that

$$(2.16) 0 \le wAw^T = 2u^T B^T v + u^T C u.$$

If $B^T v \neq 0$, then we define $u := -\alpha B^T v$ where $\alpha > 0$ is an arbitrary positive real number, and plug into (2.16) to get

(2.17)
$$0 \le -2\alpha \left\| B^T v \right\|^2 + \alpha^2 v^T B C B^T v = \alpha (\alpha v^T B C B^T v - 2 \left\| B^T v \right\|^2) =: \alpha S(\alpha)$$

Since $\lim_{\alpha \to 0} S(\alpha) = -2 \|B^T v\|^2 < 0$, (2.17) cannot be true for α small enough. Hence $B^T v = 0$, which proves (2.15).

ALJAŽ ZALAR

2.6. Subsequences of one-dimensional moment sequences.

Proposition 2.13. Let $k \in \mathbb{N}$ and $\beta = (\beta_0, \ldots, \beta_{2k})$ with $\beta_0 > 0$ be a sequence which admits a representing measure supported on $K = \mathbb{R}$. Then for every $i, j \in \mathbb{N}$, where $0 \le i \le j \le k$, a subsequence $\beta^{(i,j)} := (\beta_{2i}, \ldots, \beta_{2j})$ also admits a representing measure supported on $K = \mathbb{R}$.

Proof. Note that A_{β} is of the form

$$A_{\beta} = \begin{pmatrix} A_{\beta^{(0,i-1)}} & * & * \\ * & A_{\beta^{(i,j)}} & * \\ * & * & A_{\beta^{(j+1,k)}} \end{pmatrix}.$$

By Theorem 2.9, $A_{\beta} \succeq 0$ and hence $A_{\beta^{(i,j)}} \succeq 0$. For i = j the statement is clear, i.e., the representing atom is β_{2i} with density 1. Assume that i < j. We separate two cases according to the invertibility of $A_{\beta^{(i,j)}}$.

- (1) If $A_{\beta^{(i,j)}} \succ 0$, then rank $A_{\beta^{(i,j)}} = \operatorname{rank} \beta^{(i,j)} = j i + 1$ and by Theorem 2.9, $\beta^{(i,j)}$ admits a measure.
- (2) Else

$$A_{\beta^{(i,j)}} = \left(\begin{array}{cc} A_{\beta^{(i,j-1)}} & v^T \\ v & \beta_{2j} \end{array}\right)$$

is singular, where $v = (\beta_j \cdots \beta_{2j-1})$. We separate two cases according to the invertibility of $A_{\beta^{(i,j-1)}}$.

- If $A_{\beta^{(i,j-1)}}$ is invertible, then rank $A_{\beta^{(i,j-1)}} = \operatorname{rank} A_{\beta^{(i,j)}}$.
- Else A_β(i,j-1) is singular and by Corollary 2.7 used for β^(i,j) as β, we get rank A_β(i,j-2) = rank A_β(i,j-1). This implies that the last column of A_β(i,j-1) is in the span of the other columns of A_β(i,j-1). By Lemma 2.12, the *j*-th column of A_β is in the span of the columns *i*+1,...,*j*-1. Since β is positively recursively generated, the (*j*+1)-th column of A_β is in the span of the columns *i*+2,...,*j* and in particular the last column of A_β(i,j). In both subcases of (2), rank A_β(i,j-1) = rank A_β(i,j) and Corollary 2.10 implies that β^(i,j) admits a measure.

3. Truncated Hamburger moment problem of degree 2k with gap (β_{2k-1}) and $(\beta_{2k-2}, \beta_{2k-1})$

In this section we solve the THMP of degree 2k with gaps (β_{2k-1}) (see Theorem 3.1) and $(\beta_{2k-2}, \beta_{2k-1})$ (see Theorem 3.5). As a corollary of Theorem 3.1 we obtain the solution to the TMP for the curve $y = x^3$ (see Corollary 3.3), while as a corollary of Theorem 3.5 we get the solution to the TMP for the curve $y = x^4$ and an additional moment $\beta_{3,2k-2}$ given (see Corollary 3.6).

3.1. Truncated Hamburger moment problem of degree 2k with gap (β_{2k-1}) .

Theorem 3.1. *Let* $k \in \mathbb{N}$ *and*

$$\beta(x) := (\beta_0, \beta_1, \dots, \beta_{2k-2}, x, \beta_{2k})$$

be a sequence where each β_i is a real number, $\beta_0 > 0$ and x is a variable. Let

$$\beta := (\beta_0, \dots, \beta_{2k-4})$$
 and $\beta := (\beta_0, \dots, \beta_{2k-2})$

be subsequences of $\beta(x)$, $v := (\beta_k \cdots \beta_{2k-2})$ a vector and

$$\widetilde{A} := \left(\begin{array}{cc} A_{\widehat{\beta}} & v^T \\ v & \beta_{2k} \end{array} \right)$$

a matrix. Then the following statements are equivalent:

- (1) There exists $x_0 \in \mathbb{R}$ and a representing measure for $\beta(x_0)$ supported on $K = \mathbb{R}$.
- (2) There exists $x_0 \in \mathbb{R}$ and a $(\operatorname{rank} \widetilde{\beta})$ -atomic representing measure for $\beta(x_0)$.
- (3) $A_{\beta(x)}$ is partially positive semidefinite and one of the following conditions is true: (a) k = 1.
 - (b) k > 1 and one of the following conditions is true:
 (i) A_β ≻ 0.
 (ii) rank A_β = rank A_β = rank Ã.

Proof. First we prove the implication (1) \Rightarrow (3). By Theorem 2.9, $A_{\beta(x_0)} \succeq 0$ and rank $A_{\beta(x_0)} =$ rank $\beta(x_0)$. $A_{\beta(x_0)} \succeq 0$ in particular implies that $A_{\beta(x)}$ is ppsd. If k = 1, then (3a) holds. Otherwise k > 1. If $A_{\beta} \succ 0$, then (3(b)i) holds. Else A_{β} is singular and hence

(3.1)
$$\operatorname{rank} A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} \beta(x_0) = A_{\beta(x_0)},$$

where the first two equalities follow by Corollary 2.7 used for $\beta(x_0)$ as β and the last by Theorem 2.9. $A_{\hat{\beta}}$ being a principal submatrix of \tilde{A} and \tilde{A} being a principal submatrix of

$$A_{\beta(x_0)} = \begin{pmatrix} A_{\widehat{\beta}} & u^T & v^T \\ u & \beta_{2k-2} & x_0 \\ v & x_0 & \beta_{2k} \end{pmatrix},$$

where $u = (\beta_{k-1} \cdots \beta_{2k-3})$, imply together with (3.1) that (3(b)ii) holds and concludes the proof of the implication (1) \Rightarrow (3).

Second we prove the implication (3) \Rightarrow (2). We separate two cases according to k.

- k = 1. We have that $A_{\beta(x)} = \begin{pmatrix} \beta_0 & x \\ x & \beta_2 \end{pmatrix}$. For $x_0 = \sqrt{\beta_0 \beta_2}$, $A_{\beta(x_0)}$ is of rank 1 and the second column is the multiple of the first. Hence, by Corollary 2.10, a 1-atomic measure exists, proving the implication (3) \Rightarrow (2) in this case.
- k > 1. Notice that $A_{\beta(x)}$ is of the same form as A(x) from Lemma 2.11, where $A_{\widehat{\beta}}$, $A_{\widetilde{\beta}}$, \widetilde{A} correspond to A_1 , A_2 , A_3 , respectively. Since both cases (3(b)i) and (3(b)ii) satisfy the assumption (2.9), it follows by Lemma 2.11 that there exists x_0 such that $A_{\beta(x_0)} \succeq 0$ and

(3.2)
$$\operatorname{rank} A_{\beta(x_0)} = \max\left\{\operatorname{rank} A_{\widetilde{\beta}}, \operatorname{rank} \widetilde{A}\right\}.$$

Since in the case (3(b)i), it holds that rank $\widetilde{A} \leq \operatorname{rank} A_{\widetilde{\beta}}$, while in the case (3(b)ii), rank $\widetilde{A} = \operatorname{rank} A_{\widetilde{\beta}}$, we obtain from (3.2) that rank $A_{\beta(x_0)} = \operatorname{rank} A_{\widetilde{\beta}}$. By Corollary 2.10, (rank $\widetilde{\beta}$)-representing measure for $\beta(x_0)$ exists, which proves (2).

The implication $(2) \Rightarrow (1)$ is trivial.

Example 3.2. For k = 9, let

$$\begin{split} \beta^{(1)}(x) &= (1,0,1,0,2,0,5,0,14,0,42,0,132,0,429,0,2000,x,338881), \\ \beta^{(2)}(x) &= \left(14,\frac{7}{2},\frac{79}{4},-\frac{67}{8},\frac{1055}{16},-\frac{1935}{32},\frac{18195}{64},-\frac{43115}{128},\frac{336151}{256},-\frac{926695}{512},\frac{6407195}{1024},-\frac{19736547}{2048}, \\ &\frac{124731423}{4096},-\frac{419176415}{8192},\frac{2469281827}{16384},-\frac{8894873563}{32768},\frac{49568350247}{65536},x,\frac{1006568996907}{262144}\right), \\ \beta^{(3)}(x) &= (8,0,78,0,1446,0,32838,0,794886,0,19651398,0,489352326,0,12216629958,0,305262005766, \\ \end{split}$$

x, 7630169896518).

Let $\widetilde{A}^{(i)}$, i = 1, 2, 3, denote \widetilde{A} from Theorem 3.1 corresponding to $\beta^{(i)}(x)$. Using *Mathematica* [Wol] one can check that:

- $\widetilde{A}^{(i)} \succeq 0$ for i = 1, 2, 3.
- $A_{\widetilde{\beta}^{(1)}} \succ 0, A_{\widetilde{\beta}^{(2)}} \succeq 0, A_{\widetilde{\beta}^{(3)}} \succeq 0 \text{ and } \dim \left(\ker A_{\widetilde{\beta}^{(3)}} \right) = 1.$

• rank
$$A_{\widehat{\beta}^{(3)}} = \operatorname{rank} \widehat{A}^{(3)} = \operatorname{rank} A_{\widetilde{\beta}^{(3)}} = 8.$$

Therefore:

- $A_{\beta^{(1)}(x)}$ is ppsd and $\tilde{\beta}^{(1)}$ satisfies (3(b)i) of Theorem 3.1, implying that a 9-atomic measure for $\beta^{(1)}(x)$ exists.
- $A_{\beta^{(2)}(x)}$ is not ppsd and by Theorem 3.1, there is no representing measure for $\beta^{(2)}(x)$.
- $A_{\beta^{(3)}(x)}$ is ppsd and $\tilde{\beta}^{(3)}$ satisfies (3(b)ii) of Theorem 3.1, implying that an 8-atomic measure for $\beta^{(3)}(x)$ exists.

The following corollary is a consequence of Theorem 3.1 and is an alternative solution of the bivariate TMP for the curve $y = x^3$, first solved by Fialkow in [Fia11].

Corollary 3.3. Let $k \in \mathbb{N}$ and $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}^2_+, i+j \leq 2k}$ be a 2-dimensional real multisequence of degree 2k. Suppose M(k) is positive semidefinite and recursively generated. Let

$$u^{(i)} := (\beta_{0,i}, \beta_{1,i}, \beta_{2,i}) \text{ for } i = 0, \dots, 2k - 2,$$

 $\widehat{\beta} := (u^{(0)}, \dots, u^{(2k-2)})$ and $\widetilde{\beta} := (u^{(0)}, \dots, u^{(2k-2)}, \beta_{0,2k-1}, \beta_{1,2k-1})$

be subsequences of β . Then β has a representing measure supported on $y = x^3$ if and only if the following statements hold:

- (1) One of the following holds:
 - If $k \ge 3$, then $Y = X^3$ is a column relation of M(k).
 - If k = 2, then the equalities $\beta_{0,1} = \beta_{3,0}$, $\beta_{1,1} = \beta_{4,0}$, $\beta_{0,2} = \beta_{3,1}$ hold.
- (2) One of the following holds:
 - (a) $A_{\widetilde{\beta}} \succ 0$.

(b) $A_{\widetilde{\beta}} \succeq 0$ and rank $A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} M(k)$.

Moreover, if the representing measure exists, then:

- If $A_{\tilde{\beta}}$ is nonsingular, there exists a (3k)-atomic measure.
- If A_{β} is singular, then the measure is $(\operatorname{rank} M(k))$ -atomic.

Proof. For $m \in \{0, 1, \dots, 6k - 2, 6k\}$ we define the numbers $\widetilde{\beta}_m$ by the following rule

$$\beta_m := \beta_m \pmod{3}, \lfloor \frac{m}{3} \rfloor.$$

Claim 1. Every number $\widetilde{\beta}_m$ is well-defined.

We have to prove that $m \pmod{3} + \lfloor \frac{m}{3} \rfloor \leq 2k$. We separate three cases according to m.

- $m \le 6k 4$: $\lfloor \frac{m}{3} \rfloor + m \pmod{3} \le (2k 2) + 2 = 2k$.
- $m \in \{6k 3, 6k 2\}$: $\lfloor \frac{m}{3} \rfloor + m \pmod{3} \le (2k 1) + 1 = 2k$. m = 6k: $\lfloor \frac{m}{3} \rfloor + m \pmod{3} = 2k + 0 = 2k$.

Claim 2. Let $t \in \mathbb{N}$. The atoms $(x_1, x_1^3), \ldots, (x_t, x_t^3)$ with densities $\lambda_1, \ldots, \lambda_t$ are the $(y - x^3)$ representing measure for β if and only if the atoms x_1, \ldots, x_t with densities $\lambda_1, \ldots, \lambda_t$ are the \mathbb{R} -representing measure for $\widetilde{\beta}(x) = (\widetilde{\beta}_0, \dots, \widetilde{\beta}_{2k-2}, x, \widetilde{\beta}_{2k}).$

The if part follows from the following calculation:

$$\widetilde{\beta}_m = \beta_m \pmod{3}, \lfloor \frac{m}{3} \rfloor = \sum_{\ell=1}^t \lambda_\ell x_\ell^m \pmod{3} x_\ell^{3\lfloor \frac{m}{3} \rfloor} = \sum_{\ell=1}^t \lambda_\ell x_\ell^m \pmod{3} + 3\lfloor \frac{m}{3} \rfloor = \sum_{\ell=1}^t \lambda_\ell x_\ell^m,$$

where $m = 0, 1, \dots, 6k - 2, 6k$.

The only if part follows from the following calculation:

$$\begin{aligned} \beta_{i,j} &= \beta_{i-3,j+1} = \dots = \beta_{i \pmod{3},j+\lfloor\frac{i}{3}\rfloor} \\ &= \widetilde{\beta}_{i \pmod{3}+3(j+\lfloor\frac{i}{3}\rfloor)} = \sum_{\ell=1}^t \lambda_\ell x_\ell^{i \pmod{3}+3(j+\lfloor\frac{i}{3}\rfloor)} = \sum_{\ell=1}^t \lambda_\ell x_\ell^{i \pmod{3}+3\lfloor\frac{i}{3}\rfloor} x_\ell^{3j} = \sum_{\ell=1}^t \lambda_\ell x_\ell^{i} (x_\ell^3)^j, \end{aligned}$$

where the equalities in the first line follow by M(k) being rg.

Using Claim 2 and a theorem of Bayer and Teichmann [BT06], implying that if a finite sequence has a K-representing measure, then it has a finitely atomic K-representing measure, the statement of the Corollary follows by Theorem 3.1. \square

- (1) Corollary 3.3 in case k = 1 is an improvement of [Fial1, Proposition 5.6.ii)] Remark 3.4. by decreasing the number of atoms from 6 to 3.
 - (2) For $M(1) \succ 0$ and $A_{\beta} \not\succeq 0$, (2) of Corollary 3.3 is not satisfied and hence the measure does not exist. Since this is the case under the assumptions of [Fial1, Proposition 5.6.iii)], the additional conditions in [Fia11, Proposition 5.6.iii)] are never satisfied.
 - (3) Examples in the Example 3.2 above are derived from [Fia11, Example 5.2], [Fia08, Example 4.18], [Fia08, Example 3.3], which demonstrate the solution of the moment problem for the curve $y = x^3$.

3.2. Truncated Hamburger moment problem of degree 2k with gaps $(\beta_{2k-2}, \beta_{2k-1})$.

Theorem 3.5. Let $k \in \mathbb{N}$, k > 1, and

$$\beta(x,y) := (\beta_0, \beta_1, \dots, \beta_{2k-3}, y, x, \beta_{2k})$$

be a sequence, where each β_i is a real number, $\beta_0 > 0$ and x, y are variables. Let

 $\widehat{\beta} := (\beta_0, \dots, \beta_{2k-6})$ and $\widetilde{\beta} := (\beta_0, \dots, \beta_{2k-4})$

be subsequences of $\beta(x, y)$,

$$u := \left(\begin{array}{ccc} \beta_k & \cdots & \beta_{2k-3} \end{array} \right), \quad s := \left(\begin{array}{ccc} \beta_{k-1} & \cdots & \beta_{2k-3} \end{array} \right) \quad and \quad w := \left(\begin{array}{ccc} \beta_{k-2} & \cdots & \beta_{2k-5} \end{array} \right)$$

vectors and

$$\widetilde{A} := \left(\begin{array}{cc} A_{\widehat{\beta}} & u^T \\ u & \beta_{2k} \end{array}\right)$$

a matrix. Then the following statements are equivalent:

- (1) There exist $x_0, y_0 \in \mathbb{R}$ and a representing measure for $\beta(x_0, y_0)$ supported on $K = \mathbb{R}$.
- (2) There exist $x_0, y_0 \in \mathbb{R}$ and a $(\operatorname{rank} \widetilde{\beta})$ or $(\operatorname{rank} \widetilde{\beta} + 1)$ -atomic representing measure for $\beta(x_0, y_0)$.
- (3) $A_{\beta(x,y)}$ is partially positive semidefinite and one of the following conditions holds: (a) k = 2 and $\frac{\beta_1^2}{\beta_0} \le \sqrt{\beta_0 \beta_4}$.

(b)
$$k > 2$$
, the inequality

(3.3)
$$sA_{\widetilde{\beta}}^{+}s^{T} \leq uA_{\widehat{\beta}}^{+}w^{T} + \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})}.$$

holds and one of the following conditions is true:

(i) $A_{\tilde{\beta}} \succ 0$.

(*ii*) rank
$$A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} (A_{\widetilde{\beta}} \ s^T) = \operatorname{rank} \widetilde{A}$$
.

Moreover, if the representing measure for β exists, then:

- If k = 2, then there is a 1-atomic measure if $\frac{\beta_1^2}{\beta_0} = \sqrt{\beta_0 \beta_4}$. Otherwise there is a 2-atomic measure.
- If k > 2, there exists a $(\operatorname{rank} \widetilde{\beta})$ -atomic if and only if one of the equalities

$$(3.4) \quad sA_{\widehat{\beta}}^{+}s^{T} = uA_{\widehat{\beta}}^{+}w^{T} - \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})} \quad or \quad sA_{\widetilde{\beta}}^{+}s^{T} = uA_{\widehat{\beta}}^{+}w^{T} + \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})}$$
$$holds.$$

Proof. Note that $\beta(x, y)$ admits a measure if and only if there exist $y_0 \in \mathbb{R}$ such that $\beta(x, y_0)$ admits a measure. Theorem 3.1 implies that the following claim holds.

Claim 1. $\beta(x, y_0)$ admits a measure if and only if the following conditions hold:

- (1) $A_{\beta(x,y_0)}$ is ppsd.
- (2) One of the following is true: (a) $A_{(\tilde{a}, q)} \succ 0$, where

$$A_{(\tilde{\beta},\beta_{2k-3},y)} = \begin{cases} \begin{pmatrix} \beta_0 & \beta_1 \\ \beta_1 & y \end{pmatrix}, & \text{if } k = 2, \\ \begin{pmatrix} A_{\tilde{\beta}} & s^T \\ s & y_0 \end{pmatrix} = \begin{pmatrix} A_{\hat{\beta}} & w^T & s_1^T \\ w & \beta_{2k-4} & \beta_{2k-3} \\ s_1 & \beta_{2k-3} & y_0 \end{pmatrix} & \text{where} \quad s_1^T = \begin{pmatrix} \beta_{k-1} \\ \vdots \\ \beta_{2k-4} \end{pmatrix}, \text{ otherwise}$$

(b) rank $A_{\tilde{\beta}} = \operatorname{rank} \widehat{A}(y_0)$, where

$$\widehat{A}(y) := \begin{cases} \begin{pmatrix} \beta_0 & y \\ y & \beta_4 \end{pmatrix}, & \text{if } k = 2, \\ \begin{pmatrix} A_{\widetilde{\beta}} & u(y)^T \\ u(y) & \beta_{2k} \end{pmatrix} = \begin{pmatrix} A_{\widehat{\beta}} & w^T & u^T \\ w & \beta_{2k-4} & y \\ u & y & \beta_{2k} \end{pmatrix} & \text{and} \quad u(y) := \begin{pmatrix} u & y \end{pmatrix}, \text{ otherwise}$$

Claim 2. Let k > 2. Assume $A_{\widehat{\beta}} \succ 0$ or rank $A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}}$. Then $\widehat{A}(y_0) \succeq 0$ if and only if (3.5) $\widehat{A}(y)$ is ppsd and $y_0 \in \left[uA_{\widehat{\beta}}^+ w^T - \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})}, uA_{\widehat{\beta}}^+ w^T + \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})} \right] =: [y_-, y_+].$

Moreover,

(3.6)
$$\operatorname{rank} \widehat{A}(y_0) = \begin{cases} \max \{ \operatorname{rank} A_{\widetilde{\beta}}, \operatorname{rank} \widetilde{A} \}, & \text{for } y_0 \in \{ y_-, y_+ \}, \\ \max \{ \operatorname{rank} A_{\widetilde{\beta}}, \operatorname{rank} \widetilde{A} \} + 1, & \text{for } y_0 \in (y_-, y_+). \end{cases}$$

The assumption (2.9) of Lemma 2.11 used for $\widehat{A}(y)$, $A_{\widehat{\beta}}$, $A_{\widetilde{\beta}}$, \widetilde{A} as A(x), A_1 , A_2 , A_3 , respectively, are by the assumption of Claim 2 satisfied and hence Claim 2 follows by Lemma 2.11.

Claim 3. Let k > 2. Assume $A_{\widehat{\beta}} \succ 0$ or rank $A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}}$. Then $A_{\beta(x,y_0)}$ is ppsd for some $y_0 \in \mathbb{R}$ if and only if $A_{\beta(x,y)}$ is ppsd, $s^T \in \mathcal{C}(A_{\widetilde{\beta}})$ and (3.3) holds.

Note that $A_{\beta(x,y_0)}$ is ppsd if and only if $A_{(\tilde{\beta},\beta_{2k-3},y_0)} \succeq 0$ and $\hat{A}(y_0) \succeq 0$. By Theorem 2.2, $A_{(\tilde{\beta},\beta_{2k-3},y_0)} \succeq 0$ if and only if

(3.7) $A_{\widetilde{\beta}} \succeq 0, \qquad s^T \in \mathcal{C}(A_{\widetilde{\beta}}) \qquad \text{and} \qquad A_{(\widetilde{\beta},\beta_{2k-3},y_0)}/A_{\widetilde{\beta}} = y_0 - sA_{\widetilde{\beta}}^+ s^T \ge 0,$

By Claim 2, $\hat{A}(y_0)$ is psd if and only if (3.5) holds. Now note that the first condition of (3.5) (which also includes the first condition of (3.7)) is equivalent to $A_{\beta(x,y)}$ being ppsd and that y_0 satisfying the third condition of (3.7) and the second condition of (3.5) exists if and only if (3.3) holds. This proves Claim 3.

First we prove the implication (1) \Rightarrow (3). By Claim 1, in particular $A_{\beta(x,y_0)}$ is ppsd.

If k = 2, then $A_{(\tilde{\beta},\beta_1,y)} \succeq 0$, which implies that $y_0 \ge \frac{\beta_1^2}{\beta_0}$, and $\hat{A}(y_0) \succeq 0$, which implies that $y_0 \le \sqrt{\beta_0 \beta_4}$. Hence, $\frac{\beta_1^2}{\beta_0} \le \sqrt{\beta_0 \beta_4}$, which is (3a). Since $A_{\beta(x,y_0)}$ being ppsd implies that also $A_{\beta(x,y)}$ is ppsd, this proves the implication (1) \Rightarrow (3) in this case.

It remains to prove (1) \Rightarrow (3) in the case k > 2. We separate two cases according to the invertibility of $A_{\tilde{\beta}}$.

- A_β ≻ 0: Using Claim 3, A_{β(x,y)} is ppsd, (3.3) and (3(b)i) holds, which proves the implication (1) ⇒ (3) in this case.
- $A_{\widetilde{\beta}} \not\succeq 0$: It follows that $A_{(\widetilde{\beta},\beta_{2k-3},y_0)} \not\succeq 0$ and hence (2b) of Claim 1 must hold. Corollary 2.7 used for $(\widetilde{\beta},\beta_{2k-3},y_0)$ as β implies that

(3.8)
$$\operatorname{rank} A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}}.$$

By Proposition 2.13, $(\tilde{\beta}, \beta_{2k-3}, y_0)$ also admits a measure and Corollary 2.10 used for $(\tilde{\beta}, \beta_{2k-3}, y_0)$ as β implies that

(3.9)
$$\operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} A_{(\widetilde{\beta},\beta_{2k-3},y_0)}$$

(2b) of Claim 1 together with (3.8) implies that all the inequalities in the estimate rank $A_{\hat{\beta}} \leq \operatorname{rank} \widetilde{A} \leq \operatorname{rank} \widehat{A}(y_0)$ are equalities and in particular,

(3.10)
$$\operatorname{rank} A_{\widehat{\beta}} = \operatorname{rank} A$$

(3.8), (3.9), (3.10) and Claim 3 imply that $A_{\beta(x,y)}$ is ppsd, (3.3) and (3(b)ii) holds, which proves the implication (1) \Rightarrow (3) in this case.

Second we prove the implication $(3) \Rightarrow (1)$. We separate two cases according to k.

If k = 2, then we are in the case (3a). For $y_0 = \sqrt{\beta_0 \beta_4}$, $\beta(x, y_0)$ is ppsd and satisfies (2a) of Claim 1 if $\frac{\beta_1^2}{\beta_0} < \sqrt{\beta_0 \beta_4}$ and (2b) if $\frac{\beta_1^2}{\beta_0} = \sqrt{\beta_0 \beta_4}$. In both cases Claim 1 implies the implication (3) \Rightarrow (1) is true in this case.

Else k > 2. If (3(b)i) holds, then in particular $A_{\widehat{\beta}} > 0$. Otherwise (3(b)ii) holds and in particular rank $A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}}$. In both cases the assumptions of Claims 2 and 3 are fulfilled. By Claim 3, the matrix $A_{\beta(x,y_+)}$ is ppsd and by (3.6) of Claim 2, rank $\widehat{A}(y_+) = \max\{\operatorname{rank} A_{\widetilde{\beta}}, \operatorname{rank} \widetilde{A}\}$. If (3(b)i) holds, then rank $\widehat{A}(y_+) = \operatorname{rank} A_{\widetilde{\beta}} = k - 1$. Else (3(b)ii) holds and rank $\widehat{A}(y_+) = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} \widetilde{A}$. In both cases, $\beta(x, y_+)$ satisfies (1) and (2b) of Claim 1 above and thus the measure exists which proves the implication (3) \Rightarrow (1).

The implication $(2) \Rightarrow (1)$ is trivial.

Now we prove the implication (1) \Rightarrow (2). If $\beta(x, y_0)$ has a representing measure, then:

- By Theorem 3.1 it has a $(rank(\tilde{\beta}, \beta_{2k-3}, y_0))$ -atomic representing measure.
- By Proposition 2.13, β̃ and (β̃, β_{2k-3}, y₀) also have measures and hence by Theorem 2.9, rank A_{β̃} = rank β̃ and rank(β̃, β_{2k-3}, y₀) = rank A_(β̃,β_{2k-3},y₀).

Since rank $A_{(\tilde{\beta},\beta_{2k-3},y_0)} \in \{\operatorname{rank} A_{\tilde{\beta}}, \operatorname{rank} A_{\tilde{\beta}} + 1\}$, the implication (1) \Rightarrow (2) is true.

It remains to prove the moreover part. We separate two cases according to k.

- If k = 2, then rank A_{β̃} = rank(β₀) = 1. So 1-atomic measure exists if and only if rank A_(β₀,β₁,y₀) = rank Â(y₀) = 1 for some y₀. But from the form of A_{(β₀,β₁,y⟩} and Â(y) this is possible only if y₀ = β₁/β₀ = √β₀β₄. Otherwise there is a 2-atomic measure.
 Else k > 2. By Proposition 2.3 and (3.7) above, rank A_(β̃,β₂k−3,y₀) = rank A_{β̃} if and only x̃
- Else k > 2. By Proposition 2.3 and (3.7) above, rank A_(β̃,β_{2k-3},y₀) = rank A_{β̃} if and only if y₀ = sA⁺_{β̃}s^T. In the proof of the implication (3) ⇒ (1) we see that rank A_{β̃} ≥ rank Ã. Using this in (3.6) above, it follows that sA⁺_{β̃}s^T must be equal to y₋ or y₊, which is exactly (3.4).

This concludes the proof of the theorem.

The following corollary is a consequence of Theorem 3.5 and solves the bivariate TMP for the curve $y = x^4$ where also $\beta_{3,2k-2}$ is given.

Corollary 3.6. Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}^2_+, i+j \leq 2k}$ be a 2-dimensional real multisequence of degree 2k and let $\beta_{3,2k-2}$ be also given. Suppose M(k) is positive semidefinite and recursively generated. Let

$$u^{(i)} = (\beta_{0,i}, \beta_{1,i}, \beta_{2,i}, \beta_{3,i}) \text{ for } i = 0, \dots, 2k - 1,$$

 $\widehat{\beta} := (u^{(0)}, \dots, u^{(2k-3)}, \beta_{0,2k-2}, \beta_{1,2k-2}, \beta_{2,2k-2}) \quad and \quad \widetilde{\beta} := (\widehat{\beta}, \beta_{3,2k-2}, \beta_{0,2k-1})$

be subsequences of β ,

$$u := \begin{pmatrix} u^{(k)} & \cdots & u^{(2k-1)} & \beta_{1,2k-1} \end{pmatrix}, \quad s := \begin{pmatrix} \beta_{3,k-1} & u^{(k)} & \cdots & u^{(2k-1)} & \beta_{1,2k-1} \end{pmatrix},$$
$$w := \begin{pmatrix} \beta_{2,k-1} & \beta_{3,k-1} & u^{(k)} & \cdots & u^{(2k-2)} & \beta_{1,2k-2} & \beta_{2,2k-2} & \beta_{3,2k-2} \end{pmatrix}$$

vectors and

$$\widetilde{A} := \left(\begin{array}{cc} A_{\widehat{\beta}} & u^T \\ u & \beta_{0,2k} \end{array} \right)$$

a matrix. Then β has a representing measure supported on $y = x^4$ if and only if

$$sA_{\widetilde{\beta}}^+s^T \le uA_{\widehat{\beta}}^+w^T + \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})}.$$

one of the following statements hold:

(1) One of the following holds:

- If $k \ge 4$, then $Y = X^4$ is a column relation of M(k).
- If k = 3, then the equalities $\beta_{0,1} = \beta_{4,0}$, $\beta_{1,1} = \beta_{5,0}$, $\beta_{2,1} = \beta_{6,0}$ hold.
- If k = 2, then the equality $\beta_{0,1} = \beta_{4,0}$ holds.
- k = 1.

(2) One of the following conditions holds:

(a) $A_{\tilde{\beta}} \succ 0$.

(b)
$$A_{\widetilde{\beta}} \succeq 0$$
 and rank $A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} \begin{pmatrix} A_{\widetilde{\beta}} & s^T \end{pmatrix} = \operatorname{rank} \widetilde{A}$.

Moreover, if the representing measure exists, then there is a $(\operatorname{rank} \widetilde{\beta})$ -atomic measure if

$$sA_{\widetilde{\beta}}^{+}s^{T} \in \left\{ uA_{\widehat{\beta}}^{+}w^{T} - \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})}, uA_{\widehat{\beta}}^{+}w^{T} + \sqrt{(A_{\widetilde{\beta}}/A_{\widehat{\beta}})(\widetilde{A}/A_{\widehat{\beta}})} \right\}.$$

and $(\operatorname{rank} \widetilde{\beta} + 1)$ -atomic otherwise.

Proof. For $m \in \{0, 1, ..., 8k - 3, 8k\}$ we define the numbers $\widetilde{\beta}_m$ by the following rule

$$\beta_m := \beta_m \pmod{4}, \lfloor \frac{m}{4} \rfloor.$$

Claim 1. Every number $\widetilde{\beta}_m$ is well-defined.

We will prove that $m \pmod{4} + \lfloor \frac{m}{4} \rfloor \leq 2k$ if $m \neq 8k - 5$, while for m = 8k - 5 we have $\widetilde{\beta}_{8k-5} = \beta_{3,2k-2}$. We separate three cases according to m.

- m < 8k 8: $\lfloor \frac{m}{4} \rfloor + m \pmod{4} \le (2k 3) + 3 = 2k$.
- $m \in \{8k 8, 8k 7, 8k 6\}: \lfloor \frac{m}{4} \rfloor + m \pmod{4} \le (2k 2) + 2 = 2k.$
- $m \in \{8k 4, 8k 3\}$: $\lfloor \frac{m}{4} \rfloor + m \pmod{4} \le (2k 1) + 1 = 2k$.
- m = 8k: $\left|\frac{m}{4}\right| + m \pmod{3} = 2k + 0 = 2k$.

Claim 2. Let $t \in \mathbb{N}$. The atoms $(x_1, x_1^4), \ldots, (x_t, x_t^4)$ with densities $\lambda_1, \ldots, \lambda_t$ are the $(y - x^4)$ -representing measure for β and $\beta_{3,2k-2}$ if and only if the atoms x_1, \ldots, x_t with densities $\lambda_1, \ldots, \lambda_t$ are the \mathbb{R} -representing measure for $\tilde{\beta}(x, y) = (\tilde{\beta}_0, \ldots, \tilde{\beta}_{2k-2}, y, x, \tilde{\beta}_{2k})$.

The if part follows from the following calculation:

$$\widetilde{\beta}_m = \beta_m \pmod{4}, \lfloor \frac{m}{4} \rfloor = \sum_{\ell=1}^t \lambda_\ell x_\ell^m \pmod{4} x_\ell^{4 \lfloor \frac{m}{4} \rfloor} = \sum_{\ell=1}^t \lambda_\ell x_\ell^m \pmod{4} + 4 \lfloor \frac{m}{4} \rfloor = \sum_{\ell=1}^t \lambda_\ell x_\ell^m,$$

where m = 0, ..., 8k - 3, 8k.

The only if part follows from the following calculation for $i + j \le 2k$:

$$\beta_{i,j} = \beta_{i-4,j+1} = \dots = \beta_i \pmod{4}, j+\lfloor \frac{i}{4} \rfloor$$
$$= \widetilde{\beta}_i \pmod{4} + 4(j+\lfloor \frac{i}{4} \rfloor) = \sum_{\ell=1}^t \lambda_\ell x_\ell^i \pmod{4} + 4(j+\lfloor \frac{i}{4} \rfloor) = \sum_{\ell=1}^t \lambda_\ell x_\ell^i \pmod{4} + 4\lfloor \frac{i}{4} \rfloor x_\ell^{4j} = \sum_{\ell=1}^t \lambda_\ell x_\ell^i (x_\ell^4)^j$$

where the equalities in the first line follow by M(k) being rg, and

$$\beta_{3,2k-2} = \widetilde{\beta}_{8k-5} = \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{8k-5} = \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{3} (x_{\ell}^{4})^{2k-2}.$$

Using Claim 2 and a theorem of Bayer and Teichmann [BT06], implying that if a finite sequence has a K-representing measure, then it has a finitely atomic K-representing measure, the statement of the Corollary follows by Theorem 3.5.

4. TRUNCATED HAMBURGER MOMENT PROBLEM OF DEGREE 2k WITH GAP(S) (β_1), (β_1 , β_2)

In this section we solve the THMP of degree 2k with gaps (β_1) (see Theorem 4.1) and (β_1, β_2) (see Theorem 4.5). As a corollary of Theorem 4.1 we obtain the solution to the TMP for the curve $y^2 = x^3$ (see Corollary 4.4), while as a corollary of Theorem 4.5 we get the solution to the TMP for the curve $y^3 = x^4$ and an additional moment $\beta_{\frac{5}{2},0}$ given (see Corollary 4.7).

4.1. Truncated Hamburger moment problem of degree 2k with gaps (β_1) .

Theorem 4.1. Let $k \in \mathbb{N}$, k > 1, and

$$\beta(x) := (\beta_0, x, \beta_2, \dots, \beta_{2k})$$

be a sequence where each β_i is a real number, $\beta_0 > 0$ and x is a variable. Let

$$\widehat{\beta} := (\beta_2, \dots, \beta_{2k-2}), \quad \widetilde{\beta} := (\beta_2, \dots, \beta_{2k}), \quad \overline{\beta} := (\beta_4, \dots, \beta_{2k-2}) \quad and \quad \widecheck{\beta} := (\beta_4, \dots, \beta_{2k})$$

be subsequences of $\beta(x)$,

$$v := (\beta_2 \cdots \beta_{k-1}) \text{ and } u := (\beta_2 \cdots \beta_k)$$

vectors, and

$$\widetilde{A} := \begin{pmatrix} \beta_0 & v \\ v^T & A_{\overline{\beta}} \end{pmatrix} \quad and \quad \widehat{A} := \begin{pmatrix} \beta_0 & u \\ u^T & A_{\overline{\beta}} \end{pmatrix}$$

matrices. Then the following statements are equivalent:

- (1) There exists $x_0 \in \mathbb{R}$ and a representing measure for $\beta(x_0)$ supported on $K = \mathbb{R}$.
- (2) There exists $x_0 \in \mathbb{R}$ and a $(\operatorname{rank} \widetilde{\beta})$ or a $(\operatorname{rank} \widetilde{\beta} + 1)$ -atomic representing measure for $\beta(x_0)$.
- (3) $A_{\beta(x)}$ is partially positive semidefinite and one of the following conditions is true: (a) (i) k = 2 and $A_{\tilde{\beta}} \succ 0$.
 - (*ii*) k > 2, $A_{\widetilde{\beta}} \succ 0$ and $\widetilde{A} \succ 0$. (*b*) rank $A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} A_{\widetilde{\beta}}$.

Moreover, if the representing measure exists, then there does not exist a $(\operatorname{rank} \widetilde{\beta})$ -atomic measure if and only if (3b) holds and $\operatorname{rank} A_{\widehat{\beta}} < \operatorname{rank} \widehat{A}$.

Proof. First we prove the implication (1) \Rightarrow (3). By Theorem 2.9, $A_{\beta(x_0)} \succeq 0$ and rank $A_{\beta(x_0)} = \operatorname{rank} \beta(x_0)$. The condition $A_{\beta(x_0)} \succeq 0$ implies that $A_{\beta(x)}$ is ppsd. We separate two cases according to the invertibility of A_{β} .

A_β ≻ 0: Since A_β is a principal submatrix of A_{β(x0)}, we conclude that rank A_{β(x0)} ≥ rank A_β = k, and hence A_{β(x0)} is either invertible or rank A_{β(x0)} is singular and by Corollary 2.10 used for β(x0) as β, rank A_{β(x0)} = rank A_(β0,x0,β). In both cases

$$A_{(\beta_0,x_0,\widehat{\beta})} = \begin{cases} \begin{pmatrix} \beta_0 & x_0 \\ x_0 & \beta_2 \end{pmatrix}, & \text{if } k = 2, \\ \begin{pmatrix} \beta_0 & x_0 & v \\ x_0 & \beta_2 & v_1 \\ v^T & v_1^T & A_{\overline{\beta}} \end{pmatrix} & \text{where} \quad v_1 = \begin{pmatrix} \beta_3 & \cdots & \beta_k \end{pmatrix}, & \text{if } k > 2, \end{cases}$$

is invertible. If k > 2, \widetilde{A} is a principal submatrix of $A_{(\beta_0, x_0, \widehat{\beta})}$ and it follows that $\widetilde{A} \succ 0$. Hence, (3a) holds. Together with $A_{\beta(x)}$ being ppsd, proves the implication (1) \Rightarrow (3) in this case.

• $A_{\tilde{\beta}}$ is singular: Since $\tilde{\beta}$ is a subsequence of $\beta(x_0)$ of the form from Proposition 2.13 with i = 1, j = k, it admits a measure. By Corollary 2.10 used for $\tilde{\beta}$ as β , it follows that

(4.1)
$$\operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} A_{\widehat{\beta}}.$$

By Corollary 2.8 used for $\beta(x_0)$ as β , it follows that

(4.2)
$$\operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} A_{\widetilde{\beta}}$$

Hence, (4.1) and (4.2) imply that (3b) holds. Together with $A_{\beta(x)}$ being ppsd, proves the implication (1) \Rightarrow (3) in this case.

Second we prove the implication (3) \Rightarrow (2). Let $P_1 : \mathbb{R}^{k+1} \rightarrow \mathbb{R}^{k+1}$ be the following permutation matrix

$$P_1 = \begin{pmatrix} \mathbf{0} & 0 & 1 \\ \mathbf{0} & 1 & 0 \\ I_{k-1} & 0 & 0 \end{pmatrix},$$

where **0** stands for the row of k - 2 zeros and I_{k-1} is the identity matrix of size k - 1. Then $P_1^T A_{\beta(x)} P_1$ is of the form

$$P_1^T A_{\beta(x)} P_1 = \begin{pmatrix} A_{\breve{\beta}} & w^T & u^T \\ w & \beta_2 & x \\ u & x & \beta_0 \end{pmatrix},$$

where $w = (\beta_3 \cdots \beta_{k+1})$ is a vector.

Claim. $A_{\beta(x_0)}$ is psd if and only if

$$x_0 \in \left[uA^+_{\breve{\beta}} w^T - \sqrt{(A_{\widetilde{\beta}}/A_{\breve{\beta}})(\widehat{A}/A_{\breve{\beta}})}, uA^+_{\breve{\beta}} w^T + \sqrt{(A_{\widetilde{\beta}}/A_{\breve{\beta}})(\widehat{A}/A_{\breve{\beta}})} \right] =: [x_-, x_+].$$

Moreover,

(4.3)
$$\operatorname{rank} A_{\beta(x_0)} := \begin{cases} \max\{\operatorname{rank} A_{\widetilde{\beta}}, \operatorname{rank} \widehat{A}\}, & \text{if } x_0 \in \{x_-, x_+\}, \\ \max\{\operatorname{rank} A_{\widetilde{\beta}}, \operatorname{rank} \widehat{A}\} + 1, & \text{if } x_0 \in (x_-, x_+). \end{cases}$$

Denoting the matrices

$$\mathcal{A} := \begin{pmatrix} A_{\breve{\beta}} & w^T \\ w & \beta_2 \end{pmatrix} \text{ and } \mathcal{B} := \begin{pmatrix} A_{\breve{\beta}} & u^T \\ u & \beta_0 \end{pmatrix},$$

and the permutation matrix $P_2 : \mathbb{R}^k \to \mathbb{R}^k$ by

$$P_2 = \left(\begin{array}{cc} \mathbf{0} & 1\\ I_{k-1} & 0 \end{array}\right),$$

where 0 stands for the row of k - 1 zeroes and I_{k-1} the identity matrix of size k - 1, we have that

$$\mathcal{A} = P_2^T A_{\widetilde{\beta}} P_2$$
 and $\mathcal{B} = P_2^T A P_2$

In particular,

(4.4)
$$\operatorname{rank} \mathcal{A} = \operatorname{rank} A_{\widetilde{\beta}} \quad \text{and} \quad \operatorname{rank} \mathcal{B} = \operatorname{rank} A.$$

If (3a) holds, then $A_{\tilde{\beta}} \succ 0$ implies that $A_{\check{\beta}} \succ 0$. If (3b) holds, then in particular rank $A_{\check{\beta}} = \operatorname{rank} A_{\tilde{\beta}} = \operatorname{rank} A$. Hence, the assumption (2.9) of Lemma 2.11 used for $P_1^T A_{\beta(x)} P_1, A_{\check{\beta}}, \mathcal{A}, \mathcal{B}$ as $A(x), A_1, A_2, A_3$, respectively, is satisfied and using also $\mathcal{A}/A_{\check{\beta}} = A_{\tilde{\beta}}/A_{\check{\beta}}$ and $\mathcal{B}/A_{\check{\beta}} = \widehat{A}/A_{\check{\beta}}$, Claim follows.

First assume that (3a) holds. We separate two cases according to the inverbility of A.

- ≻ 0: From A_{β̃} ≻ 0 and ≻ 0 it follows, using Proposition 2.3, that A_{β̃}/A_{β̃} > 0 and Â/A_{β̃} > 0. Hence by the definition of x_±, we have x_− < x₊ and by Claim, A_{β(x₀)} ≻ 0 for x₀ ∈ (x_−, x₊). By Theorem 2.9, (rank β(x₀)) = (rank β̃+1)-atomic representing measure for β(x₀) exists, which proves the implication (3) ⇒ (2) in this case.
- is singular: From A_β ≻ 0 it follows that A_β ≻ 0. Since is singular, Proposition 2.3 implies that Â/A_β = 0, and hence by the definition of x_±, we have x₋ = x₊. By Claim, A_{β(x±)} ≥ 0 with rank A_{β(x±)} = rank A_{β̃}. We separate two cases according to k.
 - k = 2: Since $\widehat{A} = \begin{pmatrix} \beta_0 & \beta_2 \\ \beta_2 & \beta_4 \end{pmatrix}$ and $\beta_0 > 0$, it follows that the second (also the last)

column of \widehat{A} is in the span of the first (also the others) one.

- k > 2: By assumptions $\widetilde{A} \succ 0$ and $\widehat{A} = \begin{pmatrix} \widetilde{A} & u_1^T \\ u_1 & \beta_{2k} \end{pmatrix}$ being singular, where the u_1 is equal to $u_1 = \begin{pmatrix} \beta_k & \beta_{k+2} & \cdots & \beta_{2k-1} \end{pmatrix}$, it follows that the last column of \widehat{A} is in the span of the others.

By Lemma 2.12, the last column of $A_{\beta(x_{\pm})}$ is also in the span of the others and by Corollary 2.10, we have that $(\operatorname{rank} \beta(x_{\pm})) = (\operatorname{rank} \tilde{\beta})$ -atomic representing measure for $\beta(x_{\pm})$ exists, which proves the implication (3) \Rightarrow (2) in this case.

Otherwise (3b) holds. Proposition 2.3 implies that $A_{\beta}/A_{\beta} = 0$, and hence by the definition of x_{\pm} , we have $x_{-} = x_{+}$. By Claim, $A_{\beta(x_{\pm})} \succeq 0$. The assumption rank $A_{\beta} = \operatorname{rank} A_{\beta}$, also implies that the last column of $A_{\beta} = \begin{pmatrix} A_{\beta} & u_{2}^{T} \\ u_{2} & \beta_{2k} \end{pmatrix}$, where $u_{2} = \begin{pmatrix} \beta_{k} & \cdots & \beta_{2k-1} \end{pmatrix}$, is in the span of the others. By Lemma 2.12, the last column of $A_{\beta(x_{\pm})}$ is in the span of the others. Hence, by Corollary 2.10, $(\operatorname{rank} \beta(x_{\pm}))$ -atomic measure for $\beta(x_{\pm})$ exists. Since β is a subsequence of $\beta(x_{0})$ of the form from Proposition 2.13 with i = 1, j = k, it admits a measure and hence Theorem 2.9 implies that $\operatorname{rank} A_{\beta} = \operatorname{rank} \beta$. From (4.3), it follows that:

- If rank $\widehat{A} \leq \operatorname{rank} A_{\widetilde{\beta}}$, then rank $\beta(x_{\pm}) = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} \widetilde{\beta}$.
- Else rank $\widehat{A} = \operatorname{rank} A_{\widetilde{\beta}} + 1$ and rank $\beta(x_{\pm}) = \operatorname{rank} \widehat{A} = \operatorname{rank} \widetilde{\beta} + 1$.

This proves the implication $(3) \Rightarrow (2)$ in this case.

The implication $(2) \Rightarrow (1)$ is trivial.

It remains to prove the moreover part. Observe that in the proof of the implication (3) \Rightarrow (2), $(\operatorname{rank} \widetilde{\beta})$ -atomic measure might not exist if (3a) holds with $\widehat{A} \succ 0$ and does not exist if (3b) holds with $\operatorname{rank} A_{\widetilde{\beta}} < \operatorname{rank} \widehat{A}$. We will prove that in the first case there always exists a $(\operatorname{rank} \widetilde{\beta})$ -atomic measure. Assume that $A_{\widetilde{\beta}} \succ 0$ and $\widehat{A} \succ 0$. We will prove that one of $A_{\beta(x_{\pm})}$ or $A_{\beta(x_{\pm})}$ satisfies

(4.5)
$$\operatorname{rank} A_{\beta(x_{\pm})} = \operatorname{rank} A_{\beta(x_{\pm})}(k-1),$$

and hence by Corollary 2.10, a $(\operatorname{rank} \beta(x_{\pm})) = (\operatorname{rank} \widetilde{\beta})$ -atomic measure exists. Using Proposition 2.4 with for $A_{\beta(x)}, A_{\widetilde{\beta}}, A_{\widehat{\beta}}$ as K, N, C, respectively, and denoting $u := A_{\widetilde{\beta}}/A_{\widehat{\beta}}$, we have that

$$f(x) := A_{\beta(x)} / A_{\widetilde{\beta}} = \left(\beta_0 - e(x)A_{\widehat{\beta}}^{-1}e(x)^T\right) - \frac{1}{u}\left(\beta_k - e(x)A_{\widehat{\beta}}^{-1}z^T\right)^2 =: g(x) - \frac{1}{u}h(x)^2,$$

where $e(x) := (x \ \beta_2 \ \cdots \ \beta_{k-1})$ and $z := (\beta_{k+1} \ \cdots \ \beta_{2k-1})$. From the proof of the implication (3) \Rightarrow (2), we know that $x_- < x_+$ and

(4.6)
$$f(x_{-}) = f(x_{+}) = 0.$$

Note that $g(x) = A_{(\beta_0, x, \widehat{\beta})} / A_{\widehat{\beta}}$. If

(4.7)
$$g(x_{-}) = g(x_{+}) = 0$$

then $h(x_{-}) = h(x_{+}) = 0$. But h(x) is a linear function in x, so this is possible only if h(x) = 0 for every $x \in \mathbb{R}$. This is possible only if

(4.8)
$$A_{\widehat{\beta}}^{-1}z^T = \begin{pmatrix} 0 & b_2 & \cdots & b_{k-1} \end{pmatrix}^T$$
 for some $b_2, \dots, b_k \in \mathbb{R}$ and $\beta_k = \sum_{i=2}^{k-1} \beta_i b_i$.

We write $(A_{\beta(x)})|_{S_1,S_2}$ for the restriction of $A_{\beta(x)}$ to rows from S_1 and columns from S_2 . Since $A_{\beta(x)}$ is a Hankel matrix, we have

$$(A_{\beta(x)})_{\{1,\dots,X^{k-1}\},\{X,\dots,X^k\}} = (A_{\beta(x)})_{\{X,\dots,X^k\},\{1,\dots,X^{k-1}\}}$$

which is equal to

$$\left(\begin{array}{cc} e(x) & \beta_k \\ A_{\widehat{\beta}} & z^T \end{array}\right) = \left(\begin{array}{cc} e(x)^T & A_{\widehat{\beta}} \\ \beta_k & z \end{array}\right).$$

(4.8) implies that the last column of $(A_{\beta(x)})_{\{1,...,X^{k-1}\},\{X,...,X^k\}}$ is in the span of the columns $2, \ldots, k-1$. From $(A_{\beta(x)})_{\{X,...,X^k\},\{1,...,X^{k-1}\}}$ this in particular implies that the last column of $A_{\widehat{\beta}}$ is in the span of the others and $A_{\widehat{\beta}}$ is singular, which is a contradiction with the assumption $A_{\widetilde{\beta}} \succ 0$. Therefore (4.7) cannot be true and one of $g(x_-)$ and $g(x_+)$ is positive. By Proposition 2.3, this means that $A_{(\beta_0,x_+,\widehat{\beta})} \succ 0$ or $A_{(\beta_0,x_-,\widehat{\beta})} \succ 0$ and hence rank $A_{(\beta_0,x_-,\widehat{\beta})} = k$ or rank $A_{(\beta_0,x_+,\widehat{\beta})} = k$. By Proposition 2.3 and (4.6), rank $A_{\beta(x_-)} = \operatorname{rank} A_{\beta(x_+)} = \operatorname{rank} A_{\beta(x_+)} = \operatorname{rank} A_{\beta(x_-)} = A_{\beta(x_\pm)}(k-1)$, it follows that one of x_{\pm} satisfies (4.5). This concludes the proof of the moreover part.

Remark 4.2. For k = 1, the THMP with gaps (β_1) coincides with the THMP with gaps (β_{2k-1}) and hence the case k = 1 is already covered by Theorem 3.1.

Example 4.3. For k = 9, let

$$\begin{split} \beta^{(1)}(x) &= \left(1, x, 11, 0, \frac{979}{5}, 0, 4103, 0, \frac{462979}{5}, 0, 2174855, 0, \frac{261453379}{5}, 0, 1275350087, 0, \frac{156925970179}{5}, 0, 776760884999\right), \\ \beta^{(2)}(x) &= \left(1, x, \frac{15}{2}, 0, \frac{177}{2}, 0, \frac{2445}{2}, 0, \frac{36177}{2}, 0, \frac{554325}{2}, 0, \frac{8656377}{2}, 0, \frac{136617405}{2}, 0, \frac{2169039777}{2}, 0, \frac{138214318741}{8}\right), \\ \beta^{(3)}(x) &= \left(1, x, \frac{15}{2}, 0, \frac{177}{2}, 0, \frac{2445}{2}, 0, \frac{36177}{2}, 0, \frac{554325}{2}, 0, \frac{8656377}{2}, 0, \frac{136617405}{2}, 0, \frac{2169039777}{2}, 0, \frac{34553579685}{2}\right), \\ \beta^{(4)}(x) &= \frac{1}{9}(9, x, 133, -235, 3157, -7987, 86893, -281995, 2598757, -10096867, 82154653, -362972155, 2699153557, -13062280147, 91112865613, -470199300715, 3134918735557, -16926788453827 \end{split}$$

Let $\widetilde{A}^{(i)}$ and $\widehat{A}^{(i)}$, i = 1, 2, 3, denote \widetilde{A} , \widehat{A} , respectively, from Theorem 4.1 corresponding to $\beta^{(i)}(x)$. Using *Mathematica* [Wol] one can check that:

- $\widehat{A}^{(1)} \succ 0$, while for i = 2, 3, 4 it holds that $\widehat{A}^{(i)} \succeq 0$ and dim $\left(\ker \widehat{A}^{(i)} \right) = 1$.
- For i = 1, 4 we have $\widetilde{A}^{(i)} \succ 0$ for i = 1, 4, while for i = 2, 3 it holds that $\widetilde{A}^{(i)} \succeq 0$ and $\dim (\ker \widetilde{A}^{(i)}) = 1$.
- $A_{\widetilde{\beta}^{(i)}} \succ 0$ for $i = 1, 2, 4, A_{\widetilde{\beta}^{(3)}} \succeq 0$ and dim $\left(\ker A_{\widetilde{\beta}^{(3)}}\right) = 1$.
- $A_{\widehat{\beta}^{(3)}} \succ 0$ and $A_{\widecheck{\beta}^{(3)}} \succ 0$.

Therefore:

- $A_{\beta^{(1)}(x)}$ is ppsd and (3a) of Theorem 4.1 is true, implying that a 9-atomic measure for $\beta^{(1)}(x)$ exists.
- $\beta^{(2)}(x)$ does not satisfy (3a) neither (3b) of Theorem 4.1, implying there is no representing measure for $\beta^{(2)}(x)$.
- $A_{\beta^{(3)}(x)}$ is ppsd and $\tilde{\beta}^{(3)}$ satisfies (3b) of Theorem 4.1 together with rank $A_{\hat{\beta}^{(3)}} = \operatorname{rank} \hat{A}^{(3)}$, implying that an 8-atomic measure for $\beta^{(3)}(x)$ exists.
- $A_{\beta^{(4)}(x)}$ is ppsd and $\tilde{\beta}^{(4)}$ satisfies (3a) of Theorem 4.1, implying that a 9-atomic measure for $\beta^{(4)}(x)$ exists.

The following corollary is a consequence of Theorem 4.1 and gives the solution of the bivariate TMP for the curve $y^2 = x^3$.

Corollary 4.4. Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}^2_+, i+j \leq 2k}$ be a 2-dimensional real multisequence of degree 2k. Suppose M(k) is positive semidefinite and recursively generated. Let

$$u^{(i)} := (\beta_{1,i}, \beta_{0,i+1}, \beta_{2,i}) \quad for \ i = 0, \dots, 2k - 2,$$
$$\widehat{\beta} := (u^{(0)}, \dots, u^{(2k-2)}), \quad \widetilde{\beta} := (\widehat{\beta}, \beta_{1,2k-1}, \beta_{0,2k}), \quad \overline{\beta} := (\beta_{2,0}, u^{(1)}, \dots, u^{(2k-2)})$$
$$and \quad \widecheck{\beta} := (\overline{\beta}, \beta_{1,2k-1}, \beta_{0,2k})$$

be subsequences of β ,

$$v := (u^{(0)} \cdots u^{(k-2)} \beta_{1,k-1})$$

a vector and

$$\widetilde{A} := \left(\begin{array}{cc} \beta_0 & v \\ v^T & A_{\overline{\beta}} \end{array}\right)$$

a matrix. Then β has a representing measure supported on $y^2 = x^3$ if and only if the following statements hold:

(1) One of the following holds:

- If k > 3, then $Y^2 = X^3$ is a column relation of M(k).
- If k = 2, then the equalities $\beta_{0,2} = \beta_{3,0}$, $\beta_{1,2} = \beta_{4,0}$, $\beta_{0,3} = \beta_{3,1}$ hold.
- k = 1.

(2) One of the following holds:

(a) $A_{\widetilde{\beta}} \succ 0$ and $\widetilde{A} \succ 0$.

(b) $A_{\widetilde{\beta}} \succeq 0$ and rank $A_{\widehat{\beta}} = \operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} A_{\widetilde{\beta}}$.

Moreover, if the representing measure exists, then there exists a $(\operatorname{rank} \widetilde{\beta})$ -atomic measure if (2a) is true or (2b) holds with rank $A_{\widehat{\beta}} = \operatorname{rank} M(k)$. Otherwise there is a $(\operatorname{rank} \widetilde{\beta} + 1)$ -atomic measure.

Proof. For $m \in \{0, 2..., 6k\}$ we define the numbers $\widetilde{\beta}_m$ by the following rule

$$\widetilde{\beta}_m := \begin{cases} \beta_{0,\frac{m}{3}}, & \text{if } m \pmod{3} = 0, \\ \beta_{2,\lfloor\frac{m}{3}\rfloor - 1}, & \text{if } m \pmod{3} = 1, \\ \beta_{1,\lfloor\frac{m}{3}\rfloor}, & \text{if } m \pmod{3} = 2. \end{cases}$$

Claim 1. Every number $\widetilde{\beta}_m$ is well-defined.

We have to prove that $i + j \leq 2k$, where i, j are indices of $\beta_{i,j}$ used in the definition of β_m . We separate three cases according to *m*:

- $m \pmod{3} = 0: \frac{m}{3} \le 2k.$ $m \pmod{3} = 1: 2 + (\lfloor \frac{m}{3} \rfloor 1) \le 2 + (2k 2) = 2k.$ $m \pmod{3} = 2: 1 + \lfloor \frac{m}{3} \rfloor \le 1 + (2k 1) = 2k.$

Claim 2. Let $t \in \mathbb{N}$. The atoms $(x_1^2, x_1^3), \ldots, (x_t^2, x_t^3)$ with densities $\lambda_1, \ldots, \lambda_t$ are the $(y^2 - x^3)$ representing measure for β if and only if the atoms x_1, \ldots, x_t with densities $\lambda_1, \ldots, \lambda_t$ are the \mathbb{R} -representing measure for $\widetilde{\beta}(x) = (\widetilde{\beta}_0, x, \widetilde{\beta}_2, \dots, \widetilde{\beta}_{2k}).$

The if part follows from the following calculation:

$$\widetilde{\beta}_{m} = \begin{cases} \beta_{0,\frac{m}{3}}, & \text{if } m \pmod{3} = 0, \\ \beta_{2,\lfloor\frac{m}{3}\rfloor-1}, & \text{if } m \pmod{3} = 1, \\ \beta_{1,\lfloor\frac{m}{3}\rfloor}, & \text{if } m \pmod{3} = 2, \end{cases} = \begin{cases} \sum_{\ell=1}^{t} \lambda_{\ell} (x_{\ell}^{3})^{\frac{m}{3}}, & \text{if } m \pmod{3} = 0, \\ \sum_{\ell=1}^{t} \lambda_{\ell} (x_{\ell}^{2})^{2} (x_{\ell}^{3})^{\lfloor\frac{m}{3}\rfloor-1}, & \text{if } m \pmod{3} = 1, \\ \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{2} (x_{\ell}^{3})^{\lfloor\frac{m}{3}\rfloor}, & \text{if } m \pmod{3} = 2, \end{cases} = \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{m},$$

where m = 0, 2, ..., 6k.

The only if part follows from the following calculation:

$$\begin{aligned} \beta_{i,j} &= \beta_{i-3,j+2} = \dots = \beta_i \pmod{3}_{j+2\lfloor \frac{i}{3} \rfloor} = \beta_{2(i \pmod{3})+3(j+2\lfloor \frac{i}{3} \rfloor)} \\ &= \sum_{\ell=1}^t \lambda_\ell x_\ell^{2(i \pmod{3})+3(j+2\lfloor \frac{i}{3} \rfloor)} = \sum_{\ell=1}^t \lambda_\ell x_\ell^{2(i \pmod{3})+3\lfloor \frac{i}{3} \rfloor} x_\ell^{3j} = \sum_{\ell=1}^t \lambda_\ell (x_\ell^2)^i (x_\ell^3)^j, \end{aligned}$$

where the first three equalities in the first line follow by M(k) being rg.

Using Claim 2 and a theorem of Bayer and Teichmann [BT06], implying that if a finite sequence has a K-representing measure, then it has a finitely atomic K-representing measure, the statement of the Corollary follows by Theorem 4.1.

4.2. Truncated Hamburger moment problem of degree 2k with gaps (β_1, β_2) .

Theorem 4.5. Let $k \in \mathbb{N}$, k > 2, and

$$\beta(x,y) := (\beta_0, x, y, \beta_3, \dots, \beta_{2k})$$

be a sequence, where each β_i is a real number, $\beta_0 > 0$ and x, y are variables. Let

$$\widetilde{\beta} := (\beta_4, \dots, \beta_{2k-2}), \ \overline{\beta} := (\beta_4, \dots, \beta_{2k}), \ \breve{\beta} := (\beta_6, \dots, \beta_{2k-2}), \ \overline{\beta} := (\beta_6, \dots, \beta_{2k})$$

be subsequences of $\beta(x, y)$,

$$v := \left(\begin{array}{ccc} \beta_3 & \dots & \beta_{k-1}\end{array}\right), \quad u := \left(\begin{array}{ccc} \beta_3 & \cdots & \beta_k\end{array}\right), \quad s := \left(\begin{array}{ccc} \beta_3 & \cdots & \beta_{k+1}\end{array}\right),$$
$$w := \left(\begin{array}{ccc} \beta_5 & \cdots & \beta_{k+2}\end{array}\right),$$

vectors, and

$$\overline{A} := \begin{pmatrix} \beta_0 & v \\ v^T & A_{\breve{\beta}} \end{pmatrix} \quad and \quad \widetilde{A} := \begin{pmatrix} \beta_0 & u \\ u^T & A_{\overline{\overline{\beta}}} \end{pmatrix}$$

matrices. Then the following statements are equivalent:

- (1) There exist $x_0, y_0 \in \mathbb{R}$ and a representing measure for $\beta(x_0, y_0)$ supported on $K = \mathbb{R}$.
- (2) There exist $x_0, y_0 \in \mathbb{R}$ and a $(\operatorname{rank} \overline{\beta})$ or $(\operatorname{rank} \overline{\beta} + 1)$ -atomic representing measure for $\beta(x_0, y_0)$.
- (3) $A_{\beta(x,y)}$ is partially positive semidefinite,

(4.9)
$$sA_{\overline{\beta}}^{+}s^{T} \leq uA_{\overline{\beta}}^{+}w^{T} + \sqrt{(A_{\overline{\beta}}/A_{\overline{\beta}})(\widetilde{A}/A_{\overline{\beta}})}$$

and one of the following statements is true:

- (a) $A_{\overline{\beta}} \succ 0$ and one of the following holds:
 - (i) (A) k = 3 and the inequality in (4.9) is strict.
 - (B) k > 3, $\overline{A} \succ 0$ and the inequality in (4.9) is strict.
 - (ii) The following inequalities holds:

$$uA_{\widetilde{\beta}}^+ u^T < sA_{\overline{\beta}}^+ s^T \quad and \quad uA_{\overline{\beta}}^+ w^T - \sqrt{(A_{\overline{\beta}}/A_{\overline{\beta}})(\widetilde{A}/A_{\overline{\beta}})} \leq sA_{\overline{\beta}}^+ s^T.$$

(b) $\operatorname{rank} A_{\widetilde{\beta}} = \operatorname{rank} A_{\overline{\beta}} = \operatorname{rank} \left(\begin{array}{cc} s^T & A_{\overline{\beta}} \end{array} \right).$

Moreover, if the representing measure exists, then there is a $(\operatorname{rank} \overline{\beta})$ -atomic if and only if (3(a)ii) or (3b) holds.

Proof. Note that $\beta(x, y)$ admits a measure if and only if there exist $y_0 \in \mathbb{R}$ such that $\beta(x, y_0)$ admits a measure. Theorem 4.1 implies the following claim holds.

Claim 1. $\beta(x, y_0)$ admits a measure if and only if the following conditions hold:

- (1) $A_{\beta(x,y_0)}$ is ppsd.
- (2) Denoting

$$\widetilde{\beta}(y_0) := (y_0, \beta_3, \dots, \beta_{2k-2}) \quad \text{and} \quad \overline{\beta}(y_0) := (y_0, \beta_3, \dots, \beta_{2k}),$$

one of the following is true:

$$(a) \ A_{\overline{\beta}(y_0)} = \begin{pmatrix} y_0 & s \\ s^T & A_{\overline{\beta}} \end{pmatrix} \succ 0 \text{ and } \overline{A}(y_0) \succ 0, \text{ where} \\ \\ \left\{ \begin{array}{c|c} \left(\begin{array}{c|c} \beta_0 & y & \beta_3 \\ \hline y & A_{\overline{\beta}} \end{array} \right), & \text{if } k = 3, \\ \left(\begin{array}{c|c} \beta_0 & v(y) \\ v(y)^T & A_{\overline{\beta}} \end{array} \right) = \begin{pmatrix} \beta_0 & y & v \\ y & \beta_4 & w_1 \\ v^T & w_1^T & A_{\overline{\beta}} \end{array} \end{pmatrix}, & v(y)^T = \begin{pmatrix} y \\ v \end{pmatrix} \text{ and } w_1^T = \begin{pmatrix} \beta_5 \\ \vdots \\ \beta_{k+2} \end{array} \end{pmatrix}, \text{ otherwise} \\ (b) \ \operatorname{rank} A_{\overline{\beta}(y_0)} = \operatorname{rank} A_{\overline{\beta}(y_0)} = \operatorname{rank} A_{\overline{\beta}}. \end{cases}$$

We denote by

$$\widetilde{A}(y) := \begin{pmatrix} \beta_0 & u(y) \\ u(y)^T & A_{\overline{\beta}} \end{pmatrix} \text{ where } u(y) = \begin{pmatrix} y & u \end{pmatrix}.$$

Claim 2. Assume $A_{\overline{\beta}} \succ 0$ or rank $A_{\overline{\beta}} = \operatorname{rank} A_{\overline{\beta}}$. Then $\widetilde{A}(y_0) \succeq 0$ if and only if (4.10) $\widetilde{A}(y)$ is ppsd and $y_0 \in \left[uA_{\overline{\beta}}^+ w^T - \sqrt{(A_{\overline{\beta}}/A_{\overline{\beta}})}(\widetilde{A}/A_{\overline{\beta}}), uA_{\overline{\beta}}^+ w^T + \sqrt{(A_{\overline{\beta}}/A_{\overline{\beta}})}(\widetilde{A}/A_{\overline{\beta}}) \right] =: [y_-, y_+].$ Moreover,

(4.11)
$$\operatorname{rank} \widetilde{A}(y_0) = \begin{cases} \max\{\operatorname{rank} A_{\overline{\beta}}, \operatorname{rank} \widetilde{A}\}, & y \in \{y_-, y_+\}, \\ \max\{\operatorname{rank} A_{\overline{\beta}}, \operatorname{rank} \widetilde{A}\} + 1, & y \in (y_-, y_+). \end{cases}$$

Let P_2 be the permutation matrix as in the proof of Theorem 4.1. We have that $P_2^T \widetilde{A}(y) P_2$ is of the form

(4.12)
$$P_2^T \widetilde{A}(y) P_2 = \begin{pmatrix} A_{\overline{\beta}} & w^T & u^T \\ w & \beta_4 & y \\ u & y & \beta_0 \end{pmatrix},$$

and denoting the matrices

$$\mathcal{A} := \begin{pmatrix} A_{\overline{\beta}} & w^T \\ w & \beta_4 \end{pmatrix} \text{ and } \mathcal{B} := \begin{pmatrix} A_{\overline{\beta}} & u^T \\ u & \beta_0 \end{pmatrix},$$

and the permuation matrix $P_3: \mathbb{R}^{k-1} \to \mathbb{R}^{k-1}$ by

$$P_3 = \left(\begin{array}{cc} \mathbf{0} & 1\\ I_{k-2} & 0 \end{array}\right),$$

where 0 stands for the row of k - 2 zeros and I_{k-2} is the identity matrix of size k - 2, we have that

(4.13)
$$\mathcal{A} = P_3^T A_{\overline{\beta}} P_3 \quad \text{and} \quad \mathcal{B} = P_3^T \widetilde{A} P_3.$$

By the assumptions in Claim 2 and (4.13), $A_{\overline{\beta}} \succ 0$ or rank $A_{\overline{\beta}} = \operatorname{rank} \mathcal{A}$. Hence, the assumption (2.9) of Lemma 2.11 used for $P_2^T \widetilde{A}(y) P_2$, $A_{\overline{\beta}}$, \mathcal{A}, \mathcal{B} as $A(x), A_1, A_2, A_3$, respectively, is satisfied and using also $\mathcal{A}/A_{\overline{\beta}} = A_{\overline{\beta}}/A_{\overline{\beta}}$, $\mathcal{B}/A_{\overline{\beta}} = \widetilde{A}/A_{\overline{\beta}}$, Claim 2 follows.

Theorem 2.2 implies the following claim.

Claim 3. It is true that:

(1) $A_{\overline{\beta}(y_0)} \succeq 0$ if and only if

(4.14) $A_{\overline{\beta}} \succeq 0, \quad s^T \in \mathcal{C}(A_{\overline{\beta}}) \quad \text{and} \quad A_{\overline{\beta}(y_0)}/A_{\overline{\beta}} = y_0 - sA_{\overline{\beta}}^+ s^T \ge 0.$

(2) $A_{\widetilde{\beta}(y_0)} \succeq 0$ if and only if

(4.15) $A_{\widetilde{\beta}} \succeq 0, \quad u^T \in \mathcal{C}(A_{\widetilde{\beta}}) \quad \text{and} \quad A_{\widetilde{\beta}(y_0)}/A_{\widetilde{\beta}} = y_0 - uA_{\widetilde{\beta}}^+ u^T \ge 0.$

Claim 4. Assume $A_{\overline{\beta}} \succ 0$ or rank $A_{\overline{\beta}} = \operatorname{rank} A_{\overline{\beta}}$. Then $A_{\beta(x,y_0)}$ is ppsd for some $y_0 \in \mathbb{R}$ if and only if $A_{\beta(x,y)}$ is ppsd, $s^T \in \mathcal{C}(A_{\overline{\beta}})$ and (4.9) holds.

Note that $A_{\beta(x,y_0)}$ is ppsd if and only if $A_{\overline{\beta}(y_0)} \succeq 0$ and $\widetilde{A}(y_0) \succeq 0$. The first condition of (4.10) (which also includes the first condition of (4.14)) is equivalent to $A_{\beta(x,y)}$ being ppsd. Further on, y_0 satisfying the third condition of (4.14) and the second condition of (4.10) exists if and only if (4.9) holds. This proves Claim 4.

First we prove the implication (1) \Rightarrow (3). By Claim 1, in particular $A_{\beta(x,y_0)}$ (and hence also $A_{\beta(x,y)}$) is ppsd. Since $\tilde{\beta}(y_0)$ also admits a measure by Proposition 2.13, we either have $A_{\tilde{\beta}(y_0)} \succ 0$ and in particular $A_{\overline{\beta}} \succ 0$, or $A_{\tilde{\beta}(y_0)}$ is singular and it follows by Corollary 2.8 that $A_{\overline{\beta}} \succ 0$ or rank $A_{\overline{\beta}} = \operatorname{rank} A_{\overline{\beta}}$.

If (2a) of Claim 1 holds, then in particular $A_{\overline{\beta}} \succ 0$ and if k > 3 also $\overline{A} \succ 0$. Since $A_{\overline{\beta}}(y_0) \succ 0$, it follows using Proposition 2.3 that $A_{\overline{\beta}}(y_0)/A_{\overline{\beta}} > 0$ or equivalently $y_0 > sA_{\overline{\beta}^+}s^T$. Since by Claim 2, $y_0 \in [y_-, y_+]$, this implies that $sA_{\overline{\beta}^+}s^T < y_+$ which means that the inequality in (4.9) is strict. Hence, $A_{\beta(x,y)}$ is ppsd, $A_{\overline{\beta}} \succ 0$ and (3(a)i) holds. This proves the implication (1) \Rightarrow (3) in this case.

Assume now that (2b) of Claim 1 holds. There are two cases to consider:

A_β > 0: It follows that rank A_{β̃(y0)} = rank A_{β̃(y0)} = k − 1, which implies that:
 A_{β̃(y0)} > 0 since A_{β̃(y0)} is of size k − 1.

- By Proposition 2.3,
$$y_0 = sA_{\overline{\beta}}^+ s^T$$
 since $A_{\overline{\beta}(y_0)} = \begin{pmatrix} y_0 & s \\ s^T & A_{\overline{\beta}} \end{pmatrix}$ is singular.
- $k - 1 \leq \operatorname{rank} A_{\beta(x_0, y_0)} \leq k$ for some $x_0 \in \mathbb{R}$ such that $A_{\beta(x_0, y_0)} \succeq 0$, since

$$A_{\beta(x_0,y_0)} = \begin{pmatrix} \beta_0 & u(x_0,y_0) \\ u(x_0,y_0)^T & A_{\overline{\beta}(y_0)} \end{pmatrix} \quad \text{where} \quad u(x_0,y_0) = \begin{pmatrix} x_0 & y_0 & u \end{pmatrix}.$$

From $A_{\widetilde{\beta}(y_0)} \succ 0$ and $y_0 = sA_{\overline{\beta}}^+ s^T$, it follows by Proposition 2.3 and (4.15) that $uA_{\widetilde{\beta}}^+ u^T < sA_{\overline{\beta}}^+ s^T$. Further on, $y_- \leq sA_{\overline{\beta}}^+ s^T$ since by Claim 2, $\widetilde{A}(y_0) \succeq 0$ implies that $y_0 \in [y_-, y_+]$. Hence, $A_{\beta(x,y)}$ is ppsd, $A_{\overline{\beta}} \succ 0$ and (3(a)ii) holds. This proves the implication (1) \Rightarrow (3) in this case.

A_β ≯ 0: By Lemma 2.13, β also admits a measure and hence by Corollary 2.10 used for β as β, rank A_β = rank A_β. Together with the second condition in (4.14), this implies that (3b) holds. Since A_{β(x,y)} is ppsd and (4.9) holds, this proves the implication (1) ⇒ (3) in this case.

Second we prove the implication (3) \Rightarrow (1). If (3a) holds, then $A_{\overline{\beta}} \succ 0$ and in particular $A_{\overline{\beta}} \succ 0$. Else (3b) holds and in particular $A_{\overline{\beta}}$ is singular. By Claim 3, $A_{\overline{\beta}(sA_{\overline{\beta}}^+s^T)} \succeq 0$ and hence by Corollary 2.8 used for $\overline{\beta}(sA_{\overline{\beta}}^+s^T)$ as β we conclude that rank $A_{\overline{\beta}} = \operatorname{rank} A_{\overline{\beta}}$. Hence the assumption of Claims 2 and 4 is satisfied and $A_{\beta(x,y_0)}$ is ppsd for every y_0 from the interval $[\max\{y_-, sA_{\overline{\beta}}^+s^T\}, y_+]$. We separate cases three cases according to the assumptions:

- Case (3(a)i): We separate two cases according to the invertibility of \widetilde{A} .
 - à ≻ 0: Since A_{β̃} ≻ 0 and à ≻ 0, it follows that A_{β̃}/A_{β̃} > 0 and Ã/A_{β̃} > 0. By the form of y_± given in Claim 2, we have that y_− < y₊. Since by assumption also the inequality (4.9) is strict, the interval (max{y_−, sA⁺_βs^T}, y₊) is not empty and hence for every y₀ ∈ (max{y_−, sA⁺_βs^T}, y₊), A_{β(x,y₀)} satisfies (2a) above by Claims 2 and 3. This proves the implication (3) ⇒ (1) in this case.
 - A in singular: First we show that the last column of \tilde{A} is in the span of others. We separate two cases according to k.
 - * k = 3: Since $\widetilde{A} = \begin{pmatrix} \beta_0 & \beta_3 \\ \beta_3 & \beta_6 \end{pmatrix}$ and $\beta_0 > 0$, it follows that the second (also the last) solumn of \widetilde{A} is a multiple of the first (also it the energy of the others)
 - last) column of \widetilde{A} is a multiple of the first (also it the span of the others). * k > 3: Since $\overline{A} \succ 0$, the last column of $\widetilde{A} = \begin{pmatrix} \overline{A} & r^T \\ r & \beta_{2k} \end{pmatrix}$ is in the span of the others, where $r = \begin{pmatrix} \beta_k & \beta_{k+2} & \cdots & \beta_{2k-1} \end{pmatrix}$.

Since $A_{\overline{\beta}} \succ 0$, it follows that $A_{\overline{\beta}} \succ 0$ and $\widetilde{A}/A_{\overline{\beta}} = 0$. By the form of y_{\pm} given in Claim 2, we have that $y_{-} = y_{+}$. By Claim 2, $\widetilde{A}(y_{+}) = \begin{pmatrix} \overline{A}(y_{+}) & r_{1}^{T} \\ r_{1} & \beta_{2k} \end{pmatrix} \succeq 0$, where $r_{1} = \begin{pmatrix} \beta_{k} & \cdots & \beta_{2k-1} \end{pmatrix}$, and rank $\widetilde{A}(y_{+}) = \operatorname{rank} A_{\overline{\beta}} = k - 1$. By Lemma 2.12, the last column of $\widetilde{A}(y_{+})$ is in the span of the others and hence $\overline{A}(y_{+}) \succ 0$. Since by assumption also the inequality (4.9) is strict, $A_{\overline{\beta}(y_{+})} \succ 0$ by Claim 3. Hence, (2a) of Claim 1 holds for $y_{0} = y_{+}$, which proves the implication (3) \Rightarrow (1) in this case. • Case (3(a)ii): $\beta(x, sA_{\overline{\beta}} s^{T})$ is ppsd. Since

$$A_{\overline{\beta}(sA_{\overline{\beta}}^{+}s^{T})} = \begin{pmatrix} A_{\widetilde{\beta}(sA_{\overline{\beta}}^{+}s^{T})} & r_{2}^{T} \\ r_{2} & \beta_{\beta_{2k}} \end{pmatrix}$$

where $r_2 = (\beta_{k+1} \cdots \beta_{2k-1})$, is singular, the assumption $uA_{\tilde{\beta}}^+ u^T < sA_{\tilde{\beta}}^+ s^T$ and Claim 3 imply that $A_{\tilde{\beta}(sA_{\tilde{\beta}}^+ s^T)} \succ 0$, hence rank $A_{\tilde{\beta}(sA_{\tilde{\beta}}^+ s^T)} = \operatorname{rank} A_{\overline{\beta}(sA_{\tilde{\beta}}^+ s^T)} = \operatorname{rank} A_{\overline{\beta}}$. Hence, (2b) of Claim 1 for $y_0 = sA_{\tilde{\beta}}^+ s^T$ holds, which proves the implication (3) \Rightarrow (1) in this case.

Case (3b): By assumption rank A_{β̃} = rank A_{β̃}, it follows that the last column of A_{β̃} is in the span of the others. There exists x₀ ∈ ℝ such that A_{β(x₀,y₊)} is psd and by Lemma 2.12, the last column of A_{β̃(y+}) is in the span of the others and hence rank A_{β̃(y+}) = rank A_{β̃(y+}). Since A_{β̃(y+}) is singular, using Corollary 2.8 with β equal to β(x₀, y₊), we get rank A_{β̃(y+}) = rank A_{β̃}, which in particular implies that y₊ = sA⁺_{β̃}s^T. Hence, rank A_{β̃(y+}) = rank A_{β̃(y+}) = rank A_{β̃}, which is (2b) of Claim 1. This proves the implication (3) ⇒ (1) in this case.

It remains to prove the implication (1) \Rightarrow (2). By Theorem 4.1, if $\beta(x, y_0)$ has a representing measure, then there is a $(\operatorname{rank} \overline{\beta}(y_0))$ or $(\operatorname{rank} \overline{\beta}(y_0) + 1)$ -atomic representing measure. By Corollary 2.8, $\operatorname{rank} \overline{\beta}(y_0) = \operatorname{rank} A_{\overline{\beta}(y_0)} = \operatorname{rank} A_{\overline{\beta}(y_0)}$ is singular and $\operatorname{rank} \overline{\beta}(y_0) = \operatorname{rank} A_{\overline{\beta}} + 1 = \operatorname{rank} \overline{\beta} + 1$ otherwise.

For the moreover part, note from the previous paragraph that $(\operatorname{rank} \overline{\beta})$ -atomic measure exists if and only if $A_{\overline{\beta}(y_0)} = \operatorname{rank} A_{\overline{\beta}}$ for some y_0 such the $\beta(x, y_0)$ admits a measure. The only $y_0 \in \mathbb{R}$ satisfying $\operatorname{rank} A_{\overline{\beta}(y_0)} = \operatorname{rank} A_{\overline{\beta}}$ is $sA_{\overline{\beta}}^+s^T$ and hence a $(\operatorname{rank} \overline{\beta})$ -atomic measure exists if and only if $\beta(x, sA_{\overline{\beta}}^+s^T)$ admits a measure. From the proof of the implication (3) \Rightarrow (1) we see that this is true in the cases (3(a)ii) and (3b). Finally, if (3(a)i) holds, then we see that:

- If $\widetilde{A} \succ 0$, then we must have $y_{-} \leq sA_{\overline{\beta}}^{+}s^{T}$ and $uA_{\overline{\beta}}^{+}u^{T} < sA_{\overline{\beta}}^{+}s^{T}$ (see the proof of (3(a)ii)), which means that (3(a)ii) holds.
- If \widetilde{A} is singular, then $sA_{\overline{\beta}}^+s^T < y_- = y_+$ and $\beta(x, sA_{\overline{\beta}}^+s^T)$ does not admit a $(\operatorname{rank} \overline{\beta})$ -atomic measure.

This establishes the proof of the moreover part.

Remark 4.6. For k = 2, the THMP with gaps (β_1, β_2) coincides with the THMP with gaps $(\beta_{2k-2}, \beta_{2k-1})$ and hence the case k = 2 is already covered by Theorem 3.5.

The following corollary is a consequence of Theorem 4.5 and solves the bivariate TMP for the curve $y^3 = x^4$ where also $\beta_{\frac{5}{3},0}$ is given. Here $\beta_{\frac{5}{3},0}$ stands for the integral of $x^{\frac{5}{3}}$ w.r.t. μ , i.e., $\int_{K} x^{\frac{5}{3}} d\mu$.

Corollary 4.7. Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}^2_+, i+j \leq 2k} \beta$ be a 2-dimensional real multisequence of degree 2k and let $\beta_{\frac{5}{3},0}$ be also given. Suppose M(k) is positive semidefinite and recursively generated. Let

$$u^{(1)} = (\beta_{0,1}, \beta_{\frac{5}{3},0}, \beta_{2,0}, \beta_{1,2}), \quad u^{(i)} = (\beta_{0,i}, \beta_{3,i-2}, \beta_{2,i-1}, \beta_{1,i}) \quad \text{for} \quad i = 2, \dots, 2k-1,$$

$$\widetilde{\beta} := (u^{(1)}, \dots, u^{(2k-2)}, \beta_{0,2k-1}, \beta_{3,2k-3}, \beta_{2,2k-2}), \quad \overline{\beta} := (\widetilde{\beta}, \beta_{1,2k-1}, \beta_{0,2k}),$$

$$\breve{\beta} := (\widehat{\beta}, \beta_{3,2k-3}, \beta_{2,2k-2}) \quad \text{and} \quad \overline{\overline{\beta}} := (\breve{\beta}, \beta_{3,2k-1}, \beta_{0,2k})$$

be subsequences of β ,

$$v := \left(\begin{array}{cccc} \beta_{1,0} & u^{(1)} & \cdots & u^{(k-2)} & \beta_{0,k-1} & \beta_{3,k-3} & \beta_{2,k-2} & \beta_{1,k-1}\end{array}\right), \quad u := \left(\begin{array}{cccc} v & \beta_{0,k}\end{array}\right),$$
$$s := \left(\begin{array}{cccc} u & \beta_{3,k-2}\end{array}\right), \quad w := \left(\begin{array}{cccc} \beta_{\frac{5}{3},0} & \beta_{2,0} & \beta_{1,1} & u^{(2)} & \cdots & u^{(k-1)} & \beta_{0,k} & \beta_{3,k-2} & \beta_{2,k-1}\end{array}\right)$$

vectors and

$$\overline{A} := \begin{pmatrix} \beta_0 & v \\ v^T & A_{\breve{\beta}} \end{pmatrix} \quad and \quad \widetilde{A} := \begin{pmatrix} \beta_0 & u \\ u^T & A_{\overline{\overline{\beta}}} \end{pmatrix}$$

matrices. Then β has a representing measure supported on $y^3 = x^4$ if and only if

(4.16)
$$sA_{\overline{\beta}}^{+}s^{T} \leq uA_{\overline{\beta}}^{+}w^{T} + \sqrt{(A_{\overline{\beta}}/A_{\overline{\beta}})(\widetilde{A}/A_{\overline{\beta}})}$$

one of the following statements hold:

- (1) One of the following holds:
 - If $k \ge 4$, then $Y^3 = X^4$ is a column relation of M(k).
 - If k = 3, then the equalities $\beta_{0,3} = \beta_{4,0}$, $\beta_{1,3} = \beta_{5,0}$, $\beta_{2,3} = \beta_{6,0}$, $\beta_{0,4} = \beta_{4,1}$, $\beta_{0,5} = \beta_{4,2}$.
 - If k = 2, then the equality $\beta_{0,3} = \beta_{4,0}$ holds.

•
$$k =$$

- (2) One of the following holds:
 - (a) $A_{\overline{\beta}} \succ 0$, $A \succ 0$ and the inequality in (4.16) is strict.
 - (b) $A_{\overline{\beta}} \succ 0$ and the following inequalities holds:

$$uA_{\widetilde{\beta}}^+u^T < sA_{\overline{\beta}}^+s^T \quad and \quad uA_{\overline{\beta}}^+w^T - \sqrt{(A_{\overline{\beta}}/A_{\overline{\beta}})(\widetilde{A}/A_{\overline{\beta}})} \leq sA_{\overline{\beta}}^+s^T.$$

(c)
$$A_{\overline{\beta}} \succeq 0$$
 and rank $A_{\widetilde{\beta}} = \operatorname{rank} A_{\overline{\beta}} = \operatorname{rank} (s^T A_{\overline{\beta}}).$

Moreover, if the representing measure exists, then there exists a $(\operatorname{rank} \overline{\beta})$ -atomic measure if and only if (2b) or (2c) holds. Otherwise there is a $(\operatorname{rank} \overline{\beta} + 1)$ -atomic measure

Proof. For $\{0, 3, 4, 6, \dots, 8k\}$ we define the numbers $\widetilde{\beta}_m$ by the following rule

$$\widetilde{\beta}_m := \begin{cases} \beta_{0,\frac{m}{4}}, & \text{if } m \pmod{4} = 0, \\ \beta_{3,\lfloor\frac{m}{4}\rfloor - 2}, & \text{if } m \pmod{4} = 1, \\ \beta_{2,\lfloor\frac{m}{4}\rfloor - 1}, & \text{if } m \pmod{4} = 2, \\ \beta_{1,\lfloor\frac{m}{4}\rfloor}, & \text{if } m \pmod{4} = 3. \end{cases}$$

Claim 1. Every number $\tilde{\beta}_m$ is well-defined.

We have to prove that $i + j \leq 2k$, where i, j are indices of $\beta_{i,j}$ used in the definition of β_m . We separate four cases according to m:

- $m \pmod{4} = 0: \frac{m}{4} \le 2k.$ $m \pmod{4} = 1: \lfloor \frac{m}{4} \rfloor 2 + 3 \le (2k 1) + 1 = 2k.$ $m \pmod{4} = 2: \lfloor \frac{m}{4} \rfloor 1 + 2 \le (2k 1) + 1 = 2k.$ $m \pmod{4} = 3: \lfloor \frac{m}{4} \rfloor + 1 \le (2k 1) + 1 = 2k.$

We also define $\tilde{\beta}_5 := \beta_{\frac{5}{2},0}$.

Claim 2. Let $t \in \mathbb{N}$. The atoms $(x_1^3, x_1^4), \ldots, (x_t^3, x_t^4)$ with densities $\lambda_1, \ldots, \lambda_t$ are the $(y^3 - x^4)$ representing measure for β with $\beta_{\frac{5}{2},0}$ known if and only if the atoms x_1,\ldots,x_t with densities $\lambda_1, \ldots, \lambda_t$ are the \mathbb{R} -representing measure for $\widetilde{\beta}(x, y) = (\widetilde{\beta}_0, x, y, \widetilde{\beta}_3, \ldots, \widetilde{\beta}_{2k})$.

The if part follows from the following calculation:

$$\widetilde{\beta}_{m} = \begin{cases} \beta_{0,\frac{m}{4}}, & \text{if } m \pmod{4} = 0, \\ \beta_{3,\lfloor\frac{m}{4}\rfloor-2}, & \text{if } m \pmod{4} = 1, \\ \beta_{2,\lfloor\frac{m}{4}\rfloor-1}, & \text{if } m \pmod{4} = 2, \\ \beta_{1,\lfloor\frac{m}{4}\rfloor}, & \text{if } m \pmod{4} = 3, \end{cases} = \begin{cases} \sum_{\ell=1}^{t} \lambda_{\ell}(x_{\ell}^{4})^{\frac{m}{4}}, & \text{if } m \pmod{4} = 0, \\ \sum_{\ell=1}^{t} \lambda_{\ell}(x_{\ell}^{3})^{3}(x_{\ell}^{4})^{\lfloor\frac{m}{4}\rfloor-2}, & \text{if } m \pmod{4} = 1, \\ \sum_{\ell=1}^{t} \lambda_{\ell}(x_{\ell}^{3})^{2}(x_{\ell}^{4})^{\lfloor\frac{m}{4}\rfloor-1}, & \text{if } m \pmod{4} = 2, \\ \sum_{\ell=1}^{t} \lambda_{\ell}x_{\ell}^{3}(x_{\ell}^{4})^{\lfloor\frac{m}{4}\rfloor}, & \text{if } m \pmod{4} = 3, \end{cases} = \sum_{\ell=1}^{t} \lambda_{\ell}x_{\ell}^{m}$$

where $m = 0, 3, 4, 6, \dots, 8k$ and

$$\widetilde{\beta}_{5} = \beta_{\frac{5}{3},0} = \sum_{\ell=1}^{t} \lambda_{\ell} (x_{\ell}^{3})^{\frac{5}{3}} = \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{5}$$

The only if part follows from the following calculation:

$$\beta_{i,j} = \beta_{i-4,j+3} = \dots = \beta_{i \pmod{4}, j+3\lfloor\frac{i}{4}\rfloor} = \beta_{3(i \pmod{4})+4(j+3\lfloor\frac{i}{4}\rfloor)}$$
$$= \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{3(i \pmod{4})+4(j+3\lfloor\frac{i}{4}\rfloor)} = \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{3(i \pmod{4})+4\lfloor\frac{i}{4}\rfloor} x_{\ell}^{4j} = \sum_{\ell=1}^{t} \lambda_{\ell} (x_{\ell}^{3})^{i} (x_{\ell}^{4})^{j}$$

where the first three equalities in the first line follow by M(k) being rg and

$$\beta_{\frac{5}{3},0} = \widetilde{\beta}_{5} = \sum_{\ell=1}^{t} \lambda_{\ell} x_{\ell}^{5} = \sum_{\ell=1}^{t} \lambda_{\ell} (x_{\ell}^{3})^{\frac{5}{3}}.$$

Using Claim 2 and a theorem of Bayer and Teichmann [BT06], implying that if a finite sequence has a K-representing measure, then it has a finitely atomic K-representing measure, the statement of the Corollary follows by Theorem 4.5.

REFERENCES

- [Akh65] N. I. Akhiezer, *The classical moment problem and some related questions in analysis*. Hafner Publishing Co., New York, 1965.
- [AhK62] N. I. Akhiezer, M. Krein, *Some questions in the theory of moments*. Transl. Math. Monographs 2, American Math. Soc. Providence, 1962.
- [Alb69] A. Albert, *Conditions for positive and nonnegative definiteness in terms of pseudoinverses*. SIAM J. Appl. Math. **17** (1969), 434–440.
- [BW11] M. Bakonyi, H. J. Woerdeman, *Matrix Completions, Moments, and Sums of Hermitian Squares*. Princeton University Press, Princeton, 2011.
- [BT06] C. Bayer, J. Teichmann, *The proof of Tchakaloff's theorem*. Proc. Amer. Math. Soc. **134** (2006), 3035–3040.
- [Ble15] G. Blekherman, *Positive Gorenstein ideals*. Proc. Amer. Math. Soc. **143** (2015), 69–86.
- [BF20] G. Blekherman, L. Fialkow, *The core variety and representing measures in the truncated moment problem*. J. of Operator Theory. **84** (2020), 185–209.
- [Bol96] V. Bolotnikov, On degenerate Hamburger moment problem and extensions of nonnegative Hankel block matrices. Integral equations Operator Theory **25** (1996), 253–276.
- [BK10] Burgdorf, S., Klep, I.: Trace-positive polynomials and the quartic tracial moment problem. C. R. Math. Acad. Sci. Paris 348, 721–726 (2010)
- [BK12] Burgdorf, S., Klep, I.: The truncated tracial moment problem. J. Oper. Theory 68, 141–163 (2012)
- [CH69] D. Crabtree, E. Haynsworth, *An identity for the Schur complement of a matrix*. Proc. Am. Math. Soc. **22** (1969), 364–366.
- [CF91] R. Curto, L. Fialkow, *Recursiveness, positivity, and truncated moment problems.* Houston J. Math. **17** (1991), 603–635.
- [CF96] R. Curto, L. Fialkow, *Solution of the truncated complex moment problem for flat data*. Mem. Amer. Math. Soc. **119** (1996).
- [CF98a] R. Curto, L. Fialkow, *Flat extensions of positive moment matrices: relations in analytic or conjugate terms.* Oper. Theory Adv. Appl. **104** (1998), 59–82.

- [CF98b] R. Curto, L. Fialkow, *Flat extensions of positive moment matrices: recursively generated relations*. Mem. Amer. Math. Soc. **136** (1998).
- [CF02] R. Curto, L. Fialkow, *Solution of the singular quartic moment problem*. J. Operator Theory **48** (2002), 315–354.
- [CF04] R. Curto, L. Fialkow, *Solution of the truncated parabolic moment problem*. Integral Equations Operator Theory **50** (2004), 169–196.
- [CF05] R. Curto, L. Fialkow, *Solution of the truncated hyperbolic moment problem*. Integral Equations Operator Theory **52** (2005), 181–218.
- [CF08] R. Curto, L. Fialkow, An analogue of the Riesz-Haviland theorem for the truncated moment problem. J. Funct. Anal. 225 (2008), 2709–2731.
- [CF13] R. Curto, L. Fialkow, *Recursively determined representing measures for bivariate truncated moment sequences*. J. Operator theory **70** (2013), 401–436.
- [CFM08] R. Curto, L. Fialkow, H. M. Möller, *The extremal truncated moment problem*. Integral Equations Operator Theory **60 (2)** (2008), 177-200.
- [CS15] R. Curto, S. Yoo, Non-extremal sextic moment problems. J. Funct. Anal. 269 (3) (2015), 758–780.
- [CS16] R. Curto, S. Yoo, *Concrete solution to the nonsingular quartic binary moment problem*. Proc. Amer. Math. Soc. **144** (2016), 249–258.
- [Dan92] J. Dancis, *Positive semidefinite completions of partial hermitian matrices*. Linear Algebra Appl. **175** (1992), 97–114.
- [DS18] P. di Dio, K. Schmüdgen, *The multidimensional truncated Moment Problem: Atoms, Determinacy, and Core Variety*. J. Funct. Anal. **274** (2018), 3124–3148.
- [DW05] M. Dritschel, H. Woerdeman, *Outer factorizations in one and several variables*. Trans. Amer. Math. Soc. **357** (2005), 4661–4679.
- [DU18] M. Dritschel, B. Undrakh, *Rational dilation problems associated with constrained algebras*. J. Math. Anal. Appl. **467** (2018), 95–131.
- [Hav35] E. K. Haviland, On the momentum problem for distribution functions in more than one dimension II. Amer. J. Math. **58** (2006), 164–168.
- [FN10] L. Fialkow, J. Nie, *Positivity of Riesz functionals and solutions of quadratic and quartic moment problems*. J. Funct. An. **258** (2010), 328–356.
- [Fia08] L. Fialkow, *Truncated multivariable moment problems with finite variety*. J. of Operator Theory **60** (2008), 343–377.
- [Fia11] L. Fialkow, Solution of the truncated moment problem with variety $y = x^3$. Trans. Amer. Math. Soc. 363 (2011), 3133–3165.
- [Fia14] L. Fialkow, *The truncated moment problem on parallel lines*. The Varied Landscape of Operator Theory (2014), 99–116.
- [Fia17] L. Fialkow, *The core variety of a multisequence in the truncated moment problem*. J. Math. Anal. Appl. **456** (2017), 946–969.
- [GJSW84] R. Grone, C. R. Johnson, E. M. Sá, H. Wolkowicz, Positive definite completions of partial hermitian matrices. Linear Algebra Appl. 58 (1984), 109–124.
- [IKLS17] M. Infusino, T. Kuna, J. L. Lebowitz, E. R. Speer, *The truncated moment problem on* \mathbb{N}_0 . J. Math. Anal. Appl. **452** (2017), 443–468.
- [Ioh82] I. S. Iohvidov, Hankel and Toeplitz matrices and forms: Algebraic theory. Birkhäuser Verlag, Boston, 1982.
- [KW13] D. Kimsey, H. Woerdeman, *The multivariable matrix valued K-moment problem on* \mathbb{R}^d , \mathbb{C}^d , \mathbb{T}^d . Trans. Amer. Math. Soc. **365** (2013), 5393–5430.
- [KN77] M. G. Krein, A. A. Nudelman, *The Markov moment problem and extremal problems*. Translations of Mathematical Monographs, Amer. Math. Soc., 1977.
- [KN77] M. G. Krein, A. A. Nudelman, *The Markov moment problem and extremal problems*. Translation of Mathematical Monographs, Amer. Math. Soc., 1977.
- [Las01] J. B. Lasserre, *Global optimization with polynomials and the problem of moments*. SIAM J. Optim. **11** (3) (2001), 796–817.
- [Las09] J. B. Lasserre, Moments, positive polynomials and their applications. Imperial College Press, 2009.
- [Lau05] M. Laurent, *Revising two theorems of Curto and Fialkow on moment matrices*. Proc. Amer. Math. Soc. **133** (2005), 2965–2976.

ALJAŽ ZALAR

- [Lau09] M. Laurent, Sums of squares, moment matrices and optimization over polynomials. In: Emerging Applications of Algebraic Geometry, Vol. 149 of IMA Volumes in Mathematics and its Applications, pp. 157–270, Springer-Verlag, 2009.
- [Mar08] M. Marshall, *Positive polynomials and sums of squares*. Mathematical Surveys and Monographs **146**, Amer. Math. Soc., 2008.
- [Nie14] . Nie, The A-truncated K-moment problem. Found. Comput. Math. 14 (2014), 1243–1276.
- [PS01] V. Powers, C. Scheiderer, *The moment problem for non-compact semialgebraic sets*. Adv. Geom. **1** (2001), 71–88.
- [Put93] M. Putinar, Positive polynomials on compact semi-algebraic sets. Indiana Univ. Math. J. 42 (1993), 969–984.
- [PS06] M. Putinar, C. Scheiderer, Multivariate moment problems: Geometry and indeterminateness. Ann. Sc. Norm. Super. Pisa Cl. Sci. 5 (2006), 137–157.
- [PS08] M. Putinar, K. Schmüdgen, *Multivariate determinateness*. Indiana Univ. Math. J. 57 (2008), 2931–2968.
- [PV99] M. Putinar, F. H. Vasilescu, *Solving moment problems by dimensional extension*. Ann. of Math. **149** (1999), 1087–1107.
- [Sch91] K. Schmüdgen, The K-moment problem for compact semi-algebraic sets. Math. Ann. 289 (1991), 203–206.
- [Sch03] K. Schmüdgen, On the moment problem for closed semi-algebraic sets. J. Reine Angew. Math. 588 (2003), 225–234.
- [Sch17] K. Schmüdgen, The moment problem. Graduate Texts in Mathematics 277, Springer, Cham, 2017.
- [Sto01] J. Stochel, Solving the truncated moment problem solves the moment problem. Glasgow J. Math. **43** (2001) 335–341.
- [Wol] Wolfram Research, Inc., Mathematica, Version 10.0, Wolfram Research, Inc., Champaign, IL, 2019.
- [Zha05] F. Zhang, The Schur Complement and Its Applications. Springer-Verlag, New York, 2005.