

NONLINEAR DIFFUSION IN THE KELLER-SEGEL MODEL OF PARABOLIC-PARABOLIC TYPE

XIANGSHENG XU

Department of Mathematics & Statistics
Mississippi State University
Mississippi State, MS 39762, USA

ABSTRACT. In this paper we study the initial boundary value problem for the system $u_t - \Delta u^m = -\operatorname{div}(u^q \nabla v)$, $v_t - \Delta v + v = u$. This problem is the so-called Keller-Segel model with nonlinear diffusion. Our investigation reveals that nonlinear diffusion can prevent overcrowding. To be precise, we show that solutions are bounded as long as $m > q > 0$, thereby substantially generalizing the known results in this area. Furthermore, our result seems to imply that the Keller-Segel model can have bounded solutions and blow-up ones simultaneously.

1. INTRODUCTION

Theoretical and mathematical modeling of chemotaxis dates back to the works of Patlak in the 1950s [16] and Keller and Segel in the 1970s [12]. The general form of the model reads:

$$(1.1) \quad \partial_t u = \operatorname{div}(k_1(u, v) \nabla u - k_2(u, v) \nabla v) + k_3(u, v) \quad \text{in } \Omega_T \equiv \Omega \times (0, T],$$

$$(1.2) \quad \partial_t v = k_c \Delta v + k_4(u, v) \quad \text{in } \Omega_T,$$

$$(1.3) \quad \frac{\partial u}{\partial \mathbf{n}} = \frac{\partial v}{\partial \mathbf{n}} = 0 \quad \text{on } \Sigma_T \equiv \partial \Omega \times (0, T],$$

$$(1.4) \quad (u, v)|_{t=0} = (u_0(x), v_0(x)) \quad \text{on } \Omega.$$

Here u denotes the cell density and v is the concentration of the chemical signal. The function k_1 is the diffusivity of the cells, k_2 is the chemotactic sensitivity, k_3 describes the cell growth and death. In the signal concentration model, k_4 describes the net effect of the production and degradation of the chemical signal. As for the remaining terms in the problem, Ω is a bounded domain in \mathbb{R}^N with $C^{1,1}$ boundary $\partial \Omega$, \mathbf{n} the unit outward normal to $\partial \Omega$, and T any positive number.

Motivated by applications, various assumptions on the given data were suggested to further simplify the model [6, 19]. In this paper we focus our attention on the so-called nonlinear-diffusion model. In this case,

$$k_1 = m u^{m-1}, \quad k_2 = u^q, \quad k_3 = 0, \quad k_c = 1, \quad k_4 = u - v,$$

where

$$m, q \in (0, \infty).$$

The resulting problem is:

$$(1.5) \quad u_t - \Delta u^m = -\operatorname{div}(u^q \nabla v) \quad \text{in } \Omega_T,$$

$$(1.6) \quad v_t - \Delta v + v = u \quad \text{in } \Omega_T,$$

$$(1.7) \quad \frac{\partial u^m}{\partial \mathbf{n}} = \frac{\partial v}{\partial \mathbf{n}} = 0 \quad \text{on } \Sigma_T,$$

$$(1.8) \quad (u, v)|_{t=0} = (u_0(x), v_0(x)) \quad \text{on } \Omega.$$

1991 *Mathematics Subject Classification*. Primary: 35B45, 35B65, 35Q92, 35K51.

Key words and phrases. The Keller-Segel model, chemotaxis equations, non-linear diffusion model, global existence, uniform gradient bounds .

It is certainly beyond the scope of this paper to give a comprehensive review for the Keller-Segel model. In this regard, we would like to refer the reader to [8, 9]. A problem similar to (1.5)-(1.8) was investigated in [10, 11, 20] under the assumptions that $N \geq 2, m \geq 1, q \geq 1$ (note that our q here is their $q - 1$). The global existence of a weak solution was established if, in addition, $m > q + 1 - \frac{2}{N}$. When this inequality fails, one obtains local existence and the global existence only holds for small data. Hölder continuity and uniqueness of weak solutions were considered in [13]. Some relevance of nonlinear diffusion in chemotaxis was discussed in [2].

The objective of this paper is to show that the results in the preceding papers can be substantially improved. Before stating our results, let us define our notion of a weak solution.

Definition 1.1. We say that (u, v) is a weak solution to (1.5)-(1.8) if

$$\begin{aligned} u &\in L^\infty(\Omega_T), \quad u \geq 0, \quad u^m \in L^2(0, T; W^{1,2}(\Omega)), \\ v &\in L^\infty(0, T; W^{1,\infty}(\Omega)), \quad v \geq 0 \end{aligned}$$

and

$$\begin{aligned} - \int_{\Omega_T} u \partial_t \xi dxdt + \int_{\Omega_T} \nabla u^m \nabla \xi dxdt &= \int_{\Omega} u_0 \xi(x, 0) dx + \int_{\Omega_T} u^q \nabla v \nabla \xi dxdt, \\ - \int_{\Omega_T} v \partial_t \eta dxdt + \int_{\Omega_T} \nabla v \nabla \eta dxdt &= \int_{\Omega} v_0 \eta(x, 0) dx + \int_{\Omega_T} (u - v) \eta dxdt \end{aligned}$$

for each pair of smooth functions (ξ, η) with $\xi(x, T) = \eta(x, T) = 0$.

Our main result is:

Theorem 1.2 (Main theorem). *Assume:*

- (H1) Ω is a bounded domain in \mathbb{R}^N , $N \geq 3$, with $C^{1,1}$ boundary $\partial\Omega$;
- (H2) $u_0 \in L^\infty(\Omega)$, $v_0 \in W^{1,\infty}(\Omega)$ with $u_0 \geq 0, v_0 \geq 0$;

Then there is a weak solution (u, v) to (1.5)-(1.8), provided that one of the following conditions holds.

- (H3) $m > 0$, $q > 0$, and $m > q$;
- (H4) $m > 0$, $1 \geq q > 0$, and $q + \frac{q-1}{N+1} \leq m \leq q$.

Note that (H4) allows the possibility that $m = q = 1$. This is the classical Keller-Segel system, which is well known to have blow-up solutions. Thus our theorem actually implies that the Keller-Segel model can have bounded solutions and blow-up ones simultaneously. As far as we know, this is the first result in this direction. Our method seems to suggest that solutions blow up as $m \rightarrow q^+$, while solutions remain bounded as $m \rightarrow q^-$ with $q \leq 1$. All the results are established under the assumption $N > 2$. But it is not difficult to see that Theorem 1.2 remains true for $N = 2$.

Motivated by numerical and modeling issues, the question of how blow-up of cells can be avoided has received a lot of attention. One way of doing this is to add a cross-diffusion term to the equation for v [7]. A second way is to alter the cell diffusion [1]. There are other related works. See, e.g., [2] in the context of volume effects. Here we show that nonlinear diffusion can also prevent blow-up.

Throughout this paper the letter c is always used to represent a positive number whose value is determined by the given data. The norm of a function in $L^p(\Omega)$ is denoted by $\|\cdot\|_{p,\Omega}$. The Lebesgue measure of a set D in \mathbb{R}^N is represented by $|D|$. Whenever there is no confusion, we suppress the dependence of a function on its variables, e.g., we write u for $u(x, t)$.

2. PRELIMINARIES

In this section we collect a few preparatory results. The first one deals with sequences of non-negative numbers which satisfy certain recursive inequalities.

Proposition 2.1. *Let $\{y_n\}, n = 0, 1, 2, \dots$, be a sequence of positive numbers satisfying the recursive inequalities*

$$y_{n+1} \leq cb^n y_n^{1+\alpha} \quad \text{for some } b > 1, c, \alpha \in (0, \infty).$$

If

$$y_0 \leq c^{-\frac{1}{\alpha}} b^{-\frac{1}{\alpha^2}},$$

then $\lim_{n \rightarrow \infty} y_n = 0$.

This proposition can be found in ([4], p.12).

The following proposition plays a key role in the proof of our main theorem. It can be viewed as a continuous version of Lemma 3.1 in [15, 17].

Proposition 2.2. *Let $h(\tau)$ be a continuous non-negative function defined on $[0, T_0]$ for some $T_0 > 0$. Suppose that there exist three positive numbers ε, δ, b such that*

$$(2.1) \quad h(\tau) \leq \varepsilon h^{1+\delta}(\tau) + b \quad \text{for each } \tau \in [0, T_0].$$

Then

$$(2.2) \quad h(\tau) \leq \frac{1}{[\varepsilon(1+\delta)]^{\frac{1}{\delta}}} \equiv s_0 \quad \text{for each } \tau \in [0, T_0],$$

provided that

$$(2.3) \quad \varepsilon \leq \frac{\delta^\delta}{(b+\delta)^\delta (1+\delta)^{1+\delta}} \quad \text{and } h(0) \leq s_0.$$

Proof. Consider the function $f(s) = \varepsilon s^{1+\delta} - s + b$ on $[0, \infty)$. Then condition (2.1) simply says

$$(2.4) \quad f(h(\tau)) \geq 0 \quad \text{for each } \tau \in [0, T_0].$$

It is easy to check that the function f achieves its minimum value at $s_0 = \frac{1}{[\varepsilon(1+\delta)]^{\frac{1}{\delta}}}$. The minimum value

$$(2.5) \quad \begin{aligned} f(s_0) &= \frac{\varepsilon}{[\varepsilon(1+\delta)]^{\frac{1+\delta}{\delta}}} - \frac{1}{[\varepsilon(1+\delta)]^{\frac{1}{\delta}}} + b \\ &= b - \frac{\delta}{\varepsilon^{\frac{1}{\delta}} (1+\delta)^{\frac{1+\delta}{\delta}}}. \end{aligned}$$

By the first inequality in (2.3), $f(s_0) \leq -\delta$. Consequently, the equation $f(s) = 0$ has exactly two solutions $0 < s_1 < s_2$ with s_0 lying in between. Evidently, f is positive on $[0, s_1)$, negative on (s_1, s_2) , and positive again on (s_2, ∞) . The range of h is a closed interval because of its continuity, and this interval is either contained in $[0, s_1)$ or (s_2, ∞) due to (2.4). The latter cannot occur due to the second inequality in (2.3). Thus the proposition follows. \square

Proposition 2.3. *Let v be the solution of the problem*

$$(2.6) \quad v_t - \Delta v + v = u \quad \text{in } \Omega_T,$$

$$(2.7) \quad \frac{\partial v}{\partial \mathbf{n}} = 0 \quad \text{on } \Sigma_T,$$

$$(2.8) \quad v(x, 0) = v_0(x) \quad \text{on } \Omega.$$

If (H1) holds, then for each $p > \frac{N+2}{2}$ there is a positive number c such that

$$(2.9) \quad \sup_{0 \leq t \leq T} \|\nabla v\|_{W^{1,\infty}(\Omega)} \leq c \|\nabla v_0\|_{W^{1,\infty}(\Omega)} + c \|u\|_{2p, \Omega_T}.$$

Proof. We do not believe that this result is new. However, we cannot find a good reference to it. So we offer a proof here. First we obtain a local interior estimate. The boundary estimate is achieved by flattening the relevant portion of the boundary.

Now fix a point $z_0 = (x_0, t_0) \in \Omega_T$. Then pick a number R from $(0, \min\{\text{dist}(x_0, \partial\Omega), \sqrt{t_0}\})$. Define a sequence of cylinders $Q_{R_n}(z_0)$ in Ω_T as follows:

$$Q_{R_n}(z_0) = B_{R_n}(x_0) \times (t_0 - R_n^2, t_0],$$

where

$$R_n = \frac{R}{2} + \frac{R}{2^{n+1}} \quad n = 0, 1, 2, \dots$$

Choose a sequence of smooth functions θ_n so that

$$\begin{aligned} \theta_n(x, t) &= 1 \quad \text{in } Q_{R_n}(z_0), \\ \theta_n(x, t) &= 0 \quad \text{outside } B_{R_{n-1}}(x_0) \text{ and } t < t_0 - R_{n-1}^2, \\ |\partial_t \theta_n(x, t)| &\leq \frac{c4^n}{R^2} \quad \text{on } Q_{R_{n-1}}(z_0), \\ |\nabla \theta_n(x, t)| &\leq \frac{c2^n}{R} \quad \text{on } Q_{R_{n-1}}(z_0), \quad \text{and} \\ 0 &\leq \theta_n(x, t) \leq 1 \quad \text{on } Q_{R_{n-1}}(z_0). \end{aligned}$$

Let p be given as in the lemma. Select

$$(2.10) \quad K \geq R^{1 - \frac{N+2}{2p}} \|u\|_{2p, Q_R(z_0)}$$

as below. Set

$$K_n = K - \frac{K}{2^{n+1}}, \quad n = 0, 1, 2, \dots$$

Fix an $i \in \{1, \dots, N\}$. Define

$$(2.11) \quad w = v_{x_i}.$$

Then w satisfies the equation

$$(2.12) \quad w_t - \Delta w + w = u_{x_i} \quad \text{in } \Omega_T.$$

Without loss of generality, assume $\sup_{\Omega_T} w = \|w\|_{\infty, \Omega_T}$. We use $\theta_{n+1}^2 (w - K_{n+1})^+$ as a test function in (2.12) to derive

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\Omega} \theta_{n+1}^2 [(w - K_{n+1})^+]^2 dx + \int_{\Omega} \theta_{n+1}^2 |\nabla (w - K_{n+1})^+|^2 dx + \int_{\Omega} w \theta_{n+1}^2 (w - K_{n+1})^+ dx \\ &= \int_{\Omega} \theta_{n+1} \partial_t \theta_{n+1} [(w - K_{n+1})^+]^2 dx - 2 \int_{\Omega} \theta_{n+1} \nabla \theta_{n+1} \nabla w (w - K_{n+1})^+ dx \\ (2.13) \quad &- \int_{\Omega} u \theta_{n+1}^2 \partial_{x_i} (w - K_{n+1})^+ - 2 \int_{\Omega} u \theta_{n+1} \partial_{x_i} \theta_{n+1} (w - K_{n+1})^+ dx, \end{aligned}$$

from whence follows

$$\begin{aligned} &\sup_{0 \leq t \leq t_0} \int_{\Omega} \theta_{n+1}^2 [(w - K_{n+1})^+]^2 dx + \int_0^{t_0} \int_{\Omega} \theta_{n+1}^2 |\nabla (w - K_{n+1})^+|^2 dx dt \\ &\leq \frac{c4^n}{R^2} \int_{Q_{R_n}(z_0)} [(w - K_{n+1})^+]^2 dx dt + c \int_{A_{n+1}} u^2 \theta_{n+1}^2 dx dt \\ (2.14) \quad &\leq \frac{c4^n}{R^2} y_n + c \|u^2\|_{p, Q_R(z_0)} |A_{n+1}|^{1 - \frac{1}{p}}, \end{aligned}$$

where

$$(2.15) \quad y_n = \int_{Q_{R_n}(z_0)} [(w - K_n)^+]^2 dxdt,$$

$$(2.16) \quad A_{n+1} = \{(x, t) \in Q_{R_n}(z_0) : w(x, t) \geq K_{n+1}\}.$$

By Poincaré's inequality,

$$(2.17) \quad \begin{aligned} & \int_0^{t_0} \int_{\Omega} [\theta_{n+1}(w - K_{n+1})^+]^{\frac{4}{N}+2} dxdt \\ & \leq \int_0^{t_0} \left(\int_{\Omega} [\theta_{n+1}(w - K_{n+1})^+]^2 dx \right)^{\frac{2}{N}} \left(\int_{\Omega} [\theta_{n+1}(w - K_{n+1})^+]^{\frac{2N}{N-2}} dx \right)^{\frac{N-2}{N}} dt \\ & \leq \left(\sup_{0 \leq t \leq t_0} \int_{\Omega} [\theta_{n+1}(w - K_{n+1})^+]^2 dx \right)^{\frac{2}{N}} \int_0^{t_0} \int_{\Omega} |\nabla (\theta_{n+1}(w - K_{n+1})^+)|^2 dxdt \\ & \leq c(1 + 4^n) \left(\frac{c4^n}{R^2} y_n + c \|u^2\|_{p, Q_R(z_0)} |A_{n+1}|^{1-\frac{1}{p}} \right)^{\frac{N+2}{N}}. \end{aligned}$$

Subsequently,

$$(2.18) \quad \begin{aligned} y_{n+1} & = \int_{Q_{R_{n+1}}(z_0)} [(w - K_{n+1})^+]^2 dxdt \\ & \leq \int_0^{t_0} \int_{\Omega} [\theta_{n+1}(w - K_{n+1})^+]^2 dxdt \\ & \leq \left(\int_0^{t_0} \int_{\Omega} [\theta_{n+1}(w - K_{n+1})^+]^{2\frac{N+2}{N}} dxdt \right)^{\frac{N}{N+2}} |A_{n+1}|^{\frac{2}{N+2}} \\ & \leq c4^{\frac{Nn}{N+2}} \left(\frac{c4^n}{R^2} y_n + c \|u^2\|_{p, Q_R(z_0)} |A_{n+1}|^{1-\frac{1}{p}} \right) |A_{n+1}|^{\frac{2}{N+2}} \\ & \leq c4^{\frac{Nn}{N+2}} \left(\frac{c4^n}{R^2} y_n + cR^{\frac{N+2}{p}-2} K^2 |A_{n+1}|^{1-\frac{1}{p}} \right) |A_{n+1}|^{\frac{2}{N+2}}. \end{aligned}$$

The last step is due to (2.10). We also have

$$(2.19) \quad y_n \geq \int_{A_{n+1}} (K_{n+1} - K_n)^2 dxdt = \frac{K^2}{4^{n+1}} |A_{n+1}|.$$

It immediately follows that

$$(2.20) \quad \begin{aligned} y_n |A_{n+1}|^{\frac{2}{N+2}} & = y_n |A_{n+1}|^{\frac{1}{p}} |A_{n+1}|^{\frac{2}{N+2} - \frac{1}{p}} \\ & \leq \frac{cR^{\frac{N+2}{p}} 4^{\frac{(n+1)[2p-(N+2)]}{p(N+2)}}}{K^{2\frac{2p-(N+2)}{p(N+2)}}} y_n^{1 + \frac{2p-(N+2)}{p(N+2)}}, \end{aligned}$$

$$(2.21) \quad \begin{aligned} K^2 |A_{n+1}|^{1-\frac{1}{p}} |A_{n+1}|^{\frac{2}{N+2}} & = K^2 |A_{n+1}| |A_{n+1}|^{\frac{2}{N+2} - \frac{1}{p}} \\ & \leq \frac{c4^{n+1 + \frac{(n+1)[2p-(N+2)]}{p(N+2)}}}{K^{2\frac{2p-(N+2)}{p(N+2)}}} y_n^{1 + \frac{2p-(N+2)}{p(N+2)}}. \end{aligned}$$

Use these in (2.18) to derive

$$(2.22) \quad y_{n+1} \leq \frac{cb^n R^{\frac{N+2}{p}-2} 1 + \frac{2p-(N+2)}{p(N+2)}}{K^{2\frac{2p-(N+2)}{p(N+2)}}} y_n.$$

By Proposition 2.1, if we choose K so large that

$$(2.23) \quad y_0 \leq cK^2 R^{N+2},$$

then

$$(2.24) \quad \sup_{Q_{\frac{R}{2}}(z_0)} w \leq K.$$

In view of (2.10), it is enough for us to take

$$(2.25) \quad K = c \left(\frac{y_0}{R^{N+2}} \right)^{\frac{1}{2}} + R^{1-\frac{N+2}{2p}} \|u\|_{2p, Q_R(z_0)}.$$

Recall that

$$(2.26) \quad y_0 = \int_{Q_R(z_0)} \left[\left(w - \frac{K}{2} \right)^+ \right]^2 dxdt \leq \int_{Q_R(z_0)} (w^+)^2 dxdt.$$

Hence,

$$(2.27) \quad \sup_{Q_{\frac{R}{2}}(z_0)} w \leq c \left(\int_{Q_R(z_0)} (w^+)^2 dxdt \right)^{\frac{1}{2}} + R^{1-\frac{N+2}{2p}} \|u\|_{2p, Q_R(z_0)}.$$

This is the so-called local interior estimate. Now we proceed to derive the boundary estimate. Suppose $x_0 \in \partial\Omega$. Our assumption on the boundary implies that there exist a neighborhood $U(x_0)$ of x_0 and a $C^{1,1}$ diffeomorphism \mathbb{T} defined on $U(x_0)$ such that the image of $U(x_0) \cap \Omega$ under \mathbb{T} is the half ball $B_\delta^+(y_0) = \{y : |y - y_0| < \delta, y_i > 0\}$, where $\delta > 0, y_0 = \mathbb{T}(x_0)$, and i is given as in (2.11). This implies that we have flatten $U(x_0) \cap \partial\Omega$ into a region in the plane $y_i = 0$ in the y space [3]. Set

$$\tilde{v} = v \circ \mathbb{T}^{-1}, \quad \tilde{w} = \tilde{v}_{y_i}.$$

We can choose \mathbb{T} so that $\tilde{w} = \tilde{w}(y, t)$ satisfies the boundary condition

$$(2.28) \quad \tilde{w} |_{y_i=0} = \tilde{v}_{y_i} |_{y_i=0} = \frac{\partial \tilde{v}}{\partial \mathbf{n}} |_{y_i=0} = 0.$$

One way of doing this is to pick $\mathbb{T} = \begin{pmatrix} f_1(x) \\ \vdots \\ f_N(x) \end{pmatrix}$ so that the graph of $f_1(x) = 0$ is $U(x_0) \cap \partial\Omega$ and

the set of vectors $\{\nabla f_1, \dots, \nabla f_N\}$ is orthogonal. By a result in [21], \tilde{w} satisfies the equation

$$\partial_t \tilde{w} - \operatorname{div} [(J_{\mathbb{T}}^T J_{\mathbb{T}}) \circ \mathbb{T}^{-1} \nabla \tilde{w}] + \tilde{w} = (\mathbf{h} J_{\mathbb{T}}) \circ \mathbb{T}^{-1} \nabla \tilde{w} + (J_{\mathbb{T}} \circ \mathbb{T}^{-1} \nabla \tilde{u})_i \quad \text{in } B_\delta^+(y_0),$$

where $J_{\mathbb{T}}$ is the Jacobian matrix of \mathbb{T} , i.e.,

$$J_{\mathbb{T}} = \nabla \mathbb{T},$$

$(J_{\mathbb{T}} \circ \mathbb{T}^{-1} \nabla \tilde{u})_i$ is the i -th component of the vector $J_{\mathbb{T}} \circ \mathbb{T}^{-1} \nabla \tilde{u}$, and the row vector \mathbf{h} is roughly $\operatorname{div}(J_{\mathbb{T}}^T J_{\mathbb{T}})$ and is, therefore, bounded by our assumption on \mathbb{T} . In view of (2.28), the method employed to prove (2.27) still works here. The only difference is that we use $B_{R_n}^+(y_0)$ instead of $B_{R_n}(y_0)$ in the proof. If $t_0 = 0$, then we just need to change $Q_{R_n}(z_0)$ to $B_{R_n} \times [0, R_n^2]$ and require

$$K \geq 2 \|\nabla v_0\|_{\infty, \Omega},$$

in addition to (2.10) in the proof. Subsequently, (2.27) follows.

Finally, use v as a test function in (3.14) to derive

$$(2.29) \quad \frac{1}{2} \frac{d}{dt} \int_{\Omega} v^2 + \int_{\Omega} |\nabla v|^2 dx + \int_{\Omega} v^2 dx = \int_{\Omega} uv dx.$$

It immediately follows that

$$(2.30) \quad \int_{\Omega_T} |\nabla v|^2 dx dt \leq c \int_{\Omega_T} u^2 dx dt + c \int_{\Omega} v_0^2 dx.$$

Finally, we have

$$(2.31) \quad \|v\|_{\infty, \Omega_T} \leq c \|u\|_{p, \Omega_T} + \|v_0\|_{\infty, \Omega}.$$

This completes the proof. □

3. PROOF OF THEOREM 1.2

A solution to (1.5)-(1.8) is constructed as the limit of a sequence of approximate solutions. Our approximate problems are formulated as follows (also see [20]):

$$(3.1) \quad \partial_t U - m \operatorname{div}((U^+ + \sigma)^{m-1} \nabla U) = -\operatorname{div}((U^+)^q \nabla V) \quad \text{in } \Omega_T,$$

$$(3.2) \quad \partial_t V - \Delta V + V = U \quad \text{in } \Omega_T,$$

$$(3.3) \quad \frac{\partial U}{\partial \mathbf{n}} = \frac{\partial V}{\partial \mathbf{n}} = 0 \quad \text{on } \Sigma_T,$$

$$(3.4) \quad (U, V)|_{t=0} = (u_0, v_0) \quad \text{on } \Omega,$$

where $\sigma > 0$. The existence of a solution to the above problem can be established via the Leray-Schauder fixed point theorem ([5], p.280). To this end, we define an operator $\mathbb{T}: L^\infty(\Omega_T) \rightarrow L^\infty(\Omega_T)$ as follows: Let $U \in L^\infty(\Omega_T)$. We say $w = \mathbb{T}(U)$ if w is the unique solution of the problem

$$\begin{aligned} \partial_t w - m \operatorname{div}((U^+ + \sigma)^{m-1} \nabla w) &= -\operatorname{div}((U^+)^q \nabla V) \quad \text{in } \Omega_T, \\ \frac{\partial w}{\partial \mathbf{n}} &= 0 \quad \text{on } \Sigma_T, \\ w|_{t=0} &= u_0 \quad \text{on } \Omega, \end{aligned}$$

where V solves the problem

$$\begin{aligned} \partial_t V - \Delta V + V &= U \quad \text{in } \Omega_T, \\ \frac{\partial V}{\partial \mathbf{n}} &= 0 \quad \text{on } \Sigma_T, \\ V|_{t=0} &= v_0 \quad \text{on } \Omega. \end{aligned}$$

To see that \mathbb{T} is well-defined, we conclude from Proposition 2.3 that $|\nabla V| \in L^\infty(\Omega_T)$. Moreover, the two initial boundary value problems in the definition of \mathbb{T} are both linear and uniformly parabolic. We can infer from ([14], Chap. III) that w is Hölder continuous in $\overline{\Omega_T}$. It follows that \mathbb{T} is continuous and maps bounded sets into precompact ones. We still need to show that there is a positive number c such that

$$(3.5) \quad \|U\|_{\infty, \Omega_T} \leq c$$

for all $U \in L^\infty(\Omega_T)$ and $\eta \in (0, 1)$ satisfying $U = \eta \mathbb{T}(U)$. This equation is equivalent to the following problem

$$(3.6) \quad \partial_t U - m \operatorname{div}((U^+ + \sigma)^{m-1} \nabla U) = -\eta \operatorname{div}((U^+)^q \nabla V) \quad \text{in } \Omega_T,$$

$$(3.7) \quad \partial_t V - \Delta V + V = U \quad \text{in } \Omega_T,$$

$$(3.8) \quad \frac{\partial U}{\partial \mathbf{n}} = \frac{\partial V}{\partial \mathbf{n}} = 0 \quad \text{on } \Sigma_T,$$

$$(3.9) \quad (U, V)|_{t=0} = (\eta u_0, v_0) \quad \text{on } \Omega.$$

Use U^- as a test function in (3.6) to get

$$-\frac{1}{2} \frac{d}{dt} \int_{\Omega} (U^-)^2 dx - m \int_{\Omega} (U^+ + \sigma)^{m-1} |\nabla U^-|^2 dx = 0.$$

Integrate to get

$$(3.10) \quad U \geq 0 \quad \text{a.e. on } \Omega_T.$$

This implies that

$$(3.11) \quad V \geq 0 \quad \text{a.e. on } \Omega_T.$$

We introduce the following change of dependent variables

$$(3.12) \quad u = U + \sigma, \quad v = V + \sigma.$$

Then (u, v) satisfies the problem

$$(3.13) \quad u_t - \Delta u^m = -\eta \operatorname{div}(u - \sigma)^q \nabla v \quad \text{in } \Omega_T,$$

$$(3.14) \quad v_t - \Delta v + v = u \quad \text{in } \Omega_T,$$

$$(3.15) \quad \frac{\partial u}{\partial \mathbf{n}} = \frac{\partial v}{\partial \mathbf{n}} = 0 \quad \text{on } \Sigma_T,$$

$$(3.16) \quad (u, v)|_{t=0} = (\eta u_0(x) + \sigma, v_0(x) + \sigma) \quad \text{on } \Omega.$$

There is no loss of generality for us to assume that $T \leq 1$. Otherwise, we simply consider $(u(x, Tt), v(x, Tt))$ on $[0, 1]$. From here on we will do that, and also let

$$(3.17) \quad \sigma \in (0, 1).$$

We already have $\eta \in (0, 1)$. The generic positive number c will be independent of all three of them.

Lemma 3.1. *Let (H3) hold. Then for each s sufficiently large there is a positive number c such that*

$$(3.18) \quad \sup_{0 \leq t \leq T} \int_{\Omega} u^{s+1} dx + \int_{\Omega_T} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx dt \leq c \|\nabla v\|_{\infty, \Omega_T}^{\frac{m+s}{m-q}} + c.$$

Proof. First remember that

$$(3.19) \quad \sigma \leq u \in L^\infty(\Omega_T).$$

Thus for each $r \in \mathbb{R}$, we have

$$(3.20) \quad u^r \in L^2(0, T; W^{1,2}(\Omega)).$$

Now pick a number

$$(3.21) \quad s > \max\{0, m - 2q\}.$$

Use u^s as a test function in (3.13) to derive

$$(3.22) \quad \begin{aligned} & \frac{1}{s+1} \frac{d}{dt} \int_{\Omega} u^{s+1} dx + ms \int_{\Omega} u^{m+s-2} |\nabla u|^2 dx \\ &= s\eta \int_{\Omega} (u - \sigma)^q u^{s-1} \nabla v \nabla u dx \leq s \int_{\Omega} u^{q+s-1} |\nabla v \nabla u| dx \\ &\leq \frac{1}{2} ms \int_{\Omega} u^{m+s-2} |\nabla u|^2 dx + \frac{s \|\nabla v\|_{\infty, \Omega_T}^2}{2m} \int_{\Omega} u^{2q-m+s} dx. \end{aligned}$$

To estimate the last integral, first notice that

$$(3.23) \quad \int_{\Omega} u^{m+s-2} |\nabla u|^2 dx = \frac{4}{(m+s)^2} \int_{\Omega} |\nabla u^{\frac{m+s}{2}}|^2 dx.$$

Recall the Sobolev embedding theorem which states that for each $r \in [1, N)$ there is a positive number c such that

$$(3.24) \quad \|w\|_{\frac{Nr}{N-r}, \Omega} \leq c \|\nabla w\|_{r, \Omega} + c \|w\|_{1, \Omega} \quad \text{for each } w \in W^{1, r}(\Omega).$$

We wish to apply this inequality with $w = u^{\frac{m+s}{2}}$ and $r = 2$. For this purpose, we further require

$$(3.25) \quad \frac{2q - m + s}{m + s} \leq \frac{N}{N - 2}.$$

Or equivalently,

$$(3.26) \quad s \geq \frac{(2q - m)(N - 2) - Nm}{2}.$$

We derive from Hölder's inequality and (3.24) that

$$(3.27) \quad \begin{aligned} \int_{\Omega} u^{2q-m+s} dx &= \int_{\Omega} (u^{m+s})^{\frac{2q-m+s}{m+s}} dx \\ &\leq c \left(\int_{\Omega} (u^{m+s})^{\frac{N}{N-2}} dx \right)^{\frac{(2q-m+s)(N-2)}{(m+s)N}} \\ &\leq c \left(\int_{\Omega} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx + \left(\int_{\Omega} u^{\frac{m+s}{2}} dx \right)^2 \right)^{\frac{2q-m+s}{m+s}} \\ &\leq c \left(\int_{\Omega} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx \right)^{\frac{2q-m+s}{m+s}} + c \left(\int_{\Omega} u^{\frac{m+s}{2}} dx \right)^{\frac{2(2q-m+s)}{m+s}}. \end{aligned}$$

We integrate (3.13) over Ω to get

$$\frac{d}{dt} \int_{\Omega} u dx = 0.$$

Subsequently,

$$(3.28) \quad \int_{\Omega} u(x, t) dx = \int_{\Omega} (\eta u_0(x) + \sigma) dx \leq c \quad \text{for each } t > 0.$$

If we further assume that

$$(3.29) \quad \frac{m+s}{2} < 2q - m + s,$$

then we can appeal to the interpolation inequality ([5], p.146), thereby deriving

$$(3.30) \quad \|u\|_{\frac{m+s}{2}, \Omega} \leq \varepsilon \|u\|_{2q-m+s, \Omega} + \frac{1}{\varepsilon^{\mu}} \|u\|_{1, \Omega} \leq \varepsilon \|u\|_{2q-m+s, \Omega} + \frac{c}{\varepsilon^{\mu}},$$

where $\varepsilon > 0$, $\mu = \left(1 - \frac{2}{m+s}\right) / \left(\frac{2}{m+s} - \frac{1}{2q-m+s}\right)$. Condition (3.29) is equivalent to

$$(3.31) \quad s > 3m - 4q.$$

Use (3.30) in (3.27) and choose ε suitably small in the resulting inequality to obtain

$$(3.32) \quad \int_{\Omega} u^{2q-m+s} dx \leq c \left(\int_{\Omega} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx \right)^{\frac{2q-m+s}{m+s}} + c.$$

Plug this into (3.22) to get

$$\begin{aligned}
& \frac{1}{s+1} \frac{d}{dt} \int_{\Omega} u^{s+1} dx + \frac{2ms}{(m+s)^2} \int_{\Omega} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx \\
& \leq c \left(\int_{\Omega} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx \right)^{\frac{2q-m+s}{m+s}} \|\nabla v\|_{\infty, \Omega_T}^2 + c \|\nabla v\|_{\infty, \Omega_T}^2 \\
(3.33) \quad & \leq \varepsilon \int_{\Omega} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx + c(\varepsilon) \|\nabla v\|_{\infty, \Omega_T}^{\frac{m+s}{m-q}} + c \|\nabla v\|_{\infty, \Omega_T}^2.
\end{aligned}$$

The last step is due to the assumption $m > q$ and Young's inequality ([5], p. 145). Once again, by taking ε suitably small, we arrive at

$$(3.34) \quad \sup_{0 \leq t \leq T} \int_{\Omega} u^{s+1} dx + \int_{\Omega_T} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx dt \leq c \|\nabla v\|_{\infty, \Omega_T}^{\frac{m+s}{m-q}} dt + c.$$

Here we have used the fact $\frac{m+s}{m-q} > 2$ due to (3.21). That is to say, the lemma is valid for any s that satisfies (3.21), (3.26), and (3.31). This completes the proof. \square

Lemma 3.2. *Let (H3) hold and s be given as in Lemma 3.1. Then there is a positive number c such that*

$$(3.35) \quad \|u\|_{\infty, \Omega_T} \leq c \|\nabla v\|_{\infty, \Omega_T}^{\gamma} + c,$$

where

$$(3.36) \quad \gamma = \frac{[(s+1)(N+2) + N(m-1)^+](m+s) + (s+1)N(m-q)(N+2)}{(s+1)(m-q)[(N+2)(s+1) + 2N(m-q)]}.$$

Proof. Let

$$(3.37) \quad K \geq 2(\|u_0\|_{\infty, \Omega} + 1)$$

be selected as below. Define

$$(3.38) \quad K_n = K - \frac{K}{2^{n+1}}, n = 0, 1, \dots.$$

Obviously,

$$(3.39) \quad \frac{K}{2} \leq K_n \leq K.$$

Set

$$(3.40) \quad S_n(t) = \{x \in \Omega : u(x, t) \geq K_n\},$$

$$(3.41) \quad A_n = \cup_{0 \leq t \leq T} S_n(t) = \{(x, t) \in \Omega_T : u(x, t) \geq K_n\}.$$

Subsequently,

$$(3.42) \quad \int_0^T |S_{n+1}(t)| dt = |A_{n+1}|.$$

To simplify our presentation, we also introduce two parameters

$$(3.43) \quad m_s = (s+1) \frac{2}{N} + m + s,$$

$$(3.44) \quad q_s = m_s - (2q - m + s),$$

where s is given as in Lemma 3.1, i.e., s is sufficiently large. Then use $(u^s - K_{n+1}^s)^+$ as a test function in (3.13) to derive

$$(3.45) \quad \begin{aligned} & \frac{d}{dt} \int_{\Omega} \int_0^u (\tau^s - K_{n+1}^s)^+ d\tau dx + ms \int_{S_{n+1}(t)} u^{m+s-2} |\nabla u|^2 dx \\ &= s\eta \int_{S_{n+1}(t)} (u - \sigma)^q u^{s-1} \nabla v \nabla u dx \leq s \int_{S_{n+1}(t)} u^{q+s-1} |\nabla v| |\nabla u| dx. \end{aligned}$$

After a suitable application of Cauchy's inequality ([14], p. 58), we integrate to obtain

$$(3.46) \quad \begin{aligned} & \sup_{0 \leq t \leq T} \int_{\Omega} \int_0^u (\tau^s - K_{n+1}^s)^+ d\tau dx + \int_{A_{n+1}} \left| \nabla u^{\frac{m+s}{2}} \right|^2 dx dt \\ & \leq c \int_{A_{n+1}} u^{2q-m+s} |\nabla v|^2 dx dt \leq c \|\nabla v\|_{\infty, \Omega_T}^2 \int_{A_{n+1}} u^{2q-m+s} dx dt. \end{aligned}$$

Since $s > 1$, we have

$$(3.47) \quad \begin{aligned} \int_{K_{n+1}}^u (\tau^s - K_{n+1}^s)^+ d\tau \chi_{A_{K_{n+1}}} & \geq \int_{K_{n+1}}^u [(\tau - K_{n+1})^+]^s d\tau \\ & = \frac{1}{s+1} [(u - K_{n+1})^+]^{s+1}. \end{aligned}$$

Recall that $m_s = (s+1)\frac{2}{N} + m + s$. We estimate, with the aid of Hölder's inequality and (3.24), that

$$(3.48) \quad \begin{aligned} y_{n+1} & \equiv \int_0^T \int_{\Omega} [(u - K_{n+1})^+]^{m_s} dx dt \\ & \leq \int_0^T \left(\int_{\Omega} [(u - K_{n+1})^+]^{s+1} dx \right)^{\frac{2}{N}} \left(\int_{\Omega} [(u - K_{n+1})^+]^{\frac{(m+s)N}{N-2}} dx \right)^{\frac{N-2}{N}} dt \\ & \leq c \left(\sup_{0 \leq t \leq T} \int_{\Omega} [(u - K_{n+1})^+]^{s+1} dx \right)^{\frac{2}{N}} \\ & \quad \cdot \int_0^T \left(\int_{\Omega} \left| \nabla [(u - K_{n+1})^+]^{\frac{m+s}{2}} \right|^2 dx + \left(\int_{\Omega} [(u - K_{n+1})^+]^{\frac{m+s}{2}} dx \right)^2 \right) dt \end{aligned}$$

We can easily verify that

$$(3.49) \quad \begin{aligned} \left| \nabla [(u - K_{n+1})^+]^{\frac{m+s}{2}} \right| &= \frac{m+s}{2} [(u - K_{n+1})^+]^{\frac{m+s}{2}-1} |\nabla u| \\ &\leq \frac{m+s}{2} u^{\frac{m+s}{2}-1} |\nabla u| \chi_{S_{n+1}(t)} = \left| \nabla u^{\frac{m+s}{2}} \right| \chi_{S_{n+1}(t)}, \end{aligned}$$

$$(3.50) \quad \int_{\Omega} [(u - K_{n+1})^+]^{\frac{m+s}{2}} dx \leq \left(\int_{\Omega} [(u - K_{n+1})^+]^{m_s} dx \right)^{\frac{m+s}{2m_s}} |S_{n+1}(t)|^{1-\frac{m+s}{2m_s}}.$$

The latter yields

$$\begin{aligned}
& \int_0^T \left(\int_{\Omega} [(u - K_{n+1})^+]^{\frac{m+s}{2}} dx \right)^2 dt \\
& \leq \int_0^T \left(\int_{\Omega} [(u - K_{n+1})^+]^{m_s} dx \right)^{\frac{m+s}{m_s}} |S_{n+1}(t)|^{2-\frac{m+s}{m_s}} dt \\
& \leq \left(\int_{\Omega_T} [(u - K_{n+1})^+]^{m_s} dx dt \right)^{\frac{m+s}{m_s}} \left(\int_0^T |S_{n+1}(t)|^{1+\frac{Nm_s}{2(s+1)}} dt \right)^{\frac{2(s+1)}{Nm_s}} \\
(3.51) \quad & \leq c|A_{n+1}|^{\frac{2(s+1)}{Nm_s}} y_n^{\frac{m+s}{m_s}}.
\end{aligned}$$

Here we have used the fact that $\{y_n\}$ is a decreasing sequence. Use (3.49) and (3.51) in (3.48) and take (3.46) into account to derive

$$\begin{aligned}
y_{n+1} & \leq c \|\nabla v\|_{\infty, \Omega_T}^{\frac{2(N+2)}{N}} \left(\int_{A_{n+1}} u^{2q-m+s} dx dt \right)^{\frac{N+2}{N}} \\
(3.52) \quad & + c \|\nabla v\|_{\infty, \Omega_T}^{\frac{4}{N}} \left(\int_{A_{n+1}} u^{2q-m+s} dx dt \right)^{\frac{2}{N}} |A_{n+1}|^{\frac{2(s+1)}{Nm_s}} y_n^{\frac{m+s}{m_s}}.
\end{aligned}$$

The first integral on the right-hand side of (3.52) can be estimated as follows:

$$\begin{aligned}
\left(\int_{A_{n+1}} u^{2q-m+s} dx dt \right)^{\frac{N+2}{N}} & = K_{n+1}^{\frac{(N+2)(2q-m+s)}{N}} \left(\int_{A_{n+1}} \left(\frac{u}{K_{n+1}} \right)^{2q-m+s} dx dt \right)^{\frac{N+2}{N}} \\
& \leq \frac{1}{K_{n+1}^{\frac{(N+2)m_s}{N} - \frac{(N+2)(2q-m+s)}{N}}} \left(\int_{A_{n+1}} u^{m_s} dx dt \right)^{\frac{N+2}{N}} \\
(3.53) \quad & = \frac{1}{K_{n+1}^{\frac{(N+2)q_s}{N}}} \left(\int_{A_{n+1}} u^{m_s} dx dt \right)^{1+\frac{2}{N}}.
\end{aligned}$$

Similarly,

$$(3.54) \quad \left(\int_{A_{n+1}} u^{2q-m+s} dx dt \right)^{\frac{2}{N}} \leq \frac{1}{K_{n+1}^{\frac{2q_s}{N}}} \left(\int_{A_{n+1}} u^{m_s} dx dt \right)^{\frac{2}{N}}.$$

Recall that

$$K_{n+1} - K_n = \frac{K}{2^{n+2}}, \quad \frac{K_{n+1} - K_n}{K_{n+1}} = \frac{1}{2^{n+2} - 1} > \frac{1}{2^{n+2}}.$$

With the aid of the preceding two results, we obtain

$$(3.55) \quad y_n \geq \int_{A_{n+1}} [(K_{n+1} - K_n)^+]^{m_s} dx dt = \frac{K^{m_s}}{2^{(n+2)m_s}} |A_{n+1}|,$$

$$\begin{aligned}
y_n & \geq \int_{A_{n+1}} u^{m_s} \left[\left(1 - \frac{K_n}{u} \right)^+ \right]^{m_s} dx dt \\
(3.56) \quad & \geq \int_{A_{n+1}} u^{m_s} \left(1 - \frac{K_n}{K_{n+1}} \right)^{m_s} dx dt \geq \frac{1}{2^{(n+2)m_s}} \int_{A_{n+1}} u^{m_s} dx dt.
\end{aligned}$$

By (3.55),

$$|A_{n+1}|^{\frac{2(s+1)}{Nm_s}} \leq \frac{2^{\frac{2(n+2)(s+1)}{N}}}{K^{\frac{2(s+1)}{N}}} y_n^{\frac{2(s+1)}{Nm_s}}.$$

Keeping this, (3.54), (3.39), and (3.56) in mind, we derive from (3.52) that

$$\begin{aligned} y_{n+1} &\leq \frac{c \|\nabla v\|_{\infty, \Omega_T}^{\frac{2(N+2)}{N}} 2^{\frac{(n+2)m_s(N+2)}{N}}}{K_{n+1}^{\frac{(N+2)q_s}{N}}} y_n^{1+\frac{2}{N}} \\ &\quad + \frac{c 2^{\frac{[2(s+1)+2m_s](n+2)}{N}} \|\nabla v\|_{\infty, \Omega_T}^{\frac{4}{N}}}{K^{\frac{2(s+1)}{N} + \frac{2q_s}{N}}} y_n^{1+\frac{2}{N}} \\ (3.57) \quad &\leq cb^n \left(\frac{\|\nabla v\|_{\infty, \Omega_T}^{\frac{2(N+2)}{N}}}{K^{\frac{(N+2)q_s}{N}}} + \frac{\|\nabla v\|_{\infty, \Omega_T}^{\frac{4}{N}}}{K^{\frac{2(s+1)+2q_s}{N}}} \right) y_n^{1+\frac{2}{N}}, \end{aligned}$$

where

$$b = \max \left\{ 2^{\frac{m_s(N+2)}{N}}, 2^{\frac{2(s+1)+2m_s}{N}} \right\}.$$

We can easily check from (3.43) and (3.44) that

$$(N+2)q_s \geq 2(s+1) + 2q_s \quad \text{if and only if} \quad m \geq q.$$

Recall that $K_n \geq 1$. Thus if (H3) holds, we can deduce from (3.57) that

$$y_{n+1} \leq \frac{cb^n \left(\|\nabla v\|_{\infty, \Omega_T}^{\frac{2(N+2)}{N}} + \|\nabla v\|_{\infty, \Omega_T}^{\frac{4}{N}} \right)}{K^{\frac{2(s+1)+2q_s}{N}}} y_n^{1+\frac{2}{N}}.$$

According to Proposition 2.1, if we choose K so large that

$$\begin{aligned} y_0 &= \int_{\Omega_T} \left[\left(u - \frac{K}{2} \right)^+ \right]^{m_s} dxdt \leq \int_{\Omega_T} u^{m_s} dxdt \\ &\leq \frac{cK^{s+1+q_s}}{\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2}, \end{aligned}$$

then

$$\sup_{\Omega_T} u \leq K.$$

In view of (3.37), it is enough for us to take

$$\begin{aligned} (3.58) \quad K &= c \left(\int_{\Omega_T} u^{m_s} dxdt \right)^{\frac{1}{s+1+q_s}} \left(\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2 \right)^{\frac{1}{s+1+q_s}} \\ &\quad + 2\|u_0\|_{\infty, \Omega} + 2. \end{aligned}$$

In light of (3.48), (3.51), and (3.18), we have

$$\begin{aligned}
\int_{\Omega_T} u^{m_s} dx dt &\leq \left(\sup_{0 \leq t \leq T} \int_{\Omega} u^{s+1} dx \right)^{\frac{2}{N}} \int_{\Omega_T} |\nabla u^{\frac{m+s}{2}}|^2 dx dt \\
&\quad + c \left(\sup_{0 \leq t \leq T} \int_{\Omega} u^{s+1} dx \right)^{\frac{2}{N}} \left(\int_{\Omega_T} u^{m_s} dx dt \right)^{\frac{m+s}{m_s}} \\
&\leq c \|\nabla v\|_{\infty, \Omega_T}^{\frac{(N+2)(m+s)}{N(m-q)}} + c + \varepsilon \int_{\Omega_T} u^{m_s} dx dt + c(\varepsilon) \left(\sup_{0 \leq t \leq T} \int_{\Omega} u^{s+1} dx \right)^{\frac{m_s}{s+1}}.
\end{aligned}$$

Choosing ε suitably small, we arrive at

$$\int_{\Omega_T} u^{m_s} dx dt \leq c \|\nabla v\|_{\infty, \Omega_T}^{\frac{(N+2)(m+s)}{N(m-q)}} + c \|\nabla v\|_{\infty, \Omega_T}^{\frac{m_s(m+s)}{(s+1)(m-q)}} + c.$$

Substituting this into (3.58) yields

$$\begin{aligned}
\|u\|_{\infty, \Omega_T} &\leq c \left[\left(\|\nabla v\|_{\infty, \Omega_T}^{\frac{(N+2)(m+s)}{N(m-q)}} + \|\nabla v\|_{\infty, \Omega_T}^{\frac{m_s(m+s)}{(s+1)(m-q)}} + 1 \right) \left(\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2 \right) \right]^{\frac{1}{s+1+q_s}} + c \\
&\leq c \left[\left(\|\nabla v\|_{\infty, \Omega_T}^{\left(\frac{N+2}{N} + \frac{(m-1)^+}{s+1} \right) \frac{m+s}{m-q}} + 1 \right) \left(\|\nabla v\|_{\infty, \Omega_T}^{N+2} + 1 \right) \right]^{\frac{1}{s+1+q_s}} + c \\
&\leq c \left(\|\nabla v\|_{\infty, \Omega_T}^{\frac{[(s+1)(N+2)+N(m-1)^+](m+s)}{(s+1)N(m-q)} + N+2} + 1 \right)^{\frac{1}{s+1+q_s}} + c \\
&\leq c \|\nabla v\|_{\infty, \Omega_T}^{\frac{[(s+1)(N+2)+N(m-1)^+](m+s)+(s+1)N(m-q)(N+2)}{(s+1)N(m-q)(s+1+q_s)}} + c.
\end{aligned}$$

This together with (3.44) implies the lemma. \square

Proof of Theorem 1.2 under (H3). We wish to show

$$(3.59) \quad \|v\|_{L^\infty(0,T;W^{1,\infty}(\Omega))} + \|u\|_{\infty, \Omega \times [0,T]} \leq c.$$

Let γ be given as in Lemma 3.2. Note that

$$\lim_{s \rightarrow \infty} \gamma = \lim_{s \rightarrow \infty} \frac{[(s+1)(N+2) + N(m-1)^+](m+s) + (s+1)N(m-q)(N+2)}{(s+1)(m-q)[(N+2)(s+1) + 2N(m-q)]} = \frac{1}{m-q}.$$

If $\frac{1}{m-q} > 1$, then there is a $\beta > 0$ such that

$$(3.60) \quad \gamma = 1 + \beta \quad \text{for some suitably large } s.$$

Fix this s and let p be given as in Proposition 2.3. We can derive from (2.9) and Lemma 3.2

$$\begin{aligned}
\|v\|_{L^\infty(0,T;W^{1,\infty}(\Omega))} + \|u\|_{\infty, \Omega_T} &\leq c \|u\|_{2p, \Omega_T} + c \|\nabla v\|_{\infty, \Omega_T}^{1+\beta} + c \\
&\leq c T^{\frac{1}{2p}} \|u\|_{\infty, \Omega_T} + c \|u\|_{2p, \Omega_T}^{1+\beta} + c \\
&\leq c T^{\frac{1}{2p}} \|\nabla v\|_{\infty, \Omega_T}^{1+\beta} + c T^{\frac{1+\beta}{2p}} \|u\|_{\infty, \Omega_T}^{1+\beta} + c \\
(3.61) \quad &\leq c T^{\frac{1}{2p}} \left(\|v\|_{L^\infty(0,T;W^{1,\infty}(\Omega))} + \|u\|_{\infty, \Omega_T} \right)^{1+\beta} + c.
\end{aligned}$$

Here we have used the fact that $T \leq 1$. Set

$$(3.62) \quad h(\tau) = \|v\|_{L^\infty(0,\tau;W^{1,\infty}(\Omega))} + \|u\|_{\infty, \Omega \times [0,\tau]}.$$

Let $T_0 \in (0, T]$ be selected below. It follows from (3.61) that

$$(3.63) \quad h(\tau) \leq cT_0^{\frac{1}{2p}} h^{1+\beta}(\tau) + c \text{ for each } \tau \in [0, T_0].$$

It is not difficult for us to see from the proof of Proposition 2.3 that ∇v is actually Hölder continuous on $\overline{\Omega_T}$, so is u for each fixed $\sigma > 0$. Thus $h(\tau)$ is a continuous function of τ . In view of Proposition 2.2, if we choose T_0 so that

$$(3.64) \quad cT_0^{\frac{1}{2p}} \leq \frac{\beta^\beta}{(c+\beta)^\beta(1+\beta)^{1+\beta}} \text{ and } \|\nabla(v_0+1)\|_{W^{1,\infty}(\Omega)} + \|u_0+1\|_{\infty,\Omega} \leq \frac{1}{\left[cT_0^{\frac{1}{2p}}(1+\beta)\right]^{\frac{1}{\beta}}}$$

then

$$(3.65) \quad \|\nabla v\|_{\infty,\Omega \times [0, T_0]} \leq \frac{1}{\left[cT_0^{\frac{1}{2p}}(1+\beta)\right]^{\frac{1}{\beta}}}.$$

By setting $T = 0$ in (3.61), we see that $\|\nabla(v_0+1)\|_{W^{1,\infty}(\Omega)} + \|u_0+1\|_{\infty,\Omega} \leq c$. If we take

$$(3.66) \quad cT_0^{\frac{1}{2p}} = \frac{\beta^\beta}{(c+\beta)^\beta(1+\beta)^{1+\beta}},$$

then the second inequality in (3.64) is automatically satisfied. Upon doing so, we arrive at

$$(3.67) \quad \|v\|_{L^\infty(0, T_0; W^{1,\infty}(\Omega))} + \|u\|_{\infty,\Omega \times [0, T_0]} \leq \frac{(c+\beta)(1+\beta)}{\beta}.$$

Set $k = \lfloor \frac{T}{T_0} \rfloor$, the integer part of the number $\frac{T}{T_0}$. If $k \geq 1$, we consider

$$(3.68) \quad u_{T_0}(x, t) = u(t + T_0, x), \quad v_{T_0}(x, t) = v(t + T_0, x) \text{ on } [0, T_0].$$

Obviously, (u_{T_0}, v_{T_0}) satisfies the same conditions as (u, v) on $\Omega \times (0, T_0)$. Thus we can repeat the previous arguments to yield (3.67) for (u_{T_0}, v_{T_0}) . After a finite number of steps, we obtain (3.59). Of course, in the last step, we will have to use $\min\{T_0, T - kT_0\}$ instead of T_0 .

If $\frac{1}{m-q} < 1$, an application of Young's inequality is enough to reach (3.59).

If $\frac{1}{m-q} = 1$, this can also be handled easily. We verify that $\frac{d\gamma}{ds}$ changes signs at least three times. Thus either γ decreases toward 1 as $s \rightarrow \infty$, which can be treated like the first case, or γ increases toward 1 as $s \rightarrow \infty$, which is essentially the second case.

Clearly, (3.5) is a consequence of (3.59). Thus we can conclude from the Leray-Schauder fixed point theorem that (3.1)-(3.4) has a solution. Denote the solution by (U_σ, V_σ) . In view of (3.10), we can rewrite (3.1)-(3.4) as

$$(3.69) \quad \partial_t U_\sigma - m \operatorname{div}((U_\sigma + \sigma)^{m-1} \nabla U_\sigma) = -\operatorname{div}(U_\sigma^q \nabla V_\sigma) \text{ in } \Omega_T,$$

$$(3.70) \quad \partial_t V_\sigma - \Delta V_\sigma + V_\sigma = U_\sigma \text{ in } \Omega_T,$$

$$(3.71) \quad \frac{\partial U_\sigma}{\partial \mathbf{n}} = \frac{\partial V_\sigma}{\partial \mathbf{n}} = 0 \text{ on } \Sigma_T,$$

$$(3.72) \quad (U_\sigma, V_\sigma)|_{t=0} = (u_0, v_0) \text{ on } \Omega.$$

Furthermore,

$$U_\sigma \geq 0, \quad V_\sigma \geq 0, \quad \text{and} \quad \|V_\sigma\|_{L^\infty(0, T; W^{1,\infty}(\Omega))} + \|U_\sigma\|_{\infty, \Omega_T} \leq c.$$

We wish to show that we can take $\sigma \rightarrow 0$ in (3.69)-(3.72). For this purpose, we use $(U_\sigma + \sigma)^m$ as a test function in (3.69) to derive

$$(3.73) \quad \frac{1}{m+1} \sup_{0 \leq t \leq T} \int_\Omega (U_\sigma + \sigma)^{m+1} dx + \int_{\Omega_T} |\nabla(U_\sigma + \sigma)^m|^2 dx dt \leq c.$$

We compute

$$\begin{aligned}
\partial_t(U_\sigma + \sigma)^{m+1} &= (m+1)(U_\sigma + \sigma)^m \partial_t U_\sigma \\
&= (m+1) \operatorname{div}((U_\sigma + \sigma)^m \nabla(U_\sigma + \sigma)^m) - (m+1) |\nabla(U_\sigma + \sigma)^m|^2 \\
&\quad - (m+1) \operatorname{div}((U_\sigma + \sigma)^m U_\sigma^q \nabla V_\sigma) + (m+1) U_\sigma^q \nabla V_\sigma \cdot \nabla(U_\sigma + \sigma)^m, \\
\nabla(U_\sigma + \sigma)^{m+1} &= \frac{m+1}{m} (U_\sigma + \sigma) \nabla(U_\sigma + \sigma)^m.
\end{aligned}$$

Thus the sequence $\{\partial_t(U_\sigma + \sigma)^{m+1}\}$ is bounded in $L^2(0, T; (W^{1,2}(\Omega))^*) + L^1(\Omega_T) \equiv \{\psi_1 + \psi_2 : \psi_1 \in L^2(0, T; (W^{1,2}(\Omega))^*), \psi_2 \in L^1(\Omega_T)\}$ and the sequence $\{(U_\sigma + \sigma)^{m+1}\}$ is bounded in $L^2(0, T; W^{1,2}(\Omega))$. This puts us in a position to apply the Lions-Aubin lemma [19]. Upon doing, we obtain the pre-compactness of $\{(U_\sigma + \sigma)^{m+1}\}$ in $L^2(\Omega_T)$. We can extract a subsequence of $\{U_\sigma + \sigma\}$, still denoted by $\{U_\sigma + \sigma\}$, such that $U_\sigma + \sigma$ converges a.e. on Ω_T . This is enough to justify passing to the limit in (3.69)-(3.72). The proof is complete. \square

We would like to remark that as $m \rightarrow q^+$ the upper bound in (3.67) deteriorates. This foretells the possibility that solutions blow up if $m = q$.

Proof of Theorem 1.2 under (H4). We will show that an estimate like (3.35) remains true even without the benefit of Lemma 3.1. Let s be given as before, i.e., s is large enough. With the aid of (H4), we can derive from (3.57) that

$$y_{n+1} \leq \frac{cb^n \left(\|\nabla v\|_{\infty, \Omega_T}^{\frac{2(N+2)}{N}} + \|\nabla v\|_{\infty, \Omega_T}^{\frac{4}{N}} \right)}{K^{\frac{(N+2)q_s}{N}}} y_n^{1+\frac{2}{N}}.$$

In light of Proposition 2.1, if K is so chosen that

$$y_0 \leq \frac{cK^{\frac{(N+2)q_s}{2}}}{\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2},$$

then

$$(3.74) \quad \sup_{\Omega_T} u \leq K.$$

In view of (3.37), it is enough for us to take

$$(3.75) \quad K = c \left(\int_{\Omega_T} u^{m_s} dx dt \right)^{\frac{2}{(N+2)q_s}} \left(\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2 \right)^{\frac{2}{(N+2)q_s}} + 2\|u_0\|_{\infty, \Omega} + 2.$$

If

$$(3.76) \quad \frac{2m_s}{(N+2)q_s} < 1,$$

or equivalently,

$$q < 1 \quad \text{and} \quad m > q + \frac{q-1}{N+1},$$

then Young's inequality asserts

$$\begin{aligned}
K &\leq \varepsilon \|u\|_{m_s, \Omega_T} + c(\varepsilon) \left(\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2 \right)^{\frac{2}{(N+2)q_s - 2m_s}} \\
&\quad + 2\|u_0\|_{\infty, \Omega} + 2.
\end{aligned}$$

Use this in (3.74) to derive

$$(3.77) \quad \|u\|_{\infty, \Omega_T} \leq c \|\nabla v\|_{\infty, \Omega_T}^{\frac{N+2}{(N+1)m - (N+2)q+1}} + c\|u_0\|_{\infty, \Omega} + c.$$

If

$$(3.78) \quad \frac{2m_s}{(N+2)q_s} = 1,$$

we can appeal to the interpolation inequality ([5], p. 146) to obtain

$$(3.79) \quad \|u\|_{m_s, \Omega_T} \leq \varepsilon \|u\|_{\infty, \Omega_T} + \frac{1}{\varepsilon^{m_s-1}} \|u\|_{1, \Omega_T} \leq \varepsilon \|u\|_{\infty, \Omega_T} + \frac{c}{\varepsilon^{m_s-1}}.$$

With this in mind, we derive from (3.75) that

$$\begin{aligned} K &\leq c \left(\varepsilon \|u\|_{\infty, \Omega_T} + \frac{c}{\varepsilon^{m_s-1}} \right) \left(\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2 \right)^{\frac{2}{(N+2)q_s}} \\ &\quad + 2\|u_0\|_{\infty, \Omega} + 2 \\ &= \alpha \|u\|_{\infty, \Omega_T} + \frac{c}{\alpha^{m_s-1}} \left(\|\nabla v\|_{\infty, \Omega_T}^{N+2} + \|\nabla v\|_{\infty, \Omega_T}^2 \right)^{\frac{2m_s}{(N+2)q_s}} \\ &\quad + 2\|u_0\|_{\infty, \Omega} + 2. \end{aligned}$$

Plug this into (3.74) and choose α suitably small in the resulting inequality to derive

$$(3.80) \quad \|u\|_{\infty, \Omega_T} \leq c \|\nabla v\|_{\infty, \Omega_T}^{N+2} + c \|u_0\|_{\infty, \Omega} + c.$$

The rest of the proof is similar to that under (H3). That is, (3.59) can be inferred from either (3.77) or (3.80). \square

REFERENCES

- [1] M. Burger, M. Di Francesco, and Y. DolakStruss, The KellerSegel model for chemotaxis with prevention of overcrowding: linear vs. nonlinear diffusion, *SIAM J. Math. Anal.*, **38**(2006),12881315.
- [2] V. Calvez and J. A. Carrillo, Volume effects in the KellerSegel model: energy estimates preventing blow-up, *J. Math. Pures Appl.*, **86**(2006), 155175.
- [3] F. Chiarenza, M. Frasca, and P. Longo, $W^{2,p}$ -solvability of the Dirichlet problem for nondivergence elliptic equations with VMO coefficients, *Trans. Amer. Math. Soc.*, **336**(1993), 841-853.
- [4] E. DiBenedetto, *Degenerate Parabolic Equations*, Springer-Verlag, New York, 1993.
- [5] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, 1983.
- [6] T. Hillen and K. J. Painter, A users guide to PDE models for chemotaxis, *J. Math. Biol.*, **58**(2009), 183217.
- [7] S. Hittmeir and A. Jüngel, Cross diffusion preventing blow-up in the two-dimensional Keller-Segel model, *SIAM J. Math. Anal.*, **43**(2011), 997-1022.
- [8] D. Horstmann, From 1970 until present: The Keller-Segel model in chemotaxis and its consequences, Part I, *Jahresbericht der DMV*, **105**(2003), no. 3, 103165.
- [9] D. Horstmann, From 1970 until present: The Keller-Segel model in chemotaxis and its consequences, Part II, *Jahresbericht der DMV*, **106**(2004), no. 2, 5169.
- [10] S. Ishida and T. Yokota, Global existence of weak solutions to quasilinear degenerate Keller-Segel systems of parabolic-parabolic type, *J. Differential Equations*, **252**(2012), 1421-1440.
- [11] S. Ishida and T. Yokota, Global existence of weak solutions to quasilinear degenerate Keller-Segel systems of parabolic-parabolic type with small data, *J. Differential Equations*, **252**(2012), 2469-2491.
- [12] E.F. Keller and L.A. Segel, Initiation of slime mold aggregation viewed as an instability, *J. Theor. Biol.*, **26** (1970), 399-415.
- [13] S. Kim and Ki-Ahm Lee, Hölder regularity and uniqueness theorem on weak solutions to the degenerate KellerSegel system, *Nonlinear Anal.*, **138**(2016), 229-252.
- [14] Q.A. Ladyzenskaja, V.A. Solonnikov, and N.N. Ural'ceva, *Linear and Quasi-linear Equations of Parabolic Type*, Tran. Math. Monographs, Vol. 23, AMS, Providence, RI, 1968.
- [15] C. Miao, Weak solution of class of nonlinear heat equation systems and application to the NavierStokes system, *J. Differential Equations*, **61**(1986), 141151.
- [16] C. S. Patlak, Random walk with persistence and external bias, *Bull. Math: Biophys.*, **15** (1953), 311-338.
- [17] A.N. Sandjo, S. Moutari, and Y. Gningue, Solutions of fourth-order parabolic equation modeling thin film growth, *J. Differential Equations*, **259**(2015), 7260-7283.
- [18] J. Simon, Compact sets in the space $L^p(0, T; B)$, *Ann. Mat. Pura Appl.*, **146**(1987), 65-96.

- [19] A. Stevens, The derivation of chemotaxis equations as limit dynamics of moderately interacting stochastic many-particle systems, *SIAM J. Appl. Math.*, **61**(2000), 183212.
- [20] Y. Sugiyama and H. Kunii, Global existence and decay properties for a degenerate Keller-Segel model with a power factor in drift term, *J. Differential Equations*, **227**(2006), 333-364.
- [21] X. Xu, Global existence of strong solutions to a groundwater flow problem, arXiv:1912.03793 [math.AP], 2019. *Z. angew. Math. Phys.*, **71**(2020), Art# 127.

E-mail address: xxu@math.msstate.edu