

Conjectures on congruences involving binomial coefficients and Apéry-like numbers

Zhi-Hong Sun

School of Mathematics and Statistics

Huaian, Jiangsu 223300, P.R. China

Email: zhsun@hytc.edu.cn

Homepage: <http://maths.hytc.edu.cn/szh1.htm>

Abstract

In this paper, we pose many challenging conjectures on congruences involving binomial coefficients and Apéry-like numbers.

MSC: Primary 11A07, Secondary 11B65,11E25

Keywords: congruence, binomial coefficients, Apéry-like numbers

1. Introduction

We first illustrate the notation in the paper. Let \mathbb{Z} be the set of integers, and for a prime p let \mathbb{Z}_p be the set of rational numbers whose denominator is not divisible by p . Let $[x]$ be the greatest integer not exceeding x . For $a \in \mathbb{Z}$ and given odd prime p let $(\frac{a}{p})$ denote the Legendre symbol. For positive integers a, b and n , if $n = ax^2 + by^2$ for some integers x and y , we briefly write that $n = ax^2 + by^2$.

Let p be an odd prime. It is clear that

$$\binom{2k}{k} = \frac{2k(2k-1)\cdots(k+1)}{k!} \equiv 0 \pmod{p} \quad \text{for } k = \frac{p+1}{2}, \dots, p-1$$

and so $\frac{1}{2k-1} \binom{2k}{k} \in \mathbb{Z}_p$ for $k = 0, 1, \dots, p-1$. Actually, for $k \geq 1$,

$$\frac{1}{2k-1} \binom{2k}{k} = 2 \left(\binom{2k-2}{k-1} - \binom{2k-2}{k} \right) = \frac{2}{k} \binom{2k-2}{k-1} = 2C_{k-1} \in \mathbb{Z},$$

where $C_k = \frac{1}{k+1} \binom{2k}{k}$ is the k -th Catalan number.

Let $p > 3$ be a prime. In 1987, Beukers[B] conjectured a congruence equivalent to

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{64^k} \equiv \begin{cases} 0 \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \\ 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}. \end{cases}$$

This congruence was proved by several authors including Ishikawa[I] ($p \equiv 1 \pmod{4}$), van Hamme[vH] ($p \equiv 3 \pmod{4}$) and Ahlgren[A]. In 2003, Rodriguez-Villegas[RV] posed 22 conjectures on supercongruences modulo p^2 . In particular, the following congruences are equivalent to conjectures due to Rodriguez-Villegas:

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{108^k} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ 0 \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases}$$

$$\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{12^{3k}} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ 0 \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

These conjectures have been solved by Mortenson[M] and Zhi-Wei Sun[Su2]. Since

$$C_k = \frac{\binom{2k}{k}}{k+1} = \binom{2k}{k} - \binom{2k}{k+1},$$

from [Su2] one may deduce that

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{64^k(k+1)} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -(2p+2-2^{p-1}) \binom{(p-1)/2}{(p+1)/4}^2 \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-8)^k(k+1)} \equiv 6x^2 - 4p \pmod{p^2} \quad \text{for } p = x^2 + 4y^2 \equiv 1 \pmod{4},$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{108^k(k+1)} \equiv 4x^2 - 2p \pmod{p^2} \quad \text{for } p = x^2 + 3y^2 \equiv 1 \pmod{3},$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k(k+1)} \equiv 4x^2 - 2p \pmod{p^2} \quad \text{for } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8},$$

$$\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{12^{3k}(k+1)} \equiv 4x^2 - 2p \pmod{p^2} \quad \text{for } p = x^2 + 4y^2 \equiv 1 \pmod{4}.$$

Let p be an odd prime, $m \in \mathbb{Z}$ and $p \nmid m$. In [Su1,Su3], Z.W. Sun posed many conjectures concerning congruences modulo p^2 involving the sums

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{m^k}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{m^k}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{m^k}.$$

For 13 similar conjectures see [S1]. Most of these congruences modulo p were proved by the author in [S2-S5]. In [S10], the author conjectured many congruences modulo p^3 involving the above sums. For instance, for any prime $p \neq 2, 3, 7$,

$$\sum_{k=0}^{p-1} \binom{2k}{k}^3 \equiv \begin{cases} 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} & \text{if } p \equiv 1, 2, 4 \pmod{7} \text{ and so } p = x^2 + 7y^2, \\ -\frac{11}{4}p^2 \binom{3[p/7]}{[p/7]}^{-2} \equiv -11p^2 \binom{[3p/7]}{[p/7]}^{-2} \pmod{p^3} & \text{if } 7 \mid p-3, \\ -\frac{99}{64}p^2 \binom{3[p/7]}{[p/7]}^{-2} \equiv -11p^2 \binom{[6p/7]}{[2p/7]}^{-2} \pmod{p^3} & \text{if } 7 \mid p-5, \\ -\frac{25}{176}p^2 \binom{3[p/7]}{[p/7]}^{-2} \equiv -11p^2 \binom{[3p/7]}{[p/7]+1}^{-2} \pmod{p^3} & \text{if } 7 \mid p-6. \end{cases}$$

If we replace $\binom{3[p/7]}{[p/7]}$ with $\binom{[3p/7]}{[p/7]}$, for $p \equiv 3, 5, 6 \pmod{7}$ we have

$$\sum_{k=0}^{p-1} \binom{2k}{k}^3 \equiv \begin{cases} -\frac{11p^2}{(\binom{3p/7}{[p/7]})^2} \pmod{p^3} & \text{if } p \equiv 3 \pmod{7}, \\ -\frac{11p^2}{16(\binom{3p/7}{[p/7]})^2} \pmod{p^3} & \text{if } p \equiv 5 \pmod{7}, \\ -\frac{11p^2}{4(\binom{3p/7}{[p/7]})^2} \pmod{p^3} & \text{if } p \equiv 6 \pmod{7}. \end{cases}$$

In Section 2, with the help of Maple, we pose many conjectures on congruences modulo p^2 involving the sums

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{m^k(k+1)^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{m^k(2k-1)^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{m^k(k+1)}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{m^k(k+1)^2}, \\ & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{m^k(2k-1)}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{m^k(2k-1)^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{m^k(2k-1)^3}, \\ & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k(k+1)}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k(k+1)^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k(2k-1)}, \\ & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{m^k(k+1)}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{m^k(k+1)^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{m^k(2k-1)}, \\ & \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{m^k(k+1)}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{m^k(k+1)^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{m^k(2k-1)}. \end{aligned}$$

As typical examples, we conjecture that for any prime $p > 3$,

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k(k+1)} \equiv \begin{cases} \frac{3}{2}x^2 - 4p \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and so } 4p = x^2 + 27y^2, \\ 2(2p+1)\binom{[2p/3]}{[p/3]}^2 + p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ & \sum_{k=0}^{p-1} \left(\frac{\binom{2k}{k}}{2k-1} \right)^3 \equiv \begin{cases} -804y^2 - 18p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -\frac{192}{7} \binom{[3p/7]}{[p/7]}^2 \pmod{p} & \text{if } p \equiv 3 \pmod{7}, \\ -\frac{3072}{7} \binom{[3p/7]}{[p/7]}^2 \pmod{p} & \text{if } p \equiv 5 \pmod{7}, \\ -\frac{768}{7} \binom{[3p/7]}{[p/7]}^2 \pmod{p} & \text{if } p \equiv 6 \pmod{7}, \end{cases} \\ & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-8)^k(2k-1)^2} \equiv \begin{cases} -4x^2 + 2p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ 6R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ & \left(\frac{p}{5} \right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{54000^k(k+1)^2} \end{aligned}$$

$$\begin{aligned}
&\equiv \begin{cases} \frac{2504}{25}x^2 - \frac{1414 + 25(\frac{p}{5})}{25}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -184R_3(p) + \frac{162 - 25(\frac{p}{5})}{25}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3} \text{ and } p \neq 5, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{1458^k (2k-1)} &\equiv \begin{cases} -\frac{61}{81}(4x^2 - 2p - \frac{p^2}{4x^2}) - \frac{44}{243}(-1)^{\frac{p-1}{2}}p \pmod{p^3} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{32}{81}R_3(p) - \frac{44}{243}(-1)^{\frac{p-1}{2}}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k (2k-1)} &\equiv \begin{cases} -\frac{5}{8}(4x^2 - 2p) \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{5 - 4(-1)^{\frac{p-1}{2}}}{8}R_2(p) \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases}
\end{aligned}$$

where

$$\begin{aligned}
R_1(p) &= (2p+2-2^{p-1}) \left(\frac{\frac{p-1}{2}}{\frac{p-3}{4}} \right)^2, \\
R_3(p) &= \left(1 + 2p + \frac{4}{3}(2^{p-1}-1) - \frac{3}{2}(3^{p-1}-1) \right) \left(\frac{\frac{p-1}{2}}{\left[\frac{p}{6} \right]} \right)^2, \\
R_2(p) &= \left(1 + (4 + 2(-1)^{\frac{p-1}{2}})p - 4(2^{p-1}-1) - \frac{p}{2} \sum_{k=1}^{[p/8]} \frac{1}{k} \right) \left(\frac{\frac{p-1}{2}}{\left[\frac{p}{8} \right]} \right)^2.
\end{aligned}$$

In 2009, Zagier [Z] studied the Apéry-like numbers $\{u_n\}$ satisfying

$$u_0 = 1, \quad u_1 = b \quad \text{and} \quad (n+1)^2 u_{n+1} = (an(n+1) + b)u_n - cn^2 u_{n-1} \quad (n \geq 1),$$

where $a, b, c \in \mathbb{Z}$, $c \neq 0$ and $u_n \in \mathbb{Z}$ for $n = 1, 2, 3, \dots$. Let

$$\begin{aligned}
A'_n &= \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}, \quad f_n = \sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}, \\
S_n &= \sum_{k=0}^{[n/2]} \binom{2k}{k}^2 \binom{n}{2k} 4^{n-2k} = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k}, \\
a_n &= \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}, \quad Q_n = \sum_{k=0}^n \binom{n}{k} (-8)^{n-k} f_k, \\
W_n &= \sum_{k=0}^{[n/3]} \binom{2k}{k} \binom{3k}{k} \binom{n}{3k} (-3)^{n-3k}, \\
G_n &= \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k} 4^{n-k} = \sum_{k=0}^n \binom{n}{k} (-1)^k \binom{2k}{k}^2 16^{n-k}.
\end{aligned}$$

According to [Z] and [AZ], $\{A'_n\}$, $\{f_n\}$, $\{S_n\}$, $\{a_n\}$, $\{Q_n\}$, $\{W_n\}$ and $\{G_n\}$ are Apéry-like sequences with $(a, b, c) = (11, 3, -1), (7, 2, -8), (12, 4, 32), (10, 3, 9), (-17, -6, 72), (-9, -3, 27)$ and $(32, 12, 256)$, respectively. The sequence $\{f_n\}$ is called Franel numbers. In [S7-S12] the author systematically investigated congruences for sums involving S_n , f_n , W_n

and G_n . For $\{A'_n\}$, $\{f_n\}$, $\{S_n\}$, $\{a_n\}$, $\{Q_n\}$, $\{W_n\}$ and $\{G_n\}$ see A005258, A000172, A081085, A002893, A093388, A291898 and A143583 in Sloane's database "The On-Line Encyclopedia of Integer Sequences".

Let p be an odd prime. In [Su3], Z.W. Sun posed many congruences modulo p^2 involving Apéry-like numbers. In [S9,S10,S12], the author conjectured many congruences modulo p^3 involving Apéry-like numbers. In Section 3, we pose some conjectures on $\sum_{k=0}^{p-1} \frac{\binom{2k}{k} u_k}{m^k (2k-1)}$ modulo p^2 , where $u_n \in \{A'_n, f_n, S_n, a_n, Q_n, W_n, G_n\}$. As typical examples,

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{64^k (2k-1)} &\equiv 2(-1)^{\frac{p-1}{2}} p^2 \pmod{p^3}, \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{16^k (2k-1)} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } p \equiv 1 \pmod{4}, \\ -R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{(-12)^k (2k-1)} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } p \equiv 1 \pmod{3}, \\ -2(2p+1) \left(\frac{[2p/3]}{[p/3]}\right)^2 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{(-4)^k (2k-1)} &\equiv \begin{cases} 2p - 4x^2 \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -8R_3(p) \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

2. Conjectures on congruences involving binomial coefficients

By doing calculations with Maple, we form the following challenging conjectures.

Conjecture 2.1. Let $p > 3$ be a prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k (k+1)} &\equiv \begin{cases} \frac{3}{2}x^2 - 4p \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and so } 4p = x^2 + 27y^2, \\ 2(2p+1) \left(\frac{[2p/3]}{[p/3]}\right)^2 + p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k (k+1)^2} &\equiv \begin{cases} \frac{x^2}{4} - 3p \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and so } 4p = x^2 + 27y^2, \\ 13(2p+1) \left(\frac{[2p/3]}{[p/3]}\right)^2 + \frac{p}{2} \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k (2k-1)} &\equiv \begin{cases} -\frac{3}{4}x^2 + \frac{9}{8}p \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and } 4p = x^2 + 27y^2, \\ -\frac{1}{2}(2p+1) \left(\frac{[2p/3]}{[p/3]}\right)^2 + \frac{3}{8}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

Conjecture 2.2. Let $p > 3$ be a prime. Then

$$\begin{aligned} \left(\frac{10}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-12288000)^k (k+1)} \\ \equiv \begin{cases} \frac{966}{5}x^2 - \frac{3854}{5}p \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and so } 4p = x^2 + 27y^2, \\ 1280(2p+1) \left(\frac{[2p/3]}{[p/3]}\right)^2 + \frac{1922}{5}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \end{aligned}$$

$$\begin{aligned}
& \left(\frac{10}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-12288000)^k (k+1)^2} \\
& \equiv \begin{cases} -\frac{112604}{25}x^2 + \left(\frac{293656}{25} - \left(\frac{10}{p}\right)p\right) p \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and so } 4p = x^2 + 27y^2, \\ 11776(2p+1) \left(\frac{[2p/3]}{[p/3]}\right)^2 - \left(\frac{68448}{25} + \left(\frac{10}{p}\right)p\right) p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
& \left(\frac{10}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-12288000)^k (2k-1)} \\
& \equiv \begin{cases} -\frac{177}{200}x^2 + \frac{53199}{32000}p \pmod{p^2} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 27y^2, \\ -\frac{1}{20}(2p+1) \left(\frac{[2p/3]}{[p/3]}\right)^2 + \frac{3441}{32000}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3} \text{ and } p \neq 5. \end{cases}
\end{aligned}$$

Remark 2.1 Let p be a prime with $p > 5$. In [S10], the author conjectured that if $p \equiv 1 \pmod{3}$ and so $4p = x^2 + 27y^2$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k} \equiv \left(\frac{10}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-12288000)^k} \equiv x^2 - 2p - \frac{p^2}{x^2} \pmod{p^3};$$

if $p \equiv 2 \pmod{3}$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k} \equiv \frac{800}{161} \left(\frac{10}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-12288000)^k} \equiv \frac{3}{4} p^2 \left(\frac{[2p/3]}{[p/3]}\right)^{-2} \pmod{p^3}.$$

The congruence for $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k} \pmod{p^2}$ was conjectured by Z.W. Sun[Su1] earlier.

Let $p > 3$ be a prime. In [Su1,Su3], Z.W. Sun conjectured congruences for $\sum_{k=0}^{p-1} \binom{2k}{k}^3 / m^k \pmod{p^2}$ with $m = 1, -8, 16, -64, 256, -512, 4096$. Such conjectures were proved by the author in [S2]. In [S10], the author conjectured congruences for $\sum_{k=0}^{p-1} \binom{2k}{k}^3 / m^k \pmod{p^3}$ in the cases $m = 1, -8, 16, -64, 256, -512, 4096$.

Conjecture 2.3. Let p be an odd prime. Then

$$\begin{aligned}
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-8)^k (k+1)} \equiv \frac{1}{2} R_1(p) + p \pmod{p^2} \quad \text{for } p \equiv 3 \pmod{4}, \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-8)^k (k+1)^2} \equiv \begin{cases} 8x^2 - 8p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ 3R_1(p) + 2p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-8)^k (2k-1)} \equiv \begin{cases} -4x^2 \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ 2p - 2R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-8)^k (2k-1)^2} \equiv \begin{cases} -4x^2 + 2p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ 6R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-8)^k (2k-1)^3} \equiv \begin{cases} 48y^2 \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -6p - 12R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases}
\end{aligned}$$

Conjecture 2.4. Let p be an odd prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{64^k(k+1)^2} &\equiv \begin{cases} 8x^2 - 5p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -6R_1(p) - p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{64^k(2k-1)} &\equiv \begin{cases} p - 2x^2 \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{1}{2}R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{64^k(2k-1)^2} &\equiv \begin{cases} 2x^2 - p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ \frac{3}{2}R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{64^k(2k-1)^3} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -3R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

Conjecture 2.5. Let p be an odd prime. Then

$$\begin{aligned} (-1)^{\left[\frac{p}{4}\right]} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-512)^k(k+1)} &\equiv \begin{cases} 8x^2 - 6p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -4R_1(p) - 2p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ (-1)^{\left[\frac{p}{4}\right]} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-512)^k(k+1)^2} &\equiv \begin{cases} -16x^2 + (7 - (-1)^{\left[\frac{p}{4}\right]})p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -24R_1(p) - (-1)^{\left[\frac{p}{4}\right]}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \\ (-1)^{\left[\frac{p}{4}\right]} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-512)^k(2k-1)} &\equiv \begin{cases} \frac{5}{4}p - 3x^2 \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{p}{4} + \frac{1}{4}R_1(p) \pmod{p^2} & \text{if } 4 \mid p - 3, \end{cases} \\ (-1)^{\left[\frac{p}{4}\right]} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-512)^k(2k-1)^2} &\equiv \begin{cases} 2x^2 - \frac{5}{8}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{3}{4}R_1(p) + \frac{3}{8}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ (-1)^{\left[\frac{p}{4}\right]} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-512)^k(2k-1)^3} &\equiv \begin{cases} -\frac{3}{2}x^2 + \frac{3}{8}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ \frac{3}{2}R_1(p) - \frac{3}{8}p \pmod{p^2} & \text{if } 4 \mid p - 3. \end{cases} \end{aligned}$$

Conjecture 2.6. Let $p > 3$ be a prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{648^k(k+1)} &\equiv \begin{cases} \frac{10}{3}x^2 - \frac{4}{3}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{3}{2}R_1(p) - \frac{p}{3} \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{648^k(k+1)^2} &\equiv \begin{cases} \frac{112}{9}x^2 - \frac{64}{9}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -11R_1(p) - \frac{10}{9}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{648^k(2k-1)} &\equiv \begin{cases} -\frac{76}{27}x^2 + \frac{104}{81}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{2}{9}R_1(p) + \frac{10}{81}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

Conjecture 2.7. Let $p > 3$ be a prime. Then

$$\begin{aligned} \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{12^{3k}(k+1)} &\equiv \frac{3}{5} R_1(p) \pmod{p^2} \quad \text{for } p \equiv 3 \pmod{4}, \\ \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{12^{3k}(k+1)^2} &\equiv \begin{cases} 8x^2 - (4 + (\frac{p}{3}))p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ \frac{138}{25}R_1(p) - (\frac{p}{3})p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{12^{3k}(2k-1)} &\equiv \begin{cases} -\frac{26}{9}x^2 + \frac{13}{9}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ \frac{1}{6}R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

Conjecture 2.8. Let p be a prime with $p \neq 2, 3, 11$. Then

$$\begin{aligned} \left(\frac{33}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{66^{3k}(k+1)} &\equiv \begin{cases} -26x^2 + 28p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{363}{10}R_1(p) - 15p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \left(\frac{33}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{66^{3k}(k+1)^2} &\equiv \begin{cases} 488x^2 - (295 + (\frac{33}{p}))p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{8349}{25}R_1(p) + (51 - (\frac{33}{p}))p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \left(\frac{33}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{66^{3k}(2k-1)} &\equiv \begin{cases} -\frac{3716}{1089}x^2 + \frac{18848}{11979}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{2}{33}R_1(p) + \frac{530}{3993}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

Conjecture 2.9. Let $p > 3$ be a prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{16^k(k+1)} &\equiv \begin{cases} \frac{16}{3}x^2 - \frac{10}{3}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{4}{3}R_3(p) - \frac{2}{3}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{16^k(k+1)^2} &\equiv \begin{cases} 8x^2 - 7p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -8R_3(p) - 3p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{16^k(2k-1)} &\equiv \begin{cases} 4y^2 \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{8}{3}R_3(p) + \frac{2}{3}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{16^k(2k-1)^2} &\equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ 8R_3(p) \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{16^k(2k-1)^3} &\equiv \begin{cases} -12y^2 \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -16R_3(p) - 2p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

Conjecture 2.10. Let $p > 3$ be a prime. Then

$$(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{256^k(k+1)} \equiv \begin{cases} \frac{8}{3}x^2 - \frac{2}{3}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ \frac{16}{3}R_3(p) + \frac{2}{3}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases}$$

$$\begin{aligned}
& (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{256^k (k+1)^2} \\
& \equiv \begin{cases} 16x^2 - (8 + (-1)^{\frac{p-1}{2}})p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ 32R_3(p) - (-1)^{\frac{p-1}{2}}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
& (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{256^k (2k-1)} \equiv \begin{cases} 8y^2 - \frac{3}{2}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ \frac{2}{3}R_3(p) - \frac{p}{6} \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
& (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{256^k (2k-1)^2} \equiv \begin{cases} 2x^2 - \frac{3}{4}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -2R_3(p) + \frac{p}{4} \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
& (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{256^k (2k-1)^3} \equiv \begin{cases} \frac{p}{4} - x^2 \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ 4R_3(p) - \frac{p}{4} \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}
\end{aligned}$$

Moreover,

$$(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{256^k (2k-1)} \equiv -\frac{1}{4} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{16^k (2k-1)} \pmod{p^3} \quad \text{for } p \equiv 2 \pmod{3}.$$

Conjecture 2.11. Let $p > 3$ be a prime. Then

$$\begin{aligned}
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{108^k (k+1)} \equiv -2R_3(p) \pmod{p^2} \quad \text{for } p \equiv 2 \pmod{3}, \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{108^k (k+1)^2} \equiv \begin{cases} 8x^2 - 5p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -13R_3(p) - p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{108^k (2k-1)} \equiv \begin{cases} -\frac{5}{9}(4x^2 - 2p) \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{8}{9}R_3(p) \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases}
\end{aligned}$$

Conjecture 2.12. Let $p > 3$ be a prime. Then

$$\begin{aligned}
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-144)^k (k+1)} \equiv \begin{cases} \frac{16}{3}x^2 - \frac{10}{3}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ \frac{4}{3}R_3(p) + \frac{2}{3}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-144)^k (k+1)^2} \equiv \begin{cases} \frac{40}{9}x^2 - \frac{43}{9}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ \frac{88}{9}R_3(p) + \frac{5}{9}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\
& \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-144)^k (2k-1)} \equiv \begin{cases} -\frac{28}{9}x^2 + \frac{8}{9}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{8}{9}R_3(p) + \frac{2}{3}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3} \end{cases}
\end{aligned}$$

Conjecture 2.13. Let $p > 5$ be a prime. Then

$$\begin{aligned} \left(\frac{p}{5}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{54000^k (k+1)} &\equiv \begin{cases} -\frac{16}{5}x^2 + \frac{26}{5}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -20R_3(p) - \frac{18}{5}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \left(\frac{p}{5}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{54000^k (k+1)^2} &\equiv \begin{cases} \frac{2504}{25}x^2 - \frac{1414 + 25(\frac{p}{5})}{25}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -184R_3(p) + \frac{162 - 25(\frac{p}{5})}{25}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \left(\frac{p}{5}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{54000^k (2k-1)} &\equiv \begin{cases} -\frac{748}{225}x^2 + \frac{1708}{1125}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{8}{45}R_3(p) + \frac{18}{125}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

Conjecture 2.14. Let $p > 3$ be a prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{1458^k (k+1)} &\equiv \begin{cases} 2(-1)^{\frac{p-1}{2}}p - p^2 \pmod{p^3} & \text{if } p \equiv 1 \pmod{3}, \\ -12R_3(p) + 2(-1)^{\frac{p-1}{2}}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{1458^k (k+1)^2} &\equiv \begin{cases} 42x^2 - (22 + 2(-1)^{\frac{p-1}{2}})p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -78R_3(p) - (1 + 2(-1)^{\frac{p-1}{2}})p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{1458^k (2k-1)} &\equiv \begin{cases} -\frac{61}{81}(4x^2 - 2p - \frac{p^2}{4x^2}) - \frac{44}{243}(-1)^{\frac{p-1}{2}}p \pmod{p^3} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{32}{81}R_3(p) - \frac{44}{243}(-1)^{\frac{p-1}{2}}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

Conjecture 2.15. Let p be an odd prime. Then

$$\begin{aligned} (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-64)^k (k+1)} &\equiv \begin{cases} 6x^2 - 4p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{5 - 4(-1)^{\frac{p-1}{2}}}{2}R_2(p) - p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases} \\ (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-64)^k (k+1)^2} &\equiv \begin{cases} 4x^2 - 5p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1 \pmod{8}, \\ 4x^2 - 3p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 3 \pmod{8}, \\ -3R_2(p) - 3p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ -27R_2(p) - p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases} \\ (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-64)^k (2k-1)} &\equiv \begin{cases} p - 3x^2 \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{5 - 4(-1)^{\frac{p-1}{2}}}{4}R_2(p) - \frac{p}{2} \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases} \\ (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-64)^k (2k-1)^2} &\equiv \begin{cases} x^2 \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{3}{4}(5 - 4(-1)^{\frac{p-1}{2}})R_2(p) + \frac{p}{2} \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases} \end{aligned}$$

$$(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(-64)^k (2k-1)^3} \equiv \begin{cases} p - 2x^2 \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{3}{2}(5 - 4(-1)^{\frac{p-1}{2}})R_2(p) \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Conjecture 2.16. Let p be an odd prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k (k+1)} &\equiv \begin{cases} -\frac{1}{3}R_2(p) \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ -3R_2(p) \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k (k+1)^2} &\equiv \begin{cases} 8x^2 - 5p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{22}{9}R_2(p) - p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ -22R_2(p) - p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k (2k-1)} &\equiv \begin{cases} -\frac{5}{8}(4x^2 - 2p) \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{5 - 4(-1)^{\frac{p-1}{2}}}{8}R_2(p) \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8} \end{cases} \end{aligned}$$

Conjecture 2.17. Let p be a prime with $p \neq 2, 7$. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{28^{4k} (k+1)} &\equiv \begin{cases} -284x^2 + (142 + 144(\frac{p}{3}))p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -147R_2(p) + 144(\frac{p}{3})p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ -1323R_2(p) + 144(\frac{p}{3})p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{28^{4k} (k+1)^2} &\equiv \begin{cases} 5576x^2 - (3652 + (\frac{p}{3}))p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1 \pmod{8}, \\ 5576x^2 - (2789 + 864(\frac{p}{3}))p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 3 \pmod{8}, \\ -1078R_2(p) - (1 + 864(\frac{p}{3}))p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ -9702R_2(p) - (1 + 864(\frac{p}{3}))p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{28^{4k} (2k-1)} &\equiv \begin{cases} -\frac{2363}{686}x^2 + \frac{16541 - 1224(\frac{p}{3})}{9604}p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{45 - 36(-1)^{(p-1)/2}}{392}R_2(p) - \frac{306}{2401}(\frac{p}{3})p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases} \end{aligned}$$

Conjecture 2.18. Let p be an odd prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{8^k (k+1)} \equiv \begin{cases} \frac{11}{2}x^2 - \frac{7}{2}p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{5 - 4(-1)^{\frac{p-1}{2}}}{8}R_2(p) - \frac{3}{4}p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{8^k (k+1)^2} \equiv \begin{cases} \frac{31}{4}x^2 - \frac{29}{4}p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{13}{16}R_2(p) - \frac{27}{8}p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ -\frac{117}{16}R_2(p) - \frac{27}{8}p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{8^k (2k-1)} \equiv \begin{cases} \frac{7}{2}p - x^2 \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{3}{4}(5 - 4(-1)^{\frac{p-1}{2}})R_2(p) + 3p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases}$$

Conjecture 2.19. Let p be a prime with $p \neq 2, 5$. Then

$$\left(\frac{-5}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{20^{3k} (k+1)} \equiv \begin{cases} \frac{14}{5}x^2 - \frac{4}{5}p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{5}{6}R_2(p) + \frac{3}{5}p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ \frac{15}{2}R_2(p) + \frac{3}{5}p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases}$$

$$\left(\frac{-5}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{20^{3k} (k+1)^2}$$

$$\equiv \begin{cases} \frac{484}{25}x^2 - \left(\frac{244}{25} + \left(\frac{-5}{p}\right)\right)p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{23}{3}R_2(p) - \left(\frac{2}{25} + \left(\frac{-5}{p}\right)\right)p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8}, \\ 69R_2(p) - \left(\frac{2}{25} + \left(\frac{-5}{p}\right)\right)p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}, \end{cases}$$

$$\left(\frac{-5}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{20^{3k} (2k-1)} \equiv \begin{cases} -\frac{79}{25}x^2 + \frac{181}{125}p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{5 - 4(-1)^{\frac{p-1}{2}}}{20}R_2(p) - \frac{33}{250}p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Remark 2.2 Let p be a prime with $p > 7$ and $p \neq 71$. In [S10], the author conjectured that if $p \equiv 1, 3 \pmod{8}$ and so $p = x^2 + 2y^2$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k} \equiv \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{28^{4k}} \equiv \left(\frac{-5}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{20^{3k}} \equiv 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3};$$

if $p \equiv 5, 7 \pmod{8}$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{256^k} \equiv -\frac{441}{71} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{28^{4k}} \equiv -\frac{25}{7} \left(\frac{-5}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{20^{3k}}$$

$$\equiv \begin{cases} \frac{p^2}{3} \left(\frac{[p/4]}{[p/8]}\right)^{-2} \pmod{p^3} & \text{if } p \equiv 5 \pmod{8}, \\ -\frac{3}{2}p^2 \left(\frac{[p/4]}{[p/8]}\right)^{-2} \pmod{p^3} & \text{if } p \equiv 7 \pmod{8}. \end{cases}$$

The conjecture for $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{28^{4k}} \pmod{p^2}$ was due to Z.W. Sun [Su1].

For any odd prime p let

$$R_7(p) = \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1}.$$

Conjecture 2.20. Let p be a prime with $p \neq 2, 7$. Then

$$\begin{aligned} R_7(p) &= \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1} \equiv \begin{cases} -44y^2 + 2p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -\frac{1}{7} \left(\frac{[3p/7]}{[p/7]} \right)^2 \pmod{p} & \text{if } p \equiv 3 \pmod{7}, \\ -\frac{16}{7} \left(\frac{[3p/7]}{[p/7]} \right)^2 \pmod{p} & \text{if } p \equiv 5 \pmod{7}, \\ -\frac{4}{7} \left(\frac{[3p/7]}{[p/7]} \right)^2 \pmod{p} & \text{if } p \equiv 6 \pmod{7}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(k+1)^2} &\equiv \begin{cases} -68y^2 \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ 6R_7(p) + 2p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{2k-1} &\equiv \begin{cases} 14p - 36y^2 \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ 32R_7(p) + 48p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(2k-1)^2} &\equiv \begin{cases} -284y^2 + 34p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -96R_7(p) - 96p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{(2k-1)^3} &\equiv \begin{cases} -804y^2 - 18p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ 192R_7(p) + 144p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}. \end{cases} \end{aligned}$$

Conjecture 2.21. Let p be a prime with $p \neq 2, 7$. Then

$$\begin{aligned} (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{4096^k(k+1)} &\equiv \begin{cases} 72y^2 + 2p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -64R_7(p) - 66p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\ (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{4096^k(k+1)^2} &\equiv \begin{cases} -1136y^2 + (64 - (-1)^{\frac{p-1}{2}})p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -384R_7(p) - (456 + (-1)^{\frac{p-1}{2}})p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\ (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{4096^k(2k-1)} &\equiv \begin{cases} -\frac{7}{4}p + 22y^2 \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -\frac{1}{2}R_7(p) - \frac{3}{4}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\ (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{4096^k(2k-1)^2} & \end{aligned}$$

$$\begin{aligned}
&\equiv \begin{cases} -17y^2 + \frac{97}{64}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{3}{2}R_7(p) - \frac{3}{64}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\
&(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{4096^k (2k-1)^3} \\
&\equiv \begin{cases} \frac{201}{16}y^2 + \frac{81}{64} \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -3R_7(p) - \frac{243}{64}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7} \end{cases}
\end{aligned}$$

and

$$(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{4096^k (2k-1)} \equiv -\frac{1}{64} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{2k-1} \pmod{p^3} \quad \text{for } p \equiv 3, 5, 6 \pmod{7}.$$

Conjecture 2.22. Let p be a prime with $p \neq 2, 3, 7$. Then

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{81^k (k+1)} &\equiv \begin{cases} -\frac{100}{3}y^2 + 2p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{3}{2}R_7(p) + \frac{4}{3}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{81^k (k+1)^2} &\equiv \begin{cases} -\frac{436}{9}y^2 + \frac{4}{3}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ 11R_7(p) + \frac{94}{9}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{81^k (2k-1)} &\equiv \begin{cases} \frac{436}{27}y^2 - \frac{50}{81}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{16}{9}R_7(p) + \frac{208}{81}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}. \end{cases}
\end{aligned}$$

Conjecture 2.23. Let p be a prime with $p \neq 2, 3, 7$. Then

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-3969)^k (k+1)} &\equiv \begin{cases} \frac{28}{3}x^2 - \frac{22}{3}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -21R_7(p) - \frac{64}{3}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-3969)^k (k+1)^2} &\equiv \begin{cases} -\frac{356}{9}x^2 + \frac{188}{9}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -154R_7(p) - \frac{1612}{9}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-3969)^k (2k-1)} &\equiv \begin{cases} \frac{2102}{189}y^2 - \frac{7090}{3969}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{32}{63}R_7(p) + \frac{3088}{3969}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}. \end{cases}
\end{aligned}$$

Conjecture 2.24. Let p be a prime with $p > 7$. Then

$$\left(\frac{-15}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-15)^{3k} (k+1)}$$

$$\begin{aligned}
&\equiv \begin{cases} -\frac{188}{5}y^2 + 2p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{15}{128}R_7(p) - \frac{81}{40}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\
&\left(\frac{-15}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-15)^{3k}(k+1)^2} \\
&\equiv \begin{cases} \frac{172}{25}y^2 - \left(\frac{7}{5} + \left(\frac{-15}{p}\right)\right)p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{69}{64}R_7(p) - \left(\frac{1323}{100} + \left(\frac{-15}{p}\right)\right)p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases} \\
&\left(\frac{-15}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-15)^{3k}(2k-1)} \\
&\equiv \begin{cases} \frac{2542}{225}y^2 - \frac{2138}{1125}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -\frac{8}{15}R_7(p) - \frac{112}{125}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}. \end{cases}
\end{aligned}$$

Conjecture 2.25. Let $p > 3$ be a prime. Then

$$\begin{aligned}
&\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-12288)^k(k+1)} \\
&\equiv \begin{cases} 16x^2 - 14p \pmod{p^2} & \text{if } p \equiv 1 \pmod{12} \text{ and so } p = x^2 + 9y^2, \\ -8x^2 + 14p \pmod{p^2} & \text{if } p \equiv 5 \pmod{12} \text{ and so } 2p = x^2 + 9y^2, \\ -24 \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 7 \pmod{12}, \\ 48 \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 11 \pmod{12}, \end{cases} \\
&\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-12288)^k(k+1)^2} \\
&\equiv \begin{cases} -128x^2 + 75p \pmod{p^2} & \text{if } p \equiv 1 \pmod{12} \text{ and so } p = x^2 + 9y^2, \\ 64x^2 - 77p \pmod{p^2} & \text{if } p \equiv 5 \pmod{12} \text{ and so } 2p = x^2 + 9y^2, \\ -176 \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 7 \pmod{12}, \\ 352 \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 11 \pmod{12}, \end{cases} \\
&\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-12288)^k(2k-1)}
\end{aligned}$$

$$\equiv \begin{cases} -\frac{13}{4}x^2 + \frac{93}{64}p \pmod{p^2} & \text{if } p \equiv 1 \pmod{12} \text{ and so } p = x^2 + 9y^2, \\ \frac{13}{8}x^2 - \frac{93}{64}p \pmod{p^2} & \text{if } p \equiv 5 \pmod{12} \text{ and so } 2p = x^2 + 9y^2, \\ \frac{3}{16} \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 7 \pmod{12}, \\ -\frac{3}{8} \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 11 \pmod{12}. \end{cases}$$

Conjecture 2.26. Let p be a prime with $p \neq 2, 5$. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k} \equiv \begin{cases} 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} & \text{if } p \equiv 1, 9 \pmod{20} \text{ and so } p = x^2 + 5y^2, \\ 2p - 2x^2 + \frac{p^2}{2x^2} \pmod{p^3} & \text{if } p \equiv 3, 7 \pmod{20} \text{ and so } 2p = x^2 + 5y^2, \\ \frac{2p^2}{\binom{(p-1)/2}{[p/20]} \binom{(p-1)/2}{[3p/20]}} \pmod{p^3} & \text{if } p \equiv 11 \pmod{20}, \\ \frac{2p^2}{9 \binom{(p-1)/2}{[p/20]} \binom{(p-1)/2}{[3p/20]}} \pmod{p^3} & \text{if } p \equiv 13 \pmod{20}, \\ \frac{6p^2}{7 \binom{(p-1)/2}{[p/20]} \binom{(p-1)/2}{[3p/20]}} \pmod{p^3} & \text{if } p \equiv 17 \pmod{20}, \\ \frac{2p^2}{21 \binom{(p-1)/2}{[p/20]} \binom{(p-1)/2}{[3p/20]}} \pmod{p^3} & \text{if } p \equiv 19 \pmod{20}. \end{cases}$$

Remark 2.3 For any prime $p \neq 2, 5$, the congruence for $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k}$ modulo p^2 was first conjectured by Z.W. Sun in [Su1].

Conjecture 2.27. Let p be a prime with $p \neq 2, 5$. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (k+1)} \equiv \begin{cases} \frac{8}{5} \left(\frac{p-1}{[p/20]} \right) \left(\frac{p-1}{[3p/20]} \right) \pmod{p} & \text{if } p \equiv 1, 3, 7, 9 \pmod{20}, \\ \frac{4}{5} \left(\frac{p-1}{[p/20]} \right) \left(\frac{p-1}{[3p/20]} \right) \pmod{p} & \text{if } p \equiv 11 \pmod{20}, \\ \frac{36}{5} \left(\frac{p-1}{[p/20]} \right) \left(\frac{p-1}{[3p/20]} \right) \pmod{p} & \text{if } p \equiv 13 \pmod{20}, \\ \frac{28}{15} \left(\frac{p-1}{[p/20]} \right) \left(\frac{p-1}{[3p/20]} \right) \pmod{p} & \text{if } p \equiv 17 \pmod{20}, \\ \frac{84}{5} \left(\frac{p-1}{[p/20]} \right) \left(\frac{p-1}{[3p/20]} \right) \pmod{p} & \text{if } p \equiv 19 \pmod{20}. \end{cases}$$

Moreover, if $(\frac{-5}{p}) = 1$, then

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (k+1)} \\ & \equiv \begin{cases} \frac{32}{5}x^2 - \frac{22}{5}p \pmod{p^2} & \text{if } p \equiv 1, 9 \pmod{20} \text{ and so } p = x^2 + 5y^2, \\ -\frac{16}{5}x^2 + \frac{22}{5}p \pmod{p^2} & \text{if } p \equiv 3, 7 \pmod{20} \text{ and so } 2p = x^2 + 5y^2, \end{cases} \\ & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (k+1)^2} \\ & \equiv \begin{cases} -\frac{32}{5}x^2 + \frac{7}{5}p \pmod{p^2} & \text{if } p \equiv 1, 9 \pmod{20} \text{ and so } p = x^2 + 5y^2, \\ \frac{16}{5}x^2 - \frac{17}{5}p \pmod{p^2} & \text{if } p \equiv 3, 7 \pmod{20} \text{ and so } 2p = x^2 + 5y^2, \end{cases} \\ & \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (2k-1)} \\ & \equiv \begin{cases} -\frac{31}{10}x^2 + \frac{103}{80}p \pmod{p^2} & \text{if } p \equiv 1, 9 \pmod{20} \text{ and so } p = x^2 + 5y^2, \\ \frac{31}{20}x^2 - \frac{103}{80}p \pmod{p^2} & \text{if } p \equiv 3, 7 \pmod{20} \text{ and so } 2p = x^2 + 5y^2; \end{cases} \end{aligned}$$

if $(\frac{-5}{p}) = -1$, then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (k+1)^2} & \equiv \frac{22}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (k+1)} + (8(-1)^{\frac{p-1}{2}} - 1)p \pmod{p^2}, \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (2k-1)} & \equiv -\frac{3}{32} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (k+1)} - \frac{3}{8}(-1)^{\frac{p-1}{2}}p \pmod{p^2}. \end{aligned}$$

Conjecture 2.28. Let p be a prime with $(\frac{-6}{p}) = -1$. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k} \equiv \begin{cases} -\frac{7}{5} \cdot \frac{p^2}{\binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]}} \pmod{p^3} & \text{if } p \equiv 13 \pmod{24}, \\ \frac{7}{5} \cdot \frac{p^2}{\binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]}} \pmod{p^3} & \text{if } p \equiv 17 \pmod{24}, \\ \frac{1}{11} \cdot \frac{p^2}{\binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]}} \pmod{p^3} & \text{if } p \equiv 19 \pmod{24}, \\ -\frac{1}{11} \cdot \frac{p^2}{\binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]}} \pmod{p^3} & \text{if } p \equiv 23 \pmod{24} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}} \equiv \begin{cases} \frac{1}{5} \cdot \frac{p^2}{\binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]}} \pmod{p^3} & \text{if } p \equiv 13, 17 \pmod{24}, \\ -\frac{1}{77} \cdot \frac{p^2}{\binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]}} \pmod{p^3} & \text{if } p \equiv 19, 23 \pmod{24}. \end{cases}$$

Conjecture 2.29. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (k+1)} \equiv \begin{cases} \frac{7}{8} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 1, 11 \pmod{24}, \\ -\frac{7}{8} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 5, 7 \pmod{24}, \\ -\frac{5}{8} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 13 \pmod{24}, \\ \frac{5}{8} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 17 \pmod{24}, \\ \frac{77}{8} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 19 \pmod{24}, \\ -\frac{77}{8} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 23 \pmod{24}. \end{cases}$$

Moreover, if $(\frac{-6}{p}) = 1$, then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (k+1)} &\equiv \begin{cases} \frac{7}{2}x^2 - \frac{3}{2}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ 7x^2 - 2p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (k+1)^2} &\equiv \begin{cases} \frac{43}{4}x^2 - \frac{25}{4}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ \frac{43}{2}x^2 - \frac{13}{2}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (2k-1)} &\equiv \begin{cases} -\frac{23}{9}x^2 + \frac{7}{6}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ -\frac{46}{9}x^2 + \frac{25}{18}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}; \end{cases} \end{aligned}$$

if $(\frac{-6}{p}) = -1$, then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (k+1)^2} &\equiv \frac{13}{2} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (k+1)} - \left(1 + \frac{3}{2} \binom{p}{3}\right)p \pmod{p^2}, \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (2k-1)} &\equiv \frac{2}{9} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (k+1)} - \frac{1}{6} \binom{p}{3}p \pmod{p^2}. \end{aligned}$$

Conjecture 2.30. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k} (k+1)} \equiv \begin{cases} \frac{1}{3} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 1, 5 \pmod{24}, \\ -\frac{1}{3} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 7, 11 \pmod{24}, \\ \frac{5}{3} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 13, 17 \pmod{24}, \\ -\frac{77}{3} \binom{(p-1)/2}{[p/24]} \binom{(p-1)/2}{[5p/24]} \pmod{p} & \text{if } p \equiv 19, 23 \pmod{24}. \end{cases}$$

Moreover, if $(\frac{-6}{p}) = 1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}(k+1)} \equiv \begin{cases} \frac{4}{3}x^2 + \frac{2}{3}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ -\frac{8}{3}x^2 - \frac{2}{3}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}(k+1)^2} \equiv \begin{cases} \frac{280}{9}x^2 - \frac{157}{9}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ -\frac{560}{9}x^2 + \frac{139}{9}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}(2k-1)} \equiv \begin{cases} -\frac{55}{18}x^2 + \frac{49}{36}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ \frac{55}{9}x^2 - \frac{49}{36}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}; \end{cases}$$

if $(\frac{-6}{p}) = -1$, then

$$\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}(k+1)} \equiv -\frac{8}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k(k+1)} + \frac{p}{3} \left(2\left(\frac{p}{3}\right) + 4\right) \pmod{p^2},$$

$$\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}(k+1)^2} \equiv -\frac{176}{9} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k(k+1)} + \frac{p}{9} \left(35\left(\frac{p}{3}\right) - 8\right) \pmod{p^2},$$

$$\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}(2k-1)} \equiv -\frac{1}{9} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k(k+1)} + \frac{p}{36} \left(\left(\frac{p}{3}\right) - 6\right) \pmod{p^2}.$$

Remark 2.4 Let p be a prime with $p > 3$. In [S10], the author conjectured that if $p \equiv 1, 5, 7, 11 \pmod{24}$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k} \equiv \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}}$$

$$\equiv \begin{cases} 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} & \text{if } p \equiv 1, 7 \pmod{24} \text{ and so } p = x^2 + 6y^2, \\ 8x^2 - 2p - \frac{p^2}{8x^2} \pmod{p^3} & \text{if } p \equiv 5, 11 \pmod{24} \text{ and so } p = 2x^2 + 3y^2; \end{cases}$$

if $p \equiv 13, 17, 19, 23 \pmod{24}$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k} \equiv -7\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{48^{2k}} \pmod{p^3}.$$

The corresponding congruences modulo p^2 were conjectured by Z.W. Sun in [Su1].

Conjecture 2.31. Let $p > 5$ be a prime.

(i) If $p \equiv 1, 19 \pmod{30}$ and so $p = x^2 + 15y^2$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k(k+1)} \equiv -84y^2 + 2p \pmod{p^2},$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k(k+1)^2} \equiv -96y^2 \pmod{p^2},$$

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (2k-1)} &\equiv \frac{148}{3} y^2 - \frac{26}{9} p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (k+1)} &\equiv 60y^2 + 2p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (k+1)^2} &\equiv -1320y^2 + 36p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (2k-1)} &\equiv \frac{3508}{75} y^2 - \frac{1954}{1125} p \pmod{p^2}.
\end{aligned}$$

(ii) If $p \equiv 17, 23 \pmod{30}$ and so $p = 3x^2 + 5y^2$, then

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)} &\equiv 28y^2 - 2p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)^2} &\equiv 32y^2 - 2p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (2k-1)} &\equiv -\frac{148}{9} y^2 + \frac{26}{9} p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (k+1)} &\equiv -20y^2 - 2p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (k+1)^2} &\equiv 440y^2 - 38p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (2k-1)} &\equiv -\frac{3508}{225} y^2 + \frac{1954}{1125} p \pmod{p^2}.
\end{aligned}$$

(iii) If $(\frac{-15}{p}) = -1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)} \equiv \begin{cases} \frac{2}{5} \cdot 5^{-[\frac{p}{3}]} \left(\frac{[p/3]}{[p/15]} \right)^2 \pmod{p} & \text{if } p \equiv 7 \pmod{30}, \\ \frac{1}{10} \cdot 5^{-[\frac{p}{3}]} \left(\frac{[p/3]}{[p/15]} \right)^2 \pmod{p} & \text{if } p \equiv 11 \pmod{30}, \\ \frac{32}{5} \cdot 5^{-[\frac{p}{3}]} \left(\frac{[p/3]}{[p/15]} \right)^2 \pmod{p} & \text{if } p \equiv 13 \pmod{30}, \\ \frac{8}{5} \cdot 5^{-[\frac{p}{3}]} \left(\frac{[p/3]}{[p/15]} \right)^2 \pmod{p} & \text{if } p \equiv 29 \pmod{30}. \end{cases}$$

Moreover,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)^2} \equiv \frac{13}{2} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)} + \left(3 \left(\frac{p}{3} \right) - 1 \right) p \pmod{p^2},$$

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (2k-1)} &\equiv -\frac{16}{9} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)} - \frac{8}{3} \left(\frac{p}{3}\right) p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (k+1)} &\equiv -20 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)} - 12 \left(\frac{p}{3}\right) p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (k+1)^2} &\equiv -130 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)} - \left(1 + 111 \left(\frac{p}{3}\right)\right) p \pmod{p^2}, \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k} (2k-1)} &\equiv -\frac{64}{225} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k (k+1)} - \frac{152}{375} \left(\frac{p}{3}\right) p \pmod{p^2}.
\end{aligned}$$

Conjecture 2.32. Let p be a prime with $p > 5$ and

$$t_p = \frac{\binom{(p-1)/2}{[7p/40]} \binom{(p-1)/2}{[9p/40]} \binom{[3p/40]}{[p/40]}}{\binom{[19p/40]}{[p/20]}}.$$

(i) If $\left(\frac{-10}{p}\right) = 1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k} (k+1)} \equiv \begin{cases} -\frac{49}{15} t_p \pmod{p} & \text{if } p \equiv 1, 23 \pmod{40}, \\ -\frac{49}{5} t_p \pmod{p} & \text{if } p \equiv 7 \pmod{40}, \\ \frac{7}{5} t_p \pmod{p} & \text{if } p \equiv 9 \pmod{40}, \\ \frac{833}{195} t_p \pmod{p} & \text{if } p \equiv 11 \pmod{40}, \\ -\frac{833}{285} t_p \pmod{p} & \text{if } p \equiv 13 \pmod{40}, \\ -\frac{98}{555} t_p \pmod{p} & \text{if } p \equiv 19 \pmod{40}, \\ -\frac{98}{55} t_p \pmod{p} & \text{if } p \equiv 37 \pmod{40}. \end{cases}$$

Moreover,

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k} (k+1)} &\equiv \begin{cases} -\frac{196}{15} x^2 + \frac{226}{15} p \pmod{p^2} & \text{if } p = x^2 + 10y^2 \equiv 1, 9, 11, 19 \pmod{40}, \\ \frac{392}{15} x^2 - \frac{226}{15} p \pmod{p^2} & \text{if } p = 2x^2 + 5y^2 \equiv 7, 13, 19, 37 \pmod{40}, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k} (k+1)^2} &\equiv \begin{cases} \frac{10088}{45} x^2 - \frac{6113}{45} p \pmod{p^2} & \text{if } p = x^2 + 10y^2 \equiv 1, 9, 11, 19 \pmod{40}, \\ -\frac{20176}{45} x^2 + \frac{6023}{45} p \pmod{p^2} & \text{if } p = 2x^2 + 5y^2 \equiv 7, 13, 19, 37 \pmod{40}, \end{cases} \\
\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k} (2k-1)} &
\end{aligned}$$

$$\equiv \begin{cases} -\frac{883}{270}x^2 + \frac{2393}{1620}p \pmod{p^2} & \text{if } p = x^2 + 10y^2 \equiv 1, 9, 11, 19 \pmod{40}, \\ \frac{883}{135}x^2 - \frac{2393}{1620}p \pmod{p^2} & \text{if } p = 2x^2 + 5y^2 \equiv 7, 13, 23, 37 \pmod{40} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}} \equiv \begin{cases} 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} & \text{if } p \equiv 1, 9, 11, 19 \pmod{40} \text{ and so } p = x^2 + 10y^2, \\ 2p - 8x^2 + \frac{p^2}{8x^2} \pmod{p^3} & \text{if } p \equiv 7, 13, 23, 37 \pmod{40} \text{ and so } p = 2x^2 + 5y^2. \end{cases}$$

(ii) If $(\frac{-10}{p}) = -1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}(k+1)} \equiv \begin{cases} -\frac{21}{5}t_p \pmod{p} & \text{if } p \equiv 3 \pmod{40}, \\ -\frac{4446}{155}t_p \pmod{p} & \text{if } p \equiv 17 \pmod{40}, \\ -\frac{189}{5}t_p \pmod{p} & \text{if } p \equiv 21 \pmod{40}, \\ -\frac{702}{5}t_p \pmod{p} & \text{if } p \equiv 27 \pmod{40}, \\ \frac{66}{5}t_p \pmod{p} & \text{if } p \equiv 29 \pmod{40}, \\ \frac{1026}{5}t_p \pmod{p} & \text{if } p \equiv 31 \pmod{40}, \\ -\frac{462}{5}t_p \pmod{p} & \text{if } p \equiv 33 \pmod{40}, \\ -\frac{858}{85}t_p \pmod{p} & \text{if } p \equiv 39 \pmod{40}. \end{cases}$$

Moreover,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}(k+1)^2} \equiv \frac{22}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}(k+1)} + \left(\frac{256}{3} \left(\frac{p}{5} \right) - 1 \right)p \pmod{p^2},$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}(2k-1)} \equiv \frac{1}{216} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}(k+1)} + \frac{16}{81} \left(\frac{p}{5} \right)p \pmod{p^2}$$

and

$$\left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}(k+1)} \right) \left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}} \right) \equiv -\frac{49}{15}p^2 \pmod{p^3}.$$

Remark 2.5 Let $p > 3$ be a prime. In [Su1], Z.W. Sun conjectured the congruence for $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{12^{4k}} \pmod{p^2}$.

Conjecture 2.33. Let p be a prime with $p \neq 2, 11$ and $r_p = \binom{[3p/11]}{[p/11]} \binom{[6p/11]}{[3p/11]} / \binom{[4p/11]}{[2p/11]}$.

(i) If $p \equiv 1, 3, 4, 5, 9 \pmod{11}$ and so $4p = x^2 + 11y^2$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} \equiv \begin{cases} \frac{25}{22} r_p^2 \pmod{p} & \text{if } p \equiv 1, 4, 5, 9 \pmod{11}, \\ \frac{2}{11} r_p^2 \pmod{p} & \text{if } p \equiv 3 \pmod{11}. \end{cases}$$

Moreover,

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} &\equiv -\frac{25}{2} y^2 + 2p \pmod{p^2}, \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)^2} &\equiv -\frac{83}{4} y^2 + 2p \pmod{p^2}, \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(2k-1)} &\equiv \frac{23}{4} y^2 - \frac{7}{8} p \pmod{p^2}, \\ \left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}(k+1)} &\equiv -26y^2 + 2p \pmod{p^2}, \\ \left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}(k+1)^2} &\equiv 148y^2 - \left(24 + \left(\frac{-2}{p}\right)\right)p \pmod{p^2}, \\ \left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}(2k-1)} &\equiv \frac{73}{8} y^2 - \frac{467}{256} p \pmod{p^2}. \end{aligned}$$

(ii) If $p \equiv 2, 6, 7, 8, 10 \pmod{11}$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} \equiv \begin{cases} -\frac{50}{11} r_p^2 \pmod{p} & \text{if } p \equiv 2 \pmod{11}, \\ -\frac{32}{11} r_p^2 \pmod{p} & \text{if } p \equiv 6 \pmod{11}, \\ -\frac{2}{11} r_p^2 \pmod{p} & \text{if } p \equiv 7 \pmod{11}, \\ -\frac{72}{11} r_p^2 \pmod{p} & \text{if } p \equiv 8 \pmod{11}, \\ -\frac{18}{11} r_p^2 \pmod{p} & \text{if } p \equiv 10 \pmod{11}. \end{cases}$$

Moreover,

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)^2} &\equiv \frac{13}{2} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} \pmod{p^2}, \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(2k-1)} &\equiv \frac{3}{4} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} + \frac{3}{8} p \pmod{p^2}, \\ \left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}(k+1)} &\equiv \frac{128}{15} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} - \frac{2}{5} p \pmod{p^2}, \end{aligned}$$

$$\left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}(k+1)^2} \equiv \frac{5888}{75} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} + \left(\frac{608}{25} - \left(\frac{-2}{p}\right)\right)p \pmod{p^2},$$

$$\left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}(2k-1)} \equiv -\frac{1}{8} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)} - \frac{51}{256}p \pmod{p^2}$$

and

$$\left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k(k+1)}\right) \left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k}\right) \equiv \frac{25}{22}p^2 \pmod{p^3}.$$

Remark 2.6 Let p be a prime with $p \neq 2, 3, 11, 13$. In [S10] the author conjectured that if $p \equiv 1, 3, 4, 5, 9 \pmod{11}$ and so $4p = x^2 + 11y^2$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k} \equiv \left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}} \equiv x^2 - 2p - \frac{p^2}{x^2} \pmod{p^3};$$

if $p \equiv 2, 6, 7, 8, 10 \pmod{11}$ and $f = [\frac{p}{11}]$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k} \equiv \frac{160}{39} \left(\frac{-2}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}}$$

$$\equiv \begin{cases} -p^2 \left(\frac{5 \binom{4f}{2f}}{2 \binom{3f}{f} \binom{6f}{3f}}\right)^2 \pmod{p^3} & \text{if } p \equiv 2 \pmod{11}, \\ -p^2 \left(\frac{13 \binom{4f}{2f}}{30 \binom{3f}{f} \binom{6f}{3f}}\right)^2 \pmod{p^3} & \text{if } p \equiv 6 \pmod{11}, \\ -p^2 \left(\frac{85 \binom{4f}{2f}}{558 \binom{3f}{f} \binom{6f}{3f}}\right)^2 \pmod{p^3} & \text{if } p \equiv 7 \pmod{11}, \\ -p^2 \left(\frac{7 \binom{4f}{2f}}{148 \binom{3f}{f} \binom{6f}{3f}}\right)^2 \pmod{p^3} & \text{if } p \equiv 8 \pmod{11}, \\ -p^2 \left(\frac{29 \binom{4f}{2f}}{756 \binom{3f}{f} \binom{6f}{3f}}\right)^2 \pmod{p^3} & \text{if } p \equiv 10 \pmod{11}. \end{cases}$$

The corresponding congruence modulo p^2 was conjectured by Z.W. Sun [Su3].

Conjecture 2.34. Let $p > 3$ be a prime and $s_p = \binom{[8p/19]}{[p/19]} \binom{[10p/19]}{[4p/19]} / \binom{[5p/19]}{[2p/19]}$.

(i) If $(\frac{p}{19}) = 1$ and so $4p = x^2 + 19y^2$, then

$$\left(\frac{-6}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}(2k-1)} \equiv \begin{cases} -\frac{1183}{1368} s_p^2 \pmod{p} & \text{if } p \equiv 1, 7, 11 \pmod{19}, \\ -\frac{1183}{342} s_p^2 \pmod{p} & \text{if } p \equiv 4 \pmod{19}, \\ -\frac{1183}{2432} s_p^2 \pmod{p} & \text{if } p \equiv 5, 6 \pmod{19}, \\ -\frac{1183}{18392} s_p^2 \pmod{p} & \text{if } p \equiv 9 \pmod{19}, \\ -\frac{1183}{27702} s_p^2 \pmod{p} & \text{if } p \equiv 16 \pmod{19}, \\ -\frac{57967}{12312} s_p^2 \pmod{p} & \text{if } p \equiv 17 \pmod{19}. \end{cases}$$

Moreover,

$$\begin{aligned} \left(\frac{-6}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}(2k-1)} &\equiv \frac{1183}{72} y^2 - \frac{4273}{2304} p \pmod{p^2}, \\ \left(\frac{-6}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}(k+1)} &\equiv -394y^2 + 2p \pmod{p^2}, \\ \left(\frac{-6}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}(k+1)^2} &\equiv 6772y^2 - \left(536 + \left(\frac{-6}{p}\right)\right)p \pmod{p^2}. \end{aligned}$$

(ii) If $(\frac{p}{19}) = -1$, then

$$\left(\frac{-6}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}(2k-1)} \equiv \begin{cases} \frac{49}{228} s_p^2 \pmod{p} & \text{if } p \equiv 2 \pmod{19}, \\ \frac{1}{228} s_p^2 \pmod{p} & \text{if } p \equiv 3 \pmod{19}, \\ \frac{27}{14896} s_p^2 \pmod{p} & \text{if } p \equiv 8 \pmod{19}, \\ \frac{3}{19} s_p^2 \pmod{p} & \text{if } p \equiv 10 \pmod{19}, \\ \frac{121}{57} s_p^2 \pmod{p} & \text{if } p \equiv 12 \pmod{19}, \\ \frac{4}{57} s_p^2 \pmod{p} & \text{if } p \equiv 13 \pmod{19}, \\ \frac{121}{228} s_p^2 \pmod{p} & \text{if } p \equiv 14 \pmod{19}, \\ \frac{27}{76} s_p^2 \pmod{p} & \text{if } p \equiv 15 \pmod{19}, \\ \frac{49}{57} s_p^2 \pmod{p} & \text{if } p \equiv 18 \pmod{19}. \end{cases}$$

Moreover,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}(k+1)} \equiv -\frac{9216}{5} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}(2k-1)} - 270 \left(\frac{-6}{p}\right) p \pmod{p^2},$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k} (k+1)^2} \equiv \frac{46}{5} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k} (k+1)} + \left(540 \left(\frac{-6}{p} \right) - 1 \right) p \pmod{p^2}$$

and

$$\left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k} (2k-1)} \right) \left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}} \right) \equiv -\frac{985}{87552} p^2 \pmod{p^3}.$$

Conjecture 2.35. Let $p > 5$ be a prime. If $(\frac{p}{43}) = 1$ and so $4p = x^2 + 43y^2$, then

$$\left(\frac{-15}{p} \right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-960)^{3k} (2k-1)} \equiv \frac{140501}{3600} y^2 - \frac{4384321}{2304000} p \pmod{p^2}.$$

If $(\frac{p}{43}) = -1$, then

$$\left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-960)^{3k} (2k-1)} \right) \left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-960)^{3k}} \right) \equiv -\frac{933889}{198144000} p^2 \pmod{p^3}.$$

Conjecture 2.36. Let $p > 5$ be a prime. If $(\frac{p}{67}) = -1$, then

$$\left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-5280)^{3k} (2k-1)} \right) \left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-5280)^{3k}} \right) \equiv -\frac{155357161}{51365952000} p^2 \pmod{p^3}.$$

Remark 2.7 Suppose that $p > 3$ is a prime. In [Su3], Z.W. Sun conjectured that

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-32)^{3k}} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } (\frac{p}{11}) = -1, \\ (\frac{-2}{p})(x^2 - 2p) \pmod{p^2} & \text{if } (\frac{p}{11}) = 1 \text{ and so } 4p = x^2 + 11y^2. \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k}} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } (\frac{p}{19}) = -1, \\ (\frac{-6}{p})(x^2 - 2p) \pmod{p^2} & \text{if } (\frac{p}{19}) = 1 \text{ and so } 4p = x^2 + 19y^2, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-960)^{3k}} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } (\frac{p}{43}) = -1 \text{ and } p \neq 5, \\ (\frac{p}{15})(x^2 - 2p) \pmod{p^2} & \text{if } (\frac{p}{43}) = 1 \text{ and so } 4p = x^2 + 43y^2, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-5280)^{3k}} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } (\frac{p}{67}) = -1 \text{ and } p \neq 5, 11, \\ (\frac{-330}{p})(x^2 - 2p) \pmod{p^2} & \text{if } (\frac{p}{67}) = 1 \text{ and so } 4p = x^2 + 67y^2 \end{cases} \end{aligned}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-640320)^{3k}} \equiv \begin{cases} 0 \pmod{p^2} & \text{if } (\frac{p}{163}) = -1 \text{ and } p \neq 5, 23, 29, \\ (\frac{-10005}{p})(x^2 - 2p) \pmod{p^2} & \text{if } (\frac{p}{163}) = 1 \text{ and so } 4p = x^2 + 163y^2. \end{cases}$$

These congruences modulo p were proved by the author in [S4]

Conjecture 2.37. Let p be an odd prime. If $p \equiv 1 \pmod{4}$ and so $p = x^2 + 4y^2$ with $4 \mid x - 1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{32^k(2k-1)^2} \equiv x - \frac{p}{4x} \pmod{p^2} \quad \text{and} \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{32^k(k+1)^2} \equiv 8x - 7 \pmod{p}.$$

If $p \equiv 3 \pmod{4}$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{32^k(2k-1)^2} \equiv \frac{1}{2}(2p+3-2^{p-1}) \binom{\frac{p-1}{2}}{\frac{p-3}{4}} \pmod{p^2}.$$

Conjecture 2.38. Let p be an odd prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{(-16)^k(2k-1)^2} \equiv \begin{cases} (-1)^{\frac{p-1}{4}} \frac{p}{x} \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4} \text{ and } 4 \mid x - 1, \\ 2(-1)^{\frac{p+1}{4}} \binom{(p-1)/2}{(p-3)/4} \pmod{p} & \text{if } p \equiv 3 \pmod{4} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{(-16)^k(k+1)^2} \equiv 5 \pmod{p} \quad \text{for } p \equiv 1 \pmod{4}.$$

Conjecture 2.39. Let p be an odd prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{8^k(2k-1)^2} \equiv \begin{cases} 2(-1)^{\frac{p-1}{4}} x \pmod{p} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4} \text{ and } 4 \mid x - 1, \\ 2(-1)^{\frac{p+1}{4}} \binom{\frac{p-1}{2}}{\frac{p-3}{4}} \pmod{p} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Remark 2.8 Let p be an odd prime. In [Su4], Z.W. Sun established the congruences for $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{m^k(2k-1)} \pmod{p^2}$ in the cases $m = -16, 8, 32$.

Conjecture 2.40. Let $a, x \in \mathbb{Q}$ and $d \in \mathbb{Z}$ with $adx \neq 0$, where \mathbb{Q} is the set of rational numbers. Suppose that $\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} x^k \equiv 0 \pmod{p}$ for all odd primes p satisfying $a, x \in \mathbb{Z}_p$ and $(\frac{d}{p}) = -1$. Then there is a constant $c \in \mathbb{Q}$ such that for all odd primes p with $c \in \mathbb{Z}_p$ and $(\frac{d}{p}) = -1$,

$$\sum_{k=0}^{p-1} \binom{2k}{k} \binom{a}{k} \binom{-1-a}{k} (x(1-x))^k \equiv c \left(\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} x^k \right)^2 \pmod{p^3}.$$

Remark 2.9 By [S6], for any odd prime p and $a, x \in \mathbb{Z}_p$,

$$\left(\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} x^k \right)^2 \equiv \sum_{k=0}^{p-1} \binom{2k}{k} \binom{a}{k} \binom{-1-a}{k} (x(1-x))^k \pmod{p^2}.$$

Conjecture 2.41. Let $a, m \in \mathbb{Q}$ and $d \in \mathbb{Z}$ with $adm \neq 0$. Suppose that

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{a}{k} \binom{-1-a}{k}}{m^k} \equiv 0 \pmod{p^2}$$

for all odd primes p satisfying $a, m \in \mathbb{Z}_p$, $p \nmid m$ and $(\frac{d}{p}) = -1$. Then there is a constant $c \in \mathbb{Q}$ such that for all odd primes p with $c, m \in \mathbb{Z}_p$, $p \nmid m$ and $(\frac{d}{p}) = -1$,

$$\left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{a}{k} \binom{-1-a}{k}}{(2k-1)m^k} \right) \left(\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{a}{k} \binom{-1-a}{k}}{m^k} \right) \equiv cp^2 \pmod{p^3}.$$

Moreover, there is a constant $C \in \mathbb{Q}$ such that for all odd primes p with $C, m \in \mathbb{Z}_p$, $p \nmid m$ and $(\frac{d}{p}) = 1$,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{a}{k} \binom{-1-a}{k}}{(2k-1)m^k} \equiv C \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{a}{k} \binom{-1-a}{k}}{m^k} \pmod{p}.$$

3. Conjectures for congruences involving Apéry-like numbers

With the help of Maple, we discover the following conjectures involving Apéry-like numbers.

Conjecture 3.1. Let p be an odd prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{16^k (k+1)} &\equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ 0 \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{16^k (2k-1)} &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } p \equiv 1 \pmod{4}, \\ -R_1(p) \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

Conjecture 3.2. Let $p > 3$ be a prime. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{32^k (2k-1)} &\equiv \begin{cases} (-1)^{\frac{p-1}{2}} \left(\frac{p}{2} - x^2\right) \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{5(-1)^{\frac{p-1}{2}} - 4}{4} R_2(p) \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{800^k (2k-1)} &\equiv \begin{cases} -\frac{73}{100} (-1)^{\frac{p-1}{2}} (4x^2 - 2p) - \frac{24}{125} \left(\frac{3}{p}\right)p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{45(-1)^{\frac{p-1}{2}} - 36}{100} R_2(p) + \frac{24}{125} \left(\frac{3}{p}\right)p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8} \text{ and } p \neq 5, \end{cases} \\ \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{(-768)^k (2k-1)} &\equiv \begin{cases} -\frac{73}{96} \left(\frac{3}{p}\right) (4x^2 - 2p) - \frac{11}{48} (-1)^{\frac{p-1}{2}} p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{15 - 12(-1)^{\frac{p-1}{2}}}{32} \left(\frac{3}{p}\right) R_2(p) - (-1)^{\frac{p-1}{2}} \frac{11}{48} p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases} \end{aligned}$$

Conjecture 3.3. Let p be an odd prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{7^k(2k-1)} \equiv \begin{cases} \frac{124}{49}x^2 - \frac{46}{49}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{64}{7} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1} + \frac{496}{49}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{25^k(2k-1)} \equiv \begin{cases} -\frac{124}{175}x^2 + \frac{326}{875}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{448}{175} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1} + \frac{2576}{875}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7} \text{ and } p > 7. \end{cases}$$

Conjecture 3.4. Let $p > 3$ be a prime. Then

$$(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{(-32)^k(2k-1)} \equiv \begin{cases} -\frac{11}{3}x^2 + \frac{7}{6}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ \frac{22}{3}x^2 - \frac{7}{6}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}, \\ \frac{2}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k(k+1)} + \frac{p}{2} \pmod{p^2} & \text{if } p \equiv 13, 19 \pmod{24}, \\ -\frac{2}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k(k+1)} - \frac{5}{6}p \pmod{p^2} & \text{if } p \equiv 17, 23 \pmod{24} \end{cases}$$

and

$$(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} S_k}{64^k(2k-1)} \equiv \begin{cases} -\frac{11}{6}x^2 + \frac{5}{6}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ -\frac{11}{3}x^2 + \frac{5}{6}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}, \\ -\frac{1}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k(k+1)} \pmod{p^2} & \text{if } p \equiv 13, 19 \pmod{24}, \\ -\frac{1}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k(k+1)} - \frac{p}{6} \pmod{p^2} & \text{if } p \equiv 17, 23 \pmod{24}. \end{cases}$$

Conjecture 3.5. Let p be a prime with $p > 3$. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{(-12)^k(2k-1)} \equiv \begin{cases} 0 \pmod{p^2} & \text{if } p \equiv 1 \pmod{3}, \\ -2(2p+1) \left(\frac{[2p/3]}{[p/3]} \right)^2 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Conjecture 3.6. Let p be a prime with $p > 3$. Then

$$\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{54^k(2k-1)} \equiv \begin{cases} -\frac{28}{9}x^2 + \frac{10}{9}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ \frac{2}{3}R_1(p) - \frac{4}{9}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Conjecture 3.7. Let p be an odd prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{8^k (2k-1)} \equiv \begin{cases} -\frac{11}{2}x^2 + \frac{5}{4}p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{45 - 36(-1)^{\frac{p-1}{2}}}{8} R_2(p) + \frac{3}{2}p \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Conjecture 3.8. Let p be a prime with $p \neq 2, 3, 7$. Then

$$\begin{aligned} \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{(-27)^k (2k-1)} \\ \equiv \begin{cases} -\frac{2}{3}p + \frac{76}{9}y^2 \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -\frac{8}{3} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1} - \frac{28}{9}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7} \end{cases} \end{aligned}$$

and

$$\begin{aligned} \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{243^k (2k-1)} \\ \equiv \begin{cases} \frac{1676}{81}y^2 - \frac{142}{81}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ -\frac{32}{27} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1} - \frac{44}{27}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}. \end{cases} \end{aligned}$$

Conjecture 3.9. Let p be a prime with $p \neq 2, 11$. Then

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{(-44)^k (2k-1)} \\ \equiv \begin{cases} \frac{52}{11}y^2 - \frac{116}{121}p \pmod{p^2} & \text{if } p \equiv 1, 3, 4, 5, 9 \pmod{11} \text{ and so } 4p = x^2 + 11y^2, \\ \frac{9}{11} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k (k+1)} + \frac{39}{121}p \pmod{p^2} & \text{if } p \equiv 2, 6, 7, 8, 10 \pmod{11} \end{cases} \end{aligned}$$

Conjecture 3.10. Let p be a prime with $p \neq 2, 3, 19$. Then

$$\begin{aligned} \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} W_k}{(-108)^k (2k-1)} \\ \equiv \begin{cases} -152y^2 + \frac{4849}{288}p \pmod{p^2} & \text{if } \left(\frac{p}{19}\right) = 1 \text{ and so } 4p = x^2 + 19y^2, \\ 8\left(\frac{-6}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k} (2k-1)} + \frac{241}{288}p \pmod{p^2} & \text{if } \left(\frac{p}{19}\right) = -1. \end{cases} \end{aligned}$$

Conjecture 3.11. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{(-4)^k (2k-1)} \equiv \begin{cases} 2p - 4x^2 \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -8R_3(p) \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases}$$

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{50^k (2k-1)} \\ & \equiv \begin{cases} -\frac{13}{25}(4x^2 - 2p) - \frac{12}{125}(-1)^{\frac{p-1}{2}} p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{32}{25}R_3(p) - \frac{12}{125}(-1)^{\frac{p-1}{2}} p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3} \text{ and } p \neq 5. \end{cases} \end{aligned}$$

Conjecture 3.12. Let $p > 3$ be a prime. Then

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{96^k (2k-1)} \\ & \equiv \begin{cases} -\frac{29}{48} \left(\frac{p}{3}\right) (4x^2 - 2p) - \frac{p}{6} \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ \frac{15 - 12(-1)^{(p-1)/2}}{16} \left(\frac{p}{3}\right) R_2(p) + \frac{p}{6} \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases} \end{aligned}$$

Conjecture 3.13. Let p be a prime with $p \neq 2, 5$. If $(\frac{-5}{p}) = 1$, then

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{16^k (2k-1)} \\ & \equiv \begin{cases} -\frac{7}{5}x^2 + \frac{9}{10}p \pmod{p^2} & \text{if } p \equiv 1, 9 \pmod{20} \text{ and so } p = x^2 + 5y^2, \\ \frac{14}{5}x^2 - \frac{p}{2} \pmod{p^2} & \text{if } p \equiv 3, 7 \pmod{20} \text{ and so } 2p = x^2 + 5y^2; \end{cases} \end{aligned}$$

if $(\frac{-5}{p}) = -1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{16^k (2k-1)} \equiv -6(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-1024)^k (2k-1)} - \frac{11}{8}p \pmod{p^2}.$$

Conjecture 3.14. Let $p > 3$ be a prime. If $(\frac{-6}{p}) = 1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{32^k (2k-1)} \equiv \begin{cases} -\frac{7}{4}x^2 + \frac{7}{8}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ -\frac{7}{2}x^2 + \frac{7}{8}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}; \end{cases}$$

if $(\frac{-6}{p}) = -1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{32^k (2k-1)} \equiv \frac{9}{4} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (2k-1)} + \frac{1}{4} \left(\frac{p}{3}\right)p \pmod{p^2}.$$

Conjecture 3.15. Let $p > 3$ be a prime. Then

$$(-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} a_k}{54^k (2k-1)}$$

$$\equiv \begin{cases} \frac{52}{9}y^2 - \frac{26 + 2(-1)^{\frac{p-1}{2}}}{27}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ \frac{32}{27}R_3(p) + \frac{2}{27}(-1)^{\frac{p-1}{2}}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Conjecture 3.16. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} a_k}{100^k (2k-1)} \equiv \begin{cases} -\frac{58}{25}x^2 + \frac{145 - 18(\frac{p}{3})}{125}p \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{9}{50}R_2(p) - \frac{18}{125}(\frac{p}{3})p \pmod{p^2} & \text{if } p \equiv 5 \pmod{8} \text{ and } p \neq 5, \\ -\frac{81}{50}R_2(p) - \frac{18}{125}(\frac{p}{3})p \pmod{p^2} & \text{if } p \equiv 7 \pmod{8}. \end{cases}$$

Conjecture 3.17. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} a_k}{(-12)^k (2k-1)} \equiv \begin{cases} p - 4x^2 \pmod{p^2} & \text{if } 12 \mid p-1 \text{ and so } p = x^2 + 9y^2, \\ 2x^2 - p \pmod{p^2} & \text{if } 12 \mid p-5 \text{ and so } 2p = x^2 + 9y^2, \\ 3 \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 7 \pmod{12}, \\ -6 \left(\frac{[p/3]}{[p/12]} \right)^2 \pmod{p} & \text{if } p \equiv 11 \pmod{12}. \end{cases}$$

Conjecture 3.18. Let $p > 3$ be a prime. If $(\frac{-6}{p}) = 1$, then

$$\left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} a_k}{36^k (2k-1)} \equiv \begin{cases} -\frac{14}{9}x^2 + \frac{7}{9}p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1, 7 \pmod{24}, \\ -\frac{28}{9}x^2 + \frac{7}{9}p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}; \end{cases}$$

if $(\frac{-6}{p}) = -1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} a_k}{36^k (2k-1)} \equiv -2 \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k (2k-1)} - \frac{2}{9}p \pmod{p^2}.$$

Conjecture 3.19. Let p be an odd prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{64^k (k+1)} \equiv (-1)^{\frac{p-1}{2}} \pmod{p^2} \text{ and } \sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{64^k (2k-1)} \equiv 2(-1)^{\frac{p-1}{2}} p^2 \pmod{p^3}.$$

Conjecture 3.20. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{72^k (2k-1)} \equiv \begin{cases} -\frac{4}{9}x^2 + \frac{8}{27}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{2}{9}R_1(p) - \frac{2}{27}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{576^k (2k-1)} \equiv \begin{cases} -\frac{8}{3}x^2 + \frac{32}{27}p \pmod{p^2} & \text{if } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \\ -\frac{2}{9}R_1(p) + \frac{4}{27}p \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Conjecture 3.21. Let p be an odd prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{128^k (2k-1)} \equiv \begin{cases} -\frac{3}{8}(4x^2 - 2p) \pmod{p^2} & \text{if } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ -\frac{5 - 4(-1)^{(p-1)/2}}{8} R_2(p) \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Conjecture 3.22. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{48^k (2k-1)} \equiv \begin{cases} \frac{4}{9}x^2 \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{8}{9}R_3(p) - \frac{2}{9}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{(-192)^k (2k-1)} \equiv \begin{cases} -\frac{32}{9}x^2 + \frac{4}{3}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{8}{9}R_3(p) + \frac{4}{9}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Conjecture 3.23. Let p be a prime with $p \neq 2, 3, 7$. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{63^k (2k-1)} \equiv \begin{cases} \frac{4}{7}y^2 + \frac{26}{1323}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{32}{63} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1} + \frac{1024}{1323}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} G_k}{(-4032)^k (2k-1)} \equiv \begin{cases} \frac{1408}{63}y^2 - \frac{2368}{1323}p \pmod{p^2} & \text{if } p = x^2 + 7y^2 \equiv 1, 2, 4 \pmod{7}, \\ \frac{32}{63} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^3}{k+1} + \frac{688}{1323}p \pmod{p^2} & \text{if } p \equiv 3, 5, 6 \pmod{7}. \end{cases}$$

Conjecture 3.24. Let $p > 3$ be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} Q_k}{(-36)^k (2k-1)} \equiv \begin{cases} -\frac{4}{9}x^2 + \frac{2}{9}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{8}{9}R_3(p) \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} Q_k}{18^k (2k-1)} \equiv \begin{cases} -\frac{52}{9}x^2 + \frac{26 - 12(-1)^{(p-1)/2}}{9}p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ -\frac{32}{9}R_3(p) - \frac{4}{3}(-1)^{\frac{p-1}{2}}p \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Conjecture 3.25. Let p be a prime with $p \neq 2, 3, 11$.

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} A'_k}{4^k (2k-1)}$$

$$\equiv \begin{cases} 2p - 2y^2 \pmod{p^2} & \text{if } p \equiv 1, 3, 4, 5, 9 \pmod{11} \text{ and so } 4p = x^2 + 11y^2, \\ 5 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k (k+1)} + 3p \pmod{p^2} & \text{if } p \equiv 2, 6, 7, 8, 10 \pmod{11}. \end{cases}$$

Conjecture 3.26. Let p be a prime with $p \neq 2, 3, 19$. If $(\frac{p}{19}) = 1$ and so $4p = x^2 + 19y^2$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} A'_k}{36^k (2k-1)} \equiv \frac{74}{9} y^2 - \frac{22}{27} p \pmod{p^2}.$$

If $(\frac{p}{19}) = -1$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} A'_k}{36^k (2k-1)} \equiv -\frac{40}{3} \left(\frac{-6}{p}\right) \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k} \binom{6k}{3k}}{(-96)^{3k} (2k-1)} \pmod{p}.$$

Remark 3.1 In [Su3], Z.W. Sun conjectured that for any prime $p > 3$,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} A'_k}{4^k} \equiv \begin{cases} x^2 - 2p \pmod{p^2} & \text{if } (\frac{p}{11}) = 1 \text{ and so } 4p = x^2 + 11y^2, \\ 0 \pmod{p^2} & \text{if } (\frac{p}{11}) = -1, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} A'_k}{36^k} \equiv \begin{cases} x^2 - 2p \pmod{p^2} & \text{if } (\frac{p}{19}) = 1 \text{ and so } 4p = x^2 + 19y^2, \\ 0 \pmod{p^2} & \text{if } (\frac{p}{19}) = -1. \end{cases}$$

References

- [A] S. Ahlgren, *Gaussian hypergeometric series and combinatorial congruences*, in: Symbolic computation, number theory, special functions, physics and combinatorics (Gainesville, FL, 1999), pp. 1-12, Dev. Math., Vol. 4, Kluwer, Dordrecht, 2001.
- [AZ] G. Almkvist and W. Zudilin, *Differential equations, mirror maps and zeta values*, in: Mirror Symmetry AMS/IP Stud. Adv. Math. **38**(2006), International Press & Amer. Math. Soc., 481-515.
- [B] F. Beukers, *Another congruence for the Apéry numbers*, J. Number Theory **25**(1987), 201-210.
- [I] T. Ishikawa, *Super congruence for the Apéry numbers*, Nagoya Math. J. **118**(1990), 195-202.
- [M] E. Mortenson, *Supercongruences for truncated ${}_n+1F_n$ hypergeometric series with applications to certain weight three newforms*, Proc. Amer. Math. Soc. **133**(2005), 321-330.
- [RV] F. Rodriguez-Villegas, *Hypergeometric families of Calabi-Yau manifolds*, in: Calabi-Yau Varieties and Mirror Symmetry (Yui, Noriko (ed.) et al., Toronto, ON, 2001), 223-231, Fields Inst. Commun., 38, Amer. Math. Soc., Providence, RI, 2003.

- [S1] Z. H. Sun, *Congruences concerning Legendre polynomials*, Proc. Amer. Math. Soc. **139**(2011), 1915-1929.
- [S2] Z. H. Sun, *Congruences concerning Legendre polynomials II*, J. Number Theory **133**(2013), 1950-1976.
- [S3] Z. H. Sun, *Congruences involving $\binom{2k}{k}^2 \binom{3k}{k}$* , J. Number Theory **133**(2013), 1572-1595.
- [S4] Z. H. Sun, *Legendre polynomials and supercongruences*, Acta Arith. **159**(2013), 169-200.
- [S5] Z. H. Sun, *Congruences for Domb and Almkvist-Zudilin numbers*, Integral Transforms Spec. Funct. **26**(2015), 642-659.
- [S6] Z. H. Sun, *Note on super congruences modulo p^2* , arXiv:1503.0341, 2015.
- [S7] Z.H. Sun, *Identities and congruences for Catalan-Larcombe-French numbers*, Int. J. Number Theory **13**(2017), 835-851.
- [S8] Z.H. Sun, *Congruences for sums involving Franel numbers*, Int. J. Number Theory **14**(2018), 123-142.
- [S9] Z.H. Sun, *Super congruences for two Apéry-like sequences*, J. Difference Equ. Appl. **24**(2018), 1685-1713.
- [S10] Z.H. Sun, *Congruences involving binomial coefficients and Apéry-like numbers*, Publ. Math. Debrecen **96**(2020), 315-346.
- [S11] Z.H. Sun, *New congruences involving Apéry-like numbers*, arXiv:2004.07172.
- [S12] Z.H. Sun, *Congruences for two types of Apéry-like sequences*, arXiv:2005.02081.
- [Su1] Z.W. Sun, *Super congruences and Euler numbers*, Sci. China Math. **54**(2011), 2509-2535.
- [Su2] Z.W. Sun, *On sums involving products of three binomial coefficients*, Acta Arith. **156**(2012), 123-141.
- [Su3] Z.W. Sun, *Conjectures and results on $x^2 \bmod p^2$ with $4p = x^2 + dy^2$* , in: Number Theory and Related Areas (eds., Y. Ouyang, C. Xing, F. Xu and P. Zhang), Higher Education Press & International Press, Beijing and Boston, 2013, pp.149-197.
- [Su4] Z.W. Sun, *Two new kinds of numbers and related divisibility results*, Colloq. Math. **154**(2018), 241-273.
- [vH] L. van Hamme, *Proof of a conjecture of Beukers on Apéry numbers*, Proceedings of the Conference on p-adic Analysis (N. De Grande-De Kimpe and L. van Hamme, ed., Houthalen, 1987), pp. 189-195, Vrije Univ. Brussel, Brussels, 1986.

- [Z] D. Zagier, *Integral solutions of Apéry-like recurrence equations*, in: Groups and Symmetries: From Neolithic Scots to John McKay, J. Harnad and P. Winternitz, eds., CRM Proceedings and Lecture Notes, Vol. 47 (American Mathematical Society, Providence, RI, 2009), pp.349-366.