

A SYSTOLIC INEQUALITY WITH REMAINDER IN THE REAL PROJECTIVE PLANE

MIKHAIL G. KATZ AND TAHL NOWIK

ABSTRACT. The first paper in systolic geometry was published by Loewner’s student P. M. Pu over half a century ago. Pu proved an inequality relating the systole and the area of an arbitrary metric on the real projective plane. We prove a stronger version of Pu’s systolic inequality with a remainder term.

1. INTRODUCTION

Loewner’s systolic inequality for the torus and Pu’s inequality [1] for the real projective plane were historically the first results in systolic geometry. Great stimulus was provided in 1983 by Gromov’s paper [2], and later by his book [3].

Our goal is to prove a strengthened version with a remainder term of Pu’s systolic inequality $\text{sys}^2(g) \leq \frac{\pi}{2} \text{area}(g)$ (for an arbitrary metric g on \mathbb{RP}^2), analogous to Bonnesen’s inequality $L^2 - 4\pi A \geq \pi^2(R - r)^2$, where L is the length of a Jordan curve in the plane, A is the area of the region bounded by the curve, R is the circumradius and r is the inradius.

Note that both the original proof in Pu ([1], 1952) and the one given by Berger ([4], 1965, pp. 299–305) proceed by averaging the metric and showing that the averaging process decreases the area and increases the systole. Such an approach involves a 5-dimensional integration (instead of a 3-dimensional one given here) and makes it harder to obtain an explicit expression for a remainder term. Analogous results for the torus were obtained in [5] with generalisations in [6] - [17].

2. THE RESULTS

We define a closed 3-dimensional manifold $M \subseteq \mathbb{R}^3 \times \mathbb{R}^3$ by setting

$$M = \{(v, w) \in \mathbb{R}^3 \times \mathbb{R}^3 : v \cdot v = 1, w \cdot w = 1, v \cdot w = 0\}$$

2010 *Mathematics Subject Classification.* Primary 53C23; Secondary 53A30 .

Key words and phrases. Systole; geometric inequality; Riemannian submersion; Cauchy-Schwarz theorem; probabilistic variance.

where $v \cdot w$ is the scalar product on \mathbb{R}^3 . We have a diffeomorphism $M \rightarrow SO(3, \mathbb{R})$, $(v, w) \mapsto (v, w, v \times w)$, where $v \times w$ is the vector product on \mathbb{R}^3 . Given a point $(v, w) \in M$, the tangent space $T_{(v,w)}M$ can be identified by differentiating the three defining equations of M along a path through (v, w) . Thus

$$T_{(v,w)}M = \{(X, Y) \in \mathbb{R}^3 \times \mathbb{R}^3 : X \cdot v = 0, Y \cdot w = 0, X \cdot w + Y \cdot v = 0\}.$$

We define a Riemannian metric g_M on M as follows. Given a point $(v, w) \in M$, let $n = v \times w$ and declare the basis $(0, n), (n, 0), (w, -v)$ of $T_{(v,w)}M$ to be orthonormal. This metric is a modification of the metric restricted to M from $\mathbb{R}^3 \times \mathbb{R}^3 = \mathbb{R}^6$. Namely, with respect to the Euclidean metric on \mathbb{R}^6 the above three vectors are orthogonal and the first two have length 1. However, the third vector has Euclidean length $\sqrt{2}$, whereas we have defined its length to be 1. Thus if $A \subseteq T_{(v,w)}M$ denotes the span of $(0, n)$ and $(n, 0)$, and $B \subseteq T_{(v,w)}M$ is spanned by $(w, -v)$, then the metric g_M on M is obtained from the Euclidean metric g on \mathbb{R}^6 (viewed as a quadratic form) as follows:

$$g_M = g \downarrow_A + \frac{1}{2} g \downarrow_B. \quad (2.1)$$

Each of the natural projections $p, q: M \rightarrow S^2$ given by $p(v, w) = v$ and $q(v, w) = w$, exhibits M as a circle bundle over S^2 .

Lemma 2.1. *The maps p and q on (M, g_M) are Riemannian submersions, over the unit sphere $S^2 \subseteq \mathbb{R}^3$.*

Proof. For the projection p , given $(v, w) \in M$, the vector $(0, n)$ as defined above is tangent to the fiber $p^{-1}(v)$. The other two vectors, $(n, 0)$ and $(w, -v)$, are thus an orthonormal basis for the subspace of $T_{(v,w)}M$ normal to the fiber, and are mapped by dp to the orthonormal basis n, w of $T_v S^2$. \square

The projection p maps the fiber $q^{-1}(w)$ onto a great circle of S^2 . This map preserves length since the unit vector $(n, 0)$, tangent to the fiber $q^{-1}(w)$ at (v, w) , is mapped by dp to the unit vector $n \in T_v S^2$. The same comments apply when the roles of p and q are reversed.

In the following proposition, integration takes place respectively over great circles $C \subseteq S^2$, over the fibers in M , over S^2 , and over M . The integration is always with respect to the volume element of the given Riemannian metric. Since p and q are Riemannian submersions by Lemma 2.1, we can use Fubini's Theorem to integrate over M by integrating first over the fibers of either p or q , and then over S^2 ; cf. [18, Lemma 4]. By the remarks above, if $C = p(q^{-1}(w))$ and $f: S^2 \rightarrow \mathbb{R}$ then $\int_{q^{-1}(w)} f \circ p = \int_C f$.

Proposition 2.2. *Given a continuous function $f: S^2 \rightarrow \mathbb{R}^+$, we define $m \in \mathbb{R}$ by setting $m = \min\{\int_C f: C \subseteq S^2 \text{ a great circle}\}$. Then*

$$\frac{m^2}{\pi} \leq \frac{1}{4\pi} \left(\int_{S^2} f \right)^2 \leq \int_{S^2} f^2,$$

where equality in the second inequality occurs if and only if f is constant.

Proof. Using the fact that M is the total space of a pair of Riemannian submersions, we obtain

$$\begin{aligned} \int_{S^2} f &= \int_{S^2} \left(\frac{1}{2\pi} \int_{p^{-1}(v)} f \circ p \right) \\ &= \frac{1}{2\pi} \int_M f \circ p \\ &= \frac{1}{2\pi} \int_{S^2} \left(\int_{q^{-1}(w)} f \circ p \right) \\ &\geq \frac{1}{2\pi} \int_{S^2} m = 2m, \end{aligned}$$

proving the first inequality. By the Cauchy–Schwarz inequality, we have

$$\left(\int_{S^2} 1 \cdot f \right)^2 \leq 4\pi \int_{S^2} f^2,$$

proving the second inequality. Here equality occurs if and only if f and 1 are linearly dependent, i.e., if and only if f is constant. \square

We define the quantity V_f by setting $V_f = \int_{S^2} f^2 - \frac{1}{4\pi} \left(\int_{S^2} f \right)^2$. Then Proposition 2.2 can be restated as follows.

Corollary 2.3. *Let $f: S^2 \rightarrow \mathbb{R}^+$ be continuous. Then*

$$\int_{S^2} f^2 - \frac{m^2}{\pi} \geq V_f \geq 0,$$

and $V_f = 0$ if and only if f is constant.

Proof. The proof is obtained from Proposition 2.2 by noting that $a \leq b \leq c$ if and only if $c - a \geq c - b \geq 0$. \square

We can assign a probabilistic meaning to V_f as follows. Divide the area measure on S^2 by 4π , thus turning it into a probability measure μ . A function $f: S^2 \rightarrow \mathbb{R}^+$ is then thought of as a random variable with expectation $E_\mu(f) = \frac{1}{4\pi} \int_{S^2} f$. Its variance is thus given by

$$\text{Var}_\mu(f) = E_\mu(f^2) - (E_\mu(f))^2 = \frac{1}{4\pi} \int_{S^2} f^2 - \left(\frac{1}{4\pi} \int_{S^2} f \right)^2 = \frac{1}{4\pi} V_f.$$

The variance of a random variable f is non-negative, and it vanishes if and only if f is constant. This reproves the corresponding properties of V_f established above via the Cauchy–Schwarz inequality.

Now let g_0 be the metric of constant Gaussian curvature $K = 1$ on \mathbb{RP}^2 . The double covering $\rho: S^2 \rightarrow (\mathbb{RP}^2, g_0)$ is a local isometry. Each projective line $C \subseteq \mathbb{RP}^2$ is the image under ρ of a great circle of S^2 .

Proposition 2.4. *Given a function $f: \mathbb{RP}^2 \rightarrow \mathbb{R}^+$, we define $\bar{m} \in \mathbb{R}$ by setting $\bar{m} = \min\{\int_C f: C \subseteq \mathbb{RP}^2 \text{ a projective line}\}$. Then*

$$\frac{2\bar{m}^2}{\pi} \leq \frac{1}{2\pi} \left(\int_{\mathbb{RP}^2} f \right)^2 \leq \int_{\mathbb{RP}^2} f^2,$$

where equality in the second inequality occurs if and only if f is constant.

Proof. We apply Proposition 2.2 to the composition $f \circ \rho$. Note that we have $\int_{\rho^{-1}(C)} f \circ \rho = 2 \int_C f$ and $\int_{S^2} f \circ \rho = 2 \int_{\mathbb{RP}^2} f$. The condition for f to be constant holds since f is constant if and only if $f \circ \rho$ is constant. \square

For \mathbb{RP}^2 we define $\bar{V}_f = \int_{\mathbb{RP}^2} f^2 - \frac{1}{2\pi} \left(\int_{\mathbb{RP}^2} f \right)^2 = \frac{1}{2} V_{f \circ \rho}$. We obtain the following restatement of Proposition 2.4.

Corollary 2.5. *Let $f: \mathbb{RP}^2 \rightarrow \mathbb{R}^+$ be a continuous function. Then*

$$\int_{\mathbb{RP}^2} f^2 - \frac{2\bar{m}^2}{\pi} \geq \bar{V}_f \geq 0,$$

where $\bar{V}_f = 0$ if and only if f is constant.

Relative to the probability measure induced by $\frac{1}{2\pi}g_0$ on \mathbb{RP}^2 , we have $E(f) = \frac{1}{2\pi} \int_{\mathbb{RP}^2} f$, and therefore $\text{Var}(f) = \frac{1}{2\pi} \bar{V}_f$, providing a probabilistic meaning for the quantity \bar{V}_f , as before.

By the uniformization theorem, every metric g on \mathbb{RP}^2 is of the form $g = f^2 g_0$ where g_0 is of constant Gaussian curvature $+1$, and the function $f: \mathbb{RP}^2 \rightarrow \mathbb{R}^+$ is continuous. The area of g is $\int_{\mathbb{RP}^2} f^2$, and the g -length of a projective line C is $\int_C f$. Let L be the shortest length of a noncontractible loop. Then $L \leq \bar{m}$ where \bar{m} is defined in Proposition 2.4, since a projective line in \mathbb{RP}^2 is a noncontractible loop. Then Corollary 2.5 implies $\text{area}(\mathbb{RP}^2, g) - \frac{2L^2}{\pi} \geq \bar{V}_f \geq 0$. If $\text{area}(\mathbb{RP}^2, g) = \frac{2L^2}{\pi}$ then $\bar{V}_f = 0$, which implies that f is constant, by Corollary 2.5. Conversely, if f is a constant c , then the only geodesics are the projective lines, and therefore $L = c\pi$. Hence $\frac{2L^2}{\pi} = 2\pi c^2 =$

$\text{area}(\mathbb{RP}^2)$. We have thus completed the proof of the following result strengthening Pu's inequality.

Theorem 2.6. *Let g be a Riemannian metric on \mathbb{RP}^2 . Let L be the shortest length of a noncontractible loop in (\mathbb{RP}^2, g) . Let $f: \mathbb{RP}^2 \rightarrow \mathbb{R}^+$ be such that $g = f^2 g_0$ where g_0 is of constant Gaussian curvature $+1$. Then*

$$\text{area}(g) - \frac{2L^2}{\pi} \geq 2\pi \text{Var}(f),$$

where the variance is with respect to the probability measure induced by $\frac{1}{2\pi}g_0$. Furthermore, equality $\text{area}(g) = \frac{2L^2}{\pi}$ holds if and only if f is constant.

REFERENCES

- [1] Pao Ming Pu, Some inequalities in certain nonorientable Riemannian manifolds, *Pacific J. Math.* 2 (1952), 55–71.
- [2] Mikhael Gromov, Filling Riemannian manifolds. *J. Differential Geom.* 18 (1983), no. 1, 1–147.
- [3] Mikhael Gromov, Metric structures for Riemannian and non-Riemannian spaces, Based on the 1981 French original with appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates, *Progress in Mathematics*, 152, Birkhäuser Boston, Boston, MA, 1999.
- [4] Marcel Berger, Lectures on geodesics in Riemannian geometry, *Tata Institute of Fundamental Research Lectures on Mathematics*, No. 33 Tata Institute of Fundamental Research, Bombay 1965.
- [5] Charles Horowitz, Karin Usadi Katz, Mikhail Katz, Loewner's torus inequality with isosystolic defect, *J. Geom. Anal.* 19 (2009), no. 4, 796–808.
- [6] Ivan Babenko, Florent Balacheff, and Guillaume Bulteau, Systolic geometry and simplicial complexity for groups, *J. Reine Angew. Math.* 757 (2019), 247–277.
- [7] Florent Balacheff, A local optimal diastolic inequality on the two-sphere, *J. Topol. Anal.* 2 (2010), no. 1, 109–121.
- [8] Viktor Bangert, Christopher Croke, Sergei Ivanov, and Mikhail Katz, Boundary case of equality in optimal Loewner-type inequalities, *Transactions of the American Mathematical Society* **359** (2007) (1), 1–17.
- [9] Paul Creutz, Rigidity of the Pu inequality and quadratic isoperimetric constants of normed spaces, preprint (2020). See <https://arxiv.org/pdf/2004.01076.pdf>
- [10] Chady El Mir and Zeina Yassine, Conformal geometric inequalities on the Klein bottle, *Conform. Geom. Dyn.* 19 (2015), 240–257.
- [11] Larry Guth, Systolic inequalities and minimal hypersurfaces, *Geom. Funct. Anal.* 19 (2010), no. 6, 1688–1692.
- [12] James Hebda, The primitive length spectrum of 2-D tori and generalized Loewner inequalities, *Trans. Amer. Math. Soc.* 372 (2019), no. 9, 6371–6401.
- [13] Sergei Ivanov and Mikhail Katz, Generalized degree and optimal Loewner-type inequalities, *Israel J. Math.* 141 (2004), 221–233.

- [14] Mikhail Katz, *Systolic geometry and topology*, with an appendix by Jake P. Solomon, *Mathematical Surveys and Monographs*, 137, American Mathematical Society, Providence, RI, 2007.
- [15] Mikhail Katz and Stephane Sabourau, *Hyperelliptic surfaces are Loewner*, *Proceedings of the American Mathematical Society* 134 (2006), no. 4, 1189–1195.
- [16] Yevgeny Liokumovich, Alexander Nabutovsky, and Regina Rotman, *Lengths of three simple periodic geodesics on a Riemannian 2-sphere*, *Math. Ann.* 367 (2017), no. 1–2, 831–855.
- [17] Stephane Sabourau, *Isosystolic genus three surfaces critical for slow metric variations*, *Geom. Topol.* 15 (2011), no. 3, 1477–1508.
- [18] Balázs Csikós and Márton Horváth, *Harmonic manifolds and tubes*, *J. Geom. Anal.* 28 (2018), no. 4, 3458–3476.

M. KATZ, DEPARTMENT OF MATHEMATICS, BAR ILAN UNIVERSITY, RAMAT GAN 5290002 ISRAEL

E-mail address: `katzmik@macs.biu.ac.il`

T. NOWIK, DEPARTMENT OF MATHEMATICS, BAR ILAN UNIVERSITY, RAMAT GAN 5290002 ISRAEL

E-mail address: `tahl@math.biu.ac.il`