

# THE THIRD LOGARITHMIC COEFFICIENT FOR THE SUBCLASSES OF CLOSE-TO-CONVEX FUNCTIONS

NAJLA M. ALARIFI

ABSTRACT. Let  $\mathcal{A}$  denote the set of all analytic functions  $f$  in the unit disk  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$  normalized by  $f(0) = 0$  and  $f'(0) = 1$ . The logarithmic coefficients  $\gamma_n$  of  $f \in \mathcal{A}$  are defined by  $\log f(z)/z = 2 \sum_{n=1}^{\infty} \gamma_n z^n$ . In the present paper, the upper bound of the third logarithmic coefficient in general case of  $f''(0)$  was computed when  $f$  belongs to some familiar subclasses of close-to-convex functions.

## 1. INTRODUCTION AND PRELIMINARIES

Let  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$  denote the open unit disk in the complex plane  $\mathbb{C}$ . Let  $\mathcal{A}$  denote the set of all analytic functions  $f$  in the open unit disk of the form

$$(1.1) \quad f(z) = \sum_{n=1}^{\infty} a_n z^n, \quad \text{with } a_1 = 1,$$

and  $\mathcal{S}$  be its subclass consisting functions that are univalent in  $\mathbb{D}$ . Given a function  $f \in \mathcal{S}$ , the coefficients  $\gamma_n$  defined by

$$(1.2) \quad \log \frac{f(z)}{z} = 2 \sum_{n=1}^{\infty} \gamma_n z^n, \quad z \in \mathbb{D} \setminus \{0\}, \quad \log 1 := 0.$$

are called the logarithmic coefficients of  $f$ . As it is commonly known, the logarithmic coefficients takes a leading role in Milin conjecture ([8], [6, p.155]), that is  $f \in \mathcal{S}$ ,

$$\sum_{m=1}^n \sum_{k=1}^m \left( k |\gamma_k|^2 - \frac{1}{k} \right) \leq 0.$$

Milin conjecture was confirmed to be the famous Bieberbach conjecture (e.g., [6, p.37]) by De Branges [2]. Sharp estimates for the class  $\mathcal{S}$  are known only for the first two coefficients,

$$|\gamma_1| \leq 1, \quad |\gamma_2| \leq \frac{1}{2} + \frac{1}{e} = 0.635 \dots$$

However, Obradović and Tuneski [9] obtained an upper bound of  $|\gamma_3|$  for the class  $\mathcal{S}$ .

The problem of estimating the modulus of the first three logarithmic coefficients is significantly studied for the subclasses of  $\mathcal{S}$  and in some cases sharp bounds are obtained. For instance, sharp estimates for the class of starlike functions  $\mathcal{S}^*$  are given by the inequality  $|\gamma_n| \leq 1/n$  holds for  $n \in \mathbb{N}$  [13, p.42]. Furthermore, for  $f \in \mathcal{SS}^*$ , the class of strongly starlike function of order  $\beta$ , ( $0 \leq \beta \leq 1$ ), it holds that  $|\gamma_n| \leq \beta/n$  ( $n \in \mathbb{N}$ ) [11]. The bounds of  $\gamma_n$  for functions in subclasses of  $\mathcal{S}$  has been widely studied in recent years, sharp estimates for the class of strongly starlike functions of certain order and gamma-starlike functions and different subclasses of close-to-convex functions are given in [11], [13, p.116]

---

2010 *Mathematics Subject Classification*. Primary: 30C45, 30C50; Secondary: 30C80.

*Key words and phrases*. Analytic function; univalent functions; Close-to-convex functions; Logarithmic coefficients.

and [12] respectively, while non-sharp estimates for the class of Bazilevic, close-to-convex are given in [1, 5, 10] respectively.

Let  $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$  denote the subclasses of  $\mathcal{S}$  satisfying respectively the next condition

$$\begin{aligned} \operatorname{Re}\{(1-z)f'(z)\} &> 0, \quad z \in \mathbb{D}, \\ \operatorname{Re}\{(1-z^2)f'(z)\} &> 0, \quad z \in \mathbb{D}, \\ \operatorname{Re}\{(1-z+z^2)f'(z)\} &> 0, \quad z \in \mathbb{D}. \end{aligned}$$

Account that each class defined above is the subclass of the well known class of close-to-convex functions, consequently families  $\mathcal{F}_i$ ,  $i = 1, 2, 3$ , contain only univalent functions [6, Vol.II, p. 2]. The sharp bounds of  $\gamma_1, \gamma_2$  and partial results for  $\gamma_3$  of the subclasses  $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$  of  $\mathcal{S}$  were determined by Kumar and Ali [7]. Moreover, Cho *et al.* [4] computed the sharp upper bounds for the third logarithmic coefficient  $\gamma_3$  of  $f$  when  $a_2$  is real number.

Differentiating (1.2) and comparing the coefficients with (1.1), we get  $\gamma_1 = \frac{1}{2}a_1$ ,  $\gamma_2 = \frac{1}{2}(a_3 - \frac{1}{2}a_2^2)$  and

$$(1.3) \quad \gamma_3 = \frac{1}{2} \left( a_4 - a_2a_3 + \frac{1}{3}a_2^3 \right).$$

The main aim of this paper is to determine upper bound of the third logarithmic coefficient in general case of  $a_2$ .

The following lemma are needed to prove our main results

**Lemma 1.1.** [3] *Let  $w(z) = c_1z + c_2z^2 + \dots$  be a Schwarz function. Then*

$$|c_1| \leq 1, \quad |c_2| \leq 1 - |c_1|^2 \quad \text{and} \quad |c_3| \leq 1 - |c_1|^2 - \frac{|c_2|^2}{1 + |c_1|}.$$

## 2. MAIN RESULTS

**Theorem 2.1.** *Let  $f \in \mathcal{F}_1$ . Then*

$$|\gamma_3| \leq \frac{15.75}{48} = 0.328125.$$

*Proof.* Since  $f \in \mathcal{F}_1$ , an analytic self-map  $w$  of  $\mathbb{D}$  with  $w(0) = 0$  exists and

$$(2.1) \quad (1-z)f'(z) = \frac{1+w(z)}{1-w(z)} = 1 + 2w(z) + 2w^2(z) + \dots$$

Writing

$$(2.2) \quad w(z) = c_1z + c_2z^2 + \dots,$$

then by using (2.1) along with (2.2) lead to

$$(2.3) \quad \begin{aligned} a_2 &= \frac{1}{2}(1 + 2c_1), \\ a_3 &= \frac{1}{3}(1 + 2c_1 + 2c_1^2 + 2c_2), \\ a_4 &= \frac{1}{4}(1 + 2c_1 + 2c_2 + 2c_3 + 2c_1^2 + 4c_1c_2 + 2c_1^3). \end{aligned}$$

From (1.3) and (2.3) after some calculations, the following was obtained

$$\gamma_3 = \frac{1}{48}(3 + 2c_1 + 4c_2 + 12c_3 + 8c_1c_2 + 4c_1^3),$$

and from here by using Lemma 1.1

$$(2.4) \quad \begin{aligned} 48|\gamma_3| &\leq 3 + 2|c_1| + 4|c_2| + 12|c_3| + 8|c_1||c_2| + 4|c_1|^3 \\ &\leq 3 + 2|c_1| + 4|c_2| + 12 \left( 1 - |c_1|^2 - \frac{|c_2|^2}{1 + |c_1|} \right) + 8|c_1||c_2| + 4|c_1|^3 =: f_1(|c_1|, |c_2|), \end{aligned}$$

where

$$\begin{aligned} f_1(x, y) &= 3 + 2x + 4y + 12 \left( 1 - x^2 - \frac{y^2}{1 + x} \right) + 8xy + 4x^3, \\ (x, y) \in E &: \quad 0 \leq x \leq 1, \quad 0 \leq y \leq 1 - x^2. \end{aligned}$$

The system

$$\begin{aligned} \frac{\partial f_1(x, y)}{\partial x} &= 2 - 24x + 12 \left( \frac{y}{1 + x} \right)^2 + 8y + 12x^2 = 0, \\ \frac{\partial f_1(x, y)}{\partial y} &= 4 - \frac{24y}{1 + x} + 8x = 0 \end{aligned}$$

has unique solution  $(x_1, y_1) = (1/4, 5/16) \in E \setminus \partial E$  with

$$(2.5) \quad f_1(x_1, y_1) = 15.75.$$

The max  $f_1(x_1, y_1)$ , needs to be found when  $(x, y)$  belongs to the boundary of  $E$ . In that sense, we have

$$(2.6) \quad \begin{aligned} f_1(x, 0) &= 15 + 2x - 12x^2 + 4x^3 \leq 9 + \frac{10\sqrt{30}}{9} = 15.08580 \dots \quad \text{for } 0 \leq x \leq 1, \\ f_1(0, y) &= 15 + 4y - 12y^2 \leq \frac{46}{3} = 15.33 \dots \quad \text{for } 0 \leq y \leq 1, \\ f_1(x, 1 - x^2) &= 7 + 22x - 4x^2 - 16x^3 \leq 15.304035 \dots \end{aligned}$$

Using (2.4), (2.5) and (2.6), concludes that

$$48|\gamma_3| \leq 15.75, \quad \text{i.e., } |\gamma_3| \leq 0.328125.$$

This completes the proof. □

**Remark 2.1.** If  $f \in \mathcal{F}_1$ , where  $f''(0)$  is real number, then [4]

$$|\gamma_3| \leq \frac{1}{288} \left( 11 + 15\sqrt{30} \right) = 0.323466 \dots$$

**Theorem 2.2.** Let  $f \in \mathcal{F}_2$ . Then

$$|\gamma_3| \leq 0.258765 \dots$$

*Proof.* Since  $f \in \mathcal{F}_2$ , there exists an analytic self-map  $w$  of  $\mathbb{D}$  with  $w(0) = 0$  and

$$(2.7) \quad (1 - z^2)f'(z) = \frac{1 + w(z)}{1 - w(z)} = 1 + 2w(z) + 2w^2(z) + \dots$$

The coefficients can be found by comparing the notations given in (2.2) and (2.7)

$$(2.8) \quad \begin{aligned} a_2 &= c_1, \\ a_3 &= \frac{1}{3}(1 + 2c_2 + 2c_1^2), \\ a_4 &= \frac{1}{2}(c_1 + c_3 + 2c_1c_2 + c_1^3). \end{aligned}$$

From (1.3) and (2.8) after some calculations, the following was obtained

$$\gamma_3 = \frac{1}{12}(c_1 + 3c_3 + 2c_1c_2 + c_1^3),$$

and from here by using Lemma 1.1

$$(2.9) \quad \begin{aligned} 12|\gamma_3| &\leq |c_1| + 3|c_3| + 2|c_1||c_2| + |c_1|^3 \\ &\leq |c_1| + 3 \left( 1 - |c_1|^2 - \frac{|c_2|^2}{1 + |c_1|} \right) + 2|c_1||c_2| + |c_1|^3 =: f_2(|c_1|, |c_2|), \end{aligned}$$

where

$$\begin{aligned} f_2(x, y) &= x + 3 \left( 1 - x^2 - \frac{y^2}{1 + x} \right) + 2xy + x^3, \\ (x, y) \in E &: \quad 0 \leq x \leq 1, \quad 0 \leq y \leq 1 - x^2. \end{aligned}$$

From the system

$$\begin{aligned} \frac{\partial f_2(x, y)}{\partial x} &= 1 - 6x + 3 \left( \frac{y}{1 + x} \right)^2 + 2y + 3x^2 = 0, \\ \frac{\partial f_2(x, y)}{\partial y} &= -\frac{6y}{1 + x} + 2x = 0, \end{aligned}$$

only one solution  $(x_2, y_2)$  lies in the interior of  $E$ , where

$$\begin{aligned} x_2 &= \frac{4 - \sqrt{7}}{6} = 0.22570 \dots, \\ y_2 &= \frac{47 - 14\sqrt{7}}{108} = 0.092217 \dots, \end{aligned}$$

and

$$(2.10) \quad f_2(x_2, y_2) = 3.10518 \dots$$

On the boundary of  $E$ , we have the next property

$$(2.11) \quad \begin{aligned} f_2(x, 0) &= 3(1 - x^2) + x + x^3 \leq 2 + \frac{4}{9}\sqrt{6} = 3.08866 \quad \text{for } 0 \leq x \leq 1, \\ f_2(0, y) &= 3(1 - y^2) \leq 3 \quad \text{for } 0 \leq y \leq 1, \\ f_2(x, 1 - x^2) &= 6x - 4x^3 \leq 2\sqrt{2} = 2.82842 \dots \end{aligned}$$

Consequently (2.9), (2.10) and (2.11) yield

$$12|\gamma_3| \leq 3.10518 \dots, \quad \text{i.e.,} \quad |\gamma_3| \leq 0.258765 \dots$$

□

**Remark 2.2.** If  $f \in \mathcal{F}_2$ , where  $f''(0)$  is real number, then [4]

$$|\gamma_3| \leq \frac{1}{972} \left( 95 + 23\sqrt{46} \right) = 0.258223 \dots$$

**Theorem 2.3.** Let  $f \in \mathcal{F}_3$ . Then

$$|\gamma_3| \leq \frac{17.75}{48} = 0.36979 \dots$$

*Proof.* Proceeding similarly as in the previous proofs, there exists an analytic self-map  $w$  of  $\mathbb{D}$  with  $w(0) = 0$  and

$$(2.12) \quad (1 - z + z^2)f'(z) = \frac{1 + w(z)}{1 - w(z)} = 1 + 2w(z) + 2w^2(z) + \dots$$

Substituting (2.2) into (2.12) and after comparing coefficients leads to

$$(2.13) \quad \begin{aligned} a_2 &= \frac{1}{2}(1 + 2c_1), \\ a_3 &= \frac{2}{3}(c_1 + c_2 + c_1^2), \\ a_4 &= \frac{1}{4}(2c_2 + 2c_3 + 2c_1^2 + 2c_1^3 + 4c_1c_2 - 1). \end{aligned}$$

By using (1.3) along with (2.13), upon simplification

$$\gamma_3 = \frac{1}{48}(-5 - 2c_1 + 4c_2 + 12c_3 + 8c_1c_2 + 4c_1^3),$$

and from here by using Lemma 1.1

$$(2.14) \quad \begin{aligned} 48|\gamma_3| &\leq 5 + 2|c_1| + 4|c_2| + 12|c_3| + 8|c_1||c_2| + 4|c_1|^3 \\ &\leq 5 + 2|c_1| + 4|c_2| + 12 \left( 1 - |c_1|^2 - \frac{|c_2|^2}{1 + |c_1|} \right) + 8|c_1||c_2| + 4|c_1|^3 =: f_3(|c_1|, |c_2|), \end{aligned}$$

where

$$\begin{aligned} f_3(x, y) &= 5 + 2x + 4y + 12 \left( 1 - x^2 - \frac{y^2}{1 + x} \right) + 8xy + 4x^3, \\ (x, y) \in E : \quad &0 \leq x \leq 1, \quad 0 \leq y \leq 1 - x^2. \end{aligned}$$

The system

$$\begin{aligned} \frac{\partial f_3(x, y)}{\partial x} &= 2 - 24x + 12 \left( \frac{y}{1 + x} \right)^2 + 8y + 12x^2 = 0, \\ \frac{\partial f_3(x, y)}{\partial y} &= 4 - \frac{24y}{1 + x} + 8x = 0 \end{aligned}$$

has unique solution  $(x_3, y_3) = (1/4, 5/16)$  belongs to the interior of  $E$  and

$$(2.15) \quad f_1(x_3, y_3) = 17.75.$$

On the boundary of  $E$ , the following cases are given

$$(2.16) \quad \begin{aligned} f_3(x, 0) &= 17 + 2x - 12x^2 + 4x^3 \leq 11 + \frac{10\sqrt{30}}{9} = 17.08580\dots \quad \text{for } 0 \leq x \leq 1, \\ f_3(0, y) &= 17 + 4y - 12y^2 \leq 17.33\dots \quad \text{for } 0 \leq y \leq 1, \\ f_3(x, 1 - x^2) &= 9 + 22x - 4x^2 - 20x^3 \leq 16.56455\dots \end{aligned}$$

Equations (2.14), (2.15) and (2.16) show that

$$|\gamma_3| \leq \frac{17.75}{48} = 0.36979\dots$$

□

**Remark 2.3.** Let  $f \in \mathcal{F}_3$ , where  $f''(0)$  is real number. Then [4]

$$|\gamma_3| \leq \frac{1}{7776} \left( 743 + 131\sqrt{262} \right) = 0.368238\dots$$

## REFERENCES

- [1] M. F. Ali and A. Vasudevarao, On logarithmic coefficients of some close-to-convex functions, Proc. Amer. Math. Soc. **146** (2018), no. 3, 1131–1142.
- [2] L. de Branges, A proof of the Bieberbach conjecture, Acta Math. **154** (1985), no. 1-2, 137–152.
- [3] F. Carlson, Sur les coefficients d'une fonction bornée dans le cercle unité, Ark. Mat. Astr. Fys. **27A** (1940), no. 1, 8 pp.
- [4] N. E. Cho et al., On the third logarithmic coefficient in some subclasses of close-to-convex functions, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **114** (2020), no. 2, Paper No. 52, 14 pp.
- [5] Q. Deng, On the logarithmic coefficients of Bazilevič functions, Appl. Math. Comput. **217** (2011), no. 12, 5889–5894.
- [6] P.T. Duren, Univalent Functions. Springer, New York (1983).
- [7] U. P. Kumar and A. Vasudevarao, Logarithmic coefficients for certain subclasses of close-to-convex functions, Monatsh. Math. **187** (2018), no. 3, 543–563.
- [8] I. M. Milin, Univalent functions and orthonormal systems, (Russian), Izdat. “Nauka”, Moscow, 1971.
- [9] M. Obradović and N. Tuneski, The third logarithmic coefficient for the class  $\mathcal{S}$ , arXiv preprint arXiv:2002.12865, 2020.
- [10] D. K. Thomas, On the coefficients of Bazilevič functions with logarithmic growth, Indian J. Math. **57** (2015), no. 3, 403–418.
- [11] D. K. Thomas, On the coefficients of strongly starlike functions, Indian J. Math. **58** (2016), no. 2, 135–146.
- [12] D. K. Thomas, On the logarithmic coefficients of close to convex functions, Proc. Amer. Math. Soc. **144** (2016), no. 4, 1681–1687.
- [13] D. K. Thomas, N. Tuneski and A. Vasudevarao, *Univalent functions*, De Gruyter Studies in Mathematics, 69, De Gruyter, Berlin, 2018.

DEPARTMENT OF MATHEMATICS, IMAM ABDULRAHMAN BIN FAISAL UNIVERSITY, DAMMAM 31113,  
KINGDOM OF SAUDI ARABIA

*E-mail address:* najarifi@gmail.com