

Isometric immersions with flat normal bundle between space forms

M. Dajczer, C.-R. Onti and Th. Vlachos

Abstract

We investigate the behavior of the second fundamental form of an isometric immersion of a space form with negative curvature into a space form so that the extrinsic curvature is negative. If the immersion has flat normal bundle, we prove that its second fundamental form grows exponentially.

It is a long-standing problem if the complete hyperbolic space \mathbb{H}^n can be isometrically immersed in the Euclidean space \mathbb{R}^{2n-1} . In fact, the non-existence of such an immersion has been frequently conjectured; see Yau [12], Moore [10] and Gromov [8]. A positive answer to the conjecture would be a natural generalization to higher dimensions of the classical result from 1901 by Hilbert for the hyperbolic plane. On one hand, Cartan [4] in 1920 showed that \mathbb{H}^n , $n \geq 3$, cannot be isometrically immersed in \mathbb{R}^{2n-2} even locally. On the other hand, he proved that there exists an abundance of local isometric immersions of \mathbb{H}^n into \mathbb{R}^{2n-1} and that these have all flat normal bundle.

Nikolayevsky [11] proved that complete non-simply connected Riemannian manifolds of constant negative sectional curvature cannot be isometrically immersed into Euclidean space with flat normal bundle. Let \mathbb{Q}_c^m denote a complete simply connected m -dimensional Riemannian manifold of constant sectional curvature c , that is, the Euclidean space \mathbb{R}^m , the Euclidean sphere \mathbb{S}_c^m or the hyperbolic space \mathbb{H}_c^m according to whether $c = 0, c > 0$ or $c < 0$, respectively. It was observed in [7] that the proof by Nikolayevsky gives, in fact, the following slightly more general result:

If there exists an isometric immersion $f: M_c^n \rightarrow \mathbb{Q}_{\tilde{c}}^{n+p}$, $n \geq 2$ and $c < 0$, with flat normal bundle of a complete Riemannian manifold M_c^n of constant sectional curvature c with $c < \tilde{c}$, then $M_c^n = \mathbb{H}_c^n$.

In view of Nikolayevsky's result, the following weaker version of the problem discussed above has already been considered by Brander [3].

PROBLEM: Do isometric immersions with flat normal bundle of \mathbb{H}_c^n into $\mathbb{Q}_{\tilde{c}}^{n+p}$ for $n \geq 2$ and $c < \tilde{c}$ exist?

In this paper, we analyze the behavior of the second fundamental form of a possible submanifold as in the problem above, and conclude that it must have exponential growth, as defined next.

Let $f: M^n \rightarrow \mathbb{Q}_{\tilde{c}}^{n+p}$ be an isometric immersion of a complete non-compact Riemannian manifold M^n . It is said that the second fundamental form $\alpha_f: TM \times TM \rightarrow N_f M$ of f has *exponential growth* if there exist $x_0 \in M^n$ and positive constants $k, \ell \in \mathbb{R}$ such that

$$\max \{ \|\alpha_f(x)\| : x \in D_r(x_0) \} \geq ke^{\ell r}$$

for any $r > r_0$ for some $r_0 > 0$, where $D_r(x_0)$ denotes the closed geodesic ball of M^n of radius r centered at x_0 and $\|\alpha_f\|$ is the norm of the second fundamental form given by

$$\|\alpha_f(x)\|^2 = \sum_{i,j} \|\alpha_f(X_i, X_j)(x)\|^2$$

where $X_1, \dots, X_n \in T_x M$ is an orthonormal basis.

Theorem 1. *If a complete n -dimensional Riemannian manifold M_c^n , $n \geq 2$ and $c < 0$, admits an isometric immersion $f: M_c^n \rightarrow \mathbb{Q}_{\tilde{c}}^{n+p}$, $c < \tilde{c}$, with flat normal bundle then $M_c^n = \mathbb{H}_c^n$ and the second fundamental form of f has exponential growth.*

The above gives as corollary the result due to Bolotov [2] that there is no isometric immersion of \mathbb{H}_c^n into \mathbb{R}^{n+p} with mean curvature vector field of bounded length.

The conclusion of Theorem 1 does not hold if the assumption of having flat normal bundle is dropped. For instance, it was shown by Aminov [1] that the example constructed by Rozendorn of an isometric immersion of \mathbb{H}^2 in \mathbb{R}^5 has no flat normal bundle and that the norm of its second fundamental form is globally bounded.

The aforementioned result for codimension $p = n - 1$ due to Cartan has the following immediate consequence:

Corollary 2. *If there exists an isometric immersion $f: \mathbb{H}_c^n \rightarrow \mathbb{Q}_{\tilde{c}}^{2n-1}$ with $c < \tilde{c}$ then the second fundamental form of f has exponential growth.*

1 The proof

Let $f: M^n \rightarrow \mathbb{Q}_c^{n+p}$ be an isometric immersion of a Riemannian manifold M^n into the space form \mathbb{Q}_c^{n+p} . If the immersion f has flat normal bundle, that is, if at any point the curvature tensor of the normal connection vanishes, then it is a standard fact (cf. [7]) that at any point $x \in M^n$ there exists a set of unique pairwise distinct normal vectors $\eta_i(x) \in N_f M(x)$, $1 \leq i \leq s(x)$, called the *principal normals* of f at x , and an associate orthogonal splitting of the tangent space as

$$T_x M = E_{\eta_1}(x) \oplus \cdots \oplus E_{\eta_s}(x),$$

where

$$E_{\eta_i}(x) = \{X \in T_x M : \alpha_f(X, Y) = \langle X, Y \rangle \eta_i \text{ for all } Y \in T_x M\}.$$

The *multiplicity* of a principal normal $\eta_i \in N_f M(x)$ of f at $x \in M^n$ is the dimension of the tangent subspace $E_{\eta_i}(x)$. If $s(x) = k$ is constant on M^n , then the maps $x \in M^n \mapsto \eta_i(x)$, $1 \leq i \leq k$, are smooth vector fields, called the *principal normal vector fields* of f . Moreover, also the distributions $x \in M^n \mapsto E_{\eta_i}(x)$, $1 \leq i \leq k$, are smooth.

In the sequel, let $f: M_c^n \rightarrow \mathbb{Q}_c^{n+p}$, $c < \tilde{c}$, be an isometric immersion with flat normal bundle. Since $C = \tilde{c} - c > 0$, it follows from the Gauss equation that any principal normal has multiplicity one. Thus, there exist exactly n nonzero principal normal vector fields η_1, \dots, η_n satisfying

$$\langle \eta_i, \eta_j \rangle = C, \quad 1 \leq i \neq j \leq n. \quad (1)$$

If $X_i \in \Gamma(E_{\eta_i})$, $1 \leq i \leq n$, is a unit local vector field then the local orthonormal frame X_1, \dots, X_n diagonalizes the second fundamental form of f , that is,

$$\alpha_f(X_i, X_j) = \delta_{ij} \eta_i, \quad 1 \leq i, j \leq n,$$

where δ_{ij} is the Kronecker delta. Such a frame is called a *principal frame*.

Lemma 3. *The following holds:*

$$\nabla_{X_i} X_j = -\lambda_i X_j (1/\lambda_i) X_i, \quad 1 \leq i \neq j \leq n, \quad (2)$$

where $\lambda_i = 1/\sqrt{\|\eta_i\|^2 + C}$.

Proof: The Codazzi equation is equivalent to

$$\nabla_{X_j}^\perp \eta_i = \langle \nabla_{X_i} X_j, X_j \rangle (\eta_i - \eta_j) \quad (3)$$

and

$$\langle \nabla_{X_\ell} X_j, X_i \rangle (\eta_i - \eta_j) = \langle \nabla_{X_j} X_\ell, X_i \rangle (\eta_i - \eta_\ell) \quad (4)$$

for all $1 \leq i \neq j \neq \ell \neq i \leq n$.

The vectors $\eta_i - \eta_j$ and $\eta_i - \eta_\ell$, $1 \leq i \neq j \neq \ell \neq i \leq n$ are linearly independent. Suppose otherwise that $\eta_i - \eta_j = \mu(\eta_i - \eta_\ell)$. Taking the inner product with η_i and using (1) gives $\|\eta_i\|^2 = C < 0$, a contradiction.

It now follows from (4) that

$$\nabla_{X_i} X_j = \Gamma_{ij}^i X_i + \Gamma_{ij}^j X_j, \quad i \neq j,$$

where $\Gamma_{ij}^k = \langle \nabla_{X_i} X_j, X_k \rangle$. Since $\Gamma_{ij}^j = \langle \nabla_{X_i} X_j, X_j \rangle = 0$, then

$$\nabla_{X_i} X_j = \Gamma_{ij}^i X_i = -\Gamma_{ii}^j X_i.$$

On the other hand, taking the inner product of (3) with η_i and using (1) is easily seen to give that $\Gamma_{ii}^j = \lambda_i X_j (1/\lambda_i)$, as we wished. ■

Lemma 4. *For each $x_0 \in M_c^n$ there exists a diffeomorphism $F: U \rightarrow V$ from an open subset $U \subset \mathbb{R}^n$ endowed with coordinates $\{u_1, \dots, u_n\}$ onto an open neighborhood $V \subset M_c^n$ of x_0 such that the tangent frame*

$$\sqrt{\|\eta_1\|^2 + CF_*(\partial/\partial u_1)}, \dots, \sqrt{\|\eta_n\|^2 + CF_*(\partial/\partial u_n)} \quad (5)$$

is orthonormal and principal. Moreover, if M_c^n is complete and simply connected then $F: \mathbb{R}^n \rightarrow M_c^n$ is a diffeomorphism.

Proof: For the local existence, observe that Lemma 3 implies

$$[\lambda_i X_i, \lambda_j X_j] = 0, \quad 1 \leq i \neq j \leq n.$$

For the proof of the global part, we follow a similar argument as in the proof of Theorem 3 in [9] or Proposition 5.6 in [7]. Assume that M_c^n is complete and simply connected. Set $Y_i = \lambda_i X_i$ and let $\varphi_i(x, t)$, $x \in M_c^n$, $t \in \mathbb{R}$ be the one-parameter group of diffeomorphisms generated by Y_i . Since the vector fields Y_i , $1 \leq i \leq n$, have bounded lengths, it follows that $\varphi_i(x, t)$ is defined for all values of x and t . Thus, for any $x \in M_c^n$, the map $t \mapsto \varphi_i(x, t)$

is the integral curve of Y_i with $\varphi_i(x, 0) = x$. Let x_0 be a fixed point in M_c^n and define a function $F = F_{x_0}: \mathbb{R}^n \rightarrow M_c^n$ by

$$F(t_1, t_2, \dots, t_n) = \varphi_n(\varphi_{n-1}(\dots \varphi_2(\varphi_1(x_0, t_1), t_2), \dots), t_n).$$

Since the Lie bracket $[Y_i, Y_j]$ vanishes the parameter groups φ_i and φ_j commute. This implies that

$$F_{x_0}(t+s) = \varphi_n(\varphi_{n-1}(\dots \varphi_2(\varphi_1(F_{x_0}(s), t_1), t_2), \dots), t_n) = F_{F_{x_0}(s)}(t) \quad (6)$$

where $t = (t_1, \dots, t_n)$ and $s = (s_1, \dots, s_n)$. Thus

$$F_*(s)\partial_i = \frac{d}{dt}\Big|_{t=0} F(s_1, \dots, s_i + t, \dots, s_n) = \frac{d}{dt}\Big|_{t=0} \varphi_i(F(s), t) = Y_i(F(s)).$$

We claim that F is a covering map. Then this and that M_c^n is simply connected yields that F is a diffeomorphism, which gives the proof.

Given $x \in M_c^n$, let $\tilde{B}_{2\varepsilon}(0)$ be an open ball of radius 2ε centered at the origin such that $F_x|_{\tilde{B}_{2\varepsilon}(0)}$ is a diffeomorphism onto $B_{2\varepsilon}(x) = F_x(\tilde{B}_{2\varepsilon}(0))$. Set $\{\tilde{x}_\alpha\}_{\alpha \in A} = F^{-1}(x)$ and denote by $\tilde{B}_{2\varepsilon}(\tilde{x}_\alpha)$ the open ball of radius 2ε centered at \tilde{x}_α . Define a map $\phi_\alpha: B_{2\varepsilon}(x) \rightarrow \tilde{B}_{2\varepsilon}(\tilde{x}_\alpha)$ by

$$\phi_\alpha(y) = \tilde{x}_\alpha + F_x^{-1}(y).$$

From (6) we obtain

$$F_{x_0}(\phi_\alpha(y)) = F_{x_0}(\tilde{x}_\alpha + F_x^{-1}(y)) = F_{F_{x_0}(\tilde{x}_\alpha)}(F_x^{-1}(y)) = F_x(F_x^{-1}(y)) = y$$

for all $y \in B_{2\varepsilon}(x)$. Thus F_{x_0} is a diffeomorphism from $\tilde{B}_{2\varepsilon}(\tilde{x}_\alpha)$ onto $B_{2\varepsilon}(x)$ having ϕ_α as its inverse. In particular, this implies that $\tilde{B}_{2\varepsilon}(\tilde{x}_\alpha)$ and $\tilde{B}_{2\varepsilon}(\tilde{x}_\beta)$ are disjoint if $\alpha, \beta \in A$ are distinct indices. Finally, it remains to check that if $\tilde{y} \in F_{x_0}^{-1}(B_\varepsilon(x))$, then $\tilde{y} \in \tilde{B}_\varepsilon(\tilde{x}_\alpha)$ for some $\alpha \in A$. This follows from the fact that

$$F_{x_0}(\tilde{y} - F_x^{-1}(F_{x_0}(\tilde{y}))) = F_{F_{x_0}(\tilde{y})}(-F_x^{-1}(F_{x_0}(\tilde{y}))) = x.$$

For the last equality, observe from (6) that for all $x, y \in M_c^n$ we have $F_x(t) = y$ if and only if $F_y(-t) = x$. ■

The *third fundamental form* $\text{III}_f(x)$ of f at $x \in M^n$ is given by

$$\text{III}_f(X, Y)(x) = \text{tr} \langle \alpha_f(X, \cdot), \alpha_f(Y, \cdot) \rangle, \quad X, Y \in T_x M.$$

Since α_f has no kernel (that is, positive index of relative nullity), then $\text{III}_f(x)$ is a positive definite inner product.

Lemma 5. *The Riemannian metric $g^0 = Cg + \text{III}_f$ is flat where g is the metric of M_c^n . Moreover, the metric g^0 is complete if g is complete.*

Proof: In terms of the system of principal coordinates $\{u_1, \dots, u_n\}$ given by Lemma 4, we have

$$g_{ij}^0 = Cg_{ij} + \text{III}_f(\partial/\partial u_i, \partial/\partial u_j) = \frac{C}{\|\eta_i\|^2 + C} \delta_{ij} + \frac{\|\eta_i\|^2}{\|\eta_i\|^2 + C} \delta_{ij} = \delta_{ij}.$$

Moreover, the metric g^0 is complete since $g_{ij}^0 > Cg_{ij}$. ■

Proof of Theorem 1. By Nikolayevsky's result we have that $M_c^n = \mathbb{H}_c^n$. Let $F: \mathbb{R}^n \rightarrow \mathbb{H}_c^n$ be the global diffeomorphism given by Lemma 4. We endow \mathbb{R}^n with the pullbacks of the two metrics considered in Lemma 5 that are still denoted by g and g^0 . Notice that (\mathbb{R}^n, g^0) is the standard flat Euclidean space.

Given a smooth curve $\gamma: [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}^n$ set

$$\hat{S}(\gamma) = \max_{t \in [a, b]} \|\alpha_f\|^2(F(\gamma(t))).$$

We have from (5) that

$$g_{ij} = \frac{1}{\|\eta_i\|^2 + C} \delta_{ij} \geq \frac{1}{\hat{S}(\gamma) + C} \delta_{ij}.$$

Then, the lengths of γ satisfy

$$L_{g^0}(\gamma) < (\hat{S}(\gamma) + C)^{1/2} L_g(\gamma). \quad (7)$$

Let $\gamma: [a, b] \rightarrow \mathbb{R}^n$ and $\tilde{\gamma}: [a, b] \rightarrow \mathbb{R}^n$ be the unique Euclidean and hyperbolic geodesics, respectively, joining $\gamma(a) = \tilde{\gamma}(a)$ to $\gamma(b) = \tilde{\gamma}(b)$. From (7) we have

$$L_{g^0}(\gamma) \leq L_{g^0}(\tilde{\gamma}) < (\hat{S}(\tilde{\gamma}) + C)^{1/2} L_g(\tilde{\gamma}).$$

Thus, if $\gamma_{x,y}$ is the unique hyperbolic geodesic joining $x \neq y \in \mathbb{R}^n$, then the distances with respect to g^0 and g satisfy

$$d_{g^0}(x, y) < (\hat{S}(\gamma_{x,y}) + C)^{1/2} d_g(x, y). \quad (8)$$

Fix $x_0 \in \mathbb{R}^n$ and let $D_r^g(x_0)$ and $D_r^{g^0}(x_0)$ be the closed geodesic balls of radius $r > 0$ centered at x_0 with respect to g and g^0 , respectively.

It holds that

$$D_r^g(x_0) \subset \text{int} \left(D_{\psi(r)}^{g^0}(x_0) \right), \quad (9)$$

where

$$\psi(r) = r(S(r) + C)^{1/2} \quad \text{and} \quad S(r) = \max_{x \in D_r^g(x_0)} (\|\alpha_f\|^2(F(x))).$$

In fact, if $y \in D_r^g(x_0)$ we have using (8) that

$$d_{g^0}(x_0, y) < (\hat{S}(\gamma_{x_0, y}) + C)^{1/2} d_g(x_0, y) \leq r(\hat{S}(\gamma_{x_0, y}) + C)^{1/2} \leq \psi(r).$$

Then, we obtain using (9) that the volumes of the geodesic balls satisfy

$$\begin{aligned} \text{Vol}_g(D_r^g(x_0)) &\leq \text{Vol}_g(D_{\psi(r)}^{g^0}(x_0)) \\ &= \int_{D_{\psi(r)}^{g^0}(x_0)} \prod_{i=1}^n (\|\eta_i\|^2 + C)^{-1/2} du_1 \wedge \cdots \wedge du_n \\ &< \int_{D_{\psi(r)}^{g^0}(x_0)} C^{-n/2} du_1 \wedge \cdots \wedge du_n = C^{-n/2} \text{Vol}_{g^0}(D_{\psi(r)}^{g^0}(x_0)) \\ &= r^n (1 + S(r)/C)^{n/2} \omega_n, \end{aligned}$$

where ω_n is the volume of the Euclidean unit n -ball. Since $\text{Vol}_g(D_r^g(x_0))$ is well known to grow exponentially with r (for instance, see [5]), it follows that also $S(r)$ grows exponentially with r , and thus the second fundamental form of f has exponential growth. ■

Remark 6. It is worth mentioning that it was shown in [6] that there is no isometric immersion with flat normal bundle of a complete Riemannian manifold M_c^n , $c > 0$, into $\mathbb{Q}_{\tilde{c}}^{n+p}$ with $c < \tilde{c}$. Notice that this follows using Lemma 4.

References

- [1] Aminov, Y., *Extrinsic geometric properties of the Rozendorn surface, which is an isometric immersion of the Lobachevski plane into E^5* . Sb. Math. **200** (2009), 1575–1586.
- [2] Bolotov, D., *On an isometric immersion with a flat normal connection of the Lobachevsky space L^n into the Euclidean space \mathbb{R}^{n+m}* , Math. Notes **82** (2007), 10–12.

- [3] Brander, D., *Results related to generalizations of Hilbert's non-immersibility theorem for the hyperbolic plane*, Electron. Res. Announc. Math. Sci. **15** (2008), 8–16.
- [4] Cartan, E., *Sur les variétés de courbure constante d'un espace euclidien ou non-euclidien*, Bull. Soc. Math. France **47** (1919), 125–160; **48** (1920), 132–208.
- [5] Chavel, I., “Riemannian Geometry: A Modern Introduction” Cambridge University Press, Cambridge, 1993.
- [6] Dajczer M. and Tojeiro, R., *Isometric immersions and the generalized Laplace and elliptic sinh-Gordon equations*, J. Reine Angew. Math. **467** (1995), 109–147.
- [7] Dajczer M. and Tojeiro, R., “Submanifold theory beyond an introduction”. Universitext. Springer, 2019.
- [8] Gromov, M., *Geometric, algebraic, and analytic descendants of Nash isometric embedding theorems*, Bull. Amer. Math. Soc. **54** (2017), 173–245.
- [9] Moore, J. D., *Isometric immersions of space forms in space forms*, Pacific J. Math. **40** (1972), 157–166.
- [10] Moore, J. D., *Problems in the geometry of submanifolds*, Mat. Fiz. Anal. Geom. **9** (2002), 648–662,
- [11] Y. Nikolayevsky, *Non-immersion theorem for a class of hyperbolic manifolds*, Differential Geom. Appl. **9** (1998) 239–242.
- [12] Yau, S.-T., *Seminar on Differential Geometry*, Ann. of Math. Stud. 102, Princeton Univ. Press, Princeton, N.J. (1982), 669–706,

Marcos Dajczer
 IMPA – Estrada Dona Castorina, 110
 22460–320, Rio de Janeiro – Brazil
 e-mail: marcos@impa.br

Christos-Raent Onti
 Department of Mathematics and Statistics

University of Cyprus
1678, Nicosia – Cyprus

Theodoros Vlachos
University of Ioannina
Department of Mathematics
Ioannina – Greece
e-mail: tvlachos@uoi.gr