

# Looking for a vector charmonium-like state $Y$ in $e^+e^- \rightarrow \bar{D}D_1(2420) + c.c.$

Jian-Rong Zhang

*Department of Physics, College of Liberal Arts and Sciences,  
National University of Defense Technology, Changsha 410073, Hunan, People's Republic of China*

Inspired by the first observation of a vector charmonium-like state  $Y(4626)$  decaying to a meson pair  $D_s^+D_{s1}(2536)^-$ , which could be viewed as a  $P$ -wave scalar-scalar  $[cs][\bar{c}\bar{s}]$  tetraquark state, we predict a potential vector charmonium-like state  $Y$  with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration. The corresponding mass spectrum of  $Y$  state is calculated to be  $4.33_{-0.23}^{+0.16}$  GeV in the framework of QCD sum rules. We suggest that the predicted  $Y$  state could be looked for in an open-charm  $e^+e^- \rightarrow \bar{D}D_1(2420) + c.c.$  process.

PACS numbers: 11.55.Hx, 12.38.Lg, 12.39.Mk

## I. INTRODUCTION

In recent years, a series of vector charmonium-like  $Y$  states have been observed in the initial-state radiation processes  $e^+e^- \rightarrow \gamma_{ISR}\pi^+\pi^-J/\psi$  ( $\psi(2S)$ ) [1–8] or in the direct processes  $e^+e^- \rightarrow \pi^+\pi^-J/\psi$  ( $\psi(2S)$ ) [9–12]. These experiments show that  $Y$  states mainly couple to hidden-charm final states. In contrast, Belle newly reported the first observation of  $Y(4626)$  in an open-charm process  $e^+e^- \rightarrow D_s^+D_{s1}(2536)^- + c.c.$  with a significance of  $5.9\sigma$  [13], which has promptly attracted much attention [14–23]. Theoretically, some authors pointed that  $Y(4626)$  can be well interpreted as a  $P$ -wave  $[cs][\bar{c}\bar{s}]$  state with a multi-quark color flux-tube model [21]. Moreover, we studied  $Y(4626)$  from two-point QCD sum rules, and finally arrived at that it could be a  $P$ -wave scalar-scalar  $[cs][\bar{c}\bar{s}]$  state [23]. On analogy of  $Y(4626)$ 's observation in the open-charm process, we propose that a novel vector charmonium-like state  $Y$  could be looked for in an open-charm  $e^+e^- \rightarrow \bar{D}D_1(2420) + c.c.$  process. In theory, the predicted  $Y$  state could correspondingly be regarded as a  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  tetraquark state.

In this work, we endeavor to explore the charmonium-like state  $Y$  with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration. To deal with the hadronic state, one has to confront the complicated nonperturbative QCD problem. As one trusty method for evaluating nonperturbative effects, the QCD sum rule [24] is firmly founded on the basic QCD theory, and has been successfully applied to plenty of hadronic systems (for reviews see Refs. [25–28] and references therein). Accordingly, we intend to study this  $Y$  state by making use of the QCD sum rule approach.

The paper's organization is as follows. In Sec. II, the QCD sum rule is derived for  $Y$  with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  structure, along with numerical analysis and discussions in Sec. III. The last part includes a brief summary.

## II. THE QCD SUM RULE FOR $Y$ WITH $P$ -WAVE SCALAR-SCALAR $[cq][\bar{c}\bar{q}]$ STRUCTURE

Generally speaking, one could have several choices on diquarks to characterize a  $P$ -wave tetraquark state with  $J^P = 1^-$ . It is worth noting that there have been broad discussions on the so-called “good” or “bad” diquarks for the tetraquark states [29], and then the  $Y$  state with  $P$ -wave  $[cq][\bar{c}\bar{q}]$  structure could be represented basing on following considerations [30]. A “good” diquark operator in the attractive anti-triplet color channel can be  $\bar{q}_c\gamma_5q$  with  $0^+$ , and a “bad” diquark operator can be  $\bar{q}_c\gamma q$  with  $1^+$ . Similarly, operators with  $0^-$  and  $1^-$  can be written as  $\bar{q}_c q$  and  $\bar{q}_c\gamma\gamma_5q$ , respectively. Further, it is suggested that diquarks are preferably formed into spin 0 from lattice results [31]. Comparatively, the solid tetraquark candidates tend to be composed of  $0^+$  “good” diquarks. For example, the final results from QCD sum rules favor the scalar diquark-scalar antidiquark case after comparing different diquark configurations [32]. Thereby,

the predicted  $Y$  state would be dominantly structured as the  $P$ -wave scalar diquark-scalar antidiquark, which contains the flavor content  $[cq][\bar{c}\bar{q}]$  with momentum numbers  $S_{[cq]} = 0$ ,  $S_{[\bar{c}\bar{q}]} = 0$ ,  $S_{[cq][\bar{c}\bar{q}]} = 0$ , and  $L_{[cq][\bar{c}\bar{q}]} = 1$ . Here  $q$  can be  $u$  or  $d$  quark, and  $c$  is the charm quark. Considering that both light  $u$  and  $d$  quark masses are taken as current-quark masses in the paper, they are so small comparing with the heavy running charm mass  $m_c$  that they will be neglected in the calculation complying with the usual treatment of heavy hadrons. Thus it is not concretely differentiated whether  $q = u$  or  $q = d$  for brevity. The corresponding current could be constructed as

$$j_\mu = \epsilon_{def} \epsilon_{d'e'f'} (q_d^T C \gamma_5 c_e) D_\mu (\bar{q}_{d'} \gamma_5 C \bar{c}_{e'}^T), \quad (1)$$

in which the index  $T$  denotes matrix transposition,  $C$  means the charge conjugation matrix,  $D_\mu$  is the covariant derivative to generate  $L = 1$ , and  $d, e, f, d',$  and  $e'$  are color indices.

Generally, the two-point correlator  $\Pi_{\mu\nu}(q^2) = i \int d^4x e^{iq \cdot x} \langle 0 | T [j_\mu(x) j_\nu^\dagger(0)] | 0 \rangle$  can be parameterized as

$$\Pi_{\mu\nu}(q^2) = \frac{q_\mu q_\nu}{q^2} \Pi^{(0)}(q^2) + \left( \frac{q_\mu q_\nu}{q^2} - g_{\mu\nu} \right) \Pi^{(1)}(q^2). \quad (2)$$

To yield the sum rule, the part  $\Pi^{(1)}(q^2)$  can be evaluated in two different ways. At the hadronic level, it can be expressed as

$$\Pi^{(1)}(q^2) = \frac{\lambda^2}{M_H^2 - q^2} + \frac{1}{\pi} \int_{s_0}^{\infty} ds \frac{\text{Im} \Pi^{(1)}(s)}{s - q^2}, \quad (3)$$

where  $\lambda$  is the hadronic coupling constant and  $M_H$  is the hadron's mass. At the quark level, it can be written as

$$\Pi^{(1)}(q^2) = \int_{4m_c^2}^{\infty} ds \frac{\rho(s)}{s - q^2}, \quad (4)$$

for which the spectral density  $\rho(s) = \frac{1}{\pi} \text{Im} \Pi^{(1)}(s)$ .

In deriving  $\rho(s)$ , one could work at leading order in  $\alpha_s$  and consider condensates up to dimension 8. To keep the heavy-quark mass finite, one uses the heavy-quark propagator in momentum space [33]. The correlator's light-quark part is calculated in the coordinate space and Fourier-transformed to the  $D$  dimension momentum space, which is combined with the heavy-quark part and then dimensionally regularized at  $D = 4$  [28, 34, 35]. It is given by  $\rho(s) = \rho^{\text{pert}} + \rho^{\langle \bar{q}q \rangle} + \rho^{\langle g^2 G^2 \rangle} + \rho^{\langle g \bar{q} \sigma \cdot G q \rangle} + \rho^{\langle \bar{q}q \rangle^2} + \rho^{\langle g^3 G^3 \rangle} + \rho^{\langle \bar{q}q \rangle \langle g^2 G^2 \rangle} + \rho^{\langle \bar{q}q \rangle \langle g \bar{q} \sigma \cdot G q \rangle}$ , detailedly with

$$\begin{aligned} \rho^{\text{pert}} &= -\frac{1}{3 \cdot 5 \cdot 2^{11} \pi^6} \int_{\alpha_{\min}}^{\alpha_{\max}} \frac{d\alpha}{\alpha^4} \int_{\beta_{\min}}^{1-\alpha} \frac{d\beta}{\beta^4} (1 - \alpha - \beta) \kappa r^5, \\ \rho^{\langle \bar{q}q \rangle} &= \frac{m_c \langle \bar{q}q \rangle}{3 \cdot 2^6 \pi^4} \int_{\alpha_{\min}}^{\alpha_{\max}} \frac{d\alpha}{\alpha^2} \int_{\beta_{\min}}^{1-\alpha} \frac{d\beta}{\beta^2} (2 - \alpha - \beta) r^3, \\ \rho^{\langle g^2 G^2 \rangle} &= -\frac{m_c^2 \langle g^2 G^2 \rangle}{3^2 \cdot 2^{12} \pi^6} \int_{\alpha_{\min}}^{\alpha_{\max}} \frac{d\alpha}{\alpha^4} \int_{\beta_{\min}}^{1-\alpha} \frac{d\beta}{\beta^4} (1 - \alpha - \beta) (\alpha^3 + \beta^3) \kappa r^2, \\ \rho^{\langle g \bar{q} \sigma \cdot G q \rangle} &= \frac{m_c \langle g \bar{q} \sigma \cdot G q \rangle}{2^8 \pi^4} \left\{ - \int_{\alpha_{\min}}^{\alpha_{\max}} \frac{d\alpha}{\alpha^2} \int_{\beta_{\min}}^{1-\alpha} \frac{d\beta}{\beta^2} (\alpha + \beta - 4\alpha\beta) r^2 + \int_{\alpha_{\min}}^{\alpha_{\max}} d\alpha \frac{[m_c^2 - \alpha(1 - \alpha)s]^2}{\alpha(1 - \alpha)} \right\}, \\ \rho^{\langle \bar{q}q \rangle^2} &= -\frac{m_c^2 \langle \bar{q}q \rangle^2}{3 \cdot 2^3 \pi^2} \int_{\alpha_{\min}}^{\alpha_{\max}} d\alpha [m_c^2 - \alpha(1 - \alpha)s], \\ \rho^{\langle g^3 G^3 \rangle} &= -\frac{\langle g^3 G^3 \rangle}{3^2 \cdot 2^{14} \pi^6} \int_{\alpha_{\min}}^{\alpha_{\max}} \frac{d\alpha}{\alpha^4} \int_{\beta_{\min}}^{1-\alpha} \frac{d\beta}{\beta^4} (1 - \alpha - \beta) \kappa [(\alpha^3 + \beta^3)r + 4(\alpha^4 + \beta^4)m_c^2] r, \\ \rho^{\langle \bar{q}q \rangle \langle g^2 G^2 \rangle} &= \frac{m_c \langle \bar{q}q \rangle \langle g^2 G^2 \rangle}{3^2 \cdot 2^8 \pi^4} \int_{\alpha_{\min}}^{\alpha_{\max}} \frac{d\alpha}{\alpha^2} \int_{\beta_{\min}}^{1-\alpha} \frac{d\beta}{\beta^2} \left\{ (2 - \alpha - \beta) (\alpha^3 + \beta^3) m_c^2 - 3[\alpha^2(\beta - 1) + \beta^2(\alpha - 1)] r \right\}, \end{aligned}$$

and

$$\rho^{\langle \bar{q}q \rangle \langle g\bar{q}\sigma \cdot Gq \rangle} = \frac{m_c^2 \langle \bar{q}q \rangle \langle g\bar{q}\sigma \cdot Gq \rangle}{3 \cdot 2^5 \pi^2} \int_{\alpha_{min}}^{\alpha_{max}} d\alpha (6\alpha^2 - 6\alpha + 1),$$

where  $\kappa = 1 + \alpha - 2\alpha^2 + \beta + 2\alpha\beta - 2\beta^2$ ,  $r = (\alpha + \beta)m_c^2 - \alpha\beta s$ ,  $\alpha_{min} = (1 - \sqrt{1 - 4m_c^2/s})/2$ ,  $\alpha_{max} = (1 + \sqrt{1 - 4m_c^2/s})/2$ , and  $\beta_{min} = \alpha m_c^2 / (s\alpha - m_c^2)$ . For the four-quark condensate, a general factorization  $\langle \bar{q}q\bar{q}q \rangle = \varrho \langle \bar{q}q \rangle^2$  [26, 36] has been employed, in which  $\varrho$  may be equal to 1 or 2.

Equating the two expressions (3) and (4), adopting quark-hadron duality, and making a Borel transform, the sum rule can be turned into

$$\lambda^2 e^{-M_H^2/M^2} = \int_{4m_c^2}^{s_0} ds \rho e^{-s/M^2}. \quad (5)$$

Taking the derivative of Eq. (5) with respect to  $-\frac{1}{M^2}$  and then dividing the result by Eq. (5) itself, one can obtain the hadron's mass sum rule

$$M_H^2 = \frac{\int_{4m_c^2}^{s_0} ds \rho s e^{-s/M^2}}{\int_{4m_c^2}^{s_0} ds \rho e^{-s/M^2}}, \quad (6)$$

in which light  $u$  and  $d$  current-quark masses have been safely neglected as they are so small comparing with the heavy  $m_c$ .

### III. NUMERICAL ANALYSIS AND DISCUSSIONS

In the numerical analysis, the running charm mass  $m_c$  is  $1.27 \pm 0.02$  GeV [37], and other input parameters are [24, 28]:  $\langle \bar{q}q \rangle = -(0.24 \pm 0.01)^3$  GeV<sup>3</sup>,  $m_0^2 = 0.8 \pm 0.1$  GeV<sup>2</sup>,  $\langle g\bar{q}\sigma \cdot Gq \rangle = m_0^2 \langle \bar{q}q \rangle$ ,  $\langle g^2 G^2 \rangle = 0.88 \pm 0.25$  GeV<sup>4</sup>, as well as  $\langle g^3 G^3 \rangle = 0.58 \pm 0.18$  GeV<sup>6</sup>. According to the standard criterion of sum rule analysis, one could find proper work windows for the threshold parameter  $\sqrt{s_0}$  and the Borel parameter  $M^2$ . The lower bound of  $M^2$  is obtained from the OPE convergence, and the upper one is found in view of that the pole contribution should be larger than QCD continuum one. Meanwhile, the threshold  $\sqrt{s_0}$  describes the beginning of continuum state, which is about 400 ~ 600 MeV bigger than the extracted  $M_H$  empirically.

At the very start, all the input parameters are kept at their central values and the four-quark condensate factor is taken as  $\varrho = 1$ . To get the lower bound of  $M^2$ , the OPE convergence is shown in FIG. 1 by comparing the relative contributions of different condensates from sum rule (5) for  $\sqrt{s_0} = 4.9$  GeV. Numerically, some main condensates could cancel each other out to some extent and the relative contribution of perturbative could play a predominant role in OPE at  $M^2 = 2.5$  GeV<sup>2</sup>, which is increasing with the enlarging of Borel parameter  $M^2$ . In this way, it is taken as  $M^2 \geq 2.5$  GeV<sup>2</sup> with an eye to the OPE convergence analysis. Besides, the upper bound of  $M^2$  is attained with a view to the pole contribution dominance in phenomenological side. In FIG. 2, it is compared between pole contribution and continuum from sum rule (5) for  $\sqrt{s_0} = 4.9$  GeV. The relative pole contribution is close to 50% at  $M^2 = 3.0$  GeV<sup>2</sup> and descending with the Borel parameter  $M^2$ . Thus the pole contribution dominance could be fulfilled while  $M^2 \leq 3.0$  GeV<sup>2</sup>. Accordingly, the Borel window of  $M^2$  is restricted to be 2.5 ~ 3.0 GeV<sup>2</sup> for  $\sqrt{s_0} = 4.9$  GeV. Analogously, the reasonable window of  $M^2$  is acquired as 2.5 ~ 2.9 GeV<sup>2</sup> for  $\sqrt{s_0} = 4.8$  GeV, and 2.5 ~ 3.2 GeV<sup>2</sup> for  $\sqrt{s_0} = 5.0$  GeV. In the work windows, one can expect that the two sides of QCD sum rules have a good overlap and it is reliable to extract information on the resonance. The dependence on  $M^2$  for the mass  $M_H$  of  $Y$  state is shown in FIG. 3, and its value is computed to be  $4.33 \pm 0.11$  GeV in work windows.

Next varying the input parameters, the mass  $M_H$  is obtained as  $4.33 \pm 0.11_{-0.08}^{+0.05}$  GeV (the first error due to variation of  $s_0$  and  $M^2$ , and the second one resulted from the uncertainty of QCD parameters) or shortly  $4.33_{-0.19}^{+0.16}$  GeV. In the end, paying attention to the variation of four-quark condensate factor  $\varrho$ , the




FIG. 1: The OPE convergence for the  $Y$  state with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration is shown by comparing the relative contributions of perturbative, two-quark condensate  $\langle\bar{q}q\rangle$ , two-gluon condensate  $\langle g^2 G^2\rangle$ , mixed condensate  $\langle g\bar{q}\sigma\cdot Gq\rangle$ , four-quark condensate  $\langle\bar{q}q\rangle^2$ , three-gluon condensate  $\langle g^3 G^3\rangle$ ,  $\langle\bar{q}q\rangle\langle g^2 G^2\rangle$ , and  $\langle\bar{q}q\rangle\langle g\bar{q}\sigma\cdot Gq\rangle$  from sum rule (5) for  $\sqrt{s_0} = 4.9$  GeV.




FIG. 2: The phenomenological contribution in sum rule (5) for  $\sqrt{s_0} = 4.9$  GeV for the  $Y$  state with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration. The solid line is the relative pole contribution (the pole contribution divided by the total, pole plus continuum contribution) as a function of  $M^2$  and the dashed line is the relative continuum contribution.

corresponding Borel curves are presented in FIG. 4 with  $\varrho = 2$ . In comparison with Fig. 3 for  $\varrho = 1$ , one could notice the mass uncertainty when varying  $\varrho$  from 1 to 2, and could get the final mass  $4.33^{+0.16}_{-0.23}$  GeV for the  $Y$  state with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration.

In experiment, one may note that in the hidden-charm  $e^+e^- \rightarrow \gamma_{ISR}\pi^+\pi^-\psi(2S)$  process, BABAR observed a broad structure near 4.32 GeV [2], and Belle subsequently found the charmonium-like state  $Y(4360)$  [3]. Afterward, a combined fit to these cross sections measured by BABAR and Belle experiments was performed [38], and the property of  $Y(4360)$  was further studied in  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$  via initial-state radiation at BABAR [7] and at Belle [8]. Taking notice of the close masses of  $Y(4360)$  and  $Y$  state concerned here, one could conjecture that they may be the same structure attributing to different decay




FIG. 3: The dependence on  $M^2$  for the mass  $M_H$  of the  $Y$  state with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration from sum rule (6) is shown while the four-quark condensate factor  $\varrho = 1$ . The ranges of  $M^2$  are  $2.5 \sim 2.9 \text{ GeV}^2$  for  $\sqrt{s_0} = 4.8 \text{ GeV}$ ,  $2.5 \sim 3.0 \text{ GeV}^2$  for  $\sqrt{s_0} = 4.9 \text{ GeV}$ , and  $2.5 \sim 3.2 \text{ GeV}^2$  for  $\sqrt{s_0} = 5.0 \text{ GeV}$ , respectively.




FIG. 4: The dependence on  $M^2$  for the mass  $M_H$  of the  $Y$  state with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration from sum rule (6) is shown while the four-quark condensate factor  $\varrho = 2$ .

modes. If that true, it would be very important for understanding  $Y(4360)$  to search for the predicted  $Y$  state, because complementary measurements by other decay modes such as the open-charm process will provide further insights into  $Y(4360)$ 's internal structure. Whether or not, it is undoubtedly exciting and significant if one could find a vector charmonium-like  $Y$  state particularly in an open-charm decay.

Invigoratingly, there has appeared some measurement of Born cross section for  $e^+e^- \rightarrow D^- D_1(2420)^+ + c.c.$  [39], in which the cross section line shape is consistent with the previous BESIII's result based on full reconstruction method [40], and there is some indication of enhanced cross section at the location of  $Y(4360)$ . Thereby, it seems promising that the predicted  $Y$  state could be observed in the open-charm process  $e^+e^- \rightarrow \bar{D} D_1(2420) + c.c.$  via either the initial-state radiation or the direct production for the future experiments.

#### IV. SUMMARY

Activated by the first observation of a vector charmonium-like state  $Y(4626)$  in the open-charm  $D_s^+ D_{s1}^-(2536)^-$  decay mode, for which could be a  $P$ -wave scalar-scalar  $[cs][\bar{c}\bar{s}]$  tetraquark state, we predict a novel vector charmonium-like  $Y$  state with  $P$ -wave scalar-scalar  $[cq][\bar{c}\bar{q}]$  configuration. Finally, the mass of  $Y$  is presented to be  $4.33_{-0.23}^{+0.16}$  GeV from QCD sum rules. We suggest that the predicted  $Y$  state could be searched for in an open-charm  $e^+e^- \rightarrow \bar{D}D_1(2420) + c.c.$  process through the initial-state radiation or the direct production in experiments, for which virtually there has been some indication of enhanced cross section in BESIII's existing measurements.

#### Acknowledgments

This work was supported by the National Natural Science Foundation of China under Contract Nos. 11475258 and 11675263, and by the project for excellent youth talents in NUDT.

- 
- [1] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **95**, 142001 (2005).
  - [2] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **98**, 212001 (2007).
  - [3] X. L. Wang *et al.* (Belle Collaboration), Phys. Rev. Lett. **99**, 142002 (2007).
  - [4] C. Z. Yuan *et al.* (Belle Collaboration), Phys. Rev. Lett. **99**, 182004 (2007).
  - [5] J. P. Lees *et al.* (BABAR Collaboration), Phys. Rev. D **86**, 051102(R) (2012).
  - [6] Z. Q. Liu *et al.* (Belle Collaboration), Phys. Rev. Lett. **110**, 252002 (2013).
  - [7] J. P. Lees *et al.* (BABAR Collaboration), Phys. Rev. D **89**, 111103 (2014).
  - [8] X. L. Wang *et al.* (Belle Collaboration), Phys. Rev. D **91**, 112007 (2015).
  - [9] T. E. Coan *et al.* (CLEO Collaboration), Phys. Rev. Lett. **96**, 162003 (2006).
  - [10] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **114**, 092003 (2015).
  - [11] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **118**, 092001 (2017).
  - [12] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. D **96**, 032004 (2017).
  - [13] S. Jia *et al.* (Belle Collaboration), Phys. Rev. D **100**, 111103 (2019).
  - [14] Y. Xie and Z. Q. Liu, arXiv:2001.09620 [hep-ex].
  - [15] S. Jia *et al.* (Belle Collaboration), Phys. Rev. D **101**, 091101 (2020).
  - [16] J. Z. Wang, R. Q. Qian, X. Liu, and T. Matsuki, Phys. Rev. D **101**, 034001 (2020).
  - [17] J. He, Y. Liu, J. T. Zhu, and D. Y. Chen, Eur. Phys. J. C **80**, 246 (2020).
  - [18] X. K. Dong, Y. H. Lin, and B. S. Zou, Phys. Rev. D **101**, 076003 (2020).
  - [19] H. W. Ke, X. H. Liu, and X. Q. Li, arXiv:2004.03167 [hep-ph].
  - [20] Y. Tan and J. L. Ping, Phys. Rev. D **101**, 054010 (2020).
  - [21] C. R. Deng, H. Chen, and J. L. Ping, Phys. Rev. D **101**, 054039 (2020).
  - [22] J. F. Giron and R. F. Lebed, Phys. Rev. D **101**, 074032 (2020); Phys. Rev. D **102**, 014036 (2020).
  - [23] J. R. Zhang, Phys. Rev. D **102**, 054006 (2020).
  - [24] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979); B **147**, 448 (1979); V. A. Novikov, M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Fortschr. Phys. **32**, 585 (1984).
  - [25] B. L. Ioffe, in The Spin Structure of The Nucleon, edited by B. Frois, V. W. Hughes, and N. de Groot (World Scientific, Singapore, 1997).
  - [26] S. Narison, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **17**, 1 (2002).
  - [27] P. Colangelo and A. Khodjamirian, in At the Frontier of Particle Physics: Handbook of QCD, edited by M. Shifman, Boris Ioffe Festschrift Vol. 3 (World Scientific, Singapore, 2001), pp. 1495-1576.
  - [28] M. Nielsen, F. S. Navarra, and S. H. Lee, Phys. Rep. **497**, 41 (2010).
  - [29] R. L. Jaffe, Phys. Rev. D **15**, 267 (1977); R. L. Jaffe and F. E. Low, Phys. Rev. D **19**, 2105 (1979); R. L. Jaffe and F. Wilczek, Phys. Rev. Lett. **91**, 232003 (2003).

- [30] N. V. Drenska, R. Faccini, and A. D. Polosa, *Phys. Lett. B* **669**, 160 (2008); A. Ali, C. Hambrock, and M. J. Aslam, *Phys. Rev. Lett.* **104**, 162001 (2010).
- [31] C. Alexandrou, Ph. de Forcrand, and B. Lucini, *Phys. Rev. Lett.* **97**, 222002 (2006).
- [32] J. R. Zhang, J. L. Zou, and J. Y. Wu, *Chin. Phys. C* **42**, 043101 (2018); J. R. Zhang, *Phys. Lett. B* **789**, 432 (2019).
- [33] L. J. Reinders, H. R. Rubinstein, and S. Yazaki, *Phys. Rep.* **127**, 1 (1985).
- [34] H. Kim and Y. Oh, *Phys. Rev. D* **72**, 074012 (2005); M. E. Bracco, A. Lozea, R. D. Matheus, F. S. Navarra, and M. Nielsen, *Phys. Lett. B* **624**, 217 (2005); R. D. Matheus, S. Narison, M. Nielsen, and J. M. Richard, *Phys. Rev. D* **75**, 014005 (2007).
- [35] J. R. Zhang and M. Q. Huang, *JHEP* **1011**, 057 (2010); J. R. Zhang, *Phys. Rev. D* **87**, 076008 (2013); *Phys. Rev. D* **89**, 096006 (2014); *Eur. Phys. J. C* **79**, 1001 (2019).
- [36] S. Narison, *Phys. Rep.* **84**, 263 (1982); G. Launer, S. Narison, and R. Tarrach, *Z. Phys. C* **26**, 433 (1984); S. Narison, *Phys. Lett. B* **673**, 30 (2009).
- [37] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018) and 2019 update.
- [38] Z. Q. Liu, X. S. Qin, and C. Z. Yuan, *Phys. Rev. D* **78**, 014032 (2008).
- [39] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Lett. B* **804**, 135395 (2020).
- [40] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **100**, 032005 (2019).