GENERALIZED TORSION AND DEHN FILLING

TETSUYA ITO, KIMIHIKO MOTEGI, AND MASAKAZU TERAGAITO

ABSTRACT. A generalized torsion element is a non-trivial element such that some non-empty finite product of its conjugates is the identity. We construct a generalized torsion element of the fundamental group of a 3-manifold obtained by Dehn surgery along a knot in S^3 .

1. Introduction

A non-trivial element g of a group G is a generalized torsion element if there exist $x_1, \ldots, x_k \in G$ such that

$$(1.1) (x_1 g x_1^{-1})(x_2 g x_2^{-1}) \cdots (x_k g x_k^{-1}) = 1.$$

That is, some non-empty finite product of its conjugates is the identity. The *order* of a generalized torsion element is the minimum k > 1 that satisfies (1.1).

One of motivations for exploring generalized torsion elements comes from the bi-orderability of groups. A group is bi-orderable if it admits a bi-ordering, a total ordering < such that agb < ahb for any $g, h, a, b \in G$ whenever g < h holds. In this paper, we adapt the convention that the trivial group $\{1\}$ is bi-orderable.

A bi-orderable group G has no generalized torsion element. For, if 1 < g for a bi-ordering < of G, then $1 = x1x^{-1} < xgx^{-1}$ for all $x \in G$, hence

$$1 < (x_1 g x_1^{-1})(x_2 g x_2^{-1}) \cdots (x_k g x_k^{-1})$$

for any $x_1, \ldots, x_k \in G$. The case g < 1 is similar.

In many situations one refutes the bi-orderability by finding a generalized torsion element. However, in general a group without generalized torsion element, which is called an R^* -group or a Γ -torsion-free group in literatures [1, 12, 14, 15], is not necessarily bi-orderable; see [15, Chapter 4], for example.

In [13] the second and third authors conjecture the following.

Conjecture 1.1 ([13]). Let G be the fundamental group of a 3-manifold. Then, G is bi-orderable if and only if G has no generalized torsion element.

In [13] this conjecture is verified for non-hyperbolic geometric 3–manifolds and some other examples.

Since a finitely generated bi-orderable group has \mathbb{Z} as a quotient, the fundamental group of any rational homology 3-sphere is not bi-orderable; see [2] for example. Then Conjecture 1.1 says that the fundamental group of any rational homology

1

The first named author has been partially supported by JSPS KAKENHI Grant Number JP16H02145 and JP19K03490.

The second named author has been partially supported by JSPS KAKENHI Grant Number 19K03502 and Joint Research Grant of Institute of Natural Sciences at Nihon University for 2019.

The third named author has been partially supported by JSPS KAKENHI Grant Number JP16K05149.

3–sphere has a generalized torsion element. Recall that 3–manifolds obtained by Dehn surgery along a knot in S^3 are rational homology 3–spheres except when the surgery is longitudinal. In this article we concentrate our attention on such 3–manifolds.

Let N(K) be a tubular neighborhood of K and denote the knot exterior by $E(K) = S^3 - \operatorname{Int} N(K)$. The knot group G(K) means the fundamental group $\pi_1(E(K))$. Throughout the paper we take a base point of E(K) which lies on $\partial E(K)$. We denote a meridian and a (preferred) longitude by $\mu, \lambda \in G(K)$, respectively.

Denote by K(m/n) the 3-manifold obtained by m/n-Dehn surgery on a knot K in S^3 . Note that $\pi_1(K(\infty)) = \pi_1(S^3) = \{1\}$ is bi-orderable and does not have a generalized torsion element by definition. Then Conjecture 1.1 implies:

Conjecture 1.2. Let K be a non-trivial knot in S^3 . Then $\pi_1(K(r))$ admits a generalized torsion element if $r \neq 0, \infty$.

For several classes of knots, we have supporting evidences for Conjecture 1.2.

Theorem 1.3. Let K be a knot in S^3 . Then $\pi_1(K(m/n))$ admits a generalized torsion element in the following cases.

- (1) K is a non-trivial torus knot, and $m/n \neq \infty$;
- (2) K is a (p,q)-cable knot, and $|pqn-m| \neq 1$;
- (3) K is a composite knot, and $n \neq 0, 1$.

For the first, if G(K) contains a generalized torsion element g, then its image in $\pi_1(K(r))$ is also a generalized torsion element whenever g remains non-trivial in $\pi_1(K(r))$. It is shown in [8] that for any hyperbolic knot K, there are only finitely many slopes r such that g becomes trivial in $\pi_1(K(r))$. So if G(K) contains a generalized torsion element, $\pi_1(K(r))$ also contains such an element for all but finitely many slopes r.

On the other hand, some knot K, such as the figure-eight knot, has bi-orderable knot group [18], and hence G(K) has no generalized torsion element. However, according to Conjecture 1.2, $\pi_1(K(r))$ should have a generalized torsion element for $r \neq 0, \infty$. So some non-generalized torsion element becomes a generalized torsion element via Dehn fillings.

The purpose of this article is to give several constructions of a generalized torsion element of $\pi_1(K(r))$ which is not the image of a generalized torsion element in G(K). This gives new supporting evidences for Conjecture 1.2, hence, for Conjecture 1.1.

For a knot K in S^3 , a singular spanning disk of K is a smooth map $\Phi \colon D^2 \to S^3$ (or, its image, by abuse of notation) such that $\Phi(\partial D^2) = K$ and that K intersects $\Phi(\operatorname{int} D^2)$ transversely in finitely many points. Each intersection point $\Phi(\operatorname{int} D^2) \cap K$ has a sign according to the orientations after orienting K and D^2 suitably. We say that $\Phi(D^2)$ is a (p,q)-singular spanning disk if K intersects $\Phi(\operatorname{int} D^2)$ positively in p points and negatively in p points.

A singular spanning disk appeared in early 3-dimensional topology; Dehn's lemma [17, 7] says that K has a (0,0)-singular spanning disk if and only if K is the trivial knot. More generally, K has a (p,q)-singular spanning disk with $0 \le p,q \le 1$ if and only if K is still the trivial knot [9].

Theorem 1.4. Let K be a knot in S^3 . If K has a (p,0)-singular spanning disk, then the image of a meridian μ in $\pi_1(K(m/n))$ is a generalized torsion element of

order m whenever $\frac{m}{n} \geq p$. Similarly, if K has a (0,q)-singular spanning disk, then the image of a meridian μ in $\pi_1(K(m/n))$ is a generalized torsion element order m whenever $\frac{m}{n} \leq -q$.

This construction illustrates how a generalized torsion element arises via Dehn fillings. A typical and fundamental example of (p,0)— or (0,q)—singular spanning disk is a clasp disk having the same sign of clasps.

Example 1.5. Let D be a standard disk in S^3 . Attach mutually disjoint arcs a_1, a_2, \ldots, a_p to D so that $a_i \cap D = a_i \cap \partial D = \partial a_i$. These arcs may be mutually linked and knotted. See Figure 1. Then, replace each arc a_i with a band having a single clasp as illustrated in Figure 1. These bands may have several full twists. Let K be a knot bounding this disk with p clasps. We remark that all clasps have the same sign. Then Theorem 1.4 asserts that $\pi_1(K(m/n))$ has a generalized torsion element whenever $m/n \geq 2p$




FIGURE 1. A disk D with arcs a_1, a_2, \ldots, a_p (Left). Replace each arc with a band having a single clasp (Right).

We will discuss further applications of this construction in Section 5. Theorem 1.4 is used in [11] to give an example of 3-manifold with non-trivial torus (JSJ) decomposition $M = M_1 \cup M_2 \cup \cdots \cup M_n$ such that each $\pi_1(M_i)$ has no generalized torsion element, but $\pi_1(M)$ has a generalized torsion element; see Remark 2.1.

Let us take a closer look at Dehn surgery along a double twist knot. A double twist knot is a 2-bridge knot of genus one given by Conway's notation C[2p, 2q] (p > 0) as in Figure 2. Double twist knots exhaust all genus one 2-bridge knots.




FIGURE 2. C[2p, 2q] (p = 2, q = 3)

A double twist knot is a generalization of a twist knot, which provides a quite interesting example in a study of generalized torsion elements. For a twist knot $K_q := C[2, 2q]$, the knot group $G(K_q)$ has a generalized torsion element if q < 0. If q = -1, then K_{-1} is the right-handed trefoil knot, for which $G(K_{-1})$ is known to have a generalized torsion element [16, Corollary 3.4]. For q = -2, Naylor and Rolfsen [16, Theorem 4.1] show that $G(K_{-2})$ admits a generalized torsion element. Surprisingly, this was the first example of hyperbolic knot whose knot group admits such an element. The general case $q \leq -2$ was proven by the third author [20]. On the other hand, when $q \geq 0$ the knot group $G(K_q)$ is bi-orderable [4], and hence it does not have a generalized torsion element.

It is not hard to see that K = C[2p, 2q] admits a (2p, 0)-singular spanning disk, and a (0,2q)-singular spanning disk if q>0 or, a (-2q,0)-singular spanning disk if q < 0. So by Theorem 1.4 the image of a meridian in $\pi_1(K(m/n))$ is a generalized torsion if $m \ge 2np$, or $m \le -2nq$ (when q > 0), or, $m \ge -2nq$ (when q < 0). The next theorem improves these conditions.

Theorem 1.6. Let K = C[2p, 2q] (p > 0) be a genus one two-bridge knot. Then the image of the meridian of K in $\pi_1(K(m/n))$ $(n \geq 1)$ is a generalized torsion element provided when

- (1) $m \geq (2n-1)p$; or
- (2) q > 0 and $m \le -(2n-1)q$; or (3) q < 0 and $m \ge -(2n-1)q$.

Example 1.7. Let K = C[2,2], which is the figure-eight knot. By Theorem 1.6, if either $m \geq 2n-1$ or $m \leq -(2n-1)$, then $\pi_1(K(m/n))$ has a generalized torsion element. In particular, $\pi_1(K(r))$ has a generalized torsion element for all non-zero integers r. Since $\pi_1(K(0))$ is known to be bi-orderable, $\pi_1(K(0))$ has no generalized torsion element.

In the final section we will propose some questions.

2. Proof of Theorem 1.3

In this section we prove Theorem 1.3.

(1) The (p,q)-torus knot group $G(K) = G(T_{p,q})$ (0 is presented as $\langle x,y | x^p = y^q \rangle$. First we observe that the commutator [x,y] is a generalized torsion element in G(K) [16]. Note that

$$[x, y^q] = [x, y](y^{-1}[x, y]y)(y^{-2}[x, y]y^2) \cdots (y^{-(q-1)}[x, y]y^{q-1}),$$

where $[a, b] = a^{-1}b^{-1}ab$. However, $[x, y^q] = [x, x^p] = 1$ in G(K). Since $[x, y] \neq 1$, this implies that [x, y] is a generalized torsion element in G(K).

Recall that $\pi_1(K(r))$ is a quotient group of G(K). Since a non-trivial finite cyclic group obviously has a torsion element, we prove that for every non-abelian (equivalently, non-cyclic) quotient group of the knot group G(K), the image of [x,y]remains a generalized torsion element.

Let \overline{G} be a non-abelian quotient group of G(K) and we denote by \overline{g} the image of $g \in G(K)$ under the quotient map $G(K) \to \overline{G}$. Then we can assume that $\overline{[x,y]} \neq 1$, for otherwise \overline{G} is a cyclic group. Thus $\overline{[x,y]}$ remains to be a generalized torsion element in \overline{G} .

- (2) Let K be a (p,q)-cable of a knot k. We may decompose E(K) as the union of E(k) and the (p,q)-cable space $C_{p,q}$, which is a Seifert fibered manifold over the annulus with an exceptional fiber of index $q \geq 2$. Then K(m/n) is the union of E(k) and $C_{p,q} \cup_{m/n} (S^1 \times D^2)$. By the assumption $C_{p,q} \cup_{m/n} (S^1 \times D^2)$ is a Seifert fibered manifold over the disk with two exceptional fibers of indices q and $|pqn-m| \geq 2$, or the connected sum of the solid torus and a lens space if |pqn-m|=0. In the former case, [13, Lemma 3.5] shows that the subgroup $\pi_1(C_{p,q} \cup_{m/n} (S^1 \times D^2))$ of $\pi_1(K(m/n))$ has a generalized torsion element. In the latter case, $\pi_1(K(m/n))$ has a torsion element.
- (3) We may write $K = K_1 \sharp K_2$, where K_1 and K_2 are both non-trivial knots. Then E(K) consists of $E(K_1)$, $E(K_2)$ and the 2-fold composing space $C = [\text{disk with 2 holes}] \times S^1$, where $\partial = \partial E(K) \cup \partial E(K_1) \cup \partial E(K_2)$. Note that the S^1 -fiber of C on $\partial E(K)$ is a meridian of K. Thus for any non-integral slope m/n $(n \geq 2)$, $C \cup_{m/n} (S^1 \times D^2)$ is a Seifert fibered manifold over the annulus with an exceptional fiber of index $n \geq 2$. Hence it follows from [13, Lemma 3.5] that the subgroup $\pi_1(C \cup_{m/n} (S^1 \times D^2))$ of $\pi_1(K(m/n))$ has a generalized torsion element. \square

Remark 2.1. In (3) we may decompose $(K_1\sharp K_2)(r)$ as $E(K_1), E(K_2)$ and $C \cup_{m/n} (S^1 \times D^2)$. If one of $G(K_1)$ and $G(K_2)$ has a generalized torsion element g, then it remains a generalized torsion element in $\pi_1((K_1\sharp K_2)(r))$ for all $r \in \mathbb{Q}$. Even when none of $G(K_1)$ and $G(K_2)$ has a generalized torsion element, Theorem 1.3(3) says that $\pi_1((K_1\sharp K_2)(r))$ has such an element if $r \in \mathbb{Q}$ is non-integral. In this case the generalized torsion element lies in the fundamental group of a decomposing piece $C \cup_{m/n} (S^1 \times D^2)$.

In the case where r is integral, the fundamental group of $C \cup_{m/n} (S^1 \times D^2)$ does not have a generalized torsion element [2, Theorem 1.5]. So (3) does not hold in general.

Let us take $K_i = K_{q_i}$, a positive twist knot, whose knot group $G(K_i)$ has no generalized torsion element [4]. Then as above the fundamental group of each decomposing piece has no generalized torsion element. However, applying Theorem 1.4, we show that $\pi_1((K_1\sharp K_2)(r))$ has a generalized torsion element if $r \geq 4$; see [11].

3. Generalized torsion elements arising from singular spanning disks

In this section we prove theorem 1.4. First we recall the following observation in [6]:

Lemma 3.1. Suppose that K has a (p,q)-singular spanning disk, then we have a factorization of the slope element $\lambda \mu^{p-q}$ as a product of p conjugates of λ and q conjugates of λ^{-1} .

Proof. Let $\Phi \colon D^2 \to S^3$ be a (p,q)-singular spanning disk of K. By restricting Φ on a slightly smaller subdisk $D \subset D^2$, $c := \Phi(\partial D)$ is a simple closed curve on $\partial N(K)$, which has the slope p-q hence it represents $\lambda \mu^{p-q} \in G(K)$.

Let $x_1, \ldots, x_p \in D$ (resp. y_1, \ldots, y_q) be the preimage of the positive (resp. negative) intersections of K and $\Phi(D)$. We take small oriented loops c_i around x_i (resp. d_j around y_j) so that the homological sum $[c_1] + \cdots + [c_p] + [d_1] + \cdots + [d_j] = [\partial D]$ in $H_1(D - \{x_1, \ldots, x_p, y_1, \ldots, y_q\})$. Then $\Phi(c_i) = \mu_{c_i}$ and $\Phi(d_j) = \mu_{d_j}^{-1}$. See Figure 3.

Take a base point $z \in \partial D$ of D so that $\Phi(z)$ is a base point of E(K). Then we take paths a_i in D from z to a point on a loop c_i , and paths b_j in D from z to a point on a loop d_j , so that their concatenation

$$(a_1*c_1*\overline{a_1})*\cdots*(a_p*c_p*\overline{a_p})*(b_1*d_1*\overline{b_1})*\cdots*(b_q*d_q*\overline{b_q})$$

is homotopic to ∂D as a based loop in $D-\{x_1,\ldots,x_p,y_1,\ldots,y_q\}$. Here * represents the concatenation of paths, and \overline{a} means the path a with opposite orientation; see Figure 3.

Then

$$[\Phi(a_i * c_i * \overline{a_i})] = [\Phi(a_i) * \Phi(c_i) * \overline{\Phi(a_i)}] = [a_i' * \mu_{c_i} * \overline{a_i'}] = \alpha_i \mu \alpha_i^{-1} \in G(K),$$

where we slide μ_{c_i} , together with a'_i , on $\partial N(K)$, so that μ_{c_i} becomes μ and a'_i becomes a loop representing α_i . Similarly

$$[\Phi(b_j*d_j*\overline{b_j})] = [\Phi(b_j)*\Phi(d_j)*\overline{\Phi(b_j)}] = [b_j'*\mu_{d_j}^{-1}*\overline{b_j'}] = \beta_j\mu^{-1}\beta_j^{-1} \in G(K),$$

where we slide $\mu_{d_j}^{-1}$, together with b'_j , on $\partial N(K)$, so that $\mu_{d_j}^{-1}$ becomes μ^{-1} and b'_j becomes a loop representing β_i .

Therefore we obtain a factorization

$$\lambda \mu^{p-q} = (\alpha_1 \mu \alpha_1^{-1}) \cdots (\alpha_p \mu \alpha_p^{-1}) (\beta_1 \mu^{-1} \beta_1^{-1}) \cdots (\beta_q \mu^{-1} \beta_q^{-1})$$

of the slope element $\lambda \mu^{p-q}$ as a product of p conjugates of μ and q conjugates of μ^{-1} .




FIGURE 3. The singular spanning disk gives rise to a factorization of a slope element $\lambda \mu^{p-q}$.

Proof of Theorem 1.4. We prove the theorem for the case K admits a (p, 0)-singular spanning disk. The case where K admits a (0, q)-singular spanning disk is similar.

By Lemma 3.1, from a (p,0)-singular spanning disk, we get a factorization of $\lambda \mu^p$ as a product of p conjugates of meridian μ

$$\lambda \mu^p = (\alpha_1 \mu \alpha_1^{-1}) \cdots (\alpha_p \mu \alpha_p^{-1}).$$

Therefore,

$$\lambda^n \mu^m = (\lambda \mu^p)^n \mu^{m-pn} = ((\alpha_1 \mu \alpha_1^{-1}) \cdots (\alpha_p \mu \alpha_p^{-1}))^n \mu^{m-pn}.$$

In $\pi_1(K(m/n))$, $\lambda^n \mu^m = 1$. Hence

$$((\alpha_1 \mu \alpha_1^{-1}) \cdots (\alpha_p \mu \alpha_p^{-1}))^n \mu^{m-pn} = 1,$$

and the meridian μ becomes a generalized torsion element of order $\leq m$ in $\pi_1(K(m/n))$ if $m/n \geq p$. Since $H_1(K(m/n)) = \mathbb{Z}_m$ and $[\mu]$ is its generator, μ is indeed a generalized torsion element of order m.

4. Dehn surgery along genus one two-bridge knot

Let g be a non-trivial element in a group G. Denote by $\langle \langle g \rangle \rangle^+$ the semigroup consisting of non-empty finite products of conjugates of g. Then a non-trivial element g is a generalized torsion element if and only if $1 \in \langle \langle g \rangle \rangle^+$.

Since

$$y(\prod_{i=1}^{n} (x_i g x_i^{-1})) y^{-1} = \prod_{i=1}^{n} (y x_i) g(y x_i)^{-1},$$

we have the following.

• $a \in \langle\langle g \rangle\rangle^+$ implies that $\langle\langle a \rangle\rangle^+ \subset \langle\langle g \rangle\rangle^+$.

We collect some useful properties as a lemma below, which will be repeatedly used in the proof of Theorem 1.6. Recall that the commutator $g^{-1}h^{-1}gh$ is denoted by [g, h].

Lemma 4.1. Let g, h, x be elements in G. Then the following holds.

- (1) $g^n h^n \in \langle \langle gh \rangle \rangle^+$ for all n > 0.
- (2) If $gh \in \langle \langle x \rangle \rangle^+$, then $g^n h^n \in \langle \langle x \rangle \rangle^+$ for all n > 0.
- (3) If $[g,h] \in \langle\langle x \rangle\rangle^+$, then $[g^n,h^m] \in \langle\langle x \rangle\rangle^+$ for all n,m>0.

Proof. (1) follows from the equality:

$$g^n h^n = (g^{n-1}(gh)g^{-(n-1)})(g^{n-2}(gh)g^{-(n-2)})\cdots(gh).$$

- (2) If $gh \in \langle \langle x \rangle \rangle^+$, then $\langle \langle gh \rangle \rangle^+ \subset \langle \langle x \rangle \rangle^+$. Then (1) shows that $g^n h^m \in \langle \langle gh \rangle \rangle^+ \subset \langle \langle x \rangle \rangle^+$.
- (3) Assume that $[g,h] \in \langle\!\langle x \rangle\!\rangle^+$. Then $\langle\!\langle [g,h] \rangle\!\rangle^+ = \langle\!\langle g^{-1}(h^{-1}gh) \rangle\!\rangle^+ \subset \langle\!\langle x \rangle\!\rangle^+$. Apply (2) to see that $(g^{-n}h^{-1}g^n)h = g^{-n}(h^{-1}gh)^n \in \langle\!\langle x \rangle\!\rangle^+$. Then apply (2) again to see that $[g^n,h^m] = g^{-n}h^{-m}g^nh^m = (g^{-n}h^{-1}g^n)^mh^m \in \langle\!\langle x \rangle\!\rangle^+$.

Proof of Theorem 1.6. We take elements a, b, t of G(K) as indicated in Figure 4.




FIGURE 4. C[2p, 2q] (p = 2, q = 3); a, b, t are generator of the knot group G(K).

The knot group G(K) of K = C[2p, 2q] has a presentation

(4.1)
$$G(K) = \langle a, b, t \mid ta^p t^{-1} = b^{-1} a^p, \ tb^{-q} a^{-1} t^{-1} = b^{-q} \rangle.$$

(This is the so-called Lin presentation which is obtained by using the Seifert surface. See [20].)

Since the meridian and the longitude are given by t and $[b^q, a^p]$, respectively, m/n-surgery adds additional relation $t^m[b^q, a^p]^n = 1$. Hence we have:

$$\pi_1(K(m/n)) = \langle a, b, t \mid ta^p t^{-1} = b^{-1}a^p, \ tb^{-q}a^{-1}t^{-1} = b^{-q}, \ t^m[b^q, a^p]^n = 1 \rangle.$$

From the first and second relations, we have $b = [a^{-p}, t^{-1}]$ and $a = [t, b^{-q}]$.

First we prove (1). Assume that $n \ge 1$ and $m \ge (2n-1)p$. We show $a^p, a^{-p} \in \langle\!\langle t \rangle\!\rangle^+$. This then implies that $1 = a^p a^{-p} \in \langle\!\langle t \rangle\!\rangle^+$, so t is a generalized torsion element

Since $ta = b^q t b^{-q} \in \langle \langle t \rangle \rangle^+$ (from the second relation), and $b^q t b^{-q} a^{-1} = t \in \langle \langle t \rangle \rangle^+$, Lemma 4.1(2) shows that

$$(4.2) t^p a^p, b^q t^p b^{-q} a^{-p} = (b^q t b^{-q})^p (a^{-1})^p \in \langle \langle t \rangle \rangle^+.$$

Then

$$\begin{array}{lcl} t^{2p}[b^q,a^p] & = & t^{2p}b^{-q}(b^qt^{-p}b^{-q})(b^qt^pb^{-q})a^{-p}b^qt^{-p}t^pa^p \\ & = & (t^pb^{-q})(b^qt^pb^{-q}a^{-p})(t^pb^{-q})^{-1}\cdot(t^pa^p). \end{array}$$

Hence $t^{2p}[b^q, a^p] \in \langle \langle t \rangle \rangle^+$. By Lemma 4.1(2), we conclude

$$(4.3) t^{2(n-1)p}[b^q, a^p]^{n-1} \in \langle \langle t \rangle \rangle^+.$$

From the third relation for $\pi_1(K(m/n))$, we have

$$t^m[b^q,a^p]^n=t^m[b^q,a^p]^{n-1}[b^q,a^p]=t^m[b^q,a^p]^{n-1}\cdot b^{-q}a^{-p}b^qa^p=1.$$

This gives

$$a^p = b^q a^p t^m [b^q, a^p]^{n-1} b^{-q}$$
 and $a^{-p} = t^m [b^q, a^p]^{n-1} \cdot b^{-q} a^{-p} b^q$.

Then

$$\begin{array}{lll} a^p & = & b^q \cdot t^{-p} (t^p a^p) t^m t^{-2(n-1)p} (t^{2(n-1)p} [b^q, a^p]^{n-1}) b^{-q} \\ & = & b^q \left(t^{-p} (t^p a^p) t^p \cdot t^{m-(2n-1)p} \cdot (t^{2(n-1)p} [b^q, a^p]^{n-1}) \right) b^{-q}, \end{array}$$

and

$$\begin{array}{lll} a^{-p} & = & t^m \cdot t^{-2(n-1)p} (t^{2(n-1)p} [b^q, a^p]^{n-1}) \cdot b^{-q} (b^q t^{-p} b^{-q}) (b^q t^p b^{-q} a^{-p}) b^q \\ & = & t^{m-2(n-1)p} (t^{2(n-1)p} [b^q, a^p]^{n-1}) t^{-m+2(n-1)p} \cdot t^{m-(2n-1)p} \cdot b^{-q} (b^q t^p b^{-q} a^{-p}) b^q. \end{array}$$

Since
$$m - (2n - 1)p \ge 0$$
, by (4.2), (4.3) we conclude $a^p, a^{-p} \in \langle \langle t \rangle \rangle^+$.

Next we prove (2). Assume that $n \ge 1$, q > 0 and $m \le -(2n-1)q$. we show $b^q, b^{-q} \in \langle\langle t \rangle\rangle^+$.

Since $bt = a^p t a^{-p} \in \langle \langle t \rangle \rangle^+$ (from the first relation) and $b^{-1} a^p t a^{-p} = t \in \langle \langle t \rangle \rangle^+$, Lemma 4.1(2) shows that

$$(4.4) b^q t^q, b^{-q} a^p t^q a^{-p} = (b^{-1})^q (a^p t a^{-p})^q \in \langle \langle t \rangle \rangle^+.$$

Then

$$\begin{array}{lll} t^{2q}[a^p,b^q] & = & t^{2q}a^{-p}b^{-q}(a^pt^qa^{-p})(a^pt^{-q}a^{-p})a^pb^qt^qt^{-q} \\ & = & t^{2q}a^{-p}(b^{-q}a^pt^qa^{-p})a^pt^{-2q}\cdot t^q(b^qt^q)t^{-q}. \end{array}$$

Hence $t^{2q}[a^p, b^q] \in \langle \langle t \rangle \rangle^+$. By Lemma 4.1(2) we get

(4.5)
$$t^{2(n-1)q}[a^p, b^q]^{n-1} \in \langle\!\langle t \rangle\!\rangle^+.$$

From the third relation for $\pi_1(K(m/n))$, we have

$$t^m[b^q,a^p]^n=t^m[b^q,a^p][b^q,a^p]^{n-1}=t^m(b^{-q}a^{-p}b^qa^p)[b^q,a^p]^{n-1}=1.$$

This gives

$$b^{q} = a^{p}b^{q}t^{-m}[a^{p}, b^{q}]^{n-1}a^{-p}$$
 and $b^{-q} = t^{-m}[a^{p}, b^{q}]^{n-1}a^{-p}b^{-q}a^{p}$.

Therefore

$$\begin{array}{lcl} b^q & = & a^p (b^q t^q) t^{-q} t^{-m} t^{-2(n-1)q} (t^{2(n-1)q} [a^p, b^q]^{n-1}) a^{-p} \\ & = & a^p \big((b^q t^q) \cdot t^{-m - (2n-1)q} \cdot (t^{2(n-1)q} [a^p, b^q]^{n-1}) \big) a^{-p}, \end{array}$$

and

$$\begin{array}{lll} b^{-q} & = & t^{-m}t^{-2(n-1)q}t^{2(n-1)q}[a^p,b^q]^{n-1}a^{-p}b^{-q}(a^pt^qa^{-p})(a^pt^{-q}a^{-p})a^p \\ & = & t^{-m-2(n-1)q}(t^{2(n-1)q}[a^p,b^q]^{n-1})t^{-q}t^qa^{-p}(b^{-q}a^pt^qa^{-p})a^pt^{-q} \\ & = & t^{-m-(2n-1)q}\cdot t^q(t^{2(n-1)q}[a^p,b^q]^{n-1})t^{-q}\cdot t^qa^{-p}(b^{-q}a^pt^qa^{-p})a^pt^{-q}. \end{array}$$

Since $-m - (2n-1)q \ge 0$, by (4.4), (4.5) we conclude that $b^q, b^{-q} \in \langle \langle t \rangle \rangle^+$.

Finally we prove (3). Assume that $n \ge 1$, q < 0 and $m \ge -(2n-1)q$. We show $b^q, b^{-q} \in \langle\!\langle t \rangle\!\rangle^+$. As in (ii), $bt = a^p t a^{-p} \in \langle\!\langle t \rangle\!\rangle^+$ and $b^{-1} a^p t a^{-p} = t \in \langle\!\langle t \rangle\!\rangle^+$ imply that

$$b^{-q}t^{-q}, b^qa^pt^{-q}a^{-p} = (b^{-1})^{-q}(a^pta^{-p})^{-q} \in \langle \! \langle t \rangle \! \rangle^+.$$

Then

$$\begin{aligned} [b^q, a^p] t^{-2q} &= b^{-q} a^{-p} (b^q a^p t^{-q} a^{-p}) a^p t^{-q} \\ &= (b^{-q} t^{-q}) \cdot t^q a^{-p} (b^q a^p t^{-q} a^{-p}) a^p t^{-q} \in \langle\!\langle t \rangle\!\rangle^+. \end{aligned}$$

Lemma 4.1(2) gives

$$[b^q, a^p]^{n-1}t^{-2(n-1)q} \in \langle \langle t \rangle \rangle^+.$$

From the third relation for $\pi_1(K(m/n))$, we have

$$t^{m}[b^{q}, a^{p}]^{n} = t^{m}[b^{q}, a^{p}][b^{q}, a^{p}]^{n-1} = t^{m}(b^{-q}a^{-p}b^{q}a^{p})[b^{q}, a^{p}]^{n-1} = 1.$$

This gives

$$b^{q} = a^{-p}b^{q}a^{p}[b^{q}, a^{p}]^{n-1}t^{m}$$
 and $b^{-q} = a^{p}[b^{q}, a^{p}]^{n-1}t^{m}b^{-q}a^{-p}$.

Therefore

$$\begin{array}{lcl} b^q & = & a^{-p}b^qa^pt^{-q}a^{-p}a^pt^q[b^q,a^p]^{n-1}t^{-2(n-1)q}t^{2(n-1)q}t^m \\ & = & a^{-p}(b^qa^pt^{-q}a^{-p})a^p\cdot t^q([b^q,a^p]^{n-1}t^{-2(n-1)q})t^{-q}\cdot t^{(2n-1)q+m}, \end{array}$$

and

$$\begin{array}{lll} b^{-q} & = & a^p ([b^q, a^q]^{n-1} t^{-2(n-1)q}) t^{2(n-1)q} t^m b^{-q} a^{-p} \\ & = & a^p \big([b^q, a^q]^{n-1} t^{-2(n-1)q} \cdot t^{(2n-1)q+m} \cdot t^{-q} (b^{-q} t^{-q}) t^q \big) a^{-p}. \end{array}$$

Since $m + (2n - 1)q \ge 0$, we conclude that $b^q, b^{-q} \in \langle \langle t \rangle \rangle^+$.

Remark 4.2. The Alexander polynomial of K = C[2p, 2q] (p > 0) is $\Delta_K(t) = -(pq)t + (2pq - 1) - (pq)t^{-1}$.

If q > 0, then all the roots of the $\Delta_K(t)$ are positive real. Known bi-orderability criterion [18, 19, 3] says that under additional assumption (such as fiberedness), the knot group G(K) is bi-orderable if all the roots of the Alexander polynomial $\Delta_K(t)$ are positive real. In a light of this, we expect G(K) is bi-orderable, although this is confirmed only for p = 1 at present.

On the other hand, if q < 0, then $\Delta_K(t)$ has no positive real roots. By [4, Theorem 3.3] or [10], this shows that G(K) is not bi-orderable. In a light of Conjecture 1.1, G(K) would have a generalized torsion element, but this is confirmed only for p = 1 again.

5. Further examples

5.1. Generalized Whitehead doubles. Let us take a standardly embedded solid torus V in S^3 and a knot k_{ω}^{τ} in V as depicted in Figure 5 (Left). Let $f: V \to S^3$ be an orientation preserving embedding such that the core of f(V) is a non-trivial knot $k \subset S^3$ and f sends a preferred longitude of V to that of k. Then the image $f(k_{\omega}^{\tau})$ is called a τ -twisted, ω -generalized Whitehead double of k; see Figure 5. If $\omega = -1$, then it is a usual positive Whitehead double of k.




FIGURE 5. f is a faithful embedding of V into S^3 which sends the longitude of V to that of k.

Corollary 5.1. Let K be a τ -twisted, ω -generalized Whitehead double of a knot k ($\omega < 0$). Then $\pi_1(K(r))$ has a generalized torsion element whenever $r \ge 2\omega$.

Proof. As shown in Figure 5, K bounds a $(2\omega, 0)$ -singular spanning disk. Apply Theorem 1.4 to obtain the desired result.

5.2. **Montesinos knots.** A tangle R = (B, t) is a pair of a 3-ball B (which is the unit 3-ball in \mathbb{R}^3) and two disjoint arcs t properly embedded in B. We say that a tangle (B,t) is trivial if there is a pairwise homeomorphism from (B,t) to $(D^2 \times I, \{x_1, x_2\} \times I)$, where x_1, x_2 are distinct points. Two tangles (B,t) and (B,t') with $\partial t = \partial t'$ are equivalent if there is a pairwise homeomorphism $h: (B,t) \to (B,t')$ which is the identity on ∂B .

Take 4 points NW, NE, SE, SW on the boundary of B so that NW = $(0, -\alpha, \alpha)$, NE = $(0, \alpha, \alpha)$, SE = $(0, \alpha, -\alpha)$, SW = $(0, -\alpha, -\alpha)$, where $\alpha = \frac{1}{\sqrt{2}}$. A tangle (B, t)

 $(\partial t = \{\text{NW}, \text{NE}, \text{SE}, \text{SW}\})$ is rational if it is a trivial tangle. Any rational tangle can be constructed from a sequence of integers a_1, a_2, \ldots, a_n as shown in Figure 6, where only the last horizontal twist a_n may be 0. We denote the resulting tangle by $[a_1, a_2, \ldots, a_n]$. We say that a rational tangle denoted by $[a_1, a_2, \ldots, a_n]$ is odd type (resp. $even\ type$) if n is odd (resp. even).

A Montesinos knot $M(R_1, ..., R_m)$ is a knot which has a diagram in Figure 7 (Top-left), where R_i is a rational tangle $[a_{i,1}, a_{i,2}, ..., a_{i,n_i}]$.




FIGURE 6. Rational tangle $[a_1, a_2, \ldots, a_n]$




FIGURE 7. Montesinos knot with (p, 0)-singular spanning disk

We say that a Montesinos knot $K = M(R_1, ..., R_m)$ satisfies the Condition (*) if

- (1) R_1, \dots, R_{m-1} are odd type rational tangles and R_m is an even type rational tangle,
- (2) for each odd type rational tangle $R_i = [a_{i,1}, a_{i,2}, \dots, a_{i,n_i}]$ $(1 \le i \le m-1)$, $a_{i,j}$ is even and $a_{i,\text{even}} < 0$ (Bottom-left of Figure 7), and

(3) for the even type rational tangle $R_m = [a_{m,1}, a_{m,2}, \dots, a_{m,n_m}], a_{m,\text{odd}}$ is positive even and $a_{m,\text{even}}$ is odd (Top-right of Figure 7), or $a_{m,\text{odd}}$ is negative even and $a_{m,\text{even}}$ is even (Bottom-right of Figure 7).

Put

$$c(K) = c(M(R_1, \dots, R_m)) = \sum_{1 \le i \le m-1} a_{i,\text{even}} + \sum |a_{m,\text{odd}}|$$

Corollary 5.2. Let $K = M(R_1, ..., R_m)$ be a Montesinos knot which satisfies the Condition (*). Then $\pi_1(K(r))$ has a generalized torsion whenever $r \ge c(K)$.

Proof. Following Figure 7, we see that K bounds a (c(K), 0)-coherent clasp disk.

5.3. Positive knots, almost positive knots and thier slight generalization.

A knot is said to be *positive* (resp. almost positive) if it admits a diagram whose crossings are all positive (resp. all positive except one). More generally, we can handle a knot which admits a diagram whose negative crossings appear successively along a single overarc as in Figure 8. If k=0 (resp. 1), the diagram D is a positive (resp. almost positive) diagram.




FIGURE 8. Diagram D has successive k negative crossings and p-k positive crossings.

Theorem 5.3. Let K be a knot which admits a diagram D with p crossings and k negative crossings that appear successively along an overarc. If $r \geq p - k$, then $\pi_1(K(r))$ has a generalized torsion element.

Proof. In D, assign the meridian generators t_1, t_2, \ldots, t_d of G(K) for the overarcs along the knot. Here, we choose t_1 for the overarc running over k successive negative crossings. Note that any t_i ($i \neq 1$) is a conjugate of t_1 in G(K). Then if we traverse the longitude λ from the overarc running over the negative crossings, then

$$\lambda = t_1^{-(p-2k)} \cdot t_{i(1)} \cdots t_{i(s_1)}^{-1} \cdots t_{i(s_2)}^{-1} \cdots t_{i(s_k)}^{-1} \cdots t_{i(d)}^{-1},$$

where $i(s_j) = 1$ (j = 1, 2, ..., k) and the others are not equal to 1. We should remark that the writhe of D is p - 2k.

We may rewrite

$$\lambda = t_1^{-(p-2k)} \cdot U_1 t_1^{-1} U_2 t_1^{-1} \cdots U_k t_1^{-1} \cdot V$$

$$= t_1^{-(p-k)} (t_1^k U_1 t_1^{-k}) (t_1^{k-1} U_2 t_1^{-(k-1)}) \cdots (t_1^2 U_{k-1} t_1^{-2}) (t_1 U_k t_1^{-1}) \cdot V$$

$$= t_1^{-(p-k)} W,$$

where U_i , V and W are products of conjugates of t_1 . In $\pi_1(K(m/n))$, the surgery relation is $t_1^m \lambda^n = 1$. Since

$$\begin{split} t_1^m \lambda^n &= t_1^m (t_1^{-(p-k)} W)^n \\ &= t_1^m \cdot t_1^{-n(p-k)} (t_1^{(n-1)(p-k)} W t_1^{-(n-1)(p-k)}) \cdots \\ &\quad (t_1^{2(p-k)} W t_1^{-2(p-k)}) (t_1^{p-k} W t_1^{-(p-k)}) W \\ &= t_1^{m-n(p-k)} (t_1^{(n-1)(p-k)} W t_1^{-(n-1)(p-k)}) \cdots \\ &\quad (t_1^{2(p-k)} W t_1^{-2(p-k)}) (t_1^{p-k} W t_1^{-(p-k)}) W, \end{split}$$

the meridian t_1 gives a generalized torsion element if $m - n(p - k) \ge 0$.

The special case where k = 0 or 1 of Theorem 5.3 immediately gives the following.

Corollary 5.4. Let K be a positive resp.analmost positive knot in S^3 with positive resp.almost positive diagram D. Let p be the number of crossings of D. Then $\pi_1(K(r))$ admits a generalized torsion element whenever $r \geq p$ (resp. $r \geq p-1$).

Remark 5.5. For positive knots, we have another way to give the same result. Suppose that K is embedded in the boundary of a 3-ball B^3 except p over arcs. Let ∞ is the center of B^3 and take a singular disk D with center P and $\partial D = K$. Since K is positive, D is a (p,0)-singular spanning disk and Theorem 1.4 gives the desired conclusion. We should remark that this singular disk is not a clasp disk, because it has a branch point P.




Figure 9. Cone of positive diagram yields a (p,0)-singular spanning disk of K

6. Questions

As we mentioned in Section 1, there are two kinds of generalized torsion elements of $\pi_1(K(r))$:

- A generalized torsion element which is the image of a generalized torsion element of G(K)
- A generalized torsion element which is the image of a non-generalized torsion element of G(K)

For the first kind of generalized torsion elements, a generalized torsion element of G(K) always belongs to the commutator subgroup, so it always vanishes in $\pi_1(K(r))$ for a cyclic surgery slope r. However, the proof of Theorem 1.3(1) shows that for a torus knot $K = T_{p,q}$, $g = [x,y] \in [G(K), G(K)] \subset G(K)$ is a generalized torsion element, which becomes a generalized torsion element in $\pi_1(K(r))$ for all $r \in \mathbb{Q}$ except cyclic surgery slopes. This raises the following question;

Question 6.1. Let K be a non-trivial knot such that G(K) has a generalized torsion element g. Does g remain a generalized torsion element in $\pi_1(K(r))$ for all $r \in \mathbb{Q}$ except cyclic surgery slopes?

Note that if K admits a cyclic surgery, then K is an L-space knot, and hence G(K) is not bi-orderable [5]. So following our conjecture we anticipate such a knot K admits a generalized torsion element.

For the second kind of generalized torsion elements, both Theorems 1.4 and 1.6 tell us that the image of a meridian becomes a generalized torsion element in $\pi_1(K(r))$ under suitable assumptions.

Since the abelianization $H_1(G(K); \mathbb{Z}) = H_1(E(K); \mathbb{Z}) = \mathbb{Z}$ is torsion-free, any generalized torsion element of G(K) is homologically trivial. Hence a meridian, the generator of $H_1(E(K); \mathbb{Z})$, cannot be a generalized torsion element in G(K). It is quite interesting to ask the following.

Question 6.2. Let K be a non-trivial knot in S^3 . Then is the image of a meridian μ a generalized torsion element in $\pi_1(K(r))$ if $r \neq 0, \infty$? In particular, does this hold if |r| is sufficiently large?

Question 6.3. Let K be a non-trivial knot in S^3 . Then does there exist a non-trivial element $g \in G(K)$ which satisfies (1) g is homologically trivial, but not a generalized torsion element in G(K), and (2) g becomes a generalized torsion element in $\pi_1(K(r))$ for some $r \in \mathbb{Q}$?

Acknowledgements

We would like to thank Stefan Friedl for his valuable comments. We would also like to thank the referee for careful reading and useful comments.

References

- V. V. Bludov and E. S. Lapshina, On ordering groups with a nilpotent commutant (in Russian), Sibirsk. Mat. Zh. 44 (2003), no. 3, 513–520; translation in Siberian Math. J. 44 (2003), no. 3, 405–410.
- [2] S. Boyer, D. Rolfsen and B. Wiest, Orderable 3-manifold groups, Ann. Inst. Fourier 55 (2005), 243–288.
- [3] I. Chiswell, A. Glass and J. Wilson, Residual nilpotence and ordering in one-relator groups and knot groups, Math. Proc. Cambridge Philos. Soc. 158 (2015), no. 2, 275–288.

- [4] A. Clay, C. Desmarais and P. Naylor, Testing bi-orderability of knot groups, Canad. Math. Bull. 59 (2016), no. 3, 472–482.
- [5] A. Clay, and D. Rolfsen, Ordered groups, eigenvalues, knots, surgery and L-spaces, Math. Proc. Camb. Phil. Soc. 152 (2012), 115–129.
- [6] M. Greene and B. Wiest, A natural framing of knots, Geom. Topol. 2 (1998), 31–64.
- [7] T. Homma, On Dehn Lemma for three sphere, Yokohama Math. J. 5 (1957), 223-244.
- [8] K. Ichihara, K. Motegi and M. Teragaito, Vanishing non-trivial elements in a knot group by Dehn fillings, Topology Appl. 264 (2019), 223–232.
- [9] T. Ito, Framing functions and a strengthened version of Dehn's lemma, J. Knot Theory Ramifications 25 (2016), no. 6, 1650031, 5 pp.
- [10] T. Ito, Alexander polynomial obstruction of bi-orderability for rationally homologically fibered knot groups, New York J. Math 23 (2017), 497–503.
- [11] T. Ito, K. Motegi and M. Teragaito, Generalized torsion and decomposition of 3-manifolds, Proc. Amer. Math. Soc. 147 (2019) 4999–5008.
- [12] P. Longobardi, M. Maj and A. Rhemtulla, On solvable R*-groups, J. Group Theory 6 (2003), no. 4, 499–503.
- [13] K. Motegi and M. Teragaito, Generalized torsion elements and bi-orderability of 3-manifold groups, Canad. Math. Bull. 60 (2017), no. 4, 830–844.
- [14] R. Mura and A. Rhemtulla, Solvable R*-groups, Math. Z. 142 (1975), 293-298.
- [15] R. Mura and A. Rhemtulla, Orderable groups, Lecture Notes in Pure and Applied Mathematics, Vol. 27. Marcel Dekker, Inc., New York-Basel, 1977.
- [16] G. Naylor and D. Rolfsen, Generalized torsion in knot groups, Canad. Math. Bull. 59 (2016), 182–189.
- [17] D. Papakyriakopoulos, On Dehn's lemma and the sphericity of knots, Ann. Math. 66 (1957), 1–26.
- [18] P. Perron and D. Rolfsen, On orderability of fibred knot groups, Math. Proc. Cambridge Philos. Soc. 135 (2003), 147–153.
- [19] P. Perron and D. Rolfsen, Invariant ordering of surface groups and 3-manifolds which fibre over S¹, Math. Proc. Cambridge Philos. Soc. 141 (2006), 273-280.
- [20] M. Teragaito, Generalized torsion elements in the knot groups of twist knots, Proc. Amer. Math. Soc. 144 (2016), no.6, 2677–2682.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN *E-mail address*: tetitoh@math.kyoto-u.ac.jp

Department of Mathematics, Nihon University, 3-25-40 Sakurajosui, Setagaya-ku, Tokyo 156–8550, Japan

 $E ext{-}mail\ address: motegi@math.chs.nihon-u.ac.jp}$

Department of Mathematics and Mathematics Education, Hiroshima University, 1-1-1 Kagamiyama, Higashi-Hiroshima, 739–8524, Japan

 $E ext{-}mail\ address: teragai@hiroshima-u.ac.jp}$