

New Families of Solutions for the Space Time Fractional Burgers' Equation

Abaker A. Hassaballa^{1,2}, Ahmed M. A. Adam^{1,3}, Eltayeb A. Yousif^{1,4}, and Mohamed I. Nouh⁵

¹ Department of Mathematics, Faculty of Science, Northern Border University, Arar 91431, Saudi Arabia

² Department of Mathematics, College of Applied & Industrial Sciences, Bahri University, Khartoum, Sudan

³ Faculty of Engineering, Alzaiem Alazhari University, Khartoum North 13311, Sudan

⁴ Department of Applied Mathematics, Faculty of Mathematical Sciences, University of Khartoum, Khartoum 11111, Sudan

⁵ Astronomy Department, National Research Institute of Astronomy and Geophysics (NRIAG), 11421 Helwan, Cairo, Egypt

Abstract

In this paper, the hyperbolic tangent function method is applied for constructing exact solutions for space-time conformal fractional Burger's equation. Furthermore, the space-time conformal fractional Burgers' equation is tested for the Painlevé property and consequently new numerous exact solutions are generated via Bäcklund transform.

Keywords: Conformal fractional Burgers' equation, hyperbolic tangent function method, Painlevé property, Bäcklund transform, fractional differential equations.

1. Introduction

Recently, fractional Calculus has made a very important impact to the most fields of science, such as mathematics, engineering, physics, economics, etc. [1-21]. The applications of fractional calculus are contemporary [4, 19]. Historically, the fractional derivatives introduced in different ways, for example, Riemann–Liouville, Riesz, Caputo, Modified Riemann–Liouville [1, 19]. In the year (2014), a new fractional derivative is defined by the authors Khalil et al [21], named conformal fractional derivative (CFD). The definition and basic concepts of CFD is developed [22]. Furthermore, the interpretations of CFD in engineering and physics applications are introduced and discussed [23]. The definition of CFD of a function $f: (0, \infty) \rightarrow \mathbb{R}$ of order α , where $0 < \alpha \leq 1$ is mainly given by the following limit

$$D^\alpha f(t) = \lim_{\varepsilon \rightarrow 0} \frac{f(t + \varepsilon t^{1-\alpha}) - f(t)}{\varepsilon} \quad (1.1)$$

The function f is said to be α -differentiable. The fractional derivative at $t = 0$ is given as

$$f^\alpha(0) = \lim_{t \rightarrow 0^+} D^\alpha f(t) \quad (1.2)$$

The advantages of CFD is that satisfying the properties of the classical integer derivatives [21-23].

Assume that f and g are α -differentiable functions and λ, a, b are constants, then the CFD satisfies the following properties:

i. $D^\alpha t^p = p t^{p-\alpha}, p \in \mathbb{R}$.

ii. $D^\alpha \lambda = 0$.

iii. $D^\alpha f(t) = t^{1-\alpha} \frac{df}{dt}$.

iv. $D^\alpha (af + bg) = aD^\alpha f + bD^\alpha g$.

v. $D^\alpha (fg) = fD^\alpha g + gD^\alpha f$.

vi. $D^\alpha \left(\frac{f}{g} \right) = \frac{gD^\alpha f - fD^\alpha g}{g^2}$.

vii. $D^\alpha (f(g(t))) = \frac{df}{dg} D^\alpha g(t) = t^{1-\alpha} \frac{df}{dg} \frac{dg}{dt}$.

In mathematical physics and many other phenomena in various fields of applied science are described by nonlinear models, particularly by nonlinear partial differential equations (NLPDEs), such as astrophysics, optics, fluid dynamics, mathematical biology, plasma physics and so on. It is important to search for the solutions of the concerned models to understand and interpret their physical mechanism and behavior. The search for exact solutions to NLPDE has become of great interest by mathematicians and physicists and they have exerted great efforts for that. There are many powerful and efficient methods for finding exact solutions have been introduced, such as the inverse scattering method [24, 25], Hirota method [26], Bäcklund transformation [27-29], Darboux transformation [30], hyperbolic tangent function method [31, 32], Jacobi elliptic function method [33, 34], truncated Painlevé expansion method [35-38], homogenous balance method [39, 40], and there are other various methods in the literature. The Bäcklund transformations (BT) are considered as powerful tools for integrable systems to relate NLPDEs and their solutions [27, 28, 29, 41, 42]. Up to now, the research is still devoted and ongoing for finding the BT, e.g from the Painlevé property [35-38], the Ablowitz-Kaup-Newell-Segur (AKNS) system [41], the nonclassical symmetries [42], etc.

One of the simple NLPDEs in mathematical physics is the Burger's equation that arising in many areas of science such as Navier-Stokes equations, traffic flow, and acoustics [43, 44]. Consider the space-time conformable fractional Burger's equation (CFBE)

$$D_t^\alpha u + u D_x^\alpha u = \sigma D_x^{\alpha\alpha} u, \quad (1.3)$$

where σ is arbitrary constants, $\alpha \in (0,1]$, and $D_x^{\alpha\alpha} u = D_x^\alpha(D_x^\alpha u)$. In a few years ago, many researchers are introduced the solutions of CFBE by different methods. The Hopf-Cole transform is applied to a time CFBE, subsequently the approximate analytical solution is founded by applying a Homotopy Analysis Method [45]. Also, the exact solution is obtained by using Fourier transform [46]. The residual power series method is introduced for finding approximate solutions of a time CFBE [47]. Also, the residual power series method is used in finding the solution of the space-time conformable fractional KdV-Burgers equation [48]. The solution of regular and singular space-time coupled CFBEs is formulated by applying double Laplace transform [49].

The rest of the paper is organized as follows: In Sec. 2, we show that the space-time CFBE possesses the Painlevé property. In Sec. 3 the exact solutions for the space-time CFBE is constructed based on the hyperbolic tangent function method. In Sec. 4, the BT is used for generating abundant new exact solutions for the space-time CFBE. The paper is concluded in Sec. 5.

2. Painlevé property for the space-time conformal fractional Burger's equation

In this section, we intent to test the Painlevé property for the space-time CFBE given by Eq. (1.3) following the approach introduced in the Refs. [35- 37, 50]. The space-time CFBE has the Painlevé property when all the movable singularities are simple poles. For Eq. (1.3) we let

$$u = u\left(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha}\right) = \phi^n \sum_{j=0}^{\infty} \phi^j u_j, \quad (2.1)$$

where $u_0 \neq 0$, n is an integer, $\phi = \phi\left(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha}\right)$ and $u_j = u_j\left(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha}\right)$ are analytic functions of $\left(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha}\right)$ in a neighborhood of $M = \left\{\left(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha}\right) : \phi\left(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha}\right) = 0\right\}$.

To determine the values of n we consider the ansatz

$$u = \phi^n u_0 \quad (2.2)$$

By using chain rule on Eq. (2.2) we obtain

$$\left. \begin{aligned} D_t^\alpha u &= nu_0 \phi^{n-1} D_t^\alpha \phi + \phi^n D_t^\alpha u_0 \\ D_x^\alpha u &= nu_0 \phi^{n-1} D_x^\alpha \phi + \phi^n D_x^\alpha u_0 \\ D_x^{\alpha\alpha} u &= nu_0 \phi^{n-1} D_x^{\alpha\alpha} \phi + n(n-1)u_0 \phi^{n-2} (D_x^\alpha \phi)^2 + 2n\phi^{n-1} D_x^\alpha \phi D_x^\alpha u_0 + \phi^n D_x^{\alpha\alpha} u_0. \end{aligned} \right\} (2.3)$$

By substituting Eqs. (2.2) and (2.3) into Eq. (1.3), we get

$$\begin{aligned} & nu_0 \phi^{n-1} D_t^\alpha \phi + \phi^n D_t^\alpha u_0 + nu_0^2 \phi^{2n-1} D_x^\alpha \phi + u_0 \phi^{2n} D_x^\alpha u_0 \\ &= n\sigma u_0 \phi^{n-1} D_x^{\alpha\alpha} \phi + n(n-1)\sigma u_0 \phi^{n-2} (D_x^\alpha \phi)^2 + 2n\sigma \phi^{n-1} D_x^\alpha \phi D_x^\alpha u_0 + \sigma \phi^n D_x^{\alpha\alpha} u_0. \end{aligned} \quad (2.4)$$

The dominant terms of Eq. (2.4) are ϕ^{2n-1} & ϕ^{n-2} . Balancing of these gives $n = -1$. Thus,

$$u_0 = -2\sigma D_x^\alpha \phi \quad (2.5)$$

Back to Eq. (2.1), we can consider the following Ansatz with resonance r as

$$u = u_0 \phi^{-1} + b\phi^{r-1} = -2\sigma \phi^{-1} D_x^\alpha \phi + b\phi^{r-1} \quad (2.6)$$

From Eq. (2.6) we get

$$\left. \begin{aligned} D_x^\alpha u &= 2\sigma \phi^{-2} (D_x^\alpha \phi)^2 - 2\sigma \phi^{-1} D_x^{\alpha\alpha} \phi + b(r-1)\phi^{r-2} D_x^\alpha \phi \\ &= 2\sigma \phi^{-2} (D_x^\alpha \phi)^2 + b(r-1)\phi^{r-2} D_x^\alpha \phi + \dots \\ D_x^{\alpha\alpha} u &= -4\sigma \phi^{-3} (D_x^\alpha \phi)^3 + 4\sigma \phi^{-2} D_x^\alpha \phi D_x^{\alpha\alpha} \phi + 2\sigma \phi^{-2} D_x^\alpha \phi D_x^{\alpha\alpha} \phi - 2\sigma \phi^{-1} D_x^{\alpha\alpha\alpha} \phi \\ &\quad + b(r-1)(r-2)\phi^{r-3} (D_x^\alpha \phi)^2 + b(r-1)\phi^{r-2} D_x^{\alpha\alpha} \phi \\ &= -4\sigma \phi^{-3} (D_x^\alpha \phi)^3 + b(r-1)(r-2)\phi^{r-3} (D_x^\alpha \phi)^2 + \dots \end{aligned} \right\} (2.7)$$

The fractional derivatives are rearranged in terms of powers of $D_x^\alpha \phi$. Now, By substituting Eqs.

(2.6) and (2.7) into $uD_x^\alpha u = \sigma D_x^{\alpha\alpha} u$, we have

$$\begin{aligned} & (-2\sigma \phi^{-1} D_x^\alpha \phi + b\phi^{r-1})(2\sigma \phi^{-2} (D_x^\alpha \phi)^2 + b(r-1)\phi^{r-2} D_x^\alpha \phi + \dots) \\ &= \sigma(-4\sigma \phi^{-3} (D_x^\alpha \phi)^3 + b(r-1)(r-2)\phi^{r-3} (D_x^\alpha \phi)^2 + \dots) \\ \Rightarrow & -2b(r-1)\sigma \phi^{r-3} (D_x^\alpha \phi)^2 + 2b\sigma \phi^{r-3} (D_x^\alpha \phi)^2 = \sigma b(r-1)(r-2)\phi^{r-3} (D_x^\alpha \phi)^2 + \dots \\ \Rightarrow & (-2r+2+2-r^2+3r-2)\sigma b\phi^{r-3} (D_x^\alpha \phi)^2 = \dots \\ \Rightarrow & -(r+1)(r-2)\sigma b\phi^{r-3} (D_x^\alpha \phi)^2 = \dots \\ \Rightarrow & r = -1 \text{ and } r = 2. \end{aligned}$$

If M is a singularity manifold, it is obtained that $n = -1$. By leading order analysis,

$$u = \phi^{-1} \sum_{j=0}^{\infty} \phi^j u_j = \sum_{j=0}^{\infty} \phi^{j-1} u_j, \quad (2.8)$$

From (2.8) we get

$$D_t^\alpha u = \left[\sum_{j=0}^{\infty} [(j-1)\phi^{j-2} u_j D_t^\alpha \phi + \phi^{j-1} D_t^\alpha u_j] \right] = \left[\sum_{j=0}^{\infty} [(j-2)\phi^{j-3} u_{j-1} D_t^\alpha \phi + \phi^{j-3} D_t^\alpha u_{j-2}] \right],$$

$$\begin{aligned}
D_x^\alpha u &= \left[\sum_{j=0}^{\infty} [(j-1)\phi^{j-2}u_j D_x^\alpha \phi + \phi^{j-1} D_x^\alpha u_j] \right] = \left[\sum_{j=0}^{\infty} [(j-1)\phi^{j-2}u_j D_x^\alpha \phi + \phi^{j-2} D_x^\alpha u_{j-1}] \right], \\
D_x^{\alpha\alpha} u &= \left[\sum_{j=0}^{\infty} [(j-1)(j-2)u_j \phi^{j-3} (D_x^\alpha \phi)^2 + (j-1)\phi^{j-2} u_j D_x^{\alpha\alpha} \phi + 2(j-1)\phi^{j-2} D_x^\alpha \phi D_x^\alpha u_j \right. \\
&\quad \left. + \phi^{j-1} D_x^{\alpha\alpha} u_j] \right], \\
&= \left[\sum_{j=0}^{\infty} [(j-1)(j-2)u_j \phi^{j-3} (D_x^\alpha \phi)^2 + (j-2)\phi^{j-3} u_{j-1} D_x^{\alpha\alpha} \phi \right. \\
&\quad \left. + 2(j-2)\phi^{j-3} D_x^\alpha \phi D_x^\alpha u_{j-1} + \phi^{j-3} D_x^{\alpha\alpha} u_{j-2}] \right].
\end{aligned}$$

The recursion relations for u_j are found to be

$$\begin{aligned}
&D_t^\alpha u_{j-2} + (j-2)u_{j-1} D_t^\alpha \phi + \sum_{m=0}^j u_{j-m} [D_x^\alpha u_{m-1} + (m-1)u_m D_x^\alpha \phi] \\
&= \sigma [(j-1)(j-2)u_j (D_x^\alpha \phi)^2 + (j-2)u_{j-1} D_x^{\alpha\alpha} \phi + 2(j-2)D_x^\alpha \phi D_x^\alpha u_{j-1} + D_x^{\alpha\alpha} u_{j-2}], \quad (2.9)
\end{aligned}$$

where $u_k = 0$ for $k = -1, -2, -3, \dots$

From the recurrence formula (2.9) we obtain for:

$$\begin{aligned}
j = 0 &\Rightarrow -u_0^2 D_x^\alpha \phi = 2\sigma u_0 (D_x^\alpha \phi)^2 \\
&\Rightarrow u_0 = -2\sigma D_x^\alpha \phi. \quad (2.10)
\end{aligned}$$

Eq. (2.10) is identical to Eq. (2.5).

$$\begin{aligned}
j = 1 &\Rightarrow -u_0 D_t^\alpha \phi + \sum_{m=0}^1 u_{1-m} [D_x^\alpha u_{m-1} + (m-1)u_m D_x^\alpha \phi] = -\sigma [u_0 D_x^{\alpha\alpha} \phi + 2D_x^\alpha \phi D_x^\alpha u_0] \\
&\Rightarrow -u_0 D_t^\alpha \phi - u_0 u_1 D_x^\alpha \phi + u_0 D_x^\alpha u_0 = -\sigma [u_0 D_x^{\alpha\alpha} \phi + 2D_x^\alpha \phi D_x^\alpha u_0] \\
&\quad = \sigma [2\sigma D_x^\alpha \phi D_x^{\alpha\alpha} \phi + 4\sigma D_x^\alpha \phi D_x^{\alpha\alpha} \phi] = 6\sigma^2 D_x^\alpha \phi D_x^{\alpha\alpha} \phi \\
&\Rightarrow -u_0 D_t^\alpha \phi - u_0 u_1 D_x^\alpha \phi + 4\sigma^2 D_x^\alpha \phi D_x^{\alpha\alpha} \phi = 6\sigma^2 D_x^\alpha \phi D_x^{\alpha\alpha} \phi \\
&\Rightarrow -u_0 D_t^\alpha \phi - u_0 u_1 D_x^\alpha \phi = 2\sigma^2 D_x^\alpha \phi D_x^{\alpha\alpha} \phi = -u_0 \sigma D_x^{\alpha\alpha} \phi \Rightarrow \\
&\Rightarrow D_t^\alpha \phi + u_1 D_x^\alpha \phi - \sigma D_x^{\alpha\alpha} \phi = 0 \quad (2.11)
\end{aligned}$$

$j = 2$ and considering Eqs. (2.10, 2.11), then from Eq. (2.9) we get

$$\begin{aligned}
D_t^\alpha u_0 + \sum_{m=0}^2 u_{2-m} [D_x^\alpha u_{m-1} + (m-1)u_m D_x^\alpha \phi] &= \sigma D_x^{\alpha\alpha} u_0 \\
\Rightarrow D_t^\alpha u_0 + u_1 D_x^\alpha u_0 + u_0 D_x^\alpha u_1 &= \sigma D_x^{\alpha\alpha} u_0 \\
\Rightarrow D_x^\alpha (D_t^\alpha \phi + u_1 D_x^\alpha \phi - \sigma D_x^{\alpha\alpha} \phi) &= 0
\end{aligned} \tag{2.12}$$

By Eq. (2.11) the compatibility condition Eq. (2.12) at $j = 2$ is satisfied identically. For $j = 3$, then from recurrence formula given by Eq. (2.9) we have

$$\begin{aligned}
D_t^\alpha u_1 + u_2 D_t^\alpha \phi + \sum_{m=0}^3 u_{3-m} [D_x^\alpha u_{m-1} + (m-1)u_m D_x^\alpha \phi] \\
= \sigma [2u_3 (D_x^\alpha \phi)^2 + u_2 D_x^{\alpha\alpha} \phi + 2D_x^\alpha \phi D_x^\alpha u_2 + D_x^{\alpha\alpha} u_1] \\
\Rightarrow [D_t^\alpha u_1 + u_1 D_x^\alpha u_1 - \sigma D_x^{\alpha\alpha} u_1] + u_2 [D_t^\alpha \phi + u_1 D_x^\alpha \phi - \sigma D_x^{\alpha\alpha} \phi] \\
+ [u_2 D_x^\alpha u_0 + u_0 D_x^\alpha u_2 - 2\sigma D_x^\alpha \phi D_x^\alpha u_2] + u_3 D_x^\alpha \phi [u_0 - 2\sigma D_x^\alpha \phi] = 0 \\
\Rightarrow [D_t^\alpha u_1 + u_1 D_x^\alpha u_1 - \sigma D_x^{\alpha\alpha} u_1] + [u_2 D_x^\alpha u_0 + 2u_0 D_x^\alpha u_2] + 2u_3 u_0 D_x^\alpha \phi = 0
\end{aligned} \tag{2.13}$$

Since the resonances occur at $r = -1, 2$, and (ϕ, u_2) are arbitrary functions of $(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha})$ in the expansion (2.13). If we let the arbitrary functions $u_2 = u_3 = 0$, then we get

$$D_t^\alpha u_1 + u_1 D_x^\alpha u_1 = \sigma D_x^{\alpha\alpha} u_1 \tag{2.14}$$

For $j = 3$ then Eq. (2.14) is automatically satisfied for u_1 .

for $j = 4$, $u_0 = -2\sigma D_x^\alpha \phi$, and $u_2 = u_3 = 0$ then from Eq. (2.9) we obtain

$$\begin{aligned}
\sum_{m=0}^4 u_{4-m} [D_x^\alpha u_{m-1} + (m-1)u_m D_x^\alpha \phi] &= 6\sigma u_4 (D_x^\alpha \phi)^2 \\
\Rightarrow u_4 D_x^\alpha \phi (2u_0 - 6\sigma D_x^\alpha \phi) &= 0 \Rightarrow u_4 = 0.
\end{aligned}$$

Thus, we conclude that all

$$u_j = 0, j \geq 2, \tag{2.15}$$

providing u_1 satisfies space-time CFBE (Eq. (2.14)). Hence, the space-time CFBE possesses the Painlevé property and the truncation of the Painlevé expansion Eq. (2.8) then takes the form

$$u = -\frac{2\sigma}{\phi} D_x^\alpha \phi + u_1, \tag{2.16}$$

which is the Bäcklund transform for the space-time CFBE. When $u_1 = 0$ then from Eq. (2.16) we obtain

$$u = -\frac{2\sigma}{\phi} D_x^\alpha \phi, \tag{2.17}$$

Eq. (2.17) yields the fractional Cole-Hopf transform. When $u_1 = \phi$ then we have

$$u = -\frac{2\sigma}{\phi} D_x^\alpha \phi + \phi, \quad (2.18)$$

where

$$D_t^\alpha \phi + \phi D_x^\alpha \phi = \sigma D_x^{\alpha\alpha} \phi \quad (2.19)$$

Eqs. (2.18) and (2.19) represent the Bäcklund transform for the space-time CFBE.

3. Exact solutions for the space-time conformal fractional Burger's equation

The hyperbolic tangent function method (tanh-method) is an efficient method that used for constructing the exact traveling wave solutions for NLPDEs. The Ansatz is considered as a power series in \tanh , where \tanh is introduced as a new variable. Moreover, the derivatives of \tanh are also given in terms of \tanh itself [31, 32]. In the following steps we seek to describe the general tanh-method for constructing the solutions of the space-time conformal fractional partial differential equations (CFPDEs). Consider a general space-time conformal fractional partial differential equation (space-time CFPDE):

$$H(u, D_t^\alpha u, D_x^\alpha u, D_t^{\alpha\alpha} u, D_t^\alpha (D_x^\alpha u), D_x^{\alpha\alpha} u, \dots) = 0. \quad (3.1)$$

In the following steps, we summarize the tanh-method for solving Eq. (3.1):

Step 1: consider the traveling wave solution of Eq. (3.1) as

$$u(x, t) = u(\xi) \quad \xi = \frac{k}{\alpha} (x^\alpha - \omega t^\alpha), \quad (3.2)$$

where k and ω are the wave number and wave velocity, respectively. Substitution of Eq. (3.2) into Eq. (3.1) produces the following ordinary differential equation for $u(\xi)$

$$\tilde{H}(u, u', u'', \dots) = 0 \quad u' = \frac{du}{d\xi}, \dots \text{ etc.} \quad (3.3)$$

Step 2: Assume that the solution of Eq. (3.3) can be expressed as a finite power series of $F(\xi)$

$$u(\xi) = a_0 + \sum_{j=1}^s a_j F^j(\xi), \quad a_s \neq 0, \quad (3.4)$$

where $s \in \mathbb{N}$, which is determined by balancing the highest power of the linear term with the highest power of the nonlinear term in Eq. (3.3), and a_j are constants to be determined. The new exact solutions of the space-time CFPDE can be obtained via the solutions of the Riccati equation that satisfied by \tanh function. Consider the required Riccati equation to be

$$F' = A + BF + CF^2, \quad ' \equiv \frac{d}{d\xi}, \quad (3.5)$$

where A, B and C are constants.

Step 3: Substitution of Eq. (3.4) into Eq. (3.3), generates system of algebraic equations for $a_0, a_1, \dots, a_s, \omega$, and k .

Step 4: Solution of the system obtained in step 3, produces the values of $a_0, a_1, \dots, a_s, \omega$ and k in terms of A, B and C . By substituting these results into Eq. (3.4), we obtain the general form of traveling wave solution of Eq. (3.1).

Choosing of each proper value for A, B and C in Eq. (3.5) corresponds a solution $F(\xi)$ of Eq. (3.5) that is could be one of the hyperbolic function or triangular function as follows.

Case 1: If $A = C = 1$, and $B = 0$, then Eq. (3.5) has the solutions, $\tan\xi$.

Case 2: If $A = C = -1$, and $B = 0$, then Eq. (3.5) has the solutions, $\cot\xi$.

Case 3: If $A = 1$, $B = 0$, and $C = -1$, then Eq. (3.5) has the solutions, $\tanh\xi, \coth\xi$.

Case 4: If $A = \frac{1}{2}$, $B = 0$ and, $C = -\frac{1}{2}$, then Eq. (3.5) has the solutions, $\tanh\xi \pm i \operatorname{sech}\xi$,

$$\coth\xi \pm \operatorname{csch}\xi, \frac{\tanh\xi}{1 \pm \operatorname{sech}\xi}, \frac{\coth\xi}{1 \pm i \operatorname{csch}\xi}, i^2 = -1.$$

Case 5: If $A = C = \frac{1}{2}$, and $B = 0$, then Eq. (3.5) has the solutions, $\tan\xi \pm \sec\xi, \csc\xi - \cot\xi$,

$$\frac{\tan\xi}{1 \pm \sec\xi}.$$

Case 6: If $A = C = -\frac{1}{2}$, and $B = 0$, then Eq. (3.5) has the solutions, $\cot\xi \pm \csc\xi, \sec\xi - \tan\xi$,

$$\frac{\cot\xi}{1 \pm \csc\xi}.$$

Case 7: If $A = 1, C = -4$, and $B = 0$, then Eq. (3.5) has the solutions, $\frac{\tanh\xi}{1 + \tanh^2\xi}$.

Case 8: If $A = 1, C = 4$, and $B = 0$, then Eq. (3.5) has the solutions, $\frac{\tan\xi}{1 - \tan^2\xi}$.

Case 9: If $A = -1, C = -4$, and $B = 0$, then Eq. (3.5) has the solutions, $\frac{\cot\xi}{1 - \cot^2\xi}$.

Case 10: If $A = 1, B = -2$, and $C = 2$, then Eq. (3.5) has the solutions, $\frac{\tan\xi}{1 + \tan\xi}$.

Case 11: If $A = 1$ and $B = C = 2$, then Eq. (3.5) has the solutions, $\frac{\tan\xi}{1 - \tan\xi}$.

Case 12: If $A = -1, B = 2$, and $C = -2$, then Eq. (3.5) has the solutions, $\frac{\cot\xi}{1 + \cot\xi}$.

Case 13: If $A = -1, B = C = -2$, then Eq. (3.5) has the solutions, $\frac{\cot\xi}{1 - \cot\xi}$.

Case 14: If $A = B = 0$ and $C \neq 0$, then Eq. (3.5) has the solutions, $\frac{-1}{C\xi + C_0}$.

Case 15: If $A \neq 0$, $C = 0$, and $B \neq 0$, then Eq. (3.5) has the solutions, $\frac{1}{B}(\exp(B\xi) - A)$.

Case 16: If $A = 0$ and $B = C = 1$, then Eq. (3.5) has the solutions, $\frac{\exp(\xi)}{1 - \exp(\xi)}$.

Case 17: If $A = 0$, and $B = C = \frac{1}{2}$, then Eq. (3.5) has the solutions, $\frac{\exp(\frac{\xi}{2})}{1 - \exp(\frac{\xi}{2})}$.

Case 18: If $A = -\frac{1}{2}$, $B = 0$, and $C = \frac{1}{2}$, then Eq. (3.5) has the solutions, $-\tanh\left(\frac{\xi}{2}\right)$, $-\coth\left(\frac{\xi}{2}\right)$.

Case 19: If $A = -1$, and $B = C = 2$, then Eq. (3.5) has the solutions, $-\frac{1}{2} - \frac{\sqrt{3}\tanh(\sqrt{3}\xi)}{2}$,
 $-\frac{1}{2} - \frac{\sqrt{3}\coth(\sqrt{3}\xi)}{2}$.

Case 20: If $A = 1$, $B = 1$, and $C = 1$, then Eq. (3.5) has the solutions, $-\frac{1}{2} + \frac{\sqrt{3}\tan\left(\frac{\sqrt{3}\xi}{2}\right)}{2}$,
 $-\frac{1}{2} - \frac{\sqrt{3}\cot\left(\frac{\sqrt{3}\xi}{2}\right)}{2}$.

Case 21: If $A = -4$, $B = 0$, and $C = 4$, then Eq. (3.5) has the solutions, $-\tanh(4\xi)$, $-\coth(4\xi)$.

Case 22: If $A = \frac{1}{2}$, $B = -1$, and $C = 1$, then Eq. (3.5) has the solutions, $\frac{1}{2} + \frac{1}{2}\tan\left(\frac{\xi}{2}\right)$,
 $\frac{1}{2} - \frac{1}{2}\cot\left(\frac{\xi}{2}\right)$.

Now, we need to implement the tanh-method technique into the space-time CFBE given by Eq. (1.3) to generate new exact solutions. Firstly, substituting the traveling wave solution given by Eq. (3.2) into Eq. (1.3) we obtain

$$-\omega u' + uu' - k\sigma u'' = 0, \quad (3.6)$$

where $D_t^\alpha u = -\omega ku'$, $D_x^\alpha u = ku'$, $D_x^{\alpha\alpha} u = k^2 u''$, by balancing u'' with uu' gives $s = 1$. Use $s = 1$ in Eq. (3.4), then the solution of Eq. (1.3) can be expressed as

$$u = a_0 + a_1 F, \quad (3.7)$$

substituting Eq. (3.7) into Eq. (3.6) and using Eq. (3.5), then we obtain a set of algebraic equations with respect to $F^i (i = 0, 1, 2, 3)$. Equating the coefficients of $F^i (i = 0, 1, 2, 3)$ to zero. The solution of the resulting system is given by

$$a_0 = \omega + \sigma Bk, \quad a_1 = 2\sigma Ck, \quad (3.8)$$

with ω and k are arbitrary constants. Inserting Eq. (3.8) into Eq. (3.7) and using the special solutions of Eq. (3.5), we obtain the following soliton like-solution and triangular periodic solutions of the space-time CFBE:

$$u_1 = \omega + 2\sigma k \tan \xi, \quad (3.9)$$

$$u_2 = \omega - 2\sigma k \cot \xi, \quad (3.10)$$

$$u_3 = \omega - 2\sigma k \tanh \xi, \quad (3.11)$$

$$u_4 = \omega - 2\sigma k \coth \xi, \quad (3.12)$$

$$u_5 = \omega - \sigma k (\tanh \xi \pm i \operatorname{sech} \xi), \quad (3.13)$$

$$u_6 = \omega - \sigma k (\coth \xi \pm \operatorname{csch} \xi), \quad (3.14)$$

$$u_7 = \omega + \sigma k (\tan \xi \pm \sec \xi), \quad (3.15)$$

$$u_8 = \omega - \sigma k (\cot \xi \pm \csc \xi), \quad (3.16)$$

$$u_9 = \omega - \frac{8\sigma k \tanh \xi}{1 + \tanh^2 \xi}, \quad (3.17)$$

$$u_{10} = \omega + \frac{8\sigma k \tan \xi}{1 - \tan^2 \xi}, \quad (3.18)$$

$$u_{11} = \omega - \frac{8\sigma k \cot \xi}{1 - \cot^2 \xi}, \quad (3.19)$$

$$u_{12} = \omega - \frac{\sigma k \tanh \xi}{1 + \operatorname{sech} \xi}, \quad (3.20)$$

$$u_{13} = \omega - \frac{\sigma k \coth \xi}{1 + i \operatorname{csch} \xi}, \quad (3.21)$$

$$u_{14} = \omega + \frac{\sigma k \tan \xi}{1 + \sec \xi}, \quad (3.22)$$

$$u_{15} = \omega - \frac{\sigma k \cot \xi}{1 + \csc \xi}, \quad (3.23)$$

$$u_{16} = \omega - 2\sigma k + \frac{4\sigma k \tan \xi}{1 + \tan \xi}, \quad (3.24)$$

$$u_{17} = \omega + 2\sigma k - \frac{4\sigma k \cot \xi}{1 + \cot \xi}, \quad (3.25)$$

$$u_{18} = \frac{(\sigma k + \omega) + (\sigma k - \omega)e^\xi}{1 - e^\xi}, \quad (3.26)$$

$$u_{19} = \frac{(\sigma k/2 + \omega) + (\sigma k/2 - \omega)e^{\xi/2}}{1 - e^{\xi/2}}, \quad (3.27)$$

$$u_{20} = \omega - \sigma k \tanh \frac{\xi}{2}, \quad (3.28)$$

$$u_{21} = \omega - \sigma k \coth \frac{\xi}{2}, \quad (3.29)$$

$$u_{22} = \omega - 2\sqrt{3}\sigma k \tanh \sqrt{3}\xi, \quad (3.30)$$

$$u_{23} = \omega - 2\sqrt{3}\sigma k \coth h\sqrt{3}\xi, \quad (3.31)$$

$$u_{24} = \omega + \sqrt{3}\sigma k \tan\left(\frac{\sqrt{3}}{2}\xi\right), \quad (3.32)$$

$$u_{25} = \omega - \sqrt{3}\sigma k \cot\left(\frac{\sqrt{3}}{2}\xi\right), \quad (3.33)$$

$$u_{26} = \omega - 8\sigma k \tanh(4\xi), \quad (3.34)$$

$$u_{27} = \omega - 8\sigma k \coth(4\xi), \quad (3.35)$$

$$u_{28} = \omega + \sigma k \tan\left(\frac{\xi}{2}\right), \quad (3.36)$$

$$u_{29} = \omega - \sigma k \cot\left(\frac{\xi}{2}\right), \quad (3.37)$$

Remarks:

1- Making the transformation $k \rightarrow 2k$ then Eq. (3.28) and Eq. (3.29) transformed to Eq. (3.11) and Eq. (3.12) respectively, and if $k \rightarrow 2ki$ where $i = \sqrt{-1}$, they transformed to Eq. (3.9) and Eq. (3.10) respectively.

2- Making the transformation $k \rightarrow \frac{k}{\sqrt{3}}$ then Eq. (3.30) and Eq. (3.31) transformed to Eq. (3.11) and Eq. (3.12) respectively.

3- Making the transformation $k \rightarrow \frac{2k}{\sqrt{3}}$ then Eq. (3.32) and Eq. (3.33) transformed to Eq. (3.9) and Eq. (3.10) respectively.

4- Making the transformation $k \rightarrow 2k$ then Eq. (3.27) transformed to Eq. (3.26).

5- Making the transformation $k \rightarrow \frac{k}{4}$ then Eq. (3.34) and Eq. (3.35) transformed to Eq. (3.11) and Eq. (3.12) respectively.

6- Making the transformation $k \rightarrow 2k$ then Eq. (3.36) and Eq. (3.37) transformed to Eq. (3.9) and Eq. (3.10) respectively.

Also, we can get two new exact solutions

$$u_{30} = \omega - 2\sigma k (\tanh\xi + \coth\xi), \quad (3.38)$$

$$u_{31} = \omega + 2\sigma k (\tan\xi - \cot\xi), \quad (3.39)$$

with $\xi = \frac{k}{\alpha}(x^\alpha - \omega t^\alpha)$.

In the following section we use the obtained solutions Eqs. (3.9) - (3.39) to generate new abundant exact solutions via BT.

4. BT and abundant exact solutions

From section 2, we show that the space-time CFBE possesses the Painlevé property and it has a BT in the form

$$u = -\frac{2\sigma}{\phi} D_x^\alpha \phi + w, \quad (4.1)$$

where $\phi = \phi\left(\frac{x^\alpha}{\alpha}, \frac{t^\alpha}{\alpha}\right)$ is the singular manifold variable, w is a function of $\frac{x^\alpha}{\alpha}$ and $\frac{t^\alpha}{\alpha}$. Also, the function w solves the space-time CFBE given by Eq. (1.3) and the function ϕ satisfies the FDE

$$D_t^\alpha \phi + w D_x^\alpha \phi = \sigma D_x^{\alpha\alpha} \phi. \quad (4.2)$$

Now, if we take $w = \phi$ then the function ϕ satisfies also the space-time CFBE

$$D_t^\alpha \phi + \phi D_x^\alpha \phi = \sigma D_x^{\alpha\alpha} \phi, \quad (4.3)$$

thus the BT for the space-time CFBE takes the following recurrence form

$$u_{n+1} = -\frac{2\sigma}{u_n} \frac{\partial^\alpha u_n}{\partial x^\alpha} + u_n. \quad (4.4)$$

We turn to the application of the BT for the FDEs. Their power lies in that they may be used to generate additional solutions of the FDEs. Here u_{n+1} quantities refer to new solution and u_n quantities refer to old solution. This means that, on the basis of a known solution to the space-time CFBE, we are able to find new solution of space-time CFBE. To construct the new solution of the space-time CFBE one can start with the solution u_1 obtained in Eq. (3.9) and using BT given in Eq. (4.4) we get the following set of new solutions:

$$u_{32} = \frac{-4\sigma^2 k^2 + \omega^2 + 4\sigma k \omega \tan \xi}{\omega + 2\sigma k \tan \xi}, \quad (4.5)$$

Inserting u_{32} into the BT given in Eq. (4.4) we have

$$u_{33} = \frac{12\sigma^2 k^2 \omega - \omega^3 + 2\sigma k (4\sigma^2 k^2 - 3\omega^2) \tan \xi}{4\sigma^2 k^2 - \omega^2 - 4\sigma k \omega \tan \xi}, \quad (4.6)$$

furthermore, using u_{33} and BT given in Eq. (4.4) we get

$$u_{34} = \frac{-16\sigma^4 k^4 + 24\sigma^2 k^2 \omega^2 - \omega^4 + (32\sigma^3 k^3 \omega - 8\omega^3 \sigma k) \tan \xi}{12\sigma^2 k^2 \omega - \omega^3 + 2\sigma k (4\sigma^2 k^2 - 3\omega^2) \tan \xi}, \quad (4.7)$$

and so on, we can get a sequences of exact solutions generated by the known tan-function solution Eq. (3.9) of the space-time CFBE. Starting from u_2 obtained in Eq. (3.10) and using BT given in Eq. (4.4) we get

$$u_{35} = \frac{-4\sigma^2 k^2 + \omega^2 - 4\sigma k \omega \cot \xi}{\omega - 2\sigma k \cot \xi}, \quad (4.8)$$

Inserting u_{35} into the BT given in Eq. (4.4) we have

$$u_{36} = \frac{-12\sigma^2 k^2 \omega + \omega^3 + 2\sigma k(4\sigma^2 k^2 - 3\omega^2) \cot \xi}{-4\sigma^2 k^2 + \omega^2 - 4\sigma k \omega \cot \xi}, \quad (4.9)$$

furthermore, using u_{36} and BT given in Eq. (4.4) we get

$$u_{37} = \frac{16\sigma^4 k^4 - 24\sigma^2 k^2 \omega^2 + \omega^4 + (32\sigma^3 k^3 \omega - 8\omega^3 \sigma k) \cot \xi}{-12\sigma^2 k^2 \omega + \omega^3 + 2\sigma k(4\sigma^2 k^2 - 3\omega^2) \cot \xi}, \quad (4.10)$$

and so on, we can get a sequences of exact solutions generated by the cot-function solution Eq. (3.10) of the space time fractional CFBE. Starting from u_3 obtained in Eq. (3.11) and using BT given in Eq. (4.4) we get

$$u_{38} = \frac{-4\sigma^2 k^2 - \omega^2 + 4\sigma k \omega \tanh \xi}{-\omega + 2\sigma k \tanh \xi}, \quad (4.11)$$

Inserting u_{38} into the BT given in Eq. (4.4) we have

$$u_{39} = \frac{12\sigma^2 k^2 \omega + \omega^3 - 2\sigma k(4\sigma^2 k^2 + 3\omega^2) \tanh \xi}{4\sigma^2 k^2 + \omega^2 - 4\sigma k \omega \tanh \xi}, \quad (4.12)$$

furthermore, using u_{39} and BT given in Eq. (4.4) we get

$$u_{40} = \frac{16\sigma^4 k^4 + 24\sigma^2 k^2 \omega^2 + \omega^4 - (32\sigma^3 k^3 \omega + 8\omega^3 \sigma k) \tanh \xi}{12\sigma^2 k^2 \omega + \omega^3 - 2\sigma k(4\sigma^2 k^2 + 3\omega^2) \tanh \xi}, \quad (4.13)$$

and so on, we can get a sequences of exact solutions generated by the tanh-function solution Eq. (3.11) of the space-time CFBE. Starting from u_4 obtained in Eq. (3.12) and using BT given in Eq. (4.4) we get

$$u_{41} = \frac{-4\sigma^2 k^2 - \omega^2 + 4\sigma k \omega \coth \xi}{-\omega + 2\sigma k \coth \xi}, \quad (4.14)$$

Inserting u_{41} into the BT given in Eq. (4.4) we have

$$u_{42} = \frac{12\sigma^2 k^2 \omega + \omega^3 - 2\sigma k(4\sigma^2 k^2 + 3\omega^2) \coth \xi}{4\sigma^2 k^2 + \omega^2 - 4\sigma k \omega \coth \xi}, \quad (4.15)$$

furthermore, using u_{42} and BT given in Eq. (4.4) we get

$$u_{43} = \frac{16\sigma^4 k^4 + 24\sigma^2 k^2 \omega^2 + \omega^4 - (32\sigma^3 k^3 \omega + 8\omega^3 \sigma k) \coth \xi}{12\sigma^2 k^2 \omega + \omega^3 - 2\sigma k(4\sigma^2 k^2 + 3\omega^2) \coth \xi}, \quad (4.16)$$

and so on, we can get a sequences of exact solutions generated by the coth-function solution Eq. (3.12) of the space-time CFBE. Starting from u_5 obtained in Eq. (3.13) and using BT given in Eq. (4.4) we get

$$u_{44} = \frac{\sigma^2 k^2 + \omega^2 - 2\sigma k \omega (\tanh \xi \pm i \operatorname{sech} \xi)}{\omega - \sigma k (\tanh \xi \pm i \operatorname{sech} \xi)}, \quad (4.17)$$

and so on. Starting from u_6 obtained in Eq. (3.14) and using BT given in Eq. (4.4) we get

$$u_{45} = \frac{\sigma^2 k^2 + \omega^2 - 2\sigma k \omega (\coth \xi \pm \operatorname{csch} \xi)}{\omega - \sigma k (\coth \xi \pm \operatorname{csch} \xi)}, \quad (4.18)$$

and so on. Starting from u_7 obtained in Eq. (3.15) and using BT given in Eq. (4.4) we get

$$u_{46} = \frac{\sigma^2 k^2 - \omega^2 - 2\sigma k \omega (\tan \xi \pm \sec \xi)}{-\omega - \sigma k (\tan \xi \pm \sec \xi)}, \quad (4.19)$$

and so on. Starting from u_8 obtained in Eq. (3.16) and using BT given in Eq. (4.4) we get

$$u_{47} = \frac{-\sigma^2 k^2 + \omega^2 - 2\sigma k \omega (\cot \xi \pm \csc \xi)}{\omega - \sigma k (\cot \xi \pm \csc \xi)}, \quad (4.20)$$

and so on. Starting from u_9 obtained in Eq. (3.17) and using BT given in Eq. (4.4) we get

$$u_{48} = \frac{(16\sigma^2 k^2 + \omega^2) \tanh^2 \xi - 16\sigma k \omega \tanh \xi + 16\sigma^2 k^2 + \omega^2}{\omega \tanh^2 \xi - 8\sigma k \tanh \xi + \omega}, \quad (4.21)$$

Inserting u_{48} into the BT given in Eq. (4.4) we have

$$u_{49} = \frac{(48\sigma^2 k^2 \omega + \omega^3) \tanh^2 \xi - (24\sigma k \omega^2 + 128\sigma^3 k^3) \tanh \xi + 48\sigma^2 k^2 \omega + \omega^3}{(16\sigma^2 k^2 + \omega^2) \tanh^2 \xi - 16\sigma k \omega \tanh \xi + 16\sigma^2 k^2 + \omega^2}, \quad (4.22)$$

and so on. Starting from u_{10} obtained in Eq. (3.18) and using BT given in Eq. (4.4) we get

$$u_{50} = \frac{(16\sigma^2 k^2 - \omega^2) \tan^2 \xi + 16\sigma k \omega \tan \xi - 16\sigma^2 k^2 + \omega^2}{\omega - \omega \tan^2 \xi + 8\sigma k \tan \xi}, \quad (4.23)$$

Inserting u_{50} into the BT given in Eq. (4.4) we have

$$u_{51} = \frac{(48\sigma^2 k^2 \omega - \omega^3) \tan^2 \xi + (24\sigma k \omega^2 - 128\sigma^3 k^3) \tan \xi - 48\sigma^2 k^2 \omega + \omega^3}{(16\sigma^2 k^2 - \omega^2) \tan^2 \xi + 16\sigma k \omega \tan \xi - 16\sigma^2 k^2 + \omega^2}, \quad (4.24)$$

and so on. Starting from u_{11} obtained in Eq. (3.19) and using BT given in Eq. (4.4) we get

$$u_{52} = \frac{(16\sigma^2 k^2 - \omega^2) \cot^2 \xi - 16\sigma k \omega \cot \xi - 16\sigma^2 k^2 + \omega^2}{\omega - \omega \cot^2 \xi - 8\sigma k \cot \xi}, \quad (4.25)$$

Inserting u_{52} into the BT given in Eq. (4.4) we have

$$u_{53} = \frac{(\omega^3 - 48\sigma^2 k^2 \omega) \cot^2 \xi + (24\sigma k \omega^2 - 128\sigma^3 k^3) \cot \xi + 48\sigma^2 k^2 \omega - \omega^3}{(\omega^2 - 16\sigma^2 k^2) \cot^2 \xi + 16\sigma k \omega \cot \xi + 16\sigma^2 k^2 - \omega^2}, \quad (4.26)$$

and so on. Starting from u_{12} obtained in Eq. (3.20) and using BT given in Eq. (4.4) we get

$$u_{54} = \frac{\sigma^2 k^2 + \omega^2 + (\sigma^2 k^2 + \omega^2) \operatorname{sech} \xi - 2\sigma k \omega \tanh \xi}{\omega + \omega \operatorname{sech} \xi - \sigma k \tanh \xi}, \quad (4.27)$$

and so on. Starting from u_{13} obtained in Eq. (3.21) and using BT given in Eq. (4.4) we get

$$u_{55} = \frac{\sigma^2 k^2 + \omega^2 + (\sigma^2 k^2 + \omega^2) i \operatorname{csch} \xi - 2\sigma k \omega \coth \xi}{\omega + \omega i \operatorname{csch} \xi - \sigma k \coth \xi}, \quad (4.28)$$

and so on. Starting from u_{14} obtained in Eq. (3.22) and using BT given in Eq. (4.4) we get

$$u_{56} = -\frac{(\sigma^2 k^2 - \omega^2) \cos \xi + \sigma^2 k^2 - \omega^2 - 2\sigma k \omega \sin \xi}{\omega + \omega \cos \xi + \sigma k \sin \xi}, \quad (4.29)$$

and so on. Starting from u_{15} obtained in Eq. (3.23) and using BT given in Eq. (4.4) we get

$$u_{57} = \frac{(\sigma^2 k^2 - \omega^2) \sin \xi + \sigma^2 k^2 - \omega^2 + 2\sigma k \omega \cos \xi}{-\omega - \omega \sin \xi + \sigma k \cos \xi}, \quad (4.30)$$

and so on. Starting from u_{16} obtained in Eq. (3.24) and using BT given in Eq. (4.4) we get

$$u_{58} = \frac{\omega^2 - 4\sigma^2 k^2 - 4\sigma k \omega + (\omega^2 - 4\sigma^2 k^2 + 4\sigma k \omega) \tan \xi}{(\omega + 2\sigma k) \tan \xi + \omega - 2\sigma k}, \quad (4.31)$$

Inserting u_{58} into the BT given in Eq. (4.4) we have

$$u_{59} = \frac{\omega^3 + 8\sigma^3 k^3 - 6\sigma k \omega^2 - 12\sigma^2 k^2 \omega + (\omega^3 - 8\sigma^3 k^3 + 6\sigma k \omega^2 - 12\sigma^2 k^2 \omega) \tan \xi}{\omega^2 - 4\sigma^2 k^2 - 4\sigma k \omega + (\omega^2 - 4\sigma^2 k^2 + 4\sigma k \omega) \tan \xi}, \quad (4.32)$$

and so on. Starting from u_{17} obtained in Eq. (3.25) and using BT given in Eq. (4.4) we get

$$u_{60} = \frac{\omega^2 - 4\sigma^2 k^2 + 4\sigma k \omega + (\omega^2 - 4\sigma^2 k^2 - 4\sigma k \omega) \cot \xi}{(\omega - 2\sigma k) \cot \xi + \omega + 2\sigma k}, \quad (4.33)$$

Inserting u_{60} into the BT given in Eq. (4.4) we have

$$u_{61} = \frac{\omega^3 - 8\sigma^3 k^3 + 6\sigma k \omega^2 - 12\sigma^2 k^2 \omega + (\omega^3 + 8\sigma^3 k^3 - 6\sigma k \omega^2 - 12\sigma^2 k^2 \omega) \cot \xi}{\omega^2 - 4\sigma^2 k^2 + 4\sigma k \omega + (\omega^2 - 4\sigma^2 k^2 - 4\sigma k \omega) \cot \xi}, \quad (4.34)$$

and so on. Starting from u_{18} obtained in Eq. (3.26) and using BT given in Eq. (4.4) we get

$$u_{62} = \frac{(\sigma k + \omega)^2 - (\sigma k - \omega)^2 e^\xi}{\sigma k + \omega + (\sigma k - \omega) e^\xi}, \quad (4.35)$$

Inserting u_{62} into the BT given in Eq. (4.4) we have

$$u_{63} = \frac{(\sigma k + \omega)^3 + (\sigma k - \omega)^3 e^\xi}{(\sigma k + \omega)^2 - (\sigma k - \omega)^2 e^\xi}, \quad (4.36)$$

Furthermore, using u_{63} and BT given in Eq. (4.4) we get

$$u_{64} = \frac{(\sigma k + \omega)^4 - (\sigma k - \omega)^4 e^\xi}{(\sigma k + \omega)^3 + (\sigma k - \omega)^3 e^\xi}, \quad (4.37)$$

and so on, we can get a new sequences of exact solution of the space-time CFBE.

Starting from u_{30} obtained in Eq. (3.38) and using BT given in Eq. (4.4) we get

$$u_{65} = \frac{-(\omega^2 + 16\sigma^2 k^2) \cosh \xi \sinh \xi + 8\sigma k \omega \cosh^2 \xi - 4\sigma k \omega}{-\omega \cosh \xi \sinh \xi + 4\sigma k \cosh^2 \xi - 2\sigma k}, \quad (4.38)$$

and so on, we can get a sequences of solutions generated by the addition of two functions tanh and coth-function solution of the space-time CFBE. Starting from u_{31} obtained in Eq. (3.39) and using BT given in Eq. (4.4) we get

$$u_{66} = -\frac{(\omega^2 - 16\sigma^2 k^2) \cos \xi \sin \xi - 8\sigma k \omega \cos^2 \xi + 4\sigma k \omega}{-\omega \cos \xi \sin \xi + 4\sigma k \cos^2 \xi - 2\sigma k}, \quad (4.39)$$

and so on, we can get a sequences of solutions generated by the addition of two functions tan and cot-function solution of the space-time CFBE.

By means of the variational iteration method [35] and the Adomian decomposition method [36] the solution of the space-time CFBE in closed form is

$$u_{67} = \frac{(\omega + k) + (\omega - k) \exp(\xi/\sigma)}{1 + \exp(\xi/\sigma)}, \quad (4.40)$$

From u_{67} obtained in Eq. (4.40) we can get a new sequences of exact solution for the space-time CFBE by using BT given in Eq. (4.4) in the form

$$u_{68} = \frac{(\omega+k)^2+(\omega-k)^2 \exp(\xi/\sigma)}{(\omega+k)+(\omega-k) \exp(\xi/\sigma)}, \quad (4.41)$$

$$u_{69} = \frac{(\omega+k)^3+(\omega-k)^3 \exp(\xi/\sigma)}{(\omega+k)^2+(\omega-k)^2 \exp(\xi/\sigma)}, \quad (4.42)$$

$$u_{70} = \frac{(\omega+k)^4+(\omega-k)^4 \exp(\xi/\sigma)}{(\omega+k)^3+(\omega-k)^3 \exp(\xi/\sigma)}, \quad (4.43)$$

$$u_{71} = \frac{(\omega+k)^5+(\omega-k)^5 \exp(\xi/\sigma)}{(\omega+k)^4+(\omega-k)^4 \exp(\xi/\sigma)}, \quad (4.44)$$

and so on, we can get a new sequences of exact solution of the space-time CFBE.

5. Conclusion

In this work, we discuss the Painlevé property for non-linear conformal fractional differential equations for the first time . We apply the desired method to the space time conformal fractional Burger's equation. Also, we derive the Bäcklund transform. The general solutions of the space-time conformal CFDEs is described based on the tanh-method, accordingly the method is successively implement to space-time CFBE. Moreover, the space-time CFBE is found to possess the Painlevé property and then Bäcklund transform. Also, we introduced a new recurrence formulae based on Bäcklund transform, that is enable us to derive analytical solution from known solution or old solution to give new solution. New numerous exact solutions are generated based on the Bäcklund transform.

Acknowledgments

The authors gratefully acknowledge the approval and the support of this research study by the grant no. SCI-2019-1-10-F-8306 from the Deanship of Scientific Research at Northern Border University, Arar, Saudi Arabia.

References

- [1] Kilbas, A. A., Srivastava, H. M., and Trujillo J. J., Theory and applications of fractional differential equations, Amsterdam: Elsevier B.V., 2006.
- [2] Hilfer, R., Applications of Fractional Calculus in Physics, World Science, Singapore, 2000.

- [3] Miller, K. S., and Ross, B., An Introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley and Sons, New York, NY, USA, 1993.
- [4] Abdel-Salam, E. A.-B., Yousif, E. A., and El-Aasser, M. A., Analytical Solution of The space-time fractional nonlinear Schrödinger equation. Reports on Mathematical Physics 2016; 19:77.
- [5] Abdel-Salam, E. A-B, Al-Muhiameed Z. I. A., Analytic solutions of the space-time fractional combined KdV-mKdV equation. Mathematical Problems in Engineering 2015; Article ID 871635.
- [6] Abdel-Salam, E. A-B, Hassan G.F., Solutions to class of linear and nonlinear fractional differential equations. Communications in Theoretical Physics 2016; 65: 127.
- [7] Abdel-Salam, E. A-B, Yousif E. A., Solution of nonlinear space-time fractional differential equations using the fractional Riccati expansion method, Mathematical Problems in Engineering 2013; Article ID 846283.
- [8] Nouh, M.I., Abdel-Salam, E .A-B., Approximate Solution to the Fractional Lane–Emden Type Equations. Iranian Journal of Science Technology, Transactions Science 2018;42: 2199-2206.
- [9] Ahmed, S. E., Mansour, M. A., Abdel-Salam, E. A-B, Mohamed, E.F, Studying the fractional derivative for natural convection in slanted cavity containing porous media. SN Applied Sciences 2019;1: 1117.
- [10] Abdel-Salam, E.A-B, Nouh, M.I., Conformable fractional polytropic gas spheres. New Astronomy 2020; 76: 101322.
- [11] Nouh, M.I., Abdel-Salam, E. A-B., Analytical solution to the fractional polytropic gas spheres. European Physics Journal Plus 2018;133: 149.
- [12] Abdel-Salam, E. A-B, Nouh, M.I., Approximate solution to the fractional second-type lane- emden equation. Astrophysics 2016; 59: 398.
- [13] Yousif, E. A., Abdel-Salam, E. A-B, El-Aasser, M. A., On the solution of the space-time fractional cubic nonlinear Schrödinger equation. Results In Physics 2018; 8: 702.
- [14] Mainardi F., Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models. Imperial College Press, London, 2010.
- [15] Abdel-Salam, E. A-B, Mourad, M. F., Fractional quasi AKNS-technique for nonlinear space–time fractional evolution equations. Mathematical Methods in the Applied Sciences 2018; 42: 5953-5968.

- [16] Nouh, M.I., Azzam, Y.A. & Abdel-Salam, E. A-B. Modeling fractional polytropic gas spheres using artificial neural network. *Neural Comput & Applic* (2020).
<https://doi.org/10.1007/s00521-020-05277-9>
- [17] Abo-Dahab, S. M., Kilany A. A., Abdel-Salam E A-B. and Hatem A, Fractional derivative order analysis and temperature-dependent properties on p- and SV-waves reflection under initial stress and three-phase-lag model, *Results In Physics*, 18 (2020) 103270.
<https://doi.org/10.1016/j.rinp.2020.103270>.
- [18] J. Sabatier, O. P. Agrawal, and J. A. Tenreiro Machado, *Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering*, Springer, 2007.
- [19] Hong Guang Sun, Yong Zhang, Dumitru Baleanu, Wen Chen, Yang Quan Chen, A new collection of real world applications of fractional calculus in science and engineering, *Communications in Nonlinear Science and Numerical Simulation* **64**, 213-231 (2018).
- [20] D. Valério, J. Machado, V. Kiryakova, Some pioneers of the applications of fractional calculus. *Fract. Calc. Appl. Anal.* **17**(2), 552–578 (2014).
- [21] R. Khalil, M. A. Horani, A. Yousef, M. Sababheh. A new definition of fractional derivative, *J. Comput. Appl. Math.* **264**, 65–70 (2014).
- [22] T. Abdeljawad, on conformable fractional calculus, *Journal of Computational and Applied Mathematics* **279**, 57-66 (2015).
- [23] D. Zhao, M. Luo, General conformable fractional derivative and its physical interpretation. *Calcolo* **54**, 903–917 (2017).
- [24] Mark J. Ablowitz, David J. Kaup, Alan C. Newell, and Harvey Segur, Nonlinear-Evolution Equations of Physical Significance, *Phys. Rev. Lett.* 31, 125 – 127 (1973).
- [25] V.O. Vakhnenko, E.J. Parkes, the singular solutions of a nonlinear evolution equation taking continuous part of the spectral data into account in inverse scattering method, *Chaos, Solitons & Fractals* 45, 846-852 (2012).
- [26] Chen, H., Zhang, H. Extended Jacobin elliptic function method and its applications. *Journal of Applied Mathematics and Computing* **10**, 119–130 (2002).
- [27] C. Rogers and W. F. Shadwick, *Bäcklund Transformations and Their Applications*, Academic Press (1982).

- [28] Liu Xi-Zhong, Yu Jun, Ren Bo, and Yang Jian-Rong, Bäcklund transformations for the Burgers equation via localization of residual symmetries, *Chin. Phys. B* Vol. 23, No. 11 (2014) 110203.
- [29] Mei-na Sun, Shu-fang Deng, and Deng-yuan Chen, The Bäcklund transformation and novel solutions for the Toda lattice, *Chaos, Solitons and Fractals* 23 (2005) 1169–1175.
- [30] Matveev, V.B. and Salle, M.A, *Darboux Transformations and Solitons*, Springer Verlag (1991).
- [31] Malfliet, W, Solitary wave solutions of nonlinear wave equations, *Amer. J. Phys.* **60**, 650-654, (1992).
- [32] Malfliet, W. and Hereman, W., The Tanh Method: I. Exact Solutions of Nonlinear Evolution and Wave Equations, *Phys. Scr.* **54**, 563 – 568 (1996).
- [33] Liu, S.K. Fu, Z.T. Liu, S.D. and Zhao, Q, Jacobi elliptic function expansion method and periodic wave solutions of nonlinear wave equations, *Phys. Lett. A.* **289**, 69-74 (2001).
- [34] Yan, Z, Jacobi elliptic function solutions of nonlinear wave equations via the new sinh-Gordon equation expansion method, *J. Phys. A: Math. Gen.* **36**, 1961-1962 (2003).
- [35] Wies, J. Tabor, M. and Garnevale, G, The Painlevé property for partial differential equations, *J. Math. Phys.* **24**, 522-526 (1983).
- [36] Peter A. Clarkson, the Painlevé conjecture, the Painlevé property for partial differential equations and complete integrability, *Physica* 18D (1986) 209-210.
- [37] M. Tabor and J.D. Gibbon, aspects of the Painlevé property for partial differential equations, *Physica* 18D (1986) 180-189.
- [38] Xing Lu, Tao Geng, Cheng Zhang, Hong-Wu Zhu, Xiang-Hua Meng, and Bo Tian, Multi-soliton solutions and their interactions for the (2+1)-dimensional Sawada-Kotera model with truncated Painlevé expansion, Hirota Bilinear method and symbolic computation, *International Journal of Modern Physics B* Vol. 23, No. 25 (2009) 5003–5015.
- [39] Wang, M.L. Zhou, Y.B. and Li, Z.B., Application of a homogeneous balance method to exact solutions of nonlinear equations in mathematical physics, *Phys. Lett. A* **216** (1996) 67-75.
- [40] Yang, L. Zhu, Z. and Wang, Y., Exact solutions of nonlinear equations, *Phys. Lett. A* **260** (1999) 55-59.
- [41] Konno, K. and Wadati, M., Simple Derivation of Bäcklund Transformation from Riccati Form of Inverse Method, *Prog. Theor. Phys.* **53** (1975) 1652-1656.

- [42] Nucci, M.C., Pseudopotentials, Lax equations and Bäcklund transformations for nonlinear evolution equations, *J. Phys. A: Math. Gen.* **21** (1988) 73-79.
- [43] Orlandi P., The Burgers equation. Fluid Flow Phenomena. Fluid Mechanics and Its Applications, vol 55. 40-50 (2000).
- [44] Bonkile, M.P., Awasthi, A., Lakshmi, C. *et al.* A systematic literature review of Burgers' equation with recent advances. *Pramana - J Phys* **90**, 69 (2018).
- [45] Ali Kurt, Yücel Çenesiz, and Orkun Tasbozan, On the Solution of Burgers' Equation with the New Fractional Derivative, *Open Phys.* 2015; 13:355–360.
- [46] Ali Kurt, Yücel Cenesiz, and Orkun Tasbozan, Exact Solution for the Conformable Burgers' Equation by the Hopf-Cole Transform, *Cankaya University Journal of Science and Engineering*, Volume 13, No. 2 (2016) 018–023.
- [47] Mehmet Senol, Orkun Tasbozan, and Ali Kurt, Numerical Solutions of Fractional Burgers' Type Equations with Conformable Derivative, *Chinese Journal of Physics*, Volume 58 (2019) 75-84.
- [48] Ahmad El-Ajou, Zeyad Al-Zhour, Moa'ath Oqielat, Shaher Momani, and Tasawar Hayat, Series solutions of nonlinear conformable fractional KdV-Burgers' equation with some applications, *Eur. Phys. J. Plus* (2019) 134: 402.
- [49] Hassan Eltayeb, Imed Bachar, and Adem Kılıçman, On Conformable Double Laplace Transform and One Dimensional Fractional Coupled Burgers' Equation, *Symmetry* 2019, 11, 417; doi:10.3390/sym11010417.
- [50] Gui-qiong Xu, Zhi-bin Li, Symbolic computation of the Painlevé test for nonlinear partial differential equations using Maple, *Computer Physics Communications* 161 (2004) 65–75.