

THE DIXMIER-DOUADY CLASS AND AN ABELIAN EXTENSION OF THE HOMEOMORPHISM GROUP

SHUHEI MARUYAMA

ABSTRACT. Let X be a connected topological space and $c \in H^2(X; \mathbb{Z})$ a non-zero cohomology class. A $\text{Homeo}(X, c)$ -bundle is a fiber bundle with fiber X whose structure group reduces to the group $\text{Homeo}(X, c)$ of c -preserving homeomorphisms of X . If $H^1(X; \mathbb{Z}) = 0$, then a characteristic class for $\text{Homeo}(X, c)$ -bundles called the Dixmier-Douady class is defined via the Serre spectral sequence. We show a relation between the universal Dixmier-Douady class for foliated $\text{Homeo}(X, c)$ -bundles and the gauge group extension of $\text{Homeo}(X, c)$. Moreover, under some assumptions, we construct a central S^1 -extension and a group two-cocycle on $\text{Homeo}(X, c)$ corresponding to the Dixmier-Douady class.

1. INTRODUCTION

1.1. Main results. Let X be a connected topological space whose homeomorphism group $\text{Homeo}(X)$ is a topological group with respect to the compact-open topology. Assume that there exists a non-zero cohomology class $c \in H^2(X; \mathbb{Z})$. Let $G = \text{Homeo}(X, c)$ be the group of c -preserving homeomorphisms of X , and G^δ the group G with the discrete topology. A fiber bundle $X \rightarrow E \rightarrow B$ is called a G -bundle if the structure group reduces to G . A G -bundle is said to be *foliated* if the structure group reduces to the discrete group G^δ .

Assume that $H^1(X; \mathbb{Z}) = 0$. For a G -bundle $X \rightarrow E \rightarrow B$, we define a characteristic class $D_c(E)$ in $H^3(B; \mathbb{Z})$ as the image of c under the transgression map (see Subsection 2.1). We call this class $D_c(E)$ the *Dixmier-Douady class with respect to c* , or simply the *Dixmier-Douady class* if c is understood. Let $D_c \in H^3(BG; \mathbb{Z})$ denote the universal Dixmier-Douady class. Under the map

$$(1.1) \quad \iota^*: H^*(BG; \mathbb{Z}) \rightarrow H^*(BG^\delta; \mathbb{Z})$$

induced from the canonical map $G^\delta \rightarrow G$ and the isomorphism $H^3(BG^\delta; \mathbb{Z}) \cong H_{\text{grp}}^3(G; \mathbb{Z})$, we obtain the third group cohomology class $\iota^* D_c$ in $H_{\text{grp}}^3(G; \mathbb{Z})$, which we call the *discrete Dixmier-Douady class*.

In this paper we give algebraic interpretations of the discrete Dixmier-Douady class. Throughout this paper, we regard the circle S^1 as the quotient group $\mathbb{R}/\mathbb{Z}$. Let $P \rightarrow X$ be a principal S^1 -bundle whose first Chern class is equal to $c \in H^2(X; \mathbb{Z})$. Then the group $\text{Aut}(P)$ of bundle automorphisms of P gives rise to a group extension

$$(1.2) \quad 0 \rightarrow \text{Gau}(P) \rightarrow \text{Aut}(P) \rightarrow G \rightarrow 1.$$

2020 *Mathematics Subject Classification.* 37E45; 37E10.

Key words and phrases. Dixmier-Douady class; Gauge group.

Here $\text{Gau}(P)$, which is called the *gauge group of P* , is the kernel of the natural projection. Since the group S^1 is abelian, the gauge group is also abelian. Hence group extension (1.2) is an abelian extension. In general, an abelian extension $1 \rightarrow A \rightarrow \Gamma \rightarrow G \rightarrow 1$ determines a second group cohomology class $e(\Gamma) \in H_{\text{grp}}^2(G; A)$. Here A is regarded as a right G -module by the conjugation Γ -action on A . In particular, abelian extension (1.2) defines a group cohomology class $e(\text{Aut}(P))$ in $H_{\text{grp}}^2(G; \text{Gau}(P))$.

Here we obtain two group cohomology classes, the class $e(A_G(P)) \in H_{\text{grp}}^2(G; \text{Gau}(P))$ corresponding to (1.2) and the discrete Dixmier-Douady class $\iota^* D_c \in H_{\text{grp}}^3(G; \mathbb{Z})$. Our main theorem (Theorem 1.1) describes the relation between these two classes.

To state Theorem 1.1, we need to define a map $\mathbf{d}: H_{\text{grp}}^2(G; \text{Gau}(P)) \rightarrow H_{\text{grp}}^3(G; \mathbb{Z})$. Note that, since S^1 is abelian, the gauge group $\text{Gau}(P)$ is isomorphic to the group $C(X, S^1)$ of S^1 -valued continuous maps on X . Let $C(X, \mathbb{R})$ denote the group of continuous maps from X to $\mathbb{R}$. By the assumption $H^1(X; \mathbb{Z}) = 0$ and the identification $H^1(X; \mathbb{Z}) = \text{Hom}(\pi_1(X), \mathbb{Z}) = \text{Hom}(\pi_1(X), \pi_1(S^1))$, all elements in $C(X, S^1)$ can be lifted to elements in $C(X, \mathbb{R})$. Hence the sequence

$$(1.3) \quad 0 \rightarrow \mathbb{Z} \rightarrow C(X, \mathbb{R}) \rightarrow C(X, S^1) \rightarrow 0$$

is exact. By (1.3) and the identification $\text{Gau}(P) \cong C(X, S^1)$, we have the connecting homomorphism

$$\mathbf{d}: H_{\text{grp}}^2(G; \text{Gau}(P)) \rightarrow H_{\text{grp}}^3(G; \mathbb{Z}).$$

We are now ready to state Theorem 1.1.

Theorem 1.1. Let $\mathbf{d}: H_{\text{grp}}^2(G; \text{Gau}(P)) \rightarrow H_{\text{grp}}^3(G; \mathbb{Z})$ denote the connecting homomorphism. Then the following holds:

$$\mathbf{d}(e(\text{Aut}(P))) = \iota^* D_c.$$

From here to the end of this subsection, we further assume that the cohomology class $c \in H^2(X; \mathbb{Z})$ is equal to zero in $H^2(X; \mathbb{R})$. By abuse of notation, we use the symbol $\mathbf{d}$ to denote the connecting homomorphism $\mathbf{d}: H_{\text{grp}}^2(G; S^1) \rightarrow H_{\text{grp}}^3(G; \mathbb{Z})$ induced from the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow S^1 \rightarrow 1$ of coefficients. Since the sequence

$$\dots \rightarrow H^1(X; S^1) \xrightarrow{\mathbf{d}} H^2(X; \mathbb{Z}) \rightarrow H^2(X; \mathbb{R}) \rightarrow \dots$$

is exact, there exists a cohomology class $\rho \in H^1(X; S^1)$ satisfying $\mathbf{d}(\rho) = c$. By using ρ , we can construct a central S^1 -extension

$$(1.4) \quad 1 \rightarrow S^1 \rightarrow \text{Aut}(P_\rho^\delta) \rightarrow G \rightarrow 1$$

(see Section 4). This central S^1 -extension determines a second cohomology class $e(\text{Aut}(P_\rho^\delta))$ in $H_{\text{grp}}^2(G; S^1)$. Then the following theorem holds, which is analogous to Theorem 1.1.

Theorem 1.2. Let $\mathbf{d}: H_{\text{grp}}^2(G; S^1) \rightarrow H_{\text{grp}}^3(G; \mathbb{Z})$ be the connecting homomorphism with respect to the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow S^1 \rightarrow 1$ of coefficients. Then the following holds:

$$\mathbf{d}(e(\text{Aut}(P_\rho^\delta))) = \iota^* D_c.$$

Moreover, we describe a group two-cocycle $\mathfrak{G}_{x,\alpha} \in C_{\text{grp}}^2(G; S^1)$ which represents the class $e(\text{Aut}(P_\rho^\delta))$ (see (4.4)).

1.2. Background. To give a context to our work, we briefly describe a related results. Our main theorem can be seen as a higher dimensional analogue of the relation between the Euler class of foliated S^1 -bundles and the universal covering of the group $\text{Homeo}_+(S^1)$ of orientation preserving homeomorphisms of the circle.

One way to define the Euler class of (orientable) S^1 -bundle $S^1 \rightarrow E \rightarrow B$ is to use the transgression map $d_2^{0,1}: H^1(S^1; \mathbb{Z}) \rightarrow H^2(B; \mathbb{Z})$, that is, the Euler class $e(E)$ is defined by $e(E) = -d_2^{0,1}(c)$ for the generator $c \in H^1(S^1; \mathbb{Z})$. The universal Euler class e is the class in $H^2(B \text{Homeo}_+(S^1); \mathbb{Z})$, where $\text{Homeo}_+(S^1)$ is the group of orientation preserving homeomorphisms. The universal Euler class of foliated S^1 -bundles (discrete Euler class) is defined by ι^*e , where $\iota^*: H^2(B \text{Homeo}_+(S^1); \mathbb{Z}) \rightarrow H^2(B \text{Homeo}_+(S^1)^\delta; \mathbb{Z}) \cong H_{\text{grp}}^2(\text{Homeo}_+(S^1); \mathbb{Z})$ is the canonical map. As an algebraic interpretation of the discrete Euler class, it is known that the central $\mathbb{Z}$ -extension

$$0 \rightarrow \mathbb{Z} \rightarrow \widetilde{\text{Homeo}}_+(S^1) \rightarrow \text{Homeo}_+(S^1) \rightarrow 1$$

induced from the universal covering of $\text{Homeo}_+(S^1)$ corresponds to the universal Euler class of foliated S^1 -bundles (see [Fri17] for example).

This correspondence between the Euler class of S^1 -bundles and the central extension is generalized in [GK11] as follows. Let X be a connected topological space and $c \in H^1(X; \mathbb{Z})$ a non-zero *first* cohomology class. Let $\text{Homeo}(X, c)$ be the group of c -preserving homeomorphisms of X and $X \rightarrow E \rightarrow B$ a $\text{Homeo}(X, c)$ -bundle. A characteristic class $e_c(E) \in H^2(B; \mathbb{Z})$ of the G -bundle is defined as the image of c under the transgression map. The universal characteristic class e_c is defined as an element of $H^2(B \text{Homeo}(X, c); \mathbb{Z})$ and the universal characteristic class of foliated $\text{Homeo}(X, c)$ -bundles is given by ι^*e_c , where $\iota^*: H^*(BG; \mathbb{Z}) \rightarrow H^*(BG^\delta; \mathbb{Z})$ is the canonical map. It was shown in [GK11, Section 5] that the class ι^*e_c corresponds to a central $\mathbb{Z}$ -extension of G given by the group of homeomorphisms of the covering space of X associated to $c \in H^1(X; \mathbb{Z})$ which commute with the deck transformations.

Theorems 1.1 and 1.2 can be seen as results analogous to the above ones. We note that in our setting, the class e_c cannot be defined since there are no non-zero class in $H^1(X; \mathbb{Z})$.

There is also an analogous result in symplectic geometry ([Mar20]). Let (X, ω) be a one-connected integral symplectic manifold, that is, the class $[\omega] \in H^2(X; \mathbb{R})$ is in the image of the map $H^2(X; \mathbb{Z}) \rightarrow H^2(X; \mathbb{R})$. Then, if we replace $\text{Homeo}(X, c)$ with the *symplectomorphism group* $\text{Symp}(X, \omega)$, we obtain the Dixmier-Douady class for symplectic fibrations. The Dixmier-Douady class for symplectic fibration (or Hamiltonian fibration) is studied in [SS20]. It is known that there exists a central S^1 -extension of $\text{Symp}(X, \omega)$ called the *prequantum extension*, and this central extension corresponds to the discrete Dixmier-Douady class of foliated symplectic fibrations in the same manner as in Theorem 1.2 ([Mar20]). Note that the existence of the prequantum extension is guaranteed without the assumption that the class $[\omega] \in H^2(X; \mathbb{R})$ is zero, which corresponds to the condition in Theorem 1.2.

2. PRELIMINARIES

2.1. The Dixmier-Douady class. The following description of the Dixmier-Douady class is based on [CCM98]. For the basics of the Dixmier-Douady class, see [Bry93] for example.

Let X be a connected topological space satisfying $H^1(X; \mathbb{Z}) = 0$. Take a non-zero cohomology class c in $H^2(X; \mathbb{Z})$. Let $X \rightarrow E \rightarrow B$ denote a G -bundle with a connected base space B , and $(E_r^{p,q}, d_r^{p,q})$ the Serre spectral sequence of the bundle $X \rightarrow E \rightarrow B$. By the assumption $H^1(X; \mathbb{Z}) = 0$, we have

$$E_3^{0,2} = E_2^{0,2} = H^0(B; \mathcal{H}^2(X; \mathbb{Z}))$$

and

$$E_3^{3,0} = E_2^{3,0} = H^3(B; \mathcal{H}^0(X; \mathbb{Z})) = H^3(B; \mathbb{Z}),$$

where $\mathcal{H}^*(X; \mathbb{Z})$ is the local system over B induced from the bundle $X \rightarrow E \rightarrow B$. It is known that the cohomology group $H^0(B; \mathcal{H}^2(X; \mathbb{Z}))$ is isomorphic to the invariant part $H^2(X; \mathbb{Z})^{\pi_1(B)}$ of the monodromy action of $\pi_1(B)$ on $H^2(X; \mathbb{Z})$ (see [DK01, Proposition 5.14] for example). Since the structure group of the bundle E is contained in $G = \text{Homeo}(X, c)$, the monodromy action preserves the class c , and hence the class c can be seen as an element of $E_3^{0,2} = H^0(B; \mathcal{H}^2(X; \mathbb{Z}))$. By using the transgression map

$$d_3^{0,2}: E_3^{0,2} \rightarrow E_3^{3,0} = H^3(B; \mathbb{Z}),$$

we obtain the cohomology class $d_3^{0,2}c$ in $H^3(B; \mathbb{Z})$.

Definition 2.1. For a G -bundle $X \rightarrow E \rightarrow B$, the cohomology class

$$D_c(E) = -d_3^{0,2}c \in H^3(B; \mathbb{Z})$$

is called the *Dixmier-Douady class with respect to c* (or simply called the *Dixmier-Douady class* if c is understood).

By the naturality of the Serre spectral sequence, the cohomology class $d_3^{0,2}c$ has also the naturality with respect to bundle maps. Hence the class $D_c(E)$ gives rise to a characteristic class of G -bundles. Let $D_c \in H^3(BG; \mathbb{Z})$ denote the universal Dixmier-Douady class of G -bundles. By the canonical map $\iota^*: H^3(BG; \mathbb{Z}) \rightarrow H^3(BG^\delta; \mathbb{Z})$, we obtain the universal Dixmier-Douady class ι^*D_c of foliated G -bundles.

In section 5, we give examples of G -bundles whose Dixmier-Douady class is non-zero.

2.2. Group cohomology and Hochschild-Serre spectral sequence. Let G be a group and A a right G -module. A *group p -cochain* on G is a function from p -fold product G^p to A . Let $C_{\text{grp}}^p(G; A)$ be the set of all group p -cochains, and the coboundary map $\delta: C_{\text{grp}}^p(G; A) \rightarrow C_{\text{grp}}^{p+1}(G; A)$ is defined by

$$\begin{aligned} \delta c(g_1, \dots, g_{p+1}) = & c(g_2, \dots, g_{p+1}) + \sum_{i=1}^p (-1)^i c(g_1, \dots, g_i g_{i+1}, \dots, g_{p+1}) \\ & + (-1)^{p+1} c(g_1, \dots, g_p) \cdot g_{p+1} \end{aligned}$$

for $p > 0$ and $\delta = 0$ for $p = 0$. The cohomology of the cochain complex $(C_{\text{grp}}^*(G; A), \delta)$ is called the *group cohomology of G with coefficients in A* and denoted by $H_{\text{grp}}^*(G; A)$.

The group cohomology $H_{\text{grp}}^*(G; A)$ is known to be isomorphic to the singular cohomology $H^*(BG^\delta; \mathcal{A})$ of the classifying space BG^δ , where G^δ denote the group G with discrete topology and $\mathcal{A}$ is the local system on BG^δ induced from the G -module A (see [Bro82]).

For an exact sequence $1 \rightarrow K \xrightarrow{i} \Gamma \xrightarrow{\pi} G \rightarrow 1$ of groups, there is a spectral sequence $(E_r^{p,q}, d_r^{p,q})$ with $E_2^{p,q} \cong H_{\text{grp}}^p(G; H_{\text{grp}}^q(K; A))$ which converges to $H_{\text{grp}}^*(\Gamma; A)$. This spectral sequence is called the Hochschild-Serre spectral sequence ([HS53]). Here, $H_{\text{grp}}^q(K; A)$ is seen as a right G -module as follows. For a q -cochain c on K , we define a Γ -action by

$$(c \cdot \gamma)(k_1, \dots, k_q) = c(\gamma k_1 \gamma^{-1}, \dots, \gamma k_q \gamma^{-1}) \cdot \gamma.$$

This action induces a G -action on $H_{\text{grp}}^q(K; A)$. Let $H_{\text{grp}}^q(K; A)^G$ denote the G -invariant part.

The transgression map $d_r^{0,r-1}: E_r^{0,r-1} \rightarrow E_r^{0,r}$ is described as follows:

Proposition 2.2 ([HS53, p.129]). Let $f \in C_{\text{grp}}^r(\Gamma; A)$ be a cochain. Assume that there exists a cocycle $c \in C_{\text{grp}}^{r+1}(G; A)$ such that the pullback π^*c is equal to δf . Then, the cochain f defines an element $[f]_r$ of $E_r^{0,r-1}$ and the cohomology class $d_r^{0,r-1}([f]_r)$ is equal to $[\pi^*c]_r \in E_r^{r,0}$. Moreover, if $E_r^{0,r-1} \cong E_2^{0,r-1} \cong H_{\text{grp}}^{r-1}(K; A)$ and $E_r^{r,0} \cong E_2^{r,0} \cong H_{\text{grp}}^r(G; A)$, then the cohomology classes of $H_{\text{grp}}^{r-1}(K; A)$ and $H_{\text{grp}}^r(G; A)$ corresponding to $[f]_r$ and $d_r^{0,r-1}([f]_r)$ are represented by $f|_{K^{r-1}}$ and c , respectively.

Proposition 2.2 for the low degree cases $d_2^{0,1}$ and $d_3^{0,2}$ is described as follows. By definition, the cohomology $E_2^{0,1} = H_{\text{grp}}^0(G; H_{\text{grp}}^1(K; A))$ is isomorphic to $H_{\text{grp}}^1(K; A)^G$.

Corollary 2.3. Let $f \in C_{\text{grp}}^1(\Gamma; A)$ be a cochain. Assume that there exists a cocycle $c \in C_{\text{grp}}^2(G; A)$ satisfying $\delta f = \pi^*c$. Then, the pullback $i^*f \in C_{\text{grp}}^1(K; A)$ is a cocycle representing an element of $H_{\text{grp}}^1(K; A)^G$ and

$$d_2^{0,1}([i^*f]) = [c] \in H_{\text{grp}}^2(G; A).$$

If $H_{\text{grp}}^1(K; A)$ is trivial, then so is $E_2^{p,1} = H_{\text{grp}}^p(G; H_{\text{grp}}^1(K; A))$ for any $p \geq 0$. Hence the transgression map $d_3^{0,2}$ induces a map $d_3^{0,2}: H_{\text{grp}}^2(K; A)^G = E_3^{0,2} \rightarrow E_3^{3,0} = H_{\text{grp}}^3(G; A)$. By Proposition 2.2, we have the following.

Corollary 2.4. Assume that $H_{\text{grp}}^1(K; A)$ is trivial. Let $f \in C_{\text{grp}}^2(\Gamma; A)$ be a cochain. Assume that there exists a cocycle $c \in C_{\text{grp}}^3(G; A)$ satisfying $\delta f = \pi^*c$. Then, the pullback $i^*f \in C_{\text{grp}}^2(K; A)$ is a cocycle representing an element of $H_{\text{grp}}^2(K; A)^G$ and

$$d_3^{0,2}([i^*f]) = [c] \in H_{\text{grp}}^3(G; A).$$

2.3. Abelian extension and second group cohomology. An exact sequence $1 \rightarrow A \xrightarrow{i} \Gamma \xrightarrow{\pi} G \rightarrow 1$ of groups is called an *abelian extension of G* if the group A is abelian. The group Γ acts on A by conjugation, and this action induces a right G -action on A . Thus we consider the abelian group A as a right G -module. It is known that the second group cohomology $H_{\text{grp}}^2(G; A)$ is isomorphic to the set of equivalence classes of abelian A -extensions of G (see [Bro82]).

For an abelian A -extension Γ , the corresponding cohomology class $e(\Gamma)$ is defined as follows. Take a section $s: G \rightarrow \Gamma$ of the projection $p: \Gamma \rightarrow G$. For any $g, h \in G$, the value $s(g)s(h)s(gh)^{-1}$ is in $i(A) \cong A$. We define a group two-cochain $c \in C_{\text{grp}}^2(G; A)$ by

$$(2.1) \quad c(g, h) = s(g)s(h)s(gh)^{-1}.$$

This cochain c is a cocycle, and its cohomology class $[c]$ does not depend on the choice of sections. This cocycle represents the class $e(\Gamma)$.

By the definition of the derivations of the Hochschild-Serre spectral sequence, the cohomology class $e(\Gamma)$ can be described as follows (see [NSW08, (2.4.4) Theorem] for example).

Lemma 2.5. Let $1 \rightarrow A \rightarrow \Gamma \xrightarrow{\pi} G \rightarrow 1$ be an abelian A -extension of G and $(E_r^{p,q}, d_r^{p,q})$ the Hochschild-Serre spectral sequence of it. Then the cohomology class $e(\Gamma)$ is equal to the negative of $d_2^{0,1}(\text{id}_A)$, where $d_2^{0,1}: H_{\text{grp}}^1(A; A)^G = E_2^{0,1} \rightarrow E_2^{2,0} = H_{\text{grp}}^2(G; A)$ is the derivation of the spectral sequence and $H_{\text{grp}}^1(A; A)^G$ is the set of G -equivariant homomorphisms.

2.4. Cohomology of the gauge group. By the fibration

$$0 \rightarrow \mathbb{Z} \rightarrow C(X, \mathbb{R}) \xrightarrow{\pi} C(X, S^1) (\cong \text{Gau}(P)) \rightarrow 0$$

and the contractibility of $C(X, \mathbb{R})$, the gauge group $\text{Gau}(P)$ is turned out to be an Eilenberg-MacLane space $K(\mathbb{Z}, 1)$. Hence the classifying space $B\text{Gau}(\mathbb{Z})$ is $K(\mathbb{Z}, 2)$, and therefore we have

$$H^1(B\text{Gau}(P); \mathbb{Z}) = 0 \quad \text{and} \quad H^2(B\text{Gau}(P); \mathbb{Z}) = \mathbb{Z}.$$

Next we show $H_{\text{grp}}^1(\text{Gau}(P); \mathbb{Z}) = 0$. Note that the first cohomology $H_{\text{grp}}^1(\text{Gau}(P); \mathbb{Z})$ is isomorphic to the group $\text{Hom}(\text{Gau}(P), \mathbb{Z})$ of homomorphisms from $\text{Gau}(P)$ to $\mathbb{Z}$. Since $\text{Gau}(P)$ is a divisible group, we have $H_{\text{grp}}^1(\text{Gau}(P); \mathbb{Z}) = \text{Hom}(\text{Gau}(P), \mathbb{Z}) = 0$.

Let $i: \mathbb{R} \rightarrow C(X, \mathbb{R})$ and $j: S^1 \rightarrow \text{Gau}(P) = C(X, S^1)$ be the inclusions, and $\text{ev}_x^{\mathbb{R}}: C(X, \mathbb{R}) \rightarrow \mathbb{R}$ and $\text{ev}_x: C(X, S^1) \rightarrow S^1$ denote the evaluation maps at $x \in X$. Then we have the following commutative diagram of fibrations

$$\begin{array}{ccccc} B\mathbb{Z} & \longrightarrow & B\mathbb{R} & \longrightarrow & BS^1 \\ \parallel & & \downarrow i & & \downarrow j \\ B\mathbb{Z} & \longrightarrow & BC(X, \mathbb{R}) & \longrightarrow & B\text{Gau}(P) \\ \parallel & & \downarrow \text{ev}_x^{\mathbb{R}} & & \downarrow \text{ev}_x \\ B\mathbb{Z} & \longrightarrow & B\mathbb{R} & \longrightarrow & BS^1, \end{array}$$

where we use, by abuse of notation, the symbols $i, j, \text{ev}_x^{\mathbb{R}}$, and ev_x to denote the induced maps between the classifying spaces. Since the composition of j and ev_x is equal to the identity homomorphism, so is the composition of $\text{ev}_x^*: H^*(BS^1; \mathbb{Z}) \rightarrow H^*(B\text{Gau}(P); \mathbb{Z})$ and $j^*: H^*(B\text{Gau}(P); \mathbb{Z}) \rightarrow H^*(BS^1; \mathbb{Z})$. Hence, the first Chern class of $B\mathbb{Z} \rightarrow BC(X, \mathbb{R}) \rightarrow B\text{Gau}(P)$ is equal to $\text{ev}_x^* c_1$ and non-zero, where $c_1 \in H^2(BS^1; \mathbb{Z})$ is the universal first Chern class.

Let $e(\mathbb{R}) \in H_{\text{grp}}^2(S^1; \mathbb{Z})$ be the group cohomology class corresponding to the central extension $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow S^1 \rightarrow 1$. It is known that the class $c_1 \in H^2(BS^1; \mathbb{Z})$ corresponds to

$e(\mathbb{R})$ under the canonical isomorphism

$$\iota^*: H^2(BS^1; \mathbb{Z}) \rightarrow H^2(B(S^1)^\delta; \mathbb{Z}) = H_{\text{grp}}^2(S^1; \mathbb{Z}),$$

that is, $\iota^*c_1 = e(\mathbb{R})$.

3. PROOFS

Recall that $c \in H^2(X; \mathbb{Z})$ is a non-zero class and $G = \text{Homeo}(X, c)$ is the group of c -preserving homeomorphisms of X .

Lemma 3.1. Let $P \rightarrow X$ be a principal S^1 -bundle with the first Chern class $c \in H^2(X; \mathbb{Z})$. Let $\text{Aut}(P)$ be the group of bundle automorphisms of P . Then the canonical projection $p: \text{Aut}(P) \rightarrow \text{Homeo}(X)$ gives the surjection $\text{Aut}(P) \rightarrow G$. In particular, we have the following exact sequence

$$(3.1) \quad 0 \rightarrow \text{Gau}(P) \rightarrow \text{Aut}(P) \rightarrow G \rightarrow 1.$$

Remark 3.2. The exact sequence above gives rise to a Serre fibration with respect to the compact-open topology (see [Yam95]).

Proof of lemma 3.1. For a bundle automorphism $F: P \rightarrow P$, the induced homeomorphism $p(F): X \rightarrow X$ preserves the class c by the naturality of the Chern class. Hence the homeomorphism $p(F)$ is in G . For a homeomorphism $f \in G$, let $f^*P \rightarrow X$ denote the pullback bundle of P by f . Since the first Chern class of $f^*P \rightarrow X$ is equal to $f^*c = c$ and the first Chern class completely determines the isomorphism class of principal S^1 -bundles, the bundles $f^*P \rightarrow X$ is isomorphic to $P \rightarrow X$. Hence we have the following diagram

$$\begin{array}{ccccc} P & \xrightarrow{\cong} & f^*P & \longrightarrow & P \\ \downarrow & & \downarrow & & \downarrow \\ X & \xrightarrow{\text{id}_X} & X & \xrightarrow{f} & X, \end{array}$$

and this gives a bundle automorphism of P that covers the homeomorphism f . $\square$

Remark 3.3. The G -action on $C(X, S^1) = \text{Gau}(P)$ induced from the $\text{Aut}(P)$ -conjugation action is given by the pullback, that is, $\lambda \cdot g = g^*\lambda = \lambda \circ g$ for $\lambda \in C(X, S^1)$ and $g \in G$.

Let $(E_r'^{p,q}, d_r'^{p,q})$ and $(E_r''^{p,q}, d_r''^{p,q})$ be the Hochschild-Serre spectral sequences of exact sequence (3.1) with coefficients in $\text{Gau}(P)$ and $\mathbb{Z}$, respectively. Here $\mathbb{Z}$ is regarded as a trivial module. Then

$$E_2'^{0,1} = H_{\text{grp}}^0(G; H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P))) = H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P))^G,$$

where $H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P))^G$ is the group of G -equivariant homomorphisms on $\text{Gau}(P)$. Since $H_{\text{grp}}^1(\text{Gau}(P); \mathbb{Z})$ is trivial (Subsection 2.4), so is $E_2''^{p,1} = H_{\text{grp}}^p(G; H_{\text{grp}}^1(\text{Gau}(P); \mathbb{Z}))$ for all $p \geq 0$. Hence $d_2''^{0,2} = 0$, and we have

$$E_3''^{0,2} = E_2''^{0,2} = H_{\text{grp}}^0(G; H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})) = H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})^G,$$

where $H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})^G$ is the group of G -invariant cohomology classes.

Lemma 3.4. The connecting homomorphism

$$\mathbf{d}: H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P)) \rightarrow H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})$$

with respect to exact sequence (1.3) of coefficients induces the map

$$\mathbf{d}: H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P))^G \rightarrow H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})^G.$$

Proof. Let $[\phi]$ be an element of $H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P))^G$. Note that the cocycle ϕ is a G -equivariant homomorphism, that is, the following holds

$$\phi(\lambda \cdot g) = \phi(\lambda) \cdot g = \phi(\lambda) \circ g$$

for any $\lambda \in \text{Gau}(P)$ and $g \in G$. For a point x in X , let $s_x(\lambda): X \rightarrow \mathbb{R}$ denote the lift of $\lambda: X \rightarrow S^1 = \mathbb{R}/\mathbb{Z}$ satisfying $s_x(\lambda)(x) \in [0, 1)$. This map s_x defines a section

$$s_x: (\text{Gau}(P) =) C(X, S^1) \rightarrow C(X, \mathbb{R}).$$

Then, by the definition of the connecting homomorphism, a cocycle $c_x \in C_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})$ representing $\mathbf{d}[\phi] \in H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})$ is given by

$$\begin{aligned} c_x(\lambda_1, \lambda_2) &= \delta(s_x \circ \phi)(\lambda_1, \lambda_2) \\ &= s_x(\phi(\lambda_2)) - s_x(\phi(\lambda_1 \lambda_2)) + s_x(\phi(\lambda_1)) \end{aligned}$$

for $\lambda_1, \lambda_2 \in \text{Gau}(P)$. For $g \in G$, we set $y = g(x)$. Note that $s_x(\lambda \circ g) = (s_y(\lambda)) \circ g$ for any $\lambda \in \text{Gau}(P)$. Since the G -action $\text{Gau}(P)$ is given by pullback (Remark 3.3), we have

$$\begin{aligned} (c_x \cdot g)(\lambda_1, \lambda_2) &= c_x(\lambda_1 \circ g, \lambda_2 \circ g) \\ &= s_x(\phi(\lambda_2 \circ g)) - s_x(\phi((\lambda_1 \lambda_2) \circ g)) + s_x(\phi(\lambda_1 \circ g)) \\ &= s_x(\phi(\lambda_2) \circ g) - s_x(\phi(\lambda_1 \lambda_2) \circ g) + s_x(\phi(\lambda_1) \circ g) \\ &= (s_y(\phi(\lambda_2)) - s_y(\phi(\lambda_1 \lambda_2)) + s_y(\phi(\lambda_1))) \circ g \\ &= s_y(\phi(\lambda_2)) - s_y(\phi(\lambda_1 \lambda_2)) + s_y(\phi(\lambda_1)) \\ &= c_y(\lambda_1, \lambda_2). \end{aligned}$$

Since the image $\mathbf{d}[\phi]$ of the connecting homomorphism is independent of the choice of sections, we have

$$(\mathbf{d}[\phi]) \cdot g = [c_x \cdot g] = [c_y] = \mathbf{d}[\phi].$$

Hence the cohomology class $\mathbf{d}[\phi]$ is in $H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})^G$. $\square$

Remark 3.5. Let $e(C(X, \mathbb{R})) \in H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})$ be the class corresponding to the central extension $0 \rightarrow \mathbb{Z} \rightarrow C(X, \mathbb{R}) \rightarrow \text{Gau}(P) \rightarrow 0$. Then, by the direct computation, we can show that

$$-\mathbf{d}(\text{id}_{\text{Gau}(P)}) = e(C(X, \mathbb{R})).$$

By the commutative diagram of central extensions

$$\begin{array}{ccccc} \mathbb{Z} & \longrightarrow & C(X, \mathbb{R}) & \longrightarrow & \text{Gau}(P) \\ \parallel & & \downarrow \text{ev}_x^{\mathbb{R}} & & \downarrow \text{ev}_x \\ \mathbb{Z} & \longrightarrow & \mathbb{R} & \longrightarrow & S^1, \end{array}$$

we have $e(C(X, \mathbb{R})) = \text{ev}_x^* e(\mathbb{R}) = \text{ev}_x^* \iota^* c_1$.

Lemma 3.6. The following diagram

$$(3.2) \quad \begin{array}{ccc} H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P))^G = E_2'^{0,1} & \xrightarrow{d_2'^{0,1}} & E_2'^{2,0} = H_{\text{grp}}^2(G; \text{Gau}(P)) \\ \downarrow \mathbf{d} & & \downarrow \mathbf{d} \\ H_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})^G = E_3''^{0,2} & \xrightarrow{-d_3''^{0,2}} & E_3''^{3,0} = H_{\text{grp}}^3(G; \mathbb{Z}) \end{array}$$

commutes, where the vertical maps $\mathbf{d}$ are the connecting homomorphisms.

Proof. Let $[\phi]$ be an element of $H_{\text{grp}}^1(\text{Gau}(P); \text{Gau}(P))^G$ and $s: G \rightarrow \text{Aut}(P)$ a section such that $s(\text{id}_X) = \text{id}_P$. First, we give a cocycle representing $\mathbf{d}d_2'^{0,1}[\phi]$. Let us define a cochain $\phi_s: \text{Aut}(P) \rightarrow \text{Gau}(P)$ by

$$\phi_s(F) = \phi(F \circ (s(p(F)))^{-1}) \in \text{Gau}(P),$$

where $p: \text{Aut}(P) \rightarrow G$ is the projection. Then, a cocycle $c \in C_{\text{grp}}^3(G; \text{Gau}(P))$ is defined by

$$p^*c = \delta\phi_s.$$

Indeed, $\delta\phi_s(F_1, F_2) \in \text{Gau}(P)$ does depend on $p(F_1)$ and $p(F_2) \in G$ since the restriction $\phi_s|_{\text{Gau}(P)}$ is equal to a homomorphism ϕ . Hence, by Corollary 2.3, the cocycle c represents $d_2'^{0,1}[\phi]$, and a cocycle representing $\mathbf{d}d_2'^{0,1}[\phi]$ is given by

$$(3.3) \quad \delta(s_x \circ c) \in C_{\text{grp}}^3(G; \mathbb{Z}),$$

where $s_x: \text{Gau}(P) \rightarrow C(X, \mathbb{R})$ is the section defined in the proof of Lemma 3.4.

Next, we give a cocycle representing $d_3''^{0,2}\mathbf{d}[\phi]$. A cocycle of $\mathbf{d}[\phi]$ is given by the coboundary $c_x = \delta(s_x \circ \phi) \in C_{\text{grp}}^2(\text{Gau}(P); \mathbb{Z})$. We set

$$c'_x = \delta(s_x \circ \phi_s) - p^*(s_x \circ c) \in C_{\text{grp}}^2(\text{Aut}(P); C(X, \mathbb{R})).$$

Then, the cochain c'_x is in $C_{\text{grp}}^2(\text{Aut}(P); \mathbb{Z})$ since it is trivial under the projection $C(X, \mathbb{R}) \rightarrow C(X, S^1)$, and the restriction of c'_x on $\text{Gau}(P)$ is equal to c_x . Moreover, since the coboundary $\delta c'_x$ is equal to $-p^*\delta(s_x \circ c)$, a cocycle of $d_3''^{0,2}\mathbf{d}[\phi]$ is given by

$$(3.4) \quad -\delta(s_x \circ c) \in C_{\text{grp}}^3(G; \mathbb{Z})$$

by Corollary 2.4. By (3.3) and (3.4), we have $\mathbf{d}d_2'^{0,1}[\phi] = -d_3''^{0,2}\mathbf{d}[\phi]$ and the lemma follows. $\square$

Lemma 3.7. Let $X \rightarrow E \rightarrow BG$ be the universal G -bundle and $B\text{Gau}(P) \rightarrow B\text{Aut}(P) \rightarrow BG$ the fibration of classifying spaces of exact sequence (1.2). Then there exists a commutative diagram

$$(3.5) \quad \begin{array}{ccccc} X & \longrightarrow & E & \longrightarrow & BG \\ \downarrow f & & \downarrow \phi & & \parallel \\ B\text{Gau}(P) & \longrightarrow & B\text{Aut}(P) & \longrightarrow & BG, \end{array}$$

where the map $f: X \rightarrow B\text{Gau}(P)$ is the composition of the classifying map $X \rightarrow BS^1$ and the map $BS^1 \rightarrow B\text{Gau}(P)$ induced from the inclusion $S^1 \rightarrow \text{Gau}(P)$.

Proof. Let $P_G = P \times_{\text{Gau}(P)} \text{Gau}(P) \rightarrow X$ be the associated bundle. Then the bundle

$$E = EG \times_{\text{Aut}(P)} P_G \rightarrow BG$$

gives one of the models of the universal G -bundle with fiber X . Since $EG \times P_G \rightarrow E$ is a principal $\text{Aut}(P)$ -bundle, there exists a bundle map to the universal principal $\text{Aut}(P)$ -bundle

$$\begin{array}{ccc} EG \times P_G & \xrightarrow{\Psi} & E \text{Aut}(P) \\ \downarrow & & \downarrow \\ E & \xrightarrow{\psi} & B \text{Aut}(P). \end{array}$$

Define a map $\Phi: EG \times P_G \rightarrow EG \times E \text{Aut}(P)$ by $\Phi(a, p) = (a, \Psi(a, p))$. Then the map Φ gives a bundle map from $EG \times P_G \rightarrow E$ to $EG \times E \text{Aut}(P) \rightarrow EG \times_{\text{Aut}(P)} E \text{Aut}(P) = B \text{Aut}(P)$. Let $\phi: E \rightarrow B \text{Aut}(P)$ be the classifying map that is covered by Φ . Then the map ϕ covers the identity on BG . Moreover, the restriction $\phi|_X = f: X \rightarrow B \text{Gau}(P)$ gives rise to the classifying map of the bundle $P_G \rightarrow X$. $\square$

Proof of Theorem 1.1. Let us consider the Serre spectral sequences $E_r^{p,q}$ and $E_r^{\prime\prime\prime p,q}$ of the fibrations $X \rightarrow E \rightarrow BG$ and $B \text{Gau}(P) \rightarrow B \text{Aut}(P) \rightarrow BG$ respectively. Since $H^1(B \text{Gau}(P); \mathbb{Z}) = 0$ by Subsection 2.4, we have $E_3^{\prime\prime\prime 0,2} = E_2^{\prime\prime\prime 0,2} = H^2(B \text{Gau}(P); \mathbb{Z})$ and $E_3^{\prime\prime\prime 3,0} = E_2^{\prime\prime\prime 3,0} = H^3(BG; \mathbb{Z})$. By the naturality of the Serre spectral sequence with respect to commutative diagram (3.5), we have the commutative diagram

$$\begin{array}{ccc} H^2(B \text{Gau}(P); \mathbb{Z}) = E_3^{\prime\prime\prime 0,2} & \xrightarrow{d_3^{\prime\prime\prime 0,2}} & E_3^{\prime\prime\prime 3,0} = H^3(BG; \mathbb{Z}) \\ \downarrow f^* & & \parallel \\ H^2(X; \mathbb{Z}) = E_3^{0,2} & \xrightarrow{d_3^{0,2}} & E_3^{3,0} = H^3(BG; \mathbb{Z}). \end{array}$$

Then we have $f^* \text{ev}_x^* c_1 = c \in H^2(X; \mathbb{Z})$ since $f: X \rightarrow B \text{Gau}(P)$ is the composite of the classifying map $X \rightarrow BS^1$ and the map $j: BS^1 \rightarrow B \text{Gau}(P)$ (Subsection 2.4). Hence, the universal characteristic class $D_c = d_3^{0,2}(c) \in H^3(BG; \mathbb{Z})$ is equal to $-d_3^{\prime\prime\prime 0,2} \text{ev}_x^* c_1$. Let $E_r^{\prime\prime p,q}$ denote the Serre spectral sequence of the fibration $B \text{Gau}(P)^\delta \rightarrow B \text{Aut}(P)^\delta \rightarrow BG^\delta$ (or, equivalently, the Hochschild-Serre spectral sequence of $1 \rightarrow \text{Gau}(P) \rightarrow \text{Aut}(P) \rightarrow G \rightarrow 1$). By the commutative diagram

$$\begin{array}{ccccc} B \text{Gau}(P)^\delta & \longrightarrow & B \text{Aut}(P)^\delta & \longrightarrow & BG^\delta \\ \downarrow & & \downarrow & & \downarrow \\ B \text{Gau}(P) & \longrightarrow & B \text{Aut}(P) & \longrightarrow & BG \end{array}$$

and the naturality, we have the commutative diagram

$$\begin{array}{ccc} H^2(B \operatorname{Gau}(P); \mathbb{Z}) & \xrightarrow{d_3'''^{0,2}} & H^3(BG; \mathbb{Z}) \\ \downarrow \iota^* & & \downarrow \iota^* \\ H_{\operatorname{grp}}^2(\operatorname{Gau}(P); \mathbb{Z}) & \xrightarrow{d_3''^{0,2}} & H_{\operatorname{grp}}^3(G; \mathbb{Z}), \end{array}$$

where we identify the group cohomology with the singular cohomology of classifying space of discrete groups. Then we have

$$\iota^* D_c = -\iota^* d_3'''^{0,2} \operatorname{ev}_x^* c_1 = -d_3''^{0,2} \iota^* \operatorname{ev}_x^* c_1 = -d_3''^{0,2} \operatorname{ev}_x^* \iota^* c_1.$$

By commutative diagram (3.2) and lemma 2.5, we have

$$\mathbf{de}(\operatorname{Aut}(P)) = -\mathbf{d} d_2'^{0,1}(\operatorname{id}_{\operatorname{Gau}(P)}) = d_3''^{0,2} \mathbf{d}(\operatorname{id}_{\operatorname{Gau}(P)}).$$

Since the class $\operatorname{ev}_x^* \iota^* c_1$ is equal to $-\mathbf{d}(\operatorname{id}_{\operatorname{Gau}(P)})$ by Remark 3.5, we have

$$\iota^* D_c = \mathbf{de}(A_G(P)).$$

□

4. CENTRAL S^1 -EXTENSION OF G

In this section, we assume that the connected topological space X admits the universal covering space $\tilde{X}$. We take a non-zero cohomology class $c \in H^2(X; \mathbb{Z})$ which is equal to zero in $H^2(X; \mathbb{R})$.

4.1. Construction of a central S^1 -extension. By the cohomology long exact sequence

$$\cdots \rightarrow H^1(X; \mathbb{R}) \rightarrow H^1(X; S^1) \xrightarrow{\mathbf{d}} H^2(X; \mathbb{Z}) \rightarrow H^2(X; \mathbb{R}) \rightarrow \cdots,$$

there exists an element $\rho \in H^1(X; S^1)$ such that $\mathbf{d}(\rho) = c$. By the isomorphism $H^1(X; S^1) \cong \operatorname{Hom}(\pi_1(X); S^1)$, we regard the class ρ as a homomorphism $\rho: \pi_1(X) \rightarrow S^1$. We set $P_\rho^\delta = \tilde{X} \times_\rho (S^1)^\delta$. Then $P_\rho^\delta \rightarrow X$ is a foliated S^1 -bundle with the holonomy homomorphism ρ . Note that the group $\operatorname{Homeo}(X, \rho)$ of ρ -preserving homeomorphisms of X is equal to G since $\mathbf{d}$ is injective.

Lemma 4.1. Let $\operatorname{Aut}(P_\rho^\delta)$ denote the bundle automorphisms of P_ρ^δ . Then the canonical projection $\pi: \operatorname{Aut}(P_\rho^\delta) \rightarrow \operatorname{Homeo}(X)$ induces a surjection $\operatorname{Aut}(P_\rho^\delta) \rightarrow G$ and its kernel is isomorphic to S^1 . In particular, we obtain an exact sequence

$$(4.1) \quad 1 \rightarrow S^1 \rightarrow \operatorname{Aut}(P_\rho^\delta) \rightarrow G \rightarrow 1.$$

Proof. We regard G as the group of ρ -preserving homeomorphisms of X . First, we show that the image $\pi(\operatorname{Aut}(P_\rho^\delta))$ is contained in G . Take a bundle automorphism $F \in \operatorname{Aut}(P_\rho^\delta)$ that covers a homeomorphism $f: X \rightarrow X$. Let $\gamma: [0, 1] \rightarrow X$ be a loop and $\tilde{\gamma}: [0, 1] \rightarrow P_\rho^\delta$ a lift of γ . Then $F \circ \tilde{\gamma}: [0, 1] \rightarrow P_\rho^\delta$ is a lift of the loop $f \circ \gamma$. Since $\rho: \pi_1(X) \rightarrow (S^1)^\delta$ is the holonomy of P_ρ^δ , the value $\rho(f \circ \gamma) \in (S^1)^\delta$ is given by

$$(4.2) \quad F(\tilde{\gamma}(1)) = F(\tilde{\gamma}(0)) \cdot \rho(f \circ \gamma).$$

Since the left-hand side of (4.2) is equal to $F(\tilde{\gamma}(0) \cdot \rho(\gamma)) = F(\tilde{\gamma}(0)) \cdot \rho(\gamma)$, we have $\rho(\gamma) = \rho(f \circ \gamma) = f^* \rho(\gamma)$. Hence the homeomorphism f preserves the holonomy ρ and we have $\pi(\text{Aut}(P_\rho^\delta)) \subset G$.

Next we show that the map $\pi: \text{Aut}(P_\rho^\delta) \rightarrow G$ is surjective. For a homeomorphism $f: X \rightarrow X$ that preserves ρ , we take a lift $\tilde{f}: \tilde{X} \rightarrow \tilde{X}$. Then the map $\tilde{X} \times (S^1)^\delta \rightarrow \tilde{X} \times (S^1)^\delta$ defined by $(x, u) \rightarrow (\tilde{f}(x), u)$ induces a bundle automorphism $F: P_\rho^\delta \rightarrow P_\rho^\delta$ that covers f . Hence the map $\pi: \text{Aut}(P_\rho^\delta) \rightarrow G$ is surjective.

The kernel of $\pi: \text{Aut}(P_\rho^\delta) \rightarrow G$ is the gauge group $\text{Gau}(P_\rho^\delta)$ and this is isomorphic to $C(X, (S^1)^\delta) \cong (S^1)^\delta$ since the fiber of P_ρ^δ is the discrete group $(S^1)^\delta$ and X is connected. $\square$

Note that the group S^1 is in the center of $\text{Aut}(P_\rho^\delta)$, that is, (4.1) is a central S^1 -extension of G . Let $e(\text{Aut}(P_\rho^\delta)) \in H_{\text{grp}}^2(G; S^1)$ be the cohomology class corresponding to (4.1). Let $P_\rho \rightarrow X$ be a principal S^1 -bundle defined by $P_\rho = \tilde{X} \times_\rho S^1$. Then we have the following commutative diagram of groups

$$(4.3) \quad \begin{array}{ccccccc} 1 & \longrightarrow & S^1 & \longrightarrow & \text{Aut}(P_\rho^\delta) & \longrightarrow & G \longrightarrow 1 \\ & & \downarrow j & & \downarrow k & & \parallel \\ 1 & \longrightarrow & \text{Gau}(P_\rho) & \longrightarrow & \text{Aut}(P_\rho) & \longrightarrow & G \longrightarrow 1, \end{array}$$

where $j: S^1 \rightarrow \text{Gau}(P_\rho)$ is the inclusion. Let $e(\text{Aut}(P_\rho)) \in H_{\text{grp}}^2(G; \text{Gau}(P_\rho))$ be the cohomology class corresponding to the abelian extension $\text{Aut}(P_\rho)$.

Lemma 4.2. Let $j_*: H_{\text{grp}}^2(G; S^1) \rightarrow H_{\text{grp}}^2(G; \text{Gau}(P_\rho))$ be the change of coefficients homomorphism induced by j . Then we have $j_*(e(\text{Aut}(P_\rho^\delta))) = e(\text{Aut}(P_\rho))$ in $H_{\text{grp}}^2(G; \text{Gau}(P_\rho))$.

Proof. Recall that a cocycle c representing $e(\text{Aut}(P_\rho^\delta))$ is given by

$$c(g, h) = s(g)s(h)s(gh)^{-1},$$

where $s: G \rightarrow \text{Aut}(P_\rho^\delta)$ is a section. Since $k \circ s: G \rightarrow \text{Aut}(P_\rho)$ is a section of $\text{Aut}(P_\rho) \rightarrow G$, a cocycle c' representing $e(\text{Aut}(P_\rho))$ is given by

$$c'(g, h) = k(s(g))k(s(h))k(s(gh))^{-1} = k(s(g)s(h)s(gh)^{-1}) = j(c(g, h)).$$

Hence we have $j_*(e(\text{Aut}(P_\rho^\delta))) = e(\text{Aut}(P_\rho))$. $\square$

Proof of Theorem 1.2. By the commutative diagram of coefficients

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{R} & \longrightarrow & S^1 \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow j \\ 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & C(X, \mathbb{R}) & \longrightarrow & C(X, S^1) = \text{Gau}(P_\rho) \longrightarrow 1, \end{array}$$

we have the commutative diagram

$$\begin{array}{ccc} H_{\text{grp}}^2(G; S^1) & \xrightarrow{\mathbf{d}} & H_{\text{grp}}^3(G; \mathbb{Z}) \\ \downarrow j^* & & \parallel \\ H_{\text{grp}}^2(G; \text{Gau}(P_\rho)) & \xrightarrow{\mathbf{d}} & H_{\text{grp}}^3(G; \mathbb{Z}), \end{array}$$

where the maps $\mathbf{d}$ are the connecting homomorphisms. Together with Theorem 1.1 and Lemma 4.2, we obtain

$$\mathbf{d}e(A_G(P_\rho^\delta)) = \mathbf{d}j_*e(A_G(P_\rho^\delta)) = \mathbf{d}e(A_G(P_\rho)) = \iota^*D_c.$$

□

4.2. Cocycle description of $e(\text{Aut}(P_\rho^\delta))$. In this subsection, we give a geometric description of a group two-cocycle representing the class $e(\text{Aut}(P_\rho^\delta))$. Let $c \in H^2(X; \mathbb{Z})$ and $\rho \in H^1(X; S^1)$ be a cohomology class as above. Let $\alpha \in C^1(X; S^1)$ be a cocycle representing the cohomology class ρ , where $(C^*(X; S^1), d)$ denotes the singular cochain complex of X with coefficients in S^1 . The singular cochain $C^*(X; S^1)$ and the cohomology $H^*(X; S^1)$ are the right G -modules by pullback. Since $g^*\rho = \rho$ for any homeomorphism $g \in G$, the cochain $g^*\alpha - \alpha$ is a coboundary. Hence there exists a cochain $\mathfrak{K}_\alpha(g) \in C^0(X; S^1)$ satisfying

$$g^*\alpha - \alpha = d\mathfrak{K}_\alpha(g).$$

Then a two-cochain $\mathfrak{G}_{x,\alpha}$ in $C^2(G; S^1)$ is defined by

$$(4.4) \quad \mathfrak{G}_{x,\alpha}(g, h) = \int_x^{hx} g^*\alpha - \alpha = \mathfrak{K}_\alpha(g)(hx) - \mathfrak{K}_\alpha(g)(x),$$

where $x \in X$. Here the symbol $\int_x^{hx}$ denotes the pairing of the cocycle $g^*\alpha - \alpha$ and a path from x to hx . By the arguments same as in [ILM06, Theorem 3,1], we have the following proposition.

Proposition 4.3. The two-cochain $\mathfrak{G}_{x,\alpha} \in C_{\text{grp}}^2(G; S^1)$ is a cocycle and the cohomology class $[\mathfrak{G}_{x,\alpha}]$ is independent of the choice of the point $x \in X$ and the singular cochain α .

Then the following theorem holds.

Theorem 4.4. The cohomology class $[\mathfrak{G}_{x,\alpha}]$ is equal to $e(\text{Aut}(P_\rho)) \in H_{\text{grp}}^2(G; S^1)$.

For the proof of Theorem 4.4, we need the following lemmas. Let us recall that $p: P_\rho^\delta \rightarrow X$ is the principal $(S^1)^\delta$ -bundle with the holonomy ρ .

Lemma 4.5. The pullback $p^*\rho \in H^1(P_\rho^\delta; S^1)$ is equal to zero.

Proof. Let $\gamma: [0, 1] \rightarrow P_\rho^\delta$ be a based loop. Then γ is a lift of the loop $p \circ \gamma$ in X . Since $\rho: \pi(X) \rightarrow S^1$ is the holonomy homomorphism, we have

$$p^*\rho(\gamma) = \rho(p \circ \gamma) = \gamma(1) - \gamma(0) = 0,$$

and the lemma follows. □

By lemma 4.5, we take a singular cochain $\theta' \in C^0(P_\rho^\delta; S^1)$ satisfying $d\theta' = p^*\alpha$. We take a base point $y \in P_\rho^\delta$ and set $p(y) = x$. For any $u \in S^1$, let $P_\rho^\delta(u)$ denote the connected component of P_ρ^δ which contains the point $y \cdot u$. Then we set

$$z(u) = \theta'(y) - \theta'(y \cdot u) + u \in S^1.$$

By a straightforward calculation, we have the equality $z(u) = z(u + \rho(\gamma))$ for any $\gamma \in \pi_1(X)$. Hence, the value $z(u)$ gives a continuous function $z: P_\rho^\delta \rightarrow (S^1)^\delta$. Note that $z(y \cdot u) = z(u)$ and $z(y) = 0$. We set $\theta = \theta' + z \in C^0(X; \mathbb{Z})$. Since $dz = 0$, we have $d\theta = p^*\alpha$. Let $\tau \in C_{\text{grp}}^1(A_G(P_\rho^\delta); S^1)$ be a cochain defined by

$$\tau(\phi) = \int_y^{\phi y} \theta = \theta(\phi y) - \theta(y).$$

Lemma 4.6. The restriction $\tau|_{S^1}: S^1 \rightarrow S^1$ is equal to the identity.

Proof. By the definition of τ , we have

$$\tau(u) = \theta(y \cdot u) - \theta(y) = \theta'(y \cdot u) + z(y \cdot u) - \theta'(y) - z(y) = u$$

for any $u \in S^1$. □

Lemma 4.7. The equality

$$-\delta\tau = \pi^*\mathfrak{G}_{x,\alpha} \in C_{\text{grp}}^2(\text{Aut}(P_\rho^\delta); S^1)$$

holds, where $\pi: \text{Aut}(P_\rho^\delta) \rightarrow G$ is the projection.

Proof. For the pullback $p^*\mathfrak{K}_\alpha(g) \in C^0(P_\rho^\delta; S^1)$, we have

$$\begin{aligned} d(p^*\mathfrak{K}_\alpha(g)) &= p^*(d\mathfrak{K}_\alpha(g)) = p^*g^*\alpha - p^*\alpha \\ &= \phi^*p^*\alpha - p^*\alpha = \phi^*d\theta - d\theta = d(\phi^*\theta - \theta), \end{aligned}$$

where $\phi \in \text{Aut}(P_\rho^\delta)$ is a lift of $g \in G$. Hence, for $\phi, \psi \in \text{Aut}(P_\rho^\delta)$ which cover $g, h \in G$, respectively, we have

$$\begin{aligned} \pi^*\mathfrak{G}_{x,\alpha}(\phi, \psi) &= \mathfrak{G}_{p(y),\alpha}(g, h) = \mathfrak{K}_\alpha(g)(hp(y)) - \mathfrak{K}_\alpha(g)(p(y)) \\ &= \mathfrak{K}_\alpha(g)(p\psi(y)) - \mathfrak{K}_\alpha(g)(p(y)) \\ &= p^*\mathfrak{K}_\alpha(g)(\psi(y)) - p^*\mathfrak{K}_\alpha(g)(y) \\ &= (\phi^*\theta - \theta)(\psi(y)) - (\phi^*\theta - \theta)(y) \\ &= (\theta(\phi\psi(y)) - \theta(y)) - (\theta(\phi(y)) - \theta(y)) - (\theta(\psi(y)) - \theta(y)) \\ &= -\delta\tau(\phi, \psi), \end{aligned}$$

and the lemma follows. □

Proof of theorem 4.4. Let $E_r^{p,q}$ denote the Hochschild-Serre spectral sequence of $0 \rightarrow S^1 \rightarrow \text{Aut}(P_\rho^\delta) \rightarrow G \rightarrow 1$ with coefficients in S^1 . Then the derivation $d_2^{0,1}$ defines a map

$$d_2^{0,1}: H_{\text{grp}}^1(S^1; S^1) = E_2^{0,1} \rightarrow E_2^{2,0} = H_{\text{grp}}^2(G; S^1).$$

By Corollary 2.3, Lemma 4.6, and Lemma 4.7, we have $d_2^{0,1}(\text{id}_{S^1}) = -[\mathfrak{G}_{x,\alpha}]$. On the other hand, by Lemma 2.5, we have $d_2^{0,1}(\text{id}_{S^1}) = -e(\text{Aut}(P_\rho^\delta))$. Hence we have $[\mathfrak{G}_{x,\alpha}] = e(\text{Aut}(P_\rho^\delta))$. $\square$

5. EXAMPLES

In this section, we give two examples that the universal characteristic classes D_c and $\iota^* D_c$ are non-trivial. The first example is the complex projective space $\mathbb{C}P^n$. The proof of the following theorem is similar to the proof of [Mar20, Theorem 7.1] (and also similar to the proof of Theorem 5.2 below), so we will omit it.

Theorem 5.1. Let n be a positive integer and $c \in H^2(\mathbb{C}P^n; \mathbb{Z})$ the generator. Let G denote the group $\text{Homeo}(\mathbb{C}P^n, c)$ of c -preserving homeomorphisms. Then the universal characteristic classes $D_c \in H^3(BG; \mathbb{Z})$ and $\iota^* D_c \in H^3(BG^\delta; \mathbb{Z})$ are non-zero.

The cohomology class c in Theorem 5.1 is non-zero in $H^2(\mathbb{C}P^n; \mathbb{R})$. Hence this example does not satisfy the assumption in Section 4. An example which satisfies the assumption in Section 4 is given as follows.

Theorem 5.2. Let c be the non-zero element in $H^2(\mathbb{R}P^3; \mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$. Let G denote the group $\text{Homeo}(\mathbb{R}P^3, c)$ of c -preserving homeomorphisms. Then the universal characteristic classes $D_c \in H^3(BG; \mathbb{Z})$ and $\iota^* D_c \in H^3(BG^\delta; \mathbb{Z})$ are non-zero.

Proof. Since $\mathbb{R}P^3$ is homeomorphic to $SO(3)$, the group $SO(3)$ is included in G as the left translations. Hence we have the commutative diagram

$$\begin{array}{ccccc} SO(3) & \longrightarrow & ESO(3) & \longrightarrow & BSO(3) \\ \downarrow & & \downarrow F & & \downarrow f \\ G & \longrightarrow & EG & \longrightarrow & BG. \end{array}$$

Recall that a universal G -bundle $\mathbb{R}P^3 \rightarrow E \rightarrow BG$ is given as $E = EG \times_G \mathbb{R}P^3$. For the base point $b \in \mathbb{R}P^3$ corresponding to the unit $1 \in SO(3)$, we define a map $\phi: ESO(3) \rightarrow E$ by $\Phi(x) = [F(x), b]$ for $x \in ESO(3)$. Then we have the commutative diagram

$$(5.1) \quad \begin{array}{ccccc} SO(3) & \longrightarrow & ESO(3) & \longrightarrow & BSO(3) \\ \parallel & & \downarrow \Phi & & \downarrow f \\ SO(3) \cong \mathbb{R}P^3 & \longrightarrow & E & \longrightarrow & BG. \end{array}$$

Let us consider the Serre spectral sequences of the fibrations in (5.1). By the naturality of the Serre spectral sequence, the pullback $f^* D_c \in H^3(BSO(3); \mathbb{Z})$ is equal to the transgression image of $c \in H^2(\mathbb{R}P^3; \mathbb{Z}) \cong H^2(SO(3); \mathbb{Z})$ with respect to the fibration $SO(3) \rightarrow ESO(3) \rightarrow BSO(3)$. Since $ESO(3)$ is contractible, the transgression map is injective. Hence the class $f^* D_c$ is non-zero, and so is D_c .

Let us consider the commutative diagram

$$\begin{array}{ccc} H^3(BG; \mathbb{Z}) & \xrightarrow{\iota^*} & H^3(BG^\delta; \mathbb{Z}) \\ \downarrow f^* & & \downarrow \\ H^3(BSO(3); \mathbb{Z}) & \xrightarrow{\iota^*} & H^3(BSO(3)^\delta; \mathbb{Z}). \end{array}$$

Since the map $\iota^*: H^3(BSO(3); \mathbb{Z}) \rightarrow H^3(BSO(3)^\delta; \mathbb{Z})$ is injective [Mil83], the class $\iota^* f^* D_c \in H^3(BSO(3)^\delta; \mathbb{Z})$ is non-zero. Hence the class $\iota^* D_c \in H^3(BG^\delta; \mathbb{Z})$ is also non-zero. $\square$

Remark 5.3. By Theorem 1.2 and Theorem 5.2, the class $e(\text{Aut}(P_\rho))$ is also non-zero for $G = \text{Homeo}(\mathbb{R}P^3, c)$. Hence, this gives an example that cocycle (4.4) is cohomologically non-trivial.

Remark 5.4. The non-triviality of the class c does not imply the non-triviality of D_c nor $\iota^* D_c$. For example, we consider the real projective plane $\mathbb{R}P^2$. Let $c \in H^2(\mathbb{R}P^2; \mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ be a non-zero element and set $G = \text{Homeo}(\mathbb{R}P^2, c)$. Then the class $D_c \in H^3(BG; \mathbb{Z})$ is equal to zero. To see this, we consider the inclusion $i: SO(3) \hookrightarrow \text{Homeo}_0(\mathbb{R}P^2)$ induced from the natural $SO(3)$ -action on $\mathbb{R}P^2$. This inclusion is a homotopy equivalence ([Dob21, Corollary 1.2]), and hence the map $i^*: H^3(BG; \mathbb{Z}) \rightarrow H^3(BSO(3); \mathbb{Z})$ is an isomorphism. Let us consider the commutative diagram of fibrations

$$(5.2) \quad \begin{array}{ccccc} SO(3) & \longrightarrow & ESO(3) & \longrightarrow & BSO(3) \\ \downarrow f & & \downarrow & & \downarrow \\ \mathbb{R}P^2 & \longrightarrow & EG \times_G \mathbb{R}P^2 & \longrightarrow & BG. \end{array}$$

Since $f: SO(3) \rightarrow \mathbb{R}P^2$ is induced from the natural $SO(3)$ -action on S^2 , the map f factors through S^2 . Since $H^2(SO(3); \mathbb{Z}) \cong H^2(\mathbb{R}P^2; \mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ and $H^2(S^2; \mathbb{Z}) \cong \mathbb{Z}$, the map f induces the zero map on second cohomology. The naturality of the Dixmier-Douady class with respect to (5.2), together with the fact that the induced map $i^*: H^3(BG; \mathbb{Z}) \rightarrow H^3(BSO(3); \mathbb{Z})$ is an isomorphism, implies that the Dixmier-Douady class D_c is equal to zero.

REFERENCES

- [Bro82] Kenneth S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York-Berlin, 1982.
- [Bry93] Jean-Luc Brylinski, *Loop spaces, characteristic classes and geometric quantization*, Progress in Mathematics, vol. 107, Birkhäuser Boston, Inc., Boston, MA, 1993.
- [CCM98] Alan L. Carey, Diarmuid Crowley, and Michael K. Murray, *Principal bundles and the Dixmier-Douady class*, Comm. Math. Phys. **193** (1998), no. 1, 171–196.
- [DK01] James F. Davis and Paul Kirk, *Lecture notes in algebraic topology*, Graduate Studies in Mathematics, vol. 35, American Mathematical Society, Providence, RI, 2001.
- [Dob21] Michael Gene Dobbins, *A strong equivariant deformation retraction from the homeomorphism group of the projective plane to the special orthogonal group*, arXiv:2108.02134 (2021).
- [Fri17] Roberto Frigerio, *Bounded cohomology of discrete groups*, Mathematical Surveys and Monographs, vol. 227, American Mathematical Society, Providence, RI, 2017.
- [GK11] Światosław R. Gal and Jarek Kędra, $\int_x^{hx} (g^* \alpha - \alpha)$, arXiv:1105.0825 (2011).

- [HS53] G. Hochschild and J.-P. Serre, *Cohomology of group extensions*, Trans. Amer. Math. Soc. **74** (1953), 110–134.
- [ILM06] Rais S. Ismagilov, Mark Losik, and Peter W. Michor, *A 2-cocycle on a symplectomorphism group*, Mosc. Math. J. **6** (2006), no. 2, 307–315, 407.
- [Mar20] Shuhei Maruyama, *The dixmier-douady class, the action homomorphism, and group cocycles on the symplectomorphism group*.
- [Mil83] J. Milnor, *On the homology of Lie groups made discrete*, Comment. Math. Helv. **58** (1983), no. 1, 72–85.
- [NSW08] Jürgen Neukirch, Alexander Schmidt, and Kay Wingberg, *Cohomology of number fields*, second ed., Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 323, Springer-Verlag, Berlin, 2008.
- [SS20] Yasha Savelyev and Egor Shelukhin, *K-theoretic invariants of Hamiltonian fibrations*, J. Symplectic Geom. **18** (2020), no. 1, 251–289.
- [Yam95] Tsuneyo Yamanoshita, *On the group of S^1 -equivariant homeomorphisms of the 3-sphere*, Publ. Res. Inst. Math. Sci. **31** (1995), no. 5, 953–958.

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, JAPAN

Email address: m17037h@math.nagoya-u.ac.jp