
NONPARAMETRIC CAUSAL MEDIATION ANALYSIS FOR STOCHASTIC INTERVENTIONAL (IN)DIRECT EFFECTS

Nima S. Hejazi

Graduate Group in Biostatistics, and
Center for Computational Biology,
University of California, Berkeley
nhejazi@berkeley.edu

Kara E. Rudolph

Department of Epidemiology,
Mailman School of Public Health,
Columbia University
kr2854@cumc.columbia.edu

Mark J. van der Laan

Division of Biostatistics,
School of Public Health, and
Department of Statistics,
University of California, Berkeley
laan@berkeley.edu

Iván Díaz

Division of Biostatistics,
Department of Population Health Sciences,
Weill Cornell Medicine
ild2005@med.cornell.edu

August 3, 2022

ABSTRACT

Causal mediation analysis has historically been limited in two important ways: (i) a focus has traditionally been placed on binary treatments and static interventions, and (ii) direct and indirect effect decompositions have been pursued that are only identifiable in the absence of intermediate confounders affected by treatment. We present a theoretical study of an (in)direct effect decomposition of the population intervention effect, defined by stochastic interventions jointly applied to the treatment and mediators. In contrast to existing proposals, our causal effects can be evaluated regardless of whether a treatment is categorical or continuous and remain well-defined even in the presence of intermediate confounders affected by treatment. Our (in)direct effects are identifiable without a restrictive assumption on cross-world counterfactual independencies, allowing for substantive conclusions drawn from them to be validated in randomized controlled trials. Beyond the novel effects introduced, we provide a careful study of nonparametric efficiency theory relevant for the construction of flexible, multiply robust estimators of our (in)direct effects, while avoiding undue restrictions induced by assuming parametric models of nuisance parameter functionals. To complement our nonparametric estimation strategy, we introduce inferential techniques for constructing confidence intervals and hypothesis tests, and discuss open source software, the `medshift` R package, implementing the proposed methodology. Application of our (in)direct effects and their nonparametric estimators is illustrated using data from a comparative effectiveness trial examining the direct and indirect effects of pharmacological therapeutics on relapse to opioid use disorder.

1 Introduction

In myriad applications, one is often interested in the effect of a treatment on an outcome only through a particular pathway between the two. Indeed, efforts in defining and identifying such *path-specific* effects have come to constitute a rich history in not only philosophy but also in the sciences of statistics, causal inference, epidemiology, economics, and psychology. In each of these disciplines, as well as in many others among the biomedical and social sciences, developing a mechanistic understanding of the complexities that admit representations as path-specific effects remains a central goal; examples include elucidating the biological mechanism by which a vaccine reduces infection risk (e.g., [Corey et al., 2015](#); [Hejazi et al., 2020](#)), assessing the effect on preterm birth of maternal exposure to environmental

toxins (e.g., [Ferguson et al., 2017](#)), and ascertaining the effect of novel pharmacological therapies on substance abuse disorder relapse.

The latter serves as our motivating example as we consider how exposure to a buprenorphine dose schedule characterized by successive increases toward a maximum dose early in treatment (versus static dose) affects the risk of relapse to opioid use disorder, both directly and indirectly through mediating factors such as depression and pain. Developing a detailed mechanistic understanding of the process by which such therapeutics modulate intermediary states is necessarily a *causal* question — one central to designing and successively improving upon available therapies in a manner targeted towards the mitigation of the risk of substance abuse relapse. In comparative effectiveness trials of promising opioid use disorder therapeutics, detailed dissections of the complex neurological and psychiatric pathways involved in the development of addiction disorders is of clinical interest ([Lee et al., 2018](#); [Rudolph et al., 2020a](#)). The ability to define and evaluate causal effects along paths involving or avoiding mediating neuropsychiatric sequela would facilitate drug efficacy assessments; moreover, the ability to further examine and refine scientific conclusions based on statistical evidence through randomized controlled trials remains key to furthering clinical progress.

To carefully study complex mediation relationships, a wealth of techniques rooted in statistical causal inference have been formulated. Path analysis ([Wright, 1921, 1934](#)), perhaps the earliest example of such methodology, directly inspired the development of subsequent techniques that leveraged parametric structural equation models (e.g., [Goldberger, 1972](#); [Baron & Kenny, 1986](#)) for mediation analysis. More recently, the advent of modern frameworks and formalisms for causal inference, including nonparametric structural equation models, directed acyclic graphs, and their underlying do-calculus ([Pearl, 1995, 2000](#)), provided the necessary foundational tools to express causal mechanisms without reliance on the more restrictive approach of parametric modeling.

In tandem with the developments of [Pearl \(2000\)](#), similar approaches spearheaded by [Robins \(1986\)](#), [Spirtes et al. \(2000\)](#), [Dawid \(2000\)](#), and [Richardson & Robins \(2013\)](#) both allowed nonparametric formulations of mediation analysis and uncovered significant limitations of the earlier efforts focused on parametric structural equation models ([Pearl, 1998](#); [Imai et al., 2010](#)). Recent applications of modern causal models have illustrated the failings of popular parametric modeling strategies (i.e., [Baron & Kenny, 1986](#)), in the presence of intermediate confounders of the mediator-outcome relationship ([Cole & Hernán, 2002](#)). Consequently, the usually implausible assumptions that underlie such restrictive structural equation models make these approaches of limited applicability for the examination of complex phenomena in the biomedical, health, and social sciences.

Modern approaches to causal inference have allowed for significant advances over the methodology of traditional mediation analysis, overcoming the significant restrictions imposed by the use of parametric structural equation modeling. For example, both [Robins & Greenland \(1992\)](#), using the potential outcomes framework, and [Pearl \(2006\)](#), using nonparametric structural equation models, provided equivalent nonparametric decompositions of the average treatment effect (for binary treatments) into *natural* direct and indirect effects, which quantify all effects of the treatment on the outcome through paths avoiding the mediator and all paths involving the mediator, respectively. Such advances were not without their own limitations, however. A key assumption of the nonparametric decomposition of the average treatment effect is the requirement of *cross-world* counterfactual independencies (i.e., the condition that counterfactuals indexed by distinct intervention assignments be independent). Unfortunately, such an assumption limits the scientific relevance of the natural (in)direct effects by making them unidentifiable in randomized trials, directly implying that corresponding scientific claims cannot be falsified through experimentation ([Popper, 1934](#); [Dawid, 2000](#); [Robins & Richardson, 2010](#)). Importantly, such cross-world independencies are also unsatisfied in the presence of intermediate confounders affected by treatment ([Avin et al., 2005](#); [Tchetgen Tchetgen & VanderWeele, 2014](#)). As such confounders are often present in practice, the natural (in)direct effects are fundamentally of limited applicability.

To circumvent the limitations imposed by cross-world counterfactual independencies, a rich literature has recently developed — with alternative definitions of the (in)direct effects ([Petersen et al., 2006](#); [van der Laan & Petersen, 2008](#); [Vansteelandt & VanderWeele, 2012](#); [VanderWeele et al., 2014](#)). Such developments have led directly to the formulation of the new and promising family of *interventional* (in)direct effects ([Didelez et al., 2006](#); [VanderWeele et al., 2014](#); [Lok, 2016](#); [Vansteelandt & Daniel, 2017](#); [Zheng & van der Laan, 2017](#); [Rudolph et al., 2017](#); [Lok, 2019](#); [Nguyen et al., 2019](#)), for which nonparametric effect decompositions and corresponding efficiency theory have only recently become available ([Díaz et al., 2020](#); [Benkeser, 2020](#)). While resolving the issues arising from requiring cross-world counterfactual independencies, these interventional (in)direct effects are limited by their lack of applicability beyond binary treatments.

A related thread of the literature has considered stochastic interventions. The framework of stochastic interventions generalizes many intervention classes — for example, within this framework, static interventions result in post-intervention exposures that have degenerate distributions. [Stock \(1989\)](#) first considered the estimation of the total effects of stochastic interventions, while many others (e.g., [Robins et al., 2004](#); [Didelez et al., 2006](#); [Tian, 2008](#); [Pearl, 2009](#); [Stitelman et al., 2010](#); [Haneuse & Rotnitzky, 2013](#); [Díaz & van der Laan, 2013](#); [Dudík et al., 2014](#); [Young et al.,](#)

2014) provided careful studies that expanded the underlying theory of stochastic interventions and demonstrated their numerous applications. Uniquely, stochastic interventions can be applied to define causal effects of continuous-valued treatments, with an interpretation echoing that of regression adjustment. For example, [Díaz & van der Laan \(2012\)](#) and [Haneuse & Rotnitzky \(2013\)](#) described modified treatment policies, which assign post-intervention counterfactuals based on the natural value of the treatments; their methods were demonstrated in the context of increasing leisure physical activity in the elderly and reducing surgical time for non-small-cell lung cancer operations. Stochastic interventions have also successfully been applied to binary treatments: [Kennedy \(2019\)](#) proposed incremental propensity score interventions and demonstrated their use in assessing their corresponding causal effects in longitudinal studies while circumventing identifiability and estimation issues arising from positivity violations.

Contemporaneously, [Díaz & Hejazi \(2020\)](#) proposed a decomposition of the total effect of stochastic interventions into the population intervention (in)direct effects, which are endowed with interpretations analogous to that of the natural (in)direct effects. Prior related attempts at the same (e.g., [Vansteelandt et al., 2012](#)) introduced parametric modeling assumptions to lessen reliance on the assumption of cross-world counterfactual independencies, introducing flexibility at the cost of bias and restrictive assumptions on post-intervention distributions. In a similar vein, the stochastic (in)direct effects of [Díaz & Hejazi \(2020\)](#) do not require cross-world counterfactual independencies but simultaneously allow for nonparametric estimation strategies to be pursued. Consequently, these population intervention (in)direct effects may be estimated without restrictive assumptions and yield scientific results that can be tested through randomization of both the treatment and mediator. Despite these advances, current theory on the population intervention (in)direct effects cannot accommodate the presence of intermediate confounders affected by treatment, limiting their use to settings in which the absence of such confounders is assured.

In the present work, we provide a general framework for assessing causal mediation effects defined by stochastic interventions. Our proposed effects are the first to simultaneously avoid the requirement of cross-world counterfactual independencies; leverage the flexibility of stochastic interventions in order to ensure their applicability to binary, categorical, and continuous-valued treatments; and remain identifiable regardless of the presence of intermediate confounders affected by treatment. As such, our contributions may be applied to a broader class of treatments than the interventional effects (e.g., [Díaz et al., 2020](#); [Benkeser, 2020](#)) while generalizing prior work (e.g., [Díaz & Hejazi, 2020](#)) to accommodate the presence of intermediate confounders. To complement our (in)direct effects, we derive the efficient influence function (EIF) in the nonparametric model, using this to develop two efficient estimators based on the frameworks of one-step ([Pfanzagl & Wefelmeyer, 1985](#); [Bickel et al., 1993](#)) and targeted minimum loss estimation ([van der Laan & Rubin, 2006](#); [van der Laan & Rose, 2011, 2018](#)). Our flexible estimators remain regular and asymptotically linear even when their relevant nuisance parameters are estimated via data adaptive (machine learning) strategies; moreover, by utilizing the form of the EIF, these estimators are multiply robust, producing consistent point estimates even under certain configurations of nuisance parameter misspecification. Lastly, we provide implementations of our methodological advances in our free and open source `medshift` ([Hejazi & Díaz, 2020](#)) package, for the R language and environment for statistical computing ([R Core Team, 2020](#)).

2 Mediation analysis for population intervention effects

Let A denote a continuous or categorical treatment variable, let Y denote a continuous or binary outcome, let Z denote a mediator, let W denote a vector of observed pre-treatment covariates, and let L denote an intermediate (mediator-outcome) confounder affected by treatment. We formalize the causal inference problem using the following nonparametric structural equation model (NPSEM). Assume

$$\begin{aligned} W &= f_W(U_W); A = f_A(W, U_A); L = f_L(A, W, U_L); \\ Z &= f_Z(L, A, W, U_Z); Y = f_Y(Z, L, A, W, U_Y). \end{aligned} \quad (1)$$

In the NPSEM (1), $U = (U_W, U_A, U_L, U_Z, U_Y)$ is a vector of exogenous factors, and the functions f are assumed deterministic but unknown. This mechanistic model is assumed to generate the observed data O ; it encodes several fundamental assumptions. First, an implicit temporal ordering $W \rightarrow A \rightarrow L \rightarrow Z \rightarrow Y$ is assumed. Second, each variable (i.e., $\{W, A, L, Z, Y\}$) is assumed to be generated from the corresponding deterministic function of the observed variables that precede it temporally, plus an exogenous variable denoted by U . Each exogenous variable is assumed to contain all unobserved causes of the corresponding observed variable. Unlike the mediation effects for stochastic interventions presented in [Díaz & Hejazi \(2020\)](#), we consider an intermediate confounder L that is affected by treatment; our results generalize the methodology of [Díaz & Hejazi \(2020\)](#), extending their (in)direct effects to settings with intermediate confounding. For a random variable X , let X_a denote the counterfactual outcome observed in a hypothetical world in which $P(A = a) = 1$. For example, we have $L_a = f_L(a, W, U_L)$, $Z_a = f_Z(L_a, a, W, U_Z)$, and $Y_a = f_Y(Z_a, L_a, a, W, U_Y)$. Likewise, we let $Y_{a,z} = f_Y(z, L_a, a, W, U_Y)$ denote the value of the outcome in a hypothetical world where $P(A = a, Z = z) = 1$. Figure 1 represents model (1) in terms of a directed acyclic graph.




Figure 1: Directed Acyclic Graph of NPSEM (1).

Letting $O = (W, A, Z, L, Y)$ represent a random variable with distribution P , we denote by O_1, \dots, O_n a sample of n i.i.d. observations of O . We let $Pf = \int f(o)dP(o)$ for a given function $f(o)$. We use P_c to denote the joint distribution of (O, U) , and let E and E_c denote corresponding expectation operators. We use P_n to denote the empirical distribution of O_1, \dots, O_n , and assume $P \in \mathcal{M}$, where \mathcal{M} is the nonparametric statistical model defined as all continuous densities on O with respect to a dominating measure ν . Let p denote the corresponding probability density function. We use $g(a | w)$ and $e(a | z, w)$ to denote the probability density function or the probability mass function of A conditional on $W = w$ and (Z, W) , respectively; $m(z, l, a, w)$ to denote the outcome regression function $E(Y | Z = z, L = l, A = a, W = w)$. Let $g(\cdot | w)$ and $e(\cdot | z, w)$ be dominated by a measure $\kappa(a)$ (e.g., the counting measure for binary A and the Lebesgue measure for continuous A). We will use the parameterizations

$$\frac{p(z | w)}{p(z | a, w)} = \frac{g(a | w)}{e(a | z, w)}; \quad \frac{p(z | a, w)}{p(z | l, a, w)} = \frac{p(l | a, w)}{p(l | z, a, w)} \quad (2)$$

in constructing our estimators, as such parameterizations allow for estimation and integration with respect to multivariate conditional densities on the mediator Z to be avoided. We use $\mathcal{W}, \mathcal{A}, \mathcal{Z}, \mathcal{L}$, and \mathcal{Y} to denote the support of the corresponding random variables.

Causal effects are defined in terms of hypothetical interventions on the NPSEM (1). In particular, consider an intervention in which the structural equation corresponding to A is removed, with the treatment drawn instead from a user-specified distribution $g_\delta(a | w)$, which may itself depend on the natural exposure distribution and a user-specified parameter δ . Going forward, we let A_δ denote a draw from $g_\delta(a | w)$.

Alternatively, such modifications can occasionally be described in terms of an intervention in which the structural equation corresponding to A is removed and the treatment is set equal to a hypothetical regime $d(A, W)$. Regime d depends on the treatment level A that would be assigned in the absence of the regime as well as on W . The latter intervention has been referred to as depending on the *natural value of treatment*, or as a *modified treatment policy* (Haneuse & Rotnitzky, 2013). For such interventions, Haneuse & Rotnitzky (2013) introduced the assumption of *piecewise smooth invertibility*, which ensures that the change of variable formula can be used when computing integrals over \mathcal{A} :

A1 (Piecewise smooth invertibility). For each $w \in \mathcal{W}$, assume that the interval $\mathcal{I}(w) = (l(w), u(w))$ may be partitioned into subintervals $\mathcal{I}_{\delta,j}(w) : j = 1, \dots, J(w)$ such that $d(a, w)$ is equal to some $d_j(a, w)$ in $\mathcal{I}_{\delta,j}(w)$ and $d_j(\cdot, w)$ has inverse function $h_j(\cdot, w)$ with derivative $h'_j(\cdot, w)$.

Assumption A1 can be used to show that the intervention may also be represented as a change in which the structural equation f_A is removed from the NPSEM, with A being drawn from the post-intervention distribution $g_\delta(a | w)$. Young et al. (2014) provide a discussion comparing and contrasting the interpretation and identification of these two interventions. Such stochastic interventions can be used to define the *population intervention effect* (PIE) of A on Y .

To illustrate, consider continuous-valued A and assume the distribution of A conditional on $W = w$ is supported in the interval $(l(w), u(w))$. Then, one may define

$$d(a, w) = \begin{cases} a - \delta & \text{if } a > l(w) + \delta \\ a & \text{if } a \leq l(w) + \delta, \end{cases} \quad (3)$$

where $0 < \delta < u(w)$ is an arbitrary prespecified value. We can alternatively define a tilted intervention distribution as

$$g_\delta(a | w) = \frac{\exp(\delta a)g(a | w)}{\int \exp(\delta a)g(a | w)d\kappa(a)}, \quad (4)$$

for $\delta \in \mathbb{R}$. Kennedy (2019) proposed a form of exponential tilting (4) under the parameterization $\delta' = \exp(\delta)$, appropriate for incremental interventions on the propensity score for binary A . Díaz & Hejazi (2020) provide a careful

study of the interventions 3 and 4 in the context of mediation, introducing novel (in)direct effects and corresponding efficiency theory; however, their contributions assume the absence of intermediate confounding.

2.1 Stochastic mediation effects

Díaz & Hejazi (2020) defined the (in)direct effect of A on Y in terms of a decomposition of the total effect of a stochastic intervention. In particular, the total effect $E(Y - Y_{A_\delta})$ may be decomposed as the sum of the population intervention direct effect (PIDE) and the population intervention indirect effect (PIIE):

$$\begin{aligned} \text{PIDE} &= E_c\{f_Y(Z, L, A, W, U_Y) - f_Y(Z, L_{A_\delta}, A_\delta, W, U_Y)\} \\ \text{PIIE} &= E_c\{f_Y(Z, L_{A_\delta}, A_\delta, W, U_Y) - f_Y(Z_{A_\delta}, L_{A_\delta}, A_\delta, W, U_Y)\}. \end{aligned}$$

Upon inspection, the definitions above reveal that the direct effect measures the effect through paths *not* involving the mediator (i.e., $A \rightarrow Y$ and $A \rightarrow L \rightarrow Y$), whereas the indirect effect measures the effect through paths involving the mediator (i.e., $A \rightarrow Z \rightarrow Y$ and $A \rightarrow L \rightarrow Z \rightarrow Y$).

Unfortunately, the population intervention direct and indirect effects are not generally identified in the presence of an intermediate confounder affected by treatment such as in the DAG in Figure (1) (Díaz & Hejazi, 2020). This is due to the dual role of L as a confounder of the relation between Z and Y , which requires adjustment, and a variable on the path from A to Y , which precludes adjustment. Next, we present a solution to this complication using a joint stochastic intervention on the treatment A and mediator Z . We also show that the effects defined in this paper are a generalization of the effects of Díaz & Hejazi (2020) in the sense that the former reduce to the latter in the absence of intermediate confounding.

2.2 Interventional stochastic mediation effects

To introduce (in)direct effects robust to the presence of intermediate confounders, we draw upon ideas first outlined by Didelez et al. (2006), Petersen et al. (2006), and van der Laan & Petersen (2008), all subsequently formalized by VanderWeele et al. (2014) and Vansteelandt & Daniel (2017). Owing to their definition in terms of stochastic interventions on the mediator, these (in)direct effects have been collectively termed *interventional effects*. We leverage two types of stochastic interventions: one on the treatment A , which defines the intervention of interest, and one on the mediator Z , which is used to achieve identifiability of the effects. Following the convention of the literature, we term stochastic interventions on the mediator *interventional*, while reserving the label of *stochastic* to refer only to interventions on the treatment A . To proceed, let G_δ denote a random draw from the distribution of Z_{A_δ} conditional on (A_δ, W) , and let G denote a random draw from the distribution of Z conditional on (A, W) . We consider the effect defined by $\psi_\delta = E_c\{Y_{A,G} - Y_{A_\delta,G_\delta}\}$. Note that the effect ψ_δ is distinct from the effect considered by Díaz & Hejazi (2020), which may be expressed $E_c\{Y_{A,Z} - Y_{A_\delta,Z_\delta}\}$. The effect ψ_δ arises from fixing the mediator to a random value chosen from its distribution among all those with a particular treatment level, rather than fixing it to what it would have been under a particular (fixed) treatment. Defining the effect in this way aids in achieving an identifiable decomposition into direct and indirect effects. In particular, we may decompose this effect in terms of interventional stochastic *direct effects* (DE) and *indirect effects* (IE):

$$\psi(\delta) = \overbrace{E\{Y_{A,G} - Y_{A_\delta,G}\}}^{\text{DE}} + \overbrace{E\{Y_{A_\delta,G} - Y_{A_\delta,G_\delta}\}}^{\text{IE}}. \quad (5)$$

Decomposition as the sum of direct and indirect effects affords an interpretation analogous to the corresponding standard decomposition of the average treatment effect into the natural direct and indirect effects (Pearl, 2006). In particular, the direct effect arises from drawing a counterfactual value of A from a post-intervention distribution while keeping the distribution of Z fixed. The indirect effect arises from replacing the distribution of Z with a candidate post-intervention distribution while holding A fixed. Our proposed stochastic interventional effects have an interpretation similar to the interventional effects of VanderWeele et al. (2014); moreover, while both effect definitions account for the presence of an intermediate confounder, our direct and indirect effects accommodate flexible, stochastic interventions on the exposure while those of VanderWeele et al. (2014) are limited to static interventions.

2.3 Identification

In order to construct estimators of our proposed causal direct and indirect effects, we turn to examining assumptions needed to estimate components of the post-intervention quantities matching the counterfactual variables of interest. Towards this end, we introduce the following identification assumptions:

A2 (Common support). Assume $\text{supp}\{g_\delta(\cdot | w)\} \subseteq \text{supp}\{g(\cdot | w)\}$ for all $w \in \mathcal{W}$.

A3 (No unmeasured exposure-outcome confounder). Assume $Y_{a,z} \perp\!\!\!\perp A \mid W$.

A4 (No unmeasured mediator-outcome confounder). Assume $Y_{a,z} \perp\!\!\!\perp Z \mid (L, A, W)$.

A5 (No unmeasured exposure-mediator confounder). Assume $Z_a \perp\!\!\!\perp A \mid W$.

Under these assumptions, we have the following identification results. A proof is available in the [Supplementary Materials](#).

Theorem 1 (Identification). *Define*

$$\begin{aligned}\theta_{1,\delta} &= \int m(z, l, a, w) p(l \mid a, w) p(z \mid a, w) g_\delta(a \mid w) p(w) d\nu(a, z, l, w), \\ \theta_{2,\delta} &= \int m(z, l, a, w) p(l \mid a, w) p(z \mid w) g_\delta(a \mid w) p(w) d\nu(a, z, l, w).\end{aligned}$$

Under **A2–A5**, the direct effect $\psi_{D,\delta}$ and indirect effect $\psi_{I,\delta}$ (5) are identified and given, respectively, by

$$\begin{aligned}\psi_{D,\delta} &= \theta_{1,0} - \theta_{2,\delta} \\ \psi_{I,\delta} &= \theta_{2,\delta} - \theta_{1,\delta}.\end{aligned}\tag{6}$$

Assumption **A3** states that, conditional on W , there is no unmeasured confounding of the relation between A and Y ; assumption **A5** states that conditional on W there is no unmeasured confounding of the relation between A and Z ; and assumption **A4** states that conditional on (W, A, L) there is no unmeasured confounding of the relation between Z and Y . These assumptions are standard in causal mediation analysis. In addition to these assumptions, standard mediation analyses (e.g., [VanderWeele et al., 2014](#)) require positivity assumptions on the treatment and mediation mechanisms. The stochastic intervention framework we adopt does not require such assumptions, as positivity can be arranged by definition of g_δ . For example, the interventions in expressions (3) and (4) satisfy assumption **A2** by definition. The interested reader is encouraged to consult [Kennedy \(2019\)](#) and [Díaz & Hejazi \(2020\)](#) for a discussion on this topic.

Another consequence of this identification result is that the definitions (6) reduce to the stochastic (in)direct effects of [Díaz & Hejazi \(2020\)](#) in the absence of intermediate confounders L . Importantly, this implies that our estimators can be safely used in the absence of intermediate confounders; furthermore, it implies that the corresponding estimates may be interpreted in terms of a decomposition of the population intervention effect $E_c\{Y - Y_{A_\delta}\}$, which is arguably of more scientific interest than the interventional effect $\psi_\delta = E_c\{Y_{A,G} - Y_{A_\delta,G_\delta}\}$.

As is clear from the definition (6), evaluation of $\psi_{D,\delta}$ and $\psi_{I,\delta}$ requires access to $\theta_{1,0}$, the population mean in the absence of any intervention on the treatment mechanism, as well as both of $\theta_{1,\delta}$ and $\theta_{2,\delta}$, which are based on the post-intervention treatment mechanism $g_\delta(a \mid w)$. Consequently, we next turn our attention to developing efficiency theory for estimation of the statistical parameter $\theta_{j,\delta} : j = 1, 2$, which depends on the observed data distribution P .

3 Optimality theory for estimation of the direct effect

Thus far, we have discussed the decomposition of the effect of a stochastic intervention into direct and indirect effects and have provided identification results under standard identifiability assumptions. We consider the development of efficiency theory for the estimation of $\theta_{1,\delta}$ and $\theta_{2,\delta}$ in the nonparametric model \mathcal{M} . To do so, we introduce the *efficient influence function* (EIF), which characterizes the asymptotic behavior of all regular and asymptotically linear estimators ([Bickel et al., 1993](#); [van der Vaart, 2002](#)). Three common approaches exist for constructing local efficient estimators from the EIF: (i) using the EIF as an estimating equation (e.g., [van der Laan & Robins, 2003](#)), (ii) using the EIF in a one-step bias correction (e.g., [Pfanzagl & Wefelmeyer, 1985](#); [Bickel et al., 1993](#)), and targeted minimum loss estimation ([van der Laan & Rubin, 2006](#); [van der Laan & Rose, 2011, 2018](#)).

As a consequence of its representation in terms of orthogonal score equations, the EIF estimating equation makes it possible to construct consistent estimators of the target parameter even when certain components of its distribution are inconsistently estimated. Thirdly, second-order bias terms may be derived from asymptotic analysis of estimators constructed based on the EIF — often, these estimators require slow convergence rates (e.g., $n^{-1/4}$) for the nuisance parameters involved. This latter property enables the use of flexible, data adaptive regression techniques in estimating these quantities.

For simplicity, we focus on the case of a binary intermediate confounder L , though our general approach requires only that either L or Z be low-dimensional. In [Theorem 2](#), we present the EIF for a general stochastic intervention. Although the components of the EIF associated with (Y, Z, L, W) are the same, the component associated with the model for the distribution of A must be computed on a case-by-case basis, that is, for each intervention of interest.

Lemmas 1 and 2 present such components for modified treatment policies satisfying assumption A1 and for exponential tilting, respectively. In theorem 2 below, we present a representation of the EIF that avoids the computation of multivariate integrals over Z . To introduce the EIF, we define the following auxiliary nuisance parameters:

$$\begin{aligned} u(z, a, w) &= \int m(z, l, a, w) dP(l | a, w); & \bar{u}(a, w) &= \int u(z, a, w) dP(z | a, w) \\ v(l, a, w) &= \int m(z, l, a, w) dP(z | a, w); & \bar{v}(a, w) &= \int v(l, a, w) dP(l | a, w) \\ s(l, a, w) &= \int m(z, l, a, w) dP(z | w); & \bar{s}(a, w) &= \int s(l, a, w) dP(l | a, w) \end{aligned} \quad (7)$$

Proofs for the following results are detailed in the [Supplementary Materials](#).

Theorem 2 (Efficient influence functions).

$$H_{P,\delta}^1(a, z, l, w) = \frac{g_\delta(a | w)}{g(a | w)} \frac{p(z | a, w)}{p(z | a, l, w)}; \quad H_{P,\delta}^2(a, z, l, w) = \frac{g_\delta(a | w)}{g(a | w)} \frac{p(z | w)}{p(z | a, l, w)} \quad (8)$$

The efficient influence functions for $\theta_{j,\delta} : j = 1, 2$ in the nonparametric model are equal to $D_{P,\delta}^j(o) - \theta_{j,\delta}$, where $D_{P,\delta}^j(o) = S_{P,\delta}^j(o) + S_{P,\delta}^{j,A}(o)$ and

$$S_{P,\delta}^1(o) = H_{P,\delta}^1(a, z, l, w) \{y - m(z, l, a, w)\} \quad (9)$$

$$+ \frac{g_\delta(a | w)}{g(a | w)} [v(l, a, w) - \bar{v}(a, w) + u(z, a, w) - \bar{u}(a, w)] \quad (10)$$

$$+ \int \bar{u}(a, w) g_\delta(a | w) d\kappa(a)$$

$$S_{P,\delta}^2(o) = H_{P,\delta}^2(a, z, l, w) \{y - m(z, l, a, w)\} \quad (11)$$

$$+ \frac{g_\delta(a | w)}{g(a | w)} \{s(l, a, w) - \bar{s}(a, w)\} \quad (12)$$

$$+ \int u(z, a, w) g_\delta(a | w) d\kappa(a),$$

and $S_{P,\delta}^{1,A}(o)$, $S_{P,\delta}^{2,A}(o)$ are the respective efficient score functions corresponding to the model for $g(a | w)$.

An immediate consequence of Theorem 2 is that, in a randomized trial, $S_{P,\delta}^{j,A}(o) = 0$ for $j = 1, 2$; however, even in such trials, covariate adjustment can improve the efficiency of the resultant estimator ([van der Laan & Robins, 2003](#)). We now present the efficient scores $S_{P,\delta}^{j,A}(o)$ for modified treatment policies and exponentially tilted stochastic interventions. To do so, we define the parameter $q(a, w) = \int u(z, a, w) dP(z | w)$.

Lemma 1 (Modified treatment policies). *If the modified treatment policy $d(A, W)$ satisfies assumption A1, then*

$$S_{P,\delta}^{1,A}(o) = \bar{u}(d(a, w), w) - \int \bar{u}(d(a, w), w) g_\delta(a | w) d\kappa(a)$$

$$S_{P,\delta}^{2,A}(o) = q(d(a, w), w) - \int q(d(a, w), w) g_\delta(a | w) d\kappa(a).$$

Lemma 2 (Exponential tilt). *If the stochastic intervention is the exponential tilt (4), then*

$$S_{P,\delta}^{1,A}(o) = \frac{g_\delta(a | w)}{g(a | w)} \left\{ \bar{u}(a, w) - \int \bar{u}(a, w) g_\delta(a | w) d\kappa(a) \right\} \quad (13)$$

$$S_{P,\delta}^{2,A}(o) = \frac{g_\delta(a | w)}{g(a | w)} \left\{ q(a, w) - \int q(a, w) g_\delta(a | w) d\kappa(a) \right\} \quad (14)$$

For binary treatments, the EIF corresponding to the incremental propensity score intervention may be simplified as per the following corollary.

Corollary 1 (Efficient influence function for incremental propensity score interventions). *Let A take values on $\{0, 1\}$, and let the exponentially tilted intervention $g_{\delta,0}(1 | W)$ be based on (4) under the parameterization $\delta' = \exp(\delta)$. Then, the EIF of Lemma 2 may be simplified as follows. Define the nuisance parameters*

$$\begin{aligned} q^1(w) &= \bar{u}(1, w) - \bar{u}(0, w), \\ q^2(w) &= E \{ u(Z, 1, W) - u(Z, 0, W) | W = w \}, \end{aligned} \quad (15)$$

Then

$$S_{\eta, \delta}^{j, A}(o) = \frac{\delta q^j(w) \{a - g(1 | w)\}}{\{\delta g(1 | w) + 1 - g(1 | w)\}^2}.$$

In contrast to the efficient influence function for the interventional (in)direct effects (Díaz et al., 2020), the contribution of the treatment process to the EIF for the stochastic interventional effects defined here is non-zero. This is a direct consequence of the fact that the parameter of interest depends on g ; moreover, this implies that the efficiency bound in observational studies differs from the efficiency bound in randomized studies. This further implies that it is not generally possible to obtain estimating equations that are robust to inconsistent estimation of g . Such robustness will only be possible if the stochastic intervention is also a modified treatment policy satisfying assumption A1.

The form of Theorem 2 makes it clear that estimation of multivariate or continuous conditional density functions on the mediators Z or intermediate confounders L , as well as integrals with respect to these density functions, is generally necessary for computation of the EIF. This poses a significant challenge from the perspective of estimation, due to both the curse of dimensionality and the practical computational complexity inherent in solving multivariate numerical integrals. A simplification is possible when either Z or L is low-dimensional; this is achieved by reparameterizing the densities as conditional expectations (or low-dimensional conditional densities) that take other nuisance parameters as pseudo-outcomes is possible. To demonstrate, we assume L is univariate (e.g., binary as in our illustrative application), though similar parameterizations may be achieved if Z is low-dimensional. In cases where L or Z is low-dimensional, our proposed re-parameterizations allow for the conditional density to be estimated via appropriate semiparametric estimators (e.g., Díaz & van der Laan, 2011).

Lemma 3 (Low-dimensional L and Z). *If L is low-dimensional (e.g., binary and univariate) and Z is multivariate, we can choose a representation of v , s , and \bar{u} in terms of conditional expectations in order to facilitate their estimation. Denote $b(l | a, w)$ and $d(l | z, a, w)$ the probability that $L = l \in \{0, 1\}$ conditional on (A, W) and (Z, A, W) , respectively. Then, using (2), we have*

$$\begin{aligned} v(l, a, w) &= E \left[m(z, l, a, w) \frac{b(L | A, W)}{d(L | Z, A, W)} \mid L = l, A = a, W = w \right], \\ s(l, a, w) &= E \left[m(z, l, a, w) \frac{b(L | A, W)}{d(L | Z, A, W)} \frac{g(A | W)}{e(A | Z, W)} \mid L = l, A = a, W = w \right], \\ \bar{u}(a, w) &= E \left[u(Z, A, W) \mid A = a, W = w \right]. \end{aligned} \quad (16)$$

Likewise,

$$H_{P, \delta}^1(a, z, l, w) = \frac{g_\delta(a | w)}{g(a | w)} \frac{b(l | a, w)}{d(l | z, a, w)}; \quad H_{P, \delta}^2(a, z, l, w) = \frac{g_\delta(a | w)}{e(a | z, w)} \frac{b(l | a, w)}{d(l | z, a, w)},$$

and

$$q(a, w) = E \left\{ \frac{g(A | W)}{e(A | Z, W)} u(Z, A, W) \mid A = a, W = w \right\}.$$

Analogous representations may be constructed for \bar{v} , \bar{s} , and u based on the parameterizations (2) if L is multivariate and Z is of low dimension. We note, however, that at least one of Z or L must be of small dimensionality so that its density may be estimated and integrals over its range may be computed with ease.

In what follows, we assume L is univariate, denote $\eta = (m, g, b, \bar{u}, v, d, e, s, q)$ and let $D_{P, \delta}^j(o) = D_{\eta, \delta}^j(o)$. The choice of parameterization in Lemma 3 has important consequences for the purpose of estimation, as it helps to bypass estimation of the (possibly high-dimensional) conditional density of the mediators, instead allowing for regression methods, far more readily available throughout the statistics literature and software, to be used for estimation of the relevant quantities. Similar ideas have been used by Zheng & van der Laan (2017), Díaz & Hejazi (2020), and Díaz et al. (2020). In addition to the expression for the efficient influence function in Lemma 3, it is important to understand the behavior of the difference $PD_{\eta_1} - \theta$, which is expected to yield a second order term in differences $\eta_1 - \eta$, so that consistent estimation of θ is possible under consistent estimation of certain configurations of the parameters in η . As we will see in Theorems 3 and 4, this second-order term is fundamental in the construction of asymptotically linear estimators. Lemmas 5 and 6, found in the Supplementary Materials, delineate these second-order terms. The following lemma is a direct consequence.

Lemma 4 (Multiple robustness for modified treatment policies). *Let the modified treatment policy satisfy A1, and let η_1 be such that one of the following conditions hold:*

	$m_1 = m$	$g_1 = g$	$b_1 = b$	$\bar{u}_1 = \bar{u}$	$v_1 = v$	$d_1 = d$	$e_1 = e$	$s_1 = s$	$q_1 = q$
Cond. 1	×	×	×						
Cond. 2	×	×			×			×	
Cond. 3		×	×			×	×		
Cond. 4		×		×	×	×	×		
Cond. 5	×		×	×					×
Cond. 6	×			×	×			×	×

Table 1: Different configurations of consistency for nuisance parameters

Then $PD_{\eta_1, \delta}^1 = \theta_{1, \delta}$ and $PD_{\eta_1, \delta}^2 = \theta_{2, \delta}$, with $D_{\eta, \delta}^1$ and $D_{\eta, \delta}^2$ as defined in Theorem 2 and Lemma 1.

The above lemma implies that it is possible to construct consistent estimators for the (in)direct effects under consistent estimation of subsets of the nuisance parameters in η , in the configurations described in the lemma. Lemma 4 follows directly from Lemma 5, found in the [Supplementary Materials](#). Some readers may find it surprising that estimation of $\theta_{j, \delta}$ may be robust to inconsistent estimation of g , even when the parameter definitions are explicitly dependent on g . We offer some intuition into this result by noting that assumption A1 allows use of the change of variable formula to obtain

$$\theta_{2, \delta} = E \left\{ \int m(z, l, d(A, W), W) p(l | d(A, W), W) p(z | W) d\nu(z, l) \right\}.$$

Estimation of this parameter without relying on g may be carried out by consistently estimating $m(z, l, a, w)$, $p(l | a, w)$, and $p(z | w)$ and using the empirical distribution as an estimator of the outer expectation. This behavior has been previously observed for related stochastic intervention effects under assumption A1 (Díaz & van der Laan, 2012; Haneuse & Rotnitzky, 2013; Díaz & Hejazi, 2020).

The robustness result for the case an exponentially tilted intervention (1), which does not satisfy assumption A1, is presented in the following lemma

Lemma 5 (Multiple robustness for exponential tilting). *Let g_δ be defined as in (4). Let η_1 be such that at least one of Cond. 1-4 in Table 1 holds. Then $PD_{\eta_1, \delta}^1 = \theta_{1, \delta}$ and $PD_{\eta_1, \delta}^2 = \theta_{2, \delta}$, with $D_{\eta, \delta}^1$ and $D_{\eta, \delta}^2$ as defined in Theorem 2 and Lemma 2*

Lemma 5 is a direct consequence of Lemma 6 in the [Supplementary Materials](#). The corresponding proof reveals that the EIF for the binary distribution is not robust to inconsistent estimation of g — that is, the intervention fails to satisfy assumption A1 and integrals over the range of A cannot be computed using the change of variable formula. This behavior has been previously observed for other interventions that do not satisfy assumption A1 (e.g., Díaz & van der Laan, 2013). Even though this lemma implies that consistent estimation of g is required, the bias terms remain second-order; thus, an estimator of g converging at rate $n^{1/4}$ or faster is sufficient.

4 Efficient estimation and statistical inference

We discuss two efficient estimators that rely on the efficient influence function $D_{\eta, \delta}$, in order to build an estimator that is both efficient and robust to model misspecification. We discuss an asymptotic linearity result for the doubly robust estimator that allows computation of asymptotically correct confidence intervals and hypothesis tests. In the sequel, we assume that preliminary estimators of the components of η are available. These estimators may be obtained from flexible regression techniques such as support vector machines, regression trees, boosting, neural networks, splines, or ensembles thereof (Wolpert, 1992; Breiman, 1996; van der Laan et al., 2007). As previously discussed, the consistency of these estimators will determine the consistency of our estimators of $\theta_{j, \delta}$.

Both of our proposed efficient estimators make use of the EIF $D_{\eta, \delta}$ to revise an initial substitution estimator through a bias correction step. As such, estimation proceeds by first constructing initial estimators of the nuisance parameters in η ; then, each of the efficient estimators is constructed by application of distinct bias-correction steps. In constructing these efficient estimators, we advocate for the use of cross-fitting (Klaassen, 1987; Zheng & van der Laan, 2011; Chernozhukov et al., 2018) in order to avoid imposing entropy conditions on the initial estimators of the nuisance parameters in η . Let $\mathcal{V}_1, \dots, \mathcal{V}_J$ denote a random partition of the index set $\{1, \dots, n\}$ into J prediction sets of

approximately the same size. That is, $\mathcal{V}_j \subset \{1, \dots, n\}$; $\bigcup_{j=1}^J \mathcal{V}_j = \{1, \dots, n\}$; and $\mathcal{V}_j \cap \mathcal{V}_{j'} = \emptyset$. In addition, for each j , the associated training sample is given by $\mathcal{T}_j = \{1, \dots, n\} \setminus \mathcal{V}_j$. Denote by $\hat{\eta}_j$ the estimator of η , obtained by training the corresponding prediction algorithm using only data in the sample \mathcal{T}_j . Further, let $j(i)$ denote the index of the validation set which contains observation i .

4.1 Efficient one-step estimator

To construct a robust and efficient estimator using the efficient influence function $D_{\eta, \delta}$, the one-step bias correction (Pfanzagl & Wefelmeyer, 1985; Bickel et al., 1993) adds the empirical mean of the estimated EIF $D_{\hat{\eta}, \delta}$ to an initial substitution estimator. The estimators are thus defined

$$\begin{aligned}\hat{\psi}_{D, \delta}^{\text{os}} &= \frac{1}{n} \sum_{i=1}^n \{D_{\hat{\eta}_{j(i)}, 0}^1(O_i) - D_{\hat{\eta}_{j(i)}, \delta}^2(O_i)\} \\ \hat{\psi}_{I, \delta}^{\text{os}} &= \frac{1}{n} \sum_{i=1}^n \{D_{\hat{\eta}_{j(i)}, \delta}^2(O_i) - D_{\hat{\eta}_{j(i)}, \delta}^1(O_i)\}\end{aligned}\tag{17}$$

Asymptotic linearity and efficiency of the estimators for modified treatment policies is detailed below.

Theorem 3 (Weak convergence of one-step estimators). *Let $\|\cdot\|$ denote the $L_2(\mathbb{P})$ -norm defined as $\|f\|^2 = \int f^2 d\mathbb{P}$. Define the following assumptions.*

- (i) $\mathbb{P}\{|D_{\eta, \delta}^j(O)| \leq C\} = \mathbb{P}\{|D_{\hat{\eta}, \delta}^j(O)| \leq C\} = 1$ for $j = 1, 2$ and for some $C < \infty$.
- (ii) *The following second-order terms converge at the specified rate*
 - $\|\hat{m} - m\| \{\|\hat{g} - g\| + \|\hat{e} - e\| + \|\hat{d} - d\|\} = o_{\mathbb{P}}(n^{-1/2})$
 - $\|\hat{g} - g\| \{\|\hat{u} - u\| + \|\hat{q} - q\|\} = o_{\mathbb{P}}(n^{-1/2})$
 - $\|\hat{b} - b\| \{\|\hat{v} - v\| + \|\hat{s} - s\|\} = o_{\mathbb{P}}(n^{-1/2})$, and
- (iii) *The effect is defined in terms of modified treatment policy $d(a, w)$, which is piecewise smooth invertible (A1).*
- (iv) *The intervention g_δ is an exponential tilting intervention and $\mathbb{P}\{\int (\hat{g} - g) d\kappa\}^2 = o_{\mathbb{P}}(n^{-1/2})$.*

If assumptions (i) and (ii) hold, and one of assumptions (iii) and (iv) holds, then:

$$\begin{aligned}\sqrt{n}\{\hat{\psi}_{D, \delta}^{\text{os}} - \psi_{D, \delta}\} &\rightsquigarrow N(0, \sigma_{D, \delta}^2), \text{ and} \\ \sqrt{n}\{\hat{\psi}_{I, \delta}^{\text{os}} - \psi_{I, \delta}\} &\rightsquigarrow N(0, \sigma_{I, \delta}^2),\end{aligned}$$

where $\sigma_{D, \delta}^2 = \text{Var}\{D_{\eta, 0}^1(O) - D_{\eta, \delta}^2(O)\}$ and $\sigma_{I, \delta}^2 = \text{Var}\{D_{\eta, \delta}^2(O) - D_{\eta, \delta}^1(O)\}$ are the respective efficiency bounds.

Theorem 3 establishes the weak convergence of $\hat{\psi}_{D, \delta}^{\text{os}}$ and $\hat{\psi}_{I, \delta}^{\text{os}}$ pointwise in δ . This convergence is useful to derive confidence intervals in situations where the modified treatment policy has a suitable scientific interpretation for a given realization of δ . Under the assumptions of the theorem, an estimator $\hat{\sigma}_{D, \delta}^2$ of $\sigma_{D, \delta}^2$ may be obtained as the empirical variance of $D_{\hat{\eta}_{j(i)}, 0}^1(O_i) - D_{\hat{\eta}_{j(i)}, \delta}^2(O_i)$, and a Wald-type confidence interval may be constructed as $\hat{\psi}_{D, \delta}^{\text{os}} \pm z_{1-\alpha/2} \hat{\sigma}_{D, \delta}^2(\delta) / \sqrt{n}$. Similar considerations apply to $\hat{\psi}_{I, \delta}^{\text{os}}$.

Although the one-step estimator has optimal asymptotic performance, its finite-sample behavior may be affected by the inverse probability weighting involved in the computation of the efficient influence functions $D_{\hat{\eta}}^j(O_i) : j = 1, 2$. In particular, it is not guaranteed that $\hat{\psi}_{D, \delta}^{\text{os}}$ and $\hat{\psi}_{I, \delta}^{\text{os}}$ will remain within the bounds of the parameter space. This issue may be attenuated by performing weight stabilization. The estimated EIF $D_{\hat{\eta}_{j(i)}}^1(O_i)$ can be weight-stabilized by dividing (9) and (11) by the empirical mean of $H_{\hat{\eta}_{j(i)}, \delta}^1(A_i, Z_i, L_i, W_i)$ and $H_{\hat{\eta}_{j(i)}, \delta}^2(A_i, Z_i, L_i, W_i)$, respectively; as well as dividing (10), (12), (13), and (14) by the empirical mean of $\hat{g}_{j(i), \delta}(A_i | W_i) / \hat{g}_{j(i)}(A_i | W_i)$.

4.2 Efficient targeted minimum loss estimator

Although corrections may be applied to the one-step estimator, a more principled way to obtain estimators that remain in the parameter space may be derived from the targeted minimum loss (TML) estimation framework. The TML estimator is constructed by tilting an initial data adaptive estimator $\hat{\eta}$ towards a solution $\tilde{\eta}$ of the estimating equations

$$\begin{aligned} P_n\{D_{\eta,0}^1 - D_{\eta,\delta}^2\} &= \psi_{D,\delta}(\tilde{\eta}) \\ P_n\{D_{\eta,\delta}^2 - D_{\eta,\delta}^1\} &= \psi_{I,\delta}(\tilde{\eta}), \end{aligned} \quad (18)$$

where $\psi_{D,\delta}(\tilde{\eta})$ and $\psi_{I,\delta}(\tilde{\eta})$ are the substitution estimators in formula (19) obtained by plugging in the estimates $\tilde{\eta}$ in the parameter definition (6). Thus, a TML estimator is guaranteed to remain in the parameter space by virtue of its being a substitution estimator. The fact that the nuisance estimators solve the relevant estimating equation is used to obtain a weak convergence result analogous to Theorem 3. Thus, while the TML estimator is expected to attain the same optimal asymptotic behavior as the one-step estimator, its finite-sample behavior may be better. An algorithm to compute a TML estimator $\tilde{\eta}$ is presented in the [Supplementary Materials](#). Roughly, the algorithm proceeds by projecting the EIF into score functions for the model of each nuisance parameter, and fitting appropriate parametric submodels (van der Laan & Rose, 2011, 2018). For example, the following model is fitted for m :

$$\text{logit } m_\beta(a, z, l, w) = \text{logit } \hat{m}(z, l, a, w) + \beta_I H_I(o) + \beta_D H_D(o),$$

where

$$\begin{aligned} H_D(o) &= \frac{\hat{b}(l | a, w)}{\hat{d}(l | z, a, w)} \left\{ 1 - \frac{\hat{g}_\delta(a | w)}{\hat{e}(a | z, w)} \right\} \\ H_I(o) &= \frac{\hat{b}(l | a, w)}{\hat{d}(l | z, a, w)} \left\{ \frac{\hat{g}_\delta(a | w)}{\hat{e}(a | z, w)} - \frac{\hat{g}_\delta(a | w)}{\hat{g}(a | w)} \right\}, \end{aligned}$$

and $\text{logit}(p) = \log\{p(1-p)^{-1}\}$. Here, the initial estimator $\text{logit } \hat{m}(z, l, a, w)$ is considered a fixed offset variable (i.e., a variable with known parameter value equal to one). The score of these tilting models is equal to the corresponding component of the efficient influence function. The parameter $\beta = (\beta_I, \beta_D)$ may be estimated by running standard logistic regression of Y on $(H_D(O), H_I(O))$ with no intercept and an offset term equal to $\text{logit } \hat{m}(z, l, a, w)$. Let $\hat{\beta}$ denote the MLE, and let $\tilde{m} = m_{\hat{\beta}}$ denote the updated estimates. Fitting this model ensures that \tilde{m} solves the relevant score equations. Models like this are estimated iteratively for all parameters in a way that guarantees that the estimating equations (18) are solved up to a term that converges to zero in probability at rate faster than $n^{-1/2}$. After the iteration ends, the TML estimators are defined as

$$\begin{aligned} \hat{\psi}_{D,\delta}^{\text{tmle}} &= \frac{1}{n} \int \sum_{i=1}^n \{ \tilde{u}(a, W_i) \tilde{g}(a | W_i) - \tilde{u}(Z_i, a, W_i) \tilde{g}_\delta(a | W_i) \} d\kappa(a) \\ \hat{\psi}_{I,\delta}^{\text{tmle}} &= \frac{1}{n} \int \sum_{i=1}^n \{ \tilde{u}(Z_i, a, W_i) - \tilde{u}(a, W_i) \} \tilde{g}_\delta(a | W_i) d\kappa(a). \end{aligned} \quad (19)$$

The fact that the TML estimator solves estimating equations (18) is a fundamental step in proving the following theorem.

Theorem 4 (Weak convergence of TML estimator). *Assume (i) and (ii) hold, and one of (iii), (iv) defined in Theorem 3 holds, then:*

$$\begin{aligned} \sqrt{n}\{\hat{\psi}_{D,\delta}^{\text{tmle}} - \psi_{D,\delta}\} &\rightsquigarrow N(0, \sigma_{D,\delta}^2), \text{ and} \\ \sqrt{n}\{\hat{\psi}_{I,\delta}^{\text{tmle}} - \psi_{I,\delta}\} &\rightsquigarrow N(0, \sigma_{I,\delta}^2), \end{aligned}$$

where $\sigma_{D,\delta}^2 = \text{Var}\{D_{\eta,0}^1(O) - D_{\eta,\delta}^2(O)\}$ and $\sigma_{I,\delta}^2 = \text{Var}\{D_{\eta,\delta}^2(O) - D_{\eta,\delta}^1(O)\}$.

Using Theorem 4, asymptotically valid variance estimators, p-values, and confidence intervals for the (in)direct effects may be obtained in a manner analogous to those for the one-step estimator. The proof of the theorem proceeds using similar arguments as the proof of Theorem 3 for the one-step estimator, using empirical process theory and leveraging cross-fitting to avoid entropy conditions on the initial estimators of η . Since the estimators now depend on the full sample through the estimates of the parameters β of the logistic tilting models, the empirical process treatment is slightly different from that of Theorem 3. The proof is detailed in the [Supplementary Materials](#).

5 Simulation study

We used simulation experiments to assess our two proposed efficient estimators of the (in)direct effects. On account of computational considerations, we focus on binary exposures and intermediate confounders in this example; however, as noted in the prior, our proposed methodology is general enough to be readily applicable in the presence of continuous-valued covariates, treatment, mediators, intermediate confounders, and outcome. We made use of the following data-generating mechanism for the joint distribution of O to generate synthetic data for evaluation of the two estimators:

$$\begin{aligned}
 W_1 &\sim \text{Bernoulli}(p = 0.6); W_2 \sim \text{Bernoulli}(p = 0.3); \\
 W_3 \mid (W_1, W_2) &\sim \text{Bernoulli}(p = 0.2 + 1/3 \cdot (W_1 + W_2)); \\
 A \mid W &\sim \text{Bernoulli}(p = \text{expit}(2 + (5/(W_1 + W_2 + W_3)))); \\
 L \mid (A, W) &\sim \text{Bernoulli}(p = \text{expit}(1/3(W_1 + W_2 + W_3) - A - \log(2) + 0.2)); \\
 Z \mid (L, A, W) &\sim \text{Bernoulli}(p = \text{expit}(\log(3) \cdot (W_1 + W_2) + A - L)); \\
 Y \mid (Z, L, A, W) &\sim \text{Bernoulli}\left(p = \text{expit}\left(1 - \frac{3 \cdot (3 - L - 3A + Z)}{2 + (W_1 + W_2 + W_3)}\right)\right),
 \end{aligned}$$

where $\text{expit}(x) := \{1 + \exp(x)\}^{-1}$. For each of the sample sizes $n \in \{200, 800, 1800, 3200, 5000, 7200, 9800, 12800, 16200\}$, 500 datasets were generated. For every dataset, six variations of each of the two efficient estimators was applied — five variants were based on misspecification of a single nuisance parameter among $\{e, m, d, g, b\}$ while the sixth variant was constructed based on consistent estimation of all five nuisance parameters. An intercept-only logistic regression model provided inconsistent estimation of each of the nuisance parameters $\{e, m, d, g, b\}$, while a Super Learner ensemble (van der Laan et al., 2007) was used to achieve consistent estimation. The Super Learner ensemble was constructed with a library of algorithms composed of intercept-only logistic regression; main-terms logistic regression; and several variants of the highly adaptive lasso (Benkeser & van der Laan, 2016; van der Laan, 2017; Coyle et al., 2020), a nonparametric regression approach capable of flexibly estimating arbitrary functional forms at a fast convergence rate under only a global smoothness assumption (van der Laan & Bibaut, 2017; Bibaut & van der Laan, 2019). Note that we do not consider cases of misspecified estimation of $\{v, s, q, \bar{u}\}$, for their consistent estimation depends on consistent estimation of a subset of the nuisance parameters $\{e, m, d, g, b\}$. Generally, based on Lemmas 4 and 5, robustness of the direct and indirect effect estimators to misspecification of $\{e, m, d\}$ is to be expected, but the same is not true under misspecification of $\{g, b\}$.

Figure 2 summarizes the results of our investigations of the relative performance of the estimator variants enumerated above. Specifically, we assess the relative performance of our proposed estimators in terms of absolute bias, scaled (by $n^{1/2}$) bias, standard error and scaled (by n) mean squared error relative to the efficiency bound for the data-generating model, the empirical coverage of 95% confidence intervals, and relative efficiency. In terms of both raw (unscaled) bias and scaled bias, the estimator variants appear to conform to the predictions of Lemmas 4 and 5 — specifically, raw bias vanishes and scaled bias stabilizes to a small value (providing evidence for rate-consistency) under misspecification of any of $\{e, m, d\}$ as well as in the case of no nuisance parameter misspecification. In the same vein, when either of $\{g, b\}$ are estimated inconsistently, some of the estimator variants display diverging asymptotic (scaled) bias, in agreement with expectations based upon theory. The consistency of other estimator variants (e.g., the one-step estimator under misspecification of b) is likely an artifact of this data-generating mechanism, not to be taken as a general indication of robust performance. In terms of their relative mean squared error, the estimators of the (in)direct effects exhibit convergence to the efficiency bound under misspecification of $\{e, m, d\}$ and under no misspecification; this also appears to hold for a subset of the estimator variants under misspecification of $\{g, b\}$. We stress that aspects of this are likely to be a particularity of the given data-generating mechanism or on account of the irregularity of misspecified estimator variants, for the regularity and asymptotically linearity of the estimators is only to be expected under consistent estimation of all nuisance parameters. Finally, the empirical coverage of 95% confidence intervals is as expected: under a lack of nuisance parameter misspecification, both the one-step and TML estimators of the direct and indirect effect achieve 95% coverage in larger sample sizes. We note that misspecification of e leads to over-coverage for all estimator variants, implying an overly inflated variance estimate, while the confidence intervals fail to attain the nominal rate in most other instances. Notably, several of the estimator variants generate confidence intervals that are liable to converge to 0% coverage in larger samples under misspecification of $\{g, b\}$, very much in line with theoretical expectations.




Figure 2: Comparison of efficient estimators across different nuisance parameter configurations.

Importantly, the TML estimator appears to generally outperform the one-step estimator throughout several scenarios. This comes in several forms, including lower bias, relative standard deviation, or relative mean squared error under misspecification of $\{e, m, d\}$ or under no misspecification; however, under inconsistent estimation of $\{g, b\}$, the irregularity of the estimators complicates this comparison. Interestingly, under misspecification of g , the TML estimators of the direct and indirect effects appear unbiased and efficient, a result unpredictable from theory given the irregularity of the estimators under this configuration. Altogether, results of our numerical experiments indicate that our proposed estimators exhibit properties that align with the theoretical results of Lemmas 4 and 5.

6 Application to the X:BOT Trial

We now consider the application of our proposed stochastic interventional direct and indirect effects to decompose the causal effect of a strategy where buprenorphine dose is successively increased early in treatment (regardless of opioid use) on relapse among those with opioid use disorder (OUD). Data for our illustrative analysis come from the X:BOT trial, a 24-week, multi-site randomized controlled trial designed to examine the comparative effectiveness of extended-release naltrexone (XR-NTX) and sublingual buprenorphine-naloxone (BUP-NX) on relapse (Lee et al., 2018, 2016; Nunes et al., 2016). The X:BOT trial enrolled 570 participants, all of whom were 18 years or older, had OUD (as per the Diagnostic and Statistical Manual of Mental Disorders-5; American Psychiatric Association, 2013), and had used non-prescribed opioids in the 30 days preceding enrollment. Participants were randomized to receive either XR-NTX or BUP-NX using a stratified permuted block design; 287 of the 570 were randomized to receive

BUP-NX. Prior analytic efforts have established a protective effect of BUP-NX administration (versus placebo) on OUD relapse (Mattick et al., 2014). For each participant assigned to receive BUP-NX, the prescribed dose was based on both clinical indication (Lee et al., 2018) and clinician judgment. Some clinicians tended to hold dose constant over time (i.e., a static dose regimen), while others increased dose — either based on clinical assessment or based on the hypothesis that higher doses would result in better outcomes (Nutt, 2015; Greenwald et al., 2003; Comer et al., 2005; Heikman et al., 2017). In this analysis, we estimated hypothetical stochastic interventional direct and indirect effects to assess the mechanism by which universally ramping up BUP-NX dose early in treatment (defined as three or more dose increases in the first four weeks of treatment) could mitigate the risk of OUD relapse.

Baseline covariates (W) available in the data included site; gender; age; race/ethnicity; homeless status; educational attainment; employment status; marital status; current intravenous drug use; alcohol use disorder; cocaine use disorder; age at start of heroin use; severity of current opioid use; indicator of prior OUD treatment; past withdrawal discomfort level; histories of amphetamine use, sedative use, and cannabis use; weekly cost of primary drug; whether or not living with an individual currently using drugs or with alcohol use disorder; histories of psychiatric illnesses; randomization timing; baseline pain level; baseline depression symptoms. The exposure (A) was taken to be successive increases in dose of BUP-NX versus static dose, measured during the first four weeks of treatment. Mediating factors (Z) included depression and pain, measured from week 6 until relapse or week 24 (end of follow-up). Abstinence from illicit opioid use early in the treatment schedule, measured between weeks 4 and 6, acted as an intermediate confounder affected by exposure (L). OUD relapse status at the X:BOT trial's end of follow-up was the outcome of interest (Y). To examine the effect of exposure to successive increases in BUP-NX dose, we consider an incremental propensity score intervention, which, for binary A , replaces the propensity score $g(1 | w)$ with a shifted variant constructed from multiplying the odds of treatment by a user-specified parameter δ (Kennedy, 2019). Specifically, we considered a grid $\log(\delta) \in \{-10.0, -9.5, -9.0, \dots, 9.5, 10.0\}$ of the exposure odds observed in the X:BOT trial. Across all such estimates in the odds δ of exposure, the stochastic interventional direct and indirect effects that we estimated may be interpreted in terms of the overall effect of increasingly encouraging ramping up BUP-NX dose early in treatment on the counterfactual risk of OUD relapse; thus, the results of our analysis may be informative of the mechanisms by which increasing BUP-NX dose can alter the risk of OUD relapse. The direct and indirect effect estimates, across the grid in δ , are presented in Figure 3.




Figure 3: Stochastic interventional direct (upper panel) and indirect (lower panel) effect estimates of a hypothetical intervention increasing the odds of exposure to a BUP-NX dose schedule in which dose is successively increased early in treatment on OUD relapse across a grid of shifts δ in the odds.

We applied both of our cross-fitted, efficient one-step and TML estimators to examine the stochastic interventional direct and indirect effects of increasing the odds of ramping up BUP-NX dose. Both estimation strategies produced results that were generally in very close agreement as to the magnitude of the direct and indirect effects. For each point estimate, standard error estimates and 95% Wald-style confidence intervals were constructed based on the conclusions of Theorem 4. In order to ensure the flexibility of our estimators, each component of the vector of nuisance parameters $\eta = (e, m, d, g, b)$ was estimated via ensemble machine learning, using the Super Learner algorithm (van der Laan et al., 2007). The library of machine learning algorithms from which the Super Learner ensemble was constructed included intercept-only logistic regression, logistic regression with Bayesian priors on parameters, multi-variate adaptive regression splines (Friedman, 1991), generalized additive models (Hastie & Tibshirani, 1990), random forests (Breiman, 2001), gradient boosted machines (Friedman, 2001), and the highly adaptive lasso (Benkeser & van der Laan, 2016; Coyle et al., 2020).

From examination of the point estimates and confidence intervals of the direct and indirect effects in Figure 3, two conclusions may be drawn. Firstly, there appears to be little to no indirect effect of successively increasing BUP-NX dose on risk of OUD relapse, revealing that any effect of BUP-NX dose does not appear to operate through mediating factors such as depression or pain. Secondly, the direct effect of successively increasing BUP-NX dose varies considerably across changes in the odds of the introduction of such a dose schedule. Importantly, it appears that decreasing the odds of increasing dose could lead to as much as a 5% increase in the OUD relapse risk, with a plateau emerging at odds lower than $\approx 0.1\%$, suggesting that static dose can lead to increased relapse risk relative to successive dose increases. Continuing this pattern, OUD relapse risk appears to decrease by $\approx 10\%$ with increasing odds of successively increasing BUP-NX dose, with the risk plateauing at odds higher than 33%. This decrease in the counterfactual risk of OUD relapse suggests a protective effect of BUP-NX dose schedules where dose is successively increased early in treatment relative to static dose.

The conclusions that may be drawn from our re-analysis using the stochastic interventional direct and indirect effects complement those previously reported in the investigations of Lee et al. (2018), who evaluated the total effect of BUP-NX (versus XR-NTX) treatment on OUD relapse, and Rudolph et al. (2020a), who used the interventional mediation analysis approach of Díaz et al. (2020) (limited to static interventions on A) to examine differences in relapse risk between homeless and non-homeless participants. Importantly, our substantive conclusion — that dose increases directly lower the risk of relapse — agree generally with those of Rudolph et al. (2020b), who found that dose increases directly lowered risk of OUD relapse when such dose increases followed opioid use. Notably, our proposed (in)direct effects and estimation approach differ from previous efforts in three important ways: (i) our causal effect definitions remain unaltered in the presence intermediate confounders affected by exposure and may be re-evaluated in randomized trials, (ii) the flexible estimators we introduce eschew restrictive modeling assumptions by incorporating state-of-the-art machine learning in the estimation of nuisance parameters, and (iii) our strategy provides an analog to a dose-response analysis by allowing for the risk of OUD relapse to be traced out across changes in the odds of exposure to a schedule in which BUP-NX dose is increased repeatedly early in treatment.

7 Discussion

We have proposed a class of novel direct and indirect effects for causal mediation analysis, as well as two efficient estimators of these effects in the nonparametric statistical model. Importantly, our proposed estimation framework allows for data adaptive estimation of nuisance parameters, while still preserving the benefits associated with similar classical techniques — that is, our estimators are regular and asymptotically linear, provide unbiased point estimates, are multiply robust, allow the construction of asymptotically valid confidence intervals, and are capable of attaining the nonparametric efficiency bound. Our (in)direct effects have interpretations that echo those of the classical natural (in)direct effects; however, our effects remain well-defined even in the presence of intermediate confounders affected by exposure. Further, any scientific conclusions drawn based upon our proposed (in)direct effects may be readily interrogated in trials that randomize both the exposure and mediators. Such flexible effect definitions and estimators seem necessary both to cope with the design complexity exhibited by modern epidemiological and biomedical studies and to take appropriate advantage of the ever-growing number of data adaptive regression techniques.

The challenge of leveraging data adaptive regression methodology to construct robust estimators that accommodate valid statistical inference is not a new one. It has been considered in great detail as early as the work of Pfanzagl & Wefelmeyer (1985) as well in numerous recent advances, most notably by van der Laan & Rose (2011, 2018) and Chernozhukov et al. (2018); related work by these authors presents a wealth of extensions and applications. In the present work, we derive multiply robust, efficient estimators based on both the one-step and targeted minimum loss estimation frameworks. Following Klaassen (1987) and Zheng & van der Laan (2011), our estimators leverage cross-validation to avoid imposing possibly restrictive assumptions on nuisance function estimators. We demonstrated the properties of our estimators in simulation experiments that illustrated their ability to yield unbiased point estimates,

attain the nonparametric efficiency bound, and build confidence interval covering at the nominal rate across several nuisance parameter configurations, all within the context of a problem in which classical mediation effects are ill-defined. Finally, we demonstrated the application of our novel (in)direct effects in dissecting the mechanism by which increasing the odds of adopting a dose schedule corresponding to universal successive increases in buprenorphine dose early in treatment affects relapse to OUD (Lee et al., 2018; Rudolph et al., 2020a).

Several significant extensions and refinements are left for future consideration. Firstly, our proposed estimation strategy for the direct and indirect effects leverages re-parameterizations of factors of the likelihood in order to simplify the estimation of nuisance parameters. This approach works particularly well when either mediators or intermediate confounders are of low dimension; however, improving this approach to accommodate moderate dimensionality of both mediators and intermediate confounders would surely widen the range of scenarios to which the methodology may be applied. Secondly, when defining effects based upon stochastic interventions indexed by the user-specified parameter δ , an important consideration is choosing *a priori* a particular value of δ . One solution is to evaluate a set of causal effects indexed by a grid in δ . In such cases, aggregate effects (across δ) may be summarized via working marginal structural models (e.g., Hejazi et al., 2020) or the construction of uniform tests of the null hypothesis of no direct effect (e.g., Díaz & Hejazi, 2020). Developments of these distinct summarization strategies would enrich the range of scientific applications on which these robust and flexible (in)direct effects could be brought to bear.

Acknowledgements

The authors thank John Rotrosen, Edward Nunes, and Marc Fishman for raising the research question in the Application and for helpful feedback. KER’s time was supported by a grant from the National Institute on Drug Abuse (award number R00-DA042127), while MJvdL’s time was supported by a grant from the National Institute of Allergy and Infectious Diseases (award number R01-AI074345). The X:BOT trial was supported by the National Institute on Drug Abuse, Clinical Trials Network (award numbers U10DA013046, UG1/U10DA013035, UG1/U10DA013034, U10DA013045, UG1/U10DA013720, UG1/U10DA013732, UG1/U10DA013714, UG1/U10DA015831, U10DA015833, HHSN271201200017C, and HHSN271201500065C).

References

- AMERICAN PSYCHIATRIC ASSOCIATION (2013). *Diagnostic and statistical manual of mental disorders (DSM-5®)*. American Psychiatric Pub.
- AVIN, C., SHPITSER, I. & PEARL, J. (2005). Identifiability of path-specific effects. In *IJCAI International Joint Conference on Artificial Intelligence*.
- BARON, R. M. & KENNY, D. A. (1986). The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology* **51**, 1173.
- BENKESER, D. (2020). Nonparametric inference for interventional effects with multiple mediators .
- BENKESER, D. & VAN DER LAAN, M. J. (2016). The highly adaptive lasso estimator. In *2016 IEEE International Conference on Data Science and Advanced Analytics (DSAA)*. IEEE.
- BIBAUT, A. F. & VAN DER LAAN, M. J. (2019). Fast rates for empirical risk minimization over càdlàg functions with bounded sectional variation norm. *arXiv preprint arXiv:1907.09244* .
- BICKEL, P. J., KLAASSEN, C. A., RITOV, Y. & WELLNER, J. A. (1993). *Efficient and adaptive estimation for semiparametric models*. Johns Hopkins University Press Baltimore.
- BREIMAN, L. (1996). Stacked regressions. *Machine Learning* **24**, 49–64.
- BREIMAN, L. (2001). Random forests. *Machine learning* **45**, 5–32.
- CHERNOZHUKOV, V., CHETVERIKOV, D., DEMIRER, M., DUFLO, E., HANSEN, C., NEWWEY, W. & ROBINS, J. M. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal* **21**, C1–C68.

- COLE, S. R. & HERNÁN, M. A. (2002). Fallibility in estimating direct effects. *International Journal of Epidemiology* **31**, 163–165.
- COMER, S. D., WALKER, E. A. & COLLINS, E. D. (2005). Buprenorphine/naloxone reduces the reinforcing and subjective effects of heroin in heroin-dependent volunteers. *Psychopharmacology* **181**, 664–675.
- COREY, L., GILBERT, P. B., TOMARAS, G. D., HAYNES, B. F., PANTALEO, G. & FAUCI, A. S. (2015). Immune correlates of vaccine protection against HIV-1 acquisition. *Science Translational Medicine* **7**, 310rv7–310rv7.
- COYLE, J. R., HEJAZI, N. S. & VAN DER LAAN, M. J. (2020). *hal9001: The scalable highly adaptive lasso*. R package version 0.2.6.
- DAWID, A. P. (2000). Causal inference without counterfactuals. *Journal of the American Statistical Association* **95**, 407–424.
- DÍAZ, I. & HEJAZI, N. S. (2020). Causal mediation analysis for stochastic interventions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **82**, 661–683.
- DÍAZ, I., HEJAZI, N. S., RUDOLPH, K. E. & VAN DER LAAN, M. J. (2020). Non-parametric efficient causal mediation with intermediate confounders .
- DÍAZ, I. & VAN DER LAAN, M. J. (2011). Super learner based conditional density estimation with application to marginal structural models. *The International Journal of Biostatistics* **7**, 1–20.
- DÍAZ, I. & VAN DER LAAN, M. J. (2012). Population intervention causal effects based on stochastic interventions. *Biometrics* **68**, 541–549.
- DÍAZ, I. & VAN DER LAAN, M. J. (2013). Assessing the causal effect of policies: an example using stochastic interventions. *The International Journal of Biostatistics* **9**, 161–174.
- DIDELEZ, V., DAWID, P. & GENELETTI, S. (2006). Direct and indirect effects of sequential treatments. In *Proceedings of the 22nd Annual Conference on Uncertainty in Artificial Intelligence*.
- DUDÍK, M., ERHAN, D., LANGFORD, J. & LI, L. (2014). Doubly robust policy evaluation and optimization. *Statistical Science* **29**, 485–511.
- FERGUSON, K. K., CHEN, Y.-H., VANDERWEELE, T. J., MCEL RATH, T. F., MEEKER, J. D. & MUKHERJEE, B. (2017). Mediation of the relationship between maternal phthalate exposure and preterm birth by oxidative stress with repeated measurements across pregnancy. *Environmental Health Perspectives* **125**, 488–494.
- FRIEDMAN, J. H. (1991). Multivariate adaptive regression splines. *Annals of Statistics* , 1–67.
- FRIEDMAN, J. H. (2001). Greedy function approximation: a gradient boosting machine. *Annals of Statistics* , 1189–1232.
- GOLDBERGER, A. S. (1972). Structural equation methods in the social sciences. *Econometrica: Journal of the Econometric Society* , 979–1001.
- GREENWALD, M. K., JOHANSON, C.-E., MOODY, D. E., WOODS, J. H., KILBOURN, M. R., KOEPPE, R. A., SCHUSTER, C. R. & ZUBIETA, J.-K. (2003). Effects of buprenorphine maintenance dose on μ -opioid receptor availability, plasma concentrations, and antagonist blockade in heroin-dependent volunteers. *Neuropsychopharmacology* **28**, 2000–2009.
- HANEUSE, S. & ROTNITZKY, A. (2013). Estimation of the effect of interventions that modify the received treatment. *Statistics in Medicine* **32**, 5260–5277.
- HASTIE, T. J. & TIBSHIRANI, R. J. (1990). *Generalized additive models*, vol. 43. CRC press.
- HEIKMAN, P. K., MUHONEN, L. H. & OJANPERÄ, I. A. (2017). Polydrug abuse among opioid maintenance treatment patients is related to inadequate dose of maintenance treatment medicine. *BMC psychiatry* **17**, 1–11.

- HEJAZI, N. S. & DÍAZ, I. (2020). *medshift: Causal mediation analysis for stochastic interventions*. R package version 0.1.4.
- HEJAZI, N. S., VAN DER LAAN, M. J., JANES, H. E., GILBERT, P. B. & BENKESER, D. C. (2020). Efficient nonparametric inference on the effects of stochastic interventions under two-phase sampling, with applications to vaccine efficacy trials .
- IMAI, K., KEELE, L. & TINGLEY, D. (2010). A general approach to causal mediation analysis. *Psychological Methods* **15**, 309.
- KENNEDY, E. H. (2019). Nonparametric causal effects based on incremental propensity score interventions. *Journal of the American Statistical Association* **114**, 645–656.
- KLAASSEN, C. A. (1987). Consistent estimation of the influence function of locally asymptotically linear estimators. *Annals of Statistics* , 1548–1562.
- LEE, J. D., NUNES, E. V., BAILEY, G. L., BRIGHAM, G. S., COHEN, A. J., FISHMAN, M., LING, W., LINDBLAD, R., SHMUELI-BLUMBERG, D., STABLEIN, D. et al. (2016). Nida clinical trials network ctn-0051, extended-release naltrexone vs. buprenorphine for opioid treatment (x: Bot): study design and rationale. *Contemporary clinical trials* **50**, 253–264.
- LEE, J. D., NUNES JR, E. V., NOVO, P., BACHRACH, K., BAILEY, G. L., BHATT, S., FARKAS, S., FISHMAN, M., GAUTHIER, P., HODGKINS, C. C., KING, J., LINDBLAD, R., LIU, D., MATTHEWS, A. G., MAY, J., PEAVY, K. M., ROSS, S., SALAZAR, D., SCHKOLNIK, P., SHMUELI-BLUMBERG, D., STABLEIN, D., SUBRAMANIAM, G. & ROTROSEN, J. (2018). Comparative effectiveness of extended-release naltrexone versus buprenorphine-naloxone for opioid relapse prevention (X:BOT): a multicentre, open-label, randomised controlled trial. *The Lancet* **391**, 309–318.
- LOK, J. J. (2016). Defining and estimating causal direct and indirect effects when setting the mediator to specific values is not feasible. *Statistics in Medicine* **35**, 4008–4020.
- LOK, J. J. (2019). Causal organic direct and indirect effects: closer to baron and kenny. *arXiv preprint arXiv:1903.04697* .
- MATTICK, R. P., BREEN, C., KIMBER, J. & DAVOLI, M. (2014). Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. *Cochrane database of systematic reviews* .
- NGUYEN, T. Q., SCHMID, I. & STUART, E. A. (2019). Clarifying causal mediation analysis for the applied researcher: Defining effects based on what we want to learn. *arXiv preprint arXiv:1904.08515* .
- NUNES, E. V., LEE, J. D., SISTI, D., SEGAL, A., CAPLAN, A., FISHMAN, M., BAILEY, G., BRIGHAM, G., NOVO, P., FARKAS, S. et al. (2016). Ethical and clinical safety considerations in the design of an effectiveness trial: A comparison of buprenorphine versus naltrexone treatment for opioid dependence. *Contemporary clinical trials* **51**, 34–43.
- NUTT, D. J. (2015). Considerations on the role of buprenorphine in recovery from heroin addiction from a uk perspective. *Journal of Psychopharmacology* **29**, 43–49.
- PEARL, J. (1995). Causal diagrams for empirical research. *Biometrika* **82**, 669–688.
- PEARL, J. (1998). Graphs, causality, and structural equation models. *Sociological Methods & Research* **27**, 226–284.
- PEARL, J. (2000). *Causality*. Cambridge university press.
- PEARL, J. (2006). Direct and indirect effects. In *Proceedings of the 17th Annual Conference on Uncertainty in Artificial Intelligence*.
- PEARL, J. (2009). Myth, Confusion, and Science in Causal Analysis. Tech. Rep. R-348, Cognitive Systems Laboratory, Computer Science Department, University of California, Los Angeles, Los Angeles, CA.

- PETERSEN, M. L., SINISI, S. E. & VAN DER LAAN, M. J. (2006). Estimation of direct causal effects. *Epidemiology*, 276–284.
- PFANZAGL, J. & WEFELMEYER, W. (1985). Contributions to a general asymptotic statistical theory. *Statistics & Risk Modeling* **3**, 379–388.
- POPPER, K. (1934). *The logic of scientific discovery*. Routledge.
- R CORE TEAM (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- RICHARDSON, T. S. & ROBINS, J. M. (2013). Single world intervention graphs (swigs): A unification of the counterfactual and graphical approaches to causality. *Center for the Statistics and the Social Sciences, University of Washington Series. Working Paper* **128**, 2013.
- ROBINS, J. M. (1986). A new approach to causal inference in mortality studies with sustained exposure periods — application to control of the healthy worker survivor effect. *Mathematical Modelling* **7**, 1393–1512.
- ROBINS, J. M. & GREENLAND, S. (1992). Identifiability and exchangeability for direct and indirect effects. *Epidemiology*, 143–155.
- ROBINS, J. M., HERNÁN, M. A. & SIEBERT, U. (2004). Effects of multiple interventions. In *Comparative Quantification of Health Risks*, vol. 1. Citeseer, pp. 2191–2230.
- ROBINS, J. M. & RICHARDSON, T. S. (2010). Alternative graphical causal models and the identification of direct effects. *Causality and Psychopathology: Finding the Determinants of Disorders and Their Cures*, 103–158.
- RUDOLPH, K. E., DÍAZ, I., HEJAZI, N. S., VAN DER LAAN, M. J., LUO, S. X., SHULMAN, M., CAMPBELL, A., ROTROSEN, J. & NUNES, E. V. (2020a). Explaining differential effects on opioid use disorder treatment using a novel causal approach incorporating mediating and intermediate variables .
- RUDOLPH, K. E., SHULMAN, M., FISHMAN, M., DÍAZ, I., ROTROSEN, J. & NUNES, E. V. (2020b). Association between dynamic dose adjustment of buprenorphine for treatment of opioid use disorder and risk of relapse .
- RUDOLPH, K. E., SOFRYGIN, O., ZHENG, W. & VAN DER LAAN, M. J. (2017). Robust and flexible estimation of stochastic mediation effects: a proposed method and example in a randomized trial setting. *Epidemiologic Methods* **7**.
- SPIRITES, P., GLYMOUR, C. N., SCHEINES, R., HECKERMAN, D., MEEK, C., COOPER, G. & RICHARDSON, T. (2000). *Causation, prediction, and search*. MIT press.
- STITELMAN, O. M., HUBBARD, A. E. & JEWELL, N. P. (2010). The impact of coarsening the explanatory variable of interest in making causal inferences: Implicit assumptions behind dichotomizing variables .
- STOCK, J. H. (1989). Nonparametric policy analysis. *Journal of the American Statistical Association* **84**, 567–575.
- TCHETGEN TCHETGEN, E. J. & VANDERWEELE, T. J. (2014). On identification of natural direct effects when a confounder of the mediator is directly affected by exposure. *Epidemiology* **25**, 282.
- TIAN, J. (2008). Identifying dynamic sequential plans. In *Proceedings of the Twenty-Fourth Conference on Uncertainty in Artificial Intelligence*. AUAI Press.
- VAN DER LAAN, M. J. (2017). A generally efficient targeted minimum loss based estimator based on the highly adaptive lasso. *The International Journal of Biostatistics* **13**.
- VAN DER LAAN, M. J. & BIBAUT, A. F. (2017). Uniform consistency of the highly adaptive lasso estimator of infinite-dimensional parameters. *arXiv preprint arXiv:1709.06256* .
- VAN DER LAAN, M. J. & PETERSEN, M. L. (2008). Direct effect models. *The International Journal of Biostatistics* **4**.

- VAN DER LAAN, M. J., POLLEY, E. C. & HUBBARD, A. E. (2007). Super learner. *Statistical Applications in Genetics and Molecular Biology* **6**.
- VAN DER LAAN, M. J. & ROBINS, J. M. (2003). *Unified Methods for Censored Longitudinal Data and Causality*. Springer Science & Business Media.
- VAN DER LAAN, M. J. & ROSE, S. (2011). *Targeted Learning: Causal Inference for Observational and Experimental Data*. Springer Science & Business Media.
- VAN DER LAAN, M. J. & ROSE, S. (2018). *Targeted Learning in Data Science: Causal Inference for Complex Longitudinal Studies*. Springer Science & Business Media.
- VAN DER LAAN, M. J. & RUBIN, D. (2006). Targeted maximum likelihood learning. *The International Journal of Biostatistics* **2**.
- VAN DER VAART, A. W. (2002). Semiparametric statistics. *Lectures on Probability Theory and Statistics* .
- VANDERWEELE, T. J., VANSTEELANDT, S. & ROBINS, J. M. (2014). Effect decomposition in the presence of an exposure-induced mediator-outcome confounder. *Epidemiology* **25**, 300.
- VANSTEELANDT, S., BEKAERT, M. & LANGE, T. (2012). Imputation strategies for the estimation of natural direct and indirect effects. *Epidemiologic Methods* **1**, 131–158.
- VANSTEELANDT, S. & DANIEL, R. M. (2017). Interventional effects for mediation analysis with multiple mediators. *Epidemiology* **28**, 258.
- VANSTEELANDT, S. & VANDERWEELE, T. J. (2012). Natural direct and indirect effects on the exposed: effect decomposition under weaker assumptions. *Biometrics* **68**, 1019–1027.
- WOLPERT, D. H. (1992). Stacked generalization. *Neural Networks* **5**, 241–259.
- WRIGHT, S. (1921). Correlation and causation. *Journal of Agricultural Research* **20**, 557–585.
- WRIGHT, S. (1934). The method of path coefficients. *The Annals of Mathematical Statistics* **5**, 161–215.
- YOUNG, J. G., HERNÁN, M. A. & ROBINS, J. M. (2014). Identification, estimation and approximation of risk under interventions that depend on the natural value of treatment using observational data. *Epidemiologic Methods* **3**, 1–19.
- ZHENG, W. & VAN DER LAAN, M. J. (2011). Cross-validated targeted minimum-loss-based estimation. In *Targeted Learning: Causal Inference for Observational and Experimental Data*. Springer, pp. 459–474.
- ZHENG, W. & VAN DER LAAN, M. J. (2017). Longitudinal mediation analysis with time-varying mediators and exposures, with application to survival outcomes. *Journal of Causal Inference* **5**.

Supplementary Materials for

NONPARAMETRIC CAUSAL MEDIATION ANALYSIS FOR STOCHASTIC INTERVENTIONAL (IN)DIRECT EFFECTS

Nima S. Hejazi
Graduate Group in Biostatistics, and
Center for Computational Biology,
University of California, Berkeley
nhejazi@berkeley.edu

Kara E. Rudolph
Department of Epidemiology,
Mailman School of Public Health,
Columbia University
kr2854@cumc.columbia.edu

Mark J. van der Laan
Division of Biostatistics,
School of Public Health, and
Department of Statistics,
University of California, Berkeley
laan@berkeley.edu

Iván Díaz
Division of Biostatistics,
Department of Population Health Sciences,
Weill Cornell Medicine
ild2005@med.cornell.edu

August 3, 2022

S1 Theorem 1

Proof First, we have

$$\begin{aligned} & \mathbb{E}\{Y_{A_\delta, G_\delta}\} \\ &= \int \mathbb{E}\{Y_{a,z} \mid A_\delta = a, G_\delta = z, W = w\} g_\delta(a \mid w) P(G_\delta = z \mid A_\delta = a, W = w) p(w) d\nu(a, z, w) \\ &= \int \mathbb{E}\{Y_{a,z} \mid W = w\} g_\delta(a \mid w) P(Z(a) = z \mid A_\delta = a, W = w) p(w) d\nu(a, z, w) \end{aligned} \quad (\text{S1})$$

$$= \int \mathbb{E}\{Y_{a,z} \mid A = a, W = w\} g_\delta(a \mid w) P(Z(a) = z \mid W = w) p(w) d\nu(a, z, w) \quad (\text{S2})$$

$$= \int \mathbb{E}\{Y_{a,z} \mid A = a, W = w\} g_\delta(a \mid w) P(Z(a) = z \mid A = a, W = w) p(w) d\nu(a, z, w) \quad (\text{S3})$$

$$\begin{aligned} &= \int \mathbb{E}\{Y_{a,z} \mid A = a, W = w, L = l\} b(l \mid a, w) g_\delta(a \mid w) p(z \mid a, w) p(w) d\nu(a, z, l, w) \\ &= \int m(a, z, l, w) b(l \mid a, w) g_\delta(a \mid w) p(z \mid a, w) p(w) d\nu(a, z, l, w), \end{aligned} \quad (\text{S4})$$

where (S1) follows by definition of (A_δ, G_δ) , (S2) follows by A3 and definition of A_δ , (S3) follows by A5, and (S4) follows by A4. Similar arguments yield

$$\mathbb{E}\{Y_{A,G}\} = \int m(a, z, l, w) b(l \mid a, w) g(z \mid w) p(z \mid a, w) p(w) d\nu(a, z, l, w).$$

We also have

$$\begin{aligned}
& \mathbb{E}\{Y_{A_\delta, G}\} \\
&= \int \mathbb{E}\{Y_{a,z} \mid A_\delta = a, G = z, W = w\} g_\delta(a \mid w) P(G = z \mid A_\delta = a, W = w) p(w) d\nu(a, z, w) \\
&= \int \mathbb{E}\{Y_{a,z} \mid W = w\} g_\delta(a \mid w) P(G = z \mid W = w) p(w) d\nu(a, z, w) \\
&= \int \mathbb{E}\{Y_{a,z} \mid A = a, W = w\} g_\delta(a \mid w) p(z \mid w) p(w) d\nu(a, z, w) \\
&= \int \mathbb{E}\{Y_{a,z} \mid A = a, W = w\} g_\delta(a \mid w) p(z \mid w) p(w) d\nu(a, z, w) \\
&= \int \mathbb{E}\{Y_{a,z} \mid A = a, W = w, L = l\} b(l \mid a, w) g_\delta(a \mid w) p(z \mid w) p(w) d\nu(a, z, l, w) \\
&= \int m(a, z, l, w) b(l \mid a, w) g_\delta(a \mid w) p(z \mid w) p(w) d\nu(a, z, l, w).
\end{aligned}$$

Subtracting gives the expressions for the PIIE and PIDE in the theorem. \square

S2 Efficient influence functions (Theorem 2)

Proof In this proof we will use $\Theta_j(P) : j = 1, 2$ to denote a parameter as a functional that maps the distribution P in the model to a real number. We will assume that the measure ν is discrete so that integrals can be written as sums, and will omit the dependence on δ . It can be checked algebraically that the resulting influence function will also correspond to the influence function of a general measure ν . The true parameter value for θ_1 is thus given by

$$\theta_1 = \Theta_1(P) = \sum_{y,a,z,m,w} y p(y \mid a, z, l, w) p(l \mid a, w) p(z \mid a, w) g_\delta(a \mid w) p(w).$$

The non-parametric MLE of θ_1 in the model of g_δ known is given by

$$\Theta(P_n) = \sum_{y,a,z,m,w} y \frac{P_n f_{y,a,z,l,w}}{P_n f_{a,z,l,w}} \frac{P_n f_{l,a,w}}{P_n f_{a,w}} \frac{P_n f_{z,a,w}}{P_n f_{a,w}} g_\delta(a \mid w) P_n f_w, \quad (\text{S5})$$

where we remind the reader of the notation $Pf = \int f dP$. Here $f_{y,a,z,l,w} = \mathbb{1}(Y = y, A = a, Z = z, M = m, W = w)$, and $\mathbb{1}(\cdot)$ denotes the indicator function. The other functions f are defined analogously.

We will use the fact that the efficient influence function in a non-parametric model corresponds with the influence curve of the NPMLE. This is true because the influence curve of any regular estimator is also a gradient, and a non-parametric model has only one gradient. The Delta method (see, e.g., Appendix 18 of [van der Laan & Rose, 2011](#)) shows that if $\hat{\Theta}_1(P_n)$ is a substitution estimator such that $\theta_1 = \hat{\Theta}_1(P)$, and $\hat{\Theta}_1(P_n)$ can be written as $\hat{\Theta}_1^*(P_n f : f \in \mathcal{F})$ for some class of functions \mathcal{F} and some mapping Θ_1^* , the influence function of $\hat{\Theta}_1(P_n)$ is equal to

$$\text{IF}_P(O) = \sum_{f \in \mathcal{F}} \frac{d\hat{\Theta}_1^*(P)}{dPf} \{f(O) - Pf\}.$$

Applying this result to (S5) with $\mathcal{F} = \{f_{y,a,z,l,w}, f_{a,z,l,w}, f_{z,a,w}, f_{a',w}, f_{l,a,w}, f_{a,w}, f_w : y, a, z, l, w\}$ and rearranging terms gives the result of the theorem. The algebraic derivations involved here are lengthy and not particularly illuminating, and are therefore omitted from the proof. Similar analyses may be performed for the model where only g_δ is unknown, as well as θ_2 . \square

S3 Targeted minimum loss estimation algorithm

To simplify notation, in the remaining of this section we will denote $\tilde{\eta}_{j(i)}(O_i)$ with $\tilde{\eta}(O_i)$. If L is binary, the efficient influence functions in Theorem 2 may be simplified using the following identity:

$$v(l, a, w) - \bar{v}(a, w) = \{v(1, a, w) - v(0, a, w)\} \{l - b(1 \mid a, w)\},$$

which also holds for v replaced by s and \bar{v} by \bar{s} .

Step 1. Initialize $\tilde{\eta} = \hat{\eta}$. Compute \tilde{v} , \tilde{s} , and \tilde{q}^j by plugging in \tilde{m} , \tilde{g} , \tilde{e} , \tilde{d} into equations (7), (16) and (15) if Z is multivariate, and fitting data-adaptive regression algorithms as appropriate.

Step 2. For each subject, compute the auxiliary covariates

$$\begin{aligned} H_{D,i} &= \frac{\tilde{b}(L_i | A_i, W_i)}{\tilde{d}(L_i | Z_i, A_i, W_i)} \left\{ 1 - \frac{\tilde{g}_\delta(A_i | W_i)}{\tilde{e}(A_i | Z_i, W_i)} \right\} \\ H_{I,i} &= \frac{\tilde{b}(L_i | A_i, W_i)}{\tilde{d}(L_i | Z_i, A_i, W_i)} \left\{ \frac{\tilde{g}_\delta(A_i | W_i)}{\tilde{e}(A_i | Z_i, W_i)} - \frac{\tilde{g}_\delta(A_i | W_i)}{\tilde{g}(A_i | W_i)} \right\} \\ K_{D,i} &= \tilde{v}(1, A_i, W_i) - \tilde{v}(0, A_i, W_i) - \frac{\tilde{g}_\delta(A_i | W_i)}{\tilde{g}(A_i | W_i)} \{ \tilde{s}(1, A_i, W_i) - \tilde{s}(0, A_i, W_i) \} \\ K_{I,i} &= \frac{\tilde{g}_\delta(A_i | W_i)}{\tilde{g}(A_i | W_i)} \{ \tilde{s}(1, A_i, W_i) - \tilde{s}(0, A_i, W_i) - \tilde{v}(1, A_i, W_i) + \tilde{v}(0, A_i, W_i) \} \\ M_{D,i} &= -\frac{\tilde{g}_\delta(1 | w)(1 - \tilde{g}_\delta(1 | w))}{\tilde{g}(1 | w)(1 - \tilde{g}(1 | w))} \tilde{q}^2(w) \\ M_{I,i} &= \frac{\tilde{g}_\delta(1 | w)(1 - \tilde{g}_\delta(1 | w))}{\tilde{g}(1 | w)(1 - \tilde{g}(1 | w))} \{ \tilde{q}^2(w) - \tilde{q}^1(w) \} \end{aligned}$$

Step 3. Fit the logistic tilting models

$$\begin{aligned} \text{logit } m_\beta(A_i, Z_i, L_i, W_i) &= \text{logit } \tilde{m}(A_i, Z_i, L_i, W_i) + \beta_I H_{I,i} + \beta_D H_{D,i} \\ \text{logit } b_\alpha(1 | A_i, W_i) &= \text{logit } \tilde{b}(1 | A_i, W_i) + \alpha_I K_{I,i} + \alpha_D K_{D,i} \\ \text{logit } g_\gamma(1 | W_i) &= \text{logit } \tilde{g}(1 | W_i) + \gamma_I M_{I,i} + \gamma_D M_{D,i} \end{aligned}$$

where $\text{logit}(p) = \log\{p(1-p)^{-1}\}$. Here, $\text{logit } \tilde{m}(a, z, l, w)$ is an offset variable (i.e., a variable with known parameter value equal to one). The parameter $\beta = (\beta_I, \beta_D)$ may be estimated by running standard logistic regression of Y_i on $(H_{D,i}, H_{I,i})$ with no intercept and an offset term equal to $\text{logit } \tilde{m}(A_i, Z_i, L_i, W_i)$. Let $\hat{\beta}$ denote the estimate, and let $\tilde{m} = m_{\hat{\beta}}$ denote the updated estimates. Perform analogous computations for b and g .

Step 4. Compute \tilde{u} according to equation (7) by plugging in \tilde{m} and \tilde{b} . Compute the covariate

$$J_i = \frac{\tilde{g}_\delta(A_i | W_i)}{\tilde{g}(A_i | W_i)},$$

and fit the model

$$\text{logit } \bar{u}_\kappa(A_i, W_i) = \text{logit } \tilde{u}(A_i, W_i) + \kappa_D + \kappa_I J_i$$

by running a logistic regression of $\tilde{u}(Z_i, A_i, W_i)$ on J_i with an intercept and offset $\text{logit } \tilde{u}(A_i, W_i)$. Let $\hat{\kappa}$ denote the MLE, and update $\tilde{u} = \bar{u}_{\hat{\kappa}}$.

Step 5. The TMLE of the direct and indirect effects are defined as:

$$\begin{aligned} \hat{\psi}_{D,\delta}^{\text{tmle}} &= \frac{1}{n} \int \sum_{i=1}^n \{ \tilde{u}(a, W_i) \tilde{g}(a | W_i) - \tilde{u}(Z_i, a, W_i) \tilde{g}_\delta(a | W_i) \} d\kappa(a) \\ \hat{\psi}_{I,\delta}^{\text{tmle}} &= \frac{1}{n} \int \sum_{i=1}^n \{ \tilde{u}(Z_i, a, W_i) - \tilde{u}(a, W_i) \} \tilde{g}_\delta(a | W_i) d\kappa(a) \end{aligned}$$

S4 Proof of Theorem 3

Proof Let $P_{n,j}$ denote the empirical distribution of the prediction set \mathcal{V}_j , and let $G_{n,j}$ denote the associated empirical process $\sqrt{n/J}(P_{n,j} - P)$. For simplicity we denote a general parameter ψ with influence function D_η , the proof applies equally to the direct and indirect effect parameters. Note that

$$\hat{\psi}_\delta^{\text{os}} = \frac{1}{J} \sum_{j=1}^J P_{n,j} D_{\hat{\eta}_j, \delta}, \quad \psi_\delta = P D_\eta.$$

Thus,

$$\sqrt{n}\{\hat{\psi}_\delta^{\text{os}} - \psi_\delta\} = G_n\{D_{\eta,\delta} - \psi_\delta\} + R_{n,1}(\delta) + R_{n,2}(\delta),$$

where

$$R_{n,1}(\delta) = \frac{1}{\sqrt{J}} \sum_{j=1}^J G_{n,j}(D_{\hat{\eta}_j,\delta} - D_{\eta,\delta}), \quad R_{n,2}(\delta) = \frac{\sqrt{n}}{J} \sum_{j=1}^J P\{D_{\hat{\eta}_j,\delta} - \psi_\delta\}.$$

It remains to show that $R_{n,1}(\delta)$ and $R_{n,2}(\delta)$ are $o_P(1)$. Lemmas 4 and 5 together with the Cauchy-Schwartz inequality and assumption (ii) of the theorem shows that $\|R_{n,2}\|_\Delta = o_P(1)$. For $\|R_{n,1}\|_\Delta$ we use empirical process theory to argue conditional on the training sample \mathcal{T}_j . In particular, Lemma 19.33 of van der Vaart (2000) applied to the class of functions $\mathcal{F} = \{D_{\hat{\eta}_j,\delta} - D_{\eta,\delta}\}$ (which consists of one element) yields

$$E \left\{ \left| G_{n,j}(D_{\hat{\eta}_j,\delta} - D_{\eta,\delta}) \right| \middle| \mathcal{T}_j \right\} \lesssim \frac{2C \log 2}{n^{1/2}} + \|D_{\hat{\eta}_j,\delta} - D_{\eta,\delta}\|(\log 2)^{1/2}$$

By assumption (ii), the left hand side is $o_P(1)$. Lemma 6.1 of Chernozhukov et al. (2018) may now be used to argue that conditional convergence implies unconditional convergence, concluding the proof. \square

S5 Theorem 4

Proof Let $P_{n,j}$ denote the empirical distribution of the prediction set \mathcal{V}_j , and let $G_{n,j}$ denote the associated empirical process $\sqrt{n/J}(P_{n,j} - P)$. For simplicity we denote a general parameter ψ with influence function D_η , the proof applies equally to the direct and indirect effect parameters. By definition, the sum of the scores of the sub-models $\{\mathbf{m}_\beta, \mathbf{b}_\alpha, \mathbf{g}_\gamma, \bar{\mathbf{u}}_\kappa : (\beta, \alpha, \gamma, \kappa)\}$ at the last iteration of the TMLE procedure is equal to $n^{-1} \sum_{i=1}^n D_{\hat{\eta}}(O_i) = o_P(n^{-1/2})$. Thus, we have

$$\hat{\psi}_\delta^{\text{tmle}} = \frac{1}{J} \sum_{j=1}^J P_{n,j} D_{\hat{\eta}_j} + o_P(n^{-1/2}).$$

Thus,

$$\sqrt{n}(\hat{\psi}_\delta^{\text{tmle}} - \theta) = G_n(D_\eta - \theta) + R_{n,1} + R_{n,2} + o_P(n^{-1/2}),$$

where

$$R_{n,1} = \frac{1}{\sqrt{J}} \sum_{j=1}^J G_{n,j}(D_{\hat{\eta}_j} - D_\eta), \quad R_{n,2} = \frac{\sqrt{n}}{J} \sum_{j=1}^J P(D_{\hat{\eta}_j} - \theta).$$

As in the proof of Theorem 3, Lemmas 4 and 5 together with the Cauchy-Schwartz inequality and the assumptions of the theorem shows that $R_{n,2} = o_P(1)$.

Since $D_{\hat{\eta}_j}$ depends on the full sample through the estimates of the parameters β of the logistic tilting models, the empirical process treatment of $R_{n,1}$ needs to be slightly from that in the proof of Theorem 3. To make this dependence explicit, we introduce the notation $D_{\hat{\eta}_j,\beta} = D_{\hat{\eta}_j}$ and $R_{n,1}(\beta)$. Let $\mathcal{F}_n^j = \{D_{\hat{\eta}_j,\beta} - D_\eta : \beta \in B\}$. Because the function $\hat{\eta}_j$ is fixed given the training data, we can apply Theorem 2.14.2 of van der Vaart & Wellner (1996) to obtain

$$E \left\{ \sup_{f \in \mathcal{F}_n^j} |G_{n,j}f| \middle| \mathcal{T}_j \right\} \lesssim \|F_n^j\| \int_0^1 \sqrt{1 + N_{[]}(\epsilon \|F_n^j\|, \mathcal{F}_n^j, L_2(P))} d\epsilon,$$

where $N_{[]}(\epsilon \|F_n^j\|, \mathcal{F}_n^j, L_2(P))$ is the bracketing number and we take $F_n^j = \sup_{\beta \in B} |D_{\hat{\eta}_j,\beta} - D_\eta|$ as an envelope for the class \mathcal{F}_n^j . Theorem 2.7.2 of van der Vaart & Wellner (1996) shows

$$\log N_{[]}(\epsilon \|F_n^j\|, \mathcal{F}_n^j, L_2(P)) \lesssim \frac{1}{\epsilon \|F_n^j\|}.$$

This shows

$$\begin{aligned} \|F_n^j\| \int_0^1 \sqrt{1 + N_{[]}(\epsilon \|F_n^j\|, \mathcal{F}_n^j, L_2(P))} d\epsilon &\lesssim \int_0^1 \sqrt{\|F_n^j\|^2 + \frac{\|F_n^j\|}{\epsilon}} d\epsilon \\ &\leq \|F_n^j\| + \|F_n^j\|^{1/2} \int_0^1 \frac{1}{\epsilon^{1/2}} d\epsilon \\ &\leq \|F_n^j\| + 2\|F_n^j\|^{1/2}. \end{aligned}$$

Since $\|F_n^j\| = o_P(1)$, this shows $\sup_{f \in \mathcal{F}_n^j} G_{n,j}f = o_P(1)$ for each j , conditional on \mathcal{T}_j . Thus $\sup_{\beta \in B} R_{n,1}(\beta) = o_P(1)$. Lemmas 4 and 5 together with the Cauchy-Schwartz inequality and the assumptions of the theorem show that $R_{n,2} = o_P(1)$, concluding the proof of the theorem. \square

S6 Additional results

Lemma 5 (Second order terms for modified treatment policies). *Let $d\xi(o)$ denote $d\nu(a, l, z)dP(w)$, and let $r(z \mid a, w)$ denote $p(z \mid a, w)$, and let $h(z \mid w)$ denote $p(z \mid w)$. Let $d(a, w)$ denote a modified treatment policy satisfying A1. We have*

$$PD_{\eta_1, \delta}^1 - \psi_1(\delta) = \int \left(\frac{g}{g_1} \frac{d}{d_1} - 1 \right) (m - m_1) b_1 r g_{\delta, 1} d\xi \quad (S6)$$

$$- \int \left(\frac{g}{g_1} - 1 \right) (\bar{u}_1 - \bar{u}) g_{\delta, 1} d\xi \quad (S7)$$

$$+ \int \left(\frac{g}{g_1} - 1 \right) (m_1 - m) b_1 r g_{\delta, 1} d\xi \quad (S8)$$

$$- \int \frac{g}{g_1} (b_1 - b) (v_1 - v) g_{\delta, 1} d\xi \quad (S9)$$

$$- \int (\bar{u}_1 - \bar{u}) (g_{\delta, 1} - g_\delta) d\xi \quad (S10)$$

and

$$PD_{\eta_1, \delta}^2 - \psi_2(\delta) = \int \left(\frac{e}{e_1} \frac{d}{d_1} - 1 \right) (m - m_1) b_1 h g_{\delta, 1} d\xi$$

$$+ \int \frac{g}{g_1} (b_1 - b) (s_1 - s) g_{\delta, 1} d\xi$$

$$- \int (q_1 - q) (g_{\delta, 1} - g_\delta) d\xi.$$

Proof Note that

$$\begin{aligned}
PS_{\eta_1, \delta}^1 - \psi_1(\delta) &= \int \left(\frac{g}{g_1} \frac{d}{d_1} - 1 \right) (m - m_1) b_1 r g_{\delta, 1} d\xi + \int (m - m_1) b_1 r g_{\delta, 1} d\xi \\
&\quad - \int \bar{u}(g_\delta - g_{\delta, 1}) d\xi - \int \bar{u} g_{\delta, 1} d\xi \\
&\quad + \int \frac{g}{g_1} (b - b_1) v_1 g_{\delta, 1} d\xi + \int \frac{g}{g_1} (u_1 r - \bar{u}_1) g_{\delta, 1} d\xi + \int \bar{u}_1 g_{\delta, 1} d\xi \\
&= (S6) + \int (\bar{u}_1 - \bar{u}) g_{\delta, 1} d\xi \\
&\quad + \int (m - m_1) b_1 r g_{\delta, 1} d\xi + \int \frac{g}{g_1} (b - b_1) v_1 g_{\delta, 1} d\xi \\
&\quad + \int \frac{g}{g_1} u_1 r g_{\delta, 1} d\xi - \int \frac{g}{g_1} \bar{u}_1 g_{\delta, 1} d\xi \\
&\quad - \int \bar{u}(g_\delta - g_{\delta, 1}) d\xi \\
&= (S6) - \int \left(\frac{g}{g_1} - 1 \right) (\bar{u}_1 - \bar{u}) g_{\delta, 1} d\xi \\
&\quad + \int \frac{g}{g_1} m_1 b_1 r g_{\delta, 1} d\xi - \int \frac{g}{g_1} m b r g_{\delta, 1} d\xi \\
&\quad + \int (m - m_1) b_1 r g_{\delta, 1} d\xi + \int \frac{g}{g_1} (b - b_1) v_1 g_{\delta, 1} d\xi \\
&\quad - \int \bar{u}(g_\delta - g_{\delta, 1}) d\xi \\
&= (S6) - (S7) \\
&\quad + \int (m - m_1) b_1 r g_{\delta, 1} d\xi + \int \frac{g}{g_1} (b - b_1) v_1 g_{\delta, 1} d\xi \\
&\quad + \int \frac{g}{g_1} (m_1 b_1 + m b_1 - m b_1 - m b) r g_{\delta, 1} d\xi \\
&\quad - \int \bar{u}(g_\delta - g_{\delta, 1}) d\xi \\
&= (S6) - (S7) \\
&\quad + \int (m - m_1) b_1 r g_{\delta, 1} d\xi + \int \frac{g}{g_1} (b - b_1) v_1 g_{\delta, 1} d\xi \\
&\quad + \int \frac{g}{g_1} (m_1 - m) b_1 r g_{\delta, 1} d\xi + \int \frac{g}{g_1} (b_1 - b) m r g_{\delta, 1} d\xi \\
&\quad - \int \bar{u}(g_\delta - g_{\delta, 1}) d\xi \\
&= (S6) - (S7) + (S8) - (S9) \\
&\quad - \int \bar{u}(g_\delta - g_{\delta, 1}) d\xi.
\end{aligned} \tag{S11}$$

Using A1 we can change variables to obtain

$$PS_{\eta_1, \delta}^{A, 1} = \int \bar{u}_1 (g_\delta - g_{\delta, 1}) d\xi.$$

The proof for ψ_2 is analogous. This completes the proof of the theorem. \square

Lemma 6 (Second order terms for exponential tilting.). Define $c(w) = \{\int_a \exp(\delta a) \mathbf{g}(a \mid w)\}^{-1}$, and let $c_1(w)$ be defined analogously. Let $b(a) = \exp(\delta a)$. Using the same notation as in Lemma 4, we have

$$\begin{aligned} PD_{\eta_1, \delta}^1 - \psi_1(\delta) &= \int \left(\frac{\mathbf{g}}{\mathbf{g}_1} \frac{\mathbf{d}}{\mathbf{d}_1} - 1 \right) (\mathbf{m} - \mathbf{m}_1) \mathbf{b}_1 \mathbf{r} \mathbf{g}_{\delta, 1} d\xi \\ &\quad - \int \left(\frac{\mathbf{g}}{\mathbf{g}_1} - 1 \right) (\bar{\mathbf{u}}_1 - \bar{\mathbf{u}}) \mathbf{g}_{\delta, 1} d\xi \\ &\quad + \int \left(\frac{\mathbf{g}}{\mathbf{g}_1} - 1 \right) (\mathbf{m}_1 - \mathbf{m}) \mathbf{b}_1 \mathbf{r} \mathbf{g}_{\delta, 1} d\xi \\ &\quad - \int \frac{\mathbf{g}}{\mathbf{g}_1} (\mathbf{b}_1 - \mathbf{b}) (\mathbf{v}_1 - \mathbf{v}) \mathbf{g}_{\delta, 1} d\xi \\ &\quad + \int (\bar{\mathbf{u}}_1 - \bar{\mathbf{u}}) (\mathbf{g}_{\delta, 1} - \mathbf{g}_\delta) d\xi \\ &\quad - \int \left\{ (c_1 - c)^2 \int b \mathbf{g}_1 \bar{\mathbf{u}}_1 d\kappa \int b \mathbf{g} d\kappa \right\} d\xi \\ &\quad + \int \left\{ (c_1 - c) \int b \bar{\mathbf{u}}_1 (\mathbf{g} - \mathbf{g}_1) d\kappa \right\} d\xi, \end{aligned}$$

and

$$\begin{aligned} PD_{\eta_1, \delta}^2 - \psi_2(\delta) &= \int \left(\frac{\mathbf{e}}{\mathbf{e}_1} \frac{\mathbf{d}}{\mathbf{d}_1} - 1 \right) (\mathbf{m} - \mathbf{m}_1) \mathbf{b}_1 \mathbf{h} \mathbf{g}_{\delta, 1} d\xi \\ &\quad + \int \frac{\mathbf{g}}{\mathbf{g}_1} (\mathbf{b}_1 - \mathbf{b}) (\mathbf{s}_1 - \mathbf{s}) \mathbf{g}_{\delta, 1} d\xi \\ &\quad - \int (\mathbf{q}_1 - \mathbf{q}) (\mathbf{g}_{\delta, 1} - \mathbf{g}_\delta) d\xi \\ &\quad - \int \left\{ (c_1 - c)^2 \int b \mathbf{g}_1 \bar{\mathbf{q}}_1 d\kappa \int b \mathbf{g} d\kappa \right\} d\xi \\ &\quad + \int \left\{ (c_1 - c) \int b \bar{\mathbf{q}}_1 (\mathbf{g} - \mathbf{g}_1) d\kappa \right\} d\xi. \end{aligned}$$

Proof In this proof, (S11) is also valid. We have

$$PS_{\eta_1, \delta}^{1, A} - \int \bar{\mathbf{u}} (\mathbf{g}_\delta - \mathbf{g}_{\delta, 1}) d\xi = PS_{\eta_1, \delta}^{1, A} - \int \bar{\mathbf{u}}_1 (\mathbf{g}_\delta - \mathbf{g}_{\delta, 1}) d\xi + \int (\bar{\mathbf{u}}_1 - \bar{\mathbf{u}}) (\mathbf{g}_{\delta, 1} - \mathbf{g}_\delta) d\xi$$

It thus remains to prove that

$$\begin{aligned} PS_{\eta_1, \delta}^{1, A} - \int \bar{\mathbf{u}}_1 (\mathbf{g}_\delta - \mathbf{g}_{\delta, 1}) d\xi &= - \int \left\{ (c_1 - c)^2 \int b \mathbf{g}_1 \bar{\mathbf{u}}_1 d\kappa \int b \mathbf{g} d\kappa \right\} d\xi \\ &\quad + \int \left\{ (c_1 - c) \int b \bar{\mathbf{u}}_1 (\mathbf{g} - \mathbf{g}_1) d\kappa \right\} d\xi. \end{aligned}$$

We have

$$\begin{aligned}
PS_{\eta_i}^{1,A} &= \int \bar{u}_1 (g_\delta - g_{\delta,1}) d\xi \\
&= \int \left\{ \int \frac{g_{1,\delta}}{g_1} \bar{u}_1 g d\kappa - \int \frac{g_{1,\delta}}{g_1} g d\kappa \int \bar{u}_1 g_{1,\delta} d\kappa + \int (g_{1,\delta} - g_\delta) \bar{u}_1 d\kappa \right\} d\xi \\
&= \int \left\{ \frac{g_{1,\delta}}{g_1} g \bar{u}_1 d\kappa - \int g_\delta \bar{u}_1 d\kappa + \int g_{1,\delta} \bar{u}_1 d\kappa \left[1 - \int \frac{g_{1,\delta}}{g_1} g d\kappa \right] \right\} d\xi \\
&= \int \left\{ c_1 \int b \bar{u}_1 g d\kappa - c_1 \int b \bar{u}_1 g d\kappa + c_1 \int b g \bar{u}_1 d\kappa \int (c - c_1) b g d\kappa \right\} d\xi \\
&= \int (c_1 - c) \left\{ \int b \bar{u}_1 g d\kappa - c_1 \int b g_1 \bar{u}_1 d\kappa \int b g d\kappa \right\} d\xi \\
&= \int (c_1 - c) \left\{ \int b \bar{u}_1 g d\kappa - c \int b g_1 \bar{u}_1 d\kappa \int b g d\kappa - (c_1 - c) \int b g_1 \bar{u}_1 d\kappa \int b g d\kappa \right\} d\xi \\
&= \int \left\{ -(c_1 - c)^2 \int b g_1 \bar{u}_1 d\kappa \int b g d\kappa + (c_1 - c) \left[\int b \bar{u}_1 g d\kappa - \int b g_1 \bar{u}_1 d\kappa \right] \right\} d\xi \quad (S12) \\
&= \int \left\{ -(c_1 - c)^2 \int b g_1 \bar{u}_1 d\kappa \int b g d\kappa + (c_1 - c) \int b \bar{u}_1 (g - g_1) d\kappa \right\} d\xi,
\end{aligned}$$

where (S12) follows from $c \int b g d\kappa = 1$. The proof for ψ_2 is analogous. \square

References

- CHERNOZHUKOV, V., CHETVERIKOV, D., DEMIRER, M., DUFLO, E., HANSEN, C., NEWEY, W. & ROBINS, J. M. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal* **21**, C1–C68.
- VAN DER LAAN, M. J. & ROSE, S. (2011). *Targeted Learning: Causal Inference for Observational and Experimental Data*. Springer Science & Business Media.
- VAN DER VAART, A. W. (2000). *Asymptotic statistics*, vol. 3. Cambridge university press.
- VAN DER VAART, A. W. & WELLNER, J. A. (1996). *Weak convergence and empirical processes*. Springer.