

# Fundamental physical constants at low and high redshifts

Sergei A. Levshakov

A. F. Ioffe Physical-Technical Institute, St. Petersburg, Russia,  
[lev@astro.ioffe.ru](mailto:lev@astro.ioffe.ru)

**Abstract.** Spacetime variations of physical constants can be associated with the existence of Higgs-like scalar field(s) that couple non-universally to the baryonic matter. Recent results of astronomical spectral measurements of the fractional changes in the electron-to-proton mass ratio,  $\mu = m_e/m_p$ , at low ( $z \sim 0$ ) and high ( $z \sim 6.5$ ) redshifts are discussed. It is shown that the distribution of the most accurate estimates of  $\Delta\mu/\mu = (\mu_{\text{obs}} - \mu_{\text{lab}})/\mu_{\text{lab}}$  ranging between  $z = 0$  and  $z \sim 1100$  can be approximated by a power law  $\Delta\mu/\mu = k_\mu(1+z)^p$ , with  $k_\mu = (1.7 \pm 0.3) \times 10^{-8}$  and  $p = 1.99 \pm 0.03$ , implying a dynamical nature of the scalar field(s).

**Keywords:** elementary particles – techniques: spectroscopic – cosmology: observations

## 1 Introduction

The suggestion that the physical constants may vary on the cosmological time scale can be traced as far back as 1937, when Milne (1) and Dirac (2) argued about possible changes in the gravitational constant  $G$  during the lifetime of the universe. Modern physics considers the masses of quarks and leptons as a result of their Yukawa coupling to the scalar Higgs field. Other scalar fields, which are proposed to explain the phenomena of dark energy and dark matter, can also couple to the standard baryonic matter and, hence, change the masses of elementary particles. Since the proton and neutron masses are mainly determined by the binding energy of quarks (3), the changes of their masses due to dynamical scalar fields are much smaller compared to the effect of altering  $m_e$  (4). Thus, values of the fine structure constant,  $\alpha = e^2/\hbar c$ , and the electron to proton mass ratio,  $\mu = m_e/m_p$ , measured at different physical and cosmological conditions can be used to probe the hidden scalar fields. Nowadays, the search for hypothetical variations of  $\alpha$  and  $\mu$  is among the most extensively studying problems in laboratory and cosmic physics.

The unprecedentedly sensitive limits on the temporal drift of these dimensionless constants achieved for the passed decades in laboratory experiments are of one part in  $10^{15} - 10^{16}$  per year (5; 6; 7).

Astrophysical studies of extragalactic objects at low ( $z \sim 0$ ) and high ( $z \sim 1 - 7$ ) redshifts constrain the fractional changes in  $\Delta\alpha/\alpha = (\alpha_{\text{obs}} - \alpha_{\text{lab}})/\alpha_{\text{lab}}$  and  $\Delta\mu/\mu = (\mu_{\text{obs}} - \mu_{\text{lab}})/\mu_{\text{lab}}$  at the same order of magnitude (8; 9; 10; 11) if we assume a linear drift of the physical constants with cosmic time. New optical spectral observations with two high-resolution spectrographs ESPRESSO/VLT (12) and HIRES/E-ELT (13) are planning to put ever deeper constraints on  $\alpha$ - and  $\mu$ -variations.

A recent analysis of the ionization processes at redshift  $z \sim 1100$  which are responsible for the temperature and polarization anisotropies of the cosmic microwave background (CMB) radiation has shown that the value of the Hubble constant  $H_0$  correlates with the electron mass (14). As a result, an increased effective electron mass at the epoch of recombination,  $m_{e,z} = (1.0190 \pm 0.0055)m_{e,0}$ , yields  $H_0 \simeq 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$  — the value shifted with respect to  $H_0 = 67.4 \pm 0.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$  which is based on the standard  $\Lambda$ CDM model (15). Such a 2% increasing of the electron mass at  $z \sim 1100$  may alleviate the difference between the Hubble constant measured at  $z \lesssim 1$ ,  $H_0 = 74.03 \pm 1.42 \text{ km s}^{-1} \text{ Mpc}^{-1}$  (16), and the CMB value.


## 2 Constraints on the time variation of $\mu$

Note that the spacetime variation of the electromagnetic fine structure constant  $\alpha$  should be much smaller than that of  $\mu$  (17). In the following we will consider  $\alpha$  unchangeable and put  $\Delta\alpha/\alpha = 0$ .

The variation of  $\mu$  can be probed by comparing the relative frequencies of different molecular transitions as was originally suggested in 1975 by Thompson (18). Later on, it was shown that transitions within the molecular bands have different sensitivities to the variation of  $\mu$  (19). The corresponding sensitivity coefficients  $Q$  for the Lyman and Werner transitions of  $\text{H}_2$  are typically  $\sim 0.01$  and differ in sign (19; 20). The  $\text{H}_2$  absorption lines detected in quasar spectra (21; 22) give the following estimation of  $\Delta\mu/\mu$ :

$$\Delta\mu/\mu = (R_{ij} - R_{ij}^0)/[R_{ij}^0(Q_j - Q_i)] , \quad (1)$$

where  $R_{ij} = (\lambda_i/\lambda_j)_z$  is the ratio of two  $\text{H}_2$  lines at the epoch  $z$ ,  $R_{ij}^0$  — at  $z = 0$ , and  $Q_i, Q_j$  are the sensitivities of the wavelengths  $\lambda_i, \lambda_j$  to the variation of  $\mu$ . This so-called  $\text{H}_2$  method, being applied to optical spectra of quasars, provides the mean value of  $\langle \Delta\mu/\mu \rangle = (-2.5 \pm 5.3) \times 10^{-6}$  in the interval from  $z = 2.05$  to  $z = 4.22$  (23).



The microwave and submillimeter astronomical spectra are essentially more sensitive to variations in  $\mu$  than the H<sub>2</sub> lines (24). For instance, the *NH<sub>3</sub> method* (25) yields

$$\Delta\mu/\mu = (V_{\text{rot}} - V_{\text{inv}})/[c(Q_{\text{inv}} - Q_{\text{rot}})] , \quad (2)$$

where the observed radial velocity  $V_{\text{inv}}$  of the inversion transition of NH<sub>3</sub>(1,1) with  $Q_{\text{inv}} = 4.46$  is compared with a suitable rotational radial velocity  $V_{\text{rot}}$  of another molecule co-spatially distributed with ammonia and having  $Q_{\text{rot}} = 1$ , and  $c$  is the speed of light. In the Milky Way disk, the *NH<sub>3</sub> method* provides  $\langle\Delta\mu/\mu\rangle = (2 \pm 6) \times 10^{-9}$  (26).

A wide range of  $Q$ -values (from  $\simeq -15$  to  $\simeq 45$ ) of methanol CH<sub>3</sub>OH lines was deduced in (27; 28). Any two transitions of CH<sub>3</sub>OH with different sensitivity coefficients  $Q_i$  and  $Q_j$  give the following estimate of  $\Delta\mu/\mu$ :

$$\Delta\mu/\mu = (V_j - V_i)/[c(Q_i - Q_j)] , \quad (3)$$

where  $V_j$  and  $V_i$  are the apparent radial velocities of the corresponding methanol transitions. The deepest limits on  $\Delta\mu/\mu$  were obtained by the *CH<sub>3</sub>OH method* for the Galactic cloud L1498 (29) and the  $z = 0.89$  gravitational lens (10):  $\Delta\mu/\mu = (-3 \pm 2) \times 10^{-8}$  and  $(3 \pm 6) \times 10^{-8}$ , respectively.

Finally, to probe  $\Delta\mu/\mu$  by spectroscopic methods at very high redshifts ( $z \gtrsim 6$ ) a fine structure transition method (FST) was proposed in (30). It is based on the analysis of the radial velocity offsets between CO rotational lines and [C I] and/or [C II] fine-structure transitions. Under assumption that  $\Delta\alpha/\alpha = 0$ , it gives (23):

$$\Delta\mu/\mu = \Delta V/c , \quad (4)$$

where  $\Delta V$  is the radial velocity offset,  $\Delta V = V_{\text{fs}} - V_{\text{rot}}$ . Three estimates of  $\Delta\mu/\mu$  at  $z = 6.003$ , 6.419, and 6.519 by the FST method provide a weighted mean value of  $\langle\Delta\mu/\mu\rangle = (0.7 \pm 1.2) \times 10^{-5}$  at  $\bar{z} = 6.3$  (23).

All available values of  $\Delta\mu/\mu$ , ranging between  $z = 0$  and  $z \sim 1100$ , were put together in Fig. 3 in (23). Here, in Fig. 1, we reproduce this compilation with a slightly updated dataset: instead of one  $\Delta\mu/\mu$  value averaged over the whole Triangulum galaxy M33 ( $D \simeq 800$  kpc) we use three  $\Delta\mu/\mu$  values calculated on base of the CO(2-1) and [C II] emission lines in regions at different galactocentric distances:  $R_c = 0, 2$ , and 3.5 kpc. The distribution of the updated data points (shown by dots with  $1\sigma$  error bars in Fig. 1) was approximated by a simple power law

$$\Delta\mu/\mu = k_\mu(1 + z)^p , \quad (5)$$

which gives  $k_\mu = (1.7 \pm 0.3) \times 10^{-8}$  and  $p = 1.99 \pm 0.03$  ( $1\sigma$ ). The corresponding regression curve is shown by red in Fig. 1.

An important point to note is that the revealed  $z$ -dependence of  $\mu$  crucially depends on the CMB estimate of the electron mass at  $z \sim 1100$ .

It is obvious that to verify the redshift dependence of  $\mu$  a more refined analysis of the CMB anisotropies is needed along with more accurate spectral measurements at lower redshifts. The required uncertainty of the molecular line position measurements should be less or about  $10 \text{ m s}^{-1}$  which can be achieved with the existing radio astronomical facilities in observations of objects in the local Universe ( $z \lesssim 1$ ).

## Bibliography

- [1] Milne, E.A.: Kinematics, dynamics, and the scale of time. Proc. Roy. Soc. A. 158, 324-348 (1937). doi:10.1098/rspa.1937.0023
- [2] Dirac, P.A.M.: The cosmological constants. Nature. 139, 323 (1937). doi:10.1038/139323a0
- [3] Uzan, J.-P.: Varying constants, gravitation and cosmology. Living Reviews Relativity. 14, 2-155 (2011). doi:10.12942/lrr-2011-2
- [4] Kujat, J., Scherrer, R.J.: Effect of time variation in the Higgs vacuum expectation value on the cosmic microwave background. Phys. Rev. D. 62, 023510 (2000). doi:10.1103/PhysRevD.62.023510
- [5] Blatt, S., et al.: New limits on coupling of fundamental constants to gravity using Sr87 optical lattice clocks. Phys. Rev. Lett. 100, 140801 (2008). doi:10.1103/PhysRevLett.100.140801
- [6] M.C. Ferreira, M.D. Julião, C.J.A.P. Martins, A.M.R.V.L. Monteiro.: Probing unification scenarios with atomic clocks. Phys. Rev. D. 86, 125025 (2012). doi:10.1103/PhysRevD.86.125025
- [7] Schwarz, R. et al.: Long term measurement of the Sr87 clock frequency at the limit of primary Cs clocks. Phys. Rev. Res. 2, 033243 (2020). doi:10.1103/PhysRevResearch.2.033243
- [8] Molaro, P., Reimers, D., Agafonova, I.I., Levshakov, S.A.: Bounds on the fine structure constant variability from FeII absorption lines in QSO spectra. European Phys. J. Special Topics. 163, 173-189 (2008). doi:10.1140/epjst/e2008-00818-4
- [9] Agafonova, I.I., Molaro, P., Levshakov, S.A., Hou, J.L.: First measurement of Mg isotope abundances at high redshifts and accurate estimate of  $\Delta\alpha/\alpha$ . Astron. Astrophys. 529, A28 (2011). doi:10.1051/0004-6361/201016194
- [10] Kanekar, N., et al.: Constraints on changes in the proton-to-electron mass ratio using methanol lines. Mon. Not. R. Astron. Soc. 448, L104-L108 (2015). doi:10.1093/mnras/slu206

- [11] Levshakov, S.A., et al.: Testing the weak equivalence principle by differential measurements of fundamental constants in the Magellanic Clouds. *Mon. Not. R. Astron. Soc.* 487, 5175-5187 (2019). doi:10.1093/mnras/stz1628
- [12] Lee, C.-C., Webb, J.K., Carswell, R.F., Dinko, M.: Artificial intelligence and quasar absorption system modeling; application to fundamental constants at high redshift. Eprint arXiv: 2008.02583 (2020).
- [13] Marconi, A., et al.: ELT-HIRES, the high resolution spectrograph for the ELT: results from the Phase A study. In: Proceedings of the SPIE. 10702, 107021 (2018). doi:10.1117/12.2311664
- [14] Hart, L., Chluba, J.: Updated fundamental constant constraints from Planck 2018 data and possible relations to the Hubble tension. *Mon. Not. R. Astron. Soc.* 493, 3255-3263 (2020). doi:10.1093/mnras/staa412
- [15] Planck Collaboration; Aghanim, N., et al.: Planck 2018 results. VI. Cosmological parameters. *Astron. Astrophys.* 641, A6 (2020). doi:10.1051/0004-6361/201833910
- [16] Riess, A.G., Casertano, S., Yuan, W., Macri, L.M., Scolnic, D.: Large Magellanic Cloud Cepheid standards provide a 1% foundation for the determination of the Hubble constant and stronger evidence for physics beyond  $\Lambda$ CDM. *ApJ.* 876, 85 (2019). doi:10.3847/1538-4357/ab1422
- [17] Yoo, J.J., Scherrer, R.J.: Big bang nucleosynthesis and cosmic microwave background constraints on the time variation of the Higgs vacuum expectation value. *Phys. Rev. D.* 67, 043517 (2003). doi:10.1103/PhysRevD.67.043517
- [18] Thompson, R.I.: The determination of the electron to proton inertial mass via molecular transitions. *Astrophys. Lett.* 16, 3-4 (1975).
- [19] Varshalovich, D.A., Levshakov, S.A.: On a time dependence of physical constants. *JETP Lett.* 58, 237-240 (1993).
- [20] Ubachs, W., Buning, R., Eikema, K.S.E., Reinhold, E.: On a possible variation of the proton-to-electron mass ratio: H<sub>2</sub> spectra in the line of sight of high-redshift quasars and in the laboratory. *J. Mol. Spec.* 241, 155-179 (2007). doi:10.1016/JMoSp.241..155U
- [21] Levshakov, S.A., Varshalovich, D.A.: Molecular hydrogen in the z=2.811 absorbing material toward the quasar PKS 0528-250. *Mon. Not. R. Astron. Soc.* 212, 517-521 (1985). doi:10.1093/mnras/212.3.517
- [22] Balashev, S.A., et al.: Nature of the DLA towards Q 0528-250: high pressure and strong UV field revealed by excitation of CI, H<sub>2</sub>, and SiII. *Mon. Not. R. Astron. Soc.* 497, 1946-1956 (2020). doi:10.1093/mnras/staa2108
- [23] Levshakov, S.A., Kozlov, M.G., Agafonova, I.I.: Constraints on the electron-to-proton mass ratio variation at the epoch of reionization. *Mon. Not. R. Astron. Soc.* 498, 3624-3632 (2020). doi:10.1093/mnras/staa2635

- [24] Kozlov, M.G., Levshakov, S.A.: Microwave and submillimeter molecular transitions and their dependence on fundamental constants. *Annalen der Physik.* 525, 453-471 (2013). doi:10.1002/andp.201300010
- [25] Flambaum, V.V., Kozlov, M.G.: Limit on the cosmological variation of  $m_p/m_e$  from the inversion spectrum of ammonia. *Phys. Rev. Lett.* 98, 240801 (2007). doi:10.1103/PhysRevLett.98.240801
- [26] Levshakov, S.A., et al.: Limits on the spatial variations of the electron-to-proton mass ratio in the Galactic plane. *Astron. Astrophys.* 559, A91 (2013). doi:10.1051/0004-6361/201322535
- [27] Jansen, P., et al.: Methanol as a sensitive probe for spatial and temporal variations of the proton-to-electron mass ratio. *Phys. Rev. Lett.* 106, 100801 (2011). doi:10.1103/PhysRevLett.106.100801
- [28] Levshakov, S.A., Kozlov, M.G., Reimers, D.: Methanol as a tracer of fundamental constants. *ApJ.* 738, 26-30 (2011). doi:10.1088/0004-637X/738/1/26
- [29] Dapprà, M., et al.: Testing the variability of the proton-to-electron mass ratio from observations of methanol in the dark cloud core L1498. *Mon. Not. R. Astron. Soc.* 472, 4434-4443 (2017). doi:10.1093/mnras/stx2308
- [30] Levshakov, S.A., et al.: A new approach for testing variations of fundamental constants over cosmic epochs using FIR fine-structure lines. *Astron. Astrophys.* 479, 719-723 (2008). doi:10.1051/0004-6361:20079116