

PRERESOLVING SUBCATEGORIES AND DERIVED EQUIVALENCES

RUBEN HENRARD AND ADAM-CHRISTIAAN VAN ROOSMALEN

ABSTRACT. It is well known that a resolving subcategory \mathcal{A} of an abelian subcategory \mathcal{E} induces several derived equivalences: the equivalence $\mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{E})$ holds in general and furthermore restricts to equivalence $\mathbf{D}^b(\mathcal{A}) \rightarrow \mathbf{D}^b(\mathcal{E})$ if $\text{res. dim}_{\mathcal{A}}(E) < \infty$ for any object $E \in \mathcal{E}$. If the category \mathcal{E} is uniformly bounded, i.e. $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) < \infty$, one obtains a derived equivalence $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$.

In this paper, we show that all of the above statements hold for preresolving subcategories of (one-sided) exact categories. By passing to a one-sided language, one can remove the assumption that $\mathcal{A} \subseteq \mathcal{E}$ is extension-closed completely from the classical setting, yielding easier criteria and more examples. To illustrate this point, we consider the Isbell category \mathcal{I} and show that $\mathcal{I} \subseteq \mathbf{Ab}$ is preresolving but \mathcal{I} cannot be realized as an extension-closed subcategory of an exact category.

We also consider a criterion given by Keller to produce derived equivalences of fully exact subcategories. We show that this criterion fits into the framework of preresolving subcategories by considering the relative weak idempotent completion of said subcategory.

CONTENTS

1. Introduction	1
2. Preliminaries	2
2.1. One-sided exact categories	2
2.2. Derived categories of one-sided exact categories	4
3. Preresolving subcategories and bounded derived equivalences	4
3.1. Definitions and basic properties	4
3.2. Induced bounded derived equivalences	6
4. The resolution dimension and the unbounded derived equivalence	7
5. Examples and applications	10
5.1. Examples	10
5.2. Application: the exact hull	10
6. A comparison to Keller's theorem	11
References	13

1. INTRODUCTION

A basic technique in homological algebra is replacing an object in an abelian or exact category with a well-chosen resolution. For example, if an abelian category \mathcal{E} has enough projective objects, one can replace any object $E \in \mathcal{E}$ with a projective resolution $P^\bullet \rightarrow E$. The existence of such resolutions provides a triangle equivalence $\mathbf{K}^-(\text{Proj } \mathcal{E}) \rightarrow \mathbf{D}^-(\mathcal{E})$.

The notion of a resolving subcategory [2] generalizes this setting: a resolving subcategory \mathcal{A} of an abelian or exact category \mathcal{E} allows one to take resolutions of all objects in \mathcal{E} by objects in \mathcal{A} . For example, if S is a ring, then the subcategory \mathcal{F} of flat S -modules is a resolving subcategory of $\text{Mod } S$. As \mathcal{F} is extension-closed in $\text{Mod } S$, the category \mathcal{F} inherits an exact structure (so that the derived category of \mathcal{F} is defined, see [20]) and we again find a triangle equivalence $\mathbf{D}^-(\mathcal{F}) \rightarrow \mathbf{D}^-(\text{Mod } S)$.

The definition of a preresolving subcategory \mathcal{A} of \mathcal{E} is obtained from the definition of a resolving subcategory by removing the assumption that \mathcal{A} is extension-closed in \mathcal{E} . Specifically, let \mathcal{E} be an exact category. A full additive subcategory \mathcal{A} of \mathcal{E} is called *preresolving* if \mathcal{A} is closed under kernels of deflations (we say that \mathcal{A} lies *deflation-closed* in \mathcal{E} , see Definition 3.1), and for every $E \in \mathcal{E}$ there is a deflation $A \rightarrow E$ for some $A \in \mathcal{A}$.

Given a preresolving subcategory $\mathcal{A} \subseteq \mathcal{E}$, we may attach an \mathcal{A} -resolution dimension to each object $E \in \mathcal{E}$ as the smallest number $n \in \mathbb{N} \cup \{\infty\}$ such that there is an exact sequence $0 \rightarrow A^{-n} \rightarrow A^{-n-1} \rightarrow$

2020 *Mathematics Subject Classification*. Primary: 18E05, 18G10, 18G80.

Key words and phrases. (Pre)resolving subcategories, (one-sided) exact categories, derived categories.

$\cdots \rightarrow A^0 \rightarrow E$ with each $A^i \in \mathcal{A}$. We write $\text{res. dim}_{\mathcal{A}}(\mathcal{E})$ for the supremum of $\text{res. dim}_{\mathcal{A}}(E)$, ranging over all $E \in \mathcal{E}$.

As \mathcal{A} need not be extension-closed in \mathcal{E} , the category \mathcal{A} need not inherit an exact structure from \mathcal{E} . However, using the fact that \mathcal{A} lies deflation-closed in \mathcal{E} , the subcategory \mathcal{A} still inherits a rich homological structure from \mathcal{E} : it is a deflation-exact category (see Proposition 3.6 in the text). Deflation-exact categories still admit a natural (bounded) derived category [3, 11], and we obtain the following theorem (see Theorem 3.11 in the text).

Theorem 1.1. *Let \mathcal{A} be a preresolving subcategory of a (deflation-)exact category \mathcal{E} .*

- (1) *The embedding $\mathcal{A} \subseteq \mathcal{E}$ lifts to a triangle equivalence $\mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{E})$.*
- (2) *If $\text{res. dim}_{\mathcal{A}}(E) < \infty$ for all $E \in \mathcal{E}$, then the embedding lifts to a triangle equivalence $\mathbf{D}^b(\mathcal{A}) \rightarrow \mathbf{D}^b(\mathcal{E})$.*
- (3) *If $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) < \infty$, then the embedding lifts to a triangle equivalence $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$.*

Similar results can be found in various places in the literature (see for example [10, Lemma I.4.6], [16, Proposition 13.2.2], [28, Proposition 5.14] or [29, Lemma 3.8] and Remark 3.14 for a short comparison). An example of a resolving subcategory of an abelian category \mathcal{E} is given by a cotilting torsion-free class \mathcal{F} ; in this case, $\text{res. dim}_{\mathcal{F}}(\mathcal{E}) \leq 1$. In this setting, Theorem 1.1 can be recovered from [6, Appendix B], generalizing the well-known tilting construction from [9, Theorem 3.3]. Theorem 1.1 allows us to generalize this equivalence to the case where \mathcal{F} is a cotilting pretorsion-free class in \mathcal{E} . The examples in §5 are of this form.

In Theorem 1.1, the preresolving subcategory \mathcal{A} is endowed with the following set of conflations: a sequence $A \rightarrow B \rightarrow C$ in \mathcal{A} is a conflation in \mathcal{A} if and only if it is a conflation in \mathcal{E} . In [18, Theorem 12.1], a variant of Theorem 1.1 was shown for subcategories \mathcal{A} which are not deflation-closed subcategories of \mathcal{E} , and hence, not resolving. Removing the assumption of being deflation-closed, it is not immediately clear how to obtain a meaningful deflation-exact structure on \mathcal{A} .

In section 6, we describe a more involved conflation structure on \mathcal{A} (namely, we endow \mathcal{A} with the largest deflation-exact structure such that the embedding $\mathcal{A} \rightarrow \mathcal{E}$ maps conflations to conflations). Modifying condition **C2** accordingly, we find a version of [18, Theorem 12.1] for subcategories \mathcal{A} which are not necessarily fully exact in \mathcal{E} .

Theorem 1.2. *Let \mathcal{E} be a deflation-exact category and let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory satisfying axioms **C1** and **C2'**. If we endow \mathcal{A} with the largest deflation-exact structure such that the embedding $\mathcal{A} \rightarrow \mathcal{E}$ maps conflations in \mathcal{A} to conflations in \mathcal{E} , then the obvious functor $\mathbf{D}^-(\mathcal{A}) \xrightarrow{\Phi} \mathbf{D}^-(\mathcal{E})$ is a triangle equivalence.*

Moreover, if $\text{res. dim}_{\mathcal{A}}(E) < \infty$ for all $E \in \mathcal{E}$, the functor $\mathbf{D}^b(\mathcal{A}) \rightarrow \mathbf{D}^b(\mathcal{E})$ is an equivalence.

When \mathcal{E} is exact and \mathcal{A} is a fully exact subcategory, then Theorem 1.2 reduces to [18, Theorem 12.1].

Acknowledgements. The authors wish to thank Francesco Genovese and Jan Šťovíček for their useful comments and references. The second author is currently a postdoctoral researcher at FWO (12.M33.16N).

2. PRELIMINARIES

In these preliminaries, we summarize results of [3, 11, 12] in a convenient form.

2.1. One-sided exact categories.

Definition 2.1. (1) A *conflation category* is an additive category \mathcal{E} together with a chosen class of kernel-cokernel pairs, called *conflations*, such that this class is closed under isomorphisms. The kernel part of a conflation is called an *inflation* and the cokernel part of a conflation is called a *deflation*. We depict inflations by the symbol \rightrightarrows and deflations by \dashrightarrow .

- (2) An additive functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between conflation categories is called (*conflation-*)*exact* if conflations are mapped to conflations.
- (3) A map $f: X \rightarrow Y$ in a conflation category \mathcal{E} is called *admissible* if f admits a deflation-inflation factorization, i.e. f factors as $X \dashrightarrow I \rightrightarrows Y$. An admissible morphism is depicted by $\dashrightarrow \rightrightarrows$.
- (4) A cochain complex (X^\bullet, d^\bullet) in $\mathbf{C}(\mathcal{E})$ with \mathcal{E} a conflation category is called *acyclic* or *exact* if, for each $i \in \mathbb{Z}$, the map $d_X^i: X^i \rightarrow X^{i+1}$ is admissible and $\ker(d_X^{i+1}) \cong \text{im}(d_X^i)$.

Remark 2.2. Let \mathcal{E} be a conflation category and let $f: X \rightarrow Y$ be a map. If f is admissible, then the deflation-inflation factorization is unique up to isomorphism. Indeed, given a deflation-inflation factorization $X \dashrightarrow I \rightrightarrows Y$ of f , it is clear that f admits a kernel and a cokernel, moreover, $\text{coker}(\ker(f)) = \text{coim}(f) \cong I \cong \text{im}(f) = \ker(\text{coker}(f))$.

Definition 2.3. A *deflation-exact category* \mathcal{E} is a conflation category satisfying the following axioms:

- R0** For each $X \in \mathcal{E}$, the map $X \rightarrow 0$ is a deflation.
- R1** The composition of two deflations is a deflation.
- R2** The pullback of a deflation along any morphism exists and deflations are stable under pullbacks.

Dually, an *inflation-exact category* is a conflation category \mathcal{E} such that the inflations satisfy the following axioms:

- L0** For each $X \in \mathcal{E}$, the map $0 \rightarrow X$ is an inflation.
- L1** The composition of two inflations is an inflation.
- L2** The pushout of an inflation along any morphism exists and inflations are stable under pushouts.

Definition 2.4. Let \mathcal{E} be a conflation category. In addition to the properties listed in definition 2.3, we also consider the following axioms:

- R3** If $i: A \rightarrow B$ and $p: B \rightarrow C$ are morphisms in \mathcal{E} such that p has a kernel and pi is a deflation, then p is a deflation.
- R3⁺** If $i: A \rightarrow B$ and $p: B \rightarrow C$ are morphisms in \mathcal{E} such that pi is a deflation, then p is a deflation.

The axioms **L3** and **L3⁺** are defined dually. A deflation-exact category satisfying **R3** is called *strongly deflation-exact*. Dually, an inflation-exact category satisfying axiom **L3** is called a strongly inflation-exact category.

Remark 2.5. (1) Inflation-exact categories have also been called left exact categories in [3, 25] and right exact categories in [24]. Similarly, deflation-exact categories have also been referred to as right exact or left exact categories. As the use of left and right is not consistent, we prefer to use the above terminology.

- (2) A deflation-exact category has the structure of a Grothendieck pretopology (see for example [15, 24]).
- (3) Axiom **R0** above is axiom **R0*** in [3, 12]. This axiom ensures that all split kernel-cokernel pairs are conflations in a deflation-exact category (this is also required in [25]).
- (4) An exact category in the sense of Quillen (see [23]) is a conflation category \mathcal{E} satisfying axioms **R0** through **R3** and **L0** through **L3**. In [17, Appendix A], Keller shows that axioms **R0**, **R1**, **R2**, and **L2** suffice to define an exact category.
- (5) Axioms **R3** and **L3** are sometimes referred to as Quillen's *obscure axioms* (see [7, 27]).
- (6) Axiom **R3⁺** implies axiom **R3**, dually, axiom **L3⁺** implies axiom **L3**.
- (7) For a deflation-exact category \mathcal{E} , axiom **R3⁺** is equivalent to \mathcal{E} being weakly idempotent complete and satisfying axiom **R3**.

The following proposition is used multiple times throughout this text (see [3, Proposition 5.7] or [12, Proposition 3.7]).

Proposition 2.6. *Let \mathcal{E} be a deflation-exact category. For a commutative diagram*

$$\begin{array}{ccc} Y' & \xrightarrow{p'} & Z' \\ \downarrow f & & \downarrow g \\ Y & \xrightarrow{p} & Z \end{array}$$

whose horizontal arrows are deflation, the following are equivalent:

- (1) the square is a pullback square;
- (2) the square is bicartesian, i.e. both a pullback and pushout square;
- (3) the induced sequence $Y' \xrightarrow{\begin{pmatrix} f \\ -p' \end{pmatrix}} Y \oplus Z' \xrightarrow{(p \ g)} Z$ is a conflation.

The following theorem (see [14, Theorem 1.1 and Theorem 1.2]) highlights the importance of axioms **R3** and **R3⁺**.

Theorem 2.7. *Let \mathcal{E} be a deflation-exact category.*

- (1) *The category \mathcal{E} satisfies axiom **R3** if and only if the Nine Lemma holds.*
- (2) *The category \mathcal{E} satisfies axiom **R3⁺** if and only if the Snake Lemma holds.*

2.2. Derived categories of one-sided exact categories. Let \mathcal{E} be a one-sided exact category. The following is a small extension of [3, Lemma 7.2].

Lemma 2.8. *The mapping cone $\text{cone}(f^\bullet)$ of a map $f^\bullet: X^\bullet \rightarrow Y^\bullet$ between acyclic complexes $X^\bullet, Y^\bullet \in \mathbf{C}(\mathcal{E})$ is acyclic. Furthermore, for each $n \in \mathbb{Z}$, there are conflations:*

$$\text{im}(d_X^{n-1}) \twoheadrightarrow X^n \oplus \text{im}(d_X^{n-1}) \twoheadrightarrow \text{im}(d_{\text{cone}(f)}^{n-1}) \quad \text{and} \quad \text{im}(d_{\text{cone}(f)}^{n-1}) \twoheadrightarrow Y^n \oplus \text{im}(d_X^n) \twoheadrightarrow \text{im}(d_Y^n).$$

Proof. The proof is as in [3, Lemma 7.2], where it was noted that, for each $n \in \mathbb{Z}$, there is a commutative diagram:

$$\begin{array}{ccccc} \text{im}(d_X^{n-1}) & \twoheadrightarrow & X^n & \twoheadrightarrow & \text{im}(d_X^n) \\ \downarrow & \text{BC} & \downarrow & & \parallel \\ \text{im}(d_Y^{n-1}) & \twoheadrightarrow & \text{im}(d_{\text{cone}(f)}^{n-1}) & \twoheadrightarrow & \text{im}(d_Y^n) \\ \parallel & & \downarrow & \text{BC} & \downarrow \\ \text{im}(d_Y^{n-1}) & \twoheadrightarrow & Y^n & \twoheadrightarrow & \text{im}(d_Y^n) \end{array}$$

where the rows are conflations and the squares labeled BC are bicartesian. The required conflations are obtained by [12, Proposition 3.7]. \square

Write $\mathbf{Ac}(\mathcal{E}) \subseteq \mathbf{C}(\mathcal{E})$ for the unbounded acyclic complexes. Analogous to exact categories [20], one can define the unbounded derived category $\mathbf{D}(\mathcal{E})$ as the Verdier localization $\mathbf{K}(\mathcal{E}) / \langle \mathbf{Ac}(\mathcal{E}) \rangle_{\text{thick}}$ of the homotopy category $\mathbf{K}(\mathcal{E})$ by the thick closure of the triangulated subcategory of acyclic complexes. Note that $\mathbf{D}(\mathcal{E})$ is the localization of $\mathbf{K}(\mathcal{E})$ with respect to the saturated multiplicative system \mathcal{S} of those maps $f^\bullet \in \text{Mor}(\mathbf{K}(\mathcal{E}))$ such that $\text{cone}(f^\bullet) \in \langle \mathbf{Ac}(\mathcal{E}) \rangle_{\text{thick}}$. The right bounded derived category $\mathbf{D}^r(\mathcal{E})$ and bounded derived category $\mathbf{D}^b(\mathcal{E})$ can be defined analogously.

Convention 2.9. We use the following convention: a map $f^\bullet \in \text{Mor}(\mathbf{C}(\mathcal{E}))$ such that $\text{cone}(f^\bullet) \in \mathbf{Ac}(\mathcal{E})$ is called a *quasi-isomorphism*. Note that \mathcal{S} is simply the multiplicative system in $\mathbf{K}(\mathcal{E})$ obtained as the saturated closure of the quasi-isomorphisms, in particular, the derived category $\mathbf{D}(\mathcal{E})$ is equivalent to the localization $\mathbf{K}(\mathcal{E})[\mathcal{S}^{-1}]$.

Following the above convention, a composition of quasi-isomorphisms need not be a quasi-isomorphism but it does produce a morphism in $\mathcal{S} \subseteq \text{Mor}(\mathbf{K}(\mathcal{E}))$.

The following proposition summarizes some useful properties of the derived category (see [11]).

Proposition 2.10. *Let \mathcal{E} be a deflation-exact category and let $* \in \{-, b\}$*

- (1) *The natural embedding $i: \mathcal{E} \rightarrow \mathbf{D}(\mathcal{E})$ is fully faithful.*
- (2) *For all $X, Y \in \mathcal{E}$ and $n > 0$, $\text{Hom}_{\mathbf{D}(\mathcal{E})}(\Sigma^n iX, iY) = 0$.*
- (3) *Every conflation $X \twoheadrightarrow Y \twoheadrightarrow Z$ in \mathcal{E} maps to a triangle $iX \rightarrow iY \rightarrow iZ \rightarrow \Sigma iX$ in $\mathbf{D}(\mathcal{E})$.*
- (4) *Axiom **R3** holds if and only if every sequence $X \rightarrow Y \rightarrow Z$ that lifts to a triangle in $\mathbf{D}(\mathcal{E})$ is a conflation in \mathcal{E} .*
- (5) *The category $\mathbf{Ac}^*(\mathcal{E})$ is a thick triangulated subcategory of $\mathbf{K}^*(\mathcal{E})$ if and only if \mathcal{E} satisfies axiom **R3**⁺.*
- (6) *The category $\mathbf{Ac}(\mathcal{E})$ is a thick triangulated subcategory of $\mathbf{K}(\mathcal{E})$ if and only if \mathcal{E} is idempotent complete and satisfies axiom **R3**.*

3. PRERESOLVING SUBCATEGORIES AND BOUNDED DERIVED EQUIVALENCES

Throughout this section, \mathcal{E} denotes a deflation-exact category (all results can be dualized to inflation-exact categories, see Remark 3.4).

3.1. Definitions and basic properties. We start by recalling the notion of a preresolving subcategory and the resolution dimension (see [28, Definition 2.1] or [26, Definition 2]).

Definition 3.1. A full additive subcategory $\mathcal{A} \subseteq \mathcal{E}$ is called *preresolving* if the following two conditions are met:

PR1 For every $E \in \mathcal{E}$ there exists a deflation $A \twoheadrightarrow E$ with $A \in \mathcal{A}$.

PR2 The subcategory $\mathcal{A} \subseteq \mathcal{E}$ is *deflation-closed*, i.e. for every conflation $X \twoheadrightarrow Y \twoheadrightarrow Z$, if $Y, Z \in \mathcal{A}$, then $X \in \mathcal{A}$.

Let $\mathcal{A} \subseteq \mathcal{E}$ be an additive subcategory satisfying axiom **PR1**. The *resolution dimension* of an object $E \in \mathcal{E}$ with respect to \mathcal{A} , denoted $\text{res. dim}_{\mathcal{A}}(E)$, is the smallest integer $n \geq 0$ such that there exists an exact sequence

$$0 \rightarrow A^{-n} \rightarrow A^{-n+1} \rightarrow \cdots \rightarrow A^0 \rightarrow E \rightarrow 0$$

in \mathcal{E} with each $A^i \in \mathcal{A}$. If such an n does not exist, we write $\text{res. dim}_{\mathcal{A}}(E) = \infty$.

Furthermore, we define $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) = \sup\{\text{res. dim}_{\mathcal{A}}(E) \mid \forall E \in \mathcal{E}\} \in \mathbb{Z}^+ \cup \{\infty\}$.

A preresolving subcategory $\mathcal{A} \subseteq \mathcal{E}$ is called *finitely* preresolving if for each object $E \in \mathcal{E}$, $\text{res. dim}_{\mathcal{A}}(E) < \infty$, and is called *uniformly* preresolving subcategory if $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) < \infty$. Furthermore, a preresolving subcategory $\mathcal{A} \subseteq \mathcal{E}$ is called *resolving* if $\mathcal{A} \subseteq \mathcal{E}$ is extension-closed.

Remark 3.2. It follows from axiom **PR1** that each object in \mathcal{E} has a resolution by objects in \mathcal{A} .

Notation 3.3. For an object $E \in \mathcal{E}$, we often write $A^\bullet \rightarrow E$ for the complex $\cdots \rightarrow A^{-n} \rightarrow A^{-n+1} \rightarrow \cdots \rightarrow A^0 \rightarrow E \rightarrow 0 \rightarrow \cdots$. Here, we assume that E has degree 1.

Remark 3.4. The notions of a coresolving and a precoresolving subcategory of an exact or inflation-exact category are dual.

Remark 3.5. The category \mathcal{P} of projectives of [4, Example 4.10] is a finitely resolving subcategory of $\text{mod}(\mathfrak{b})$, but not uniformly resolving.

Proposition 3.6. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a deflation-closed subcategory. The category \mathcal{A} inherits a deflation-exact structure from \mathcal{E} (the conflations are simply those conflations in \mathcal{E} that lie in \mathcal{A}). Moreover, if \mathcal{E} satisfies axiom **R3⁺** or **R3**, then \mathcal{A} satisfies axiom **R3⁺** or **R3**, respectively.*

Proof. Axioms **R0** and **R1** follow directly from the definition. To show axiom **R2**, let $X \twoheadrightarrow Y \twoheadrightarrow Z$ be a conflation in \mathcal{A} and let $A \rightarrow Z$ be any map in \mathcal{A} . Applying axiom **R2** in \mathcal{E} , we obtain the following commutative diagram:

$$\begin{array}{ccccc} X & \twoheadrightarrow & P & \twoheadrightarrow & A \\ \parallel & & \downarrow & & \downarrow \\ X & \twoheadrightarrow & Y & \twoheadrightarrow & Z \end{array}$$

By Proposition 2.6, the induced kernel-cokernel pair $P \twoheadrightarrow A \oplus Y \twoheadrightarrow Z$ is a conflation in \mathcal{E} . As $\mathcal{A} \subseteq \mathcal{E}$ is deflation-closed, it follows that $P \in \mathcal{A}$ as required.

That axiom **R3⁺** is preserved is straightforward to show. That axiom **R3** is preserved is shown in [12, Proposition 3.11]. \square

If $\mathcal{A} \subseteq \mathcal{E}$ is deflation-closed, Proposition 3.6 endows \mathcal{A} with a natural deflation-exact structure; in this case, the derived category $\mathbf{D}(\mathcal{A})$ has been defined in [3]. It is clear that $\mathbf{Ac}(\mathcal{A}) \subseteq \mathbf{Ac}(\mathcal{E})$; however, in general, $\mathbf{Ac}(\mathcal{A}) \neq \mathbf{C}(\mathcal{A}) \cap \mathbf{Ac}(\mathcal{E})$. Thus, a complex $A^\bullet \in \mathbf{C}(\mathcal{A})$ which is acyclic in \mathcal{E} need not be acyclic in \mathcal{A} , see Remark 3.10 below. For this reason, we make the following distinction.

Definition 3.7. As in Convention 2.9, let \mathcal{S} denote the collection of morphisms $f^\bullet \in \text{Hom}_{\mathbf{C}(\mathcal{E})}(X^\bullet, Y^\bullet)$ such that $\text{cone}(f^\bullet) \in \mathbf{Ac}(\mathcal{E})$. If $\mathcal{A} \subseteq \mathcal{E}$ is a deflation-closed subcategory, we write \mathcal{T} for those morphisms $f^\bullet \in \text{Hom}_{\mathbf{C}(\mathcal{A})}(A^\bullet, B^\bullet)$ such that $\text{cone}(f^\bullet) \in \mathbf{Ac}(\mathcal{A})$. The bounded variants \mathcal{T}^- , \mathcal{T}^b , \mathcal{S}^- and \mathcal{S}^b are defined analogously.

Remark 3.8. If $\mathcal{A} \subseteq \mathcal{E}$ is deflation-closed, and hence deflation-exact by Proposition 3.6, then $\mathbf{D}(\mathcal{A}) = \mathbf{K}(\mathcal{A})[\mathcal{T}^{-1}]$.

Proposition 3.9. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a deflation-closed subcategory. Then $\mathbf{Ac}^-(\mathcal{E}) \cap \mathbf{C}^-(\mathcal{A}) = \mathbf{Ac}^-(\mathcal{A})$. In particular, $\mathcal{T}^- = \mathcal{S}^- \cap \text{Mor}(\mathbf{K}^-(\mathcal{A}))$ and $\mathcal{T}^b = \mathcal{S}^b \cap \text{Mor}(\mathbf{K}^b(\mathcal{A}))$.*

Proof. Let $A^\bullet \in \mathbf{C}^-(\mathcal{A})$. Without loss of generality, assume that $A^k = 0$ for all $k > 0$. It suffices to show that A^\bullet is acyclic in \mathcal{A} if and only if A^\bullet is acyclic in \mathcal{E} .

If $A^\bullet \in \mathbf{Ac}(\mathcal{A})$, then we have that $A^\bullet \in \mathbf{Ac}(\mathcal{E})$ (as Proposition 3.6 yields that the embedding $\mathcal{A} \rightarrow \mathcal{E}$ is conflation-exact). Conversely, assume that $A^\bullet \in \mathbf{Ac}(\mathcal{E})$. In this case, $d_A^{-1}: A^{-1} \rightarrow A^0$ is a deflation and hence its kernel $K^{-1} \twoheadrightarrow A^{-1}$ belongs to \mathcal{A} as $\mathcal{A} \subseteq \mathcal{E}$ is deflation-closed. Moreover, since the complex A^\bullet is acyclic, we have $K^{-1} \cong \text{im}(d_A^{-2})$. Repeating this argument on the deflation $A^{-2} \twoheadrightarrow \text{im}(d_A^{-2})$, one finds that $\text{im}(d_A^{-3}) \in \mathcal{A}$. Proceeding in this fashion yields that A^\bullet is acyclic in \mathcal{A} . \square

Remark 3.10. Proposition 3.9 does not need to hold for unbounded complexes. Indeed, consider the category $\mathcal{E} = \text{mod}(A)$ of finitely generated k -modules over the k -algebra $A = k[X]/(X^2)$ and let $\mathcal{A} = \text{add}(\mathcal{E})$ be the full additive subcategory generated by A . The complex

$$\dots \xrightarrow{\cdot X} A \xrightarrow{\cdot X} A \xrightarrow{\cdot X} A \xrightarrow{\cdot X} \dots$$

belongs to \mathcal{A} and is acyclic in \mathcal{E} . Clearly, the images of the differentials do not belong to \mathcal{A} and thus the above sequence is not acyclic in \mathcal{A} .

3.2. Induced bounded derived equivalences. The goal of this section is to prove the following theorem.

Theorem 3.11. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory and write $\phi: \mathcal{A} \hookrightarrow \mathcal{E}$ for the embedding.*

- (1) *If $\mathcal{A} \subseteq \mathcal{E}$ is preresolving, the functor ϕ lifts to a triangle equivalence $\Phi: \mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{E})$.*
- (2) *If $\mathcal{A} \subseteq \mathcal{E}$ is finitely preresolving, the functor ϕ lifts to a triangle equivalence $\Phi: \mathbf{D}^b(\mathcal{A}) \rightarrow \mathbf{D}^b(\mathcal{E})$.*

That the above functors are essentially surjective is settled by the following lemma.

Lemma 3.12. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory satisfying axiom **PR1**.*

- (1) *For each $E^\bullet \in \mathbf{C}^-(\mathcal{E})$, there exists a complex $A^\bullet \in \mathbf{C}^-(\mathcal{A})$ and a map $f^\bullet: A^\bullet \rightarrow E^\bullet$ in $\mathbf{C}^-(\mathcal{E})$ such that $f^\bullet \in \mathcal{S}^-$.*
- (2) *Assume that $\text{res. dim}_{\mathcal{A}}(E) < \infty$ for all $E \in \mathcal{E}$. For each $E^\bullet \in \mathbf{C}^b(\mathcal{E})$, there exists a complex $A^\bullet \in \mathbf{C}^b(\mathcal{A})$ and a map $f^\bullet: A^\bullet \rightarrow E^\bullet$ in $\mathbf{C}^b(\mathcal{E})$ such that $f^\bullet \in \mathcal{S}^b$.*

Proof. We only show the right bounded case as the bounded case is similar.

Let $E \in \mathbf{C}^-(\mathcal{E})$. Without loss of generality we assume that $E^i = 0$ for all $i > 1$. By axiom **PR1**, there exists a deflation $A \twoheadrightarrow E^0$ with $A \in \mathcal{A}$. By axiom **R2**, the pullback P^{-1} of $d_E^{-1}: E^{-1} \rightarrow E^0$ along $A \twoheadrightarrow E^0$ exists. By the pullback property, we obtain the following commutative diagram:

$$\begin{array}{ccccccccccc} A_1^\bullet & & & \dots & \longrightarrow & E^{-2} & \longrightarrow & P^{-1} & \longrightarrow & A^0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & & & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \\ & & & & & E^{-2} & \longrightarrow & E^{-1} & \longrightarrow & E^0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & & & & \downarrow f_1^\bullet & & & & & & & & \end{array}$$

By Proposition 2.6, $\text{cone}(f_1^\bullet) \in \mathbf{Ac}(\mathcal{E})$ and thus f_1^\bullet is a quasi-isomorphism. Similarly, by axiom **PR1**, there is a deflation $A^{-1} \twoheadrightarrow P^{-1}$. Taking the pullback P^{-2} along $E^{-2} \rightarrow P^{-1}$, we obtain the following commutative diagram:

$$\begin{array}{ccccccccccc} A_2^\bullet & & & \dots & \longrightarrow & E^{-3} & \longrightarrow & P^{-2} & \longrightarrow & A^{-1} & \longrightarrow & A^0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & & & & \parallel & & \downarrow & & \downarrow & & \parallel & & & & \\ & & & & & E^{-3} & \longrightarrow & E^{-2} & \longrightarrow & P^{-1} & \longrightarrow & A^0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & & & & \downarrow f_2^\bullet & & & & & & & & & & \end{array}$$

As above, f_2^\bullet is a quasi-isomorphism. Iterating this procedure, one constructs a complex A^\bullet from right to left; moreover, the natural map $f^\bullet: A^\bullet \rightarrow E^\bullet$ belongs to \mathcal{S}^- as one can check degreewise. \square

We are now in a position to prove Theorem 3.11.

Proof of Theorem 3.11. We only prove the first statement, the second is similar.

As $\mathcal{A} \subseteq \mathcal{E}$ is deflation-closed, \mathcal{A} inherits a deflation-exact structure from \mathcal{E} by Proposition 3.6. Hence $\phi: \mathcal{A} \rightarrow \mathcal{E}$ is an exact functor and thus lifts to a triangle functor $\Phi: \mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{E})$.

That Φ is essentially surjective follows from Lemma 3.12. By Lemma 3.12 and Proposition 3.9, the conditions of (the dual of) [16, Proposition 7.2.1 (ii)] are satisfied and thus Φ is fully faithful. \square

Remark 3.13. That Φ is fully faithful also follows from [16, Proposition 10.2.7] or [22, Lemma in section 1.6].

Remark 3.14. Theorem 3.11 is akin to [29, Lemma 3.8] (or [16, Proposition 13.2.2]), i.e. given an abelian category \mathcal{E} and a full additive subcategory $\mathcal{A} \subseteq \mathcal{E}$ such that every object of \mathcal{E} admits a finite \mathcal{A} -resolution, the embedding lifts to a triangle equivalence $\mathbf{D}^b(\mathcal{A}; \mathcal{E}) \rightarrow \mathbf{D}^b(\mathcal{E})$. Here $\mathbf{D}^b(\mathcal{A}; \mathcal{E})$ is the derived category of \mathcal{A} relative to \mathcal{E} (see [29, Definition 3.6]), that is, $\mathbf{D}^b(\mathcal{A}; \mathcal{E}) := \mathbf{K}^b(\mathcal{A}) / \mathbf{Ac}^b(\mathcal{A}; \mathcal{E})$ where $\mathbf{Ac}^b(\mathcal{A}; \mathcal{E}) := \mathbf{K}^b(\mathcal{A}) \cap \mathbf{Ac}^b(\mathcal{E})$ are the bounded cochain complexes in \mathcal{A} which are acyclic in \mathcal{E} . If $\mathcal{A} \subseteq \mathcal{E}$ is finitely preresolving, Proposition 3.9 yields that $\mathbf{D}^b(\mathcal{A}; \mathcal{E}) \simeq \mathbf{D}^b(\mathcal{A})$ are triangle equivalent. As such, we recover [29, Lemma 3.8].

Remark 3.15. The bounded derived ∞ -category of a one-sided exact category is defined in [11], following [1]. Given an preresolving subcategory \mathcal{A} of a (one-sided) exact category \mathcal{E} , one obtains a functor $\Phi_\infty: \mathbf{D}_\infty^b(\mathcal{A}) \rightarrow \mathbf{D}_\infty^b(\mathcal{E})$ of stable ∞ -categories. If \mathcal{A} is finitely preresolving in \mathcal{E} , then Φ_∞ is an equivalence (this follows Theorem 3.11, using [21, Theorem I.3.3] or [5, Proposition 5.14]). In particular, for any additive or localizing invariant \mathbb{K} (in the sense of [5], such as non-connective K -theory) we have $\mathbb{K}(\mathbf{D}_\infty^b(\mathcal{A})) \cong \mathbb{K}(\mathbf{D}_\infty^b(\mathcal{E}))$.

4. THE RESOLUTION DIMENSION AND THE UNBOUNDED DERIVED EQUIVALENCE

Throughout this section, \mathcal{E} denotes a deflation-exact category. The goal of this section is to prove Theorem 4.1 below, which is the unbounded analogue of Theorem 3.11. The strategy of the proof is identical to the proof of Theorem 3.11. Lemma 3.12 directly generalizes to the unbounded setting with the additional assumption that $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) < \infty$ (see Lemma 4.2 below). However, Proposition 3.9 does not directly carry over to the unbounded setting. Nonetheless, Corollary 4.6 below provides a suitable generalization such that the proof of Theorem 3.11 carries over verbatim.

Theorem 4.1. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a uniformly preresolving subcategory. The embedding $\phi: \mathcal{A} \rightarrow \mathcal{E}$ lifts to a triangle equivalence $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$.*

The embedding $\mathcal{A} \rightarrow \mathcal{E}$ lifts to a triangle functor $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$ by Proposition 3.6. The first step of our proof is to establish that Φ is essentially surjective. This will be done in Lemma 4.2 below. The proof is an adaptation of [10, Lemma I.4.6] to the (one-sided) exact setting.

Lemma 4.2. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a preresolving subcategory with $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) = n \in \mathbb{N}$. Let $E^\bullet \in \mathbf{C}(\mathcal{E})$. There exists a map $f^\bullet: A^\bullet \rightarrow E^\bullet$ in $\mathbf{K}(\mathcal{E})$ such that $A^\bullet \in \mathbf{C}(\mathcal{A})$ and $f^\bullet \in \mathcal{S}$.*

Proof. Let $E^\bullet \in \mathbf{C}(\mathcal{E})$. From Lemma 3.12, we obtain the following commutative diagram

$$\begin{array}{cccccccccccc} A_1^\bullet & & \dots & \longrightarrow & A^{-2} & \longrightarrow & A^{-1} & \longrightarrow & A^0 & \longrightarrow & E^1 & \longrightarrow & E^2 & \longrightarrow & \dots \\ & & & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \parallel & & \\ E^\bullet & & \dots & \longrightarrow & E^{-2} & \longrightarrow & E^{-1} & \longrightarrow & E^0 & \longrightarrow & E^1 & \longrightarrow & E^2 & \longrightarrow & \dots \end{array}$$

such that $A^i \in \mathcal{A}$ and f_1^\bullet is a quasi-isomorphism. We provide an n -step procedure to construct a morphism $h^\bullet: A_2^\bullet \rightarrow A_1^\bullet$ in \mathcal{S} satisfying the following properties: $A_2^i \in \mathcal{A}$ for $i < 2$, and $h^i: A_1^i \rightarrow A_2^i$ is the identity whenever $i \notin \{-n, \dots, 0, 1\}$.

Let $C^\bullet \in \mathbf{C}^b(\mathcal{A})$ be a finite \mathcal{A} -resolution of E^1 exhibiting that $\text{res. dim}(E) \leq n$. Consider the following commutative diagram:

$$\begin{array}{ccccccccccccccc} & & & & & & & & \text{im}(d_C^{-1}) & & & & & & & \\ & & & & & & & & \downarrow & & & & & & & \\ \dots & \longrightarrow & A^{-2} & \longrightarrow & B^{-1} & \longrightarrow & B^0 & \longrightarrow & P & \longrightarrow & C^0 & \longrightarrow & E^2 & \longrightarrow & E^3 & \longrightarrow & \dots \\ & & \parallel & & \downarrow & \text{PB} & \downarrow & & \downarrow & \text{PB} & \downarrow & & \parallel & & \parallel & & \\ \dots & \longrightarrow & A^{-2} & \longrightarrow & A^{-1} & \longrightarrow & A^0 & \longrightarrow & A^0 & \longrightarrow & E^1 & \longrightarrow & E^2 & \longrightarrow & E^3 & \longrightarrow & \dots \end{array}$$

Here, the above diagram is constructed from right to left as follows: the squares labeled PB are pullback squares, which exist by axiom **R2**; the deflation $B^0 \rightarrow P$ (with $B^0 \in \mathcal{A}$) exists by axiom **PR1**. Note that $B^{-1} \in \mathcal{A}$ as it fits into the conflation $B^{-1} \rightarrow A^{-1} \oplus B^0 \rightarrow A^0$ and $\mathcal{A} \subseteq \mathcal{E}$ is deflation-closed.

Consider now the following commutative diagram:

$$\begin{array}{ccccccccccccccc} A_{1,-1}^\bullet & & \dots & \longrightarrow & A^{-2} & \longrightarrow & B^{-1} & \longrightarrow & \text{im}(d_C^{-1}) \oplus C^0 & \longrightarrow & C^0 & \longrightarrow & E^2 & \longrightarrow & \dots \\ & & & & \parallel & & \parallel & & \downarrow & \text{PB} & \downarrow & & \parallel & & \\ & & & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \\ & & & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \\ & & & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \\ A_1^\bullet & & \dots & \longrightarrow & A^{-2} & \longrightarrow & A^{-1} & \longrightarrow & A^0 & \longrightarrow & E^1 & \longrightarrow & E^2 & \longrightarrow & \dots \end{array}$$

Here, the top pullback square is a split pullback as $B^0 \rightarrow E^1$ factors through $C^0 \rightarrow E^1$. Note that all terms up to and including degree 1 of the complex $A_{1,-1}^\bullet$ belong to \mathcal{A} , with the possible exception of

$\text{im}(d_C^{-1}) \oplus C^0$. By construction, $\text{res. dim}(\text{im}(d_C^{-1}) \oplus C^0) < \text{res. dim}(E^1) \leq n$. Hence, iterating the above procedure (the next step is to now taking a strictly shorter resolution of $\text{im}(d_C^{-1}) \oplus C^0$), one obtains a complex $A_2^\bullet = A_{1,-n}^\bullet$ such that all terms belong to \mathcal{A} up to degree 1 and A_2^\bullet maps to A_1^\bullet via a map in \mathcal{S} . Furthermore, $A_2^{-n-l} = A_1^{-n-l}$ for all $l \geq 1$, $A_2^1 = C^0$ and $A_2^k = E^k$ for all $k \geq 2$. Iterating this entire procedure, one produces a string of morphisms $E^\bullet \leftarrow A_1^\bullet \leftarrow A_2^\bullet \leftarrow \dots$ in \mathcal{S} that stabilizes in each degree. The result follows. \square

In order to prove Theorem 4.1, it suffices to show that $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$ is fully faithful. The following proposition already indicates that in many cases, fully faithfulness readily holds.

Proposition 4.3. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a uniformly preresolving subcategory.*

- (1) *If $\mathcal{A} \subseteq \mathcal{E}$ is closed under normal subobjects, i.e. $X \in \mathcal{A}$ if $X \twoheadrightarrow A$ is an inflation with $A \in \mathcal{A}$, then the natural map $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$ is a triangle equivalence.*
- (2) *If $\mathcal{A} \subseteq \mathcal{E}$ is extension-closed and \mathcal{E} satisfies axiom **R3**, then the natural map $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$ is a triangle equivalence.*

Proof. As $\mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{E})$ is already essentially surjective by Lemma 4.2, it suffices to show that if $A^\bullet \in \mathbf{C}(\mathcal{A}) \cap \mathbf{Ac}(\mathcal{E})$, then $A^\bullet \in \mathbf{Ac}(\mathcal{A})$. We show this property in both cases.

- (1) Assume that $\mathcal{A} \subseteq \mathcal{E}$ is closed under normal subobjects. Let $A^\bullet \in \mathbf{C}(\mathcal{A}) \cap \mathbf{Ac}(\mathcal{E})$. Clearly $\text{im}(d_A^i) \in \mathcal{A}$ for all $i \in \mathbb{Z}$ as $\text{im}(d_A^i)$ is a normal subobject of A^{i+1} . The result follows.
- (2) Assume that $\mathcal{A} \subseteq \mathcal{E}$ is extension-closed and \mathcal{E} satisfies axiom **R3**. The reader may verify that the proof of [28, Proposition 2.3] only uses that $\mathcal{A} \subseteq \mathcal{E}$ is extension-closed and the Nine Lemma. By Theorem 2.7, the Nine Lemma holds in \mathcal{E} as \mathcal{E} satisfies axiom **R3**. Now let $A^\bullet \in \mathbf{C}(\mathcal{A}) \cap \mathbf{Ac}(\mathcal{E})$ and consider $\text{im}(d_A^i) \in \mathcal{E}$ for some $i \in \mathbb{Z}$. The following sequence yields a resolution of $\text{im}(d_A^i)$:

$$\text{im}(d_A^{i-n-1}) \rightarrow A^{i-n} \dots \rightarrow A^{i-1} \rightarrow A^i \rightarrow \text{im}(d_A^i) \rightarrow 0.$$

As $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) = n$, [28, Proposition 2.3] yields that $\text{im}(d_A^{i-n-1}) \in \mathcal{A}$. As i is chosen arbitrarily, $A^\bullet \in \mathbf{Ac}(\mathcal{A})$. The result follows. \square

In order to prove Theorem 4.1 without the additional assumptions from Proposition 4.3, more work is needed. We start with the following lemma.

Lemma 4.4. *Let \mathcal{A} be a preresolving subcategory in \mathcal{E} . Let $A^\bullet \rightarrow E$ and $B^\bullet \rightarrow E$ be resolutions of $E \in \mathcal{E}$ such that there is a commutative diagram*

$$\begin{array}{ccc} B^\bullet & \longrightarrow & E \\ \downarrow f^\bullet & & \parallel \\ A^\bullet & \longrightarrow & E \end{array}$$

If $\ker d_A^k \in \mathcal{A}$ for some $k \leq 0$, then $\ker d_B^k \in \mathcal{A}$ (where $d_A^0: A^0 \twoheadrightarrow E$). Moreover, $\text{cone}(f^\bullet) \in \mathbf{Ac}^-(\mathcal{A})$.

Proof. Assume that $\ker(d_A^k) \in \mathcal{A}$. Replacing the complex A^\bullet by the truncated complex $\tau^{\leq k-1} A^\bullet$ with $\ker(d_A^k)$ in degree $k-1$ and taking the cone of the above diagram, we obtain the following complex:

$$\dots \rightarrow B^{k-2} \rightarrow B^{k-1} \xrightarrow{\begin{pmatrix} -d_B^{k-1} \\ \alpha \end{pmatrix}} B^k \oplus \ker(d_A^k) \rightarrow B^{k+1} \oplus A^k \rightarrow \dots \rightarrow B^0 \oplus A^{-1} \rightarrow E \oplus A^0 \rightarrow E \rightarrow 0 \rightarrow \dots$$

By Lemma 2.8, the above complex is acyclic. It follows that $\text{coker}(-d_B^{k-2}) \cong \ker(-d_B^k)$ is isomorphic to

the image of the admissible map $B^{k-1} \xrightarrow{\begin{pmatrix} -d_B^{k-1} \\ \alpha \end{pmatrix}} B^k \oplus \ker(d_A^k)$. One readily verifies that removing the contractible $E \xrightarrow{1_E} E$ from the above acyclic complex yields $\text{cone}(f^\bullet) \in \mathbf{Ac}^-(\mathcal{E}) \cap \mathbf{C}^-(\mathcal{A})$. Proposition 3.9 now implies that $\text{cone}(f^\bullet) \in \mathbf{Ac}^-(\mathcal{A})$. It follows that $\ker(d_B^k) \in \mathcal{A}$. \square

Proposition 4.5. *Let \mathcal{A} be a preresolving subcategory in \mathcal{E} . Let $E \in \mathcal{E}$ with $\text{res. dim}_{\mathcal{A}}(E) = n < \infty$. For any \mathcal{A} -resolution $A^\bullet \rightarrow E$, there is a commutative diagram*

$$\begin{array}{ccc} B^\bullet & \longrightarrow & E \\ \downarrow f^\bullet & & \parallel \\ A^\bullet & \longrightarrow & E. \end{array}$$

where the top row is an \mathcal{A} -resolution and $\ker d_B^{-n+1} \in \mathcal{A}$. Moreover, $\text{cone}(f^\bullet) \in \mathbf{Ac}^-(\mathcal{A})$.

Proof. Let $C^\bullet \rightarrow E$ be an \mathcal{A} -resolution of E exhibiting that $\text{res. dim}_{\mathcal{A}} E = n$ (so $C^{-n-1} = 0$ and hence $\ker(d^{-n+1}) = C^{-n} \in \mathcal{A}$). By axiom **R2**, we obtain the pullback P^0 of $A^0 \rightarrow E$ along $C^0 \rightarrow E$. By axiom **PR1**, there is a deflation $B^0 \rightarrow P^0$ with $B^0 \in \mathcal{A}$. By axiom **R1**, we obtain the following commutative diagram

$$\begin{array}{ccccc} \text{im}(d_C^{-1}) & \longrightarrow & C^0 & \twoheadrightarrow & E \\ \uparrow \text{dotted} & & \uparrow & & \parallel \\ I_B^{-1} & \longrightarrow & B^0 & \twoheadrightarrow & E \\ \downarrow \text{dotted} & & \downarrow & & \parallel \\ \text{im}(d_A^{-1}) & \longrightarrow & A^0 & \twoheadrightarrow & E \end{array}$$

whose rows are conflations. Note that the left squares are pullback squares and, by axiom **R2**, the induced vertical maps are deflations.

We now iteratively construct B^\bullet as follows: Applying axiom **R2** thrice, we obtain the following commutative diagram

$$\begin{array}{ccccc} C^{-1} & \longrightarrow & \text{im}(d_C^{-1}) & & \\ & \swarrow & \text{PB} & \nearrow & \\ & & Q^{-1} & & \\ B^{-1} \twoheadrightarrow R^{-1} & \nearrow & \text{PB} & \searrow & I_B^{-1} \\ & \searrow & P^{-1} & \nearrow & \\ A^{-1} & \longrightarrow & \text{im}(d_A^{-1}) & & \end{array}$$

where the deflation $B^{-1} \rightarrow R^{-1}$ is obtained by axiom **PR1**. By axiom **R1**, one obtains the conflation $I_B^{-2} \rightarrow B^{-1} \rightarrow I_B^{-1}$ and the induced maps $\text{im}(d_C^{-2}) \leftarrow I_B^{-2} \rightarrow \text{im}(d_A^{-2})$. Iterating this procedure, one obtains morphisms $A^\bullet \xleftarrow{f^\bullet} B^\bullet \rightarrow C^\bullet$ such that B^\bullet is an \mathcal{A} -resolution of E . By Lemma 4.4, $\ker(d_B^{-n+1}) \in \mathcal{A}$. That f^\bullet is a quasi-isomorphism in $\mathbf{C}(\mathcal{A})$ is shown in the proof of Lemma 4.4. \square

Corollary 4.6. *Let \mathcal{A} be a preresolving subcategory in \mathcal{E} with $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) = n \in \mathbb{N}$. For each $E^\bullet \in \mathbf{Ac}(\mathcal{E}) \cap \mathbf{C}(\mathcal{A})$, there is a map $\alpha^\bullet: A^\bullet \rightarrow E^\bullet$ in \mathcal{T} where $A^\bullet \in \mathbf{Ac}(\mathcal{A})$. In particular, $A^\bullet \cong E^\bullet$ in $\mathbf{D}(\mathcal{A})$.*

Proof. Choose $k \in \mathbb{Z}$. The complex E^\bullet yields an \mathcal{A} -resolution $E_L^\bullet \rightarrow \text{im}(d_E^k)$ of $\text{im}(d_E^k) \in \mathcal{E}$ where E_L^\bullet is the brutal truncation of E^\bullet in degree k . By Proposition 4.5, we obtain a commutative diagram

$$\begin{array}{ccc} B^\bullet & \longrightarrow & \text{im}(d_E^k) \\ \downarrow f^\bullet & & \parallel \\ E_L^\bullet & \longrightarrow & \text{im}(d_E^k) \end{array}$$

such that $\text{cone}(f^\bullet) \in \mathbf{Ac}^-(\mathcal{A})$ and such that $\text{im}(d_B^{k-n}) \in \mathcal{A}$. Consider now the following map

$$\begin{array}{ccccccc} \Sigma^{-1} \text{cone}(f^\bullet) & \cdots & \longrightarrow & B^l \oplus E_L^{l-1} & \longrightarrow & B^{l+1} \oplus E_L^l & \longrightarrow & B^{l+2} \oplus E_L^{l+1} & \longrightarrow & \cdots \\ \downarrow g^\bullet & & & \downarrow (1 \ 0) & & \downarrow (1 \ 0) & & \downarrow (1 \ 0) & & \\ B^\bullet & \cdots & \longrightarrow & B^l & \longrightarrow & B^{l+1} & \longrightarrow & B^{l+2} & \longrightarrow & \cdots \\ \downarrow & & & \downarrow & & \downarrow & & \downarrow & & \\ \text{cone}(g^\bullet) & \cdots & \longrightarrow & B^{l+1} \oplus E_L^l \oplus B^l & \longrightarrow & B^{l+2} \oplus E_L^{l+1} \oplus B^{l+1} & \longrightarrow & B^{l+3} \oplus E_L^{l+2} \oplus B^{l+2} & \longrightarrow & \cdots \end{array}$$

It is well known (and straightforward to verify) that $\text{cone}(g) \cong \text{cone}(1_{B^\bullet}) \oplus E_L^\bullet$ in $\mathbf{C}(\mathcal{E})$. By Lemma 2.8, we have the following conflation

$$\text{im}(d_{\text{cone}(g)}^l) \twoheadrightarrow \text{im}(d_{\text{cone}(f)}^{l+1}) \oplus B^{l+1} \twoheadrightarrow \text{im}(d_B^{l+1})$$

for each $l \in \mathbb{Z}$. When $l = k - n - 1$, then $\text{im}(d_B^{l+1}) \in \mathcal{A}$ and, by axiom **PR2**, we have $\text{im}(d_{\text{cone}(g)}^l) \cong \text{im}(d_E^l) \oplus B^{l+1} \in \mathcal{A}$. It follows that one can add a null-homotopic complex $C^\bullet \in \mathbf{C}(\mathcal{A})$ to E^\bullet such that $C^\bullet \oplus E^\bullet \in \mathbf{Ac}(\mathcal{A})$. this concludes the proof. \square

Remark 4.7. Let \mathcal{A} be a preresolving subcategory in \mathcal{E} with $\text{res. dim}_{\mathcal{A}}(\mathcal{E}) = n \in \mathbb{N}$. The above proof yields that $\text{im}(d_E^l) \oplus B^{l+1} \in \mathcal{A}$ for all $l \in \mathbb{Z}$, it follows that $\text{res. dim}_{\mathcal{A}}(\text{im}(d_E^l)) \leq 1$.

Using the above corollary, one readily verifies that the conditions of [16, Proposition 7.2.1] are satisfied. This completes the proof of Theorem 4.1.

5. EXAMPLES AND APPLICATIONS

5.1. Examples.

Example 5.1. Consider the Isbell category \mathcal{I} (see [19, Section 2]), that is, \mathcal{I} is the full subcategory of the category \mathbf{Ab} of abelian groups containing no element of order p^2 (for a fixed prime p). Clearly $\mathcal{I} \subseteq \mathbf{Ab}$ is deflation-closed (in fact, $\mathcal{I} \subseteq \mathbf{Ab}$ is closed under subobjects) and thus \mathcal{I} is a deflation-exact category satisfying axiom $\mathbf{R3}^+$ (see Proposition 3.6).

Clearly, \mathcal{I} contains all projectives and thus $\mathcal{I} \subseteq \mathbf{Ab}$ satisfies condition $\mathbf{PR1}$. Hence $\mathcal{I} \subseteq \mathbf{Ab}$ is preresolving. Furthermore, as $\mathbf{Ab} = \text{Mod}(\mathbb{Z})$ global dimension one, we find that $\text{res. dim}_{\mathcal{I}}(\mathbf{Ab}) = 1$. By Theorems 3.11 and 4.1, we know that the natural functors $\mathbf{D}^*(\mathcal{I}) \simeq \mathbf{D}^*(\mathbf{Ab})$ are triangle equivalences (for $*$ $\in \{-, b, \emptyset\}$).

Remark 5.2. (1) Note that the Isbell category \mathcal{I} is not an extension-closed subcategory of \mathbf{Ab} , as evidenced by the conflation $\mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{Z}/p^2\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$. In particular, $\mathcal{I} \subseteq \mathbf{Ab}$ is not a (finitely) resolving subcategory.

(2) By [3, Example 4.7], the category \mathcal{I} is pre-abelian but does not satisfy axioms $\mathbf{L1}$, $\mathbf{L2}$ and $\mathbf{L3}$. In particular, \mathcal{I} is not exact.

Example 5.3. Let Γ be an ordered group. Given a Γ -filtered ring FR and a subset $\Lambda \subseteq \Gamma$, one can define the category of FR -glider representations $\text{Glid}_{\Lambda}(FR)$ (see [8, 13]). The category of FR -glider representations is a subcategory of the category of FR -prefragments $\text{Prefrag}_{\Lambda}(FR)$. By [13, Corollary 5.11], $\text{Glid}_{\Lambda}(FR) \subseteq \text{Prefrag}_{\Lambda}(FR)$ is closed under subobjects (in particular, $\text{Glid}_{\Lambda}(FR) \subseteq \text{Prefrag}_{\Lambda}(FR)$ is deflation-closed), but in general not under extensions. Moreover, by [13, Corollary 8.5], $\text{Glid}_{\Lambda}(FR)$ has enough projectives and the projectives belong to $\text{Prefrag}_{\Lambda}(FR)$. It follows that

$$\text{res. dim}_{\text{Glid}_{\Lambda}(FR)}(\text{Prefrag}_{\Lambda}(FR)) = 1.$$

By Theorems 3.11 and 4.1, we find that the natural functors $\mathbf{D}^*(\text{Glid}_{\Lambda}(FR)) \simeq \mathbf{D}^*(\text{Prefrag}_{\Lambda}(FR))$ are triangle equivalences (for $*$ $\in \{-, b, \emptyset\}$). This recovers [13, Proposition 8.6].

5.2. Application: the exact hull. Throughout this section, let \mathcal{E} denote a deflation-exact category satisfying axiom $\mathbf{R3}^+$. Following [11] and [24], there is an exact category \mathcal{E}^{ex} , called the *exact hull* of \mathcal{E} , together with an exact embedding $j: \mathcal{E} \rightarrow \mathcal{E}^{\text{ex}}$ which is 2-universal among exact functors to exact categories. Explicitly, the exact hull \mathcal{E}^{ex} can be constructed as the extension-closure of $i(\mathcal{E})$ in $\mathbf{D}^b(\mathcal{E})$ and inherits a conflation structure from the triangulated structure of $\mathbf{D}^b(\mathcal{E})$. The aim of this section is to provide an alternative proof for the fact that j lifts to a derived equivalence on the bounded derived categories (see [11, Theorem 7.15]) using Theorem 3.11.

The following is [11, Corollary 7.5].

Proposition 5.4. *For every $Z \in \mathcal{E}^{\text{ex}}$, there exists a conflation $X \rightarrow Y \rightarrow Z$ in \mathcal{E}^{ex} such that $X, Y \in \mathcal{E}$.*

The above proposition yields that $\mathcal{E} \subseteq \mathcal{E}^{\text{ex}}$ satisfies condition $\mathbf{PR1}$, in fact, $\text{res. dim}_{\mathcal{E}}(\mathcal{E}^{\text{ex}}) \leq 1$. Thus, to show that $j: \mathcal{E} \rightarrow \mathcal{E}^{\text{ex}}$ lifts to a derived equivalence $\mathbf{D}^*(\mathcal{E}) \rightarrow \mathbf{D}^*(\mathcal{E}^{\text{ex}})$ (where $*$ $\in \{-, b, \emptyset\}$), it suffices to show that $\mathcal{E} \subseteq \mathcal{E}^{\text{ex}}$ is deflation-closed.

Proposition 5.5. *A deflation-exact category \mathcal{E} satisfying axiom $\mathbf{R3}^+$ lies deflation-closed in its exact hull \mathcal{E}^{ex} .*

Proof. Let $X \xrightarrow{f} Y \xrightarrow{g} Z$ be a conflation in \mathcal{E}^{ex} with $Y, Z \in \mathcal{E}$. Viewing g as a map between stalk complexes in $\mathcal{E}^{\text{ex}} \subseteq \mathbf{D}^b(\mathcal{E})$, its cone is given by the two-term complex $Y \rightarrow Z$ with Y in degree -1 . Thus X is isomorphic (in $\mathbf{D}^b(\mathcal{E})$) to the two term complex X^{\bullet} given by $Y \rightarrow Z$ with Y in degree zero. On the other hand, Proposition 5.4 yields a conflation $A \rightarrow B \rightarrow X$ in \mathcal{E}^{ex} with $A, B \in \mathcal{E}$. It follows that X is also isomorphic to the two-term complex U^{\bullet} given by $A \rightarrow B$ with A in degree -1 .

Hence there is an isomorphism $U^{\bullet} \rightarrow X^{\bullet}$ in $\mathbf{D}^b(\mathcal{E})$ which can be represented by a roof $U^{\bullet} \xleftarrow{\alpha} V^{\bullet} \xrightarrow{\beta} X^{\bullet}$ in $\mathbf{K}^b(\mathcal{E})$ where both α and β are quasi-isomorphisms (here we used axiom $\mathbf{R3}^+$ and Proposition 2.10 to

see that all isomorphisms are quasi-isomorphisms). By [11, Proposition 3.18], we may assume that U^\bullet is a two-term complex $U^{-1} \rightarrow U^0$. It follows that the cone of β is the acyclic complex

$$0 \rightarrow U^{-1} \rightarrow U^0 \rightarrow Y \xrightarrow{g} Z \rightarrow 0$$

in $\mathbf{K}^b(\mathcal{E})$. It follows that g is a deflation in \mathcal{E} and thus g admits a kernel $\ker(g) \in \mathcal{E}$. One readily verifies that $\ker(g) \cong X$ in \mathcal{E}^{ex} as j is an exact embedding. \square

Remark 5.6. We now have two conflation structures on \mathcal{E} : the original conflation structure, and the conflation structure induced from \mathcal{E}^{ex} . It follows from Proposition 2.10.(4), and the assumption that \mathcal{E} satisfies axiom $\mathbf{R3}^+$, that these conflation structures coincide.

We have now proved the following theorem.

Theorem 5.7. *Let \mathcal{E} be a deflation-exact category. If \mathcal{E} satisfies axiom $\mathbf{R3}^+$, then \mathcal{E} is a preresolving subcategory of the exact category \mathcal{E}^{ex} such that $\text{res. dim}_{\mathcal{E}}(\mathcal{E}^{\text{ex}}) \leq 1$. In particular, the embedding $j: \mathcal{E} \rightarrow \mathcal{E}^{\text{ex}}$ lifts to a triangle equivalence $\mathbf{D}^*(\mathcal{E}) \rightarrow \mathbf{D}^*(\mathcal{E}^{\text{ex}})$, where $*$ is in $\{-, b, \emptyset\}$.*

Remark 5.8. In general, a deflation-exact category \mathcal{E} satisfying axiom $\mathbf{R3}^+$ need not be a resolving subcategory of \mathcal{E}^{ex} (since then, as an extension-closed subcategory of an exact category, \mathcal{E} would be exact itself). If \mathcal{E} does not satisfy axiom $\mathbf{R3}^+$, \mathcal{E} need not be a deflation-closed subcategory of \mathcal{E}^{ex} (as is illustrated in [14, Example 7.6]).

6. A COMPARISON TO KELLER'S THEOREM

For a preresolving subcategory \mathcal{A} of a deflation-exact category \mathcal{E} , we have used axiom $\mathbf{PR2}$ to show that the obvious induced conflation structure on \mathcal{A} is a deflation-exact structure. In this section, we weaken the conditions required on \mathcal{A} , following [18, Theorem 12.1]. In this case, as \mathcal{A} is not deflation-closed in \mathcal{E} , we cannot use Proposition 3.6) to endow \mathcal{A} with a deflation-exact structure. In Definition 6.4, we present a different conflation structure on \mathcal{A} and we show in Proposition 6.5 that this conflation structure endows \mathcal{A} with the structure of a deflation-exact category. The derived category of \mathcal{A} in Theorem 6.11 is taken with respect to this conflation structure.

We recall the following (dualized) definition from [18].

Definition 6.1. Let \mathcal{E} be an exact category and let $\mathcal{A} \subseteq \mathcal{E}$ be a fully exact subcategory.

C1 For each $E \in \mathcal{E}$, there exists a deflation $A \twoheadrightarrow E$ with $A \in \mathcal{A}$.

C2 For each conflation $X \twoheadrightarrow Y \twoheadrightarrow C$ with $C \in \mathcal{A}$, there exists a commutative diagram

$$\begin{array}{ccccc} A & \longrightarrow & B & \twoheadrightarrow & C \\ \downarrow & & \downarrow & & \parallel \\ X & \twoheadrightarrow & Y & \twoheadrightarrow & C \end{array}$$

where the top row is a conflation in \mathcal{A} .

The following is [18, Theorem 12.1].

Theorem 6.2. *Let $\mathcal{A} \subseteq \mathcal{E}$ be a fully exact subcategory of an exact category. If $\mathcal{A} \subseteq \mathcal{E}$ satisfies axioms **C1** and **C2**, the induced functor $\mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{E})$ is a triangle equivalence.*

Remark 6.3. (1) Axiom **C1** is equal to axiom $\mathbf{PR1}$.

(2) Keller remarks (see the paragraph above [18, Theorem 12.1]) that if $\mathcal{A} \subseteq \mathcal{E}$ is a fully exact subcategory of an exact category \mathcal{E} and $\mathcal{A} \subseteq \mathcal{E}$ is deflation-closed and satisfies axiom **C1**, then axiom **C2** is implied.

(3) As Keller requires $\mathcal{A} \subseteq \mathcal{E}$ to be extension-closed, \mathcal{A} automatically inherits an exact structure from \mathcal{E} (in particular, $\mathbf{D}(\mathcal{A})$ is defined).

Inspired by the above remarks, we seek to weaken axiom $\mathbf{PR2}$ in such a way that one obtains a natural deflation-exact structure on \mathcal{A} and such that we obtain an analogue of Proposition 3.9.

The following definition and proposition yield a natural deflation-exact structure on $\mathcal{A} \subseteq \mathcal{E}$.

Definition 6.4. Let \mathcal{E} be a deflation-exact subcategory and let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory. A conflation $A \xrightarrow{f} B \xrightarrow{g} C$ in \mathcal{E} with $A, B, C \in \mathcal{A}$ is called an \mathcal{A} -conflation if g admits all pullbacks in \mathcal{A} , i.e. for any $h: D \rightarrow C$ in \mathcal{A} , the pullback in \mathcal{E} of h along g belongs to \mathcal{A} .

Proposition 6.5. *Let \mathcal{E} be a deflation-exact category and let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory. The \mathcal{A} -conflations define a deflation-exact structure on \mathcal{A} .*

Proof. Axiom **R0** is clearly satisfied. Axiom **R2** holds by definition of an \mathcal{A} -conflation and the Pullback Lemma. It remains to show axiom **R1**, to that end, consider two \mathcal{A} -conflations $K \twoheadrightarrow B \twoheadrightarrow C$ and $L \twoheadrightarrow A \twoheadrightarrow C$. Consider the following commutative diagram in \mathcal{E} :

$$\begin{array}{ccccc} L & \xlongequal{\quad} & L & & \\ \downarrow & & \downarrow & & \\ P & \twoheadrightarrow & A & \twoheadrightarrow & C \\ \downarrow & & \downarrow & & \parallel \\ K & \twoheadrightarrow & B & \twoheadrightarrow & C \end{array}$$

Here the lower-left square is a pullback square in \mathcal{E} (which exists by axiom **R2**). As $A \twoheadrightarrow B$ is an \mathcal{A} -deflation, the pullback P belongs to \mathcal{A} . The Pullback Lemma now shows that $P \twoheadrightarrow A \twoheadrightarrow C$ is an \mathcal{A} -conflation. \square

Remark 6.6. The conflation structure from Definition 6.4 is the largest conflation structure on \mathcal{A} such that the embedding $\mathcal{A} \rightarrow \mathcal{E}$ is conflation-exact. Indeed, the class of \mathcal{A} -conflations is the largest subclass for which axiom **R2** is satisfied.

We now modify axiom **C2** accordingly.

Definition 6.7. Let \mathcal{E} be a deflation-exact category and let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory. We define the following axiom:

C2' For each conflation $X \twoheadrightarrow Y \twoheadrightarrow C$ with $C \in \mathcal{A}$, there exists a commutative diagram

$$\begin{array}{ccccc} A & \twoheadrightarrow & B & \twoheadrightarrow & C \\ \downarrow & & \downarrow & & \parallel \\ X & \twoheadrightarrow & Y & \twoheadrightarrow & C \end{array}$$

where the top row is an \mathcal{A} -conflation.

The following proposition clarifies the difference between an \mathcal{A} -conflation and an \mathcal{E} -conflation with terms in \mathcal{A} .

Proposition 6.8. Let \mathcal{A} be a full additive subcategory of a deflation-exact category \mathcal{E} such that axiom **C2'** holds. Let $X \xrightarrow{i} Y \xrightarrow{p} Z$ be a conflation in \mathcal{E} .

- (1) If $Y, Z \in \mathcal{A}$, then there exists a $B \in \mathcal{A}$ such that $X \oplus B \in \mathcal{A}$.
- (2) If X, Y and $Z \in \mathcal{A}$, then there exists an $B \in \mathcal{A}$ such that

$$X \oplus B \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1_B \end{pmatrix}} Y \oplus B \xrightarrow{(p \ 0)} Z$$

is an \mathcal{A} -conflation.

Proof. Assume that $Z \in \mathcal{A}$. By axiom **C2'**, we obtain an \mathcal{A} -conflation $A \twoheadrightarrow B \twoheadrightarrow Z$ that maps to (i, p) via maps $f: A \rightarrow X$ and $g: B \rightarrow Y$. By axiom **R2**, we obtain the following pullback square:

$$\begin{array}{ccccc} A & \xlongequal{\quad} & A & & \\ \downarrow & & \downarrow & & \\ X & \twoheadrightarrow & X \oplus B & \xrightarrow{(0 \ 1_B)} & B \\ \parallel & & \downarrow & \swarrow g & \downarrow \\ X & \twoheadrightarrow & Y & \twoheadrightarrow & Z \end{array}$$

- (1) If $Y \in \mathcal{A}$, then using that $A \twoheadrightarrow B \twoheadrightarrow Z$ is an \mathcal{A} -conflation, we find that $B \oplus X \in \mathcal{A}$.
- (2) Assume that $X, Y \in \mathcal{A}$. As $A \twoheadrightarrow B \twoheadrightarrow Z$ is an \mathcal{A} -conflation and the \mathcal{A} -conflations form a deflation-exact structure (Proposition 6.5), we can use Proposition 2.6 to obtain the \mathcal{A} -conflation

$$X \oplus B \xrightarrow{\begin{pmatrix} i & g \\ 0 & -1_B \end{pmatrix}} Y \oplus B \xrightarrow{(p \ pg)} Z .$$

The following diagram show that this \mathcal{A} -conflation is isomorphic to the desired \mathcal{A} -conflation:

$$\begin{array}{ccccc} X \oplus B & \xrightarrow{\begin{pmatrix} i & g \\ 0 & -1_B \end{pmatrix}} & Y \oplus B & \xrightarrow{\begin{pmatrix} p & pg \end{pmatrix}} & Z \\ \begin{pmatrix} 1_X & 0 \\ 0 & -1_B \end{pmatrix} \downarrow & & \downarrow \begin{pmatrix} 1_Y & g \\ 0 & 1_B \end{pmatrix} & & \parallel \\ X \oplus B & \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1_B \end{pmatrix}} & Y \oplus B & \xrightarrow{\begin{pmatrix} p & 0 \end{pmatrix}} & Z \end{array}$$

This concludes the proof. \square

Corollary 6.9. *Let \mathcal{A} be a full additive subcategory of a deflation-exact category \mathcal{E} satisfying axiom $\mathbf{C2}'$.*

- (1) *If $E^\bullet \in \mathbf{Ac}^-(\mathcal{E}) \cap \mathbf{C}^-(\mathcal{A})$, then E^\bullet is homotopic to a complex $A^\bullet \in \mathbf{Ac}^-(\mathcal{A})$.*
- (2) *If $E^\bullet \in \mathbf{Ac}^b(\mathcal{E}) \cap \mathbf{C}^b(\mathcal{A})$, then E^\bullet is homotopic to a complex $A^\bullet \in \mathbf{Ac}^b(\mathcal{A})$.*

Proof. This is similar to Proposition 3.9 and uses Proposition 6.8 to add in a controlled manner objects to each $\text{im}(d_E^k)$ so that the resulting complex is the desired complex. \square

Let \mathcal{E} be a deflation-exact category and let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory. The *relative weak idempotent completion* $\widehat{\mathcal{A}}_{\mathcal{E}}$ is the full subcategory of \mathcal{E} generated by all kernels in \mathcal{E} of retractions $r: A \rightarrow B$ in \mathcal{E} with $A, B \in \mathcal{A}$.

If $\mathcal{A} \subseteq \mathcal{E}$ is an additive subcategory satisfying axiom $\mathbf{C1}$, then $\widehat{\mathcal{A}}_{\mathcal{E}} \subseteq \mathcal{E}$ also satisfies axiom $\mathbf{C1}$.

Corollary 6.10. *Let \mathcal{A} be a full additive subcategory of a deflation-exact category \mathcal{E} satisfying axiom $\mathbf{C2}'$.*

- (1) *The subcategory $\widehat{\mathcal{A}}_{\mathcal{E}} \subseteq \mathcal{E}$ is deflation-closed and the embedding $\mathcal{A} \rightarrow \widehat{\mathcal{A}}_{\mathcal{E}}$ lifts to a triangle equivalence $\mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\widehat{\mathcal{A}}_{\mathcal{E}})$.*
- (2) *Assume that $\mathcal{A} \subseteq \mathcal{E}$ satisfies axiom $\mathbf{C1}$. If $\text{res. dim}_{\mathcal{A}}(E) = n$ for $E \in \mathcal{E}$, then $\text{res. dim}_{\widehat{\mathcal{A}}_{\mathcal{E}}}(E) \leq n$.*

Proof. This follows from Proposition 6.8. \square

Theorem 6.11. *Let \mathcal{E} be a deflation-exact category and let $\mathcal{A} \subseteq \mathcal{E}$ be a full additive subcategory. If $\mathcal{A} \subseteq \mathcal{E}$ satisfies axioms $\mathbf{C1}$ and $\mathbf{C2}'$, then the induced functor $\mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{E})$ is a triangle equivalence. Moreover, if $\text{res. dim}_{\mathcal{A}}(E) < \infty$ for all $E \in \mathcal{E}$, the functor $\mathbf{D}^b(\mathcal{A}) \rightarrow \mathbf{D}^b(\mathcal{E})$ is essentially surjective.*

Proof. This follows directly from Lemma 3.12 and Corollary 6.9. Alternatively, the functor $\mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{E})$ factors as $\mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\widehat{\mathcal{A}}_{\mathcal{E}}) \rightarrow \mathbf{D}^-(\mathcal{E})$. The first map is a triangle equivalence by Corollary 6.10 and the second map is a triangle equivalence by Theorem 3.11. \square

REFERENCES

- [1] Benjamin Antieau, David Gepner, and Jeremiah Heller, *K-theoretic obstructions to bounded t-structures*, Invent. Math. **216** (2019), no. 1, 241–300.
- [2] Maurice Auslander and Mark Bridger, *Stable module theory*, Memoirs of the American Mathematical Society, No. 94, American Mathematical Society, Providence, R.I., 1969.
- [3] Silvana Bazzoni and Septimiu Crivei, *One-sided exact categories*, J. Pure Appl. Algebra **217** (2013), no. 2, 377–391.
- [4] Carl Fredrik Berg and Adam-Christiaan van Roosmalen, *Representations of thread quivers*, Proc. Lond. Math. Soc. (3) **108** (2014), no. 2, 253–290.
- [5] Andrew J. Blumberg, David Gepner, and Gonalo Tabuada, *A universal characterization of higher algebraic K-theory*, Geom. Topol. **17** (2013), no. 2, 733–838.
- [6] A. Bondal and M. van den Bergh, *Generators and representability of functors in commutative and noncommutative geometry*, Mosc. Math. J. **3** (2003), no. 1, 1–36, 258.
- [7] Theo B  uhler, *Exact categories*, Expo. Math. **28** (2010), no. 1, 1–69.
- [8] Frederik Caenepeel and Fred Van Oystaeyen, *Glider representations*, CRC Press, 2019.
- [9] Dieter Happel, Idun Reiten, and Sverre O. Smal , *Tilting in abelian categories and quasitilted algebras*, Mem. Amer. Math. Soc. **120** (1996), no. 575, viii+ 88.
- [10] Robin Hartshorne, *Residues and duality*, Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20, Springer-Verlag, Berlin-New York, 1966.
- [11] Ruben Henrard and Adam-Christiaan van Roosmalen, *Derived categories of (one-sided) exact categories and their localizations*, 2019.
- [12] ———, *Localizations of (one-sided) exact categories*, 2019.
- [13] ———, *A categorical framework for glider representations*, 2020.
- [14] ———, *On the obscure axiom for one-sided exact categories*, 2020.
- [15] Dmitry Kaledin and Wendy Lowen, *Cohomology of exact categories and (non-)additive sheaves*, Adv. Math. **272** (2015), 652–698.

- [16] Masaki Kashiwara and Pierre Schapira, *Categories and sheaves*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 332, Springer-Verlag, Berlin, 2006.
- [17] Bernhard Keller, *Chain complexes and stable categories*, Manuscripta Math. **67** (1990), no. 4, 379–417.
- [18] ———, *Derived categories and their uses*, Handbook of algebra, Vol. 1, Handb. Algebr., vol. 1, Elsevier/North-Holland, Amsterdam, 1996, pp. 671–701.
- [19] G. M. Kelly, *Monomorphisms, epimorphisms, and pull-backs*, J. Austral. Math. Soc. **9** (1969), 124–142.
- [20] Amnon Neeman, *The derived category of an exact category*, J. Algebra **135** (1990), no. 2, 388–394.
- [21] Thomas Nikolaus and Peter Scholze, *On topological cyclic homology*, Acta Math. **221** (2018), no. 2, 203–409.
- [22] Leonid Positselski, *Two kinds of derived categories, Koszul duality, and comodule-contramodule correspondence*, Mem. Amer. Math. Soc. **212** (2011), no. 996, vi+133.
- [23] Daniel Quillen, *Higher algebraic K-theory. I*, Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), 1973, pp. 85–147. Lecture Notes in Math., Vol. 341.
- [24] Alexander Rosenberg, *Homological algebra of noncommutative spaces, 'i'*, preprint Max Planck, Bonn: MPIM 11-69.
- [25] Wolfgang Rump, *On the maximal exact structure on an additive category*, Fund. Math. **214** (2011), no. 1, 77–87.
- [26] ———, *The abelian closure of an exact category*, J. Pure Appl. Algebra **224** (2020), no. 10, 106395, 29.
- [27] R. W. Thomason and Thomas Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, The Grothendieck Festschrift, Vol. III, Progr. Math., vol. 88, Birkhäuser Boston, Boston, MA, 1990, pp. 247–435.
- [28] Jan Šťovíček, *Derived equivalences induced by big cotilting modules*, Adv. Math. **263** (2014), 45–87.
- [29] Jan Šťovíček, Otto Kerner, and Jan Trlifaj, *Tilting via torsion pairs and almost hereditary noetherian rings*, J. Pure Appl. Algebra **215** (2011), no. 9, 2072–2085.

RUBEN HENRARD, UNIVERSITEIT HASSELT, CAMPUS DIEPENBEEK, DEPARTEMENT WNI, 3590 DIEPENBEEK, BELGIUM
Email address: Ruben.Henrard@uhasselt.be

ADAM-CHRISTIAAN VAN ROOSMALEN, UNIVERSITEIT HASSELT, CAMPUS DIEPENBEEK, DEPARTEMENT WNI, 3590 DIEPENBEEK, BELGIUM
Email address: Adam-Christiaan.vanRoosmalen@UHasselt.be