## DECIMATION AND INTERLEAVING OPERATIONS IN ONE-SIDED SYMBOLIC DYNAMICS

WILLIAM C. ABRAM, JEFFREY C. LAGARIAS, AND DANIEL J. SLONIM

ABSTRACT. This paper studies subsets of one-sided shift spaces on a finite alphabet. Such subsets arise in symbolic dynamics, in fractal constructions, and in number theory. We study a family of decimation operations, which extract subsequences of symbol sequences in infinite arithmetic progressions, and show they are closed under composition. We also study a family of *n*-ary interleaving operations, one for each  $n \ge 1$ . Given subsets  $X_0, X_1, \dots, X_{n-1}$  of the shift space, the *n*-ary interleaving operator produces a set whose elements combine individual elements  $x_i$ , one from each  $X_i$ , by interleaving their symbol sequences cyclically in arithmetic progressions  $(\mod n)$ . We determine algebraic relations between decimation and interleaving operators and the shift operator. We study set-theoretic n-fold closure operations  $X \mapsto X^{[n]}$ , which interleave decimations of X of modulus level n. A set is n-factorizable if  $X = X^{[n]}$ . The n-fold interleaving operators are closed under composition and are idempotent. To each X we assign the set  $\mathcal{N}(X)$  of all values  $n \geq 1$  for which  $X = X^{[n]}$ . We characterize the possible sets  $\mathcal{N}(X)$  as nonempty sets of positive integers that form a distributive lattice under the divisibility partial order and are downward closed under divisibility. We show that all sets of this type occur. We introduce a class of weakly shift-stable sets and show that this class is closed under all decimation, interleaving, and shift operations. We study two notions of entropy for subsets of the full one-sided shift and show that they coincide for weakly shift-stable X, but can be different in general. We give a formula for entropy of interleavings of weakly shift-stable sets in terms of individual entropies.

## CONTENTS

1. Introduction	2
1.1. Summary	3
1.2. Background	4
1.3. Related work	5
2. Results	5
2.1. Decimation operations	6
2.2. Interleaving operations	6
2.3. Interleaving closure operations	7
2.4. Structure of interleaving factorizations	9
2.5. Shift-stability and weak shift-stability	10
2.6. Entropy of interleavings	10
2.7. Composition of interleavings and operad structure	11
2.8. Contents of paper	12
3. Decimations of arbitrary subsets of $\mathcal{A}^{\mathbb{N}}$	12
3.1. Compositions of decimations	12
3.2. Decimations and the shift	13
4. Interleaving for arbitrary subsets of $\mathcal{A}^{\mathbb{N}}$	13
4.1. Interleaving and decimation	13
4.2. <i>n</i> -fold interleaving closure operations	14
4.3. Shuffle identities for interleaving operators	15
4.4. Composition identities for interleaving closure operations	16

Date: November 9, 2020.

<sup>2020</sup> Mathematics Subject Classification. Primary 37B10; Secondary 11B25, 18M65, 37B40.

Key words and phrases. symbolic dynamics, entropy, infinite words, nonsymmetric operad.

The research of the second author was supported by NSF grant DMS-1701229, and a 2018 Simons Fellowship in Mathematics .

4.5. Interleaving commutes with set intersection	18
4.6. Shift action on interleavings	19
4.7. Topological closure	20
5. Interleaving factorizations and divisibility	21
5.1. Divisibility for interleaving factorizations	21
5.2. Structure of interleaving factorizations	22
5.3. Divisibility for self-interleaving factorizations	22
6. Infinitely factorizable closed subsets of $\mathcal{A}^{\mathbb{N}}$	24
6.1. Characterization of infinitely factorizable closed sets	24
6.2. Consequences of infinite factorizability	25
7. Iterated interleaving factorizations of general closed subsets of $\mathcal{A}^{\mathbb{N}}$	26
7.1. Iterated interleaving factorization trees	26
7.2. Arbitrary depth factorizations	27
8. Shift-stable and weakly shift-stable sets	29
8.1. Shift-stable sets	29
8.2. Closed shift-stable sets	30
8.3. Weakly shift-stable sets	32
9. Entropy of interleavings for general sets	33
9.1. Topological entropy and prefix topological entropy	33
9.2. Entropy and the shift operator	34
9.3. Entropy and decimations	34
9.4. Prefix entropy upper bound for interleaving	35
9.5. Stable prefix entropy and interleaving entropy equality	36
10. Concluding remarks	37
10.1. General interleaving operations	37
10.2. Iterated interleaving closure operations	38
10.3. Characterizing closed weakly shift-stable sets	38
Appendix A. Interleaving operad	38
References	40

#### 1. INTRODUCTION

Let  $\mathcal{A}$  be a finite alphabet of symbols, and suppose  $|\mathcal{A}| \geq 2$ . A basic object in one-sided symbolic dynamics is the full one-sided shift space  $\mathcal{A}^{\mathbb{N}}$ , which is the space of all one-sided infinite strings of symbols drawn from  $\mathcal{A}$ . Here  $\mathbb{N} = \{0, 1, 2, ...\}$  denotes the natural numbers, and  $\mathbb{N}^+ = \mathbb{N} \setminus \{0\}$  denotes the positive integers. We view  $\mathcal{A}^{\mathbb{N}} = \prod_{j \in \mathbb{N}} \mathcal{A}$  as a compact topological space carrying the product topology, with each copy of  $\mathcal{A}$  carrying the discrete topology; we call this topology of  $\mathcal{A}^{\mathbb{N}}$  the symbol topology. The dynamics in one-sided symbolic dynamics is the action of the *(one-sided) shift operator*  $S : \mathcal{A}^{\mathbb{N}} \to \mathcal{A}^{\mathbb{N}}$  on individual symbol sequences  $\mathbf{x} = a_0 a_1 a_2 a_3 \cdots$  by

$$S(\mathbf{x}) := a_1 a_2 a_3 a_4 \cdots . \tag{1.1}$$

In contrast, two-sided symbolic dynamics (treated in Lind and Marcus [33]) uses the two-sided shift operator  $S : \mathcal{A}^{\mathbb{Z}} \to \mathcal{A}^{\mathbb{Z}}$  with  $S((a_i)_{i \in \mathbb{Z}}) = (b_i)_{i \in \mathbb{Z}}$  with  $b_i = a_{i+1}$ . It focuses on sets  $X \subseteq \mathcal{A}^{\mathbb{Z}}$  that are invariant under the (two-sided) shift operator: SX = X. Such sets arise as discretizations of continuous dynamical systems such as geodesic flow, and led to the original formulation of symbolic dynamics by Morse and Hedlund [39]. In one-sided symbolic dynamics on subsets of  $\mathcal{A}^{\mathbb{N}}$  the spaces X can encode initial conditions. Initial conditions can break shift-invariance, so it is natural to consider spaces that are stable under the shift:  $SX \subseteq X$ .

This paper studies the action of decimation and interleaving operations acting on sets X in the framework of symbolic dynamics and coding theory. Decimation operations are important in digital signal processing and coding theory, and interleaving operations form a kind of inverse operation to them, see (1.4).

(1) At the level of individual symbol sequences, the *jth decimation operation at level* n, for  $i \ge 0$  and  $n \ge 1$ , denoted  $\psi_{i,n} : \mathcal{A}^{\mathbb{N}} \to \mathcal{A}^{\mathbb{N}}$ , for an individual symbol sequence  $\mathbf{x} = a_0 a_1 a_2 a_3 \cdots$  is

$$\psi_{i,n}(\mathbf{x}) := a_i a_{i+n} a_{i+2n} a_{i+3n} \cdots . \tag{1.2}$$

This operator extracts symbol subsequences having indices in an arithmetic progression given by  $i \pmod{n}$ , starting at initial index j.

(2) The *n*-fold interleaving operation  $\circledast_n : \mathcal{A}^{\mathbb{N}} \times \mathcal{A}^{\mathbb{N}} \times \cdots \times \mathcal{A}^{\mathbb{N}} \to \mathcal{A}^{\mathbb{N}}$  is an *n*-ary operation whose action on *n* individual symbol sequences  $\mathbf{x}_i = a_{i,0}a_{i,1}a_{i,2}\cdots$  for  $0 \le i \le n-1$  is defined by

$$(\mathbf{x}_0, \mathbf{x}_1, \cdots, \mathbf{x}_{n-1}) \mapsto \mathbf{y} := (\circledast_n)_{i=0}^{n-1} \mathbf{x}_i = \mathbf{x}_0 \circledast \mathbf{x}_1 \circledast \cdots \circledast \mathbf{x}_{n-1} = b_0 b_1 b_2 \cdots$$
(1.3)

in which the output sequence  $\mathbf{y} := b_0 b_1 b_2 \dots$  interleaves the symbols in arithmetic progressions of symbol indices (mod n), so that

 $b_{i+jn} = a_{i,j}$  for  $j \ge 0, \ 0 \le i \le n-1$ .

That is, the output y has in its symbol positions  $i \pmod{n}$  the symbols of  $x_i$  given in order.

Decimation and interleaving operations defined pointwise extend by set union to define set-valued operators acting on arbitrary subsets X of  $\mathcal{A}^{\mathbb{N}}$  (resp. of  $(\mathcal{A}^{\mathbb{N}})^n$ ). For examples, see Sections 2.1 and 2.2.

All individual symbol sequences x are constructible as n-fold interleavings of suitable decimations:

$$\mathbf{x} = (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(\mathbf{x}) \quad \text{for} \quad \mathbf{x} \in \mathcal{A}^{\mathbb{N}},$$
(1.4)

see Section 4.1.

1.1. **Summary.** This paper treats two topics.

1.1.1. The first topic studies algebraic properties of the algebraic structure of decimation and interleaving operators under composition. The set of all decimation operations is closed under composition, and the decimation and shift actions are compatible in a sense we describe in Section 3. Decimation operators are closed under composition.

We define the *n*-fold interleaving closure  $X^{[n]}$  of a set X in Section 2 as  $X^{[n]} = (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(X)$ , an operation that combines both decimations and interleavings. We show the operation sending X to  $X^{[n]}$  is a set-valued closure operation in the Moore sense, in particular  $X \subseteq X^{[n]}$ . A main result is that interleaving closure operators under composition satisfy

$$(X^{[m]})^{[n]} = (X^{[n]})^{[m]} = X^{[\operatorname{lcm}(m,n)]},$$
(1.5)

where lcm(m, n) denotes the least common multiple of m and n. Thus these operators are closed under composition, commute under composition, and are idempotent.

We show X is closed under n-fold interleaving closure, meaning  $X = X^{[n]}$ , if and only if X factorizes as  $X = (\circledast_n)_{j=0}^{n-1} X_j$  under the n-fold interleaving operation for some  $X_j$ . We study the allowable sets M of integers  $n \ge 1$  for which there exists some set X that has  $X = X^{[n]}$  if and only if  $n \in M$ . That is, letting  $\mathcal{N}(X) = \{n : X = X^{[n]}\}$ , we classify the sets  $M \subset \mathbb{N}^+$  such that  $M = \mathcal{N}(X)$  for some  $X \subset \mathcal{A}^{\mathbb{N}}$ . We show that if finite, the set  $\mathcal{N}(X)$  consists of the set of all divisors of an integer  $n_0$ , and all such  $n_0$  may occur. A new phenomenon is the existence of *infinitely factorizable* X, which necessarily have  $X = X^{[n]}$ for all n in an infinite distributive sublattice of  $\mathbb{N}^+$  under the divisibility partial order, downward closed under divisibility. We show all such infinite sublattices may occur for non-closed X, but if X is closed, we show the only allowed infinite sublattice is  $\mathbb{N}^+$ .

There is an additional algebraic structure consisting of the collection of all operations obtained from combining interleaving operations of different arities under composition. These form a nonsymmetric operad in the category of sets, and we term it the *interleaving nonsymmetric operad*. We give a series of

universal shuffle identities under composition satisfied by this operad. We discuss the operad formalism in Section 2.7 and in Appendix A.

1.1.2. The second topic studies symbolic dynamics aspects of decimation and interleaving operations. The shift operation acts compatibly with decimations and with *n*-fold interleaving, and we give commutation identities describing its action. The class of shift-invariant sets (those with SX = X) and the class of shift-stable sets (those with  $SX \subseteq X$ ) are not preserved under interleaving. We introduce an enlarged class of sets better adapted to these operations.

A set  $X \subseteq \mathcal{A}^{\mathbb{N}}$  is said to be *weakly shift-stable* if there are integers  $k > j \ge 0$  such that  $S^k X \subseteq S^j X$ . The set X need not be a closed set in the symbol topology. We show the class of all weakly shift-stable sets, denoted  $\mathcal{W}(\mathcal{A})$ , is closed under the shift and under all decimation and interleaving operations, as is the subclass  $\overline{\mathcal{W}}(\mathcal{A})$  of all closed sets in  $\mathcal{W}(\mathcal{A})$ .

The complexity of a set X can be measured using various notions of the entropy of X, which provide invariants that distinguish dynamical systems. The paper [2] studied two concepts of of entropy for X, the topological entropy  $H_{top}(X)$  and *path topological entropy*  $H_p(X)$ , which we term here *prefix topological entropy*. For general X one has  $H_p(X) \leq H_{top}(X)$ , and strict inequality may occur. We obtain an inequality relating the prefix topological entropy of an n-fold interleaving  $X = (\circledast_n)_{i=0}^{n-1} X_i$  to that of its factors  $X_i$ :

$$H_p(X) \le \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i),$$
(1.6)

and strict inequality may occur. A main result is that the class of weakly shift-stable sets  $\mathcal{W}(\mathcal{A})$  has good properties for both entropies; the two entropies are equal and equality holds in the interleaving inequality (1.6). In consequence, for weakly shift-stable sets we obtain a formula for topological entropy under interleaving:

$$H_{\rm top}(X) = \frac{1}{n} \sum_{i=0}^{n-1} H_{\rm top}(X_i)$$

1.1.3. Most results in this paper apply to general sets X, but for symbolic dynamics applications we are most interested in closed sets X in the product topology on  $\mathcal{A}^{\mathbb{N}}$ . These satisfy:

- (1) If X is a closed set, then all decimations  $\psi_{j,n}(X)$  are closed sets for  $j \ge 0$  and  $n \ge 1$ .
- (2) If X<sub>0</sub>, X<sub>1</sub>, ..., X<sub>n-1</sub> are closed sets, then their n-fold interleaving X = (⊛<sub>n</sub>)<sup>n-1</sup><sub>i=0</sub>X<sub>i</sub> is a closed set.
  (3) Conversely, if X is a closed set and has an n-fold interleaving factorization X = (⊛<sub>n</sub>)<sup>n-1</sup><sub>i=0</sub>X<sub>i</sub>, then
- (3) Conversely, if X is a closed set and has an *n*-fold interleaving factorization  $X = (\circledast_n)_{i=0}^{n-1} X_i$ , then each  $X_i$  is a closed set.

The decimation, interleaving, and shift operators all commute with topological closure operation  $X \mapsto \overline{X}$ . In consequence all *n*-fold interleaving closure operations commute with topological closure.

Detailed statements of results are made in Section 2. The main results concerning properties of n-fold interleaving closure operations of a set X are Theorems 2.10, 2.12 and 2.13. The main results concerning weakly shift-stable sets X are Theorems 2.15 and 2.20.

1.2. **Background.** This study was motivated by work on path sets initiated in [2]. Path sets are a class of closed sets in  $\mathcal{A}^{\mathbb{N}}$  that form a generalization of shifts of finite type and of sofic shifts in symbolic dynamics, and which also include sets not invariant under the shift operator. Path sets are described by finite automata, and have an automata-theoretic characterization as the closed sets in  $\mathcal{A}^{\mathbb{N}}$  that are the set of all infinite paths in some deterministic Büchi automaton. This class includes interesting sets arising in fractal constructions and in study of radix expansions in number theory (see [3], [4]) arising from a number theory problem of Erdős ([31]) The paper [2] considered decimation operations on path sets and showed that decimations of path sets are also path sets. The *p*-adic integers with the *p*-adic topology form a shift space with *p*-symbols, and interleaving operations on path sets arose in this context in [1].

The authors recently studied the action of interleaving operations on path sets, in [5]. Interleaving operations already lead to the breaking of shift-invariance even if all sets  $X_i$  used in the interleaving are

shift-invariant. The paper [5] shows that the class C(A) of all path sets on a finite alphabet A is closed under all interleaving operators.

This paper obtains results valid for general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , which provide perspective on results on path sets proved in [5]. The concept of weakly shift-stable closed sets  $\overline{\mathcal{W}}(\mathcal{A})$  supplies a good generalization of the class  $\mathcal{C}(\mathcal{A})$  of path sets to more general closed sets. The paper [5] shows that all path sets are weakly shift-invariant, which implies they are weakly shift-stable. In consequence, the entropy equalities of the present paper under weak shift-stability apply to interleaving of path sets. The present paper includes examples showing that various finiteness results given in [5] for path sets are not valid for general closed sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , see Remark 7.3.

1.3. **Related work.** Decimation operations play an important role in sampling and interpolation operations in digital signal processing ("downsampling"), and in multi-scale analysis and wavelets (e.g., [16], [29]). Interleaving constructions have been used in coding theory as a method for improving the burst error correction capability of a code (cf. [48, Section 7.5]). They are also considered in formal language theory; see Krieger et al. ([30]). The analogue of n-fold interleaving for finite codes is referred to by coding theorists as *block interleaving of depth* n. Decimation and interleaving operators together have been considered both in cryptography and cryptanalysis, cf. Rueppel [45] and Cardell et al [10]. Since methods of encoding and decoding can be viewed as dynamical processes, it is of interest to view these operations in a dynamical context.

1.3.1. There has been prior work on interleaving operations in the automata theory literature, typically for finite words. In 1974 Eilenberg [19, Chap. II.3, page 20] introduced a notion of *internal shuffle product*  $A \coprod B$  of two recognizable sets (= regular language) which corresponds to 2-fold interleaving. A more general notion is *alphabetic shuffle*. The shuffle product has been characterized in the context of finite automata by Duchamp et al [18, Sect. 4]. In this paper we are considering such operations on infinite words, which differ in nature from the finite word case. For infinite words viewed in an automata-theoretic context, see Perrin and Pin [40]. We are not aware of prior work studying the algebraic operator structure of interleaving operations.

1.3.2. Regarding dynamics, one-sided shift-stable sets have their dynamics partially classified by  $C^*$ algebra invariants. The work of Cuntz and Krieger [15] and Cuntz [14] was seminal in attaching such invariants to topological Markov chains (= two-sided shifts of finite type). Carlsen ([11], [12]) attached  $C^*$ algebras to one-sided subshifts and studied their properties. Shift-stable sets X are studied in the context of partial isometry actions and  $C^*$  algebras attached to them by Dokuchaev and Exel [17]. See Exel [22] for related background. One may ask whether there are generalizations of these constructions to weakly shift-stable sets introduced in this paper.

1.3.3. In the ongoing development of operad theory and *n*-categories, interleaving operations have recently played a role at a categorical level, see Leinster [32] and Cottrell [13]. General references for operads are Markl, Shnider and Stasheff [35], and more recently Loday and Vallette [34] and Bremner and Dotsenko [9] for algebraic operads.

Acknowledgments. We thank the reviewers for helpful comments and references. We thank I. Kriz for a discussion on operads. Some work of W. Abram and D. Slonim was facilitated by the Hillsdale College LAUREATES program, done by D. Slonim under the supervision of W. Abram. Work of J. Lagarias.was partially supported by NSF grants DMS-1401224 and DMS-1701229, and by a Simons Fellowship in Mathematics in 2019.

#### 2. Results

We give formal definitions with examples, and then state results.

#### 2.1. Decimation operations.

**Definition 2.1.** (Decimation operations) Let  $\mathcal{A}$  be a finite alphabet of symbols.

(1) For individual sequences  $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$  the *i*-th decimation operation at level *n*, denoted  $\psi_{i,n} : \mathcal{A}^{\mathbb{N}} \to \mathcal{A}^{\mathbb{N}}$ , for  $i \geq 0$  is defined for  $\mathbf{x} = a_0 a_1 a_2 a_3 \cdots$  by

$$\psi_{i,n}(\mathbf{x}) := a_i a_{i+n} a_{i+2n} a_{i+3n} \cdots$$

This operator extracts symbol subsequences having indices in an arithmetic progression given by i (mod n), which starts at initial index i. The principal n-decimations are those with 0 ≤ i ≤ n − 1.
(2) For sets X ⊆ A<sup>N</sup> the *i*-th decimation at level n, denoted ψ<sub>in</sub>(X), is the set union

$$\psi_{i,n}(X) := \left\{ \psi_{i,n}(\mathbf{x}) : \mathbf{x} \in X \right\}.$$

$$(2.1)$$

**Example 2.2.** For the alphabet  $\mathcal{A} = \{0, 1, 2, 3\}$  consider the sets<sup>1</sup>

$$X = {\mathbf{x}_1 = (01)^\infty, \mathbf{x}_2 = (10)^\infty}, \text{ and } Y = {\mathbf{y}_1 = (323)^\infty, \mathbf{y}_2 = (332)^\infty},$$

containing two periodic infinite words of period 2 and two periodic infinite words of period 3, respectively.

The principal 2-decimations of the elements of X are

$$\psi_{0,2}((01)^{\infty}) = 0^{\infty}, \psi_{1,2}((01)^{\infty}) = 1^{\infty}, \text{ and } \psi_{0,2}((10)^{\infty}) = 1^{\infty}, \psi_{1,2}((10)^{\infty}) = 0^{\infty}.$$

Thus  $\psi_{0,2}(X) := \{0^{\infty}, 1^{\infty}\}$  and  $\psi_{1,2}(X) = \{1^{\infty}, 0^{\infty}\} = \psi_{0,2}(X).$ 

The principal 2-decimations of the elements of Y are

$$\begin{split} \psi_{0,2}((323)^{\infty}) &= (332)^{\infty}, \psi_{1,2}((323)^{\infty}) = (233)^{\infty}, \text{ and } \psi_{0,2}((332)^{\infty}) = (323)^{\infty}, \psi_{1,2}((332)^{\infty}) = (332)^{\infty}. \end{split}$$
We obtain  $\psi_{0,2}(Y) &:= \{(332)^{\infty}, (323)^{\infty}\} = Y \text{ and } \psi_{1,2}(Y) = \{(233)^{\infty}, (332)^{\infty}\} \neq Y. \end{split}$ 

In Section 3 we show:

(1) The set of all decimation operations are closed under composition. For  $X \subseteq \mathcal{A}^{\mathbb{N}}$ ,

$$\psi_{j,m} \circ \psi_{i,n}(X) = \psi_{i+jn,mn}(X).$$

- This identity on subscripts matches an action of the (ax + b)-group on  $\mathbb{Z}$ .
- (2) The shift action is compatible with the decimation action: For  $X \subseteq \mathcal{A}^{\mathbb{N}}$ ,

$$\psi_{i,n}(SX) = \psi_{i+1,n}(X).$$

and

$$S\psi_{i,n}(X) = \psi_{i,n}(S^n X).$$

2.2. Interleaving operations. Interleaving operations comprise an infinite collection of *n*-ary operations  $(n \ge 1)$ , defined for arbitrary subsets X of the shift space  $\mathcal{A}^{\mathbb{N}}$ .

**Definition 2.3.** (Interleaving operations) Let  $\mathcal{A}$  be a finite alphabet of symbols.

(1) For individual sequences  $\mathbf{x}_i = a_{i,0}a_{i,1}a_{i,2}\cdots \in \mathcal{A}^{\mathbb{N}}$ ,  $(0 \leq i \leq n-1)$  the *n*-fold interleaving operation  $\circledast_n : \mathcal{A}^{\mathbb{N}} \times \mathcal{A}^{\mathbb{N}} \times \cdots \times \mathcal{A}^{\mathbb{N}} \to \mathcal{A}^{\mathbb{N}}$ , denoted either  $(\circledast_n)_{i=0}^{n-1}\mathbf{x}_i$  or  $\mathbf{x}_0 \circledast \mathbf{x}_1 \circledast \cdots \circledast \mathbf{x}_{n-1}$ , combines these sequences by

 $(\mathbf{x}_0, \mathbf{x}_1, \cdots, \mathbf{x}_{n-1}) \mapsto \mathbf{x}_0 \circledast \mathbf{x}_1 \cdots \circledast \mathbf{x}_n = \mathbf{y} := (a_{0,0} a_{1,0} \cdots a_{n-1,0}) \circ (a_{0,1} a_{1,1} \cdots a_{n-1,1}) \circ (a_{0,2} \cdots a_{n-1,0}) \circ (a_{0,1} a_{1,1} \cdots a_{n-1,1}) \circ (a_{0,2} \cdots a_{n-1,0}) \circ (a_{0$ 

where  $\circ$  denotes concatenation of sequences. That is,  $\mathbf{y} = b_0 b_1 b_2 \cdots$  with

$$b_{i+kn} = a_{i,j}$$
 for  $0 \le i \le n-1$ , and  $j \ge 0$ ,

so that the symbols of y in symbol positions  $i \pmod{n}$  are the symbols of  $\mathbf{x}_i$ ,  $(0 \le i \le n-1)$ .

(2) For sets  $X_j \subseteq \mathcal{A}^{\mathbb{N}}$ ,  $(0 \le j \le n-1)$ , their *n*-fold interleaving, denoted  $(\circledast_n)_{i=0}^{n-1} X_i$  or  $X_0 \circledast_n X_1 \circledast_n X_2 \circledast_n \cdots \circledast_n X_{n-1}$ , is defined by the set union:

$$(\circledast_n)_{i=0}^{n-1}X_i = \{\mathbf{x}_0 \circledast \mathbf{x}_1 \circledast \cdots \circledast \mathbf{x}_{n-1} : \mathbf{x}_j \in X_j \text{ for all } 0 \le j \le n-1\}.$$

<sup>1</sup>Here  $\mathbf{x}_1 = (01)^\infty = 010101...$ 

A set  $X = (\circledast_n)_{i=0}^{n-1} X_i$  is said to have an *n*-fold interleaving factorization. The sets  $X_i$  are called *n*-fold interleaving factors of X, or just interleaving factors. One can express *n*-fold interleavings in terms of principal decimations of level *n* as:  $(\circledast_n)_{i=0}^{n-1} X_i = \{\mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{j,n}(\mathbf{x}) \in X_j \text{ for all } 0 \le j \le n-1\}$ , see Proposition 4.1.

Example 2.4. Continuing Example 2.2, the 2-fold interleaving of X with itself is

$$X \circledast X = \{ \mathbf{x}_1 \circledast \mathbf{x}_1, \mathbf{x}_1 \circledast \mathbf{x}_2, \mathbf{x}_2 \circledast \mathbf{x}_1, \mathbf{x}_2 \circledast \mathbf{x}_2 \}$$
  
=  $\{ (0011)^{\infty}, (0110)^{\infty}, (1001)^{\infty}, (1100)^{\infty} \}.$ 

It contains four periodic words of period 4.

The 2-fold interleaving of Y with itself is

$$Y \circledast Y = \{ \mathbf{y}_1 \circledast \mathbf{y}_1, \mathbf{y}_1 \circledast \mathbf{y}_2, \mathbf{y}_2 \circledast \mathbf{y}_1, \mathbf{y}_2 \circledast b y_2 \}$$
  
=  $\{ (332233)^{\infty}, (332332)^{\infty}, (333223)^{\infty}, (333322)^{\infty} \}.$ 

It contains four periodic words of period 6. The 2-fold interleavings of X and Y are

$$\begin{aligned} X \otimes Y &:= \{ \mathbf{x}_1 \otimes \mathbf{y}_1, \mathbf{x}_1 \otimes \mathbf{y}_2, \mathbf{x}_2 \otimes \mathbf{y}_1, \mathbf{x}_2 \otimes \mathbf{y}_2 \} \\ &= \{ (031203130213)^{\infty}, (031302130312)^{\infty}, (130213031203)^{\infty}, (130312031302)^{\infty}. \} \end{aligned}$$

Each of them contains four periodic words of period 12. We have  $X \circledast Y \neq Y \circledast X$ .

A basic relation between interleaving and decimation is an identity, valid at the pointwise level, stating that ordered n-fold decimations post composed with n-fold interleaving give the identity map:

$$(\circledast_n)_{i=0}^{n-1}\psi_{i,n}(\mathbf{x}) = \mathbf{x} \quad \text{for} \quad \mathbf{x} \in \mathcal{A}^{\mathbb{N}}.$$
(2.2)

For this reason we call the decimations  $\psi_{i,n}$  for  $0 \le i \le n-1$ , the *principal decimations*. The remaining decimations  $i \ge n$  may be obtained by applying the one-sided shift operator to these decimation sets; see Proposition 3.2.

2.3. Interleaving closure operations. The interleaving operations together with principal decimations define a family of set-theoretic closure operations on general subsets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ . These operators are a main focus of this paper.

**Definition 2.5.** The *n*-fold interleaving closure operation  $X \mapsto X^{[n]}$  is defined for each  $X \subseteq \mathcal{A}^{\mathbb{N}}$  by

$$X^{[n]} := (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(X).$$
(2.3)

**Example 2.6.** For  $X = \{(10)^{\infty}, (01)^{\infty}\}$ , the 2-fold interleaving closure  $X^{[2]} := \psi_{0,2}(X) \circledast \psi_{1,2}(X)$  is

$$X^{[2]} = \{0^{\infty}, (01)^{\infty}, (10)^{\infty}, 1^{\infty}\}.$$

We have  $X \subseteq X^{[2]}$ .

**Example 2.7.** (Interleaving and *n*-fold interleaving closure) Let  $\mathcal{A} = \{0, 1\}$  and let  $X_0 \subset \mathcal{A}^{\mathbb{N}}$  be the onesided Fibonacci shift consisting of all words that do not contain the pattern 11 in two consecutive digits. Let  $X_1 = \mathcal{A}^{\mathbb{N}}$  be the full shift. Then:

- (1)  $X_0 \otimes X_1 \subset \mathbb{A}^{\mathbb{N}}$  consists of all words that do not contain a 1 in digits *i* and *i* + 2 for any *i* even. That is, there can be no 1's in consecutive even digits, but there are no other restrictions on the word. Here  $X_0$  and  $X_1$  are each invariant under the shift operator, i.e.,  $S(X_i) = X_i$ , but  $X_0 \otimes X_1$  is not shift-invariant.
- (2) Interleaving any number of copies of  $X_1$  gives  $X_1$ . That is,  $(\circledast_n)_{i=0}^{n-1}X_1 = X_1$  for  $n \ge 1$ .
- (3) The *n*-fold interleaving closure of  $X_0$  is  $X_1$  for all  $n \ge 2$ , that is,  $X_0^{[n]} = X_1 = \mathcal{A}^{\mathbb{N}}$ . This holds because  $\psi_{j,n}(X_0) = X_1$  for all  $j \ge 0$  when  $n \ge 2$ .

(4) Likewise,  $X_1^{[n]} = X_1$  for  $n \ge 1$ . So  $X^1$  has *n*-fold interleavings for all  $n \ge 1$ .

In Section 4.1 we show the existence of an n-fold interleaving factorization of a set X corresponds to its invariance under n-fold interleaving closure, and that its interleaving factors are its principal decimations.

#### **Theorem 2.8.** (Decimations and interleaving factorizations)

(1) A subset X of  $\mathcal{A}^{\mathbb{N}}$  has an n-fold interleaving factorization  $X = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1}$  if and only if  $X = X^{[n]}$ .

(2) If  $X = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1}$  has an n-fold interleaving factorization, then its ordered set of n-fold interleaving factors is unique, given by its principal decimations

$$X_i = \psi_{i,n}(X)$$
 for  $0 \le i \le n-1$ .

Regarding (2), there typically are many sets Y such that  $\psi_{i,n}(Y) = \psi_{i,n}(X)$  for  $0 \le i \le n-1$ , and we show X contains every such set Y in Theorem 4.2.

In Section 4.2 we justify the name *n*-fold interleaving closure by showing that  $X \mapsto X^{[n]}$  is a set-theoretic closure operation, as formalized in Grätzer [26, Chap. I, Sect. 3.12, Defn.26], and that  $X^{[n]}$  is characterized as the maximal set Z having the property that  $\psi_{i,n}(Z) = \psi_{i,n}(X)$  for  $0 \le i \le n - 1$ .

In Section 4.3 we establish universal algebraic identities relating certain compositions of n-fold interleavings for different n.

**Proposition 2.9.** (Interleaving shuffle identities) For each  $m, n \ge 2$  and arbitrary sets  $\{X_i : 0 \le i \le mn-1\}$  contained in the one-sided shift  $\mathcal{A}^{\mathbb{N}}$ , one has the identity of sets

$$(\circledast_n)_{i=0}^{n-1} \left( (\circledast_m)_{j=0}^{m-1} X_{i+jn} \right) = (\circledast_{mn})_{k=0}^{mn-1} X_k.$$
(2.4)

These identities are termed *shuffle identities* because the n-fold interleaving operation acts like a shuffling of n decks of cards together, taking the top cards in a particular order from each of the n decks, where the cards correspond to positions of symbols in the expansion.

In Section 4.4 we establish a main result determining the action of composition of interleaving closure operators. The shuffle identities play a crucial role in proving this result.

**Theorem 2.10.** (Composition of interleaving closures) For all  $m, n \ge 1$ , and all  $X \subseteq \mathcal{A}^{\mathbb{N}}$ ,

$$(X^{[m]})^{[n]} = (X^{[n]})^{[m]} = X^{[\operatorname{lcm}(m,n)]},$$
(2.5)

where lcm(m, n) denotes the least common multiple of m and n.

In Section 4.5 we show interleaving commutes with intersection:

$$\bigcap_{j=0}^{m-1} ((\circledast_n)_{i=0}^{n-1} X_{jn+i}) = (\circledast_n)_{i=0}^{n-1} (\bigcap_{j=0}^{m-1} X_{jn+i}).$$

In Section 4.6 we determine the action of the shift operator on n-fold interleavings and interleaving closures. In particular we show that

$$SX^{[n]} = (SX)^{[n]}.$$

In Section 4.7 we show that the topological closure operation commutes with both decimations and interleaving operations. In particular it commutes under composition with n-fold interleaving closure:

$$\overline{X}^{[n]} = \overline{X^{[n]}}$$

Thus if X is a closed then its n-fold interleaving closure  $X^{[n]}$  is closed.

2.4. Structure of interleaving factorizations. We study the possible structure of the set of all interleaving factorizations of a fixed set  $X \subseteq \mathcal{A}^{\mathbb{N}}$ .

**Definition 2.11.** Let X in  $\mathcal{A}^{\mathbb{N}}$  be a fixed set, with  $\mathcal{A}$  a finite alphabet.

(1) The *interleaving closure set*  $\mathcal{N}(X) \subseteq \mathbb{N}_+$  of X is the set of integers

 $\mathcal{N}(X) := \{ n : n \ge 1 \quad \text{and} \quad X = X^{[n]} \}.$ 

(2) The *interleaving factor set*  $\mathfrak{F}(X)$  consists of all *n*-ary interleaving factors,  $X_{i,n}$ , for all  $n \in \mathcal{N}(X)$ , i.e.

$$\mathfrak{F}(X) = \{\psi_{i,n}(X) : n \in \mathcal{N}(X), 0 \le i \le n-1\}$$

(3) The (full) decimation set  $\mathfrak{D}(X)$  consists of all decimations of X.

$$\mathfrak{D}(X) = \{ \psi_{i,n}(X) : i \ge 0, \ n \ge 1 \}.$$

The principal decimation set  $\mathfrak{D}_{prin}(X)$  consists of all principal decimations

$$\mathfrak{D}_{prin}(X) := \{\psi_{i,n}(X) : n \ge 1, 0 \le i \le n-1\}.$$

The interleaving factor set is a subset of the set of all principal decimations:  $\mathfrak{F}(X) \subseteq \mathfrak{D}_{prin}(X)$  We always have  $X \in \mathfrak{F}(X)$  and  $1 \in \mathcal{N}(X)$ .

An important feature of factorizations is that some X are *infinitely factorizable* in the sense that they have *n*-fold interleaving factorizations for infinitely many *n*, i.e.  $\mathcal{N}(X)$  is infinite. The full one-sided shift  $X = \mathcal{A}^{\mathbb{N}}$  on the alphabet  $\mathcal{A}$  is an example; it has *m*-fold factorizations for all  $m \ge 1$ , and  $\mathcal{N}(\mathcal{A}^{\mathbb{N}}) = \mathbb{N}^+$ , while its interleaving factor set  $\mathfrak{F}(\mathcal{A}^{\mathbb{N}}) = \{\mathcal{A}^{\mathbb{N}}\}$  contains one element. We term all the remaining ones *finitely factorizable*. There exist closed sets X having infinite  $\mathcal{N}(X)$  and having an infinite interleaving factor set  $\mathfrak{F}(X)$ , see example 6.5.

**Theorem 2.12.** (Structure of interleaving closure sets) Let  $\mathcal{N}(X) = \{n \ge 1 : X = X^{[n]}\}$ . Then  $\mathcal{N}(X)$  is nonempty and has the following properties.

(1) If  $n \in \mathcal{N}(X)$  and d divides n, then  $d \in \mathcal{N}(X)$ .

(2) If  $m, n \in \mathcal{N}(X)$  then their least common multiple  $\operatorname{lcm}(m, n) \in \mathcal{N}(X)$ .

Conversely, if a subset  $N \subset \mathbb{N}^+$  is nonempty and has properties (1) and (2), then there exists  $X \subseteq \mathcal{A}^{\mathbb{N}}$  with  $N = \mathcal{N}(X)$ .

This result is proved separately in the direct and converse directions as Theorem 5.1 and Theorem 5.3, respectively. A nonempty structure N having properties (1), (2) is abstractly characterized as any nonempty subset of  $\mathbb{N}^+$  that is a sublattice under the divisibility partial order, which is also downward closed under divisibility, see 5.1 (3). The notion of lattice here is that of G. Birkhoff, see Grätzer [26].

In Section 5.3 we also treat *self-interleaving factorizations*, which are interleaving factorizations in which all factors are identical. For a general set X we define the *self-interleaving closure set* 

$$\mathcal{N}_{\text{self}}(X) := \{ n \in \mathbb{N} : X = (\circledast_n)_{i=0}^{n-1} Y \text{ for some } Y \subseteq \mathcal{A}^{\mathbb{N}} \}$$

as the set of *n* such that *X* has an *n*-fold self-interleaving factorization. Self-interleavings naturally occur in circumstances given in We show that the structure of  $\mathcal{N}_{self}(X)$  has exactly the same collection allowed structure as  $\mathcal{N}(X)$  in Theorem 2.12; however for individual *X* the set of values  $\mathcal{N}_{self}(X)$  can be strictly smaller than  $\mathcal{N}(X)$ .

In Section 6 we study infinitely factorizable sets X in the special case that X is a closed set.

**Theorem 2.13.** (Classification of infinitely factorizable closed X) For a closed set  $X \subseteq A^{\mathbb{N}}$ , where A is a finite alphabet, the following properties are equivalent.

- (i) X is infinitely factorizable; i.e.,  $\mathcal{N}(X)$  is an infinite set.
- (ii) X has an n-fold interleaving factorization for all  $n \ge 1$ , ie.  $\mathcal{N}(X) = \mathbb{N}^+$ .
- (iii) For each  $k \ge 0$  there are nonempty subsets  $\mathcal{A}_k \subseteq \mathcal{A}$  such that  $X = \prod_{k=0}^{\infty} \mathcal{A}_k$  is a countable product of finite sets with the product topology.

In view of Theorem 2.12, the assumption that X is closed is necessary for these three equivalences to hold. The important consequence for closed sets X is that if they are infinitely factorizable then  $\mathcal{N}(X) = \mathbb{N}^+$ .

In Section 7 we study an iterated interleaving factorization process for a closed set X. If X is infinitely factorizable, we "freeze" it. If it is finitely factorizable, we decompose it to its maximal factorization, and then repeat the process on each of these factors. We show by example that this factorization process can go to infinite depth.

2.5. Shift-stability and weak shift-stability. We consider several classes of sets X having different transformation properties under the shift action.

Definition 2.14. (Shift-invariance, shift-stability, weak shift-stability)

(1) A set  $X \subseteq \mathcal{A}^{\mathbb{N}}$  is shift-invariant if SX = X.

(2) A set X is *shift-stable* if  $SX \subseteq X$ .

(3) A set X is weakly shift-invariant if there are integers  $k > j \ge 0$  such that  $S^k X = S^j X$ . (4) A set X is weakly shift-stable if there are integers  $k > j \ge 0$  such that  $S^k X \subseteq S^j X$ .

These definitions do not require the set X to be closed in the symbol topology.

In Section 8 we show consequences of these properties. We show that for shift-invariant sets, all interleaving factorizations are self-interleaving factorizations; that is, if X is shift-invariant, then  $\mathcal{N}(X) = \mathcal{N}_{self}(X)$ . We show that closed shift-stable sets have a forbidden blocks characterization paralleling the two-sided shift case. Example 8.5 constructs a closed set X having an infinite, strictly descending chain of sets under the shift operation.

An important property introduced here is weak shift-stability. The usefulness of this property is that the class  $\mathcal{W}(\mathcal{A})$  of all weakly shift-stable sets on a finite alphabet  $\mathcal{A}$  is closed under all decimation, interleaving and shift operations. This is not the case for properties (1)-(3) above.

**Theorem 2.15.** Let  $\mathcal{A}$  be finite alphabet and let  $X \subseteq \mathcal{A}^{\mathbb{N}}$  be a general set.

(1) If X is weakly shift-stable, then all decimations  $\psi_{j,n}(X)$  for  $j \ge 0$ ,  $n \ge 1$  are weakly shift-stable.

(2) If  $X_0, X_1, \dots, X_{n-1}$  are weakly shift-stable, then their n-fold interleaving  $Y := (\circledast_n)_{i=0}^{n-1} X_i$  is also weakly shift-stable.

(3) If X is weakly shift-stable, then its n-fold interleaving closure  $X^{[n]}$  is weakly shift-stable for each n > 1.

A parallel result holds for the class  $\overline{W}(\mathcal{A})$  of all closed weakly shift-stable sets on the finite alphabet  $\mathcal{A}$ . This latter class of sets includes the path sets studied in [2], as shown in [5].

2.6. Entropy of interleavings. In Section 9 we study two notions of entropy for general sets X.

**Definition 2.16.** (Topological entropy) The topological entropy  $H_{top}(X)$  is given by

$$H_{top}(X) := \limsup_{k \to \infty} \frac{1}{k} \log N_k(X)$$

where  $N_k(X)$  counts the number of distinct blocks of length k to be found across all words  $\mathbf{x} \in X$ .

The topological entropy is defined here as a limsup, however the limit always exists, as a consequence of a submultiplicativity property of block counting functions  $N_{k_1+k_2}(X) \leq N_{k-1}(X)N_{k_2}(X)$  of  $N_k(X)$ , see [6, Property 8]. Here  $\log$  denotes the natural logarithm; in information theory  $\log_2$  is used instead.

**Definition 2.17.** (Prefix entropy and stable prefix entropy)

(1) The prefix entropy (or path topological entropy of  $H_p(X)$  of a general set X, is defined by

$$H_p(X) := \limsup_{k \to \infty} \frac{1}{k} \log N_k^I(X),$$
(2.6)

where  $N_k^I(X)$  counts the number of distinct prefix blocks.  $b_0 b_1 \cdots b_{k-1}$  of length k found across all words  $\mathbf{x} \in X$ .

(2) The limit in (2.6) does not always exist, and we say that X has *stable prefix entropy* if the limit does exist:

$$H_p(X) := \lim_{k \to \infty} \frac{1}{k} \log N_k^I(X), \tag{2.7}$$

The prefix entropy was introduced in [2] under the name *path topological entropy* for a class of sets called *path sets*. In that paper symbol sequences were labels attached to paths of edges in a directed labeled graph. Prefix blocks were termed there *initial blocks* (for path sets) i because they represented the initial steps along a path in a directed labeled graph defining the path set. Since  $N_k^I(X) \le N_k(X)$  we always have  $H_p(X) \le H_{top}(X)$ , and strict inequality may hold.

In Section 9.2 we show the shift operator preserves both entropies. Decimation operations need not preserve entropy and Section 9.3 gives inequalities such entropies must satisfy. In Section 9.4 we establish an inequality for prefix entropy of interleavings of general sets,

**Theorem 2.18.** (Prefix entropy bound under interleaving) Let  $X_0, X_1, \dots, X_{n-1}$  be arbitrary subsets of  $\mathcal{A}^{\mathbb{N}}$ . The prefix entropy of the set  $X = X_0 \circledast X_1 \circledast \dots \circledast X_{n-1}$  is bounded above by the arithmetic mean of the prefix entropies of  $X_0, \dots, X_{n-1}$ . That is:

$$H_p(X_0 \circledast \cdots \circledast X_{n-1}) \le \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i).$$
 (2.8)

Example 9.6 shows that strict inequality in (2.8) may occur.

In Section 9.5 we show that the assumption of stable prefix entropy for each of the sets  $X_0, X_1, ..., X_{n-1}$  implies equality in this formula. and that the *n*-fold interleaving  $X = X_0 \circledast \cdots \circledast X_{n-1}$  itself has stable prefix entropy.

**Theorem 2.19.** (Stable prefix entropy interleaving formula) If each of the sets  $X_0, X_1, ..., X_{n-1}$  has stable prefix entropy, then the n-fold interleaving  $X = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1}$  also has stable prefix entropy. In addition

$$H_p(X) = \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i).$$
(2.9)

In contrast to this result for interleaving, decimations of a set X having stable prefix entropy need not have stable prefix entropy, see Remark 9.7.

We also deduce in Section 9.5 that all weakly shift-stable sets X have good entropy properties.

**Theorem 2.20.** (Weak shift-stability implies stable prefix entropy) If X is weakly shift-stable, then X has stable prefix entropy. and in addition  $H_p(X) = H_{top}(X)$ . Consequently n-fold interleaving of weakly-shift stable sets  $X_i$  has

$$H_{\rm top}(X) = \frac{1}{n} \sum_{i=0}^{n-1} H_{\rm top}(X_i).$$
(2.10)

Finally we observe that since all decimations of weakly shift-stable sets are weakly shift-stable, they will have stable prefix entropy.

2.7. Composition of interleavings and operad structure. In Section 7 we consider factorizations of a set X under iterated composition. We give examples of sets X having iterated factorizations going to infinite depth. This behavior differs from interleaving restricted to the class of all path sets on the finite alphabet A, as we show in [5] that the iterated factorization of any path set terminates at some finite depth.

Abstractly, the family of operations obtained under iterated composition from interleaving operations of all arities determines a *non-symmetric operad* (also called a *non-\Sigma operad*) in the sense of May [38], see also Markl et al. [35, Part I, Sect. 1.3] and Markl [36, Sect. 1]. Non-symmetric operads arise in many combinatorial constructions, see work of Giraudo [24], [25]. Iterated interleaving operations satisfy nontrivial universal identities under composition, examples being the shuffle identities given in Theorem 2.9.

These identities show that certain nested compositions of interleaving operations give equivalent operations. However most nestings of compositions yield distinct operations. In particular, interleaving operations do not satisfy the associative law when acting on collections of subsets X of  $\mathcal{A}^{\mathbb{N}}$ . For instance, each of the 3-ary operations  $X_0 \otimes X_1 \otimes X_2$  and  $X_0 \otimes (X_1 \otimes X_2)$  and  $(X_0 \otimes X_1) \otimes X_2$  are distinct.

Operads in general are characterized as (universal) algebraic objects satisfying a given set of universal identities. We shall consider the *interleaving non-symmetric operad* to be the non-symmetric operad whose universal identities are all the identities satisfied on the collection of all sets  $X \subseteq A^{\mathbb{N}}$  with alphabet size  $|\mathcal{A}| = 2$ . These universal identities include the shuffle identities in Theorem 2.9. This set of identities may be a generating set for all universal identities for this operad; we leave it as an open question to determine a generating set.

In Appendix A we provide details checking the operad structure associated to interleaving.

2.8. Contents of paper. The contents of the remainder of the paper are as follows:

Section 3 relates decimation operations and shows these operations are closed under composition and under the shift operator.

Section 4 studies interleaving operations and the interleaving closure operation  $X \to X^{[n]}$  for general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , proving Theorem 2.8 and the shuffle identities.

Section 5 establishes divisibility properties of n-fold factorizations of a closed set X.

Section 6 classifies infinitely factorizable closed sets X. These sets have more restricted factorizations than for non-closed sets.

Section 7 studies iterated interleaving factorizations of closed sets X. It shows by example that such iterated factorizations can continue to infinite depth.

Section 8 studies shift-stability and weak shift-stability of sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ . It gives a forbidden-blocks characterization of shift-stable closed sets. It shows that the class of weakly shift-stable sets is closed under all decimation and interleaving operations.

Section 9 defines and discusses topological entropy and prefix (topological) entropy, proving Theorems 2.18, through 2.20.

Section 10 discusses further directions for research.

Appendix A studies an operad structure generated by interleaving operations.

# 3. Decimations of arbitrary subsets of $\mathcal{A}^{\mathbb{N}}$

This section studies decimations and interleaving for subsets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ . All results in this section apply to arbitrary subsets X of  $\mathcal{A}^{\mathbb{N}}$ .

3.1. Compositions of decimations. The set of all decimation operators is closed under composition of operators. This composition action is a representation of the discrete ax+b semigroup given by the nonnegative integer matrices  $\begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}$  with  $a \ge 1$  and  $b \ge 0$ .

**Proposition 3.1.** (Composition of decimations) Let  $X \subseteq \mathcal{A}^{\mathbb{N}}$  be an arbitrary set. For all  $j, k \geq 0$  and  $m, n \geq 1$  we have

$$\psi_{j,m} \circ \psi_{k,n}(X) := \psi_{j,m}(\psi_{k,n}(X)) = \psi_{jn+k,mn}(X) \tag{3.1}$$

*Proof.* The result is verified separately for each element  $\mathbf{x} = x_0 x_1 x_2 \cdots \in X$  We set  $\mathbf{y} := \psi_{k,n}(\mathbf{x}) = x_k x_{k+n} x_{k+2n} x_{k+3n} \cdots$  where  $\mathbf{y} = y_0 y_1 y_2 \cdots$  has  $y_j = x_{k+jn}$ . Now

$$\psi_{j,m} \circ \psi_{k,n}(\mathbf{x}) = \psi_{j,m}(\mathbf{y}) = y_j y_{j+m} y_{j+2m} y_{j+3m} \cdots$$
$$= x_{k+jn} x_{k+jn+mn} x_{k+jn+2mn} \cdots,$$
$$= \psi_{jn+k,mn}(\mathbf{x}),$$

as asserted.

3.2. Decimations and the shift. The decimation operations also transform nicely under the one-sided shift  $S(a_0a_1a_2...) = a_1a_2a_3\cdots$ .

**Proposition 3.2.** (Shift of decimations) Let  $X \subseteq \mathcal{A}^{\mathbb{N}}$  be an arbitrary set. (1) For all  $j \ge 0$  and  $m \ge 1$  the one-sided shift S acts as

$$\psi_{j,m}(SX) = \psi_{j+1,m}(X).$$
 (3.2)

(2) In addition

$$S(\psi_{j,m}(X)) = \psi_{j+m,m}(X) = \psi_{j,m}(S^m X).$$
(3.3)

*Proof.* For a single element  $\mathbf{x} \in X$ , (3.2) is equivalent to the assertion

$$\psi_{j,m}(S\mathbf{x}) = \psi_{j,m}(S(x_0x_1x_2\cdots)) = \psi_{j,m}(x_1x_2x_3\cdots) = x_{j+1}x_{m+(j+1)}x_{2m+(j+1)}\cdots = \psi_{j+1,m}(\mathbf{x})$$

In this case (3.3) for  $\mathbf{x} \in X$  is equivalent to  $S(x_j x_{m+j} x_{2m+j} \cdots) = x_{m+j} x_{2m+j} x_{3m+j} \cdots$ . (2) For  $\mathbf{x} \in X$  we have

$$S\psi_{j,m}(\mathbf{x}) = S(x_j x_{j+m} x_{j+2m} \cdots) = x_{j+m} x_{j+2m} \cdots = \psi_{j+m,m}(\mathbf{x}) = \psi_{j,m}(S^m \mathbf{x}),$$

where the last equality used (1) iterated m times.

# 4. Interleaving for arbitrary subsets of $\mathcal{A}^{\mathbb{N}}$

4.1. Interleaving and decimation. Interleaving operations can be characterized in terms of the principal decimations of their output. The criterion (2) below could be used as an alternate definition of n-fold interleaving of sets.

**Proposition 4.1.** (Decimation characterization of interleavings)

(1) Every  $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$  has an n-fold interleaving factorization  $\mathbf{x} = (\circledast_n)_{i=0}^{n-1} \mathbf{x}_i$  for all  $n \ge 1$ . This factorization is unique, with  $\mathbf{x}_i = \psi_{i,n}(\mathbf{x})$  ( $0 \le i \le n-1$ ), so that

$$\mathbf{x} = \psi_{0,n}(\mathbf{x}) \circledast \psi_{1,n}(\mathbf{x}) \circledast \cdots \circledast \psi_{n-1,n}(\mathbf{x}) = (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(\mathbf{x}).$$
(4.1)

(2) If  $X \subseteq \mathcal{A}^{\mathbb{N}}$  has an n-fold interleaving factorization  $X = (\circledast_n)_{i=0}^{n-1} X_i$ , then

$$X = \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{i,n}(\mathbf{x}) \in X_i \quad \text{for all} \quad 0 \le i \le n-1 \}.$$

$$(4.2)$$

This factorization is unique with  $X_i = \psi_{i,n}(X) \quad (0 \le i \le n-1)$ , so that

$$X = \psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X) = (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(X).$$
(4.3)

*Proof.* The identity (4.1) is immediate from the definition of interleaving product, checking it symbol by symbol. This *n*-fold interleaving factorization of **x** is unique because if  $\mathbf{x} = (\circledast_n)_{i=0}^{n-1} \mathbf{x}_i$ , then the (i + kn)th symbol of **x** is by definition the *k*th symbol of  $\mathbf{x}_i$ , so that each symbol of  $\mathbf{x}_i$  is determined by a symbol of **x**.

(2) Let  $X = X_0 \circledast X_{\circledast} \cdots \circledast X_{n-1}$ . By definition

$$X = \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \mathbf{z} = (\circledast_n)_{i=0}^{n-1} \mathbf{x}_i : \mathbf{x}_i \in X_i, \text{ for all } 0 \le i \le n-1 \}$$
$$= \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{i,n}(\mathbf{x}) = \mathbf{x}_i, \text{ with } \mathbf{x}_i \in X_i, \text{ for all } 0 \le i \le n-1 \}$$
$$= \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{i,n}(\mathbf{x}) \in X_i \text{ for all } 0 \le i \le n-1 \},$$

which is (4.2); we used (1) to deduce the second equality.

To show (4.3), it suffices to show  $\psi_{i,n}(X) = X_i$ . We have  $\psi_{i,n}(X) \subseteq X_i$  by (4.2). To show the map is onto, for any  $\mathbf{x}_i$  we can pick arbitrary  $\mathbf{x}_j \in X_j$  for  $j \neq i$  and then (1) gives  $\mathbf{x} := (\circledast_n)_{j=0}^{n-1} \mathbf{x}_j \in Z$  has  $\psi_{i,n}(\mathbf{x}) = \mathbf{x}_i$ , as required.

We deduce Theorem 2.8 from the proposition.

*Proof of Theorem 2.8.* We are to show X has an interleaving factorization if and only if  $X = X^{[n]}$ .

(1) Suppose  $X = X^{[n]}$ . By definition  $X^{[n]} = \psi_{0,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X)$  has an interleaving factorization, so X does too. Conversely if  $X = X_0 \otimes X_1 \otimes \cdots \otimes X_{n-1}$  is an interleaving factorization then by Proposition 4.1 (2)  $X_i = \psi_{i,n}(X)$  whence  $X^{[n]} = X_0 \otimes X_1 \otimes \cdots \otimes X_{n-1} = X$ . 

(2) This is Proposition 4.1 (2).

4.2. *n*-fold interleaving closure operations. We show that the family of closure operations  $X \to X^{[n]}$  on sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  commutes with topological closure, and the equality  $X = X^{[n]}$  corresponds to X having an *n*-fold interleaving factorization.

The following result shows this operation is a closure operation in the set-theoretic sense.

**Theorem 4.2.** (*n*-fold interleaving closure) The *n*-fold interleaving closure operation  $X^{[n]}$  of sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ has the following properties:

(1) (Projection property) The n-fold interleaving closure  $X^{[n]}$  is characterized by the property that it is the maximal set Z such that its principal decimations at level n satisfy

$$\psi_{i,n}(Z) = \psi_{i,n}(X) \quad \text{for} \quad 0 \le i \le n-1.$$
 (4.4)

(2) (Extension property) Any set  $X \subseteq \mathcal{A}^{\mathbb{N}}$  satisfies

$$K \subseteq X^{[n]}.\tag{4.5}$$

- (3) (Idempotent property) The operation  $X \mapsto X^{[n]}$  is idempotent; i.e.,  $(X^{[n]})^{[n]} = X^{[n]}$  for all X.
- (4) (Isotone property) If  $X \subseteq Y$  then  $X^{[n]} \subset Y^{[n]}$ .

Remark 4.3. (Set theory closure property) Properties (2), (3), and (4) comprise the axioms of a Moore closure property (See Schechter [46, Sec. 4.1-4.12]). These axioms are known to be equivalent to the property of being closed under arbitrary intersections. The n-fold interleaving closure operation does not satisfy all of Kuratowski's axioms defining the closed sets of a topology; it does not satisfy the set union property  $(X \cup Y)^{[n]} = X^{[n]} \cup Y^{[n]}$ . It does satisfy the inclusion

$$X^{[n]} \cup Y^{[n]} \subseteq \left(X \cup Y\right)^{[n]}.$$
(4.6)

As an example showing the inclusion can be strict, take  $X = X^{[2]} = \{0^{\infty}\}, Y = Y^{[2]} = \{1^{\infty}\}$ . Then  $X^{[2]} \cup Y^{[2]} \subsetneq (X \cup Y)^{[2]} = \{0^{\infty}, 1^{\infty}, (01)^{\infty}, (10)^{\infty}\}$ . Relations between the interleaving closure operations and topological closure in  $\mathcal{A}^{\mathbb{N}}$  are given in Section 4.7.

*Proof.* (1) If a collection of sets each have property (4.4) then so does their union, and X has property (4.4), so there exists a maximal set Z with property (4.4). By definition

$$X^{[n]} := \psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X)$$

Then by Proposition 4.1(2),

$$X^{[n]} = \{ \mathbf{z} \in \mathcal{A}^{\mathbb{N}} : \psi_{i,n}(\mathbf{z}) \in \psi_{i,n}(X) \text{ for all } 0 \le i \le n-1 \}$$

$$(4.7)$$

The statement  $\psi_{i,n}(Z) = \psi_{i,n}(X)$  means that  $\psi_{i,n}(\mathbf{z})in\psi_{i,n}(X)$  for all  $\mathbf{z} \in Z$ . From (4.7), one sees that  $Z = X^{[n]}$  is precisely the maximal set such that (4.4) holds for all  $0 \le i \le n-1$ .

(2) It follows from (1). Alterntively, by Proposition 4.1(1) given  $\mathbf{x} \in X$  we have

$$\mathbf{x} = \psi_{0,n}(\mathbf{x}) \circledast \psi_{1,n}(\mathbf{x}) \circledast \cdots \circledast \psi_{n-1,n}(\mathbf{x}) \in \psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X)$$

which certifies  $\mathbf{x} \in X^{[n]}$ , proving (4.5).

(3) Idempotence follows from (1) and (2): By (1)  $X^{[n]}$  is the maximal set having  $\psi_{i,n}(X^{[n]}) \subseteq \psi_{i,n}(X)$ holds for  $0 \le i \le n-1$ . Now by (2)  $(X^{[n]})^{[n]}$  contains  $X^{[n]}$ . But  $\psi_{i,n}((X^{[n]})^{[n]}) \subseteq \psi_{i,n}((X^{[n]}) \subseteq \psi_{i,n}(X)$ for  $0 \le i \le n-1$ , so it is also maximal, so  $(X^{[n]})^{[n]} = X^{[n]}$ .

# (4) Suppose that $X \subseteq Y$ . Using the projection property (2) for X and Y separately shows $\psi_{i,n}(Y^{[n]} \cup X^{[n]}) = \psi_{i,n}(Y^{[n]}) \cup \psi_{i,n}(X^{[n]}) = \psi_{i,n}(X) \cup \psi_{i,n}(Y) = \psi_{i,n}(Y) = \psi_{i,n}(Y^{[n]}) \quad 0 \le i \le n-1.$

The projection property now gives  $Y^{[n]} \cup X^{[n]} \subseteq Y^{[n]}$ , whence  $X^{[n]} \subseteq Y^{[n]}$ .

We now prove for general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  the characterization of interleaving factorizations given in Theorem 2.8.

Proof of Theorem 2.8. (1) Suppose  $X = X^{[n]}$ . Then by definition  $X = X^{[n]} = \psi_{0,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X)$ is an interleaving factorization. Conversely if  $X = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1}$  is an interleaving factorization then by Proposition 4.1 (2)  $X_i = \psi_{i,n}(X) = X_i$  whence  $X^{[n]} = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1} = X$ . This proves both (1) and the uniqueness of factorization (2).

4.3. Shuffle identities for interleaving operators. The family of interleaving operations satisfy universal algebraic identities under particular compositions of operations, acting on general subsets of  $\mathcal{A}^{\mathbb{N}}$ . We now prove Proposition 2.9, which asserts

$$(\circledast_n)_{i=0}^{n-1} \left( (\circledast_m)_{j=0}^{m-1} X_{i+jn} \right) = (\circledast_{mn})_{k=0}^{mn-1} X_k.$$
(4.8)

One reads the interleaving of interleavings on the left side of (4.8) as

$$(X_0 \circledast X_n \circledast \cdots \circledast X_{(m-1)n}) \circledast (X_1 \circledast X_{n+1} \circledast \cdots \circledast X_{(m-1)n+1}) \circledast \cdots \\ \cdots \circledast (X_{n-1} \circledast X_{2n-1} \circledast \cdots \circledast X_{mn-1}),$$

with parentheses indicating composition of m-fold interleavings given as input to an n-fold interleaving. The right side of (4.8) is an mn-fold interleaving,

$$X_0 \circledast X_1 \circledast X_2 \circledast \cdots \circledast X_{n-1} \circledast X_n \circledast X_{n+1} \circledast \cdots \circledast \cdots \circledast X_{(m-1)n+n-2} \circledast X_{(m-1)n+n-1},$$
(4.9)

with factors taken in linear order.

Proof of Proposition 2.9. Using Proposition 4.1(2) we obtain

$$(\circledast_{n})_{i=0}^{n-1} \left( (\circledast_{m})_{j=0}^{m-1} X_{i+jn} \right) = \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{i,n}(\mathbf{x}) \in (\circledast_{m})_{j=0}^{m-1} X_{i+jn} \text{ for all } 0 \le i \le n-1 \}$$
$$= \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{j,m} \left( \psi_{i,n}(\mathbf{x}) \right) \in X_{i+jn} \text{ for all } 0 \le j \le n-1, 0 \le i \le n-1 \}$$
$$= \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{i+jn,mn}(\mathbf{x}) \in X_{i+jn} \text{ for all } 0 \le j \le n-1, 0 \le i \le n-1 \}$$
$$= (\circledast_{mn})_{k=0}^{mn-1} X_{k}.$$

The interleaving definition gives the second and fourth equality and Proposition 3.1 the third equality.  $\Box$ 

Shuffle identities are useful in studying self-interleavings of sets X.

**Definition 4.4.** Give  $X \subseteq \mathcal{A}^{\mathbb{N}}$  let  $X^{(\circledast n)}$  denote the *n*-fold self-interleaving defined by

$$X^{(\circledast n)} := (\circledast_n)_{i=0}^{n-1} X = X \circledast X \circledast \cdots \circledast X \quad (n \text{ factors in product})$$

The special case of self-interleaving under composition satisfies identities similar to that of exponentiation, a consequence of the shuffle identities.

**Proposition 4.5.** (Composition of self-interleavings) For any natural numbers  $m, n \ge 1$ , and any subset X of  $\mathcal{A}^{\mathbb{N}}$ , the following set-theoretic identity holds for n-fold, m-fold and mn-fold self-interleaving.

$$(X^{(\circledast n)})^{(\circledast m)} = (X^{(\circledast m)})^{(\circledast n)} = X^{(\circledast mn)}.$$
(4.10)

*Proof.* In Theorem 2.9 choose all  $X_k = X$  for  $0 \le k \le mn - 1$  and obtain  $(X^{(\circledast m)})^{(\circledast n)} = X^{(\circledast mn)}$ . Then interchange m and n.

4.4. Composition identities for interleaving closure operations. We prove Theorem 2.10 determining the composition of self-interleaving closure operations:  $(X^{[m]})^{[n]} = (X^{[n]})^{[m]} = X^{[\text{lcm}(m,n)]}$ .

We first establish a preliminary result giving formulas and inclusions for compositions of interleaving closure operations.

**Proposition 4.6.** (Composition formulas) (1) For all  $m, n \ge 1$ , and all  $X \subseteq \mathcal{A}^{\mathbb{N}}$ ,

$$(X^{[m]})^{[n]} = (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(X^{[m]}).$$
(4.11)

(2) For all  $m, n \geq 1$ , and all  $X \subseteq \mathcal{A}^{\mathbb{N}}$ ,

$$X^{[mn]} = (\circledast_n)_{i=0}^{n-1} (\psi_{i,n}(X)^{[m]}).$$
(4.12)

(3) For all  $m, n \geq 1$ 

$$X^{[m]} \subseteq X^{[mn]}. \tag{4.13}$$

(4) If gcd(m, n) = 1 then

$$(X^{[m]})^{[n]} = (X^{[n]})^{[m]} = X^{[mn]}$$

*Proof.* (1) This assertion is the definition of the *n*-fold interleaving closure of  $X^{[m]}$ .

(2) We set  $X_k := \psi_{k,mn}(X)$  for  $0 \le k \le mn - 1$  in the shuffle identity (2.4), obtaining

$$X^{[mn]} = (\circledast_n)_{i=0}^{n-1} \left( (\circledast_m)_{j=0}^{m-1} \psi_{i+jn,mn}(X) \right)$$

The right side of this equation contains terms  $Z_i := (\circledast_m)_{j=0}^{m-1} \psi_{jn+i,mn}(X)$ , and we must show  $Z_i = \psi_{i,n}(X)^{[m]}$ . We have

$$\psi_{i,n}(X)^{[m]} := \left(\psi_{0,m} \circ \psi_{i,n}(X)\right) \circledast \left(\psi_{1,m} \circ \psi_{i,n}(X)\right) \circledast \cdots \circledast \left(\psi_{m-1,m} \circ \psi_{i,n}(X)\right)$$
$$= \psi_{i,mn}(X) \circledast \psi_{i+n,mn}(X) \circledast \cdots \circledast \psi_{i+(m-1)n,mn}(X) = Z_i.$$

as required.

(3) We have  $X^{[m]} \subseteq (X^{[m]})^{[n]}$  by the extension property of *n*-fold interleaving. We claim that

$$(X^{[m]})^{[n]} \subseteq X^{[mn]}.$$
 (4.14)

To prove the claim, comparing the now proved (4.11) and (4.12), it suffices to show

$$\psi_{i,n}(X^{[m]}) \subseteq \psi_{i,n}(X)^{[m]} \quad \text{for} \quad 0 \le i \le n-1.$$
 (4.15)

For fixed *i*, the right side of this inclusion is an *m*-fold interleaving

$$\psi_{i,n}(X)^{[m]} = \left(\psi_{0,m} \circ \psi_{i,n}(X)\right) \circledast \left(\psi_{1,m} \circ \psi_{i,n}(X)\right) \circledast \cdots \circledast \left(\psi_{m-1,m} \circ \psi_{i,n}(X)\right)$$

The composition rule for decimations (Proposition 3.1) shows that

$$\psi_{i,n}(X)^{[m]} = \psi_{i,mn}(X) \circledast \psi_{i+n,mn}(X) \circledast \cdots \circledast \psi_{i+(m-1)n,mn}(X).$$
(4.16)

To evaluate the left side of the inclusion (4.15), suppose  $\mathbf{x} = \psi_{i,n}(\mathbf{z}) \in \psi_{i,n}(X^{[m]})$  with  $\mathbf{z} \in X^{[m]}$ . Now by Proposition 4.1 (1),  $\mathbf{x}$  has an *m*-fold interleaving factorization

$$\mathbf{x} = (\circledast_m)_{j=0}^{m-1} \mathbf{w}_j = \mathbf{w}_0 \circledast \mathbf{w}_1 \circledast \cdots \circledast \mathbf{w}_{m-1},$$

where

$$\mathbf{w}_j = \psi_{j,m}(\mathbf{x}) = \psi_{j,m}(\psi_{i,n}(\mathbf{z})) = \psi_{i+jn,mn}(\mathbf{z})$$

Therefore

$$\mathbf{x} = \psi_{i,n}(\mathbf{z}) = (\circledast_m)_{j=0}^{m-1} \psi_{i+jn,mn}(\mathbf{z}) = \psi_{i,mn}(\mathbf{z}) \circledast \psi_{i+n,mn}(\mathbf{z}) \circledast \cdots \circledast \psi_{i+(m-1)n,mn}(\mathbf{z}).$$
(4.17)

We are to show  $\mathbf{x} \in \psi_{i,n}(X)^{[m]}$  . It suffices to show

$$\psi_{i+jn,mn}(\mathbf{z}) \in \psi_{i+jn,mn}(X) \quad \text{for} \quad 0 \le j \le m-1,$$
(4.18)

since (4.17) then asserts  $\mathbf{x} \in (\circledast_m)_{j=0}^{m-1} \psi_{i+jn,mn}(X)$  whence (4.16) shows  $\mathbf{x} \in \psi_{i,n}(X)^{[m]}$ .

To show (4.18), any  $\mathbf{z} \in X^{[m]}$  has, for  $0 \le k \le m - 1$ ,

$$\psi_{k,m}(\mathbf{z}) \in \psi_{k,m}(X^{[m]}) = \psi_{k,m}(X),$$

where the equality of sets holds by definition of *m*-fold interleaving. Thus there exists some  $\tilde{\mathbf{z}}_k \in X$  with  $\psi_{k,m}(\mathbf{z}) = \psi_{k,m}(\tilde{\mathbf{z}}_k)$ . Now for  $0 \le i \le n - 1, 0 \le j \le m - 1$ , there exist unique  $(k, \ell)$  satisfying

$$i + jn = k + \ell m, \tag{4.19}$$

with  $0 \le k \le m-1, 0 \le \ell \le n-1$ . Here k = k(i, j) is determined by  $k \equiv i + jn \pmod{m}$ . We have

$$\psi_{i+jn,mn}(\mathbf{z}) = \psi_{k+\ell m,mn}(\mathbf{z}) = \psi_{\ell,n}(\psi_{k,m}(\mathbf{z}))$$
$$= \psi_{\ell,n}(\psi_{k,m}(\tilde{\mathbf{z}}_k)) = \psi_{k+\ell m,mn}(\tilde{\mathbf{z}}_k)$$
$$= \psi_{i+jn,mn}(\tilde{\mathbf{z}}_k) \in \psi_{i+jn,mn}(X),$$

showing (4.18).

(4) It suffices to show  $(X^{[m]})^{[n]} = X^{[mn]}$  if gcd(m, n) = 1; interchanging m and n then gives the other case. The proof of (3) showed that  $(X^{[m]})^{[n]} \subseteq X^{[mn]}$  holds (with no gcd restriction), so it suffices to show the reverse inclusion  $X^{[mn]} \subseteq (X^{[m]})^{[n]}$ . By the already proved (4.11) and (4.12) this assertion is

$$X^{[mn]} = (\circledast_n)_{i=0}^{n-1}(\psi_{i,n}(X)^{[m]}) \subseteq (\circledast_n)_{i=0}^{n-1}(\psi_{i,n}(X^{[m]})) = (X^{[m]})^{[n]}.$$
(4.20)

It therefore suffices to prove the individual set equalities

$$\psi_{i,n}(X^{[m]}) = \psi_{i,n}(X)^{[m]} \quad \text{for} \quad 0 \le i \le n-1.$$
 (4.21)

hold when gcd(m, n) = 1.

Now suppose we are given an arbitrary  $\mathbf{x} \in \psi_{i,n}(X)^{[m]}$ . We wish to show  $\mathbf{x} \in \psi_{i,n}(X^{[m]})$ . To begin,  $\mathbf{x}$  has an *m*-fold interleaving factorization

$$\mathbf{x} = (\circledast_m)_{j=0}^{m-1} \psi_{j,m}(\mathbf{x}_j),$$

in which each  $\mathbf{x}_j = \psi_{i,n}(\mathbf{z}_j) \in \psi_{i,n}(X)$  with  $\mathbf{z}_j \in X$ . Thus we have

$$\psi_{j,m}(\mathbf{x}_j) = \psi_{j,m}(\psi_{i,n}(\mathbf{z}_j)) = \psi_{i+jn,mn}(\mathbf{z}_j).$$
(4.22)

As in (3) there are  $(k, \ell)$  with

$$\psi_{i+jn,mn}(\mathbf{z}_j) = \psi_{k+\ell m,mn}(\mathbf{z}_j) = \psi_{\ell,n}(\psi_{k,m}(\mathbf{z}_j)).$$

Here, for fixed *i*, the value k = k(i, j) is given by  $k \equiv i + jn \pmod{m}$ . The values k(i, j) are all distinct as *j* ranges from 0 to m - 1 with *i* fixed, because gcd(m, n) = 1. It follows that the inverse map j = j(i, k)is well defined. By definition of *m*-fold interleaving closure, there will exist a value  $z \in X^{[m]}$  having

$$\psi_{k,m}(\mathbf{z}) = \psi_{k,m}(\mathbf{z}_j) \quad \text{for} \quad 0 \le k \le m - 1, \tag{4.23}$$

with  $\mathbf{z}_j \in X$  and j = j(i, k) runs over all  $0 \le j \le m - 1$  as k varies.

We claim that  $\psi_{i,n}(\mathbf{z}) = \mathbf{x}$ . We have

$$\begin{aligned} \psi_{i,n}(\mathbf{z}) &= (\circledast_m)_{j=0}^{m-1} \psi_{i+jn,mn}(\mathbf{z}) & \text{by (4.17)} \\ &= (\circledast_m)_{j=0}^{m-1} \psi_{k(i,j)+\ell(i,j)m,mn}(\mathbf{z}) & \text{by (4.19)} \\ &= (\circledast_m)_{j=0}^{m-1} \psi_{\ell(i,j),n}(\psi_{k(i,j),m}(\mathbf{z})) \\ &= (\circledast_m)_{j=0}^{m-1} \psi_{\ell(i,j),n}(\psi_{k(i,j),m}(\mathbf{z}_j)), & \text{by (4.23)} \end{aligned}$$

where the last line used the defining property of  $z \in X^{[m]}$ . Now we simplify

( ) m 1

$$\psi_{i,n}(\mathbf{z}) = (\circledast_m)_{j=0}^{m-1} \psi_{k(i,j)+\ell(i,j)m,mn}(\mathbf{z}_j))$$
  
=  $(\circledast_m)_{j=0}^{m-1} \psi_{i+jn,mn}(\mathbf{z}_j) = (\circledast_m)_{j=0}^{m-1} \psi_{j,m}(\mathbf{x}_j)$  by (4.22)  
=  $\mathbf{x}$ ,

so  $\mathbf{x} \in \psi_{i,n}(X^{[m]})$ .

We now prove the Theorem 2.10 formulas for composition of interleaving closures. .

Proof of Theorem 2.10. It suffices to prove  $(X^{[m]})^{[n]} = X^{[\operatorname{lcm}(m,n)]}$ , because its right side is symmetric in m and n; we may then exchange m and n to establish  $(X^{[n]})^{[m]} = X^{[\operatorname{lcm}(m,n)]}$ . We have already proven  $(X^{[m]})^{[n]} = X^{[mn]}$  for the case  $\operatorname{gcd}(m,n) = 1$  in Proposition 4.6.

For general n, m we let  $d = \gcd(m, n)$ , the greatest common divisor. One can always find e, f with d = ef such that e|m and f|n and  $\gcd(\frac{m}{e}, \frac{n}{f}) = 1$ . To see this, let  $d = \prod_p p^{e(p,d)}$  denote the prime factorization of d; then the choice  $e = \prod_{p^{e(p,d)}||m} p^{e(p,d)}$  and  $f = \prod_{p^{e(p,d)+1}|m} p^{e(p,d)}$  will work. Note that if  $p^{e(p,d)+1}|m$ , then necessarily  $p^{e(p,d)}||n$ , so that f|n. By construction, e|m, ef = d, and  $\gcd(\frac{m}{e}, \frac{n}{f}) = 1$ .

We then have

$$X^{[\operatorname{lcm}(m,n)]} = X^{[mn/ef]} = (X^{[m/e]})^{[n/f]} \subseteq (X^{[m/e]})^{[n]} \subseteq (X^{[m]})^{[n]}$$

Reading from left to right the second equality comes from Proposition 4.6 (4), the first inclusion follows from Proposition 4.6(3), and the final inclusion follows from the isotone property (4) in Theorem 4.2.

It remains to show that

$$(X^{[m]})^{[n]} \subset X^{[\operatorname{lcm}(m,n)]}$$

Now let  $d = \gcd(m, n)$ , so that  $\ell = \operatorname{lcm}(m, n) = \frac{mn}{d}$ . By Proposition 4.6(1) we have  $(X^{[m]})^{[n]} = (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(X^{[m]})$  (without any gcd restriction).

Now consider 
$$\mathbf{x} = \mathbf{x}_0 \circledast \mathbf{x}_1 \circledast \cdots \circledast \mathbf{x}_{n-1} \in (X^{[m]})^{[n]}$$
, and write  $\mathbf{x} = b_0 b_1 b_2 \cdots$ . Here for  $0 \le i \le n-1$ ,

$$\mathbf{x}_i := \psi_{i,n}(\mathbf{x}) = b_i b_{i+n} b_{i+2n} b_{i+3n} \cdots$$

We are to show that  $\mathbf{x} \in X^{[\operatorname{lcm}(m,n)]}$ . To begin, we have

$$\mathbf{x}_i = \psi_{i,n}(\mathbf{z}_{i,0} \otimes \mathbf{z}_{i,1} \otimes \cdots \otimes \mathbf{z}_{i,m-1}) \in \psi_{i,n}(X^{[m]}).$$

where each  $\mathbf{z}_{i,j} \in \psi_{j,m}(X)$  for  $0 \le j \le m - 1$ , so that

$$\mathbf{z}_{i,j} = \psi_{j,m}(\mathbf{w}_{i,j}) \quad \text{with} \quad \mathbf{w}_{i,j} \in X$$

Because gcd(m, n) = d, the application of  $\psi_{i,n}(\cdot)$  to  $\mathbf{z}_i = (\circledast_m)_{j=0}^{m-1} \mathbf{z}_{i,j} \in X^{[m]}$  only hits those words  $\mathbf{z}_{i,j}$  having subscripts j falling in  $\frac{m}{d}$  different residue classes  $(\mod m)$ , and it visits each such class exactly d times, as j varies over  $0 \le j \le m-1$ . These  $\frac{m}{d}$  classes  $(\mod m)$  comprise distinct residue classes  $(\mod \frac{m}{d})$ , again because gcd(m, n) = d. These classes are exactly  $i + jn \pmod{\frac{m}{d}}$  for  $0 \le j \le \frac{m}{d} - 1$ . We can therefore rewrite  $\mathbf{x}_i = \mathbf{y}_{i,0} \circledast \mathbf{y}_{i,1} \circledast \cdots \circledast \mathbf{y}_{i,\frac{m}{d}-1}$  with  $\mathbf{y}_{i,j} = b_{i+jn}b_{i+jn+mn/d}b_{i+jn+2mn/d} \cdots$  for  $0 \le j \le \frac{m}{d} - 1$ . We have  $lcm(m, n) = \frac{mn}{d}$  different elements  $\mathbf{y}_{i,j} \in \psi_{k,\frac{m}{d}}(X)$ . The key point is that for k = i + jn we have

$$\mathbf{y}_{i,j} = \psi_{i+jn \pmod{m/d}, m/d}(\mathbf{w}_{i+jn}) \in \psi_{k, \frac{m}{d}}(X) \text{ for } 0 \le i \le n-1, \ 0 \le j \le \frac{m}{d} - 1.$$

Here k = i + jn varies over the interval  $0 \le k \le \frac{mn}{d} - 1$ . Consequently,

$$\mathbf{x} = (\circledast_n)_{i=0}^{n-1} \mathbf{x}_i = (\circledast_n)_{i=0}^{n-1} \left( (\circledast_{m/d})_{j=0}^{m/d-1} \mathbf{y}_{i,j} \right) = (\circledast_{\frac{mn}{d}})_{k=0}^{mn/d-1} \psi_{k,\frac{mn}{d}} (\mathbf{w}_k) \right).$$

where the last equality uses the shuffle identity (2.4). We also find that k = i + jn runs through the residue classes  $(\mod mn/d)$  in the correct order. We conclude that  $\mathbf{x} \in X^{[mn/d]} = X^{[lcm(m,n)]}$ , establishing the desired inclusion.

4.5. **Interleaving commutes with set intersection.** Interleaving also behaves well with respect to intersection.

**Proposition 4.7.** (Interleaving commutes with intersection) For  $m, n \ge 2$  and subsets  $X_0, X_1, \dots, X_{mn-1}$  of  $\mathcal{A}^{\mathbb{N}}$ , the following set-theoretic identity holds:

$$\bigcap_{j=0}^{m-1} ((\circledast_n)_{i=0}^{n-1} X_{jn+i}) = (\circledast_n)_{i=0}^{n-1} (\bigcap_{j=0}^{m-1} X_{jn+i}).$$
(4.24)

*Proof.* By Proposition 4.1 (2), we have  $\mathbf{x} \in Z_j := (\circledast_n)_{i=0}^{n-1} X_{jn+i}$  if and only if  $\psi_i(\mathbf{x}) \in X_{jn+i}$  for  $0 \le i \le n-1$ . Consequently:

$$\mathbf{x} \in \bigcap_{j=0}^{m-1} ((\circledast_n)_{i=0}^{n-1} X_{jn+i}) \quad \Leftrightarrow \quad \psi_i(\mathbf{x}) \in X_{jn+i} \quad \text{for} \quad 0 \le i \le n-1, \ 0 \le j \le m-1$$
$$\Leftrightarrow \quad \psi_i(\mathbf{x}) \in \bigcap_{j=0}^{m-1} X_{jn+i} \quad \text{for} \quad 0 \le i \le n-1$$
$$\Leftrightarrow \quad \mathbf{x} \in (\circledast_n)_{i=0}^{n-1} (\bigcap_{j=0}^{m-1} X_{jn+i}),$$

verifying (4.24).

**Corollary 4.8.** Let  $X, Y \subseteq \mathcal{A}^{\mathbb{N}}$ . Then their *n*-fold interleaving closures satisfy

$$X^{[n]} \cap Y^{[n]} = Z^{[n]} \tag{4.25}$$

where  $Z := (\circledast_n)_{i=0}^{n-1} (\psi_{i,n}(X) \cap \psi_{i,n}(Y)) = Z^{[n]}.$ 

*Proof.* In Proposition 4.7 take m = 2 and  $n \ge 2$  and choose  $X_i = \psi_{i,n}(X)$  and  $X_{n+i} = \psi_{i,n}(Y)$  for  $0 \le i \le n-1$ . The left side of (4.24) is  $X^{[n]} \cap Y^{[n]}$  and the right side is Z. Here  $Z = Z^{[n]}$  holds because Z is defined as an n-fold interleaving of  $\psi_{i,n}(Z) := \psi_{i,n}(X) \cap \psi_{i,n}(Y)$  by construction. (In general  $\psi_{i,n}(X \cap Y) \subseteq \psi_{i,n}(X) \cap \psi_{i,n}(Y)$ , so that  $(X \cap Y)^{[n]} \subseteq Z^{[n]}$ , and strict inequality can hold.)  $\Box$ 

*Remark* 4.9. (Intersection of general interleaving closures) For intersection of two interleaving closures of different arities of a single set X we have, for all  $m, n \ge 1$ ,

$$X^{[\operatorname{gcd}(m,n)]} \subseteq X^{[m]} \cap X^{[n]}.$$
(4.26)

Equality always holds trivially when m = n, but need not hold when  $m \neq n$ . As an example, for m = 2, n = 3 take  $X = {\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3} = {(010100)^{\infty}, (111111)^{\infty}, (110111)^{\infty}}$ . Then  $(01)^{\infty}$  is contained in both  $X^{[2]}$  via the 2-fold interleaving  $\psi_{0,2}(\mathbf{x}_1) \circledast \psi_{1,2}(\mathbf{x}_2)$ , and  $X^{[3]}$  via the 3-fold interleaving  $\psi_{0,3}(\mathbf{x}_1) \circledast \psi_{1,3}(\mathbf{x}_1) \circledast \psi_{2,3}(\mathbf{x}_3)$ . We conclude  $X \subsetneq X^{[2]} \cap X^{[3]}$ . Note this example is closed and weakly shift-stable, having  $S^6 X = X$ .

## 4.6. Shift action on interleavings. The one-sided shift map acts as

$$S(a_0a_1a_2a_3\cdots)=a_1a_2a_3a_4\cdots$$

We show the one-sided shift S action preserves the property of having an n-fold interleaving factorization.

**Proposition 4.10.** (Interleaving and the shift map) Suppose that X has an n-fold interleaving factorization  $X = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-2} \circledast X_{n-1}$ .

(1) The one-sided shift map S acts by

$$S(X) = X_1 \circledast X_2 \circledast \cdots \circledast X_{n-1} \circledast S(X_0).$$
(4.27)

*Consequently* 

$$S^{n}(X) = S(X_{0}) \circledast S(X_{1}) \circledast \cdots \circledast S(X_{n-2}) \circledast S(X_{n-1}).$$
 (4.28)

(2) All iterates  $S^k(X)$  possess n-fold interleaving factorizations

$$S^{k}(X) = \psi_{k,n}(X) \circledast \psi_{k+1,n}(X) \circledast \cdots \circledast \psi_{k+n-1,n}(X).$$

*Proof.* (1), (2). It suffices to prove (4.27). The other assertions in (1) and assertion (2) then follow easily by induction on  $k \ge 1$ .

To begin, for all infinite words  $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$  we have

$$\psi_{j,n}(S\mathbf{x}) = \psi_{j+1,n}(\mathbf{x}) \quad \text{for all} \quad j \ge 0.$$
(4.29)

By Theorem 2.8 we have  $X_i = \psi_{i,n}(X)$  for  $0 \le i \le n-1$ . We set  $X_n = \psi_{n,n}(X)$ . By Proposition 3.1 (2),

$$\psi_{n,n}(X) = \psi_{0,n}(SX) = S^n \psi_{0,n}(X) = S(X_0).$$

We assert  $S(X) = X_1 \circledast X_2 \circledast \cdots \circledast X_{n-1} \circledast S(X_0)$ . We have the inclusion

$$S(X) \subseteq (\circledast_n)_{i=0}^{n-1} \psi_{i,n}(S(X)) = (\circledast_n)_{i=0}^{n-1} \psi_{i+1,n}(X) = (\circledast_n)_{i=1}^n X_i.$$

To show the opposite inclusion

$$X_1 \circledast X_2 \circledast \cdots \circledast X_{n-1} \circledast S(X_0) \subseteq SX,$$

let  $\mathbf{y} = \mathbf{y}_1 \circledast \mathbf{y}_2 \circledast \cdots \circledast \mathbf{y}_n \in X_1 \circledast X_2 \circledast \cdots \circledast X_n$ ; then  $\mathbf{y}_i \in X_i$  for  $1 \le i \le n$ . For  $\mathbf{y}_n \in X_n$  by definition there exists  $\mathbf{x} \in X$  such that  $\psi_{0,n}(\mathbf{x}) = x_0 \circ \mathbf{y}_n \in X_0$ , for some  $x_0$ , where  $x_0 \circ \mathbf{y}_n$  denotes the concatenation of the letter  $x_0$  and the infinite word  $\mathbf{y}_n$ . By the *n*-fold factorization hypothesis on X one may choose this  $\mathbf{x}$  so that also  $\psi_{i,n}(\mathbf{x}) = \mathbf{y}_i$  holds for  $1 \le i \le n - 1$ . Now one checks using (4.29) that

$$\psi_{i,n}(S(\mathbf{x})) = \mathbf{y}_{i+1} \quad \text{for} \quad 0 \le i \le n-1.$$

**Proposition 4.11.** (Shift map and *n*-fold interleaving closure) The shift map commutes with *n*-fold interleaving closure. For each  $n \ge 1$ , and a general set  $X \subseteq A^{\mathbb{N}}$ , there holds

$$S(X^{[n]}) = (SX)^{[n]}.$$
(4.30)

*Proof.* By definition the *n*-fold interleaving closure  $X^{[n]}$  has an *n*-fold interleaving factorization. Applying parts (1) and (2) of Proposition 4.10 we have

$$S(X^{[n]}) = S(\psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-2,n}(X) \circledast \psi_{n-1,n}(X))$$
  
=  $\psi_{1,n}(X) \circledast \psi_{2,n}(X) \circledast \cdots \circledast \cdots \circledast \psi_{n-1,n}(X) \circledast S \psi_{0,n}(X)$   
=  $\psi_{1,n}(X) \circledast \psi_{2,n}(X) \circledast \cdots \circledast \cdots \circledast \psi_{n-1,n}(X) \circledast \psi_{n,n}(X) = (SX)^{[n]}.$ 

4.7. Topological closure. Decimation and interleaving operations and the shift operation all commute with topological closure in  $\mathcal{A}^{\mathbb{N}}$ .

**Theorem 4.12.** (*n*-fold interleaving closure) Given a subset X of  $\mathcal{A}^{\mathbb{N}}$ , let  $\overline{X}$  denote its topological closure in the shift topology (product topology) in  $\mathcal{A}^{\mathbb{N}}$ .

(1) For each  $n \ge 1$  and  $j \ge 1$ ,

$$\psi_{j,n}(\overline{X}) = \overline{\psi_{j,n}(X)}.$$

In particular if X is a closed set in  $\mathcal{A}^{\mathbb{N}}$  then each decimation  $X_{j,n} = \psi_{j,n}(X)$  is a closed set. (2) For  $X_0, X_1, \ldots, X_{n-1} \subseteq \mathcal{A}^{\mathbb{N}}$ , there holds

$$(\circledast_n)_{j=0}^{n-1}\overline{X}_j = \overline{(\circledast_n)_{j=0}^{n-1}X_j}.$$

In particular the *n*-fold interleaving of closed sets is a closed set.

(3) The n-fold interleaving closure operation commutes with the closure operation on the product topology on  $\mathcal{A}^{\mathbb{N}}$ ,

$$(\overline{X})^{[n]} = \overline{X^{[n]}}.$$

(4) The shift operator commutes with topological closure,

$$S\overline{X} = \overline{SX}.$$

*Proof.* (1) Given a sequence  $\{\psi_{j,n}(\mathbf{x}_k) : k \ge 1\}$  in  $\psi_{j,n}(X)$ , with each  $\mathbf{x}_k = x_{0,k}x_{1,k}x_{2,k} \dots \in X$ , because closed sets X are compact in  $\mathcal{A}^{\mathbb{N}}$ , there exists a convergent subsequence of the  $\mathbf{x}_k \in X$ , having limit  $\mathbf{x} = x_0x_1x_2 \dots \in X$ , say. (Convergence is defined by eventual stability of each symbol  $x_{\ell,k}$  as  $k \to \infty$ , having  $x_{\ell,k} = x_\ell$  for all sufficiently large k.) It is easy to see that if  $\mathbf{x}_k \to \mathbf{x}$  in X then necessarily  $\psi_{j,n}(\mathbf{x}_k) \to \psi_{j,n}(\mathbf{x})$  is a convergent subsequence in  $\psi_{j,n}(X)$ , establishing that  $\psi_{j,n}(X)$  is closed.

(2) For interleavings of closed sets the convergence of symbols in a given position  $\ell = n\ell' + j'$  in  $X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1}$  depends only on  $X_{j'}$ , hence the closure property of X is inherited from that of the individual factors  $X_{j'}$ .

(3) The closure equality follows from (1), since  $X^{[n]} = \psi_{0,n}(X) \circledast \psi_{1,n} \circledast \cdots \circledast \psi_{n-1,n}$  and both sides of the equality add in all sequence limits taken in each symbol position separately.

(4) We have  $\mathbf{x} \in \overline{X}$  if there is a sequence  $\{\mathbf{x}_k : k \ge 1\}$  in X converging to  $\mathbf{x}$ . Then the sequence  $\mathbf{y}_k := S\mathbf{x}_k \in SX$  converges to  $S\mathbf{x}$  in  $S\overline{X}$  so  $S\overline{X} \subseteq \overline{SX}$ . Take now  $\mathbf{y}_k \in SX$  converging to  $\mathbf{x}$ . By definition of X there exists  $\mathbf{x}_k \in S$  with  $S\mathbf{x}_k = \mathbf{y}_k$ . Since the alphabel  $\mathcal{A}$  is finite, infinitely many of the  $\mathbf{x}_k$  have a fixed letter  $a_0$  as initial symbol. These define a subsequence  $\mathbf{x}_{k_i}$  that converges in X to a limit word  $\mathbf{x}$  and necessarily  $S\mathbf{x} = \mathbf{y}$ . Thus  $\overline{SX} \subseteq S\overline{X}$ .

## 5. INTERLEAVING FACTORIZATIONS AND DIVISIBILITY

We classify the possible values of n in n-fold interleaving factorizations for different n of arbitrary subsets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ .

# 5.1. Divisibility for interleaving factorizations.

**Theorem 5.1.** (Divisibility structure for interleaving factorizations) Let  $\mathcal{N}(X) = \{n : X = X^{[n]}\}$ .

(1) If  $n \in \mathcal{N}(X)$  and d divides n, then  $d \in \mathcal{N}(X)$ .

(2) If  $m, n \in \mathcal{N}(X)$  then their least common multiple  $\operatorname{lcm}(m, n) \in \mathcal{N}(X)$ .

(3) The interleaving closure set  $\mathcal{N}(X)$  of X has the structure of a distributive lattice with respect to the divisibility partial order, being closed under the join operation (least common multiple lcm), and the meet operation (greatest common divisor (gcd)). It is downward closed under divisibility, and contains the minimal element 1.

Proof of Theorem 5.1. (1) If  $n \in \mathcal{N}(X)$  then  $X = X^{[n]}$ . Suppose d divides n, so n = de. Now  $X \subseteq X^{[d]}$  by the extension property of Theorem 4.2. However  $X^{[d]} \subseteq X^{[de]} = X^{[n]}$  by Proposition 4.6 (2). Since  $X^{[n]} = X$  we conclude  $X^{[d]} = X$ , so  $d \in \mathcal{N}(X)$ .

(2) Suppose  $m, n \in \mathcal{N}(X)$  so that  $X = X^{[m]}$  and  $X = X^{[n]}$ . Then

$$X = X^{[n]} = (X^{[m]})^{[n]} = X^{[\operatorname{lcm}(m,n)]}$$

where, reading from the left, the second equality substituted  $X^{[m]}$  for X and the last equality is Theorem 2.10. Thus  $lcm(m,n) \in \mathcal{N}(X)$ .

(3) The set  $\mathcal{N}(X)$  is downward closed under divisibility by (1). If  $m, n \in \mathcal{N}(X)$  then  $gcd(m, n) \in \mathcal{N}(X)$  since it divides m. It is closed under the join operation lcm by (2). Thus  $\mathcal{N}(X)$  is a sublattice of the distributive lattice of integers  $\mathbb{N}_+$  under divisibility. It always has minimal element 1.

A corollary of part (2) says that interleaving factors of infinitely factorizable sets are infinitely factorizable.

**Corollary 5.2.** Let X be infinitely factorizable. Then every interleaving factor of X is also infinitely factorizable.

*Proof.* Suppose X is infinitely factorizable, and  $X = (\circledast_n)_{i=1}^{n-1} X_i$ . We show that  $X_i$  is infinitely factorizable for each  $0 \le i \le n-1$ . Since X has an m-fold interleaving factorization for infinitely many m. Theorem 2.12 implies that X has an lcm(m, n)-fold interleaving factorization for infinitely many m. Thus, X has an ne-fold interleaving factorization for infinitely many m. Thus, X has an ne-fold interleaving factorization for infinitely many m. Thus,  $X = (\circledast_{ne})_{k=0}^{ne-1}$ ,

then the shuffle identities of Theorem 2.9, combined with uniqueness of *ne*-fold interleaving factorizations, imply that each  $X_i$  has the *e*-fold interleaving factorization  $X_i = (\circledast_e)_{i=0}^{e-1} X_{i+jn}$ .

#### 5.2. Structure of interleaving factorizations.

**Theorem 5.3.** (Converse divisibility structure for interleaving factorizations)

Let  $N \subseteq \mathbb{N}^+$  be a nonempty set with the following properties:

- (1) If  $n \in N$  and d divides n, then  $d \in N$ .
- (2) If  $m, n \in N$ , then  $lcm(m, n) \in N$ .

If the alphabet  $\mathcal{A}$  has at least two letters, then  $N = \mathcal{N}(X)$  for some  $X \subseteq \mathcal{A}^{\mathbb{N}}$ .

*Proof.* Given a set N satisfying (1), (2) we construct a set  $\widetilde{X}$  on  $\mathcal{A} = \{0, 1\}$  with  $\mathcal{N}(\widetilde{X}) = N$ . Enumerate the elements of N as  $n_1, n_2, \ldots$ . Let  $\ell_1 = n_1$ , and for i > 1, let  $\ell_i = \operatorname{lcm}(n_1, \ldots, n_i)$ . Notice for  $i \leq j$ , that  $\ell_j = \operatorname{lcm}(\ell_i, \ell_j)$ , hence for any set  $X \subseteq \mathcal{A}^{\mathbb{N}}$  we have  $X^{[\ell_i]} \subseteq (X^{[\ell_i]})^{[\ell_j]} = X^{[\operatorname{lcm}(\ell_i, \ell_j)]} = X^{[\ell_j]}$ , using Theorem 2.10. Thus,  $(X^{[\ell_j]})_i$  is an increasing sequence of sets.

Choose  $X = \{0^{\infty}, 1^{\infty}\}$ . Notice that for any  $n \in \mathbb{N}$ ,  $X^{[n]}$  is precisely the set of all sequences in  $\mathcal{A}$  that are periodic with period dividing n. Now set  $\widetilde{X} := \lim_{j \to \infty} X^{[\ell_j]} = \bigcup_{j=1}^{\infty} X^{[\ell_j]}$ , so that  $\widetilde{X}$  is the set of sequences in  $\mathcal{A}$  that are periodic and have a period  $p \in N$  (since N is precisely the set  $\{n : n | \ell_j \text{ for some } j \geq 1\}$ ).

Claim.  $N = \mathcal{N}(\tilde{X}).$ 

(1) We show that if  $n \in N$ , then  $\widetilde{X}^{[n]} = \widetilde{X}$ . We already know  $\widetilde{X} \subseteq \widetilde{X}^{[n]}$ . Let  $\mathbf{x} \in \widetilde{X}^{[n]}$ . Then  $\mathbf{x} = (\circledast_n)_{i=0}^{n-1} \mathbf{x}_i$  for  $\mathbf{x}_1, \ldots, \mathbf{x}_n \in \widetilde{X}$ . Since there are finitely many of these  $\mathbf{x}_i$ , there is an  $\ell_j$  large enough that  $\mathbf{x}_1, \ldots, \mathbf{x}_n \in X^{[\ell_j]}$ . Choose  $\ell_j$  with j large enough that  $n|\ell_j$ . Then  $\operatorname{lcm}(n, \ell_j) = \ell_j$ , so  $X^{[\ell_j]}$  is closed under n-fold interleaving, and thus  $\mathbf{x} \in X^{[\ell_j]} \subseteq \widetilde{X}$ . Hence  $\widetilde{X}^{[n]} = \widetilde{X}$ , and so  $n \in \mathcal{N}(\widetilde{X})$ .

(2) We show that if  $n \notin N$ , then  $\widetilde{X}^{[n]} \neq \widetilde{X}$ . Since  $X \subseteq \widetilde{X}$ , we have  $X^{[n]} \subseteq \widetilde{X}^{[n]}$  by the extension property in Theorem 4.2. Let  $\mathbf{x}$  be any sequence in  $\mathcal{A}$  that is periodic with period n. Then  $\mathbf{x} \in X^{[n]}$ , and so  $\mathbf{x} \in \widetilde{X}^n$ . However, for any  $\ell_j$  we have  $\ell_j \in N$  by the structure of N, and since M is closed under divisibility,  $n \notin N$  implies n does not divide  $\ell_j$ ; hence  $\mathbf{x} \notin X^{[\ell_j]}$ . Since this is the case for all  $\ell_j$ ,  $\mathbf{x} \notin \widetilde{X}$ , and so  $n \notin \mathcal{N}(\widetilde{X})$ .

*Remark* 5.4. The sets  $\widetilde{X}$  constructed in the proof of Theorem 5.3 are all shift-invariant:  $S\widetilde{X} = \widetilde{X}$ . To show this, we note that a word  $\mathbf{x}$  on alphabet  $\mathcal{A} = \{0, 1\}$  is in  $\widetilde{X}$  if and only if it is fully periodic with a minimal period p belonging to  $N \subseteq \mathbb{N}^+$ , since N is downward closed under divisibility. The word  $S\mathbf{x}$  is also periodic with the same period, so  $S\mathbf{x} \in \widetilde{X}$ , hence  $S\widetilde{X} \subseteq \widetilde{X}$ . Since  $S^p\mathbf{x} = \mathbf{x}$ , we have  $\mathbf{y} = S^{p-1}\mathbf{x}$  is periodic with the same period , so  $\mathbf{y} \in \widetilde{X}$ , and  $S\mathbf{y} = S^p\mathbf{x} = \mathbf{x} \in S\widetilde{X}$ . It follows that  $S\widetilde{X} = \widetilde{X}$ .

#### 5.3. Divisibility for self-interleaving factorizations.

**Definition 5.5.** An *n*-fold interleaving factorization  $X = (\circledast_n)_{i=0}^{n-1} X_{i,n}$  is self-interleaving (or *n*-fold self-interleaving), if all factors are identical, i.e.  $X_{i,n} = X_{0,n}$  holds for  $1 \le i \le n-1$ . We sometimes write  $Z_n := X_{0,n}$  for the unique factor in this case.

There exist many sets X for which every interleaving factorization is a self-interleaving. We will show later, in Proposition 8.2, that if X is shift-invariant, then  $X = X^{[n]}$  implies that  $X_{0,n} = X_{i,n}$  holds for all  $n \ge 1$ . In addition there the exist examples with X having an n-fold self-interleaving, so that  $X_{0,n} = X_{in}$ for  $0 \le i \le n - 1$ , but with  $X_{0,n} \ne X_{i,n}$  for all  $i \ge n$ , see Example 5.7. The latter sets X can have a mixture of self-interleaving factorizations and non-self interleaving factorizations.

We show set of values of n for which a given X has an n-self-interleaving has divisibility properties parallel to those described in Theorem 2.12.

#### **Theorem 5.6.** (Structure of self-interleaving closure sets)

Let  $\mathcal{N}_{self}(X) = \{n \ge 1 : X = (\circledast_n)_{i=0}^{n-1} Z_n \text{ for some } Z_n \subseteq \mathcal{A}^{\mathbb{N}} \}$ . Then  $\mathcal{N}_{self}(X)$  is nonempty and has the following properties.

(1) If  $n \in \mathcal{N}_{self}(X)$  and d divides n, then  $d \in \mathcal{N}_{self}(X)$ .

(2) If  $m, n \in \mathcal{N}_{self}(X)$  then their least common multiple  $lcm(m, n) \in \mathcal{N}_{self}(X)$ .

Conversely, if a subset  $N \subseteq \mathbb{N}^+$  is nonempty and has properties (1) and (2), then there exists  $X \subseteq \mathcal{A}^{|nn|}$  with  $N = \mathcal{N}(X)$ .

*Proof.* (1) If d divides n we have n = de and now  $X = (\bigotimes_n)_{k=0}^{de-1} Z_n$  and  $Z_n = \psi_{k,de}(X)$  for  $0 \le k \le de-1$ . By the shuffle product identities in Theorem 2.9,

$$X = (\circledast_d)_{i=0}^{d-1} ((\circledast_e)_{j=0}^{e-1} (X_{jd+i})) = (\circledast_d)_{i=0}^{d-1} \left( (\circledast_e)_{j=0}^{e-1} Z_n \right).$$

We deduce  $X = (\circledast_d)_{i=0}^{d-1} Z_d$  where  $Z_d = (\circledast_e)_{j=0}^{e-1} Z_n$ , so X has a d-fold self-interleaving.

(2) Suppose that X has both an n-fold and an m-fold self-interleaving factorization. We wish to show it has an lcm(m, n)-fold self-interleaving factorization. Let d = gcd(m, n), and recall that there exist e, f with e|m, f|n having d = ef and  $gcd(\frac{m}{e}, \frac{n}{f}) = 1$  (shown in the proof of Theorem 2.10). By (1) the set is of self-interleaving factorizations is downward closed under divisibility, so that it has an  $\frac{m}{e}$ -fold self-interleaving factorization and an  $\frac{n}{f}$ -fold self-interleaving factorization, and now  $lcm(\frac{m}{e}, \frac{n}{f}) = \frac{mn}{ef} = \frac{mn}{d} = lcm(m, n)$ . We have therefore reduced proving (2) to proving it in the special case where gcd(m, n) = 1, with lcm(m, n) = mn.

In this case we are given that X has an m-fold and an n-fold self-interleaving factorization. We now have gcd(m, n) = 1 so by Theorem 2.12 we have an mn-fold interleaving factorization  $X = (\circledast_{mn})_{k=0}^{mn-1} X_{k,mn}$ . We wish to show it is self-interleaving, i.e. that

$$X_{k_1,mn} = X_{k_2,mn}$$
 for  $0 \le k_1 < k_2 \le mn - 1.$  (5.1)

We assert that for each  $0 \le i \le n-1$ ,

$$X_{j_1+im,mn} = X_{j_2+im,mn}$$
 for  $0 \le j_1 < j_2 \le m-1.$  (5.2)

To see this, note that by the shuffle identities in Theorem 2.9,

$$(\circledast_{mn})_{k=0}^{mn-1}X_{k,mn} = (\circledast_m)_{j=0}^{m-1}\Big((\circledast_n)_{i=0}^{n-1}X_{j+im,mn}\Big).$$

Since *m*-fold factorizations are unique, the right-hand side is a self-interleaving factorization, so for all  $0 \le j_1 < j_2 \le m - 1$ ,  $(\circledast_n)_{i=0}^{n-1} X_{j_1+im,mn} = (\circledast_n)_{i=0}^{n-1} X_{j_2+im,mn}$ . This implies, again by uniqueness, that  $X_{j_1+im,mn} = X_{j_2+im,mn}$  for all  $0 \le i \le n - 1$ .

Similarly, using the shuffle identity with the *n*-fold interleaving on the outside and the *m*-fold interleaving on the inside, we obtain for  $0 \le i_1 < i_2 \le n - 1$  that for each  $0 \le j \le m - 1$ ,

$$X_{i_1+jn,mn} = X_{i_2+jn,mn}$$
 for  $0 \le i_1 < i_2 \le n-1.$  (5.3)

Now we assert that when gcd(m, n) = 1 that (5.2) and (5.3) imply (5.1). Now (5.2) implies the equalities for all members in consecutive values of k in groups of length m,  $B_i = \{k = j + in : 0 \le j \le m - 1\}$ (for fixed i) but with no connection between between blocks for different i. Now (5.3) implies the same for blocks of consecutive values of k in groups of length n,  $C_j = \{k = i + jn : 0 \le i \le n - 1\}$  (for fixed j). Now the condition gcd(m, n) = 1 implies that for each  $0 \le i \le m - 2$  some group  $C_j$  includes members of both  $B_i$  and  $B_{i+1}$ , because the broken connection between blocks  $B_i$  is between  $k \equiv m - 1 \pmod{m}$ and  $k \equiv 0 \pmod{m}$ , and all multiples  $jn \not\equiv 0 \pmod{m}$  for  $1 \le j \le m - 1$ . Hence we get enough pairwise equalities to force (5.1) to hold. (This argument for equality of different  $X_{k,mn}$  stops at position mn, where the groups  $B_i$  and  $C_j$  both line up to create a barrier going to higher k.) This proves property (2) when gcd(m, n) = 1.

For the converse, it remains to show that if a subset  $N \subseteq \mathbb{N}^+$  is nonempty and has properties (1) and (2), then there exists  $X \subseteq \mathcal{A}^{\mathbb{N}}$  with  $N = \mathcal{N}_{self}(X)$ . For this, we use the fact that the sequences  $\widetilde{X}$  constructed in Theorem 5.3. that achieve  $N = \mathcal{N}(\widetilde{X})$  are shift-invariant, see Remark 5.4. Now Proposition 8.2 (which will be proved in Section 8) asserts that any shift- invariant X has the property that all of its interleaving factorizations will be self-interleaving factorizations. Thus,  $\mathcal{N}(\widetilde{X}) = \mathcal{N}_{self}(\widetilde{X})$ . We have already shown in the proof of Theorem 5.3 that  $\mathcal{N}(\widetilde{X}) = N$ .

**Example 5.7.** For a general set X the set of n giving self-interleaving factorization can be a strict subset of all interleaving factorizations of X. For the binary alphabet  $\mathcal{A} = \{0, 1\}$ , take  $X = \{00\{0, 1\}^{\mathbb{N}}\}$ , i.e. all infinite words beginning with 00.

We first show X has a n-fold interleaving factorization for all  $n \ge 1$ , so  $\mathcal{N}(X) = \mathbb{N}_+$ . For n = 1and n = 2 the factorization is self-interleaving with  $X_{0,2} = X_{1,2} = \{0\{0,1\}^{\mathbb{N}}\}$ . (Note that for  $j \ge 2$ one has  $X_{j,2} = \{0,1\}^{\mathbb{N}}$ .) In contrast we show  $\mathcal{N}_{self}(X) = \{1,2\}$  is finite. For  $n \ge 3$  its interleaving factorization has  $X_{0,n} = X_{1,n} = \{0\{0,1\}^{\mathbb{N}}\}$ , while  $X_{j,n} = \{0,1\}^{\mathbb{N}}$  is the full shift, for all  $j \ge 2$ , so it is not self-interleaving.

# 6. Infinitely factorizable closed subsets of $\mathcal{A}^{\mathbb{N}}$

**Definition 6.1.** A subset  $X \subseteq \mathcal{A}^{\mathbb{N}}$  is *infinitely factorizable* (under interleaving) if it has an *n*-fold interleaving factorization

$$X = X^{[n]} = \psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X)$$

for infinitely many  $n \ge 1$ .

6.1. Characterization of infinitely factorizable closed sets. We now characterize infinitely factorizable closed sets X by the properties given in Theorem 2.13. Property (iii) shows there are uncountably many different infinitely factorizable closed sets when the alphabet size  $|\mathcal{A}| \ge 2$ .

*Proof of Theorem 2.13.* We prove  $(iii) \Rightarrow (ii) \Rightarrow (i) \Rightarrow (iii)$ .

 $(iii) \Rightarrow (ii)$ . Suppose property (iii) holds. Write an element of X as  $\mathbf{x} = x_0 x_1 x_2 \cdots$ . Property (iii) says that the allowable symbols in location  $x_k$  of  $\mathbf{x}$  may be chosen arbitrarily while holding all other  $x_j, j \neq k$  fixed. In consequence all elements of  $X_k = \psi_{k,n}(X)$  may be chosen arbitrarily from  $\mathcal{A}_k$  while holding all the other  $X_j = \psi_{j,n}(X)$  constant. This may be done for each value of k, which implies property (ii) holds.  $(iii) \Rightarrow (ii)$ . Suppose property (iii) holds, and let  $n \geq 1$ . Then, using Proposition 4.1(2), we have,

$$X = \{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : x_k \in \mathcal{A}_k \text{ for all } k \ge 0 \}$$
  
=  $\{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : x_{j+kn} \in \mathcal{A}_{j+kn} \text{ for all } k \ge 0, \ 0 \le j \le n-1 \}$   
=  $\{ \mathbf{x} \in \mathcal{A}^{\mathbb{N}} : \psi_{j,n}(\mathbf{x}) \in \prod_{k=0}^{\infty} \mathcal{A}_{j+kn} \text{ for all } 0 \le j \le n-1 \}$   
=  $(\circledast_n)_{j=0}^{n-1} \prod_{k=0}^{\infty} \mathcal{A}_{j+kn},$ 

which is an n-fold interleaving factorization.

 $(ii) \Rightarrow (i)$ . Immediate.

 $(i) \Rightarrow (iii)$ . We prove the contrapositive. Suppose Property (iii) does not hold for X, we are to show property (i) does not hold. Let  $\mathcal{A}_k$  denote the letters that occur in the kth position of some word in X; it is a finite nonempty subset of the (finite) alphabet  $\mathcal{A}$ . For each  $k \ge 0$  all letter patterns  $\mathcal{A}_k \times \mathcal{A}_{k+1} \cdots \times \mathcal{A}_{k+\ell}$ may occur for each  $\ell \ge 1$ , then by the assumption  $\mathcal{A}$  is closed, we would have  $X = \prod_{k=0}^{\infty} \mathcal{A}_k$  which has Property (iii), contradicting our assumption. Therefore there must exist some finite  $k, \ell \ge 1$  and a finite set of consecutive  $\mathcal{A}_k, \mathcal{A}_{k+1}, \mathcal{A}_{k+2}, \dots \mathcal{A}_{k+\ell}$  such that there is a block  $a_k a_{k+1} \cdots a_{k+\ell}$  with each  $a_{k+i} \in \mathcal{A}_{k+i}$ for  $0 \le i \le \ell$  that does not occur in any element of X. We call this situation a  $(k, \ell)$ -missing-configuration.

To show Property (i) does not hold for this X we argue by contradiction. If Property (i) held for X, then there would exist some  $n \ge k + \ell + 1$  such that it had an n-fold interleaving factorization

$$X = \psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X).$$

Each word  $\mathbf{x} \in X$  has symbol  $x_{k+i}$  in position k+i lying as the first symbol in a word in the *n*-decimation set  $\psi_{k+i,n}(X)$ . We can find an infinite word, call it  $\mathbf{w}(k+i) \in X$  such that in position j := k+i it has the symbol in position k+i being  $a_i \in \mathcal{A}_{k+i}$ , for  $0 \le i \le \ell$  (by definition of  $\mathcal{A}_{k+i}$ ). For all remaining positions,  $0 \le j \le n-1$ , with  $j \notin \{k, k+1, \dots, k+\ell\}$  we pick a word  $\mathbf{w}(j) \in X$  arbitrarily.

Now the symbol sequence  $\mathbf{w} := \circledast_{j=0}^{n-1} \psi_{j,n}(w(j)) \in \psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X)$  contains the forbidden block  $a_k a_{k+1} \cdots a_{k+\ell}$  in positions k through  $k+\ell$ , showing that  $\mathbf{w} \notin X$ , the desired contradiction.

*Remark* 6.2. An important finiteness feature of the proof of Theorem 2.13 is that it shows that that existence of a  $(k, \ell)$ -missing-configuration certifies that X has no n-fold interleaving factorization with  $n \ge k + \ell + 1$  when X is closed.

The following example shows the hypothesis of X being closed set is necessary in the statement of Theorem 2.13.

**Example 6.3.** (Non-closed infinitely factorizable sets) Let X be the countable subset of  $\mathcal{A}^{\mathbb{N}}$  consisting of all infinite sequences having a finite number of 1's. Then X is infinitely factorizable, and all decimations  $\psi_{j,n}(X) = X$  are copies of itself. It is not a closed set; its closure in  $\mathcal{A}^{\mathbb{N}}$  is the full one-sided shift. It satisfies properties (i) and (ii) of Theorem 2.13 but fails to satisfy property (iii). (The set X can be viewed as the set of teminating binary expansions of all nonnegative dyadic rationals  $\frac{k}{2^m}$ .)

The construction of Theorem 5.3 produces infinitely factorizable X having  $\mathcal{N}(X) \subsetneq \mathbb{N}^+$ . Such sets satisfy property (i), and do not satisfy properties (ii), (iii) of Theorem 2.13.

#### 6.2. Consequences of infinite factorizability.

**Corollary 6.4.** Let X be an infinitely factorizable closed subset of  $\mathcal{A}^{\mathbb{N}}$ . Then its factor set  $\mathfrak{F}(X)$  consists of all decimations  $\psi_{j,n}(X)$  for  $n \ge 1$  and  $0 \le j \le n-1$ . Each decimated set  $\psi_{j,n}(X)$  is also infinitely factorizable.

*Proof.* By property (ii) of Theorem 2.13 X is factorizable for each  $n \ge 1$ , and its n-fold factors are  $\psi_{j,n}(X)$  for  $0 \le j \le n - 1$ . Now the property (iii) is preserved under decimations of all orders, hence all  $\psi_{j,n}(X)$  must be infinitely factorizable.

**Example 6.5.** (Infinitely factorizable closed subsets X of  $\mathcal{A}^{\mathbb{N}}$  having all decimations  $\psi_{j,n}(X)$  distinct) For  $\mathcal{A} = \{0, 1\}$  define  $\mathcal{A}_k \subset \mathcal{A}$  for  $0 \leq k < \infty$  as follows. Let  $\mathcal{A}_k = \{0\}$  for all indices  $k \in A$  with

$$A := \{k \ge 0 : 0 \le \{k\sqrt{2}\} < \frac{1}{2}\}$$
 where  $\{x\} = x - \lfloor x \rfloor$ .

(No special properties other than irrationality of  $\sqrt{2}$  are used). This set of indices is aperiodic (and has natural density  $\frac{1}{2}$ , using Weyl's equidistribution theorem.) Set  $\mathcal{A}_k = \{0, 1\}$  for all other integers  $k \notin A$ , which is also an aperiodic set (of natural density  $\frac{1}{2}$ ).

Set  $X = \prod_{k=0}^{\infty} \mathcal{A}_k$ . By Theorem 2.13 property (iii) it is a closed set and is infinitely factorizable, i.e.  $\mathcal{N}(X) = \mathbb{N}_+$ . Each decimation  $\psi_{j,n}(X)$  is also an infinite product space of the same kind whose set of indices k that have reduced alphabet  $\{0\}$  is exactly

$$A(j,n) := \{k \ge 0 : 0 \le \left\{ (nk+j)\sqrt{2} \right\} < \frac{1}{2} \}$$

Each  $\psi_{j,n}(X)$  is closed and infinitely factorizable. Consider now two distinct decimations  $\psi_{j,n}(X)$  and  $\psi_{\ell,m}(X)$ , where we may suppose  $1 \le n \le m$  and  $0 \le j, \ell < \infty$ , with  $j \ne \ell$  if n = m. To show distinctness we must show  $A(j,n) \ne A(\ell,m)$ . We use the well known fact that for each  $n \ge 1$  the sequence of fractional parts  $x_k = \{k(n\sqrt{2})\}$  ( $k \ge 1$ ) is dense modulo 1. (In fact, since  $n\sqrt{2}$  is irrational, Weyl's theorem implies that the sequence  $x_k$  is uniformly distributed modulo 1.) The argument has two cases.

*Case 1.* n = m. We write  $x_k := \{(kn + j)\sqrt{2}\}$ , and  $y_k := \{(kn + \ell)\sqrt{2}\}$ , where  $j \neq \ell$ . Now  $x_k = \{a_k + \theta\}$  for all k, where  $\theta = \{(\ell - j)\sqrt{2}\}$ . Because  $\sqrt{2}$  is irrational,  $\theta \in (0, 1)$ ; hence there must

be an open interval  $(a, b) \subset [0, \frac{1}{2})$  such that  $(a + \theta, b + \theta) \subset (\frac{1}{2}, 1]$ . Since  $x_k$  takes values dense in (0, 1), we will have infinitely many k with  $x_k \in (a, b)$ , and thus with  $y_k \in (a + \theta, b + \theta)$ . Therefore, there are infinitely many k with  $x_k \in [0, \frac{1}{2})$  and  $y_k \in (\frac{1}{2}, 1]$ . For these k,  $\mathcal{A}_{kn+j} = \{0\}$ , while  $\mathcal{A}_{kn+\ell} = \{0, 1\}$ , so all sequences in  $\psi_{j,n}(X)$  must have kth symbol 0, while  $\psi_{n,\ell}(X)$  has sequences with kth symbol taking both values 0 or 1 (i.e.,  $k \in A(j, n)$  but  $k \notin A(\ell, n)$  for these k). Thus  $\psi_{j,n}(X) \neq \psi_{\ell,n}(X)$ .

*Case 2.* n < m. We write  $x_k := \{(nk+j)\sqrt{2}\}$  and  $y_k := \{(mk+\ell)\sqrt{2}\}$ . A calculation shows that  $y_k = \{\frac{m}{n}x_k + \theta\}$ , where  $\theta = \{(\ell - \frac{jm}{n})\sqrt{2}\}$ . Again,  $\theta \in (0, 1)$ . There is an open interval  $(c, d) \subset (\frac{1}{2}, 1]$  such that  $(c - \theta, d - \theta) \subset (0, \frac{1}{2})$ . Letting  $(a, b) = \frac{n}{m}(c - \theta, d - \theta)$ , we see that if  $x_k \in (a, b)$ , then  $y_k \in (c, d)$ . Again, by positive density of  $x_k$ , this happens infinitely often, and so there are infinitely many k with  $x_k \in [0, \frac{1}{2})$  and  $y_k \in (\frac{1}{2}, 1]$ . We conclude as in Case 1 that  $\psi_{j,n}(X) \neq \psi_{\ell,m}(X)$ .

We conclude that the interleaving factor set  $\mathfrak{F}(X)$  consists of all principal decimations, and they are all distinct. Therefore  $\mathfrak{F}(X)$  is infinite.

**Example 6.6.** (A closed X with an infinite factor set  $\mathfrak{F}(X)$ ) The set X constructed in Example 6.5 has infinitely many distinct decimations so its decimation set  $\mathfrak{D}(X)$  and its principal decimation set  $\mathfrak{D}_{prin}(X)$  are infinite. In addition all principal decimations are interleaving factors, so that its factor set  $\mathfrak{F}(X)$  is also infinite.

**Corollary 6.7.** The set  $\mathcal{Y}(\mathcal{A})$  of all infinitely factorizable closed subsets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  is closed under *n*-fold interleaving operations of all  $n \geq 1$ . That is, if  $X_0, X_1, \dots, X_{n-1} \in \mathcal{Y}(\mathcal{A})$ , then

$$(\circledast_n)_{i=0}^{n-1}X_i = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1} \in \mathcal{Y}(\mathcal{A}).$$

*Proof.* The corollary follows using the characterization of membership in  $\mathcal{Y}(\mathcal{A})$  by property (iii) of Theorem 2.13. Property (iii) is inherited under *n*-fold interleaving of sets  $X_i$  that have it.

# 7. Iterated interleaving factorizations of general closed subsets of $\mathcal{A}^{\mathbb{N}}$

We consider iterated interleaving factorizations for general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ . If a set X factors as  $X = X_0 \circledast \cdots \circledast X_{n-1}$ , it is possible that one or more of the factors  $X_j$  can itself be factored. However, unlike with factorizations of positive integers, for example, the further factors that appear at lower levels may not be interleaving factors of the original set X. We therefore name them *iterated interleaving factors*. We define an *iterated interleaving factorization* as follows: The iterated interleaving factorization of depth 0 of a set X is the equation X = X (or the right hand side of such an equation). An iterated interleaving factorization of depth 1 is a single n-fold factorization  $X = (Y_0 \circledast Y_1 \circledast \cdots \circledast Y_{n-1})$  (with parentheses). The  $Y_i$  are iterated interleaving factorizable sets Y on the right hand side of depth k being replaced by interleaving factorizations  $Y = (Y_0 \circledast Y_1 \circledast \cdots \circledast Y_{n-1})$  (with parentheses), for  $n \ge 2$  (allowing different n for different Y). The new added internal factors on the right are assigned depth k + 1; they are inside a nested set of k + 1 parentheses.

7.1. Iterated interleaving factorization trees. An iterated interleaving factorization can be visually represented by a rooted tree, as pictured in Figure 7.1. It has root node X, leaf nodes corresponding to the factors in the iterated interleaving factorization and internal nodes corresponding to intermediate factors.

In our definition of iterated interleaving factorizations, each step is a finite factorization. If an iterated interleaving factor Y at level k has n-fold interleaving factorizations for multiple values of n, it is natural to choose the n-fold factorization with the largest n because this factorization refines all the other possible factorizations of Y, by the divisibility properties of  $\mathcal{N}(X)$  from Theorem 2.12.

How should one treat infinitely factorizable factors? We will adopt the convention in this factorization process that we "freeze" any infinitely factorizable factors encountered, and do not further factorize them. We do this for two reasons. First, for infinitely factorizable Y, no natural choice of n exists for a n-factorization at the next level. Secondly, all interleaving factors of infinitely factorizable sets are also



FIGURE 7.1. Iterated interleaving tree for  $X = (X_{0,4} \circledast ((Z_{0,3} \circledast Z_{1,3} \circledast Z_{2,3}) \circledast Y_{1,2}) \circledast X_{2,4} \circledast X_{3,4})$ , an iterated interleaving factorization of depth 3.

infinitely factorizable by Corollary 5.2, so the factorization process would necessarily proceed forever if we did not freeze any infinitely factorizable factors.

This raises the question: If one factorizes only finitely factorizable sets, will the iterated interleaving factorization process always terminate at a finite depth? We show below that the answer is: there are closed X where the iteration process can go on forever.

7.2. Arbitrary depth factorizations. We show, by construction, that there exist closed sets X having iterated interleaving factorizations of all depths  $k \ge 1$ , with all factors at all depths being finitely factorizable. (Thus so the "freezing" property is never needed).

**Theorem 7.1.** (Infinite depth interleaving factorizations) There exist uncountably many closed sets  $Z_I \subseteq \mathcal{A}^{\mathbb{N}}$ with  $\mathcal{A} = \{0, 1\}$ , indexed by  $I \in \mathcal{A}^{\mathbb{N}}$ , that possess iterated interleaving factorizations of every depth  $k \ge 1$ . They each have a unique iterated interleaving factorization of depth k, for all  $k \ge 1$ . Each  $Z_I$  has an interleaving factor set  $\mathfrak{F}(Z_I)$  containing at most three elements. There exist such I for which the principal decimation set  $\mathfrak{D}_{prin}(Z_I)$  is infinite.

*Proof.* Let  $X_0$  and  $X_1$  be two distinct closed sets in  $\mathcal{A}^{\mathbb{N}}$  having trivial interleaving set  $\mathcal{N}(X_0) = \mathcal{N}(X_1) = \{1\}$ . For definiteness consider  $X_0 = X_F$  the *Fibonacci shift*, consisting of all words which do not have two consecutive 1's, and  $X_2 = X_{AF}$  the *anti-Fibonacci shift*, which consists of all one-sided infinite words which do not contain two consecutive 0's. Example 2.7 showed  $X_F$  has no *n*-fold interleaving factorizations for  $n \geq 2$ , and the proof applies to  $X_{AF}$ . Given an index set  $I = i_0 i_1 i_2 \cdots \in \mathcal{A}^{\mathbb{N}}$ , we define a set

$$Z_{I} = \{ \mathbf{z} \in \mathcal{A}^{\mathbb{N}} : \psi_{2^{r}-1,2^{r+1}}(\mathbf{z}) \in X_{i_{r}} \text{ for } r \ge 0 \}.$$
(7.1)

Let  $\mathbf{z} = z_0 z_1 z_2 \cdots$ . The decimations determine the values of  $z_i$  for subscripts in arithmetic progressions. We represent an arithmetic progression as  $\operatorname{AP}(a; d) = \{n \ge 0 : n \equiv a \pmod{d}\}$ . Then the values  $\mathbf{z}_i$  for  $i \in \operatorname{AP}(2^r - 1; 2^{r+1})$  are restricted by  $\psi_{2^r - 1, 2^{r+1}}(\mathbf{z}) \in X_{i_r}$ . We first show that  $Z_I$  is well-defined.

**Claim 1.** The set of arithmetic progressions  $AP(2^r - 1; 2^{r+1})$  for  $r \ge 0$  form a partition of  $\mathbb{N}$ .

We show by induction on  $r \ge 0$  that  $N_m := \bigsqcup_{r=0}^m \operatorname{AP}(2^r - 1; 2^{r+1}) = \mathbb{N} \smallsetminus \operatorname{AP}(2^{r+1} - 1; 2^{r+1})$ , a disjoint union. The base case r = 0 asserts  $AP[0; 2) = \mathbb{N} \smallsetminus AP(1; 2)$ . The induction step uses  $AP(2^{m+1} - 1; 2^{m+1}) = AP(2^{m+1} - 1, 2^{m+2}) \sqcup \operatorname{AP}(2^{m+2} - 1; 2^{m+2})$ . Finally, the set  $N_m$  contains the interval  $[0, 2^m - 2]$ , so the infinite set union covers  $\mathbb{N}$ , proving claim 1.

**Claim 2.** If  $I \neq J$  then  $Z_I \neq Z_J$ .

If  $I \neq J$  then some  $i_r \neq j_r$ . Then  $\psi_{2^r-1,2^{r+1}}(Z_I) = X_{i_r}$  and  $\psi_{2^r-1,2^{r+1}}(Z_J) = X_{j_r}$  which are distinct since  $X_1 \neq X_2$ . thus  $Z_I \neq Z_J$ , proving claim 2.

**Claim 3.** Each  $Z_I$  is a closed set in  $\mathcal{A}^{\mathbb{N}}$ .

It suffices to show each convergent subsequence of elements of  $Z_I$  has a limit in  $Z_I$ . Convergence is  $\mathcal{A}^{\mathbb{N}}$  is pointwise on each index separately. Suppose  $\mathbf{x}_k \to \mathbf{y}$  in  $\mathcal{A}^{\mathbb{N}}$   $(k \in \mathbb{N})$  as  $k \to \infty$  with each  $\mathbf{x}_k \in Z_I$ . We then have  $\psi_{2^r-1,2^{r+1}}(\mathbf{x}_k) \to \psi_{2^r-1,2^{r+1}}(\mathbf{y})$  in  $\mathcal{A}^{\mathbb{N}}$ . For each  $r \ge 0$  we have  $\psi_{2^r-1,2^{r+1}}(\mathbf{x}_k) \in X_{i_r}$ , hence  $\psi_{2^r-1,2^{r+1}}(\mathbf{x}_k) \to \psi_{2^r-1,2^{r+1}}(\mathbf{y}) \in X_{i_r}$ , since  $X_{i_r}$  is a closed set. The property  $\psi_{2^r-1,2^{r+1}}(\mathbf{y}) \in X_{i_r}$  for all  $r \ge 0$  certifies that  $\mathbf{y} \in Z_I$ , proving claim 3.

Claim 4. Each  $Z_I$  has a 2-fold interleaving factorization

$$Z_I = X_{i_0} \circledast Z_{SI},$$

where  $SI = i_1 i_2 i_3 \cdots$  denotes the one-sided shift of  $I \in \mathcal{A}^{\mathbb{N}}$ .

Using Proposition 3.1 we find

$$\psi_{2^r-1,2^{r+1}}(\mathbf{z}) = \psi_{0,2} \circ \underbrace{\psi_{1,2} \circ \cdots \circ \psi_{1,2}}_{r \text{ times}}(\mathbf{z}),$$

and one proves it by induction on  $r \ge 0$ . Letting  $\mathbf{w} = \psi_{1,2}(\mathbf{z})$ , we have for  $r \ge 1$ 

$$\psi_{2^{r}-1,2^{r+1}}(\mathbf{z}) = \psi_{0,2} \circ \underbrace{\psi_{1,2} \circ \cdots \circ \psi_{1,2}}_{r-1 \text{ times}}(\mathbf{w}) = \psi_{2^{r-1}-1,2^{r}}(\mathbf{w}).$$
(7.2)

By definition

$$Z_{SI} = \{ \mathbf{w} \in \mathcal{A}^{\mathbb{N}} : \psi_{2^{r}-1,2^{r+1}}(\mathbf{w}) \in X_{i_{r+1}} \text{ for } r \ge 0 \}.$$

Now we have, using (7.2),

$$\begin{aligned} Z_I &= \{ \mathbf{z} \in \mathcal{A}^{\mathbb{N}}; \, \psi_{0,2}(\mathbf{z}) \in X_{i_0} \text{ and } \mathbf{w} = \psi_{1,2}(\mathbf{z}) \text{ has } \psi_{2^r - 1, 2^{r+1}}(\mathbf{w}) \in X_{i_{r+1}} \text{ for } r \ge 1 \} \\ &= \{ \mathbf{z} \in \mathcal{A}^{\mathbb{N}}: \, \psi_{0,2}(\mathbf{z}) \in X_{i_0} \text{ and } \psi_{1,2}(\mathbf{z}) \in Z_{SI} \} = X_{i_0} \circledast Z_{SI}, \end{aligned}$$

proving Claim 4.

At this point we obtain an iterated interleaving factorization for  $Z_I$  to arbitrary depth  $k \ge 1$ , by iterating the factorization given in Claim 4. This can be done since one factor is again of the form  $Z_I$  (with a different I). Given I, using the notation  $Z_0 := Z_I$  and  $Z_k := Z_{S^kI}$  we have the depth k factorization

$$Z_{k} = X_{i_{0}} \circledast \left( X_{i_{1}} \circledast \left( \cdots \left( X_{i_{k-2}} \circledast \left( X_{i_{k-1}} \circledast Z_{k} \right) \right) \cdots \right) \right)$$

Figure 7.2 shows a tree corresponding to such an iterated factorization after the fourth level of factoring.



FIGURE 7.2. Iterated interleaving tree for  $Z_0 = (X_{i,0} \circledast (X_{i_1} \circledast (X_{i_2} \circledast (X_{i_3} \circledast Z_4)))).$ 

The remaining part of the proof will show this factorization tree is unique at every level k. Finally a suitable choice of I will lead to  $Z_I$  having infinitely many different principal decimations.

**Claim 5.** The interleaving closure set  $\mathcal{N}(Z_I) = \{1, 2\}$  with associated factor set  $\mathfrak{F}(Z_I) = \{Z_I, X_{i_0}, Z_{SI}\}$ .

It suffices to show that  $Z_I$  has no *n*-fold interleavings with  $n \ge 3$ , in view of Claim 4. We argue by contradiction. Given an *n*-fold interleaving for  $n \ge 3$ , by Theorem 2.12(2), it would also have an lcm(2, n)-fold interleaving, and we set 2m := lcm(2, n) with  $m \ge 2$ . A shuffle identity from Proposition 2.9 gives

$$Z_I = (\circledast_{2m})_{j=0}^{2m-1} X_{i,2m} = \left( (\circledast_m)_{i=0}^{m-1} X_{2i,2m} \right) \circledast \left( (\circledast_m)_{i=0}^{m-1} X_{2i+1,2m} \right)$$

Since 2-fold interleaving factorizations are unique, and  $Z_I = X_{i_0} \otimes Z_{SI}$ , we must have

$$X_{i_0} = (\circledast_m)_{i=0}^{m-1} X_{2i,2m}.$$

This contradicts the fact that  $X_0$  and  $X_1$  have no nontrivial interleaving factorizations, proving claim 5.

**Claim 6.** For  $k \ge 1$ , each  $Z_I$  has a unique iterated interleaving factorization of depth k, whose iterated interleaving factors are  $X_{I_r}$  for  $0 \le r \le k - 1$  and  $Z_{S^k}(I)$ .

This claim follows by induction on  $k \ge 1$ , the base case being the factorization in Claim 4. For the induction step from k to k + 1, all but one of the leaves of the tree (iterated interleaving factors) are of form  $X_i$ , which have no non-trivial interleaving factors, and the remaining factor  $Z_J$ , with  $J = S^k I$ , which has only a 2-fold interleaving factorization  $Z_{S^kI} = X_{i_k} \circledast Z_{S^{k+1}(I)}$ . By updating the list of iterated interleaving factors we complete the induction step. This proves claim 6.

**Claim 7.** If I is strongly aperiodic, meaning that all its shifts  $S^kI$  for  $k \ge 0$  are distinct, then all the decimations of  $Z_I$  of form  $\psi_{2^r-1,2^{r+1}}(Z_I)$  for  $r \ge 0$  are distinct. In particular, the principal decimation set  $\mathfrak{D}_{prin}(Z_I)$  of  $Z_I$  is an infinite set.

We have  $\psi_{2^r-1,2^{r+1}}(Z_I) = Z_{S^rI}$ . By Claim 2 distinct  $S^I$  give distinct  $Z_{S^rI}$ . The strongly aperiodic assumption then makes all  $\psi_{2^r-1,2^{r+1}}(Z_I)$  distinct. They are principal decimations, so  $\mathfrak{D}_{prin}(Z_I)$  is infinite. This proves Claim 7.

**Example 7.2.** (A closed set with an infinite principal decimation set but a finite factor set) Theorem 7.1 exhibited  $Z_I$  that have infinitely many distinct principal decimations;  $\mathfrak{D}_{prin}(Z_I) \subseteq \mathfrak{D}(Z_I)$ . However Claim 5 showed the factor set  $\mathfrak{F}(Z_I)$  is always finite.

*Remark* 7.3. The sets  $Z_I$  in Example 7.2 exhibit the failure of two finiteness properties possessed by all path sets studied in [5]. First, interleaving factorizations of path sets  $\mathcal{P}$  always halt at finite depth (under the freezing convention), while  $Z_I$  never does. Second, path sets  $\mathcal{P}$  always have finitely many different decimations, i.e.  $\mathfrak{D}(\mathcal{P})$  is finite, while this example does not. Example 6.6 gave another example having infinitely many different decimations.

## 8. SHIFT-STABLE AND WEAKLY SHIFT-STABLE SETS

Classical symbolic dynamics is concerned with properties of sets  $X \subseteq A^{\mathbb{N}}$  invariant under the shift operator. The class of such sets is not preserved under decimation or interleaving operations. We study two weaker notions of sets X compatible with the shift operation—*shift-stable sets* and *weakly shift-stable sets* with better properties. Shift-stable sets naturally arise in one-sided dynamics that encode initial conditions, and we show they are closed under all decimations, but not closed under interleaving operations. The wider class of weakly shift-stable sets is closed under all decimation and interleaving operations.

8.1. Shift-stable sets. Recall from Definition 2.14 that a general set  $X \subseteq \mathcal{A}^{\mathbb{N}}$  is *shift-stable* if  $SX \subseteq X$ , and it is *shift-invariant* if SX = X. These definitions allows non-closed sets. Shift-stability is a strictly weaker condition than shift-invariance; see Example 8.5 below.

Shift-stable and shift-invariant sets satisfy the following closure properties under decimation and interleaving closure operations: **Theorem 8.1.** Let  $\mathcal{A}$  be finite alphabet and let  $X \subseteq \mathcal{A}^{\mathbb{N}}$  be a general set (not necessarily closed).

(1) If X is shift-stable (resp. shift-invariant), then all decimations  $\psi_{j,n}(X)$  for  $j \ge 0$ ,  $n \ge 1$  are shift stable (resp. shift-invariant).

(2) If X is shift-stable (resp. shift-invariant) then all n-fold interleaving closures  $X^{[n]}$  with  $n \ge 1$  are shift-stable (resp. shift-invariant).

*Proof.* (1) Shift-stability of X implies  $S^m X \subseteq S^{m-1} X$  whence  $S^m X \subseteq X$  for all  $m \ge 0$ . Now Proposition 3.2 gives

$$S\psi_{j,n}(X) = \psi_{j,n}(S^n X) \subseteq \psi_{j,n}(X)$$

If X is shift invariant, then  $S^m X = X$  for all  $m \ge 0$  and equality holds.

(2) If X is shift stable, then we have, by Proposition 4.10, Proposition 3.2, and (1):

$$SX^{[n]} = S(\psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X))$$
  
$$= \psi_{1,n}(X) \circledast \psi_{2,n}(X) \circledast \cdots \circledast \psi_{n,n}(X)$$
  
$$= \psi_{0,n}(SX) \circledast \psi_{1,n}(SX) \circledast \cdots \circledast \psi_{n-1,n}(SX)$$
  
$$\subseteq \psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X) = X^{[n]}.$$

If X is shift invariant, then all steps hold with equality, as required.

The shift-invariant property restricts the form of interleaving factorizations.

**Proposition 8.2.** (Shift invariance implies self-interleaving) If a general set  $X \subseteq A^{\mathbb{N}}$  is shift-invariant, then all of its interleaving factorizations will be self-interleaving factorizations.

*Proof.* We have for each  $n \ge 1$ , that for  $j \ge 0$ 

$$\psi_{j+1,n}(X) = \psi_{j,n}(SX) = \psi_{j,n}(X)$$

with the leftmost equality generally true by Proposition 3.1 (2) and the second equality from shift invariance. We now have

$$\psi_{j,n}(X) = \psi_{0,n}(X) \quad \text{for} \quad j \ge 0.$$

But by Theorem 2.8 any *n*-fold interleaving  $X = (\circledast_n)_{i=0}^{n-1} X_{i,n}$  has  $X_{i,n} = \psi_{i,n}(X)$ , hence it is a self-interleaving with  $Z_n = \psi_{0,n}(X)$ .

8.2. Closed shift-stable sets. An important feature of closed shift-stable sets is that they are characterized by forbidden blocks, paralleling the definition of two-sided shift spaces in [33, Sec. 1.2]. Let  $\mathcal{A}^*$  denote the set of all finite words in the alphabet  $\mathcal{A}$ , including the empty word. A *block* in an infinite word  $\mathbf{x} = a_0 a_1 a_2 \cdots$  is a finite sequence of consecutive symbols  $a_k a_{k+1} \cdots a_{k+\ell}$ .

**Proposition 8.3.** (Forbidden block characterization of shift-stability) *The following statements about a set*  $X \subseteq A^{\mathbb{N}}$  are equivalent.

- (1) X is closed and shift-stable, i.e. X is closed and  $SX \subseteq X$ .
- (2) X is the set of all infinite words avoiding a (finite or infinite) set  $\mathcal{B}^{\perp} \subseteq \mathcal{A}^*$  of forbidden blocks.

*Remark* 8.4. An analogous result holds in two-sided symbolic dynamics for subsets of  $\mathcal{A}^{\mathbb{Z}}$ , ([33, Theorem 6.1.21]), where shift-stability is replaced by shift invariance, proved with a similar argument. The difference between shift-stability and shift-invariance is discussed in Example 8.7.

*Proof.* (2)  $\Rightarrow$  (1). The set X is closed, since any limit word in the sequence topology will not contain any forbidden block. Now SX is a closed set of infinite words, which do not contain any of the forbidden blocks. It follows that  $SX \subseteq X$ .

 $(1) \Rightarrow (2)$ . The hypothesis  $SX \subseteq X$  implies  $S^k X \subseteq S^{k-1} X \subseteq X$  for all  $k \ge 1$  by induction on k. We let  $\mathcal{B}^{\perp}(X) \subseteq \mathcal{A}^*$  denote all the finite words that do not appear anywhere in any word in X. Let Y denote the set of all infinite words that avoid any block in  $\mathcal{B}^{\perp}(X)$ . By definition  $X \subseteq Y$ . To complete the proof we show the reverse inclusion  $Y \subseteq X$ . Let  $\mathbf{y} = b_0 b_1 b_2 \cdots \in Y$ . By hypothesis the initial word  $b_0 b_1 \cdots b_k \in Y$  does

not contain any element of  $\mathcal{B}^{\perp}(X)$ , so it must occur as a block inside some word  $\mathbf{x} = a_0 a_1 a_2 \cdots \in X$ , for if it did not this would contradict maximality of  $\mathcal{B}^{\perp}(X)$ . Say it is positions  $a_j a_{j+1} \cdots a_{j+k} = b_0 b_1 \cdots b_k$ . Now  $\mathbf{y}_k := S^j \mathbf{x} = .b_0 b_1 \cdots b_k a_{k+1} \cdots \in S^k X \subseteq X$ . We now have a sequence  $\{\mathbf{y}_k : k \ge 0\}$  with  $\mathbf{y}_k \in X$ that converges in the sequence topology to  $\mathbf{y} \in Y$ . Since X is closed, we deduce  $\mathbf{y} \in X$  as required.  $\Box$ 

We give examples of allowed behavior and of non-behavior of closed shift-stable sets.

**Example 8.5.** There exists a shift-stable closed set X which yields an infinite strictly descending chain of inclusions under application of the shift; i.e.:

$$X \underset{\neq}{\supseteq} SX \underset{\neq}{\supseteq} S^2X \underset{\neq}{\supseteq} S^3X \underset{\neq}{\supseteq} \cdots$$

To construct X, define for each  $k \ge 4$  the set  $X_k := (0^{k}1)^k \{000, 111\}^{\mathbb{N}}$ . That is,  $X_k$  has a fixed finite prefix  $(0^k 1)^k$  of length k(k+1) followed by a full 2-block shift

$$Y = \{000, 111\}^{\mathbb{N}}.$$

Note that  $S^3Y = Y$ . We now set

$$X := \bigcup_{k=4}^{\infty} \big(\bigcup_{n=0}^{\infty} S^n X_k\big).$$

The set X is shift-stable, since

$$SX = \bigcup_{k=4}^{\infty} \left( \bigcup_{n=1}^{\infty} S^n X_k \right) \subseteq X.$$

Every element of X is an (eroded) finite prefix followed by a member of Y, SY, or  $S^2Y$ . The set X is closed because the only limit point obtainable in  $\mathcal{A}^{\mathbb{N}}$  from repeated shifts of blocks in the finite prefixes alone is the vector  $0^{\infty}$ , which already belongs to Y.

To show all inclusions are strict, we note for  $0 \le j \le 3$  the set  $S^j X$  contains the word  $0^{4-j}1(0^41)^3(000)^{\infty}$ , which is not contained in any  $S^m X$  for  $m \ge j + 1$ . For  $j \ge 4$  each set  $S^j X$  contains the word  $1(0^j1)^{j-1}(000)^{\infty}$ , which is not contained in any  $S^m X$  for  $m \ge j + 1$ .

**Example 8.6.** (Shift-stability is not always preserved under interleavings) The one-sided Fibonacci shift  $X_F$  having 11 as a forbidden block and the one-sided anti-Fibonacci shift  $X_{AF}$  having 00 as a forbidden block are both closed, shift-invariant sets. We show their 2-fold interleaving  $Y = X_{AF} \otimes X_F$  is not shift-stable. Indeed  $X_{AF}$  allows the initial block 0110, and  $X_F$  allows the initial block 010, whence  $X_{AF} \otimes X_F$  allows the initial block 0011100, so SY contains the initial block 011100. If  $SY \subseteq Y$ , then there is a  $\mathbf{y} = \mathbf{y}_1 \otimes \mathbf{y}_2 \in Y$  with initial block 011100. But this means  $\mathbf{y}_2 \in X_F$  has initial block 110, which is a forbidden block of the Fibonacci shift, a contradiction showing that  $SY \not\subseteq Y$ . (We do have  $S^2Y = Y$ .)

**Example 8.7.** (One-sided shifts) The notion of *one-sided shift* X defined by Lind and Marcus [33, Sect. 12.8] consists of those sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  that are the restriction to positions  $k \ge 0$  of all sequences in a two-sided shift  $X_{\pm}$  described by forbidden blocks. One-sided shifts X are necessarily closed and shift-invariant: SX = X, so they form a strict subclass of closed shift-stable X.

The difference between one-sided shifts and closed shift-stable sets is visible at the level of *minimal* forbidden blocks, which are forbidden blocks that do not contain any other forbidden block as a strict subblock. For a one-sided shift-stable set X we let  $\mathcal{B}_{\min}^{\perp}(X)$  denote its minimal forbidden block set. For a two-sided shift  $X_{\pm}$  we let  $\mathcal{B}_{\min,\pm}^{\perp}(X_{\pm})$  denote its minimal forbidden block set. Now consider the closed set  $Y = \{001^{\infty}, 01^{\infty}, 1^{\infty}\}$  which has  $SY = \{01^{\infty}, 1^{\infty}\} \subset X$ , so is shift-stable but not shift-invariant. Here  $S^2Y = \{1^{\infty}\}$  is shift-invariant. It is easy to check that  $\mathcal{B}_{\min}^{\perp}(Y) = \{1001, 101, 000\}$ . The two-sided shift  $Y^{\pm}$  determined by this set of forbidden blocks is  $Y^{\pm} = \{1^{\mathbb{Z}}\} \in \mathcal{A}^{\mathbb{Z}}$ , because any bi-infinite word that contains a 0 must also contain one of the patterns 101, 1001, 000 and so is excluded. However  $Y_{\pm}$  has minimal forbidden block set  $\mathcal{B}_{\min,\pm}^{\perp}(Y_{\pm}) = \{0\}$  viewed as a two-sided shift. The one-sided shift  $\tilde{Y}$ 

Y and SY cannot be obtained by the Lind and Marcus prescription' their minimal forbidden block sets are not minimal forbidden block sets of any two-sided shift.

8.3. Weakly shift-stable sets. The notion of *weak shift-stability* provides a large class of sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  which respect the shift operator and are closed under all decimation and interleaving operators. This class of sets includes all path sets studied in [2], see [5].

**Definition 8.8.** A general set  $X \subseteq \mathcal{A}^{\mathbb{N}}$  is *weakly shift-stable* if there are  $\ell > k \ge 0$  such that  $S^{\ell}X \subseteq S^{k}X$ . We call  $p = \ell - k$  an *eventual period* for this shift semi-stable set.

The notion of *eventual period* of X reflects the inclusion

 $S^{\ell+j}X = S^{(k+j)+p}X \subseteq S^{k+j}X \quad \text{ for all } \quad j \ge 0.$ 

Theorem 2.15 shows that the class W(A) of all weakly shift-stable sets is closed under all decimation and interleaving operations:

*Proof of Theorem 2.15.* (1) Weak shift-stability  $S^{\ell}X \subseteq S^kX$  gives  $S^{\ell+j}X \subseteq S^{k+j}X$  for all  $j \ge 0$ . Setting  $p = \ell - k$ , we deduce for  $m \ge k$  that

$$S^{m+jp}(X) \subseteq S^m X$$
 whenever  $j \ge 1$ . (8.1)

By Proposition 3.2 we have, for  $j \ge 0, n \ge 1$ ,

$$S^{\ell p}\psi_{j,n}(X) = \psi_{j+\ell pn,n}(X) = \psi_{j,n}(S^{\ell pn}X) \subseteq \psi_{j,n}(S^{kpn}X) = S^{kp}\psi_{j,n}(X)$$

the inclusion holding because  $S^{\ell pn}(X) \subseteq S^{kpn}(X)$  by (8.1), since the difference of iterations is a multiple of p and  $kpn \ge k$ .

(2) Let  $X_j$  be weakly shift-stable with parameters  $(\ell_j, k_j)$ , for  $0 \le j \le n - 1$ , and  $p_j = \ell_j - k_j$ . We assert that  $Y = (\circledast_n)_{i=0}^{n-1} X_i$  is weakly shift-stable with an eventual period  $p = p_0 p_1 \cdots p_j$ . Indeed, setting  $k = \max_j(k_j)$  and  $\ell = k + 1$ , we have, using Proposition 4.10:

$$S^{\ell pn}Y = S^{\ell pn}(X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1})$$
  
=  $(S^{\ell p}X_0) \circledast (S^{\ell p}X_1) \circledast \cdots \circledast (S^{\ell p}X_{n-1})$   
 $\subseteq (S^{kp}X_0) \circledast (S^{kp}X_1) \circledast \cdots \circledast (S^{kp}X_{n-1}) = S^{kpn}X^{[n]}.$ 

The third line above used the inclusions  $S^{\ell p}X_i \subseteq S^{kp}X_i$  for  $0 \le i \le n-1$ , which follow from (8.1), since  $k \ge k_i$ , and  $p_i$  divides p.

(3) We have by Proposition 4.10 and Proposition 3.2:

$$S^{\ell n}X^{[n]} = S^{\ell n} (\psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X))$$
  
$$= S^{\ell} \psi_{0,n}(X) \circledast S^{\ell} \psi_{1,n}(X) \circledast \cdots \circledast S^{\ell} \psi_{n-1,n}(X)$$
  
$$= \psi_{0,n}(S^{\ell n}X) \circledast \psi_{1,n}(S^{\ell n}X) \circledast \cdots \circledast \psi_{n-1,n}(S^{\ell n}X)$$

Now, applying the hypothesis  $S^{\ell}X \subseteq S^kX$ :

$$S^{\ell n} X^{[n]} = \psi_{0,n}(S^{\ell n} X) \circledast \psi_{1,n}(S^{\ell n} X) \circledast \cdots \circledast \psi_{n-1,n}(S^{\ell n} X)$$

$$\subseteq \psi_{0,n}(S^{kn} X) \circledast \psi_{1,n}(S^{kn} X) \circledast \cdots \circledast \psi_{n-1,n}(S^{kn} X)$$

$$= S^{kn}(\psi_{0,n}(X) \circledast \psi_{1,n}(X) \circledast \cdots \circledast \psi_{n-1,n}(X))$$

$$= S^{kn} X^{[n]}.$$

Thus  $X^{[n]}$  is weakly shift-stable.

*Remark* 8.9. Path sets, studied in [2], are closed subsets of  $\mathcal{A}^{\mathbb{N}}$  describable as infinite paths in graphs of finite automata. Such sets are not always shift-stable. In [5] it is shown they are always weakly shift-invariant, so they are weakly shift-stable.

#### 9. ENTROPY OF INTERLEAVINGS FOR GENERAL SETS

We study studies two notions of entropy for general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , topological entropy H(X) and prefix entropy  $H_p(X)$ , defined for all sets X, and we also study a notion of stable prefix topological entropy which only certain sets X possess.

9.1. Topological entropy and prefix topological entropy. We recall two notions of topological entropy for general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , following the paper [2], given in Definition 2.16 and Definition 2.17(1).

(1) The topological entropy of X is

$$H_{\rm top}(X) := \limsup_{k \to \infty} \frac{1}{k} \log N_k(X)$$

where  $N_k(X)$  counts the number of distinct blocks of length k to be found across all words  $\mathbf{x} \in X$ . It is defined as a limsup, but the limit always exists.

(2) The prefix entropy (or path topological entropy) of X is

$$H_p(X) := \limsup_{k \to \infty} \frac{1}{k} \log N_k^I(X)$$

where  $N_k^I(X)$  counts the number of distinct prefix blocks  $b_0b_1 \cdots b_{k-1}$  of length k found across all words  $\mathbf{x} \in X$ .

As remarked in Section 2.6 for  $H_{top}(X)$  the lim sup is always a limit. However the limsup is needed in the definition of prefix entropy, as shown by the next example.

**Example 9.1.** (The limit of  $\frac{1}{k} \log N_k^I(X)$  may not exist) Take  $X_0 = \prod_{j=0}^{\infty} A_j$  where  $A_j = \{0\}$  for  $0 \leq 1$  $j \leq 3$  and, for  $m \geq 1$ ,

- (i)  $\mathcal{A}_j = \{0\}$  for  $2^{2m} \le j \le 2^{2m+1} 1$ (ii)  $\mathcal{A}_j = \{0, 1\}$  for  $2^{2m+1} \le j \le 2^{2m+2} 1$ .

Then  $X_0$  is a closed subset of  $\mathcal{A}^{\mathbb{N}}$  having values  $\frac{1}{k} \log N_k^I(X_0)$  that oscillate between  $\frac{1}{3} \log 2$  and  $\frac{2}{3} \log 2$  infinitely often as  $k \to \infty$ , with minima at  $k = 2^{2m+1}$  and maxima at  $k = 2^{2m+2}$ . Here the lim sup gives  $H_p(X) = \frac{2}{3}\log 2$ . On the other hand, property (ii) implies  $N_k(X_0) = 2^k$  so  $H_{top}(X_0) = \log 2$ .

Example 9.1 shows, first, that  $H_p(X)$  cannot in general be defined as a limit, and second, that  $H_p(X)$ and  $H_{top}(X)$  need not be equal.

**Proposition 9.2.** For general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , the following hold.

(1) Let  $\overline{X}$  denote the closure of X in the natural topology on  $\mathcal{A}^{\mathbb{N}}$ . One has  $H_{top}(X) = H_{top}(\overline{X})$  and  $H_p(X) = H_p(\overline{X}).$ 

(2) One has

 $H_p(X) \leq H_{top}(X) \leq \log |\mathcal{A}|,$ 

*Proof.* (1) The definitions of  $H_{top}(X)$  and  $H_p(X)$  depend only on finite symbol sequences (resp. finite initial symbol sequences) that occur in X. However all infinite words in  $\overline{X} \setminus X$  have all finite symbol sequences (resp. finite initial symbol sequences) occurring for some word in X.

(2) The bounds follow from  $N_k^I(X) \le N_k(X) \le |\mathcal{A}|^k$ . **Example 9.3.** (Strict inequality  $H_p(X) < H_{top}(X)$  may occur for general X) Let  $\mathcal{A} = \{0, 1\}$ , and let the closed set X consist of all words which, for  $m \ge 1$ ,

(i) have symbol 0 in each position  $2^m \le k \le 2^{m+1} - m$ ,

(ii) allow arbitrary symbols  $\{0,1\}$  in positions  $2^{m+1} - (m-1) \le k \le 2^{m+1} - 1$ .

Then  $N_k = 2^k$  for all  $k \ge 1$ , because (ii) gives arbitrarily long blocks of the full shift, whence  $H_{top}(X) =$  $\log 2$ .

On the other hand, for a given symbol position k there are at most  $(\log_2 k)^2$  symbol positions of type (ii), so we obtain  $N_k^I(X) \le 2^{(\log_2 k)^2}$ . It follows that  $H_p(X) = 0$ .

9.2. Entropy and the shift operator. The shift operator preserves both entropies  $H_{top}(X)$  and  $H_p(X)$  separately.

**Proposition 9.4.** For general sets  $X \subseteq A^{\mathbb{N}}$  on a finite alphabet A the following hold. (1) The shift operator S preserves topological entropy:

$$H_{\text{top}}(SX) = H_{\text{top}}(X).$$

(2) The shift operator S preserves prefix entropy:

$$H_p(SX) = H_p(X).$$

Proof. (1) We have, for a finite alphabet,

$$N_k(X) \ge N_k(SX) \ge \frac{1}{|\mathcal{A}|} N_{k+1}(X),$$

since there are at most  $|\mathcal{A}|$  choices for a letter that is dropped. Using a limsup definition for  $H_{top}(X)$  (although the limit always exists) we have

$$H_{\rm top}(SX) = \limsup_{k \to \infty} \frac{1}{k} \log N_k(SX) \le \limsup_{k \to \infty} \frac{1}{k} \log N_k(X) = H_{\rm top}(X).$$

On the other hand,

$$H_{top}(SX) = \limsup_{k \to \infty} \frac{1}{k} \log N_k(SX)$$
  

$$\geq \limsup_{k \to \infty} \left( \frac{1}{k} \log N_{k+1}(X) - \frac{1}{k} \log |\mathcal{A}| \right)$$
  

$$= \limsup_{k \to \infty} \frac{1}{k+1} \log N_{k+1}(X) = H_{top}(X).$$

(2) For a finite alphabet  $\mathcal{A}$  we have

$$N_{k+1}^{I}(X) \ge N_{k}^{I}(SX) \ge \frac{1}{|\mathcal{A}|} N_{k+1}^{I}(X).$$
(9.1)

The result  $H_p(SX) = H_p(X)$  is proved similarly to (1).

9.3. Entropy and decimations. Entropies may change under decimation, subject to the following inequalities.

**Proposition 9.5.** For general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  on a finite alphabet  $\mathcal{A}$  the following hold. Then for all  $n \geq 1$  and all  $i \geq 0$ ,

$$0 \le H_{top}(\psi_{i,n}(X)) \le \min(nH_{top}(X), \log |\mathcal{A}|).$$

and

 $0 \le H_p(\psi_{i,n}(X)) \le \min(nH_p(X), \log |\mathcal{A}|).$ 

All equalities can be attained.

*Proof.* The lower bounds are trivial, and the upper bounds  $\log |\mathcal{A}|$  are trivial. For the upper bounds, the symbols any block of size k of  $\psi_{i,n}(X)$  are contained ( in successive positions with index  $i \pmod{n}$ ) inside a block of length nk of X with the first symbol aligned, hence  $N_k(\psi_{i,n}(X)) \leq N_{nk}(X)$ . We have

$$H_{top}(\psi_{i,n}(X)) = \limsup_{k \to \infty} \frac{1}{k} \log N_k(\psi_{i,n}(X))$$
  
$$\leq \limsup_{k \to \infty} \frac{1}{k} \log N_{nk}(X) \leq n \left(\limsup_{k \to \infty} \frac{1}{k} \log N_k(X)\right) = n H_{top}(X).$$

For the corresponding prefix entropy upper bound we use the bound  $N_k^I(\psi_{i,n}(X)) \leq |\mathcal{A}|^i N_{nk}^I(X)$ , obtained by containment of a prefix of length k in  $\psi_{i,n}(X)$  inside a prefix of X of length nk + i.

To show the bounds are attained, take the interleaved set  $X = (\circledast_n)_{i=0}^{n-1} X_i$  where  $X_0 = \mathcal{A}^{\mathbb{N}}$  and each  $X_i = \{0^{\infty}\}$  for  $1 \le i \le n-1$ . We have  $H_{\text{top}}(X) = H_p(X) = \frac{1}{n} \log |\mathcal{A}|$  (by counting blocks). For the upper bound we have  $H_{\text{top}}(\psi_{0,n}(X)) = H_p(\psi_{0,n}(X)) = \log |\mathcal{A}|$ . For the lower bound  $H_{\text{top}}(\psi_{1,n}(X)) = H_p(\psi_{1,n}(X)) = 0$ .

9.4. **Prefix entropy upper bound for interleaving.** We prove a general upper bound for the prefix entropy of an *n*-fold interleaving in terms of the prefix entropies of its factors, which is Theorem 2.18.

Proof of Theorem 2.18. By definition

$$H_p(X_0 \circledast \cdots \circledast X_{n-1}) = \limsup_{k \to \infty} \frac{1}{k} \log \left( N_k^I(X_0 \circledast \cdots \circledast X_{n-1}) \right), \tag{9.2}$$

where  $N_k^I(X)$  is the number of distinct initial blocks of length k occurring in the symbol sequences of X. Now we partition into subsequences  $\{nk + j : k \ge 0\}$  for  $0 \le j \le n - 1$  to obtain:

$$H_p(X_0 \circledast \cdots \circledast X_{n-1}) = \max_{0 \le j \le n-1} \limsup_{k \to \infty} \frac{1}{nk+j} \log \left( N_{nk+j}^I(X_0 \circledast \cdots \circledast X_{n-1}) \right).$$

Call the terms on the right side

$$H_{p,j}(X) := \limsup_{k \to \infty} \frac{1}{nk+j} \log \left( N_{nk+j}^I(X_0 \circledast \cdots \circledast X_{n-1}) \right)$$

for  $0 \le j \le n-1$ . The number of distinct initial (nk+j)-blocks in  $X_0 \circledast \cdots \circledast X_{n-1}$  is simply the product of the number of distinct initial (k+1)-blocks in each of  $X_0, X_1, \ldots, X_{j-1}$  and of the the distinct initial k-blocks in  $X_j, X_{j+1}, \cdots , X_{n-1}$ . Thus we obtain, for a fixed  $j, 0 \le j \le n-1$ ,

$$H_{p,j}(X) = \limsup_{k \to \infty} \frac{1}{nk+j} \log \left( N_{nk+j}^{I}(X_0 \circledast \cdots \circledast X_{n-1}) \right)$$
  
$$= \limsup_{k \to \infty} \frac{1}{nk+j} \log \left( \prod_{i=0}^{j-1} N_{k+1}^{I}(X_i) \cdot \prod_{i=j}^{n-1} N_{k}^{I}(X_i) \right)$$
  
$$= \limsup_{k \to \infty} \frac{1}{nk+j} \left( \sum_{i=1}^{j-1} \log N_{k+1}^{I}(X_i) + \sum_{i=j}^{n-1} \log N_{k}^{I}(X_i) \right)$$

By (9.1), which applies to general sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , each  $\log N_{k+1}^{I}(X_i)$  differs from  $\log N_{k}^{I}(X_i)$  by no more than  $\log |\mathcal{A}|$ . Since the entire sum is divided by nk + j, this difference does not affect the limsup, so:

$$H_{p,j}(X) = \limsup_{k \to \infty} \frac{1}{nk+j} \sum_{i=0}^{n-1} \log N_k^I(X_i) = \frac{1}{n} \limsup_{k \to \infty} \frac{1}{k} \sum_{i=0}^{n-1} \log N_k^I(X_i)$$
  
$$\leq \frac{1}{n} \sum_{i=0}^{n-1} \limsup_{k \to \infty} \frac{1}{k} \log N_k^I(X_i) = \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i).$$

Thus, all the  $H_{p,j}(X)$  are bounded above by  $\frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i)$ . It follows that  $H_p = \max_{0 \le j \le n-1} H_{p,j}(X)$  obeys the same bound.

**Example 9.6.** (X may have full topological entropy and zero prefix entropy) We start with the closed set  $X_0$  with alphabet  $\mathcal{A} = \{0, 1\}$  defined in Example 9.1. Let a second closed set  $X_1$  consist of all words that allow  $\{0\}$  in index positions where  $X_0$  allows  $\{0, 1\}$ , and allow  $\{0, 1\}$  in all index positions where  $X_0$  allows only  $\{0\}$ ; i.e.,  $X_1 = \prod_{j=0}^{\infty} \mathcal{A}'_j$  where  $\mathcal{A}'_j = \{0, 1\}$  for  $0 \le j \le 3$  and, for  $m \ge 1$ ,

(i) 
$$\mathcal{A}'_{i} = \{0, 1\}$$
 for  $2^{2m} \le j \le 2^{2m+1} - 1$ 

(ii)  $\mathcal{A}'_j = \{0\}$  for  $2^{2m+1} \le j \le 2^{2m+2} - 1$ .

Then  $\mathcal{B}_k(X_1) = \{0,1\}^k$  for all  $k \ge 1$ , since (ii) has arbitrarily long blocks of the full shift, whence  $H_{\text{top}}(X) = \log 2$ . We have  $H_p(X_0) = H_p(X_1) = \frac{2}{3} \log 2$ , by the same calculation as in Example 9.1. We assert that the interleaved set  $X := X_0 \circledast X_1$  has

$$H_p(X) = \frac{1}{2}\log 2 < \frac{1}{2} \left( H_p(X_0) + H_p(X_1) \right) = \frac{2}{3}\log 2.$$

To compute  $H_p(X)$ , note that in each pair of consecutive symbol positions (2j, 2j + 1) the words in X have one symbol frozen to be 0 and the other symbol free to be chosen in  $\{0, 1\}$ , where the frozen symbol is the symbol in position 2j for  $2^{2m} \le j < 2^{2m+1}$  and is the symbol in position 2j + 1 for  $2^{2m+1} \le j < 2^{2m+2}$ . Thus  $2^{k/2-1} \le N_k^I(X) \le 2^{k/2+1}$  for all  $k \ge 0$ , whence  $H_p(X) = \lim_{k \to \infty} \frac{1}{k} \log N_k^I(X) = \frac{1}{2} \log 2$ .

9.5. Stable prefix entropy and interleaving entropy equality. We study the concept of stable prefix entropy and show its consequences for the behavior of entropy under interleaving. Recall from Definition 2.17 (2) that a set  $X \subseteq \mathcal{A}^{\mathbb{N}}$  has *stable prefix entropy*, if the prefix entropy can defined as a limit. That is, the following limit exists:

$$H_p(X) := \lim_{k \to \infty} \frac{1}{k} \log N_k^I(X).$$

Recall that Theorem 2.19 asserts that stable prefix entropy is preserved under interleaving, and that stable prefix entropy of all the interleaving factors implies equality in the prefix entropy formula of Theorem 2.18.

Proof of Theorem 2.19. Let  $X = (\circledast_n)_{i=0}^{n-1} X_i$ . The inequality  $H_p(X) \leq \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i)$  in Theorem 2.18 arose in interchanging a finite sum with a lim sup. Using the stable prefix hypothesis for each  $X_i$ , we obtain a matching lower bound.

By definition  $H_p(X) := \limsup_{k \to \infty} \frac{1}{k} \log N_k^I(X)$ . Let  $H'_p(X) := \liminf_{k \to \infty} \frac{1}{k} N_k(X)$ . It suffices to show that  $H'_p(X) \ge \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i)$  to conclude that  $H'_p(X) = H_p(X)$  has a limit which is the desired value  $\frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i)$ .

Partitioning into subsequences  $\{nk + j : k \ge 0\}$  for  $0 \le j \le n - 1$  as in the proof of Theorem 2.18, we get:

$$H'_p(X) = \min_{0 \le j \le n-1} \left( \liminf_{k \to \infty} \frac{1}{nk+j} \log \left( N^I_{nk+j}(X_0 \circledast \cdots \circledast X_{n-1}) \right) \right).$$

Call the right side values  $H'_{p,i}(X)$ . We have

$$\begin{aligned} H'_{p,j}(X) &\geq \frac{1}{n} \liminf_{k \to \infty} \left( \sum_{i=0}^{n-1} \frac{1}{k} \log N_k^I(X_i) \right). \\ &\geq \frac{1}{n} \sum_{i=0}^{n-1} \liminf_{k \to \infty} \frac{1}{k} \log N_k^I(X_i) \\ &= \frac{1}{n} \sum_{i=0}^{n-1} \lim_{k \to \infty} \frac{1}{k} \log N_k^I(X_i) = \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i), \end{aligned}$$

where stable prefix entropy was used in the last line. We conclude  $H'_p(X) \ge \frac{1}{n} \sum_{i=0}^{n-1} H_p(X_i)$ .

**Example 9.7.** (Stable prefix entropy is not always preserved under decimation) The set  $X = X_0 \circledast X_1$  of Example 9.6 has stable prefix entropy, but  $X_0 = \psi_{0,2}(X)$  does not, as shown in Example 9.1. The set  $X_1 = \psi_{1,2}(X)$  does not have stable prefix entropy by a similar analysis.

Recall that Theorem 2.20 asserts weak shift-stability implies both stable prefix entropy and equality of the two notions of entropy,  $H_p(X)$  and  $H_{top}(X)$ .

*Proof of Theorem 2.20.* For any set X we have  $N_m^I(X) \leq N_m(X)$ . By hypothesis,  $S^{\ell}X \subseteq S^kX$  for some  $\ell \geq k \geq 0$ . Since  $X \subseteq Y$  implies  $S(X) \subseteq S(Y)$ , an easy induction argument shows that  $S^{\ell+j}X \subseteq S^{k+j}X$ holds for all  $j \ge 0$ . Since any block of length m in X, starting in any position n, is an initial block of  $S^n(X)$ , we may conclude that it is an initial block of  $S^{\ell'}(X)$ , for some  $\ell' \leq \ell$ . Consequently all such blocks are counted among the initial blocks of  $X, SX, \dots S^{\ell-1}(X)$  of length m. To each such block one can associate an initial block of length  $m + \ell$  of X which contains the given block in positions  $\ell'$  through  $\ell' + m - 1$ . Any initial block of length  $m + \ell$  can be counted this way at most  $\ell + 1$  times, one for each prefix  $\ell' \leq \ell$ , so we obtain the upper bound

$$N_m(X) \le (\ell+1)N_{m+\ell}^I(X).$$

We then obtain the bounds

$$N_m^I(X) \le N_m(X) \le (\ell+1)|\mathcal{A}|^\ell N_m^I(X),$$

since  $N^I_{m+\ell}(X) \leq |\mathcal{A}|^{\ell} N^I_m(X)$ . It follows that

$$\log N_m^I(X) \le \log N_m(X) \le \log N_m^I(X) + C,$$

for an absolute constant C. Thus

$$\lim_{m \to \infty} \frac{1}{m} \left( \log N_m^I(X) - \log N_m(X) \right) = 0.$$

Since the limit  $\lim_{m\to\infty} \frac{1}{m} \log N_m(X)$  exists for topological entropy, it must also exist for prefix entropy, showing stability. Moreover, since the limits are the same,  $H_p(X) = H_{top}(X)$ . Finally, since weak shiftstability is preserved under n-fold interleaving, the entropy equation (2.10) for topological entropy follows from Theorem 2.19. 

#### **10. CONCLUDING REMARKS**

10.1. General interleaving operations. Iterated interleaving factorizations are a special case of factorizations of closed sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  into a product of closed sets obtained by projections onto subsets of indices  $I_j \subseteq \mathbb{N}$ , where the index sets  $\{I_j : 0 \le j \le n-1\}$  form a partition of  $\mathbb{N}$ . Iterated interleaving factorizations project onto a partition of  $\mathbb{N}$  in which each  $I_i$  is a complete arithmetic progression in  $\mathbb{N}$ .

*Exact covering systems* are partitions of  $\mathbb{N}$  into a finite set of disjoint complete arithmetic progressions (of various moduli). They have been extensively studied, see [21], [42] and [43] for surveys. There are interesting necessary and sufficient conditions for a finite set of complete arithmetic progressions to be an exact cover of  $\mathbb{N}$ , starting with Fraenkel [23], see also Beebee [7] and Porubský and Schónheim [44]. The exact covers determined by iterated interleaving are the set of *natural exact covering systems* introduced by Porubský [41], who credits the construction to an unpublished paper of Znam. It is known that not all exact covers can be obtained by iterated interleaving constructions. An example due to Znam (cf. Guy [28, Problem F14]) is :

$$\{0 \pmod{6}; 1 \pmod{10}; 2 \pmod{15}; 3, 4, 5, 7 - 10, 13 - 16, 19, 20, 22, 23, 25 - 29 \pmod{30}\}$$

This set of arithmetic progressions has gcd(6, 10, 15) = 1, while any iterated interleaving factorization with an initial *n*-fold interleaving necessarily has all arithmetic progressions in any refinement having periods divisible by n. The natural exact covering systems play a special role in the reversion (inversion under composition) of the Möbius function power series, see Goulden et al [27].

One can introduce more general interleaving operations, which might include arbitrary exact covering systems. For a set  $X \subseteq \mathcal{A}^{\mathbb{N}}$ , one can ask which decimations  $\psi_{j,n}(X)$  have the property that X can be written as a topological product  $\psi_{j,n}(X) \times Y$ , where Y is the projection of X onto the set I of all indices having  $i \neq j \pmod{n}$ ? Call such a decimation  $\psi_{i,n}(X)$  with this property a generalized factor of X. Can one characterize the possible sets of all generalized factors of X, as X varies?

10.2. Iterated interleaving closure operations. One may ask for a given set X, what are the set of all interleaving closures of it:  $\{X^{[n]} : n \ge 1\}$ . We can define a filtered limit as  $n \to \infty$  as follows. Letting  $p_k$  denote the kth prime in increasing order, we can define

$$X^{[\infty]} := \lim_{n_k = (p_1 p_2 \cdots p_k)^k \to \infty} X^{[n_k]}$$

where the limit exists since  $X^{[n_k]} \subseteq X^{[n_{k+1}]}$  by Proposition 4.6(3), and for each n one has n divides  $n_k$  for all sufficiently large k. The set  $X^{[\infty]}$  will be infinitely factorizable. What can one say about the possible forms of  $X^{[\infty]}$ ?

10.3. Characterizing closed weakly shift-stable sets. Is there a characterization of closed weakly shift-stable sets  $X \subseteq \mathcal{A}^{\mathbb{N}}$  having a parallel with the characterization by forbidden blocks of closed shift-stable sets given in Proposition 8.3?

## APPENDIX A. INTERLEAVING OPERAD

Operads were systematically developed by Boardman and Vogt [8] and May [37] and as a vehicle to study iterated loop spaces in stable homotopy theory. More recently, operads have been used by researchers in homological algebra, category theory, algebraic geometry, and mathematical physics; see [47] for a brief introduction. Interleaving operations determine a certain kind of operad, giving an application of the operad concept to symbolic dynamics. In this Appendix we only define operads over the category of sets, although they can be defined over any symmetric monoidal category.

Non-symmetric operads (as in [34], [25]) are a weak version of operads which do not require equivariance under actions of symmetric groups on factors. They provide a convenient framework to keep track of properties of an infinite family of n-ary operations under iterated composition.

**Definition A.1.** A *non-symmetric operad* (or *plain operad*)  $\underline{\mathcal{O}}$  consists of a set  $\underline{\mathcal{O}}(n)$  for each natural number n satisfying the following conditions:

(a) (composition) for all positive integers  $n, k_1, \ldots, k_n$ , there is a composition function

$$\circ: \underline{\mathcal{O}}(n) \times \underline{\mathcal{O}}(k_1) \times \cdots \times \underline{\mathcal{O}}(k_n) \to \underline{\mathcal{O}}(k_1 + \cdots + k_n),$$

written as  $(f, f_1, \ldots, f_n) \mapsto f \circ (f_1, \ldots, f_n)$  for elements  $f \in \mathcal{O}(n)$  and  $f_i \in \underline{\mathcal{O}}(k_i)$ ;

(b) (identity) there is an element  $1 \in \underline{\mathcal{O}}(1)$ , called the *identity*, such that

$$f \circ (1, \dots, 1) = f = 1 \circ f$$

for all f;

(c) (associativity) there holds

$$f \circ (f_1 \circ (f_{1,1}, \dots, f_{1,k_1}), f_n \circ (f_{n,1}, \dots, f_{n,k_n})) = = (f \circ (f_1, \dots, f_n)) \circ (f_{1,1}, \dots, f_{1,k_1}, \dots, f_{n,1}, \dots, f_{n,k_n})$$

for all  $f \in \underline{\mathcal{O}}(n)$ ,  $f_i \in \underline{\mathcal{O}}(k_i)$  and  $f_{i,j}$ .

For a non-symmetric operad  $\underline{\mathcal{O}}$ , we think of the elements of  $\underline{\mathcal{O}}(n)$  as *n*-ary operations. An *operad* is a non-symmetric operad that also possesses a right-action of the symmetric group  $\Sigma_n$  on the set of operations of arity *n* for each *n*, satisfying an equivariance condition, as described in the definition below.

Following [35], we use an underline to denote non-symmetric operads  $\underline{O}$  and remove the underline for (symmetric) operads O.

**Definition A.2.** An *operad* (or *symmetric operad*)  $\mathcal{O}$  is a non-symmetric operad together with a right action of the symmetric group  $\Sigma_n$  on each  $\mathcal{O}(n)$  satisfying the following equivariance conditions for each  $\sigma \in \Sigma_n$ ,  $\tau_i \in \Sigma_{k_i}, f \in \mathcal{O}(n)$ , and  $f_i \in \mathcal{O}(k_i)$  for  $1 \le i \le n$ :

(A)  $(f \cdot \sigma) \circ (f_1, \dots, f_n) = (f \circ (f_1, \dots, f_n)) \cdot \sigma;$ (B)  $f \circ (f_1 \cdot \tau_1, \dots, f_n \cdot \tau_n) = (f \circ (f_1, \dots, f_n) \cdot (\tau_1, \dots, \tau_n).$  Here the action of  $\sigma$  on the right-half of (A) is defined as the action of the permutation  $\tilde{\sigma} \in \sigma_{k_1+\dots+k_n}$  that permutes consecutive blocks of length  $k_1, \dots, k_n$ , respectively, according to the permutation  $\sigma$ .

We let  $S(\mathcal{A})$  denote any class of subsets of  $\mathcal{A}^{\mathbb{N}}$  that is closed under all decimation and interleaving operations, combining *n* sets in  $S(\mathcal{A})$  in any order in any *n*-fold interleaving. Examples of such classes include the collection  $\mathcal{W}(\mathcal{A})$  of all weakly shift-stable sets (Theorem 2.15), the sub-collection  $\overline{\mathcal{W}}(\mathcal{A})$  of all closed weakly shift-stable sets (since the property of being closed is preserved under all decimation and interleaving operations), and the class  $C(\mathcal{A})$  of path sets studied in [2], which is shown to satisfy weak shift-stability in [5].

We first construct a non-symmetric operad  $\underline{\mathcal{I}}$  such that each element of  $\underline{\mathcal{I}}(n)$  is an *n*-ary operation acting on  $\mathcal{S}(\mathcal{A}) \times \mathcal{S}(\mathcal{A}) \times \cdots \times \mathcal{S}(\mathcal{A})$  (*n* times). Although the non-symmetric operad  $\underline{\mathcal{I}}$  will be built up from the *n*-fold interleaving operations, the resulting set  $\underline{\mathcal{I}}(n)$  of operations at level *n* will contain many more operations. For notational convenience, let  $\circledast_n$  denote the *n*-fold interleaving operation on  $\mathcal{S}(\mathcal{A})$ . We let  $\underline{\mathcal{I}}(1) = \{\circledast_1\}$ , where of course  $\circledast_1 = id_{\mathcal{S}(\mathcal{A})}$  is the trivial "1-fold interleaving". Also let  $\underline{\mathcal{I}}(2) = \{\circledast_2\}$ . However, it will not be sufficient for  $\underline{\mathcal{I}}(3)$  to be a singleton set. Rather,

$$\underline{\mathcal{I}}(3) = \{ \circledast_3, \circledast_2 \circ (\circledast_1, \circledast_2), \circledast_2 \circ (\circledast_2, \circledast_1) \},\$$

where, for instance,

$$[\circledast_2 \circ (\circledast_1, \circledast_2)](X_1, X_2, X_3) = X_1 \circledast (X_2 \circledast X_3)$$

for general sets  $X_1, X_2, X_3 \in \mathcal{S}(\mathcal{A})$ .  $\underline{\mathcal{I}}(n)$  for n > 3 is defined analogously, so as to satisfy the composition condition of Definition A.1. It is easy to see that  $\circledast_1$  serves as an identity for  $\underline{\mathcal{I}}$  with respect to the various compositions, as in (b). Since the compositions of  $\underline{\mathcal{I}}$  are genuine function composition, associativity in  $\underline{\mathcal{I}}$ follows from the associativity of function composition. Therefore,  $\underline{\mathcal{I}}$  is a non-symmetric operad. We call  $\underline{\mathcal{I}}$ the *interleaving non-symmetric operad*, and refer to operations from  $\underline{\mathcal{I}}$  as *compound interleaving operations*.

The non-symmetric operad  $\underline{\mathcal{I}}$  can be upgraded to a symmetric operad by adding a right action of the symmetric group permuting the interleaving factors. This requires adding additional *n*-ary operations for each *n*. In particular, for  $\sigma \in \Sigma_n$  and an operation  $f \in \mathcal{I}(n)$ , we need to admit the operation  $f \cdot \sigma$  where  $(f \cdot \sigma)(X_1, \ldots, X_n) = f(X_{\sigma(1)}, \ldots, X_{\sigma(n)})$ . Note that, like the interleaving operations themselves, this is also a function  $\mathcal{S}(\mathcal{A}) \times \cdots \times \mathcal{S}(\mathcal{A}) \to \mathcal{S}(\mathcal{A})$ , given by a (possibly compound) interleaving of some permutation of the input sets. Denote by  $\mathcal{I}(n)$  the set of *n*-ary operations expanded to include the operations  $f \cdot \sigma$  defined above, which permute the inputs prior to any (compound) interleaving. Note that we can think of an element  $f \in \underline{\mathcal{I}}(n)$  as corresponding to  $f \cdot \epsilon \in \mathcal{I}(n)$ , where  $\epsilon \in \Sigma_n$  is the identity element. We can then extend the compositions for the  $\underline{\mathcal{I}}(k)$  to

$$\circ: \mathcal{I}(n) \times \mathcal{I}(k_1) \times \cdots \times \mathcal{I}(k_n) \to \mathcal{I}(k_1 + \cdots + k_n)$$

by genuine function composition. Then it is natural to define a right action of  $\Sigma_n$  on  $\mathcal{I}(n)$  by  $(f \cdot \sigma) \cdot \tau = f \cdot (\sigma \tau)$  for  $f \cdot \sigma \in \mathcal{I}(n)$  and  $\tau \in \Sigma_n$ . Note that the equivariance conditions (A) and (B) of Definition A.2 apply generally to an action permuting the inputs of genuine functions with respect to genuine composition. Thus, since the *n*-ary operations in  $\mathcal{I}(n)$  are genuine functions on sets and the compositions are function composition, these conditions hold. We call the resulting (symmetric) operad the *interleaving symmetric operad* and denote it by  $\mathcal{I}$ .

**Proposition A.3.** Let  $\mathcal{I}$  be the interleaving symmetric operad acting on a collection of sets  $\mathcal{S}(\mathcal{A})$  closed under all decimation and interleaving operations. Then for any  $f \in \mathcal{I}(n)$  and any sets  $X_0, \ldots, X_{n-1} \in \mathcal{S}(\mathcal{A})$ , we have also  $f(X_0, \ldots, X_{n-1}) \in \mathcal{S}(\mathcal{A})$ .

*Proof.* Every  $f \in \mathcal{I}(n)$  is just a composition of interleavings of various *n*-arities, where possibly the input sets have their order permuted. Since  $\mathcal{S}(\mathcal{A})$  is closed under the interleaving operations, it follows that it is closed under all composition operations from  $\mathcal{I}$ .

Generally, we recall below the notion of an algebra over an operad. We will see that the descriptions given above for the nonsymmetric operad  $\underline{\mathcal{I}}$  and the (symmetric) operad  $\mathcal{I}$  were really given in terms of

certain algebras over those operads. This approach has helped to keep the exposition concretely rooted in the examples of interest, but differs from the more typical, categorical exposition.

The following definition matches [35, Definition 1.20], restricted to operads in the category of sets. For a set X, let  $\mathcal{E}nd_X(n)$  denote the set of all functions  $X^n \to X$ , and let  $\mathcal{E}nd_X = \bigcup_{n=1}^{\infty} \mathcal{E}nd_X(n)$ . Then  $\mathcal{E}nd_X$  has the structure of an operad, and is called the *Endomorphism Operad* (of sets), see [35, Definition 1.7].

**Definition A.4.** Let  $\mathcal{O}$  be an operad in the category of sets, and let X be a set. An  $\mathcal{O}$ -algebra structure on X is a morphism of operads  $\alpha_X : \mathcal{O} \to \mathcal{E}nd_X$ , that is, a family of  $\Sigma_n$ -equivariant morphisms  $\alpha_X(n) :$  $\mathcal{O}(n) \to \mathcal{E}nd_X(n), n \ge 1$ , compatible with the identity, composition, and equivariance structures of  $\mathcal{O}$  and  $\mathcal{E}nd_X$ .

If we omit the equivariance structure from the above definition, then we get the notion of an *algebra over a nonsymmetric operad*.

#### **Example A.5.** (Algebras over interleaving nonsymmetric operad $\mathcal{I}$ )

The sets  $\mathcal{W}(\mathcal{A})$  of all weakly shift-stable sets on the finite alphabet  $\mathcal{A}, \overline{\mathcal{W}}(\mathcal{A})$  of all closed weakly shiftstable sets on  $\mathcal{A}$ , and  $\mathcal{C}(\mathcal{A})$  of path sets on  $\mathcal{A}$  are all algebras over the interleaving nonsymmetric operad  $\underline{\mathcal{I}}$ . If the set  $\mathcal{S}(\mathcal{A})$  is any of these sets, and for any  $n \in \mathbb{N}$ , the maps  $\alpha_{\mathcal{S}(\mathcal{A})}$  of Definition A.4 are built up from

$$\alpha_{\mathcal{S}(\mathcal{A})}(n)(\circledast_n)[(X_0,\ldots,X_{n-1})] := (\circledast_n)_{j=0}^{n-1} X_j = X_0 \circledast X_1 \circledast \cdots \circledast X_{n-1}$$

by function composition, where  $(X_0, \ldots, X_{n-1}) \in \mathcal{S}(\mathcal{A})^n$ .

#### REFERENCES

- W.C. Abram, A. Bolshakov, J.C. Lagarias, Intersection of multiplicative translates of 3-adic Cantor sets II: Two infinite families, Exp. Math. 26 (2017) 468–489.
- [2] W.C. Abram, J.C. Lagarias, Path sets in one-sided symbolic dynamics, Adv. Appl. Math. 56 (2014) 109-134.
- [3] W.C. Abram, J.C. Lagarias, p-Adic path set fractals and arithmetic, J. Fractal Geom. 1 (2014) 45-81.
- [4] W.C.Abram, J.C. Lagarias, Intersections of multiplicative translates of 3-adic Cantor sets, J. Fractal Geom. 1 (2014) 349–390.
- [5] W.C. Abram, J.C. Lagarias, D.J. Slonim, Interleaving of path sets, paper in preparation.
- [6] R. Adler, A.G. Konheim, M.H. McAndrew, Topological entropy, Trans. Amer. Math. Soc. 114 (1965) 309–319.
- [7] J.Beebee, Bernoulli numbers and exact covering systems, Amer. Math. Monthly 99 (1992) 946–948.
- [8] J.M. Boardman, R.M. Vogt, Homotopy invariant algebraic structures on topological spaces, Lecture Notes in Math. 347, Springer, Berlin, 1973.
- [9] M.R. Bremner, V. Dotsenko, Algebraic Operads: An Algorithmic Companion, CRC Press, Boca Raton, FL, 2014.
- [10] S.D. Cardell, A. Fuster-Sabatier, A.H. Ranea, Linearity in decimation-based generators: An improved cryptanalysis of the shrinking generator, Open Math. 16 (2018), 646–655.
- [11] T.M. Carlsen, Cuntz-Pimsner C\*-algebras associated with subshifts, Internat. J. Math. 9 (2008) 47–70.
- [12] T.M. Carlsen, An introduction to the C\*-algebra of a one-sided shift space, Operator algebras and dynamics, 63–88, Springer Proc. Math. Stat. 58, Springer, Heidelberg 2013.
- [13] T. Cottrell, A study of Penon weak n-categories, Part 1. Monad interleaving, Cah. Topol. Géom Différ. Catég. 59 (2018) 197–259.
- [14] J. Cuntz, A class of  $C^*$  algebras and topological Markov chains II. Reducible chains and the Ext-function for  $C^*$ -algebras, Invent. Math. 63 (1981) 25–40.
- [15] J. Cuntz, W. Krieger, A class of C\*- algebras and topological Markov chains, Invent. Math. 56 (1980) 251–268.
- [16] I. Daubechies, Ten Lectures on Wavelets, CBMS-NSF Regional Conf. Ser. in Appl. Math. 61, SIAM, Philadelphia, PA, 1992.
- [17] M. Dokuchaev, R. Exel, Partial actions and subshifts, J. Funct. Anal. 272 (2017) 5038–5106.
- [18] G. Duchamp, M. Flouret, É. Laugerotte, J.G. Luque, Direct and dual laws for automata with multiplicities, Theoret. Comput. Sci. 267 (2001), 105–120.
- [19] S. Eilenberg, Automata, languages and machines, Volume A, Pure and Applied Mathematics, Vl. 59A, Academic Press, New York, 1974.
- [20] P. Erdős, Some unconventional problems in number theory, Math. Mag. 52 (1979), 67–70.
- [21] P. Erdős, R.L. Graham, Old and new problems and results in combinatorial number theory, Monographic No. 28 de L'Enseignements Mathématique Univ. Genéve, 1980.
- [22] R. Exel, Partial dynamical systems, Fell bundles and applications, Math. Surveys Monogr. vol. 224, Amer. Math. Soc. Providence, RI, 2017.
- [23] A. Fraenkel, A characterization of exactly covering congruences, Discrete Math. 4 (1973) 359–366.

- [24] S. Giraudo, Operads in algebraic combinatorics, Habilitation 2017, Univsité Paris-Est Marne-la-Vallée, 388 pages. eprint: arXiv:1712.03782.
- [25] S. Giraudo, Nonsymmetric operads in combinatorics, Springer, Cham, 2018.
- [26] G. Grätzer, Lattice theory: Foundation, Birkhäuser, Berlin, 2011.
- [27] I.P. Goulden, A. Granville, L.B. Richmond, J. Shallit, Natural exact covering systems and the reversion of the Möbius series, Ramanujan J. 50 (2019) 211–235.
- [28] R.K. Guy, Unsolved Problems in Number Theory: Third Edition, Springer-Verlag, New York, 2004.
- [29] Zhiqing Hui, J. Baccou, J. Liandrat, On the coupling of decimation operator with subdivision schemes for multi-scale analysis, Mathematical methods for curves and surfaces, 162–185, Lecture Notes in Computer Science 10521, Springer, Cham 2017.
- [30] D. Krieger, A. Miller, N. Rampersad, B. Ravikumar, J.O. Shallit, Decimations of languages and state complexity, Theoret. Comput. Sci. 410 (2009) 2401–2409.
- [31] J.C. Lagarias, Ternary expansions of powers of 2. J. London Math. Soc. 79 (2009) 562–588.
- [32] T. Leinster, Higher operads, higher categories, London Math. Soc. Lecture Note Series, 298. Cambridge University Press, Cambridge, 2004.
- [33] D. Lind, B. Marcus, An Introduction to Symbolic Dynamics and Coding, Cambridge University Press, New York, 1995. (Reprinted 1999 with corrections.)
- [34] Jean-Louis Loday, Bruno Vallette, Algebraic operads, Grundlehren der mathematischen Wissenschaften, Vol. 346, Springer-Verlag, New York 2012.
- [35] M. Markl, S. Shnider, J. Stasheff, Operads in algebra, topology and physics. Math. Surveys Monogr. 96. American Mathematical Society, Providence, R,I 2002.
- [36] M. Markl, Operads and PROPS, pp. 87–140 in: Handbook of algebra, Vol. 5, Elsevier/North Holland, Amsterdam, 2008.
- [37] J.P. May, The Geometry of Iterated Loop Spaces. Springer Lecture Notes in Mathematics, Vol. 271, Springer, 1972.
- [38] J.P. May, Operads, algebras and modules, in Operads: Proceedings of Renaissance Conferences (Hartford, CT/Luminy, 1995), 15–31, Contemp. Math., 202, Amer. Math. Soc., Providence, RI, 1997.
- [39] M. Morse, G. A.Hedlund, Symbolic dynamics Amer. J. Math. 60 (1938) 815-866.
- [40] D. Perrin, J-E. Pin, Infinite Words: Automata, Semigroups, Logic and Games, Elesevier, Dordrecht, 2004.
- [41] S. Porubský, Natural exactly covering systems of congruences, Czechoslovak Math. J. 24 (1974) 598-606.
- [42] S. Porubský, Results and problem on covering systems of residue classes, Mitteilungen Math. Sem. Giessen 150 (1981) 1–85.
- [43] S. Porubský, J. Schönheim, Covering systems of Paul Erdős. Past, present and future, pp. 581–627 in: Paul Erdős and his mathematics I. (Budapest, 1999), J. Bolyai Math. Soc., Budapest 2002.
- [44] S. Porubský, J. Schönheim, Old and new necessary and sufficient conditions on  $(a_i, m_i)$  in order that  $n \equiv a_i \pmod{m_i}$  be a covering system, Math. Slovaca 53 (2003) 341–349.
- [45] R.A. Rueppel, Analysis and Design of Stream Ciphers, Springer-Verlag, Berlin, 1986.
- [46] E. Schechter, Handbook of Analysis and Its Foundations, Academic Press, New York, 1996. (Elsevier, Dordrecht, 1997.)
- [47] J. Stasheff, What is an operad?, Notices Amer. Math. Soc. 51 (2004) 630-631.
- [48] S.A. Vanstone, P. van Oorschot, An Introduction to Error Correcting Codes with Applications, Kluwer Academic Publishers, Boston, 1989.

ALLEN BAILEY & ASSOCIATES, AUSTIN, TX 78731 *Email address*: abramwc@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1043, USA *Email address*: lagarias@umich.edu

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47906 *Email address*: dslonim@purdue.edu