BOUNDS FOR THE ORDER OF AUTOMORPHISM GROUPS OF CYCLIC COVERING FIBRATIONS OF A RULED SURFACE

HIROTO AKAIKE

ABSTRACT. We study the order of automorphism groups of cyclic covering fibrations of a ruled surface. Arakawa and later Chen studied it for hyperelliptic fibrations and gave the upper bound. The purpose of present paper is to pursue the analog for cyclic covering fibrations.

INTRODUCTION

Let $f: S \to B$ be a surjective morphism from a complex smooth projective surface S to a smooth projective curve B with connected fibers. We call it a *fibration* of genus g when a general fiber is a curve of genus g. A fibration is called *relatively minimal*, when any (-1)-curve is not contained in fibers. Here we call a smooth rational curve C with $C^2 = -n$ a (-n)-curve. A fibration is called *smooth* when all fibers are smooth, *isotrivial* when all of the smooth fibers are isomorphic, *locally trivial* when it is smooth and isotrivial.

An automorphism of the fibration $f: S \to B$ is a pair of automorphisms $(\kappa_S, \kappa_B) \in$ Aut $(S) \times$ Aut(B) satisfying $f \circ \kappa_S = \kappa_B \circ f$, that is, the diagram

$$\begin{array}{ccc} S & \xrightarrow{\kappa_S} & S \\ f & & & \downarrow f \\ B & \xrightarrow{\kappa_B} & B \end{array}$$

is commutative. We denote by $\operatorname{Aut}(f)$ the group of all automorphisms of f. Arakawa [1] and later Chen [2] studied it for hyperelliptic fibrations and gave the upper bound of the order of a finite subgroup of $\operatorname{Aut}(f)$ in terms of g, g(B) (the genus of B) and K_f^2 (the square of the relative canonical bundle). The purpose of the present paper is to pursue the analog for primitive cyclic covering fibrations of type (g, 0, n) introduced by Enokizono in [3]. Roughly, they are *n*-fold cyclic branched coverings of ruled surfaces and hyperelliptic fibrations are nothing more than those with n = 2.

Let $f: S \to B$ be a primitive cyclic covering fibration of type (g, 0, n) and G a finite subgroup of $\operatorname{Aut}(f)$. Then G can be expressed as the extension of its horizontal part Hby its vertical part K, that is, $1 \to K \to G \to H \to 1$ (exact). Here H is a subgroup of $\operatorname{Aut}(B)$ consisting of κ_B such that one can find a $\kappa_S \in \operatorname{Aut}(S)$ satisfying $(\kappa_S, \kappa_B) \in G$,

²⁰²⁰ Mathematical Subject Classification. Primary 14J50, Secondary 14J29.

Key words and phrases. fibration, automorphism group.

while K naturally acts on fibers of f as a subgroup of $\operatorname{Aut}(S/B)$. We can study H by using known results for the automorphism groups of curves. Hence it suffices to manage K. Note that we have the canonical cyclic subgroup of $\operatorname{Aut}(S/B)$ of order n, the covering transformation group, with a (singular) ruled surface as the quotient by its action. We may assume that K contains it. Then K induces a subgroup \widetilde{K} of the automorphism group of the ruled surface preserving the branch locus and our task is reduced to estimating the order of \widetilde{K} . For this purpose, we use the localization of the invariant K_f^2 . It is known that K_f^2 can be localized to fibers of f (Lemma 1.7 of [3], or Lemma 2.5). We refine the localized K_f^2 further by using the quantity defined at a point on a fiber, and obtain the new expression of it in Proposition 2.6. Then, we estimate the order of \widetilde{K} from above with the localized K_f^2 multiplied by an explicit function in g and n (Propositions 4.1, 4.2 and 4.3).

Now, the main results can be stated as follows.

Theorem 0.1 (Theorem 4.4). Let $f: S \to B$ be a non-isotrivial primitive cyclic covering fibration of type (g, 0, n) with $n \ge 4$ and $r := \frac{2g}{n-1} + 2$. Assume $\begin{cases} r \ge 3n & (n \ge 5), \\ r \ge 13 & (n = 4). \end{cases}$

Put

$$\mu_{g,n} := \frac{2n^3(2g+n-1)(g+n-1)}{(n-1)\left(n(n+1)g+n(n-1)-4(n-1)^2\right)(g-1)}$$

Let G be a finite subgroup of Aut(f). Then it holds

$$\sharp G \leq \begin{cases} 6(2g(B) - 1)\mu_{g,n}K_f^2 & (g(B) \ge 1) \\ 5\mu_{g,n}K_f^2 & (g(B) = 0). \end{cases}$$

Theorem 0.2 (Corollary 4.5). Let $f : S \to B$ be a non-locally trivial primitive cyclic covering fibration of type (g, 0, n) with $n \ge 4$. Assume $\begin{cases} r \ge 3n & (n \ge 5), \\ r \ge 13 & (n = 4). \end{cases}$

Then it holds

$$\# \operatorname{Aut}(S/B) \le \frac{n^3(2g+n-1)(g+n-1)}{(n-1)\left(n(n+1)g+n(n-1)-4(n-1)^2\right)(g-1)} K_f^2.$$

Unfortunately, ours cannot cover the case n = 3. So, we leave it as a future problem.

In Sections 1 and 2, we recall basic properties of primitive cyclic covering fibrations mainly due to [3]. Among other things, we recast Section 5 of [3] and consider the refined localization of K_f^2 . In Section 3, we study automorphisms of primitive cyclic covering fibrations of type (g, 0, n) to reduce the problem to that on a ruled surface. Especially we discuss whether it is possible to perform a sequence of equivariant blowing-downs (with respect to \tilde{K}) in order to get a geometrically ruled surface on which the branch locus has only singular points of standardized multiplicities. In Section 4, we show our main result, Theorem 4.4. Section 5 is devoted to construct an example that shows, at least for fiber germs, that our estimate is almost optimal.

Acknowledgement

The author is deeply grateful to Professors Kazuhiro Konno and Osamu Fujino for their valuable advices and supports. The author also thanks to Doctor Makoto Enokizono for his precious advices. The research is supported by JSPS KAKENHI No. 20J20055.

CONTENTS

| Int | 1 | |
|------------|--------------------------------------|----|
| 1. | Primitive cyclic covering fibrations | 3 |
| 2. | Localization of K_f^2 | 5 |
| 3. | Automorphism of a fibered surface | 12 |
| 4. | Upper bounds of the order. | 15 |
| 5. | An example | 26 |
| References | | |

1. PRIMITIVE CYCLIC COVERING FIBRATIONS

We recall basic properties of primitive cyclic covering fibrations, most of which can be found in [3].

Definition 1.1. Let $f: S \to B$ be a relatively minimal fibration of genus $g \ge 2$. We call it a primitive cyclic covering fibration of type (g, 0, n), when there are a (not necessarily relatively minimal) fibration $\tilde{\varphi}: \tilde{P} \to B$ of genus 0 (*i.e.* ruled surface) and a classical *n*-cyclic covering

$$\widetilde{\theta}: \widetilde{S} = \operatorname{Spec}_{\widetilde{P}}\left(\bigoplus_{j=0}^{n-1} \mathcal{O}_{\widetilde{P}}(-j\widetilde{\mathfrak{d}})\right) \to \widetilde{P}$$

branched over a smooth curve $\widetilde{R} \in |n\widetilde{\mathfrak{d}}|$ for some $n \geq 2$ and $\widetilde{\mathfrak{d}} \in \operatorname{Pic}(\widetilde{P})$ such that f is the relatively minimal model of $\widetilde{f} = \widetilde{\varphi} \circ \widetilde{\theta}$.

In addition, we employ the following notation. We denote by $\Sigma = \langle \tilde{\sigma} \rangle$ the covering transformation group of $\tilde{\theta}$ and by $\varphi : P \to B$ a relatively minimal model of $\tilde{\varphi} : \tilde{P} \to B$ with the natural contraction map $\tilde{\psi} : \tilde{P} \to P$. Furthermore, $\tilde{F}, F, \tilde{\Gamma}$ and Γ will denote general fibers of $\tilde{f}, f, \tilde{\varphi}$ and φ , respectively. To specify it is a fiber over $p \in B$, we write F_p , etc.

Remark 1.2. We note important properties of primitive cyclic covering fibrations in the following, which can be found in [3].

• If σ is the automorphism of S over B induced by a generator $\tilde{\sigma}$ of Σ , then we can assume that the natural morphism $\tilde{S} \to S$ is a minimal succession of blowing-ups that resolves all isolated fixed points of σ .

- $\tilde{\varphi}$ -vertical components of \tilde{R} is a disjoint union of non-singular rational curves each of which is a (-an)-curve for some positive integer a.
- Arbitrary $\tilde{\varphi}$ -vertical (-1)-curve and \tilde{R} meet.

From now on, we let $f: S \to B$ be a primitive cyclic covering fibration of type (g, 0, n)and freely use the above notation and convention.

Since the restriction map $\tilde{\theta}|_{\widetilde{F}} : \widetilde{F} \to \widetilde{\Gamma} \simeq \mathbb{P}^1$ is a classical *n*-cyclic covering branched over $\widetilde{R} \cap \widetilde{\Gamma}$, the Hurwitz formula for $\tilde{\theta}|_{\widetilde{F}}$ gives us

(1.1)
$$r := \widetilde{R}\widetilde{\Gamma} = \frac{2(g+n-1)}{n-1}.$$

From $\widetilde{R} \in |n\widetilde{\mathfrak{d}}|$, it follows that r is a multiple of n.

Since $\tilde{\psi}: \tilde{P} \to P$ is a composite of blowing-ups, we can write $\tilde{\psi} = \psi_1 \circ \cdots \psi_N$, where $\psi_i: P_i \to P_{i-1}$ denotes the blowing-up at $x_i \in P_{i-1}$ $(i = 1, \cdots, N), P_0 = P$ and $P_N = \tilde{P}$. We define a reduced curve R_i inductively as $R_{i-1} = (\psi_i)_* R_i$ starting from $R_N = \tilde{R}$ down to $R_0 = R$. We call R_i branch locus on P_i . We also put $E_i = \psi_i^{-1}(x_i)$ and $m_i = \text{mult}_{x_i} R_{i-1}$ $(i = 1, \cdots, N)$.

Lemma 1.3. In the above situation, the following hold for any $i = 1, \dots, N$.

- (1) Either $m_i \in n\mathbb{Z}_{\geq 1}$ or $n\mathbb{Z}_{\geq 1} + 1$, where $\mathbb{Z}_{\geq 1}$ is the set of positive integers. Furthermore, $m_i \in n\mathbb{Z}_{\geq 1}$ if and only if E_i is not contained in R_i .
- (2) $R_i = \psi_i^* R_{i-1} n[\frac{m_i}{n}] E_i$, where [t] denotes the greatest integer not exceeding t.
- (3) There exists a $\mathfrak{d}_i \in \operatorname{Pic}(P_i)$ such that $\mathfrak{d}_i = \psi_i^* \mathfrak{d}_{i-1} [\frac{m_i}{n}] E_i$ and $R_i \sim n \mathfrak{d}_i, \ \mathfrak{d}_N = \widetilde{\mathfrak{d}}$.

We say that a singular point of an R_i is of type $n\mathbb{Z}$ (resp. $n\mathbb{Z}+1$) if its multiplicity is in $n\mathbb{Z}_{\geq 1}$ (resp. $n\mathbb{Z}_{\geq 1}+1$).

Lemma 1.4. Let $f: S \to B$ be a primitive cyclic covering fibration of type (g, 0, n). Then there is a relatively minimal model $\varphi: P \to B$ of $\tilde{\varphi}: \tilde{P} \to B$ such that

$$\operatorname{mult}_{x} R_{h} \le \frac{r}{2} = \frac{g}{n-1} + 1$$

for all $x \in R_h$, where R_h denotes the φ -horizontal part of R. Moreover if $\operatorname{mult}_x R > \frac{r}{2}$, then $\operatorname{mult}_x R \in n\mathbb{Z} + 1$.

Later, we will need to pay a special attention to the action of a finite subgroup of $\operatorname{Aut}(\widetilde{P}/B)$ preserving \widetilde{R} . We show in Lemma 3.3 that there is a relatively minimal model $\varphi: P \to B$, compatible with such an action, which satisfies the inequality in Lemma 1.4. So we assume that $\varphi: P \to B$ always satisfies Lemma 1.4 in what follows.

2. Localization of K_f^2

We reconsider the argument in Section 5 of [3] and give localizations of K_f^2 . Let $f : S \to B$ be a primitive cyclic covering fibration of type (g, 0, n).

First of all, let us recall the singularity index [3]. For any fixed $p \in B$, we consider all singular points (including infinitely near ones) of R on Γ_p . For any positive integer k, we let $\alpha_k(\Gamma_p)$ be the number of singular points of multiplicity either kn or kn + 1 among them. We put $\alpha_k := \sum_{p \in B} \alpha_k(\Gamma_p)$ and call it the *k*-th singularity index of the fibration. Let A be the sum of all (-n)-curves contained in \widetilde{R} and put $\widetilde{R}_0 := \widetilde{R} - A$. Then the 0-th singularity index is defined by $\alpha_0 := (K_{\tilde{\varphi}} + \widetilde{R}_0)\widetilde{R}_0$.

For any $p \in B$, we denote the number of isolated (-n)-curves in $\widetilde{\Gamma}_p$ by $\varepsilon(\Gamma_p)$ and put

$$\varepsilon := \sum_{p \in B} \varepsilon(\Gamma_p).$$

Let $\widehat{P} \to B$ be any intermediate ruled surface between \widetilde{P} and P, and regard $\widetilde{\psi} : \widetilde{P} \to P$ as the composite of the natural birational morphisms $\check{\psi} : \widetilde{P} \to \widehat{P}$ and $\widehat{\psi} : \widehat{P} \to P$ as $\widetilde{\psi} = \widehat{\psi} \circ \check{\psi}$:



We put $\widehat{R} = \widehat{\psi}_* \widetilde{R}$. The fiber of $\widehat{\varphi}$ over $p \in B$ will be denoted by $\widehat{\Gamma}_p$.

Let $\widehat{\alpha}_k(\Gamma_p)$ and $\check{\alpha}_k(\Gamma_p)$ be the number of the singular points of R on Γ_p appearing in $\widehat{\psi}$ and $\check{\psi}$, respectively. We note that the number of singular points of \widehat{R} on $\widehat{\Gamma}_p$ is counted by $\widehat{\alpha}_k(\Gamma_p)$. Then $\alpha_k(\Gamma_p) = \widehat{\alpha}_k(\Gamma_p) + \check{\alpha}_k(\Gamma_p)$. We put $\widehat{\alpha}_k := \sum_{p \in B} \widehat{\alpha}_k(\Gamma_p)$ and $\check{\alpha}_k := \sum_{p \in B} \check{\alpha}_k(\Gamma_p)$.

For an effective vertical divisor T and $p \in B$, we denote the biggest subdivisor of T whose support is in the fiber over p by T(p). Then $T = \sum_{p \in B} T(p)$. We sometimes write $\sharp T$ to indicate the number of irreducible components of T.

Choose and fix $p \in B$ and $\hat{z} \in \widehat{\Gamma}_p$. We consider the vertical part $\widetilde{R}_v = \sum_{p \in B} \widetilde{R}_v(p)$ of \widetilde{R} with respect to $\widetilde{\varphi} : \widetilde{P} \to B$. We let $\widetilde{R}_v(p)_{\hat{z}}$ be the biggest subdivisor of $\widetilde{R}_v(p)$ contracted to \hat{z} by $\hat{\psi}$. Note that we have $\widetilde{R}_v(p)_{\hat{z}} \neq 0$ only when there is a singular point (of the branch locus) whose multiplicity is in $n\mathbb{Z} + 1$ (and > 1) and which is infinitely near to \hat{z} (including \hat{z} itself). Note also that $\widetilde{R}_v(p)_{\hat{z}}$ is a disjoint union of non-singular rational curves each of which is a (-an)-curve for some positive integer a.

To decompose $\widetilde{R}_v(p)_{\widehat{z}}$, we define a family $\{L_i\}_i$ consisting of vertical irreducible curves in $\widetilde{R}_v(p)_{\widehat{z}}$ as follows.

(i) Choose and fix a (-1)-curve E_1 over \hat{z} on a ruled surface between \tilde{P} and \hat{P} , and let L_1 be the proper transform of E_1 on \tilde{P} .

(ii) For $i \ge 2$, L_i is the proper transform of an exceptional (-1)-curve E_i that is contracted to a "point" x_i on E_k or its proper transform for some k < i ("point" does NOT include an infinitely near point on them).

Put $\mathfrak{L} := \{\{L_i\}_i \mid \{L_i\}_i \text{ satisfies (i) and (ii)}\}$. If $\{L_i\}_i$ and $\{L'_j\}_j$ $(\{L_i\}_i, \{L'_j\}_j \in \mathfrak{L})$ have a common curve, one can show the union of them $\{L_i, L'_j\}_{i,j}$ is in \mathfrak{L} . We define a partially order $\{L_i\}_i \leq \{L'_j\}_j$ if $\{L_i\}_i \subset \{L'_j\}_j$. Then (\mathfrak{L}, \leq) is a partially order set, so there exist maximal elements $\{L_{1,k_1}\}_{k_1}, \cdots, \{L_{\eta_{\widehat{z}},k_{\eta_{\widehat{z}}}}\}_{k_{\eta_{\widehat{z}}}}$, where $\eta_{\widehat{z}}$ is the number of maximal elements. It's an abuse of the symbol, but we replace indices $k_1, \cdots, k_{\eta_{\widehat{z}}}$ to k. We note that each two $\{L_{1,k}\}_k, \cdots, \{L_{\eta_{\widehat{z}},k}\}_k$ have a no common curve. We put $D_t := \sum_{k\geq 1} L_{t,k}$ for $t = 1, \cdots, \eta_{\widehat{z}}$.

We can describe $\widetilde{R}_v(p)_{\widehat{z}}$ by above D_t 's as the following. $\widetilde{R}_v(p)_{\widehat{z}}$ is decomposed a disjoint sum consisted of such sum uniquely. We denote as

(2.1)
$$\widetilde{R}_v(p)_{\widehat{z}} = D_1 + \dots + D_{\eta_{\widehat{z}}} \quad D_t = \sum_{k \ge 1} L_{t,k}.$$

Let $C_{t,k}$ be the exceptional (-1)-curve whose proper transform to \widetilde{P} is $L_{t,k}$.

For a positive integer a, we denote by $j_a^t(\Gamma_p)_{\hat{z}}$ the number of (-an)-curves in D_t , and put

$$(2.2) j_a(\Gamma_p)_{\widehat{z}} := \sum_{t=1}^{\eta_{\widehat{z}}} j_a^t(\Gamma_p)_{\widehat{z}}, \quad j^t(\Gamma_p)_{\widehat{z}} := \sum_{a\geq 1} j_a^t(\Gamma_p)_{\widehat{z}}, \quad j(\Gamma_p)_{\widehat{z}} := \sum_{t=1}^{\eta_{\widehat{z}}} j^t(\Gamma_p)_{\widehat{z}}.$$

Let $\widetilde{\Gamma}_{p,\widehat{z}}$ be the biggest subdivisor of $\widetilde{\Gamma}_p$ which is contracted to \widehat{z} by $\widehat{\psi}$. We put (2.3)

$$\alpha_0^+(\Gamma_p)_{\widehat{z}} := \begin{cases} \widetilde{R}_h \widetilde{\Gamma}_{p,\widehat{z}} - \sharp(\operatorname{Supp}(\widetilde{R}_h) \cap \operatorname{Supp}(\widetilde{\Gamma}_{p,\widehat{z}})) & \text{(if } \widehat{R} \text{ is singular at } \widehat{z}), \\ (\text{The ramification index of } \widehat{\varphi}|_{\widehat{R}_h} : \widehat{R}_h \to B \text{ at } \widehat{z}) - 1 & \text{(if } \widehat{R} \text{ is smooth at } \widehat{z}), \\ \widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} := \alpha_0^+(\Gamma_p)_{\widehat{z}} - 2\sharp(\widetilde{R}_0)_v(p)_{\widehat{z}}. \end{cases}$$

Let $\check{\varepsilon}(\Gamma_p)$ be the number of irreducible curves of $\widehat{R}_v(p)$ whose proper transforms on \widetilde{P} are (-n) curves. Then we have

(2.4)
$$\varepsilon(\Gamma_p) = \sum_{\widehat{z}\in\widehat{\Gamma}_p} j_1(\Gamma_p)_{\widehat{z}} + \check{\varepsilon}(\Gamma_p), \quad \alpha_0(\Gamma_p) = \sum_{\widehat{z}\in\widehat{\Gamma}_p} \widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} - 2(\sharp\widehat{R}_v(p) - \check{\varepsilon}(\Gamma_p)).$$

We let $\iota^t(\Gamma_p)_{\widehat{z}}$ and $\kappa^t(\Gamma_p)_{\widehat{z}}$ denote the numbers of singular points (including infinitely near one) of branch locus over \widehat{z} of types $n\mathbb{Z}$ and $n\mathbb{Z} + 1$, respectively, at which the proper transforms of two curves from $\{C_{t,k}\}_k$ meet, and put $\iota(\Gamma_p)_{\widehat{z}} = \sum_{t=1}^{\eta_{\widehat{z}}} \iota^t(\Gamma_p)_{\widehat{z}}$ and $\kappa(\Gamma_p)_{\widehat{z}} = \sum_{t=1}^{\eta_{\widehat{z}}} \kappa^t(\Gamma_p)_{\widehat{z}}$. By the definition, we note that $\iota(\Gamma_p)_{\widehat{z}} = \kappa(\Gamma_p)_{\widehat{z}} = 0$ if \widehat{R} is smooth at \widehat{z} .

We consider and recast Lemma 5.2 in [3]. Let $\widehat{\alpha}_k(\Gamma_p)_{\widehat{z}}$ be the number of singular points contributing $\widehat{\alpha}_k(\Gamma_p)$ over \widehat{z} .

Lemma 2.1. The following hold:

(1) $\iota(\Gamma_p)_{\widehat{z}} = j(\Gamma_p)_{\widehat{z}} - \eta_{\widehat{z}}.$ (2) $\alpha_0^+(\Gamma_p)_{\widehat{z}} \ge (n-2)(j(\Gamma_p)_{\widehat{z}} - \eta_{\widehat{z}} + \kappa(\Gamma_p)_{\widehat{z}}).$ (3) $\sum_{k\ge 1} \widehat{\alpha}_k(\Gamma_p)_{\widehat{z}} \ge \sum_{a\ge 1} (an-2)j_a(\Gamma_p)_{\widehat{z}} + 2\eta_{\widehat{z}} - \kappa(\Gamma_p)_{\widehat{z}}.$

Proof. For each t, we consider the graph \mathbb{G}_t corresponding to D_t as follows. The vertex set $V(\mathbb{G}_t)$ and the edge set $E(\mathbb{G}_t)$ are respectively the sets of symbols $\{v_{t,k}\}_{k=1}^{j^t(\Gamma_p)_{\widehat{z}}}$ and $\{e_x\}_x$, where x runs over all the singular points contributing to $\iota^t(\Gamma_p)_{\widehat{z}}$. If $C_{t,k}$ or a proper transform of it meets that of $C_{t,k'}$ at the singular point x of type $n\mathbb{Z}$, the edge e_x connects $v_{t,k}$ and $v_{t,k'}$. By the definition of D_t , we see that \mathbb{G}_t is a connected tree. Counting the numbers of vertices and edges, we have $\iota^t(\Gamma_p)_{\widehat{z}} = j^t(\Gamma_p)_{\widehat{z}} - 1$. Thus, we get (1) by summing it up for t.

When the component $L_{t,k}$ of D_t is a (-an)-curve, it is obtained by blowing $C_{t,k}$ up an-1 times. Thus, D_t is obtained from the bunch $\{C_{t,k}\}_k$ of (-1)-curves after performing $\sum_{a\geq 1}(an-1)j_a^t(\Gamma_p)_{\widehat{z}}$ times of blowing-ups, disregarding overlaps. Taking into the account the duplication and the first blowing-up creating the (-1)-curve for $L_{t,1}$, we see that the number of blowing-ups to obtain D_t is not less than

$$\sum_{a\geq 1} (an-1)j_a^t(\Gamma_p)_{\widehat{z}} - \iota^t(\Gamma_p)_{\widehat{z}} - \kappa^t(\Gamma_p)_{\widehat{z}} + 1 = \sum_{a\geq 2} (an-2)j_a^t(\Gamma_p)_{\widehat{z}} + 2 - \kappa^t(\Gamma_p)_{\widehat{z}},$$

since $\iota^t(\Gamma_p)_{\widehat{z}} = j^t(\Gamma_p)_{\widehat{z}} - 1$. Then, the number of singular points blown up to obtain $\widetilde{R}_v(p)_{\widehat{z}}$ is at least

$$\sum_{t=1}^{\eta_{\widehat{z}}} \left(\sum_{a \ge 2} (an-2) j_a^t (\Gamma_p)_{\widehat{z}} + 2 - \kappa^t (\Gamma_p)_{\widehat{z}} \right) = \sum_{a \ge 2} (an-2) j_a (\Gamma_p)_{\widehat{z}} + 2\eta_{\widehat{z}} - \kappa (\Gamma_p)_{\widehat{z}}.$$

This gives (3).

It remains to show (2). We may assume \widehat{R} is singular at \widehat{z} . Let $\widetilde{\Gamma}_{p,\widehat{z}} = \sum m_i G_i$ be the irreducible decomposition. Then it follows from (2.3) that

$$\alpha_0^+(\Gamma_p)_{\widehat{z}} = \sum_i m_i \widetilde{R}_h G_i - \sharp(\operatorname{Supp}(\widetilde{R}_h) \cap \operatorname{Supp}(\cup_i G_i)) \ge \sum_i (m_i - 1) \widetilde{R}_h G_i.$$

We consider a directed graph \mathbb{F} whose vertex set $V(\mathbb{F})$ is the set of symbols $\{v_x\}$, where x runs over all the singular points which contribute to either $\iota(\Gamma_p)_{\widehat{z}}$ or $\kappa(\Gamma_p)_{\widehat{z}}$. We define the directed edge from v_x to $v_{x'}$ if x' is a singular point infinitely near to x and any singular point between x and x' contributes to neither $\iota(\Gamma_p)_{\widehat{z}}$ nor $\kappa(\Gamma_p)_{\widehat{z}}$. Let $\mathbb{T}_1, \dots, \mathbb{T}_s$ be connected components of the graph \mathbb{F} . We note that \mathbb{T}_j $(j = 1, \dots, s)$ is a directed, rooted tree graph. We denote the leaf set of \mathbb{T}_j by $L(\mathbb{T}_j)$. By the definition of \mathbb{F} , any vertex in $L(\mathbb{T}_j)$ corresponds to a singular point contributing $\iota(\Gamma_p)_{\widehat{z}}$. Let E^x be the exceptional curve obtained by blowing up at $x \in L(\mathbb{T}_j)$. If there is a singular point of type $n\mathbb{Z}$ on E^x , we blow up at this point and forget E^x to call the new exceptional curve E^x . Repeating

such a procedure, we may assume that there are no singular points of type $n\mathbb{Z}$ on E^x . Since any singular points over x contribute to neither $\iota(\Gamma_p)_{\widehat{z}}$ nor $\kappa(\Gamma_p)_{\widehat{z}}$, we may assume that there are no singular points of type $n\mathbb{Z} + 1$ on E^x as well. Then we write it as \widetilde{E}^x to emphasize such properties, i.e., \widetilde{E}^x has no singular points of the branch locus on it. Since \widetilde{E}^x arises from a singular point of type $n\mathbb{Z}$, we see that $\widetilde{R}\widetilde{E}^x$ is a positive multiple of n, but it may be possible that \widetilde{E}^x meets two vertical components of \widetilde{R} . Thus we have $\widetilde{R}_h \widetilde{E}^x \ge n-2$. Letting m^x be the multiplicity along \widetilde{E}^x of $\widetilde{\Gamma}_p$, we have

$$\sum_{i} (m_i - 1)\widetilde{R}_h G_i \ge \sum_{j=1}^s \sum_{x \in L(\mathbb{T}_j)} (m^x - 1)\widetilde{R}_h \widetilde{E}^x.$$

We will show that

$$\sum_{x \in L(\mathbb{T}_j)} (m^x - 1) \ge \iota(\mathbb{T}_j) + \kappa(\mathbb{T}_j),$$

where $\iota(\mathbb{T}_j)$ (resp. $\kappa(\mathbb{T}_j)$) denotes the number of singular points of type $n\mathbb{Z}$ (resp. $n\mathbb{Z}+1$) in \mathbb{T}_j . Let $P^x(\mathbb{T}_j)$ be the set consisting of vertices appearing in the path connecting the root of \mathbb{T}_j and $x \in L(\mathbb{T}_j)$. We denote the number of singular points of type $n\mathbb{Z}$ (resp. $n\mathbb{Z}+1$) in $P^x(\mathbb{T}_j)$ by $\iota(P^x(\mathbb{T}_j))$ (resp. $\kappa(P^x(\mathbb{T}_j))$). We claim that $m^x \geq 2\iota(P^x(\mathbb{T}_j)) + \kappa(P^x(\mathbb{T}_j))$. This can be seen as follows. Put $P^x(\mathbb{T}_j) = \{v_1, \cdots, v_l = v_x\}$ and let m_1, \cdots, m_l be the multiplicities of the fiber $\widetilde{\Gamma}_p$ along the proper transforms of the exceptional curves arising from v_1, \cdots, v_l , respectively. If v_1 is of type $n\mathbb{Z}$, then we have $m_2 \geq m_1 + 2$. If v_1 is of type $n\mathbb{Z} + 1$, then we have $m_2 \geq m_1 + 1$. So we get $m^x = m_l \geq 2\iota(P^x(\mathbb{T}_j)) + \kappa(P^x(\mathbb{T}_j))$ inductively. Then,

$$\begin{split} \sum_{x \in L(\mathbb{T}_j)} (m^x - 1) &\geq \sum_{x \in L(\mathbb{T}_j)} \left(2\iota(P^x(\mathbb{T}_j)) + \kappa(P^x(\mathbb{T}_j)) \right) - \sharp L(\mathbb{T}_j) \\ &\geq 2\iota(\mathbb{T}_j) + \kappa(\mathbb{T}_j) - \sharp L(\mathbb{T}_j) \\ &\geq \iota(\mathbb{T}_j) + \kappa(\mathbb{T}_j) \end{split}$$

as wished.

Now, we get

$$\sum_{x \in L(\mathbb{T}_j)} (m^x - 1) \widetilde{R}_h G_i \ge \sum_{j=1}^s \sum_{x \in L(\mathbb{T}_j)} (m^x - 1) \widetilde{R}_h \widetilde{E}^x$$
$$\ge (n - 2) \sum_{j=1}^s \sum_{x \in L(\mathbb{T}_j)} (m^x - 1)$$
$$\ge (n - 2) (\iota(\Gamma_p)_{\widehat{z}} + \kappa(\Gamma_p)_{\widehat{z}}).$$

Plugging the inequalities thus obtained to (2.3), we conclude that $\alpha_0^+(\Gamma_p)_{\widehat{z}} \ge (n-2)(\iota(\Gamma_p)_{\widehat{z}} + \kappa(\Gamma_p)_{\widehat{z}})$. Then, since $\iota(\Gamma_p)_{\widehat{z}} = j(\Gamma_p)_{\widehat{z}} - \eta_{\widehat{z}}$ by (1), we get (2).

Remark 2.2. Lemma 5.2, (2) in [3] insists on

$$\alpha_0^+(\Gamma_p)_{\widehat{z}} \ge (n-2)(j(\Gamma_p)_{\widehat{z}} - \eta_{\widehat{z}} + 2\kappa(\Gamma_p)_{\widehat{z}}),$$

which is stronger than Lemma 2.1, (2) we have just shown. But it seems that the proof of the inequality in [3] has a gap. In that paper, it is used to show the upper bound of the slope of primitive cyclic covering fibrations of type (g, 0, n). Though we have failed to fill the gap itself, our inequality in Lemma 2.1, (2) is sufficient to save his bound.

Lemma 2.3. Assume that $n \ge 4$. Then it hold that

$$\widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} + \frac{2}{n} \sum_{k \ge 1} \widehat{\alpha}_k(\Gamma_p)_{\widehat{z}} - 2j_1(\Gamma_p)_{\widehat{z}} \ge 0.$$

Proof. From $\widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} = \alpha_0^+(\Gamma_p)_{\widehat{z}} - 2\sharp(\widetilde{R}_0)_v(p)_{\widehat{z}} = \alpha_0^+(\Gamma_p)_{\widehat{z}} - 2\sum_{a\geq 2} j_a(\Gamma_p)_{\widehat{z}}$ and Lemma 2.1, (2), we have

$$\widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} \ge (n-2)(j(\Gamma_p)_{\widehat{z}} - \eta_{\widehat{z}} + \kappa(\Gamma_p)_{\widehat{z}}) - 2\sum_{a \ge 2} j_a(\Gamma_p)_{\widehat{z}}$$
$$= (n-4)\sum_{a \ge 2} j_a(\Gamma_p)_{\widehat{z}} - (n-2)\eta_{\widehat{z}} + (n-2)\kappa(\Gamma_p)_{\widehat{z}} + (n-2)j_1(\Gamma_p)_{\widehat{z}}.$$

By Lemma 2.1, (3), we have

$$\frac{2}{n}\sum_{k\geq 1}\widehat{\alpha}_k(\Gamma_p)_{\widehat{z}} \geq \sum_{a\geq 1}\frac{2}{n}(an-2)j_a(\Gamma_p)_{\widehat{z}} + \frac{4}{n}\eta_{\widehat{z}} - \frac{2}{n}\kappa(\Gamma_p)_{\widehat{z}}$$

Therefore we get

$$\begin{aligned} \widehat{\alpha}_{0}(\Gamma_{p})_{\widehat{z}} &+ \frac{2}{n} \sum_{k \ge 1} \widehat{\alpha}_{k}(\Gamma_{p})_{\widehat{z}} \\ &\geq \sum_{a \ge 2} \left((n-4) + \frac{2}{n} (an-2) \right) j_{a}(\Gamma_{p})_{\widehat{z}} + \left(\frac{4}{n} - (n-2) \right) \eta_{\widehat{z}} \\ &+ \left(n-2 - \frac{2}{n} \right) \kappa(\Gamma_{p})_{\widehat{z}} + \left((n-2) + \frac{2}{n} (n-2) \right) j_{1}(\Gamma_{p})_{\widehat{z}} \\ &= \sum_{a \ge 2} \left(-2 + 2a \right) j_{a}(\Gamma_{p})_{\widehat{z}} + \left(n-2 - \frac{4}{n} \right) \left(j(\Gamma_{p})_{\widehat{z}} - \eta_{\widehat{z}} \right) \\ &+ \left(n-2 - \frac{2}{n} \right) \kappa(\Gamma_{p})_{\widehat{z}} + 2j_{1}(\Gamma_{p})_{\widehat{z}}. \end{aligned}$$

By the assumption that $n \ge 4$, the coefficients of $j_a(\Gamma_p)_{\widehat{z}}$, $j(\Gamma_p)_{\widehat{z}} - \eta_{\widehat{z}} = \iota(\Gamma_p)_{\widehat{z}}$ and $\kappa(\Gamma_p)_{\widehat{z}}$ are all non-negative. Therefore we get $\widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} + \frac{2}{n}\sum_{k\ge 1}\widehat{\alpha}_k(\Gamma_p)_{\widehat{z}} - 2j_1(\Gamma_p)_{\widehat{z}} \ge 0$.

Remark 2.4. When $n \ge 4$, we can show $\alpha_0(\Gamma_p) + \frac{2}{n} \sum_{k \ge 1} \alpha_k(\Gamma_p) - 2\varepsilon(\Gamma_p) \ge 0$ similarly.

We discuss the localization of the invariant K_f^2 . We recall the following lemma.

Lemma 2.5 ([3]). Let $f : S \to B$ be a primitive cyclic covering fibration of type (g, 0, n). Then it holds

$$K_{f}^{2} = \frac{n-1}{r-1} \left(\frac{(n-1)r - 2n}{n} (\alpha_{0} - 2\varepsilon) + (n+1) \sum_{k \ge 1} k(r-nk)\alpha_{k} \right) - n \sum_{k \ge 1} \alpha_{k} + \varepsilon.$$

So if we put

$$(2.5)$$

$$K_f^2(\Gamma_p) := \frac{n-1}{r-1} \left(\frac{(n-1)r-2n}{n} (\alpha_0(\Gamma_p) - 2\varepsilon(\Gamma_p)) + (n+1) \sum_{k \ge 1} k(r-nk) \alpha_k(\Gamma_p) \right)$$

$$-n \sum_{k \ge 1} \alpha_k(\Gamma_p) + \varepsilon(\Gamma_p)$$

for $p \in B$, then we get $K_f^2 = \sum_{p \in B} K_f^2(\Gamma_p)$. Since we have $K_f^2(\Gamma_p) = 0$ except for a finite number of points $p \in B$, we see that the invariant K_f^2 has been localized to a finite number of fibers. We will further localize a part of $K_f^2(\Gamma_p)$ to points on the fiber $\widehat{\Gamma}_p$.

For a fiber $\widehat{\Gamma}_p$ and a point $\widehat{z} \in \widehat{\Gamma}_p$, we put

(2.6)
$$K_{f}^{2}(\Gamma_{p})_{\widehat{z}} := \frac{n-1}{r-1} \frac{(n-1)r-2n}{n} (\widehat{\alpha}_{0}(\Gamma_{p})_{\widehat{z}} - 2j_{1}(\Gamma_{p})_{\widehat{z}}) + j_{1}(\Gamma_{p})_{\widehat{z}} + \frac{1}{r-1} \sum_{k \ge 1} ((n^{2}-1)k(r-nk) - (r-1)n) \widehat{\alpha}_{k}(\Gamma_{p})_{\widehat{z}}.$$

Then we get the following:

Proposition 2.6. Let $f: S \to B$ be a primitive cyclic covering fibration of type (g, 0, n). Let $\tilde{\psi} = \hat{\psi} \circ \check{\psi}$ be an arbitrary decomposition of $\tilde{\psi}: \tilde{P} \to P$ to have the commutative diagram



Denote a fiber of $\widehat{\varphi}$ over $p \in B$ by $\widehat{\Gamma}_p$. Then it holds

$$\begin{split} K_f^2(\Gamma_p) &= \sum_{\widehat{z} \in \widehat{\Gamma}_p} K_f^2(\Gamma_p)_{\widehat{z}} - 2 \frac{n-1}{r-1} \frac{(n-1)r - 2n}{n} \sharp \widehat{R}_v(p) + \check{\varepsilon}(\Gamma_p) \\ &+ \frac{1}{r-1} \sum_{k \ge 1} \left((n^2 - 1)k(r - nk) - (r - 1)n \right) \check{\alpha}_k(\Gamma_p). \end{split}$$

Proof. By (2.5), (2.4), we get

$$\begin{split} K_f^2(\Gamma_p) &= \frac{n-1}{r-1} \frac{(n-1)r-2n}{n} (\alpha_0(\Gamma_p) - 2\varepsilon(\Gamma_p)) + \varepsilon(\Gamma_p) \\ &+ \frac{1}{r-1} \sum_{k\geq 1} \left((n^2-1)k(r-nk) - (r-1)n \right) \alpha_k(\Gamma_p) \\ &= \frac{n-1}{r-1} \frac{(n-1)r-2n}{n} (\sum_{\widehat{z}\in\widehat{\Gamma}_p} \widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} - 2(\sharp \widehat{R}_v(p) - \check{\varepsilon}(\Gamma_p)) - 2\sum_{\widehat{z}\in\widehat{\Gamma}_p} j_1(\Gamma_p)_{\widehat{z}} - 2\check{\varepsilon}(\Gamma_p)) \\ &+ \sum_{\widehat{z}\in\widehat{\Gamma}_p} j_1(\Gamma_p)_{\widehat{z}} + \check{\varepsilon}(\Gamma_p) \\ &+ \frac{1}{r-1} \sum_{k\geq 1} \left((n^2-1)k(r-nk) - (r-1)n \right) \left(\sum_{\widehat{z}\in\widehat{\Gamma}_p} \widehat{\alpha}_k(\Gamma_p)_{\widehat{z}} + \check{\alpha}_k(\Gamma_p) \right) \\ &= \sum_{\widehat{z}\in\widehat{\Gamma}_p} K_f^2(\Gamma_p)_{\widehat{z}} - 2\frac{n-1}{r-1} \frac{(n-1)r-2n}{n} \sharp \widehat{R}_v(p) + \check{\varepsilon}(\Gamma_p) \\ &+ \frac{1}{r-1} \sum_{k\geq 1} \left((n^2-1)k(r-nk) - (r-1)n \right) \check{\alpha}_k(\Gamma_p) \end{split}$$
as wished.

as wished.

Lemma 2.7. Assume that $n \ge 4$. Then the following hold:

$$\begin{split} (r-1)K_{f}^{2}(\Gamma_{p})_{\widehat{z}} &\geq \sum_{k\geq 1} \left(-2\frac{n-1}{n^{2}} \left((n-1)r - 2n \right) + (n^{2}-1)k(r-nk) - n(r-1) \right) \widehat{\alpha}_{k}(\Gamma_{p})_{\widehat{z}} \\ &+ (r-1)j_{1}(\Gamma_{p})_{\widehat{z}}. \\ (r-1)K_{f}^{2}(\Gamma_{p}) &\geq \sum_{k\geq 1} \left(-2\frac{n-1}{n^{2}} \left((n-1)r - 2n \right) + (n^{2}-1)k(r-nk) - n(r-1) \right) \alpha_{k}(\Gamma_{p}) \\ &+ (r-1)\varepsilon(\Gamma_{p}). \end{split}$$

In particular, $K_f^2(\Gamma_p)_{\widehat{z}}$ and $K_f^2(\Gamma_p)$ are both non-negative.

Proof. From Lemma 2.1, we have

$$\widehat{\alpha}_0(\Gamma_p)_{\widehat{z}} - 2j_1(\Gamma_p)_{\widehat{z}} \ge -\frac{2}{n} \sum_{k \ge 1} \widehat{\alpha}_k(\Gamma_p)_{\widehat{z}}.$$

Then plugging it to (2.6), we get

$$\begin{split} (r-1)K_f^2(\Gamma_p)_{\widehat{z}} &\geq \sum_{k\geq 1} \left(-2\frac{n-1}{n^2} \big((n-1)r - 2n \big) + (n^2 - 1)k(r - nk) - n(r-1) \right) \widehat{\alpha}_k(\Gamma_p)_{\widehat{z}} \\ &+ (r-1)j_1(\Gamma_p)_{\widehat{z}}. \end{split}$$

By Remark 2.4, one can show the assertion for $K_f^2(\Gamma_p)$ similarly.

3. Automorphism of a fibered surface

It is classically known that finite subgroups of $Aut(\mathbb{P}^1)$ can be classified as in the following table:

| | | Order | Length of non-trivial orbits |
|----------------|-------------------|------------------|------------------------------|
| \mathbb{Z}_l | Cyclic group | l | 1, 1 |
| D_{2l} | Dihedral group | $2l \ (l \ge 2)$ | l, l, 2 |
| T_{12} | Tetrahedral group | 12 | 4, 4, 6 |
| O_{24} | Octahedral group | 24 | 6, 8, 12 |
| I_{60} | Icosahedral group | 60 | 12, 20, 30 |

In the table, the entry "4, 4, 6" for example means the action of the tetrahedral group has one orbit of length 6, two orbits of length 4 and the other orbits are of length 12. Note that the group which has a fixed point is necessarily a cyclic group.

For a fibration $f: S \to B$, we define the automorphism group of f as

$$\operatorname{Aut}(f) := \{ (\kappa_S, \kappa_B) \in \operatorname{Aut}(S) \times \operatorname{Aut}(B) \mid f \circ \kappa_S = \kappa_B \circ f \}.$$

Let $f: S \to B$ be a primitive cyclic covering fibration of type (g, 0, n). Let Σ be the covering transformation group of $S \to P$, the cyclic group of order n generated by σ . Let G be an arbitrary finite subgroup of $\operatorname{Aut}(f)$. Since $\operatorname{Aut}(S/B)$ is a finite group, we may assume $\sigma \in G$ to estimate the order of it. We have the exact sequence

$$1 \to K \to G \to H \to 1,$$

where $K := \{(\kappa_S, id) \in G\}$ and $H := \{\kappa_B \in \operatorname{Aut}(B) \mid (\kappa_S, \kappa_B) \in G, \exists \kappa_S \in \operatorname{Aut}(S)\}$. K can be considered as a subgroup of $\operatorname{Aut}(F)$ for a general fiber F of f.

Lemma 3.1. Assume that $r \ge 3n$. Take a general fiber F of f and any point $z \in F$. For any $\kappa \in Aut(F)$, it holds

$$\kappa(\Sigma \cdot z) = \Sigma \cdot \kappa(z),$$

where $\Sigma \cdot z$ denotes the Σ -orbit of z.

Proof. The subgroup $\Sigma \subset \operatorname{Aut}(F)$ induces the morphism $\theta : F \to \mathbb{P}^1$ of degree n. From the assumption that $r \geq 3n$, we know that F is an n-gonal curve with the unique gonality pencil θ ([3], Lemma 3.3). So there is a projective transformation $\rho : \mathbb{P}^1 \to \mathbb{P}^1$ such that the diagram

$$\begin{array}{c} F \xrightarrow{\kappa} F \\ \theta \\ \downarrow & \downarrow \theta \\ \mathbb{P}^1 \xrightarrow{\rho} \mathbb{P}^1 \end{array}$$

commutes. Note that $\theta(\kappa(\Sigma \cdot z)) = \rho(\theta(\Sigma \cdot z))$, we get

$$\kappa(\Sigma \cdot z) \subset \theta^{-1}(\theta(\kappa(\Sigma \cdot z))) = \theta^{-1}(\rho(\theta(\Sigma \cdot z))) = \theta^{-1}(\rho(\theta(z))) = \theta^{-1}(\theta(\kappa(z))) = \Sigma \cdot \kappa(z).$$

Arguing with $(\kappa^{-1}, \rho^{-1}, \kappa(z))$ instead of (κ, ρ, z) , we also have $\kappa^{-1}(\Sigma \cdot \kappa(z)) \subset \Sigma \cdot \kappa^{-1}(\kappa(z)) = \Sigma \cdot z$. Hence we obtain $\kappa(\Sigma \cdot z) = \Sigma \cdot \kappa(z)$.

In what follows, we tacitly assume that $r \geq 3n$.

Lemma 3.2. Let F be a general fiber of $f : S \to B$. Regard Σ as a subgroup of Aut(F). Then Σ is a normal subgroup of Aut(F).

Proof. For any $\kappa \in \operatorname{Aut}(F)$ and $z \in \operatorname{Fix}(\sigma)$, we have $\{\kappa(z)\} = \kappa(\Sigma \cdot z) = \Sigma \cdot \kappa(z)$ from Lemma 3.1. Hence the Σ -orbit of $\kappa(z)$ consists of the one point and we have $\kappa(z) \in \operatorname{Fix}(\sigma)$. Therefore, regarding κ as a permutation of $\operatorname{Fix}(\sigma)$, there is the group homomorphism

$$\lambda : \operatorname{Aut}(F) \to \mathfrak{S}_r$$

where $\mathfrak{S}_r = \operatorname{Aut}(\operatorname{Fix}(\sigma))$ is the *r*-th symmetric group. It suffices for our purpose to show that Ker $\lambda = \Sigma$.

Since it is clear that $\Sigma \subset \text{Ker } \lambda$, we shall show $\text{Ker } \lambda \subset \Sigma$. If $\kappa \in \text{Ker } \lambda$, there is a $\rho \in \text{Aut}(\mathbb{P}^1)$ such that the diagram

$$\begin{array}{c} F & \stackrel{\kappa}{\longrightarrow} F \\ \theta & \downarrow \rho \\ \mathbb{P}^1 & \stackrel{\rho}{\longrightarrow} \mathbb{P}^1. \end{array}$$

commutes from the uniqueness of the n-gonal pencil of F.

Since the action κ fixes every point in $\operatorname{Fix}(\sigma)$, we see that ρ fixes more than two points. So ρ is the identity, i.e., $\kappa \in \operatorname{Aut}(F/\mathbb{P}^1)$. Since $\theta : F \to \mathbb{P}^1$ is a Galois covering with Galois group Σ , we obtain $\kappa \in \Sigma$.

From Lemma 3.2, we get $\Sigma \triangleleft K$. Hence the action of $K \subset \operatorname{Aut}(S/B)$ on S can be lifted to the one on \widetilde{S} and we can regard K as a subgroup of $\operatorname{Aut}(\widetilde{S}/B)$. If we put $\widetilde{K} = K/\Sigma$, then we have the exact sequence

$$1 \to \Sigma \to K \to \widetilde{K} \to 1$$

Note that $\widetilde{K} \subset \operatorname{Aut}(\widetilde{P}/B)$ and \widetilde{R} is \widetilde{K} -stable (namely $\widetilde{K}(\widetilde{R}) = \widetilde{R}$).

Lemma 3.3. There is a relatively minimal model $\varphi : P \to B$ of $\tilde{\varphi}$ satisfying the following two conditions.

- $\operatorname{mult}_{x} R_{h} \leq r/2$ holds for any point x of the φ -horizontal part R_{h} of R.
- There exists a finite (possibly empty) subset Δ of B such that the action of K descends down faithfully on P \ φ⁻¹(Δ) but not over Δ. One can find one point on each fiber of φ in φ⁻¹(Δ) and, if P denotes the blowing-up at such \$\$ Δ points on P, then the action of K descends down faithfully on P which can switch two irreducible components on each singular fiber of the natural ruling \$\$ φ : P → B.

Proof. Let $\widetilde{\Gamma}_p$ be an arbitrary singular fiber of $\widetilde{P} \to B$, and E be any (-1)-curve contained in $\widetilde{\Gamma}_p$. Then its \widetilde{K} -orbit $\widetilde{K} \cdot E$ satisfies one of the following:

- $\widetilde{K} \cdot E$ consists of disjoint union of (-1)-curves in $\widetilde{\Gamma}_p$,
- We can find two curves in $\widetilde{K} \cdot E = \{E_1, E_2, \cdots, E_t\}$ meeting at a point.

In the former case, contracting $\widetilde{K} \cdot E$ to points, the action of \widetilde{K} descends down to the action on the new fiber obtained by the contraction.

In the latter case, we may assume that E_1 and E_2 meet. Since the intersection form on $E_1 \cup E_2$ is negative semi-definite, $E_1^2 = E_2^2 = -1$ and $E_1E_2 > 0$, the only possibility is: $E_1E_2 = 1$ and $(E_1 + E_2)^2 = 0$. Since any ruled surface has no multiple fibers, we get t = 2 and

$$\widetilde{\Gamma}_p = E_1 + E_2, \quad E_1 E_2 = 1$$

by Zariski's Lemma. If the action of \widetilde{K} never switch E_1 and E_2 , \widetilde{K} induces the action on the new fiber obtained by contracting either E_1 or E_2 . Otherwise, we stop.

Hence, repeating the above procedure, we finally get a fiber without (-1)-curves or a fiber of type $E_1 + E_2$ (and \tilde{K} switches E_1 and E_2).

Let $\varphi': P' \to B$ be the model at which the above algorithm terminates for all singular fibers of $\tilde{\varphi}$.



Denote the image of \widetilde{R} to P' by R'. Note that any singular fibers of φ' are of the form $E_1 + E_2$.

Let Γ' be a smooth fiber of $P' \to B$. We will show that $\operatorname{mult}_{x} R'_{h} \leq r/2$ can be assumed to hold for all singular points x of R'_{h} on Γ' . When \widetilde{K} is not cyclic, the length of \widetilde{K} orbits is not less than 2. So there are at least two singular points of R'_{h} on Γ' which are analytically equivalent. Therefore the multiplicity of the singular point is not bigger than r/2. Assume that \widetilde{K} is cyclic. If R'_{h} has a singular point on a \widetilde{K} -orbit of length $\sharp \widetilde{K}$, then there are at least two singular points of R'_{h} on Γ' and we can argue as in the previous case. So we assume that the singular point of R'_{h} of multiplicity greater than r/2 is fixed by \widetilde{K} . We perform the elementally transformation at the fixed point. Then \widetilde{K} acts naturally on the new fiber and one can check easily that the multiplicity of the new branch locus at the resulting fixed point is less than r/2 (cf. Lemma 3.1 of [3]).

Let $\overline{\varphi}: \overline{P} \to B$ be the model obtained from $\varphi': P' \to B$ as above, and let \overline{R} be the image of \widetilde{R} to \overline{P} . Note that the multiplicity of any singular point of \overline{R}_h on smooth fibers is at most r/2. Recall that any singular fiber of $\overline{\varphi}$ is of the form $E_1 + E_2$, and \widetilde{K} can switch E_1 and E_2 . Hence the (singular) points of \overline{R}_h on E_1 and on E_2 which are switched

by \widetilde{K} are analytically equivalent. In particular, $\overline{R}_h E_2$ has to coincide with $\overline{R}_h E_1$. Then, from $\overline{R}_h(E_1 + E_2) = r$, we get $\overline{R}_h E_1 = \overline{R}_h E_2 = r/2$. Therefore, we get the desired model $\varphi: P \to B$ after contracting one of E_1 and E_2 .

In what follows, we consider the model $\overline{\varphi}: \overline{P} \to B$ introduced in the above lemma. We denote by \overline{K} the subgroup of $\operatorname{Aut}(\overline{P}/B)$ induced by \widetilde{K} . We remark the following which has been shown in the proof of the above lemma.

Lemma 3.4. Assume that $\Delta \neq \emptyset$ and let Γ_p be a fiber over $p \in \Delta$. Then there is a singular point of R_h of multiplicity r/2 on Γ_p .

We close the section by giving two remarks.

Lemma 3.5. For any fiber $\overline{\Gamma}$ of $\overline{\varphi}$, let $\overline{K} \to \operatorname{Aut}(\overline{\Gamma})$ be the restriction of \overline{K} to $\overline{\Gamma}$. Then $\overline{K} \to \operatorname{Aut}(\overline{\Gamma})$ is an inclusion.

Proof. The assertion is clearly true for a general fiber of $\overline{\varphi}$. We will prove it by deriving a contradiction. Let $\overline{\Gamma}$ be a fiber of $\overline{\varphi}$. Assume that there is a non-trivial $\kappa \in \overline{K}$ such that $\kappa|_{\overline{\Gamma}} = id_{\overline{\Gamma}}$. The cyclic subgroup $\langle \kappa \rangle \subset \overline{K}$ fixes two points when κ is restricted to a general fiber of $\overline{\varphi}$. Hence $\operatorname{Fix}(\kappa)$ contains the $\overline{\varphi}$ -horizontal curve C_{κ} (which is not necessarily an irreducible curve) such that the degree of the morphism $\overline{\varphi}|_{C_{\kappa}} : C_{\kappa} \to B$ is 2. By the assumption, we have $\overline{\Gamma} \subset \operatorname{Fix}(\kappa)$. Then $C_{\kappa} \cap \overline{\Gamma}$ gives us singular points of 1-dimensional fixed locus of κ . This is impossible, since κ is not the identity. \Box

Lemma 3.6. Assume that $n \ge 3$ and let $f : S \to B$ be a primitive cyclic covering fibration of type (g, 0, n). If R_h is étale over B, then f is locally trivial and R is φ -horizontal.

Proof. Since f is a primitive cyclic covering fibration of type (g, 0, n), the multiplicity of any singular point of R is either in $n\mathbb{Z}$ or in $n\mathbb{Z} + 1$. From $n \geq 3$, we conclude that R has no double points. On the other hand, since R_h is étale over B, R_h is smooth and no fibers of φ have a contact to R_h . So we get $R = R_h$. Since R is étale over B, f is locally trivial.

4. Upper bounds of the order.

Let $f: S \to B$ a primitive cyclic covering fibration of type (g, 0, n) with $r \ge 3n$ if $n \ge 5, r \ge 13$ if n = 4 and assume that f is not locally trivial. We put

(4.1)
$$\mu_{r,n} := \frac{4n^3 r(r-1)}{(n(n+1)r - 2n^2 - 8(n-1))((n-1)r - 2n)}$$

Since r = 2(g + n - 1)/(n - 1), we can rewrite it as

$$\mu_{g,n} = \frac{2n^3(2g+n-1)(g+n-1)}{(n-1)\left(n(n+1)g+n(n-1)-4(n-1)^2\right)(g-1)}$$

as a function in g, n.

Let $K_f^2(\Gamma_p)$ be as in (2.5). We show that

$$\frac{n \sharp \overline{K}}{K_f^2(\Gamma_p)} \le \mu_{r,n}$$

or equivalently

(4.2)
$$4n^2r(r-1)K_f^2(\Gamma_p) \ge \left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1)\right) \sharp \overline{K}$$

holds for any $p \in B$ satisfying $K_f^2(\Gamma_p) \neq 0$. There are three possibilities: Assume that R is smooth. Since f is not locally trivial, Lemma 3.6 assures us that there is a fiber Γ_p which is tangent to R. Assume that R has a singular point on a fiber Γ_p over $p \in B$. Then either we can find such a p in $B \setminus \Delta$, or not, that is, all such p's are in Δ .

Since the problem is local, we may consider $f: S \to B$ locally around p.

Proposition 4.1. Let the notation and the assumptions be as above. If R is smooth in a neighborhood of Γ_p and is tangent to Γ_p , then it holds that

In particular, (4.2) holds true.

Proof. Since R is smooth around Γ_p , we may assume that $\overline{P} = P$. Note that we have $\alpha_k(\Gamma_p) = 0$ for $k \ge 1$ and $\varepsilon(\Gamma_p) = 0$. Furthermore, $\alpha_0(\Gamma_p)$ is nothing but the degree of the ramification divisor of $R \to B$ over p. Let $z \in R \cap \Gamma_p$ be a point of contact and $\operatorname{Stab}_{\overline{K}}(z) := \{\kappa \in \overline{K} \mid \kappa(z) = z\}$ the stabilizer of z in \overline{K} .

Case 1. $\operatorname{Stab}_{\overline{K}}(z)$ is not the unit group.

We know that $\operatorname{Stab}_{\overline{K}}(z)$ is a cyclic group, since it has z as a fixed point. Put $s = \sharp\operatorname{Stab}_{\overline{K}}(z)$. Note that the fixed curve (that is, the curve traced by the fixed points on fibers) of $\operatorname{Stab}_{\overline{K}}(z)$ is a section of $\varphi : P \to B$ locally around z and is not a part of R, since R is smooth and tangent to Γ_p at z. Then the $\operatorname{Stab}_{\overline{K}}(z)$ -orbit of any point on R sufficiently close to z consists of s distinct points. Since R and Γ_p are preserved by the action of $\operatorname{Stab}_{\overline{K}}(z)$, we get $(R, \Gamma_p)_z \ge s$, where $(R, \Gamma_p)_z$ denotes the local intersection number of R and Γ_p at z. Note that $(R, \Gamma_p)_z$ is the ramification index of $R \to B$ at z in the present case. It follows that z contributes at least s - 1 to $\alpha_0(\Gamma_p)$ and so does any point in the \overline{K} -orbit $\overline{K} \cdot z$ of z. Hence

$$\alpha_0(\Gamma_p) \ge \sharp(\overline{K} \cdot z)(s-1) = \frac{\sharp K}{s}(s-1)$$

and we infer readily from (2.5) that

$$4nr(r-1)K_f^2(\Gamma_p) \ge 4(n-1)r\big((n-1)r-2n\big)\frac{(s-1)}{s} \sharp \overline{K}.$$

Since $s \ge 2$, we get the desired inequality.

Case 2. Stab_{\overline{K}}(z) is the unit group.

By the assumption, we have $\#(\overline{K} \cdot z) = \#\overline{K}$, which implies that there are at least $\#\overline{K}$ points of contact in $R \cap \Gamma_p$. Since the ramification index of $\varphi|_R : R \to B$ at each point of contact is not less than 2, we have $\alpha_0(\Gamma_p) \ge (2-1)\#\overline{K} = \#\overline{K}$ and

$$4nr(r-1)K_f^2(\Gamma_p) \ge 4(n-1)r((n-1)r-2n) \sharp \overline{K}$$

from (2.5), which is stronger than what we want.

Proposition 4.2. Let $f: S \to B$ be a primitive cyclic covering fibration of type (g, 0, n)with $\begin{cases} r \geq 3n & (n \geq 5), \\ r \geq 13 & (n = 4). \end{cases}$ Assume that there exists a point $p \in B \setminus \Delta$ such that R has a singular point on Γ_p . Then it holds that

$$4n^2r(r-1)K_f^2(\Gamma_p) \ge 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1)\right) \sharp \overline{K}.$$

In particular, (4.2) holds true.

Proof. Let z be a singular point of R on Γ_p . Put $s = \sharp \operatorname{Stab}_{\overline{K}}(z)$.

Case 1. Stab_{\overline{K}}(z) is not the unit group.

Note that the fixed curve of $\operatorname{Stab}_{\overline{K}}(z)$ is a section of $\varphi : P \to B$ locally around z. Since we have $\operatorname{mult}_z R \geq 4$ by the assumption $n \geq 4$, there exist a local analytic branch of R_h which is not fixed by $\operatorname{Stab}_{\overline{K}}(z)$. Let D be such a local analytic branch of R and \mathbb{D} the $\operatorname{Stab}_{\overline{K}}(z)$ -orbit of D. Put $m_{\mathbb{D}} = \operatorname{mult}_z \mathbb{D}$ and $m := \min\{m' \mid m' \geq m_{\mathbb{D}}, m' \in$ $n\mathbb{Z}$ or $n\mathbb{Z} + 1\}$. Since $\operatorname{Stab}_{\overline{K}}(z)$ does not fix D, we get $(\mathbb{D}, \Gamma_p)_z \geq s$.

Case 1-1. The case that $m_{\mathbb{D}} = (\mathbb{D}, \Gamma_p)_z$.

We note that $m \geq s$ in this case. There exist $\sharp(\overline{K} \cdot z)$ singular points of R on Γ_p at the \overline{K} -oribit $\overline{K} \cdot z$. The multiplicity of R at each of such singular points is at least m. If $m \in n\mathbb{Z}$, we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m}{n}(r-m)-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)r.$$

from Lemma 2.7. We need to show

$$\left(-8(n-1)\left((n-1)r-2n\right) + 4n^2(n^2-1)\frac{m}{n}(r-m) - 4n^3(r-1) \right) \sharp(\overline{K} \cdot z)r - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1) \right) \sharp\overline{K} \ge 0.$$

in order to obtain the desired inequality. By $m \sharp(\overline{K} \cdot z) \ge s \sharp(\overline{K} \cdot z) = \sharp \overline{K}$, it is sufficient to show

(4.3)
$$\left(-8(n-1)\left((n-1)r-2n\right) + 4n^2(n^2-1)\frac{m}{n}(r-m) - 4n^3(r-1)\right)r - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1)\right)m \ge 0.$$

The left hand side of (4.3) can be considered as a quadratic function in m defined in the interval $n \leq m \leq r/2$. Since it is downward concave, it suffices to consider its value at the endpoints. When m = n, (4.3) is

$$(2n^4 - 4n^3 - 10n^2 + 16n - 8)r^2 + (-4n^5 + 8n^4 + 24n^3 - 16n^2)r - 8n^4 - 32n^3 + 32n^2 \ge 0$$

From $n \ge 4$ and $r \ge 3n$, we can check it by a simple calculation. When m = r/2, the left hand side of (4.3) becomes 0 as is easily seen. If $m \in n\mathbb{Z} + 1$, we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m-1}{n}(r-(m-1))-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)r.$$

from Lemma 2.7. We need to show

$$\left(-8(n-1)\left((n-1)r-2n\right)+4n^2(n^2-1)\frac{m-1}{n}(r-(m-1))-4n^3(r-1)\right)\sharp(\overline{K}\cdot z)r-2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^2-1)r^2-4n^3(r-1)\right)\sharp\overline{K}\geq 0.$$

in order to obtain the desired inequality. By $m \sharp(\overline{K} \cdot z) \ge s \sharp(\overline{K} \cdot z) = \sharp \overline{K}$, it is sufficient to show

$$(4.4) \quad \left(-8(n-1)\left((n-1)r-2n\right)+4n^2(n^2-1)\frac{m-1}{n}(r-(m-1))-4n^3(r-1)\right)r \\ -2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^2-1)r^2-4n^3(r-1)\right)m \ge 0.$$

From $m \in n\mathbb{Z} + 1$ and $r \in n\mathbb{Z}$, we have $n + 1 \le m \le (r - n)/2 + 1$.

The left hand side of (4.4) can be considered as a quadratic function in m defined in the interval $n \le m \le r/2$. Since it is downward concave, it suffices to consider its value at the endpoints. When m = n + 1, (4.4) is

$$(2n^4 - 6n^3 - 10n^2 + 18n - 8)r^2$$
$$+(-4n^5 + 8n^4 + 32n^3 - 32n + 16)r$$
$$-8n^4 - 40n^3 + 32n \ge 0.$$

From $n \ge 4$ and $r \ge 3n$, we can check it by a simple calculation. When m = (r-n)/2 + 1, we can check it from $r \ge 3n$ if $n \ge 5$ and $r \ge 13$ if n = 4. Hence we are done in this case.

Case 1-2. The case that $m_{\mathbb{D}} < (\mathbb{D}, \Gamma_p)_z$.

By the argument of Case 1-1, we may assume that if there exists a local analytic branch of R_h which is not tangent to Γ_p at z, then it is fixed by $\operatorname{Stab}_{\overline{K}}(z)$.

We let $\mathbb{D}_1 (:= \mathbb{D}), \mathbb{D}_2, \cdots, \mathbb{D}_d$ be the $\operatorname{Stab}_{\overline{K}}(z)$ -orbits of local analytic branches of R which are tangent to Γ_p at z, where d is the number of $\operatorname{Stab}_{\overline{K}}(z)$ -orbits of such local analytic branches of R.

- (i) The case that d = 1.
- We put $\mathbb{D} := \{ D = D_1, \cdots D_\delta \}.$
- (i-1) The case that D has a cusp at z.

We blow up at the point of contact as long as the proper transforms of D and Γ_p have a common tangent. Assume that we need v times of blowing-ups until the proper transform of D is not tangent to that of Γ_p . Let m_D be the multiplicity of D at z.

We assume that the proper transform of D is not tangent to the exceptional curve of the v-th blowing-up. Then we have

$$(D, \Gamma_p)_z = (v+1)m_D.$$

Since there are at least (v + 1) singular points of \mathbb{D} (infinitely near to z) of multiplicity $m_{\mathbb{D}}$, there exist at least $\sharp(\overline{K} \cdot z)(v + 1)$ singular points of R of multiplicity at least m. If $m \in n\mathbb{Z}$, we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m}{n}(r-m)-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)(v+1)r.$$

from Lemma 2.7 Then we only have to show

$$\left(-8(n-1)\left((n-1)r-2n\right) + 4n^2(n^2-1)\frac{m}{n}(r-m) - 4n^3(r-1) \right) \sharp(\overline{K} \cdot z)(v+1)r - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1)\right) \sharp\overline{K} \ge 0.$$

By $(v+1)m \ge (\mathbb{D}, \Gamma_p)_z \ge s$, we have $\sharp(\overline{K} \cdot z)(v+1)m \ge \sharp\overline{K}$. So it is sufficient to show

$$\left(-8(n-1)\left((n-1)r-2n\right) + 4n^2(n^2-1)\frac{m}{n}(r-m) - 4n^3(r-1) \right) r - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1) \right) m \ge 0,$$

which is nothing more than (4.3). If $m \in n\mathbb{Z} + 1$, we can show similarly as in the case $m \in n\mathbb{Z}$ by (4.4).

We assume that the proper transform of D is tangent to the exceptional curve of the v-th blowing-up. Then we have

$$(D,\Gamma_p)_z = vm_D + u_D \quad (0 \le u_D \le m_D - 1),$$

where u_D denotes the multiplicity of the proper transform of D at the point of contact to the exceptional curve of the *v*-th blowing-up. Denote by $u_{\mathbb{D}}$ the multiplicity of the proper transform of \mathbb{D} at the point of contact to the exceptional curve of the *v*-th blowing-up. We put

$$u = \min\{u' \mid u' \ge u_{\mathbb{D}}, \ u' \in n\mathbb{Z} \text{ or } n\mathbb{Z} + 1\}.$$

We consider the following two cases separately. The first is that the branch locus is singular at the point of contact to the exceptional curve of the v-th blowing-up. The second is that the branch locus is smooth at such a point. This latter case does not happen when $\Gamma_p \subset R$.

(i-1-1) In the first case, there are at least $\sharp(\overline{K} \cdot z)v$ singular points of R (infinitely near to z) of multiplicity at least m and $\sharp(\overline{K} \cdot z)$ singular points of multiplicity are at least u. If $m \in n\mathbb{Z}$ and $u \in n\mathbb{Z}$, we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m}{n}(r-m)-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)vr + \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{u}{n}(r-u)-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)r.$$

from Lemma 2.7. Hence we only have to show

$$\left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m}{n}(r-m)-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)vr + \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{u}{n}(r-u)-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)r - 2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^{2}-1)r^{2}-4n^{3}(r-1)\right)\sharp\overline{K}\geq 0.$$

By $(vm+u) \ge (\mathbb{D}, \Gamma_p)_z \ge s$, we get $\sharp(\overline{K} \cdot z)(vm+u) \ge \sharp \overline{K}$. So it suffices to show that

$$\left(-8(n-1)\left((n-1)r-2n\right)+4n^2(n^2-1)\frac{m}{n}(r-m)-4n^3(r-1)\right)vr + \left(-8(n-1)\left((n-1)r-2n\right)+4n^2(n^2-1)\frac{u}{n}(r-u)-4n^3(r-1)\right)r - 2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^2-1)r^2-4n^3(r-1)\right)(vm+u) \ge 0.$$

We can check it by (4.3). If $m \in n\mathbb{Z} + 1$ or $u \in n\mathbb{Z} + 1$, we can show similarly as in the above by (4.3) and (4.4).

(i-1-2) Next, we consider the second case. Note that we have $\Gamma_p \not\subset R$, $\delta = u = 1$ by the assumption of (i). Put $\overline{K} \cdot z := \{z(1)(:=z), \cdots, z(\sharp(\overline{K} \cdot z))\}$. Let $\mathbb{D}(j)$ be the \overline{K} -orbit of \mathbb{D} such that $z(j) \in \mathbb{D}(j)$ for $j = 1, \cdots, \sharp(\overline{K} \cdot z)$. Note that $\mathbb{D}(1) = \mathbb{D}$. Since $\mathbb{D}(j)$ is analytically equivalent to \mathbb{D} , after v-times blowing-ups over z(j), the proper transform of $\mathbb{D}(j)$ is tangent to the exceptional curve of the v-th blowing-up (let $\hat{z}(j)$ be such a tangent point) over z(j).

Let $\check{\psi}: \widehat{P} \to P$ be the composite of the above *v*-times blowing-ups for $j = 1, \cdots, \sharp(\overline{K} \cdot z)$, \widehat{R} the branch locus on \widehat{P} and let $\widehat{\Gamma}_p$ be a fiber of $\widehat{\varphi} := \varphi \circ \check{\psi}$ over *p*.

We will show $\sharp \widehat{R}_v(p) = 0$. Let $\widehat{R}_v(p)_z$ be the biggest subdivisor of $\widehat{R}_v(p)$ contracted to z by $\check{\psi}$. It is sufficient to show $\widehat{R}_v(p)_z = 0$, since $\mathbb{D}(j)$ is analytically equivalent to \mathbb{D} . We prove $\operatorname{mult}_z R \in n\mathbb{Z}$ by deriving a contradiction. Assume that $\operatorname{mult}_z R \in n\mathbb{Z} + 1$. Recall that we can assume that if there exist a local analytic branch of R which is not tangent to Γ_p at z, then it is fixed by $\operatorname{Stab}_{\overline{K}}(z)$. We blow-up R at z. If R has a local analytic

branch at z which is fixed by $\operatorname{Stab}_{\overline{K}}(z)$, then there exist a singular point of the branch locus whose multiplicity is 2. If R has no local analytic branch at z which is fixed by $\operatorname{Stab}_{\overline{K}}(z)$, then there exist a singular point of the branch locus whose multiplicity is in $n\mathbb{Z} + 2$. These are impossible by Lemma 1.3 (1). Hence we have $\operatorname{mult}_z R \in n\mathbb{Z}$. Therefore also we find that R has only one local analytic branch at z which is tangent to Γ_p at z if $v \geq 2$. So we get $\sharp \widehat{R}_v(p)_z = 0$ from Lemma 1.3 (1). We also have $m \in n\mathbb{Z}$.

There are at least $\sharp(\overline{K} \cdot z)v$ singular points of R (counting infinitely near ones) of multiplicity at least m. From Proposition 2.6, we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq 4n^{2}r(r-1)\sum_{j=1}^{\sharp(\overline{K}\cdot z)}K_{f}^{2}(\Gamma_{p})_{\widehat{z}(j)} + \left(4n^{2}(n^{2}-1)\frac{m}{n}(r-m) - 4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)vr.$$

Since \widehat{R} is smooth and ramified at $\widehat{z}(j)$, we get

$$4n^2 r(r-1) \sum_{j=1}^{\sharp(\overline{K}\cdot z)} K_f^2(\Gamma_p)_{\widehat{z}(j)} \ge 4n(n-1)\left((n-1)r-2n\right) \sharp(\overline{K}\cdot z)r$$

Then we only have to show

$$4n(n-1)\left((n-1)r-2n\right)\sharp(\overline{K}\cdot z)r + \left(4n^2(n^2-1)\frac{m}{n}(r-m)-4n^3(r-1)\right)\sharp(\overline{K}\cdot z)vr - 2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^2-1)r^2-4n^3(r-1)\right)\sharp\overline{K} \ge 0.$$

By $(vm+1) \ge (\mathbb{D}, \Gamma_p)_z \ge s$, we have $\sharp(\overline{K} \cdot z)(vm+1) \ge \sharp \overline{K}$. So it is sufficient to show

$$4n(n-1)\left((n-1)r-2n\right)r + \left(4n^2(n^2-1)\frac{m}{n}(r-m) - 4n^3(r-1)\right)vr \\ - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1)\right)(vm+1) \ge 0.$$

Since the coefficient of v is non-negative, we may assume v = 1 to show the above inequality. Hence it is sufficient to show

$$4n(n-1)\left((n-1)r-2n\right)r + \left(4n^2(n^2-1)\frac{m}{n}(r-m) - 4n^3(r-1)\right)r - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1)\right)(m+1) \ge 0.$$

We can check it by a simple calculation.

(i-2) The case that D is smooth at z.

We show that $\delta \geq 2$ by deriving a contradiction. Assume that $\delta = 1$. Recall that if there exist a local analytic branch of R which is not tangent to Γ_p at z, then it is fixed by $\operatorname{Stab}_{\overline{K}}(z)$. By d = 1 and $\delta = 1$, we get $\operatorname{mult}_z R \leq 3$, which is impossible when $n \geq 4$ by Lemma 1.3. Hence we have $\delta \geq 2$.

By $(D, \Gamma_p)_z \ge s/\delta$, the proper transforms of D and Γ_p meet until (s/δ) -times of blowingups over z. By $\delta \ge 2$, there exist at least $\sharp(\overline{K} \cdot z) \cdot s/\delta$ singular points of R of multiplicity at least $m_{\delta} := \min\{m' \mid m' \geq \delta, m' \in n\mathbb{Z} \text{ or } n\mathbb{Z}+1\}$. If $m_{\delta} \in n\mathbb{Z}$, we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m_{\delta}}{n}(r-m_{\delta})-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)\frac{s}{\delta}r.$$

from Lemma 2.7. Then we only have to show

$$\left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m_{\delta}}{n}(r-m_{\delta})-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)\frac{s}{\delta}r-2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^{2}-1)r^{2}-4n^{3}(r-1)\right)\sharp\overline{K}\geq0.$$

By $\#\overline{K} = \#(\overline{K} \cdot z)s$ and $m_{\delta} \geq \delta$, it is sufficient to show

$$\left(-8(n-1)\left((n-1)r-2n\right) + 4n^2(n^2-1)\frac{m_\delta}{n}(r-m_\delta) - 4n^3(r-1) \right) r - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1) \right) m_\delta \ge 0,$$

which is nothing more than (4.3). If $m_{\delta} \in n\mathbb{Z} + 1$, we can show similarly as in the case $m_{\delta} \in n\mathbb{Z}$ by (4.4).

(ii) The case that $d \geq 2$.

We put $\mathbb{D}_i := \{D_1^i, \cdots D_{\delta_i}^i\}$, where δ_i is the number of local analytic branches in \mathbb{D}_i . Changing the numbering of $\mathbb{D}_1, \dots, \mathbb{D}_d$ if necessary, we may assume that the proper transforms of D_1^1 and Γ_p become disjoint after the smallest number of blowing-ups over z among $D_1^1, \cdots D_1^d$.

(ii-1)The local analytic branch D_1^1 has a cusp at z.

Considering \mathbb{D}_1 as in (i-1), we can get the desired inequality by a similar calculation. We note that the case (i-1-2) can not happen by the assumption \mathbb{D}_1

(ii-2) The local analytic branch D_1^1 is smooth at z.

There are at least $\sharp(\overline{K} \cdot z)(D_1^1, \Gamma_p)_z$ singular points of R of multiplicity at least $m_1 :=$ $\min\{m' \mid m' \ge \delta_1, m' \in n\mathbb{Z} \text{ or } n\mathbb{Z}+1\}$. If $m_1 \in n\mathbb{Z}$, we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m_{1}}{n}(r-m_{1})-4n^{3}(r-1)\right)\sharp(\overline{K}\cdot z)(D_{1}^{1},\Gamma_{p})_{z}r.$$
from Lemma 2.7. Then we only have to show

 $2(2 1) m_1$

$$\left(-8(n-1)\left((n-1)r-2n\right) + 4n^2(n^2-1)\frac{m_1}{n}(r-m_1) - 4n^3(r-1) \right) \sharp(\overline{K} \cdot z)(D_1^1, \Gamma_p)_z r - 2\left(-8(n-1)\left((n-1)r-2n\right) + n(n^2-1)r^2 - 4n^3(r-1) \right) \sharp\overline{K} \ge 0.$$

By $\delta_1(D_1^1,\Gamma_p)_z = (\mathbb{D}_1,\Gamma_p)_z \ge s$, we get $m_1(D_1^1,\Gamma_p)_z \sharp(\overline{K}\cdot z) \ge \sharp \overline{K}$. So it is sufficient to show

$$\left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)\frac{m_{1}}{n}(r-m_{1})-4n^{3}(r-1)\right)r - 2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^{2}-1)r^{2}-4n^{3}(r-1)\right)m_{1} \ge 0,$$

which is nothing more than (4.3). If $m_1 \in n\mathbb{Z} + 1$, we can show similarly as in the case $m_1 \in n\mathbb{Z}$ by (4.4).

Case 2. Stab $\overline{K}(z)$ is the unit group.

By the assumption, we have $\#(\overline{K} \cdot z) = \#\overline{K}$, which implies that there are at least $\#\overline{K}$ singular points of R which are analytically equivalent to z. So we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)(r-n)-4n^{3}(r-1)\right) \sharp \overline{K}r$$

from Lemma 2.7. Then we only have to show

$$\left(-8(n-1)\left((n-1)r-2n\right)+4n^{2}(n^{2}-1)(r-n)-4n^{3}(r-1)\right)r - 2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^{2}-1)r^{2}-4n^{3}(r-1)\right) \ge 0,$$

which is clear from (4.3).

Proposition 4.3. Let
$$f: S \to B$$
 be a primitive cyclic covering fibration of type $(g, 0, n)$
with $\Delta \neq \emptyset$ and $\begin{cases} r \geq 3n & (n \geq 5), \\ r \geq 13 & (n = 4). \end{cases}$ Then it holds that
 $4n^2r(r-1)K_f^2(\Gamma_p) \geq (-8(n-1)((n-1)r-2n) + n(n^2-1)r^2 - 4n^3(r-1)) \sharp \overline{K}$

for any fiber $\Gamma_p \ (p \in \Delta)$.

Proof. Let Γ_p be the fiber over $p \in \Delta$. On the model \overline{P} , $\overline{\Gamma}_p$ consists of two (-1)-curves $\Gamma_i(i = 1, 2)$ with $\Gamma_1\Gamma_2 = 1$ and there is an element $\kappa_{sw} \in \overline{K}$ such that $\kappa_{sw}(\Gamma_1) = \Gamma_2$, $\kappa_{sw}(\Gamma_2) = \Gamma_1$ from Lemma 3.3. We put $\Gamma_1 \cap \Gamma_2 = \{z_0\}$.

We consider the subgroup

$$N := \{ \kappa \in \overline{K} \mid \kappa(\Gamma_i) = \Gamma_i \quad (i = 1, 2) \}.$$

For any $\kappa' \in \overline{K}$, we have $\kappa_{sw}\kappa' \in N$ if $\kappa' \notin N$. Hence, $\overline{K} = N \cup \kappa_{sw}N$. It follows that N is a normal subgroup of \overline{K} with the quotient group $\overline{K}/N \cong \mathbb{Z}_2$. Since N acts on $\Gamma_i \cong \mathbb{P}^1$ and fixes $z_0 \in \Gamma_i$ (i = 1, 2), we see that N is a cyclic group. Hence \overline{K} is cyclic or dihedral.

We claim that \overline{K} is either dihedral or $\overline{K} \cong \mathbb{Z}_2$. We assume that \overline{K} is cyclic and N is not the unit group, and show that this eventually leads us to a contradiction. Let $\overline{C}_{\overline{K}}$ be the fixed curve of \overline{K} on \overline{P} , i.e., the curve traced out by fixed points of \overline{K} on fibers. Since κ_{sw} fixes $\overline{C}_{\overline{K}}$ and the fixed point of κ_{sw} on $\overline{\Gamma}$ is only z_0 , we see that $\overline{C}_{\overline{K}}$ passes through z_0 . Since $\overline{C}_{\overline{K}}\overline{\Gamma} = 2$ and $\kappa_{sw}(\Gamma_1) = \Gamma_2$, we have $\overline{C}_{\overline{K}}\Gamma_i = 1$ for i = 1, 2. In particular, $\overline{C}_{\overline{K}}$ is smooth at z_0 . Since N is not the unit group, the fixed curve \overline{C}_N of N must coincides with $\overline{C}_{\overline{K}}$. We blow \overline{P} down to P. Then the action of N descends down to P locally around Γ_p and the fixed curve on N on P is tangent to Γ_p . This means, however, that N acts trivially on P, contradicting that it is not the unit group. Hence \overline{K} is dihedral. Note further that we have shown that \overline{C}_N does not pass through z_0 , when N is non-trivial. As Lemma 3.4 assures, we have the singular point z of R_h on Γ_p of multiplicity r/2. Let \overline{R} be the branch locus on \overline{P} .

Case 1. The group \overline{K} is a dihedral group.

Since N acts $\varphi: P \to B$ around Γ_p , We get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq 2\left(-8(n-1)\left((n-1)r-2n\right)+n(n^{2}-1)r^{2}-4n^{3}(r-1)\right)\sharp\overline{N}$$

similarly as in Proposition 4.2. So we get

$$4n^{2}r(r-1)K_{f}^{2}(\Gamma_{p}) \geq \left(-8(n-1)\left((n-1)r-2n\right)+n(n^{2}-1)r^{2}-4n^{3}(r-1)\right)\sharp\overline{K}.$$

Case 2. The group \overline{K} is the cyclic group \mathbb{Z}_2 .

Since z is a singular point of R of multiplicity at least r/2, we have

$$4n^2r(r-1)K_f^2(\Gamma_p) \ge \left(-8(n-1)((n-1)r-2n) + n(n^2-1)r^2 - 4n^3(r-1)\right)r.$$

So we only have to show

$$(-8(n-1)((n-1)r-2n) + n(n^2-1)r^2 - 4n^3(r-1))r - (-8(n-1)((n-1)r-2n) + n(n^2-1)r^2 - 4n^3(r-1)) 2 \ge 0,$$

which is trivial.

Now, we consider the upper bound of the order of the automorphism group of f. Let Γ_p be a fiber which is tangent to R_h or passes through a singular point of R or is over $p \in \Delta$. Put $r_p = \sharp \operatorname{Stab}_H(p)$. From $r \geq 3n$ and Lemma 2.7, we have

$$K_f^2 \ge \frac{\sharp H}{r_p} K_f^2(\Gamma_p).$$

Since $\sharp G = \sharp K \cdot \sharp H = n \sharp \overline{K} \cdot \sharp H$, we get

$$\sharp G \leq \frac{n \sharp \overline{K}}{K_f^2(\Gamma_p)} r_p K_f^2$$

From Propositions 4.1, 4.2 and 4.3, we have

$$\frac{n \sharp \overline{K}}{K_f^2(\Gamma_p)} \le \mu_{r,n}$$

and, hence,

$$\sharp G \le \mu_{r,n} r_p K_f^2.$$

So we only have to manage r_p .

Theorem 4.4. Let $f : S \to B$ be a primitive cyclic covering fibration of type (g, 0, n)with $\begin{cases} r \ge 3n & (n \ge 5), \\ r \ge 13 & (n = 4). \end{cases}$ Put $\mu_{r,n} := \frac{4n^3r(r-1)}{(n(n+1)r-2n^2-8(n-1))((n-1)r-2n)}.$ Assume furthermore that f is not locally trivial and, when g(B) = 0, f has at least 3 singular fibers (e.g., f is not iso-trivial). Let G be a finite subgroup of Aut(f). Then it holds

$$\sharp G \leq \begin{cases} 6(2g(B) - 1)\mu_{r,n}K_f^2 & (g(B) \ge 1) \\ 5\mu_{r,n}K_f^2 & (g(B) = 0). \end{cases}$$

Proof. Put $r_p = \sharp \operatorname{Stab}_H(p)$ for $p \in B$. Let $\pi : B \to B/H$ be the quotient map of B by H. Note that r_p be the ramification index of π at p.

(i) The case of $g(B) \ge 2$.

We denote the genus of B/H by g(B/H). From the Hurwitz formula, we get

$$2g(B) - 2 = \# H\left(2g(B/H) - 2 + \sum_{i=1}^{s} \frac{r_i - 1}{r_i}\right),$$

where s is the number of ramification points and r_i is the ramification index. Put

$$T := 2g(B/H) - 2 + \sum_{i=1}^{s} \frac{r_i - 1}{r_i}$$

which must be positive in the present case. If $g(B/H) \ge 2$, then we get $\#H \le g(B) - 1$ by $T \ge 2$, and it follows that $r_i \le \#H \le g(B) - 1$ for any $i = 1, \dots s$.

Assume that g(B/H) = 1. Then we get s > 0 by T > 0. By $r_i \ge 2$ for any $i = 1, \dots, s$, we get $1 - 1/r_i \ge 1/2$. Therefore we obtain $r_i \le \sharp H \le 4(g(B) - 1)$ for any $i = 1, \dots, s$ by $T \ge 1/2$.

Assume that g(B/H) = 0. When $s \ge 5$, we get $r_i \le \#H \le 4(g(B)-1)$ for any $i = 1, \dots s$ by $T \ge 1/2$. When s = 4, one of r_i is not less than 3. So we get $r_i \le \#H \le 12(g(B)-1)$ for any $i = 1, \dots s$ by $T \ge 1/6$. When s = 3, we may assume $r_1 \ge r_2 \ge r_3$. By the definition of T, we get

$$r_1 \le \sharp H = \frac{2g(B) - 2}{1 - \frac{1}{r_1} - \frac{1}{r_2} - \frac{1}{r_3}},$$

so we obtain

$$r_1 - 1 - \frac{r_1}{r_2} - \frac{r_1}{r_3} \le 2g(B) - 2.$$

If $r_3 = 2$, then $r_2 \ge 3$, so we get

$$r_1 - 1 - \frac{r_1}{3} - \frac{r_1}{2} \le 2g(B) - 2.$$

Hence we get $r_1 \leq 6(2g(B) - 1)$. If $r_2 \geq r_3 \geq 3$, we get $r_1 \leq 3(2g(B) - 1)$ Therefore we obtain $r_p \leq 6(2g(B) - 1)$ for any $p \in B$.

(ii) The case of g(B) = 1.

We need not consider a translation, since it has no fixed points. Then it is well known that the order of the automorphism group of B which fixes a point is at most 6. So the order of stabilizer of H is at most 6, and we get $r_p \leq 6$. (iii) The case of g(B) = 0

If H is neither a cyclic group nor a dihedral group, the order of a stabilizer of H is at most 5. So we may assume H is either a cyclic group or a dihedral group. It is well known that a rational pencil with at most two singular fibers is iso-trivial. If the number of singular fibers of f is at least three, then there exist a point $p \in B$ such that $K_f^2(\Gamma_p) > 0$ and $r_p \leq 2$. Hence we may assume $r_p \leq 2$.

Corollary 4.5. Let
$$f: S \to B$$
 be a non-locally trivial primitive cyclic covering fibration
of type $(g, 0, n)$ with $\begin{cases} r \ge 3n & (n \ge 5), \\ r \ge 13 & (n = 4). \end{cases}$ Then it holds
 $\#\operatorname{Aut}(S/B) \le \frac{n^3(2g+n-1)(g+n-1)}{(n-1)(n(n+1)g+n(n-1)-4(n-1)^2)(g-1)}K_f^2.$

Proof. We can apply argument of Section 3 and 4 for $G = \operatorname{Aut}(S/B)$. Note that K is $\operatorname{Aut}(S/B)$ and H is the unit group. Since f is not locally trivial, there exist at least two singular fibers F_p and $F_{p'}$ $(p, p' \in B)$ of f. So there exist at least two fibers Γ_p and $\Gamma_{p'}$ with $K_f^2(\Gamma_p), K_f^2(\Gamma_{p'}) > 0$. We may assume $K_f^2(\Gamma_p) \leq K_f^2(\Gamma_{p'})$. Therefore we get

$$\sharp \operatorname{Aut}(S/B) \le \frac{n \sharp \overline{K}}{2K_f^2(\Gamma_p)} K_f^2$$

where \overline{K} is defined by K as in Section 3. From Propositions 4.1, 4.2 and 4.3, we get the desired inequality.

5. An example

We construct a fiber germ of a primitive cyclic covering fibration of type (g, 0, n) and the automorphism group \widetilde{K} which satisfy

$$\frac{n \sharp \overline{K}}{K_f^2(\Gamma_p)} = \frac{4n^2 r(r-1)}{(n^2-1)r^2 - 4n^2(r-1)}$$

where \overline{K} is defined by \widetilde{K} as in Section 3.

We consider the projection $\Phi : \mathbb{P}^2_{\mathbb{C}} \times \mathbb{C} \to \mathbb{C}$ defined by $\Phi([w_0 : w_1 : w_2], z) = z$ where $[w_0 : w_1 : w_2]$ is a system of homogeneous coordinates on $\mathbb{P}^2_{\mathbb{C}}$ and z is a coordinate of \mathbb{C} . Let \overline{P} be a hypersurface of $\mathbb{P}^2_{\mathbb{C}} \times \mathbb{C}$ defined by the equation $w_0w_1 - w_2^2z = 0$. We consider the morphism $\overline{\varphi} : \overline{P} \hookrightarrow \mathbb{P}^2_{\mathbb{C}} \times \mathbb{C} \xrightarrow{\Phi} \mathbb{C}$, then $\overline{\varphi}$ is a conic bundle over \mathbb{C} . We have the commutative diagram:

$$\begin{array}{c|c} \overline{P} & & \mathbb{P}^2_{\mathbb{C}} \times \mathbb{C} \\ \hline \varphi & & & & \\ \hline \varphi & & & & \\ \mathbb{C} & \stackrel{\cong}{\longrightarrow} & \mathbb{C} \end{array}$$

We denote the fiber of $\overline{\varphi}$ over $z \in \mathbb{C}$ by $\overline{\Gamma}_z$. Note that $\overline{\varphi}$ has only one singular fiber $\overline{\Gamma}_0$ defined by the equation $w_0 w_1 = 0$ in $\mathbb{P}^2_{\mathbb{C}}$, which is the fiber of Φ over $0 \in \mathbb{C}$.

Let l be a positive integer divisible by n. We define automorphisms of $\overline{\varphi}: \overline{P} \to \mathbb{C}$ as follows:

$$\begin{aligned} \zeta : ([w_0 : w_1 : w_2], z) &\mapsto ([\zeta_l w_0 : \zeta_l^{-1} w_1 : w_2], z), \\ \kappa_{sw} : ([w_0 : w_1 : w_2], z) &\mapsto ([w_1 : w_0 : w_2], z), \end{aligned}$$

where ζ_l denotes a primitive *l*-th root of unity. Let \overline{K} be the automorphism group on \overline{P} generated by ζ and κ_{sw} , then \overline{K} is a dihedral group with order 2*l*. Since κ_{sw} switches irreducible components of $\overline{\Gamma}_0$, κ_{sw} does not preserve a relatively minimal model of $\overline{\varphi}$.

We define a primitive cyclic covering fibration over $\overline{\varphi}: \overline{P} \to \mathbb{C}$. Let \overline{D} be a smooth curve on \overline{P} defined by equations $w_2 - (w_0 + w_1) = 0$ and $w_0w_1 - zw_2^2 = 0$ in $\mathbb{P}^2_{\mathbb{C}} \times \mathbb{C}$. We find that $\overline{\varphi}|_{\overline{D}}: \overline{D} \to \mathbb{C}$ is degree 2 and $\sharp(\overline{D} \cap \overline{\Gamma}_0) = 2$. So \overline{D} passes $\overline{\Gamma}_0$ transversally. Let \overline{R} be the \overline{K} -orbits of \overline{D} . Then \overline{R} is smooth and reduced. Put $r := \overline{R}.\overline{\Gamma}_0$. From $\alpha(D) = D$, we have $r = \sharp \overline{K} = 2l$. Since l is divisible by n, there is a divisor $\overline{\mathfrak{d}}$ on \overline{P} such that $\overline{R} = n\overline{\mathfrak{d}}$. We consider the classical cyclic n-covering

$$\overline{\theta}: \overline{S} := \operatorname{Spec}_{\overline{P}} \left(\bigoplus_{j=0}^{n-1} \mathcal{O}_{\overline{P}}(-j\overline{\mathfrak{d}}) \right) \to \overline{P}$$

branched along \overline{R} . We will show that there exist an action $\kappa_{\overline{S}} \in \operatorname{Aut}(\overline{S}/\mathbb{C})$ for any $\kappa_{\overline{W}} \in \overline{K}$ which commutes the diagram



For any $\kappa_{\overline{W}} \in \overline{K}$, we have $\kappa_{\overline{W}}^* \overline{D} \sim \overline{D}$. So we have $\kappa_{\overline{W}}^* \overline{\mathfrak{d}} \sim \overline{\mathfrak{d}}$. Moreover, the defining equation of \overline{R} is invariant under the \overline{K} -action. Therefore there exist an action $\kappa_{\overline{S}}$ for any $\kappa_{\overline{W}}$. So we can construct a subgroup K of $\operatorname{Aut}(\overline{S}/\mathbb{C})$ generated by $\kappa_{\overline{S}}$'s and $\operatorname{Aut}(\overline{S}/\overline{W})$. It is nothing more than the desired group K. By the construction of $\overline{\varphi} : \overline{P} \to \mathbb{C}$, we get

$$(r-1)K_f^2(\Gamma_0) = \frac{n^2 - 1}{4n}r^2 - n(r-1).$$

Therefore we get

$$\frac{n \sharp \overline{K}}{K_f^2(\Gamma_0)} = \frac{4n^2 r(r-1)}{(n^2 - 1)r^2 - 4n^2(r-1)}.$$

References

- T. Arakawa, Bounds for the order of automorphism groups of hyperelliptic fibrations, Tôhoku. Math. J. 50 (1998), 317–323.
- [2] Z. Chen, Best bounds of automorphism groups of hyperelliptic fibrations, Tôhoku. Math. J. 50 (1998), 469–489.

[3] M. Enokizono, Slopes of fibered surfaces with a finite cyclic automorphism, Michigan Math. J. 66 (2017), 125–154.

Hiroto Akaike

Department of Mathematics, Graduate School of Science, Osaka University,

1-1 Machikaneyama, Toyonaka, Osaka 560-0043, Japan

e-mail: u802629d@ecs.osaka-u.ac.jp