

Refined universality for critical KCM: lower bounds

Ivailo Hartarsky^{*,1} and Laure Maréché^{†,2}

¹CEREMADE, CNRS, UMR 7534, Université Paris-Dauphine, PSL University
Place du Maréchal de Lattre de Tassigny, 75016 Paris, France

²Institut de Recherche Mathématique Avancée
UMR 7501 Université de Strasbourg et CNRS
7 rue René-Descartes, 67000 Strasbourg, France

November 16, 2020

Abstract

We study a general class of interacting particle systems called kinetically constrained models (KCM) in two dimensions tightly linked to the monotone cellular automata called bootstrap percolation. There are three classes of such models [4], the most studied being the critical one. In a recent series of works [17, 19, 25] it was shown that the KCM counterparts of critical bootstrap percolation models with the same properties [3] split into two classes with different behaviour.

Together with the companion paper by the first author [15], our work determines the logarithm of the infection time up to a constant factor for all critical KCM, which were previously known only up to logarithmic corrections. This improves all previous results except for the Duarte-KCM, for which we give a new proof of the best result known [24]. We establish that on this level of precision critical KCM have to be classified into seven categories instead of the two in bootstrap percolation [3]. In the present work we establish lower bounds for critical KCM in a unified way, also recovering the universality result of Toninelli and the authors [17] and the Duarte model result of Martinelli, Toninelli and the second author [24].

MSC2010: Primary 60K35; Secondary 82C22, 60J27, 60C05

Keywords: Kinetically constrained models, bootstrap percolation, universality, classification, Glauber dynamics, spectral gap.

^{*}hartarsky@ceremade.dauphine.fr

[†]laure.mareche@math.unistra.fr

1 Introduction and results

1.1 Kinetically constrained models

Kinetically constrained models (KCM) are a class of interacting particle systems used since the 80s to model the liquid-glass transition [12, 13] (see [14, 29] for reviews). We will be interested in the very general class of \mathcal{U} -KCM first introduced by Cancrini, Martinelli, Roberto and Toninelli in 2008 [5], which includes all previously considered cases on \mathbb{Z}^d .

Let us start by introducing these models directly on the two-dimensional lattice, to which we restrict our attention. A KCM is a Markov process with state space $\Omega = \{0, 1\}^{\mathbb{Z}^2}$. For a configuration $\eta \in \Omega$ and a site $x \in \mathbb{Z}^2$, we denote η_x the value of η at x . We say that x is *empty* or *infected* if $\eta_x = 0$ and that it is *occupied* or *healthy* if $\eta_x = 1$. We thus naturally identify a configuration $\eta \in \Omega$ with the set of its infected sites, so that $\eta \subset \mathbb{Z}^2$.

A \mathcal{U} -KCM is specified by two parameters—an *update family* \mathcal{U} and an *equilibrium measure* μ . The measure μ on Ω is chosen to be the product Bernoulli measure such that each site is infected with probability $q > 0$ and healthy with probability $1 - q$. All asymptotics hereafter are taken as $q \rightarrow 0$. The update family \mathcal{U} is a finite set of finite nonempty subsets of $\mathbb{Z}^2 \setminus \{0\}$ called *update rules*. The dynamics is the following. For each site $x \in \mathbb{Z}^2$ at rate 1 we attempt to update η_x by replacing it by an independent Bernoulli($1 - q$) variable. However, the update is only performed if there exists an update rule $U \in \mathcal{U}$ such that $\eta \supset (x + U)$, while otherwise the configuration remains unchanged. In more formal terms the generator of the Markov process is the following for any function $f : \Omega \mapsto \mathbb{R}$ depending on a finite number of sites.

$$\mathcal{L}(f)(\eta) = \sum_{x \in \mathbb{Z}^2} \mathbb{1}_{\{\exists U \in \mathcal{U}, \eta_{x+U}=0\}} (\mu_x(f) - f)(\eta),$$

where η_X denotes the restriction of η to $X \cap \mathbb{Z}^2$ for $X \subset \mathbb{R}^2$ and $\mu_x(f)$ denotes the average of f with respect to η_x conditionally on all other occupation variables. We further view η_X as the element of $\{0, 1\}^{\mathbb{Z}^2}$ equal to η in $X \cap \mathbb{Z}^2$ and to 1 elsewhere. For background on the theoretical foundations of such interacting particle systems the reader is referred to [22]. In particular, μ is indeed a reversible invariant measure for the \mathcal{U} -KCM.

For any KCM arguably the most natural quantity of interest describing the speed at which memory of the initial state is lost is the *first infection time* of the origin

$$\tau_0 = \inf\{t \geq 0, (\eta(t))_0 = 0\} \in [0, \infty],$$

where $(\eta(t))_{t \geq 0}$ denotes the \mathcal{U} -KCM process. As we will concentrate on the equilibrium properties of the KCM, we will rather be concerned with $\mathbb{E}_\mu[\tau_0]$,

that is the expectation of τ_0 with respect to the law of the stationary \mathcal{U} -KCM with initial condition distributed according to its equilibrium measure μ .

1.2 Bootstrap percolation and universality

Bootstrap percolation is a close relative of KCM, though the two fields remained relatively independent for decades. Formally, the continuous time version of \mathcal{U} -bootstrap percolation is the \mathcal{U} -KCM with $q = 1$. However, bootstrap percolation has important additional properties and has attracted a great deal of attention with different motivation, also in non-lattice settings, as well as from computer science and sociological perspectives. We direct the reader to [8, 28] and the references therein for an overview of this rich field.

In \mathcal{U} -bootstrap percolation (the update family \mathcal{U} being as above), for each integer $t \geq 0$, a set A_t of infected sites at time t is constructed as follows. Given an initial set of infected sites $A_0 \subset \mathbb{Z}^2$ we set for all integers $t \geq 0$

$$A_{t+1} = A_t \cup \{x \in \mathbb{Z}^2 : \exists U \in \mathcal{U}, x + U \subset A_t\}.$$

That is, at each discrete time step the sites such that a rule translated at them is already fully infected also become infected, while infections never heal. Thus, given the initial condition, \mathcal{U} -bootstrap percolation is a monotone deterministic cellular automaton. For any set $A_0 \subset \mathbb{Z}^2$ we denote $[A_0] = \bigcup_{t \geq 0} A_t$ its *closure* with respect to the \mathcal{U} -bootstrap percolation process, i.e. the sites that are eventually infected when the initial infection is A_0 (we may also use this notation when $A_0 \not\subset \mathbb{Z}^2$; it will then mean $[A_0 \cap \mathbb{Z}^2]$). The best-studied setting, which is also the one relevant to us, is taking A_0 random with law μ , so that each site is infected independently with probability q . In this case the most prominent questions are for which values of q we have $[A_0] = \mathbb{Z}^2$ a.s. and, for such values of q , what is the typical order of magnitude of the infection time of the origin τ_0 defined as for KCM.

The general \mathcal{U} -bootstrap percolation framework only gained visibility after the work of Bollobás, Smith and Uzzell [4]. They introduced several crucial notions, which we discuss next. We invite the reader unfamiliar with these notions to systematically consult the examples in Figure 1 and apply the definitions to them.

We denote by $S^1 = \{z \in \mathbb{R}^2, \|z\| = 1\}$ the unit circle, which we standardly identify with $\mathbb{R}/2\pi\mathbb{Z}$, where $\|\cdot\|$ is the Euclidean norm of \mathbb{R}^2 (all distances in this work will be Euclidean, denoted by $d(\cdot, \cdot)$). A direction $u \in S^1$ will be called *rational* when $\tan u \in \mathbb{Q} \cup \{\infty\}$. For a direction $u \in S^1$ and a scalar $x \in \mathbb{R}$, we define the open half-plane $\mathbb{H}_u(x) = \{y \in \mathbb{R}^2, \langle u, y \rangle < x\}$ directed

by u translated by x , and $\bar{\mathbb{H}}_u(x) = \{y \in \mathbb{R}^2, \langle u, y \rangle \leq x\}$ the corresponding closed half-plane. We further set $\mathbb{H}_u = \mathbb{H}_u(0)$. A direction $u \in S^1$ is said to be *unstable* (for \mathcal{U}) if there exists $U \in \mathcal{U}$ such that $U \subset \mathbb{H}_u$ and *stable* otherwise. It is not hard to see (Theorem 1.10 of [4], Lemma 2.6 of [3]) that the set of stable directions is a finite union of closed intervals of S^1 with rational endpoints. Bollobás, Smith and Uzzell [4] introduced the following partition of update families into three classes. In order to state their results and others, we will need the following standard asymptotic notation. For any real functions $f(q)$, $g(q)$ defined for $q > 0$ sufficiently small, with $g > 0$, we write

- $f(q) = \Theta(g(q))$ when $cg(q) \leq f(q) \leq Cg(q)$,
- $f(q) = \Omega(g(q))$ when $f(q) \geq cg(q)$,
- $f(q) = O(g(q))$ when $|f(q)| \leq Cg(q)$

for some constants $0 < c \leq C < +\infty$ when $q > 0$ is sufficiently small. Finally, we write $f(q) = o(g(q))$ when $\frac{|f(q)|}{g(q)} \rightarrow 0$ when $q \rightarrow 0$. Let us note that all such implicit constants are allowed to depend on the update family \mathcal{U} and all (finite sets of) directions considered, but never on q .

Definition 1.1 (Definition 1.3 of [4]). An update family \mathcal{U} is called

- *supercritical* if there exists an open semicircle of unstable directions,
- *critical* if it is not supercritical, but there exists an open semicircle with a finite number of stable directions,
- *subcritical* otherwise.

In [4] it was proved that for supercritical models $\tau_0 = q^{-\Theta(1)}$ with high probability¹ as $q \rightarrow 0$, while for critical ones $\tau_0 = \exp(q^{-\Theta(1)})$. Completing the justification of Definition 1.1, Balister, Bollobás, Przykucki and Smith [2] proved that for subcritical models $\tau_0 = \infty$ with positive probability for q small enough. Albeit very general, these results were much less precise than what was known for most specific models studied previously, so one may view them as only qualitatively identifying the three different possible behaviours.

Supercritical models are fairly simple from the bootstrap percolation point of view, while still very little is known in general about subcritical ones [16]. Consequently, the critical models are the most pursued. That is

¹This means there exist constants $0 < c < C < +\infty$ such that $\mu(q^{-c} \leq \tau_0 \leq q^{-C}) \rightarrow 1$ when $q \rightarrow 0$.

why Bollobás, Duminil-Copin, Morris and Smith [3] established much more precise quantitative results for this class. To state their results, we need some more notation.

Definition 1.2 (Definition 1.2 of [3]). Let \mathcal{U} be a critical update family and $u \in S^1$ be a direction. Then the *difficulty* of u , $\alpha(u)$, is defined as follows.

- If u is unstable, then $\alpha(u) = 0$.
- If u is an isolated stable direction (isolated in the topological sense), then
$$\alpha(u) = \min\{n \in [1, \infty) : \exists Z \subset \mathbb{Z}^2, |Z| = n, |[\mathbb{Z}^2 \cap (\mathbb{H}_u \cup Z)] \setminus \mathbb{H}_u| = \infty\},$$
i.e. the minimal number of infections allowing \mathbb{H}_u to grow infinitely.
- Otherwise, $\alpha(u) = \infty$.

We define the *difficulty* of \mathcal{U} by

$$\alpha(\mathcal{U}) = \min_{C \in \mathcal{C}} \max_{u \in C} \alpha(u) \in [1, \infty), \quad (1)$$

where $\mathcal{C} = \{\mathbb{H}_u \cap S^1 : u \in S^1\}$ is the set of open semicircles of S^1 .

Definition 1.3 (Definition 1.3 of [3]). A critical update family with difficulty α is *balanced* if there exists a closed semicircle in which all directions have difficulty at most α and is *unbalanced* otherwise.

With these definitions, the main result of Bollobás, Duminil-Copin, Morris and Smith [3, Theorem 1.5], states

$$\tau_0 = \begin{cases} \exp\left(\frac{\Theta(1)}{q^{\alpha(\mathcal{U})}}\right) & \text{for critical balanced models,} \\ \exp\left(\frac{\Theta((\log q)^2)}{q^{\alpha(\mathcal{U})}}\right) & \text{for critical unbalanced models} \end{cases} \quad (2)$$

with high probability as $q \rightarrow 0$. These are the best general estimates currently known, though for some specific choices of \mathcal{U} sharper results are available [9, 10, 20].

1.3 Universality for KCM

Supercritical KCM As we shall see, the situation for \mathcal{U} -KCM is far more complex. Some new features already appear for supercritical models. The general setting was treated only recently by Martinelli, Morris and Toninelli [25] and Martinelli, Toninelli and the second author [24], who identified the two relevant classes of models.

Definition 1.4 (Definition 2.11 of [25]). A supercritical update family \mathcal{U} is *rooted* if there exist two non-opposite stable directions and *unrooted* otherwise.

In [25] the upper bounds in the following result for \mathcal{U} -KCM were established: as $q \rightarrow 0$,

$$\mathbb{E}_\mu[\tau_0] = \begin{cases} \exp(\Theta(\log(1/q))) & \text{supercritical unrooted,} \\ \exp(\Theta((\log(1/q))^2)) & \text{supercritical rooted,} \end{cases} \quad (3)$$

while the lower ones were supplied in [24], the lower bound for unrooted models being trivial. The lower bound for rooted models relied mostly on a combinatorial result by the second author [23] roughly stating that in order to infect the origin starting from a configuration in which the infection closest to the origin is at distance d from it, the dynamics has to go through a configuration with at least $\log d$ infections at distance at most d from the origin. This combinatorial bottleneck was originally identified in [30] (see also [7]) for the archetypal one-dimensional supercritical rooted model, which is, perhaps, the simplest and best-studied KCM—the East model (see [11] for a review). Extending the one-dimensional result to higher dimensions in [23] required the development of a new approach, which will be the starting point for our analysis, although we will not be able to apply the result of [23] itself.

Critical KCM Turning to critical models, a first observation is that bootstrap percolation provides an automatic lower bound for $\mathbb{E}_\mu[\tau_0]$ in the \mathcal{U} -KCM with the same update family. In particular, it is easy to show (see [26, Lemma 4.3] and its improved version from [18, Section 2]) that the expressions in (2) are lower bounds for $\mathbb{E}_\mu[\tau_0]$ for the corresponding \mathcal{U} -KCM.

Led by the intuition from the results for supercritical models, Morris [27], presenting his work with Martinelli and Toninelli introduced the following notion (we simplify his terminology slightly).

Definition 1.5 (Definition 2.3 of [27]). A critical update family of difficulty α is called *rooted* if there exist two non-opposite directions of difficulty strictly larger than α and *unrooted* otherwise.

Initially it was believed, as explained in [27], that if a critical model is unrooted, it satisfies $\mathbb{E}_\mu[\tau_0] = \exp(q^{-\alpha(\mathcal{U})+o(1)})$ as $q \rightarrow 0$, but not if it is rooted. This conjecture was made more precise by the same authors in [25], who then suggested a different definition of rooted/unrooted reflecting the upper bounds they proved and conjectured to be tight.

However, both conjectures were disproved by Martinelli, Toninelli and the first author [19], who proved stronger upper bounds than the ones in [25], in particular refuting the above conjectures for some rooted models and showing that the automatic bootstrap percolation lower bound is essentially sharp for them as well. In a parallel work Toninelli and the present authors [17] proved for all other models lower bounds essentially matching the upper ones from [25]. Hence, the combined results of [17, 19, 25] proved the following universality picture featuring yet a different partition of critical models. As $q \rightarrow 0$,

$$\mathbb{E}_\mu[\tau_0] = \begin{cases} \exp\left(\frac{(\log(1/q))^{O(1)}}{q^{\alpha(\mathcal{U})}}\right) & \text{critical, finitely many stable directions,} \\ \exp\left(\frac{(\log(1/q))^{O(1)}}{q^{2\alpha(\mathcal{U})}}\right) & \text{critical, infinitely many stable directions.} \end{cases}$$

This result shows that each universality class in bootstrap percolation (models with the same value of $\alpha(\mathcal{U})$) splits into two universality classes of KCM. This should indeed be viewed as the critical counterpart of (3). While the lower bound of [17] establishing the result for models with infinitely many stable directions does reflect (and actually uses) the combinatorial bottleneck of the one-dimensional supercritical rooted East model, the upper bound of [19] is based on a very peculiar efficient mechanism that uses a ‘quasi-local’ East-type movement resulting in a supercritical unrooted dynamics on larger length scales. However, it remained unclear whether this fairly unnatural mechanism of [19] and the purely East-like one used in [25] are indeed the correct ones for all models concerned.

1.4 Results

In order to identify the relevant mechanisms for different models, together with the companion work by the first author [15], we determine $\log \mathbb{E}_\mu[\tau_0]$ up to a constant factor. In the present paper we establish all lower bounds by identifying the correct bottleneck (see Section 2), while [15] provides the remaining matching upper bounds by exhibiting efficient relaxation mechanisms for all critical \mathcal{U} -KCM.

In order to state our results we introduce the last bit of notation needed to define the seven refined universality classes of critical \mathcal{U} -KCM, which we identify.

Definition 1.6. A critical update family of difficulty α is called

- *isotropic* if there is no direction of difficulty strictly greater than α ,

	infinite stable directions	finite stable directions	
		rooted	unrooted
unbalanced	(a) 2, 4, 0	(c) 1, 3, 0	(d) 1, 2, 0
balanced	(b) 2, 0, 0	(e) 1, 1, 0	(f) 1, 0, 1
			s.-dir. iso. (g) 1, 0, 0

Table 1: Classification of critical \mathcal{U} -KCM with difficulty α . Assuming [15], for each class $\mathbb{E}_\mu[\tau_0] = \exp\left(\Theta(1) \left(\frac{1}{q^\alpha}\right)^\beta \left(\log \frac{1}{q}\right)^\gamma \left(\log \log \frac{1}{q}\right)^\delta\right)$ as $q \rightarrow 0$. The label of the class and the exponents β, γ, δ are indicated in that order.

- *semi-directed* if there exists exactly one direction of difficulty strictly greater than α .

Notice that the isotropic/semi-directed critical models form a partition of unrooted balanced critical ones.

Theorem 1. *Let \mathcal{U} be a critical update family with difficulty α and infinite number of stable directions. We have the following alternatives as $q \rightarrow 0$.*

- (a) *If \mathcal{U} is unbalanced, i.e. there exist two opposite directions $u \in S^1$ with $\alpha(u) > \alpha$, then*

$$\mathbb{E}_\mu[\tau_0] = \exp\left(\frac{\Theta((\log(1/q))^4)}{q^{2\alpha}}\right).$$

- (b) *If \mathcal{U} is balanced, i.e. there do not exist two opposite directions $u \in S^1$ with $\alpha(u) > \alpha$, then*

$$\mathbb{E}_\mu[\tau_0] = \exp\left(\frac{\Omega(1)}{q^{2\alpha}}\right).$$

The upper bound in (a) was proved in [25, Theorem 2(a)]. Furthermore, for one specific model of the class (a), the Duarte model, this result was known from [24]. A different proof for case (b) was given in [17]. Finally, a matching upper bound for case (b) will be proved in the companion paper [15].

Theorem 2. *Let \mathcal{U} be a critical update family with difficulty α and finite number of stable directions. We have the following alternatives as $q \rightarrow 0$.*

- (c) If \mathcal{U} is unbalanced rooted, i.e. there exist at least three directions $u \in S^1$ with $\alpha(u) > \alpha$, two of which are opposite, then

$$\mathbb{E}_\mu[\tau_0] = \exp \left(\frac{\Theta((\log(1/q))^3)}{q^\alpha} \right).$$

- (d) If \mathcal{U} is unbalanced unrooted, i.e. there exist exactly two directions $u \in S^1$ such that $\alpha(u) > \alpha$ and they are opposite, then

$$\mathbb{E}_\mu[\tau_0] = \exp \left(\frac{\Omega((\log(1/q))^2)}{q^\alpha} \right).$$

- (e) If \mathcal{U} is balanced rooted, i.e. there exist at least two directions $u \in S^1$ with $\alpha(u) > \alpha$, but not two opposite ones, then

$$\mathbb{E}_\mu[\tau_0] = \exp \left(\frac{\Omega(\log(1/q))}{q^\alpha} \right).$$

- (f) If \mathcal{U} is semi-directed, i.e. there exists exactly one direction $u \in S^1$ such that $\alpha(u) > \alpha$, then

$$\mathbb{E}_\mu[\tau_0] = \exp \left(\frac{\Omega(\log \log(1/q))}{q^\alpha} \right).$$

- (g) If \mathcal{U} is isotropic, i.e. there exists no direction $u \in S^1$ such that $\alpha(u) > \alpha$, then

$$\mathbb{E}_\mu[\tau_0] = \exp \left(\frac{\Omega(1)}{q^\alpha} \right).$$

The upper bound in (c) was proved in [19, Theorem 1]. The lower bounds for cases (d) and (g) are only stated for completeness and follow automatically from bootstrap percolation results and [26, Lemma 4.3] as discussed in the previous section. Finally, matching upper bounds for cases (d), (e), (f) and (g) are proved in the companion paper [15].

The classification results, assuming the remaining matching bounds of [15], are summarised in Table 1. In addition, a simple representative of each class is given in Figure 1 for the reader's convenience.

Remark 1.7. The lower bounds we prove in Theorems 1 and 2 for $\mathbb{E}_\mu[\tau_0]$ also hold for another important characteristic timescale of the corresponding \mathcal{U} -KCM, the *relaxation time* T_{rel} (see e.g. [17, Definition 2.5]), which is another measure of the speed at which the memory of the initial state is lost. Indeed, [25, Equation (2.8)] yields $T_{\text{rel}} \geq q\mathbb{E}_\mu[\tau_0]$.




Figure 1: Representative models of each of the seven refined universality classes of critical \mathcal{U} -KCM. For each one the update rules are depicted on the left with 0 marked by a cross and the sites of the rule denoted by dots. The figure on the right gives the stable directions, which are thickened and have their difficulties next to them. The isolated stable directions are marked by dots. In all cases the difficulty α of the model is 1, as witnessed by the right-hand open semicircle.

1.5 Comments

Relationship with previous results Let us emphasise that, contrary to what is the case in bootstrap percolation, where the exact asymptotics and corrective terms of $\log \tau_0$ are sometimes known, for KCM there exists only one model (the Duarte model treated by Martinelli, Toninelli and the second author [24]) for which $\log \mathbb{E}_\mu[\tau_0]$ is determined up to a constant factor and none for which the exact asymptotics is known. Thereby, the combined results of [15, 19, 25] and the present work improve the best known results for all critical \mathcal{U} -KCM except the Duarte-KCM and give a new proof of the best known results for that last model. In view of the state of the art in the simpler setting of \mathcal{U} -bootstrap percolation, it does not seem currently feasible to pursue higher precision in full generality for critical \mathcal{U} -KCM. Theorem 1 establishes the lower bounds of [17, Conjecture 7.1], while Theorem 2 proves the ones of [19, Conjecture 6.1] (that conjecture was intentionally not as precise concerning the semi-directed case (f), which seemed quite mysterious at the time of writing of the conjecture).

Unification and extensions What is more, the present work treats all lower bounds in a systematic and comprehensive way, by showing that the main obstacle for the dynamics in all cases corresponds to a combinatorial bottleneck similar to the one of the two-dimensional East model (see Section 2 for more details). Moreover, the core of our argument carries over to higher dimensions with no further difficulty, thus reducing such lower bounds for higher-dimensional KCM to estimates for their bootstrap percolation counterparts.

Symmetry The arguments in this paper allow dealing with the lack of symmetry that many classes of critical models suffer from. Indeed, the models for which the present treatment is relevant (all classes except (d) and (g)) do not exhibit a lot of symmetry. One can see that an update family invariant by rotation by $\pi/2$ is necessarily isotropic (g) and that an update family invariant by rotation by π (i.e. centrally symmetric) is in class (d) or (g). We still note that all seven classes have representatives with one axial symmetry. The classification established in [15] and the present work allows understanding the precise impact of the symmetry exhibited by the update families on the timescales of the corresponding KCM.

1.6 Organisation of the paper

The remainder of the paper is organised as follows. In Section 2 we provide an outline of the ideas of the proof. The core of the proof is Section 3, where we establish the general combinatorial bottleneck for KCM dynamics. In Section 4 we deduce our main results, Theorems 1 and 2, very directly from Proposition 3.7. It is not before Section 4 that the distinction between the different classes of models becomes relevant. In addition to the central Proposition 3.7, Section 4 requires several estimates regarding bootstrap percolation, which are established in the appendices. Appendix A is rather standard and provides bounds on the probability of ‘spanning’ by uniting arguments of Bollobás, Duminil-Copin, Morris and Smith [3] and of Toninelli and the authors [17] without significant new input. This appendix may still be of interest to bootstrap percolation specialists, as it gives a notable simplification of the most technical part of the proof of the main result of [3]. Finally, in Appendix B we establish bounds on the probability of a notion of ‘crossing’ inspired from [3] but modified to suit our setting (see Section 2), which are not particularly difficult given Appendix A.

2 Outline of the proof

Let us outline the main highlights of the proof, emphasising new ideas with respect to our main sources of inspiration [3, 15, 17, 23]. The differences between the various universality classes will be much more apparent in [15], where efficient mechanisms for the infection of the origin are implemented for each class. Since one of the main virtues of our work is that the core argument is independent of the universality class, we invite the reader willing to understand the origin of the choices of length scales appearing in Section 4 to consult [15], which is also our inspiration for choosing them. For the sake of concreteness, unless otherwise indicated, in this section we restrict our attention to the representative of the class (e) of rooted balanced models with finitely many stable directions given in Figure 1e.

The proof relies on the notion of ‘bottleneck’: we show that before the origin can be infected, the dynamics has to go through a set of configurations with a probability small enough so that this does not happen for a very long time.

Morally speaking, in this model the smallest mobile entity (‘droplet’) is an infected square of size roughly $1/q$. Indeed, typically on its right and top sides one can find an infection, which allows it to infect the column of sites on its right and the row of sites above it. However, it is essentially impossible for

the infection to grow down or left, as this requires two consecutive infections and those are typically only available at distance $1/q^2$ from the droplet. We will only work in a region R of size $1/q^{7/4}$ around the droplet, so, morally, such couples of infections are not available. Thus, we can think of the droplet as performing the following simpler dynamics. If there is a droplet present at a certain position, it may create/destroy another one above it and to its right. Therefore, the droplets follow the dynamics of the two-dimensional supercritical rooted KCM called East model.

As mentioned in the introduction, for supercritical rooted models, including the (two-dimensional) East one, the second author established in [23] the following bottleneck: in order for an infection (representing a droplet in our original model) to reach the center of a box R of size 3^n initially fully healthy it is necessary to visit a configuration with at least n infections simultaneously present in the box. This was achieved by an inductive argument that we describe next. It is enough to show that if R is initially fully healthy and we can only have strictly less than n infections at the same time in R , we cannot reach a configuration with all infections in the middle of R . Indeed, this implies that there are strictly less than $n - 1$ infections at the same time in the middle of R , and one can make an induction on n . To show that we cannot reach a configuration with all infections in the middle of R , by reversibility we may instead prove that if we start with infections only in the middle of R , but we are never allowed to have n infections simultaneously in R , we cannot reach a configuration fully healthy in R . The idea is to ensure that for any path of the dynamics starting with all infections in the middle of R in which we never have n infections at the same time in R , the following two conditions remain true at all times. Firstly, a buffer zone (see the shaded frame B in Figure 2) with no infections remains intact. Secondly, there is always an infection in the internal region encircled by the buffer, so the dynamics cannot reach a configuration completely healthy in R .

In order to achieve that, we use a second induction, on the step of the path. We know that so far an infection remains trapped in the internal region encircled by the buffer, so we only have $n - 1$ infections available for disrupting the buffer from the outside, which is impossible by induction on n . Therefore, it suffices to show that we may not disrupt the buffer from the inside either. By projecting the two-dimensional East model on each axis it is clear that no infection can enter the left and bottom parts of the buffer from the inside, and the projections of the lowest and leftmost particles in the region inside the buffer need to remain where they were initially. The right part of the buffer (and similarly the top one) cannot be reached from the inside, because at least one infection needs to remain as far left as the leftmost initial one was, so we only have $n - 1$ infections with which to reach

the right part of the buffer, which is impossible by induction on n .

On a very high level we will proceed in the same way for critical models. However, there are several obvious problems in making the above reasoning rigorous for these models (in fact we do not think that a direct mapping to the two-dimensional East model can be made rigorous for our purposes). Firstly, we said above that the smallest mobile entities, ‘droplets’, were infected squares of size $1/q$, but the smallest mobile entities are actually more complicated. One needs to identify an event, which says whether or not something is a droplet and this event should be both deterministically necessary for infection to spread and sufficiently unlikely, so that having many droplets at the same time has probability small enough to be a good bottleneck. It turns out that the notion of ‘spanning’ introduced in [3], following [6] is flexible enough for us. Roughly speaking (see Definition 3.1), a droplet is spanned if the infections present inside it are sufficient to infect a connected set touching all its sides. We call a droplet critical if it has size roughly $1/q$. It is known from [3] and obtained again in Appendix A in a more adapted form that the probability of a specific critical droplet being spanned is roughly $\exp(-1/q)$. Unfortunately, given a configuration, spanned critical droplets may overlap, so in order to obtain good bounds on the probability of the configuration, one needs to consider disjointly occurring ones. We may then define the number of spanned critical droplets as the maximal number of disjointly occurring ones.

Having fixed these notions, we encounter a more significant issue—the (spanned critical) droplets may move a bit without creating another droplet, by changing their internal structure. Worse yet, they are not really forbidden to move left or down, but simply are not likely to be able to do so wherever they want: it depends on the dynamic environment. Indeed, being able to move by a single step down is allowed by the presence of a couple of infections on the side of the droplet, which has probability only as small as q and is by far not something we can ensure never happens up to time $T = \exp(\log(1/q)/q)$.

In order to handle these problems we introduce the crucial notion of crossing (not to be confused e.g. with the one of [3]). Consider a vertical strip S of width $1/q^{3/2}$ of our domain, R , which is a square of size $1/q^{7/4}$. Roughly speaking (see Definition 3.5 for a more precise statement), we say that S has a crossing if the following two events occur. Firstly, the infections in S together with the entire half-plane to the right of S are enough to infect a path from right to left in S (this is essentially the notion of crossing in [3]). Secondly, S does not contain a spanned critical droplet. Notice that these two events go opposite ways—the former is favoured by infections, while the latter is not. In Appendix B we show that the probability of a crossing decays

exponentially with the width of S at our scales.

Having established such a bound on the probability of crossings, we may safely assume (it happens with high probability) that they never occur until time T and this is the property we use to formalise the intuition that ‘moving down or left is impossible’. More precisely, assume that initially the only critical droplet is on the right of S and S never has a crossing. Then, simply because the KCM dynamics can never infect more than what bootstrap percolation can, starting from the same initial condition, the droplet will not be able to reach the left side of S . Indeed, if it could, there would be a ‘trail’ of infectable sites from the right of S to its left, which would imply a crossing.

3 Supercritical rooted dynamics of droplets

3.1 Setting and preliminaries

Let \mathcal{U} be an update family. Assuming they exist, we further fix two non-collinear rational stable directions u_1 and u_2 . We set $u_3 = u_1 + \pi$, $u_4 = u_2 + \pi$ and $\mathcal{T} = \{u_1, u_2, u_3, u_4\}$. We will simply call *parallelogram* a set of the form

$$\begin{aligned} R(a, b; c, d) &= \{x \in \mathbb{R}^2 \mid \langle x, u_3 \rangle \in [a, c], \langle x, u_4 \rangle \in [b, d]\} \\ &= \bar{\mathbb{H}}_{u_1}(-a) \cap \bar{\mathbb{H}}_{u_2}(-b) \cap \bar{\mathbb{H}}_{u_3}(c) \cap \bar{\mathbb{H}}_{u_4}(d) \end{aligned}$$

for real numbers $a \leq b, c \leq d$ and denote by $\mathring{R}(a, b; c, d)$ its topological interior. For parallelograms we will systematically extend definitions by translation and interchange of u_1 and u_2 (resp. u_3 and u_4).

Finally,

$$C_6 \gg C_5 \gg C_2' \gg C_1 \gg r = \max\{\|s - s'\| \mid s, s' \in U \cup \{0\}, U \in \mathcal{U}\}$$

are constants not depending on q , but only on \mathcal{U} and \mathcal{T} , each one sufficiently large with respect to functions of the next.² Furthermore, we systematically assume that q is small enough, as we are interested in $q \rightarrow 0$.

Definition 3.1 ([3, Definitions 2.3 and 2.4]). A set $Z \subset \mathbb{Z}^2$ is *strongly connected* if it is connected in the graph with vertex set \mathbb{Z}^2 defined by $x \sim y$ if $\|x - y\| \leq C_2'$.

Given $K \geq C_1^2 C_2'$ to be specified in Section 4, we say a parallelogram is *critical* when its diameter is contained between K/C_1 and K .

A parallelogram D is *spanned* in η if there exists a strongly connected set $X \subset [D \cap \eta]$ such that the smallest parallelogram containing X is D .

²We use C_2' instead of C_2 and avoid C_3 and C_4 for coherence with the appendices.

If η, η' are two configurations, we say that $\eta \leq \eta'$ when $\eta_s \leq \eta'_s$ for all s . For instance, if a parallelogram D is spanned for η , then it is also spanned for any $\eta' \leq \eta$. This order should not be confused with the (inverted) one induced by inclusion when viewing η as its set of infections.

Notice that the event that a given parallelogram D is spanned depends only on η_D and does not occur when η_D contains no infections. We further state two immediate consequences of Definition 3.1 for future reference.

Observation 3.2. Let $R \subset \mathbb{R}^2$. Then every parallelogram D spanned in η_R intersects R .

Observation 3.3. Let η be a configuration and X be a strongly connected component of $[\eta]$. Then $X = [\eta \cap X]$.

Proof. By maximality of a strongly connected component, $[\eta \cap X] \subset X$ is at distance at least $C'_2 > 2r$ from other strongly connected components X of $[\eta]$. Thus,

$$[\eta] = \bigsqcup_Y [\eta \cap Y],$$

where the union is on all strongly connected components of $[\eta]$. \square

Another standard fact is the following Aizenman-Lebowitz lemma originating from [1], whose proof can be found in the appendix (Lemma A.9).

Lemma 3.4. *Let D be a spanned parallelogram with diameter $d \geq C_1 C'_2$ and let $C_1 d \geq k \geq C_1 C'_2$. Then there exists a spanned parallelogram with diameter d' such that $k/C_1 \leq d' \leq k$. In particular, if $d \geq K/C_1$, then there exists a spanned critical parallelogram.*

We next import and adapt the notion of crossing from [3, Definition 8.17].

Definition 3.5 (Crossing). We say that a parallelogram $R = R(a, b; 0, d)$ is u_1 -crossed if there exists a strongly connected set in $[\mathbb{H}_{u_1} \cup (R \cap \eta)]$ intersecting both \mathbb{H}_{u_1} and $\bar{\mathbb{H}}_{u_3}(a)$.

Let $C_R^{u_1}$ denote the event that there exists $\eta' \geq \eta$ such that R is u_1 -crossed for η' , but there is no spanned critical parallelogram for η'_R .

We say that a parallelogram $\Lambda = R(0, 0; L, H)$ has no (ℓ, h) -crossing (or simply crossing) if the event $C_R^{u_1}$ does not occur for any $R \subset \Lambda$ of the form $R(a, 0; a + \ell, H)$ and the event $C_R^{u_2}$ does not occur for any $R \subset \Lambda$ of the form $R(0, b; L, b + h)$.

Finally, we say that a site $s \in \mathbb{Z}^2$ is *locally infectable* in a configuration η if $s \in [\eta \cap (s + R(-2K, -2K; 2K, 2K))]$. We also denote η^s the configuration that is equal to η everywhere except at s , i.e. $\eta_s^s = 1 - \eta_s$ and $\eta_{s'}^s = \eta_{s'}$ for any $s' \neq s$. We then have the following property, originating from [18, Section 2].

Lemma 3.6. *Let $\eta \in \{0, 1\}^{\mathbb{Z}^2}$, $s \in \mathbb{Z}^2$, $U \in \mathcal{U}$ be such that $s + U \subset \eta$ and let $R = R(-2K, -2K; 2K, 2K)$. Assume that the origin is not locally infectable in η , but is locally infectable in η^s . Then there exists a critical parallelogram D spanned in η_R^s such that $D \subset R(-3K, -3K; 3K, 3K)$.*

Proof. By definition, $0 \in [\eta^s \cap R] \setminus [\eta \cap R]$. Therefore, $s \in R$ and $s + U \not\subset R$. In particular, $d(s, 0) > K$. Let X be the strongly connected component of 0 in $[\eta^s \cap R]$. By Observation 3.3 we have that $s \in X$, since otherwise we would have $0 \in [X] \subset [\eta \cap R]$. Therefore, the smallest parallelogram containing X is spanned and has diameter at least K . In particular, by Lemma 3.4 there exists a critical parallelogram D spanned in η_R^s and we are done by Observation 3.2, since $\text{diam}(D) \leq K$ by Definition 3.1. \square

3.2 The combinatorial bottleneck

With the notation above we are ready to prove a very general deterministic bottleneck (Lemma 3.10 below), constituting the core of our work, which relatively straightforwardly translates into the following bound on $\mathbb{E}_\mu[\tau_0]$.

Proposition 3.7. *Let T, L, H, K, ℓ, h be positive real numbers sufficiently large with respect to C'_2 . Denote*

$$\rho = \max_D \mu(D \text{ is spanned}),$$

where the max is over all critical parallelograms. Also set

$$\begin{aligned} p_{\leftarrow} &= \max_{a, b \in \mathbb{R}} \mu \left(C_{R(a, b; a+\ell, b+H)}^{u_1} \right) \\ p_{\downarrow} &= \max_{a, b \in \mathbb{R}} \mu \left(C_{R(a, b; a+L, b+h)}^{u_2} \right). \end{aligned}$$

Assume that for some integer $n \geq 0$ we have the following inequalities on geometry:

$$L \geq 3^n(11K + \ell) \qquad H \geq 3^n(11K + h),$$

and probability:

$$\begin{aligned} 1/8 &\geq \mu(0 \text{ is locally infectable}) \\ 1 &\geq T(LH)^2 \max(p_{\downarrow}, p_{\leftarrow}) \\ 1 &\geq TLH(LHK^3\rho)^{n+1}. \end{aligned}$$

Then the \mathcal{U} -KCM on \mathbb{Z}^2 satisfies $\mathbb{E}_\mu[\tau_0] \geq T$.

Remark 3.8. Although the bootstrap percolation estimates needed to make use of this statement in higher dimensions are not yet available, let us mention that our argument is not dimension sensitive.

The proof of Proposition 3.7 will occupy the rest of the present section. We start by fixing ℓ, h, K as in the statement and introducing the following definitions.

Recall that for any $R \subset \mathbb{Z}^2$, we identify configurations $\eta \in \{0, 1\}^R$ with their zero set $\{x \in R, \eta_x = 0\}$. Unless otherwise specified, configurations $\eta \in \{0, 1\}^R$ are extended to $\{0, 1\}^{\mathbb{Z}^2}$ by keeping the same zero set.

Definition 3.9 (Good paths and configurations). For any parallelogram $R \subset \mathbb{R}^2$, configuration $\eta \in \{0, 1\}^{R \cap \mathbb{Z}^2}$ and integer $n \geq 0$, we say that η is *n-good* when the maximum number of critical parallelograms that are disjointly³ spanned in η is at most n and R has no crossing for η .

A *legal path* in R is a sequence $(\eta^{(j)})_{0 \leq j \leq m}$ of configurations in $\{0, 1\}^{R \cap \mathbb{Z}^2}$ such that for every $j \in \{0, \dots, m-1\}$, there exists $s \in R \cap \mathbb{Z}^2$ such that $\eta^{(j+1)} = (\eta^{(j)})^s$ and $(s + U) \cap R \subset \eta^{(j)}$ for some $U \in \mathcal{U}$. For any integer $n \geq 0$, the path is *n-good* if for every $j \in \{0, \dots, m\}$, $\eta^{(j)}$ is *n-good*. For any $A, B \subset \{0, 1\}^{R \cap \mathbb{Z}^2}$, we say $(\eta^{(j)})_{0 \leq j \leq m}$ is a path from A to B when $\eta^{(0)} \in A$ and $\eta^{(m)} \in B$ (if A or $B = \{\eta\}$, we will write η to simplify).

We denote by $G(R)$ the set of configurations in $\{0, 1\}^{R \cap \mathbb{Z}^2}$ that contain no spanned critical parallelogram and such that R contains no crossing, i.e. the 0-good configurations. For any $n \in \mathbb{N}$, we define

$$V(n, R) = \{\eta \in \{0, 1\}^{R \cap \mathbb{Z}^2} \mid \text{there is an } n\text{-good legal path from } G(R) \text{ to } \eta\}.$$

Finally, we define our domain sizes for the induction to come:

$$L_n = \frac{3^n - 1}{2} (9K + \ell) + 3^n K \quad H_n = \frac{3^n - 1}{2} (9K + h) + 3^n K,$$

so that $L_n - L_{n-1} = 2L_{n-1} + 9K + \ell$ and $H_n - H_{n-1} = 2H_{n-1} + 9K + h$.

Lemma 3.10. *For any non-negative integer n , for any parallelogram $R = R(a, b; c, d)$ such that $c - a \geq 2L_n$ and $d - b \geq 2H_n$, we have that for all $\eta \in V(n, R)$, there is no spanned critical parallelogram in η intersecting $R(a + L_n, b + H_n; c - L_n, d - H_n)$.*

We first deduce Proposition 3.7 from Lemma 3.10.

³As is standard [31], we say that the parallelograms R_1, \dots, R_k are disjointly spanned in η if one can find disjoint sets $X_1, \dots, X_k \subset \eta$ such that $\eta'_{X_i} = 0$ implies that R_i is spanned in η' for all $1 \leq i \leq k$.

Proof of Proposition 3.7, assuming Lemma 3.10. Clearly, it suffices to prove that $\mathbb{P}_\mu(\tau_0 \geq 2T) \geq 1/2$. Let $\tau' = \inf\{t \geq 0, 0 \text{ is locally infectable in } \eta(t)\}$. Clearly, $\tau' \leq \tau_0$. We denote $R = R(-L/2, -H/2; L/2, H/2)$. By Lemmas 3.6 and 3.10 and Definition 3.9 it follows that the event $\tau' \geq 2T$ contains the event G defined as the intersection of the following.

G_1 : $\tau' > 0$, i.e. 0 is not locally infectable in $\eta(0)$.

G_2 : There is no critical parallelogram spanned in $(\eta(0))_R$.

G_3 : For all $0 \leq t \leq 2T$, no $n+1$ critical parallelograms are disjointly spanned in $(\eta(t))_R$.

G_4 : For all $0 \leq t \leq 2T$, R has no crossing for $(\eta(t))_R$.

We next brutally bound the probability of these four events.

By assumption $1 - \mathbb{P}_\mu(G_1) \leq 1/8$. By the union bound on all (discrete) critical parallelograms intersecting R (recall Observation 3.2) we have

$$1 - \mathbb{P}_\mu(G_2) \leq O(LHK^2)\rho \leq \frac{LHK^3\rho}{8} \leq \frac{1}{8(TLH)^{1/(n+1)}} \leq \frac{1}{8}.$$

Let N denote the number of clock rings in R between 0 and $2T$ (the clock rings are the times at which updates are attempted by the KCM, see Section 1.1, at each site these times form a Poisson point process of parameter 1 independent from those of other sites). Let $\eta^{(j)}$ denote the restriction of the configuration to $R \cap \mathbb{Z}^2$ after the j -th such clock ring. By the weak law of large numbers we have $\mathbb{P}_\mu(N \geq O(TLH)) \leq 1/16$. If for any $\eta \in \{0, 1\}^{R \cap \mathbb{Z}^2}$ we write $\mathcal{D}_{n+1}(\eta) = \{\text{there are } n+1 \text{ parallelograms disjointly spanned in } \eta\}$, then by stationarity, the BK inequality [31] and the union bound, we get

$$\begin{aligned} 1 - \mathbb{P}_\mu(G_3) &\leq \mathbb{P}_\mu(N \geq O(TLH)) + \sum_{j=0}^{O(TLH)} \mathbb{P}_\mu(\mathcal{D}_{n+1}(\eta^{(j)})) \\ &\leq \frac{1}{16} + O(TLH)\mu(\mathcal{D}_{n+1}((\eta)_R)) \\ &\leq \frac{1}{16} + TLH(O(LHK^2)\rho)^{n+1} \leq \frac{1}{8}. \end{aligned}$$

Similarly, we have

$$1 - \mathbb{P}_\mu(G_4) \leq \frac{1}{16} + TLH \cdot O(L+H) \max(p_\downarrow, p_\leftarrow) \leq \frac{1}{8}.$$

We then conclude that

$$1 - \mathbb{P}_\mu(\tau_0 \geq 2T) \leq \sum_{i=1}^4 (1 - \mathbb{P}_\mu(G_i)) \leq \frac{1}{2}$$

by the assumptions of the proposition. \square

Proof of Lemma 3.10. We will prove the lemma by induction on n . For any n let us call \mathcal{H}_n the statement of the lemma for n . \mathcal{H}_0 holds by definition. Let $n \geq 1$ and assume that \mathcal{H}_{n-1} holds. Let $R = R(a, b; c, d)$ be such that $c - a \geq 2L_n$ and $d - b \geq 2H_n$. We will prove Lemma 3.10 by showing \mathcal{H}_n using the following result, whose proof we postpone for the moment.

Lemma 3.11. *For all $\eta \in V(n, R) \setminus G(R)$, there exists a critical parallelogram not intersecting*

$$R' = R(a + L_n - L_{n-1}, b + H_n - H_{n-1}; c - (L_n - L_{n-1}), d - (H_n - H_{n-1}))$$

that is spanned in η (see Figure 2).

From Lemma 3.11 we have that a spanned critical parallelogram remains outside R' , so there are at most $n-1$ critical parallelograms disjointly spanned in $\eta_{R'}$ for any $\eta \in V(n, R)$. Hence, for any $\eta \in V(n, R)$ we have $\eta_{R'} \in V(n-1, R')$, so we can apply \mathcal{H}_{n-1} to R' , which directly yields \mathcal{H}_n and concludes the proof of Lemma 3.10. \square

Consequently, it is enough to prove Lemma 3.11.

Proof of Lemma 3.11. Let us start by introducing the following geometric regions, represented in Figure 2.

$$\begin{aligned} R_\ell &= \overset{\circ}{R}(a + 2L_{n-1} + 7K, b + 2H_{n-1} + 7K; c - L_{n-1} - 7K, d - 2H_{n-1} - 7K) \\ R_{\ell, \downarrow} &= R(a, b; a + 2L_{n-1} + 7K, d) \\ R_{\ell, \uparrow} &= R(c - 2L_{n-1} - 7K, b; c, d) \\ R'_{\ell, \downarrow} &= R(a + L_{n-1}, b + H_{n-1}, a + L_{n-1} + 7K, d - H_{n-1}) \\ R'_{\ell, \uparrow} &= R(c - L_{n-1} - 7K, b + H_{n-1}, c - L_{n-1}, d - H_{n-1}) \end{aligned}$$

We also define similar regions with index h instead of ℓ . Moreover, we recall R' from the statement of Lemma 3.11. We further define the two frames (see Figure 2)

$$\begin{aligned} B &= R'_{\ell, \downarrow} \cup R'_{\ell, \uparrow} \cup R'_{h, \downarrow} \cup R'_{h, \uparrow} \\ B' &= R(a + L_{n-1} + 3K, b + H_{n-1} + 3K; c - L_{n-1} - 3K, d - H_{n-1} - 3K) \setminus \\ &\quad \overset{\circ}{R}(a + L_{n-1} + 4K, b + H_{n-1} + 4K; c - L_{n-1} - 4K, d - H_{n-1} - 4K). \end{aligned}$$

We next define some more subtle regions depending on the configuration.




Figure 2: The setting of the proof of Lemma 3.11. For the figure we assume that $u_3 = 0$ and $u_4 = \pi/2$. B' is the frame with thickened boundary, R_ℓ and R_h are the overlapping regions in dark gray. The regions $R'_{\ell,\downarrow}$, $R'_{h,\downarrow}$, $R'_{\ell,\uparrow}$ and $R'_{h,\uparrow}$ are in lighter gray and the frame formed by their union is B . The horizontally (resp. vertically) hatched regions are $R_{h,\downarrow}$ and $R_{h,\uparrow}$ (resp. $R_{\ell,\downarrow}$ and $R_{\ell,\uparrow}$). The contour inside B' is γ and its diagonally hatched interior is $\dot{\gamma}$. All the regions drawn are closed subsets of \mathbb{R}^2 with the exception of R_ℓ , R_h and $\dot{\gamma}$, which are open. The thicker version, $\bar{\gamma}$, of γ and the set $F \subset \mathbb{Z}^2$ defined in (4) are not drawn.

Claim 3.12. Let $\eta \in \{0, 1\}^{R \cap \mathbb{Z}^2}$ be such that every critical parallelogram spanned in η intersects R' . Then there exists a closed contour $\gamma \subset \mathbb{R}^2$ (that is, a self-avoiding and closed path obtained by connecting sites of \mathbb{Z}^2 by straight lines linking a site to its left, top, right or bottom neighbour) satisfying the following properties:

- $\gamma \subset B'$.
- $d(\gamma, R \setminus B') \geq C_1^2$.
- Every $s \in \bar{\gamma}$ is not locally infectable in η , where

$$\bar{\gamma} = \{s \in \mathbb{Z}^2 \mid d(s, \gamma) \leq C_1\}.$$

- The (topologically open) interior, $\mathring{\gamma} \subset \mathbb{R}^2$, defined by γ contains R' .

Proof. We proceed by renormalization. Let H denote the regular hexagon centred at the origin with diameter C_1^3 and having two horizontal sides. Consider the tiling of the plane with translates of H and denote by \mathbb{T} the triangular lattice formed by their centres. Let $T = \{t \in \mathbb{T} \mid H + t \subset B'\}$ be the sites of \mathbb{T} corresponding to B' . We say that a site $t \in T$ is *open* if no site in $(t + H) \cap \mathbb{Z}^2$ is locally infectable in η .

If there exists a contour of open sites in T surrounding R' (where a contour in \mathbb{T} is a self-avoiding and closed path in the graph $(\mathbb{T}, \{(t, t') \in \mathbb{T}^2 \mid t + H \text{ and } t' + H \text{ share a side}\})$), we may choose γ approximating this contour, which clearly satisfies the conditions of the claim. Assume that such a contour does not exist. Then there is a path of closed sites in T from the inner to the outer boundary of T . In particular, this path yields a strongly connected set X of sites of \mathbb{Z}^2 that are locally infectable in η , with diameter at least $K - 4C_1^3$, contained in either the “left part” of the frame B' , defined as

$$R''_{\ell, \downarrow} = R(a + L_{n-1} + 3K, b + H_{n-1} + 3K; a + L_{n-1} + 4K, d - H_{n-1} - 3K),$$

(see Figure 2) or in the top, right or bottom part of B' , defined similarly. Without loss of generality, assume that X is contained in $R''_{\ell, \downarrow}$. Since the sites of X are locally infectable in η , they are infectable in $\eta_{R''_{\ell, \downarrow}}$, where

$$R'''_{\ell, \downarrow} = R(a + L_{n-1} + K, b + H_{n-1} + K; a + L_{n-1} + 6K, d - H_{n-1} - K).$$

We denote X' the strongly connected component of $[\eta_{R'''_{\ell, \downarrow}}]$ containing X , and we consider the smallest parallelogram containing X' . By Observation 3.3, it is spanned in $\eta_{R'''_{\ell, \downarrow}}$, thus by Lemma 3.4 there exists a critical parallelogram spanned in $\eta_{R'''_{\ell, \downarrow}}$, hence spanned in η and contained in $R'_{\ell, \downarrow}$. We deduce the existence of a critical parallelogram spanned in η not intersecting R' , hence a contradiction. \square

We fix $\eta \in \{0, 1\}^{R \cap \mathbb{Z}^2} \setminus G(R)$ such that R has no crossing and every critical parallelogram spanned in η intersects R' . Let

$$F = \{s \in B' \cap \mathbb{Z}^2 \mid s \text{ is not locally infectable in } \eta\} \quad (4)$$

and fix a contour γ , its thickened version $\bar{\gamma} \subset F$ and its interior $\mathring{\gamma}$ as provided by Claim 3.12 (see Figure 2).

We will prove that there is no n -good legal path from η to $G(R)$. Since legal paths can be reversed, this implies that there $\eta \notin V(n, R)$, which proves Lemma 3.11.

Let $(\eta^{(j)})_{0 \leq j \leq m}$ be an n -good legal path with $\eta^{(0)} = \eta$. We will use an induction on $j \in \{0, \dots, m\}$ to prove that $\eta^{(m)} \notin G(R)$. More precisely, we will prove by induction on j that the following properties hold for $j \in \{0, \dots, m\}$.

\mathcal{P}_j^1 For every $\zeta \in \{\ell, h\}$, there exists a critical parallelogram contained in R_ζ spanned in $\eta^{(j)}$.

\mathcal{P}_j^2 The sites of F are not locally infectable in $\eta^{(j)}$.

\mathcal{P}_j^3 For every $(\zeta, \xi) \in \{\ell, h\} \times \{\downarrow, \uparrow\}$, $\eta_{R_{\zeta, \xi}}^{(j)} \in V(n-1, R_{\zeta, \xi})$.

\mathcal{P}_j^4 $[\eta_{\mathring{\gamma}}^{(j)}] = [\eta_{\mathring{\gamma}}^{(0)}]$.

If \mathcal{P}_m^1 is satisfied, then there exists a critical parallelogram spanned in $\eta^{(m)}$, so $\eta^{(m)} \notin G(R)$, which proves Lemma 3.11, so it suffices to establish the induction.

Base: $j=0$. Firstly, by assumption, every critical parallelogram spanned in η intersects R' , so it is contained in $R_\ell \cap R_h$ and therefore \mathcal{P}_0^1 holds, since $\eta \notin G(R)$. Secondly, the definition of F , (4), implies \mathcal{P}_0^2 . In addition, \mathcal{P}_0^4 is trivial, so we only need to prove \mathcal{P}_0^3 .

Let $(\zeta, \xi) \in \{\ell, h\} \times \{\downarrow, \uparrow\}$. By the assumption on η and Observation 3.2 there is no critical parallelogram spanned in $\eta_{R_{\zeta, \xi}}$. Moreover, $\eta_{R_{\zeta, \xi}}$ has no crossing in $R_{\zeta, \xi}$, since η has no crossing in R . Thus, $\eta_{R_{\zeta, \xi}} \in G(R_{\zeta, \xi})$, hence $\eta_{R_{\zeta, \xi}} \in V(n-1, R_{\zeta, \xi})$, so \mathcal{P}_0^3 holds.

Induction step. Let $j \in \{0, \dots, m-1\}$, and suppose that $\mathcal{P}_j^1, \mathcal{P}_j^2, \mathcal{P}_j^3$ and \mathcal{P}_j^4 hold. Since $(\eta^{(k)})_{0 \leq k \leq m}$ is a legal path, we have $\eta^{(j+1)} = (\eta^{(j)})^s$ and $(s+U) \cap R \subset \eta^{(j)}$ for some $s \in R \cap \mathbb{Z}^2$ and $U \in \mathcal{U}$.

Our first claim means that “ F shields $\mathring{\gamma}$ from the influence of the exterior.”

Claim 3.13. \mathcal{P}_{j+1}^4 holds.

Proof. If $s \notin \mathring{\gamma}$, then $\eta_{\mathring{\gamma}}^{(j+1)} = \eta_{\mathring{\gamma}}^{(j)}$, so $[\eta_{\mathring{\gamma}}^{(j+1)}] = [\eta_{\mathring{\gamma}}^{(j)}] = [\eta_{\mathring{\gamma}}^{(0)}]$ by \mathcal{P}_j^4 . Moreover, if $s \in \mathring{\gamma}$, then $s+U \subset \mathring{\gamma} \cup \bar{\gamma}$. Furthermore, the sites of $\bar{\gamma}$ are in F , so by

\mathcal{P}_j^2 they are not locally infectable in $\eta^{(j)}$ and, in particular, $\eta_{\hat{\gamma}}^{(j)} = 1$. Thus, $s + U \subset \hat{\gamma}$ and so $[\eta_{\hat{\gamma}}^{(j+1)}] = [\eta_{\hat{\gamma}}^{(j)}] = [\eta_{\hat{\gamma}}^{(0)}]$ by \mathcal{P}_j^4 . \square

We next establish that at least one spanned critical parallelogram of $\eta_{\hat{\gamma}}^{(j+1)}$ remains “to the left” of $R_{\ell, \uparrow}$ (see Figure 2), as well as one “below” $R_{h, \uparrow}$ (these two parallelograms may be the same).

Claim 3.14. There exists a critical parallelogram contained in $\mathbb{H}_{u_3}(c - 2L_{n-1} - 7K)$ that is spanned in $\eta_{\hat{\gamma}}^{(j+1)}$ and similarly for $\mathbb{H}_{u_4}(d - 2H_{n-1} - 7K)$.

Proof. We will only treat $H = \mathbb{H}_{u_3}(c - 2L_{n-1} - 7K)$ as the argument for the other half-plane is the same. Assume for a contradiction that there is no critical parallelogram contained in H that is spanned in $\eta_{\hat{\gamma}}^{(j+1)}$.

By Claim 3.13 we have that $[\eta_{\hat{\gamma}}^{(j+1)}] = [\eta_{\hat{\gamma}}^{(0)}]$ and by assumption there exists a critical parallelogram D spanned in $\eta^{(0)}$ intersecting R' . D is then contained in $\hat{\gamma}$, hence spanned in $\eta_{\hat{\gamma}}^{(0)}$. We consider X a strongly connected set of $[D \cap \eta_{\hat{\gamma}}^{(0)}]$ such that the smallest parallelogram containing X is D , then X intersects $\bar{\mathbb{H}}_{u_3}(c - 2L_{n-1} - 9K - \ell)$. Therefore, if we call X' the strongly connected component of $[\eta_{\hat{\gamma}}^{(j+1)}]$ containing X , then X' intersects $\bar{\mathbb{H}}_{u_3}(c - 2L_{n-1} - 9K - \ell)$, and the smallest parallelogram D' containing X' contains X , thus it contains D hence it has diameter at least K/C_1 , since D is critical. By Observation 3.3, D' is then spanned in $\eta_{\hat{\gamma} \cap X'}^{(j+1)}$, hence by Lemma 3.4 there exists a critical parallelogram spanned for $\eta_{\hat{\gamma} \cap X'}^{(j+1)}$. By the assumption it cannot be contained in H , so X' intersects $\bar{\mathbb{H}}_{u_1}(-(c - 2L_{n-1} - 8K))$. Now, if we denote $a_0 = \max\{a' \mid X' \subset \bar{\mathbb{H}}_{u_1}(-a')\}$, then $a_0 \geq a$, as $X' \subset [\eta_{\hat{\gamma}}^{(j+1)}]$ and u_1 is a stable direction, and $a_0 \leq c - 2L_{n-1} - 9K - \ell$. However, by Observation 3.3, $X' = [\eta_{\hat{\gamma} \cap X'}^{(j+1)}]$, so $R_{X'} = R(a_0, b, a_0 + \ell, d)$ is u_1 -crossed for $\eta_{\hat{\gamma}}^{(j+1)}$. Furthermore, any critical parallelogram intersecting $R_{X'}$ is contained in H , so by Observation 3.2 and the assumption, there is no critical parallelogram spanned for $\eta_{\hat{\gamma}}^{(j+1)} \cap R_{X'}$. This yields that $C_{R_{X'}}^{u_1}$ occurs for $\eta^{(j+1)}$, which is impossible since $\eta^{(j+1)}$ has no crossing as a configuration in a n -good legal path. \square

Since at least one spanned critical parallelogram is “to the left” of $R_{\ell, \uparrow}$, we will see that there are at most $n - 1$ inside it (and similarly for $R_{h, \uparrow}$), which will yield the following.

Claim 3.15. $\eta_{R_{\ell, \uparrow}}^{(j+1)} \in V(n - 1, R_{\ell, \uparrow})$ and $\eta_{R_{h, \uparrow}}^{(j+1)} \in V(n - 1, R_{h, \uparrow})$.

Proof. We will only prove $\eta_{R_{\ell,\uparrow}}^{(j+1)} \in V(n-1, R_{\ell,\uparrow})$, as the other proof is similar. By \mathcal{P}_j^3 it suffices to prove that $\eta_{R_{\ell,\uparrow}}^{(j+1)}$ is $(n-1)$ -good.

$R_{\ell,\uparrow}$ has no crossing for $\eta_{R_{\ell,\uparrow}}^{(j+1)}$ because R has no crossing for $\eta^{(j+1)}$. Let D_1, \dots, D_k be critical parallelograms that are disjointly spanned in $\eta_{R_{\ell,\uparrow}}^{(j+1)}$ with maximal k . By Claim 3.14, there exists a critical parallelogram $D \subset \mathbb{H}_{u_3}(c - 2L_{n-1} - 7K)$ that is spanned in $\eta_{\tilde{\gamma}}^{(j+1)}$ and, therefore, also in $\eta^{(j+1)}$. Since D is disjoint from $R_{\ell,\uparrow}$, we deduce that D_1, \dots, D_k, D are disjointly spanned in $\eta^{(j+1)}$, so $\eta^{(j+1)}$ contains $k+1$ disjointly spanned critical parallelograms. Since $\eta^{(j+1)}$ is n -good, we get $k \leq n-1$. \square

We are now ready to prove that a spanned critical parallelogram in $\mathbb{H}_{u_4}(d - 2H_{n-1} - 7K)$ provided by Claim 3.14 is in fact in R_ℓ (and similarly for R_h). In a nutshell, Claim 3.14 prevents the parallelogram from being “too far up,” being “too far left or down” induces a crossing and if the parallelogram is “too far right” we may apply \mathcal{H}_{n-1} by Claim 3.15.

Claim 3.16. \mathcal{P}_{j+1}^1 holds.

Proof. We only treat R_ℓ , as R_h is similar. Assume for a contradiction that there is no critical parallelogram contained in R_ℓ spanned in $\eta^{(j+1)}$.

By Claim 3.14, there exists a critical parallelogram D such that $D \cap R_{h,\uparrow} = \emptyset$ that is spanned in $\eta_{\tilde{\gamma}}^{(j+1)}$. By assumption $D \not\subset R_\ell$. There are three possibilities:

- a) $D \cap R_{\ell,\downarrow} \neq \emptyset$, i.e. D is “too far left;”
- b) $D \cap R_{h,\downarrow} \neq \emptyset$, i.e. D is “too far down;”
- c) $D \cap \bar{\mathbb{H}}_{u_1}(-(c - L_{n-1} - 7K)) \neq \emptyset$, i.e. D is “too far right.”

We first assume for a contradiction that case c) occurs. Since D is spanned in $\eta_{\tilde{\gamma}}^{(j+1)}$, by Observation 3.2 D intersects $\tilde{\gamma}$, hence since D is critical, the intersection of D and $\bar{\mathbb{H}}_{u_1}(-(c - L_{n-1} - 7K))$ is in $R'_{\ell,\uparrow}$. However, this contradicts \mathcal{H}_{n-1} applied to $\eta_{R_{\ell,\uparrow}}^{(j+1)} \in V(n-1, R_{\ell,\uparrow})$ by Claim 3.15.

Cases a) and b) being analogous, we assume for a contradiction that case a) occurs. Since D is spanned in $\eta_{\tilde{\gamma}}^{(j+1)}$, there exists a strongly connected set $X \subset [D \cap \eta_{\tilde{\gamma}}^{(j+1)}]$ such that D is the smallest parallelogram containing X . Let X' be the strongly connected component of $[\eta_{\tilde{\gamma}}^{(j+1)}]$ containing X , and D' be the smallest parallelogram containing X' . Then D' contains X , hence it contains D and $\text{diam}(D') \geq K/C_1$, since D is critical. Furthermore, X' is a strongly connected component of $[\eta_{\tilde{\gamma}}^{(j+1)}] = [\eta_{\tilde{\gamma}}^{(0)}]$ by Claim 3.13, so

$X' = [\eta_{\hat{\gamma}}^{(0)} \cap X']$ by Observation 3.3, hence D' is spanned in $\eta_{\hat{\gamma} \cap X'}^{(0)}$. By Lemma 3.4 there exists a critical parallelogram D'' that is spanned in $\eta_{\hat{\gamma} \cap X'}^{(0)}$. Then D'' is spanned in $\eta^{(0)} = \eta$ and therefore intersects R' , so by Observation 3.2, X' intersects $\bar{\mathbb{H}}_{u_1}(-a - 2L_{n-1} - 8K - \ell)$. Since D intersects $R_{\ell, \downarrow}$ by assumption, X intersects $\bar{\mathbb{H}}_{u_3}(a + 2L_{n-1} + 7K)$, thus X' intersects this half-plane as well. Now, if we denote $a_0 = \max\{a' \mid X' \subset \bar{\mathbb{H}}_{u_1}(-a')\}$, then $a_0 \geq a$, as $X' \subset [\eta_{\hat{\gamma}}^{(0)}]$ and u_1 is a stable direction, and $a_0 \leq a + 2L_{n-1} + 7K$. However, $X' = [\eta_{\hat{\gamma} \cap X'}^{(0)}]$, so $R_{X'} = R(a_0, b, a_0 + \ell, d)$ is u_1 -crossed for $\eta^{(0)}$. However, $\eta^{(0)}$ has no crossing, so $C_{R_{X'}}^{u_1}$ does not occur for $\eta^{(0)}$. Hence, recalling Definition 3.5 and Observation 3.2, there exists a critical parallelogram spanned in $\eta_{R_{X'}}^{(0)}$, hence in $\eta^{(0)}$, intersecting $R_{X'}$. Since critical parallelograms have diameter at most K , this contradicts the fact that all critical parallelograms spanned in $\eta^{(0)} = \eta$ intersect R' . \square

We are now able to use an argument similar to that of Claim 3.15: having a spanned critical parallelogram in R_ℓ entails that at most $n - 1$ of them are in $R_{\ell, \downarrow}$ (and similarly for $R_{h, \downarrow}$), hence the following.

Claim 3.17. \mathcal{P}_{j+1}^3 holds.

Proof. Thanks to Claim 3.15, it remains only to prove $\eta_{R_{\ell, \downarrow}}^{(j+1)} \in V(n - 1, R_{\ell, \downarrow})$ and $\eta_{R_{h, \downarrow}}^{(j+1)} \in V(n - 1, R_{h, \downarrow})$. The proof is actually the same as in Claim 3.15, replacing $\bar{\mathbb{H}}_{u_3}(c - 2L_{n-1} - 7K)$ by R_ℓ and Claim 3.14 by Claim 3.16. \square

Finally, we will see that for a site of F to be locally infectable in $\eta^{(j+1)}$, by Lemma 3.6 one would need a critical parallelogram spanned in $\eta^{(j+1)}$ near the site, which is impossible by \mathcal{H}_{n-1} since F is contained in the middle parts of the $R_{\zeta, \xi}$ and \mathcal{P}_{j+1}^3 holds.

Claim 3.18. \mathcal{P}_{j+1}^2 holds.

Proof. Assume for a contradiction that there exists $s' \in F \subset B'$ that is locally infectable in $\eta^{(j+1)}$. Then \mathcal{P}_j^2 and Lemma 3.6 imply that there exist $\zeta \in \{\ell, h\}$, $\xi \in \{\downarrow, \uparrow\}$ and a critical parallelogram $D \subset R'_{\zeta, \xi}$ spanned in $\eta_{R_{\zeta, \xi}}^{(j+1)}$. However, by Claim 3.17 $\eta_{R_{\zeta, \xi}}^{(j+1)} \in V(n - 1, R_{\zeta, \xi})$, so \mathcal{H}_{n-1} yields the desired contradiction. \square

Claims 3.13, 3.16, 3.17 and 3.18 together establish the induction step, which completes the proof of Lemma 3.11. \square

4 Application of Proposition 3.7

In this section we derive Theorems 1 and 2 from Proposition 3.7. For that purpose we require some estimates on the probabilities appearing in the statement of the proposition, which are mostly proved in the appendices. We restate those results below as needed. Throughout the section \mathcal{U} is a critical update family with difficulty α subject to further assumptions recalled in each subsection. Such a family admits two non-collinear rational stable directions. We set u_1 and u_2 to be two arbitrary such directions, which will be chosen differently for each class of update families. We will use the definitions of Section 3 with u_1 and u_2 .

Let us start with the easiest estimate.

Lemma 4.1. *Let $q^{-1/2} \leq K \leq q^{-2\alpha}$. Then,*

$$\mu(0 \text{ is locally infectable}) \leq 1/8.$$

Proof. Let $R = R(-2K, -2K; 2K, 2K)$. Since \mathcal{U} is critical, $\text{diam}([\eta_R]) \leq C_1 K$, so, starting the bootstrap percolation dynamics with η_R , the origin is either infected in time at most $C_1^3 K^2$ or not at all. We conclude using e.g. [4, Theorem 1.4]. \square

Turning to the probability of spanning there are two cases to consider. For unbalanced models the following is essentially a reformulation of the most difficult result of [3].

Lemma 4.2. *Assume that \mathcal{U} is unbalanced and $K = q^{-\alpha-1/4}$. Then, for any critical parallelogram D we have*

$$\mu(D \text{ is spanned}) \leq \exp\left(-\frac{(\log(1/q))^2}{C_5 q^\alpha}\right).$$

Proof. By definition if D is a spanned critical parallelogram, then there exists a strongly connected set $X \subset [\eta \cap D]$ with diameter at least $\text{diam}(D)/C_1$. If X' is the strongly connected component of X in $[\eta \cap D]$, then by Observation 3.3, $X' = [\eta \cap D \cap X']$. Therefore, if D' is the smallest S_U -droplet in the sense of [3, Definition 2.1] with the S_U defined in [3, Lemma 6.2] such that D' contains X' , then D' is internally spanned by $\eta \cap D$ (in the sense of [3, Definition 2.4]⁴), and $\text{diam}(D') \geq \text{diam}(D)/C_1 \geq K/C_1^2$. Repeating the proof of [3, Lemma 8.37], we get that there is a critical droplet in the sense of [3, Definition 2.5] internally spanned by $\eta \cap D$. Then the union bound over such droplets and [3, Lemma 8.36] yield the desired result. \square

⁴Technically, it is not exactly the case, as [3] uses a different choice of constants. However, their results that we use still hold in our setting.

Concerning balanced models, in Appendix A (Corollary A.11) we establish the following.

Lemma 4.3. *Assume that \mathcal{U} is balanced and $q^{-\alpha}/C_5 \leq K \leq q^{-2\alpha}$. Then for any critical parallelogram D we have*

$$\mu(D \text{ is spanned}) \leq \exp\left(-\frac{1}{q^\alpha C_5}\right).$$

For the remaining conditions of Proposition 3.7 we need to distinguish the different classes of models.

4.1 Proof of Theorem 1

In this section we assume that \mathcal{U} has an infinite number of stable directions. We then choose two rational directions $u_1 < u_2 < u_1 + \pi$ sufficiently close to each other, such that all directions in $[2u_1 - u_2, 2u_2 - u_1]$ are stable and u_1, u_2 satisfy a technical condition which the reader is advised to ignore, namely that u_1 and u_2 are constructed like the eponymous directions in the proof of [17, Lemma 4.1].

4.1.1 Proof of Theorem 1(a)

For this section we further assume that \mathcal{U} is unbalanced. We fix the values of the parameters of Proposition 3.7 as follows.

$$\begin{aligned} K &= q^{-\alpha-1/4} & \ell &= q^{-4\alpha} & L &= \exp\left(\frac{(\log(1/q))^2}{C_6 q^\alpha}\right) \\ T &= \exp\left(\frac{(\log(1/q))^4}{C_6^2 q^{2\alpha}}\right) & h &= q^{-4\alpha} & H &= \exp\left(\frac{(\log(1/q))^2}{C_6 q^\alpha}\right). \end{aligned}$$

Then Theorem 1(a) follows directly from Proposition 3.7 and the upper bound of [25, Theorem 2(a)]. The hypotheses of the proposition follow from the choice of parameters, Lemmas 4.1 and 4.2, and the following bound on the crossing probabilities proved in Appendix B (Lemma B.4).

Lemma 4.4. *With the notation and assumptions above we have*

$$\max(p_{\leftarrow}, p_{\downarrow}) \leq \exp(-q^{-3\alpha}).$$

4.1.2 Proof of Theorem 1(b)

For this section we further assume that \mathcal{U} is balanced. We fix the values of the parameters of Proposition 3.7 as follows.

$$\begin{aligned} K &= q^{-\alpha} & \ell &= q^{-4\alpha} & L &= \exp\left(\frac{1}{C_6 q^\alpha}\right) \\ T &= \exp\left(\frac{1}{C_6^2 q^{2\alpha}}\right) & h &= q^{-4\alpha} & H &= \exp\left(\frac{1}{C_6 q^\alpha}\right). \end{aligned}$$

Then Theorem 1(b) follows directly from Proposition 3.7. The hypotheses of the proposition follow from the choice of parameters, Lemmas 4.1 and 4.3, and Lemma 4.4, which still applies.

4.2 Proof of Theorem 2

4.2.1 Proof of Theorem 2(c)

In this section we assume that \mathcal{U} is unbalanced, rooted and has a finite number of stable directions. Therefore, we can find rational directions u_1, u_2, u_3 such that $u_1 + \pi = u_3$, $u_2 \in (u_1, u_3)$ and $\alpha(u_i) \geq \alpha + 1$ for all $i \in \{1, 2, 3\}$.

We fix the values of the parameters of Proposition 3.7 as follows.

$$\begin{aligned} K &= q^{-\alpha-1/4} & \ell &= q^{-\alpha-5/8} & L &= q^{-\alpha-3/4} \\ T &= \exp\left(\frac{(\log(1/q))^3}{C_6 q^\alpha}\right) & h &= q^{-\alpha-5/8} & H &= q^{-\alpha-3/4}. \end{aligned}$$

Then Theorem 2(c) follows directly from Proposition 3.7 and the upper bound of [19, Theorem 1]. The hypotheses of the proposition follow from the choice of parameters, Lemmas 4.1 and 4.2, and the following bound on the crossing probabilities proved in Appendix B (Lemma B.3).

Lemma 4.5. *With the notation and assumptions above we have*

$$\max(p_{\leftarrow}, p_{\downarrow}) \leq \exp(-q^{-\alpha-1/4}).$$

4.2.2 Proof of Theorem 2(e)

In this section we assume that \mathcal{U} is balanced, rooted and with a finite number of stable directions. Therefore, we can find non-opposite rational directions u_1, u_2 such that $\alpha(u_1) \geq \alpha + 1$ and $\alpha(u_2) \geq \alpha + 1$. We fix the values of the

parameters of Proposition 3.7 as follows.

$$\begin{aligned} K &= 1/(C_5 q^\alpha) & \ell &= q^{-\alpha-1/2} & L &= q^{-\alpha-3/4} \\ T &= \exp\left(\frac{\log(1/q)}{C_6 q^\alpha}\right) & h &= q^{-\alpha-1/2} & H &= q^{-\alpha-3/4}. \end{aligned}$$

Then Theorem 2(e) follows directly from Proposition 3.7. The hypotheses of the proposition follow from the choice of parameters, Lemmas 4.1 and 4.3, and Lemma 4.5, which still holds.

4.2.3 Proof of Theorem 2(f)

In this section we assume that \mathcal{U} is semi-directed. Therefore, we can find non-opposite rational directions u_1, u_2 such that $\alpha(u_1) = \alpha$ and $\alpha(u_2) \geq \alpha + 1$. We fix the values of the parameters of Proposition 3.7 as follows.

$$\begin{aligned} K &= 1/(C_5 q^\alpha) & \ell &= q^{-\alpha-1/2} & L &= q^{-\alpha-3/4} \\ T &= \exp\left(\frac{\log \log(1/q)}{C_6^3 q^\alpha}\right) & h &= \frac{\log \log(1/q)}{q^\alpha} & H &= \frac{(\log(1/q))^{1/4}}{q^\alpha}. \end{aligned}$$

Then Theorem 2(f) follows directly from Proposition 3.7. The hypotheses of the proposition follow immediately from the choice of parameters, Lemmas 4.1 and 4.3, and the following bound on the crossing probabilities proved in Appendix B (Lemma B.3).

Lemma 4.6. *With the notation and assumptions above we have*

$$p_{\leftarrow} \leq \exp(-q^{-\alpha-1/4}) \quad p_{\downarrow} \leq \exp\left(-\frac{\log \log(1/q)}{2C_6^2 q^\alpha}\right).$$

Acknowledgements

This work has been supported by ERC Starting Grant 680275 MALIG. We wish to thank Cristina Toninelli and Fabio Martinelli for enlightening discussions on the universality of \mathcal{U} -KCM. The first author would like to thank ÉPFL for the hospitality during his visit.

A Bounds on spanning

Relation to previous works Let us start by explaining why additional arguments are needed, as specialists would probably expect such bounds to

be automatic. In [3] two main algorithms were used—the covering and the spanning ones. The former provides bounds of the type we need but for a notion of covered droplet invoking only the initial configuration. Furthermore, its treatment involves a somewhat technical geometric result on subadditivity of the size of droplets. On the other hand, the spanning algorithm works with the closure of the initial configuration inside droplets, which obstructs obtaining results analogous to those for the covering algorithm in the same way. Yet, it is the spanning algorithm which is the most useful and particularly so for unbalanced models. In [3] an inductive multi-scale scheme was used to bootstrap the bounds on the probability of droplets being spanned from a size which is easily controlled by the more rudimentary predecessor of the covering algorithm developed in [4]. This fairly technical procedure can be circumvented using our method. Indeed, if one has bounds analogous to the ones for covered droplets up to size $1/q^{\alpha(\mathcal{U})}$, one can directly prove the result of [3] in one step, which was made there as well.

The reason why in [3] one could not directly transfer the easier bounds on covering, which were established there anyway, to spanning is that the covering algorithm there lacks the key property of being essentially closure-invariant in a sense made precise below. This property was one of the main features gained in [17] by using a less wasteful notion of cluster. Therefore, we accomplish our goal as follows. We carry through (a simplified version of) the scheme of [17] to obtain general bounds for droplets covered in the sense of [17] and we use the key closure lemma (see below) to directly transfer those to spanning. On the more technical level, we should mention that analogous bounds on spanning were established in [3] in the course of their induction, but the proof needlessly uses that the model is unbalanced and constrains the choice of directions used for defining droplets, which we will need to choose freely. Moreover, [17] made unnecessary use of the existence of strongly stable directions⁵, which is only needed for treating the algorithm with boundary condition. We are thus obliged to review the proofs. The reader familiar with the details of [17] would probably be satisfied by skipping directly to Appendix B and consulting the statements as needed there.

Outline The appendix is structured as follows. In section A.1 we recall several results from [17], leading up to Lemma A.6 providing good bounds on the probability of being covered (in a sense made precise below) and to the Closure lemma A.7 relating the results of the covering algorithm for a set and for its closure. In section A.2, using the latter lemma we transfer

⁵Strongly stable directions are those contained in the topological interior of the set of stable directions.

the bounds of the former one to the notion of spanning used throughout the body of the paper. Section A.3 establishes, yet again, the same bounds on the probability of spanned droplets occurring, but in the presence of an infected boundary, following the same reasoning and relying more closely on [17].

Notation For the remainder of the paper we fix an arbitrary critical update family \mathcal{U} with difficulty α . Following [17] we consider constants

$$1 \ll C_1 \ll C'_2 \ll C_2 \ll C_3 \ll C'_4 \ll C_4 \ll C_5 \ll C_6$$

such that each one is larger than a suitable function of the previous ones, depending on \mathcal{T} , \mathcal{T}_0 , \mathcal{S}_u , etc. to be defined below and on \mathcal{U} . These constants do not depend on q , which is always assumed small enough, as we are interested in $q \rightarrow 0$.

For any finite set of directions $\mathcal{V} \subset S^1$ a \mathcal{V} -droplet is a set of the form $\bigcap_{v \in \mathcal{V}} \bar{\mathbb{H}}_v(a_v)$ for some $a_v \in \mathbb{R}$.

A.1 Covering and closure

We start by studying the covering algorithm in the spirit of [17, Section 5] (but without the boundary and rugged edge present there). The reader is invited to consult that work for most proofs and more details, as indicated below. By definition 1.2 we can fix a set of non semi-isolated rational stable directions⁶ \mathcal{T}_0 with difficulty at least α , such that the convex envelope of the elements of \mathcal{T}_0 contain 0 in its interior and either

- $|\mathcal{T}_0| = 3$ or
- $|\mathcal{T}_0| = 4$ and one has $\mathcal{T}_0 = \{u, v, u + \pi, v + \pi\}$ for some $u, v \in S^1$.

Let Γ be the graph with vertex set \mathbb{Z}^2 but with $x \sim y$ iff $\|x - y\| \leq C_2$.

Definition A.1 ([17, Definitions 5.1 and 5.3]). Fix a finite set $Z \subset \mathbb{Z}^2$. Let κ be a connected component of the subgraph of Γ induced by the vertex set Z .

- κ is a *crumb* for Z if there exists a set $P_\kappa \subset \mathbb{Z}^2$ such that $[P_\kappa] \supset \kappa$ and $|P_\kappa| = \alpha - 1$.
- If κ is not a crumb for Z , we say that a $C \subset \kappa$ is a α -*cluster* (or simply cluster) of Z if the following conditions hold

⁶Semi-isolated stable directions are the endpoints of intervals of stable directions with nonempty interior.

- $\text{diam}(C) \leq C_3$.
- C is connected in Γ .
- For all $x \in \kappa \setminus C$ and $y \in C$ such that $x \sim y$ in Γ we have $\text{diam}(C \cup \{x\}) > C_3$.

It can be proved [17, Observation 5.4] that any cluster contains at least α sites. Moreover, Corollary 5.17 of [17] yields that a crumb has diameter at most αC_2 . For a cluster C we denote by $Q(C)$ the smallest \mathcal{T}_0 -droplet containing the set $\{x \in \mathbb{R}^2 : d(x, C) \leq C_4\}$.

We next define the covering algorithm we will use. It is an adaptation of the droplet algorithm of [17] and should not be confused with the covering algorithms of [3, 4].

Definition A.2 (Covering algorithm). Given a finite set $Z \subset \mathbb{Z}^2$ of infections the *covering algorithm* outputs a set \mathcal{D} of disjoint \mathcal{T}_0 -droplets as follows.

- Form an initial collection \mathcal{D} of \mathcal{T}_0 -droplets consisting of $Q(C)$ for all clusters C of Z .
- Whenever there exist $D_1, D_2 \in \mathcal{D}$ with $D_1 \cap D_2 \neq \emptyset$, replace them with the smallest \mathcal{T}_0 -droplet containing their union, which we denote by $D_1 \vee D_2$.
- Output the collection \mathcal{D} obtained when all \mathcal{T}_0 droplets in \mathcal{D} are disjoint.

Equivalently, \mathcal{D} is the minimal collection (with respect to inclusion of the union of its elements) of disjoint \mathcal{T}_0 -droplets containing the union of $Q(C)$ for all clusters C of Z . In particular, \mathcal{D} does not depend on the order in which droplets are merged.

We say that a \mathcal{T}_0 -droplet D is *covered* by a set Z of infections if the above algorithm for $Z \cap D$ outputs a \mathcal{T}_0 -droplet containing D .

We make the convention that all \mathcal{T}_0 -droplets have diameter at least C'_4 and contain a site of \mathbb{Z}^2 .

We next state some properties of the covering algorithm.

Lemma A.3 (Lemma 4.6 of [4]). *Let D_1 and D_2 be \mathcal{T}_0 -droplets such that $D_1 \cap D_2 \neq \emptyset$. Then*

$$\text{diam}(D_1 \vee D_2) \leq \text{diam}(D_1) + \text{diam}(D_2).$$

This immediately implies the Aizenman-Lebowitz lemma (see e.g. [4, Lemma 4.8]).

Lemma A.4 (Aizenman-Lebowitz). *Let Z be a set of infections and D be a \mathcal{T}_0 -droplet covered by Z . Then for all $C_1 C_4 \leq k \leq \text{diam}(D)$ there exists a \mathcal{T}_0 -droplet D' covered by Z with $k \leq \text{diam}(D') \leq 2k$.*

A further consequence of Lemma A.3 is the following.

Lemma A.5 (Lemma 5.14 of [17]). *Let Z be a set of infections and D be a \mathcal{T}_0 -droplet covered by Z . Then D contains at least $\lceil \text{diam}(D)/C_4^2 \rceil$ disjoint clusters of $Z \cap D$.*

We are now able to deduce the relevant bounds on covering following [17, Lemma 5.15].

Lemma A.6. *Let D be a \mathcal{T}_0 -droplet with $d = \text{diam}(D)$. Let $1 > \epsilon > 0$. Then we have*

$$\mu(D \text{ is covered}) \leq \begin{cases} q^{d\epsilon/(3C_4^2)} & \text{if } d \leq \frac{C_1}{q^{\alpha-\epsilon}} \\ e^{-C_1 C_4 d} & \text{if } \frac{1}{C_1 q^{\alpha-\epsilon}} \leq d \leq \frac{C_1}{e^{C_4^4} q^\alpha} \\ d^2 e^{-C_1/(C_5 q^\alpha)} & \text{if } \frac{1}{C_1 e^{C_4^4} q^\alpha} \leq d. \end{cases} \quad (5)$$

Proof. Let Z be the (random) set of infections in D . By Lemma A.5 we have that if D is covered, it contains at least $\lceil d/C_4^2 \rceil$ disjoint clusters of Z , each one having diameter at most C_3 and at least α sites. Thus, the union bound gives

$$\mu(D \text{ is covered}) \leq \binom{C_3^{2\alpha} d^2}{\lceil d/C_4^2 \rceil} q^{\alpha \lceil d/C_4^2 \rceil}.$$

For $d \leq C_4^2$ this gives $C_3^{2\alpha} d^2 q^\alpha$, which concludes the proof. For $C_4^2 \leq d \leq C_1/(e^{C_4^4} q^\alpha)$ we use the inequality $\binom{n}{k} \leq (ne/k)^k$ to obtain the desired bounds. For the case $d \geq 1/(e^{C_4^4} q^\alpha)$ we use Lemma A.4 to extract a smaller \mathcal{T}_0 -droplet D' covered by Z (hence intersecting D) with $1/(2e^{C_4^4} q^\alpha) \leq \text{diam}(D') \leq 1/(e^{C_4^4} q^\alpha)$. We then apply the second bound to D' and use the union bound to conclude. \square

We would now like to use analogous bounds on the probability of \mathcal{T}_0 -droplets being covered with initial condition $[Z]$ instead of Z . Unfortunately, we do not have access to the law of $[Z]$ when Z follows μ . Therefore, we rather bound the output of the covering algorithm for the closure using the original output. For that purpose, we define parallel notions of Γ' , *modified clusters* and *modified covering*, by replacing C_2 by C'_2 and C_4 by C'_4 .

We then have the following key property, whose proof is identical to the one of [17, Proposition 5.20], up to the relevant simplifications (we do not have rugged edges and there is no boundary).

Lemma A.7 (Closure). *Let $Z \subset \mathbb{Z}^2$ be a finite set and let \mathcal{D}' be the collection of \mathcal{T}_0 -droplets given by the modified covering algorithm with input $[Z]$. Let \mathcal{D} be the output of the covering algorithm for Z . Then*

$$\forall D' \in \mathcal{D}' \exists D \in \mathcal{D}, D' \subset D.$$

A.2 Spanning

Let \mathcal{T} be an arbitrary finite set of rational directions containing the origin in the interior of its convex envelope. We then generalise the notion of spanning from Definition 3.1.

Definition A.8 (Spanning). Let D be a \mathcal{T} -droplet. We say that D is *spanned* by $Z \subset \mathbb{Z}^2$ if there exists a set $C \subset [Z \cap D]$ connected in Γ' such that the smallest \mathcal{T} -droplet containing C is D .

We will need the following Aizenman-Lebowitz type lemma. Though this is a very classical result, some additional arguments are needed to prove it, because \mathcal{T} is not composed of stable directions.

Lemma A.9 (Aizenman-Lebowitz). *Let $Z \subset \mathbb{Z}^2$ and D be a \mathcal{T} -droplet spanned by Z with $\text{diam}(D) \geq C_1 C'_2$. Then for any $C'_2 \leq k \leq \text{diam}(D)$, there exists a \mathcal{T} -droplet D' spanned by $Z \cap D$ with $k \leq \text{diam}(D') \leq C_1 k$.*

Proof. If $k \geq \text{diam}(D)/C_1$ there is nothing to prove, as $D' = D$ is as desired. Assume $k \leq \text{diam}(D)/C_1$. Let C be a connected component of $[Z \cap D]$ in Γ' with maximal diameter. By Definition A.8 $\text{diam}(C) \geq \text{diam}(D)/\sqrt{C_1}$. By Observation 3.3 and [3, Lemma 6.18] (we use it although definitions slightly differ from [3], see Footnote 4) there exists $C' \subset C$ connected in Γ' such that $C' \subset [C' \cap Z \cap D]$ and $k \leq \text{diam}(C') \leq \sqrt{C_1} k$. Denoting D' the smallest \mathcal{T} -droplet containing C' , we are done. \square

Observation A.10. Let D be a \mathcal{T} -droplet spanned by $Z \subset \mathbb{Z}^2$ with $\text{diam}(D) \geq C_4$. Then there exists a \mathcal{T}_0 -droplet \bar{D} covered by Z , intersecting D and such that $\text{diam}(\bar{D}) = \Theta(\text{diam}(D))$.

Proof. Let C be as in Definition A.8. Notice that, since $\text{diam}(D) \geq C_3$, we can find modified clusters for $[Z \cap D]$ whose union is a connected set in Γ' containing C . Then there is a \mathcal{T}_0 -droplet in the output of the modified covering algorithm for $[Z \cap D]$ containing C . By Lemma A.7 there is also a \mathcal{T}_0 -droplet \bar{D} in the output of the covering algorithm for $Z \cap D$ containing C , so that $\text{diam}(\bar{D}) \geq \text{diam}(C) = \Omega(\text{diam}(D))$. But \bar{D} is at most the smallest \mathcal{T}_0 -droplet containing $\{x \in \mathbb{R}^2 : d(x, D) \leq C_4\}$, so $\text{diam}(\bar{D}) = \Theta(\text{diam}(D))$. Moreover, since \bar{D} is in the output of the covering algorithm for $Z \cap D$, it is covered by Z and intersects D . \square

We immediately deduce from this observation and Lemma A.6 the desired bounds on spanning.

Corollary A.11. *Let D be a \mathcal{T} -droplet with $d = \text{diam}(D)$ and let $1 > \epsilon > 0$. Then*

$$\mu(D \text{ is spanned}) \leq \begin{cases} q^{d\epsilon/C_5} & \text{if } d \leq q^{-\alpha+\epsilon} \\ e^{-2C_4d} & \text{if } q^{-\alpha+\epsilon} \leq d \leq \frac{C_1}{C_5q^\alpha} \\ d^{O(1)}e^{-2/(C_5q^\alpha)} & \text{if } \frac{1}{C_5q^\alpha} \leq d. \end{cases} \quad (6)$$

A.3 Boundary and spanning

We next turn to the treatment of an infinite infected boundary condition, following [17], which is applicable only for models with an infinite number of stable directions. Indeed, for a model with a finite number of stable directions a bounded set of infections next to the boundary can induce a set of supplementary infections and, thereby, a droplet of the size of the boundary, making similar algorithms useless. We therefore fix an update family \mathcal{U} with an infinite number of stable directions and difficulty α , to which the treatment of [17] applies.

For the rest of this section let $\mathcal{S}_u = \{u^-, u^+, v_1, v_2\}$ be a set of 4 directions chosen as in [17, Lemma 4.1]⁷ (we rename (u_1, u_2) from that work into (u^-, u^+) to avoid notational conflict) with $u = (u^- + u^+)/2$. The proof of [17] allows us to choose u^- and u^+ as close as we want, even depending on v_1 and v_2 . We will choose them close enough for our results to hold. Let $\partial = \mathbb{H}_u$. For any set $Z \subset \mathbb{Z}^2$ we write $[Z]_\partial = [Z \cup \partial] \setminus \partial$. We will use the term *cluster* in the sense of [17, Definition 5.3], extending Definition A.1 (crumbs close to ∂ are considered as clusters instead and Z is replaced by $Z \setminus \partial$). We replace the notion of DYD from [17] by that of \mathcal{S}_u -droplet and the notion of CDYD becomes that of *cut \mathcal{S}_u -droplet*—a nonempty set of the form

$$(\bar{\mathbb{H}}_{u^-}(x) \cap \bar{\mathbb{H}}_{u^+}(y)) \setminus \partial \quad (7)$$

for some $x, y \in \mathbb{R}$, which is a geometric triangle. We further replace the use of the diameter by considering the size $|\cdot|$ from [17, Definition 5.7]. Namely for a cut \mathcal{S}_u -droplet D we denote $|D| = \text{diam}(D)/C_1$, while for an \mathcal{S}_u -droplet D , $|D|$ denotes the length of its projection parallel to v_1 . We then define correspondingly an extension of the covering algorithm as in [17, Section 5.4] and a notion of covered (cut) \mathcal{S}_u -droplet. For the reader unfamiliar with

⁷It is not hard to see that in [17, Lemma 4.1], with a finite number of exceptions, given any rational strongly stable direction $u \in S^1$ we can define \mathcal{S}_u correspondingly.

[17], let us indicate that the change with respect to the covering algorithm of Definition A.2 corresponds to replacing at each stage of the algorithm any \mathcal{S}_u -droplet D intersecting ∂ by the smallest cut \mathcal{S}_u -droplet containing $D \setminus \partial$. The properties of [17, Section 5.5], analogous to Lemmas A.3-A.5 and A.7, remain valid for this setting. Furthermore, combining the proofs of Lemma A.6 and [17, Lemma 5.15] shows that the following holds.

Lemma A.12. *Let D be a cut \mathcal{S}_u -droplet or an \mathcal{S}_u -droplet not intersecting ∂ with $d = |D|$. Let $1 > \epsilon > 0$. Then (5) holds.*

We similarly extend Definition A.8 to the setting with boundary.

Definition A.13 (Spanning with boundary). We call *whole \mathcal{S}_u -droplet* any \mathcal{S}_u -droplet at distance at least C_3 from ∂ and, by abuse, we call collectively *\mathcal{S}_u -droplet* any cut or whole \mathcal{S}_u -droplet. We say that an \mathcal{S}_u -droplet D is *spanned* by $Z \subset \mathbb{Z}^2$ if there exists a set $C \subset [Z \cap D]_\partial$ connected in Γ' such that the smallest \mathcal{S}_u -droplet containing C is D .

We next recall several properties of the spanning algorithm following closely [3].

Definition A.14 (Definition 6.15 of [3]). Let $Z = \{z_1, \dots, z_{k_0}\}$ be a finite set of infections. Set $\mathcal{Z}^0 = \{Z_1^0, \dots, Z_{k_0}^0\}$ with $Z_i^0 = \{z_i\}$. For each $t \geq 0$ do the following.

- If there exist Z_i^t and Z_j^t such that $[Z_i^t]_\partial \cup [Z_j^t]_\partial$ is connected in Γ' , then set $\mathcal{Z}^{t+1} = (\mathcal{Z}^t \setminus \{Z_i^t, Z_j^t\}) \cup \{Z_i^t \cup Z_j^t\}$.
- Otherwise, define the *span* of Z by $\langle Z \rangle = \{D(Z^t), Z^t \in \mathcal{Z}^t\}$, where $D(Z')$ denotes the smallest \mathcal{S}_u -droplet containing Z' , and terminate the algorithm.

Similarly, for any $A \subset \mathbb{R}^2$ we denote $\langle A \rangle = \langle A \cap \mathbb{Z}^2 \rangle$.

Observation A.15 (Lemma 6.16 of [3]). We have $\langle Z \rangle = \{D(\kappa_1), \dots, D(\kappa_k)\}$, where the κ_i are the connected components of $[Z]_\partial$ in Γ' .

Observation A.16 (Lemma 6.17 of [3]). A nonempty \mathcal{S}_u -droplet is spanned iff $D \in \langle D \cap Z \rangle$.

Lemma A.17 (Lemma 6.21 of [3]). *Let Z be a finite set of at least two infections such that $[Z]_\partial$ is connected in Γ' . Then there exists a nontrivial partition $Z = Z_1 \sqcup Z_2$ such that $[Z_1]_\partial$, $[Z_2]_\partial$ and $[Z_1]_\partial \cup [Z_2]_\partial$ are connected in Γ' .*

The next lemma follows from the definition of size and [17, Lemma 5.12].

Lemma A.18. *For any \mathcal{S}_u -droplets D, D_1, D_2 with $|D_1| \geq C_3$ or $|D_2| \geq C_3$ such that $\langle D_1 \rangle = \{D_1\}$, $\langle D_2 \rangle = \{D_2\}$ and $\langle D_1 \cup D_2 \rangle = \{D\}$ we have $|D_1|/C_1 \leq |D| \leq |D_1| + |D_2| + O(C'_2)$.*

This standardly implies (see e.g. [3, Lemma 6.18]) the following.

Lemma A.19 (Aizenman-Lebowitz). *Let D be a spanned \mathcal{S}_u -droplet and $C_3 \leq k \leq |D|$. Then there exists a spanned \mathcal{S}_u -droplet $D' \subset D$ with $k \leq |D'| \leq 3k$.*

Similarly to Corollary A.11 we obtain the following.

Corollary A.20. *Let D be an \mathcal{S}_u -droplet with $d = |D| \geq 1/(C_5 q^\alpha)$. Then*

$$\mu(D \text{ is spanned}) \leq d^{O(1)} e^{-2/(C_5 q^\alpha)}.$$

Remark A.21. Let us note that the results of this section remain valid if ∂ is replaced by any sufficiently regular boundary condition. Namely, if $u_\perp = u + \pi/2$ and f is a δ -Lipschitz function for $\delta < \tan((u^+ - u^-)/2)$, then we can use any ∂ with topological interior

$$\{x \in \mathbb{R}^2, \langle x, u \rangle < f(\langle x, u_\perp \rangle)\}$$

such that $\partial, \partial \cup D$ are stable for any cut \mathcal{S}_u -droplet.

Finally, one can also remove the boundary by considering infections sufficiently far from it to recover the setting of the previous section for the directions under consideration.

B Bound on crossing

For this appendix we place ourselves in the context of Section 3 (in particular, \mathcal{T} -droplets will be parallelograms). In sections B.1 and B.2 we show that crossings are unlikely in directions with respectively finite and infinite difficulty. Of course, though we treat u_1 , the results are also valid for u_2 .

B.1 Crossing in a direction with finite difficulty

One can use Corollary A.11 to show that if u_1 has finite difficulty, a u_1 -crossing without large droplets is extremely unlikely. To do that, we will use a concept of *partition* close to the one from [3, Definition 8.20].

Definition B.1. Assume that $0 < \alpha(u_1) < \infty$. Let $R = R(a, b; c, d)$ be a parallelogram and $Z \subset R \cap \mathbb{Z}^2$. Set $m = \lfloor (c - a)/(C_1 C_6) \rfloor \geq 1$ and

$$S_i = \mathbb{H}_{u_1}(-(c - iC_1 C_6)) \cap \bar{\mathbb{H}}_{u_2}(-b) \cap \bar{\mathbb{H}}_{u_3}(c - (i - 1)C_1 C_6) \cap \bar{\mathbb{H}}_{u_4}(d)$$

for $1 \leq i \leq m - 1$ and $S_m = R(a, b; c - (m - 1)C_1 C_6, d)$. A u_1 -partition of R for Z is a sequence a_1, \dots, a_k of positive integers with $m = a_1 + \dots + a_k$ such that, setting $t_j = a_1 + \dots + a_j$, we have either

- $a_j = 1$ and S_{t_j} contains an $\alpha(u_1)$ -cluster for Z (see Definition A.1) or
- there exists a \mathcal{T} -droplet D spanned by $Z \cap \bigcup_{i=t_{j-1}+1}^{t_j} S_i$, with $C_1 C_6 a_j \geq \text{diam}(D) \geq a_j C_6$.

The following lemma is close to [3, Lemma 8.21].

Lemma B.2. *Let R be a parallelogram. If $0 < \alpha(u_1) < \infty$ and R is u_1 -crossed then there exists a u_1 -partition for $\eta \cap R$.*

Proof. For notational convenience we assume that $R = R(-a, 0; 0, d)$. In this proof, all clusters and crumbs are with respect to $\alpha(u_1)$. The proof is by induction on m .

Suppose that the property holds for any $m' \leq m - 1$. If S_1 contains a cluster of $\eta \cap R$, we set $a_1 = 1$ and we are done, since $R(-a, 0; -C_1 C_6, d)$ is u_1 -crossed. Let us assume S_1 contains no cluster of $\eta \cap R$. Then $S'_1 = R(-C_1 C_6 + C_1 C_3, 0; 0, d)$ intersects no cluster of $\eta \cap R$, so if \mathcal{K} is the set of connected components of $\eta \cap S'_1$ in Γ , each $\kappa \in \mathcal{K}$ is a crumb of $\eta \cap S'_1$. In particular, all elements of $[\kappa]$ and $[\kappa \cup \mathbb{H}_{u_1}] \setminus \mathbb{H}_{u_1}$ are at distance at most C_1 of κ (see Observation 5.16 and the proof of Corollary 5.17 in [17]). As elements of \mathcal{K} are at distance at least C_2 from one another, this means that $[\eta \cap S'_1] = \bigcup_{\kappa \in \mathcal{K}} [\kappa]$, and that $\bar{Z} = \bigcup_{\kappa \in \mathcal{K}'} [\kappa \cup \mathbb{H}_{u_1}]$ is closed, where $\mathcal{K}' = \{\kappa \in \mathcal{K} : d(\kappa, \mathbb{H}_{u_1}) \leq C_2\}$. Moreover, the diameter of a crumb is at most $\alpha(u_1)C_2$, so all elements of \bar{Z} are at distance at most $(\alpha(u_1) + 2)C_2$ of \mathbb{H}_{u_1} . Since R is u_1 -crossed, this implies that there exists $z \in \bar{Z}$ and $w \in [(\eta \cap R) \setminus \bar{Z}]$ such that $d(z, w) \leq C'_2$. Then $d(w, \mathbb{H}_{u_1}) \leq (\alpha(u_1) + 2)C_2 + C'_2$.

Let X be the connected component in Γ' of $[(\eta \cap R) \setminus \bar{Z}]$ containing w . If $X \subset [\eta \cap S'_1]$, then $X \subset \bigcup_{\kappa \in \mathcal{K}} [\kappa]$, so $X \subset [\kappa]$ for some $\kappa \in \mathcal{K}$, since they are at distance more than C'_2 from one another. Moreover, by Observation 3.3, $X = [((\eta \cap R) \setminus \bar{Z}) \cap X]$, so $X \not\subset \bar{Z}$, so $\kappa \notin \mathcal{K}'$. However, this contradicts the fact that $d(w, z) \leq C'_2$, as $d(\bar{Z}, [\kappa]) \geq C_2 - 2C_1$.

Therefore, $X \not\subset [\eta \cap S'_1]$, so X intersects $R \setminus S'_1$. Let $a_1 = \max\{i \geq 1, X \cap S_i \neq \emptyset\}$ and D be the smallest \mathcal{T} -droplet containing X . Clearly, $\text{diam}(D) \geq \text{diam}(X) \geq a_1 C_6$, since $d(w, \mathbb{H}_{u_1}) \leq C_3$. Furthermore, since

$X = [((\eta \cap R) \setminus \bar{Z}) \cap X]$, D is spanned by $\eta \cap \bigcup_{i=1}^{a_1} S_i$. We then conclude by Lemma A.9 and the induction hypothesis for $R(-a, 0; -a_1 C_1 C_6, b)$. \square

We next require a more sophisticated version of [3, Lemma 8.23].

Lemma B.3. *Fix K in Definitions 3.1 and 3.5 by*

$$K = \begin{cases} 1/(C_5 q^\alpha) & \text{if } \mathcal{U} \text{ is balanced} \\ q^{-\alpha-1/4} & \text{if } \mathcal{U} \text{ is unbalanced.} \end{cases}$$

Assume that $0 < \alpha(u_1) < \infty$. Let $R = R(a, b; c, d)$ with $d - b \leq \frac{\log(1/q)}{C_6^3 q^{\alpha(u_1)}}$ and $1/q^{C_1} \geq c - a \geq 1/q$. Then

$$\mu(C_R^{u_1}) \leq \begin{cases} \exp(-(c-a) \exp(-2C_6^2(d-b)q^{\alpha(u_1)})/C_6^2) & \text{if } \mathcal{U} \text{ is balanced} \\ \exp(-(c-a)q^{1/4}/C_6) & \text{if } \mathcal{U} \text{ is unbalanced.} \end{cases}$$

Proof. For notational convenience, we assume that $R = R(-a, 0; 0, d)$. If $C_R^{u_1}$ holds, there exists $\eta' \geq \eta$ such that R is u_1 -crossed for η' and there is no spanned critical \mathcal{T} -droplet for $\eta' \cap R$. By Lemma B.2, there exists a u_1 -partition for $\eta' \cap R$ and, by Lemma A.9, all corresponding spanned \mathcal{T} -droplets have diameter at most K/C_1 . We notice that any empty site or spanned droplet for η' is still an empty site or spanned droplet for η .

We first assume that \mathcal{U} is balanced. Given a partition \mathcal{P} we define its numbers and total sizes of big/small/cluster parts by

$$\begin{aligned} \mathcal{B} &= \{j : 1/\sqrt{q} < a_j \leq 1/(C_5 C_6 q^\alpha)\} & b &= |\mathcal{B}| & B &= \sum_{j \in \mathcal{B}} a_j \\ \mathcal{S} &= \{j : 1 < a_j \leq 1/\sqrt{q}\} & s &= |\mathcal{S}| & S &= \sum_{j \in \mathcal{S}} a_j \\ \mathcal{C} &= \{j : a_j = 1\} & c &= |\mathcal{C}|. \end{aligned}$$

We denote by $\mathcal{P}(b, s, c, B, S)$ the set of partitions \mathcal{P} with the corresponding numbers and total sizes of parts.

Then, using Corollary A.11, we get that the probability of a given \mathcal{P} occurring is at most

$$\begin{aligned} \Pi(\mathcal{P}) &= \prod_{j \in \mathcal{C}} (1 - (1 - q^{\alpha(u_1)})^{C_6^2 d}) \prod_{j \in \mathcal{S}} q^{a_j \sqrt{C_6}} \prod_{j \in \mathcal{B}} e^{-C_3 C_6 a_j} \\ &= (1 - (1 - q^{\alpha(u_1)})^{C_6^2 d})^c q^{S \sqrt{C_6}} e^{-C_3 C_6 B} \end{aligned}$$

by the union bound on all possible droplets and their positions, recalling that $d = q^{-O(1)}$. Indeed, the probability that there is no set of $\alpha(u_1)$ zeroes

connected in Γ' in a given S_i is the probability that for any possible such set C , $\eta_C \neq 0$, which, by the Harris inequality, is bigger than the product of this probability for each set C .

Assuming for simplicity that $1/\sqrt{q}$ and $1/(C_5 C_6 q^\alpha)$ are integers, we can count $\mathcal{P}(b, s, c, B, S)$ in the following way (the first binomial coefficient corresponds to the decomposition of \mathcal{B} into ordered parts, the second one to the decomposition of \mathcal{S} , and the last two to the ordering of the parts of $\mathcal{B}, \mathcal{S}, \mathcal{C}$):

$$\begin{aligned} |\mathcal{P}(b, s, c, B, S)| &\leq \binom{B - b/\sqrt{q} - 1}{b - 1} \binom{S - s - 1}{s - 1} \binom{b + s + c}{b} \binom{s + c}{s} \\ &\leq 2^{B+S} (b + s + c)^b (s + c)^s \leq e^{B+S} q^{-C_1 s}, \end{aligned}$$

recalling that $C_6(B + S + c) < a \leq 1/q^{C_1}$. Therefore, denoting by $m = \lfloor a/(C_1 C_6) \rfloor = B + S + c$ the total number of strips, we have

$$\begin{aligned} \sum_{B, S, b, s} \sum_{\mathcal{P} \in \mathcal{P}(b, s, m - B - S, B, S)} \Pi(\mathcal{P}) &\leq m^4 \max_{B, S} \left(1 - (1 - q^{\alpha(u_1)})^{C_6^2 d} \right)^{m - B - S} q^{S\sqrt{C_6}/2} e^{-C_3 C_6 B/2} \\ &\leq m^4 \max_{0 \leq c \leq m} e^{-c \exp(-2C_6^2 d q^{\alpha(u_1)})} e^{-C_2 C_6 (m - c)} \\ &\leq \exp \left(-\frac{m}{2} \exp(-2C_6^2 d q^{\alpha(u_1)}) \right), \end{aligned}$$

which concludes the proof in the balanced case, recalling the hypotheses of the lemma.

We next consider \mathcal{U} to be unbalanced. Notice that, since $K = q^{-\alpha-1/4}$, there may be droplets with diameter larger than $1/(C_5 q^\alpha)$. Therefore, we further set

$$\mathcal{H} = \{j : 1/(C_5 C_6 q^\alpha) < a_j \leq 1/(C_6 q^{\alpha+1/4})\} \quad h = |\mathcal{H}| \quad H = \sum_{j \in \mathcal{H}} a_j.$$

Then Corollary A.11 gives that the probability of a given \mathcal{P} occurring is at most

$$\Pi(\mathcal{P}) \times (q^{O(1)} e^{-2/(C_5 q^\alpha)})^h \leq \Pi(\mathcal{P}) \times \exp(-H q^{1/4} C_6 / C_5).$$

We further easily check that

$$\binom{H - h/(C_5 C_6 q^\alpha) - 1}{h - 1} \leq e^{H\sqrt{q}} \quad \binom{h + b + s + c}{h} \leq e^{H\sqrt{q}},$$

so, as above the probability of any \mathcal{P} occurring is at most

$$m^6 \exp(-m \cdot \min(C_6 q^{1/4}/(2C_5), \exp(-2C_6^2 d q^{\alpha(u_1)}))) \leq \exp\left(-\frac{C_6 m q^{1/4}}{3C_5}\right),$$

which concludes the proof. \square

B.2 Crossing in a direction with infinite difficulty

If \mathcal{U} has an infinite number of stable directions, we need to treat an infected boundary condition. This is essential, as we will work in exponentially large regions, for which the bounds from the previous section cannot be applied.

We place ourselves in the setting of Section 4.1. We will write (cut/whole) droplet for (cut/whole) \mathcal{S}_{u_1} -droplet in the sense of Definition A.13, with u_1^- and u_1^+ sufficiently close to u_1 . These should not be confused with \mathcal{T} -droplets, which are called parallelograms to avoid any confusion.

We will seek to apply Corollary A.20 rather than A.11 to prove the following.

Lemma B.4. *Fix*

$$K = \begin{cases} q^{-\alpha} & \text{if } \mathcal{U} \text{ is balanced} \\ q^{-\alpha-1/4} & \text{if } \mathcal{U} \text{ is unbalanced} \end{cases}$$

for Definitions 3.1 and 3.5. Let $R = R(a, b; c, d)$ with $C_1 \leq d - b \leq \exp(q^{-3\alpha})$ and $c - a \geq q^{-4\alpha}$. Then

$$\mu(C_R^{u_1}) \leq \exp(-q^{-3\alpha}).$$

Our strategy is as follows. Instead of considering u_1 -partitions, we directly retrace the spanning algorithm to obtain a hierarchy of droplets reaching a cut droplet of size roughly $c - a$. We reassure the reader familiar with [21] that our hierarchies will be very simple and imprecise, as the a priori hypothesis that there are no critical parallelograms removes the metastability (it is no longer easy for large droplets to grow) together with the need of fine tuning. Namely, their seeds will be of size roughly K which will also be the increment of the size of unary vertices (the reader unfamiliar with hierarchies is invited to consult the definitions below). The lack of critical parallelograms entails that all droplets in the hierarchy are cut (so they are simply very flat triangles). The bound on the probability of seeds being spanned is provided by Corollary A.20 and entropy is easily subdominant, so we can focus on the probability that the infections around a cut droplet are such that if that droplet is infected, the infection can expand to fill a slightly larger cut droplet. However, this would imply that there is a (smaller scale) u_1^\pm -crossing from the side of the smaller one to side of the larger one (see Figure 3b). The probability of this event is again bounded directly by Corollary A.20, taking into account Remark A.21.

Let us begin by introducing our hierarchies following Holroyd [21]. Let $T = q^{-\alpha-1/4}$. Fix a droplet D . A *hierarchy* \mathcal{H} for D is a rooted unary-binary

tree with each vertex x labelled by a droplet $D_x \subset D$, so that the label of the root is D . We denote by $N(x)$ the set of children of $x \in V(\mathcal{H})$, so that $|N(x)| \in \{0, 1, 2\}$ for all x . The leaves are called *seeds* and the binary vertices are called *splitters*. We alert the reader that in reality there will only be cut droplets in our hierarchies, but for technical reasons we define them in general. A hierarchy is defined to satisfy the following conditions.

- If $y \in N(x)$, then $D_y \subset D_x$.
- If D_x is a whole droplet, then x is a seed and $T/3 \leq |D_x|$.
- If D_x is a cut droplet, then $T/3 \leq |D_x| \leq T$ if and only if x is a seed.
- If $N(x) = \{y\}$ and $|N(y)| = 1$, then $T < |D_x| - |D_y| \leq 2T$.
- If $N(x) = \{y\}$, then $|D_x| - |D_y| \leq 2T$.
- If $N(x) = \{y, z\}$, then $|D_x| - |D_y| > T$ and $\langle D_y \cup D_z \rangle = \{D_x\}$.

We set

$$\begin{aligned} S(\mathcal{H}) &= \{x \in V(\mathcal{H}) : |N(x)| = 0\} \\ N(\mathcal{H}) &= \{(x, y) \in (V(\mathcal{H}))^2 : N(x) = \{y\}, |N(y)| = 1\} \end{aligned}$$

and remark that $|S(\mathcal{H})| - 1$ is the number of splitters. We say that a hierarchy \mathcal{H} *occurs* if the following events occur disjointly (are witnessed by disjoint sets of infections, see [31]).

- For every seed $x \in S(\mathcal{H})$ we have that D_x is spanned.
- For every $x \in V(\mathcal{H})$ such that $N(x) = \{y\}$ we have $D_x \in \langle D_y \cup (\eta \cap D_x) \rangle$.

Lemma B.5. *If D is a spanned droplet with $|D| \geq T/3$, then some hierarchy for D occurs.*

Proof. The proof is very similar e.g. to [3, Lemma 8.7]. Assuming that D_0 is a spanned droplet with $|D_0| \geq T/3$, we construct an occurring hierarchy by induction on D_0 with respect to inclusion. If $|D_0| \leq T$ or D_0 is a whole droplet, the hierarchy with only vertex labelled by D_0 is as desired.

Assume that D_0 is a cut droplet and $|D_0| > T$. Let Z be a connected component for Γ' of $[D_0 \cap \eta]_\partial$ such that the smallest droplet containing Z is D_0 , and let $Z_0 = Z \cap \eta$. We then have $[Z_0]_\partial = Z$. By Lemma A.17 there exist sequences Z_1, \dots, Z_m and Z'_1, \dots, Z'_m of subsets of Z_0 and D_1, \dots, D_m and D'_1, \dots, D'_m such that the following conditions hold for all $0 < i \leq m$.

- $Z_{i-1} = Z_i \sqcup Z'_i$, $[Z_i]_{\partial}$, $[Z'_i]_{\partial}$ and $[Z_i]_{\partial} \cup [Z'_i]_{\partial}$ are connected in Γ' .
- $D_i = D([Z_i]_{\partial})$ and $D'_i = D([Z'_i]_{\partial})$.
- $\langle D_i \cup D'_i \rangle = \{D_{i-1}\}$ and $|D_i| \geq |D'_i|$
- $m \geq 1$ is the minimal index such that one of the following holds:
 1. $|D_0| - |D_m| > T$;
 2. D_m is a whole droplet;
 3. $|D_m| \leq T$.

If 1 does not hold, then we attach a seed labelled by D_m to the root and we are done, as $|D_m| \geq |D_{m-1}|/3 > T/3$ by Lemma A.18 and minimality of m . Indeed, D_m being spanned is witnessed by Z_m , while $D_0 \in \langle D_m \cup (\eta \cap D_0) \rangle$ is witnessed by $Z_0 \setminus Z_m$.

Assume that 1 holds. Then we consider two cases. If $T < |D_0| - |D_m| \leq 2T$, we attach a hierarchy for D_m (occurring for Z_m) to the root D_0 and we are done using Lemma A.18 to get that $|D_m| > T/3$ as above. Otherwise we attach a splitter labelled by D_{m-1} to the root D_0 (if $m = 1$, then D_0 is the splitter) and hierarchies for D_m and D'_m to that splitter. Then we are done, recalling Lemma A.18, to get that $|D_m| \geq |D'_m| \geq |D_{m-1}| - |D_m| - O(C'_2) \geq T - O(C'_2)$. \square

In order to bound the probability that a hierarchy occurs, we will need the following result.

Lemma B.6. *Let $D_1 \subset D_2$ be two cut droplets for $\partial = \mathbb{H}_{u_1}$ such that $T < |D_2| - |D_1| \leq 2T$ and $|D_2| \leq q^{-4\alpha}$. Then*

$$\mu(D_2 \in \langle D_1 \cup (\eta \cap D_2) \rangle) \leq e^{-q^{-\alpha}/C_5}.$$

Proof. The proof is illustrated in Figure 3. Let us denote $D_i = (\bar{\mathbb{H}}_{u_1^+}(x_i) \cap \bar{\mathbb{H}}_{u_1^-}(y_i)) \setminus \mathbb{H}_{u_1}$ for $i \in \{1, 2\}$. Define the strips $X = \bar{\mathbb{H}}_{u_1^+}(x_2) \setminus \bar{\mathbb{H}}_{u_1^+}(x_1)$ and $Y = \bar{\mathbb{H}}_{u_1^-}(y_2) \setminus \bar{\mathbb{H}}_{u_1^-}(y_1)$ and assume without loss of generality that $y_2 - y_1 = \Omega(T)$. Assume that $D_2 \in \langle D_1 \cup (\eta \cap D_2) \rangle$ occurs. Setting $\eta' = \eta \cap Y \cap D_2$, this implies $D_2 \in \langle (D_2 \setminus Y) \cup \eta' \rangle$. We consider two cases.

Assume that $D_2 \in \langle \eta' \rangle$. By Corollary A.20 the probability of this event is at most $q^{-O(1)} e^{-2/(C_5 q^\alpha)}$.

Assume that, on the contrary, $D_2 \notin \langle \eta' \rangle$ and set $\partial' = \mathbb{H}_{u_1} \cup \bar{\mathbb{H}}_{u_1^-}(y_1)$. Then by Observation 3.3 there exists a set $C \subset [D_2 \cap \eta]_{\partial'}$ connected in Γ' such that $d(C, \mathbb{H}_{u_1^-}(y_1)) \leq C'_2$ and $C \not\subset \mathbb{H}_{u_1^-}(y_2)$. By definition this implies the




Figure 3: Illustration of the proof of Lemma B.6 bounding the probability that the infections around a cut droplet, D_1 , allow an infection filling D_1 to grow and fill the slightly larger cut droplet, D_2 .

existence of a cut $\mathcal{S}_{u_1^-}$ -droplet spanned by $D_2 \cap \eta$ with boundary ∂' , where the two directions of cut droplets in $\mathcal{S}_{u_1^-}$ are u_1^+ and $u_1^{--} = 2u_1^- - u_1^+$ (recall Remark A.21). Hence, by the union bound over all possible such droplets and Corollary A.20 we obtain the desired result. \square

We are now ready to assemble the proof of Lemma B.4 as outlined at the beginning of the section.

Proof of Lemma B.4. Assume that $C_R^{u_1}$ occurs and let $\eta' \geq \eta$ be as in Definition 3.5. Then there exists a spanned cut droplet for $\eta' \cap R$ with boundary \mathbb{H}_{u_1} of diameter at least $c - a$. By Lemma A.19 this implies the existence of a droplet D spanned for $\eta' \cap R$ with $q^{-4\alpha}/C_1 \leq |D| \leq 3q^{-4\alpha}/C_1$. We set $Z = \eta' \cap R \cap D$.

Let us assume for a contradiction that there exists a whole droplet of size at least $q^{-\alpha-1/4}/3$ spanned for Z . It is easy to check that there exists a parallelogram of diameter at least $q^{-\alpha-1/4}/C_1$ spanned by Z (consider a connected component satisfying Definition A.13, take the smallest parallelogram containing it and use Observation 3.3). By Lemma 3.4 this contradicts the absence of spanned critical parallelograms for $\eta' \cap R$.

Therefore, D is a cut droplet and by Lemma B.5 there exists a hierarchy for D occurring for the zero set Z , whose labels are all cut droplets. Let

$\mathcal{H}(D)$ denote the set of such hierarchies. Now, by the BK inequality [31], for any hierarchy \mathcal{H} we have the following analogue of [21, Equation (37)]:

$$\mu(\mathcal{H} \text{ occurs}) \leq \prod_{x \in S(\mathcal{H})} \mu(D_x \text{ is spanned}) \prod_{(x,y) \in N(\mathcal{H})} \mu(D_x \in \langle D_y \cup (\eta \cap D_x) \rangle).$$

Thanks to Corollary A.20 and Lemma B.6, we deduce

$$\mu(C_R^{u_1}) \leq \sum_D \sum_{\mathcal{H} \in \mathcal{H}(D)} \exp(-q^{-\alpha}(|S(\mathcal{H})| + |N(\mathcal{H})|)/C_5).$$

The number of choices for D is $O((d-b)q^{-4\alpha})$. We separate the sum over hierarchies according to their number of vertices $v(\mathcal{H}) = \Theta(|S(\mathcal{H})| + |N(\mathcal{H})|)$. By Lemma A.18 we have that $v(\mathcal{H}) = \Omega(|D|/T) = \Omega(q^{-3\alpha+1/4}/C_1)$. Finally, the number of hierarchies for a given cut droplet D with v vertices is at most $q^{-O(v)}$. Combining these bounds we have

$$\mu(C_R^{u_1}) \leq (d-b) \sum_{v=\Omega(q^{-3\alpha+1/4}/C_1)} \exp(-q^{-\alpha}\Omega(v)/C_5),$$

which concludes the proof. \square

References

- [1] M. Aizenman and J. L. Lebowitz, *Metastability effects in bootstrap percolation*, J. Phys. A **21** (1988), no. 19, 3801–3813. MR968311
- [2] P. Balister, B. Bollobás, M. Przykucki, and P. Smith, *Subcritical \mathcal{U} -bootstrap percolation models have non-trivial phase transitions*, Trans. Amer. Math. Soc. **368** (2016), no. 10, 7385–7411. MR3471095
- [3] B. Bollobás, H. Duminil-Copin, R. Morris, and P. Smith, *Universality of two-dimensional critical cellular automata*, Proc. Lond. Math. Soc. (To appear).
- [4] B. Bollobás, P. Smith, and A. Uzzell, *Monotone cellular automata in a random environment*, Combin. Probab. Comput. **24** (2015), no. 4, 687–722. MR3350030
- [5] N. Cancrini, F. Martinelli, C. Roberto, and C. Toninelli, *Kinetically constrained spin models*, Probab. Theory Related Fields **140** (2008), no. 3–4, 459–504. MR2365481
- [6] R. Cerf and E. N. M. Cirillo, *Finite size scaling in three-dimensional bootstrap percolation*, Ann. Probab. **27** (1999), no. 4, 1837–1850. MR1742890
- [7] F. Chung, P. Diaconis, and R. Graham, *Combinatorics for the East model*, Adv. Appl. Math. **27** (2001), no. 1, 192–206. MR1835679
- [8] P. de Gregorio, A. Lawlor, and K. A. Dawson, *Bootstrap percolation*, Encyclopedia of complexity and systems science, 2009, pp. 608–626.

- [9] H. Duminil-Copin and A. Holroyd, *Finite volume bootstrap percolation with balanced threshold rules on \mathbb{Z}^2* (2012). Preprint available at <http://www.ihes.fr/~duminil/>.
- [10] H. Duminil-Copin and A. C. D. van Enter, *Sharp metastability threshold for an anisotropic bootstrap percolation model*, Ann. Probab. **41** (2013), no. 3A, 1218–1242. MR3098677
- [11] A. Faggionato, F. Martinelli, C. Roberto, and C. Toninelli, *The East model: recent results and new progresses*, Markov Process. Related Fields **19** (2013), no. 3, 407–452. MR3156959
- [12] G. H. Fredrickson and H. C. Andersen, *Kinetic Ising model of the glass transition*, Phys. Rev. Lett. **53** (1984), no. 13, 1244–1247.
- [13] G. H. Fredrickson and H. C. Andersen, *Facilitated kinetic Ising models and the glass transition*, J. Chem. Phys. **83** (1985), no. 11, 5822–5831.
- [14] P. Garrahan, P. Sollich, and C. Toninelli, *Kinetically constrained models*, Dynamical heterogeneities in Glasses, colloids and granular media and jamming transitions, 2011, pp. 341–369.
- [15] I. Hartarsky, *Refined universality for critical KCM: upper bounds* (In preparation).
- [16] I. Hartarsky, *\mathcal{U} -bootstrap percolation: critical probability, exponential decay and applications*, Ann. Inst. Henri Poincaré Probab. Stat. (To appear).
- [17] I. Hartarsky, L. Marêché, and C. Toninelli, *Universality for critical KCM: infinite number of stable directions*, Probab. Theory Related Fields **178** (2020), no. 1, 289–326. MR4146539
- [18] I. Hartarsky, F. Martinelli, and C. Toninelli, *FA-2f kinetically constrained model: sharp results* (In preparation).
- [19] I. Hartarsky, F. Martinelli, and C. Toninelli, *Universality for critical KCM: finite number of stable directions*, Ann. Probab. (To appear).
- [20] I. Hartarsky and R. Morris, *The second term for two-neighbour bootstrap percolation in two dimensions*, Trans. Amer. Math. Soc. **372** (2019), no. 9, 6465–6505. MR4024528
- [21] A. E. Holroyd, *Sharp metastability threshold for two-dimensional bootstrap percolation*, Probab. Theory Related Fields **125** (2003), no. 2, 195–224. MR1961342
- [22] T. M. Liggett, *Interacting particle systems*, Classics in mathematics, Springer Berlin, Berlin, Heidelberg, 2005. Reprint of the 1985 original. MR2108619
- [23] L. Marêché, *Combinatorics for general kinetically constrained spin models*, SIAM J. Discrete Math. **34** (2020), no. 1, 370–384. MR4062795
- [24] L. Marêché, F. Martinelli, and C. Toninelli, *Exact asymptotics for Duarte and supercritical rooted kinetically constrained models*, Ann. Probab. **48** (2020), no. 1, 317–342. MR4079438
- [25] F. Martinelli, R. Morris, and C. Toninelli, *Universality results for kinetically constrained spin models in two dimensions*, Comm. Math. Phys. **369** (2019), no. 2, 761–809. MR3962008
- [26] F. Martinelli and C. Toninelli, *Towards a universality picture for the relaxation to equilibrium of kinetically constrained models*, Ann. Probab. **47** (2019), no. 1, 324–361. MR3909971

- [27] R. Morris, *Bootstrap percolation, and other automata*, European J. Combin. **66** (2017), 250–263. MR3692148
- [28] R. Morris, *Monotone cellular automata*, Surveys in combinatorics 2017, 2017, pp. 312–371. MR3728111
- [29] F. Ritort and P. Sollich, *Glassy dynamics of kinetically constrained models*, Adv. Phys. **52** (2003), no. 4, 219–342.
- [30] P. Sollich and M. R. Evans, *Glassy time-scale divergence and anomalous coarsening in a kinetically constrained spin chain*, Phys. Rev. Lett. **83** (1999), no. 16, 3238–3241.
- [31] J. van den Berg and H. Kesten, *Inequalities with applications to percolation and reliability*, J. Appl. Probab. **22** (1985), no. 3, 556–569. MR799280