

Folding-model approach to reaction cross section of $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at 790 MeV

Shingo Tagami

Department of Physics, Kyushu University, Fukuoka 819-0395, Japan

Tomotsugu Wakasa

Department of Physics, Kyushu University, Fukuoka 819-0395, Japan

Maya Takechi

Niigata University, Niigata 950-2181, Japan

Jun Matsui

Department of Physics, Kyushu University, Fukuoka 819-0395, Japan

Masanobu Yahiro*

Department of Physics, Kyushu University, Fukuoka 819-0395, Japan

(Dated: December 15, 2021)

Tanihata *et al.* determined matter radii $r_m(\sigma_I)$ for $^{4,6,8}\text{He}$ from interaction cross sections σ_I of $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at 790 MeV per nucleon, using the optical limit of the Glauber model. Lu *et al.* determined proton radii $r_p(\text{AIS})$ for $^{4,6,8}\text{He}$ with the atomic isotope shifts (AIS). We investigate whether the Love-Franey t -matrix folding model is good for $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at 790 MeV per nucleon.

I. INTRODUCTION AND CONCLUSION

Background: The matter radius r_m , the neutron skin r_{skin} and halo structure are important properties of nuclei. When a nucleus has one or more loosely-bound nucleons surrounding a tightly bound core, it is considered that the nucleus has a halo structure. Eventually, we may consider that $^{6,8}\text{He}$ have the halo structure.

Lu *et al.* measured the atomic isotope shifts (AIS) along $^{4,6,8}\text{He}$ by performing laser spectroscopy on individual trapped atoms and determined proton radii $r_p(\text{AIS}) = 1.462(6), 1.934(9), 1.881(17)$ fm for $^{4,6,8}\text{He}$ [1].

For He isotopes, Tanihata *et al.* determined r_m from interaction cross sections σ_I [2, 3]; $r_m(\sigma_I) = 1.57(4), 2.48(3), 2.52(3)$ fm for $^{4,6,8}\text{He}$ by assuming the harmonic-oscillator distribution and using the optical limit of the Glauber model [4]; see Eq. (4.8) of Ref. [3]. The Glauber model is constructed with the adiabatic and the eikonal approximation in the frame work of many-body Schrödinger equation [4, 5].

In Ref. [3], the experimental values of r_m are accumulated from ^4He to ^{32}Mg . For $^{4,6,8}\text{He}+^{12}\text{C}$ and $^{12}\text{C}+^{12}\text{C}$ scattering, the data on interaction cross section $\sigma_I(\text{exp})$ are presented at 790 MeV with high accuracy of 1% [3].

Aim: We investigate reliability of the folding model based on Love-Franey (LF) t -matrix [6] for $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at 790 MeV per nucleon.

Results: First, we apply the LF t -matrix folding model for $^{12}\text{C}+^{12}\text{C}$ scattering at $E_{\text{lab}} = 790$ MeV per nucleon, and determine the ratio $F = \sigma_R(\text{LF})/\sigma_I(\text{exp})$ at $E_{\text{lab}} = 790$ MeV per nucleon. The resulting value is $F = 0.9356$. Secondly, for $^{6,8}\text{He}$, we deduce neutron radius $r_n(\sigma_I) = 2.71, 2.70$ fm

from the $r_m(\sigma_I) = 2.48, 2.52$ fm and the $r_p(\text{AIS}) = 1.934, 1.881$ fm. For ^4He , we assume $r_n(\text{AIS}) = r_p(\text{AIS})$, leading to $r_m(\text{AIS}) = r_n(\text{AIS}) = r_p(\text{AIS})$. Thirdly, we use Gogny-D1S HFB (GHFB), and scale the proton and neutron GHFB densities so as to satisfy $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\text{AIS})$ for ^4He and $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\sigma_I)$ for $^{6,8}\text{He}$. For $^{4,6,8}\text{He}+^{12}\text{C}$ scattering, the reaction cross section $\sigma_R(\text{scaling})$ calculated with the scaled densities reproduce the $\sigma_I(\text{exp})$.

Conclusion: Our conclusion is that the Love-Franey (LF) t -matrix folding model is good for $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at 790 MeV per nucleon.

II. MODEL

In the t -matrix folding model, the potential U consists of the direct part (U^{DR}) and the exchange part (U^{EX}) defined by [7, 8]

$$U^{\text{DR}}(\mathbf{R}) = \sum_{\mu, \nu} \rho_{\text{P}}^{\mu}(\mathbf{r}_{\text{P}}) \rho_{\text{T}}^{\nu}(\mathbf{r}_{\text{T}}) t_{\mu\nu}^{\text{DR}}(s) d\mathbf{r}_{\text{P}} d\mathbf{r}_{\text{T}}, \quad (1)$$

$$U^{\text{EX}}(\mathbf{R}) = \sum_{\mu, \nu} \rho_{\text{P}}^{\mu}(\mathbf{r}_{\text{P}}, \mathbf{r}_{\text{P}} - \mathbf{s}) \rho_{\text{T}}^{\nu}(\mathbf{r}_{\text{T}}, \mathbf{r}_{\text{T}} + \mathbf{s}) \times t_{\mu\nu}^{\text{EX}}(s) \exp[-i\mathbf{K}(\mathbf{R}) \cdot \mathbf{s}/M] d\mathbf{r}_{\text{P}} d\mathbf{r}_{\text{T}}, \quad (2)$$

where $\mathbf{s} = \mathbf{r}_{\text{P}} - \mathbf{r}_{\text{T}} + \mathbf{R}$ for the coordinate \mathbf{R} between a projectile (P) and a target (T). The coordinate \mathbf{r}_{P} (\mathbf{r}_{T}) denotes the location for the interacting nucleon measured from the center-of-mass of P (T). Each of μ and ν stands for the z -component of isospin; $1/2$ means neutron and $-1/2$ does proton. The original form of U^{EX} is a non-local function of \mathbf{R} , but it has been localized in Eq. (2) with the local semi-classical approximation [9] in which P is assumed to propagate as a plane wave with the local momentum $\hbar\mathbf{K}(\mathbf{R})$ within a short range of the

* orion093g@gmail.com

nucleon-nucleon interaction, where $M = AA_T/(A + A_T)$ for the mass number A (A_T) of P (T). The validity of this localization is shown in Ref. [10].

The direct and exchange parts, $t_{\mu\nu}^{\text{DR}}$ and $t_{\mu\nu}^{\text{EX}}$, of the t matrix are described by

$$t_{\mu\nu}^{\text{DR}}(s) = \frac{1}{4} \sum_S \hat{S}^2 t_{\mu\nu}^{S1}(s) \quad \text{for } \mu + \nu = \pm 1, \quad (3)$$

$$t_{\mu\nu}^{\text{DR}}(s) = \frac{1}{8} \sum_{S,T} \hat{S}^2 t_{\mu\nu}^{ST}(s) \quad \text{for } \mu + \nu = 0, \quad (4)$$

$$t_{\mu\nu}^{\text{EX}}(s) = \frac{1}{4} \sum_S (-1)^{S+1} \hat{S}^2 t_{\mu\nu}^{S1}(s) \quad \text{for } \mu + \nu = \pm 1, \quad (5)$$

$$t_{\mu\nu}^{\text{EX}}(s) = \frac{1}{8} \sum_{S,T} (-1)^{S+T} \hat{S}^2 t_{\mu\nu}^{ST}(s) \quad \text{for } \mu + \nu = 0, \quad (6)$$

where $\hat{S} = \sqrt{2S + 1}$ and $t_{\mu\nu}^{ST}$ are the spin-isospin components of the t -matrix interaction. As for the t -matrix interaction, we take the LF version. We apply the LF t -matrix folding model for $^{12}\text{C}+^{12}\text{C}$ and $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at $E_{\text{lab}} = 790$ MeV per nucleon.

As proton and neutron densities, we use Gogny D1S HFBB (GHFB) and scale the GHFB densities; see the center-of-mass (cm) corrections for Ref. [12]. In order to make the scaling, for $^{6,8}\text{He}$, we determine $r_n(\sigma_I) = 2.71, 2.70$ fm from the $r_m(\sigma_I) = 2.48, 2.52$ fm and the $r_p(\text{AIS}) = 1.934, 1.881$ fm. For ^4He , we assume $r_n(\text{AIS}) = r_p(\text{AIS})$, leading to $r_m(\text{AIS}) = r_n(\text{AIS}) = r_p(\text{AIS})$. Next, we scale the GHFB proton and neutron densities so as to $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\sigma_I)$ for $^{6,8}\text{He}$ and $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\text{AIS})$ for ^4He . The scaled densities only are used in the folding model.

Also for ^{12}C , we construct the densities with GHFB with the angular momentum projection and scale the densities so as to $r_m(\sigma_I) = 2.35$ fm [2] and the $r_p(\text{exp}) = 2.33$ fm [13] of electron scattering.

III. RESULTS

Firstly, we apply the LF t -matrix folding model for $^{12}\text{C}+^{12}\text{C}$ scattering at $E_{\text{lab}} = 790$ MeV per nucleon, and

determine the ratio $F = \sigma_R(\text{LF})/\sigma_I(\text{exp})$ as $F = 0.9356$, where the $\sigma_I(\text{exp}) = 856$ mb was measured by Tanihata *et al.* [3].

Figure 1 shows reaction cross sections σ_R as a function of mass number A for $^{4,6,8}\text{He}+^{12}\text{C}+^{4,6,8}\text{He}$ scattering at 790 MeV. The reaction cross sections $F\sigma_R(\text{scaling})$ obtained with the scaled densities reproduce $\sigma_I(\text{exp})$ [3]; note that the data have high accuracy of 1 %. This success indicates that the LF t -matrix folding model is good for $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at 790 MeV per nucleon.

We determine $r_m(\text{exp})$ from the $\sigma_I(\text{exp})$; the results are $r_m(\text{exp}) = 1.462(6), 2.48(3), 2.53(3)$ fm for $^{4,6,8}\text{He}$. Our results are close to $r_m(\sigma_I) = 1.57(4)$ fm [2] for ^4He and agree with $r_m(\sigma_I) = 2.48(3), 2.52(3)$ fm [2] for $^{6,8}\text{He}$.




FIG. 1. Reaction cross sections σ_R for $^{4,6,8}\text{He}+^{12}\text{C}$ scattering at 790 MeV. Closed circles denote results $F\sigma_R(\text{scaling})$. The data (crosses) on σ_I are taken from Ref. [3].

ACKNOWLEDGMENTS

We would like to thank Dr. Toyokawa for providing his code.

[1] Z. T. Lu, P. Mueller, G. W. F. Drake, W. Noerthershaeuser, S. C. Pieper and Z. C. Yan, *Rev. Mod. Phys.* **85**, no.4, 1383-1400 (2013) doi:10.1103/RevModPhys.85.1383 [arXiv:1307.2872 [nucl-ex]].
[2] I. Tanihata, T. Kobayashi, O. Yamakawa, S. Shimoura, K. Ekuni, K. Sugimoto, N. Takahashi, T. Shimoda and H. Sato, *Phys. Lett. B* **206**, 592-596 (1988).
[3] A. Ozawa, T. Suzuki and I. Tanihata, *Nucl. Phys. A* **693**, 32-62 (2001).
[4] R.J. Glauber, *in Lectures in Theoretical Physics* (Interscience, New York, 1959), Vol. 1, p.315.
[5] M. Yahiro, K. Minomo, K. Ogata, and M. Kawai, *Prog. Theor.*

Phys. **120**, 767 (2008).
[6] W.G. Love and M.A. Franey, *Phys. Rev. C* **24**, 1073 (1981).
M.A. Franey and W.G. Love, *Phys. Rev. C* **31**, 488 (1985).
[7] B. Sinha, *Phys. Rep.* **20**, 1 (1975).
B. Sinha and S. A. Moszkowski, *Phys. Lett.* **B81**, 289 (1979).
[8] T. Furumoto, Y. Sakuragi, and Y. Yamamoto, *Phys. Rev.* **C82**, 044612 (2010).
[9] F.A. Brieva and J.R. Rook, *Nucl. Phys.* **A291**, 299 (1977); *ibid.* **291**, 317 (1977); *ibid.* **297**, 206 (1978).
[10] K. Minomo, K. Ogata, M. Kohno, Y. R. Shimizu, and M. Yahiro, *J. Phys. G* **37**, 085011 (2010) [arXiv:0911.1184 [nucl-th]].

- [11] K. Egashira, K. Minomo, M. Toyokawa, T. Matsumoto and M. Yahiro, Phys. Rev. C **89**, no. 6, 064611 (2014) doi:10.1103/PhysRevC.89.064611 [arXiv:1404.2735 [nucl-th]].
- [12] T. Sumi, K. Minomo, S. Tagami, M. Kimura, T. Matsumoto, K. Ogata, Y. R. Shimizu and M. Yahiro, Phys. Rev. C **85**, 064613 (2012), [arXiv:1201.2497 [nucl-th]].
- [13] I. Angeli and K. P. Marinova, Atom. Data Nucl. Data Tabl. **99**, 69 (2013).
- [14] E. Hiyama *et al.*, Physical Review C **53**, 2075 (1996).