

On the quasiconvex hull for a three-well problem in two dimensional linear elasticity

Capella, Antonio¹ and Morales, Lauro²

¹Instituto de Matemáticas-UNAM , lmm@ciencias.unam.mx

²Instituto de Matemáticas-UNAM, capella@im.unam.mx

December 2020

Abstract

We provide quantitative inner and outer bounds for the symmetric quasiconvex hull $Q^e(\mathcal{U})$ on linear strains generated by three-well sets \mathcal{U} in $\mathbb{R}_{sym}^{2 \times 2}$. In our study, we consider all possible compatible configurations for three wells and prove that if there exist two matrices in \mathcal{U} that are rank-one compatible then $Q^e(\mathcal{U})$ coincides with its symmetric lamination convex hull $L^e(\mathcal{U})$. We complete this result by providing an explicit characterization of $L^e(\mathcal{U})$ in terms of the wells in \mathcal{U} . Finally, we discuss the optimality of our outer bound and its relationship with quadratic polyconvex functions.

1 Introduction

In this work, we are concerned with variational problems of the form

$$\min_{y=y_0 \text{ on } \partial\Omega} I(y), \quad \text{where} \quad I(y) = \int_{\Omega} f(\nabla y(x)) dx, \quad (1)$$

the function $y : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$, and $f : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is continuous, bounded from below, and satisfies mild growth conditions (see, e.g. section 9 in [10]). For integral functionals like (1), other properties of f besides continuity and coercivity are needed to assure the sequential weak lower semicontinuity (swlsc) of I , *i.e.*, the weak convergence of minimizing sequences to minimizers, see for example theorems 1.13 and 1.15 in [10] pp. 15 and 17, and section 3 in [3]. The appropriate additional condition is some kind of convexity of f . In the 1950s, Morrey, see theorems 2.1 and 2.2 in [16] pp. 5 and 8, respectively, introduced quasiconvexity as the necessary and sufficient condition for swlsc (see also theorem 3.4 and remark 3.3 in [3]). A locally Borel-measurable function f is *quasiconvex* if for every $M \in \mathbb{R}^{n \times n}$ and each smooth function $\phi : \Omega \rightarrow \mathbb{R}$ compactly supported in Ω ,

$$f(M) \leq \frac{1}{|\Omega|} \int_{\Omega} f(M + \nabla \phi) dx.$$

In materials science, energy models like (1) arise in the study of martensitic phase transformations. Such models satisfy frame indifference and multi-well structure assumptions. By multi-well structure, we mean that $f \geq 0$ everywhere but $f = 0$ only on a prescribed set K . The connected components of K are the energy wells that represent the material's different phases. In most cases, these energy densities are not quasiconvex under reasonable conditions, see [2] p. 401, and [13] p. 1. Thus, minimizing sequences may develop oscillations on their gradient, and only weak convergence

is to be expected. In these models, oscillations correspond to the microstructures observed in real materials, see [1] p. 14. There are three types of sets K relevant to models of martensitic materials, depending on f 's symmetry. First, f has no symmetry and $K \subset \mathbb{R}^{n \times n}$ is a set of *matrices*. Second, f is left-invariant by orthogonal matrices, i.e., $f(OM) = f(M)$ for every $M \in \mathbb{R}^{n \times n}$ and $O \in SO(n)$. In this case, $K = \{SO(n)M_1, \dots, SO(n)M_k\}$, where $SO(n)M = \{QM \mid Q \in SO(n)\}$ is the elastic well associated with the symmetric positive-defined matrix M . Third, f is invariant under the addition of skew-symmetric matrices. In this case, $K = \{M_1 \oplus \mathbb{R}_{skew}^{n \times n}, \dots, M_k \oplus \mathbb{R}_{skew}^{n \times n}\}$ is a set of *linearized elastic wells* or *linear strains*, where $M \oplus \mathbb{R}_{skew}^{n \times n} := \{M + W \mid W \in \mathbb{R}_{skew}^{n \times n}\}$ is the linearized elastic well of M . We also say that M represents the well $M \oplus \mathbb{R}_{skew}^{n \times n}$. Elements in K can be used to construct minimizing sequences in terms of sequential lamination provided they are compatible, see for example [13] pp. 219 – 220 and theorem 3 in [2] pp. 25 – 26. As in [7] p. 846, we say that two *matrices* $M_1, M_2 \in \mathbb{R}^{n \times n}$, are compatible if they are *rank-one connected*, i.e., $\text{Rank}(M_1 - M_2) \leq 1$. In the opposite case they are called *incompatible*. Two elastic wells $SO(n)M_1$ and $SO(n)M_2$, are *compatible* if there exists $Q_1, Q_2 \in SO(n)$ such that Q_1M_1 and Q_2M_2 are rank-one connected. They are called incompatible otherwise. The corresponding notion of compatibility in linear elasticity is that: two linear strains $M_1 \oplus \mathbb{R}_{sym}^{n \times n}$ and $M_2 \oplus \mathbb{R}_{sym}^{n \times n}$ are compatible if there exists a skew symmetric matrix W such that $\text{Rank}(M_1 - M_2 + W) \leq 1$. In the particular case where $M_2 = 0$, we simply say that M_1 is compatible or incompatible if $\det M_1 \leq 0$ or $\det M_1 > 0$, respectively. As before, they are called incompatible otherwise. Moreover, we say that two matrices $M_1, M_2 \in \mathbb{R}_{sym}^{2 \times 2}$ which represent linear strains are *rank-one compatible* if $\det(M_1 - M_2) = 0$. Moreover, if M_1 and M_2 represent two compatible linear strains, then $M_1 - M_2 + W = a \otimes n$ for some $W \in \mathbb{R}_{skew}^{2 \times 2}$, $a \in \mathbb{R}^2$, and $n \in \mathbb{S}^1$.

If two matrices, elastic wells or linear strains, are compatible, layering structures can be constructed and nontrivial oscillating minimizing sequences emerge in suitable spaces, see for example proposition 2 in [1] p. 23, and section 3 in [7] pp. 855 to 868. Young measures are an effective tool to capture the nonlinear functional's asymptotic behavior along an oscillating sequence. While the weak limit carries information on average values in an oscillating sequence, the Young measure contains information on “where” these oscillations occur. If the energy achieves its minimum, the minimizing sequence does not oscillate, and the corresponding Young measure is trivial; namely, it is a Dirac mass for almost every point in the domain. For one-well and two-incompatible-wells problems, Kinderlehrer [12] (pp. 15 and 16) showed that the Young measure limit of gradients (case of elastic wells above) is trivial and furthermore constant in the domain. The case for two linear strains was studied by Kohn in [13]. Bhattacharya et al. [7], studied the problem of which measures arise as Young measures limits of gradients, linear strains, and matrices. In the case of k pairwise incompatible elastic wells or linear strains in two dimensions (see also Lemma 3 in [22], p. 408), they show that the corresponding Young measures' limits are trivial. The situation changes as soon as we consider four matrices in two space dimensions, see remark 6 in [24] pp. 194 and 195, three elastic wells, or three linear strains in three dimensions. The authors in [7] showed that in the later three cases there exist nontrivial Young measures in the limit under the assumption of pairwise incompatibility. They called this phenomenon “mutual compatibility”, see [7] p. 849.

Depending on the space dimension and the number of compatible –or incompatible– pairs among the matrices, wells, or linear strains in K , we may have the existence of nontrivial Young measures in the limit. A closely related problem is to identify the set of values taken by the (constant) weak limits of minimizing sequences. This set is known as the quasiconvex hull $Q(K)$ of K . If the corresponding Young measures are trivial, then $Q(K) = K$. As stated before, that is the case for any number of pairwise incompatible linear strains in two dimensions. If there exists a nontrivial Young measure in the limit, as in the above examples, we have that K is strictly contained in $Q(K)$. In the fully pairwise compatible case, Bhattacharya [4] p. 231 –see also Lemma 4 in [17] p.46 – proved that the quasiconvex and convex hulls are equal for any number linear strains in K .

Quasiconvex hulls are generically very difficult (if not impossible) to compute for a particular

choice of \mathcal{U} . Explicit examples of quasiconvex functions [10, 20] and quasiconvex hulls [5, 7, 6, 25] are scarce and most of the nontrivial examples are not known explicitly. One approach to this problem is to compute the quasiconvex envelope of the energy density and to study its zero energy level set. This is a difficult problem, and solutions are only known in few cases, for example see theorems 3.4 and 3.5 in [13] pp 204 and 205. There is also extensive literature on the exploration of upper and lower bounds for the quasiconvex envelope of a function, see for example [11, 15, 18].

An alternative approach is to consider quasiconvex hulls' inner and outer bounds, such as the polyconvex, rank-one, and lamination convex hulls. Following [10] pp. 156 and 157, we introduce some standard notions from semi-convex analysis. We say that a function $f : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is *polyconvex* if there exists a convex function $G : \mathbb{R}^{m_n} \rightarrow \mathbb{R}$, such that $f(M) = G \circ T(M)$, where $T : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{m_n}$ is given by

$$T(M) = (M, \text{adj}_2(M), \text{adj}_3(M), \dots, \det M), \quad (2)$$

$\text{adj}_k(M)$ stands for the $k \times k$ matrix of sub-determinants of M , $m_n = C(2n, n) - 1$, and $C(2n, n)$ is the binomial coefficient between $2n$ and n . The function $f : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is *rank-one convex* if for every $M_1, M_2 \in \mathbb{R}^{n \times n}$ such that $\text{Rank}(M_1 - M_2) \leq 1$, and every $\lambda \in [0, 1]$,

$$f(\lambda M_1 + (1 - \lambda)M_2) \leq \lambda f(M_1) + (1 - \lambda)f(M_2).$$

It can be proved that if f is convex, it is polyconvex, if f is polyconvex, it is quasiconvex, and if f is quasiconvex, it is rank-one convex, but the reverse implications are false in general, see sections 5.3.2 to 5.3.9 in [10] and [21].

For any compact set $K \subset \mathbb{R}^{n \times n}$, its semi-convex hulls are defined by means of cosets, more precisely

$$S(K) = \left\{ A \in \mathbb{R}^{n \times n} \mid f(A) \leq \sup_{B \in K} f(B), \text{ for every semi-convex } f \right\}.$$

It is clear that $R(K) \subset Q(K) \subset P(K) \subset C(K)$ where these sets correspond to the rank-one, quasiconvex, polyconvex and convex hulls, respectively. These inclusions may not be proper, see for example Theorems 7.7 and 7.28 in [10].

In the geometrically linear regime, the relevant quantities [13] are given by the *displacement* $u(x) = y(x) - x$ and the *linear strain*

$$e(\nabla u) := \frac{\nabla u + (\nabla u)^T}{2}.$$

In this framework, the energy density becomes invariant under the addition of skew-symmetric matrices in its argument, *i.e.*, $f(M) = f(M + S)$ for every $S \in \mathbb{R}_{skew}^{n \times n}$, and it vanishes on the set of linearized elastic wells represented by $\mathcal{U} = \{U_1, U_2, \dots, U_k\} \in \mathbb{R}_{sym}^{n \times n}$. We use the notation $K = \mathcal{U} \oplus \mathbb{R}_{skew}^{n \times n}$ to indicate that the symmetric part of every element in K belongs to \mathcal{U} . Also, similar ideas about semi-convexity are available. Following Boussaid *et. al* [8] (section 2, pp 423 to 427), we say that a function $f : \mathbb{R}_{sym}^{n \times n} \rightarrow \mathbb{R}$ is symmetric semi-convex if $f(e(\cdot)) : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is semi-convex. It is straightforward to see that $f : \mathbb{R}_{sym}^{n \times n} \rightarrow \mathbb{R}$ is symmetric quasiconvex if and only if for every $U \in \mathbb{R}_{sym}^{n \times n}$

$$f(U) \leq \inf \left\{ \frac{1}{|\Omega|} \int_{\Omega} f(U + e(D\phi)) dx \mid \phi \in C_0^\infty(\Omega, \mathbb{R}^n) \right\},$$

and it is symmetric rank-one convex if and only if for every U_1 and U_2 compatible symmetric matrices and $\lambda \in [0, 1]$,

$$f(\lambda U_1 + (1 - \lambda)U_2) \leq \lambda f(U_1) + (1 - \lambda)f(U_2).$$

Characterization of symmetric polyconvexity for any dimension is rather challenging. Boussaid *et. al* [8] gave an explicit characterization of symmetric polyconvex functions in the cases of two and three dimensions. In the two-dimensional case, they proved that $f : \mathbb{R}_{sym}^{2 \times 2} \rightarrow \mathbb{R}$ is symmetric polyconvex if and only if there exists $g : \mathbb{R}_{sym}^{2 \times 2} \times \mathbb{R} \rightarrow \mathbb{R}$ convex, non-increasing on its second argument, and such that $f(\cdot) = g(\cdot, \det(\cdot))$.

Also, the symmetric semi-convex hull, see [26] p. 567, for a compact set $\mathcal{U} \in \mathbb{R}_{sym}^{n \times n}$ is defined in terms of the cosets,

$$S^e(\mathcal{U}) = \left\{ A \in \mathbb{R}^{n \times n} \mid f(A) \leq \sup_{B \in \mathcal{U}} f(B), f \text{ symmetric semi-convex} \right\}, \quad (3)$$

where S must be replaced by R, Q and P for the symmetric rank-one, quasiconvex and polyconvex hull of the set \mathcal{U} . As in the nonlinear case, to determine the symmetric quasiconvex hull of a compact set \mathcal{U} is a challenging task. An inner approximation for the symmetric quasiconvex hull is given by $L^e(\mathcal{U})$, the symmetric lamination convex hull of \mathcal{U} , see [26]. This set is defined as the intersection of the symmetric lamination hulls of all ranks, namely

$$L^e(\mathcal{U}) = \bigcup_{i=0}^{\infty} L^{e,i}(\mathcal{U}),$$

where $L^{e,0}(\mathcal{U}) = \mathcal{U}$, and

$$L^{e,i+1}(\mathcal{U}) = \{ \lambda A + (1 - \lambda)B \in \mathbb{R}_{sym}^{n \times n} \mid \lambda \in [0, 1] \text{ and } A, B \in L^{e,i}(\mathcal{U}) \text{ are compatible} \}.$$

It is also known, see for example [26] p. 562, that for any compact set $\mathcal{U} \subset \mathbb{R}_{sym}^{n \times n}$,

$$\mathcal{U} \subset L^e(\mathcal{U}) \subset R^e(\mathcal{U}) \subset Q^e(\mathcal{U}) \subset P^e(\mathcal{U}) \subset C(\mathcal{U}). \quad (4)$$

In this paper, we aim to understand the effect of partial pairwise compatibility among the elements of \mathcal{U} on the algebraic restrictions on the possible values of effective linear strains. We consider sets \mathcal{U} of three linearized elastic wells in two dimensions, where the wells are not fully compatible nor incompatible. Our goal is to characterize –or bound– the symmetric quasiconvex hull $Q^e(\mathcal{U})$ of \mathcal{U} .

Smyshlyaev and Willis [19] studied the relaxation of the three-well energy in linear elasticity where the elastic wells share the same elastic modulus. In particular, they showed that the minimizing measure may be chosen as the sum of no more than three Dirac masses. In [18], Schlömerkempe *et. al.* considered an analogous three-well setting in the three dimensional linear elasticity for a particular set of wells. Here, we do not consider any restriction on the wells to get general inner and outer bounds on $Q^e(\mathcal{U})$. For the interior bound, we characterize $L^e(\mathcal{U})$ explicitly, while for the outer bound, we estimate the zero level set of a suitable quadratic function's polyconvex envelope. In some cases we are able to show that our bounds are optimal.

In the rest of the paper we will follow standard notation. For every matrix $M \in \mathbb{R}^{2 \times 2}$, $e(M)$ and $w(M)$ denote the symmetric and skew-symmetric part of M respectively. Also for every $a, b \in \mathbb{R}^2$, the tensor product $a \otimes b \in \mathbb{R}^{2 \times 2}$ is defined as $(a \otimes b)_{ij} = a_i b_j$ and its symmetric part is denoted by $a \odot b$. The 2×2 identity matrix is denoted by Id , and the bilinear form $\langle \cdot, \cdot \rangle : \mathbb{R}^{2 \times 2} \times \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ stands for the Frobenius inner product *i.e.* $\langle A, B \rangle \mapsto \text{Tr}(A^T B)$, also $\| \cdot \|$ denotes the Frobenius norm. Since the space of skew-symmetric matrices has dimension one, we will use the following representation,

$$w(M) = w_M R, \quad \text{where} \quad R = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad w_M = \frac{1}{2} \langle M, R \rangle. \quad (5)$$

Along this work and for the sake of brevity, we shall say *wells* when we actually mean *linearized elastic wells*. Moreover, we refer to the elements of \mathcal{U} as *wells* or as the representative of wells equally.

2 Main results

In this paper, we are interested in the quasiconvex hull $Q^e(\mathcal{U})$ of the set of three elastic wells $\mathcal{U} = \{U_1, U_2, U_3\} \subset \mathbb{R}_{sym}^{2 \times 2}$.

Theorem 2.1. *Let $\mathcal{U} \subset \mathbb{R}_{sym}^{2 \times 2}$ represents a three-well set such that $\text{Aff}(\mathcal{U})$ has codimension one. If there exist at least two different matrices in \mathcal{U} that represent two rank-one compatible wells, then*

$$Q^e(\mathcal{U}) = L^e(\mathcal{U}).$$

This result can be described in terms of the compatibility between the wells. Independently of the existence of a rank-one compatibility, Bhattacharya [4] p. 231 –also see [17] p. 46– proved that $Q^e(\mathcal{U}) = L^e(\mathcal{U}) = C(\mathcal{U})$ if the wells are all pairwise compatible. On the other hand, Bhattacharya *et. al.* [7] (theorem 2.3 pp. 854) showed that if the wells are pairwise incompatible, then $Q^e(\mathcal{U}) = L^e(\mathcal{U}) = \mathcal{U}$ and no microstructure can be formed. The novelty of theorem 2.1 is to consider the intermediate cases when only one or two pairs of matrices in \mathcal{U} are compatible, and one of these compatibility relation is a rank-one compatibility.

In the following result, we give an explicit characterization of the symmetric lamination convex hull when at least two wells in \mathcal{U} are incompatible. Notice that –see proposition 4.1 below– two wells U_1 and U_2 are compatible if and only if $\det(U_1 - U_2) \leq 0$, and they are incompatible if $\det(U_1 - U_2) > 0$. Conditions $\det(U_1 - U_2) = 0$ and $U_1 \neq U_2$ are equivalent to rank-one compatibility between U_1 and U_2 .

Proposition 2.1. *Let $\mathcal{U} = \{U_1, U_2, U_3\}$ be such that $\text{Aff}(\mathcal{U})$ has codimension one.*

(a) *If $\det(U_1 - U_2) > 0$, $\det(U_1 - U_3) > 0$ and $\det(U_2 - U_3) \leq 0$, then*

$$L^e(\mathcal{U}) = \{U_1\} \cup C(\{U_2, U_3\}).$$

(b) *If $\det(U_1 - U_2) \leq 0$, $\det(U_1 - U_3) \leq 0$ and $\det(U_2 - U_3) > 0$, then*

$$L^e(\mathcal{U}) = C(\{U_0, U_1, U_2\}) \cup C(\{U_0, U_1, U_3\}),$$

where $U_0 \in C(\mathcal{U})$ is uniquely characterized by $\det(U_0 - U_3) = \det(U_0 - U_2) = 0$.

Our final result is about the symmetric quasiconvex hull when there exist one incompatible and one compatible pairs of wells in \mathcal{U} , and there are no rank-one compatibilities among its elements. With this result, we cover all possibilities for three wells in two dimensions. theorem 2.2 does not give a complete characterization of $Q^e(\mathcal{U})$, but it provides explicit quantitative outer bounds.

Theorem 2.2. *Let $\mathcal{U} = \{U_1, U_2, U_3\}$ represent a three-well set and assume that $\text{Aff}(\mathcal{U})$ has codimension one. Also, assume that (a) $\det(U_1 - U_2) > 0$, $\det(U_1 - U_3) > 0$, and $\det(U_2 - U_3) < 0$, or (b) $\det(U_1 - U_2) < 0$, $\det(U_1 - U_3) < 0$, and $\det(U_2 - U_3) > 0$, then*

$$Q^e(\mathcal{U}) \subset \{U \in C(\mathcal{U}) \mid 0 \leq \hbar(U - U_0)\} \subsetneq C(\mathcal{U}),$$

where $U_0 \in C(\mathcal{U})$ is uniquely characterized by $\det(U_0 - U_2) = \det(U_0 - U_3) = 0$, $\hbar : \mathbb{R}_{sym}^{2 \times 2} \rightarrow \mathbb{R}$ is given by

$$\hbar(V) = \langle C, V - V_2 \rangle \det V_1 - \langle C, V_1 - V_2 \rangle \det V,$$

and

$$C = \frac{(V_2 - V_1) \langle V_3 - V_1, V_3 - V_2 \rangle + (V_3 - V_1) \langle V_2 - V_1, V_2 - V_3 \rangle}{\sqrt{\|V_2 - V_1\|^2 \|V_3 - V_1\|^2 - \langle V_2 - V_1, V_3 - V_1 \rangle}}.$$

The outer bound of theorem 2.2 is not necessarily optimal. Nonetheless, we show that the outer bound in theorem 2.2 coincides with the quadratic polyconvex hull of \mathcal{U} at the end of this paper. A similar bound is given by Tang & Zhang [23] (theorem 1, p. 1266) for the case of a three-dimensional three-well problem at an exposed incompatible edge of $C(\mathcal{U})$. In [23], Tang & Zhang also proved that one can chip a wedge-like slice off $C(\mathcal{U})$ at the exposed incompatible edge without touching $Q^e(\mathcal{U})$. This later estimate is independent on the diameter of the set \mathcal{U} , but it is neither explicit nor optimal. In [18], the authors estimate by numerical approximation the quasiconvex hull for special combinations of three wells in three dimensional linear elasticity that come from the cubic-to-monoclinic martensitic phase transformations model. The estimated configurations are analogous to our two-dimensional results in theorem 2.2 and theorem 2.1. For comparison, see cases 3 and 4 and figures 5 to 7 in [18] and fig. 1 below.

An important ingredient in our proofs is the geometric characterization of the subset of compatible linear strains with the null matrix. By the isomorphism between \mathbb{R}^3 and $\mathbb{R}_{sym}^{2 \times 2}$ (see eq. (6) below), this set can be identified with the exterior of a solid cone \mathcal{C}_0 , whose vertex is the null matrix, and its axis is $\{t \text{ Id} \mid t \in \mathbb{R}\}$. Moreover, the set $\partial\mathcal{C}_0$ consists of rank-one symmetric matrices. The set of compatible linear strains with respect to a given matrix $U \in \mathbb{R}_{sym}^{2 \times 2}$ can be obtained by translation of \mathcal{C}_0 , namely $\mathcal{C}_U = U + \mathcal{C}_0$, (see Lemma 4.2 and Figure 2).

Theorems 2.1 and 2.2 can be geometrically interpreted by identifying $\mathbb{R}_{sym}^{2 \times 2}$ with \mathbb{R}^3 via the linear isomorphism

$$M = \begin{pmatrix} x & z \\ z & y \end{pmatrix} \mapsto \tilde{M} = \begin{pmatrix} x \\ y \\ z\sqrt{2} \end{pmatrix}. \quad (6)$$

If $\mathcal{U} = \{U_1, U_2, U_3\} \subset \mathbb{R}_{sym}^{2 \times 2}$ satisfies that $\text{Aff}(\mathcal{U})$ has codimension one, then $\text{Aff}(\{\tilde{U}_1, \tilde{U}_2, \tilde{U}_3\})$ defines a plane in \mathbb{R}^3 . Let $U_0 \in C(\mathcal{U})$ as in lemma 4.5, then the intersection between \mathcal{C}_{U_0} and $\text{Aff}(\mathcal{U})$ determines the set of incompatible wells with U_0 in $\text{Aff}(\mathcal{U})$. Moreover, $\text{Aff}(\mathcal{U}) \cap \partial\mathcal{C}_{U_0}$ consists of matrices U such that U and U_0 represent rank-one compatible wells, and due to the isomorphism (6), this set is identified with a pair of lines in \mathbb{R}^3 . Thus we shall say that $\text{Aff}(\mathcal{U}) \cap \partial\mathcal{C}_{U_0}$ is a pair of *rank-one lines*, see fig. 1. Additionally, if we assume that \mathcal{U} meets the conditions in Proposition 2.1, then the lamination convex hull is explicitly known and can be represented in the plane $\text{Aff}(\{\tilde{U}_1, \tilde{U}_2, \tilde{U}_3\})$. Figures 1.(a) and 1.(c) display two configurations with two incompatibility relations between \mathcal{U} 's wells. Due to Proposition 2.1, the lamination convex hull equals the union between $\{U_1\}$ and the segment joining U_2 and U_3 . In the configuration presented in figure 1.(a), the outer bound given by theorem 2.2 is the union of $\{U_1\}$ with the region bounded by the curve $\Gamma = \{U \in C(\mathcal{U}) \mid h(U - U_0) = 0\}$ and the segment joining U_2 and U_3 . Additionally, the configuration shown in fig. 1.(c) has a rank-one compatibility, and the set $Q^e(\mathcal{U})$ equals $L^e(\mathcal{U})$ by Theorem 2.1.

We present fig. 1.(b) and fig. 1.(d) to display the analogous configurations when only one pair of wells in \mathcal{U} is incompatible. In the configuration presented in fig. 1.(b), $L^e(\mathcal{U})$ is the wedge-like region bounded by the polygon with vertices U_1, U_2, U_0 and U_3 by proposition 2.1, and the well $U_0 \in C(\mathcal{U})$ is rank-one compatible with U_2 and U_3 simultaneously. theorem 2.2's outer bound correspond to the region bounded by the curve Γ and the two segments joining U_1 with U_2 and U_3 , respectively. The configuration presented in fig. 1.(d) has a rank-one compatible pair, so $Q^e(\mathcal{U})$ equals $L^e(\mathcal{U})$ due to theorem 2.1, and both sets equal the blue flag-like region.

The organization of the paper is as follows. In section 3, we briefly describe the ideas for the proofs and give some further comments. In section 4, we introduce the notion of incompatible




Figure 1 Four different three-well configurations related theorem 2.1 and theorem 2.2. Black and red dots represent the wells and U_0 , respectively. Solid lines represent compatibility between wells and dashed lines incompatibility. The gray planar cone represents $\text{Aff}(\mathcal{U}) \cap \mathcal{C}_{U_0}$, namely the set of incompatible matrices with U_0 in $\text{Aff}(\mathcal{U})$. Gray lines are the rank-one lines across U_0 . In figure (a) and (b), the blue regions are the outer bounds of $Q^e(\mathcal{U})$ in theorem 2.2. Figures (c) and (d) represent two admissible configurations in theorem 2.1, and blue regions are $Q^e(\mathcal{U}) = L^e(\mathcal{U})$.

cone and study some of its properties. The proof of proposition 2.1 is presented in section 5. In section 6, we present the symmetric polyconvex conjugate and symmetric biconjugate functions, and provide its application to our particular setting in section 7. The proofs of Theorems 2.1 and 2.2 are presented section 8. Finally, in section 9 we discuss the optimality of the outer bounds in theorem 2.2 with respect to symmetric quadratic polyconvex functions.

3 Ideas of proofs and further comments

In this paper, we construct quasiconvex hulls' inner and outer bound to $Q^e(\mathcal{U})$. The inner bound is the symmetric lamination convex hull $L^e(\mathcal{U})$ that we derive explicitly in proposition 2.1. For the outer bound, let $P^e(\mathcal{U})$ be the symmetric polyconvex hull of \mathcal{U} . If f is a non-negative symmetric polyconvex function such that \mathcal{U} belongs to $\text{Ker } f := \{M \in \mathbb{R}_{sym}^{2 \times 2} \mid f(M) = 0\}$, then $Q^e(\mathcal{U}) \subset P^e(\mathcal{U}) \subset \text{Ker } f \cap \mathcal{C}(\mathcal{U})$. The later inclusion is clear from the characterization of $P^e(\mathcal{U})$ in terms of cosets, i.e., eq. (3). We do not need to construct the polyconvex function f explicitly, but only its kernel. Finally, we construct the required matrix C in terms of the elements in \mathcal{U} . With this matrix C , $\text{Ker } f^{pp}$ equals the optimal outer bound when only quadratic polyconvex functions are used to bound $Q^e(\mathcal{U})$.

There are not many examples of polyconvex functions, and we need them to vanish on \mathcal{U} . Therefore, we construct the parametric family as follows. First, we consider a function of the form

$$f_C(M) = \chi_{\bar{B}}(e(M)) (|\det e(M)| + |\langle C, M \rangle|), \quad \bar{B} = \mathbb{R}_{sym}^{2 \times 2} \setminus L^e(\mathcal{U}),$$

where $\det C < 0$, and $\chi_{\bar{B}} : \mathbb{R}_{sym}^{2 \times 2} \rightarrow \{0, 1\}$ is the indicator function of the set \bar{B} . Notice that by construction, the functions f_C vanish on $L^e(\mathcal{U})$ (hence on \mathcal{U}), and they are quadratic outside this set. Second, we compute the symmetric polyconvex envelope of f_C through its symmetric polyconvex biconjugate function f_C^{PP} . Then, we estimate $\text{Ker } f_C^{PP}$ in lemma 7.2.

In our results, we describe the geometry of the laminar convex and quasiconvex hulls in terms of the matrix U_0 . This matrix is defined as the intersection of two rank-one lines passing through two wells in \mathcal{U} (see fig. 1). These lines exist since the cone \mathcal{C}_0 centered at any element of \mathcal{U} intersects the plane defined by $\text{Aff}(\mathcal{U})$. If we denote the normal matrix to $\text{Aff}(\mathcal{U})$ by Q , we show that the later condition is equivalent to $\det Q < 0$ (see remark 4.2). This is related to the condition $\det C < 0$ in the construction of f_C . Therefore, in our arguments, the geometric relationship between the incompatible cone and $\text{Aff}(\mathcal{U})$ is important.

In the last section, we show that the outer bound in theorem 2.2 is optimal for quadratic symmetric polyconvex functions. Indeed, we notice that this bound depends on the quadratic polyconvex function \bar{h} , and the proof of the optimality for quadratic polyconvex functions heavily depends on the recent characterization of symmetric polyconvex functions in two dimensions given by Boussaid *et. al* [8] (proposition 4.5 p.435), and it can not be extended beyond quadratic polyconvexity without an analogous result for non-quadratic functions. We believe that better estimates can be obtained by considering extensions of the form

$$f_C(M) = \chi_{\bar{B}}(e(M)) (|\det e(M)|^q + |\langle C, M \rangle|^p),$$

for some positive exponents p and q . The computation in these cases become cumbersome, and we do not pursue them further in this paper. Another consequence of the optimality in the outer bound for quadratic functions is that, if the outer bounds of $Q^e(\mathcal{U})$ in theorem 2.2 are not sharp, then the quasiconvex envelope of an energy density f (as in eq. (1)), is not quadratic.

Regarding possible extensions of the presented results, we study a generalization of theorem 2.1 to four and more co-planar wells in two dimensions in [9]. In particular, we are interested in the sufficient conditions on rank-one compatible pairs to show that $Q^e(\mathcal{U}) = L^e(\mathcal{U})$.

Finally, we make a few comments on the possible extension of our results to the three-dimensional case. The first step is to identify the corresponding geometry. We need to generalize the definition of the cone \mathcal{C}_0 and the normal Q to $\text{Aff}(\mathcal{U})$. To generalize the cone \mathcal{C}_0 , we notice that two matrices M_1 and M_2 in $\mathbb{R}_{sym}^{3 \times 3}$ represent compatible linear strains if the eigenvalues of $M_1 - M_2$ are of the form $\lambda_1 < \lambda_2 = 0 < \lambda_3$. Now, the cone's boundary is a "surface" that represents the rank-one compatible directions. Therefore, by the isomorphism between $\mathbb{R}_{sym}^{3 \times 3}$ and \mathbb{R}^6 , this boundary should be an object of four dimensions in \mathbb{R}^6 . Consequently, the tangent space to the cone's boundary is of dimension four, and we conjecture that we need at most five wells to set up the analogous geometry. The challenge, in this case, is to define the analogous to Q and the generalization of condition $\det Q < 0$ that identifies the intersection of the incompatible cone and the affine space. In some cases, we believe that it is possible that for a particular set of wells in three dimensions, the cone geometry reduces to its two-dimensional counterpart, and we recover results similar to the ones presented in this paper. As a piece of evidence in this direction, we recall the results in [18] quoted before.

4 On the geometry of the incompatible cone

We begin with an equivalence result on compatibility.

Proposition 4.1. *Let $M_1, M_2 \in \mathbb{R}^{2 \times 2}$. The following statements are equivalent:*

- (a) *There exist $v \in \mathbb{R}^2$, $a \in S^1$ and $W \in \mathbb{R}_{skew}^{2 \times 2}$ such that $M_1 - M_2 + W = a^\perp \otimes v$.*

(b) There exist $v \in \mathbb{R}^2$ and $a \in S^1$ such that $e(M_1) = e(M_2) + v \odot a^\perp$.

(c) There exists $a \in S^1$ such that $\langle e(M_1) - e(M_2), a \otimes a \rangle = 0$.

(d) $\det(e(M_1) - e(M_2)) \leq 0$.

Proof. That (a) implies (b) follows directly from the symmetric parts of the equation in the statement (a). Since a^\perp is orthogonal to a , we easily conclude that (b) implies (c). To show that (c) implies (d), we proceed by contradiction and assume $\det(e(M_1) - e(M_2)) > 0$, meanwhile (c) holds. Then it follows that $a \mapsto (e(M_1) - e(M_2))a \cdot a$ is either a positive or negative defined quadratic form. Thus, the unique solution to $(e(M_1) - e(M_2))a \cdot a = 0$ is $a = 0$, a contradiction to $a \in S^1$ by hypothesis.

Now, we show that (d) implies (a). Since $\det M = \det(e(M)) + \det(w(M))$ for every $M \in \mathbb{R}^{2 \times 2}$ and (d) is assumed, we obtain

$$\det(M_1 - M_2 + \mu R) = \det(e(M_1) - e(M_2)) + \left(\frac{1}{2} \langle M_1 - M_2, R \rangle + \mu \right)^2.$$

Thus, the equation $\det(M_1 - M_2 + \mu R) = 0$ is satisfied by some $\mu^* \in \mathbb{R}$. So, the statement (a) follows, and the proof is finished. \blacksquare

The following lemma states that if $M \in \mathbb{R}_{sym}^{2 \times 2}$ has non-positive determinant, then it is the symmetric part of a tensor product, this result is used in the proof of Lemma 4.3.

Lemma 4.1. *Let $Q \in \mathbb{R}_{sym}^{2 \times 2}$ be such that its determinant is non-positive. There exist $a, n \in S^1$ and $\nu \in \mathbb{R}$ such that $Q = \nu a \odot n$ and $\det Q = -\nu^2(a \cdot n^\perp)^2/4$. Moreover, $v_+ = a + n$, and $v_- = a - n$ are eigenvectors of Q with eigenvalues $\lambda_+ = \nu[a \cdot e + 1]/2$, and $\lambda_- = \nu[a \cdot e - 1]/2$, respectively.*

Proof. The existence of $a, n \in S^1$, and $\nu \in \mathbb{R}$ such that $Q = \nu a \odot n$ follows by setting $M_1 = Q$, $M_2 = 0$ into the equivalence between (b) and (d) in proposition 4.1. The statement concerning the eigenvalues and eigenvectors follows by a direct computation

$$Qv_\pm = \frac{\nu}{2}(a \otimes n + n \otimes a)(a \pm n) = \frac{\nu}{2}[(a \cdot n)(a \pm n) + (n \pm a)] = \frac{\nu}{2}[a \cdot n \pm 1](a \pm n).$$

Thus, $\det Q = \lambda_+ \lambda_- = -\nu^2[1 - (a \cdot n)^2]/4 = -\nu^2(a \cdot n^\perp)^2/4$. \blacksquare

The next lemma characterizes the set of symmetric incompatible matrices as elements in a solid cone. In particular, the set of symmetric matrices that are rank-one compatible is identified with the boundary of this cone.

Lemma 4.2. *Let $M \in \mathbb{R}_{sym}^{2 \times 2}$, then $\det M = 0$ if and only if $|\langle M, \text{Id} \rangle| = \|M\|$. Moreover, $\det M < 0$ if and only if $|\langle M, \text{Id} \rangle| < \|M\|$.*

Proof. Let $M \in \mathbb{R}_{sym}^{2 \times 2}$ be given by

$$M = \begin{pmatrix} M_{11} & M_{12} \\ M_{12} & M_{22} \end{pmatrix}$$

with $M_{11}, M_{12}, M_{22} \in \mathbb{R}$, and consider the square of its Frobenius norm

$$\|M\|^2 = M_{11}^2 + M_{22}^2 + 2M_{12}^2 = (M_{11} + M_{22})^2 + 2(M_{12}^2 - M_{11}M_{22}).$$

Hence, $\|M\|^2 = \langle M, \text{Id} \rangle^2 - 2 \det M$, and the proof follows. \blacksquare

Because of proposition 4.1 and lemma 4.2, the set of incompatible linear strains with $U \in \mathbb{R}_{sym}^{2 \times 2}$ is given by the interior of the cone

$$\begin{aligned} \mathcal{C}_U &:= \left\{ V \in \mathbb{R}_{sym}^{2 \times 2} \text{ such that } \|V - U\| < |\langle V - U, \text{Id} \rangle| \right\}, \\ &= \left\{ V \in \mathbb{R}_{sym}^{2 \times 2} \text{ such that } \frac{1}{\sqrt{2}} < \left| \left\langle \frac{V - U}{\|V - U\|}, \frac{1}{\sqrt{2}} \text{Id} \right\rangle \right| \right\} \end{aligned}$$

Let e^1 , e^2 , and e^3 stand for the canonical vectors in \mathbb{R}^3 . Then $e^1 \otimes e^1$ and $e^2 \otimes e^2$ belong to $\partial \mathcal{C}_0$, and the cone's aperture angle, 2θ , is $\pi/2$ since $e^1 \otimes e^1$, $e^2 \otimes e^2$ and Id are coplanar, see fig. 2.

Remark 4.1. Let $\mathcal{S} : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}^{2 \times 2}$ stand for the linear map $M \mapsto RMR^T$, with R as in (5). The function \mathcal{S} maps the symmetric (skew-symmetric) subspace into itself. Moreover, a simple calculation shows that $\text{adj}(M) = SM$ for every $M \in \mathbb{R}^{2 \times 2}$, and we get the following identities:

- (a) $2 \det M = \langle SM, M \rangle$.
- (b) $\det M = \det e(M) + \det w(M) = \det e(M) + w_M^2$.
- (c) $\det(N + M) = \det M + \det M + \langle SM, N \rangle$, $N \in \mathbb{R}^{2 \times 2}$.
- (d) If $M = \xi a^\perp \otimes a^\perp + \eta n^\perp \otimes n^\perp$ for some $a, n \in S^1$ and $\xi, \eta \in \mathbb{R}$, then $\det M = \xi\eta|a \times n|^2$.




Figure 2 The incompatible cone \mathcal{C} represented in \mathbb{R}^3 by the isomorphism (6)

The next lemma describes a coordinate system defined by the intersection between the boundary of the incompatible cone and $\text{Aff}(\mathcal{U})$. This system of coordinates is convenient and simplifies many of the computations in our proofs.

Lemma 4.3. Let $V, Q \in \mathbb{R}_{sym}^{2 \times 2}$ such that $\det Q < 0$ and denote by $\Pi_Q(V)$ the affine set of codimension one that contains V and is normal to Q , namely

$$\Pi_Q(V) = \{U \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle Q, U - V \rangle = 0\}.$$

Then, there exist two nonparallel vectors $a, n \in S^1$, such that

$$\Pi_Q(V) = \{V + \xi a^\perp \otimes a^\perp + \eta n^\perp \otimes n^\perp \mid (\xi, \eta) \in \mathbb{R}^2\}. \quad (7)$$

Moreover, $U \in \Pi_Q(V)$ and V represent compatible linear strains if and only if $U - V = \xi a^\perp \otimes a^\perp + \eta n^\perp \otimes n^\perp$ with $\xi\eta \leq 0$.

Proof. By Lemma 4.1, there exist $a, n \in S^1$, and $\nu \in \mathbb{R} \setminus \{0\}$ such that $Q = \nu a \odot n$. The vectors a and n are linearly independent since $\det Q < 0$. Moreover, by proposition 4.1 (b) and (c), and the symmetric role between vectors a and n , we have that $a^\perp \otimes a^\perp$ and $n^\perp \otimes n^\perp$ are two linearly independent rank-one matrices such that

$$\langle Q, a^\perp \otimes a^\perp \rangle = \langle Q, n^\perp \otimes n^\perp \rangle = 0.$$

Hence, $\{a^\perp \otimes a^\perp, n^\perp \otimes n^\perp\}$ is a basis for the two-dimensional subspace $\Pi_Q(V) - V$ and (7) follows. Now, let $W \in \Pi_Q(V)$ and $(\xi, \eta) \in \mathbb{R}^2$ such that $W - V = \xi a^\perp \otimes a^\perp + \eta n^\perp \otimes n^\perp$. Thus $\det(W - V) = \xi\eta|a \times n|^2$, and the second part of the statement follows straight forward from item (d) in proposition 4.1. \blacksquare

Remark 4.2. *It is well known, see [7, 10] pages p.191 and p.825 respectively, that if $f : \mathbb{R}^{m \times m} \rightarrow \mathbb{R}$ is quadratic and $f(a \otimes n) \geq 0$ for every $a, n \in \mathbb{R}^m$ then f is rank-one convex. Thus, $-\det(\cdot) : \mathbb{R}_{sym}^{2 \times 2} \rightarrow \mathbb{R}$ is a symmetric rank-one convex function because item (a) in remark 4.1 implies that it is quadratic and the equivalence between items (b) and (d) in proposition 4.1 implies that $-\det(e(a \otimes n)) \geq 0$ for every $a, n \in \mathbb{R}^2$.*

Lemma 4.4. *Let $U, V, W \in \mathbb{R}_{sym}^{2 \times 2}$ such that V, W are compatible and U is incompatible with both of them. Then U is incompatible with any point in $C(\{U, V, W\}) \setminus \{U\}$.*

Proof. Let $M \in C(\{U, V, W\}) \setminus \{U\}$. Then, $M = \lambda_1 U + \lambda_2 V + \lambda_3 W$ for some $\lambda_1 \in [0, 1)$ and $\lambda_2, \lambda_3 \in [0, 1]$, such that $\lambda_1 + \lambda_2 + \lambda_3 = 1$. Since $\lambda_1 \neq 1$, we have that $\lambda_2 + \lambda_3 > 0$ and

$$M - U = (\lambda_2 + \lambda_3)(s(V - U) + (1 - s)(W - U)),$$

where $s = \lambda_2/(\lambda_2 + \lambda_3)$, and $(1 - s) = \lambda_3/(\lambda_2 + \lambda_3)$. Hence, $\det(M - U) = (\lambda_2 + \lambda_3)^2 \det(s(V - U) + (1 - s)(W - U))$. Now, since $V - U$ and $W - U$ are compatible, and $-\det(e(\cdot))$ is rank-one convex, we have that

$$\det(s(V - U) + (1 - s)(W - U)) \geq s \det(V - U) + (1 - s) \det(W - U) > 0.$$

Therefore, U and M are incompatible and the proof is complete. \blacksquare

In Proposition 2.1 and Theorem 2.2, we consider two possible configurations of wells depending on the number of compatible pairs among its elements. We use the following definition to keep our presentation compact.

Definition 4.1. *Let \mathcal{U} represent a three-well set such that $\text{Aff}(\mathcal{U})$ has codimension one. We say that \mathcal{U} is type one if up to a relabeling $\mathcal{U} = \{U_1, U_2, U_3\}$, where*

$$\det(U_1 - U_2) > 0, \det(U_1 - U_3) > 0, \text{ and } \det(U_2 - U_3) \leq 0, \quad (8)$$

and \mathcal{U} is type two if up to a relabeling $\mathcal{U} = \{U_1, U_2, U_3\}$, where

$$\det(U_1 - U_2) \leq 0, \det(U_1 - U_3) \leq 0, \text{ and } \det(U_2 - U_3) > 0. \quad (9)$$

Remark 4.3. *We claim that if \mathcal{U} is either type one or type two and Q is normal to $\text{Aff}(\mathcal{U})$, then $\det Q < 0$. We argue by contradiction. Indeed, if $\det Q \geq 0$, then $\text{Aff}(\mathcal{U}) \cap \mathcal{C}_{U_1} = \{U_1\}$ and U_1 is compatible with U_2 and U_3 , see paragraph after lemma 4.2. We conclude that \mathcal{U} is pairwise compatible by arguing the same for the two remaining wells, but this is a contradiction to eq. (8) and eq. (9).*

In the following lemma, we apply the coordinate system in lemma 4.3 to the geometry of our three-well problem.

Lemma 4.5. *Let $\mathcal{U} \subset \mathbb{R}_{sym}^{2 \times 2}$ represent a three-well set where \mathcal{U} is either type one or type two. Then, there exist $U_2, U_3 \in \mathcal{U}$ and $U_0 \in C(\mathcal{U})$ such that $\det(U_2 - U_0) = \det(U_3 - U_0) = 0$.*

Proof. First, by remark 4.3, $\det Q < 0$, and by lemma 4.1, there exist $a, n \in S^1$ linearly independent vectors such that $Q = \nu a \odot n$ for some $\nu \in \mathbb{R} \setminus \{0\}$. Hence, by Lemma 4.3,

$$U_2 = U_1 + \xi_2 a^\perp \otimes a^\perp + \eta_2 n^\perp \otimes n^\perp, \quad \text{and} \quad U_3 = U_1 + \xi_3 a^\perp \otimes a^\perp + \eta_3 n^\perp \otimes n^\perp, \quad (10)$$

for some $(\xi_2, \eta_2), (\xi_3, \eta_3) \in \mathbb{R}^2$. We notice that (ξ_2, η_2) and (ξ_3, η_3) belong to the same quadrant in the \mathbb{R}^2 . Otherwise, all elements in \mathcal{U} are either compatible or incompatible and this contradicts either eq. (8) or eq. (9).

Second, we claim that

$$U_0 = U_1 + \alpha a^\perp \otimes a^\perp + \beta n^\perp \otimes n^\perp, \quad \text{with} \quad \begin{cases} \alpha = \arg \min\{|\xi| \mid \xi \in \{\xi_2, \xi_3\}\}, \\ \beta = \arg \min\{|\eta| \mid \eta \in \{\eta_2, \eta_3\}\}, \end{cases}$$

is the desired matrix. Indeed, without loss of generality we assume \mathcal{U} is type one. Then, either $\alpha = \xi_3$ or $\beta = \eta_3$, but not both. Otherwise, by Remark 4.1 (d) and eq. (10), we get $\det(U_3 - U_2) = (\xi_3 - \xi_2)(\eta_3 - \eta_2)|a \times n|^2 \geq 0$, a contradiction with \mathcal{U} being of type one. Hence, $\det(U_3 - U_0) = (\xi_3 - \alpha)(\eta_3 - \beta)|a \times n|^2 = 0$, and by an analogous argument, we also have $\det(U_2 - U_0) = 0$. Now, since U_1 is incompatible with both U_2 and U_3 (namely, $\xi_2 \eta_2 > 0$ and $\xi_3 \eta_3 > 0$), and (ξ_2, η_2) and (ξ_3, η_3) belong to the same quadrant, we have that $\det(U_1 - U_0) = \alpha \beta |a \times n|^2 \geq 0$. We let

$$\lambda_1 = \frac{(\alpha - \xi_2)(\beta - \eta_3) - (\alpha - \xi_3)(\beta - \eta_2)}{\xi_2 \eta_3 - \xi_3 \eta_2}, \quad \lambda_2 = \frac{(\alpha - \xi_3)\beta - (\beta - \eta_3)\alpha}{\xi_2 \eta_3 - \xi_3 \eta_2}, \quad \text{and} \quad \lambda_3 = \frac{(\beta - \eta_2)\alpha - (\alpha - \xi_2)\beta}{\xi_2 \eta_3 - \xi_3 \eta_2}.$$

By a straight forward computation, we get that $U_0 = \lambda_1 U_1 + \lambda_2 U_2 + \lambda_3 U_3$ with $\lambda_1, \lambda_2, \lambda_3 \in (0, 1)$, and $\lambda_1 + \lambda_2 + \lambda_3 = 1$. Hence, $U_0 \in C(\mathcal{U})$ as claimed. The case where \mathcal{U} is type two can be treated analogously, and the result follows. \blacksquare

Corollary 4.0.1. *Let $\mathcal{U} \subset \mathbb{R}_{sym}^{2 \times 2}$ represent a three-well set where \mathcal{U} is either type one or type two. Then, there exist $U_0 \in C(\mathcal{U})$, $a, n \in S^1$, and $\xi, \eta, \gamma, \zeta \in \mathbb{R}$ such that*

$$U_1 = U_0 + \xi a^\perp \otimes a^\perp + \eta n^\perp \otimes n^\perp, \quad U_2 = U_0 + \gamma a^\perp \otimes a^\perp, \quad \text{and} \quad U_3 = U_0 + \zeta n^\perp \otimes n^\perp, \quad (11)$$

where

$$\begin{cases} \eta > 0 \geq \zeta, \quad \xi > 0 \geq \gamma \quad \text{or} \quad \zeta \geq 0 > \eta, \quad \gamma \geq 0 > \xi & \text{if } \mathcal{U} \text{ is type one,} \\ \eta \geq 0 > \zeta, \quad \gamma > 0 \geq \xi \quad \text{or} \quad \zeta > 0 \geq \eta, \quad \xi \geq 0 > \gamma & \text{if } \mathcal{U} \text{ is type two.} \end{cases} \quad (12)$$

Proof. By lemma 4.5, there exists $U_0 \in C(\mathcal{U})$ that is rank-one compatible with U_2 and U_3 . From, remark 4.3, the normal Q to $\text{Aff}(\mathcal{U})$ is such that $\det Q < 0$. Then, (11) follows from lemma 4.1, lemma 4.3, and that $\Pi_Q(U_0) = \text{Aff}(\mathcal{U})$ has codimension one.

Next, we prove eq. (12). Without loss of generality we assume that \mathcal{U} is of type two. Hence, $\det(U_2 - U_3) > 0$, and $\det(U_j - U_1) \leq 0$ for $j = 0, 2, 3$. Therefore, by (11) and remark 4.1 item (d), we get

$$-\gamma \zeta > 0, \quad \xi \eta \leq 0, \quad (\xi - \gamma)\eta \leq 0, \quad \text{and} \quad (\eta - \zeta)\xi \leq 0.$$

Now, the conditions in (12) follows since $U_0 \in C(\mathcal{U})$. The case where \mathcal{U} is type one can be proved by a similar argument. \blacksquare

5 The symmetric lamination convex hull of \mathcal{U}

In this section, we determine the symmetric lamination convex hull when \mathcal{U} is either type one or type two, *i.e.* there is one out of the three wells in \mathcal{U} that is compatible with one of the two remaining wells and incompatible with the other.




Figure 3 Auxiliary wells used in the proof of proposition 2.1. This figure shows a three-well set with strict inequalities in item (b). The incompatible cone of U_0 restricted to $\text{Aff}(\mathcal{U})$ is represented by the shaded region.

Proof. of proposition 2.1: We prove the item (a). By hypothesis, U_2 is compatible with U_3 , and U_1 is incompatible with both of them. Then, $L^{e,1}(\mathcal{U}) = \{U_1\} \cup C(\{U_2, U_3\})$. By lemma 4.4 there is not $U \in C(\mathcal{U})$ compatible with U_1 . Therefore, no laminate of degree two is admissible and $L^{e,2}(\mathcal{U}) = L^{e,1}(\mathcal{U})$. By the same argument, $L^{e,n}(\mathcal{U}) = L^{e,1}(\mathcal{U})$ for every $n \geq 1$, and item (a) follows from the definition of $L^e(\mathcal{U})$.

Now, we prove the item (b). In this case \mathcal{U} is of type two. Then, lemma 4.5 implies the existence of $U_0 \in C(\mathcal{U})$ such that

$$\det(U_2 - U_0) = \det(U_3 - U_0) = 0,$$

and we assume that the wells in \mathcal{U} are given by eq. (11) in corollary 4.0.1. Now, by the compatibility relations, \mathcal{U} 's first symmetric lamination is $L^{e,1}(\mathcal{U}) = C(\{U_1, U_2\}) \cup C(\{U_1, U_3\})$. For the laminations of degree greater than one, first, we assume that $\det(U_1 - U_2) < 0$ and $\det(U_1 - U_3) < 0$ and consider the continuous function

$$t \mapsto \det(tU_1 + (1-t)U_3 - U_0) = t^2 \det(U_1 - U_3) + t \langle \mathcal{S}(U_3 - U_1), U_3 - U_0 \rangle.$$

By remark 4.1, eq. (11), and (12) (b) for the type two case, we have that

$$\langle \mathcal{S}(U_3 - U_1), U_3 - U_0 \rangle = -\xi\zeta|n \times a|^2 > 0.$$

Thus, there exists $t_0 > 0$ such that $U_{1,3} = t_0U_1 + (1-t_0)U_3 \in C(\{U_1, U_3\})$ and $\det(U_{1,3} - U_0) = 0$ since $\det(U_1 - U_3) < 0$. Analogously, there exists $U_{1,2} \in C(\{U_1, U_2\})$ such that $\det(U_{1,2} - U_0) = 0$. Now, we define two triangular regions with vertices $K_2 = \{U_1, U_2, U_{1,3}\}$ and $K_3 = \{U_1, U_3, U_{1,2}\}$ (see fig. 3), and divide the proof in three steps:

(i) We claim that $C(K_2), C(K_3) \subset L^{e,2}(\mathcal{U})$. Indeed, if we assume that M belongs to the relative boundary of $C(K_2)$, then either $M \in C(\{U_1, U_2\})$ or $M \in C(\{U_2, U_{1,3}\}) \cup C(\{U_1, U_{1,3}\})$. Since $U_{1,3} \in L^{e,1}(\mathcal{U})$ and $C(\{U_1, U_2\}) \subset L^{e,1}(\mathcal{U})$ then $M \in L^{e,2}(\mathcal{U})$. Now, we assume that M belongs to the relative interior of $C(K_2)$. By construction (see lemma 4.2), any two matrices on the line ℓ joining U_2 and $U_{1,3}$, or on any other parallel line, are rank-one compatible. Then, there exists a line ℓ_M parallel to ℓ passing through M , such that ℓ_M intersects $C(\{U_1, U_2\})$ and $C(\{U_1, U_{1,3}\})$ at some points V and W respectively. Thus, $M \in C(\{V, W\})$, $\det(V - W) = 0$, and $V, W \in L^{e,1}(\mathcal{U})$. Hence, $M \in L^{e,2}(\mathcal{U})$. By a similar argument $C(K_3) \subset L^{e,2}(\mathcal{U})$ and the claim follows.

(ii) Now, we claim that $C(K_2) \cup C(K_3) = L^{e,2}(\mathcal{U})$. By contradiction, we assume there exists $U \in L^{e,2}(\mathcal{U}) \setminus (C(K_2) \cup C(K_3))$ (see fig. 3). The intersection between $\partial\mathcal{C}_U$ and the relative

boundary of $C(\mathcal{U})$ yields the existence of $V_2, V_3 \in C(\{U_2, U_3\})$, $V_{1,2} \in C(\{U_1, U_2\})$, and $V_{1,3} \in C(\{U_1, U_3\})$ such that $\det(V_2 - U) = \det(V_{1,2} - U) = 0$ and $\det(V_3 - U) = \det(V_{1,3} - U) = 0$. Therefore, each matrix in the set \mathcal{U} that is compatible with U belongs to either the set $C_1 = C(\{U, V_{1,3}, U_1, V_{1,2}\})$ or the set $C_2 = C(\{U, V_2, V_3\})$. Now, since $U \in L^{e,2}(\mathcal{U})$, there exist two compatible matrices $V, W \in L^{e,1}(\mathcal{U})$ such that $U = \lambda V + (1 - \lambda)W$ for some $\lambda \in (0, 1)$. Notice that C_1 and C_2 are convex sets and $C_1 \cap C_2 = \{U\}$, so V and W cannot belong both to C_1 or C_2 simultaneously. The contradiction follows from the fact $C_2 \cap L^{e,1}(\mathcal{U}) = \emptyset$ since U_2 and U_3 are incompatible.

- (iii) We claim that $L^{e,2}(\mathcal{U}) = L^{e,3}(\mathcal{U})$. The proof follows by a contradiction argument similar to the one used before. Indeed, if there exists $U \in L^{e,3}(\mathcal{U}) \setminus L^{e,2}(\mathcal{U})$, then U is a convex combination of two compatible matrices $V, W \in L^{e,2}(\mathcal{U})$. Since $C_1 \cap C_2 = \{U\}$, V and W cannot belong both to C_1 or C_2 . Again the contradiction emerged from the fact that $C_2 \cap L^{e,2}(\mathcal{U}) = \emptyset$. Hence, $L^{e,3}(\mathcal{U}) \subset L^{e,2}(\mathcal{U})$. The reverse inclusion follows from the definition of $L^{e,3}(\mathcal{U})$.

Second, we assume that there is only one rank-one compatibility among the elements of \mathcal{U} and without loss of generality, let $\det(U_1 - U_2) < 0$ and $\det(U_1 - U_3) = 0$. In this case, we define $K_2 = C(\{U_1, U_0, U_2\})$ and $K_3 = C(\{U_1, U_3\})$. Repeating the argument in step (i), we easily get that $K_2 \subset L^{e,2}(\mathcal{U})$. Moreover, by definition $K_3 \subset L^{e,1}(\mathcal{U}) \subset L^{e,2}(\mathcal{U})$, we obtain the claim in step (i). The steps (ii) and (iii) are exactly the same.

Third, we assume that there are two rank-one compatible pairs in \mathcal{U} , so without loss of generality, $\det(U_2 - U_1) = \det(U_3 - U_1) = 0$. For this case, we choose $K_2 = C(\{U_1, U_2\})$ and $K_3 = C(\{U_1, U_3\})$. Under these assumptions, $K_1 \cup K_2 = C(K_1) \cup C(K_2) = L^{e,1}(\mathcal{U})$. Hence, by step (ii), $L^{e,1}(\mathcal{U}) = L^{e,2}(\mathcal{U})$.

Finally, due to $L^{e,3}(\mathcal{U}) = L^{e,2}(\mathcal{U})$ we get that $L^{e,n}(\mathcal{U}) = L^{e,2}(\mathcal{U})$ for every $n \geq 2$ in all cases, and item (b) follows from the definition of $L^e(\mathcal{U})$. \blacksquare

6 Results about polyconvex conjugate and biconjugate functions

In this section we presents a method to construct non-negative polyconvex functions through the polyconvex conjugate and biconjugate functions $f^p(\xi^*) : \mathbb{R}^{m_n} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ and $f^{pp}(\xi) : \mathbb{R}^{n \times n} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ of a given function $f : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$, respectively. These functions are defined as

$$f^p(\xi^*) := \sup\{(T(\eta), \xi^*) - f(\eta) \mid \eta \in \mathbb{R}^{n \times n}\}, \quad \text{and} \quad f^{pp}(\xi) := (f^p)^*(T(\xi)),$$

where $T : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{m_n}$ is given by (2), and $(f^p)^* : \mathbb{R}^{m_n} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is the dual convex of f^p , which is given by $(f^p)^*(\eta) := \sup\{(\eta, \xi^*) - f^p(\xi^*) \mid \xi^* \in \mathbb{R}^{m_n}\}$. The following theorem was proved by Kohn & Strang [14] (lemma 3.4 p, 133) also, see theorem 6.6 [10] p.268.

Theorem 6.1. *Let $f : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ and $f^{pp}(M) := (f^p)^* \circ T(M)$. If $g : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is a polyconvex function such that $f(\xi) \geq g(\xi)$ for every $\xi \in \mathbb{R}^{n \times n}$, then $f^{pp} = Pf$, the polyconvex envelope of f .*

In the two-dimensional framework, $f : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ and, the above expressions become

$$f^p(M^*, \delta^*) = \sup_{M \in \mathbb{R}^{2 \times 2}} \langle M^*, M \rangle + \delta^* \det M - f(M), \quad \text{with } M^* \in \mathbb{R}^{2 \times 2} \text{ and } \delta^* \in \mathbb{R}, \quad (13)$$

and $f^{pp}(M) = (f^p)^*(M, \det M)$, where

$$(f^p)^*(M, \lambda) = \sup_{\substack{\delta^* \in \mathbb{R} \\ M^* \in \mathbb{R}^{2 \times 2}}} (\langle M, M^* \rangle + \delta^* \lambda - f^p(M^*, \delta^*)), \quad \text{for } M \in \mathbb{R}^{2 \times 2} \text{ and } \lambda \in \mathbb{R}.$$

In the proof of Theorems 2.1 and 2.2, we strongly use the existence of the matrix U_0 given in Lemma 4.5. By the self-similarity of the incompatible cone and Lemma 4.3, we set U_0 as the center of coordinates. The following lemma allow us to do that.

Proposition 6.1. *Let $f : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ and $g(M) = f(M - M_0)$, then $g^{pp}(M) = f^{pp}(M - M_0)$.*

Proof. From relation (13),

$$\begin{aligned} g^p(M^*, \delta^*) &= \sup_{M \in \mathbb{R}^{2 \times 2}} (\langle M^*, M \rangle + \delta^* \det M - f(M - M_0)) \\ &= \sup_{\bar{M} \in \mathbb{R}^{2 \times 2}} (\langle M^*, \bar{M} + M_0 \rangle + \delta^* \det(\bar{M} + M_0) - f(\bar{M})). \end{aligned}$$

The last expression for $g^p(M^*, \delta^*)$ is easily simplified by Remark 4.1 (c). Hence

$$g^p(M^*, \delta^*) = \langle M^*, M_0 \rangle + \delta^* \det M_0 + f^p(M^* + \delta^* \mathcal{S}M_0, \delta^*).$$

Moreover,

$$\begin{aligned} (g^p)^*(M, \delta) &= \sup_{\substack{M^* \in \mathbb{R}^{2 \times 2} \\ \delta^* \in \mathbb{R}}} \{ \langle M - M_0, M^* \rangle + \delta^*(\delta - \det M_0) - f^p(M^* + \delta^* \mathcal{S}M_0, \delta^*) \}, \\ &= \sup_{\substack{\bar{M} \in \mathbb{R}^{2 \times 2} \\ \delta^* \in \mathbb{R}}} \{ \langle M - M_0, \bar{M} \rangle + \delta^*(\delta - \det M_0 - \langle M - M_0, \mathcal{S}M_0 \rangle) - f^p(\bar{M}, \delta^*) \}. \end{aligned}$$

Thus, $(g^p)^*(M, \delta) = (f^p)^*(M - M_0, \delta - \det M_0 - \langle M - M_0, \mathcal{S}M_0 \rangle)$. Finally, the affirmation follows from $g^{pp}(M) = (g^p)^*(M, \det M)$, and $\langle M - M_0, \mathcal{S}M_0 \rangle = -2 \det M_0 + \langle \mathcal{S}M_0, M \rangle$. \blacksquare

The next proposition determines the polyconvex conjugate function of $f : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ when it depends on the symmetric part of the argument.

Proposition 6.2. *Let $f : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ be such that $f(M) = f(e(M))$ for every $M \in \mathbb{R}_{sym}^{2 \times 2}$. Then, the polyconvex conjugate function of f is given by*

$$f^p(M^*, \delta^*) = \begin{cases} \infty, & \text{if } \delta^* > 0 \text{ or } \delta^* = 0, w_{M^*} \neq 0, \\ -\frac{w_{M^*}^2}{\delta^*} + \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(e(M^*), \delta^*, U), & \text{if } \delta^* < 0, \\ \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(e(M^*), 0, U), & \text{if } \delta^* = w_{M^*} = 0, \end{cases} \quad (14)$$

where $w_M \in \mathbb{R}$ satisfies $w(M) = w_M R$ and

$$\mathcal{L}(V, \delta, U) := \langle V, U \rangle + \delta \det U - f(U). \quad (15)$$

Proof. From (13) and Remark 4.1 (b), we get

$$\begin{aligned} f^p(M^*, \delta^*) &= \sup_{M \in \mathbb{R}^{2 \times 2}} [\langle M^*, M \rangle + \delta^* \det M - f(e(M))] \\ &= \sup_{M \in \mathbb{R}^{2 \times 2}} [2w_M w_{M^*} + \delta^* w_M^2 + \langle e(M^*), e(M) \rangle + \delta^* \det e(M) - f(e(M))]. \end{aligned}$$

Since w_M and $e(M)$ are independent, we conclude that

$$f^p(M^*, \delta^*) = \sup_{w \in \mathbb{R}} (2ww_{M^*} + \delta^* w^2) + \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(e(M^*), \delta^*, U). \quad (16)$$

It is not difficult to see that $f^p(M^*, \delta^*) = \infty$ if either $\delta^* > 0$ or $\delta^* = 0$, and $w_{M^*} \neq 0$. Moreover, if $\delta^* < 0$, the supremum in the first term on the right-hand side of (16) is attained at $w = -w_{M^*}/\delta^*$. Thus

$$f^p(M^*, \delta^*) = -\frac{w_{M^*}^2}{\delta^*} + \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(e(M^*), \delta^*, U).$$

Finally, the third condition in (14) follows directly from (16), and this finishes the proof. \blacksquare

The following proposition characterizes the polyconvex biconjugate function of symmetric function, that is of the form $f(M) = f(e(M))$.

Proposition 6.3. *Let $f : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$. If $f(M) = f(e(M))$ for every $M \in \mathbb{R}^{2 \times 2}$, then $f^{pp}(M) = f^{pp}(e(M))$ and*

$$f^{pp}(M) = \sup_{V \in \mathbb{R}_{sym}^{2 \times 2}, \delta^* \leq 0} \left\{ \langle e(M), V \rangle + \delta^* \det e(M) - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U) \right\}.$$

Proof. Since $f(M) = f(e(M))$, $f^p(M^*, \delta^*)$ is given by (14). Hence,

$$(f^p)^*(M, \delta) = \max\{g_1(M), g_2(M, \delta)\}$$

where

$$g_1(M) = \sup_{V \in \mathbb{R}_{sym}^{2 \times 2}} \left\{ \langle e(M), V \rangle - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, 0, U) \right\}, \text{ and}$$

$$g_2(M, \delta) = \sup_{\substack{w \in \mathbb{R}, \delta^* < 0 \\ V \in \mathbb{R}_{sym}^{2 \times 2}}} \left\{ \langle e(M), V \rangle + 2w_M w + \frac{w^2}{\delta^*} + \delta \delta^* - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U) \right\},$$

with $\mathcal{L}(V, \delta^*, U)$ as in (15). Now, the function

$$w \mapsto \langle e(M), V \rangle + 2w_M w + \frac{w^2}{\delta^*} + \delta \delta^* - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U)$$

is concave in w for every $\delta^* < 0$. Thus,

$$2w_M w + \frac{w^2}{\delta^*} + \langle e(M), V \rangle + \delta \delta^* - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U) \leq \langle e(M), V \rangle + (\delta - w_M^2) \delta^* - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U),$$

with equality at $w = -\delta^* w_M$. Therefore,

$$(f^p)^*(M, \delta) = \sup_{V \in \mathbb{R}_{sym}^{2 \times 2}, \delta^* \leq 0} \left\{ \langle e(M), V \rangle + (\delta - w_M^2) \delta^* - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U) \right\}.$$

Next, we notice that $w_M^2 = \det w(M)$. Thus, since $f^{pp}(M) = (f^p)^*(e(M), \det e(M))$, by remark 4.1 (b) we obtain

$$f^{pp}(M) = \sup_{V \in \mathbb{R}_{sym}^{2 \times 2}, \delta^* \leq 0} \left\{ \langle e(M), V \rangle + \delta^* \det e(M) - \sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U) \right\}.$$

Hence, $f^{pp}(M) = f^{pp}(e(M))$ as claimed. \blacksquare

7 The polyconvex conjugate and biconjugate functions of f_C

In this section we specialize our results to a particular function f_C that we use in our proofs. Let $f : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ be such that

$$f_C(M) = \chi_{\bar{B}}(e(M)) (|\det e(M)| + |\langle C, M \rangle|), \quad (17)$$

where $C \in \mathbb{R}_{sym}^{2 \times 2}$ with negative determinant, and

$$\chi_{\bar{B}}(U) = \begin{cases} 1 & \text{if } U \in \mathbb{R}_{sym}^{2 \times 2} \setminus L^e(\mathcal{U}) \\ 0 & \text{otherwise.} \end{cases}$$

The following lemmas will be used to determine the polyconvex conjugate and biconjugate functions of f_C given as in eq. (17). Notice that f_C , restricted to $\text{Aff}(\mathcal{U})$, vanishes on $L^e(\mathcal{U})$. From now on, we denote by \mathcal{V} any three-well set where $\text{Aff}(\mathcal{V})$ is a two-dimensional subspace, and by \mathcal{U} any three-well set where $\text{Aff}(\mathcal{U})$ has codimension one.

Lemma 7.1. *Let $\mathcal{V} = \{V_1, V_2, V_3\} \subset \mathbb{R}_{sym}^{2 \times 2}$ such that $\det(V_2) = \det(V_3) = 0$, and $\text{Aff}(\mathcal{V}) \subset \mathbb{R}_{sym}^{2 \times 2}$ is a two-dimensional subspace. Also, assume that either \mathcal{V} is type one, or \mathcal{V} is type two and $0 \in C(\mathcal{V})$. Additionally, let f be defined as in eq. (17) for a fix $C \in \mathbb{R}_{sym}^{2 \times 2}$ with negative determinant, and $\mathcal{L}(V, \delta, U)$ be defined as in eq. (15). Then*

$$\sup_{U \in \mathbb{R}_{sym}^{2 \times 2}} \mathcal{L}(V, \delta^*, U) = \begin{cases} \infty & \text{if } (V, \delta^*) \in \mathcal{N}^c, \\ \max_{U \in \mathcal{V} \cup \{0\}} (k \langle C, U \rangle + \delta^* \det U) & \text{if } (V, \delta^*) \in \mathcal{N}, \end{cases} \quad (18)$$

where \mathcal{N} and \mathcal{N}^c are subsets of $\mathbb{R}_{sym}^{2 \times 2} \times (-\infty, 0]$ given by

$$\mathcal{N} = \{(kC, \delta^*) \mid -1 < k < 1, -1 \leq \delta^* \leq 0\} \quad \text{and} \quad \mathcal{N}^c = (\mathbb{R}_{sym}^{2 \times 2} \times (-\infty, 0]) \setminus \mathcal{N}.$$

Proof. We observe that $\text{Aff}(\mathcal{V}) = \Pi_Q(0)$, for some $Q \in \mathbb{R}_{sym}^{2 \times 2}$ with negative determinant since there are a compatible and an incompatible pair of wells in $\text{Aff}(\mathcal{V})$ (see remark 4.3). We divide the proof into two parts corresponding to the following claims. First, we claim that

$$\sup_{U \in \mathbb{R}_{sym}^{2 \times 2} \setminus L^e(\mathcal{V})} \mathcal{L}(V, \delta^*, U) = \begin{cases} 0 & \text{if } (V, \delta^*) \in \mathcal{N}, \\ \infty & \text{otherwise.} \end{cases} \quad (19)$$

Second, we claim that if $(V, \delta^*) \in \mathcal{N}$, then

$$\sup_{U \in L^e(\mathcal{V})} \mathcal{L}(kC, \delta^*, U) = \begin{cases} \max_{U \in \mathcal{V}} (k \langle C, U \rangle + \delta^* \det U) & \text{if } \mathcal{U} \text{ is type one,} \\ \max_{U \in \mathcal{V} \cup \{0\}} (k \langle C, U \rangle + \delta^* \det U) & \text{if } \mathcal{U} \text{ is type two.} \end{cases} \quad (20)$$

Assuming the claims, we finish the proof. Let $L := \sup\{\mathcal{L}(V, \delta^*, U) \mid U \in \mathbb{R}_{sym}^{2 \times 2}\}$ for each $(V, \delta^*) \in \mathbb{R}_{sym}^{2 \times 2} \times (-\infty, 0] = \mathcal{N} \cup \mathcal{N}^c$. Hence, if $(V, \delta^*) \in \mathcal{N}^c$, then $L = \infty$ by eq. (19), and if $(V, \delta^*) \in \mathcal{N}$, then

$$L = \max \left\{ \sup\{\mathcal{L}(V, \delta^*, U) \mid U \in L^e(\mathcal{V})\}, \sup\{\mathcal{L}(V, \delta^*, U) \mid U \in \mathbb{R}_{sym}^{2 \times 2} \setminus L^e(\mathcal{V})\} \right\}.$$

Therefore, the second part of eq. (18) is a consequence of eq. (19) and eq. (20), and the results follows.

Next, we divide the proof of the claims in two steps:

Step 1: We will prove eq. (19). Since $U \notin L^e(\mathcal{V})$, then

$$\mathcal{L}(e(M^*), \delta^*, U) = \min\{(\delta^* - 1) \det U, (\delta^* + 1) \det U\} + \min\{\langle e(M^*) - C, U \rangle, \langle e(M^*) + C, U \rangle\}.$$

We notice that

$$\mathcal{N}^c = \{(V, \delta^*) \in \mathbb{R}_{sym}^{2 \times 2} \times (-\infty, 0] \mid \delta^* < -1 \text{ or } V \not\parallel C \text{ or } V = kC, |k| > 1\}.$$

Now, we inspect each option for $(V, \delta^*) \in \mathcal{N}^c$.

- (i) Assume that $\delta^* < -1$. Since $(\delta^* - 1), (\delta^* + 1) < 0$, we choose $U(t) = t\bar{U}$ for a fix $\bar{U} \in \mathbb{R}_{sym}^{2 \times 2} \setminus L^e(\mathcal{V})$ with $\det \bar{U} < 0$, and $t \in \mathbb{R}$. We find

$$\mathcal{L}(e(M^*), \delta^*, U(t)) = t^2(\delta^* - 1) \det \bar{U} + t \min\{\langle e(M^*) - C, \bar{U} \rangle, \langle e(M^*) + C, \bar{U} \rangle\}.$$

By letting $t \rightarrow \infty$ the claim follows.

- (ii) Now, we assume $e(M^*)$ and C linearly independent. Thus, there exists $V \in \mathbb{R}_{sym}^{2 \times 2}$ with $\langle V, C \rangle = 0$ such that $e(M^*) = \alpha V + kC$ for some $\alpha, k \in \mathbb{R}$. Thus, $V \in \Pi_C(0)$. Due to $\det C < 0$ and Lemma 4.5, the boundary of $\Pi_C(0) \cap \mathcal{C}_0$ consist of two nonparallel rank-one lines, say $\ell_1 = \{tu_1 \otimes u_1 \mid t \in \mathbb{R}\}$ and $\ell_2 = \{tu_2 \otimes u_2 \mid t \in \mathbb{R}\}$ for some $u_1, u_2 \in S^1$. Therefore, we choose $\bar{U} \in \{u_1 \otimes u_1, -u_1 \otimes u_1, u_2 \otimes u_2, -u_2 \otimes u_2\}$ such that $\alpha \langle V, \bar{U} \rangle > 0$. By construction, $\det \bar{U} = 0$ and $\langle C, \bar{U} \rangle = 0$, hence $\mathcal{L}(e(M^*), \delta^*, t\bar{U}) = t\alpha \langle V, \bar{U} \rangle$ and by letting $t \rightarrow \infty$ we get the result.

- (iii) If $e(M^*) = kC$ with $|k| > 1$, then we choose $\bar{U} \in \mathcal{C}_0 \setminus \Pi_C(0)$ such that $k \langle C, \bar{U} \rangle > 0$ and $\det \bar{U} = 0$. Hence, $\mathcal{L}(e(M^*), \delta^*, t\bar{U}) = \min\{t(k-1) \langle C, \bar{U} \rangle, t(k+1) \langle C, \bar{U} \rangle\}$. Clearly, $(k-1) \langle C, \bar{U} \rangle > 0$, and $(k+1) \langle C, \bar{U} \rangle > 0$, so by letting $t \rightarrow \infty$ the result follows.

From the last three steps, we conclude that $\sup\{\mathcal{L}(V, \delta^*, U) \mid U \in \mathbb{R}_{sym}^{2 \times 2} \setminus L^e(\mathcal{V})\} = \infty$, if $(V, \delta^*) \in \mathcal{N}^c$.

Now, we assume $(V, \delta^*) \in \mathcal{N}$, and we prove that $\sup\{\mathcal{L}(e(M^*), \delta^*, U) \mid U \in \mathbb{R}_{sym}^{2 \times 2} \setminus L^e(\mathcal{V})\} = 0$. Indeed, we have that

$$\mathcal{L}(kC, \delta^*, U) = \min\{(\delta^* - 1) \det U, (\delta^* + 1) \det U\} + \min\{(k-1) \langle C, U \rangle, (k+1) \langle C, U \rangle\} \leq 0. \quad (21)$$

The last inequality follows since both terms on the right hand side of (21) are non-positive. Thus if $\bar{U} \in \mathcal{C}_0 \setminus \Pi_Q(0)$ is fixed, then we get

$$\mathcal{L}(kC, \delta^*, t\bar{U}) = \min\{t(k-1) \langle C, \bar{U} \rangle, t(k+1) \langle C, \bar{U} \rangle\} \rightarrow 0, \quad \text{as } t \rightarrow 0.$$

Therefore, the affirmation follows and the claim (i.e. eq. (19)) is proved.

Step 2: Now, we consider the optimization of \mathcal{L} on $U \in L^e(\mathcal{V})$ (i.e. eq. (20)). By lemma 4.5, for every $U \in L^e(\mathcal{V}) \subset \Pi_Q(0)$ there exist $\xi, \eta \in \mathbb{R}$ such that $U = \xi a^\perp \otimes a^\perp + \eta n^\perp \otimes n^\perp$, and

$$\mathcal{L}(kC, \delta^*, U) = k \langle C, U \rangle + \delta^* \det U = \xi \eta \delta^* |n \times a|^2 + \xi C a^\perp \cdot a^\perp + \eta C n^\perp \cdot n^\perp. \quad (22)$$

The level sets of (22) as function of ξ and η are hyperbolae (or straight lines). Hence, the supremum of $\mathcal{L}(kC, \delta^*, U)$ in $L^e(\mathcal{V})$ is attained at some point \bar{U} on the relative boundary of $L^e(\mathcal{V})$, denoted by $\partial_{ri} L^e(\mathcal{V})$. By Proposition 2.1, $\partial_{ri} L^e(\mathcal{V})$ depends on the type of \mathcal{V} , namely

$$\partial_{ri} L^e(\mathcal{V}) = \begin{cases} L^e(\mathcal{V}) & \text{if } \mathcal{V} \text{ is type one,} \\ C(\{V_1, V_2\}) \cup C(\{V_2, 0\}) \cup C(\{0, V_3\}) \cup C(\{V_3, V_1\}) & \text{if } \mathcal{V} \text{ is type two.} \end{cases}$$

First, we assume that \mathcal{V} is type two. If $\bar{U} \in C(\{V_2, 0\}) \cup C(\{0, V_3\})$, then $\det \bar{U} = 0$ and the equation (22) becomes linear. Hence, the maximum of $\mathcal{L}(kC, \delta^*, U)$ in $C(\{V_2, 0\}) \cup C(\{0, V_3\})$ is attained at either 0, V_2 or V_3 . Now, if $U \in C(\{V_1, V_2\})$, then $U = [\lambda\xi + (1-\lambda)\gamma]a^\perp \otimes a^\perp + \lambda\eta n^\perp \otimes n^\perp$ for some $\lambda \in [0, 1]$ due to corollary 4.0.1. So, eq. (22) becomes a polynomial of degree two in λ . It is readily seen that

$$\frac{d^2 \mathcal{L}(kC, \delta^*, U(\lambda))}{d\lambda^2} = 2\delta^* \eta (\xi - \gamma) |n \times a|^2 \geq 0.$$

Thus, either $\mathcal{L}(kC, \delta^*, U(\lambda))$ is linear in λ if $\xi = \gamma$ or it is quadratic in λ with a minimum if $\xi \neq \gamma$. Therefore, the maximum value of $\mathcal{L}(kC, \delta^*, U)$ on $C(\{V_1, V_2\})$ is always attained at the extremal points $\{V_1, V_2\}$. With a similar argument, the supremum of $\mathcal{L}(kC, \delta^*, U)$ on $C(\{V_1, V_3\})$ is attained on either V_1 or V_3 . Therefore, we conclude that the supremum of $\mathcal{L}(kC, \delta^*, U)$ in $L^e(\mathcal{V})$ is attained at some matrix $\bar{U} \in \{0, V_1, V_2, V_3\}$ as claimed.

Second, we assume that \mathcal{V} is of type one. In this case, the maximum value of $\mathcal{L}(kC, \delta^*, U)$ is attained either in V_1 or in $C(\{V_2, V_3\})$. If the second option happens, $d^2 \mathcal{L}/d\lambda^2 = -2\delta^* \gamma \zeta |n \times a|^2 \geq 0$, and by an analogous argument, the result follows straight forward.

Therefore, the supremum of $\mathcal{L}(kC, \delta^*, U)$ in $L^e(\mathcal{V})$ is attained at some matrix $\bar{U} \in \{V_1, V_2, V_3\}$ and eq. (20) follows. Now the proof is completed. \blacksquare

The last result of this section contains explicit bounds for $\text{Ker } f^{pp} \subset \mathbb{R}^{2 \times 2}$, the zero-level set of the polyconvex biconjugate function of $f : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$. From now on, if $g : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ is any function of the form $g(M) = g(e(M))$ for every $M \in \mathbb{R}_{sym}^{2 \times 2}$, we denote the zero-level set of $g|_{\mathbb{R}_{sym}^{2 \times 2}}$ as

$$\text{Ker}_s g := \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid g(V) = 0\}.$$

In this type of functions it readily follows that $\text{Ker } g = \{M \in \mathbb{R}^{2 \times 2} \mid e(M) \in \text{Ker}_s g\}$. In particular, if we study $\text{Ker } f^{pp}$, we can focus on $\text{Ker}_s f^{pp}$ and no information is lost.

Lemma 7.2. *Let $\mathcal{V} = \{V_1, V_2, V_3\} \subset \mathbb{R}_{sym}^{2 \times 2}$ be either type one or two such that $\text{Aff}(\mathcal{V}) \subset \mathbb{R}_{sym}^{2 \times 2}$ is a two-dimensional subspace with $\det V_2 = \det V_3 = 0$. Let $C \in \mathbb{R}_{sym}^{2 \times 2}$ with negative determinant and define $f(M)$ as in eq. (17). We have that:*

1. *If C is normal to $\text{Aff}(\mathcal{V})$, then $f^{pp}(M) \geq 0$ for every $M \in \mathbb{R}^{2 \times 2}$, and*

$$\text{Ker } f^{pp} = \{M \in \mathbb{R}^{2 \times 2} \mid e(M) \in \text{Aff}(\mathcal{V}), \det V_1 \leq \det e(M)\}. \quad (23)$$

2. *If $0 \in C(\mathcal{V})$, C satisfy $\langle C, V_1 \rangle \leq 0 < \langle C, V_3 \rangle = \langle C, V_2 \rangle$, and*

(i) *\mathcal{V} is type one, then*

$$\begin{aligned} \text{Ker } f^{pp} \subset \{M \in \mathbb{R}^{2 \times 2} \mid \langle C, e(M) - V_2 \rangle \leq 0 \leq \langle C, e(M) - V_1 \rangle, \\ 0 \leq \det e(M) \text{ and } 0 \leq \hbar(M)\}, \end{aligned}$$

(ii) *\mathcal{V} is type two, then*

$$\begin{aligned} \text{Ker } f^{pp} \subset \{M \in \mathbb{R}^{2 \times 2} \mid \langle C, e(M) - V_2 \rangle \leq 0 \leq \langle C, e(M) - V_1 \rangle, \\ \det V_1 \leq \det e(M) \text{ and } 0 \leq \hbar(M)\}. \end{aligned}$$

where

$$\hbar(M) = \langle C, e(M) - V_2 \rangle \det V_1 - \langle C, V_1 - V_2 \rangle \det e(M).$$

Moreover, f^{pp} is a non negative real-valued function and $\mathcal{V} \subset \text{Ker}_s f^{pp}$.

Proof. From proposition 6.3 and lemma 7.1 we have that

$$f^{pp}(M) = \sup_{(k, \delta^*) \in \mathcal{E}} \left(k \langle e(M), C \rangle + \delta^* \det e(M) - \max_{U \in \mathcal{V} \cup \{0\}} (k \langle C, U \rangle + \delta^* \det U) \right), \quad (24)$$

where $\mathcal{E} := [-1, 1] \times [-1, 0]$. Our goal is to estimate outer bounds on $\text{Ker}_s f^{pp}$.

First, we assume that C is normal to $\text{Aff}(\mathcal{V})$. Under these assumptions and since $0 \in \text{Aff}(\mathcal{V})$, eq. (24) becomes

$$f^{pp}(V) = \sup_{(k, \delta^*) \in \mathcal{E}} (k \langle V, C \rangle + \delta^* (\det V - \det V_1)).$$

By computing the supremum, we get

$$f^{pp}(V) = |\langle V, C \rangle| + \max\{0, \det V_1 - \det V\}. \quad (25)$$

The first term on the right hand side of (25) penalizes the distance to the plane $\text{Aff}(\mathcal{V})$, meanwhile the second term is an in-plane condition. Moreover, f^{pp} is non negative and

$$\text{Ker}_s f^{pp} = \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid V \in \text{Aff}(\mathcal{V}), \det V_1 \leq \det V\},$$

as claimed in eq. (23).

Second, we assume $\langle C, V_1 \rangle \leq 0 < \langle C, V_2 \rangle = \langle C, V_3 \rangle$. A simple computation lead us to

$$k \langle C, V_2 \rangle > \delta^* \det V_1 + k \langle C, V_1 \rangle, \quad \text{for} \quad \frac{\det V_1}{\langle C, V_2 - V_1 \rangle} \delta^* \leq k \leq 1.$$

This observation implies that

$$\max_{i=0,1,2,3} \{\delta^* \det V_i + k \langle C, V_i \rangle\} = \begin{cases} k \langle C, V_2 \rangle, & \text{if } \frac{\det V_1}{\langle C, V_2 - V_1 \rangle} \delta^* \leq k \leq 1, \\ \delta^* \det V_1 + k \langle C, V_1 \rangle, & \text{otherwise.} \end{cases} \quad (26)$$

Now, we define the sets $\mathcal{D} = \{(k, \delta^*) \in \mathcal{E} : \delta^* \det V_1 / \langle C, V_2 - V_1 \rangle \leq k \leq 1\}$, and its complement

$$\mathcal{E} \setminus \mathcal{D} = \left\{ (k, \delta^*) \in \mathcal{E} \mid -1 \leq k \leq \min \left\{ 1, \frac{-\delta^* \det V_1}{\langle C, V_1 - V_2 \rangle} \right\}, \quad -1 \leq \delta^* \leq 0 \right\},$$

relative to \mathcal{E} , see Figure 4. Let $\chi_{\mathcal{D}} : \mathcal{E} \rightarrow \{0, 1\}$ and $\chi_{\mathcal{E} \setminus \mathcal{D}} : \mathcal{E} \rightarrow \{0, 1\}$ be the characteristic functions of \mathcal{D} and $\mathcal{E} \setminus \mathcal{D}$, respectively. From eq. (24) and eq. (26), we get

$$f^{pp}(V) = \sup \{ \bar{f}_1(V, k, \delta^*) + \bar{f}_2(V, k, \delta^*) \mid (k, \delta^*) \in \mathcal{E} \},$$

where

$$\begin{cases} \bar{f}_1(V, k, \delta^*) = \chi_{\mathcal{D}} (k \langle C, V - V_2 \rangle + \delta^* \det V), \\ \bar{f}_2(V, k, \delta^*) = \chi_{\mathcal{E} \setminus \mathcal{D}} (k \langle C, V - V_1 \rangle + \delta^* (\det V - \det V_1)). \end{cases}$$

Now, \mathcal{D} and $\mathcal{E} \setminus \mathcal{D}$ are disjoint sets, hence

$$f^{pp}(V) = \max\{f_1(V), f_2(V)\}, \quad \text{where} \quad \begin{cases} f_1(V) = \sup_{(k, \delta^*) \in \mathcal{D}} \bar{f}_1(V, k, \delta^*), \\ f_2(V) = \sup_{(k, \delta^*) \in \mathcal{E} \setminus \mathcal{D}} \bar{f}_2(V, k, \delta^*). \end{cases}$$

Notice that \bar{f}_1 is linear in (k, δ^*) . Thus, its supremum in \mathcal{D} is attained on $\partial \mathcal{D}$ for each $V \in \mathbb{R}_{sym}^{2 \times 2}$. Analogously, the supremum of \bar{f}_2 in $\mathcal{E} \setminus \mathcal{D}$ is attained on $\partial(\mathcal{E} \setminus \mathcal{D})$. The common boundary between




Figure 4 Two possible subdivisions of the set \mathcal{E} when \mathcal{V} is of type two. Left and right images correspond to the cases $\langle C, V_2 - V_1 \rangle > -\det V_1$ and $\langle C, V_2 - V_1 \rangle < -\det V_1$, respectively.

\mathcal{D} and $\mathcal{E} \setminus \mathcal{D}$ depends on $\det V_1$, particularly on its sign (see Figure 4). Therefore, the computation of f_1 and f_2 also depends on V_1 . Moreover, since $0 \in C(\mathcal{V})$ and $\det V_2 = \det V_3 = 0$, corollary 4.0.1 implies that $\det V_1$ is positive or negative if \mathcal{V} is of type one or two, respectively.

Now, we assume \mathcal{V} is type two. Thus, $\det V_1 \leq 0$ and we divide the computation of f_1 into four cases:

- (a) Assume $0 < \langle C, V - V_2 \rangle$ and $0 < \det V$, then the supremum is attained at $(k, \delta^*) = (1, 0)$, and $f_1(V) = \langle C, V - V_2 \rangle > 0$.
- (b) Assume $0 < \langle C, V - V_2 \rangle$ and $\det V \leq 0$. If $\det V_1 < 0$, then \bar{f}_1 's supremum is attained at $(k, \delta^*) = (1, \max\{-1, -\langle C, V_1 - V_2 \rangle / \det V_1\})$, and if $\det V_1 = 0$ then \bar{f}_1 's supremum is attained at $(k, \delta^*) = (1, -1)$. In either case,

$$f_1(V) = \langle C, V - V_2 \rangle - \frac{1}{\max\left\{1, \frac{\det V_1}{\langle C, V_1 - V_2 \rangle}\right\}} \det V > 0.$$

- (c) Assume $\langle C, V - V_2 \rangle \leq 0$ and $\det V \leq 0$. In this case, the supremum of \bar{f}_1 is attained on the line segment

$$\left\{ \left(-\frac{\det V_1}{\langle C, V_1 - V_2 \rangle} \delta^*, \delta^* \right) \in \mathcal{E} \mid \delta^* \in [-1, 0] \right\},$$

and the function \bar{f}_1 evaluated at this segment is

$$\bar{f}_1 \left(V, -\frac{\det V_1}{\langle C, V_1 - V_2 \rangle} \delta^*, \delta^* \right) = -\frac{\bar{h}(V)}{\langle C, V_1 - V_2 \rangle} \delta^*,$$

where $\bar{h}(V) = \langle C, V - V_2 \rangle \det V_1 - \langle C, V_1 - V_2 \rangle \det V$. The last equation is linear in δ^* and increasing or decreasing depending on the sign of $\bar{h}(V) / \langle C, V_1 - V_2 \rangle$. Hence, if $\bar{h}(V)$ is non negative, $f_1(V) = 0$, meanwhile for $\bar{h}(V) < 0$,

$$f_1(V) = \min \left\{ 1, \frac{\det V_1}{\langle C, V_1 - V_2 \rangle} \right\} \langle C, V - V_2 \rangle - \frac{1}{\max \left\{ 1, \frac{\det V_1}{\langle C, V_1 - V_2 \rangle} \right\}} \det V.$$

Inserting the definition of $\bar{h}(V)$ in the last equation we get

$$f_1(V) = \frac{\bar{h}(V)}{p} > 0, \quad \text{where } p = \begin{cases} \det V_1, & \text{if } 1 \leq \det V_1 / \langle C, V_1 - V_2 \rangle, \\ \langle C, V_1 - V_2 \rangle, & \text{otherwise.} \end{cases}$$

- (d) Let $\langle C, V - V_2 \rangle \leq 0$ and $0 < \det V$. Then, the supremum of f_1 is attained at $(k, \delta^*) = (0, 0)$ and $f_1(V) = 0$. In this case, we also have that $\bar{h}(V) > 0$.

Summarizing the above cases, we conclude that

$$\text{Ker}_s f_1 = \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle C, V - V_2 \rangle \leq 0 \text{ and either } \det V \leq 0 \leq \hbar(V), \text{ or } 0 < \det V\},$$

and we easily conclude

$$\text{Ker}_s f_1 \subset \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle C, V - V_2 \rangle \leq 0 \leq \hbar(V)\}. \quad (27)$$

Next, we compute f_2 . Again, we also have four possible further cases:

- (a) If $\langle C, V - V_1 \rangle < 0$ and $0 \leq \det V - \det V_1$, then the supremum of f_2 is attained at $(k, \delta^*) = (-1, 0)$ and $f_2(V) = -\langle C, V - V_1 \rangle > 0$.
- (b) If $\langle C, V - V_1 \rangle \leq 0$ and $\det V - \det V_1 < 0$, then the supremum is attained at $(k, \delta^*) = (-1, -1)$ and $f_2(V) = -\langle C, V - V_1 \rangle - \det V + \det V_1 > 0$.
- (c) If $0 < \langle C, V - V_1 \rangle$ and $\det V - \det V_1 < 0$, then the supremum of f_2 is attained at $(k, \delta^*) = (\min\{1, \det V_1 / \langle C, V_1 - V_2 \rangle\}, -1)$ and

$$f_2(V) = \min \left\{ 1, \frac{\det V_1}{\langle C, V_1 - V_2 \rangle} \right\} \langle C, V - V_1 \rangle - \det V + \det V_1 > 0.$$

- (d) If $0 \leq \langle C, V - V_1 \rangle$ and $0 \leq \det V - \det V_1$, then the supremum is attained at a point on the segment

$$\left\{ \left(\frac{-\det V_1}{\langle C, V_1 - V_2 \rangle} \delta^*, \delta^* \right) \in \mathcal{E} \mid \delta^* \in [-1, 0] \right\},$$

and

$$\bar{f}_2 \left(V, \frac{-\det V_1}{\langle C, V_1 - V_2 \rangle} \delta^*, \delta^* \right) = \frac{-\hbar(V)}{\langle C, V_1 - V_2 \rangle} \delta^*.$$

By linearity in k , if $-\hbar(V) / \langle C, V_1 - V_2 \rangle \geq 0$, then $f_2(V) = 0$. Meanwhile, for $\hbar(V) < 0$,

$$f_2(V) = \min \left\{ 1, \frac{\det V_1}{\langle C, V_1 - V_2 \rangle} \right\} \langle C, V - V_1 \rangle - \frac{1}{\max \left\{ 1, \frac{\det V_1}{\langle C, V_1 - V_2 \rangle} \right\}} (\det V - \det V_1).$$

From the definition of $\hbar(V)$, we get

$$f_2(M) = \frac{\hbar(V)}{p} > 0, \quad \text{where } p = \begin{cases} \det V_1, & \text{if } 1 \leq \det V_1 / \langle C, V_1 - V_2 \rangle, \\ \langle C, V_1 - V_2 \rangle, & \text{otherwise.} \end{cases}$$

Hence, from the above four cases, we conclude that

$$\text{Ker}_s f_2 = \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid 0 \leq \langle C, V - V_1 \rangle, \det V_1 \leq \det V, 0 \leq \hbar(V)\}. \quad (28)$$

Now, from the above analysis, we conclude that f_C^{pp} is nonnegative and $\text{Ker}_s f^{pp} = \text{Ker}_s f_1 \cap \text{Ker}_s f_2$. Hence, by (27) and (28), we get

$$\text{Ker}_s f^{pp} \subset \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle C, V - V_2 \rangle \leq 0 \leq \langle C, V - V_1 \rangle, \det V_1 \leq \det V, 0 \leq \hbar(V)\}, \quad (29)$$

that is item (ii). A similar relation is obtained when \mathcal{V} is type one. Under this assumption, we can assume $\det V_1 > 0$ and the functions f_1 and f_2 can be computed analogously. The main observation is that \mathcal{D} and $\mathcal{E} \setminus \mathcal{D}$ are different due to the sign of $\det V_1$, see fig. 5. In fact, f_1 and f_2 are real-valued




Figure 5 Two possible subdivisions of the set \mathcal{E} when \mathcal{V} is of type one. Left and right images correspond to the cases $\langle C, V_2 - V_1 \rangle < \det V_1$ and $\langle C, V_2 - V_1 \rangle > \det V_1$, respectively.

functions, such that

$$f_1 \geq 0, \quad \text{and} \quad \text{Ker}_s f_1 = \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle C, V - V_2 \rangle \leq 0, \leq \det V, 0 \leq \hbar(V)\},$$

meanwhile,

$$f_2 \geq 0, \quad \text{and} \quad \text{Ker}_s f_2 \subset \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid 0 \leq \langle C, V - V_1 \rangle, 0 \leq \hbar(V)\}.$$

The latter equations imply that f^{pp} is such that $f^{pp} \geq 0$ and

$$\text{Ker}_s f^{pp} \subset \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle C, V - V_2 \rangle \leq 0 \leq \langle C, V - V_1 \rangle, 0 \leq \det V, 0 \leq \hbar(V)\}. \quad (30)$$

That is item (i), and the proof is complete. ■

8 Proofs of main results

This section concerns the proofs of Theorems 2.1 and 2.2.

8.1 Proof of theorem 2.1.

Proof. We consider three cases:

- (a) If all the wells in \mathcal{U} are pairwise compatible Bhattacharya [4] p. 231, proved that $Q^e(\mathcal{U}) = L^e(\mathcal{U}) = C(\mathcal{U})$, and there is nothing to prove.
- (b) We assume that \mathcal{U} is type two. Then, there exist a rank-one compatible pair of wells. Without loss of generality, we assume that

$$\det(U_1 - U_2) = 0, \det(U_1 - U_3) \leq 0, \text{ and } \det(U_2 - U_3) > 0.$$

Hence, by lemma 4.5, there exists $U_0 \in C(\mathcal{U})$ such that $\det(U_2 - U_0) = \det(U_3 - U_0) = 0$. Moreover, $U_0 \in C(\{U_1, U_2\})$ and $\det(U_1 - U_0) = 0$. In this case, the translated set $\mathcal{V} = \mathcal{U} - U_0$ satisfies that $\text{Aff}(\mathcal{V}) \subset \mathbb{R}_{sym}^{2 \times 2}$ is a two-dimensional subspace, where $\det(V_i) = 0$ for $i = 1, 2, 3$. Hence, by the first part of lemma 7.2 (namely eq. (23)), we conclude that if f is given as in eq. (17), then f^{pp} is a non negative polyconvex function such that $\mathcal{V} \subset \text{Ker}_s f^{pp}$. Therefore, by (4),

$$L^e(\mathcal{V}) \subset Q^e(\mathcal{V}) \subset \text{Ker}_s f^{pp} \cap C(\mathcal{V}) = \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid V \in C(\mathcal{V}), 0 \leq \det V\} = C(\mathcal{V}) \cap \overline{\mathcal{C}_0},$$

where the notation $\overline{\mathcal{C}_0}$ stands for the closure of the incompatible cone at 0. Now, by proposition 2.1, we have that $L^e(\mathcal{V}) = C(\{0, V_2\}) \cup C(\{0, V_1, V_3\}) = C(\mathcal{V}) \cap \overline{\mathcal{C}_0}$, and the proof follows.

- (c) We assume that there is only one compatible (in fact rank-one compatible) pair in \mathcal{U} . In this case, \mathcal{U} is of type one. Without loss of generality, we have that

$$\det(U_2 - U_1) > 0, \det(U_3 - U_1) > 0 \text{ and } \det(U_3 - U_2) = 0. \quad (31)$$

Now we claim that there exists $U_0 \notin C(\mathcal{U})$ such that

$$\det(U_0 - U_1) = \det(U_0 - U_2) = \det(U_0 - U_3) = 0. \quad (32)$$

This claim follows by noticing that

$$t \mapsto \det(tU_3 + (1-t)U_2 - U_1) = \det(U_2 - U_1) + t \langle \mathcal{S}(U_3 - U_2), U_2 - U_1 \rangle$$

is linear in t and $\langle \mathcal{S}(U_3 - U_2), U_2 - U_1 \rangle \neq 0$, since $\text{Aff}(\mathcal{U})$ is of codimension one. Thus, there exists $t_0 \in \mathbb{R}$ such that (32) holds. Moreover, $t_0 \notin [0, 1]$, and $U_0 = t_0 U_3 + (1 - t_0) U_2 \notin C(\mathcal{U})$ by (31), see fig. 1.(d). Proceeding as before, we define $\mathcal{V} = \{V_1, V_2, V_3\} = \mathcal{U} - U_0$ where $V_i = U_i - U_0$ and $\det V_i = 0$ for $i \in \{1, 2, 3\}$. Hence, by eq. (23) and the first part of lemma 7.2, we conclude that if f is given as in eq. (17), then f^{pp} is a non negative polyconvex function such that $\mathcal{V} \subset \text{Ker}_s f^{pp}$, and

$$L^e(\mathcal{V}) \subset Q^e(\mathcal{V}) \subset \text{Ker}_s f^{pp} \cap C(\mathcal{V}) \subset C(\mathcal{V}) \cap \overline{C_0},$$

but $C(\mathcal{V}) \cap \overline{C_0} = \{V_3\} \cup C(\{V_1, V_2\}) = L^e(\mathcal{V})$ by proposition 2.1.(a), and the result follows. ■

Remark 8.1. *In the proof of theorem 2.2 we are interested in $\text{Ker } f^{pp} \cap C(\mathcal{V})$. Under the assumptions of lemma 7.2 item 2, we claim that the conditions $V \in C(\mathcal{V})$ and $0 \leq \hbar(V)$ are equivalent to $\theta_1 \det V_1 \leq \det V$ for some $\theta_1 \in [0, 1]$. Assume the claim for a moment. We notice that the latter equation imply that if \mathcal{V} is of type one or type two, then $0 < \det V_1$ and $0 \leq \det V$, or $\det V_1 < 0$ and $\det V_1 < \det V$, respectively. Hence, from (30) and (29), we get*

$$\text{Ker } f^{pp} \cap C(\mathcal{V}) \subset \{M \in C(\mathcal{V}) \mid \langle C, e(M) - V_2 \rangle \leq 0 \leq \langle C, e(M) - V_1 \rangle, 0 \leq \hbar(M)\},$$

for both cases. Now, we prove the claim. By assumption $\langle C, V_1 - V_2 \rangle \neq 0$, thus from $0 \leq \hbar(V)$ we get that

$$\frac{\langle C, V - V_2 \rangle}{\langle C, V_1 - V_2 \rangle} \det V_1 \leq \det V, \quad (33)$$

Now, since $V \in C(\mathcal{V})$, then $V = \theta_1 V_1 + \theta_2 V_2 + \theta_3 V_3$ for some $\theta_1, \theta_2, \theta_3 \in [0, 1]$ such that $\theta_1 + \theta_2 + \theta_3 = 1$. Thus, $\langle C, V - V_2 \rangle = \langle C, \theta_1 V_1 - (1 - \theta_2 - \theta_3) V_2 \rangle = \theta_1 \langle C, V_1 - V_2 \rangle$ due to $\langle C, V_2 \rangle = \langle C, V_3 \rangle$. Hence, eq. (33) implies one side of the claim, and θ_1 is the volume fraction of V_1 . The reverse implication follows simple algebraic manipulations.

8.2 Proof of theorem 2.2.

Proof. We notice that under the assumptions of theorem 2.2, lemma 4.5 guarantees the existence of $U_0 \in C(\mathcal{U})$ such that $\det(U_2 - U_0) = \det(U_3 - U_0) = 0$ and $\det(U_1 - U_0) \neq 0$. Moreover, the translated set $\mathcal{V} = \{V_1, V_2, V_3\} := \mathcal{U} - U_0$ satisfies that $0 \in C(\mathcal{V})$, $\det(V_2) = \det(V_3) = 0$, and $\text{Aff}(\mathcal{V})$ is a two dimensional subspace. Hence, by lemma 7.2, we obtain explicit bound on $Q^e(\mathcal{U})$ by intersecting $C(\mathcal{U})$ with $\text{Ker}_s f^{pp}$ for admissible matrices C . Then by proposition 6.1 and eq. (4),

$$Q^e(\mathcal{U}) \subset (\text{Ker}_s f^{pp} + U_0) \cap C(\mathcal{U}), \quad (34)$$

where $\text{Ker}_s f^{pp} + U_0$ stands for the translated set $\{U_0 + V \mid V \in \text{Ker}_s f^{pp}\}$. By remark 4.3, if $Q \in \mathbb{R}_{sym}^{2 \times 2}$ is orthogonal to $\text{Aff}(\mathcal{V})$, then $\det Q < 0$, and due to corollary 4.0.1,

$$V_1 = \xi a^\perp \otimes a^\perp + \eta n^\perp \otimes n^\perp, \quad V_2 = \gamma a^\perp \otimes a^\perp, \quad \text{and} \quad V_3 = \zeta n^\perp \otimes n^\perp, \quad (35)$$

for some linearly independent $a, n \in S^1$ and constants $\xi, \eta, \gamma, \zeta \in \mathbb{R}$ that satisfy (12).

Next, in order to apply lemma 7.2, we construct a proper matrix C . To this end, we denote by \tilde{V} the \mathbb{R}^3 representation of $V \in \mathbb{R}_{sym}^{2 \times 2}$ under the isomorphism (6), and let

$$\tilde{C}_0 = \tilde{Q} \times (\tilde{V}_2 - \tilde{V}_3) \quad \text{where} \quad \tilde{Q} = \frac{(\tilde{V}_2 - \tilde{V}_1) \times (\tilde{V}_3 - \tilde{V}_1)}{\|(\tilde{V}_2 - \tilde{V}_1) \times (\tilde{V}_3 - \tilde{V}_1)\|}.$$

By construction, the matrix C_0 belongs to the affine space $\text{Aff}(\mathcal{V})$, and it is perpendicular to $V_3 - V_2$. After some direct algebraic manipulations, we get

$$\tilde{C}_0 = \frac{(\tilde{V}_2 - \tilde{V}_1) \langle \tilde{V}_3 - \tilde{V}_1, \tilde{V}_3 - \tilde{V}_2 \rangle + (\tilde{V}_3 - \tilde{V}_1) \langle \tilde{V}_2 - \tilde{V}_1, \tilde{V}_2 - \tilde{V}_3 \rangle}{|P|}, \quad (36)$$

where

$$P^2 := \|(\tilde{V}_2 - \tilde{V}_1) \times (\tilde{V}_3 - \tilde{V}_1)\|^2 = \|\tilde{V}_2 - \tilde{V}_1\|^2 \|\tilde{V}_3 - \tilde{V}_1\|^2 - \langle \tilde{V}_2 - \tilde{V}_1, \tilde{V}_3 - \tilde{V}_1 \rangle^2.$$

Notice that, since the isomorphism (6) preserves the inner product, we can drop the tildes out in eq. (36) to get the corresponding expression in $\mathbb{R}_{sym}^{2 \times 2}$. In terms of the coordinates given by (35), we get that

$$P^2 = (\eta\gamma + \xi\zeta - \gamma\zeta)^2 (1 - (a \cdot n)^4),$$

and

$$C_0 = - \left(\frac{[(\zeta - \gamma(a \cdot n)^2) a^\perp \otimes a^\perp + (\gamma - \zeta(a \cdot n)^2) n^\perp \otimes n^\perp]}{\sqrt{1 - (a \cdot n)^4}} \right) \text{sign}(\eta\gamma + \xi\zeta - \gamma\zeta).$$

By eq. (12) and corollary 4.0.1, we get

$$\text{sign}(\eta\gamma + \xi\zeta - \gamma\zeta) = \begin{cases} -1 & \mathcal{U} \text{ is type one,} \\ 1 & \mathcal{U} \text{ is type two.} \end{cases}$$

Thus,

$$\det C_0 = \frac{\gamma\zeta(1 + (a \cdot n)^2)^2 - (\gamma + \zeta)^2(a \cdot n)^2}{1 + (a \cdot n)^2}.$$

First, we assume \mathcal{U} is type two. Then, \mathcal{V} is also type two, and we let

$$C = \frac{1}{\sqrt{1 - (a \cdot n)^4}} C_0.$$

In this case, $\det C < 0$, and $\Pi_C \cap \text{Aff}(\mathcal{V})$ is a line, parallel to $V_2 - V_3$. Moreover, by an explicit computation and (12), we get

$$\langle C, V_2 \rangle = \langle C, V_3 \rangle = -\zeta\gamma > 0 \geq -(\xi\zeta + \eta\gamma) = \langle C, V_1 \rangle.$$

Therefore, the set \mathcal{V} and the matrix C meet necessary conditions to apply lemma 7.2 part (ii) and remark 8.1. Thus, we have

$$\text{Ker}_s f^{pp} = \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle C, V - V_2 \rangle \leq 0 \leq \langle C, V - V_1 \rangle, \quad 0 \leq \hbar(V)\}.$$

In view of eq. (34), we are interested in the set $\text{Ker}_s f^{pp} \cap C(\mathcal{V})$. We notice that, the sets $\{V \in \text{Aff}(\mathcal{V}) \mid \langle C, V - V_2 \rangle = 0\}$ and $\{V \in \text{Aff}(\mathcal{V}) \mid \langle C, V - V_3 \rangle = 0\}$ are supporting lines of $C(\mathcal{V})$ on V_2 and V_1 respectively. Hence,

$$Q^e(\mathcal{V}) \subset \text{Ker}_s f^{pp} \cap C(\mathcal{V}) = \{V \in C(\mathcal{V}) \mid 0 \leq \hbar(V)\}, \quad (37)$$

with

$$\hbar(V) = \langle C, V - V_2 \rangle \det V_1 - \langle C, V_1 - V_2 \rangle \det V. \quad (38)$$

Now, theorem 2.2 for \mathcal{U} of type two follows from the assignment $V_i = U_i + U_0$ and $V = U + U_0$.

Second, we assume \mathcal{U} is type one. Then, \mathcal{V} is also type one. If we let $Q \in \mathbb{R}_{sym}^{2 \times 2}$ be the orthogonal direction to $\text{Aff}(\mathcal{V})$, then we define $C(t) \in \mathbb{R}_{sym}^{2 \times 2}$ for each $t \in (-\pi, \pi]$ as

$$C(t) = Q \cos t + C \sin t,$$

where C is chosen as before in terms of C_0 . In this case, we can readily see that

$$\langle C(t), V_2 \rangle = \langle C(t), V_3 \rangle = \zeta \gamma \sin t, \quad \text{and} \quad \langle C(t), V_1 \rangle = (\xi \zeta + \eta \gamma) \sin t.$$

Thus, due to the continuity of $t \mapsto \det C(t)$ and remark 4.3, there exists $t_0 > 0$ small enough such that $\det C(t_0) < 0$ and $\langle C(t_0), V_1 \rangle \leq 0 < \langle C(t_0), V_3 \rangle = \langle C(t_0), V_2 \rangle$. Hence, by lemma 7.2 (i) and remark 8.1, we have that

$$\text{Ker}_s f^{pp} = \{V \in \mathbb{R}_{sym}^{2 \times 2} \mid \langle C(t_0), V - V_2 \rangle \leq 0 \leq \langle C(t_0), V - V_1 \rangle, \quad 0 \leq \kappa(V)\},$$

where $\kappa(V) = \langle C(t_0), V - V_2 \rangle \det V_1 - \langle C(t_0), V_1 - V_2 \rangle \det V$. By a similar argument as before, we conclude that

$$Q^e(\mathcal{V}) \subset \text{Ker}_s f^{pp} \cap C(\mathcal{V}) = \{V \in C(\mathcal{V}) \mid 0 \leq \kappa(V)\}.$$

Next, since $V \in C(\mathcal{V})$, it follows that $\kappa(V) = \sin t_0 \hbar(V)$ where \hbar is defined by eq. (38). Thus, we conclude that

$$Q^e(\mathcal{V}) \subset \{V \in C(\mathcal{V}) \mid 0 \leq \hbar(V)\}, \quad (39)$$

and theorem 2.2 follows from eq. (34). Therefore, we finish the proof of our theorem. \blacksquare

Remark 8.2. *The component-wise representation of eq. (37) and eq. (39) is straight forward. Indeed, if $V \in \text{Aff}(\mathcal{V})$, then $V = xa^\perp \otimes a^\perp + yn^\perp \otimes n^\perp$ for some real x, y , and $\hbar(V) = |n \times a|^2 h(x, y)$, where*

$$h(x, y) = xy(\eta\gamma + (\xi - \gamma)\zeta) - \xi\eta(\gamma y + (x - \gamma)\zeta).$$

Therefore, eq. (37) is rewritten as

$$Q^e(\mathcal{V}) \subset \left\{ V \in C(\mathcal{V}) \mid V = xa^\perp \otimes a^\perp + yn^\perp \otimes n^\perp, \quad 0 \leq h(x, y) \right\}.$$

Meanwhile, eq. (39) is rewritten as

$$Q^e(\mathcal{V}) \subset \left\{ V \in C(\mathcal{V}) \mid U = xa^\perp \otimes a^\perp + yn^\perp \otimes n^\perp, \quad h(x, y) \leq 0 \right\}.$$

9 Optimality of the bound for $Q^e(\mathcal{U})$ by symmetric quadratic polyconvex functions

As a final remark, we comment briefly about the optimality of the bounds in theorem 2.2. By the characterization of symmetric quadratic polyconvex functions given by Boussaid *et. al* [8] (proposition 4.5 in page 435), we can prove that our outer bound for $Q^e(\mathcal{U})$ is optimal when we restrict the analysis to quadratic functions.

Let ν be a homogeneous Young measure limit of linear strains supported on $\mathcal{V} = \mathcal{U} - U_0$, *i.e.* $\nu = \theta_1\delta_{V_1} + \theta_2\delta_{V_2} + \theta_3\delta_{V_3}$ with barycenter $V = \theta_1V_1 + \theta_2V_2 + \theta_3V_3$. Since quasiconvex functions preserves Jensen's inequality almost everywhere in Ω , see for example lemma 1.1 in [7] p. 850, we have that

$$f\left(\int_{\mathbb{R}^{2 \times 2}_{sym}} A d\nu(A)\right) \leq \int_{\mathbb{R}^{2 \times 2}_{sym}} f(A) d\nu(A).$$

Now, by proposition 4.5 in [8], $f : \mathbb{R}^{2 \times 2}_{sym} \rightarrow \mathbb{R}$ is symmetric quadratic polyconvex if and only if it has the form $f(\cdot) = g(\cdot) - \alpha \det(\cdot)$ for some convex function $g : \mathbb{R}^{2 \times 2}_{sym} \rightarrow \mathbb{R}$ and $\alpha > 0$. Therefore f is also symmetric quasiconvex. Hence,

$$g(V) - \alpha \det(V) \leq \sum_{i=1}^3 \theta_i g(V_i) - \alpha \sum_{i=1}^3 \theta_i \det(V_i).$$

Since, under the assumptions of theorem 2.2 and the definition of U_0 , $\theta_2 \det V_2 = \theta_3 \det V_3 = 0$. Therefore,

$$\det(V) - \theta_1 \det(V_1) \geq \sup \left\{ \frac{1}{\alpha} \left(g(V) - \sum_{i=1}^3 \theta_i g(V_i) \right) \mid g \text{ is convex, } \alpha > 0 \right\}, \quad (40)$$

for every $V \in Q^e(\mathcal{V})$. The supremum in the right hand side of eq. (40) is attained by the convex function $g = 0$. Hence, $\det(V) - \theta_1 \det(V_1) \geq 0$ and by remark 8.1, we recover $V \in C(\mathcal{V})$ and $0 \leq \hbar(V)$. Therefore, we obtain an outer bound for $Q^e(\mathcal{V})$ given by

$$\left\{ V = \theta_1 V_1 + \theta_2 V_2 + \theta_3 V_3 \mid \det(V) \geq \theta_1 \det(V_1), (\theta_1, \theta_2, \theta_3) \in [0, 1]^3, \sum_{i=1}^3 \theta_i = 1 \right\} \quad (41)$$

The bound in eq. (41) is equivalent to the outer in theorem 2.2.

10 Acknowledgements

The authors kindly acknowledges the support given by UNAM PAPPIT-IN106118 grant. LM acknowledge the support of CONACyT grant-590176 during his graduate studies and AC was also partially founded by CONACYT CB-2016-01-284451 grant.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1] J. M. Ball and R. D. James. Fine phase mixtures as minimizers of energy. In *Analysis and Continuum Mechanics*, pages 647–686. Springer, 1989. 2
- [2] J. M. Ball and R. D. James. Proposed experimental tests of a theory of fine microstructure and the two-well problem. *Philosophical Transactions of the Royal Society of London. Series A: Physical and Engineering Sciences*, 338(1650):389–450, 1992. 1, 2
- [3] B. Benešová and M. Kružík. Weak lower semicontinuity of integral functionals and applications. *SIAM Review*, 59(4):703–766, 2017. 1
- [4] K. Bhattacharya. Comparison of the geometrically nonlinear and linear theories of martensitic transformation. *Continuum Mechanics and Thermodynamics*, 5(3):205–242, 1993. 2, 5, 23
- [5] K. Bhattacharya and G. Dolzmann. Relaxation of some multi-well problems. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 131(2):279–320, 2001. 3
- [6] K. Bhattacharya et al. *Microstructure of martensite: why it forms and how it gives rise to the shape-memory effect*, volume 2. Oxford University Press, 2003. 3
- [7] K. Bhattacharya, N. B. Firoozye, R. D. James, and R. V. Kohn. Restrictions on microstructure. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 124(5):843–878, 1994. 2, 3, 5, 11, 27
- [8] O. Boussaid, C. Kreisbeck, and A. Schlömerkemper. Characterizations of symmetric polyconvexity. *Archive for Rational Mechanics and Analysis*, 234(1):417–451, 2019. 3, 4, 8, 27
- [9] A. Capella and L. Morales. Restriction on microstructures for n affine wells in two dimensional linear elasticity. *arXiv preprint arXiv:2012.06626*, 2020. 8
- [10] B. Dacorogna. *Direct methods in the calculus of variations*, volume 78. Springer Science & Business Media, 2007. 1, 3, 11, 14
- [11] G. Dolzmann. Numerical computation of rank-one convex envelopes. *SIAM journal on numerical analysis*, 36(5):1621–1635, 1999. 3
- [12] D. Kinderlehrer. Remarks about equilibrium configurations of crystals. *IMA preprint series*, 289, 1987. 2
- [13] R. V. Kohn. The relaxation of a double-well energy. *Continuum Mechanics and Thermodynamics*, 3(3):193–236, 1991. 1, 2, 3
- [14] R. V. Kohn and G. Strang. Optimal design and relaxation of variational problems, i. *Communications on pure and applied mathematics*, 39(1):113–137, 1986. 14
- [15] M. Kružík and M. Luskin. The computation of martensitic microstructure with piecewise laminates. *Journal of scientific computing*, 19(1):293–308, 2003. 3
- [16] C. B. Morrey et al. Quasi-convexity and the lower semicontinuity of multiple integrals. *Pacific journal of mathematics*, 2(1):25–53, 1952. 1
- [17] A. Růland. The cubic-to-orthorhombic phase transition: rigidity and non-rigidity properties in the linear theory of elasticity. *Archive for Rational Mechanics and Analysis*, 221(1):23–106, 2016. 2, 5

- [18] A. Schlömerkemper, I. V. Chenchiah, R. Fechte-Heinen, and D. Wachsmuth. Upper and lower bounds on the set of recoverable strains and on effective energies in cubic-to-monoclinic martensitic phase transformations. In *MATEC Web of Conferences*, volume 33, page 02011. EDP Sciences, 2015. [3](#), [4](#), [6](#), [8](#)
- [19] V. Smyshlyaev and J. Willis. On the relation of a three-well energy. *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 455(1983):779–814, 1999. [4](#)
- [20] V. Šverák. New examples of quasiconvex functions. *Archive for rational mechanics and analysis*, 119(4):293–300, 1992. [3](#)
- [21] V. Šverák. Rank-one convexity does not imply quasiconvexity. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 120(1-2):185–189, 1992. [3](#)
- [22] V. Šverák. On tartar’s conjecture. In *Annales de l’IHP Analyse non linéaire*, volume 10, pages 405–412, 1993. [2](#)
- [23] Q. Tang and K. Zhang. Bounds for effective strains of geometrically linear elastic multiwell model. *Journal of mathematical analysis and applications*, 339(2):1264–1276, 2008. [6](#)
- [24] L. Tartar. Some remarks on separately convex functions. In *Microstructure and phase transition*, pages 191–204. Springer, 1993. [2](#)
- [25] K. Zhang. On the structure of quasiconvex hulls. In *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*, volume 15, pages 663–686. Elsevier, 1998. [3](#)
- [26] K. Zhang. On equality of relaxations for linear elastic strains. *Communications on Pure & Applied Analysis*, 1(4):565, 2002. [4](#)