

**Obtaining a representation of the solution to the Cauchy problem for
equation high-order fractional derivative,
by the method of finding self-similar solutions**

B. Yu. Irgashev

*Namangan Engineering Construction Institute,
Namangan Branch of the Institute of Mathematics of the Republic of Uzbekistan.
E-mail: bahromirgasev@gmail.com*

Abstract. *In this paper, with the help of previously constructed self-similar solutions, a solution of the Cauchy problem for an equation of even order with a fractional Riemann-Liouville derivative of order $1 < \alpha < 2$ is obtained.*

Keywords. *Higher order equation. Fractional Riemann-Liouville derivative, self-similar solutions, fundamental solution, Cauchy problem.*

1. Construction of self-similar solutions

Consider the equation

$$L[u] \equiv D_{0y}^{\alpha} u(x, y) - dD_{0x}^{\beta} u(x, y) = 0, \quad (1)$$

here

$$q, p \in N, q - 1 < \alpha \leq q, p - 1 < \beta \leq p, q < p, d = \pm 1,$$

and $D_{0y}^{\alpha}, D_{0x}^{\beta}$ – Riemann-Liouville fractional differentiation operators, respectively, of orders α, β :

$$D_{0y}^{\alpha} u(x, y) = \frac{1}{\Gamma(q - \alpha)} \frac{d^q}{dy^q} \int_0^y \frac{u(x, \tau) d\tau}{(y - \tau)^{\alpha - q + 1}}, y > 0,$$

$$D_{0x}^{\beta} u(x, y) = \frac{1}{\Gamma(p - \beta)} \frac{d^p}{dx^p} \int_0^x \frac{u(\tau, y) d\tau}{(x - \tau)^{\beta - p + 1}}, x > 0.$$

Note that self-similar solutions of an equation with the usual derivative of the form

$$\frac{\partial^p u(x, y)}{\partial x^p} - \frac{\partial^q u(x, y)}{\partial y^q} = 0, p < q,$$

were constructed in [1].

In [2], using the methods of special operators, self-similar solutions of equation (1) were found in the cases $\frac{\beta}{2} \leq \alpha < \beta \leq 2, n - 1 < \beta \leq n \in N$.

In this paper, we construct self-similar solutions by a method that does not require knowledge of special operators.

We seek self-similar solutions to equation (1) in the form of the series

$$u(x, y) = y^b \sum_{n=0}^{\infty} c_n (x^a y^c)^{n+\gamma} = \sum_{n=0}^{\infty} c_n x^{an+a\gamma} y^{cn+c\gamma+b}, \quad (2)$$

here the parameters a, b, c, γ are to be defined.

We first introduce the notation

$$\overline{(a)}_s = a(a-1) \dots (a-(s-1)), \overline{(a)}_0 = 1, \overline{(a)}_1 = a.$$

Substitute (2) into (1). Then we will have

$$\begin{aligned}
D_{0y}^\alpha u(x, y) &= \frac{1}{\Gamma(q-\alpha)} \sum_{n=0}^{\infty} c_n x^{an+a\gamma} \frac{d^q}{dy^q} \int_0^y \frac{\tau^{cn+c\gamma+b} d\tau}{(y-\tau)^{\alpha-q+1}} = \\
&= \frac{1}{\Gamma(q-\alpha)} \sum_{n=0}^{\infty} c_n x^{an+a\gamma} \frac{d^q}{dy^q} \int_0^1 \frac{y^{cn+c\gamma+b+1} z^{cn+c\gamma+b} dz}{y^{\alpha-q+1} (1-z)^{\alpha-q+1}} = \\
&= \sum_{n=0}^{\infty} \overline{(cn+c\gamma+b-\alpha+q)_q} c_n (x^a y^c)^{n+\gamma} y^{b-\alpha} \frac{\Gamma(cn+c\gamma+b+1)}{\Gamma(cn+c\gamma+b-\alpha+q+1)}, \quad (3)
\end{aligned}$$

similarly

$$\begin{aligned}
D_{0x}^\beta u(x, y) &= \frac{1}{\Gamma(p-\beta)} \sum_{n=0}^{\infty} c_n y^{cn+c\gamma+b} \frac{d^p}{dx^p} \int_0^x \frac{\tau^{an+a\gamma} d\tau}{(x-\tau)^{\beta-p+1}} = \\
&= \frac{1}{\Gamma(p-\beta)} \sum_{n=0}^{\infty} c_n y^{cn+c\gamma+b} \frac{d^p}{dx^p} \int_0^x \frac{x^{an+a\gamma+1} z^{an+a\gamma} dz}{x^{\beta-p+1} (1-z)^{\beta-p+1}} = \\
&= \sum_{n=0}^{\infty} \overline{(an+a\gamma+p-\beta)_p} c_n (x^a y^c)^{n+\gamma} x^{-\beta} y^b \frac{\Gamma(an+a\gamma+1)}{\Gamma(an+a\gamma+1+p-\beta)}. \quad (4)
\end{aligned}$$

Substituting (3), (4) into (1), we obtain

$$\begin{aligned}
&\sum_{n=0}^{\infty} \overline{(cn+c\gamma+b-\alpha+q)_q} c_n (x^a y^c)^n x^\beta y^{-\alpha} \frac{\Gamma(cn+c\gamma+b+1)}{\Gamma(cn+c\gamma+b-\alpha+q+1)} = \\
&= d \sum_{n=0}^{\infty} \overline{(an+a\gamma+p-\beta)_p} c_n (x^a y^c)^n \frac{\Gamma(an+a\gamma+1)}{\Gamma(an+a\gamma+1+p-\beta)} \Rightarrow \\
&\quad a = \beta, c = -\alpha \Rightarrow \\
&\quad \sum_{n=0}^{\infty} c_n (x^\beta y^{-\alpha})^{n+1} \frac{\Gamma(-\alpha(n+\gamma)+b+1)}{\Gamma(-\alpha(n+\gamma+1)+b+1)} = \\
&= d \sum_{n=0}^{\infty} \overline{(\beta n + \beta\gamma + p - \beta)_p} c_n (x^a y^c)^n \frac{\Gamma(an+a\gamma+1)}{\Gamma(\beta n + \beta\gamma + 1 + p - \beta)} \Rightarrow
\end{aligned}$$

got conditions for the parameter γ :

$$\gamma_j = 1 - \frac{j}{\beta}, \quad j = 1, 2, \dots, p.$$

Substituting the found parameters into representation (2), we find the formula for the coefficients c_n . We have

$$\begin{aligned}
c_n \frac{\Gamma(\beta n + \beta\gamma + 1)}{\Gamma(\beta n + \beta\gamma + 1 - \beta)} &= d c_{n-1} \frac{\Gamma(-\alpha(n+\gamma-1)+b+1)}{\Gamma(-\alpha(n+\gamma)+b+1)} \Rightarrow \\
c_n &= d c_{n-1} \frac{\Gamma(-\alpha(n+\gamma-1)+b+1) \Gamma(\beta n + \beta\gamma + 1 - \beta)}{\Gamma(-\alpha(n+\gamma)+b+1) \Gamma(\beta n + \beta\gamma + 1)} = \\
&= d^n c_0 \frac{\Gamma(-\alpha\gamma + b + 1) \Gamma(\beta\gamma + 1)}{\Gamma(-\alpha(n+\gamma) + b + 1) \Gamma(\beta(n+\gamma) + 1)},
\end{aligned}$$

if now

$$c_0 = \frac{1}{\Gamma(-\alpha\gamma + b + 1)\Gamma(\beta\gamma + 1)},$$

then

$$\begin{aligned} c_n &= \frac{d^n}{\Gamma\left(-\alpha\left(n+1-\frac{j}{\beta}\right)+b+1\right)\Gamma\left(\beta\left(n+1-\frac{j}{\beta}\right)+1\right)} = \\ &= \frac{d^n}{\Gamma\left(-\alpha n - \alpha + \frac{\alpha}{\beta}j + b + 1\right)\Gamma(\beta n + \beta - j + 1)}. \end{aligned}$$

So we got the following self-similar solutions to equation (1):

$$\begin{aligned} u_j(x, y) &= y^b \left(x^\beta y^{-\alpha}\right)^{1-\frac{j}{\beta}} \sum_{n=0}^{\infty} \frac{(dx^\beta y^{-\alpha})^n}{\Gamma\left(-\alpha n - \alpha + \frac{\alpha}{\beta}j + b + 1\right)\Gamma(\beta n + \beta - j + 1)} = \\ &= y^b t^{1-\frac{j}{\beta}} W_{\left(-\alpha, -\alpha + \frac{\alpha}{\beta}j + b + 1\right), (\beta, \beta - j + 1)}(dt), \quad j = 1, 2, \dots, p, \quad t = x^\beta y^{-\alpha}, \quad (5) \end{aligned}$$

here

$$W_{(\mu, a), (\nu, b)}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\mu n + a)\Gamma(\nu n + b)}, \quad \mu, \nu \in R, \quad a, b \in C, \quad \mu + \nu > 0$$

- generalized Wright function [2]. Series (5) converges uniformly and it can be differentiated term by term, since $\beta - \alpha > 0$. Note that there is no condition on the b parameter. Note that solutions of the form (5) coincide exactly with the solutions obtained in [2].

Now let $\beta = p \in N$, then the solutions of equation (1) are expressions of the form:

$$u_s(x, y) = y^b \sum_{n=0}^{\infty} \frac{d^n \left(xy^{-\frac{\alpha}{p}}\right)^{pn+s}}{\Gamma\left(-\alpha n - \frac{\alpha}{p}s + b + 1\right)(pn+s)!}, \quad s = 0, 1, \dots, p-1.$$

Then their linear combination is also a solution to equation (1)

$$\begin{aligned} u(x, y) &= y^b \left(c_0 \sum_{n=0}^{\infty} \frac{d^n t^{pn}}{(pn)! \Gamma(-\alpha n + b + 1)} + c_1 \sum_{n=0}^{\infty} \frac{d^n t^{pn+1}}{(pn+1)! \Gamma\left(-\alpha\left(n+\frac{1}{p}\right)+b+1\right)} + \dots \right. \\ &\quad \left. + c_{p-1} \sum_{n=0}^{\infty} \frac{d^n t^{pn+p-1}}{(p(n+1)-1)! \Gamma\left(-\alpha\left(n+\frac{p-1}{p}\right)+b+1\right)} \right), \quad (6) \end{aligned}$$

where c_i - arbitrary real numbers, $i = 0, 1, \dots, p-1$.

Now let c_i , be such that $c_i \neq c_j$, for $i \neq j$ and $c_i^p = d$. Then from (6) we obtain

$$u(x, y) = y^b \phi\left(-\frac{\alpha}{p}, b+1, cxy^{-\frac{\alpha}{p}}\right), \quad c^p = 1, \quad (7)$$

here

$$\phi(-\delta, \varepsilon, z) = \sum_{k=0}^{\infty} \frac{z^k}{k! \Gamma(-\delta k + \varepsilon)}$$

- Wright function.

Consider the function

$$U(x - \xi, y - \eta) = (y - \eta)^b \phi\left(-\frac{\alpha}{p}, b+1, c(x - \xi)(y - \eta)^{-\frac{\alpha}{p}}\right), \quad y > \eta. \quad (8)$$

Let us note some properties of function (8) that are obtained by direct computation.

Лемма 1.

1. $D_{0y}^\gamma \left(y^b \phi \left(-\frac{\alpha}{p}, b+1, cxy^{-\frac{\alpha}{p}} \right) \right) = y^{b-\gamma} \phi \left(-\frac{\alpha}{p}, b+1-\gamma, cxy^{-\frac{\alpha}{p}} \right), \gamma \in R,$
2. $D_{y\eta}^\gamma \left((y-\eta)^b \phi \left(-\frac{\alpha}{p}, b+1, c(x-\xi)(y-\eta)^{-\frac{\alpha}{p}} \right) \right) =$
 $= (y-\eta)^{b-\gamma} \phi \left(-\frac{\alpha}{p}, b+1-\gamma, c(x-\xi)(y-\eta)^{-\frac{\alpha}{p}} \right), \gamma \in R,$
3. $\frac{\partial \gamma}{\partial x} \left(y^b \phi \left(-\frac{\alpha}{p}, b+1, c(x-\xi)y^{-\frac{\alpha}{p}} \right) \right) =$
 $= c^\gamma y^{b-\frac{\alpha}{p}\gamma} \phi \left(-\frac{\alpha}{p}, b+1-\frac{\alpha}{p}\gamma, c(x-\xi)y^{-\frac{\alpha}{p}} \right), \gamma \in \{0\} \cup N,$
4. $(D_{\eta y}^\alpha - d \frac{\partial^p}{\partial x^p}) D_{\eta y}^\gamma U(x-\xi, y-\eta) = 0,$
5. $(D_{y\eta}^\alpha - (-1)^p d \frac{\partial^p}{\partial \xi^p}) D_{y\eta}^\gamma U(x-\xi, y-\eta) = 0,$
6. $((-1)^q D_{y\eta}^{\alpha-q} \frac{\partial^q}{\partial \eta^q} - d(-1)^p \frac{\partial^p}{\partial \xi^p}) U(x-\xi, y-\eta) = 0, q-1 < \alpha \leq q.$

2. Cauchy problem

In this section, using the found particular solutions (8), we obtain an explicit form of the solution to the Cauchy problem for an equation of the form:

$$D_{0y}^\alpha u(x, y) - (-1)^{n-1} \frac{\partial^{2n} u(x, y)}{\partial x^{2n}} = f(x, y), \quad 1 < \alpha < 2. \quad (9)$$

Note that the Cauchy problem for higher order equations with a fractional derivative of order $0 < \alpha < 1$ was considered in [4], [5]. We will use ideas from these papers.

Following (7), we have

$$c_{1k}^{2n} = (-1)^{n-1} \Rightarrow c_{1k} = e^{\frac{n-1-2k}{2n}i\pi}, \quad k = \overline{0, (n-1)}, \quad \operatorname{Re} c_{1k} > 0,$$

Consider the function

$$\Gamma_b(x-\xi, y-\eta) = \begin{cases} \Gamma_b^1(x-\xi, y-\eta), & x > \xi, \\ \Gamma_b^2(x-\xi, y-\eta), & x < \xi, \end{cases} \quad (10)$$

where

$$\Gamma_b^1(x-\xi, y-\eta) = \frac{(y-\eta)^b}{2n} \sum_{k=0}^{n-1} \left(-e^{\frac{n-1-2k}{2n}i\pi} \right) \phi \left(-\frac{\alpha}{2n}, b+1, -e^{\frac{n-1-2k}{2n}i\pi} t \right),$$

$$\Gamma_b^2(x-\xi, y-\eta) = -\frac{(y-\eta)^b}{2n} \sum_{k=0}^{n-1} e^{\frac{n-1-2k}{2n}i\pi} \phi \left(-\frac{\alpha}{2n}, b+1, -e^{\frac{n-1-2k}{2n}i\pi} (-t) \right),$$

$$t = \frac{(x-\xi)}{(y-\eta)^{\frac{\alpha}{2n}}}.$$

We have

$$\frac{\partial^s \Gamma_b^1(x-\xi, y-\eta)}{\partial x^s} = \frac{1}{2n} \sum_{k=0}^{n-1} \left(-e^{\frac{n-1-2k}{2n}i\pi} \right) \frac{\partial^s \left((y-\eta)^b \phi \left(-\frac{\alpha}{2n}, b+1, -e^{\frac{n-1-2k}{2n}i\pi} t \right) \right)}{\partial x^s} =$$

$$= \frac{1}{2n} \sum_{k=0}^{n-1} \left(-e^{\frac{n-1-2k}{2n}i\pi} \right)^{s+1} (y-\eta)^{b-\frac{\alpha}{2n}s} \phi \left(-\frac{\alpha}{2n}, b+1-\frac{\alpha}{2n}s, -e^{\frac{n-1-2k}{2n}i\pi} t \right) \Rightarrow$$

$$\frac{\partial^s \Gamma_b^1(x-\xi, y-\eta)}{\partial x^s} \Big|_{x=\xi} = \frac{(y-\eta)^{b-\frac{\alpha}{2n}s}}{2n \Gamma(b+1-\frac{\alpha s}{2n})} \sum_{k=0}^{n-1} \left(-e^{\frac{n-1-2k}{2n}i\pi} \right)^{s+1}.$$

Similarly

$$\left. \frac{\partial^s \Gamma_b^2(x - \xi, y - \eta)}{\partial x^s} \right|_{x=\xi} = -\frac{(y - \eta)^{b - \frac{\alpha}{2n}s}}{2n\Gamma(b + 1 - \frac{\alpha s}{2n})} \sum_{k=0}^{n-1} \left(e^{\frac{n-1-2k}{2n}i\pi} \right)^{s+1}.$$

Hence we have if

$$s = (2n - 1) \pmod{2n},$$

then

$$\left. \frac{\partial^s \Gamma_b^1(x - \xi, y - \eta)}{\partial x^s} \right|_{x=\xi} - \left. \frac{\partial^s \Gamma_b^2(x - \xi, y - \eta)}{\partial x^s} \right|_{x=\xi} = (-1)^{n-1} \frac{(y - \eta)^{b - \frac{\alpha}{2n}s}}{\Gamma(b + 1 - \frac{\alpha s}{2n})},$$

if

$$s \neq (2n - 1) \pmod{2n},$$

then

$$\begin{aligned} & \left. \frac{\partial^s \Gamma_b^1(x - \xi, y - \eta)}{\partial x^s} \right|_{x=\xi} - \left. \frac{\partial^s \Gamma_b^2(x - \xi, y - \eta)}{\partial x^s} \right|_{x=\xi} = \\ & = \frac{(y - \eta)^{b - \frac{\alpha}{2n}s}}{2n\Gamma(b + 1 - \frac{\alpha}{2n})} \left(\sum_{k=0}^{n-1} \left(-e^{\frac{n-1-2k}{2n}i\pi} \right)^{s+1} + \sum_{k=0}^{n-1} \left(e^{\frac{n-1-2k}{2n}i\pi} \right)^{s+1} \right) = \\ & = \frac{(y - \eta)^{b - \frac{\alpha}{2n}s}}{2n\Gamma(b + 1 - \frac{\alpha}{2n})} e^{\frac{n-1}{2n}(s+1)i\pi} \sum_{k=0}^{n-1} e^{-\frac{k(s+1)}{n}i\pi} \left((-1)^{s+1} + 1 \right) = 0. \end{aligned}$$

So we have proved the following lemma.

Lemma 2. For $s \in N$, the following relation holds:

$$\begin{aligned} & \left. \frac{\partial^s \Gamma_b^1(x - \xi, y - \eta)}{\partial x^s} \right|_{x=\xi} - \left. \frac{\partial^s \Gamma_b^2(x - \xi, y - \eta)}{\partial x^s} \right|_{x=\xi} = \\ & = (-1)^{n-1} \frac{(y - \eta)^{b - \frac{\alpha}{2n}s}}{\Gamma(b + 1 - \frac{\alpha}{2n})} \begin{cases} 1, & s = (2n - 1) \pmod{2n}, \\ 0, & s \neq (2n - 1) \pmod{2n}. \end{cases} \end{aligned}$$

The results of Lemma 2 coincide with the results of [5]. In what follows, we need the asymptotics of the Wright function for large values of the variable. The main results on asymptotics were obtained by Wright [see 3]. In particular, the following theorem is proved.

Theorem 1 [see 3]. If $|\arg y| \leq \min \left\{ \frac{3}{2}\pi(1 - \sigma), \pi \right\} - \varepsilon$, $0 < \sigma < 1$, then

$$\phi(-\sigma, \beta, t) = Y^{\frac{1}{2} - \beta} e^{-Y} \left\{ \sum_{m=0}^{M-1} A_m Y^{-m} + O(Y^{-M}) \right\}, \quad |t| \rightarrow \infty, \quad (11)$$

here $Y = (1 - \sigma)(\sigma^\sigma y)^{\frac{1}{1-\sigma}}$, $y = -t$, the coefficients of A_m depend on σ, β .

If $n = 3, 4, \dots$, to $0 < \frac{\alpha}{2n} < \frac{1}{n} \leq \frac{1}{3}$, and the relation

$$|\arg y| = |\arg(-t)| = \left| \frac{n-1-2k}{2n}\pi \right| \leq \pi - \varepsilon, \quad k = 0, 1, \dots, n-1. \quad (12)$$

If $n = 2$, then for $0 < \frac{\alpha}{4} \leq \frac{1}{3}$, we have

$$|\arg y| = |\arg(-t)| = \left| \pm \frac{\pi}{4} \right| < \pi, \quad (13)$$

and at $\frac{1}{3} < \frac{\alpha}{4} < \frac{1}{2}$, get

$$|\arg y| = |\arg(-t)| = \left| \pm \frac{\pi}{4} \right| < \frac{3}{2}\pi \left(1 - \frac{\alpha}{4}\right). \quad (14)$$

Taking into account (12) - (14), we conclude that for $n = 2, 3, \dots$ we always have relation (11). Now we write (10) in the form

$$\Gamma_b(x - \xi, y - \eta) = \frac{(y - \eta)^b}{2n} \sum_{k=0}^{n-1} \left(-e^{\frac{n-1-2k}{2n}i\pi} \right) \phi \left(-\frac{\alpha}{2n}, b+1, -e^{\frac{n-1-2k}{2n}i\pi} |t| \right), \quad (15)$$

where

$$|t| = \frac{|x - \xi|}{(y - \eta)^{\frac{\alpha}{2n}}},$$

then, taking into account (11), (12) - (14), for large values of $|t|$, the estimate obtained in [5], [6] is valid for (15):

$$\left| \phi \left(-\frac{\alpha}{2n}, b+1, -e^{\frac{n-1-2k}{2n}i\pi} |t| \right) \right| \leq C |t|^{\frac{-2n}{2n-\alpha}(b+\frac{1}{2})} \exp \left(-\sigma |t|^{\frac{2n}{2n-\alpha}} \right), \quad (16)$$

here

$$\sigma = \left(1 - \frac{\alpha}{2n}\right) \left(\frac{\alpha}{2n}\right)^{\frac{\alpha}{2n-\alpha}} \cos \frac{n-1}{2n-\alpha} \pi, \quad 0 < C - \text{const},$$

or applying Lemma 1 (properties 1,3), for $x \neq 0$, $\gamma \in R$, $s \in N \cup \{0\}$, we obtain

$$\left| \frac{\partial^s}{\partial x^s} \left(D_{0y}^\gamma \Gamma_b(x, y) \right) \right| \leq M y^{b-\gamma-\frac{\alpha}{2n}(s-\theta)} |x|^{-\theta} \exp \left(-\sigma |x|^{\frac{2n}{2n-\alpha}} y^{-\frac{\alpha}{2n-\alpha}} \right), \quad (17)$$

here

$$|x| y^{-\frac{\alpha}{2n}} \rightarrow +\infty, \\ \theta = \frac{2n}{2n-\alpha} \left(\frac{1}{2} + b - \gamma - \frac{\alpha}{2n} s \right).$$

Note that estimate (17) coincides with the estimate obtained in [5], for $b = \alpha - 1 - \frac{\alpha}{2n}$.

Lemma 3. For $j, s \in N$ and any function $h(x) \in C(R)$ such that

$$|h(x)| \leq M \exp \left(c|x|^{\frac{2n}{2n-\alpha}} \right), \quad c < \sigma, \quad |x| \rightarrow \infty, \quad 0 < M - \text{const}, \quad (18)$$

equality holds

$$\lim_{y \rightarrow 0} \int_{-\infty}^{+\infty} h(\xi) D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = -h(x) \cdot \begin{cases} 0, & j \neq s, \\ 1, & j = s. \end{cases}$$

Proof. First, we calculate the integral

$$\begin{aligned} & \int_{-\infty}^{+\infty} D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = \int_{-\infty}^x D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi + \\ & \quad + \int_x^{+\infty} D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = \\ & = - \sum_{k=0}^{n-1} \frac{e^{\frac{n-1-2k}{2n}\pi i}}{2n} \int_{-\infty}^x D_{0y}^{\alpha-s} \left(y^{\alpha-\frac{\alpha}{2n}-j} \phi \left(-\frac{\alpha}{2n}, \alpha - \frac{\alpha}{2n} - j + 1, -e^{\frac{n-1-2k}{2n}\pi i} (x-\xi) y^{-\frac{\alpha}{2n}} \right) \right) d\xi - \\ & - \sum_{k=0}^{n-1} \frac{e^{\frac{n-1-2k}{2n}\pi i}}{2n} \int_x^{+\infty} D_{0y}^{\alpha-s} \left(y^{\alpha-\frac{\alpha}{2n}-j} \phi \left(-\frac{\alpha}{2n}, \alpha - \frac{\alpha}{2n} - j + 1, e^{\frac{n-1-2k}{2n}\pi i} (x-\xi) y^{-\frac{\alpha}{2n}} \right) \right) d\xi = \end{aligned}$$

$$\begin{aligned}
&= -\sum_{k=0}^{n-1} \frac{e^{\frac{n-1-2k}{2n}\pi i}}{2n} \int_{-\infty}^x y^{s-\frac{\alpha}{2n}-j} \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} (x-\xi) y^{-\frac{\alpha}{2n}}\right) d\xi \\
&\quad -\sum_{k=0}^{n-1} \frac{e^{\frac{n-1-2k}{2n}\pi i}}{2n} \int_x^{+\infty} y^{s-\frac{\alpha}{2n}-j} \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, e^{\frac{n-1-2k}{2n}\pi i} (x-\xi) y^{-\frac{\alpha}{2n}}\right) d\xi.
\end{aligned}$$

Now let's calculate each integral separately

$$\begin{aligned}
&-e^{\frac{n-1-2k}{2n}\pi i} \int_{-\infty}^x y^{s-\frac{\alpha}{2n}-j} \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} (x-\xi) y^{-\frac{\alpha}{2n}}\right) d\xi \\
&= -e^{\frac{n-1-2k}{2n}\pi i} \sum_{m=0}^{\infty} y^{s-\frac{\alpha}{2n}-j} \int_{-\infty}^x \frac{\left(-e^{\frac{n-1-2k}{2n}\pi i} y^{-\frac{\alpha}{2n}} (x-\xi)\right)^m}{m! \Gamma\left(-\frac{\alpha}{2n}m + s - \frac{\alpha}{2n} - j + 1\right)} d\xi = \\
&= -y^{s-j} \sum_{m=0}^{\infty} \frac{\left(-e^{\frac{n-1-2k}{2n}\pi i} y^{-\frac{\alpha}{2n}} (x-\xi)\right)^{m+1}}{(m+1)! \Gamma\left(-\frac{\alpha}{2n}m + s - \frac{\alpha}{2n} - j + 1\right)} \Bigg|_{\xi=-\infty}^{\xi=x} = \\
&= -y^{s-j} \left(\sum_{m=0}^{\infty} \frac{\left(-e^{\frac{n-1-2k}{2n}\pi i} y^{-\frac{\alpha}{2n}} (x-\xi)\right)^m}{m! \Gamma\left(-\frac{\alpha}{2n}m + s - j + 1\right)} - \frac{1}{\Gamma(s-j+1)} \right) \Bigg|_{\xi=-\infty}^{\xi=x} = \\
&= -y^{s-j} \left(\phi\left(-\frac{\alpha}{2n}, s-j+1, -e^{\frac{n-1-2k}{2n}\pi i} y^{-\frac{\alpha}{2n}} (x-\xi)\right) - \frac{1}{\Gamma(s-j+1)} \right) \Bigg|_{\xi=-\infty}^{\xi=x} = \\
&= -\frac{y^{s-j}}{\Gamma(s-j+1)}.
\end{aligned}$$

Similarly

$$-e^{\frac{n-1-2k}{2n}\pi i} \int_x^{+\infty} y^{s-\frac{\alpha}{2n}-j} \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, e^{\frac{n-1-2k}{2n}\pi i} (x-\xi) y^{-\frac{\alpha}{2n}}\right) d\xi = -\frac{y^{s-j}}{\Gamma(s-j+1)}.$$

from here we get

$$\int_{-\infty}^{+\infty} D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = -\frac{y^{s-j}}{\Gamma(s-j+1)}. \quad (19)$$

Note that for $s < j$ we have

$$\int_{-\infty}^{+\infty} D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = 0, \quad s < j. \quad (20)$$

Now

$$\begin{aligned}
\int_{-\infty}^{+\infty} h(\xi) D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi &= \int_{-\infty}^x h(\xi) D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi + \\
&+ \int_x^{+\infty} h(\xi) D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = I_1 + I_2,
\end{aligned}$$

We calculate each term separately

$$\begin{aligned}
I_1 &= \int_{-\infty}^x h(\xi) D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = \\
&= - \sum_{k=0}^{n-1} e^{\frac{n-1-2k}{2n}\pi i} y^{s-\frac{\alpha}{2n}-j} \int_{-\infty}^x h(\xi) \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} y^{-\frac{\alpha}{2n}}(x-\xi)\right) d\xi = \\
&= \left(t = (x-\xi) y^{-\frac{\alpha}{2n}}\right) = \\
&= - \sum_{k=0}^{n-1} e^{\frac{n-1-2k}{2n}\pi i} y^{s-j} \int_0^{+\infty} h\left(x-ty^{\frac{\alpha}{2n}}\right) \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} t\right) d\xi = \\
&= - \sum_{k=0}^{n-1} e^{\frac{n-1-2k}{2n}\pi i} y^{s-j} \int_0^{+\infty} \left(h\left(x-ty^{\frac{\alpha}{2n}}\right) - h(x)\right) \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} t\right) d\xi + \\
&\quad + h(x) \left(- \sum_{k=0}^{n-1} e^{\frac{n-1-2k}{2n}\pi i} y^{s-j} \int_0^{+\infty} \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} t\right) d\xi\right) = \\
&= J_1 + J_2,
\end{aligned}$$

Hence we have

$$\lim_{y \rightarrow +0} I_1 = \lim_{y \rightarrow +0} J_1 + \lim_{y \rightarrow +0} J_2.$$

Let's find each limit separately. Using (19), (20), we have

$$\lim_{y \rightarrow +0} J_2 = -h(x) \lim_{y \rightarrow +0} \frac{y^{s-j}}{2\Gamma(s-j+1)} = \begin{cases} -\frac{1}{2}h(x), & s = j, \\ 0, & s \neq j. \end{cases}$$

Further, if $s \neq j$, then $\lim_{y \rightarrow +0} J_1 = 0$. Let $s = j$, then

$$\begin{aligned}
&\lim_{y \rightarrow +0} \int_0^{+\infty} \left(h\left(x-ty^{\frac{\alpha}{2n}}\right) - h(x)\right) \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} t\right) d\xi = \\
&= \lim_{y \rightarrow +0} \int_0^N \left(h\left(x-ty^{\frac{\alpha}{2n}}\right) - h(x)\right) \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} t\right) d\xi + \\
&\quad + \lim_{y \rightarrow +0} \int_N^{+\infty} \left(h\left(x-ty^{\frac{\alpha}{2n}}\right) - h(x)\right) \phi\left(-\frac{\alpha}{2n}, s-\frac{\alpha}{2n}-j+1, -e^{\frac{n-1-2k}{2n}\pi i} t\right) d\xi = 0,
\end{aligned}$$

here $0 < N$. The last equality holds due to the fact that for an arbitrary small number $\epsilon > 0$, you can choose N so that the sublimit expressions will be less than $1/2\epsilon$ (the first expression, due to the continuity the function $h(x)$, and the second is due to estimates (16) and (18)).

Means

$$\lim_{y \rightarrow +0} I_1 = \begin{cases} -\frac{1}{2}h(x), & s = j, \\ 0, & s \neq j. \end{cases}$$

It is shown similarly that

$$\lim_{y \rightarrow +0} I_2 = \begin{cases} -\frac{1}{2}h(x), & s = j, \\ 0, & s \neq j. \end{cases}$$

So finally we have

$$\lim_{y \rightarrow 0} \int_{-\infty}^{+\infty} h(\xi) D_{0y}^{\alpha-s} \Gamma_{\alpha-\frac{\alpha}{2n}-j}(x-\xi, y) d\xi = -h(x) \cdot \begin{cases} 0, & j \neq s, \\ 1, & j = s. \end{cases}$$

Lemma 3 is proved.

Cauchy problem. Find the solution $u(x, y)$ to equation (9) in the region $D = \{(x, y) : -\infty < x < +\infty, 0 < y\}$, satisfying the following conditions:

- 1) $D_{0y}^{\alpha} u(x, y), \frac{\partial^{2n} u(x, y)}{\partial x^{2n}} \in C(D), D_{0y}^{\alpha-1} u(x, y), D_{0y}^{\alpha-2} u(x, y) \in C(\overline{D})$;
- 2) $\lim_{y \rightarrow +0} D_{0y}^{\alpha-1} u(x, y) = \varphi(x), \lim_{y \rightarrow +0} D_{0y}^{\alpha-2} u(x, y) = \psi(x)$.

The specified functions satisfy the constraints:

$$\begin{aligned} f(x, y) &\in C(\overline{D}), \varphi(x), \psi(x) \in C(R), \\ |\varphi(x)| &\leq M \exp\left(k|x|^{\frac{2n}{2n-\alpha}}\right), |x| \rightarrow +\infty, \\ |\psi(x)| &\leq M \exp\left(k|x|^{\frac{2n}{2n-\alpha}}\right), |x| \rightarrow +\infty, \\ |f(x, y)| &\leq M \exp\left(k|x|^{\frac{2n}{2n-\alpha}}\right), |x| \rightarrow +\infty, \\ k &< \sigma, 0 < M - \text{constant}. \end{aligned}$$

Theorem 2. The solution to the Cauchy problem has the form

$$\begin{aligned} u(x, y) &= - \int_{-\infty}^{+\infty} \varphi(\xi) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y) d\xi - \int_{-\infty}^{+\infty} \psi(\xi) \Gamma_{\alpha-\frac{\alpha}{2n}-2}(x-\xi, y) d\xi - \\ &\quad - \int_0^y \int_{-\infty}^{+\infty} f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\xi d\eta. \end{aligned}$$

Proof. It is easy to verify that the first two terms satisfy homogeneous equation (9). Consider the third term

$$\begin{aligned} &D_{0y}^{\alpha} \left(\int_0^y \int_{-\infty}^{+\infty} f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\xi d\eta \right) = \\ &= \frac{1}{\Gamma(2-\alpha)} \frac{\partial^2}{\partial y^2} \left(\int_{-\infty}^{+\infty} d\xi D_{0y}^{\alpha-2} \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\eta \right) \right) \Rightarrow \\ &D_{0y}^{\alpha-2} \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\eta \right) = \\ &= \int_0^y \frac{d\tau}{(y-\tau)^{\alpha-1}} \left(\int_0^{\tau} f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, \tau-\eta) d\eta \right) = \\ &= \int_0^y f(\xi, \eta) d\eta \left(\int_{\eta}^y \frac{\Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, \tau-\eta)}{(y-\tau)^{\alpha-1}} d\tau \right) = \end{aligned}$$

$$\begin{aligned}
&= \int_0^y f(\xi, \eta) d\eta \left(\int_0^1 \frac{(y-\eta)^{2-\alpha} \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, (y-\eta)t)}{(1-t)^{\alpha-1}} dt \right) = \\
&= \Gamma(2-\alpha) \int_0^y f(\xi, \eta) \Gamma_{1-\frac{\alpha}{2n}}(x-\xi, y-\eta) d\eta. \tag{21}
\end{aligned}$$

Hence we have

$$\begin{aligned}
&D_{0y}^\alpha \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\eta \right) = \\
&= \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\eta \right). \tag{22}
\end{aligned}$$

Note that taking into account (21), we have

$$\begin{aligned}
&\lim_{y \rightarrow +0} D_{0y}^{\alpha-2} \left(\int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha-1-\frac{\alpha}{2n}}(x-\xi, y-\eta) d\eta \right) \right) = \\
&= \lim_{y \rightarrow +0} \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{1-\frac{\alpha}{2n}}(x-\xi, y-\eta) d\eta \right) = 0, \tag{23}
\end{aligned}$$

$$\begin{aligned}
&\lim_{y \rightarrow +0} D_{0y}^{\alpha-1} \left(\int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha-1-\frac{\alpha}{2n}}(x-\xi, y-\eta) d\eta \right) \right) = \\
&= \lim_{y \rightarrow +0} \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{-\frac{\alpha}{2n}}(x-\xi, y-\eta) d\eta \right) = 0. \tag{24}
\end{aligned}$$

From (23), (24) and Lemma 3 we have

$$\begin{aligned}
&\lim_{y \rightarrow +0} D_{0y}^{\alpha-2} u(x, y) = \varphi(x), \\
&\lim_{y \rightarrow +0} D_{0y}^{\alpha-1} u(x, y) = \psi(x).
\end{aligned}$$

Now we find the partial derivatives with respect to the variable x , the third term

$$\begin{aligned}
&\frac{\partial}{\partial x} \left(\int_{-\infty}^x d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\eta \right) \right) + \\
&+ \frac{\partial}{\partial x} \left(\int_x^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\eta \right) \right) = \\
&= \lim_{\xi \rightarrow x} \int_0^y f(\xi, \eta) \left(\Gamma_{\alpha-1-\frac{\alpha}{2n}}^1(x-\xi, y-\eta) - \Gamma_{\alpha-1-\frac{\alpha}{2n}}^2(x-\xi, y-\eta) \right) d\eta + \\
&+ \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial}{\partial x} \Gamma_{\alpha-\frac{\alpha}{2n}-1}(x-\xi, y-\eta) d\eta \right) =
\end{aligned}$$

$$= \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial}{\partial x} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right),$$

continuing this process we get

$$\begin{aligned} & \frac{\partial^{2n}}{\partial x^{2n}} \left(\int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right) \right) = \\ & = \lim_{\varepsilon \rightarrow +0} \left(\frac{\partial}{\partial x} \left(\int_{-\infty}^{x-\varepsilon} d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial^{2n-1}}{\partial x^{2n-1}} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right) \right) \right) + \\ & + \lim_{\varepsilon \rightarrow +0} \left(\frac{\partial}{\partial x} \left(\int_{x-\varepsilon}^{x+\varepsilon} d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial^{2n-1}}{\partial x^{2n-1}} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right) \right) \right) + \\ & + \lim_{\varepsilon \rightarrow +0} \left(\frac{\partial}{\partial x} \left(\int_{x+\varepsilon}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial^{2n-1}}{\partial x^{2n-1}} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right) \right) \right) = \\ & = I_1 + I_2 + I_3, \end{aligned}$$

We calculate each term separately

$$\begin{aligned} I_1 &= \lim_{\varepsilon \rightarrow +0} \left(\int_0^y f(x - \varepsilon, \eta) \frac{\partial^{2n-1}}{\partial x^{2n-1}} \Gamma_{\alpha - 1 - \frac{\alpha}{2n}}^1(\varepsilon, y - \eta) d\eta \right) + \\ & + \int_{-\infty}^x d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial^{2n}}{\partial x^{2n}} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right), \end{aligned}$$

from here

$$\begin{aligned} & \lim_{\varepsilon \rightarrow +0} \int_0^y f(x - \varepsilon, \eta) \frac{\partial^{2n-1}}{\partial x^{2n-1}} \Gamma_{\alpha - 1 - \frac{\alpha}{2n}}^1(\varepsilon, y - \eta) d\eta = \\ & = \frac{(-1)^{n-1}}{2n} \sum_{k=0}^{n-1} \lim_{\varepsilon \rightarrow +0} \int_0^y f(x - \varepsilon, \eta) (y - \eta)^{-1} \sum_{m=1}^{\infty} \frac{\left(-e^{\frac{n-1-2k}{2n}\pi i} \varepsilon (y - \eta)^{-\frac{\alpha}{2n}} \right)^k}{k! \Gamma\left(-\frac{\alpha}{2n}k\right)} d\eta = \\ & \left(z = \varepsilon (y - \eta)^{-\frac{\alpha}{2n}} \Rightarrow \eta = y - \varepsilon^{\frac{2n}{\alpha}} z^{-\frac{2n}{\alpha}}, \quad d\eta = \frac{2n}{\alpha} \varepsilon^{\frac{2n}{\alpha}} z^{-\frac{2n}{\alpha}-1} dz \right) \\ & = \frac{(-1)^{n-1}}{\alpha} f(x, y) \sum_{k=0}^{n-1} \sum_{m=1}^{\infty} \frac{\left(-e^{\frac{n-1-2k}{2n}\pi i} \right)^m}{m! \Gamma\left(-\frac{\alpha}{2n}m\right)} \int_0^{+\infty} z^{m-1} dz = \\ & = \frac{(-1)^n}{2n} f(x, y) \sum_{k=0}^{n-1} \left(\sum_{m=0}^{\infty} \frac{\left(-e^{\frac{n-1-2k}{2n}\pi i} \right)^m z^m}{m! \Gamma\left(-\frac{\alpha}{2n}m + 1\right)} - 1 \right) \Big|_{z=0}^{z=\infty} = \frac{(-1)^{n-1}}{2} f(x, y). \end{aligned}$$

Means

$$I_1 = \frac{(-1)^{n-1}}{2} f(x, y) + \int_{-\infty}^x d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial^{2n}}{\partial x^{2n}} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right).$$

Similarly

$$I_3 = \frac{(-1)^{n-1}}{2} f(x, y) + \int_x^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial^{2n}}{\partial x^{2n}} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right),$$

it is easy to show that

$$I_2 = 0.$$

so

$$\begin{aligned} & \frac{\partial^{2n}}{\partial x^{2n}} \left(\int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right) \right) = \\ & = (-1)^{n-1} f(x, y) + \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \frac{\partial^{2n}}{\partial x^{2n}} \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right) = \\ & = (-1)^{n-1} f(x, y) + (-1)^{n-1} \int_{-\infty}^{+\infty} d\xi \left(\int_0^y f(\xi, \eta) \Gamma_{-\frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\eta \right). \end{aligned} \quad (25)$$

From (22) and (25) we obtain that the expression

$$- \int_0^y \int_{-\infty}^{+\infty} f(\xi, \eta) \Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) d\xi d\eta,$$

satisfies equation (9). The theorem is proved.

The question of the uniqueness of the Cauchy problem is still open. Function

$$-\Gamma_{\alpha - \frac{\alpha}{2n} - 1}(x - \xi, y - \eta) \quad (26)$$

will be called the fundamental solution of equation (9). Representation (26) coincides with the form of the fundamental solution obtained for the case $0 < \alpha < 1$ in [5].

3. Application

In [7], for the equation of beam vibration

$$u_{tt}(t, x) + a^2 u_{xxxx}(t, x) = 0,$$

a representation of the solution of the Cauchy problem was obtained using the previously constructed fundamental solution, which for $a = 1$ has the form

$$\begin{aligned} G_2(x, t, \xi) &= \sqrt{\frac{t}{\pi}} \sin \left[\frac{(\xi - x)^2}{4t} + \frac{\pi}{4} \right] + \\ &+ \frac{\xi - x}{2} \left\{ S \left[\left(\frac{(\xi - x)^2}{4t} \right) \right] - C \left[\left(\frac{(\xi - x)^2}{4t} \right) \right] \right\}, \end{aligned}$$

where

$$S(z) = \frac{1}{2\sqrt{\pi}} \int_0^z \frac{\sin t}{\sqrt{t}} dt, \quad C(z) = \frac{1}{2\sqrt{\pi}} \int_0^z \frac{\cos t}{\sqrt{t}} dt -$$

Fresnel integrals.

Our fundamental solution (27), for $\alpha = 2$, $n = 2$, has the form (y was replaced by t , $\eta = 0$)

$$-\Gamma_{\frac{1}{2}}(x - \xi, t) = -\frac{\sqrt{t}}{4} \left(-e^{\frac{\pi}{4}i} \sum_{n=0}^{\infty} \frac{(-e^{\frac{\pi}{4}i})^n \tau^n}{n! \Gamma(-\frac{1}{2}n + \frac{3}{2})} - e^{-\frac{\pi}{4}i} \sum_{n=0}^{\infty} \frac{(-e^{-\frac{\pi}{4}i})^n \tau^n}{n! \Gamma(-\frac{1}{2}n + \frac{3}{2})} \right),$$

where

$$\tau = \frac{|x - \xi|}{\sqrt{t}},$$

show that

$$G_2(x, t, \xi) = -\Gamma_{\frac{1}{2}}(x - \xi, t).$$

Indeed, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\left(-e^{\frac{\pi}{4}i}\right)^n \tau^n}{n! \Gamma\left(-\frac{1}{2}n + \frac{3}{2}\right)} &= \sum_{n=0}^{\infty} \frac{\left(-e^{\frac{\pi}{4}i}\right)^{4n} \tau^{4n}}{(4n)! \Gamma\left(-2n + \frac{3}{2}\right)} + \sum_{n=0}^{\infty} \frac{\left(-e^{\frac{\pi}{4}i}\right)^{4n+1} \tau^{4n+1}}{(4n+1)! \Gamma\left(-2n+1\right)} + \\ &+ \sum_{n=0}^{\infty} \frac{\left(-e^{\frac{\pi}{4}i}\right)^{4n+2} \tau^{4n+2}}{(4n+2)! \Gamma\left(-2n + \frac{1}{2}\right)} + \sum_{n=0}^{\infty} \frac{\left(-e^{\frac{\pi}{4}i}\right)^{4n+3} \tau^{4n+3}}{(4n+3)! \Gamma\left(-2n\right)} = \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n}}{(4n)! \Gamma\left(-2n + \frac{3}{2}\right)} + i \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n+2}}{(4n+2)! \Gamma\left(-2n + \frac{1}{2}\right)} + \left(-e^{\frac{\pi}{4}i}\right) \tau = I_1. \end{aligned}$$

Similarly

$$\sum_{n=0}^{\infty} \frac{\left(-e^{-\frac{\pi}{4}i}\right)^n \tau^n}{n! \Gamma\left(-\frac{1}{2}n + \frac{3}{2}\right)} = \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n}}{(4n)! \Gamma\left(-2n + \frac{3}{2}\right)} - i \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n+2}}{(4n+2)! \Gamma\left(-2n + \frac{1}{2}\right)} - \tau e^{-\frac{\pi}{4}i} = I_2,$$

from here

$$\begin{aligned} -\Gamma_{\frac{1}{2}}(x - \xi, t) &= -\frac{\sqrt{t}}{4} \left(-e^{\frac{\pi}{4}i} I_1 - e^{-\frac{\pi}{4}i} I_2\right) = \\ &= \frac{\sqrt{2t}}{8} \left((I_1 + I_2) + i(I_1 - I_2)\right), \end{aligned}$$

further

$$\begin{aligned} I_1 + I_2 &= 2 \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n}}{(4n)! \Gamma\left(-2n + \frac{3}{2}\right)} - \sqrt{2} \tau, \\ (I_1 - I_2) i &= -2 \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n+2}}{(4n+2)! \Gamma\left(-2n + \frac{1}{2}\right)} + \sqrt{2} \tau \Rightarrow \\ -\Gamma_{\frac{1}{2}}(x - \xi, t) &= \frac{\sqrt{2t}}{4} \left(\sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n}}{(4n)! \Gamma\left(-2n + \frac{3}{2}\right)} - \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n+2}}{(4n+2)! \Gamma\left(-2n + \frac{1}{2}\right)} \right), \end{aligned}$$

further, calculations show that

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n}}{(4n)! \Gamma\left(-2n + \frac{3}{2}\right)} &= \frac{\Gamma\left(-\frac{1}{2}\right)}{\pi} {}_1F_2 \left[-\frac{1}{4}, -\left(\frac{\tau^2}{8}\right)^2 \right], \\ \sum_{n=0}^{\infty} \frac{(-1)^n \tau^{4n+2}}{(4n+2)! \Gamma\left(-2n + \frac{1}{2}\right)} &= \frac{\Gamma\left(-\frac{1}{2}\right) \tau^2}{4\pi} {}_1F_2 \left[\frac{1}{4}, -\left(\frac{\tau^2}{8}\right)^2 \right], \end{aligned}$$

where

$${}_1F_2 \left[\begin{matrix} a, z \\ b, c \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n (c)_n} \frac{z^n}{n!}$$

-hypergeometric function,

$$(d)_n = d(d+1) \dots (d+n-1),$$

from here

$$-\Gamma_{\frac{1}{2}}(x - \xi, t) = \frac{\sqrt{2t}\Gamma(-\frac{1}{2})}{16\pi} \left({}_4F_2 \left[\begin{matrix} -\frac{1}{4}, -\left(\frac{\tau^2}{8}\right)^2 \\ \frac{3}{4}, \frac{1}{2} \end{matrix} \right] - \tau^2 {}_1F_2 \left[\begin{matrix} \frac{1}{4}, -\left(\frac{\tau^2}{8t}\right)^2 \\ \frac{3}{2}, \frac{5}{4} \end{matrix} \right] \right).$$

Now, taking into account the relations (see [8])

$${}_4F_2 \left[\begin{matrix} -\frac{1}{4}, -\left(\frac{\tau^2}{8}\right)^2 \\ \frac{3}{4}, \frac{1}{2} \end{matrix} \right] = 4 \cos \left(\frac{\tau^2}{4} \right) + 2\sqrt{2\pi} |\tau| S \left(\frac{\tau^2}{4} \right),$$

$$\tau^2 {}_1F_2 \left[\begin{matrix} \frac{1}{4}, -\left(\frac{\tau^2}{8}\right)^2 \\ \frac{3}{2}, \frac{5}{4} \end{matrix} \right] = 2|\tau| \sqrt{2\pi} C \left(\frac{\tau^2}{4} \right) - 4 \sin \frac{\tau^2}{4},$$

finally get

$$-\Gamma_{\frac{1}{2}}(x - \xi, t) = \sqrt{\frac{t}{\pi}} \sin \left(\frac{\tau^2}{4} + \frac{\pi}{4} \right) + \frac{|\xi - x|}{2} \left(S \left(\frac{\tau^2}{4} \right) - C \left(\frac{\tau^2}{4} \right) \right),$$

i.e.

$$-\Gamma_{\frac{1}{2}}(x - \xi, t) = G_2(x, t, \xi).$$

References

1. Irgashev B. Y. Construction of Singular Particular Solutions Expressed via Hypergeometric Functions for an Equation with Multiple Characteristics. //Differential Equations. 2020. T. 56. No 3, pp. 315-323.
2. L u c h k o Yu., G o r e n f l o R. Scale-invariant solutions of a partial differential equation of fractional order. // Fractional Calculus and Applied Analysis.1998. T.1, No 1. pp. 63-78.
3. Wright E. M. The generalized Bessel function of order greater than one // Quart. J. Math., Oxford Ser., 1940. T. 1, No 11. pp. 36-48.
4. Pskhu A. Fundamental solutions and cauchy problems for an odd-order partial differential equation with. // Electronic Journal of Differential Equations, Vol. 2019 (2019), No. 21, pp. 1-13.
5. Karasheva L. L. Cauchy problem for high even order parabolic equation with time fractional derivative //Sibirskie Elektronnye Matematicheskie Izvestiya [Siberian Electronic Mathematical Reports].2018. T. 15. pp. 696-706. [In Russian].
6. Karasheva L. L. On Properties of an Integer Function That Generalizes the Wright Function //Itogi Nauki i Tekhniki. Seriya"Sovremennaya Matematika i ee Prilozheniya. Tematicheskie Obzory". 2018. T. 149. pp. 38-43.[In Russian]
7. Sabitov K. B. Cauchy problem for the beam vibration equation //Differential Equations. 2017. T. 53. No. 5. pp. 658-664.
8. Dunaev A.S., Shlychkov V.I. Hypergeometric functions. Ural Federal University. Yekaterinburg. 2017. P.1273 [in Russian].