THREE CHARACTERIZATIONS OF A SELF-SIMILAR APERIODIC 2-DIMENSIONAL SUBSHIFT

SÉBASTIEN LABBÉ

ABSTRACT. The goal of this chapter is to illustrate a generalization of the Fibonacci word to the case of 2-dimensional configurations on \mathbb{Z}^2 . More precisely, we consider a particular subshift of $\mathcal{A}^{\mathbb{Z}^2}$ on the alphabet $\mathcal{A} = \{0, \ldots, 15\}$ for which we give three characterizations: as the subshift \mathcal{X}_{Φ} generated by a 2-dimensional morphism Φ defined on \mathcal{A} ; as the Wang shift $\Omega_{\mathcal{Z}}$ defined by a set \mathcal{Z} of 16 Wang tiles; as the symbolic dynamical system $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}}$ representing the orbits under some \mathbb{Z}^2 -action $R_{\mathcal{Z}}$ defined by rotations on \mathbb{T}^2 and coded by some topological partition $\mathcal{P}_{\mathcal{Z}}$ of \mathbb{T}^2 into 16 polygonal atoms. We prove their equality $\Omega_{\mathcal{Z}} = \mathcal{X}_{\Phi} = \mathcal{X}_{\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}}$ by showing they are self-similar with respect to the substitution Φ .

This chapter provides a transversal reading of results divided into four different articles obtained through the study of the Jeandel-Rao Wang shift. It gathers in one place the methods introduced to desubstitute Wang shifts and to desubstitute codings of \mathbb{Z}^2 -actions by focussing on a simple 2-dimensional self-similar subshift. SageMath code to find marker tiles and compute the Rauzy induction of \mathbb{Z}^2 -rotations is provided allowing to reproduce the computations. The chapter contains many exercises whose solutions are provided at the end.

Contents

1.	Introduction	2
2.	Preliminaries	5
3.	An aperiodic self-similar subshift	9
4.	A self-similar Wang shift	21
5.	A self-similar symbolic dynamical system	31
6.	Conclusion	46
7.	Solutions to exercises	48
Re	eferences	63

²⁰²⁰ Mathematics Subject Classification. Primary 37B51; Secondary 52C23, 28D05.

Key words and phrases. aperiodic and subshift and Sturmian and tiling and substitution and self-similar and SFT and Markov partition and self-induced and coding of rotations and Rauzy induction and polyhedron exchange transformation.

The author acknowledges financial support from the Agence Nationale de la Recherche through the projects CODYS (ANR-18-CE40-0007), PARADIS (ANR-18-CE23-0007-01) and IZES (ANR-22-CE40-0011).

S. LABBÉ

1. INTRODUCTION

The rule $s : a \mapsto ab, b \mapsto a$ defines a morphism on the monoid $\{a, b\}^*$. The succesive application of this morphism on the letter *a* defines longer and longer words covering the negative and non-negative integers:

$s^0(a)$	$s^0(a)$		<u>a</u>	a
$s^1(a)$	$s^1(a)$		\underline{ab}	ab
$s^{2}(a)$	$s^{2}(a)$		$a\underline{b}a$	aba
$s^{3}(a)$	$s^3(a)$	=	$aba\underline{ab}$	abaab
$s^{4}(a)$	$s^4(a)$		$abaaba\underline{ba}$	a ba a b a b a
$s^5(a)$	$s^5(a)$		$abaababaaba\underline{ab}$	abaabaabaabaab

The letters that change from line to line are underlined. It is an interesting exercise to show that at the limit, we obtain $\lim_{n\to\infty} s^{2n}(a)|s^{2n}(a) = \tilde{F}ba|F$ and $\lim_{n\to\infty} s^{2n+1}(a)|s^{2n+1}(a) = \tilde{F}ab|F$ where F is the well-known right-infinite Fibonacci word [Ber80]. The rule s can be seen as a substitution that we may apply on the biinfinite words $x = \tilde{F}ba|F$ and $y = \tilde{F}ab|F$ and we observe that s(x) = y and s(y) = x. Thus $s^2(x) = x$ and $s^2(y) = y$ and we say that x and y are fixed points of s^2 . The set of finite words that appear in $x = s^2(x)$ defines a language

$$\mathcal{L}_s = \{\varepsilon, a, b, aa, ab, ba, aab, aba, baa, bab, aaba, \dots\} \subset \{a, b\}^*$$

and a subshift

 $\mathcal{X}_s = \{ u \in \{a, b\}^{\mathbb{Z}} \colon \text{the finite words that appear in } u \text{ are in } \mathcal{L}_s \}.$

The subshift \mathcal{X}_s contains x, y, all shifts of x and y, and much more. Indeed, \mathcal{X}_s is a Sturmian shift which is an uncountable set. The reader will find detailed information on Sturmian sequences in [Lot02, Chapter 2] and [Fog02, Chapter 6]. In particular, \mathcal{X}_s is aperiodic, that is, none of the sequences in \mathcal{X}_s is periodic.

It is known since the early work of Morse and Hedlund in [MH40] and Coven and Hedlund in [CH73] that the 1-dimensional subshift \mathcal{X}_s , being a Sturmian subshift, has many equivalent characterizations:

- as the subshift generated by the 1-dimensional substitution s;
- as the subshift on {a, b} having exactly n + 1 factors of length n and such that the ratio of the frequency of the two letters is φ = 1+√5/2;
 as the symbolic representation of a rotation on the 1-dimensional torus T = ℝ/Z
- as the symbolic representation of a rotation on the 1-dimensional torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ through a partition into two intervals.

The goal of this chapter is to illustrate a generalization of the above to the case of 2-dimensional configurations on \mathbb{Z}^2 . More precisely, we consider a particular subshift of $[[0, 15]]^{\mathbb{Z}^2}$, first considered in [Lep24], for which we give three characterizations:

• as the subshift \mathcal{X}_{Φ} generated by the 2-dimensional morphism Φ defined on the alphabet $\mathcal{A} = \llbracket 0, 15 \rrbracket$ by the rule

$$\Phi: [0, 15]] \to [0, 15]]^{*^{2}} \\ \begin{cases} 0 \mapsto (14), & 1 \mapsto (13), & 2 \mapsto (12, 10), & 3 \mapsto (11, 8), \\ 4 \mapsto (14, 7), & 5 \mapsto (13, 7), & 6 \mapsto (12, 7), & 7 \mapsto \begin{pmatrix} 6\\12 \end{pmatrix}, \\ 8 \mapsto \begin{pmatrix} 3\\14 \end{pmatrix}, & 9 \mapsto \begin{pmatrix} 3\\13 \end{pmatrix}, & 10 \mapsto \begin{pmatrix} 2\\12 \end{pmatrix}, & 11 \mapsto \begin{pmatrix} 6&1\\12&10 \end{pmatrix}, \\ 12 \mapsto \begin{pmatrix} 6&1\\11&8 \end{pmatrix}, & 13 \mapsto \begin{pmatrix} 5&1\\15&9 \end{pmatrix}, & 14 \mapsto \begin{pmatrix} 4&1\\11&8 \end{pmatrix}, & 15 \mapsto \begin{pmatrix} 2&0\\12&7 \end{pmatrix}. \end{cases}$$

• as the Wang shift $\Omega_{\mathcal{Z}}$, that is, the set of valid configuration $\mathbb{Z}^2 \to \llbracket 0, 15 \rrbracket$ of the plane, defined by the following set \mathcal{Z} of 16 Wang tiles:

$\begin{bmatrix} O\\J & 0 \\ O\\ O\end{bmatrix}$	$\begin{bmatrix} O \\ H \ 1 \\ L \end{bmatrix}$	$\begin{bmatrix} M \\ D \stackrel{2}{_{2}} J \\ P \end{bmatrix}$	$\begin{smallmatrix} M \\ D \ 3 \ D \\ K \end{smallmatrix}$	$\begin{bmatrix} P\\J \ 4 \\ P \end{bmatrix}$	$\begin{bmatrix} P\\H {5 \atop N} H\\N \end{bmatrix}$	${\begin{smallmatrix} K\\ D \ 6 \ H\\ P \end{smallmatrix}}$	$\begin{bmatrix} 0\\ I & 7 & B\\ 0 \end{bmatrix}$
$\begin{bmatrix} L \\ 8 \\ 0 \end{bmatrix}$	$\begin{tabular}{c} L \\ C & 9 \\ L \\ I \\ L \\ end{tabular}$	L I 10A O	P I 11E P	$\begin{bmatrix} P\\I12I\\K\end{bmatrix}$	$\begin{bmatrix} K \\ B13 I \\ M \end{bmatrix}$	$ \begin{smallmatrix} K \\ A14 I \\ K \end{smallmatrix} $	$\begin{tabular}{c} N \\ I \ 15C \\ P \end{tabular}$

• as the symbolic dynamical system $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$ representing the orbits under the \mathbb{Z}^2 -action $R_{\mathcal{Z}}$ defined by rotations on \mathbb{T}^2 and coded by the topological partition $\mathcal{P}_{\mathcal{Z}}$ of \mathbb{T}^2 :



where $\varphi = \frac{1+\sqrt{5}}{2}$ is the golden ratio.

(1)

The reader may observe that while increasing the dimension from 1 to 2, we replaced the second characterization of the Fibonacci subshift \mathcal{X}_s based on the factor complexity by the notion of Wang shift or more generally subshift of finite type (SFT). It may seem counterintuitive since the Fibonacci subshift is aperiodic and 1-dimensional SFT always contain a periodic configuration [LM95], but this is not a contradiction in higher dimension since there exist aperiodic 2-dimensional SFTs [Ber66].

In this chapter, we show that $\Omega_{\mathcal{Z}}$ and $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$ are self-similar. The tools used in the proofs are completely different in each case: based on the notion of marker tiles in the former case and on Rauzy induction of \mathbb{Z}^2 -rotations in the latter. It turns out that the 2-dimensional morphism describing the self-similarities is Φ in both cases.

Theorem 1.1. [Lep24] The Wang shift $\Omega_{\mathcal{Z}} \subset [\![0,15]\!]^{\mathbb{Z}^2}$ is self-similar satisfying $\Omega_{\mathcal{Z}} = \overline{\Phi(\Omega_{\mathcal{Z}})}^{\sigma}$ where $\Phi : [\![0,15]\!] \rightarrow [\![0,15]\!]^{*^2}$ is defined in Equation (1).

S. LABBÉ

Theorem 1.1 was first proved in [Lep24]. The set \mathcal{Z} of 16 tiles was introduced in [Lep24] as a simplification of the set \mathcal{U} of 19 Wang tiles introduced in [Lab19]. Lepšová proved that $\Omega_{\mathcal{Z}}$ is topologically conjugate to $\Omega_{\mathcal{U}}$. Therefore, the Wang shift $\Omega_{\mathcal{Z}}$ is also minimal, aperiodic and self-similar as the same was known for $\Omega_{\mathcal{U}}$. The proof of Theorem 1.1 provided here is constructive and uses the tools developed in [Lab21c].

Theorem 1.2. The subshift $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}} \subset [\![0,15]\!]^{\mathbb{Z}^2}$ is self-similar satisfying $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}} = \overline{\Phi(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}})}^{\sigma}$ where $\Phi : [\![0,15]\!] \rightarrow [\![0,15]\!]^{*^2}$ is defined in Equation (1).

The proof of Theorem 1.2 provided here is also constructive and uses the tools developed in [Lab21b] to perform the Rauzy induction of toral \mathbb{Z}^2 -rotations coded by polygonal partitions.

The equality of the three subshifts follows from a criterion for the minimality of self-similar subshifts stated in Lemma 3.9.

Theorem 1.3. The three subshifts are equal: $\mathcal{X}_{\Phi} = \Omega_{\mathcal{Z}} = \mathcal{X}_{\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}}$.

The 2-dimensional subshift $\Omega_{\mathcal{U}}$ was introduced in [Lab19] and discovered during the study of the substitutive structure [Lab21c] of the Jeandel–Rao Wang shift [JR21]. Its description as the coding of a toral \mathbb{Z}^2 -action was presented in [Lab21a] and its substitutive structure was further developed in [Lab21b]. This chapter provides a transversal reading of results divided in four different articles about Jeandel–Rao tilings and gathers the methods introduced by focussing on the self-similar subshift hidden in the Jeandel–Rao Wang shift, which is more simple. Thus we avoid the difficulty raised by the Jeandel–Rao Wang shift itself which is not a minimal subshift, has a long preperiod in its substitutive description and needs the definition of other tools including the shear-conjugacy.

Structure of the chapter. Section 2 gathers preliminary notions on topological dynamical systems, subshifts and shifts of finite type and *d*-dimensional languages. In Section 3, we define a 2-dimensional self-similar subshift \mathcal{X}_{Φ} from a 2-dimensional substitution Φ defined on 16-letter alphabet. We show that \mathcal{X}_{Φ} is aperiodic. In Section 4, we introduce a Wang shift $\Omega_{\mathcal{Z}}$ defined from a set \mathcal{Z} of 16 Wang tiles and we show using the notion of marker tiles that it is self-similar and $\Omega_{\mathcal{Z}} = \mathcal{X}_{\Phi}$. In Section 5, we introduce a 2-dimensional subshift defined as the symbolic representation of a toral \mathbb{Z}^2 -rotation using a partition of $\mathbb{R}^2/\mathbb{Z}^2$ into 16 polygons. We show that it is also self-similar and equal to \mathcal{X}_{Φ} . Around 40 exercises are included in the chapter. Their solutions are gathered at the end of the chapter in Section 7.

Algorithms to find marker tiles and compute the Rauzy induction of \mathbb{Z}^2 -rotations are provided as well as the SageMath code to reproduce the computations.

Acknowledgments. The author is thankful to Jana Lepšová for her careful reading of a preliminary version of this chapter. All computations made in this chapter were made with the following versions of SageMath [Sag24] and optional package slabbe [Lab23b]:

```
sage: version()1SageMath version 10.4.beta9, Release Date: 2024-06-092sage: import importlib.metadata3sage: importlib.metadata.version("slabbe")40.7.65
```

All outputs within red boxes in this chapter are computed directly from SageMath using the sagetex package. Please contact the author if you have trouble reproducing any of the computations. It is possible to doctest (check all outputs) using the command "sage -t chapter_doctest.sage" on the file provided in the archive (166 tests, 13.20 s).

2. Preliminaries

2.1. Topological dynamical systems. Most of the notions introduced here can be found in [Wal82]. A dynamical system is a triple (X, G, T), where X is a topological space, G is a topological group and T is a continuous function $G \times X \to X$ defining a left action of G on X: if $x \in X$, e is the identity element of G and $g, h \in G$, then using additive notation for the operation in G we have T(e, x) = x and T(g + h, x) = T(g, T(h, x)). In other words, if one denotes the transformation $x \mapsto T(g, x)$ by T^g , then $T^{g+h} = T^g T^h$. In this work, we consider the Abelian group $G = \mathbb{Z} \times \mathbb{Z}$.

If $Y \subset X$, let \overline{Y} denote the topological closure of Y and let $T(Y) := \bigcup_{g \in G} T^g(Y)$ denote the T-closure of Y. Alternatively, we also use the notation $\overline{Y}^T = T(Y)$ to denote the T-closure of Y. A subset $Y \subset X$ is T-invariant if T(Y) = Y. A dynamical system (X, G, T) is called **minimal** if X does not contain any nonempty, proper, closed T-invariant subset. The left action of G on X is **free** if g = e whenever there exists $x \in X$ such that $T^g(x) = x$.

Let (X, G, T) and (Y, G, S) be two dynamical systems with the same topological group G. A **homomorphism** $\theta : (X, G, T) \to (Y, G, S)$ is a continuous function $\theta : X \to Y$ satisfying the commuting property that $S^g \circ \theta = \theta \circ T^g$ for every $g \in G$. A homomorphism $\theta : (X, G, T) \to (Y, G, S)$ is called an **embedding** if it is one-to-one, a **factor map** if it is onto, and a **topological conjugacy** if it is both one-to-one and onto and its inverse map is continuous. If $\theta : (X, G, T) \to (Y, G, S)$ is called an **extension** of (Y, G, S) is called a **factor** of (X, G, T) and (X, G, T) is called an **extension** of (Y, G, S). Two dynamical systems are **topologically conjugate** if there is a topological conjugacy between them.

A measure-preserving dynamical system is defined as a system $(X, G, T, \mu, \mathcal{B})$, where μ is a probability measure defined on the Borel σ -algebra \mathcal{B} of subsets of X, and $T^g : X \to X$ is a measurable map which preserves the measure μ for all $g \in G$, that is, $\mu(T^g(B)) = \mu(B)$ for all $B \in \mathcal{B}$. The measure μ is said to be *T*-invariant. In what follows, when it is clear from the context, we omit the Borel σ -algebra \mathcal{B} of subsets of X and write (X, G, T, μ) to denote a measure-preserving dynamical system.

The set of all *T*-invariant probability measures of a dynamical system (X, G, T) is denoted by $\mathcal{M}^T(X)$. A *T*-invariant probability measure on *X* is called **ergodic** if for every set $B \in \mathcal{B}$ such that $T^g(B) = B$ for all $g \in G$, we have that *B* has either zero or full measure. A dynamical system (X, G, T) is **uniquely ergodic** if it has only one invariant probability measure, i.e., $|\mathcal{M}^T(X)| = 1$. One can prove that a uniquely ergodic dynamical system is ergodic. A dynamical system (X, G, T) is said **strictly ergodic** if it is uniquely ergodic and minimal.

Let $(X, G, T, \mu, \mathcal{B})$ and $(X', G, T', \mu', \mathcal{B}')$ be two measure-preserving dynamical systems. We say that the two systems are **isomorphic** if there exist measurable sets $X_0 \subset X$ and $X'_0 \subset X'$ of full measure (i.e., $\mu(X \setminus X_0) = 0$ and $\mu'(X' \setminus X'_0) = 0$) with $T^g(X_0) \subset X_0$, $T'^g(X'_0) \subset X'_0$ for all $g \in G$ and there exists a map $\phi : X_0 \to X'_0$, called an **isomorphism**, that is one-to-one and onto and such that for all $A \in \mathcal{B}'(X'_0)$,

• $\phi^{-1}(A) \in \mathcal{B}(X_0),$

- $\mu(\phi^{-1}(A)) = \mu'(A)$, and
- $\phi \circ T^g(x) = T'^g \circ \phi(x)$ for all $x \in X_0$ and $g \in G$.

The role of the set X_0 is to make precise the fact that the properties of the isomorphism need to hold only on a set of full measure.

S. LABBÉ

2.2. Subshifts and shifts of finite type. In this section, we introduce multidimensional subshifts, a particular type of dynamical systems [LM95, §13.10], [Sch01, Lin04, Hoc16]. Let \mathcal{A} be a finite set, $d \geq 1$, and let $\mathcal{A}^{\mathbb{Z}^d}$ be the set of all maps $x : \mathbb{Z}^d \to \mathcal{A}$, equipped with the compact product topology. An element $x \in \mathcal{A}^{\mathbb{Z}^d}$ is called **configuration** and we write it as $x = (x_m) = (x_m : m \in \mathbb{Z}^d)$, where $x_m \in \mathcal{A}$ denotes the value of x at m. The topology on $\mathcal{A}^{\mathbb{Z}^d}$ is compatible with the metric defined for all configurations $x, x' \in \mathcal{A}^{\mathbb{Z}^d}$ by $\operatorname{dist}(x, x') = 2^{-\min\{\|n\| : x_n \neq x'_n\}}$ where $\|n\| = |n_1| + \cdots + |n_d|$. The shift action $\sigma : n \mapsto \sigma^n$ of the additive group \mathbb{Z}^d on $\mathcal{A}^{\mathbb{Z}^d}$ is defined by

(2)
$$(\sigma^n(x))_m = x_{m+n}$$

for every $x = (x_m) \in \mathcal{A}^{\mathbb{Z}^d}$ and $n \in \mathbb{Z}^d$. If $X \subset \mathcal{A}^{\mathbb{Z}^d}$, let \overline{X} denote the topological closure of X and let $\overline{X}^{\sigma} := \{\sigma^n(x) \mid x \in X, n \in \mathbb{Z}^d\}$ denote the shift-closure of X. A subset $X \subset \mathcal{A}^{\mathbb{Z}^d}$ is **shift-invariant** if $\overline{X}^{\sigma} = X$. A closed, shift-invariant subset $X \subset \mathcal{A}^{\mathbb{Z}^d}$ is a **subshift**. If $X \subset \mathcal{A}^{\mathbb{Z}^d}$ is a subshift we write $\sigma = \sigma^X$ for the restriction of the shift action (2) to X. When X is a subshift, the triple $(X, \mathbb{Z}^d, \sigma)$ is a dynamical system and the notions presented in the previous section hold.

A configuration $x \in X$ is **periodic** if there is a nonzero vector $\mathbf{n} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$ such that $x = \sigma^n(x)$ and otherwise it is **nonperiodic**. We say that a nonempty subshift X is **aperiodic** if the shift action σ on X is free.

For any subset $S \subset \mathbb{Z}^d$ let $\pi_S : \mathcal{A}^{\mathbb{Z}^d} \to \mathcal{A}^S$ denote the projection map which restricts every $x \in \mathcal{A}^{\mathbb{Z}^d}$ to S. A **pattern** is a function $p \in \mathcal{A}^S$ for some finite subset $S \subset \mathbb{Z}^d$. To every pattern $p \in \mathcal{A}^S$ corresponds a subset $\pi_S^{-1}(p) \subset \mathcal{A}^{\mathbb{Z}^d}$ called **cylinder**. A nonempty set $X \subset \mathcal{A}^{\mathbb{Z}^d}$ is a **subshift** if and only if there exists a set \mathcal{F} of **forbidden** patterns such that

(3)
$$X = \{ x \in \mathcal{A}^{\mathbb{Z}^d} \mid \pi_S \circ \sigma^n(x) \notin \mathcal{F} \text{ for every } \boldsymbol{n} \in \mathbb{Z}^d \text{ and } S \subset \mathbb{Z}^d \},$$

see [Hoc16, Prop. 9.2.4]. A subshift $X \subset \mathcal{A}^{\mathbb{Z}^d}$ is a **subshift of finite type** (SFT) if there exists a finite set \mathcal{F} such that (3) holds. In this chapter, we consider shifts of finite type on $\mathbb{Z} \times \mathbb{Z}$, that is, the case d = 2.

2.3. *d*-dimensional word. In this section, we recall the definition of *d*-dimensional word that appeared in [CKR10] and we keep the notation $u \odot^i v$ they proposed for the concatenation.

We denote by $\{e_k | 1 \le k \le d\}$ the canonical basis of \mathbb{Z}^d where $d \ge 1$ is an integer. If $i \le j$ are integers, then $[\![i, j]\!]$ denotes the interval of integers $\{i, i+1, \ldots, j\}$. Let $\mathbf{n} = (n_1, \ldots, n_d) \in \mathbb{N}^d$ and \mathcal{A} be an alphabet. We denote by \mathcal{A}^n the set of functions

$$u: \llbracket 0, n_1 - 1 \rrbracket \times \cdots \times \llbracket 0, n_d - 1 \rrbracket \to \mathcal{A}.$$

An element $u \in \mathcal{A}^n$ is called a *d*-dimensional word of shape $\boldsymbol{n} = (n_1, \ldots, n_d) \in \mathbb{N}^d$ on the alphabet \mathcal{A} . We use the notation $\text{SHAPE}(u) = \boldsymbol{n}$ when necessary. The set of all finite *d*-dimensional words is $\mathcal{A}^{*^d} = \{\mathcal{A}^n \mid \boldsymbol{n} \in \mathbb{N}^d\}$. A *d*-dimensional word of shape $\boldsymbol{e}_k + \sum_{i=1}^d \boldsymbol{e}_i$ is called a **domino in the direction** \boldsymbol{e}_k . When the context is clear, we write \mathcal{A} instead of $\mathcal{A}^{(1,\ldots,1)}$. When d = 2, we represent a *d*-dimensional word *u* of shape (n_1, n_2) as a matrix with Cartesian coordinates:

$$u = \left(\begin{array}{cccc} u_{0,n_2-1} & \dots & u_{n_1-1,n_2-1} \\ \dots & \dots & \dots \\ u_{0,0} & \dots & u_{n_1-1,0} \end{array}\right).$$

Let $n, m \in \mathbb{N}^d$ and $u \in \mathcal{A}^n$ and $v \in \mathcal{A}^m$. If there exists an index *i* such that $n_j = m_j$ for all $j \in \{1, \ldots, d\} \setminus \{i\}$, then the **concatenation** of *u* and *v* in the direction e_i is defined: it is the *d*-dimensional word $u \odot^i v$ of shape $(n_1, \ldots, n_{i-1}, n_i + m_i, n_{i+1}, \ldots, n_d) \in \mathbb{N}^d$ given as

$$(u \odot^{i} v)(\boldsymbol{a}) = \begin{cases} u(\boldsymbol{a}) & \text{if } 0 \leq a_{i} < n_{i}, \\ v(\boldsymbol{a} - n_{i}\boldsymbol{e}_{i}) & \text{if } n_{i} \leq a_{i} < n_{i} + m_{i}. \end{cases}$$

The following equation illustrates the concatenation of 2-dimensional words in the direction e_2 :

$$\begin{pmatrix} 4 & 5\\ 10 & 5 \end{pmatrix} \odot^2 \begin{pmatrix} 3 & 10\\ 9 & 9\\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 3 & 10\\ 9 & 9\\ 0 & 0\\ 4 & 5\\ 10 & 5 \end{pmatrix}$$

whereas

$$\begin{pmatrix} 3 & 10 \\ 9 & 9 \\ 0 & 0 \end{pmatrix} \odot^2 \begin{pmatrix} 4 & 5 \\ 10 & 5 \end{pmatrix} = \begin{pmatrix} 4 & 5 \\ 10 & 5 \\ 3 & 10 \\ 9 & 9 \\ 0 & 0 \end{pmatrix}$$

and in the direction e_1 :

$$\begin{pmatrix} 2 & 8 & 7 \\ 7 & 3 & 9 \\ 1 & 1 & 0 \\ 6 & 6 & 7 \\ 7 & 4 & 3 \end{pmatrix} \odot^{1} \begin{pmatrix} 3 & 10 \\ 9 & 9 \\ 0 & 0 \\ 4 & 5 \\ 10 & 5 \end{pmatrix} = \begin{pmatrix} 2 & 8 & 7 & 3 & 10 \\ 7 & 3 & 9 & 9 & 9 \\ 1 & 1 & 0 & 0 & 0 \\ 6 & 6 & 7 & 4 & 5 \\ 7 & 4 & 3 & 10 & 5 \end{pmatrix}$$

Let $n, m \in \mathbb{N}^d$ and $u \in \mathcal{A}^n$ and $v \in \mathcal{A}^m$. We say that u occurs in v at position $p \in \mathbb{N}^d$ if v is large enough, i.e., $m - p - n \in \mathbb{N}^d$ and

 $v(\boldsymbol{a} + \boldsymbol{p}) = u(\boldsymbol{a})$

for all $\boldsymbol{a} = (a_1, \ldots, a_d) \in \mathbb{N}^d$ such that $0 \leq a_i < n_i$ with $1 \leq i \leq d$. If u occurs in v at some position, then we say that u is a d-dimensional **subword** or **factor** of v.

2.4. *d*-dimensional rectangular language. A subset $L \subseteq \mathcal{A}^{*^d}$ is called a *d*-dimensional language. The factorial closure of a language L is

 $\overline{L}^{Fact} = \{ u \in \mathcal{A}^{*^d} \mid u \text{ is a } d \text{-dimensional subword of some } v \in L \}.$

A language L is **factorial** if $\overline{L}^{Fact} = L$. All languages considered in this contribution are factorial. Given a configuration $x \in \mathcal{A}^{\mathbb{Z}^d}$, the **language** $\mathcal{L}(x)$ defined by x is

$$\mathcal{L}(x) = \{ u \in \mathcal{A}^{*^d} \mid u \text{ is a } d\text{-dimensional subword of } x \}$$

The **language** of a subshift $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ is $\mathcal{L}_X = \bigcup_{x \in X} \mathcal{L}(x)$. Conversely, given a factorial language $L \subseteq \mathcal{A}^{*^d}$ we define the subshift

$$\mathcal{X}_L = \{ x \in \mathcal{A}^{\mathbb{Z}^d} \mid \mathcal{L}(x) \subseteq L \}.$$

A *d*-dimensional subword $u \in \mathcal{A}^{*^d}$ is **allowed** in a subshift $X \subset \mathcal{A}^{\mathbb{Z}^d}$ if $u \in \mathcal{L}_X$ and it is **forbidden** in X if $u \notin \mathcal{L}_X$. A language $L \subseteq \mathcal{A}^{*^d}$ is **forbidden** in a subshift $X \subset \mathcal{A}^{\mathbb{Z}^d}$ if $L \cap \mathcal{L}_X = \emptyset$.

Exercise 2.1

Let $x: \mathbb{Z}^d \to \{0, 1\}$ be the configuration defined by

 $x(\boldsymbol{n}) = \|\boldsymbol{n}\|_1 \mod 2$ for every $\boldsymbol{n} \in \mathbb{Z}^d$.

Describe $\mathcal{L}(x)$. How many elements are in the subshift $\mathcal{X}_{\mathcal{L}(x)}$?

2.5. *d*-dimensional morphisms. In this section, we generalize the definition of *d*-dimensional morphisms [CKR10] to the case where the domain and codomain are different as in the case of *S*-adic systems [BD14].

Let \mathcal{A} and \mathcal{B} be two alphabets. Let $L \subseteq \mathcal{A}^{*^d}$ be a factorial language. A function $\omega : L \to \mathcal{B}^{*^d}$ is a *d*-dimensional morphism if for every *i* with $1 \leq i \leq d$, and every $u, v \in L$ such that $u \odot^i v$ is defined and $u \odot^i v \in L$, we have that the concatenation $\omega(u) \odot^i \omega(v)$ in direction e_i is defined and

$$\omega(u \odot^i v) = \omega(u) \odot^i \omega(v).$$

Note that the left-hand side of the equation is defined since $u \odot^i v$ belongs to the domain of ω . A *d*-dimensional morphism $L \to \mathcal{B}^{*^d}$ is thus completely defined from the image of the letters in \mathcal{A} , so we sometimes denote a *d*-dimensional morphism as a rule $\mathcal{A} \to \mathcal{B}^{*^d}$ when the language *L* is unspecified.

The next lemma can be deduced from the definition. It says that when $d \geq 2$ every d-dimensional morphism defined on the whole set $L = \mathcal{A}^{*^d}$ is uniform. We say that a d-dimensional morphism $\omega : L \to \mathcal{B}^{*^d}$ is **uniform** if there exists a shape $\mathbf{n} \in \mathbb{N}^d$ such that $\omega(a) \in \mathcal{B}^n$ for every letter $a \in \mathcal{A}$. These are called block-substitutions in [Fra18].

Lemma 2.1. Let $\omega : L \to \mathcal{B}^{*^d}$ be a d-dimensional morphism. If $d \ge 2$ and $L = \mathcal{A}^{*^d}$, then ω is uniform.

Therefore, to consider non-uniform *d*-dimensional morphisms when $d \geq 2$, we need to restrict the domain to a strict subset $L \subsetneq \mathcal{A}^{*^d}$. In [CKR10] and [Moz89, p.144], they consider the case $\mathcal{A} = \mathcal{B}$ and they restrict the domain of *d*-dimensional morphisms to the language they generate.

Given a language $L \subseteq \mathcal{A}^{*^d}$ of *d*-dimensional words and a *d*-dimensional morphism $\omega : L \to \mathcal{B}^{*^d}$, we define the image of the language L under ω as the language

$$\overline{\omega(L)}^{Fact} = \{ u \in \mathcal{B}^{*^d} \mid u \text{ is a } d\text{-dimensional subword of } \omega(v) \text{ with } v \in L \} \subseteq \mathcal{B}^{*^d}.$$

Observe that some element of $\overline{\omega(L)}^{Fact}$ do not have a preimage under ω .

Let $L \subseteq \mathcal{A}^{*^d}$ be a factorial language and $\mathcal{X}_L \subseteq \mathcal{A}^{\mathbb{Z}^d}$ be the subshift generated by L. A *d*-dimensional morphism $\omega : L \to \mathcal{B}^{*^d}$ can be extended to a continuous map $\omega : \mathcal{X}_L \to \mathcal{B}^{\mathbb{Z}^d}$ in such a way that the origin of $\omega(x)$ is at zero position in the word $\omega(x_0)$ for all $x \in \mathcal{X}_L$. More precisely, the image under ω of the configuration $x \in \mathcal{X}_L$ is

$$\omega(x) = \lim_{n \to \infty} \sigma^{f(n)} \omega \left(\sigma^{-n\mathbf{1}}(x|_{[-n\mathbf{1},n\mathbf{1}[]}) \right) \in \mathcal{B}^{\mathbb{Z}^d}$$

where $\mathbf{1} = (1, \ldots, 1) \in \mathbb{Z}^d$, $f(n) = \text{SHAPE}\left(\omega(\sigma^{-n\mathbf{1}}(x|_{[-n\mathbf{1},\mathbf{0}[]}))\right)$ for all $n \in \mathbb{N}$ and $[[\mathbf{m}, \mathbf{n}[] = [[\mathbf{m}_1, n_1 - 1]] \times \cdots \times [[\mathbf{m}_d, n_d - 1]].$

In general, the closure under the shift of the image of a subshift $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ under ω is the subshift

$$\overline{\omega(X)}^{\sigma} = \{ \sigma^{k} \omega(x) \in \mathcal{B}^{\mathbb{Z}^{d}} \mid k \in \mathbb{Z}^{d}, x \in X \} \subseteq \mathcal{B}^{\mathbb{Z}^{d}}.$$

Now we show that *d*-dimensional morphisms preserve minimality of subshifts.

Lemma 2.2. Let $\omega : X \to \mathcal{B}^{\mathbb{Z}^d}$ be a d-dimensional morphism for some $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$. If X is a minimal subshift, then $\overline{\omega(X)}^{\sigma}$ is a minimal subshift.

Proof. Let $\emptyset \neq Z \subseteq \overline{\omega(X)}^{\sigma}$ be a closed shift-invariant subset. We want to show that $\overline{\omega(X)}^{\sigma} \subseteq Z$. Let $u \in \overline{\omega(X)}^{\sigma}$. Thus $u = \sigma^{k}\omega(x)$ for some $k \in \mathbb{Z}^{d}$ and $x \in X$. Since $Z \neq \emptyset$, there exists $z \in Z$. Thus $z = \sigma^{k'}\omega(x')$ for some $k' \in \mathbb{Z}^{d}$ and $x' \in X$. Since X is minimal, there exists a sequence $(\mathbf{k}_{n})_{n\in\mathbb{N}}, \mathbf{k}_{n} \in \mathbb{Z}^{d}$, such that $x = \lim_{n\to\infty} \sigma^{\mathbf{k}_{n}}x'$. For some other sequence $(\mathbf{h}_{n})_{n\in\mathbb{N}}, \mathbf{h}_{n} \in \mathbb{Z}^{d}$, we have

$$u = \sigma^{k}\omega(x) = \sigma^{k}\omega\left(\lim_{n \to \infty} \sigma^{k_{n}} x'\right) = \sigma^{k}\lim_{n \to \infty} \sigma^{h_{n}}\omega(x') = \lim_{n \to \infty} \sigma^{k+h_{n}-k'} z.$$

Since Z is closed and shift-invariant, it follows that $u \in Z$.

Exercise 2.2

We consider the 2-dimensional morphism Φ defined in Equation (1) on the alphabet $\mathcal{A} = \llbracket 0, 15 \rrbracket$. Compute $\Phi(u)$ for each 2-dimensional word u below

$$\begin{pmatrix} 11 \end{pmatrix}$$
, $(13, 7)$, $\begin{pmatrix} 6 \\ 10 \end{pmatrix}$, $\begin{pmatrix} 6 & 1 \\ 11 & 8 \end{pmatrix}$

except one for which the image $\Phi(u)$ is not well-defined.

Exercise 2.3

Let $\omega: X \to \mathcal{B}^{\mathbb{Z}^d}$ be a *d*-dimensional morphism for some $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$. Prove that if X is a subshift, then $\overline{\omega(X)}^{\sigma}$ is a subshift.

3. An Aperiodic self-similar subshift

3.1. Self-similar subshifts. In this section, we consider languages and subshifts defined from morphisms leading to self-similar structures. In this situation, the domain and codomain of morphisms are defined over the same alphabet. Formally, we consider the case of *d*-dimensional morphisms $\mathcal{A} \to \mathcal{B}^{*^d}$ where $\mathcal{A} = \mathcal{B}$.

The definition of self-similarity depends on the notion of expansiveness. It avoids the presence of lower-dimensional self-similar structure by having expansion in all directions.

Definition 3.1. We say that a d-dimensional morphism $\omega : \mathcal{A} \to \mathcal{A}^{*^d}$ is **expansive** if for every $a \in \mathcal{A}$ and $K \in \mathbb{N}$, there exists $m \in \mathbb{N}$ such that

$$\min(SHAPE(\omega^m(a))) > K.$$

Definition 3.2. A subshift $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ is **self-similar** if there exists an expansive d-dimensional morphism $\omega : \mathcal{A} \to \mathcal{A}^{*^d}$ such that $X = \overline{\omega(X)}^{\sigma}$.

Respectively, a language $L \subseteq \mathcal{A}^{*^d}$ is **self-similar** if there exists an expansive *d*-dimensional morphism $\omega : \mathcal{A} \to \mathcal{A}^{*^d}$ such that $L = \overline{\omega(L)}^{Fact}$.

Self-similar languages and subshifts can be constructed by iterative application of a morphism ω starting with the letters. The **language** \mathcal{L}_{ω} defined by an expansive *d*-dimensional morphism $\omega : \mathcal{A} \to \mathcal{A}^{*^d}$ is

 $\mathcal{L}_{\omega} = \{ u \in \mathcal{A}^{*^{d}} \mid u \text{ is a } d\text{-dimensional subword of } \omega^{n}(a) \text{ for some } a \in \mathcal{A} \text{ and } n \in \mathbb{N} \}.$ It satisfies $\mathcal{L}_{\omega} = \overline{\omega(\mathcal{L}_{\omega})}^{Fact}$ and thus is self-similar. The **substitutive shift** $\mathcal{X}_{\omega} = \mathcal{X}_{\mathcal{L}_{\omega}}$ defined from the language of ω is a self-similar subshift since $\mathcal{X}_{\omega} = \overline{\omega(\mathcal{X}_{\omega})}^{\sigma}$ holds.

Let $\Phi : [\![0, 15]\!] \to [\![0, 15]\!]^{*^2}$ be the 2-dimensional morphism defined in Exercise 2.2. At Figure 1, we compute the sequence of 2-dimensional words $(\Phi^n(12))_{n\in\mathbb{N}}$ for the first values of n. Since 12 appears in the lower left corner of $\Phi^2(12)$, the limit $\lim_{n\to\infty} \Phi^{2n}(12)$ is well-defined and it defines a configuration of the positive quadrant \mathbb{N}^2 .

This procedure can be done in each of the four quadrants. At Figure 2, we compute the sequence of 2-dimensional words $(\Phi^n \begin{pmatrix} 8 & 12 \\ 1 & 6 \end{pmatrix})_{n \in \mathbb{N}}$ for the first values of n. The limits

$$x = \lim_{n \to \infty} \Phi^{2n} \begin{pmatrix} 8 & 12\\ 1 & 6 \end{pmatrix}$$
$$y = \lim_{n \to \infty} \Phi^{2n+1} \begin{pmatrix} 8 & 12\\ 1 & 6 \end{pmatrix}$$

are well-defined and define two configurations of \mathbb{Z}^2 . They satisfy $\Phi(x) = y$ and $\Phi(y) = x$. This implies that the configurations x and y are fixed points of Φ^2 since $\Phi^2(x) = x$ and $\Phi^2(y) = y$.

Exercise 3.1

Prove that Φ defined in Equation (1) is expansive.

Exercise 3.2

Prove that $\mathcal{X}_{\Phi} \neq \emptyset$.

Exercise 3.3

Using the slabbe package of the SageMath open-source software, define the 2dimensional substitution Φ as follows. Note that 2-dimensional words are encoded using Cartesian-like coordinates instead of matrix-like coordinates.

sage:	from slabbe import Substitution2d	6
sage:	Phi = Substitution2d({0: [[14]], 1: [[13]], 2: [[12],[10]],	7
:	3: [[11],[8]], 4: [[14],[7]], 5: [[13],[7]], 6: [[12],[7]],	8
:	7: [[12,6]], 8: [[14,3]], 9: [[13,3]], 10: [[12,2]],	9
:	11: [[12,6],[10,1]], 12: [[11,6],[8,1]], 13: [[15,5],[9,1]],	10
:	14: [[11,4],[8,1]], 15: [[12,2],[7,0]]})	11

Reproduce the computations of the Figure 1 and Figure 2 in SageMath.



FIGURE 1. Building a configuration of the positive quadrant with Φ . We compute $(\Phi^n(12))_{n\in\mathbb{N}}$ for the first values of $n \in \{0, 1, 2, 3, 4, 5, 6\}$. The gray rectangles surround patterns seen two step before in the application of Φ . The limit $\lim_{n\to\infty} \Phi^{2n}(12)$ defines a configuration of the positive quadrant \mathbb{N}^2 and similarly for the limit $\lim_{n\to\infty} \Phi^{2n+1}(12)$.



FIGURE 2. Building a configuration of \mathbb{Z}^2 with Φ . We compute $(\Phi^n \begin{pmatrix} 8 & 12 \\ 1 & 6 \end{pmatrix})_{n \in \mathbb{N}}$ for the first values of $n \in \{0, 1, 2, 3, 4, 5\}$. The gray rectangles surround patterns seen two step before in the application of Φ . The limit $\lim_{n\to\infty} \Phi^{2n} \begin{pmatrix} 8 & 12 \\ 1 & 6 \end{pmatrix}$ defines a configuration of \mathbb{Z}^2 and similarly for the limit $\lim_{n\to\infty} \Phi^{2n+1} \begin{pmatrix} 8 & 12 \\ 1 & 6 \end{pmatrix}$.

Exercise 3.4

The language of horizontal and vertical dominoes that we see in Figure 1 and Figure 2 obtained from the morphism Φ are

$$H = \begin{cases} (03), (12), (13), (16), (20), (24), (36), (41), (51), \\ (61), (65), (713), (812), (911), (912), (1014), (118), \\ (127), (1210), (1211), (1215), (137), (1311), (1312), \\ (147), (1411), (159) \end{cases}$$

and

$$V = \begin{cases} \begin{pmatrix} 7 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ 1 \end{pmatrix}, \begin{pmatrix} 8 \\ 1 \end{pmatrix}, \begin{pmatrix} 10 \\ 1 \end{pmatrix}, \begin{pmatrix} 13 \\ 2 \end{pmatrix}, \begin{pmatrix} 13 \\ 3 \end{pmatrix}, \begin{pmatrix} 11 \\ 4 \end{pmatrix}, \begin{pmatrix} 11 \\ 5 \end{pmatrix}, \begin{pmatrix} 12 \\ 6 \end{pmatrix}, \begin{pmatrix} 14 \\ 6 \end{pmatrix}, \\ \begin{pmatrix} 0 \\ 7 \end{pmatrix}, \begin{pmatrix} 8 \\ 7 \end{pmatrix}, \begin{pmatrix} 10 \\ 7 \end{pmatrix}, \begin{pmatrix} 1 \\ 8 \end{pmatrix}, \begin{pmatrix} 9 \\ 8 \end{pmatrix}, \begin{pmatrix} 9 \\ 9 \end{pmatrix}, \begin{pmatrix} 1 \\ 10 \end{pmatrix}, \begin{pmatrix} 9 \\ 10 \end{pmatrix}, \begin{pmatrix} 4 \\ 11 \end{pmatrix}, \begin{pmatrix} 6 \\ 11 \end{pmatrix}, \\ \begin{pmatrix} 15 \\ 15 \end{pmatrix}, \begin{pmatrix} 2 \\ 12 \end{pmatrix}, \begin{pmatrix} 6 \\ 12 \end{pmatrix}, \begin{pmatrix} 11 \\ 12 \end{pmatrix}, \begin{pmatrix} 15 \\ 12 \end{pmatrix}, \begin{pmatrix} 3 \\ 13 \end{pmatrix}, \begin{pmatrix} 12 \\ 13 \end{pmatrix}, \begin{pmatrix} 13 \\ 13 \end{pmatrix}, \begin{pmatrix} 13 \\ 13 \end{pmatrix}, \begin{pmatrix} 14 \\ 13 \end{pmatrix}, \begin{pmatrix} 3 \\ 14 \end{pmatrix}, \\ \begin{pmatrix} 12 \\ 14 \end{pmatrix}, \begin{pmatrix} 5 \\ 15 \end{pmatrix}, \end{cases}$$

Prove that H and V are exactly the dominoes that appear in \mathcal{L}_{Φ} , that is,

$$(\mathcal{A} \odot^1 \mathcal{A}) \cap \mathcal{L}_{\Phi} = H \text{ and } (\mathcal{A} \odot^2 \mathcal{A}) \cap \mathcal{L}_{\Phi} = V$$

where $\mathcal{A} = \llbracket 0, 15 \rrbracket$ is the alphabet on which the morphism Φ is defined.

Exercise 3.5

Is the 2 × 2 word $\begin{pmatrix} 1 & 2 \\ 9 & 12 \end{pmatrix}$ in the set \mathcal{L}_{Φ} ?

Exercise 3.6

List the 45 elements of the set $\mathcal{A}^{(2,2)} \cap \mathcal{L}_{\Phi}$.

Exercise 3.7

Describe the 8 periodic points of Φ , i.e. the configurations $x \in \mathcal{A}^{\mathbb{Z}^2}$ such that $\Phi^k(x) = x$ for some $k \ge 1$.

3.2. *d*-dimensional recognizability and aperiodicity. The definition of recognizability dates back to the work of Host, Quéffelec and Mossé [Mos92]. The definition introduced below is based on work of Berthé et al. [BSTY19] on the recognizability in the case of *S*-adic systems where more than one substitution is involved.

Definition 3.3 (recognizable). Let $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ and $\omega : X \to \mathcal{B}^{\mathbb{Z}^d}$ be a d-dimensional morphism. If $y \in \overline{\omega(X)}^{\sigma}$, i.e., $y = \sigma^k \omega(x)$ for some $x \in X$ and $k \in \mathbb{Z}^d$, where σ is the d-dimensional shift map, we say that (\mathbf{k}, x) is an ω -representation of y. We say that it is centered if y_0 lies inside of the image of x_0 , i.e., if $0 \leq \mathbf{k} < \text{SHAPE}(\omega(x_0))$ coordinate-wise. We say that ω is recognizable in $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ if each $y \in \mathcal{B}^{\mathbb{Z}^d}$ has at most one centered ω -representation (\mathbf{k}, x) with $x \in X$.

Lemma 3.4. Let $\omega : X \to Y$ be some d-dimensional morphism between two subshifts X and Y.

- (1) If Y is aperiodic, then X is aperiodic.
- (2) If X is aperiodic and ω is recognizable in X, then $\overline{\omega(X)}^{\sigma}$ is aperiodic.

Proof. If X contains a periodic configuration x, then $\omega(x) \in Y$ is periodic.

S. LABBÉ

(ii) Let $y \in \overline{\omega(X)}^{\sigma}$. Then, there exist $\mathbf{k} \in \mathbb{Z}^d$ and $x \in X$ such that (\mathbf{k}, x) is a centered ω -representation of y, i.e., $y = \sigma^{\mathbf{k}}\omega(x)$. Suppose by contradiction that y has a nontrivial period $\mathbf{p} \in \mathbb{Z}^d \setminus \mathbf{0}$. Since $y = \sigma^{\mathbf{p}}y = \sigma^{\mathbf{p}+\mathbf{k}}\omega(x)$, we have that $(\mathbf{p} + \mathbf{k}, x)$ is an ω -representation of y. Since ω is recognizable, this representation is not centered. Therefore there exists $\mathbf{q} \in \mathbb{Z}^d \setminus \mathbf{0}$ such that y_0 lies in the image of $x_{\mathbf{q}} = (\sigma^q x)_0$. Therefore there exists $\mathbf{k}' \in \mathbb{Z}^d$ such that $(\mathbf{k}', \sigma^q x)$ is a centered ω -representation of y. Since ω is recognizable, we conclude that $\mathbf{k} = \mathbf{k}'$ and $x = \sigma^q x$. Then $x \in X$ is periodic which is a contradiction.

In general, $\omega(X)$ is not closed under the shift which implies that ω is not onto Y. This motivates the following definition.

Definition 3.5. Let X, Y be two subshifts and $\omega : X \to Y$ be a d-dimensional morphism. If $Y = \overline{\omega(X)}^{\sigma}$, then we say that ω is **onto up to a shift**.

The next proposition is well-known, see [Sol98, Mos92], who showed that recognizability and aperiodicity are equivalent for primitive substitutive sequences. We state and prove only one direction (the easy one) of the equivalence which does not need the notion of primitivity.

Proposition 3.6. Let $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ be a self-similar subshift satisfying $\overline{\omega(X)}^{\sigma} = X$ for some expansive d-dimensional morphism $\omega : \mathcal{A} \to \mathcal{A}^{\mathbb{Z}^d}$. If ω is recognizable in X, then X is aperiodic.

Proof. Suppose that there exists a periodic configuration $y \in X$ with period $\mathbf{p} \in \mathbb{Z}^d \setminus \{(0,0)\}$ satisfying $\sigma^{\mathbf{p}}y = y$. Since ω is expansive, let $m \in \mathbb{N}$ such that the shape of the image of every letter $a \in \mathcal{A}$ by ω^m is large enough, that is, SHAPE $(\omega^m(a)) > \mathbf{p}$ for every letter $a \in \mathcal{A}$. By hypothesis, every $y \in X$ has an ω -representation. Recursively, there exists an ω^m -representation (\mathbf{k}, x) of y satisfying $y = \sigma^k \omega^m(x)$. We may assume that it is centered since X is shift-invariant. By definition of centered representation, for every $\mathbf{u} \in \mathbb{Z}^d$ such that $\mathbf{0} \leq \mathbf{u} < \text{SHAPE}(\omega^m(x_0))$, (\mathbf{u}, x) is a centered ω^m -representation of $\sigma^{\mathbf{u}}\omega^m(x) = \sigma^{\mathbf{u}-\mathbf{k}}y$. By the choice of m, there exists $\mathbf{u} \in \mathbb{Z}^d$ such that $\mathbf{0} \leq \mathbf{u} < \text{SHAPE}(\omega^m(x_0))$ and $\mathbf{0} \leq \mathbf{u} + \mathbf{p} < \text{SHAPE}(\omega^m(x_0))$. Therefore (\mathbf{u}, x) is a centered ω^m -representation of $\sigma^{\mathbf{u}}\omega^m(x) = \sigma^{\mathbf{u}-\mathbf{k}}y$ and $(\mathbf{u} + \mathbf{p}, x)$ is a centered ω^m -representation of $\sigma^{\mathbf{u}+\mathbf{p}-\mathbf{k}}y = \sigma^{\mathbf{u}-\mathbf{k}}\sigma^{\mathbf{p}}y = \sigma^{\mathbf{u}-\mathbf{k}}y$. Therefore, ω^m is not recognizable which implies that ω is not recognizable which is a contradiction. We conclude that there is no periodic configuration $y \in X$.

Exercise 3.8

Let $\Phi : [\![0, 15]\!] \to [\![0, 15]\!]^{*^2}$ be the morphism defined in Exercise 2.2. Find periodic configurations $x, y \in [\![0, 15]\!]^{\mathbb{Z}^2}$ and $\mathbf{k} \in \mathbb{Z}^2$ such that

• (\mathbf{k}, x) is a Φ -representation of y,

- (\mathbf{k}, x) is a centered Φ -representation of y,
- (\mathbf{k}, x) is a Φ -representation of y which is not centered.

Exercise 3.9

Does there exist a configuration $y \in [\![0, 15]\!]^{\mathbb{Z}^2}$ that has more than one centered Φ -representation (\mathbf{k}, x) with $x \in [\![0, 15]\!]^{\mathbb{Z}^2}$?

Exercise 3.10

Let $\Phi : \llbracket 0, 15 \rrbracket \to \llbracket 0, 15 \rrbracket^{*^2}$ be the morphism defined in Exercise 2.2.

(1) Prove that Φ is recognizable in \mathcal{X}_{Φ} .

(2) Prove that \mathcal{X}_{Φ} is aperiodic.

3.3. Primitivity and minimality of self-similar subshifts. Substitutive shifts obtained from expansive and primitive morphisms are interesting for their properties. As in the onedimensional case, we say that ω is **primitive** if there exists $m \in \mathbb{N}$ such that for every $a, b \in \mathcal{A}$ the letter b occurs in $\omega^m(a)$.

Lemma 3.7. Let $\omega : \mathcal{A} \to \mathcal{A}^{*^d}$ be an expansive and primitive d-dimensional morphism. Then \mathcal{X}_{ω} is minimal, i.e., it contains no nonempty proper subshift.

Proof. The substitutive shift of ω is well-defined since ω is expansive and it is minimal since ω is primitive using standard arguments [Que10, §5.2].

The following two lemmas were proved in [Lab23a]. We reproduce their proof here for completeness.

Lemma 3.8. [Lab23a, Lemma 11.1] Let $\omega : \mathcal{A} \to \mathcal{A}^{*^d}$ be an expansive and primitive ddimensional morphism. Let $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ be a nonempty subshift such that $X = \overline{\omega(X)}^{\sigma}$. Then $\mathcal{X}_{\omega} \subseteq X$.

Proof. The language of X is also self-similar satisfying $\mathcal{L}(X) = \overline{\omega(\mathcal{L}(X))}^{Fact}$. Recursively, $\mathcal{L}(X) = \overline{\omega^m(\mathcal{L}(X))}^{Fact}$ for every $m \ge 1$. Thus there exists a letter $a \in \mathcal{A}$ such that for all $m \ge 1$, the d-dimensional word $\omega^m(a)$ is in the language $\mathcal{L}(X)$. From the primitivity of ω , there exists $m \ge 1$ such that $\omega^m(a)$ contains an occurrence of every letter of the alphabet \mathcal{A} . Therefore every letter is in $\mathcal{L}(X)$ and the d-dimensional word $\omega^m(a)$ is in the language $\mathcal{L}(X)$ for all letters $a \in \mathcal{A}$ and all $m \ge 1$. So we conclude that $\mathcal{L}_{\omega} \subseteq \mathcal{L}(X)$ and $\mathcal{X}_{\omega} \subseteq X$.

Recall that $\mathcal{X}_{\omega} = \overline{\omega(\mathcal{X}_{\omega})}^{\sigma}$. Thus from Lemma 3.7 and Lemma 3.8, when ω is expansive and primitive, then \mathcal{X}_{ω} is the smallest nonempty subshift $X \subseteq \mathcal{A}^{\mathbb{Z}^d}$ satisfying $X = \overline{\omega(X)}^{\sigma}$. The next result provides a criterion for the minimality of a self-similar subshift X satisfying $X = \overline{\omega(X)}^{\sigma}$.

To achieve this goal, it is convenient to consider the 2×2 patterns as well as the domino patterns of shape 1×2 and 2×1 . We use these dominoes to define two equivalence relations on the alphabet \mathcal{A} . Formally, the vertical dominoes of shape 1×2 appearing in the language $\mathcal{L}(\mathcal{X}_{\omega})$ define an equivalence relation \equiv_2 on \mathcal{A} given as the reflexive, symmetric and transitive closure of the pairs $\{(a, c) \in \mathcal{A} \times \mathcal{A} \mid {a \atop c} \in \mathcal{L}(\mathcal{X}_{\omega})\}$. Informally, $a \equiv_2 c$ for some letters $a, c \in \mathcal{A}$ means that letters a and c may appear in the same column in some configuration of \mathcal{X}_{ω} . Similarly, the horizontal dominoes of shape 2×1 appearing in the language $\mathcal{L}(\mathcal{X}_{\omega})$ define an equivalence relation \equiv_1 on \mathcal{A} given as the reflexive, symmetric and transitive closure of the pairs $\{(a, b) \in \mathcal{A} \times \mathcal{A} \mid {a \atop b} \in \mathcal{L}(\mathcal{X}_{\omega})\}$.

Using these two equivalence relations \equiv_1 and \equiv_2 on the alphabet \mathcal{A} , we consider the following graphs:

• Let $G_{\omega}^{2\times 2} = (V_{\omega}^{2\times 2}, E_{\omega}^{2\times 2})$ be the directed graph whose vertices and edges are $V_{\omega}^{2\times 2} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{A}^{2\times 2} \mid a \equiv_1 b, c \equiv_1 d, a \equiv_2 c, b \equiv_2 d \right\},$ $E_{\omega}^{2\times 2} = \left\{ \begin{pmatrix} e & f \\ g & h \end{pmatrix} \rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \middle| \begin{array}{l} a \text{ is the lower right letter of } \omega(e), \\ b \text{ is the lower left letter of } \omega(f), \\ c \text{ is the top right letter of } \omega(g), \\ d \text{ is the top left letter of } \omega(h) \end{array} \right\}.$ • Let $G_{\omega}^{2\times 1} = (V_{\omega}^{2\times 1}, E_{\omega}^{2\times 1})$ be the directed graph whose vertices and edges are $V_{\omega}^{2\times 1} = \left\{ (a & b) \in \mathcal{A}^{2\times 1} \mid a \equiv_1 b \right\},$ $E_{\omega}^{2\times 1} = \left\{ (a & b) \in \mathcal{A}^{2\times 1} \mid a \equiv_1 b \right\},$ there exists an integer j such that $0 \leq j < \text{HEIGHT}(\omega(e))$ and $a \text{ is the letter in the } j\text{-th row in the right-most column of } \omega(e), \\ b \text{ is the letter in the } j\text{-th row in the left-most column of } \omega(f) \end{array} \right\}.$ • Let $G_{\omega}^{1\times 2} = (V_{\omega}^{1\times 2}, E_{\omega}^{1\times 2})$ be the directed graph whose vertices and edges are $V_{\omega}^{1\times 2} = \left\{ \begin{pmatrix} a \\ c \end{pmatrix} \in \mathcal{A}^{1\times 2} \mid a \equiv_2 c \right\},$ $E_{\omega}^{1\times 2} = \left\{ \begin{pmatrix} e \\ g \end{pmatrix} \rightarrow \begin{pmatrix} a \\ c \end{pmatrix} \mid \text{ there exists an integer } i \text{ such that } 0 \leq i < \text{WIDTH}(\omega(e)) \text{ and} \\ a \text{ is the letter in the } i\text{-th column in the bottom-most row of } \omega(e), \\ c \text{ is the letter in the } i\text{-th column in the top-most row of } \omega(g) \end{cases} \right\}.$

RECURRENTVERTICES(G) = $\{v \in V \mid v \text{ belongs to a cycle of } G\}$.

A vertex is a recurrent vertex in a graph if and only if it belongs to a biinfinite path in the graph, thus the terminology of *recurrent*.

The following result from [Lab23a] allows to conclude that a self-similar subshift is minimal even when the 2-dimensional substitution admits more than one self-similar subshift (some made of configurations which are not uniformly recurrent). We reproduce its proof here for completeness.

Lemma 3.9. [Lab23a, Lemma 11.2] Let $X = \overline{\omega(X)}^{\sigma}$ be a nonempty self-similar subshift where $\omega : \mathcal{A} \to \mathcal{A}^{*^{d}}$ is an expansive and primitive 2-dimensional morphism. The following are equivalent:

(i) $\mathcal{L}(X) \cap \text{RECURRENTVERTICES}(G^s_{\omega}) \subset \mathcal{L}(\mathcal{X}_{\omega})$ for every shape $s \in \{2 \times 2, 2 \times 1, 1 \times 2\}$, (ii) $X = \mathcal{X}_{\omega}$,

(iii) X is minimal.

An element $u \in \mathcal{A}^n$ is called a *d*-dimensional word of shape $n = (n_1, \ldots, n_d) \in \mathbb{N}^d$ on the alphabet \mathcal{A} . We use the notation SHAPE(u) = n when necessary.

Proof. Assume that $X = \overline{\omega(X)}^{\sigma}$ for some $\emptyset \neq X \subseteq \mathcal{A}^{\mathbb{Z}^d}$.

(i) \Longrightarrow (ii) From Lemma 3.8, we have $\mathcal{X}_{\omega} \subseteq X$. Let $z \in \mathcal{L}(X)$. We want to show that $z \in \mathcal{L}(\mathcal{X}_{\omega})$. Since ω is expansive, let $m \in \mathbb{N}$ such that the image of every letter $a \in \mathcal{A}$ by ω^m is larger than z, that is, $\text{SHAPE}(\omega^m(a)) \geq \text{SHAPE}(z)$ for all $a \in \mathcal{A}$. We have $z \in \mathcal{L}(X) = \mathcal{L}(\omega^m(\mathcal{L}(X)))$. By the choice of m, z can not overlap more than two blocks $\omega^m(a)$ in the same direction. Thus, there exists a word $u \in \mathcal{L}(X)$ of shape $1 \times 1, 2 \times 1, 1 \times 2$ or 2×2 such that z is a subword of $\omega^m(u)$. If u is of shape 1×1 , then $z \in \mathcal{L}(\mathcal{X}_{\omega})$. We may assume that the word u has the smallest possible rectangular shape $s \in \{2 \times 1, 1 \times 2, 2 \times 2\}$. We have $u \in V^s_{\omega}$. Since $u \in \mathcal{L}(X)$ and X is self-similar, there exists a sequence $(u_k)_{k \in \mathbb{N}}$ with $u_k \in V^s_{\omega} \cap \mathcal{L}(X)$ for all $k \in \mathbb{N}$ such that

$$\cdots \rightarrow u_{k+1} \rightarrow u_k \rightarrow \cdots \rightarrow u_1 \rightarrow u_0 = u_0$$

is a left-infinite path in the graph G^s_{ω} . Since V^s_{ω} is finite, there exist some $k, k' \in \mathbb{N}$ with k < k' such that $u_k = u_{k'}$. Thus $u_k \in \text{RECURRENTVERTICES}(G^s_{\omega})$ and u is a subword of $\omega^k(u_k)$. From the hypothesis, we have $u_k \in \mathcal{L}(\mathcal{X}_{\omega})$. Since ω is primitive, there exists ℓ such that u_k is a subword of $\omega^{\ell}(a)$ for every $a \in \mathcal{A}$. Therefore, z is a subword of $\omega^{m+k+\ell}(a)$ for every $a \in \mathcal{A}$. Then $z \in \mathcal{L}(\mathcal{X}_{\omega})$ and $\mathcal{L}(X) \subseteq \mathcal{L}(\mathcal{X}_{\omega})$. Thus $X \subseteq \mathcal{X}_{\omega}$ and $X = \mathcal{X}_{\omega}$.

(ii) \implies (i) If $X = \mathcal{X}_{\omega}$, then $\mathcal{L}(X) = \mathcal{L}(\mathcal{X}_{\omega})$. Thus $\mathcal{L}(X) \cap \text{RECURRENTVERTICES}(G^s_{\omega}) \subset \mathcal{L}(X) = \mathcal{L}(\mathcal{X}_{\omega})$ for every shape $s \in \{2 \times 2, 2 \times 1, 1 \times 2\}$.

(ii) \implies (iii) From Lemma 3.7, the substitutive shift \mathcal{X}_{ω} is minimal.

(iii) \implies (ii) From Lemma 3.8, we have $\mathcal{X}_{\omega} \subseteq X$. Since X is minimal, we conclude that $\mathcal{X}_{\omega} = X$.

Exercise 3.11

Let $\Phi : [0, 15] \to [0, 15]^{*^2}$ be the morphism defined in Exercise 2.2.

- (1) Prove that Φ is primitive.
- (2) Prove that \mathcal{X}_{Φ} is minimal.

Exercise 3.12

Compute the sets RECURRENTVERTICES (G_{Φ}^s) for every shape $s \in \{2 \times 2, 2 \times 1, 1 \times 2\}$. Conclude that the 2-dimensional substitution Φ does not have a unique self-similar subshift $X = \overline{\Phi(X)}^{\sigma}$.

3.4. Markers. The goal of this section is to prove Theorem 3.13 which states that we can desubstitute a subshift provided that its alphabet contains a subset of markers. Markers are such that they appear on non-consecutive layers in the configurations of the subshift, see Definition 3.10. The results are stated for arbitrary dimension since their proofs are independent of the dimension, but we will use them in the 2-dimensional case afterwards.

We now define the notion of markers for subshifts $X \subset \mathcal{A}^{\mathbb{Z}^d}$ and prove that their presence allows to desubstitute uniquely the configurations in X using a *d*-dimensional morphism. Originally, those results were proved for d = 2 in order to desubstitute configurations from Wang shifts, see [Lab19] and [Lab21c]. It turns out that the notion of markers is more general and the results holds in general subshifts $X \subset \mathcal{A}^{\mathbb{Z}^d}$.

Recall that if $w : \mathbb{Z}^d \to \mathcal{A}$ is a configuration and $a \in \mathcal{A}$ is a letter, then $w^{-1}(a) \subset \mathbb{Z}^d$ is the set of positions where the letter a appears in w.

Definition 3.10. Let \mathcal{A} be an alphabet and $X \subset \mathcal{A}^{\mathbb{Z}^d}$ be a subshift. A nonempty subset $M \subset \mathcal{A}$ is called **markers in the direction** e_i , with $i \in \{1, \ldots, d\}$, if positions of the letters of M in any configuration are nonadjacent (d-1)-dimensional layers orthogonal to e_i , that is, for all configurations $w \in X$ there exists $P \subset \mathbb{Z}$ such that the positions of the markers satisfy

$$w^{-1}(M) = P \boldsymbol{e}_i + \sum_{k \neq i} \mathbb{Z} \boldsymbol{e}_k \quad with \quad 1 \notin P - P$$

where $P - P = \{b - a \mid a \in P, b \in P\}$ is the set of differences between elements of P.

Note that it follows from the definition that a subset of markers is a proper subset of \mathcal{A} as the case $M = \mathcal{A}$ is impossible.

Proving that a subset $M \subset \mathcal{A}$ is a subset of markers uses very local observations, namely the set of dominoes in the language of the subshift. It leads to the following criterion.

Lemma 3.11. Let \mathcal{A} be an alphabet and $X \subset \mathcal{A}^{\mathbb{Z}^d}$ be a subshift. A nonempty subset $M \subset \mathcal{A}$ is a subset of markers in the direction \mathbf{e}_i if and only if

 $M \odot^{i} M, \qquad M \odot^{k} (\mathcal{A} \setminus M), \qquad (\mathcal{A} \setminus M) \odot^{k} M$

are forbidden in X for every $k \in \{1, \ldots, d\} \setminus \{i\}$.

Proof. Suppose that $M \subset \mathcal{A}$ is a subset of markers in the direction \mathbf{e}_i . For any configuration $w \in X$, there exists $P \subset \mathbb{Z}$ such that $w^{-1}(M) = P\mathbf{e}_i + \sum_{k \neq i} \mathbb{Z}\mathbf{e}_k$ with $1 \notin P - P$. In any configuration $w \in X$ such that $w(p) \in M$, then $w(p \pm \mathbf{e}_k) \in M$ also belongs to M for every $k \neq i$. Therefore, $M \odot^k (\mathcal{A} \setminus M)$ and $(\mathcal{A} \setminus M) \odot^k M$ are forbidden in X for every $k \neq i$. Moreover, the fact that $1 \notin P - P$ implies that $M \odot^k M$ is forbidden in X.

Conversely, suppose that $M \odot^i M$, $M \odot^k (\mathcal{A} \setminus M)$ and $(\mathcal{A} \setminus M) \odot^k M$ are forbidden in X for every $k \neq i$. The last two conditions implies that in any configuration $w \in X$ such that $w(p) \in M$, then $w(p \pm e_k) \in M$ also belongs to M for every $k \neq i$. Therefore letters in M appears as complete layers in w, that is, $w^{-1}(M) = Pe_i + \sum_{k\neq i} \mathbb{Z}e_k$ for some $P \subset \mathbb{Z}$. Since $M \odot^i M$ is forbidden in X, it means that the layers are nonadjacent, or equivalently, $1 \notin P - P$. We conclude that M is a subset of markers in the direction e_i .

The existence of a subset of markers allows to desubstitute a subshift by "merging" each marker to the letter on its right (or on its left). This procedure creates substitution with a specific form sending a letter on a a letter or a domino. In the next lemma, we provide a sufficient condition for such substitutions to be recognizable.

Lemma 3.12. Let $d \ge 1$ and i such that $1 \le i \le d$. Let $\omega : \mathcal{B} \to \mathcal{A}^{\mathbb{Z}^d}$ be a d-dimensional morphism such that the image of letters are letters or dominoes in the direction \mathbf{e}_i . If $\omega|_{\mathcal{B}}$ is injective and there exists a subset $M \subset \mathcal{A}$ such that

(4)
$$\omega(\mathcal{B}) \subseteq (\mathcal{A} \setminus M) \cup \left((\mathcal{A} \setminus M) \odot^{i} M \right),$$

or

(5)
$$\omega(\mathcal{B}) \subseteq (\mathcal{A} \setminus M) \cup \left(M \odot^{i} (\mathcal{A} \setminus M)\right),$$

then ω is recognizable in $\mathcal{B}^{\mathbb{Z}^d}$.

If $a \odot^i b$ is a domino in the direction e_i , we say that a is on the left position and b is on the right position in the domino.

Proof. Let (\mathbf{k}, x) and (\mathbf{k}', x') be two centered ω -representations of $y \in \mathcal{A}^{\mathbb{Z}^d}$ with $\mathbf{k}, \mathbf{k}' \in \mathbb{Z}^d$ and $x, x' \in \mathcal{B}^{\mathbb{Z}^d}$. We want to show that they are equal.

Since the image of a letter under ω is a letter or a domino in the direction e_i , then $\mathbf{k}, \mathbf{k}' \in \{\mathbf{0}, e_i\}$. If $y_{\mathbf{0}} \in M$, then $y_{\mathbf{0}}$ appears as the left or right part of a domino and thus $\mathbf{k} = \mathbf{k}' = \mathbf{e}_i$ if Equation (4) holds or $\mathbf{k} = \mathbf{k}' = \mathbf{0}$ if Equation (5) holds.

Suppose now that $y_0 \in \mathcal{A} \setminus M$. If Equation (4) holds, then $\mathbf{k} = \mathbf{k}' = \mathbf{0}$. Suppose that Equation (5) holds. By contradiction, suppose that $\mathbf{k} \neq \mathbf{k}'$ and assume without lost of generality that $\mathbf{k} = \mathbf{0}$ and $\mathbf{k}' = \mathbf{e}_i$. This means that $\omega(x'_0) = y_{-\mathbf{e}_i} \odot^i y_0$ is a domino in the direction \mathbf{e}_i . Since $y_0 \in \mathcal{A} \setminus M$, we must have that $y_{-\mathbf{e}_i} \in M$ is a marker on the left. This is

impossible as $\omega(x_{-e_i}) = y_{-e_i} \in \mathcal{A} \setminus M$ or $\omega(x_{-e_i}) = y_{-2e_i} \odot^i y_{-e_i} \in M \odot^i (\mathcal{A} \setminus M)$. Therefore, we must have $\mathbf{k} = \mathbf{k}'$ and $\omega(x) = \omega(x')$.

Suppose by contradiction that $x \neq x'$. Let $\boldsymbol{a} = (a_1, \ldots, a_d) \in \mathbb{Z}^d$ be some minimal vector with respect to $\|\boldsymbol{a}\|_{\infty}$ such that $x_a \neq x'_a$. Injectivity of ω implies that $\omega(x_a)$ and $\omega(x'_a)$ must have different shapes. Suppose without lost of generality that $\omega(x_a) \in \mathcal{A}$ and $\omega(x'_a) = b \odot^i c \in \mathcal{A} \odot^i \mathcal{A}$. We need to consider two cases: $a_i \geq 0$ and $a_i < 0$.

Suppose $a_i \ge 0$. We must have that Equation (4) holds. We have $\omega(x_a) = b \in \mathcal{A} \setminus M$ and $c \in M$. But then $\omega(x_{a+e_i}) = c$ or $\omega(x_{a+e_i}) = c \odot^i d$ for some $c \in \mathcal{A} \setminus M$ and $d \in \mathcal{A}$ which is a contradiction.

Suppose $a_i < 0$. We must have that Equation (5) holds. We have $\omega(x_a) = c \in \mathcal{A} \setminus M$ and $b \in M$. But then $\omega(x_{a-e_i}) = b$ or $\omega(x_{a+e_i}) = d \odot^i b$ for some $b \in \mathcal{A} \setminus M$ and $d \in \mathcal{A}$ which is a contradiction. We conclude that x = x'.

The presence of markers allows to desubstitute uniquely the configurations of a subshift. There is even a choice to be made in the construction of the substitution. We may construct the substitution in such a way that the markers are on the left or on the right in the image of letters that are dominoes in the direction e_k . We make this distinction in the statement of the following result which was stated in the context of Wang shifts in [Lab19] and [Lab21c].

Theorem 3.13. Let \mathcal{A} be an alphabet and $X \subset \mathcal{A}^{\mathbb{Z}^d}$ be a subshift. If there exists a subset $M \subset \mathcal{A}$ of markers in the direction $\mathbf{e}_i \in \{\mathbf{e}_1, \ldots, \mathbf{e}_d\}$, then

(i) (markers on the right) there exists an alphabet \mathcal{B}_R , a subshift $Y \subset \mathcal{B}_R^{\mathbb{Z}^d}$ and a 2dimensional morphism $\omega_R : Y \to X$ such that

$$\omega_R(\mathcal{B}_R) \subseteq (\mathcal{A} \setminus M) \cup \left((\mathcal{A} \setminus M) \odot^i M \right)$$

which is recognizable and onto up to a shift and

(ii) (markers on the left) there exists an alphabet \mathcal{B}_L , a subshift $Y \subset \mathcal{B}_L^{\mathbb{Z}^d}$ and a 2dimensional morphism $\omega_L : Y \to X$ such that

 $\omega_L(\mathcal{B}_L) \subseteq (\mathcal{A} \setminus M) \cup \left(M \odot^i (\mathcal{A} \setminus M)\right)$

which is recognizable and onto up to a shift.

Proof. We do only the proof of (i) when the markers are on the right, since one case can be deduced from the other using symmetry.

Since X is a subshift, there exists a language $F \subset \mathcal{A}^{*^d}$ such that X is the set of configurations of $\mathcal{A}^{\mathbb{Z}^d}$ without any occurrence of pattern from F. Notice that since M is a set of markers in the direction e_i , we may assume $M \odot^i M \subset F$. Let $P \subset \mathcal{A} \odot^i \mathcal{A}$ and $Q \subset \mathcal{A}$ be the following sets:

$$P = \left((\mathcal{A} \setminus M) \odot^{i} M \right) \setminus F,$$

$$Q = \left\{ u \in \mathcal{A} \setminus M \mid \text{ there exists } v \in \mathcal{A} \setminus M \text{ such that } u \odot^{i} v \notin F \right\}.$$

We choose some ordering of their elements with indices starting from zero:

$$P = \{p_0, \dots, p_{|P|-1}\},\$$
$$Q = \{q_0, \dots, q_{|Q|-1}\}.$$

S. LABBÉ

We construct the alphabet $\mathcal{B} = \{0, 1, \dots, |Q| + |P| - 1\}$ and define the rule ω by

(6)
$$\omega: \mathcal{B} \to \mathcal{A}^{*^{d}}$$
$$j \mapsto \begin{cases} q_{j} & \text{if } 0 \leq j < |Q|, \\ p_{j-|Q|} & \text{if } |Q| \leq j < |Q| + |P| \end{cases}$$

We want to show that ω extends to a map from a set of configurations to X which is onto up to a shift. Let $x \in X$ be a configuration which can be seen as a function $x : \mathbb{Z}^d \to \mathcal{A}$. Consider the set $x^{-1}(M) \subset \mathbb{Z}^d$ of positions of markers in x. From the definition of markers in the direction e_i , markers appear in nonadjacent hyperplane orthogonal to e_i in the configuration x. Formally, there exists a set $H \subset \mathbb{Z}$ such that $x^{-1}(M) = \mathbb{Z}^{i-1} \times H \times \mathbb{Z}^{d-i}$ and $1 \notin H - H$. Since $1 \notin H - H$, there exists a strictly increasing sequence $(a_k)_{k \in \mathbb{Z}}$ such that $\mathbb{Z} \setminus H = \{a_k \mid k \in \mathbb{Z}\}$. We assume that $a_0 = 0$ if $0 \in \mathbb{Z} \setminus H$ and $a_0 = -1$ if $0 \in H$ which makes the sequence $(a_k)_{k \in \mathbb{Z}}$ uniquely defined.

In order to define the preimage of x under ω , it is convenient to represent x fiber by fiber. For every $\mathbf{m} \in \mathbb{Z}^{d-1}$, let $x_{\mathbf{m}} : \mathbb{Z} \to \mathcal{A}$, be the sequence such that

$$x(n_1, \ldots, n_{i-1}, n_i, n_{i+1}, \ldots, n_d) = x_{(n_1, \ldots, n_{i-1}, n_{i+1}, \ldots, n_d)}(n_i)$$

for every $\boldsymbol{n} = (n_1, \ldots, n_{i-1}, n_i, n_{i+1}, \ldots, n_d) \in \mathbb{Z}^d$. For every $\boldsymbol{m} \in \mathbb{Z}^{d-1}$, let $y_{\boldsymbol{m}} : \mathbb{Z} \to \mathcal{B}$ be defined as

$$y_{\boldsymbol{m}}: \ \mathbb{Z} \ \rightarrow \ \mathcal{B}$$

$$k \ \mapsto \begin{cases} j & \text{if } a_k + 1 \in H \text{ and } x_{\boldsymbol{m}}(a_k) = q_j, \\ j & \text{if } a_k + 1 \notin H \text{ and } x_{\boldsymbol{m}}(a_k) \odot^i x_{\boldsymbol{m}}(a_k + 1) = p_{j-|Q|}, \end{cases}$$

The function y_m is well-defined. Indeed, let $k \in \mathcal{A}$ and $m \in \mathbb{Z}^{d-1}$. If $a_k + 1 \in H$, then $x_m(a_k + 1) \in \mathcal{A} \setminus M$ and $y_m(k) = x_m(a_k) \in Q$. Also if $a_k + 1 \notin H$, then $x_m(a_k) \in \mathcal{A} \setminus M$ and $x_m(a_k + 1) \in M$. Since $x \in X$, then $x_m(a_k) \odot^i x_m(a_k + 1) \notin F$ and therefore $x_m(a_k) \odot^i x_m(a_k + 1) \in P$. We define the configuration $y : \mathbb{Z}^d \to \mathcal{B}$ by its fibers constructed above

$$y(n_1, \ldots, n_{i-1}, n_i, n_{i+1}, \ldots, n_d) = y_{(n_1, \ldots, n_{i-1}, n_{i+1}, \ldots, n_d)}(n_i)$$

for every $\boldsymbol{n} = (n_1, \dots, n_{i-1}, n_i, n_{i+1}, \dots, n_d) \in \mathbb{Z}^d$.

We may now finish the proof of surjectivity. If $0 \in H$, then the configuration x is exactly the image under ω of the configuration $y \in \mathcal{B}^{\mathbb{Z}^d}$ that we constructed: $x = \omega(y)$. If $0 \notin H$, then the configuration x is a shift of the image under ω of the configuration $y \in \mathcal{B}^{\mathbb{Z}^d}$: $x = \sigma^{e_i}\omega(y)$. Let

$$Y = \omega^{-1}(X) = \{ y \in \mathcal{B}^{\mathbb{Z}^d} \mid \omega(y) \in X \}$$

making $\omega : Y \to X$ a continuous map which is onto up to a shift. Notice that the subset Y is closed since ω is continuous and X is closed. Moreover, Y is shift-invariant since for all $y \in Y$ and $\mathbf{n} \in \mathbb{Z}^d$ there exists $\mathbf{k} \in \mathbb{Z}^d$ such that $\omega(\sigma^n(y)) = \sigma^k(\omega(y))$ meaning that $\sigma^n(y) \in Y$. Thus Y is a subshift.

The function ω is of the form

$$\omega(\mathcal{B}) \subseteq (\mathcal{A} \setminus M) \cup \left((\mathcal{A} \setminus M) \odot^{i} M \right)$$

and its restriction on \mathcal{B} is injective by construction. Therefore, we conclude from Lemma 3.12 that ω is recognizable in $\mathcal{B}^{\mathbb{Z}^d}$.

Remark that if X is an effective subshift one may also show that Y is an effective subshift. Moreover if X is a Wang shift, then Y is a Wang shift and the Wang tiles defining Y can be obtained from the Wang tiles defining X together with some fusion operation. This is what was done in [Lab19] and [Lab21c].

Exercise 3.13

Using the value of H and V from Exercise 3.4, prove that $\{0, 1, 2, 3, 4, 5, 6\}$ is a subset of markers for the direction e_2 in the subshift \mathcal{X}_{Φ} .

Exercise 3.14

Using the value of H and V from Exercise 3.4, prove that $\{0, 1, 7, 8, 9, 10\}$, is a subset of markers for the direction e_1 in the subshift \mathcal{X}_{Φ} .

Exercise 3.15

Using $M = \{0, 1, 2, 3, 4, 5, 6\}$ as subset of markers for the direction e_2 , find an alphabet \mathcal{B} , a subshift $Y \subset \mathcal{B}^{\mathbb{Z}^2}$ and a 2-dimensional morphism $\alpha : \mathcal{B} \to [\![0, 15]\!]^{*^2}$ such that

 $\alpha(\mathcal{B}) \subseteq (\llbracket 0, 15 \rrbracket \setminus M) \cup \left((\llbracket 0, 15 \rrbracket \setminus M) \odot^2 M \right)$

which extends to a recognizable continuous map $\alpha: Y \to \mathcal{X}_{\Phi}$ which is onto up to a shift.

Exercise 3.16

Using the subset $M = \{0, 1, 7, 8, 9, 10\} \subset \mathcal{A}$ of markers for the direction e_1 , find an alphabet \mathcal{C} , a subshift $Y' \subset \mathcal{C}^{\mathbb{Z}^2}$ and a 2-dimensional morphism $\xi : \mathcal{C} \to \llbracket 0, 15 \rrbracket^{*^2}$ such that

 $\xi(\mathcal{C}) \subseteq (\llbracket 0, 15 \rrbracket \setminus M) \cup \left((\llbracket 0, 15 \rrbracket \setminus M) \odot^1 M \right)$

which extends to a recognizable continuous map $\xi : Y' \to \mathcal{X}_{\Phi}$ which is onto up to a shift.

4. A Self-similar Wang shift

A Wang tile is a tuple of four colors $(a, b, c, d) \in I \times J \times I \times J$ where I is a finite set of vertical colors and J is a finite set of horizontal colors, see [Wan61, Rob71]. A Wang tile is represented as a unit square with colored edges:

$$\begin{bmatrix} c & b \\ d & a \end{bmatrix}$$

For each Wang tile $\tau = (a, b, c, d)$, let RIGHT $(\tau) = a$, TOP $(\tau) = b$, LEFT $(\tau) = c$, BOTTOM $(\tau) = d$ denote respectively the colors of the right, top, left and bottom edges of τ .

Let $\mathcal{T} = \{t_0, \ldots, t_{m-1}\}$ be a set of Wang tiles as the one shown in Figure 3. A configuration $x : \mathbb{Z}^2 \to \{0, \ldots, m-1\}$ is valid with respect to \mathcal{T} if it assigns a tile in \mathcal{T} to each position of \mathbb{Z}^2 so that contiguous edges of adjacent tiles have the same color, that is,

(7) $\operatorname{RIGHT}(t_{x(n)}) = \operatorname{LEFT}(t_{x(n+e_1)})$

(8)
$$\operatorname{TOP}(t_{x(n)}) = \operatorname{BOTTOM}(t_{x(n+e_2)})$$

$\begin{bmatrix} O \\ J & 0 \\ O \end{bmatrix} \begin{bmatrix} O \\ H & 1 \\ L \end{bmatrix}$	$ \begin{smallmatrix} M \\ D \begin{smallmatrix} 2 \\ 2 \\ P \end{smallmatrix} J$	$\begin{bmatrix} M \\ D \ 3 \ D \\ K \end{bmatrix}$	$\begin{bmatrix} P\\J & 4\\P \end{bmatrix} H$	$\stackrel{P}{\underset{N}{\overset{H}{\overset{5}{_{}}}}}_{H}$	${\rm D} \mathop{\rm D}_P^K {\rm H}$	$\begin{bmatrix} 0\\ I & 7 & B\\ O \end{bmatrix}$
$\begin{bmatrix} L \\ E & 8 \\ O \end{bmatrix} \begin{bmatrix} L \\ C & 9 \\ L \end{bmatrix}$	L I 10A O	P I 11E P	$\begin{smallmatrix} P\\I12I\\K\end{smallmatrix}$	$\overset{K}{\overset{B13I}{\overset{M}{}}}$	K A14 I K	$\begin{tabular}{c} N \\ I \ 15C \\ P \end{tabular}$

FIGURE 3. The set $\mathcal{Z} = \{u_0, \ldots, u_{15}\}$ of 16 Wang tiles is a simplification made by Jana Lepšová [Lep24] of the set \mathcal{U} of 19 Wang tiles introduced in [Lab19]. Each tile is identified uniquely by an index from the set $\{0, 1, \ldots, 15\}$ written at the center each tile.

for every $\mathbf{n} \in \mathbb{Z}^2$ where $\mathbf{e}_1 = (1,0)$ and $\mathbf{e}_2 = (0,1)$. A finite pattern which is valid with respect to \mathcal{Z} is shown in Figure 4.

	${}^{\mathrm{K}}_{\mathrm{D}\mathrm{6}\mathrm{H}}$	O H1D	${}^{\mathrm{K}}_{\mathrm{D}\mathrm{6}\mathrm{H}}$	$^{P}_{H\frac{5}{2}H}$	O H1D
(61651)	P P I11E	L L E 8 I	Р Р 1121	N N I 15C	$\frac{L}{C 9 I}$
11 8 12 15 9	P P	$\begin{array}{c} 0\\ 0\\ 170 \end{array}$	K K D121	P P	L L
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	K	$\begin{array}{c} 1 & l & B \\ 0 \\ \hline 0 \\ \end{array}$	M M	P P	$\begin{array}{c} \text{E & 1} \\ 0 \\ \end{array}$
(11 8 12 11 8)	D6H P	HĨD L	D2J P	J 4 H P	HĨD L
	P I 11E P	E ^L 8I O	Р I 12 I К	P I 11E P	$\mathrm{E}_{\mathrm{O}}^{\mathrm{L}}$ I

FIGURE 4. The finite 5×5 pattern on the left is valid with respect to \mathcal{Z} since it respects Equations (7) and (8) which we can verify on the tiling shown on the right.

Let $\Omega_{\mathcal{T}} \subset \{0, \ldots, m-1\}^{\mathbb{Z}^2}$ denote the set of all valid configurations with respect to \mathcal{T} , called the **Wang shift** of \mathcal{T} . To a configuration $x \in \Omega_{\mathcal{T}}$ corresponds a tiling of the plane \mathbb{R}^2 by the tiles \mathcal{T} where the unit square Wang tile $t_{x(n)}$ is placed at position n for every $n \in \mathbb{Z}^2$, as in Figure 4. Together with the shift action σ of \mathbb{Z}^2 , $\Omega_{\mathcal{T}}$ is a SFT of the form (3) since there exists a finite set of forbidden patterns made of all horizontal and vertical dominoes of two tiles that do not share an edge of the same color. This definition of Wang shifts allows to use the concepts of languages, 2-dimensional morphisms, recognizability introduced in the previous sections.

A configuration $x \in \Omega_{\mathcal{T}}$ is **periodic** if there exists $\mathbf{n} \in \mathbb{Z}^2 \setminus \{0\}$ such that $x = \sigma^{\mathbf{n}}(x)$. A set of Wang tiles \mathcal{T} is **periodic** if there exists a periodic configuration $x \in \Omega_{\mathcal{T}}$. Originally, Wang thought that every set of Wang tiles \mathcal{T} is periodic as soon as $\Omega_{\mathcal{T}}$ is nonempty [Wan61]. This statement is equivalent to the existence of an algorithm solving the *domino problem*, that is, taking as input a set of Wang tiles and returning *yes* or *no* whether there exists a valid configuration with these tiles. Berger, a student of Wang, later proved that the domino problem is undecidable and he also provided a first example of an aperiodic set of Wang tiles [Ber66]. A set of Wang tiles \mathcal{T} is **aperiodic** if the Wang shift $\Omega_{\mathcal{T}}$ is a nonempty aperiodic subshift. This means that in general one can not decide the emptiness of a Wang shift $\Omega_{\mathcal{T}}$. This illustrates that the behavior of *d*-dimensional SFTs when $d \geq 2$ is much different than the one-dimensional case where emptiness of a SFT is decidable [LM95]. Note that another important difference between d = 1 and $d \ge 2$ is expressed in terms of the possible values of entropy of d-dimensional SFTs, see [HM10].

The goal of this section is to prove Theorem 1.1, i.e., that the Wang shift $\Omega_{\mathcal{Z}} \subset \{0, \ldots, 15\}^{\mathbb{Z}^2}$ defined by the set of Wang tiles \mathcal{Z} shown in Figure 3 is self-similar where the self-similarity is given by the 2-dimensional morphism Φ defined in Equation (1).

Exercise 4.1

Find a valid 7×7 tiling with the set \mathcal{Z} of Wang tiles.

4.1. Markers in the context of Wang tilings. A tiling with the tiles from the set \mathcal{Z} is shown in Figure 5. It illustrates that there exists a subset $M \subset \mathcal{Z}$ of tiles such that each horizontal row of tiles in the tiling is using either tiles from M or from $\mathcal{Z} \setminus M$. Moreover, the horizontal lines using tiles from M are nonadjacent. If these conditions are satisfied for all configurations in $\Omega_{\mathcal{Z}}$, then M is a subset of markers in the direction e_2 .

$\begin{smallmatrix} P\\J \ 4 \\ P\\ \end{smallmatrix}$	$\begin{smallmatrix}&O\\H1D\\L\end{smallmatrix}$	${}^{\mathrm{K}}_{\mathrm{P}}$	${}^{O}_{L}$	M D 3 D K
P I 11E P	$\mathrm{E}_{\mathrm{O}}^{\mathrm{L}}$ I	P I 12 I K	L I 10A O	K A14 I K
$\begin{smallmatrix} P\\I 12 I\\K\end{smallmatrix}$	I 7 B O	$\mathop{\mathrm{B13I}}_{\mathrm{M}}^{\mathrm{K}}$	$\begin{smallmatrix}&0\\I&7&B\\0\end{smallmatrix}$	$\mathop{\mathrm{B13I}}_{\mathrm{M}}^{\mathrm{K}}$
${}^{\mathrm{K}}_{\mathrm{P}}{}^{\mathrm{H}}$	${}^{\rm O}_{\rm H1D}_{\rm L}$	${}^{\mathrm{M}}_{\mathrm{P}}\mathrm{J}$	$\begin{smallmatrix} 0\\J & 0\\0\\0\end{smallmatrix}$	D 3 D K
P I 11E P	$\mathbf{E} \stackrel{\mathrm{L}}{\underset{\mathrm{O}}{\overset{\mathrm{S}}{\mathrm{I}}}} \mathbf{I}$	P I 12 I K	$I \stackrel{O}{7} B$	К B13 I M
T	0		~	
P H 5 H N	O H1D L	D 6 H P	O H1D L	$\stackrel{\mathrm{M}}{\overset{\mathrm{D}}{_2}}_{\mathrm{P}}^{\mathrm{M}}$
$\frac{P}{H \frac{5}{5}H}$ $\frac{N}{I \frac{15}{2}C}$	$ \begin{array}{c} 0\\ H 1 D\\ L\\ C 9 I\\ L \end{array} $	и D 6 H P I 11E P		M D 2 J P I 12 I K
$\frac{P}{H \frac{5}{5}H}$ $\frac{N}{I \frac{15}{5}C}$ $P}{I \frac{11}{11}E}$ P	$ \begin{array}{c} $	R K D 6 H P I 11E P I 12I K	$ \begin{bmatrix} 0 \\ H \\ L \\ E \\ 8 \\ I \\ 0 \\ I \\ 7 \\ 0 \\ B \\ 0 $	M D 2 J P I 12I K B13I M
$\frac{P}{H 5 H}$ $\frac{N}{N}$ $\frac{115C}{P}$ $\frac{P}{I 11E}$ $\frac{P}{J 4 H}$	$ \begin{array}{c} 0 \\ H \stackrel{1}{} 1 \\ L \\ C \stackrel{9}{} 1 \\ L \\ E \stackrel{1}{} 8 \\ I \\ O \\ H \stackrel{1}{} 1 \\ D \\ L \end{array} $	$ \frac{K}{D \stackrel{6}{6}H} $ $ \frac{P}{I \stackrel{11E}{12I}} $ $ \frac{K}{D \stackrel{6}{6}H} $	H I D C H I D C H I D C H I D C H I D C C C C C C C C C C C C C C C C C C	M D 2 J P I 12 I K B13 I M D 3 D K

FIGURE 5. A 5 × 10 tiling with tiles from the set \mathcal{Z} . The tiles labeled from 0 to 6 (shown with yellow background) are marker tiles for the direction e_2 in the Wang shift $\Omega_{\mathcal{Z}}$ since they always appear on nonadjacent rows.

In this section we propose an algorithm to find and prove that a subset of tiles is a subset of markers in a Wang shift. We use Lemma 3.11 which provides a way to prove that a subset of Wang tiles is a subset of markers and searching for them. To use it in the context of Wang shifts and more generally in the context of SFTs, we need the following definition.

Definition 4.1 (surrounding of radius r). Let $X = SFT(\mathcal{F}) \subset \mathcal{A}^{\mathbb{Z}^2}$ be a shift of finite type for some finite set \mathcal{F} of forbidden patterns. A 2-dimensional word $u \in \mathcal{A}^n$, with n =

 $(n_1, n_2) \in \mathbb{N}^2$, admits a **surrounding of radius** $r \in \mathbb{N}$ if there exists $w \in \mathcal{A}^{n+2(r,r)}$ such that u occurs in w at position (r, r) and w contains no occurrences of forbidden patterns from \mathcal{F} .

If a word admits a surrounding of radius $r \in \mathbb{N}$, it does not mean it is in the language of the SFT. But if it admits no surrounding of radius r for some $r \in \mathbb{N}$, then for sure it is not in the language of the SFT. We state the following lemma in the context of Wang tiles.

Lemma 4.2. Let \mathcal{T} be a set of Wang tiles and $u \in \mathcal{T}^n$ be a rectangular pattern seen as a 2-dimensional word with $\mathbf{n} = (n_1, n_2) \in \mathbb{N}^2$. If u is allowed in $\Omega_{\mathcal{T}}$, then for every $r \in \mathbb{N}$ the word u has a \mathcal{T} -surrounding of radius r.

Equivalently the lemma says that if there exists $r \in \mathbb{N}$ such that u has no \mathcal{T} -surrounding of radius r, then u is forbidden in $\Omega_{\mathcal{T}}$ and this is how we use Lemma 4.2 to find markers. We propose Algorithm 1 to compute markers from a Wang tile set and a chosen surrounding radius so that the computation terminates. If the algorithm finds nothing, then maybe there are no markers or maybe one should try again after increasing the surrounding radius. We prove in the next lemma that if the output is nonempty, it contains a subset of markers.

Algorithm 1 Find markers. If no markers are found, one should try increasing the radius r.

Precondition: \mathcal{T} is a set of Wang tiles; $i \in \{1, 2\}$ is a direction e_i ; $r \in \mathbb{N}$ is some radius. 1: function FINDMARKERS (\mathcal{T}, i, r)

2: $j \leftarrow 3 - i$

3: $D_i \leftarrow \{(u, v) \in \mathcal{T}^2 \mid \text{ domino } u \odot^j v \text{ admits a } \mathcal{T}\text{-surrounding of radius } r\}$

- 4: $U \leftarrow \{\{u\} \mid u \in \mathcal{T}\}$ \triangleright Suggestion: use a union-find data structure
- 5: for all $(u, v) \in D_j$ do

6: Merge the sets containing u and v in the partition U.

7: $D_i \leftarrow \{(u, v) \in \mathcal{T}^2 \mid \text{ domino } u \odot^i v \text{ admits a } \mathcal{T}\text{-surrounding of radius } r\}$

8: **return** {set M in the partition $U \mid (M \times M) \cap D_i = \emptyset$ }

Postcondition: The output contains zero, one or more subsets of markers in the direction e_i .

Lemma 4.3. If there exists $r \in \mathbb{N}$ and $i \in \{1, 2\}$ such that the output of FINDMARKERS (\mathcal{T}, i, r) contains a set M, then $M \subset \mathcal{T}$ is a subset of markers in the direction e_i .

Proof. Suppose that i = 2, the case i = 1 being similar. The output set M is nonempty since it was created from the union of nonempty sets (see lines 4-6 in Algorithm 1). Using Lemma 4.2, lines 3 to 6 imply that $M \odot^1 (\mathcal{T} \setminus M)$ and $(\mathcal{T} \setminus M) \odot^1 M$ are forbidden in $\Omega_{\mathcal{T}}$. The lines 7 and 8 imply that $M \odot^2 M$ is forbidden in $\Omega_{\mathcal{T}}$. Then we deduce from Lemma 3.11 that $M \subset \mathcal{T}$ is a subset of markers in the direction e_2 .

We believe that if a set of Wang tiles \mathcal{T} has a subset of markers in the direction e_i then there exists a surrounding radius $r \in \mathbb{N}$ such that FINDMARKERS (\mathcal{T}, i, r) outputs this set of markers, so that Lemma 4.3 is in fact a *if and only if* but we do not provide a proof of that here. The fact that there is no upper bound for the surrounding radius is related to the undecidability of the domino problem. In practice, in the study of Jeandel–Rao tilings done in [Lab21c], a surrounding radius of 1, 2 or 3 was enough.

Exercise 4.2

Using the set \mathcal{Z} of Wang tiles defined in Figure 3, compute the sets of horizontal and vertical dominoes that admit a \mathcal{Z} -surrounding of radius 2:

 $D_1 = \left\{ (u, v) \in \mathbb{Z}^2 \mid u \odot^1 v \text{ admits a } \mathbb{Z} \text{-surrounding of radius } 2 \right\},\$

$$D_2 = \{(u, v) \in \mathbb{Z}^2 \mid u \odot^2 v \text{ admits a } \mathbb{Z}\text{-surrounding of radius } 2\}.$$

Compare them with the set H and V found in Exercise 3.4.

Exercise 4.3

Use the function FINDMARKERS defined in Algorithm 1 with a surrounding radius 2 to show that the subset

$$M = \left\{ \begin{array}{c} O \\ J & 0 \\ O \end{array} \begin{bmatrix} O \\ H & 1 \\ L \end{bmatrix} \begin{bmatrix} M \\ D & 2 \\ P \end{bmatrix} \begin{bmatrix} M \\ D & 3 \\ R \end{bmatrix} \begin{bmatrix} P \\ J & 4 \\ P \end{bmatrix} \begin{bmatrix} P \\ H & 5 \\ N \end{bmatrix} \begin{bmatrix} K \\ D & 6 \\ P \end{bmatrix} \right\}$$

of \mathcal{Z} is a subset of markers for the direction e_2 in $\Omega_{\mathcal{Z}}$.

4.2. Fusion of Wang tiles. Recall that a magma is a set \mathcal{I} equipped with a binary operation • such that for all $a, b \in \mathcal{I}$, the result of the operation $a \bullet b$ is also in \mathcal{I} . If the operation • is associative and has an identity, then \mathcal{I} is a monoid and the operation • can be omitted and represented as concatenation. But, in the general context of fusion of Wang tiles defined below where \mathcal{I} is the set of horizontal or vertical colors, we can not assume that the operation • is associative.

The fusion operation on Wang tiles is defined on pair of tiles sharing an edge in a tiling according to Equations 7 and 8. Let (\mathcal{I}, \bullet) and (\mathcal{J}, \bullet) be two magmas and let $\{A, C, Y, W\} \subset \mathcal{I}$ be some vertical colors and $\{B, D, X, Z\} \subset \mathcal{J}$ be some horizontal colors. We define two binary operations \square and \square on Wang tiles as

$$\begin{bmatrix} B \\ C \\ D \end{bmatrix} \square \begin{bmatrix} X \\ Y \\ W \\ Z \end{bmatrix} = \begin{bmatrix} B \bullet X \\ C \\ W \\ D \bullet Z \end{bmatrix} \text{ if } A = Y$$

and

$$\begin{bmatrix} B \\ C \\ D \end{bmatrix} \ \ \Box \ \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} X \\ \bullet \\ O \\ D \\ \bullet \end{bmatrix} \text{ if } B = Z.$$

If $A \neq Y$, the operation \square is not defined. Similarly, if $B \neq Z$, the operation \square is not defined. For the Wang tiles considered in this contribution, the operation \bullet is associative so we always denote it implicitly by concatenation of colors.

In what follows, we propose algorithms and results that works for both operations \square and \square . It is thus desirable to have a common notation to denote both, so we define

$$u \boxtimes^1 v = u \boxplus v$$
 and $u \boxtimes^2 v = u \boxminus v$.

If $u \boxtimes^i v$ is defined for some $i \in \{1, 2\}$, it means that tiles u and v can appear at position \boldsymbol{n} and $\boldsymbol{n} + \boldsymbol{e}_i$ in a tiling for some $\boldsymbol{n} \in \mathbb{Z}^d$. For each $i \in \{1, 2\}$, one can define a new set of tiles from two sets \mathcal{T} and \mathcal{S} of Wang tiles as

$$\mathcal{T} \boxtimes^{i} \mathcal{S} = \{ u \boxtimes^{i} v \text{ defined } \mid u \in \mathcal{T}, v \in \mathcal{S} \}.$$

Exercise 4.4

Using the set D_2 of dominoes that admits a \mathcal{Z} -surrounding of radius 2 computed in Exercise 4.2 and the subset $M \subset \mathcal{Z}$ of markers for the direction e_2 in $\Omega_{\mathcal{Z}}$ computed in Exercise 4.3, compute the set of fusion tiles:

$$\{u \boxminus v \mid (u, v) \in D_2 \text{ and } v \in M\}.$$

What is the meaning of this set?

4.3. **Desubstitution of Wang shifts.** In Theorem 3.13, we proved that the presence of markers allows to desubstitute uniquely the configurations of a subshift on \mathbb{Z}^d . In case of Wang shifts, we show in this section that the preimage is also a Wang shift and we may construct the preimage set of Wang tiles using the fusion operation defined in the previous section. We also propose an algorithm to find the desubstitution of Wang shifts when there exists a subset of marker tiles.



FIGURE 6. A 5×10 tiling with tiles from the set \mathcal{Z} is shown on the left. The tiles labeled from 0 to 6 (shown with yellow background) are marker tiles for the direction e_2 since they appear on nonadjacent rows. It can be desubstituted as a 5×6 pattern with tiles from the set \mathcal{V} using a substitution α_0 . Each marker tile (yellow background) is glued with its below tile (green background) to form a new Wang tile (blue background) using the fusion operation \boxminus . The remaining tiles are kept the same (green background) but get new indices in the set \mathcal{V} . This process is uniquely defined since the substitution α_0 is recognizable.

Before stating the result, let us see how the markers allow to desubstitute tilings. In Figure 6. we observe that markers M computed in Exercise 4.3 appear as nonadjacent rows in the Wang shift $\Omega_{\mathcal{Z}}$. Therefore the row above (and below) a row of markers is made of nonmarker tiles. Let us consider the row below. The idea is to collapse that row with the row of markers just above. Each tile is being collapsed with the above marker tile using the fusion of tiles. The set of tiles that we obtain through this process is exactly the set computed in Exercise 4.4.

Therefore to build a configuration in $\Omega_{\mathcal{Z}}$, it is sufficient to build a tiling with another set \mathcal{V} of Wang tiles obtained from the set \mathcal{Z} after removing the markers and adding the tiles obtained from the fusion operation. One may also remove the tile which always appear below of a marker tile. One may recover some configuration in $\Omega_{\mathcal{Z}}$ by applying a 2-dimensional morphism $\alpha_0 : \mathcal{V} \to \mathcal{Z}^{*^2}$ which replaces the merged tiles by their associated equivalent vertical dominoes and keeps the remaining tiles invariant, see Figure 6. It turns out that this decomposition is unique. The creation of the set \mathcal{V} from \mathcal{Z} gives the intuition on the construction of Algorithm 2 which follows the same recipe and takes any set of Wang tiles with markers as input.

We now state the result that if a set of Wang tiles \mathcal{T} has a subset of marker tiles, then there exists another set \mathcal{S} of Wang tiles and a nontrivial recognizable 2-dimensional morphism $\Omega_{\mathcal{S}} \to \Omega_{\mathcal{T}}$ that is onto up to a shift. Thus, every Wang tiling by \mathcal{T} is up to a shift the image under a nontrivial 2-dimensional morphism ω of a unique Wang tiling in $\Omega_{\mathcal{S}}$. The 2-dimensional morphism is essentially 1-dimensional as we show in the next theorem.

Theorem 4.4. [Lab19, Lab21c] Let \mathcal{T} be a set of Wang tiles and let $\Omega_{\mathcal{T}}$ be its Wang shift. If there exists a subset $M \subset \mathcal{T}$ of markers in the direction e_i for $i \in \{1, 2\}$, then

(1) there exists a set of Wang tiles S_R and a 2-dimensional morphism $\omega_R : \Omega_{S_R} \to \Omega_T$ such that

$$\omega_R(\mathcal{S}_R) \subseteq (\mathcal{T} \setminus M) \cup \left((\mathcal{T} \setminus M) \odot^i M \right)$$

which is recognizable and onto up to a shift and

(2) there exists a set of Wang tiles S_L and a 2-dimensional morphism $\omega_L : \Omega_{S_L} \to \Omega_T$ such that

$$\omega_L(\mathcal{S}_L) \subseteq (\mathcal{T} \setminus M) \cup \left(M \odot^i (\mathcal{T} \setminus M)\right)$$

which recognizable and onto up to a shift.

There exists a surrounding radius $r \in \mathbb{N}$ such that ω_R and ω_L are computed using Algorithm 2.

Proof. The existence of the recognizable 2-dimensional morphism which is onto up to a shift was done in Theorem 3.13. We only need to prove that the preimage of Ω_{τ} is a Wang shift. The proof of this fact can be found in [Lab19] and [Lab21c]. It follows the line of Algorithm 2.

In the definition of ω in Algorithm 2, given two Wang tiles u and v such that $u \boxtimes^i v$ is defined for $i \in \{1, 2\}$, the map

$$u \boxtimes^i v \quad \mapsto \quad u \odot^i v$$

can be seen as a decomposition of Wang tiles:

Algorithm 2 Find a recognizable desubstitution of $\Omega_{\mathcal{T}}$ from markers

- **Precondition:** \mathcal{T} is a set of Wang tiles; $M \subset \mathcal{T}$ is a subset of markers; $i \in \{1, 2\}$ is a direction e_i ; $r \in \mathbb{N}$ is a surrounding radius; $s \in \{\text{LEFT}, \text{RIGHT}\}$ determines whether the image of merged tiles is in $M \odot^i(\mathcal{T} \setminus M)$ (markers on the left) or in $(\mathcal{T} \setminus M) \odot^i M$ (markers on the right).
- 1: function FINDSUBSTITUTION(\mathcal{T}, M, i, r, s)
- 2: $D \leftarrow \{(u, v) \in \mathcal{T}^2 \mid \text{ domino } u \odot^i v \text{ admits a } \mathcal{T}\text{-surrounding of radius } r\}$
- 3: **if** s = LEFT then
- 4: $P \leftarrow \{(u, v) \in D \mid u \in M \text{ and } v \in \mathcal{T} \setminus M\}$
- 5: $K \leftarrow \{v \in \mathcal{T} \setminus M \mid \text{ there exists } u \in \mathcal{T} \setminus M \text{ such that } (u, v) \in D\}$
- 6: else if s = RIGHT then
- 7: $P \leftarrow \{(u, v) \in D \mid u \in \mathcal{T} \setminus M \text{ and } v \in M\}$
- 8: $K \leftarrow \{u \in \mathcal{T} \setminus M \mid \text{ there exists } v \in \mathcal{T} \setminus M \text{ such that } (u, v) \in D\}$
- 9: $K \leftarrow \text{SORT}(K), P \leftarrow \text{SORT}(P)$ \triangleright lexicographically on the indices of tiles
- 10: $\mathcal{S} \leftarrow K \cup \{ u \boxtimes^i v \mid (u, v) \in P \}$ \triangleright defines uniquely indices of tiles in \mathcal{S} from 0 to $|\mathcal{S}| 1$.

11: **return**
$$\mathcal{S}, \omega : \Omega_{\mathcal{S}} \to \Omega_{\mathcal{T}} : \begin{cases} u \boxtimes^{i} v \quad \mapsto \quad u \odot^{i} v \quad \text{if } (u, v) \in P \\ u \quad \mapsto \quad u \quad & \text{if } u \in K. \end{cases}$$

Postcondition: S is a set of Wang tiles; $\omega : \Omega_S \to \Omega_T$ is recognizable and onto up to a shift.

whether i = 1 or i = 2 and where A = Y and B = Z. The reader may wonder how the substitution decides the color A (color B if i = 2) from its input tiles. The answer is that Algorithm 2 is performing a desubstitution. Therefore the two tiles sharing the vertical side with letter A are known from the start and the algorithm just creates a new tile (W, BX, C, DZ) and claims that it will always get replaced by the two tiles with shared edge with color A.

Exercise 4.5

Using the function FINDSUBSTITUTION defined in Algorithm 2 with the subset $M \subset \mathcal{Z}$ of markers for the direction e_2 computed in Exercise 4.3, construct a set of tiles \mathcal{V} and a recognizable 2-dimensional morphism $\alpha_0 : \Omega_{\mathcal{V}} \to \Omega_{\mathcal{Z}}$ which is onto up to a shift and such that

$$\alpha_0(u) \in \mathcal{Z} \setminus M$$
 or $\alpha_0(u) \in (\mathcal{Z} \setminus M) \odot^2 M$.

4.4. Self-similarity of the Wang shift $\Omega_{\mathcal{Z}}$ defined by 16 tiles. We prove that $\Omega_{\mathcal{Z}}$ is self-similar by executing the function FINDMARKERS on the set of Wang tiles \mathcal{Z} followed by FINDSUBSTITUTION and repeating this process until we obtain a set of Wang tiles which is equivalent to the original one (two steps are needed). Each time FINDMARKERS finds at least one subset of markers using a surrounding radius of size at most 2. Thus using Theorem 4.4, we find a desubstitution of tilings with FINDSUBSTITUTION. The proof is done in SageMath [Sag24] using slabbe optional package [Lab23b].

The following result was first shown in [Lep24] where it was also proved that $\Omega_{\mathcal{Z}}$ is aperiodic and minimal. It was deduced by showing that $\Omega_{\mathcal{Z}}$ is topologically conjugate to $\Omega_{\mathcal{U}}$. Selfsimilarity, aperiodicity and minimality of $\Omega_{\mathcal{U}}$ was shown in [Lab19]. In [Lab21c], it was shown that the set $\Omega_{\mathcal{U}}$ describes the internal self-similar structure hidden in Jeandel–Rao aperiodic tilings [JR21].

A consequence of Theorem 1.1 is that the Wang shift $\Omega_{\mathcal{Z}}$ provides another description for the substitutive subshift \mathcal{X}_{Φ} , see Exercise 4.6.

Proof of Theorem 1.1. In this proof, we show there exist sets of Wang tiles \mathcal{V} and \mathcal{W} together with their associated Wang shifts $\Omega_{\mathcal{V}}$ and $\Omega_{\mathcal{W}}$ and there exist two recognizable 2-dimensional morphisms α_0 and α_1 and a bijection α_2 :

$$\Omega_{\mathcal{Z}} \xleftarrow{\alpha_0}{\leftarrow} \Omega_{\mathcal{V}} \xleftarrow{\alpha_1}{\leftarrow} \Omega_{\mathcal{W}} \xleftarrow{\alpha_2}{\leftarrow} \Omega_{\mathcal{Z}}$$

that are onto up to a shift, i.e., $\overline{\alpha_0(\Omega_{\mathcal{V}})}^{\sigma} = \Omega_{\mathcal{Z}}$, $\overline{\alpha_1(\Omega_{\mathcal{W}})}^{\sigma} = \Omega_{\mathcal{V}}$ and $\alpha_2(\Omega_{\mathcal{Z}}) = \Omega_{\mathcal{W}}$. First we define the set \mathcal{Z} of Wang tiles in SageMath:

<pre>sage: from slabbe import WangTileSet</pre>	12
<pre>sage: tiles = ["DOJO", "DOHL", "JMDP", "DMDK", "HPJP", "HPHN", "HKDP", "BOIO",</pre>	13
<pre>: "ILEO", "ILCL", "ALIO", "EPIP", "IPIK", "IKBM", "IKAK", "CNIP"]</pre>	14
<pre>sage: Z = WangTileSet([tuple(tile) for tile in tiles])</pre>	15

7 -	$\begin{pmatrix} 0 \\ J & 0 \\ 0 \\ 0 \end{pmatrix}$	$\begin{bmatrix} O \\ H \ 1 \\ L \end{bmatrix}$	$\begin{bmatrix} M \\ D \stackrel{A}{_2} J \\ P \end{bmatrix}$	$\begin{bmatrix} M \\ D \ 3 \ D \\ K \end{bmatrix}$	$\begin{bmatrix} P\\J \stackrel{A}{4}H\\P \end{bmatrix}$	$\begin{bmatrix} P\\H {5 \atop N} H\\N \end{bmatrix}$	${\rm D} \mathop{\rm G}_{\rm P}^{\rm K} {\rm H}$	$\begin{bmatrix} 0\\ I & 7 & B\\ 0 \end{bmatrix}$	
2 -	E 8 I O	$\begin{tabular}{c} L \\ C & 9 \\ L \\ I \\ L \\ end{tabular}$	L I 10A O	P I 11E P	$\begin{bmatrix} P\\I12I\\K\end{bmatrix}$	$\begin{bmatrix} K \\ B13I \\ M \end{bmatrix}$	$ \begin{smallmatrix} K \\ A14 I \\ K \end{smallmatrix} $	$\begin{bmatrix} N\\ I15C\\ P \end{bmatrix}$	

We desubstitute \mathcal{Z} with the set $\{0, 1, 2, 3, 4, 5, 6\}$ of markers in the direction e_2 :

<pre>sage: Z.find_markers(i=2,radius=2,solver="dancing_links")</pre>		
[[0, 1, 2, 3, 4, 5, 6]]	17	
sage: M = [0, 1, 2, 3, 4, 5, 6]	18	
<pre>sage: V,alpha0 = Z.find_substitution(M, i=2, radius=2, side="right",</pre>	19	
: solver="dancing_links")	20	

We obtain $\alpha_0 : \Omega_{\mathcal{V}} \to \Omega_{\mathcal{Z}}$ given as a rule of the form

$$\begin{aligned} \alpha_{0} : & [\![0,17]\!] \to [\![0,15]\!]^{*^{2}} \\ & \begin{cases} 0 \mapsto (7) \,, & 1 \mapsto (8) \,, & 2 \mapsto (10) \,, & 3 \mapsto (11) \,, & 4 \mapsto (12) \,, \\ 5 \mapsto (13) \,, & 6 \mapsto (14) \,, & 7 \mapsto \begin{pmatrix} 0 \\ 7 \end{pmatrix} \,, & 8 \mapsto \begin{pmatrix} 1 \\ 8 \end{pmatrix} \,, & 9 \mapsto \begin{pmatrix} 1 \\ 9 \end{pmatrix} \,, \\ 10 \mapsto \begin{pmatrix} 1 \\ 10 \end{pmatrix} \,, 11 \mapsto \begin{pmatrix} 4 \\ 11 \end{pmatrix} \,, 12 \mapsto \begin{pmatrix} 6 \\ 11 \end{pmatrix} \,, 13 \mapsto \begin{pmatrix} 2 \\ 12 \end{pmatrix} \,, 14 \mapsto \begin{pmatrix} 6 \\ 12 \end{pmatrix} \,, \\ 15 \mapsto \begin{pmatrix} 3 \\ 13 \end{pmatrix} \,, 16 \mapsto \begin{pmatrix} 3 \\ 14 \end{pmatrix} \,, 17 \mapsto \begin{pmatrix} 5 \\ 15 \end{pmatrix} \,. \end{aligned}$$

and the set \mathcal{V} of 18 Wang tiles



The set of tiles \mathcal{V} has three subsets of markers for the direction e_1 . We desubstitute \mathcal{V} with the subset of markers $\{0, 1, 2, 8, 9, 10, 11\}$:

<pre>sage: V.find_markers(i=1,radius=1,solver="dancing_links")</pre>		
[[0, 1, 2, 7, 8, 9, 10]]	22	
sage : $M = [0, 1, 2, 7, 8, 9, 10]$	23	
<pre>sage: W,alpha1 = V.find_substitution(M, i=1, radius=1, side="right",</pre>	24	
<pre>: solver="dancing_links")</pre>	25	

We obtain $\alpha_1 : \Omega_{\mathcal{W}} \to \Omega_{\mathcal{V}}$ given as a rule of the form

$$\begin{aligned} \alpha_1 : & \llbracket 0, 15 \rrbracket \to \llbracket 0, 17 \rrbracket^{*^2} \\ \begin{cases} 0 \mapsto (5) , & 1 \mapsto (6) , & 2 \mapsto (13) , & 3 \mapsto (14) , \\ 4 \mapsto (15) , & 5 \mapsto (16) , & 6 \mapsto (3, 1) , & 7 \mapsto (4, 0) , \\ 8 \mapsto (4, 2) , & 9 \mapsto (5, 0) , & 10 \mapsto (6, 0) , & 11 \mapsto (11, 8) , \\ 12 \mapsto (12, 8) , 13 \mapsto (13, 7) , 14 \mapsto (14, 10) , 15 \mapsto (17, 9) . \end{aligned}$$

and the set \mathcal{W} of 16 Wang tiles

It turns out that \mathcal{Z} and \mathcal{W} are equivalent. The proof can be done using graph isomorphisms:

```
sage: W.is_equivalent(Z)
True
```

26 27

The bijection **vert** between the vertical colors, the bijection **horiz** between the horizontal colors and bijection α_2 from \mathcal{Z} to \mathcal{W} is computed as follows:

```
sage: _,vert,horiz,alpha2 = Z.is_equivalent(W, certificate=True) 28
sage: vert 29
{'J': 'A', 'D': 'I', 'H': 'B', 'I': 'ID', 'B': 'IH', 'A': 'IJ', 'E': 'AD', 'C': 'BD'} 30
sage: horiz 31
{'O': 'K', 'L': 'M', 'P': 'KO', 'M': 'PL', 'K': 'PO', 'N': 'MO'} 32
```

We obtain the morphism $\alpha_2 : \Omega_{\mathcal{Z}} \to \Omega_{\mathcal{W}}$ given as a rule of the form

$$\begin{aligned} \alpha_2: & \llbracket 0, 15 \rrbracket \to \llbracket 0, 15 \rrbracket^{*^2} \\ & \begin{cases} 0 \mapsto (1), \quad 1 \mapsto (0), \quad 2 \mapsto (8), \quad 3 \mapsto (6), \\ 4 \mapsto (10), \quad 5 \mapsto (9), \quad 6 \mapsto (7), \quad 7 \mapsto (3), \\ 8 \mapsto (5), \quad 9 \mapsto (4), \quad 10 \mapsto (2), \quad 11 \mapsto (14), \\ 12 \mapsto (12), 13 \mapsto (15), 14 \mapsto (11), 15 \mapsto (13). \end{aligned}$$

We may check that $\alpha_0 \circ \alpha_1 \circ \alpha_2 = \Phi$ where the variable Phi was created in Exercise 3.3:

sage:	alpha0 * alpha1	* alpha2 ==	Phi 3	33
True			3	34

We conclude that
$$\Omega_{\mathcal{Z}} = \overline{\alpha_0(\Omega_{\mathcal{V}})}^{\sigma} = \overline{\alpha_0\alpha_1(\Omega_{\mathcal{W}})}^{\sigma} = \overline{\alpha_0\alpha_1\alpha_2(\Omega_{\mathcal{Z}})}^{\sigma} = \overline{\Phi(\Omega_{\mathcal{Z}})}^{\sigma}.$$

In the proof, we used Knuth's dancing links algorithm [Knu00] because it is faster at this particular task than the MILP solver Gurobi [GO20] or the SAT solvers Glucose [AS18] as we can see below:

```
sage: %time Z.find_markers(i=2,radius=2,solver="dancing_links")
CPU times: user 3.34 s, sys: 0 ns, total: 3.34 s
Wall time: 3.34 s
[[0, 1, 2, 3, 4, 5, 6]]
sage: %time Z.find_markers(i=2,radius=2,solver="gurobi")
CPU times: user 12.4 s, sys: 572 ms, total: 13 s
Wall time: 13 s
[[0, 1, 2, 3, 4, 5, 6]]
sage: %time Z.find_markers(i=2,radius=2,solver="glucose")
CPU times: user 50.6 s, sys: 2.53 s, total: 53.1 s
Wall time: 2min 10s
[[0, 1, 2, 3, 4, 5, 6]]
```

Note that for other tasks like finding a valid tiling of a $n \times n$ square with Wang tiles, the Glucose SAT solver [AS18] based on MiniSAT [SE05] is faster [Lab18] than Knuth's dancing links algorithm or MILP solvers.

Exercise 4.6

Using the criterion given in Lemma 3.9, prove that $\Omega_{\mathcal{Z}}$ is minimal and $\Omega_{\mathcal{Z}} = \mathcal{X}_{\Phi}$.

Exercise 4.7

Prove the self-similarity of $\Omega_{\mathcal{Z}}$ by using first markers in the direction e_1 in the set \mathcal{Z} and then using markers in the direction e_2 .

5. A self-similar symbolic dynamical system

In this section, we consider the dynamical system $(\mathbb{T}^2, \mathbb{Z}^2, R_z)$ defined on the 2-dimensional torus $\mathbb{T}^2 = (\mathbb{R}/\mathbb{Z})^2$ by the continuous \mathbb{Z}^2 -action

$$egin{array}{rcl} R_{\mathcal{Z}}: & \mathbb{Z}^2 imes \mathbb{T}^2 & o & \mathbb{T}^2 \ & (oldsymbol{n}, oldsymbol{x}) & \mapsto & oldsymbol{x} + arphi^{-2} oldsymbol{n} \end{array}$$

where $\varphi = \frac{1+\sqrt{5}}{2}$. We define a symbolic representation of that dynamical system using a wellchosen partition $\mathcal{P}_{\mathcal{Z}}$ of \mathbb{T}^2 . The partition $\mathcal{P}_{\mathcal{Z}}$ is is a simplification of the partition $\mathcal{P}_{\mathcal{U}}$ that was introduced in [Lab21a] where it was proved to be a Markov partition for the dynamical system ($\mathbb{T}^2, \mathbb{Z}^2, R_{\mathcal{Z}}$). As discovered during the PhD thesis of Jana Lepšová, it turns out that the vertical line at $x = \varphi^{-2}$ is not necessary in the partition $\mathcal{P}_{\mathcal{U}}$. Removing the vertical line at $x = \varphi^{-2}$ in the partition $\mathcal{P}_{\mathcal{U}}$ reduces the number of atoms in the partition from 19 to 16. The indices used to define the partitions are consistent with the choices made in [Lab21a] for \mathcal{U} and [Lep24] for \mathcal{Z} . As illustrated in Figure 7, the partition $\mathcal{P}_{\mathcal{Z}}$ can be defined from the following 7 segments in \mathbb{R}^2 :

$$(1,\varphi^2) \to (0,\varphi^2) \to (\varphi,0) \to (\varphi,1)$$
$$(1,1) \to (0,1) \to (1,0) \to (1,1),$$
$$\left(\frac{1}{\varphi^2},2\right) \to \left(1+\frac{1}{\varphi^2},1\right).$$



FIGURE 7. The partition $\mathcal{P}_{\mathcal{Z}}$ of \mathbb{T}^2 can be constructed from 7 segments in \mathbb{R}^2 (left) and their images under the group of translations \mathbb{Z}^2 (right).

The translations of the 7 segments under the group of translation \mathbb{Z}^2 splits the torus \mathbb{T}^2 into 16 polygonal regions indexed with integers from the set [0, 15]. The coding by the partition $\mathcal{P}_{\mathcal{Z}}$ of the orbit of a starting point in \mathbb{T}^2 by the \mathbb{Z}^2 -action of $R_{\mathcal{Z}}$ defines a configuration $w \in [0, 15]^{\mathbb{Z}^2}$, see Figure 8. The topological closure of the set of all such configurations is the symbolic dynamical system $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}}$ corresponding to $\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}$ (see Lemma 5.7). It turns out that $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}}$ is a **subshift** as it is also closed under the shift σ by integer translations.

The goal of the next sections is to prove that the symbolic dynamical system $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$ is self-similar where the self-similarity is given by the 2-dimensional morphism Φ defined in Equation (1) (Theorem 1.2).



FIGURE 8. The polygonal partition $\mathcal{P}_{\mathcal{Z}}$ of \mathbb{T}^2 with indices in the set $[\![0, 15]\!]$. The coding of the shifted lattice $\mathbf{p} + \frac{1}{\varphi^2} \mathbb{Z}^2$ by the partition defines a configuration in $[\![0, 15]\!]^{\mathbb{Z}^2}$. In the figure, the points $\mathbf{p} + \frac{1}{\varphi^2}(m, n)$ are shown in white with $\mathbf{p} = (0.1357, 0.2938)$ for each $m \in [\![0, 5]\!]$ and $n \in [\![0, 7]\!]$ and are coded by a 2-dimensional word of shape (6, 8).

Exercise 5.1

Define the polygonal partition $\mathcal{P}_{\mathcal{Z}}$ of \mathbb{T}^2 using the simplification proposed by Jana Lepšová [Lep24] of the partition $\mathcal{P}_{\mathcal{U}}$ that was introduced in [Lab21a], that is, merging the atoms labeled 6 and 7, merging the atoms labeled 12 and 13 and merging the atoms labeled 14 and 15. See Figure 9.



FIGURE 9. The partition $\mathcal{P}_{\mathcal{U}}$ introduced in [Lab21a] and the partition $\mathcal{P}_{\mathcal{Z}}$ following the simplification suggested in [Lep24].

Exercise 5.2

Figure 8 provides the construction of a 2-dimensional word of shape (6, 8). Using the same construction, extend that pattern by one unit in all directions to obtain a word of shape (8, 10).

5.1. Toral \mathbb{Z}^2 -rotations and polygon exchange transformations (PETs). Let Γ be a lattice in \mathbb{R}^2 , i.e., a discrete subgroup of the additive group \mathbb{R}^2 with 2 linearly independent generators. This defines a 2-dimensional torus $T = \mathbb{R}^2/\Gamma$. By analogy with the rotation $x \mapsto x + \alpha$ on the circle \mathbb{R}/\mathbb{Z} for an $\alpha \in \mathbb{R}$, we use the terminology of rotation (sometimes also called translation) to denote the following \mathbb{Z}^2 -action defined on a 2-dimensional torus.

Definition 5.1. Let $T = \mathbb{R}^2/\Gamma$ where Γ is a lattice in \mathbb{R}^2 . For some $\alpha, \beta \in T$, we consider the dynamical system (T, \mathbb{Z}^2, R) where $R : \mathbb{Z}^2 \times T \to T$ is the continuous \mathbb{Z}^2 -action on Tdefined by

 $R^{n}(\boldsymbol{x}) := R(\boldsymbol{n}, \boldsymbol{x}) = \boldsymbol{x} + n_{1}\boldsymbol{\alpha} + n_{2}\boldsymbol{\beta}$

for every $\mathbf{n} = (n_1, n_2) \in \mathbb{Z}^2$. We say that the \mathbb{Z}^2 -action R is a **toral** \mathbb{Z}^2 -rotation or a \mathbb{Z}^2 -rotation on \mathbf{T} which defines a dynamical system $(\mathbf{T}, \mathbb{Z}^2, R)$.

It is practical to represent a toral \mathbb{Z}^2 -rotation in terms of polygon exchange transformations [Hoo13, Sch14].

Definition 5.2. [AKY19] Let X be a polygon together with two topological partitions of X into polygons

$$X = \bigcup_{k=0}^{N} P_k = \bigcup_{k=0}^{N} Q_k$$

such that for each k, P_k and Q_k are translation equivalent, i.e., there exists $\mathbf{v}_k \in \mathbb{R}^2$ such that $P_k = Q_k + \mathbf{v}_k$. A **polygon exchange transformation** (PET) is the piecewise translation on X defined for $\mathbf{x} \in P_k$ by $T(\mathbf{x}) = \mathbf{x} + \mathbf{v}_k$. The map is not defined for points $\mathbf{x} \in \bigcup_{k=0}^N \partial P_k$.



FIGURE 10. A polygon exchange transformation defined on the triangle with vertices (0,0), (0,1) and (1,0).

A PETs can be quite complicated. For example, a polygon exchange transformation is shown in Figure 10. In this chapter, we consider pairs of commuting PETs that are much simpler given by the exchanges of two rectangles, see Figure 11.



FIGURE 11. The \mathbb{Z}^2 -action $R_{\mathcal{Z}}$ can be seen as a pair of commuting polygon exchange transformations on the unit square $[0, 1)^2$.

A \mathbb{Z}^2 -rotation R on a torus T can be decomposed into two commuting maps as follows: $R^n = R^{n_1 e_1 + n_2 e_2} = R^{n_1 e_1} \circ R^{n_2 e_2} = (R^{e_1})^{n_1} \circ (R^{e_2})^{n_2}.$

where each map R^{e_1} and R^{e_2} can be seen as polygon exchange transformation defined by the exchange of at most 4 pieces on a fundamental domain of T having for shape a parallelogram. We state this as a lemma because we use this connection two times in the proof of Theorem 1.2.

Lemma 5.3. Let $\Gamma = \ell_1 \mathbb{Z} \times \ell_2 \mathbb{Z}$ be a lattice in \mathbb{R}^2 and its rectangular fundamental domain $D = [0, \ell_1) \times [0, \ell_2)$. For every $\boldsymbol{\alpha} = (\alpha_1, \alpha_2) \in D$, the dynamical system $(\mathbb{R}^2/\Gamma, \mathbb{Z}, \boldsymbol{x} \mapsto \boldsymbol{x} + \boldsymbol{\alpha})$

is measurably conjugate to the dynamical system (D, \mathbb{Z}, T) where $T : D \to D$ is the polygon exchange transformation shown in Figure 12.



FIGURE 12. The polygon exchange transformation T of the rectangle $[0, \ell_1) \times [0, \ell_2)$ as defined on the figure can be seen as a toral rotation by the vector (α_1, α_2) on the torus $\mathbb{R}^2/(\ell_1\mathbb{Z} \times \ell_2\mathbb{Z})$.

Proof. It follows from the fact that toral rotations and such polygon exchange transformations are the Cartesian product of circle rotations and exchange of two intervals. The fact that a rotation on a circle can be seen as a exchange of two intervals is well-known as noticed for example in [Rau77]. \Box

Exercise 5.3

Recall that $R_{\mathcal{Z}}^{n}(\boldsymbol{x}) = \boldsymbol{x} + \varphi^{-2}\boldsymbol{n}$ is a \mathbb{Z}^{2} -action defined on \mathbb{T}^{2} where $\varphi = \frac{1+\sqrt{5}}{2}$. Prove that the maps $R_{\mathcal{Z}}^{e_{1}}$ and $R_{\mathcal{Z}}^{e_{2}}$ can be expressed as polygon exchange transformations on the unit square $[0, 1) \times [0, 1)$ as in Figure 11.

5.2. Symbolic dynamical systems for toral \mathbb{Z}^2 -rotations. Let Γ be a lattice in \mathbb{R}^2 and $T = \mathbb{R}^2/\Gamma$ be a 2-dimensional torus. Let (T, \mathbb{Z}^2, R) be the dynamical system given by a \mathbb{Z}^2 -rotation R on T. For some finite set \mathcal{A} , a topological partition of T is a finite collection $\{P_a\}_{a \in \mathcal{A}}$ of disjoint open sets $P_a \subset T$ such that $T = \bigcup_{a \in \mathcal{A}} \overline{P_a}$. If $S \subset \mathbb{Z}^2$ is a finite set, we say that a pattern $w \in \mathcal{A}^S$ is allowed for \mathcal{P}, R if

(9)
$$\bigcap_{k \in S} R^{-k}(P_{w_k}) \neq \emptyset$$

Let $\mathcal{L}_{\mathcal{P},R}$ be the collection of all allowed patterns for \mathcal{P}, R . The set $\mathcal{L}_{\mathcal{P},R}$ is the language of a subshift $\mathcal{X}_{\mathcal{P},R} \subseteq \mathcal{A}^{\mathbb{Z}^2}$ defined as follows, see [Hoc16, Prop. 9.2.4],

$$\mathcal{X}_{\mathcal{P},R} = \{ x \in \mathcal{A}^{\mathbb{Z}^2} \mid \pi_S \circ \sigma^n(x) \in \mathcal{L}_{\mathcal{P},R} \text{ for every } \boldsymbol{n} \in \mathbb{Z}^2 \text{ and finite subset } S \subset \mathbb{Z}^2 \}$$

Definition 5.4. We call $\mathcal{X}_{\mathcal{P},R}$ the symbolic dynamical system corresponding to \mathcal{P}, R .

For each $w \in \mathcal{X}_{\mathcal{P},R} \subset \mathcal{A}^{\mathbb{Z}^2}$ and $n \geq 0$ there is a corresponding nonempty open set

$$D_n(w) = \bigcap_{\|\boldsymbol{k}\| \le n} R^{-\boldsymbol{k}}(P_{w_{\boldsymbol{k}}}) \subseteq \boldsymbol{T}.$$

The closures $\overline{D}_n(w)$ of these sets are compact and decrease with n, so that $\overline{D}_0(w) \supseteq \overline{D}_1(w) \supseteq \overline{D}_2(w) \supseteq \ldots$. It follows that $\bigcap_{n=0}^{\infty} \overline{D}_n(w) \neq \emptyset$. In order for points in $\mathcal{X}_{\mathcal{P},R}$ to correspond to points in \mathbf{T} , this intersection should contain only one point. This leads to the following definition.

Definition 5.5. A topological partition \mathcal{P} of T gives a symbolic representation of (T, \mathbb{Z}^2, R) if for every $w \in \mathcal{X}_{\mathcal{P},R}$ the intersection $\bigcap_{n=0}^{\infty} \overline{D}_n(w)$ consists of exactly one point $x \in T$. We call w a symbolic representation of x.

In general, the existence of an atom of the partition of the torus T which is invariant only under the trivial translation is a sufficient condition for the partition to give a symbolic representation of a minimal \mathbb{Z}^2 -rotation on T.

Lemma 5.6. [Lab21a, Lemma 3.4] Let $(\mathbf{T}, \mathbb{Z}^2, R)$ be a minimal dynamical system and $\mathcal{P} = \{P_0, P_1, ..., P_{r-1}\}$ be a topological partition of \mathbf{T} . If there exists an atom P_i which is invariant only under the trivial translation in \mathbf{T} , then \mathcal{P} gives a symbolic representation of $(\mathbf{T}, \mathbb{Z}^2, R)$.

Proof. Let $\mathcal{A} = \{0, 1, \ldots, r-1\}$. Let $w \in \mathcal{X}_{\mathcal{P},R} \subset \mathcal{A}^{\mathbb{Z}^2}$. As already noticed, the closures $\overline{D}_n(w)$ are compact and decrease with n, so that $\overline{D}_0(w) \supseteq \overline{D}_1(w) \supseteq \overline{D}_2(w) \supseteq \ldots$. It follows that $\bigcap_{n=0}^{\infty} \overline{D}_n(w) \neq \emptyset$.

We show that $\bigcap_{n=0}^{\infty} \overline{D}_n(w)$ contains at most one element. Let $\boldsymbol{x}, \boldsymbol{y} \in \boldsymbol{T}$. We assume $\boldsymbol{x} \in \bigcap_{n=0}^{\infty} \overline{D}_n(w)$ and we want to show that $\boldsymbol{y} \notin \bigcap_{n=0}^{\infty} \overline{D}_n(w)$ if $\boldsymbol{x} \neq \boldsymbol{y}$. Let $P_i \subset \boldsymbol{T}$ for some $i \in \mathcal{A}$ be an atom which is invariant only under the trivial translation. Since $\boldsymbol{x} \neq \boldsymbol{y}, \overline{P_i} \setminus (\overline{P_i} - (\boldsymbol{y} - \boldsymbol{x}))$ contains an open set O. Since $(\boldsymbol{T}, \mathbb{Z}^2, R)$ is minimal, any orbit $\{R^k \boldsymbol{x} \mid \boldsymbol{k} \in \mathbb{Z}^2\}$ is dense in \boldsymbol{T} . Therefore, there exists $\boldsymbol{k} \in \mathbb{Z}^2$ such that $R^k \boldsymbol{x} \in O \subset \mathring{P_i}$. Also $\boldsymbol{x} \in \bigcap_{n=0}^{\infty} \overline{D}_n(w) \subset R^{-k} \overline{P_{w_k}}$ which implies $R^k \boldsymbol{x} \in \overline{P_{w_k}}$. Thus $\overline{P_{w_k}} \cap \mathring{P_i} \neq \emptyset$ which implies that $P_{w_k} = P_i$ and $w_k = i$ since \mathcal{P} is a topological partition. Thus

$$\bigcap_{n=0}^{\infty} \overline{D}_n(w) \subset R^{-k} \overline{P_{w_k}} = R^{-k} \overline{P_i}.$$

The fact that $R^{k} \boldsymbol{x} \in O$ also means that $R^{k} \boldsymbol{x} \notin \overline{P_{i}} - (\boldsymbol{y} - \boldsymbol{x})$ which can be rewritten as $R^{k} \boldsymbol{y} \notin \overline{P_{i}}$ or $\boldsymbol{y} \notin R^{-k} \overline{P_{i}}$ and we conclude that $\boldsymbol{y} \notin \bigcap_{n=0}^{\infty} \overline{D}_{n}(w)$. Thus \mathcal{P} gives a symbolic representation of $(\boldsymbol{T}, \mathbb{Z}^{2}, R)$.

The set

$$\Delta_{\mathcal{P},R} := \bigcup_{n \in \mathbb{Z}^2} R^n \left(\bigcup_{a \in \mathcal{A}} \partial P_a \right) \subset \boldsymbol{T}$$

is the set of points whose orbit under the \mathbb{Z}^2 -action of R intersect the boundary of the topological partition $\mathcal{P} = \{P_a\}_{a \in \mathcal{A}}$. From Baire Category Theorem [LM95, Theorem 6.1.24], the set $\mathbf{T} \setminus \Delta_{\mathcal{P},R}$ is dense in \mathbf{T} .

A topological partition $\mathcal{P} = \{P_a\}_{a \in \mathcal{A}}$ of $\mathbf{T} = \mathbb{R}^2/\Gamma$ is associated to a coding map

$$\begin{array}{rcl} \text{CODE}: & \boldsymbol{T} \setminus (\bigcup_{a \in \mathcal{A}} \partial P_a) & \to & \mathcal{A} \\ & \boldsymbol{x} & \mapsto & a & \text{if and only if} & \boldsymbol{x} \in P_a. \end{array}$$

For every starting point $\boldsymbol{x} \in \boldsymbol{T} \setminus \Delta_{\mathcal{P},R}$, the coding of its orbit under the \mathbb{Z}^2 -action of R is a configuration $\text{CONFIG}_{\boldsymbol{x}}^{\mathcal{P},R} \in \mathcal{A}^{\mathbb{Z}^2}$ defined by

$$\operatorname{CONFIG}_{\boldsymbol{x}}^{\mathcal{P},R}(\boldsymbol{n}) = \operatorname{CODE}(R^{\boldsymbol{n}}(\boldsymbol{x})).$$

for every $n \in \mathbb{Z} \times \mathbb{Z}$.

Lemma 5.7. The symbolic dynamical system $\mathcal{X}_{\mathcal{P},R}$ corresponding to \mathcal{P}, R is the topological closure of the set of configurations:

$$\mathcal{X}_{\mathcal{P},R} = \overline{\left\{ CONFIG_{\boldsymbol{x}}^{\mathcal{P},R} \mid \boldsymbol{x} \in \boldsymbol{T} \setminus \Delta_{\mathcal{P},R} \right\}}.$$

Proof. (\supseteq) Let $\boldsymbol{x} \in \boldsymbol{T} \setminus \Delta_{\mathcal{P},R}$. The patterns appearing in the configuration $\text{ConFIG}_{\boldsymbol{x}}^{\mathcal{P},R}$ are in $\mathcal{L}_{\mathcal{P},R}$. Thus $\text{ConFIG}_{\boldsymbol{x}}^{\mathcal{P},R} \in \mathcal{X}_{\mathcal{P},R}$. The topological closure of such configurations is in $\mathcal{X}_{\mathcal{P},R}$ since $\mathcal{X}_{\mathcal{P},R}$ is topologically closed.

 (\subseteq) Let $w \in \mathcal{A}^S$ be a pattern with finite support $S \subset \mathbb{Z}^2$ appearing in $\mathcal{X}_{\mathcal{P},R}$. Then $w \in \mathcal{L}_{\mathcal{P},R}$ and from Equation (9) there exists $\boldsymbol{x} \in \boldsymbol{T} \setminus \Delta_{\mathcal{P},R}$ such that $\boldsymbol{x} \in \bigcap_{k \in S} R^{-k}(P_{w_k})$. The pattern w appears in the configuration $\operatorname{CONFIG}_{\boldsymbol{x}}^{\mathcal{P},R}$. Any configuration in $\mathcal{X}_{\mathcal{P},R}$ is the limit of a sequence $(w_n)_{n \in \mathbb{N}}$ of patterns covering a ball of radius n around the origin, thus equal to some limit $\lim_{n \to \infty} \operatorname{CONFIG}_{\boldsymbol{x}_n}^{\mathcal{P},R}$ with $\boldsymbol{x}_n \in \boldsymbol{T} \setminus \Delta_{\mathcal{P},R}$ for every $n \in \mathbb{N}$.

Exercise 5.4

Prove that $\mathcal{P}_{\mathcal{Z}}$ gives a symbolic representation of the dynamical system $(\mathbb{T}^2, \mathbb{Z}^2, R_{\mathcal{Z}})$.

5.3. Factor map. An important consequence of the fact that a partition \mathcal{P} gives a symbolic representation of the dynamical system $(\mathbf{T}, \mathbb{Z}^2, R)$ is the existence of a factor map $f : \mathcal{X}_{\mathcal{P},R} \to \mathbf{T}$ which commutes the \mathbb{Z}^2 -actions. In the spirit of [LM95, Prop. 6.5.8] for \mathbb{Z} -actions, we have the following proposition whose proof can be found in [Lab21a].

Proposition 5.8. [Lab21a, Prop. 5.1] Let \mathcal{P} give a symbolic representation of the dynamical system $(\mathbf{T}, \mathbb{Z}^2, R)$. Let $f : \mathcal{X}_{\mathcal{P},R} \to \mathbf{T}$ be defined such that f(w) is the unique point in the intersection $\bigcap_{n=0}^{\infty} \overline{D}_n(w)$. The map f is a factor map from $(\mathcal{X}_{\mathcal{P},R}, \mathbb{Z}^2, \sigma)$ to $(\mathbf{T}, \mathbb{Z}^2, R)$ such that $R^{\mathbf{k}} \circ f = f \circ \sigma^{\mathbf{k}}$ for every $\mathbf{k} \in \mathbb{Z}^2$. The map f is one-to-one on $f^{-1}(\mathbf{T} \setminus \Delta_{\mathcal{P},R})$.

Using the factor map, one can prove the following lemma.

Lemma 5.9. [Lab21a, Lemma 5.2] Let \mathcal{P} give a symbolic representation of the dynamical system $(\mathbf{T}, \mathbb{Z}^2, R)$. Then

- (i) if $(\mathbf{T}, \mathbb{Z}^2, R)$ is minimal, then $(\mathcal{X}_{\mathcal{P},R}, \mathbb{Z}^2, \sigma)$ is minimal,
- (ii) if R is a free \mathbb{Z}^2 -action on T, then $\mathcal{X}_{\mathcal{P},R}$ aperiodic.

Of course, Lemma 5.9 does not hold if \mathcal{P} does not give a symbolic representation of $(\mathbf{T}, \mathbb{Z}^2, R)$.

Exercise 5.5

Prove that $(\mathcal{X}_{\mathcal{P}_{\mathbb{Z}},R_{\mathbb{Z}}},\mathbb{Z}^2,\sigma)$ is minimal and aperiodic.

5.4. Induced \mathbb{Z}^2 -actions. Renormalization schemes also known as *Rauzy induction* were originally defined for dynamical systems including interval exchange transformations (IET) [Rau79]. A natural way to generalize it to higher dimension is to consider polygon exchange transformations [Hoo13, AKY19] or even polytope exchange transformations [Sch14, Sch11] where only one map is considered. But more dimensions also allow to induce two or more (commuting) maps at the same time.

In this section, we define the induction of \mathbb{Z}^2 -actions on a sub-domain. We consider the torus $\mathbf{T} = \mathbb{R}^2/\Gamma$ where Γ is a lattice in \mathbb{R}^2 . Let $(\mathbf{T}, \mathbb{Z}^2, R)$ be a minimal dynamical system given by a \mathbb{Z}^2 -action R on \mathbf{T} . For every $\mathbf{n} \in \mathbb{Z}^2$, the toral translation R^n can be seen as a pair of polygon exchange transformations on a fundamental domain of \mathbf{T} .

Let $W \subset T$ be a set. The set of return times of $x \in T$ to the window W under the \mathbb{Z}^2 -action R is the subset of $\mathbb{Z} \times \mathbb{Z}$ defined as:

$$\delta_W(\boldsymbol{x}) = \{ \boldsymbol{n} \in \mathbb{Z} \times \mathbb{Z} \mid R^{\boldsymbol{n}}(\boldsymbol{x}) \in W \}.$$

Definition 5.10. Let $W \subset \mathbf{T}$. We say that the \mathbb{Z}^2 -action R is **Cartesian on** W if the set of return times $\delta_W(\mathbf{x})$ can be expressed as a Cartesian product, that is, for all $\mathbf{x} \in \mathbf{T}$ there exist two strictly increasing sequences $r_x^{(1)}, r_x^{(2)} : \mathbb{Z} \to \mathbb{Z}$ such that

$$\delta_W(\boldsymbol{x}) = r_{\boldsymbol{x}}^{(1)}(\mathbb{Z}) \times r_{\boldsymbol{x}}^{(2)}(\mathbb{Z}).$$

We always assume that the sequences are shifted in such a way that

$$r_{\boldsymbol{x}}^{(i)}(n) \ge 0 \iff n \ge 0 \qquad \text{for } i \in \{1, 2\}.$$

In particular, if $\boldsymbol{x} \in W$ then $(0,0) \in \delta_W(\boldsymbol{x})$, and therefore $r_{\boldsymbol{x}}^{(1)}(0) = r_{\boldsymbol{x}}^{(2)}(0) = 0$.

When the \mathbb{Z}^2 -action R is Cartesian on $W \subset \mathbf{T}$, we say that the tuple

(10)
$$(r_{\boldsymbol{x}}^{(1)}(1), r_{\boldsymbol{x}}^{(2)}(1))$$

is the **first return time** of a starting point $x \in T$ to $W \subset T$ under the action R. When the \mathbb{Z}^2 -action R is Cartesian on $W \subset T$, we may consider its return map on W and we prove in the next lemma that this induces a \mathbb{Z}^2 -action on W.

Lemma 5.11. If the \mathbb{Z}^2 -action R is Cartesian on $W \subset \mathbf{T}$, then the map $\widehat{R}|_W : \mathbb{Z}^2 \times W \to W$ defined by

$$(\hat{R}|_{W})^{n}(\boldsymbol{x}) := \hat{R}|_{W}(\boldsymbol{n}, \boldsymbol{x}) = R^{(r_{\boldsymbol{x}}^{(1)}(n_{1}), r_{\boldsymbol{x}}^{(2)}(n_{2}))}(\boldsymbol{x})$$

for every $\boldsymbol{n} = (n_1, n_2) \in \mathbb{Z}^2$ and $\boldsymbol{x} \in W$ is a well-defined \mathbb{Z}^2 -action on W.

We say that $\widehat{R}|_W$ is the **induced** \mathbb{Z}^2 -action of the \mathbb{Z}^2 -action R on W.

Proof. Let $\boldsymbol{x} \in W$. We have that

$$\widehat{R}|_{W}(\mathbf{0}, \boldsymbol{x}) = R^{(r_{\boldsymbol{x}}^{(1)}(0), r_{\boldsymbol{x}}^{(2)}(0))}(\boldsymbol{x}) = R^{(0,0)}(\boldsymbol{x}) = \boldsymbol{x}.$$

Firstly, using

$$r_{\boldsymbol{x}}^{(i)}(k+n) = r_{R^{\boldsymbol{e}_{i} \cdot r_{\boldsymbol{x}}^{(i)}(n)}(\boldsymbol{x})}(k) + r_{\boldsymbol{x}}^{(i)}(n),$$

and skipping few details, we get

$$\widehat{R}|_{W}(k\boldsymbol{e}_{i}+n\boldsymbol{e}_{i},\boldsymbol{x})=\widehat{R}|_{W}\left(k\boldsymbol{e}_{i},\left(\widehat{R}|_{W}(n\boldsymbol{e}_{i},\boldsymbol{x})\right)\right)$$

Secondly, using the fact that

$$r_{\boldsymbol{x}}^{(1)}(k_1) = r_{\boldsymbol{y}}^{(1)}(k_1)$$

whenever $y = R^{(0,r_{\boldsymbol{x}}^{(2)}(k_2))} \boldsymbol{x} = \widehat{R}|_W(k_2\boldsymbol{e}_2,\boldsymbol{x})$, we obtain

$$\widehat{R}|_{W}(\boldsymbol{k},\boldsymbol{x}) = R^{(r_{\boldsymbol{x}}^{(1)}(k_{1}),r_{\boldsymbol{x}}^{(2)}(k_{2}))}(\boldsymbol{x}) = R^{(r_{\boldsymbol{x}}^{(1)}(k_{1}),0)}R^{(0,r_{\boldsymbol{x}}^{(2)}(k_{2}))}(\boldsymbol{x})$$
$$= R^{(r_{\boldsymbol{x}}^{(1)}(k_{1}),0)}\widehat{R}|_{W}(k_{2}\boldsymbol{e}_{2},\boldsymbol{x}) = \widehat{R}|_{W}\left(k_{1}\boldsymbol{e}_{1},\widehat{R}|_{W}(k_{2}\boldsymbol{e}_{2},\boldsymbol{x})\right).$$

Therefore, for every $\boldsymbol{k}, \boldsymbol{n} \in \mathbb{Z}^2$, we have

$$(\widehat{R}|_W)^{k+n}(\boldsymbol{x}) = (\widehat{R}|_W)^{(k_1+n_1)e_1} (\widehat{R}|_W)^{(k_2+n_2)e_2}(\boldsymbol{x})$$

$$= (\widehat{R}|_W)^{k_1e_1} (\widehat{R}|_W)^{n_1e_1} (\widehat{R}|_W)^{k_2e_2} (\widehat{R}|_W)^{n_2e_2}(\boldsymbol{x})$$

$$= (\widehat{R}|_W)^{k_1e_1} (\widehat{R}|_W)^{(n_1,k_2)} (\widehat{R}|_W)^{n_2e_2}(\boldsymbol{x})$$

$$= (\widehat{R}|_W)^{k_1e_1} (\widehat{R}|_W)^{k_2e_2} (\widehat{R}|_W)^{n_1e_1} (\widehat{R}|_W)^{n_2e_2}(\boldsymbol{x})$$

$$= (\widehat{R}|_W)^k (\widehat{R}|_W)^n(\boldsymbol{x}),$$

which shows that $\widehat{R}|_W$ is a \mathbb{Z}^2 -action on W.

A consequence of the lemma is that the induced \mathbb{Z}^2 -action $\widehat{R}|_W$ is generated by two commutative maps

$$(\widehat{R}|_W)^{\boldsymbol{e}_1}(\boldsymbol{x}) = R^{(r_{\boldsymbol{x}}^{(1)}(1),0)}(\boldsymbol{x}) \text{ and } (\widehat{R}|_W)^{\boldsymbol{e}_2}(\boldsymbol{x}) = R^{(0,r_{\boldsymbol{x}}^{(2)}(1))}(\boldsymbol{x})$$

which are the first return maps of R^{e_1} and R^{e_2} to W:

$$(\widehat{R}|_W)^{\boldsymbol{e}_1}(\boldsymbol{x}) = \widehat{R^{\boldsymbol{e}_1}}|_W(\boldsymbol{x}) \quad \text{and} \quad (\widehat{R}|_W)^{\boldsymbol{e}_2}(\boldsymbol{x}) = \widehat{R^{\boldsymbol{e}_2}}|_W(\boldsymbol{x}).$$

Recall that the first return map $\widehat{T}|_W$ of a dynamical system (X,T) maps a point $x \in W \subset$ X to the first point in the forward orbit of T lying in W, i.e.

$$\widehat{T}|_W(\boldsymbol{x}) = T^{r(\boldsymbol{x})}(\boldsymbol{x}) \quad \text{where } r(\boldsymbol{x}) = \min\{k \in \mathbb{Z}_{>0} : T^k(\boldsymbol{x}) \in W\}.$$

From Poincaré's recurrence theorem, if μ is a finite T-invariant measure on X, then the first return map $\widehat{T}|_W$ is well defined for μ -almost all $\boldsymbol{x} \in W$. When T is a rotation on a torus, if the subset W is open, then the first return map is well-defined for every point $x \in W$. Moreover if W is a polygon, then the first return map $\hat{T}|_W$ is a polygon exchange transformation. An algorithm to compute the induced transformation $\widehat{T}|_W = \widehat{R^{e_i}}|_W$ of the sub-action R^{e_i} is provided in [Lab21b].

Exercise 5.6

Let $T(\boldsymbol{x}) = \boldsymbol{x} + (\frac{1}{3}, \frac{1}{2})$ be a \mathbb{Z} -action defined on \mathbb{T}^2 . Let $W \subset \mathbb{T}^2$ be the triangle with vertices (0,0), (1,0) and (0,1). Prove that $\widehat{T}|_W$ is the PET defined in Figure 10.

Exercise 5.7

Recall that $R_{\mathcal{Z}}(\boldsymbol{n}, \boldsymbol{x}) = \boldsymbol{x} + \varphi^{-2}\boldsymbol{n}$ is a \mathbb{Z}^2 -action defined on \mathbb{T}^2 where $\varphi = \frac{1+\sqrt{5}}{2}$. Let $W_0 = (0, 1) \times (0, \varphi^{-1}) + \mathbb{Z}^2$ be a subset of \mathbb{T}^2 .

- Prove that the action $R_{\mathcal{Z}}$ is Cartesian on W_0 .
- Prove that $\widehat{R_{\mathcal{Z}}}|_{W_0}: \mathbb{Z}^2 \times W_0 \to W_0$ is a well-defined induced \mathbb{Z}^2 -action.
- Describe R^{e₁}/_Z|_{W₀} and R^{e₂}/_Z|_{W₀} as polygon exchange transformations on W₀.
 Describe R_Z|_{W₀} as a toral Z²-rotation on ℝ²/Γ₁ with Γ₁ = Z × (φ⁻¹Z).

5.5. Toral partitions induced by toral \mathbb{Z}^2 -rotations. For IETs, the interval on which we define the Rauzy induction is usually given by one of the atom of the partition which defines the IET itself. In our setting, it is not the case. The partition that we use carries more information than the natural partition which allows to define the \mathbb{Z}^2 -rotation R as a pair of polygon exchange transformations. The partition is a refinement of the natural partition involving well-chosen sloped lines.

Let Γ be a lattice in \mathbb{R}^2 and $T = \mathbb{R}^2/\Gamma$ be a 2-dimensional torus. Let (T, \mathbb{Z}^2, R) be the dynamical system given by a \mathbb{Z}^2 -rotation R on T. Assuming the \mathbb{Z}^2 -rotation R is Cartesian on a window $W \subset \mathbf{T}$, then there exist two strictly increasing sequences $r_{\mathbf{x}}^{(1)}, r_{\mathbf{x}}^{(2)}: \mathbb{Z} \to \mathbb{Z}$ such that

$$(r,s) = \left(r_{\boldsymbol{x}}^{(1)}(1), r_{\boldsymbol{x}}^{(2)}(1)\right)$$

is the first return time of a starting point $x \in T$ to the window W under the action R, see Equation (10). It allows to define the **return word map** as

$$\begin{array}{ccccc} \operatorname{Return} \operatorname{Word} : & W & \to & \mathcal{A}^{*^2} \\ & & & & \\ & & & \mathbf{x} & \mapsto & \begin{pmatrix} \operatorname{Code}(R^{(0,s-1)}\boldsymbol{x}) & \cdots & \operatorname{Code}(R^{(r-1,s-1)}\boldsymbol{x}) \\ & & & \cdots & & \cdots \\ & & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & &$$

where $r, s \ge 1$ both obviously depend on \boldsymbol{x} .

The image $\mathcal{L} = \operatorname{RETURNWORD}(W) \subset \mathcal{A}^{*^2}$ is a language called the **set of return words**. We identify each return word in \mathcal{L} to a letter *b* of an alphabet \mathcal{B} in such a way that $\mathcal{L} = \{w_b\}_{b \in \mathcal{B}}$. When the return time to *W* is bounded, the set of return words \mathcal{L} is finite and the alphabet \mathcal{B} is finite. The example that we consider in this chapter are such that the return time is bounded, but this is not true in general.

Remark 5.12. The way the enumeration of \mathcal{L} is done influences the substitutions which are obtained afterward. To obtain a canonical ordering when the words are 1-dimensional, we use the total order (\mathcal{L}, \prec) defined by $u \prec v$ if |u| < |v| or |u| = |v| and $u <_{lex} v$.

The **induced partition** of \mathcal{P} by the action of R on the sub-domain W is a topological partition of W defined as the set of preimage sets under RETURNWORD:

$$\widehat{\mathcal{P}}|_W = \{ \text{RETURNWORD}^{-1}(w_b) \}_{b \in \mathcal{B}}.$$

This yields the **induced coding** on W

 $\operatorname{CODE}|_W: W \to \mathcal{B}$ $\boldsymbol{y} \mapsto \boldsymbol{b}$ if and only if $\boldsymbol{y} \in \operatorname{RETURNWORD}^{-1}(w_b).$

A natural substitution comes out of this induction procedure:

(11)
$$\begin{aligned} \omega : \ \mathcal{B} &\to \ \mathcal{A}^{*^2} \\ b &\mapsto \ w_b. \end{aligned}$$

The partition $\widehat{\mathcal{P}}|_W$ of W can be effectively computed by the refinement of the partition \mathcal{P} with translated copies of the sub-domain W under the action of R. In [Lab21b], we propose an algorithm to compute the induced partition $\widehat{\mathcal{P}}|_W$ and substitution ω . An implementation of it in SageMath is provided in the optional package slabbe [Lab23b] and is used below on an example. The next result shows that the coding of the orbit under the \mathbb{Z}^2 -rotation R is the image under the 2-dimensional substitution ω of the coding of the orbit under the \mathbb{Z}^2 -action $\widehat{R}|_W$.

Lemma 5.13. If the \mathbb{Z}^2 -action R is Cartesian on a window $W \subset \mathbf{T}$, then ω is a 2dimensional morphism, and for every $\mathbf{x} \in W$ we have

$$CONFIG_{\boldsymbol{x}}^{\mathcal{P},R} = \omega \left(CONFIG_{\boldsymbol{x}}^{\widehat{\mathcal{P}}|_{W},\widehat{R}|_{W}} \right).$$

Proof. Let $x \in W$. By hypothesis, there exist $P, Q \subset \mathbb{Z}$ such that the set of returns times satisfies $\delta_W(x) = P \times Q$ and we may write $P = \{r_i\}_{i \in \mathbb{Z}}$ and $Q = \{s_j\}_{j \in \mathbb{Z}}$ as increasing sequences such that $r_0 = s_0 = 0$. Therefore, $\text{CONFIG}_x^{\mathcal{P},R}$ may be decomposed into a lattice

of rectangular blocks. More precisely, for every $i, j \in \mathbb{Z}$, the following block is the image of a letter under ω :

$$\begin{pmatrix} \operatorname{CODE}(R^{(r_i,s_{j+1}-1)}\boldsymbol{x}) & \cdots & \operatorname{CODE}(R^{(r_{i+1}-1,s_{j+1}-1)}\boldsymbol{x}) \\ \cdots & \cdots & \cdots \\ \operatorname{CODE}(R^{(r_i,s_j)}\boldsymbol{x}) & \cdots & \operatorname{CODE}(R^{(r_{i+1}-1,s_j)}\boldsymbol{x}) \end{pmatrix} \\ = \operatorname{RETURNWORD}(R^{(r_i,s_j)}\boldsymbol{x}) = w_{b_{ij}} = \omega(b_{ij}) \end{pmatrix}$$

for some letter $b_{ij} \in \mathcal{B}$. Moreover,

$$b_{ij} = \operatorname{CODE}|_W(R^{(r_i, s_j)}\boldsymbol{x}) = \operatorname{CODE}|_W\left((\widehat{R}|_W)^{(i, j)}\boldsymbol{x}\right)$$

Since the adjacent blocks have matching dimensions, for every $i, j \in \mathbb{Z}$, the following concatenations

$$\omega \left(b_{ij} \odot^1 b_{(i+1)j} \right) = \omega \left(b_{ij} \right) \odot^1 \omega \left(b_{(i+1)j} \right) \quad \text{and} \\ \omega \left(b_{ij} \odot^2 b_{i(j+1)} \right) = \omega \left(b_{ij} \right) \odot^2 \omega \left(b_{i(j+1)} \right)$$

are well defined. Thus ω is a 2-dimensional morphism on the set $\left\{ \operatorname{CONFIG}_{\boldsymbol{x}}^{\widehat{\mathcal{P}}|_{W},\widehat{R}|_{W}} \middle| \boldsymbol{x} \in W \right\}$ and we have

$$\operatorname{Config}_{\boldsymbol{x}}^{\mathcal{P},R} = \omega \left(\operatorname{Config}_{\boldsymbol{x}}^{\widehat{\mathcal{P}}|_{W},\widehat{R}|_{W}} \right)$$

which ends the proof. Note that the domain of ω can be extended to its topological closure.

Proposition 5.14. Let \mathcal{P} be a topological partition of \mathbf{T} . If the \mathbb{Z}^2 -action R is Cartesian on a window $W \subset \mathbf{T}$, then $\mathcal{X}_{\mathcal{P},R} = \overline{\omega(\mathcal{X}_{\widehat{\mathcal{P}}|_W,\widehat{R}|_W})}^{\sigma}$.

Proof. Let

$$Y = \left\{ \operatorname{CONFIG}_{\boldsymbol{x}}^{\mathcal{P},R} \, \middle| \, \boldsymbol{x} \in \boldsymbol{T} \right\} \quad \text{and} \quad Z = \left\{ \operatorname{CONFIG}_{\boldsymbol{x}}^{\widehat{\mathcal{P}}|_{W},\widehat{R}|_{W}} \, \middle| \, \boldsymbol{x} \in W \right\},$$

(\supseteq). Let $\boldsymbol{x} \in W$. From Lemma 5.13, $\omega \left(\operatorname{CONFIG}_{\boldsymbol{x}}^{\widehat{\mathcal{P}}|_W, \widehat{R}|_W} \right) = \operatorname{CONFIG}_{\boldsymbol{x}}^{\mathcal{P}, R}$ with $\boldsymbol{x} \in W \subset \boldsymbol{T}$. (\subseteq). Let $\boldsymbol{x} \in \boldsymbol{T}$. There exist $k_1, k_2 \in \mathbb{N}$ such that $\boldsymbol{x}' = R^{-(k_1, k_2)}(\boldsymbol{x}) \in W$. Therefore, we have $\boldsymbol{x} = R^{(k_1, k_2)}(\boldsymbol{x}')$ where $0 \leq k_1 < r(\boldsymbol{x}')$ and $0 \leq k_2 < s(\boldsymbol{x}')$. Thus the shift $\boldsymbol{k} = (k_1, k_2) \in \mathbb{Z}^2$ is bounded by the maximal return time of R^{e_1} and R^{e_2} to W. We have

$$\begin{aligned} \operatorname{CONFIG}_{\boldsymbol{x}}^{\mathcal{P},R} &= \operatorname{CONFIG}_{R^{\boldsymbol{k}}\boldsymbol{x}'}^{\mathcal{P},R} = \sigma^{\boldsymbol{k}} \circ \operatorname{CONFIG}_{\boldsymbol{x}'}^{\mathcal{P},R} \\ &= \sigma^{\boldsymbol{k}} \circ \omega \left(\operatorname{CONFIG}_{\boldsymbol{x}'}^{\widehat{\mathcal{P}}|_{W},\widehat{R}|_{W}} \right) \end{aligned}$$

where we used Lemma 5.13 with $\mathbf{x}' \in W$. We conclude that $Y = \overline{\omega(Z)}^{\sigma}$. The result follows from Lemma 5.7 by taking the topological closure on both sides.

Exercise 5.8

Let $T(\boldsymbol{x}) = \boldsymbol{x} + (\frac{1}{3}, \frac{1}{2})$ be a \mathbb{Z} -action defined on \mathbb{T}^2 as in Exercise 5.6. Let $W \subset \mathbb{T}^2$ be the triangle with vertices (0,0), (1,0) and (0,1). Let \mathcal{P} be the polygonal partition of \mathbb{T}^2 into two parts separated by the positive diagonal. Compute the substitution and the partition $\widehat{\mathcal{P}}|_W$ of \mathcal{P} induced by the action of T on the sub-domain W. 5.6. Self-similarity of the subshift $\mathcal{X}_{\mathcal{P}_{\mathbb{Z}},R_{\mathbb{Z}}}$. In this section, we induce the topological partition $\mathcal{P}_{\mathbb{Z}}$ until the process loops. We need two induction steps before obtaining a topological partition which is self-induced.

The proof contains SageMath code using the slabbe optional package [Lab23b] to reproduce the computation of the induced partitions and 2-dimensional morphisms. The algorithm of the method induced_transformation and induced_partition are available in [Lab21b].

Proof of Theorem 1.2. First, we define the golden mean **phi** as an element of a number field defined by a quadratic polynomial which is more efficient when doing arithmetic operations and comparisons. We also import the necessary functions.

```
sage: z = polygen(QQ, "z")35sage: K.<phi> = NumberField(z**2-z-1, "phi", embedding=RR(1.6))36sage: from slabbe import PolyhedronExchangeTransformation as PET37sage: from slabbe.arXiv_1903_06137 import self_similar_19_atoms_partition38
```

The proof uses Proposition 5.14 two times to induce both the vertical and horizontal actions, starting with the vertical action. We begin with the lattice $\Gamma_0 = \mathbb{Z}^2$, the partition $\mathcal{P}_{\mathcal{Z}}$, the coding map $\text{CODE}_0 : \mathbb{R}^2/\Gamma_0 \to \mathcal{A}_0$, the alphabet $\mathcal{A}_0 = \llbracket 0, 15 \rrbracket$ and \mathbb{Z}^2 -action $R_{\mathcal{Z}}$ defined on \mathbb{T}^2 as shown below.

<pre>sage: Gamma0 = matrix.column([(1,0), (0,1)])</pre>	39
<pre>sage: PU = self_similar_19_atoms_partition()</pre>	40
<pre>sage: merge_dict = {0:0, 1:1, 2:2, 3:3, 4:4, 5:5, 6:6, 7:6, 8:7, 9:8, 10:9,</pre>	41
: 11:10, 12:11, 13:11, 14:12, 15:12, 16:13, 17:14, 18:15}	42
<pre>sage: PZ = PU.merge_atoms(merge_dict)</pre>	43
<pre>sage: RZe1 = PET.toral_translation(Gamma0, vector((phi^-2,0)))</pre>	44
<pre>sage: RZe2 = PET.toral_translation(Gamma0, vector((0,phi^-2)))</pre>	45



We consider the window $W_0 = (0,1) \times (0, \varphi^{-1}) + \Gamma_0$ as a subset of \mathbb{R}^2/Γ_0 . The action $R_{\mathcal{Z}}$ is Cartesian on W_0 . Thus from Lemma 5.11, $R_1 := \widehat{R_{\mathcal{Z}}}|_{W_0} : \mathbb{Z}^2 \times W_0 \to W_0$ is a well-defined \mathbb{Z}^2 -action. From Lemma 5.3, the \mathbb{Z}^2 -action R_1 can be seen as toral rotation on \mathbb{R}^2/Γ_1 with $\Gamma_1 = \mathbb{Z} \times (\varphi^{-1}\mathbb{Z})$, see Exercise 5.7. Let $\mathcal{P}_1 = \widehat{\mathcal{P}_{\mathcal{Z}}}|_{W_0}$ be the induced partition.

From Proposition 5.14, then $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}} = \overline{\beta_0(\mathcal{X}_{\mathcal{P}_1,R_1})}^{\sigma}$. The partition \mathcal{P}_1 , the action R_1 and substitution β_0 are given below with alphabet $\mathcal{A}_1 = \llbracket 0, 17 \rrbracket$:

<pre>sage: y_ineq = [phi^-1, 0, -1] # y <= phi^-1 (see Polyhedron?)</pre>	46
<pre>sage: P1,beta0 = RZe2.induced_partition(y_ineq,PZ,</pre>	47
: substitution_type="column")	48
<pre>sage: R1e1,_ = RZe1.induced_transformation(y_ineq)</pre>	49
<pre>sage: R1e2,_ = RZe2.induced_transformation(y_ineq)</pre>	50

$$\beta_{0} : \mathcal{A}_{1} \to \mathcal{A}_{0}^{*^{2}}$$

$$\begin{cases} 0 \mapsto (7), & 1 \mapsto (8), & 2 \mapsto (10), & 3 \mapsto (11), & 4 \mapsto (12), \\ 5 \mapsto (13), & 6 \mapsto (14), & 7 \mapsto \begin{pmatrix} 0 \\ 7 \end{pmatrix}, & 8 \mapsto \begin{pmatrix} 1 \\ 8 \end{pmatrix}, & 9 \mapsto \begin{pmatrix} 1 \\ 9 \end{pmatrix}, \\ 10 \mapsto \begin{pmatrix} 1 \\ 10 \end{pmatrix}, 11 \mapsto \begin{pmatrix} 4 \\ 11 \end{pmatrix}, 12 \mapsto \begin{pmatrix} 6 \\ 11 \end{pmatrix}, 13 \mapsto \begin{pmatrix} 2 \\ 12 \end{pmatrix}, 14 \mapsto \begin{pmatrix} 6 \\ 12 \end{pmatrix}, \\ 15 \mapsto \begin{pmatrix} 3 \\ 13 \end{pmatrix}, 16 \mapsto \begin{pmatrix} 3 \\ 14 \end{pmatrix}, 17 \mapsto \begin{pmatrix} 5 \\ 15 \end{pmatrix}. \end{cases}$$

$$\mathcal{P}_{1} = \begin{array}{c} \frac{1}{\varphi^{2}} & 1 & 14 & 17 \\ \frac{1}{\varphi^{2}} & 16 & 13 & 12 & 7 \\ 0 & \frac{1}{\varphi^{2}} & \frac{1}{\varphi} & 1 & 16 \\ 0 & \frac{1}{\varphi^{2}} & \frac{1}{\varphi} & 1 & 16 \\ 0 & \frac{1}{\varphi^{2}} & \frac{1}{\varphi} & 1 \\ \end{array}$$

We consider the window $W_1 = (0, \varphi^{-1}) \times (0, \varphi^{-1}) + \Gamma_1$ as a subset of \mathbb{R}^2/Γ_1 . The action R_1 is Cartesian on W_1 . Thus from Lemma 5.11, $R_2 := \widehat{R_1}|_{W_1} : \mathbb{Z}^2 \times W_1 \to W_1$ is a well-defined \mathbb{Z}^2 -action. From Lemma 5.3, the \mathbb{Z}^2 -action R_2 can be seen as toral rotation on \mathbb{R}^2/Γ_2 with $\Gamma_2 = (\varphi^{-1}\mathbb{Z}) \times (\varphi^{-1}\mathbb{Z})$. Let $\mathcal{P}_2 = \widehat{\mathcal{P}_1}|_{W_1}$ be the induced partition. From Proposition 5.14, then $\mathcal{X}_{\mathcal{P}_1,R_1} = \overline{\beta_1(\mathcal{X}_{\mathcal{P}_2,R_2})}^{\sigma}$. The partition \mathcal{P}_2 , the action R_2 and substitution β_1 are given below with alphabet $\mathcal{A}_2 = [0, 15]$:

<pre>sage: x_ineq = [phi^-1, -1, 0] # x <= phi^-1 (see Polyhedron?)</pre>	51
<pre>sage: P2,beta1 = R1e1.induced_partition(x_ineq, P1, substitution_type="row")</pre>	52
<pre>sage: R2e1,_ = R1e1.induced_transformation(x_ineq)</pre>	53
<pre>sage: R2e2,_ = R1e2.induced_transformation(x_ineq)</pre>	54

$$\beta_{1}: \mathcal{A}_{2} \to \mathcal{A}_{1}^{*^{2}}$$

$$\begin{cases}
0 \mapsto (5), & 1 \mapsto (6), & 2 \mapsto (13), & 3 \mapsto (14), \\
4 \mapsto (15), & 5 \mapsto (16), & 6 \mapsto (3, 1), & 7 \mapsto (4, 0), \\
8 \mapsto (4, 2), & 9 \mapsto (5, 0), & 10 \mapsto (6, 0), & 11 \mapsto (11, 8), \\
12 \mapsto (12, 8), 13 \mapsto (13, 7), 14 \mapsto (14, 10), 15 \mapsto (17, 9).
\end{cases}$$



Now it is appropriate to rescale the partition \mathcal{P}_2 by the factor $-\varphi$. Doing so, the new obtained action R'_2 is the same as two steps before, that is, $R_{\mathcal{Z}}$ on $\mathbb{R}^2/\mathbb{Z}^2$. More formally, let $h : (\mathbb{R}/\varphi^{-1}\mathbb{Z})^2 \to (\mathbb{R}/\mathbb{Z})^2$ be the homeomorphism defined by $h(\boldsymbol{x}) = -\varphi \boldsymbol{x}$. We define $\mathcal{P}'_2 = h(\mathcal{P}_2), \operatorname{CODE}'_2 = \operatorname{CODE}_2 \circ h^{-1}, (R'_2)^n = h \circ (R_2)^n \circ h^{-1}$ as shown below:

55

<pre>sage: P2_scaled = (-phi*F</pre>	22).translate((1,1))	
$\mathcal{P}_{2}^{\prime} = \begin{array}{c} 1 \\ \frac{1}{\varphi} \\ \frac{1}{\varphi^{2}} \\ 1 \\ 0 \\ 0 \end{array}$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$(R_2')^{m n}(m x)=m x+arphi^{-2}m n$

We observe that the scaled partition \mathcal{P}'_2 is the same as $\mathcal{P}_{\mathcal{Z}}$ up to a permutation β_2 of the indices of the atoms in such a way that $\beta_2 \circ \text{CODE}_0 = \text{CODE}'_2$. The partition $\mathcal{P}_{\mathcal{Z}}$, the action $R_{\mathcal{Z}}$ and substitution $\beta_2 : \mathcal{A}_2 \to \mathcal{A}_0$ are given below.

sage:	<pre>assert P2_scaled.is_equal_up_to_relabeling(PZ)</pre>	56
sage:	<pre>beta2 = Substitution2d.from_permutation(PZ.keys_permutation(P2_scaled))</pre>	57

$$\begin{split} \beta_2: \quad \llbracket 0, 15 \rrbracket \to \llbracket 0, 15 \rrbracket^{*^2} \\ \begin{cases} 0 \mapsto (1) \,, \quad 1 \mapsto (0) \,, \quad 2 \mapsto (8) \,, \quad 3 \mapsto (6) \,, \\ 4 \mapsto (10) \,, \quad 5 \mapsto (9) \,, \quad 6 \mapsto (7) \,, \quad 7 \mapsto (3) \,, \\ 8 \mapsto (5) \,, \quad 9 \mapsto (4) \,, \quad 10 \mapsto (2) \,, \quad 11 \mapsto (14) \,, \\ 12 \mapsto (12) \,, 13 \mapsto (15) \,, 14 \mapsto (11) \,, 15 \mapsto (13) \,. \end{split}$$

By construction, the following diagrams commute:

$$\begin{array}{cccc} (\mathbb{R}/\varphi^{-1}\mathbb{Z})^2 & \xrightarrow{h} (\mathbb{R}/\mathbb{Z})^2 & & (\mathbb{R}/\varphi^{-1}\mathbb{Z})^2 & \xrightarrow{h} (\mathbb{R}/\mathbb{Z})^2 \\ R_2^n & & \downarrow^{(R'_2)^n} & \text{and} & & \\ (\mathbb{R}/\varphi^{-1}\mathbb{Z})^2 & \xrightarrow{h} (\mathbb{R}/\mathbb{Z})^2 & & & \mathcal{A}_2 \end{array}$$

Using the above commutative properties, for every $\boldsymbol{y} \in \mathbb{R}^2/\Gamma_2$ and $\boldsymbol{n} \in \mathbb{Z}^2$, we have

$$\operatorname{CONFIG}_{\boldsymbol{y}}^{\mathcal{P}_{2},R_{2}}(\boldsymbol{n}) = \operatorname{CODE}_{2}(R_{2}^{\boldsymbol{n}}(\boldsymbol{y})) = \operatorname{CODE}_{2}^{\prime} \circ h(R_{2}^{\boldsymbol{n}}(\boldsymbol{y}))$$
$$= \beta_{2} \circ \operatorname{CODE}_{0} \circ h(R_{2}^{\boldsymbol{n}}(\boldsymbol{y})) = \beta_{2} \circ \operatorname{CODE}_{0} \circ (R_{2}^{\prime})^{\boldsymbol{n}}(h(\boldsymbol{y})))$$
$$= \beta_{2} \circ \operatorname{CODE}_{0} \circ R_{\mathcal{Z}}^{\boldsymbol{n}}(h(\boldsymbol{y}))) = \beta_{2} \left(\operatorname{CONFIG}_{h(\boldsymbol{y})}^{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}(\boldsymbol{n})\right).$$

Thus $\mathcal{X}_{\mathcal{P}_2,R_2} = \beta_2 (\mathcal{X}_{\mathcal{P}_z,R_z})$. We may check that $\beta_0 \circ \beta_1 \circ \beta_2 = \Phi$ where the variable Phi was created in Exercise 3.3:

sage: beta0 * beta1 * beta2 == Phi
True

We conclude that

$$\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}} = \overline{\beta_0(\mathcal{X}_{\mathcal{P}_1,R_1})}^{\sigma} = \overline{\beta_0\beta_1(\mathcal{X}_{\mathcal{P}_2,R_2})}^{\sigma} = \overline{\beta_0\beta_1\beta_2\left(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}\right)}^{\sigma} = \overline{\Phi\left(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}\right)}^{\sigma}.$$

58

59

Exercise 5.9

Compute the language of the dominoes and patterns of shape 2×2 within $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$: $\mathcal{L}_{1\times 2}(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}), \quad \mathcal{L}_{2\times 1}(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}) \text{ and } \mathcal{L}_{2\times 2}(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}).$

Exercise 5.10

Using the criterion given in Lemma 3.9, prove that $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$ is minimal and $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}} = \mathcal{X}_{\Phi}$.

Exercise 5.11

Prove the self-similarity of $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$ by doing the induction first horizontaly with $R_{\mathcal{Z}}^{e_1}$, and then vertically with $R_{\mathcal{Z}}^{e_2}$. Compare with the result of Exercise 4.7. See also Exercise 3.16.

6. CONCLUSION

We may now conclude with a proof of Theorem 1.3 providing three different characterizations of the same aperiodic 2-dimensional subshift.

Proof of Theorem 1.3. In Section 3, we defined the 2-dimensional subshift \mathcal{X}_{Φ} from some 2-dimensional morphism Φ .

In Section 4 we proved that $\Omega_{\mathcal{Z}} = \overline{\Phi(\Omega_{\mathcal{Z}})}^{\sigma}$ using the desubstitution of Wang shifts using marker tiles. In Exercise 4.6, we proved using the criterion given in Lemma 3.9, that $\Omega_{\mathcal{Z}} = \mathcal{X}_{\Phi}$.

In Section 5 we proved that $\mathcal{X}_{\mathcal{P}_{Z},R_{Z}} = \overline{\Phi(\mathcal{X}_{\mathcal{P}_{Z},R_{Z}})}^{\sigma}$ using induction of \mathbb{Z}^{2} -rotations. In Exercise 5.10, we proved using the criterion given in Lemma 3.9, that $\mathcal{X}_{\mathcal{P}_{Z},R_{Z}} = \mathcal{X}_{\Phi}$. Thus, we have the equality

$$\Omega_{\mathcal{Z}} = \mathcal{X}_{\Phi} = \mathcal{X}_{\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}}.$$

Since $\Omega_{\mathcal{Z}}$ is a SFT, it implies that $\mathcal{P}_{\mathcal{Z}}$ is a Markov partition for the \mathbb{Z}^2 -action $\mathbb{R}_{\mathcal{Z}}$ on \mathbb{T}^2 . Markov partitions were originally defined for one-dimensional dynamical systems $(\boldsymbol{T}, \mathbb{Z}, R)$ and were extended to \mathbb{Z}^d -actions by automorphisms of compact Abelian group in [ES97]. Following [Lab21a], we use the same terminology and extend the definition proposed in [LM95, §6.5] for dynamical systems defined by higher-dimensional actions by rotations.

Definition 6.1. A topological partition \mathcal{P} of T is a Markov partition for (T, \mathbb{Z}^2, R) if

- \mathcal{P} gives a symbolic representation of $(\mathbf{T}, \mathbb{Z}^2, R)$ and
- $\mathcal{X}_{\mathcal{P},R}$ is a shift of finite type (SFT).

The link between the subshifts $\Omega_{\mathcal{Z}}$ and $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$ can be explained directly without the 2dimensional morphism Φ by the existence of a factor map from $(\Omega_{\mathcal{Z}},\mathbb{Z}^2,\sigma)$ to $(\mathbb{T}^2,\mathbb{Z}^2,R_{\mathcal{Z}})$. It turns out that the factor is also an isomorphism of strictly ergodic measure-preserving dynamical systems. We refer the reader to [Lab21a] for more details. Moreover, there exists a 4-to-2 cut and project scheme such that the set of occurrences of any pattern in $\Omega_{\mathcal{Z}}$ is a model set, see Theorem 14.1 in [Lab21a].

Exercise 6.1

Prove that the topological partition $\mathcal{P}_{\mathcal{Z}}$ of \mathbb{T}^2 is a Markov partition for the dynamical system $(\mathbb{T}^2, \mathbb{Z}^2, R_{\mathcal{Z}})$.

Exercise 6.2

Using SageMath, verify that the equalities $\beta_0 = \alpha_0$, $\beta_1 = \alpha_1$ and $\beta_2 = \alpha_2$ hold. As a consequence, what is the relation between $\Omega_{\mathcal{V}}$ and $\mathcal{X}_{\mathcal{P}_1,R_1}$? What is the relation between $\Omega_{\mathcal{W}}$ and $\mathcal{X}_{\mathcal{P}_2,R_2}$?

Exercise 6.3

In her PhD thesis [Lep24], Jana Lepšová simplified the Wang shift $\Omega_{\mathcal{U}}$ based on 19 Wang tiles to the Wang shift $\Omega_{\mathcal{Z}}$ based on 16 Wang tiles and proved that $\Omega_{\mathcal{U}}$ and $\Omega_{\mathcal{Z}}$ are topologically conjugate.

Try merging some more labels of the set of Wang tiles \mathcal{Z} . Is the resulting Wang shift still aperiodic?

Jana did not find any more simplification of \mathcal{Z} during her PhD, but it is still an open to show that there is no further simplification. More precisely, we may state the following question.

Open Question 6.2. Prove that there is no Wang shift based on fewer than 16 tiles that is topologically conjugate to $\Omega_{\mathcal{U}}$ and $\Omega_{\mathcal{Z}}$?

More broadly, it seems 16 is a lower bound for the number of tiles of a self-similar aperiodic Wang shift. The Jeandel–Rao Wang shift based on 11 tiles is substitutive but is not self-similar [Lab21c]. Also, the Kari–Culik Wang shifts [Kar96, Cul96] (based on 14 and 13 Wang

tiles) have positive entropy thus are not self-similar [DGG17]. Another aperiodic and selfsimilar set of Wang tiles was described by Ammann and is based on 16 Wang tiles [GS87]. It was generalized to a family of sets of aperiodic and self-similar Wang tiles involving the metallic mean numbers [Lab23a, Lab24]. But 16 tiles is the minimum number of tiles in this family defined from 16, 25, 36, 49, 64, ..., $(n + 3)^2$,... Wang tiles where $n \ge 1$.

Open Question 6.3. Does there exists a self-similar and aperiodic Wang shift based on strictly less than 16 tiles?

The Ammann Wang shift based on 16 Wang tiles and the Wang shift $\Omega_{\mathcal{Z}}$ are not equivalent up to a bijection of the labels, but both are related to golden mean. This raises the following question.

Open Question 6.4. Is $\Omega_{\mathcal{Z}}$ topologically conjugate to the Ammann Wang shift?

The above question might be answered by the computation of set of slopes of nonexpansive directions in the Wang shifts, which is a topological invariant [LMMM23]. In [LMMM23], the slope of the four nonexpansive directions for the minimal subshift within the Jeandel–Rao Wang shift were computed from its associated polygonal partition.

Open Question 6.5. What are the slopes of nonexpansive directions in the Wang shift $\Omega_{\mathcal{Z}}$? in the Ammann Wang shift ?

Another open question raised by the current work is the characterization of Markov polygonal partition for toral \mathbb{Z}^2 -rotations.

Open Question 6.6. Characterize the polygonal partitions \mathcal{P} of \mathbb{T}^2 and toral \mathbb{Z}^2 -action R defined by rotations on \mathbb{T}^2 for which \mathcal{P} is a Markov partition.

7. Solutions to exercises

Solution to Exercise 2.1

There are two elements in the subshift $\mathcal{X}_{\mathcal{L}(x)}$.

Solution to Exercise 2.2

We have

 $\Phi\left(\left(11\right)\right) = \left(\begin{array}{cc} 6 & 1\\ 12 & 10 \end{array}\right), \quad \Phi\left(\left(\begin{array}{cc} 13 & 7 \end{array}\right)\right) = \left(\begin{array}{cc} 5 & 1 & 6\\ 15 & 9 & 12 \end{array}\right), \quad \Phi\left(\left(\begin{array}{cc} 6 & 1\\ 11 & 8 \end{array}\right)\right) = \left(\begin{array}{cc} 12 & 7 & 13\\ 6 & 1 & 3\\ 12 & 10 & 14 \end{array}\right).$

The image $\Phi\left(\begin{pmatrix} 6\\10 \end{pmatrix}\right)$ is not defined because the image of 6 has width 2 while the image of 10 has width 1.

Solution to Exercise 2.3

Left to the reader.

Solution to Exercise 3.1

For every letter $a \in [[0, 15]]$, the 2-dimensional word $\Phi^2(a)$ has height and width at least 2.

The configurations $x = \lim_{n \to \infty} \Phi^{2n} \begin{pmatrix} 8 & 12 \\ 1 & 6 \end{pmatrix}$ and $y = \lim_{n \to \infty} \Phi^{2n+1} \begin{pmatrix} 8 & 12 \\ 1 & 6 \end{pmatrix}$ both belong to \mathcal{X}_{Φ} . Therefore, $\mathcal{X}_{\Phi} \neq \emptyset$.

Solution to Exercise 3.3

The following computes $\Phi^n(12)$ and $\Phi^n(\frac{8}{1} \frac{12}{6})$ when n = 4.

sage: image = Phi([[12]], order=4) 60
sage: image
[[12, 6, 12, 11, 6, 12, 6, 12], [10, 1, 7, 8, 1, 10, 1, 7], [14, 3, 13, 12, 6, 62
14, 3, 13], [11, 6, 12, 15, 5, 11, 6, 12], [8, 1, 10, 9, 1, 8, 1, 7], [12,
6, 14, 12, 6, 12, 2, 13], [10, 1, 7, 10, 1, 7, 0, 7], [14, 3, 13, 14, 3, 13,
3, 13]]
sage: seed = [[1,8],[6,12]] # using Cartesian-like coordinates 63
sage: image = Phi(seed, order=4) 64

The reader may increase the power to n = 5 or n = 6 and compare the output with Figure 1 and Figure 2.

Solution to Exercise 3.4

The method list_dominoes returns the list of 1×2 or 2×1 factors in the language of the associated substitutive shift:

```
sage: XPhi_2x1 = Phi.list_dominoes(direction='horizontal')
                                                                                                                                                                                                                                                                             65
                                                                                                                                                                                                                                                                             66
sage: sorted(XPhi_2x1)
[[[0], [3]], [[1], [2]], [[1], [3]], [[1], [6]], [[2], [0]], [[2], [4]], [[3],
                                                                                                                                                                                                                                                                             67
            [6]], [[4], [1]], [[5], [1]], [[6], [1]], [[6], [5]], [[7], [13]], [[8],
            [12], [[9], [11], [[9], [12], [[10], [14], [[11], [8], [[12], [7],
            [[12], [10]], [[12], [11]], [[12], [15]], [[13], [7]], [[13], [11]], [[13],
            [12]], [[14], [7]], [[14], [11]], [[15], [9]]]
sage: XPhi_1x2 = Phi.list_dominoes(direction='vertical')
                                                                                                                                                                                                                                                                             68
sage: sorted(XPhi_1x2)
                                                                                                                                                                                                                                                                             69
[[[0, 7]], [[1, 7]], [[1, 8]], [[1, 10]], [[2, 13]], [[3, 13]], [[4, 11]], [[5, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], 
                                                                                                                                                                                                                                                                             70
            11]], [[6, 12]], [[6, 14]], [[7, 0]], [[7, 8]], [[7, 10]], [[8, 1]], [[8,
           9], [[9, 1]], [[10, 1]], [[10, 9]], [[11, 4]], [[11, 6]], [[11, 15]], [[12,
               2], [[12, 6]], [[12, 11]], [[12, 15]], [[13, 3]], [[13, 12]], [[13, 14]],
            [[14, 3]], [[14, 12]], [[15, 5]]]
```

Solution to Exercise 3.5

The letter 9 appears only in the image of 13 by Φ . Thus $\begin{pmatrix} 1 \\ 9 \end{pmatrix}$ must be obtained from an image of the letter 13. The vertical domino $\begin{pmatrix} 2 \\ 12 \end{pmatrix}$ must be obtained from the image of 10 or of 15. Therefore $\begin{pmatrix} 1 & 2 \\ 9 & 12 \end{pmatrix}$ is obtained from the horizontal dominoes (13 10) or (13 15). But, from Exercise 3.4, the only horizontal dominoes starting with 13 in the language of Φ are (13 7), (13 11) or (13 12). Therefore $\begin{pmatrix} 1 & 2 \\ 9 & 12 \end{pmatrix} \notin \mathcal{L}_{\Phi}$. S. LABBÉ

Solution to Exercise 3.6 71sage: sorted(Phi.list_2x2_factors()) [[[0, 7], [3, 13]], [[1, 7], [2, 13]], [[1, 7], [3, 13]], [[1, 8], [6, 12]],72[[1, 10], [6, 14]], [[2, 13], [0, 7]], [[2, 13], [4, 11]], [[3, 13], [6,12]], [[4, 11], [1, 8]], [[5, 11], [1, 8]], [[6, 12], [1, 7]], [[6, 12], [1, 10]], [[6, 12], [5, 11]], [[6, 14], [1, 7]], [[6, 14], [5, 11]], [[7, 0], [13, 3]], [[7, 8], [13, 12]], [[7, 10], [13, 14]], [[8, 1], [12, 2]], [[8, 1], [12, 6]], [[8, 9], [12, 11]], [[9, 1], [11, 6]], [[9, 1], [12, 6]], [[10, 1], [14, 3]], [[10, 9], [14, 12]], [[11, 4], [8, 1]], [[11, 6], [8, 1]], [[11, 15], [8, 9]], [[12, 2], [7, 0]], [[12, 2], [11, 4]], [[12, 6],[10, 1]], [[12, 6], [15, 5]], [[12, 11], [7, 8]], [[12, 15], [10, 9]], [[13, 3], [11, 6]], [[13, 3], [12, 6]], [[13, 12], [7, 10]], [[13, 12], [11, 15]], [[13, 12], [12, 11]], [[13, 12], [12, 15]], [[13, 14], [12, 11]], [[14, 3], [11, 6]], [[14, 12], [7, 10]], [[14, 12], [11, 15]], [[15, 5], [9, 1]]] Note that the 2×2 words above are written in Cartesian coordinates. For example, the list of lists [[0, 7], [3, 13]] denotes the 2-dimensional word $\begin{pmatrix} 7 & 13 \\ 0 & 3 \end{pmatrix}$.

There are 96 periodic points of Φ , i.e. the configurations $x \in \mathcal{A}^{\mathbb{Z}^2}$ such that $\Phi^k(x) = x$ for some $k \ge 1$.

73

74 75

76

77

78

```
sage: seeds = sorted(flatten(Phi.prolongable_seeds_list()))
sage: len(seeds)
96
sage: seeds[0]
[ 8 11]
[ 1 4]
```

These seeds do not all belong to the language of the substitution. If all seeds belong to the language of the substitution (computed from the letters), then there exists a unique subshift which is self-similar with respect to the substitution. In this example, it is not true:

```
sage: seeds_as_lists_of_lists = [[list(col[::-1]) for col in m.columns()] for m 79
    in seeds]
sage: XPhi_2x2 = Phi.list_2x2_factors() 80
sage: all(seed in XPhi_2x2 for seed in seeds_as_lists_of_lists) 81
False 82
```

Indeed, only 8 of the 96 seeds belong to the language of the substitution:

```
sage: [seed for seed in seeds_as_lists_of_lists if seed in XPhi_2x2] 83
[[[1, 8], [6, 12]], [[7, 8], [13, 12]], [[8, 9], [12, 11]], [[10, 9], [14, 12]], 84
      [[2, 13], [4, 11]], [[3, 13], [6, 12]], [[6, 14], [5, 11]], [[13, 14], [12,
      11]]]
```

These eight seeds

 $\left\{ \left(\begin{smallmatrix} 8 & 12 \\ 1 & 6 \end{smallmatrix}\right), \left(\begin{smallmatrix} 8 & 12 \\ 7 & 13 \end{smallmatrix}\right), \left(\begin{smallmatrix} 9 & 11 \\ 8 & 12 \end{smallmatrix}\right), \left(\begin{smallmatrix} 9 & 12 \\ 10 & 14 \end{smallmatrix}\right), \left(\begin{smallmatrix} 13 & 11 \\ 2 & 4 \end{smallmatrix}\right), \left(\begin{smallmatrix} 13 & 12 \\ 3 & 6 \end{smallmatrix}\right), \left(\begin{smallmatrix} 14 & 11 \\ 6 & 5 \end{smallmatrix}\right), \left(\begin{smallmatrix} 14 & 11 \\ 13 & 12 \end{smallmatrix}\right) \right\}$

give rise to eight different configurations which are fixed by some power of the substitution and are uniformly recurrent.

Solution to Exercise 3.8

Left to the reader.

Solution to Exercise 3.9

Left to the reader.

(1) The notion of markers is developed in Section 3.4 to help prove that a substitution is recognizable. It can be used here for Φ using the decomposition of Φ as product of simpler one-dimensional substitutions computed in Section 4 and in Section 5. (2) Aperiodicity of \mathcal{X}_{Φ} follows from the recognizability of Φ in \mathcal{X}_{Φ} .

Solution to Exercise 3.11

The 2-dimensional substitution Φ is primitive, because its incidence matrix is primitive:

sage: matrix(Phi).is_primitive()
True

85 86

From Exercise 3.1 and Lemma 3.7, \mathcal{X}_{Φ} is minimal.

The horizontal dominoes in RECURRENTVERTICES $(G_{\Phi}^{2\times 1})$ are

The vertical dominoes in RECURRENTVERTICES $(G_{\Phi}^{1\times 2})$ are

```
sage: Gv = Phi.periodic_vertical_domino_seeds_graph(clean_sources=True) 92
sage: Gv 93
Looped digraph on 36 vertices 94
sage: sorted(Gv.vertices()) 95
[(1, 7), (1, 8), (1, 9), (1, 10), (2, 13), (2, 14), (3, 13), (3, 14), (4, 11), 96
(4, 12), (5, 11), (5, 12), (5, 15), (6, 11), (6, 12), (6, 13), (6, 14), (7,
7), (7, 8), (7, 9), (7, 10), (8, 8), (8, 9), (10, 8), (10, 9), (11, 11),
(11, 15), (12, 11), (12, 12), (12, 15), (13, 11), (13, 12), (13, 13), (13, 14), (14, 11), (14, 12)]
```

There are 96 different 2×2 patterns in RECURRENTVERTICES $(G_{\Phi}^{2\times 2})$. The first of them is shown below:

97 98

99

100 101

102

```
sage: G = Phi.prolongable_seeds_graph(clean_sources=True)
sage: G
Looped digraph on 96 vertices
sage: sorted(G.vertices())[0]
[ 8 11]
[ 1 4]
```

The limit $x = \lim_{n\to\infty} \Phi^{2n} \begin{pmatrix} 8 & 11 \\ 1 & 4 \end{pmatrix}$ defines a configuration of \mathbb{Z}^2 which is fixed by the 2-dimensional morphism Φ^2 . The topological closure $X_x = \overline{\{\sigma^n(x) : n \in \mathbb{Z}^2\}}$ of the orbit under the \mathbb{Z}^2 -shift σ of the configuration x is a subshift which is self-similar with respect to Φ . From Lemma 3.8, we have $\mathcal{X}_{\Phi} \subseteq X_x$. But $\mathcal{X}_{\Phi} \neq X_x$, because $x \notin \mathcal{X}_{\Phi}$. Indeed the central pattern $\begin{pmatrix} 8 & 11 \\ 1 & 4 \end{pmatrix}$ of the configuration x is not in the language of \mathcal{X}_{Φ} , see Exercise 3.7. Also the configuration x is not uniformly recurrent because it contains only one occurrence of the pattern $\begin{pmatrix} 8 & 11 \\ 1 & 4 \end{pmatrix}$.

Let $\mathcal{A} = \llbracket 0, 15 \rrbracket$ and $M = \{0, 1, 2, 3, 4, 5, 6\} \subset \mathcal{A}$. We have $H \cap ((\mathcal{A} \setminus M) \odot^1 M) = \emptyset$ and $H \cap (M \odot^1 (\mathcal{A} \setminus M)) = \emptyset$. Also $M \odot^2 M \cap V = \emptyset$. Therefore, using Lemma 3.11, M is a subset of markers for the direction e_2 in the subshift \mathcal{X}_{Φ} .

Solution to Exercise 3.14

Let $\mathcal{A} = \llbracket 0, 15 \rrbracket$ and $M = \{0, 1, 7, 8, 9, 10\} \subset \mathcal{A}$. We have $V \cap ((\mathcal{A} \setminus M) \odot^2 M) = \emptyset$ and $V \cap (M \odot^2 (\mathcal{A} \setminus M)) = \emptyset$. Also $M \odot^1 M \cap H = \emptyset$. Therefore, using Lemma 3.11, M is a subset of markers for the direction e_1 in the subshift \mathcal{X}_{Φ} .

Solution to Exercise 3.15

Such a substitution can be found in the proof of Theorem 1.1 in Section 4.4.

Solution to Exercise 3.16

The following 2-dimensional morphism $\xi : \mathcal{C} \to [\![0, 15]\!]^{*^2}$, where $\mathcal{C} = [\![0, 15]\!]$: $\xi : \begin{cases} 0 \mapsto (2), & 1 \mapsto (3), & 2 \mapsto (6), & 3 \mapsto (12), & 4 \mapsto (13), \\ 5 \mapsto (14), & 6 \mapsto (2, 0), & 7 \mapsto (4, 1), & 8 \mapsto (5, 1), & 9 \mapsto (6, 1), \\ 10 \mapsto (11, 8), 11 \mapsto (12, 7), 12 \mapsto (12, 10), 13 \mapsto (13, 7), 14 \mapsto (14, 7), \\ 15 \mapsto (15, 9). \end{cases}$

satisfies the requirements with $M = \{0, 1, 7, 8, 9, 10\}$.

Solution to Exercise 4.1

First we define the set \mathcal{Z} of Wang tiles in SageMath:

```
sage: from slabbe import WangTileSet103sage: tiles = ["DOJO", "DOHL", "JMDP", "DMDK", "HPJP", "HPHN", "HKDP", "BOIO",104....: "ILEO", "ILCL", "ALIO", "EPIP", "IPIK", "IKBM", "IKAK", "CNIP"]105sage: Z = WangTileSet([tuple(tile) for tile in tiles])106
```

Then using the dancing links solver:

```
sage: tiling = Z.solver(7,7).solve(solver="dancing_links") 107
sage: tiling.table() # Cartesian-like coordinates 108
[[0, 7, 8, 1, 8, 9, 1], [2, 13, 12, 6, 12, 11, 6], [0, 7, 10, 1, 7, 8, 1], [3, 109
13, 14, 3, 13, 12, 2], [6, 12, 11, 6, 12, 11, 4], [1, 7, 8, 1, 7, 8, 1], [2,
13, 12, 2, 13, 12, 2]]
```

<pre>sage: Z_2x1 = [t.table() for t in Z.tilings_with_surrounding(2,1,radius=2,</pre>	110
<pre>solver="dancing_links")]</pre>	
<pre>sage: sorted(Z_2x1)</pre>	111
[[[0], [3]], [[1], [2]], [[1], [3]], [[1], [6]], [[2], [0]], [[2], [4]], [[3],	112
[6]], [[4], [1]], [[5], [1]], [[6], [1]], [[6], [5]], [[7], [13]], [[8],	
[12]], [[9], [11]], [[9], [12]], [[10], [14]], [[11], [8]], [[12], [7]],	
[[12], [10]], [[12], [11]], [[12], [15]], [[13], [7]], [[13], [11]], [[13],	
[12]], [[14], [7]], [[14], [11]], [[15], [9]]]	
<pre>sage: Z_1x2 = [t.table() for t in Z.tilings_with_surrounding(1,2,radius=2,</pre>	113
<pre>solver="dancing_links")]</pre>	
<pre>sage: sorted(Z_1x2)</pre>	114
[[[0, 7]], [[1, 7]], [[1, 8]], [[1, 10]], [[2, 13]], [[3, 13]], [[4, 11]], [[5, 10]], [[1, 10]],	115
11]], [[6, 12]], [[6, 14]], [[7, 0]], [[7, 8]], [[7, 10]], [[8, 1]], [[8,	
9]], [[9, 1]], [[10, 1]], [[10, 9]], [[11, 4]], [[11, 6]], [[11, 15]], [[12,	
2]], [[12, 6]], [[12, 11]], [[12, 15]], [[13, 3]], [[13, 12]], [[13, 14]],	
[[14, 3]], [[14, 12]], [[15, 5]]]	
	J

Solution to Exercise 4.3

sage: Z.find_markers(i=2, radius=2, solver="dancing_links")
[[0, 1, 2, 3, 4, 5, 6]]

Solution to Exercise 4.4

We compute the set of fusion tiles:

```
sage: M = [0, 1, 2, 3, 4, 5, 6]
                                                                                  118
sage: dominoes_M = sorted((a,b) for [[a,b]] in Z_1x2 if b in M)
                                                                                  119
                                                                                  120
sage: from slabbe.wang_tiles import fusion
sage: new_tiles = [fusion(Z[a],Z[b],2) for (a,b) in dominoes_M]
                                                                                  121
                                                                                  122
sage: new_tiles
[('BD', '0', 'IJ', '0'), ('ID', '0', 'EH', '0'), ('ID', '0', 'CH', 'L'), ('AD', 123
    '0', 'IH', '0'), ('EH', 'P', 'IJ', 'P'), ('EH', 'K', 'ID', 'P'), ('IJ', 'M',
    'ID', 'K'), ('IH', 'K', 'ID', 'K'), ('ID', 'M', 'BD', 'M'), ('ID', 'M', 'AD
    ', 'K'), ('CH', 'P', 'IH', 'P')]
sage: tikz = WangTileSet(new_tiles).tikz(id=None)
                                                                                  124
```

116

117

These tiles represent the result of merging every marker tile with tiles that may appear below of them:



S. LABBÉ

Solution to Exercise 4.5



In Theorem 1.1, we proved that the Wang shift $\Omega_{\mathcal{Z}} \subset \llbracket 0, 15 \rrbracket^{\mathbb{Z}^2}$ is self-similar satisfying $\Omega_{\mathcal{Z}} = \overline{\Phi(\Omega_{\mathcal{Z}})}^{\sigma}$. Thus, we may use the criterion given in Lemma 3.9. More precisely, in this case, we may show that $\mathcal{L}_s(\Omega_z) \subset \mathcal{L}(\mathcal{X}_{\Phi})$ for every shape $s \in \{2 \times 2, 2 \times 1, 1 \times 2\}$. This of course implies that $\mathcal{L}(\Omega_{\mathcal{Z}}) \cap \text{RECURRENTVERTICES}(G^s_{\Phi}) \subset \mathcal{L}(\mathcal{X}_{\Phi})$ for every shape $s \in \{2 \times 2, 2 \times 1, 1 \times 2\}$. In Exercise 3.4 and Exercise 4.2, we observed that $\mathcal{L}_{1\times 2}(\Omega_{\mathcal{Z}}) = \mathcal{L}_{1\times 2}(\mathcal{X}_{\Phi})$ and

 $\mathcal{L}_{2\times 1}(\Omega_{\mathcal{Z}}) = \mathcal{L}_{2\times 1}(\mathcal{X}_{\Phi}):$

```
sage: sorted(XPhi_2x1) == sorted(Z_2x1)
True
sage: sorted(XPhi 1x2) == sorted(Z 1x2)
True
```

129130131

132

In Exercise 3.6, we computed the patterns of shape 2×2 in the language of the 2dimensional substitution Φ . We compute the set $\mathcal{L}_{2\times 2}(\Omega_{\mathcal{Z}})$ below and we observe that $\mathcal{L}_{2\times 2}(\Omega_{\mathcal{Z}}) = \mathcal{L}_{2\times 2}(\mathcal{X}_{\Phi}).$

```
sage: tilings = Z.tilings_with_surrounding(2,2,radius=2, solver="dancing_links") 133
sage: Z_2x2 = [tiling.table() for tiling in tilings]
                                                                                     134
                                                                                     135
sage: len(Z_2x2)
                                                                                     136
45
                                                                                     137
sage: len(XPhi_2x2)
                                                                                     138
45
sage: sorted(XPhi_2x2) == sorted(Z_2x2)
                                                                                     139
True
                                                                                     140
```

We conclude from Lemma 3.9, that $\Omega_{\mathcal{Z}}$ is minimal and $\Omega_{\mathcal{Z}} = \mathcal{X}_{\Phi}$.

Solution to Exercise 4.7 Left to the reader.

Solution to Exercise 5.1

```
sage: from slabbe.arXiv_1903_06137 import self_similar_19_atoms_partition
                                                                                   141
sage: PU = self_similar_19_atoms_partition()
                                                                                   142
sage: merge_dict = {0:0, 1:1, 2:2, 3:3, 4:4, 5:5, 6:6, 7:6, 8:7, 9:8, 10:9,
                                                                                   143
....: 11:10, 12:11, 13:11, 14:12, 15:12, 16:13, 17:14, 18:15}
                                                                                   144
sage: PZ = PU.merge_atoms(merge_dict)
                                                                                   145
sage: graphics_array([PU.plot(), PZ.plot()])
                                                                                   146
Graphics Array of size 1 x 2
                                                                                   147
```

S. LABBÉ

Solution to Exercise 5.2

<pre>sage: z = polygen(QQ, "z")</pre>	148
<pre>sage: K.<phi> = NumberField(z**2-z-1, "phi", embedding=RR(1.6))</phi></pre>	149
<pre>sage: from slabbe import PolyhedronExchangeTransformation as PET</pre>	150
<pre>sage: Gamma0 = matrix.column([(1,0), (0,1)])</pre>	151
<pre>sage: RZe1 = PET.toral_translation(Gamma0, vector((phi^-2,0)))</pre>	152
<pre>sage: RZe2 = PET.toral_translation(Gamma0, vector((0,phi^-2)))</pre>	153
<pre>sage: from slabbe.arXiv_1903_06137 import self_similar_19_atoms_partition</pre>	154
<pre>sage: PU = self_similar_19_atoms_partition()</pre>	155
<pre>sage: merge_dict = {0:0, 1:1, 2:2, 3:3, 4:4, 5:5, 6:6, 7:6, 8:7, 9:8, 10:9,</pre>	156
\dots : 11:10, 12:11, 13:11, 14:12, 15:12, 16:13, 17:14, 18:15}	157
<pre>sage: PZ = PU.merge_atoms(merge_dict)</pre>	158
<pre>sage: from slabbe import PETsCoding</pre>	159
<pre>sage: X_PZ_RZ = PETsCoding((RZe1,RZe2), PZ)</pre>	160
<pre>sage: pattern = X_PZ_RZ.pattern((.1357+1/phi, .2938+1/phi), (8,10))</pre>	161
<pre>sage: m8_10 = matrix.column(col[::-1] for col in pattern)</pre>	162

The pattern of shape 8×10 is

10	1/	11	8	12	10	1/	11
10	14	11	0	14	10	14	T T
1	6	5	1	6	1	6	5
8	12	15	9	11	8	12	15
7	13	11	8	12	7	13	12
1	2	4	1	6	1	3	6
8	12	11	8	12	10	14	11
7	13	12	7	13	7	13	12
1	3	6	1	2	0	3	6
10	14	11	8	12	7	13	12
1	6	5	1	6	1	3	6

Solution to Exercise 5.3

Left to the reader.

Solution to Exercise 5.4

The dynamical system $(\mathbb{T}^2, \mathbb{Z}^2, R_z)$ is minimal. Every atom of the partition \mathcal{P}_z is invariant only under the trivial translation in \mathbb{T}^2 . Thus, from Lemma 5.6, \mathcal{P}_z gives a symbolic representation of $(\mathbb{T}^2, \mathbb{Z}^2, R_z)$.

Solution to Exercise 5.5

It follows from Lemma 5.9.

We compute the induced map $\hat{T}|_W$ in SageMath:





The first two questions are left to the reader. We compute the induced maps in SageMath:

	· · · · ·
<pre>sage: z = polygen(QQ, "z")</pre>	175
<pre>sage: K.<phi> = NumberField(z**2-z-1, "phi", embedding=RR(1.6))</phi></pre>	176
<pre>sage: from slabbe import PolyhedronExchangeTransformation as PET</pre>	177
<pre>sage: Gamma0 = matrix.column([(1,0), (0,1)])</pre>	178
<pre>sage: RZe1 = PET.toral_translation(Gamma0, vector((phi^-2,0)))</pre>	179
<pre>sage: RZe2 = PET.toral_translation(Gamma0, vector((0,phi^-2)))</pre>	180
<pre>sage: y_ineq = [phi^-1, 0, -1] # y <= phi^-1: the window W_0</pre>	181
<pre>sage: R1e1,_ = RZe1.induced_transformation(y_ineq)</pre>	182
<pre>sage: R1e2,_ = RZe2.induced_transformation(y_ineq)</pre>	183
sage: R1e1	184
Polyhedron Exchange Transformation of	185
Polyhedron partition of 2 atoms with 2 letters	186
with translations {0: (-phi + 2, 0), 1: (-phi + 1, 0)}	187
sage: R1e2	188
Polyhedron Exchange Transformation of	189
Polyhedron partition of 2 atoms with 2 letters	190
with translations {0: (0, -phi + 2), 1: (0, -2*phi + 3)}	191
P1a1 plat() P1a2 plat()	
ocmain partition image partition 0.6 0.5 0.7 0.6 0.8 0.5	



0.8 1.0

We compute the induced substitution and the induced partition $\hat{\mathcal{P}}|_W$ in SageMath:





```
203
sage: z = polygen(QQ, "z")
                                                                                                                                             204
sage: K.<phi> = NumberField(z**2-z-1, "phi", embedding=RR(1.6))
sage: from slabbe import PolyhedronExchangeTransformation as PET
                                                                                                                                             205
sage: Gamma0 = matrix.column([(1,0), (0,1)])
                                                                                                                                             206
sage: RZe1 = PET.toral_translation(Gamma0, vector((phi^-2,0)))
                                                                                                                                             207
sage: RZe2 = PET.toral_translation(Gamma0, vector((0,phi^-2)))
                                                                                                                                             208
sage: from slabbe.arXiv_1903_06137 import self_similar_19_atoms_partition
                                                                                                                                             209
sage: PU = self_similar_19_atoms_partition()
                                                                                                                                             210
                                                                                                                                             211
sage: merge_dict = {0:0, 1:1, 2:2, 3:3, 4:4, 5:5, 6:6, 7:6, 8:7, 9:8, 10:9,
....: 11:10, 12:11, 13:11, 14:12, 15:12, 16:13, 17:14, 18:15}
                                                                                                                                             212
                                                                                                                                             213
sage: PZ = PU.merge_atoms(merge_dict)
sage: from slabbe import PETsCoding
                                                                                                                                             214
                                                                                                                                             215
sage: X_PZ_RZ = PETsCoding((RZe1,RZe2), PZ)
sage: _,d22 = X_PZ_RZ.partition_for_patterns((2,2))
                                                                                                                                             216
sage: XPZRZ_2x2 = [[list(col) for col in v] for v in d22.values()]
                                                                                                                                             217
sage: XPZRZ_2x2
                                                                                                                                             218
[[[0, 7], [3, 13]], [[1, 7], [2, 13]], [[1, 7], [3, 13]], [[1, 8], [6, 12]],
                                                                                                                                             219
      [[1, 10], [6, 14]], [[2, 13], [0, 7]], [[2, 13], [4, 11]], [[3, 13], [6,
      12], [[4, 11], [1, 8]], [[5, 11], [1, 8]], [[6, 12], [1, 7]], [[6, 12], [1,
       10], [[6, 12], [5, 11]], [[6, 14], [1, 7]], [[6, 14], [5, 11]], [[7, 0],
      [13, 3]], [[7, 8], [13, 12]], [[7, 10], [13, 14]], [[8, 1], [12, 2]], [[8,
      1], [12, 6]], [[8, 9], [12, 11]], [[9, 1], [11, 6]], [[9, 1], [12, 6]],
      [[10, 1], [14, 3]], [[10, 9], [14, 12]], [[11, 4], [8, 1]], [[11, 6], [8,
      1], [[11, 15], [8, 9]], [[12, 2], [7, 0]], [[12, 2], [11, 4]], [[12, 6],
      [10, 1]], [[12, 6], [15, 5]], [[12, 11], [7, 8]], [[12, 15], [10, 9]], [[13,
       3], [11, 6]], [[13, 3], [12, 6]], [[13, 12], [7, 10]], [[13, 12], [11,
      15]], [[13, 12], [12, 11]], [[13, 12], [12, 15]], [[13, 14], [12, 11]],
      [[14, 3], [11, 6]], [[14, 12], [7, 10]], [[14, 12], [11, 15]], [[15, 5], [9,
       1]]]
                                                                                                                                             220
sage: _,d12 = X_PZ_RZ.partition_for_patterns((1,2))
sage: XPZRZ_1x2 = [[list(col) for col in v] for v in d12.values()]
                                                                                                                                             221
                                                                                                                                             222
sage: XPZRZ_1x2
                                                                                                                                             223
[[[0, 7]], [[1, 7]], [[1, 8]], [[1, 10]], [[2, 13]], [[3, 13]], [[4, 11]], [[5, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], [[1, 10]], 
      11]], [[6, 12]], [[6, 14]], [[7, 0]], [[7, 8]], [[7, 10]], [[8, 1]], [[8,
      9], [[9, 1]], [[10, 1]], [[10, 9]], [[11, 4]], [[11, 6]], [[11, 15]], [[12,
       2]], [[12, 6]], [[12, 11]], [[12, 15]], [[13, 3]], [[13, 12]], [[13, 14]],
      [[14, 3]], [[14, 12]], [[15, 5]]]
sage: _,d21 = X_PZ_RZ.partition_for_patterns((2,1))
                                                                                                                                             224
                                                                                                                                             225
sage: XPZRZ_2x1 = [[list(col) for col in v] for v in d21.values()]
                                                                                                                                             226
sage: XPZRZ_2x1
                                                                                                                                             227
[[[0], [3]], [[1], [2]], [[1], [3]], [[1], [6]], [[2], [0]], [[2], [4]], [[3],
      [6]], [[4], [1]], [[5], [1]], [[6], [1]], [[6], [5]], [[7], [13]], [[8],
      [12], [[9], [11], [[9], [12], [[10], [14], [[11], [8], [[12], [7],
      [[12], [10]], [[12], [11]], [[12], [15]], [[13], [7]], [[13], [11]], [[13],
      [12]], [[14], [7]], [[14], [11]], [[15], [9]]]
```

62

In Theorem 1.2, we proved that the symbolic dynamical system $\mathcal{X}_{\mathcal{P}_{Z},R_{Z}} \subset [\![0,15]\!]^{\mathbb{Z}^{2}}$ is self-similar satisfying $\mathcal{X}_{\mathcal{P}_{Z},R_{Z}} = \overline{\Phi(\mathcal{X}_{\mathcal{P}_{Z},R_{Z}})}^{\sigma}$. Thus, we may use the criterion given in Lemma 3.9. In Exercise 5.9, we computed $\mathcal{L}_{1\times 2}(\mathcal{X}_{\mathcal{P}_{Z},R_{Z}})$, $\mathcal{L}_{2\times 1}(\mathcal{X}_{\mathcal{P}_{Z},R_{Z}})$ and $\mathcal{L}_{2\times 2}(\mathcal{X}_{\mathcal{P}_{Z},R_{Z}})$. Comparing with Exercise 3.4 and Exercise 3.6, we observed that

 $\begin{aligned} \mathcal{L}_{1\times 2}(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}) &= \mathcal{L}_{1\times 2}(\mathcal{X}_{\Phi}), \\ \mathcal{L}_{2\times 1}(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}) &= \mathcal{L}_{2\times 1}(\mathcal{X}_{\Phi}), \\ \mathcal{L}_{2\times 2}(\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}) &= \mathcal{L}_{2\times 2}(\mathcal{X}_{\Phi}). \end{aligned}$

228229

230

231

232 233

```
sage: sorted(XPhi_2x1) == sorted(XPZRZ_2x1)
True
sage: sorted(XPhi_1x2) == sorted(XPZRZ_1x2)
True
sage: sorted(XPhi_2x2) == sorted(XPZRZ_2x2)
True
```

We conclude from Lemma 3.9, that $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}}$ is minimal and $\mathcal{X}_{\mathcal{P}_{\mathcal{Z}},R_{\mathcal{Z}}} = \mathcal{X}_{\Phi}$.

Solution to Exercise 5.11

Left to the reader.

Solution to Exercise 6.1

Since $\Omega_{\mathcal{Z}} = \mathcal{X}_{\Phi} = \mathcal{X}_{\mathcal{P}_{\mathcal{Z}}, R_{\mathcal{Z}}}$ and $\Omega_{\mathcal{Z}}$ is a subshift of finite type, it follows that $\mathcal{P}_{\mathcal{Z}}$ is a Markov partition for the dynamical system $(\mathbb{T}^2, \mathbb{Z}^2, R_{\mathcal{Z}})$.

Solution to Exercise 6.2

Left to the reader.

Solution to Exercise 6.3

Left to the reader.

References

[AKY19]	I. Alevy, R. Kenyon, and R. Yi. A family of minimal and renormalizable rectangle exchange	ge
	maps. Ergodic Theory and Dynamical Systems, pages 1-28, 2019. doi:10.1017/etds.2019.7	7.
[AS18]	G. Audemard and L. Simon. On the glucose SAT solver. International Journal on Artifica	al

- Intelligence Tools, 27(01):1840001, 2018. doi:10.1142/S0218213018400018.
- [BD14] V. Berthé and V. Delecroix. Beyond substitutive dynamical systems: S-adic expansions. In Numeration and substitution 2012, RIMS Kôkyûroku Bessatsu, B46, pages 81–123. Res. Inst. Math. Sci. (RIMS), Kyoto, 2014.
- [Ber66] R. Berger. The undecidability of the domino problem. Mem. Amer. Math. Soc. No., 66:72, 1966.
- [Ber80] J. Berstel. Mots de Fibonacci. In Séminaire d'Informatique Théorique, L.I.T.P., Paris:57–78, 1980.

S. LABBÉ

- [BSTY19] V. Berthé, W. Steiner, J. M. Thuswaldner, and R. Yassawi. Recognizability for sequences of morphisms. *Ergodic Theory Dynam. Systems*, 39(11):2896–2931, 2019. doi:10.1017/etds.2017.144.
- [CH73] E. M. Coven and G. A. Hedlund. Sequences with minimal block growth. Math. Systems Theory, 7:138–153, 1973. doi:10.1007/BF01762232.
- [CKR10] E. Charlier, T. Kärki, and M. Rigo. Multidimensional generalized automatic sequences and shape-symmetric morphic words. *Discrete Math.*, 310(6-7):1238–1252, 2010. doi:10.1016/j.disc.2009.12.002.
- [Cul96] K. Culik, II. An aperiodic set of 13 Wang tiles. Discrete Math., 160(1-3):245-251, 1996.
 doi:10.1016/S0012-365X(96)00118-5.
- [DGG17] B. Durand, G. Gamard, and A. Grandjean. Aperiodic tilings and entropy. Theoret. Comput. Sci., 666:36-47, 2017. doi:10.1016/j.tcs.2016.12.013.
- [ES97] M. Einsiedler and K. Schmidt. Markov partitions and homoclinic points of algebraic \mathbb{Z}^{d} -actions. *Tr. Mat. Inst. Steklova*, 216(Din. Sist. i Smezhnye Vopr.):265–284, 1997.
- [Fog02] N. P. Fogg. Substitutions in Dynamics, Arithmetics and Combinatorics, volume 1794 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2002. Edited by V. Berthé, S. Ferenczi, C. Mauduit and A. Siegel. doi:10.1007/b13861.
- [Fra18] N. P. Frank. Introduction to hierarchical tiling dynamical systems. February 2018.
- [GO20] L. Gurobi Optimization. Gurobi optimizer reference manual, 2020. http://www.gurobi.com.
- [GS87] B. Grünbaum and G. C. Shephard. *Tilings and patterns*. W. H. Freeman and Company, New York, 1987.
- [HM10] M. Hochman and T. Meyerovitch. A characterization of the entropies of multidimensional shifts of finite type. Ann. of Math. (2), 171(3):2011–2038, 2010. doi:10.4007/annals.2010.171.2011.
- [Hoc16] M. Hochman. Multidimensional shifts of finite type and sofic shifts. In Combinatorics, words and symbolic dynamics, volume 159 of Encyclopedia Math. Appl., pages 296–358. Cambridge Univ. Press, Cambridge, 2016. doi:10.1017/CB09781139924733.010.
- [Hoo13] W. P. Hooper. Renormalization of polygon exchange maps arising from corner percolation. Invent. Math., 191(2):255–320, 2013. doi:10.1007/s00222-012-0393-4.
- [JR21] E. Jeandel and M. Rao. An aperiodic set of 11 Wang tiles. Adv. Comb., pages Paper No. 1, 37, 2021. doi:10.19086/aic.18614.
- [Kar96] J. Kari. A small aperiodic set of Wang tiles. Discrete Math., 160(1-3):259-264, 1996. doi:10.1016/0012-365X(95)00120-L.
- [Knu00] D. E. Knuth. Dancing links. November 2000. Millenial Perspectives in Computer Science, 2000, 187–214.
- [Lab18] S. Labbé. Comparison of wang tiling solvers. http://www.slabbe.org/blogue/2018/12/ comparison-of-wang-tiling-solvers/, December 2018.
- [Lab19] S. Labbé. A self-similar aperiodic set of 19 Wang tiles. Geom. Dedicata, 201:81–109, 2019. doi:10.1007/s10711-018-0384-8.
- [Lab21a] S. Labbé. Markov partitions for toral Z²-rotations featuring Jeandel-Rao Wang shift and model sets. Ann. H. Lebesgue, 4:283–324, 2021. doi:10.5802/ahl.73.
- [Lab21b] S. Labbé. Rauzy induction of polygon partitions and toral \mathbb{Z}^2 -rotations. J. Mod. Dyn., 17:481–528, 2021. doi:10.3934/jmd.2021017.
- [Lab21c] S. Labbé. Substitutive Structure of Jeandel-Rao Aperiodic Tilings. Discrete Comput. Geom., 65(3):800–855, 2021. doi:10.1007/s00454-019-00153-3.
- [Lab23a] S. Labbé. Metallic mean Wang tiles I: self-similarity, aperiodicity and minimality. December 2023. doi:10.48550/arXiv.2312.03652. arxiv:2312.03652.
- [Lab23b] S. Labbé. Optional SageMath Package slabbe (Version 0.7.6). https://pypi.python.org/ pypi/slabbe/, 2023.
- [Lab24] S. Labbé. Metallic mean Wang tiles II: the dynamics of an aperiodic computer chip. March 2024. doi:10.48550/arXiv.2403.03197. arxiv:2403.03197.
- [Lep24] J. Lepšová. Substitutive structures in combinatorics, number theory, and discrete geometry. PhD Thesis, Université de Bordeaux, Czech Technical University in Prague, 2024. https://theses. fr/2024BORD0083.

- [Lin04] D. Lind. Multi-dimensional symbolic dynamics. In Symbolic dynamics and its applications, volume 60 of Proc. Sympos. Appl. Math., pages 61–79. Amer. Math. Soc., Providence, RI, 2004. doi:10.1090/psapm/060/2078846.
- [LM95] D. Lind and B. Marcus. An Introduction to Symbolic Dynamics and Coding. Cambridge University Press, Cambridge, 1995. doi:10.1017/CB09780511626302.
- [LMMM23] S. Labbé, C. Mann, and J. McLoud-Mann. Nonexpansive directions in the Jeandel-Rao Wang shift. Discrete Contin. Dyn. Syst., 43(9):3213-3250, 2023. doi:10.3934/dcds.2023046.
- [Lot02] M. Lothaire. Algebraic Combinatorics on Words, volume 90 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 2002.
- [MH40] M. Morse and G. A. Hedlund. Symbolic dynamics II. Sturmian trajectories. Amer. J. Math., 62:1–42, 1940. doi:10.2307/2371431.
- [Mos92] B. Mossé. Puissances de mots et reconnaissabilité des points fixes d'une substitution. Theoret. Comput. Sci., 99(2):327-334, 1992. doi:10.1016/0304-3975(92)90357-L.
- [Moz89] S. Mozes. Tilings, substitution systems and dynamical systems generated by them. J. Analyse Math., 53:139–186, 1989. doi:10.1007/BF02793412.
- [Que10] M. Queffélec. Substitution dynamical systems—spectral analysis, volume 1294 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, second edition, 2010. doi:10.1007/978-3-642-11212-6.
- [Rau77] G. Rauzy. Une généralisation du développement en fraction continue. In Séminaire Delange-Pisot-Poitou, 18e année: 1976/77, Théorie des nombres, Fasc. 1, pages Exp. No. 15, 16. Secrétariat Math., Paris, 1977.
- [Rau79] G. Rauzy. Échanges d'intervalles et transformations induites. Acta Arith., 34(4):315–328, 1979. doi:10.4064/aa-34-4-315-328.
- [Rob71] R. M. Robinson. Undecidability and nonperiodicity for tilings of the plane. Invent. Math., 12:177– 209, 1971. doi:10.1007/BF01418780.
- [Sag24] Sage Developers. SageMath, the Sage Mathematics Software System (Version 10.4), 2024. http: //www.sagemath.org.
- [Sch01] K. Schmidt. Multi-dimensional symbolic dynamical systems. In Codes, Systems, and Graphical Models (Minneapolis, MN, 1999), volume 123 of IMA Vol. Math. Appl., pages 67–82. Springer, New York, 2001. doi:10.1007/978-1-4613-0165-3_3.
- [Sch11] R. E. Schwartz. Outer billiards, quarter turn compositions, and polytope exchange transformations (preprint), 2011.
- [Sch14] R. E. Schwartz. The Octogonal PETs, volume 197 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2014. doi:10.1090/surv/197.
- [SE05] N. Sörensson and N. Een. Minisat v1.13-a SAT solver with conflict-clause minimization. International Conference on Theory and Applications of Satisfiability Testing, 01 2005.
- [Sol98] B. Solomyak. Nonperiodicity implies unique composition for self-similar translationally finite tilings. Discrete Comput. Geom., 20(2):265–279, 1998. doi:10.1007/PL00009386.
- [Wal82] P. Walters. An Introduction to Ergodic Theory, volume 79 of GTM. Springer-Verlag, New York-Berlin, 1982.
- [Wan61] H. Wang. Proving theorems by pattern recognition II. Bell System Technical Journal, 40(1):1–41, January 1961. doi:10.1002/j.1538-7305.1961.tb03975.x.

(S. Labbé) UNIV. BORDEAUX, CNRS, BORDEAUX INP, LABRI, UMR 5800, F-33400, TALENCE, FRANCE

Email address: sebastien.labbe@labri.fr