Classification of linear operators satisfying $(Au, v) = (u, A^r v)$ or $(Au, A^r v) = (u, v)$ on a vector space with indefinite scalar product

Victor Senoguchi Borges^b, Iryna Kashuba^b, Vladimir V. Sergeichuk^a, Eduardo Ventilari Sodré^b, André Zaidan^b

^aInstitute of Mathematics, Tereshchenkivska 3, Kiev, Ukraine ^bInstituto de Matemática e Estatística, Universidade de São Paulo, Brasil

Abstract

We classify all linear operators $\mathcal{A}: V \to V$ satisfying $(\mathcal{A}u, v) = (u, \mathcal{A}^r v)$ and all linear operators satisfying $(\mathcal{A}u, \mathcal{A}^r v) = (u, v)$ with r = 2, 3, ... on a complex, real, or quaternion vector space with scalar product given by a nonsingular symmetric, skew-symmetric, Hermitian, or skew-Hermitian form.

Keywords: indefinite scalar product; selfadjoint operators; quaternions 2000 MSC: 15A21, 15A63, 15B57, 46C20, 47B50

1. Introduction

Let \mathbb{F} be \mathbb{C} , \mathbb{R} , or the skew field of quaternions \mathbb{H} . Let V be a finite dimensional right vector space over \mathbb{F} with scalar product given by a nonsingular form $\mathcal{F}: V \times V \to \mathbb{F}$ that is symmetric or skew-symmetric if $\mathbb{F} \in {\mathbb{C}, \mathbb{R}}$, and Hermitian or skew-Hermitian if $\mathbb{F} \in {\mathbb{C}, \mathbb{H}}$. Let $r \in {1, 2, ...}$. A linear operator $\mathcal{A}: V \to V$ is *r*-selfadjoint if

 $\mathcal{F}(\mathcal{A}u, v) = \mathcal{F}(u, \mathcal{A}^r v) \quad \text{for all } u, v \in V;$

^{*}Linear Algebra Appl. 2021, DOI: 10.1016/j.laa.2020.12.005

Email addresses: victorsenoguchi@gmail.com (Victor Senoguchi Borges), kashuba@ime.usp.br (Iryna Kashuba), sergeich@imath.kiev.ua (Vladimir V. Sergeichuk), eduvsodre@usp.br (Eduardo Ventilari Sodré), andre.zaidan@gmail.com (André Zaidan)

 $\mathcal{A}: V \to V$ is *r*-unitary if it is nonsingular and

$$\mathcal{F}(\mathcal{A}u, \mathcal{A}^r v) = \mathcal{F}(u, v) \text{ for all } u, v \in V.$$

The 1-selfadjoint operators are selfadjoint operators on spaces with indefinite scalar product; their classification is given in [2, 6, 7, 11, 17, 18, 21, 22]. The 1-unitary operators are unitary operators on spaces with indefinite scalar product; their classification is given in [1, 6, 7, 11, 20, 22, 24].

We give canonical matrices of r-selfadjoint operators and r-unitary operators for $r \ge 2$. We use the method developed in [22], which reduces the problem of classifying systems of forms and linear mappings to the problem of classifying systems of linear mappings. This method allows to consider the problems of classifying r-selfadjoint operators and r-unitary operators as the same classification problem.

Later on, we use the term "(-r)-selfadjoint operators" instead of "runitary operators" and solve the problem of classifying r-selfadjoint operators for each $r \in \mathbb{Z} \setminus \{-1, 0, 1\}$. In matrix form, this problem is formulated as follows: we consider pairs (A, F) of $n \times n$ matrices over \mathbb{C} or \mathbb{R} satisfying

$$A^T F = F A^r, \qquad F^T = F \text{ is nonsingular}$$
(1)

and give their canonical form with respect to transformations

$$(A, F) \mapsto (S^{-1}AS, S^TFS), \qquad S \text{ is nonsingular;}$$
(2)

we also consider matrix pairs (A, F) over \mathbb{C} or \mathbb{H} satisfying

$$A^*F = FA^r, \qquad F^* = F \text{ is nonsingular} \tag{3}$$

and give their canonical form with respect to transformations

$$(A, F) \mapsto (S^{-1}AS, S^*FS), \qquad S \text{ is nonsingular}$$
(4)

(A is nonsingular if r < 0, and $S^* := \overline{S}^T$).

This research was inspired by the articles [3, 4, 5, 12, 13, 14, 15, 16], in which Catral, Lebtahi, Romero, Stuart, Thome, and Weaver study $\{R, s + 1, k\}$ -potent (respectively, $\{R, s+1, k, *\}$ -potent) matrices; i.e., those matrices $A \in \mathbb{C}^{n \times n}$ that satisfy $RA = A^{s+1}R$ (respectively, $RA^* = A^{s+1}R$), in which $R \in \mathbb{C}^{n \times n}$ is a given matrix satisfying $R^k = 1$ and s, k are positive integers; compare with (1) and (3). Each sesquilinear form $\mathcal{F}: V \times V \to \mathbb{F}$ that we consider is semilinear in the first argument and linear in the second; $\mathcal{F}: V \to V$ is *skew-Hermitian* if $\mathcal{F}(u,v) = -\overline{\mathcal{F}(v,u)}$ for all $u, v \in V$. We do not consider skew-Hermitian forms over \mathbb{C} since if $\mathcal{F}(u,v)$ is skew-Hermitian, then $i\mathcal{F}(u,v)$ is Hermitian.

Define the matrix

$$(a+bi)^{\mathbb{R}} \coloneqq \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \quad \text{for each } a+bi \in \mathbb{C} \ (a,b \in \mathbb{R}), \tag{5}$$

and the direct sum of matrix pairs

$$(A_1, F_1) \oplus (A_2, F_2) \coloneqq \left(\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}, \begin{bmatrix} F_1 & 0 \\ 0 & F_2 \end{bmatrix} \right)$$

The notation $\lambda \searrow \mu$ means that a parameter λ is determined up to replacement by μ . We write $\lambda \in \mathbb{C}^{\ddagger}$ if λ is a complex parameter that is determined up to replacement by its complex conjugate $\overline{\lambda}$. We write " $(A, \pm F)$ " instead of "(A, F) and (A, -F)". We denote by 0_n and I_n the $n \times n$ zero and identity matrices.

Our main result is the following theorem.

Theorem 1. Let $r \in \mathbb{Z} \setminus \{-1, 0, 1\}$.

(A) Let $V_{\mathbb{C}}$ be a vector space over \mathbb{C} .

(a₁) Let A be an r-selfadjoint operator on V_C with a nonsingular symmetric form F. Then there exists a basis of V_C in which the pair (A, F) is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$([\lambda], [1]), \qquad \left(\begin{bmatrix} \mu & 0 \\ 0 & \mu^r \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right),$$

in which $\lambda, \mu \in \mathbb{C}$, $\lambda^r = \lambda$, $\mu^{r^2} = \mu$, $\mu^r \neq \mu$, $\mu \searrow \mu^r$.

(a₂) Let A be an r-selfadjoint operator on V_C with a nonsingular Hermitian form F. Then there exists a basis of V_C in which the pair (A, F) is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$([\lambda], \pm [1]), \qquad \left(\begin{bmatrix} \mu & 0 \\ 0 & \bar{\mu}^r \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right),$$

in which $\lambda, \mu \in \mathbb{C}, \ \lambda^r = \lambda, \ \mu^{r^2} = \mu, \ \mu^r \neq \overline{\mu}, \ \mu \supset \overline{\mu}^r.$

(a₃) Let \mathcal{A} be an r-selfadjoint operator on $V_{\mathbb{C}}$ with a nonsingular skewsymmetric form \mathcal{F} . Then there exists a basis of $V_{\mathbb{C}}$ in which the pair $(\mathcal{A}, \mathcal{F})$ is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$\left(\begin{bmatrix} \lambda & 0 \\ 0 & \lambda^r \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right),$$

in which $\lambda \in \mathbb{C}$, $\lambda^{r^2} = \lambda$, and $\lambda \supset \lambda^r$.

- (B) Let $V_{\mathbb{R}}$ be a vector space over \mathbb{R} .
 - (b1) Let A be an r-selfadjoint operator on V_ℝ with a nonsingular symmetric form F. Then there exists a basis of V_ℝ in which the pair (A, F) is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$\begin{pmatrix} [0], \pm [1] \end{pmatrix}, \quad ([1], \pm [1]), \quad ([-1], \pm [1]) \text{ if } r \text{ is odd}, \\ \begin{pmatrix} \lambda^{\mathbb{R}}, \begin{bmatrix} 0 & 1\\ 1 & 0 \end{bmatrix} \end{pmatrix}, \quad (\mu^{\mathbb{R}}, \pm I_2), \quad \begin{pmatrix} \begin{bmatrix} \nu^{\mathbb{R}} & 0\\ 0 & (\bar{\nu}^r)^{\mathbb{R}} \end{bmatrix}, \begin{bmatrix} 0 & I_2\\ I_2 & 0 \end{bmatrix} \end{pmatrix},$$

in which $\lambda, \mu, \nu \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \lambda^{r} = \lambda, \ \mu^{r} = \overline{\mu}, \ \nu^{r^{2}} = \nu, \ \nu^{r} \neq \nu, \ \nu^{r} \neq \overline{\nu}, \ \nu \searrow \nu^{r}.$

(b₂) Let \mathcal{A} be an r-selfadjoint operator on $V_{\mathbb{R}}$ with a nonsingular skewsymmetric form \mathcal{F} . Then there exists a basis of $V_{\mathbb{R}}$ in which the pair $(\mathcal{A}, \mathcal{F})$ is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$\begin{pmatrix} 0_2, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \end{pmatrix}, \quad \begin{pmatrix} I_2, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \end{pmatrix}, \quad \begin{pmatrix} -I_2, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \end{pmatrix} \text{ if } r \text{ is odd,} \\ \begin{pmatrix} \lambda^{\mathbb{R}}, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \end{pmatrix}, \quad \begin{pmatrix} \begin{bmatrix} \mu^{\mathbb{R}} & 0 \\ 0 & (\bar{\mu}^r)^{\mathbb{R}} \end{bmatrix}, \begin{bmatrix} 0 & -I_2 \\ I_2 & 0 \end{bmatrix} \end{pmatrix},$$

 $in \ which \ \lambda, \mu \in \mathbb{C}^{\updownarrow} \smallsetminus \mathbb{R}, \ \lambda^r = \bar{\lambda}, \ \mu^{r^2} = \mu, \ \mu^r \neq \bar{\mu}, \ \mu \searrow \mu^r.$

(C) Let $V_{\mathbb{H}}$ be a right vector space over \mathbb{H} .

(c₁) Let \mathcal{A} be an r-selfadjoint operator on $V_{\mathbb{H}}$ with a nonsingular Hermitian form \mathcal{F} with respect to quaternion conjugation

$$h = a + bi + cj + dk \quad \mapsto \quad \overline{h} = a - bi - cj - dk, \qquad a, b, c, d \in \mathbb{R}.$$
(6)

Then there exists a basis of $V_{\mathbb{H}}$ in which the pair $(\mathcal{A}, \mathcal{F})$ is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$([\lambda], \pm [1]), \qquad \left(\begin{bmatrix} \mu & 0\\ 0 & \overline{\mu}^r \end{bmatrix}, \begin{bmatrix} 0 & 1\\ 1 & 0 \end{bmatrix} \right),$$

 $in \ which \ \lambda, \mu \in \mathbb{C}^{\updownarrow}, \ \lambda^r = \bar{\lambda}, \ \mu^{r^2} = \mu, \ \mu^r \neq \bar{\mu}, \ \mu \searrow \mu^r.$

(c₂) Let \mathcal{A} be an r-selfadjoint operator on $V_{\mathbb{H}}$ with a nonsingular Hermitian form \mathcal{F} with respect to quaternion semiconjugation

$$h = a + bi + cj + dk \quad \mapsto \quad \tilde{h} = a - bi + cj + dk, \qquad a, b, c, d \in \mathbb{R}.$$
(7)

Then there exists a basis of $V_{\mathbb{H}}$ in which the pair $(\mathcal{A}, \mathcal{F})$ is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$\begin{array}{l} ([\lambda], \pm [1]) \ if \ \lambda \notin \mathbb{R}, \\ ([\mu], [j]), \\ \begin{pmatrix} \nu & 0 \\ 0 & \bar{\nu}^r \end{pmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \end{pmatrix},$$

in which $\lambda, \mu, \nu \in \mathbb{C}^{\ddagger}$, $\lambda^r = \overline{\lambda}$, $\mu^r = \mu \notin \mathbb{R}$, $\nu^{r^2} = \nu$, $\nu^r \neq \nu$, $\nu^r \neq \overline{\nu}$, $\nu \supset \nu^r$.

(c₃) Let A be an r-selfadjoint operator on V_H with a nonsingular form F that is skew-Hermitian with respect to quaternion conjugation
(6). Then there exists a basis of V_H in which the pair (A, F) is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$\begin{array}{l} ([\lambda], \pm[i]) \ if \ \lambda \notin \mathbb{R}, \\ ([\mu], [j]), \\ \end{array} \begin{pmatrix} \begin{pmatrix} \nu & 0 \\ 0 & \bar{\nu}^r \end{pmatrix}, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \end{pmatrix}, \\ \end{array}$$

 $\begin{array}{l} in \ which \ \lambda, \mu, \nu \in \mathbb{C}^{\ddagger}, \ \lambda^{r} = \bar{\lambda}, \ \mu \notin \mathbb{R}, \ \mu^{r} = \mu, \ \nu^{r^{2}} = \nu, \ \nu^{r} \neq \nu, \ \nu^{r} \neq \bar{\nu}, \\ \nu \searrow \nu^{r}. \end{array}$

(c₄) Let A be an r-selfadjoint operator on V_H with a nonsingular form F that is skew-Hermitian with respect to quaternion semiconjugation (7). Then there exists a basis of V_H in which the pair (A, F) is given by a direct sum, uniquely determined up to permutations of summands, of pairs of the form

$$([\lambda], \pm[i]), \qquad \left(\begin{bmatrix} \mu & 0\\ 0 & \bar{\mu}^r \end{bmatrix}, \begin{bmatrix} 0 & -1\\ 1 & 0 \end{bmatrix} \right),$$

in which $\lambda, \mu \in \mathbb{C}^{\ddagger}, \ \lambda^r = \bar{\lambda}, \ \mu^{r^2} = \mu, \ \mu^r \neq \bar{\mu}, \ \mu \supset \mu^r.$

Each condition $\lambda^{r^2} = \lambda$, $\lambda^r = \lambda$, or $\lambda^r = \overline{\lambda}$ implies that $\lambda \neq 0$ if r < 0. Theorem 1 remains true if \mathbb{C} , \mathbb{R} , and \mathbb{H} are replaced by an algebraically closed field of zero characteristic, a real closed field, and the skew field of quaternions over a real closed field, respectively.

An involution $a \mapsto \widetilde{a}$ on a field or skew field \mathbb{F} is a bijection $\mathbb{F} \to \mathbb{F}$ satisfying

$$\widetilde{a+b} = \widetilde{a} + \widetilde{b}, \quad \widetilde{ab} = \widetilde{b} \ \widetilde{a}, \quad \widetilde{\widetilde{a}} = a \quad \text{for all } a, b \in \mathbb{F}.$$

If an involution on \mathbb{H} is not quaternion conjugation (6), then it is quaternion semiconjugation (7) in a suitable set of the fundamental units i, j, k; see [24, Lemma 2.2].

2. Reduction of the problem of classifying r-selfadjoint operators to the problem of classifying matrices under similarity

We prove Theorem 1 in the next section by the method that is developed in [22]. It reduces the problem of classifying systems of linear mappings and forms to the problem of classifying systems of linear mappings. Bilinear and sesquilinear forms, pairs of symmetric, skew-symmetric, and Hermitian forms, unitary and selfadjoint operators on a vector space with indefinite scalar product are classified in [22] over a field \mathbb{K} of characteristic not 2 up to classification of Hermitian forms over finite extensions of \mathbb{K} (and so they are fully classified over \mathbb{R} and \mathbb{C}).

The reader is expected to be familiar with this method; it is described in details in [24] and is used in [9, 10, 19, 23]. In this section, we sketchily describe it in a special case: for the problem of classifying r-selfadjoint operators. Systems consisting of vector spaces and of linear mappings and forms on them are considered as *representations of mixed graphs*; i.e., graphs with undirected and directed edges. Its vertices represent vector spaces, its undirected edges represent forms, and its directed edges represent linear mappings.

In particular, each pair $(\mathcal{A}, \mathcal{F})$ from Theorem 1 defines the representation

$$\mathcal{F}(\mathcal{A}u, v) = \mathcal{F}(u, \mathcal{A}^{r}v) \text{ if } r \ge 2,$$

$$\mathcal{F}(\mathcal{A}u, \mathcal{A}^{-r}v) = \mathcal{F}(u, v) \text{ if } r \le -2,$$

$$\mathcal{F}(u, v) = \varepsilon \widetilde{\mathcal{F}(v, u)} \text{ is nonsingular}$$
(8)

of the mixed graph $\bigcirc \bullet \bigcirc$ over \mathbb{F} with involution $a \mapsto \tilde{a}$, in which $\varepsilon \coloneqq 1$ if \mathcal{F} is symmetric or Hermitian, and $\varepsilon \coloneqq -1$ if \mathcal{F} is skew-symmetric or skew-Hermitian. Choosing a basis in V, we give (8) by its matrices

$$A \bigcap n \bigcap F \qquad A^{\star}F = FA^{r}, \quad F^{\star} = \varepsilon F \text{ is nonsingular}, \tag{9}$$

in which $n \coloneqq \dim V$, A and F are $n \times n$ matrices over \mathbb{F} , and $A^{\star} \coloneqq \widetilde{A}^T$ ($A^{\star} = A^T$ if $a \mapsto \widetilde{a}$ is the identity involution, and $A^{\star} = A^*$ otherwise). Changing the basis in V, we can reduce (A, F) by transformations

$$(A, F) \mapsto (S^{-1}AS, S^{\star}FS), \qquad S \text{ is nonsingular}$$
(10)

(see (2) and (4)). We say that the pairs (A, F) and $(S^{-1}AS, S^{\star}FS)$ are isomorphic via S.

Replacing $\mathcal{F}: V \times V \to \mathbb{F}$ in (8) by the pair of mutually adjoint linear mappings $\mathcal{F}: v \mapsto \mathcal{F}(?, v)$ and $\mathcal{F}^{\scriptscriptstyle \wedge}: u \mapsto \widetilde{\mathcal{F}(u, ?)}$, we obtain the system of linear mappings

$$\mathcal{A} \bigcap V \underbrace{\overset{\mathcal{F}}{\longrightarrow}}_{\mathcal{F}^{\lambda}} V^{\lambda} \bigcirc \mathcal{A}^{\lambda} \quad \mathcal{A}^{\lambda} \mathcal{F} = \mathcal{F} \mathcal{A}^{r}, \quad \mathcal{F}^{\lambda} = \varepsilon \mathcal{F} \text{ is nonsingular, (11)}$$

in which V^{\star} is the $^{\star}dual \ space$ (with respect to the involution $a \mapsto \tilde{a}$) consisting of semilinear forms on V, and $\mathcal{A}^{\star}: V^{\star} \to V^{\star}$ is the $^{\star}dual \ mapping$ defined by $\varphi \mapsto \varphi \mathcal{A}$. In the matrix form,

$$A \bigcap n \underbrace{\overset{F}{\underset{F^{\star}}{\longrightarrow}}} n \bigcap A^{\star} \quad A^{\star}F = FA^{r}, \quad F^{\star} = \varepsilon F \text{ is nonsingular.}$$
(12)

Thus, there is the bijective correspondence

$$A \bigcap n \bigcap F \qquad \mapsto \qquad A \bigcap n \underbrace{\overset{F}{\underset{F^{\star}}{\longrightarrow}}} n \bigcap A^{\star} \qquad (13)$$

between the matrix sets of systems (8) and (11).

Let us consider a system of linear mappings over \mathbb{F} :

$$\mathcal{M}: \ \mathcal{A}_1 \bigcap V_1 \underbrace{\overset{\mathcal{F}_1}{\longrightarrow}}_{\mathcal{F}_2} V_2 \bigcirc \mathcal{A}_2 \qquad \begin{array}{c} \mathcal{A}_2 \mathcal{F}_1 = \mathcal{F}_1 \mathcal{A}_1^r, \ \mathcal{A}_2^r \mathcal{F}_2 = \mathcal{F}_2 \mathcal{A}_1, \\ \mathcal{F}_1 = \varepsilon \mathcal{F}_2 \text{ is nonsingular,} \end{array}$$
(14)

$$M: A_1 \bigcap n \underbrace{\stackrel{F_1}{\longrightarrow}}_{F_2} n \bigcap A_2 \qquad \begin{array}{c} A_2F_1 = F_1A_1^r, \ A_2^rF_2 = F_2A_1, \\ F_1 = \varepsilon F_2 \text{ is nonsingular.} \end{array}$$
(15)

Changing bases in V_1 and V_2 , we can reduce (15) by transformations

$$M': RA_1R^{-1} \bigcap n \underbrace{\overset{SF_1R^{-1}}{\longrightarrow}}_{SF_2R^{-1}} n \bigcirc SA_2S^{-1} \qquad R, S \text{ are nonsingular.} (16)$$

We say that the matrix sets (15) and (16) are *isomorphic via* R and S and write $M \simeq M'$. This isomorphism can be shown by the commutative diagram

$$n \xrightarrow{A_{1}} n \xrightarrow{F_{1}} n \xrightarrow{A_{2}} n$$

$$R \downarrow \qquad RA_{1}R^{-1} \downarrow R \downarrow \qquad SF_{1}R^{-1} \downarrow S \qquad SA_{2}S^{-1} \downarrow S$$

$$n \xrightarrow{RA_{1}R^{-1}} n \xrightarrow{R} n \xrightarrow{SF_{2}R^{-1}} n \xrightarrow{R} n \xrightarrow{SA_{2}S^{-1}} n$$

The direct sum of systems M and M' is the system

$$M \oplus M': \quad A_1 \oplus A'_1 \bigcirc n + n' \underbrace{\xrightarrow{F_1 \oplus F'_1}}_{F_2 \oplus F'_2} n + n' \bigcirc A_2 \oplus A'_2 \tag{17}$$

A system is *indecomposable* if it is not isomorphic to a direct sum of the form (17) with nonzero n and n'.

For each system (15), we define the *dual system*

$$M^{\circ}: \quad A_{2}^{\scriptscriptstyle \wedge} \bigcap n \underbrace{\overset{F_{2}^{\scriptscriptstyle \wedge}}{\underset{F_{1}^{\scriptscriptstyle \wedge}}{\longrightarrow}}} n \bigcap A_{1}^{\scriptscriptstyle \wedge}$$

- A system M is *selfdual* if $M = M^{\circ}$, which means that it has the form (12). Suppose we know the following sets of systems of the form (15):
- $\mathfrak{M}_{\varepsilon}(\mathbb{F})$ which is a set of nonisomorphic indecomposable systems such that every indecomposable system (15) is isomorphic to exactly one system from $\mathfrak{M}_{\varepsilon}(\mathbb{F})$,
- $\mathfrak{M}'_{\varepsilon}(\mathbb{F})$ which is a set of nonisomorphic indecomposable selfdual systems such that every indecomposable selfdual system is isomorphic to exactly one system from $\mathfrak{M}'_{\varepsilon}(\mathbb{F})$,
- $\mathfrak{M}_{\varepsilon}^{\prime\prime}(\mathbb{F})$ which is a set of nonisomorphic indecomposable systems that are not isomorphic to selfdual such that every indecomposable system that is not isomorphic to selfdual is isomorphic to exactly one system from $\mathfrak{M}_{\varepsilon}^{\prime\prime}(\mathbb{F})$.

For each $M \in \mathfrak{M}'_{\varepsilon}(\mathbb{F})$ of the form (12), we define the matrix pairs

$$\mathring{M}: A \bigcap n \bigcap F \qquad \qquad \mathring{M}^{-}: A \bigcap n \bigcap -F \qquad (18)$$

and for each $N \in \mathfrak{M}_{\varepsilon}^{\prime\prime}(\mathbb{F})$ of the form (15), we define the matrix pair

$$N^{+}: \begin{bmatrix} A_{1} & 0\\ 0 & A_{2}^{\star} \end{bmatrix} \bigcirc 2n \bigcirc \begin{bmatrix} 0 & F_{2}^{\star}\\ F_{1} & 0 \end{bmatrix}.$$

Note that the natural bijection (13) takes \mathring{M} into M, and N^+ into a selfdual system that is isomorphic to $N \oplus N^\circ$.

The following lemma reduces the problem of classifying matrix pairs (9) up to transformations (10) to the problem of classifying systems (15) up to transformations (16). This lemma is a special case of [24, Theorem 3.2] about arbitrary systems of linear mappings and forms.

Lemma 1. Each pair (9) over $\mathbb{F} \in {\mathbb{C}, \mathbb{R}, \mathbb{H}}$ is isomorphic to a direct sum of pairs of the types

$$N^{+} \quad and \quad \begin{cases} \mathring{M} & \text{if } \mathring{M}^{-} \text{ and } \mathring{M} \text{ are isomorphic,} \\ \mathring{M}, \ \mathring{M}^{-} & \text{if } \mathring{M}^{-} \text{ and } \mathring{M} \text{ are not isomorphic,} \end{cases}$$

in which $M \in \mathfrak{M}'_{\varepsilon}(\mathbb{F})$ and $N \in \mathfrak{M}''_{\varepsilon}(\mathbb{F})$. This sum is uniquely determined, up to permutations of direct summands and replacements of $N \in \mathfrak{M}''_{\varepsilon}(\mathbb{F})$ by N° .

Lemma 2. If a system of the form (15) is isomorphic to a selfdual system via R and S, then it is isomorphic to some selfdual system via I and $R^{\star}S$.

Proof. The corresponding selfdual system is constructed as follows:

$$A_{1} \bigcap_{R} n \xrightarrow{F_{1}} n \bigcap_{A_{2}} A_{2}$$

$$B \bigcap_{R^{-1}} n \xrightarrow{G^{\wedge}} n \bigcap_{B^{\wedge}} B^{\wedge} - \text{selfdual}$$

$$R^{-1}BR \bigcap_{R^{\wedge}} n \xrightarrow{R^{\wedge}G^{\wedge}R} n \bigcap_{R^{\wedge}B^{\wedge}(R^{\wedge})^{-1}} - \text{selfdual}$$

3. Proof of Theorem 1

Lemma 3. Each square matrix over \mathbb{C} , \mathbb{R} , and \mathbb{H} is similar to a direct sum, uniquely determined up to permutations of summands, of matrices from the following matrix sets:

(a)
$$\mathfrak{C}(\mathbb{C}) \coloneqq \{J_n(\lambda) \mid \lambda \in \mathbb{C}\}, in which$$

$$J_n(\lambda) \coloneqq \begin{bmatrix} \lambda & 1 & 0 \\ \lambda & \ddots & \\ & \ddots & 1 \\ 0 & & \lambda \end{bmatrix} (n \text{-} by \text{-} n).$$

(b) $\mathfrak{C}(\mathbb{R}) \coloneqq \{J_n(a) \mid a \in \mathbb{R}\} \cup \{J_n(\lambda)^{\mathbb{R}} \mid \lambda \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}\}, in which$

$$J_n(\lambda)^{\mathbb{R}} \coloneqq \begin{bmatrix} \lambda^{\mathbb{R}} & I_2 & 0 \\ \lambda^{\mathbb{R}} & \ddots & \\ & \ddots & I_2 \\ 0 & & \lambda^{\mathbb{R}} \end{bmatrix} (2n - by - 2n),$$

 $\lambda^{\mathbb{R}}$ is determined in (5), and $\mathbb{C}^{\ddagger} = \{\lambda \in \mathbb{C} \mid \lambda \searrow \overline{\lambda}\}.$

(c) $\mathfrak{C}(\mathbb{H}) \coloneqq \{J_n(\lambda) \mid \lambda \in \mathbb{C}^{\ddagger}\}.$

Proof. The statement (a) is the Jordan theorem; (b) and (c) are given in [8, Theorem 3.4.1.5] and [21, Theorem 5.5.3].

Each system (15) is reduced by transformations (16) with $R = F_1$ and $S = I_n$ to a system of the form

$$M_{\varepsilon}(A): \quad A \bigcap n \underbrace{I_n}_{\varepsilon I_n} n \bigcap A^r \qquad A^{r^2} = A,$$

whose dual system is

$$M_{\varepsilon}(A)^{\circ}: \quad (A^{r})^{\star} \bigcap n \xrightarrow{\varepsilon I_{n}} n \bigcap A^{\star}$$

Clearly, $M_{\varepsilon}(A) \simeq M_{\varepsilon}(B)$ if and only if A and B are similar; and so

$$\mathfrak{M}_{\varepsilon}(\mathbb{F}) = \{ M_{\varepsilon}(A) \, | \, A \in \mathfrak{C}(\mathbb{F}) \text{ such that } A^{r^2} = A \};$$
(19)

$$M_{\varepsilon}(B) \simeq M_{\varepsilon}(A)^{\circ} \iff B \text{ is similar to } (A^r)^{\perp}.$$
 (20)

3.1. Case (A): $\mathbb{F} = \mathbb{C}$

Let $a \mapsto \tilde{a}$ be the identity involution or complex conjugation. Suppose that $J_n(\lambda)^{r^2} = J_n(\lambda)$ with $\lambda \in \mathbb{C}$. If $\lambda = 0$, then n = 1 since $r^2 \ge 4$. If $\lambda \ne 0$, then $J_n(\lambda)^{r^{2}-1} = I_n$, all entries of the first over-diagonal of $J_n(\lambda)^{r^{2}-1}$ are $(r^2 - 1)\lambda^{r^2-2}$, and so n = 1 too. Thus,

$$J_n(\lambda)^{r^2} = J_n(\lambda) \implies n = 1$$
(21)

and

$$\mathfrak{M}_{\varepsilon}(\mathbb{C}) = \{ M_{\varepsilon}(\lambda) \, | \, \lambda \in \mathbb{C}, \ \lambda^{r^2} = \lambda \}$$

(to simplify notation, we write λ instead of λI_1).

Since

$$M_{\varepsilon}(\lambda): \ \lambda \bigcap 1 \underbrace{1}_{\varepsilon} 1 \bigcap \lambda^{r} \qquad M_{\varepsilon}(\lambda)^{\circ}: \ \tilde{\lambda}^{r} \bigcap 1 \underbrace{1}_{1} \bigcap \tilde{\lambda}$$

and $\lambda^{r^2} = \lambda$, we have $M_{\varepsilon}(\lambda)^{\circ} \simeq M_{\varepsilon}(\widetilde{\lambda}^r)$. Hence,

$$M_{\varepsilon}(\mu) \simeq M_{\varepsilon}(\lambda)^{\circ} \iff \mu = \widetilde{\lambda}^r.$$
 (22)

The following cases are possible:

(a₁): $\varepsilon = 1$ and the involution on \mathbb{C} is the identity. Then

$$M_1(\lambda) \simeq M_1(\lambda)^\circ \iff M_1(\lambda) = M_1(\lambda)^\circ \iff \lambda = \lambda^r,$$

and so

$$\mathfrak{M}_{1}'(\mathbb{C}) = \{ M_{1}(\lambda) | \lambda \in \mathbb{C}, \ \lambda^{r} = \lambda \}, \\ \mathfrak{M}_{1}''(\mathbb{C}) = \{ M_{1}(\mu) | \mu \in \mathbb{C}, \ \mu^{r^{2}} = \mu, \ \mu^{r} \neq \mu \}$$

Lemma 1 and (22) ensure (a_1) .

(a₂): $\varepsilon = 1$ and the involution on \mathbb{C} is complex conjugation. Then

$$M_1(\lambda) \simeq M_1(\lambda)^\circ \iff M_1(\lambda) = M_1(\lambda)^\circ \iff \lambda = \overline{\lambda}^r,$$

and so

$$\mathfrak{M}_{1}^{\prime}(\mathbb{C}) = \{ M_{1}(\lambda) \, | \, \lambda \in \mathbb{C}, \ \lambda^{r} = \bar{\lambda} \}, \\ \mathfrak{M}_{1}^{\prime\prime}(\mathbb{C}) = \{ M_{1}(\mu) \, | \, \mu \in \mathbb{C}, \ \mu^{r^{2}} = \mu, \ \mu^{r} \neq \bar{\mu} \}.$$

Lemma 1 and (22) ensure (a_2) .

(a₃): $\varepsilon = -1$ and the involution on \mathbb{C} is the identity. The system $M_{-1}(\lambda)$ is not isomorphic to a selfdual system since there are no nonsingular 1×1 matrices R and S such that $SI_1R^{-1} = S(-I_1)R^{-1}$ (see (16)). Therefore,

$$\mathfrak{M}_{-1}'(\mathbb{C}) = \emptyset,$$

$$\mathfrak{M}_{-1}''(\mathbb{C}) = \{ M_{-1}(\lambda) \, | \, \lambda \in \mathbb{C}, \ \lambda^{r^2} = \lambda \}.$$

Lemma 1 and (22) ensure (a_3) .

3.2. Case (B): $\mathbb{F} = \mathbb{R}$

The set $\mathfrak{C}(\mathbb{R})$ is given in Lemma 3(b). The equality

$$\mathfrak{M}_{\varepsilon}(\mathbb{R}) = \{ M_{\varepsilon}(0), \ M_{\varepsilon}(1) \} \cup \{ M_{\varepsilon}(-1) | \text{ if } r \text{ is odd} \} \\ \cup \{ M_{\varepsilon}(\lambda^{\mathbb{R}}) | \lambda \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \lambda^{r^{2}} = \lambda \}$$

$$(23)$$

is proved as follows:

- Consider J_n(a) ∈ 𝔅(ℝ) with a ∈ ℝ and J_n(a)^{r²} = J_n(a). By (21), n = 1. Since a is real, a^{r²} = a implies that either a = 0, or a = ±1 if r is odd and a = 1 if r is even. Note that
 each system M_ε(λ) with λ ∈ {0, 1, -1 (if r is odd)} is selfdual if ε = 1; it is not isomorphic to selfdual if ε = -1.
- Consider $J_n(\lambda)^{\mathbb{R}} \in \mathfrak{C}(\mathbb{R})$ with $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $(J_n(\lambda)^{\mathbb{R}})^{r^2} = J_n(\lambda)^{\mathbb{R}}$. The matrix $J_n(\lambda)^{\mathbb{R}}$ is similar over \mathbb{C} to $J_n(\lambda) \oplus J_n(\overline{\lambda})$. Hence $J_n(\lambda)^{r^2} = J_n(\lambda)$, n = 1, and $\lambda^{r^2} = \lambda$.

For every $M_{\varepsilon}(\lambda^{\mathbb{R}}), M_{\varepsilon}(\mu^{\mathbb{R}}) \in \mathfrak{M}_{\varepsilon}(\mathbb{R})$, we have

$$M_{\varepsilon}(\mu^{\mathbb{R}}) \simeq M_{\varepsilon}(\lambda^{\mathbb{R}})^{\circ} \qquad \Longleftrightarrow \qquad \mu = \lambda^r \text{ or } \mu = \bar{\lambda}^r$$
 (25)

since $M_{\varepsilon}(\mu^{\mathbb{R}}) \simeq M_{\varepsilon}(\lambda^{\mathbb{R}})^{\circ}$ if and only if $\mu^{\mathbb{R}}$ is similar to $((\lambda^{\mathbb{R}})^r)^T$, if and only if diag $(\mu, \bar{\mu})$ is similar to diag $(\lambda, \bar{\lambda})^r$, if and only if $\mu = \lambda^r$ or $\mu = \bar{\lambda}^r$.

Thus, if $M_{\varepsilon}(\lambda^{\mathbb{R}})$ is isomorphic to a selfdual system, then $\lambda^r = \lambda$ or $\lambda^r = \overline{\lambda}$. Write $\lambda = a + bi$ $(a, b \in \mathbb{R}, b \neq 0)$, then $\lambda^{\mathbb{R}} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ and

$$\begin{aligned} \lambda^r &= \lambda &\implies \begin{bmatrix} a & -b \\ b & a \end{bmatrix}^r &= \begin{bmatrix} a & -b \\ b & a \end{bmatrix}, \\ \lambda^r &= \bar{\lambda} &\implies \begin{bmatrix} a & -b \\ b & a \end{bmatrix}^r &= \begin{bmatrix} a & b \\ -b & a \end{bmatrix}. \end{aligned}$$

The following two cases are possible:

(b₁): $\varepsilon = 1$. Let $\lambda = a + bi$ $(a, b \in \mathbb{R}, b \neq 0)$. If $M_1(\lambda^{\mathbb{R}})$ is isomorphic to a selfdual system, then $\lambda^r = \lambda$ or $\lambda^r = \overline{\lambda}$. If $\lambda^r = \lambda$, then $\begin{bmatrix} a & -b \\ b & a \end{bmatrix} Z = Z \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$ with $Z := \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, and so $M_1(\lambda^{\mathbb{R}})$ is isomorphic to a selfdual system:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \bigcirc 2 \xrightarrow{I_2} 2 \bigcirc \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$
$$\downarrow_{I_2} \downarrow \qquad \qquad \downarrow_{Z} \downarrow$$
$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \bigcirc 2 \xrightarrow{Z} 2 \bigcirc \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$$

If $\lambda^r = \overline{\lambda}$, then the system $M_1(\lambda^{\mathbb{R}})$ is selfdual. Thus,

$$\mathfrak{M}_{1}^{\prime}(\mathbb{R}) = \{M_{1}(0), \ M_{1}(1)\} \cup \{M_{1}(-1) | \text{ if } r \text{ is odd}\} \\ \cup \{\lambda^{\mathbb{R}} \bigcap 2 \xrightarrow{Z} 2 \bigcap \bar{\lambda}^{\mathbb{R}} | \lambda \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \lambda^{r} = \lambda\} \\ \cup \{M_{1}(\mu^{\mathbb{R}}) | \mu \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \mu^{r} = \bar{\mu}\}, \\ \mathfrak{M}_{1}^{\prime\prime}(\mathbb{R}) = \{M_{1}(\nu^{\mathbb{R}}) | \nu \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \nu^{r^{2}} = \nu, \ \nu^{r} \neq \nu, \ \nu^{r} \neq \bar{\nu}\}.$$

Each system $\lambda^{\mathbb{R}} \bigcap 2 \xrightarrow{Z} 2 \bigcap \overline{\lambda}^{\mathbb{R}}$ from $\mathfrak{M}'_1(\mathbb{R})$ defines the pairs $(\lambda^{\mathbb{R}}, Z)$ and $(\lambda^{\mathbb{R}}, -Z)$ of the form (18); they are isomorphic via $S = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ (see (10)). Each system $M_1(\mu^{\mathbb{R}}) \in \mathfrak{M}'_1(\mathbb{R})$ defines the pairs $(\mu^{\mathbb{R}}, I_2)$ and $(\mu^{\mathbb{R}}, -I_2)$; they are not isomorphic since I_2 and $-I_2$ are not congruent over \mathbb{R} .

Lemma 1 and (25) ensure (b₁) (we do not write $\nu \supset \bar{\nu}^r$ since ν is determined up to replacement by $\bar{\nu}$).

(b₂): $\varepsilon = -1$. Let $\lambda = a + bi$ $(a, b \in \mathbb{R}, b \neq 0)$. If $M_{-1}(\lambda^{\mathbb{R}})$ is isomorphic to a selfdual system, then $\lambda^r = \lambda$ or $\lambda^r = \overline{\lambda}$. If $\lambda^r = \lambda$, then $M_{-1}(\lambda^{\mathbb{R}})$ is not isomorphic to a selfdual system; otherwise by Lemma 2 there is a nonsingular P such that

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \bigcirc 2 \xrightarrow{I_2} 2 \bigcirc \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$
$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \bigcirc 2 \xrightarrow{P} 2 \bigcirc P \begin{bmatrix} a & -b \\ b & a \end{bmatrix} P^{-1} = \begin{bmatrix} a & b \\ -P = P^T \end{bmatrix} 2 \bigcirc P \begin{bmatrix} a & -b \\ b & a \end{bmatrix} P^{-1} = \begin{bmatrix} a & b \\ -P = P^T \end{bmatrix}$$

Then $P = \begin{bmatrix} 0 & -x \\ x & 0 \end{bmatrix}$ for some $x \neq 0$ since $P^T = -P$. The equality $P \begin{bmatrix} a & -b \\ b & a \end{bmatrix} = \begin{bmatrix} a & b \\ -b & a \end{bmatrix} P$ implies that b = 0, which contradicts our assumption that $\lambda \notin \mathbb{R}$. If $\lambda^r = \overline{\lambda}$, then $\begin{bmatrix} a & b \\ -b & a \end{bmatrix} L = L \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$ with $L := \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, and so $M_{-1}(\lambda^{\mathbb{R}})$ is isomorphic to a selfdual system:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \bigcirc 2 \xrightarrow{I_2} 2 \bigcirc \begin{bmatrix} a & b \\ -I_2 & \downarrow L \\ I_2 & \downarrow L \\ \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \bigcirc 2 \xrightarrow{L} 2 \bigcirc \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$$

Using (24), we obtain

$$\mathfrak{M}_{-1}'(\mathbb{R}) = \left\{ \lambda^{\mathbb{R}} \bigcap 2 \xrightarrow{L} 2 \bigcap \bar{\lambda}^{\mathbb{R}} \mid \lambda \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \lambda^{r} = \bar{\lambda} \right\},$$
$$\mathfrak{M}_{-1}''(\mathbb{R}) = \left\{ M_{-1}(0), \ M_{-1}(1) \right\} \cup \left\{ M_{1}(-1) \mid \text{if } r \text{ is odd} \right\}$$
$$\cup \left\{ M_{-1}(\mu^{\mathbb{R}}) \mid \mu \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \mu^{r^{2}} = \mu, \ \mu^{r} \neq \bar{\mu} \right\}.$$

Each system from $\mathfrak{M}'_{-1}(\mathbb{R})$ defines the pairs $(\lambda^{\mathbb{R}}, L)$ and $(\lambda^{\mathbb{R}}, -L)$; they are isomorphic via $S = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ (see (10) and (18)). Lemma 1 and (25) ensure (b₂). 3.3. Case (C): $\mathbb{F} = \mathbb{H}$.

In this case, $\varepsilon = \pm 1$, $a \mapsto \tilde{a}$ is quaternion conjugation (6) or quaternion semiconjugation (7), and $\mathfrak{C}(\mathbb{H})$ is given in Lemma 3(c). By (19) and (21),

$$\mathfrak{M}_{\varepsilon}(\mathbb{H}) = \{ M_{\varepsilon}(\lambda) \, | \, \lambda \in \mathbb{C}^{\updownarrow}, \ \lambda^{r^2} = \lambda \}$$

If $M_{\varepsilon}(\lambda), M_{\varepsilon}(\mu) \in \mathfrak{M}_{\varepsilon}(\mathbb{H})$, then

$$M_{\varepsilon}(\mu) \simeq M_{\varepsilon}(\lambda)^{\circ} \quad \iff \quad \mu = \lambda^r \text{ or } \mu = \bar{\lambda}^r.$$
 (26)

Indeed, if $M_{\varepsilon}(\mu) \simeq M_{\varepsilon}(\lambda)^{\circ}$, then the 1 × 1 matrix $[\mu]$ is similar to $[\lambda^{r}]^{\star} = [\bar{\lambda}^{r}]$. By Lemma 3(c), $\mu = \lambda^{r}$ or $\mu = \bar{\lambda}^{r}$. Conversely, let $\mu = \lambda^{r}$ or $\mu = \bar{\lambda}^{r}$. We can take $\mu = \bar{\lambda}^{r}$ since λ is determined up to replacement by $\bar{\lambda}$. Then $M_{\varepsilon}(\mu) \simeq M_{\varepsilon}(\lambda)^{\circ}$ since



If $M_{\varepsilon}(\lambda) \in \mathfrak{M}_{\varepsilon}(\mathbb{H})$ is isomorphic to a selfdual system, then by (26) $\lambda^r = \lambda$ or $\lambda^r = \overline{\lambda}$. Conversely,

• if $\lambda^r = \overline{\lambda}$, then $M_{\varepsilon}(\lambda)$ is isomorphic to a selfdual system:

$$\lambda \bigcap_{1} 1 \xrightarrow{\varepsilon} 1 \xrightarrow{\varepsilon} 1 \xrightarrow{\varepsilon} 1 \xrightarrow{\varepsilon} \lambda^{r} = \overline{\lambda}$$

$$\lambda \bigcap_{1} 1 \xrightarrow{\delta_{\varepsilon}} 1 \xrightarrow{\delta_{\varepsilon}} 1 \xrightarrow{\overline{\lambda}} \overline{\lambda}$$

$$\delta_{\varepsilon} := \begin{cases} 1 & \text{if } \varepsilon = 1, \\ i & \text{if } \varepsilon = -1; \end{cases}$$

$$(27)$$

• if $\lambda^r = \lambda$ and $\lambda \notin \mathbb{R}$ (the case $\lambda \in \mathbb{R}$ is considered in the previous paragraph), then $M_{\varepsilon}(\lambda) \in \mathfrak{M}_{\varepsilon}(\mathbb{H})$ is isomorphic to a selfdual system if and only if either $\varepsilon = 1$ and the involution is (7), or $\varepsilon = -1$ and the involution is (6). Indeed, suppose that $M_{\varepsilon}(\lambda)$ is isomorphic to a selfdual system. By Lemma 2, there exists $h \in \mathbb{H}$ such that

$$\lambda \bigcap_{1} 1 \xrightarrow{1} \varepsilon \qquad 1 \bigcap_{h} \lambda^{r} = \lambda$$

$$\lambda \bigcap_{1} 1 \xrightarrow{h} 1 \bigcap_{\overline{h}} \overline{\lambda}$$
(28)

is an isomorphism. If either $\varepsilon = 1$ and the involution is (7), or $\varepsilon = -1$ and the involution is (6), then (28) holds for h = j. If $\varepsilon = 1$ and the involution is (6), then (28) implies $h = \overline{h}, h \in \mathbb{R}, h\lambda = \overline{\lambda}h$, and so $\lambda \in \mathbb{R}$, a contradiction. If $\varepsilon = -1$ and the involution is (7), then (28) implies $-h = \widehat{h}, h \in \mathbb{R}i, h\lambda = \overline{\lambda}h$, and so $\lambda \in \mathbb{R}$, a contradiction.

The following cases are possible:

(c₁): $\varepsilon = 1$ and the involution is quaternion conjugation (6). Then

$$\mathfrak{M}_{1}^{\prime}(\mathbb{H}) = \left\{ \lambda \bigcap 1 \xrightarrow{1} 1 \bigcap_{\overline{\lambda}} |\lambda \in \mathbb{C}^{\ddagger}, \ \lambda^{r} = \overline{\lambda} \right\},$$
$$\mathfrak{M}_{1}^{\prime\prime}(\mathbb{H}) = \left\{ M_{1}(\mu) | \mu \in \mathbb{C}^{\ddagger}, \ \mu^{r^{2}} = \mu, \ \mu^{r} \neq \overline{\mu} \right\}.$$

Each system from $\mathfrak{M}'_1(\mathbb{H})$ defines the pairs $(\lambda, 1)$ and $(\lambda, -1)$; they are not isomorphic since $\bar{c} \, 1c \neq -1$ for all $c \in \mathbb{H}$.

Lemma 1 and (22) ensure (c_1) .

(c₂): $\varepsilon = 1$ and the involution is quaternion semiconjugation (7). Then

$$\mathfrak{M}_{1}'(\mathbb{H}) = \left\{ \lambda \bigcap 1 \xrightarrow{1}_{1} 1 \bigcap_{\overline{\lambda}} \left| \lambda \in \mathbb{C}^{\ddagger}, \ \lambda^{r} = \overline{\lambda} \right\} \\ \cup \left\{ \mu \bigcap 1 \xrightarrow{j}_{j} 1 \bigcap_{\overline{\mu}} \left| \mu \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \mu^{r} = \mu \right\}, \\ \mathfrak{M}_{1}''(\mathbb{H}) = \left\{ M_{1}(\nu) \left| \nu \in \mathbb{C}^{\ddagger}, \ \nu^{r^{2}} = \nu, \ \nu^{r} \neq \nu, \ \nu^{r} \neq \overline{\nu} \right\}.$$

Each system $\lambda \bigcap 1 \xrightarrow{1}{\longrightarrow} 1 \bigcap_{\overline{\lambda}} 1 \longrightarrow \overline{\lambda}$ from $\mathfrak{M}'_1(\mathbb{H})$ defines the pairs $(\lambda, 1)$ and $(\lambda, -1)$.

- If $\lambda \notin \mathbb{R}$, then $(\lambda, 1)$ and $(\lambda, -1)$ are not isomorphic. On the contrary, suppose that there is a nonzero $c \in \mathbb{H}$ such that

$$c^{-1}\lambda c = \lambda, \qquad \widehat{c} \, 1c = -1 \tag{29}$$

(see (10)). By $c^{-1}\lambda c = \lambda$, we have $c \in \mathbb{C}$, which contradicts $\widehat{c} \, 1c = -1$.

- If $\lambda \in \mathbb{R}$, then $(\lambda, 1)$ and $(\lambda, -1)$ are isomorphic since (29) holds for c = j.

The pairs (μ, j) and $(\mu, -j)$ constructed by $\mu \bigcap 1 \xrightarrow{j}{j} 1 \bigcap \overline{\mu}$ from $\mathfrak{M}'_1(\mathbb{H})$ are isomorphic via *i* since $i\mu = \mu i$ and $\hat{\imath} j i = -iji = -ki = -j$. Lemma 1 and (22) ensure (c₂).

(c₃): $\varepsilon = -1$ and the involution is quaternion conjugation (6). Then

$$\mathfrak{M}_{-1}'(\mathbb{H}) = \left\{ \lambda \bigcap 1 \xrightarrow{i}_{-i} 1 \bigcap \overline{\lambda} \mid \lambda \in \mathbb{C}^{\ddagger}, \ \lambda^{r} = \overline{\lambda} \right\}$$
$$\cup \left\{ \mu \bigcap 1 \xrightarrow{j}_{-j} 1 \bigcap \overline{\mu} \mid \mu \in \mathbb{C}^{\ddagger} \setminus \mathbb{R}, \ \mu^{r} = \mu \right\},$$
$$\mathfrak{M}_{-1}''(\mathbb{H}) = \left\{ M_{-1}(\nu) \mid \nu \in \mathbb{C}^{\ddagger}, \ \nu^{r^{2}} = \nu, \ \nu^{r} \neq \nu, \ \nu^{r} \neq \overline{\nu} \right\}.$$

If $\lambda \notin \mathbb{R}$, then the pairs (λ, i) and $(\lambda, -i)$ constructed by a system from $\mathfrak{M}'_{-1}(\mathbb{H})$ are not isomorphic. On the contrary, suppose there exists $c \in \mathbb{H}$ such that

$$c^{-1}\lambda c = \lambda, \qquad \bar{c}\,ic = -i \tag{30}$$

(see (10)). Since $\lambda = \lambda_1 + \lambda_2 i$ with $\lambda_1, \lambda_2 \in \mathbb{R}$ and $\lambda_2 \neq 0$, the equality $c^{-1}\lambda c = \lambda$ implies that ic = ci, and so $c \in \mathbb{C}$, which contradicts $\bar{c}ic = -i$. If $\lambda \in \mathbb{R}$, then (λ, i) and $(\lambda, -i)$ are isomorphic since (30) holds for c = j. The pairs (μ, j) and $(\mu, -j)$ are isomorphic via i since $i\mu = \mu i$ and $\bar{i}ji = -iji = -ki = -j$.

Lemma 1 and (22) ensure (c_3) .

(c₄): $\varepsilon = -1$ and the involution is quaternion semiconjugation (7). Then

$$\mathfrak{M}_{-1}'(\mathbb{H}) = \left\{ \lambda \bigcap 1 \xrightarrow{i}_{-i} 1 \bigcap_{\overline{\lambda}} \left| \lambda \in \mathbb{C}^{\ddagger}, \ \lambda^{r} = \overline{\lambda} \right\}, \\ \mathfrak{M}_{-1}''(\mathbb{H}) = \left\{ M_{-1}(\mu) \left| \mu \in \mathbb{C}^{\ddagger}, \ \mu^{r^{2}} = \mu, \ \mu^{r} \neq \overline{\mu} \right\}.$$

The pairs (λ, i) and $(\lambda, -i)$ constructed by a system from $\mathfrak{M}'_{-1}(\mathbb{H})$ are not isomorphic since $\widehat{c}ic = i\overline{c}c \neq -i$ for all $c \in \mathbb{H}$.

Lemma 1 and (22) ensure (c_4) .

Acknowledgements

This paper is a result of a student seminar held at the University of São Paulo during the visit of V.V. Sergeichuk in 2019 and 2020; he is grateful to the university for hospitality and to the FAPESP for financial support (2018/24089-4).

References

- Y.H. Au-Yeung, C.K. Li, L. Rodman, *H*-unitary and Lorentz matrices: a review, SIAM J. Matrix Anal. Appl. 25 (2004) 1140–1162.
- [2] J.V. Caalim, V. Futorny, Y. Tanaka, V.V. Sergeichuk, Isometric and selfadjoint operators on a vector space with nondegenerate diagonalizable form, Linear Algebra Appl. 587 (2020) 92–110.
- [3] M. Catral, L. Lebtahi, J. Stuart, N. Thome, On a matrix group constructed from an $\{R, s+1, k\}$ -potent matrix, Linear Algebra Appl. 461 (2014) 200–201.
- [4] M. Catral, L. Lebtahi, J. Stuart, N. Thome, Matrices A such that $A^{s+1}R = RA^*$ with $R^k = I$, Linear Algebra Appl. 552 (2018) 85–104.
- [5] M. Catral, L. Lebtahi, J. Stuart, N. Thome, Spectral study of $\{R, s + 1, k\}$ and $\{R, s + 1, k, *\}$ -potent matrices, J. Comput. Appl. Math. 373 (2020) 112414.
- [6] I. Gohberg, P. Lancaster, L. Rodman, Matrices and Indefinite Scalar Products, Birkhäuser Verlag, Basel, 1983.
- [7] I. Gohberg, P. Lancaster, L. Rodman, Indefinite Linear Algebra and Applications, Birkhäuser, Boston, 2006.
- [8] R.A. Horn, C.R. Johnson, Matrix Analysis, 2nd ed., Cambridge University Press, Cambridge, 2013.
- [9] R.A. Horn, V.V. Sergeichuk, Canonical matrices of bilinear and sesquilinear forms, Linear Algebra Appl. 428 (2008) 193–223.
- [10] R.A. Horn, V.V. Sergeichuk, Representations of quivers and mixed graphs, Chapter 34 in: L. Hogben (Ed.), Handbook of Linear Algebra, 2nd ed., CRC Press, 2014.

- [11] P. Lancaster, L. Rodman, Algebraic Riccati Equations, The Clarendon Press, Oxford University Press, New York, 1995.
- [12] L. Lebtahi, Ó. Romero, N. Thome, Characterizations of $\{K, s+1\}$ -potent matrices and applications, Linear Algebra Appl. 436 (2012) 293–306.
- [13] L. Lebtahi, O. Romero, N. Thome, Relations between $\{K, s+1\}$ -potent matrices and different classes of complex matrices, Linear Algebra Appl. 438 (2013) 1517–1531.
- [14] L. Lebtahi, Ó. Romero, N. Thome, Algorithms for $\{K, s + 1\}$ -potent matrix constructions, J. Comput. Appl. Math. 249 (2013) 157–162.
- [15] L. Lebtahi, J. Stuart, N. Thome, J. Weaver, Matrices A such that $RA = A^{s+1}R$ when $R^k = I$, Linear Algebra Appl. 439 (2013) 1017–1023.
- [16] L. Lebtahi, N. Thome, Properties of a matrix group associated to a $\{K, S+1\}$ -potent matrix, Electron. J. Linear Algebra 24 (2012/13) 34–44.
- [17] C. Mehl, On classification of normal matrices in indefinite inner product spaces, Electron. J. Linear Algebra 15 (2006) 50–83.
- [18] C. Mehl, Essential decomposition of polynomially normal matrices in real indefinite inner product spaces, Electron. J. Linear Algebra, 15 (2006) 84–106.
- [19] J. Meleiro, V.V. Sergeichuk, T. Solovera, A. Zaidan, Classification of linear mappings between indefinite inner product spaces, Linear Algebra Appl. 531 (2017) 356–374.
- [20] J. Milnor, On isometries of inner product spaces, Invent. Math. 8 (1969) 83–97.
- [21] L. Rodman, Topics in Quaternion Linear Algebra, Princeton University Press, 2014.
- [22] V.V. Sergeichuk, Classification problems for systems of forms and linear mappings, Math. USSR-Izv. 31 (no. 3) (1988) 481–501. Theorem 2 is corrected in: arxiv.org/abs/0801.0823.

- [23] V.V. Sergeichuk, Linearization method in classification problems of linear algebra, São Paulo J. Math. Sci. 1 (2007) 219–240.
- [24] V.V. Sergeichuk, Canonical matrices of isometric operators on indefinite inner product spaces, Linear Algebra Appl. 428 (2008) 154–192.