

EXTENSIONS OF TORIC LINE BUNDLES

KLAUS ALTMANN, AMELIE FLATT, AND LUTZ HILLE

ABSTRACT. For any two nef line bundles $\mathcal{L}^+ := \mathcal{O}_X(\Delta^+)$ and $\mathcal{L}^- := \mathcal{O}_X(\Delta^-)$ on a toric variety X represented by lattice polyhedra Δ^+ respectively Δ^- , we present the universal equivariant extension of \mathcal{L}^- by \mathcal{L}^+ under use of the connected components of the set theoretic difference $\Delta^- \setminus \Delta^+$.

1. INTRODUCTION

1.1. Spotting Cohomology. Consider a projective toric variety $X = \mathbb{P}(\Delta)$ corresponding to a lattice polytope $\Delta \subseteq M_{\mathbb{R}}$, $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ for a lattice M , over \mathbb{C} . A torus invariant Cartier divisor D on X can be represented by a pair (Δ^+, Δ^-) of lattice polytopes, where both polytopes encode nef divisors D_{Δ^+} and D_{Δ^-} on X and $D = D_{\Delta^+} - D_{\Delta^-}$. Denote the associated line bundle by $\mathcal{O}_X(D) := \mathcal{O}_X(\Delta^+ - \Delta^-)$. It is well-known that the cohomology groups $H^i(X, \mathcal{O}_X(D))$ of the line bundle $\mathcal{O}_X(D)$ of a torus invariant Cartier divisor D are M -graded (compare for example [CLS11, §9.1] or [Ful93, Section 3.5]):

$$(1.1) \quad H^i(X, \mathcal{O}_X(D)) = \bigoplus_{m \in M} H^i(X, \mathcal{O}_X(D))_m.$$

By [ABKW20, Thm. III.6] or [AP20] we can describe their homogeneous component of degree $m \in M$ by

$$(1.2) \quad H^i(X, \mathcal{O}_X(\Delta^+ - \Delta^-))_m = \tilde{H}^{i-1}(\Delta^- \setminus (\Delta^+ - m), \mathbb{C}),$$


where $\tilde{H}^{i-1}(Z, \mathbb{C})$ on the right hand side denotes the reduced singular cohomology of a topological space Z with complex coefficients, $(\Delta^+ - m)$ denotes the polytope Δ^+ shifted by the lattice point $-m \in M$ and $\Delta^- \setminus (\Delta^+ - m)$ denotes the set-theoretic difference of the two polytopes. Recall the (-1) -st reduced singular cohomology:

$$(1.3) \quad \tilde{H}^{-1}(Z, \mathbb{C}) = \begin{cases} \mathbb{C}, & \text{if } Z = \emptyset, \\ 0, & \text{otherwise.} \end{cases}$$

The 0-th reduced cohomology is the quotient $\tilde{H}^0(Z, \mathbb{C}) = H^0(Z, \mathbb{C})/H^0(\{\cdot\}, \mathbb{C})$, and thus, its dimension is the number of connected components of Z minus 1.

1.2. The Example \mathbb{F}_1 . We look at the first Hirzebruch surface \mathbb{F}_1 as a first example. Consider the projections $p_1: \mathbb{F}_1 \rightarrow \mathbb{P}^1$ as a ruled surface and $p_2: \mathbb{F}_1 \rightarrow \mathbb{P}^2$ as a blow-up. We use $\mathcal{O}_{\mathbb{F}_1}(1, 0) := p_1^* \mathcal{O}_{\mathbb{P}^1}(1)$ and $\mathcal{O}_{\mathbb{F}_1}(0, 1) := p_2^* \mathcal{O}_{\mathbb{P}^2}(1)$ as a basis for $\text{Pic } \mathbb{F}_1 = \mathbb{Z}^2$. For $i, j \geq 0$ the sheaves $\mathcal{O}_{\mathbb{F}_1}(i, j)$ correspond to lattice polytopes $\Delta_{(i,j)}$

in \mathbb{R}^2 . For example, for $\mathcal{O}_{\mathbb{F}_1}(0, 0)$, $\mathcal{O}_{\mathbb{F}_1}(1, 0)$, $\mathcal{O}_{\mathbb{F}_1}(0, 1)$, $\mathcal{O}_{\mathbb{F}_1}(0, 2)$, and the ample $\mathcal{O}_{\mathbb{F}_1}(1, 1)$, the lattice polytopes look as follows:



The red dot indicates the position of the origin in each figure, fixing the exact position of each polytope within the plane. We see an example of the result quoted in subsection (1.1). Consider the polytope $\Delta_{(0,2)}$ shaded in yellow and the polytope $\Delta_{(1,0)} + (0, 1)$, that is, $\Delta_{(1,0)}$ shifted by $(0, 1) \in \mathbb{Z}^2$, depicted in orange:



The two connected components of the set-theoretic difference $\Delta_{(0,2)} \setminus (\Delta_{(1,0)} + (0, 1))$ provide a one-dimensional 0-th reduced singular cohomology

$$(1.4) \quad \tilde{H}^0(\Delta_{(0,2)} \setminus (\Delta_{(1,0)} + (0, 1))),$$

and so by [ABKW20] and [AP20] a one-dimensional piece (in fact, the only one) of

$$(1.5) \quad H^1(\mathbb{F}_1, \mathcal{O}_{\mathbb{F}_1}(\Delta_{(1,0)} - \Delta_{(0,2)})) = H^1(\mathbb{F}_1, \mathcal{O}_{\mathbb{F}_1}(1, -2)),$$

sitting in degree $m = -(0, 1) \in \mathbb{Z}^2$.

In this paper we will take another point of view. The partition of $\Delta_{(0,2)}$ into two connected components induces the following “exact sequences of polytopes”:

$$(1.6) \quad 0 \rightarrow \text{red dot} \rightarrow \boxed{\text{blue triangle} \oplus \text{orange triangle}} \rightarrow \text{yellow triangle} \rightarrow 0.$$

The two polytopes in the middle cover the polytope $\Delta_{(0,2)} - (0, 1)$ and intersect in the polytope $\Delta_{(1,0)}$, hence they give an inclusion/exclusion sequence of polytopes. In section 3 we show that this corresponds to an exact sequence of sheaves

$$(1.7) \quad 0 \rightarrow \mathcal{O}_{\mathbb{F}_1}(1, 0) \rightarrow \mathcal{O}_{\mathbb{F}_1}(0, 1) \oplus \mathcal{O}_{\mathbb{F}_1}(1, 1) \rightarrow \mathcal{O}_{\mathbb{F}_1}(0, 2) \rightarrow 0.$$

We obtain an extension of $\mathcal{O}_{\mathbb{F}_1}(0, 2)$ by $\mathcal{O}_{\mathbb{F}_1}(1, 0)$, that is, an element of the group

$$(1.8) \quad \text{Ext}^1(\mathcal{O}_{\mathbb{F}_1}(0, 2), \mathcal{O}_{\mathbb{F}_1}(1, 0)) = H^1(\mathbb{F}_1, \mathcal{O}_{\mathbb{F}_1}(1, -2))$$

$$(1.9) \quad = H^1(\mathbb{F}_1, \mathcal{O}_{\mathbb{F}_1}(\Delta_{(1,0)} - \Delta_{(0,2)})),$$

which we know to be one-dimensional by [ABKW20] and [AP20]. We will show that the short exact extension sequence (1.7) induced by the “exact sequences of polytopes” (1.6) represents this one-dimensional vector space $\text{Ext}^1(\mathcal{O}_{\mathbb{F}_1}(0, 2), \mathcal{O}_{\mathbb{F}_1}(1, 0))$ and, moreover, that this concept works in general.

1.3. Acknowledgements. We would like to thank Christian Haase for helpful discussions throughout the work on this paper, in particular for the triangulation argument in the proof of Proposition 13. Moreover, we thank the anonymous referee for the careful reading and many valuable hints and remarks. The second author was supported by the BMS ("Berlin Mathematical School") and the third one by the DFG and Exzellenzcluster "Mathematik Münster".¹

2. TORIC GEOMETRY

We introduce some basics of toric geometry central to this paper. Readers not familiar with toric geometry can take a look at one of the numerous introductory texts, for example [CLS11], [Ful93], [Dan78], or [Dem70].

Let $M \cong \mathbb{Z}^r$ be a lattice and $N = \text{Hom}_{\mathbb{Z}}(M, \mathbb{Z}) \cong \mathbb{Z}^r$ its dual lattice. There is a natural pairing $\langle \cdot, \cdot \rangle: M \times N \rightarrow \mathbb{Z}$. We consider the algebraic torus $T = \text{Spec } \mathbb{C}[M]$. The isomorphism $M \cong \mathbb{Z}^r$ induces an isomorphism $T \cong (\mathbb{C}^*)^r$. The lattice M can be recovered as the *character lattice* $\text{Hom}(T, \mathbb{C}^*)$. We denote the character of $M \ni m \mapsto (a_1, \dots, a_r) \in \mathbb{Z}^r$ by $\chi^m: T \rightarrow \mathbb{C}^*$, $(t_1, \dots, t_r) \mapsto t_1^{a_1} \cdots t_r^{a_r}$. The dual lattice N corresponds to the *group of 1-parameter subgroups* $\text{Hom}(\mathbb{C}^*, T)$. Here $N \ni n \mapsto (b_1, \dots, b_r) \in \mathbb{Z}^r$ corresponds to $\lambda^n: \mathbb{C}^* \rightarrow T$, $t \mapsto (t^{b_1}, \dots, t^{b_r})$. A *toric variety* is an irreducible variety containing an algebraic torus $T \cong (\mathbb{C}^*)^r$ as an open dense subset, such that the action of the torus on itself by multiplication extends to an algebraic action on the whole variety [CLS11, Def. 3.1.1]. We sketch how *normal toric varieties* can be constructed from cones and fans in $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$.

By a *cone* in $N_{\mathbb{R}}$ we mean a convex subset $\sigma = \text{cone}(S) = \{\sum_{i=1}^k \lambda_i v_i \mid \lambda_i \geq 0\}$ generated by a finite set $S = \{v_1, \dots, v_k\} \subseteq N$. The *dual cone* to σ in $M_{\mathbb{R}}$ is $\sigma^\vee = \{u \in M_{\mathbb{R}} \mid \langle u, v \rangle \geq 0 \text{ for all } v \in \sigma\}$. A cone σ is *pointed* if $\sigma \cap (-\sigma) = \{0\}$. We write $\tau \preceq \sigma$ whenever τ is a *face* of σ . A pointed cone $\sigma \subseteq N_{\mathbb{R}}$ leads to an *affine toric variety* $\text{TV}(\sigma) := \text{Spec } \mathbb{C}[\sigma^\vee \cap M]$. The inclusion of a face $\tau \preceq \sigma$ induces an open embedding $\text{TV}(\tau) \hookrightarrow \text{TV}(\sigma)$. In particular, the inclusion of the origin induces an open embedding of the torus $T = \text{Spec } \mathbb{C}[M] \hookrightarrow \text{TV}(\sigma)$.

A *fan* Σ in $N_{\mathbb{R}}$ is a finite collection of pointed cones that is closed under taking faces and such that the intersection of two cones is a face of each. The affine toric varieties $U_\sigma := \text{TV}(\sigma)$ associated to the cones σ in a fan Σ glue together to a *normal separated* toric variety $\text{TV}(\Sigma)$ with open affine charts the U_σ [CLS11, Thm. 3.1.5].

A *polyhedron* Δ in $M_{\mathbb{R}}$ is the intersection of finitely many closed half spaces

$$(2.1) \quad \Delta = \{m \in M_{\mathbb{R}} \mid \langle m, v_i \rangle \geq -\lambda_i, i = 1, \dots, s\} = \bigcap_{i=1}^s H_i^+,$$

¹Gefördert durch die Deutsche Forschungsgemeinschaft (DFG) im Rahmen der Exzellenzstrategie des Bundes und der Länder EXC 2044 -390685587, Mathematik Münster: Dynamik-Geometrie-Struktur

where $H_i^+ = \{m \in M_{\mathbb{R}} \mid \langle m, v_i \rangle \geq -\lambda_i\}$ for some inward pointing normal vector $v_i \in N_{\mathbb{R}}$ and some scalar $\lambda_i \in \mathbb{R}$ for $i \in \{1, \dots, s\}$. A compact polyhedron is called a *polytope*. A polyhedron Δ can be written as the *Minkowski sum* $\Delta = \nabla + \delta$ of a polytope ∇ and its *tail cone* $\delta = \text{tail}(\Delta) = \{m \in M_{\mathbb{R}} \mid u + m \in \Delta \text{ for all } u \in \Delta\} \subseteq M_{\mathbb{R}}$. To a full-dimensional lattice polyhedron Δ we associate its *inner normal fan*

$$(2.2) \quad \mathcal{N}(\Delta) := \{\tau \mid \tau \preceq \sigma_m, m \in \text{vert}(\Delta)\} \text{ with } |\mathcal{N}(\Delta)| = \text{tail}(\Delta)^\vee,$$

whose maximal cones σ_m are given by $\sigma_m := (\text{cone}(\Delta - m))^\vee = \mathbb{R}_{\geq 0} \cdot (\Delta - m)^\vee \subseteq N_{\mathbb{R}}$. This yields a semiprojective toric variety $\mathbb{P}(\Delta) := \mathbb{T}\mathbb{V}(\mathcal{N}(\Delta))$. We will assume all polyhedra to have at least one vertex. The normal fan of such a polyhedron will have convex support of full dimension.

2.1. Divisors on Toric Varieties. Let Σ be a fan in $N_{\mathbb{R}}$ with convex support of full dimension $r = \dim N_{\mathbb{R}}$, for example, the normal fan of a full-dimensional lattice polyhedron in $M_{\mathbb{R}}$. Let $X := \mathbb{T}\mathbb{V}(\Sigma)$ be the toric variety given by Σ .

Every lattice polyhedron $\Delta \subseteq M_{\mathbb{R}}$ with tail cone $|\Sigma|^\vee$ and whose normal fan $\mathcal{N}(\Delta)$ is refined by Σ gives rise to a nef Cartier divisor D_Δ on X by the following construction: If Σ refines $\mathcal{N}(\Delta)$ then the function $|\Sigma| \rightarrow \mathbb{R}, v \mapsto \min\langle \Delta, v \rangle$ is linear on the cones of Σ . So for each $\sigma \in \Sigma$ there is some $m_\sigma \in \Delta \cap M$ such that $\min\langle \Delta, v \rangle = \langle m_\sigma, v \rangle$ for each $v \in \sigma$. The lattice point m_σ is a vertex of Δ and if $\sigma \in \Sigma(r)$ is a maximal cone, then m_σ is uniquely determined. Note that $\Delta + \sigma^\vee = m_\sigma + \sigma^\vee$. Let D_Δ be the Cartier divisor with Cartier data $\{m_\sigma\}_{\sigma \in \Sigma}$, that is, locally on $U_\sigma \subseteq X$ given by $D|_{U_\sigma} = \text{div}(\chi^{-m_\sigma})|_{U_\sigma}$. Then for the associated line bundle $\mathcal{O}_X(\Delta) := \mathcal{O}_X(D_\Delta)$:

$$(2.3) \quad \Gamma(U_\sigma, \mathcal{O}_X(\Delta)) = \bigoplus_{m \in (\Delta + \sigma^\vee) \cap M} \mathbb{C} \cdot \chi^m = \chi^{m_\sigma} \cdot \mathbb{C}[\sigma^\vee \cap M] = \mathbb{C}[m_\sigma + (\sigma^\vee \cap M)],$$

so the vertex m_σ of Δ encodes the local sections of the line bundle $\mathcal{O}_X(\Delta)$ over the affine open U_σ . The line bundle $\mathcal{O}_X(\Delta)$ is *ample* if and only if $\mathcal{N}(\Delta) = \Sigma$.

We denote the set of lattice polyhedra with prescribed tail cone δ by Pol_δ^+ and the set of lattice polyhedra *compatible* with Σ , that is, with tail cone $|\Sigma|^\vee$ and whose normal fan is refined by Σ , by $\text{Pol}^+(\Sigma)$. The set Pol_δ^+ forms a semigroup with respect to Minkowski addition. For a fan Σ the set $\text{Pol}^+(\Sigma)$ forms a finitely generated subsemigroup. The union over the $\text{Pol}^+(\Sigma)$ with $|\Sigma|^\vee = \delta$ is the semigroup Pol_δ^+ . These semigroups are cancellative, which is caused by the presence of the prescribed common tail cone δ : It makes sure that a polyhedron $\Delta \in \text{Pol}_\delta^+$ is uniquely determined by the values of $\min\langle \Delta, v \rangle$ with v running through δ^\vee . Hence, these semigroups embed into their respective Grothendieck groups of formal differences:

$$(2.4) \quad \text{Pol}_\delta^+ \hookrightarrow \text{Pol}_\delta = \{\Delta^+ - \Delta^- \mid \Delta^+, \Delta^- \in \text{Pol}_\delta^+\},$$

$$(2.5) \quad \text{Pol}^+(\Sigma) \hookrightarrow \text{Pol}(\Sigma) = \{\Delta^+ - \Delta^- \mid \Delta^+, \Delta^- \in \text{Pol}^+(\Sigma)\}.$$

On a quasiprojective toric variety $X = \mathbb{T}\mathbb{V}(\Sigma)$, every Cartier divisor D can be written (non-uniquely) as a difference $D = D^+ - D^-$ with both D^+ and D^- nef

Cartier divisors [CLS11, Thm. 6.3.22]. In particular every Cartier divisor on X can be represented by a pair of lattice polyhedra (Δ^+, Δ^-) compatible with the fan Σ , that is, an element of the Grothendieck group $\text{Pol}(\Sigma)$.

3. FROM COMPLEXES OF POLYHEDRA TO COMPLEXES OF SHEAVES

3.1. The Koszul Complex of Polyhedra. In this section we construct “exact sequences of polyhedra” that induce exact sequences of split vector bundles on a toric variety $X = \mathbb{T}\mathbb{V}(\Sigma)$ over \mathbb{C} . We start out on the polyhedral side and let Σ be a fan in $N_{\mathbb{R}}$ with convex support of full dimension. Let $\text{Pol}^+(\Sigma)$ denote the set of lattice polyhedra in $M_{\mathbb{R}}$ compatible with Σ . Also include the empty set in $\text{Pol}^+(\Sigma)$.

Definition 1. A Σ -family of polyhedra is a finite set $S = \{\nabla_i \mid i \in I\}$ of lattice polyhedra $\nabla_i \in \text{Pol}^+(\Sigma)$ satisfying the following conditions:

- (1) $\bigcup_{i \in I} \nabla_i =: \nabla \in \text{Pol}^+(\Sigma)$ is a lattice polyhedron compatible with Σ ;
- (2) all intersections $\nabla_{I'} := \bigcap_{i \in I'} \nabla_i$ with $\emptyset \neq I' \subseteq I$ are either empty or compatible with Σ , that is, $\nabla_{I'} \in \text{Pol}^+(\Sigma)$. Furthermore, set $\nabla_{\emptyset} := \nabla$.

We consider two categories and a functor associated to a Σ -family between them.

Definition 2. Let 2^I be the *poset category* associated to the power set of the finite set I , that is, objects are subsets of I and there exists a unique morphism from I' to I'' whenever $I' \subseteq I''$. Let $\text{Pol}^+(\Sigma)$ also denote the *category of lattice polyhedra compatible with the fan Σ* , that is, objects are compatible lattice polyhedra as defined above or the empty set and for two polyhedra Δ_1 and Δ_2 in $\text{Pol}^+(\Sigma)$ we define

$$(3.1) \quad \text{Hom}_{\text{Pol}^+(\Sigma)}(\Delta_1, \Delta_2) = \begin{cases} \{\text{inclusion}\} & \text{if } \Delta_1 \subseteq \Delta_2, \\ \emptyset & \text{if } \Delta_1 \not\subseteq \Delta_2. \end{cases}$$

Given an Σ -family of polyhedra S we can define a contravariant functor

$$(3.2) \quad F_S: 2^I \rightarrow \text{Pol}^+(\Sigma) \quad I' \mapsto \nabla_{I'} = \bigcap_{i \in I'} \nabla_i, \quad (I' \subseteq I'') \mapsto (\nabla_{I''} \hookrightarrow \nabla_{I'}).$$

In general, given any contravariant functor $F: 2^I \rightarrow \text{Pol}^+(\Sigma)$, we define a subcomplex of the Koszul complex $\bigwedge^\bullet \mathbb{C}^I$ as follows.

Definition 3. For a contravariant functor $F: 2^I \rightarrow \text{Pol}^+(\Sigma)$ and $p \in \mathbb{N}$ let

$$(3.3) \quad \mathcal{C}_p^F := \text{span}\{e_{I'} \mid \#I' = p \text{ and } F(I') \neq \emptyset\} \subseteq \bigwedge^p \mathbb{C}^I.$$

Here $e_{I'} := \bigwedge_{i \in I'} e_i$, where $\{e_i \mid i \in I\}$, denotes the canonical basis of \mathbb{C}^I .

The map $d: \mathcal{C}_{p+1}^F \rightarrow \mathcal{C}_p^F$ is defined as for the Koszul complex $\bigwedge^\bullet \mathbb{C}^I$. For a fixed total order on I , say $I = \{1 < \dots < \#I\}$, and $I' = \{i_0 < \dots < i_p\} \subseteq I$, we have

$$(3.4) \quad d(e_{I'}) = \sum_{j=0}^p (-1)^j e_{I' \setminus \{i_j\}} = \sum_{i \in I'} (-1)^{|i|} e_{I' \setminus \{i\}},$$

where $|i|$ refers to the index j of $i = i_j$ in I' .

Recall that the polyhedra $F(I') \in \text{Pol}^+(\Sigma)$ are contained in $M_{\mathbb{R}}$. For each $m \in M_{\mathbb{R}}$ we define the *evaluation subcomplex* (“at m ”) as

$$(3.5) \quad \mathcal{C}_p^F(m) := \text{span}\{e_{I'} \mid \#I' = p \text{ and } m \in F(I')\} \subseteq \mathcal{C}_p^F \subseteq \bigwedge^p \mathbb{C}^I$$

with the restrictions $d_{|\mathcal{C}_{p+1}^F(m)}: \mathcal{C}_{p+1}^F(m) \rightarrow \mathcal{C}_p^F(m)$ as boundary maps.

Lemma 4. *For a Σ -family S set $F := F_S$.*

- (i) *If $m \in M_{\mathbb{R}} \setminus \nabla$, then $\mathcal{C}_p^F(m) = 0$ for all $p \geq 0$. In particular, $\mathcal{C}_\bullet^F(m)$ is exact.*
- (ii) *If $m \in \nabla \subseteq M_{\mathbb{R}}$, then $\mathcal{C}_\bullet^F(m)$ is still exact.*
- (iii) *The complex $\mathcal{C}_\bullet^F = \sum_{m \in M} \mathcal{C}_\bullet^F(m)$ is exact.*

Proof. Statement (i) is clear because $F(I') \subseteq \nabla$ for all $I' \subseteq I$.

For statement (ii) set $I_m := \{i \in I \mid m \in F(i) := F(\{i\}) = \nabla_i\}$. Then for $I' \subseteq I$ we have $I' \subseteq I_m$ if and only if $m \in \bigcap_{i \in I'} F(i) = F(I')$. Therefore $\mathcal{C}_\bullet^F(m)$ equals $\bigwedge^\bullet \mathbb{C}^{I_m}$, which is exact [Wei95, Cor. 4.5.5].

For statement (iii) first note that we have $\mathcal{C}_\bullet^F = \sum_{m \in M_{\mathbb{R}}} \mathcal{C}_\bullet^F(m) = \sum_{m \in \nabla \cap M} \mathcal{C}_\bullet^F(m)$. By (i) and (ii) each of the summands $\mathcal{C}_\bullet^F(m)$ is exact. For any subset $M' \subseteq M_{\mathbb{R}}$ the intersection of the evaluation subcomplexes for $m \in M'$ is $\bigcap_{m \in M'} \mathcal{C}_\bullet^F(m) = \bigwedge^\bullet \mathbb{C}^{I_{M'}}$, where $I_{M'} := \{i \in I \mid M' \subseteq F(i) = \nabla_i\}$. Furthermore, we have for $m_1, m_2, m_3 \in M_{\mathbb{R}}$:

$$(\mathcal{C}_\bullet^F(m_1) + \mathcal{C}_\bullet^F(m_2)) \cap \mathcal{C}_\bullet^F(m_3) = (\mathcal{C}_\bullet^F(m_1) \cap \mathcal{C}_\bullet^F(m_3)) + (\mathcal{C}_\bullet^F(m_2) \cap \mathcal{C}_\bullet^F(m_3)),$$

because the $\mathcal{C}_p^F(m_i)$ are all subspaces of \mathcal{C}_p^F spanned by a subset of the prescribed basis $\{e_{I'} \mid I' \subseteq I, \#I' = p, F(I') \neq \emptyset\}$ of \mathcal{C}_p^F .

Now, we switch to a slightly more general setup. Assume that $\mathcal{C}_\bullet^i \subseteq \mathcal{C}_\bullet^F$ ($i = 1, \dots, k$) are complexes such that all their mutual intersections are exact and

$$(\sum_{i \in I} \mathcal{C}_\bullet^i) \cap \mathcal{C}_\bullet^j = \sum_{i \in I} (\mathcal{C}_\bullet^i \cap \mathcal{C}_\bullet^j)$$


for any j and $I \subseteq \{1, \dots, k\}$. Then $\sum_{i=1, \dots, k} \mathcal{C}_\bullet^i$ is exact, too. The proof uses induction by k and exploits the short exact sequence

$$0 \rightarrow \left(\sum_{i=1, \dots, k-1} \mathcal{C}_\bullet^i \right) \cap \mathcal{C}_\bullet^k \rightarrow \left(\sum_{i=1, \dots, k-1} \mathcal{C}_\bullet^i \right) \oplus \mathcal{C}_\bullet^k \rightarrow \sum_{i=1, \dots, k} \mathcal{C}_\bullet^i \rightarrow 0.$$

The generalization of the setup is needed to ensure that the induction hypothesis implies that, besides the central term, the left most complex is exact, too.

Finally, we apply the previous claim to the complexes $\mathcal{C}_\bullet^i := \mathcal{C}_\bullet^F(m)$ where $m \in \nabla \cap M$ replaces $i = 1, \dots, k$. Note that even though the set $\nabla \cap M$ may be infinite, the induction ends after finitely many steps since the Koszul complex $\bigwedge^\bullet \mathbb{C}^I$ is finite-dimensional and all $\mathcal{C}_\bullet^F(m) \subseteq \mathcal{C}_\bullet^F \subseteq \bigwedge^\bullet \mathbb{C}^I$. \square

Example 5. We revisit an example already encountered in the Introduction (1.2). Let Σ be the fan of the first Hirzebruch surface $\mathbb{F}_1 = \mathbb{P}(\Delta) = \mathbb{T}\mathbb{V}(\Sigma)$.



The fan is complete, so compatible polyhedra will be polytopes. Define the Σ -family

$$S := \left\{ \begin{array}{c} \nabla_0 \\ \text{[diagram of } \nabla_0 \text{]} \end{array}, \begin{array}{c} \nabla_1 \\ \text{[diagram of } \nabla_1 \text{]} \end{array} \right\} \text{ and set } \nabla := \nabla_0 \cup \nabla_1 = \text{[diagram of } \nabla \text{]} \in \text{Pol}^+(\Sigma).$$

The complex \mathcal{C}_\bullet^F is $0 \rightarrow \mathbb{C} \cdot (e_0 \wedge e_1) \rightarrow \mathbb{C} \cdot e_0 \oplus \mathbb{C} \cdot e_1 \rightarrow \mathbb{C} \cdot e_\emptyset \rightarrow 0$. The evaluation subcomplex $\mathcal{C}_\bullet^F(m)$ for $m \in M_\mathbb{R}$ is the subcomplex indicating whether $m \in \nabla$ for $\mathcal{C}_0^F(m)$, $m \in \nabla_0$ respectively $m \in \nabla_1$ for $\mathcal{C}_1^F(m)$, and $m \in \nabla_0 \cap \nabla_1 = \text{[diagram of } \nabla_0 \cap \nabla_1 \text{]}$ for $\mathcal{C}_2^F(m)$.

There is a particularly nice case in which exactness of the evaluation subcomplexes for all lattice points $m \in M$ is equivalent to exactness for all $m \in M_\mathbb{R}$.

Lemma 6. *Assume that Σ consists of a single cone σ and its faces, with σ^\vee smooth and full-dimensional. Then for any contravariant functor $F: 2^I \rightarrow \text{Pol}^+(\Sigma)$, the complexes $\mathcal{C}_\bullet^F(m)$ are exact for all $m \in M$ if and only if they are exact for all $m \in M_\mathbb{R}$.*

Remark 7. This situation occurs naturally when we consider smooth affine toric varieties $\mathbb{T}\mathbb{V}(\sigma)$ or an affine open subset $U_\sigma \subseteq \mathbb{T}\mathbb{V}(\Sigma)$ for a cone $\sigma \in \Sigma$. In the latter case the polyhedra $F(I')$ are changed to $F(I') + \sigma^\vee$ when considering the affine open U_σ (see section (3.2)). In the case of a functor F_S associated to a Σ -family S the ∇_i become $\nabla_i + \sigma^\vee$ and the $\nabla_{I'}$ for $I' \subseteq I$ become $\nabla_{I'} + \sigma^\vee$.

Proof of the Lemma. Because σ^\vee is smooth, we may choose coordinates such that $M = \mathbb{Z}^r$ and $\sigma^\vee = \mathbb{R}_{\geq 0}^r$. For $F(I') \in \text{Pol}^+(\Sigma)$ we either have $F(I') = \emptyset$, in which case $m \in F(I') \Leftrightarrow \lfloor m \rfloor \in F(I')$ is clear, or $F(I')$ has only one vertex and can be written as $F(I') = r_{I'} + \mathbb{R}_{\geq 0}^r$ for some $r_{I'} \in \mathbb{Z}^r$. We obtain for every $m \in \mathbb{R}^r$ that

$$m \in F(I') \Leftrightarrow m \geq r_{I'} \Leftrightarrow \lfloor m \rfloor \geq r_{I'} \Leftrightarrow \lfloor m \rfloor \in F(I'),$$

where $\lfloor m \rfloor$ and the relation \geq are meant componentwise. Hence, $\mathcal{C}_\bullet^F(m) = \mathcal{C}_\bullet^F(\lfloor m \rfloor)$, proving the claim. \square

Example 8. We give a counterexample for the global situation, that is, for Σ consisting of more than a cone σ . Let $\Sigma = \mathcal{N}(\Delta)$ be the fan of $\mathbb{P}^1 = \mathbb{P}(\Delta) = \mathbb{T}\mathbb{V}(\Sigma)$:

$$\Delta = \text{[diagram of } \Delta \text{]} \text{ in } M_\mathbb{R} = \mathbb{R}, \quad \mathcal{N}(\Delta) = \text{[diagram of } \mathcal{N}(\Delta) \text{]} \text{ in } N_\mathbb{R} = \mathbb{R}.$$

Let $I := \{0, 1\}$ and define the functor $F: 2^I \rightarrow \text{Pol}^+(\Sigma)$ as follows

$$\emptyset \mapsto \text{[diagram of } \emptyset \text{]}, \quad \{0\} \mapsto \text{[diagram of } \{0\} \text{]}, \quad \{1\} \mapsto \text{[diagram of } \{1\} \text{]}, \quad \{0, 1\} \mapsto \text{[diagram of } \{0, 1\} \text{]} = \emptyset.$$

The only non-trivial evaluation subcomplexes at lattice points are at $m = 0$, $m = 1$:

$$\begin{aligned} \mathcal{C}_\bullet^F(0): 0 &\rightarrow \mathbb{C} \cdot e_{\{0\}} \xrightarrow{\cong} \mathbb{C} \cdot e_\emptyset \rightarrow 0 \\ \mathcal{C}_\bullet^F(1): 0 &\rightarrow \mathbb{C} \cdot e_{\{1\}} \xrightarrow{\cong} \mathbb{C} \cdot e_\emptyset \rightarrow 0 \end{aligned}$$

These are exact. But at $m = \frac{1}{2} \in M_{\mathbb{R}}$ we have the non-exact evaluation subcomplex

$$\mathcal{C}_\bullet^F\left(\frac{1}{2}\right): 0 \rightarrow 0 \rightarrow \mathbb{C} \cdot e_\emptyset \rightarrow 0.$$

3.2. Localization of the Koszul Complex. We use the notation of section (3.1). In section (4) we will think about the polyhedra $\nabla_{I'}$ as representing some nef line bundles on some toric variety $\mathbb{P}(\Delta) = \mathbb{T}\mathbb{V}(\mathcal{N}(\Delta))$ over \mathbb{C} . Then the complexes constructed in subsection (3.1) describe maps between the global sections of them, and thus among the sheaves themselves. To understand the local behavior of these complexes, we will look at the affine charts $\mathbb{T}\mathbb{V}(\sigma) \subseteq \mathbb{P}(\Delta)$ for cones $\sigma \in \mathcal{N}(\Delta)$.

Definition 9. For a contravariant functor $F: 2^I \rightarrow \text{Pol}^+(\Sigma)$, where $|\Sigma|$ is convex and full-dimensional in $M_{\mathbb{R}}$, and a cone $\sigma \in \Sigma$ define the functor

$$(3.6) \quad F^\sigma: 2^I \rightarrow \text{Pol}^+(\sigma), I' \mapsto F(I') + \sigma^\vee.$$

Because $F(I') \in \text{Pol}^+(\Sigma)$ has tail cone $|\Sigma|^\vee$ or is empty and $\sigma \in \Sigma$ implies $|\Sigma|^\vee \subseteq \sigma^\vee$, the image $F(I') + \sigma^\vee$ has tail cone $|\Sigma|^\vee + \sigma^\vee = \sigma^\vee$ or is empty. If σ^\vee is full-dimensional in $M_{\mathbb{R}}$, then the normal fan of $F(I') + \sigma^\vee$ consists of just σ with its faces, that is, $F(I') + \sigma^\vee \in \text{Pol}^+(\sigma)$.

Proposition 10. For a polyhedron Δ and a finite set I , let $F: 2^I \rightarrow \text{Pol}^+(\mathcal{N}(\Delta))$ be a contravariant functor satisfying the conclusions (i) - (iii) of Lemma 4. Then the complexes $\mathcal{C}_\bullet^{F^\sigma}(m)$ are exact for all $m \in M_{\mathbb{R}}$ and for all $\sigma \in \mathcal{N}(\Delta)$.

Remark 11. Moreover, this claim remains true if we consider any polyhedra compatible with the fan $\mathcal{N}(\Delta)$, not just lattice polyhedra. This generalization will be important for the induction performed in the upcoming proof.

Proof. Step 1. Assume that $r = \text{rk } M = 1$. Then $\Delta \subseteq \mathbb{R}$ is a subset of the real line and its normal fan $\mathcal{N}(\Delta)$ can contain the origin $\sigma = \{0\}$, the rays $\sigma = \mathbb{R}_{\geq 0}$ and $\sigma = \mathbb{R}_{\leq 0}$ and the real line $\sigma = \mathbb{R}$ as cones. For $\sigma = \mathbb{R}$ and $\sigma = \{0\}$ the claim is clear from the assumptions. We deal with the case $\sigma^\vee = \mathbb{R}_{\geq 0}$ (the case $\sigma^\vee = \mathbb{R}_{\leq 0}$ is analogous). A polyhedron $F(I')$ compatible with $\mathcal{N}(\Delta)$ is either empty or an interval of the form $[a, b]$ with $-\infty < a \leq b \leq \infty$. For $F(I') \neq \emptyset$, let $a(I') \in \mathbb{R}$ denote the start point of the interval $F(I')$ and $b(I') \in \mathbb{R} \cup \{\infty\}$ the end point. Let $A(F) := \{a(I') \mid F(I') \neq \emptyset\}$ and $B(F) := \{b(I') \mid F(I') \neq \emptyset\}$ be the sets of start and end points, respectively. Since I is finite, $A(F) \subseteq \mathbb{R}$ and $B(F) \subseteq \mathbb{R} \cup \{\infty\}$ are finite.

We understand the local complex $\mathcal{C}_p^{F^\sigma}$ in terms of global evaluation subcomplexes:

$$\begin{aligned}\mathcal{C}_p^{F^\sigma}(m) &= \text{span}\{e_{I'} \mid \#I' = p \text{ and } m \in F(I') + \mathbb{R}_{\geq 0}\} \\ &= \text{span}\{e_{I'} \mid \#I' = p \text{ and } a(I') \leq m\} \\ &= \sum_{m' \leq m} \mathcal{C}_p^F(m') \subseteq \mathcal{C}_p^F.\end{aligned}$$

We make three observations:

- (1) For $m \in \mathbb{R}$ satisfying $m < b$ for all $b \in B(F)$:

$$\begin{aligned}\mathcal{C}_p^{F^\sigma}(m) &= \text{span}\{e_{I'} \mid \#I' = p \text{ and } a(I') \leq m\} \\ &= \text{span}\{e_{I'} \mid \#I' = p \text{ and } a(I') \leq m \leq b(I')\} = \mathcal{C}_p^F(m).\end{aligned}$$

So for $m \ll 0$ the local complexes $\mathcal{C}_p^{F^\sigma}(m)$ stabilize to the exact sequence $\mathcal{C}_p^F(m)$. Recall that $B(F)$ is finite, so m can be chosen small enough.

- (2) For $m' \leq m$ we have embeddings $\mathcal{C}_p^{F^\sigma}(m') \hookrightarrow \mathcal{C}_p^{F^\sigma}(m)$ of complexes.
(3) A non-trivial jump from $\mathcal{C}_p^{F^\sigma}(< m) := \sum_{m' < m} \mathcal{C}_p^{F^\sigma}(m')$ to $\mathcal{C}_p^{F^\sigma}(m)$ can only occur at a starting point $m = a(I') \in A(F)$ for some $I' \subseteq I$.

We inductively show exactness of $\mathcal{C}_p^{F^\sigma}(m)$ for all $m \in M_{\mathbb{R}}$ by investigating these non-trivial jumps. Let $m := a(I') \in A(F)$. Because I is finite we can choose an $\epsilon > 0$ such that any $a \in A(F)$ and any $b \in B(F)$ is either equal to m or has distance $|a - m| > \epsilon$ respectively $|b - m| > \epsilon$. Consider the non-trivial embedding $\mathcal{C}_p^{F^\sigma}(m - \epsilon) \hookrightarrow \mathcal{C}_p^{F^\sigma}(m)$ and denote its cokernel by \mathcal{C}_\bullet . This yields an exact sequence

$$(3.7) \quad 0 \rightarrow \mathcal{C}_p^{F^\sigma}(m - \epsilon) \rightarrow \mathcal{C}_p^{F^\sigma}(m) \rightarrow \mathcal{C}_\bullet \rightarrow 0.$$

Because $b(F) \cap [m - \epsilon, m] = \emptyset$, there is also an embedding $\mathcal{C}_p^F(m - \epsilon) \hookrightarrow \mathcal{C}_p^F(m)$ for the original functor F , also with cokernel \mathcal{C}_\bullet . This gives the short exact sequence

$$(3.8) \quad 0 \rightarrow \mathcal{C}_p^F(m - \epsilon) \rightarrow \mathcal{C}_p^F(m) \rightarrow \mathcal{C}_\bullet \rightarrow 0$$

in which the first two complexes are exact by assumption, so \mathcal{C}_\bullet is also exact.

By observation (1) above we can start from the exact complex $\mathcal{C}_p^{F^\sigma}(m) = \mathcal{C}_p^F(m)$ for $m < b$ for all $b \in B(F)$ and then inductively use the exact sequence (3.7) with exact complexes $\mathcal{C}_p^{F^\sigma}(m - \epsilon)$ and \mathcal{C}_\bullet to obtain that $\mathcal{C}_p^{F^\sigma}(m)$ is exact for all $m \in M_{\mathbb{R}}$.

Step 2. Consider a general lattice M of rank $r \in \mathbb{N}_{\geq 1}$ and let $C \subseteq M_{\mathbb{R}}$ be any ray. Replacing the polyhedra $F(I')$ by the Minkowski sums $F(I') + C$ yields the functor $F^C: 2^I \rightarrow \text{Pol}_{\delta+C}^{+, \mathbb{R}}$, where $\delta = |\mathcal{N}(\Delta)|^\vee$ is the tail cone of Δ . Now, for $m \in M_{\mathbb{R}}$ we restrict the complexes \mathcal{C}_p^F and \mathcal{C}_p^{F+C} to the affine line $m + (C - C)$, that is, consider $F(I') \cap (m + (C - C))$ and $(F(I') + C) \cap (m + (C - C))$. By assumption, the complex $\mathcal{C}_p^{F \cap (m + (C - C))}$ leads to exact evaluation subcomplexes $\mathcal{C}_p^{F \cap (m + (C - C))}(m') = \mathcal{C}_p^F(m')$

for all $m' \in m + (C - C)$ because in this case

$$\begin{aligned} \mathcal{C}_p^{F \cap (m + (C - C))}(m') &= \text{span}\{e_{I'} \mid \#I' = p \text{ and } m' \in F(I') \cap (m + (C - C))\} \\ &= \text{span}\{e_{I'} \mid \#I' = p \text{ and } m' \in F(I')\} = \mathcal{C}_p^F(m'). \end{aligned}$$

The complex $\mathcal{C}_\bullet^{(F+C) \cap (m+(C-C))}$ inherits this property by step 1 and we obtain exactness of $\mathcal{C}_\bullet^{F+C}(m')$ for all $m' \in M_{\mathbb{R}}$.

Step 3. If $\sigma^\vee \in \mathcal{N}(\Delta)$ is an arbitrary (non-trivial) polyhedral cone, then we may apply step 2 successively to all its fundamental rays. \square

3.3. Exactness of the Sequence Associated to a Σ -Family.

Theorem 12. *Let Δ be a full-dimensional lattice polyhedron in $M_{\mathbb{R}}$ with normal fan $\Sigma := \mathcal{N}(\Delta)$ and $X := \mathbb{P}(\Delta)$ the toric variety given by Δ . For $S = \{\nabla_i \mid i \in I\}$ a Σ -family with $\#I = n$, let $F_S: 2^I \rightarrow \text{Pol}^+(\Sigma)$ be the functor introduced in Definition 2. The complex $\mathcal{C}_\bullet^{F_S}$ with its exact evaluation subcomplexes $\mathcal{C}_\bullet^{F_S}(m)$ for $m \in M_{\mathbb{R}}$ induces a T -equivariant exact sequence of direct sums of nef line bundles on X*

$$(3.9) \quad 0 \rightarrow \mathcal{O}_X(\nabla_I) \rightarrow \bigoplus_{\#I'=n-1} \mathcal{O}_X(\nabla_{I'}) \rightarrow \cdots \rightarrow \bigoplus_{i \in I} \mathcal{O}_X(\nabla_i) \rightarrow \mathcal{O}_X(\nabla) \rightarrow 0.$$

Proof. The evaluation subcomplex $\mathcal{C}_\bullet^{F_S}(m)$ includes $\mathbb{C} \cdot e_{I'}$ if and only if $m \in \nabla_{I'}$. For $m \in M$, this in turn is equivalent to $\chi^m \in \Gamma(X, \mathcal{O}_X(D_{\nabla_{I'}}))$. Hence, the evaluation subcomplex $\mathcal{C}_\bullet^{F_S}(m)$ corresponds to a complex of global sections in degree $m \in M$:

$$(3.10) \quad \mathcal{C}_p^{F_S}(m) \cong \bigoplus_{\#I'=p} \Gamma(X, \mathcal{O}_X(\nabla_{I'}))_m.$$

Summing the $\mathcal{C}_\bullet^{F_S}(m)$ for lattice points $m \in M$ yields a sequence of global sections:

$$(3.11) \quad 0 \rightarrow \Gamma(X, \mathcal{O}_X(\nabla_I)) \rightarrow \cdots \rightarrow \Gamma(X, \bigoplus_{i \in I} \mathcal{O}_X(\nabla_i)) \rightarrow \Gamma(X, \mathcal{O}_X(\nabla)) \rightarrow 0.$$

The sheaves $\mathcal{O}_X(\nabla_I)$, $\mathcal{O}_X(\nabla_i)$, $i \in I$, and $\mathcal{O}_X(\nabla)$ are globally generated. By T -equivariance, they are subsheaves of $j_* \mathcal{O}_T$ for $j: T \hookrightarrow X$ the inclusion of the torus. Hence, the sequence of global sections (3.11) determines a sequence of sheaves:

$$(3.12) \quad 0 \rightarrow \mathcal{O}_X(\nabla_I) \rightarrow \cdots \rightarrow \bigoplus_{i \in I} \mathcal{O}_X(\nabla_i) \rightarrow \mathcal{O}_X(\nabla) \rightarrow 0.$$

For a cone $\sigma \in \Sigma$ and a lattice point $m \in M$, the evaluation subcomplex $\mathcal{C}_\bullet^{F_S^\sigma}(m)$ includes $e_{I'}$ for $I' \subseteq I$ if and only if $\chi^m \in \Gamma(U_\sigma, \mathcal{O}_X(\nabla_{I'}))$. The sequence of sections over $U_\sigma \subseteq X$ therefore corresponds to the direct sum of the evaluation subcomplexes $\mathcal{C}_\bullet^{F_S^\sigma}(m)$ for $m \in M$. By Proposition 10 this sequence is exact for each $\sigma \in \Sigma$. Hence, the restriction of the sequence of sheaves to each affine chart of the covering $\{U_\sigma\}_{\sigma \in \Sigma}$ is exact and consequently the constructed sequence is exact. \square

4. DISPLAYING Ext^1

Throughout this section let $X := \mathbb{P}(\Delta)$ be the toric variety over \mathbb{C} associated to the lattice polyhedron $\Delta \subseteq M_{\mathbb{R}}$, whose normal fan $\Sigma := \mathcal{N}(\Delta)$ has convex support of full dimension. The (possibly trivial) tail cone of Δ is $\delta := \text{tail}(\Delta) = |\Sigma|^\vee$. Let $\Delta^+, \Delta^- \in \text{Pol}^+(\Sigma)$ be lattice polyhedra compatible with Σ , that is, their normal fans are refined by Σ . These polyhedra correspond to T -invariant nef Cartier divisors $D^+ = D_{\Delta^+}$ and $D^- = D_{\Delta^-}$ on X . We study the space of extensions

$$(4.1) \quad \text{Ext}(\Delta^-, \Delta^+) \cong \text{Ext}^1(\Delta^-, \Delta^+) := \text{Ext}^1(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+)),$$

that is, extensions of the line bundle $\mathcal{O}_X(\Delta^-)$ by the line bundle $\mathcal{O}_X(\Delta^+)$. More specifically, we study T -equivariant extension sequences. These are elements in $\text{Ext}(\Delta^-, \Delta^+)_0$. To understand this space we start with specific extension sequences induced by inclusion/exclusion sequences of polyhedra, such as the sequence considered in the Introduction (1.2). For an n -dimensional $\text{Ext}(\Delta^-, \Delta^+)_0$ all extensions are encoded in a single sequence of the form $0 \rightarrow \mathcal{O}_X(\Delta^+)^n \rightarrow \mathcal{H} \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0$. We will show that this is the universal extension sequence for $\text{Ext}(\Delta^-, \Delta^+)_0$. First, we show that it can be constructed from the exact sequence of sheaves associated to a Σ -family (Theorem 12). We then trace the sequence along the identifications:

$$\begin{aligned} \text{Ext}(\Delta^-, (\Delta^+)^n) &\stackrel{(1)}{\cong} \text{Ext}(\mathcal{O}_X, \mathcal{O}_X(\Delta^+ - \Delta^-)^n) \\ &\stackrel{(2)}{=} \text{H}^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-))^n \\ &\stackrel{(3)}{\cong} \left(\bigoplus_{m \in M} \tilde{\text{H}}^0(\Delta^- \setminus (\Delta^+ - m)) \right)^n. \end{aligned}$$

Note that all of the above groups are M -graded and all identifications respect these M -gradings. Since we consider equivariant extensions, we will obtain an n -tuple of elements in $\text{H}^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-))_0 \cong \tilde{\text{H}}^0(\Delta^- \setminus \Delta^+)$, without an integral shift.

4.1. Inclusion of Polyhedra. Since we are relating $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$ to the reduced singular cohomology group $\tilde{\text{H}}^0(\Delta^- \setminus \Delta^+)$, the easiest case is to assume Δ^+ to be contained in Δ^- . The inclusion/exclusion sequence of polyhedra is obtained by covering Δ^- by certain polyhedra ∇_i that intersect (pairwise) in Δ^+ . These ∇_i are obtained by taking the unions of the connected components C_0, \dots, C_n of the set-theoretic difference $\Delta^- \setminus \Delta^+$ with Δ^+ , that is, $\nabla_i := C_i \cup \Delta^+$, $i = 0, \dots, n$.

Proposition 13. *For two lattice polyhedra $\Delta^+ \subseteq \Delta^- \in \text{Pol}^+(\Sigma)$ and each connected component $C = C_i$ of $\Delta^- \setminus \Delta^+$, the union $\nabla = C \cup \Delta^+$ is again a lattice polyhedron.*

Proof. It is easy to see that ∇ is closed. We first show that ∇ is convex. Assume that $x, y \in \nabla$. We know that $\overline{xy} \subseteq \Delta^-$, and this line segment might touch Δ^+ or not. If not, then $\overline{xy} \subseteq (\Delta^- \setminus \Delta^+)$. Since \overline{xy} is connected, it then has to be contained in a single connected component of $\Delta^- \setminus \Delta^+$. Since $x \in \nabla = C \cup \Delta^+$, we know that $x \in C$. Hence, $\overline{xy} \subseteq C$. If $\overline{xy} \cap \Delta^+ \neq \emptyset$, then this set is a closed

subsegment $\overline{x'y'} \subseteq \overline{xy}$. In particular, the half open ends $\overline{xx'}$ and $\overline{y'y}$ (excluding x' and y') belong to $\Delta^- \setminus \Delta^+$, and the same argument as in the first case applies again: Since $x, y \in \nabla = C \cup \Delta^+$, both of these half open ends $\overline{xx'}$ and $\overline{y'y}$ lie in C . The segment \overline{xy} thus lies in $\nabla = C \cup \Delta^+$.

Next we realize that vertices of ∇ are vertices of Δ^+ or of Δ^- . We start with the case of Δ^+ and Δ^- being compact. Choose a generic regular triangulation induced from some map $\omega: \text{vert}(\Delta^+) \rightarrow \mathbb{R}$ and then extend it generically to $\text{vert}(\Delta^-) \setminus \text{vert}(\Delta^+)$ with sufficiently independent heights. This yields a triangulation $\Delta^- = \bigcup_{i \in I} \Delta^i$ that restricts to a triangulation $\Delta^+ = \bigcup_{j \in J \subseteq I} \Delta^j$ that uses only the vertices of Δ^- and Δ^+ . In particular all Δ^i are lattice simplices. The union $\nabla = C \cup \Delta^+$ is a union of lattice simplices in I . In particular all of its vertices are lattice points.

If Δ^+ and Δ^- are not compact but have the same tail cone δ we choose a half-space H such that all vertices of Δ^+ and Δ^- are contained in the interior of H . Then we can write $\Delta^+ = P^+ + \delta$ and $\Delta^- = P^- + \delta$ with $P^+ := \Delta^+ \cap H$, $P^- := \Delta^- \cap H$. The previous discussion applied to the polytopes P^+ and P^- , all of whose relevant vertices are integral (not necessarily those lying on the boundary of H , but they do not yield vertices of $C \cup \Delta^+$), yields that all vertices of $\nabla = C \cup \Delta^+$ are integral. \square

Corollary 14. *For $\Delta^+ \subseteq \Delta^- \in \text{Pol}^+(\Sigma)$, the set $S := \{\nabla_i \mid i \in \{0, \dots, n\} =: I\}$ with $\nabla_i := C_i \cup \Delta^+$ for the connected components C_0, \dots, C_n of $\Delta^- \setminus \Delta^+$ yields a Σ' -family in the sense of Definition 1 for some refinement $\Sigma' \leq \Sigma$.*

Proof. By the previous Proposition 13 the ∇_i are lattice polyhedra. The tail cone of each ∇_i is $|\Sigma|^\vee$. Hence, each normal fan $\mathcal{N}(\nabla_i)$ has the same support as Σ . Since we are only dealing with finitely many ∇_i there is a common refinement Σ' of Σ and all $\mathcal{N}(\nabla_i)$. Then $\nabla_i \in \text{Pol}^+(\Sigma')$. The conditions for a Σ' -family follow from the assumptions on Σ' and $\nabla_{I'} = \Delta^+ \in \text{Pol}^+(\Sigma')$ for any $I' \subseteq I$ with $\#I' \geq 2$. \square

4.1.1. *Refining the Fan.* In many cases the lattice polyhedra ∇_i built from the components of $\Delta^- \setminus \Delta^+$ will already be compatible with the fan Σ we started with. We believe that if Δ^+ and Δ^- are sufficiently ample, this will always be the case. However, in general, if the ∇_i are not compatible with Σ , we can refine Σ to a fan Σ' with the same support such that all ∇_i are compatible with Σ' , as in Corollary 14. This induces a proper birational toric morphism $\pi: X' = \mathbb{T}\mathbb{V}(\Sigma') \rightarrow \mathbb{T}\mathbb{V}(\Sigma) = X$. This morphism satisfies that $\pi_* \mathcal{O}_{X'} = \mathcal{O}_X$ and $R^\nu \pi_* \mathcal{O}_{X'} = 0$ for all $\nu > 0$. The projection formula implies that $\pi_* \pi^* \mathcal{F} = \mathcal{F}$ for a locally free sheaf \mathcal{F} on X . For a short exact sequence of sheaves on X' of the form

$$(4.2) \quad 0 \rightarrow \pi^* \mathcal{F}_1 \rightarrow \mathcal{E} \rightarrow \pi^* \mathcal{F}_2 \rightarrow 0,$$

derived pushforward $R\pi_*$ yields a long exact sequence of sheaves on X :

$$(4.3) \quad 0 \rightarrow \underbrace{\pi_* \pi^* \mathcal{F}_1}_{=\mathcal{F}_1} \rightarrow \pi_* \mathcal{E} \rightarrow \underbrace{\pi_* \pi^* \mathcal{F}_2}_{=\mathcal{F}_2} \rightarrow \underbrace{R^1 \pi_* \pi^* \mathcal{F}_1}_{=\mathcal{F}_1 \otimes R^1 \pi_* \mathcal{O}_{X'} = 0} \rightarrow \dots,$$

which turns out to be a short exact sequence by the derived version of the projection formula. In particular, the vanishing $R^\nu \pi_* \pi^* \mathcal{F}_j = 0$ for $\nu \geq 1$ implies $R^\nu \pi_* \mathcal{E} = 0$, too. Hence, using both vanishings, we have for all $i \geq 0$ and for $j = 1, 2$ the isomorphisms

$$(4.4) \quad H^i(X', \mathcal{E}) \cong H^i(X, \pi_* \mathcal{E}) \text{ and } H^i(X', \pi^* \mathcal{F}_j) \cong H^i(X, \pi_* \pi^* \mathcal{F}_j) = H^i(X, \mathcal{F}_j).$$

In the study of $\text{Ext}(\Delta^-, \Delta^+) \cong H^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-)) \cong H^1(X', \mathcal{O}_{X'}(\Delta^+ - \Delta^-))$ we can therefore pull back and push forward along π without impacting the extension or cohomology classes and groups. In the following, we will assume for simplicity that the fan Σ is already refined enough so that all ∇_i are compatible with it.

4.1.2. *Two Components.* We dive into Theorem 12 in the case where $\Delta^- \setminus \Delta^+$ consists of two components C_0 and C_1 and we assume $\nabla_0 = C_0 \cup \Delta^+$ and $\nabla_1 = C_1 \cup \Delta^+$ to be compatible with the fan Σ . The complex \mathcal{C}_\bullet^{Fs} associated to $S = \{\nabla_0, \nabla_1\}$ is

$$(4.5) \quad 0 \rightarrow \mathbb{C} \cdot e_I \xrightarrow{\begin{pmatrix} -1 \\ 1 \end{pmatrix}} \mathbb{C} \cdot e_0 \oplus \mathbb{C} \cdot e_1 \xrightarrow{\begin{pmatrix} 1 & 1 \end{pmatrix}} \mathbb{C} \cdot e_\emptyset \rightarrow 0.$$

By Theorem 12 the $\mathcal{C}_\bullet^{Fs}(m)$ for $m \in M$ induce a T -equivariant exact sequence

$$(4.6) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+) \rightarrow \mathcal{O}_X(\nabla_0) \oplus \mathcal{O}_X(\nabla_1) \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0.$$

Theorem 15. *Via the identification $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0 = \tilde{H}^0(\Delta^- \setminus \Delta^+)$, the exact extension sequence (4.6) corresponds to the reduced singular 0-th cohomology class $[C_1] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$, which is equal to the class $-[C_0] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$.*

Proof. We follow the steps mentioned at the start of section 4. We first tensor the sequence (4.6) with $\mathcal{O}_X(\Delta^-)^{-1}$ and set $\mathcal{L} := \mathcal{O}_X(\Delta^+ - \Delta^-)$ and $\mathcal{E}_i := \mathcal{O}_X(\nabla_i - \Delta^-)$ for $i = 0, 1$ to obtain an extension sequence

$$(4.7) \quad 0 \rightarrow \mathcal{L} \rightarrow \mathcal{E}_0 \oplus \mathcal{E}_1 \rightarrow \mathcal{O}_X \rightarrow 0$$

in $\text{Ext}(\mathcal{O}_X, \mathcal{L})_0$. Consider the associated long exact sequence in cohomology:

$$(4.8) \quad 0 \rightarrow \Gamma(X, \mathcal{L}) \rightarrow \Gamma(X, \mathcal{E}_0) \oplus \Gamma(X, \mathcal{E}_1) \rightarrow \Gamma(X, \mathcal{O}_X) \xrightarrow[1 \mapsto \eta]{d} H^1(X, \mathcal{L}) \rightarrow \dots$$

The image of the extension sequence (4.6) in $H^1(X, \mathcal{L})$ under identification (2) is the image η of $1 \in \Gamma(X, \mathcal{O}_X)$ under d . Note that $\eta \in H^1(X, \mathcal{L})_0$ is of degree 0.

To understand identification (3) we translate $\eta \in H^1(X, \mathcal{L})$ to an element in the Čech cohomology group $\check{H}^1(\mathcal{U}, \mathcal{L})$ for $\mathcal{U} = \{U_{\sigma_i}\}_{\sigma_i \in \Sigma_{\max}}$ with $U_{\sigma_i} = \text{Spec } \mathbb{C}[\sigma_i^\vee \cap M]$ the standard toric affine open covering of X . The long exact sequence in cohomology (4.8) corresponds to a long exact sequence of Čech cohomology groups

$$(4.9) \quad 0 \rightarrow \check{H}^0(\mathcal{U}, \mathcal{L}) \rightarrow \check{H}^0(\mathcal{U}, \mathcal{E}_0) \oplus \check{H}^0(\mathcal{U}, \mathcal{E}_1) \rightarrow \check{H}^0(\mathcal{U}, \mathcal{O}_X) \xrightarrow[1 \mapsto \eta]{d} \check{H}^1(\mathcal{U}, \mathcal{L}) \rightarrow \dots$$

Order the maximal cones $\sigma \in \Sigma_{\max}$ so that $1 \in \mathcal{E}_0(U_i)$ for $i \in \{0, \dots, l\}$ and $1 \in \mathcal{E}_1(U_i)$ for $i \in \{l+1, \dots, m\}$. This is possible because the map of sheaves

$\mathcal{E}_0 \oplus \mathcal{E}_1 \rightarrow \mathcal{O}_X$ is surjective, each $U_i := U_{\sigma_i}$ is affine, and everything is T -equivariant. Note that this is not a dichotomy; it might happen that $1 \in \mathcal{E}_0(U_i) \cap \mathcal{E}_1(U_i)$ for some i . Using this order, the boundary homomorphism d the element $(1, \dots, 1) \in \prod_i \mathcal{O}_X(U_i)$ maps to the class $\eta \in \check{H}^1(\mathcal{U}, \mathcal{L})$ of

$$(4.10) \quad (\underbrace{0, \dots, 0}_{i < j \leq l}, \underbrace{1, \dots, 1}_{i \leq l < j}, \underbrace{0, \dots, 0}_{l < i < j}) \in \prod_{i < j} \mathcal{L}(U_{ij}).$$

Understanding η in terms of Čech cohomology is convenient because by [AP20] the degree 0 part of the Čech complex $\mathcal{C}^\bullet(\mathcal{U}, \mathcal{L})$ giving $\check{H}^\bullet(\mathcal{U}, \mathcal{L})$ is the same as a Čech complex $\mathcal{C}^\bullet(\Delta^-, \mathcal{S})$ defined in [AP20, (3.4)] giving the relative singular cohomology $H^\bullet(\Delta^-, \Delta^- \setminus \Delta^+)$. It is defined as $\mathcal{C}^p(\Delta^-, \mathcal{S}) := \prod_{i_0 < \dots < i_p} H^0(\Delta^-, S(\sigma_{i_0} \cap \dots \cap \sigma_{i_p}))$ for $S(\sigma) := \Delta^- \setminus (\Delta^+ + \sigma^\vee)$. Investigating the groups $H^0(U_\sigma, \mathcal{L})_0 = H^0(\Delta^-, S(\sigma))$ and $H^0(U_\sigma, \mathcal{E}_i)_0$, $i = 0, 1$, as in [AP20, (3.2)], one realizes that for $\sigma \in \Sigma_{\max}$ with $S(\sigma) \neq \emptyset$, $1 \in \mathcal{E}_0(U_\sigma)$ implies $S(\sigma) \subseteq C_0$ and $1 \in \mathcal{E}_1(U_\sigma)$ implies $S(\sigma) \subseteq C_1$. Here C_0 and C_1 denote the two connected components of $\Delta^- \setminus \Delta^+$.

The element $\eta \in \check{H}^1(\mathcal{U}, \mathcal{L})_0 \cong H^1(\Delta^-, \Delta^- \setminus \Delta^+)$ can be lifted to $H^0(\Delta^- \setminus \Delta^+)$ by considering the long exact sequence of the pair $(\Delta^-, \Delta^- \setminus \Delta^+)$. In terms of Čech complexes the boundary morphism $H^0(\Delta^- \setminus \Delta^+) \xrightarrow{d} H^1(\Delta^-, \Delta^- \setminus \Delta^+)$ is given by the snake lemma. One shows that η can be lifted to the cocycle which is 1 on the component C_1 of $\Delta^- \setminus \Delta^+$ and 0 on the component C_0 . It can also be lifted to the cocycle which is 0 on the component C_1 and -1 on the component C_0 . Modulo $H^0(\Delta^-)$ these cocycles are equivalent and we have found the image of sequence (4.6) to be $[C_1] = -[C_0] \in \check{H}^0(\Delta^- \setminus \Delta^+)$. \square

4.1.3. More than Two Components. We now deal with the case where $\Delta^- \setminus \Delta^+$ consists of $n+1$ connected components C_0, \dots, C_n for some $n \geq 2$. Set $\nabla_i := C_i \cup \Delta^+$ for $i = 0, \dots, n$ and note that $\bigcup_{i=0}^n \nabla_i = \Delta^-$ and $\nabla_{I'} = \bigcap_{i \in I'} \nabla_i = \Delta^+$ for $I' \subseteq I$ of cardinality k with $2 \leq k \leq n+1$. By subsection (4.1.1) we may assume that the set $S := \{\nabla_i \mid i \in \{0, \dots, n\}\}$ forms a Σ -family. The exact sequence of sheaves induced from this Σ -family as in Theorem 12 looks as follows:

$$(4.11) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+) \rightarrow \dots \rightarrow \mathcal{O}_X(\Delta^+)^{\binom{n+1}{2}} \rightarrow \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i) \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0.$$

We will replace this sequence (4.11) by a quasi-isomorphic short exact sequence. To this end, consider the exact Koszul complex $\bigwedge^\bullet \mathbb{C}^{n+1} \otimes \mathcal{O}_X(\Delta^+)$:

$$(4.12) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+) \xrightarrow{\bar{d}_{n+1}} \dots \xrightarrow{\bar{d}_3} \mathcal{O}_X(\Delta^+)^{\binom{n+1}{2}} \xrightarrow{\bar{d}_2} \mathcal{O}_X(\Delta^+)^{n+1} \xrightarrow{\bar{d}_1} \mathcal{O}_X(\Delta^+) \rightarrow 0,$$

where $\bar{d}_i = d_i \otimes \text{id}_{\mathcal{O}_X(\Delta^+)}$ for d_i the i -th differential in $\bigwedge^\bullet \mathbb{C}^{n+1}$. The green part is identical to part of sequence (4.11). We obtain a quasi-isomorphism from sequence (4.11):

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{O}_X(\Delta^+) & \longrightarrow & \dots & \longrightarrow & \mathcal{O}_X(\Delta^+)^{\binom{n+1}{2}} & \longrightarrow & \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i) & \longrightarrow & \mathcal{O}_X(\Delta^-) & \longrightarrow & 0 \\ & & \downarrow & & & & \downarrow & & \parallel & & \parallel & & & \\ 0 & \longrightarrow & 0 & \longrightarrow & \dots & \longrightarrow & \ker(\bar{d}_1) & \longrightarrow & \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i) & \longrightarrow & \mathcal{O}_X(\Delta^-) & \longrightarrow & 0. \end{array}$$

Denoting $K := \ker(\bar{d}_1) = \ker(d_1) \otimes \mathcal{O}_X(\Delta^+)$, we obtain a short exact sequence

$$(4.13) \quad 0 \rightarrow K \rightarrow \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i) \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0.$$

We now choose the set $\{e_i - e_0 \mid i \in \{1, \dots, n\}\}$ with respect to the standard basis $\{e_0, \dots, e_n\}$ for \mathbb{C}^{n+1} as a basis for $\ker(d_1)$. This induces an isomorphism $K \cong \mathcal{O}_X(\Delta^+)^n$ under which sequence (4.13) corresponds to the short exact sequence

$$(4.14) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+)^n \rightarrow \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i) \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0,$$

where the maps can be thought of as

$$(4.15) \quad \mathcal{O}_X(\Delta^+)^n \xrightarrow{\begin{pmatrix} -1 & -1 & \dots & -1 \\ 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & 1 \end{pmatrix}} \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i),$$

$$(4.16) \quad \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i) \xrightarrow{\begin{pmatrix} 1 & 1 & \dots & 1 \end{pmatrix}} \mathcal{O}_X(\Delta^-).$$

Theorem 16. *Via the identification $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+)^n)_0 = \tilde{H}^0(\Delta^- \setminus \Delta^+)^n$, the extension sequence (4.14) corresponds to the n -tuple $([C_1], \dots, [C_n]) \in \tilde{H}^0(\Delta^- \setminus \Delta^+)^n$ of reduced singular 0-th cohomology classes.*

Proof. The steps in this case are very similar to the case of two components. The extension sequence

$$0 \rightarrow \mathcal{L}^n \rightarrow \bigoplus_{i=0}^n \mathcal{E}_i \rightarrow \mathcal{O}_X \rightarrow 0$$

corresponds to the image η of $1 \in \Gamma(X, \mathcal{O}_X)$ in $H^1(X, \mathcal{L}^n) = H^1(X, \mathcal{L})^n$ under the long exact sequence in cohomology associated to the short exact extension sequence. In our case $\mathcal{L} = \mathcal{O}_X(\Delta^+ - \Delta^-)$ and $\mathcal{E}_i = \mathcal{O}_X(\nabla_i - \Delta^-)$. Again, this image is

understood using the sequence of Čech complexes, only looking at degree 0:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \prod_i \mathcal{L}(U_i)^n & \longrightarrow & \prod_i (\bigoplus_{s=0}^n \mathcal{E}_s(U_i)) & \longrightarrow & \prod_i \mathcal{O}_X(U_i) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \prod_{i<j} \mathcal{L}(U_{ij})^n & \longrightarrow & \prod_{i<j} (\bigoplus_{s=0}^n \mathcal{E}_s(U_{ij})) & \longrightarrow & \prod_{i<j} \mathcal{O}_X(U_{ij}) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \prod_{i<j<k} \mathcal{L}(U_{ijk})^n & \longrightarrow & \prod_{i<j<k} (\bigoplus_{s=0}^n \mathcal{E}_s(U_{ijk})) & \longrightarrow & \prod_{i<j<k} \mathcal{O}_X(U_{ijk}) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \vdots & & \vdots & & \vdots
\end{array}$$

We lift the element $(1, \dots, 1) \in \prod_i \mathcal{O}_X(U_i)$ to $\prod_i (\bigoplus_{s=0}^n \mathcal{E}_s(U_i))$, then map it to $\prod_{i<j} (\bigoplus_{s=0}^n \mathcal{E}_s(U_{ij}))$ via the Čech differential and lift it to $\prod_{i<j} \mathcal{L}(U_{ij})^n$. The resulting element $(\eta_{ij})_{i<j} \in \prod_{i<j} \mathcal{L}(U_{ij})^n$ can be viewed as an element of $\prod_{i<j} H^0(\Delta^-, S(\sigma_i \cap \sigma_j))^n$. The preimage of $[\eta]$ under the map d^n in the long exact sequence

$$0 \rightarrow H^0(\Delta^-, \Delta^- \setminus \Delta^+)^n \rightarrow H^0(\Delta^-)^n \rightarrow H^0(\Delta^- \setminus \Delta^+)^n \xrightarrow{d^n} H^1(\Delta^-, \Delta^- \setminus \Delta^+)^n \rightarrow 0.$$

is the desired element $([C_1], \dots, [C_n]) \in \tilde{H}^0(\Delta^- \setminus \Delta^+)^n$. \square

Recall that $\text{Ext}^1(\mathcal{O}_X(\Delta^-), -)$ is a covariant functor, where a map of extensions is induced by a pushout.

Corollary 17. *Given the extension sequence (4.14) in $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+)^n)_0$, we obtain n extensions in $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$, one for each $i = 1, \dots, n$:*

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathcal{O}_X(\Delta^+)^n & \longrightarrow & \bigoplus_{i=0}^n \mathcal{O}(\nabla_i) & \longrightarrow & \mathcal{O}_X(\Delta^-) \longrightarrow 0 \\
& & \downarrow \text{pr}_i & & \downarrow & & \parallel \\
0 & \longrightarrow & \mathcal{O}_X(\Delta^+) & \longrightarrow & \mathcal{H}_i & \longrightarrow & \mathcal{O}_X(\Delta^-) \longrightarrow 0,
\end{array}$$

where \mathcal{H}_i is the pushout of the left square.

Via the identification $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0 = \tilde{H}^0(\Delta^- \setminus \Delta^+)$, the i -th extension

$$(4.17) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+) \rightarrow \mathcal{H}_i \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0$$

corresponds to the class $[C_i] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$. In particular, the n extensions for $i \in \{1, \dots, n\}$ form a basis of $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$.

Proof. This follows from functoriality of $\text{Ext}(\mathcal{O}_X(\Delta^-), -)$ and naturality of the isomorphisms we identify along. The i -th projection $\mathcal{O}_X(\Delta^+)^n \rightarrow \mathcal{O}_X(\Delta^+)$ induces

the i -th projection $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0^n \rightarrow \text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$. By the previous Theorem 16 this yields $[C_i] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$. \square

Remark 18. To obtain $[C_0]$ one has to replace pr_i by $(-1, \dots, -1)$.


4.1.4. *The Universal Extension.* For two \mathcal{O}_X -modules \mathcal{F} and \mathcal{G} and their (finite-dimensional) space of extensions $E := \text{Ext}(\mathcal{F}, \mathcal{G})$, a *universal extension* is a short exact sequence of the form $0 \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow \mathcal{F} \otimes E \rightarrow 0$ such that for any $t \in E$ the induced pullback sequence is $t \in E = \text{Ext}(\mathcal{F}, \mathcal{G})$. Equivalently, it is a short exact sequence $0 \rightarrow \mathcal{G} \otimes E^\vee \rightarrow \mathcal{H}' \rightarrow \mathcal{F} \rightarrow 0$, where $E^\vee = \text{Hom}_{\mathbb{C}}(E, \mathbb{C})$, such that for any $t \in E$ the sequence induced by the pushout along $t \in E \cong \text{Hom}_{\mathbb{C}}(E^\vee, \mathbb{C})$ is $t \in E = \text{Ext}(\mathcal{F}, \mathcal{G})$. Analogously one defines a *universal T -equivariant extension* $0 \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow \mathcal{F} \otimes E_0 \rightarrow 0$ or $0 \rightarrow \mathcal{G} \otimes E_0^\vee \rightarrow \mathcal{H}' \rightarrow \mathcal{F} \rightarrow 0$ for $E_0 := \text{Ext}(\mathcal{F}, \mathcal{G})_0$.

Theorem 19. *The extension sequence (4.13) in $\text{Ext}(\mathcal{O}_X(\Delta^-), K)_0$ is a universal extension for $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$.*


Proof. The \mathbb{C} -vector space $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0 \cong \tilde{H}^0(\Delta^- \setminus \Delta^+)$ is the cokernel of the homomorphism $H^0(\{\cdot\}) \rightarrow H^0(\Delta^- \setminus \Delta^+)$. The components C_0, \dots, C_n provide a natural basis for $H^0(\Delta^- \setminus \Delta^+)$ with dual basis $C_0^\vee, \dots, C_n^\vee$ of $H^0(\Delta^- \setminus \Delta^+)^\vee$. The quotient $\tilde{H}^0(\Delta^- \setminus \Delta^+) = \frac{H^0(\Delta^- \setminus \Delta^+)}{H^0(\{\cdot\})}$ is generated by $[C_0], \dots, [C_n]$ and subject to the relation $[C_0] + \dots + [C_n] = 0$. We choose $\{C_1^\vee - C_0^\vee, \dots, C_n^\vee - C_0^\vee\}$ as a basis for the dual vector space $\tilde{H}^0(\Delta^- \setminus \Delta^+)^\vee = \ker(H^0(\Delta^- \setminus \Delta^+)^\vee \rightarrow H^0(\{\cdot\})^\vee)$. This basis is dual to the basis $\{[C_1], \dots, [C_n]\}$ of $\tilde{H}^0(\Delta^- \setminus \Delta^+)$. The isomorphism $\tilde{H}^0(\Delta^- \setminus \Delta^+)^\vee \xrightarrow{\cong} \ker(d_1)$ maps the i -th basis element $C_i^\vee - C_0^\vee$ to the i -th basis element $e_i - e_0$. Recall that $K = \ker(d_1) \otimes \mathcal{O}_X(\Delta^+)$.

We now check the universal property on the basis $\{[C_1], \dots, [C_n]\}$ of $\tilde{H}^0(\Delta^- \setminus \Delta^+)$. By definition of the dual and double dual, $[C_i] \in \tilde{H}^0(\Delta^- \setminus \Delta^+) \cong \text{Hom}_{\mathbb{C}}(\tilde{H}^0(\Delta^- \setminus \Delta^+)^\vee, \mathbb{C})$ corresponds to the projection to the i -th coordinate $\text{pr}_i: \tilde{H}^0(\Delta^- \setminus \Delta^+)^\vee \cong \mathbb{C}^n \rightarrow \mathbb{C}$. By Corollary 17 the pushout along the i -th projection homomorphism gives an extension in $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0 \cong \tilde{H}^0(\Delta^- \setminus \Delta^+)$ which corresponds precisely to the basis element $[C_i] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$. \square

4.1.5. *The Cremona Example.* Let X be the graph of the Cremona transformation $\mathbb{P}^2 \dashrightarrow \mathbb{P}^2$. In toric language, $X = \mathbb{P}(H) = \mathbb{T}\mathbb{V}(\Sigma)$, where $\Sigma = \mathcal{N}(H)$:




Here, the ray ρ_i of the inner normal fan Σ corresponds to the facet F_i of H . Let $D_i := \text{orb}(\rho_i)$. The Minkowski summands A and B of H correspond to the divisors $D_A = D_2 + D_3 + D_4$ and $D_B = D_1 + D_5 + D_6$. The interesting line bundle on X arises from $\Delta^+ = A$ and $\Delta^- = 2B + (1, -1)$ with $D_{\Delta^-} = D_1 + D_2 + 2D_3 + D_4 + D_5$. The figure below displays the set-theoretic difference $(2B + (1, -1)) \setminus A$. It shows that $H^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-)) \cong \bigoplus_{m \in M} \tilde{H}^0(\Delta^- \setminus (\Delta^+ + m))$ is 2-dimensional in degree $m = 0$, because $\dim \tilde{H}^0(\Delta^- \setminus \Delta^+) = \dim \tilde{H}^0((2B + (1, -1)) \setminus A) = 2$.



There is no other shift m such that $(2B - m) \setminus A$, or equivalently $2B \setminus (A + m)$, has non-trivial reduced 0-th cohomology. Hence, $H^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-))$ is 2-dimensional, sitting completely in degree $m = 0$.

Denote the connected components of $\Delta^- \setminus \Delta^+$ by C_0, C_1, C_2 and set $\nabla_i := C_i \cup \Delta^+$.



From the facet presentations of ∇_0, ∇_1 and ∇_2 we can read off the associated divisors $D_{\nabla_0} = D_1 + D_2 + D_3 + D_4$, $D_{\nabla_1} = D_2 + 2D_3 + D_4$, $D_{\nabla_2} = D_2 + D_3 + D_4 + D_5$. In this case, sequence (4.14) in subsection (4.1.3) is the sequence


$$(4.18) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+)^2 \xrightarrow{\begin{pmatrix} -1 & -1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}} \bigoplus_{i=0}^2 \mathcal{O}_X(\nabla_i) \xrightarrow{\begin{pmatrix} 1 & 1 & 1 \end{pmatrix}} \mathcal{O}_X(\Delta^-) \rightarrow 0$$

in $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+)^2)_0$. It corresponds to $([C_1], [C_2]) \in \tilde{H}^0(\Delta^- \setminus \Delta^+)^2$.

The pushout construction described in Corollary 17 yields two extension sequences $0 \rightarrow \mathcal{O}_X(\Delta^+) \rightarrow \mathcal{G}_1 \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0$ and $0 \rightarrow \mathcal{O}_X(\Delta^+) \rightarrow \mathcal{G}_2 \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0$ in $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$. The first is represented by $[C_1] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$ and the second by $[C_2] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$. The class of the constant cocycle on $\Delta^- \setminus \Delta^+$ is trivial, so $[C_0] = -[C_1] - [C_2] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$ and the negative of the Baer sum of the two sequences above is represented by $[C_0] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$.

4.2. General Position of Polyhedra. In the previous section we dealt with the special case of one polyhedron Δ^+ being contained in the other polyhedron Δ^- . In this section we deal with Δ^+ and Δ^- lying in general position.


4.2.1. Observing Two Problems. To illustrate the problem for the general case, we start with an example. Let Σ_0 be the 2-dimensional, singular fan made from the rays spanned by $(1, 0)$, $(0, 1)$, $(-2, -1)$, and $(0, -1)$, respectively (displayed in black below). Adding the rays spanned by $(-1, 0)$ and $(-1, -1)$ (displayed in blue below), we obtain the smooth subdivision Σ .



Denote $X := \mathbb{T}\mathbb{V}(\Sigma)$ and $f: X \rightarrow X_0 := \mathbb{T}\mathbb{V}(\Sigma_0)$, the birational contraction. The Picard group $\text{Pic}(X_0)$ is freely generated by (the sheaves represented by) the polytopes C and D displayed below. $\text{Pic}(X) = \text{Cl}(X)$ has $\{A, B, C, D\}$ as a basis:



Define $\Delta^- := D$ and $\Delta^+ := C$. The difference $\Delta^- \setminus \Delta^+$ splits into two components:



However, since Δ^+ is not contained in Δ^- , there is no map from $\mathcal{O}_X(\Delta^+)$ into the sheaves of the two components, which are subsheaves of $\mathcal{O}_X(\Delta^-)$. The solution to this problem is to replace Δ^+ by $(\Delta^+ \cap \Delta^-)$, then proceed as in section (4.1), and, finally, to use functoriality of Ext along the embedding $\mathcal{O}_X(\Delta^+ \cap \Delta^-) \hookrightarrow \mathcal{O}_X(\Delta^+)$. But here one can spot the second problem. As in the example, the intersection

$\Delta^+ \cap \Delta^-$ is not necessarily a lattice polyhedron. This will be overcome by refining the lattice. We replace the lattice M by a larger lattice $\widetilde{M} \supseteq M$ such that $\Delta^+ \cap \Delta^-$ is a lattice polyhedron with respect to \widetilde{M} . Dually, this means to consider some sublattice $\widetilde{N} \subseteq N$ of finite index and the induced finite covering $p: \mathbb{T}\mathbb{V}(\Sigma, \widetilde{N}) \rightarrow \mathbb{T}\mathbb{V}(\Sigma, N)$.

4.2.2. The Intersection is a Lattice Polyhedron. We start with the case where the intersection $\Delta^+ \cap \Delta^-$ is a lattice polyhedron with respect to the lattice M . Suppose that $\Delta^- \setminus \Delta^+ = \Delta^- \setminus (\Delta^+ \cap \Delta^-)$ consists of $n+1$ connected components C_0, \dots, C_n . Set $\nabla_i := C_i \cup (\Delta^+ \cap \Delta^-)$ for $i \in \{0, \dots, n\}$. As before, we can assume without loss of generality that the fan Σ is refined enough so that Δ^+ , Δ^- , $\Delta^+ \cap \Delta^-$ and ∇_i for $i \in \{0, \dots, n\}$ are compatible with Σ and define nef Cartier divisors on $X = \mathbb{T}\mathbb{V}(\Sigma)$.

Consider the following pushout diagram:

$$(4.19) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X(\Delta^+ \cap \Delta^-)^n & \longrightarrow & \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i) & \longrightarrow & \mathcal{O}_X(\Delta^-) \longrightarrow 0 \\ & & \downarrow h^n & & \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{O}_X(\Delta^+)^n & \longrightarrow & \mathcal{H} & \longrightarrow & \mathcal{O}_X(\Delta^-) \longrightarrow 0. \end{array}$$

Here the upper exact sequence is constructed as in subsection (4.1.3). It is an extension sequence in $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+ \cap \Delta^-)^n)_0$. By Theorem 16 this corresponds to the n -tuple $([C_1], \dots, [C_n]) \in \widetilde{H}^0(\Delta^- \setminus (\Delta^+ \cap \Delta^-))^n = \widetilde{H}^0(\Delta^- \setminus \Delta^+)^n$. The inclusion of lattice polyhedra $\Delta^+ \cap \Delta^- \hookrightarrow \Delta^+$ induces the embedding of sheaves $h: \mathcal{O}_X(\Delta^+ \cap \Delta^-) \hookrightarrow \mathcal{O}_X(\Delta^+)$ and its n -th power $h^n: \mathcal{O}_X(\Delta^+ \cap \Delta^-)^n \hookrightarrow \mathcal{O}_X(\Delta^+)^n$. The sheaf \mathcal{H} is the pushout of the left square and the universal property of the pushout induces a map $\mathcal{H} \rightarrow \mathcal{O}_X(\Delta^-)$, that makes the lower sequence exact and the right square commutative. All together we obtain a map of complexes from the upper to the lower exact sequence. This is functoriality of $\text{Ext}(\mathcal{O}_X(\Delta^-), -)$.

Proposition 20. *Via the identification $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+)^n)_0 = \widetilde{H}^0(\Delta^- \setminus \Delta^+)^n$ the short exact extension sequence*

$$(4.20) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+)^n \rightarrow \mathcal{H} \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0$$

corresponds to the n -tuple $([C_1], \dots, [C_n]) \in \widetilde{H}^0(\Delta^- \setminus \Delta^+)^n$.

Proof. Tensoring both sequences in diagram (4.19) with $\mathcal{O}_X(\Delta^-)^{-1}$ yields

$$(4.21) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X((\Delta^+ \cap \Delta^-) - \Delta^-)^n & \longrightarrow & \bigoplus_{i=0}^n \mathcal{O}_X(\nabla_i - \Delta^-) & \longrightarrow & \mathcal{O}_X \longrightarrow 0 \\ & & \downarrow (h')^n & & \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{O}_X(\Delta^+ - \Delta^-)^n & \longrightarrow & \mathcal{H}' & \longrightarrow & \mathcal{O}_X \longrightarrow 0, \end{array}$$

where $\mathcal{H}' = \mathcal{H} \otimes \mathcal{O}_X(\Delta^-)^{-1}$ is also the pushout of the left square. The map of complexes between the short exact sequences induces a map of complexes between

the long exact sequences in cohomology. In particular we obtain a commuting square

$$(4.22) \quad \begin{array}{ccc} \Gamma(\mathcal{O}_X) & \xrightarrow{d} & \mathrm{H}^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-))^n & \ni & \mu \\ \parallel & & \cong \downarrow \mathrm{H}^1((h')^n) & & \downarrow \\ \Gamma(\mathcal{O}_X) & \xrightarrow{d} & \mathrm{H}^1(X, \mathcal{O}_X((\Delta^+ \cap \Delta^-) - \Delta^-))^n & \ni & \eta, \end{array}$$

where η and μ denote the images of $1 \in \Gamma(X, \mathcal{O}_X)_0$ under the differential d . By Theorem 16 the isomorphism from $\mathrm{H}^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-))^n$ to reduced singular cohomology maps η to $([C_1], \dots, [C_n]) \in \tilde{\mathrm{H}}^0(\Delta^- \setminus (\Delta^+ \cap \Delta^-))^n = \tilde{\mathrm{H}}^0(\Delta^- \setminus \Delta^+)^n$. It is easy to see that $\mu = \mathrm{H}^1((h')^n)^{-1}(\eta)$ is also mapped to $([C_1], \dots, [C_n]) \in \tilde{\mathrm{H}}^0(\Delta^- \setminus \Delta^+)^n$. \square

4.2.3. The Intersection is Not a Lattice Polyhedron. We now deal with the case where $\Delta^+ \cap \Delta^- \subseteq M_{\mathbb{R}}$ is not a lattice polyhedron with respect to the lattice M . Take any refinement $\tilde{M} \supseteq M$ such that M is a sublattice of finite index in \tilde{M} and $\Delta^+ \cap \Delta^- \subseteq \tilde{M}_{\mathbb{R}} = M_{\mathbb{R}} \cong \mathbb{R}^r$ is a lattice polyhedron with respect to \tilde{M} . Dually, $\tilde{N} \subseteq N \cong \mathbb{Z}^r$ is a sublattice of finite index. Let $G := N/\tilde{N}$ be the finite quotient group. For a toric variety $X = \mathrm{TV}(\Sigma, N)$ we can consider the fan Σ in $N_{\mathbb{R}} = \tilde{N}_{\mathbb{R}}$ with respect to the lattice \tilde{N} and obtain a second toric variety $\tilde{X} = \mathrm{TV}(\Sigma, \tilde{N})$ realizing X as the geometric quotient $X = \tilde{X}/G$. The lattice inclusion $\iota: \tilde{N} \hookrightarrow N$ induces the toric covering morphism $p: \tilde{X} \rightarrow X$ [CLS11, Prop. 3.3.7]. We pull the sheaves $\mathcal{O}_X(\Delta^+)$ and $\mathcal{O}_X(\Delta^-)$ back to \tilde{X} via p and use the results from section (4.2.2).

Example 21. For $M = \mathbb{Z}^r$ and its dual $N = \mathbb{Z}^r$ take \tilde{M} to be $(\frac{1}{d}\mathbb{Z})^r$ and correspondingly $\tilde{N} = (d\mathbb{Z})^r$. Then each ray generator $\tilde{v}_\rho \in \tilde{N}$ of a ray $\rho \in \Sigma(1)$ is the d -multiple of the corresponding ray generator $v_\rho \in N$. The coefficients of the pullback p^*D of a Weil divisor D on X will be d -multiples of the coefficients of D .

Construction 22. Back to our general case, the refinement $\tilde{M} \supseteq M$ was chosen so that $\Delta^+ \cap \Delta^-$ is a lattice polyhedron with respect to \tilde{M} and we are in the case of section (4.2.2), however on the space \tilde{X} . We have the short exact extension sequence

$$(4.23) \quad 0 \rightarrow \mathcal{O}_{\tilde{X}}(\Delta^+ \cap \Delta^-)^n \rightarrow \bigoplus_{i=0}^n \mathcal{O}_{\tilde{X}}(\nabla_i) \rightarrow \mathcal{O}_{\tilde{X}}(\Delta^-) \rightarrow 0$$

and the short exact extension sequence

$$(4.24) \quad 0 \rightarrow \mathcal{O}_{\tilde{X}}(\Delta^+)^n \rightarrow \tilde{\mathcal{H}} \rightarrow \mathcal{O}_{\tilde{X}}(\Delta^-) \rightarrow 0$$

induced by the embedding $\mathcal{O}_{\tilde{X}}(\Delta^+ \cap \Delta^-)^n \hookrightarrow \mathcal{O}_{\tilde{X}}(\Delta^+)^n$. Under the identification with $\tilde{\mathrm{H}}^0(\Delta^- \setminus \Delta^+)^n$, both correspond to the n -tuple $([C_1], \dots, [C_n]) \in \tilde{\mathrm{H}}^0(\Delta^- \setminus \Delta^+)^n$ by Theorem 16 and Proposition 20. Since $p: \tilde{X} \rightarrow X$ is affine, the pushforward p_* is exact and we obtain a short exact extension sequence of sheaves on X :

$$(4.25) \quad 0 \rightarrow p_*\mathcal{O}_{\tilde{X}}(\Delta^+)^n \rightarrow p_*\tilde{\mathcal{H}} \rightarrow p_*\mathcal{O}_{\tilde{X}}(\Delta^-) \rightarrow 0.$$

Recall that in the first step of the identification of $\text{Ext}(\mathcal{O}_{\tilde{X}}(\Delta^-), \mathcal{O}_{\tilde{X}}(\Delta^+)^n)_0$ with $\tilde{H}^0(\Delta^- \setminus \Delta^+)^n$ we tensor sequence (4.24) with $\mathcal{O}_{\tilde{X}}(\Delta^-)^{-1}$. Its pushforward is

$$(4.26) \quad 0 \rightarrow p_*(\mathcal{O}_{\tilde{X}}(\Delta^+ - \Delta^-))^n \rightarrow p_*(\tilde{\mathcal{H}}^!) \rightarrow p_*\mathcal{O}_{\tilde{X}} \rightarrow 0.$$

By the projection formula this is equal to sequence (4.25) tensored with $\mathcal{O}_X(\Delta^-)^{-1}$. For an affine morphism $p: \tilde{X} \rightarrow X$ of noetherian separated schemes and a short exact sequence $0 \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0$ of quasi-coherent sheaves on \tilde{X} there is a commuting diagram in which all vertical morphisms are isomorphisms:

$$\begin{array}{ccccccccc} \Gamma(\tilde{X}, \mathcal{F}) & \longrightarrow & \Gamma(\tilde{X}, \mathcal{G}) & \longrightarrow & \Gamma(\tilde{X}, \mathcal{H}) & \xrightarrow{\partial} & H^1(\tilde{X}, \mathcal{F}) & \longrightarrow & H^1(\tilde{X}, \mathcal{G}) & \longrightarrow & \dots \\ \downarrow \cong & & \\ \Gamma(X, p_*\mathcal{F}) & \longrightarrow & \Gamma(X, p_*\mathcal{G}) & \longrightarrow & \Gamma(X, p_*\mathcal{H}) & \xrightarrow{\partial} & H^1(X, p_*\mathcal{F}) & \longrightarrow & H^1(X, p_*\mathcal{G}) & \longrightarrow & \dots \end{array}$$

In particular, $H^1(X, p_*\mathcal{O}_{\tilde{X}}(\Delta^+ - \Delta^-))^n = H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}(\Delta^+ - \Delta^-))^n$ and the image of $1 \in \Gamma(X, p_*\mathcal{O}_{\tilde{X}})$ under ∂ is the desired n -tuple $([C_1], \dots, [C_n])$.

While $p_*\mathcal{O}_{\tilde{X}}(\Delta^+) = p_*p^*\mathcal{O}_X(\Delta^+)$ and $p_*\mathcal{O}_{\tilde{X}}(\Delta^-) = p_*p^*\mathcal{O}_X(\Delta^-)$ are in general not equal to $\mathcal{O}_X(\Delta^+)$ and $\mathcal{O}_X(\Delta^-)$, respectively, the latter are a direct summand:

$$(4.27) \quad \mathcal{O}_X(\Delta^\pm) = \bigoplus_{m \in M} \mathcal{O}_X(\Delta^\pm)_m = \bigoplus_{m \in M} (p_*p^*\mathcal{O}_X(\Delta^\pm))_m \subseteq \bigoplus_{\tilde{m} \in \tilde{M}} (p_*p^*\mathcal{O}_X(\Delta^\pm))_{\tilde{m}}.$$

Furthermore, $\mathcal{O}_X(\Delta^\pm)$ corresponds precisely to the G -invariants of $p_*p^*\mathcal{O}_X(\Delta^\pm)$.

Example 23. For $\mathcal{O}_X(\Delta^\pm)$ and the pullback $p^*\mathcal{O}_X(\Delta^\pm) = \mathcal{O}_{\tilde{X}}(\Delta^\pm)$ we have

$$(4.28) \quad \mathcal{O}_X(\Delta^\pm)(U_\sigma) = \chi^{m_\sigma^\pm} \cdot \mathbb{C}[\sigma^\vee \cap M] \subseteq \chi^{m_\sigma^\pm} \cdot \mathbb{C}[\sigma^\vee \cap \tilde{M}] = p_*\mathcal{O}_{\tilde{X}}(\Delta^\pm)(U_\sigma),$$

where $\{m_\sigma^\pm\}_{\sigma \in \Sigma}$ is the Cartier data of D_{Δ^\pm} . All m_σ^\pm are contained in $M \subseteq \tilde{M}$. Via $\mathbb{C}[\sigma^\vee \cap M] \hookrightarrow \mathbb{C}[\sigma^\vee \cap \tilde{M}]$ we can view the above as an inclusion of $\mathbb{C}[\sigma^\vee \cap M]$ -modules. Any M -graded module obtains an \tilde{M} -grading by adding zeros in degrees $\tilde{m} \in \tilde{M} \setminus M$.

For the sheaves in the short exact sequence (4.25) we have the \tilde{M} -graded subsheaves $\mathcal{O}_X(\Delta^+)$ of $p_*\mathcal{O}_{\tilde{X}}(\Delta^+)$ and $\mathcal{O}_X(\Delta^-)$ of $p_*\mathcal{O}_{\tilde{X}}(\Delta^-)$. The inclusions are isomorphisms when restricted to $M \subseteq \tilde{M}$. Because the morphisms in sequence (4.25) are homogeneous of degree 0, taking the M -graded part of $p_*\tilde{\mathcal{H}}$ yields a subsheaf $\mathcal{H} := \bigoplus_{m \in M} (p_*\tilde{\mathcal{H}})_m \subseteq p_*\tilde{\mathcal{H}}$ that fits into a commuting diagram of exact sequences

$$(4.29) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X(\Delta^+)^n & \longrightarrow & \mathcal{H} & \longrightarrow & \mathcal{O}_X(\Delta^-) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & p_*\mathcal{O}_{\tilde{X}}(\Delta^+)^n & \longrightarrow & p_*\tilde{\mathcal{H}} & \longrightarrow & p_*\mathcal{O}_{\tilde{X}}(\Delta^-) \longrightarrow 0. \end{array}$$

Proposition 24. *Given the commuting diagram (4.29), the upper sequence*

$$(4.30) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+)^n \rightarrow \mathcal{H} \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0$$

corresponds to the n -tuple $([C_1], \dots, [C_n]) \in \tilde{H}^0(\Delta^- \setminus \Delta^+)^n$ under the identification $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+)^n)_0 = \tilde{H}^0(\Delta^- \setminus \Delta^+)^n$.

Proof. Tensoring diagram (4.29) with $\mathcal{O}_X(\Delta^-)^{-1}$ yields the commutative diagram

$$(4.31) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X(\Delta^+ - \Delta^-)^n & \longrightarrow & \mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{O}_X(\Delta^-)^{-1} & \longrightarrow & \mathcal{O}_X \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & p_*(\mathcal{O}_{\tilde{X}}(\Delta^+ - \Delta^-))^n & \longrightarrow & p_*\tilde{\mathcal{H}}' & \longrightarrow & p_*\mathcal{O}_{\tilde{X}} \longrightarrow 0, \end{array}$$

the lower sequence inducing the desired n -tuple by Construction 22. We obtain a commutative diagram of long exact sequences

$$(4.32) \quad \begin{array}{ccccccc} \dots & \longrightarrow & \Gamma(X, \mathcal{O}_X) & \longrightarrow & H^1(X, \mathcal{O}_X(\Delta^+ - \Delta^-))^n & \longrightarrow & \dots \\ & & \downarrow (1) & & \downarrow (2) & & \\ \dots & \longrightarrow & \Gamma(X, p_*\mathcal{O}_{\tilde{X}}) & \longrightarrow & H^1(X, p_*(\mathcal{O}_{\tilde{X}}(\Delta^+ - \Delta^-)))^n & \longrightarrow & \dots \end{array}$$

The homomorphisms (1) and (2) restrict to isomorphisms in degree $0 \in M$, so $1 \in \Gamma(X, \mathcal{O}_X)_0 = \Gamma(X, p_*\mathcal{O}_{\tilde{X}})_0$ also maps to $([C_1], \dots, [C_n]) \in \tilde{H}^0(\Delta^- \setminus \Delta^+)^n$. \square

Corollary 25. *Sequence (4.30) induces n extensions in $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$, one for each $i = 1, \dots, n$:*

$$(4.33) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X(\Delta^+)^n & \longrightarrow & \mathcal{H} & \longrightarrow & \mathcal{O}_X(\Delta^-) \longrightarrow 0 \\ & & \downarrow pr_i & & \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{O}_X(\Delta^+) & \longrightarrow & \mathcal{H}_i & \longrightarrow & \mathcal{O}_X(\Delta^-) \longrightarrow 0. \end{array}$$

Via the identification $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0 = \tilde{H}^0(\Delta^- \setminus \Delta^+)$, the i -th extension corresponds to the class $[C_i] \in \tilde{H}^0(\Delta^- \setminus \Delta^+)$. In particular, the n extensions for $i \in \{1, \dots, n\}$ form a basis of $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$.

Furthermore, sequence (4.30) is a universal extension for $\text{Ext}(\mathcal{O}_X(\Delta^-), \mathcal{O}_X(\Delta^+))_0$.

5. USING KLYACHKO'S DESCRIPTION OF TORIC REFLEXIVE SHEAVES

We briefly show how to construct the universal extension sequence in the case where $\Delta^+ \cap \Delta^-$ is not a lattice polyhedron in terms of Klyachko's description of toric reflexive sheaves (see [Kly90] and [Kly02]; a short summary can be found in [Pay08]; see [DJS14] and [KM] for more recent approaches).

5.1. Describing Sheaves via Filtrations. Consider a toric variety $X = \mathbb{T}\mathbb{V}(\Sigma)$ given by a fan Σ in $N_{\mathbb{R}}$. Let $1 \in T \subseteq X$ denote the neutral element. Each \mathcal{O}_X -module \mathcal{E} gives rise to a \mathbb{C} -vector space $E := \mathcal{E}(1) := \mathcal{E}_1/\mathfrak{m}_{X,1}\mathcal{E}_1$, where \mathcal{E}_1 denotes the stalk of \mathcal{E} at $1 \in X$ and $\mathfrak{m}_{X,1}$ the maximal ideal of 1 . If \mathcal{E} is a T -equivariant, torsion free sheaf on X , the sections of \mathcal{E} on the open, affine, T -invariant subsets $\mathbb{T}\mathbb{V}(\sigma) \subseteq X$ with $\sigma \in \Sigma$ are M -graded subsets of $E \otimes_{\mathbb{C}} \mathbb{C}[M]$. If, in addition, \mathcal{E} is reflexive, then \mathcal{E} is already determined by its restriction to open subsets whose complements are of codimension equal or greater than two. Via Klyachko's description [Kly90], a toric reflexive sheaf \mathcal{E} corresponds to a set of decreasing \mathbb{Z} -filtrations

$$(5.1) \quad F(\mathcal{E})_{\rho}^{\bullet} = [\dots \supseteq E_{\rho}^{\ell-1} \supseteq E_{\rho}^{\ell} \supseteq E_{\rho}^{\ell+1} \supseteq \dots] \quad (\ell \in \mathbb{Z})$$

of the vector space E which are parametrized by the rays $\rho \in \Sigma(1)$. Let v_{ρ} denote the primitive generator of the ray ρ . The filtrations encode the sections of \mathcal{E} on the T -invariant open subsets $U_{\rho} = \mathbb{T}\mathbb{V}(\rho) \subseteq X$ defined by ρ . Namely, for $u \in M$,

$$(5.2) \quad e \otimes \chi^u \in \Gamma(U_{\rho}, \mathcal{E}) \quad \Leftrightarrow \quad e \in E_{\rho}^{-\langle u, v_{\rho} \rangle} = F(\mathcal{E})_{\rho}^{-\langle u, v_{\rho} \rangle}.$$

Remark 26. The reflexive sheaf \mathcal{E} defines a toric *vector bundle* if it is subject to *Klyachko's compatibility condition* [Kly90]: For each cone $\sigma \in \Sigma$ there exists a decomposition $E = \bigoplus_{[u] \in M/M \cap \sigma^{\perp}} E_{[u]}$ so that $E_{\rho}^{\ell} = \sum_{\langle u, v_{\rho} \rangle \geq \ell} E_{[u]}$ for each $\rho \in \sigma(1)$.

5.2. Line and Tangent Bundles. Line bundles and the tangent are examples of toric bundles with filtrations as follows (compare [Kly90, Example 2.3]).

Example 27. Let $D_{\rho} = \overline{\text{orb}(\rho)}$ be the closure of the orbit defined by $\rho \in \Sigma(1)$. For $D = \sum_{\rho \in \Sigma(1)} \lambda_{\rho} D_{\rho}$, $\lambda_{\rho} \in \mathbb{Z}$, the invertible sheaf $\mathcal{O}_X(D)$ is encoded by

$$(5.3) \quad E_{\rho}^{\ell} := \left\{ \begin{array}{ll} \mathbb{C} & \text{if } \ell \leq \lambda_{\rho} \\ 0 & \text{if } \ell \geq \lambda_{\rho} + 1 \end{array} \right\} \subseteq \mathbb{C} =: E.$$

Example 28. The tangent sheaf \mathcal{T}_X corresponds to the filtration

$$(5.4) \quad T_{\rho}^{\ell} := \left\{ \begin{array}{ll} N_{\mathbb{C}} = N \otimes_{\mathbb{Z}} \mathbb{C} & \text{if } \ell \leq 0 \\ \text{span}(\rho) & \text{if } \ell = 1 \\ 0 & \text{if } \ell \geq 2. \end{array} \right\} \subseteq N_{\mathbb{C}} =: E.$$

Remark 29. Projective n -space \mathbb{P}^n is the toric variety associated to the normal fan $\mathcal{N}(\Delta_n)$ of the standard n -simplex Δ_n . The fan $\mathcal{N}(\Delta_n)$ has $n+1$ rays ρ_0, \dots, ρ_n . The direct sum of invertible sheaves $\bigoplus_{j=0}^n \mathcal{O}_{\mathbb{P}^n}(\overline{\text{orb}(\rho_j)})$ corresponds to the filtrations

$$(5.5) \quad E_{\rho_i}^{\ell} := \left\{ \begin{array}{ll} E & \text{if } \ell \leq 0 \\ \mathbb{C} \cdot e^{\rho_i} & \text{if } \ell = 1 \\ 0 & \text{if } \ell \geq 2 \end{array} \right\} \subseteq E := \bigoplus_{j=0}^n \mathbb{C} \cdot e^{\rho_j}.$$

The canonical surjection $\pi: E \rightarrow N_{\mathbb{C}}$, $e^{\rho_i} \mapsto v_{\rho_i}$, where v_{ρ_i} generates the ray ρ_i , induces the filtrations of $N_{\mathbb{C}}$ corresponding to $\mathcal{T}_{\mathbb{P}^n}$. On $\ker(\pi) \cong \mathbb{C}$ the induced filtrations are those of the structure sheaf $\mathcal{O}_{\mathbb{P}^n}$. This yields the *Euler sequence*

$$(5.6) \quad 0 \rightarrow \mathcal{O}_{\mathbb{P}^n} \rightarrow \bigoplus_{j=0}^n \mathcal{O}_{\mathbb{P}^n}(\overline{\text{orb}(\rho_j)}) \rightarrow \mathcal{T}_{\mathbb{P}^n} \rightarrow 0.$$

5.3. Pullbacks under Toric Morphisms. Let \mathcal{E} be a toric reflexive sheaf on $X := \mathbb{T}\mathbb{V}(\Sigma, N)$. Similarly to subsection (4.2.3), let $\tilde{N} \subseteq N$ be a sublattice of finite index and $\tilde{X} := \mathbb{T}\mathbb{V}(\Sigma, \tilde{N})$. Let $v_\rho \in N$ denote the primitive generator of the ray $\rho \in \Sigma(1)$ in N and let \bar{v}_ρ be its image in the quotient group $G := N/\tilde{N}$. For $d_\rho := \min\{d \geq 1 \mid d \cdot \bar{v}_\rho = 0 \text{ in } G\}$ the order of \bar{v}_ρ in G , the multiple $\tilde{v}_\rho := d_\rho \cdot v_\rho \in \tilde{N}$ is the primitive generator of ρ in \tilde{N} . Let $p: \tilde{X} \rightarrow X$ be the toric covering morphism. Analogously to [Pay06, Prop. 4.9] one can give the following description of the filtrations of a pullback of a toric reflexive sheaf on X along p .

Proposition 30. *Suppose the toric reflexive sheaf \mathcal{E} corresponds to the vector space $E = \mathcal{E}(1)$ with filtrations $(E_\rho^l)_{l \in \mathbb{Z}}$ for $\rho \in \Sigma(1)$. Then the pullback $\mathcal{F} := p^*\mathcal{E}$ of \mathcal{E} is a toric reflexive sheaf on \tilde{X} . It corresponds to the same vector space $F = E$ and the filtration for $\rho \in \Sigma(1)$ is given by $F_\rho^l = E_\rho^{\lceil \frac{l}{d_\rho} \rceil}$, $l \in \mathbb{Z}$, where $\lceil \cdot \rceil$ denotes the ceiling function, giving the smallest integer equal to or larger than the argument.*

This filtration $(F_\rho^l)_{l \in \mathbb{Z}}$ of $F = E$ can be thought of as the filtration $(E_\rho^l)_{l \in \mathbb{Z}}$ stretched by the factor d_ρ and we will refer to it as the *d_ρ -th stretching of the filtration*.

5.4. Constructing the Universal Extension using Klyachko's Description.

We are in the setting of subsection (4.2.3). Consider the two line bundles $\mathcal{O}_X(\Delta^+)$ and $\mathcal{O}_X(\Delta^-)$ on X . In Klyachko's description, let them be given by the \mathbb{C} -vector spaces with decreasing \mathbb{Z} -filtrations $(E_+, E_{+, \rho}^\bullet)_{\rho \in \Sigma(1)}$ and $(E_-, E_{-, \rho}^\bullet)_{\rho \in \Sigma(1)}$, respectively. By Proposition 30 the pullback line bundles $\mathcal{O}_{\tilde{X}}(\Delta^+) = p^*\mathcal{O}_X(\Delta^+)$ and $\mathcal{O}_{\tilde{X}}(\Delta^-) = p^*\mathcal{O}_X(\Delta^-)$ on \tilde{X} correspond to the \mathbb{C} -vector spaces and decreasing \mathbb{Z} -filtrations $(F_+, F_{+, \rho}^\bullet)_{\rho \in \Sigma(1)}$ and $(F_-, F_{-, \rho}^\bullet)_{\rho \in \Sigma(1)}$ with $F_\pm = E_\pm$ and $F_{\pm, \rho}^l = E_{\pm, \rho}^{\lceil \frac{l}{d_\rho} \rceil}$. The filtration F_ρ^\bullet for a ray ρ corresponding to a pullback sheaf $\mathcal{F} = p^*\mathcal{E}$ of \mathcal{E} on X is a *d_ρ -th stretching* in the sense that a proper inclusion can only occur every d_ρ steps in the filtration:

$$(5.7) \quad \dots = \underbrace{F_\rho^{d_\rho k}}_{=E_\rho^k} \supseteq \underbrace{F_\rho^{d_\rho k+1}}_{=E_\rho^{k+1}} = \dots = \underbrace{F_\rho^{d_\rho(k+1)}}_{=E_\rho^{k+1}} \supseteq \underbrace{F_\rho^{d_\rho(k+1)+1}}_{=E_\rho^{k+2}} = \dots$$

In sequence (4.24) the outer two sheaves are the pullbacks of line bundles on X .

Lemma 31. *The sheaf $\tilde{\mathcal{H}}$ in sequence (4.24) is a reflexive sheaf on \tilde{X} and corresponds to a \mathbb{C} -vector space with filtrations $(\tilde{H}, \tilde{H}_\rho^\bullet)_{\rho \in \Sigma(1)}$, where the filtration \tilde{H}_ρ^\bullet for the ray $\rho \in \Sigma(1)$ is a d_ρ -th stretching in the sense introduced above.*

Proof. Applying the contravariant functor $\mathcal{H}om(-, \mathcal{O}_{\tilde{X}}): \text{Coh}(\tilde{X}) \rightarrow \text{Coh}(\tilde{X})$ to the sequence (4.24) twice yields a short exact sequence of double duals (using that the outer two sheaves are locally free) with a canonical homomorphism from the original sequence. Reflexivity of the outer two sheaves and the five lemma give reflexivity of the sheaf $\tilde{\mathcal{H}}$. Let \tilde{H}_ρ^\bullet , $F_{+, \rho}^\bullet$ and $F_{-, \rho}^\bullet$, $\rho \in \Sigma(1)$, denote the filtrations

corresponding to the reflexive sheaves $\tilde{\mathcal{H}}$, $\mathcal{O}_{\tilde{X}}(\Delta^+)$ and $\mathcal{O}_{\tilde{X}}(\Delta^-)$, respectively. Since sequence (4.24) is T -equivariant, it induces a short exact sequence of filtrations $0 \rightarrow F_{+, \rho}^\bullet \rightarrow \tilde{H}_\rho^\bullet \rightarrow F_{-, \rho}^\bullet \rightarrow 0$ for each $\rho \in \Sigma(1)$. Hence, the filtration H_ρ^\bullet of H is determined by the filtrations $F_{+, \rho}^\bullet$ and $F_{-, \rho}^\bullet$ and is thus also d_ρ -th stretching. \square

Remark 32. A filtration of a vector space that is a d -th stretching can also be *squished* back: For a vector space F with filtration

$$(5.8) \quad \dots = F^{dk} \supseteq F^{dk+1} = \dots = F^{d(k+1)} \supseteq F^{d(k+1)+1} = \dots$$

the d -th *squishing* of (F, F^\bullet) is the vector space F with filtration

$$(5.9) \quad \dots \supseteq F^{d(k-1)} \supseteq F^{dk} \supseteq F^{d(k+1)} \supseteq \dots$$


Corollary 33. *The reflexive sheaf \mathcal{H} on X corresponding to (H, H_ρ^\bullet) , where $H := \tilde{H}$ and the filtration H_ρ^\bullet given by $H_\rho^l := \tilde{H}_\rho^{d_\rho l}$ ($= \tilde{H}_\rho^{d_\rho l-1} = \dots = \tilde{H}_\rho^{d_\rho l - (d_\rho - 1)}$), $l \in \mathbb{Z}$, is the d_ρ -th squishing of \tilde{H}_ρ^\bullet , $\rho \in \Sigma(1)$, fits into a short exact sequence*

$$(5.10) \quad 0 \rightarrow \mathcal{O}_X(\Delta^+)^n \rightarrow \mathcal{H} \rightarrow \mathcal{O}_X(\Delta^-) \rightarrow 0$$

that pulls back to the short exact sequence (4.24) under $p: \tilde{X} \rightarrow X$.

Sequence (5.10) is precisely the universal extension sequence (4.30).

Example 34. We continue with the example introduced in subsection (4.2.1) using Klyachko's language. Let $M = \mathbb{Z} \oplus \mathbb{Z}$, $N = \mathbb{Z} \oplus \mathbb{Z}$ and recall the smooth fan Σ in $N_{\mathbb{R}} \cong \mathbb{R}^2$ and the lattice polytopes Δ^+ and Δ^- with respect to $M = \mathbb{Z} \oplus \mathbb{Z}$:



with Δ^+ not contained in Δ^- . The intersection $\Delta^+ \cap \Delta^-$ is not a lattice polytope with respect to $\mathbb{Z} \oplus \mathbb{Z}$. Consider the sublattice $\tilde{N} = 2\mathbb{Z} \oplus \mathbb{Z} \hookrightarrow \mathbb{Z} \oplus \mathbb{Z} = N$ and dually $M = \mathbb{Z} \oplus \mathbb{Z} \hookrightarrow (\frac{1}{2}\mathbb{Z}) \oplus \mathbb{Z} = \tilde{M}$, so that $\Delta^+ \cap \Delta^-$ is a lattice polytope with respect to \tilde{M} . Denote the unions of the components of $\Delta^- \setminus (\Delta^+ \cap \Delta^-)$ with $\Delta^+ \cap \Delta^-$ by

$$\nabla_0 := \text{triangle with vertices } (0,0), (1,0), (0,-1) \text{ and } \nabla_1 := \text{triangle with vertices } (0,0), (1,0), (0,1).$$

Set $\tilde{X} := \text{TV}(\Sigma, \tilde{N})$ and denote the covering morphism by $p: \tilde{X} \rightarrow X$. The minimal ray generators \tilde{v}_{ρ_i} , $i = 1, \dots, 6$, in the lattice \tilde{N} are $\tilde{v}_{\rho_1} = (2, 0)$, $\tilde{v}_{\rho_2} = (0, 1)$, $\tilde{v}_{\rho_3} = (-2, 0)$, $\tilde{v}_{\rho_4} = (-2, -1)$, $\tilde{v}_{\rho_5} = (-2, -2)$, and $\tilde{v}_{\rho_6} = (0, -1)$. From these we determine the filtrations of the divisors $\tilde{D}_{\Delta^+ \cap \Delta^-}$, \tilde{D}_{Δ^+} , \tilde{D}_{Δ^-} , \tilde{D}_{∇_0} and \tilde{D}_{∇_1} on \tilde{X} . Taking pushouts allows us to calculate the filtrations for $\tilde{\mathcal{H}}$ on \tilde{X} . In order to obtain

the filtrations for the sheaf \mathcal{H} we need to take the d_ρ -squishings of the filtrations for $\tilde{\mathcal{H}}$, whenever $d_\rho \neq 1$. This is the case for ρ_1, ρ_3 and ρ_5 , with $d_\rho = 2$ in each case. The following table depicts the resulting filtrations for \mathcal{H} .

\mathcal{H}	0	1	2	3
ρ_1	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	0	$\supseteq 0$	$\supseteq 0$
ρ_2	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	\mathbb{C}_1	$\supseteq 0$	$\supseteq 0$
ρ_3	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	$\supseteq 0$	$\supseteq 0$
ρ_4	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	$\supseteq \mathbb{C}_+$	$\supseteq 0$
ρ_5	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	$\supseteq 0$	$\supseteq 0$
ρ_6	$\supseteq (\mathbb{C}_+ \oplus \mathbb{C}_0 \oplus \mathbb{C}_1)/\mathbb{C}$	\mathbb{C}_0	$\supseteq 0$	$\supseteq 0$

Note that the vector bundle \mathcal{H} (and even the vector bundle $\tilde{\mathcal{H}}$ on \tilde{X}) does not split. This can be seen using a criterion from Klyachko, that a vector bundle splits if and only if the vector spaces in the filtrations of all the rays form a distributive lattice, or, equivalently are given by coordinate subspaces (compare [Kly89, Cor. 2.2.3]).

REFERENCES

- [ABKW20] Klaus Altmann, Jarosław Buczyński, Lars Kastner, and Anna-Lena Winz. Immaculate line bundles on toric varieties. *Pure and Applied Mathematics Quarterly*, 16(4):1147–1217, 2020.
- [AP20] Klaus Altmann and David Ploog. Displaying the cohomology of toric line bundles. *Izvestiya: Mathematics*, 84(4):683–693, 2020.
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*. Providence, RI: American Mathematical Society (AMS), 2011.
- [Dan78] Vladimir I. Danilov. The Geometry of Toric Varieties. *Russian Mathematical Surveys*, 33(2):97–154, 1978.
- [Dem70] Michel Demazure. Sous-groupes algébriques de rang maximum du groupe de cremona. *Annales scientifiques de l'École Normale Supérieure*, 4e série, 3(4):507–588, 1970.
- [DJS14] Sandra Di Rocco, Kelly Jabbusch, and Gregory Smith. Toric vector bundles and parliaments of polytopes. *Transactions of the American Mathematical Society*, 370, 09 2014.
- [Ful93] William Fulton. *Introduction to toric varieties. The 1989 William H. Roever lectures in geometry*. Princeton, NJ: Princeton University Press, 1993.
- [Kly89] Alexander A. Klyachko. Toric bundles and problems of linear algebra. *Functional Analysis and its Applications*, 23(2):135–137, 1989.
- [Kly90] Alexander A. Klyachko. Equivariant bundles on toral varieties. *Mathematics of the USSR. Izvestija*, 35(2):337–375, 1990.
- [Kly02] Alexander A. Klyachko. Vector bundles, linear representations, and spectral problems. In *Proceedings of the international congress of mathematicians, ICM 2002, Beijing, China, August 20–28, 2002. Vol. II: Invited lectures*, pages 599–613. Beijing: Higher Education Press; Singapore: World Scientific/distributor, 2002.
- [KM] Kiumars Kaveh and Christopher Manon. Toric bundles, valuations, and tropical geometry over semifield of piecewise linear functions. arXiv:1907.00543 [math.AG].
- [Pay06] Sam Payne. Toric vector bundles, branched covers of fans, and the resolution property. *Journal of Algebraic Geometry*, 18:1–36, 2006.

- [Pay08] Sam Payne. Moduli of toric vector bundles. *Compositio Mathematica*, 144(5):1199–1213, 2008.
- [Wei95] Charles A. Weibel. *An Introduction to Homological Algebra*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1995.

INSTITUT FÜR MATHEMATIK, FU BERLIN, KÖNIGIN-LUISE-STR. 24-26, 14195 BERLIN, GERMANY

Email address: `altmann@math.fu-berlin.de`

INSTITUT FÜR MATHEMATIK, HU BERLIN, RUDOWER CHAUSSEE 25, 12489 BERLIN, GERMANY

Email address: `amelie.flatt@hu-berlin.de`

MATHEMATISCHES INSTITUT DER UNIVERSITÄT MÜNSTER, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY

Email address: `lutz.hille@uni-muenster.de`