# Derangements and the *p*-adic incomplete gamma function

#### ANDREW O'DESKY AND DAVID HARRY RICHMAN

ABSTRACT. We introduce a p-adic analogue of the incomplete gamma function. We also introduce quantities (m-values) associated to a function on natural numbers and prove a new characterization of p-adic continuity for functions with p-integral m-values. Combinatorial interpretations for the integral values of the incomplete gamma function and functions with m-values zero or one are obtained, which show that these functions count derangements in generalized symmetric groups and permutations with restricted cycle lengths.

#### 1. Introduction

The first object of this paper is to introduce a p-adic analogue of the incomplete gamma function. The incomplete gamma function  $\Gamma(s,z)$  is the holomorphic function of two variables obtained by integrating the differential  $z^{s-1}e^{-z}\,dz$  over any contour from z to  $+\infty$  disjoint from the real interval  $\{-\infty < x \le 0\}$ . For positive integers n and r,  $\Gamma(n,r)$  is a rational multiple of  $1/e^r$ . Let  $\tau_p \colon \mathbb{Q}(e) \to \mathbb{Q}_p$  denote the unique field homomorphism satisfying  $\tau_p(1/e) = 1 + p + p^2/2! + p^3/3! + \cdots$ . Then  $\tau_p\Gamma(n,r)$  is a p-adic number and we may ask for the existence of a continuous function which interpolates these values.

**Theorem 1.1.** There exists a continuous function

$$\Gamma_p \colon \mathbb{Z}_p \times (1 + p\mathbb{Z}_p) \to \mathbb{Z}_p$$

satisfying

$$\Gamma_p(n,r) = \tau_p \Gamma(n,r)$$

for any positive integers n and r satisfying  $r \equiv 1 \pmod{p}$ . Explicitly, for any  $s \in \mathbb{Z}_p$  and  $r \in 1 + p\mathbb{Z}_p$  we have the formula

$$\Gamma_p(s,r) = \exp_p(pr) \sum_{k=0}^{\infty} r^{s-1-k} k! \binom{s-1}{k}$$

where  $\exp_p$  is the p-adic exponential function.

Date: January 19, 2022.

<sup>2020</sup> Mathematics Subject Classification. 33B20, 11S80, 11B75, 05A05.

Key words and phrases. derangements, incomplete gamma function, p-adic analysis.

We also prove a curious formula for  $\Gamma_p$  involving the floor function  $\lfloor \cdot \rfloor$ . For any  $n \in \mathbb{N}$  and  $r \in \mathbb{Z}$  satisfying  $r \equiv 1 \pmod{p}$ ,

$$\Gamma_p(n+1,1/r) = r^{-n} \exp_p(p/r) \cdot \begin{cases} \lfloor e^{1/r} r^n n! + \frac{1}{2} \rfloor & \text{if } r < 0, \\ \lfloor e^{1/r} r^n n! \rfloor & \text{if } r > 0. \end{cases}$$

This formula expresses a p-adically continuous function in terms of a function which crucially uses the linear ordering on the real numbers.

1.1. p-adic continuity via m-values. The second object of this paper is to present a new type of characterization for p-adic continuity in terms of certain auxiliary quantities associated to a function  $f \colon \mathbb{N} \to \mathbb{Q}_p$ . These quantities are defined using universal polynomials from the theory of symmetric functions. Let  $h_k = \sum_{i_1 \leq \cdots \leq i_k} x_{i_1} \cdots x_{i_k}$  denote the kth complete homogeneous symmetric function. For each positive integer k there is a unique polynomial  $M_k(h_1, \ldots, h_k)$  satisfying

$$M_k(h_1,\ldots,h_k) = p_k = x_1^k + x_2^k + x_3^k + \cdots$$

The  $kth\ m$ -value of a function f is defined by

$$m_k = M_k \left( \frac{f(1)}{1!}, \frac{f(2)}{2!}, \dots, \frac{f(k)}{k!} \right).$$
 (1)

When f(0) = 1, the m-values of f satisfy the identity

$$\sum_{n=0}^{\infty} f(n) \frac{X^n}{n!} = \exp\left(\sum_{k=1}^{\infty} m_k \frac{X^k}{k}\right).$$

For instance,  $m_2 = f(2) - f(1)^2$ , and  $m_3 = \frac{1}{2}f(3) - \frac{3}{2}f(1)f(2) + f(1)^3$  (cf. §3.1).

**Theorem 1.2.** Let  $f: \mathbb{N} \to \mathbb{Q}_p$  be a function satisfying f(0) = 1. Suppose  $m_k$  is p-integral for all  $k \geq 2$ . Then f is p-adically continuous if and only if

$$m_p \equiv m_1 - 1 \pmod{p}. \tag{2}$$

One interesting function encompassed by this theorem is Morita's p-adic gamma function  $\Gamma_{p,M}(n) = (-1)^n \prod_{1 \le k < n, p \nmid k} k$ ; see Example 1.

Another example comes from the EGF-coefficients  $f_{\ell}$  of the  $\ell$ -adic Artin–Hasse exponential function  $E_{\ell}(X)$ :

$$E_{\ell}(X) = \exp\left(X + \frac{X^{\ell}}{\ell} + \frac{X^{\ell^2}}{\ell^2} + \cdots\right) = \sum_{n=0}^{\infty} f_{\ell}(n) \frac{X^n}{n!}.$$

The m-values of  $f_{\ell}$  are

$$m_k = \begin{cases} 0 & \text{if } k \text{ is not a power of } \ell, \\ 1 & \text{if } k \text{ is a power of } \ell, \end{cases}$$

<sup>&</sup>lt;sup>1</sup>For the purpose of testing continuity of f we may assume f(0) = 1 by composing with a suitable affine transformation, so this hypothesis entails no loss of generality.

and

$$m_p - m_1 + 1 = m_p = \begin{cases} 0 & \text{if } p \neq \ell, \\ 1 & \text{if } p = \ell. \end{cases}$$

Corollary 1.3. For primes p and  $\ell$ , the EGF-coefficients  $f_{\ell}$  of the  $\ell$ -adic Artin–Hasse exponential function are p-adically continuous if and only if  $p \neq \ell$ .

The same conclusion holds for  $truncated \ \ell$ -adic Artin–Hasse exponential functions (cf. [7]). For another application of Theorem 1.2 involving the counts of arrangements and derangements in generalized symmetric groups see Remark 1.

- 1.2. Generalized symmetric groups and cycle-restricted permutations. The third object of this paper is to prove combinatorial interpretations for each of the following quantities: the integral values of the incomplete gamma function and functions whose *m*-values are either zero or one.
- 1.2.1. Integral values of the incomplete gamma function. Just as the values of the gamma function at positive integers count permutations, we show that the values of the incomplete gamma function at integral values count cyclic arrangements and derangements in generalized symmetric groups.

In analogy with the classical formula  $\Gamma(n+1) = n!$  we show that

$$\Gamma(n+1, 1/r) = a(n, r)r^{-n}e^{-1/r}$$

for  $(n,r) \in \mathbb{N}^{>0} \times \mathbb{Z}$  where

$$a(n,r) := \begin{cases} (-1)^n (\# \text{ of } |r| \text{-cyclic derangements of degree } n) & \text{if } r < 0, \\ \# \text{ of } r \text{-cyclic arrangements of degree } n & \text{if } r > 0. \end{cases}$$
(3)

Here  $C_r$  (resp.  $S_n$ ) denotes the cyclic group of order r (resp. the symmetric group of degree n), and an r-cyclic derangement of degree n is an element of the generalized symmetric group  $C_r \wr S_n$  whose action on  $C_r \times [n]$  has no fixed point, while an r-cyclic arrangement of degree n is a choice of subset  $A \subset [n]$  and element of  $C_r \wr S_{|A|}$  (for details see §4.1). Assaf [3] introduced r-cyclic derangements while r-cyclic arrangements do not seem to have appeared in the combinatorics literature yet.

- 1.2.2. Functions with m-values zero or one. Our second combinatorial interpretation is for functions whose m-values (cf. (1)) are all in  $\{0,1\}$ . We show that any such function admits a combinatorial interpretation in terms of cycle-restricted permutations.
- **Theorem 1.4.** Let  $f: \mathbb{N} \to \mathbb{Q}$  be a function whose m-values are all in  $\{0, 1\}$ . Then f(n) is equal to the number  $d_n^L$  of permutations in  $S_n$  whose cycle lengths are in  $L = \{r : m_r = 1\}$ .

This theorem generalizes a classical formula of Chowla-Herstein-Scott [6],

$$\sum_{n=0}^{\infty} \# \{ \sigma^{\ell} = 1 : \sigma \in S_n \} \frac{X^n}{n!} = \exp \left( \sum_{d \mid \ell} \frac{X^d}{d} \right),$$

which is recovered when L consists of the divisors of a given integer  $\ell$ .

Combining this theorem with Theorem 1.2 leads to a vast generalization of the observation that  $n \mapsto (-1)^n d_n$  is p-adically continuous. Define the count of permutations with 'permissible' cycle lengths,

$$d_n^L := \#\{\sigma \in S_n : \text{ every cycle length of } \sigma \text{ is in } L\}$$

where  $L \subset \mathbb{N}^{>0}$  is any subset.<sup>2</sup>

**Theorem 1.5.** Let L be a set of positive integers, and let  $p \ge 3$  be a prime.

- (a)  $n \mapsto d_n^L$  is p-adically continuous if and only if  $1 \in L$  and  $p \notin L$ . (b)  $n \mapsto (-1)^n d_n^L$  is p-adically continuous if and only if  $1 \notin L$  and  $p \in L$ . (c)  $n \mapsto d_n^L$  is 2-adically continuous if and only if one of the following hold:
  - $1 \in L$  and  $2 \notin L$ , or
  - $1 \notin L$  and  $2 \in L$ .

Theorem 1.5 is deduced from a more uniform statement, which is that  $n \mapsto$  $\alpha^n d_n^L$  is p-adically continuous (for any prime p) if and only if

$$\alpha(\chi_L(1) - \chi_L(p)) \equiv 1 \pmod{p}$$

where  $\alpha$  is an integer and  $\chi_L$  is the characteristic function of L.

We mention some related articles for the interested reader.  $n \mapsto d_n^{\{1,p\}}$  has some intriguing p-adic properties (despite not being p-adically continuous) cf. [13], [7]. Arithmetic aspects of the function  $n \mapsto a(n, -1)$  counting derangements have been studied in [28], [27], [16], [26]. For any  $r \in \mathbb{Z}$  the function  $n \mapsto a(n,r)$  is a pseudopolynomial (in the sense of [11]) which is simultaneously p-adic locally analytic for every prime p (Theorem 4.3). Generalizations of pseudopolynomials have been recently studied in [14], [10]. The assertion that  $n \mapsto a(n,r)$  is p-adic locally analytic was first proven for r=-1 (counting derangements) in [4] using a formal Laplace transform, and for r=1 (counting arrangements) more recently in [22].

**Notation.** We use  $\mathbb N$  to denote the set of nonnegative integers and  $\mathbb N^{>0}$  for the set of positive integers. We let  $\mathbb{Z}_p$  (resp.  $\mathbb{Q}_p$ ) denote the ring of p-adic integers (resp. field of p-adic numbers). We use  $|\cdot|_p$  (resp.  $v_p$ ) to denote the field norm (resp. p-adic additive valuation) on  $\mathbb{Q}_p$ . We let  $\mathbb{C}_p$  denote the metric completion of an algebraic closure of  $\mathbb{Q}_p$ . The p-adic valuation  $v_p$  and the norm  $|\cdot|_p$  on  $\mathbb{Q}_p$ both extend uniquely to  $\mathbb{C}_p$  and are denoted by the same symbols.

<sup>&</sup>lt;sup>2</sup>The unique element of  $S_0$  acting on the empty set is considered to have no cycle lengths; thus if L is empty, then  $d_n^L = \delta_{0,n}$  which is never p-adically continuous.

#### 2. Preliminaries

In this section we briefly recall some standard results from p-adic analysis for the reader's convenience. For more background we refer to [24].

Given a function  $f: \mathbb{N} \to \mathbb{C}_p$ , the basic question of p-adic interpolation is whether f extends to a continuous mapping (Lipschitz function, locally analytic function, analytic function, etc.)  $\tilde{f}: \mathbb{Z}_p \to \mathbb{C}_p$ . This question is generally approached using finite differences. The *nth finite difference* c(n) of f (at zero) is

$$c(n) := \sum_{k=0}^{n} (-1)^{n-k} \binom{n}{k} f(k). \tag{4}$$

This admits an inverse relation given by  $f(n) = \sum_{k=0}^{n} {n \choose k} c(k)$  for  $n \ge 0$ .

A classical theorem of Mahler [15] says that f extends to a continuous function on  $\mathbb{Z}_p$  if and only if  $|c(n)|_p \to 0$  as  $n \to \infty$ . More generally, faster rates of p-adic decay for the finite differences c(n) correspond to stronger p-adic properties of f.

**Definition 2.1.** A function  $f: X \to Y$  of metric spaces is a metric map if  $d_Y(f(x), f(y)) \le d_X(x, y)$  for all  $x, y \in X$ . (Equivalently, f is Lipschitz continuous with constant one.) If D is an open disk of  $\mathbb{C}_p$ , a function  $f: D \to \mathbb{C}_p$  is analytic if there are elements  $a_0 \in D$  and  $c_0, c_1, \ldots \in \mathbb{C}_p$  such that  $f(x) = \sum_{n=0}^{\infty} c_n(x-a_0)^n$  for all  $x \in D$ . If U is an open subset of  $\mathbb{C}_p$ , a function  $f: U \to \mathbb{C}_p$  is locally analytic if for each  $a \in U$  there is an open disk  $D \subset U$  containing a such that  $f|_D$  is analytic.

**Theorem 2.2.** Let  $f: \mathbb{N} \to \mathbb{C}_p$  be a function and let c(n) be given by (4). Then f extends to:

(1) a continuous map on  $\mathbb{Z}_p$  if and only if

$$|c(n)|_p \to 0 \text{ as } n \to \infty,$$

(2) a metric map on  $\mathbb{Z}_p$  if and only if

$$\sup_{n\geq 1} |c(n)|_p p^{\lfloor \frac{\log n}{\log p} \rfloor} \leq 1,$$

(3) an analytic function on the closed unit disk if and only if

$$|c(n)/n!|_p \to 0 \text{ as } n \to \infty,$$

(4) a locally analytic function on the closed unit disk if and only if

$$\liminf_{n \to \infty} \left( \frac{v_p(c(n))}{n} \right) > 0.$$

*Proof.* The first assertion is due to Mahler [15]. For the second see [23, §5.1], or [21, Proposition 2] for a generalization. The third is [24, Theorem 54.4]. The fourth is due to Amice [2, III, §10, Corollaire 2, p. 162].  $\Box$ 

## 3. p-adic continuity via congruences

In this section, we give a new characterization of p-adic continuity for a class of functions satisfying certain integrality conditions. We then apply this characterization to a family of combinatorial sequences which count permutations with restricted cycle types.

Our characterization uses universal polynomials from the theory of symmetric functions. Let  $h_k = \sum_{i_1 \leq \cdots \leq i_k} x_{i_1} \cdots x_{i_k}$  denote the kth complete homogeneous symmetric function. For each positive integer k there is a unique polynomial  $M_k(h_1,\ldots,h_k)$  satisfying  $M_k(h_1,\ldots,h_k) = p_k = x_1^k + x_2^k + x_3^k + \cdots$ .

**Definition 3.1.** Let  $f: \mathbb{N} \to \mathbb{Q}_p$  be a function. The kth m-value of f is

$$m_k := M_k\left(\frac{f(1)}{1!}, \frac{f(2)}{2!}, \dots, \frac{f(k)}{k!}\right).$$

The main result of this section is the following theorem.

**Theorem 3.2** (Theorem 1.2). Let  $f: \mathbb{N} \to \mathbb{Q}_p$  be a function satisfying f(0) = 1. Suppose  $m_k$  is p-integral for all  $k \geq 2$ . Then f is p-adically continuous if and only if

$$M_p\left(\frac{f(1)}{1!}, \frac{f(2)}{2!}, \dots, \frac{f(p)}{p!}\right) \equiv f(1) - 1 \pmod{p\mathbb{Z}_p}.$$

It is rather surprising that continuity for functions with p-integral m-values can be detected using just  $m_p$  and  $m_1$ , as these m-values only depend on the values of f on the integers  $1, 2, \ldots, p$ . Thus the assumption of p-integrality of m-values rigidly couples the values of f, in the sense that if f has p-integral m-values and  $m_p \equiv m_1 - 1 \pmod{p}$ , then if any value of f on an integer greater than p is modified, the m-values of f cease to be p-integral.

**Example 1.** Morita's p-adic gamma function is defined by

$$\Gamma_{p,\mathcal{M}}(z) = \lim_{n \to z^+} (-1)^n \prod_{1 \le k < n, p \nmid k} k$$

where n approaches  $z \in \mathbb{Z}_p$  through positive integers. Set  $f(n) = -\Gamma_{p,M}(n+1)$ . Then

$$\sum_{n=0}^{\infty} f(n) \frac{X^n}{n!} = \frac{1 + X^p}{1 + X} e^{-X^p/p},$$

and from this one finds (for p odd)

$$m_k = \begin{cases} (-1)^k & \text{if } k \not\equiv 0 \pmod{p}, \\ p - 2 & \text{if } k = p, \\ (-1)^k (1 - p) & \text{if } k \equiv 0 \pmod{p} \text{ and } k > p. \end{cases}$$

Then  $m_p = p - 2 \equiv (-1) - 1 \pmod{p}$  so we conclude  $n \mapsto -\Gamma_{p,M}(n+1)$ , and therefore  $n \mapsto \Gamma_{p,M}(n)$  also, is p-adically continuous.

3.1. The universal polynomials  $M_k$ . It is well-known that  $\{h_k\}_k$  and  $\{p_k\}_k$  are algebraic bases of the ring of symmetric functions, and the universal polynomials  $M_1, M_2, M_3, \ldots$  express the change of basis from the power basis  $\{p_k\}_k$  to the complete homogeneous basis  $\{h_k\}_k$ .

The  $\{M_k\}_k$  can be computed using the classical identity

$$\sum_{k=1}^{\infty} \frac{p_k}{k} X^k = \log \left( \sum_{n=0}^{\infty} h_n X^n \right). \tag{5}$$

From this one finds

$$\sum_{k=1}^{\infty} M_k \frac{X^k}{k} = h_1 X + \left( -h_1^2 + 2h_2 \right) \frac{X^2}{2} + \left( h_1^3 - 3h_1 h_2 + 3h_3 \right) \frac{X^3}{3} + \left( -h_1^4 + 4h_1^2 h_2 - 2h_2^2 - 4h_1 h_3 + 4h_4 \right) \frac{X^4}{4} + \left( h_1^5 - 5h_1^3 h_2 + 5h_1 h_2^2 + 5h_1^2 h_3 - 5h_2 h_3 - 5h_1 h_4 + 5h_5 \right) \frac{X^5}{5} + \cdots$$

For a function f satisfying f(0) = 1, set  $h_n = f(n)/n!$  for all n, and write  $f_k = f(k)$  for readability. Then

$$\sum_{k=1}^{\infty} m_k \frac{X^k}{k} = f_1 X +$$

$$(-f_1^2 + f_2) \frac{X^2}{2} +$$

$$\left( f_1^3 - \frac{3}{2} f_1 f_2 + \frac{1}{2} f_3 \right) \frac{X^3}{3} +$$

$$\left( -f_1^4 + 2 f_1^2 f_2 - \frac{1}{2} f_2^2 - \frac{2}{3} f_1 f_3 + \frac{1}{6} f_4 \right) \frac{X^4}{4} +$$

$$\left( f_1^5 - \frac{5}{2} f_1^3 f_2 + \frac{5}{4} f_1 f_2^2 + \frac{5}{6} f_1^2 f_3 - \frac{5}{12} f_2 f_3 - \frac{5}{24} f_1 f_4 + \frac{1}{24} f_5 \right) \frac{X^5}{5} + \cdots$$

One can expand the power series for the logarithm in (5) to find

$$\frac{M_k}{k} = \sum_{\lambda \vdash k} \frac{(-1)^{\ell(\lambda)+1}}{\ell(\lambda)} \binom{\ell(\lambda)}{\lambda[1], \dots, \lambda[k]} \prod_{j=1}^k h_j^{\lambda[j]}$$

and

$$\frac{m_k}{k} = \sum_{\lambda \vdash k} \frac{(-1)^{\ell(\lambda)+1}}{\ell(\lambda)} \binom{\ell(\lambda)}{\lambda[1], \dots, \lambda[k]} \prod_{j=1}^k \left(\frac{f(j)}{j!}\right)^{\lambda[j]}$$

where the sum is taken over partitions  $\lambda$  of k,  $\ell(\lambda)$  is the number of parts of  $\lambda$ , and  $\lambda[j]$  is the number of parts of  $\lambda$  equal to j.

The *m*-values of a function may also be computed rapidly from the recursive relation  $(k \ge 2)$ 

$$m_k = \frac{f(k)}{(k-1)!} - \sum_{j=1}^{k-1} \frac{f(k-j)}{(k-j)!} m_j.$$

3.2. **Proof of Theorem 3.2.** The key technical result in the proof of Theorem 3.2 is Theorem 3.6, which gives necessary and sufficient conditions for the convergence of certain infinite products of exponentials. The convergence takes place in a p-adic Banach algebra V of exponential generating functions associated to p-adically continuous functions.

**Definition 3.3.** Let V denote the set of all formal power series  $\sum_{n=0}^{\infty} c_n \frac{X^n}{n!} \in \mathbb{Q}_p[\![X]\!]$  with the property that  $c_n = \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} f(k)$  for some continuous function  $f: \mathbb{Z}_p \to \mathbb{Q}_p$ . Define the norm of  $F = \sum_{n=0}^{\infty} c_n \frac{X^n}{n!}$  to be

$$||F|| := \sup_{n \ge 0} |c_n|_p. \tag{6}$$

We say that  $c_n$  is the *nth EGF-coefficient* of  $\sum_{n=0}^{\infty} c_n \frac{X^n}{n!}$ .

**Proposition 3.4.**  $(V, ||\cdot||)$  is a Banach algebra.

*Proof.* By Mahler's theorem, V is equal to the set of formal power series  $\sum_{n} c_n \frac{X^n}{n!}$  satisfying  $|c_n|_p \to 0$ . It is now clear that V is the completion of the polynomial ring  $\mathbb{Q}_p[X]$  for the norm (6), and is therefore complete.

We must verify that  $\|\cdot\|$  is submultiplicative. Let  $F(X) = \sum_{n=0}^{\infty} a_n \frac{X^n}{n!}$  and  $G(X) = \sum_{n=0}^{\infty} b_n \frac{X^n}{n!}$ . We have that

$$F(X)G(X) = \sum_{n>0} c_n \frac{X^n}{n!} \quad \text{where} \quad c_n = \sum_{k=0}^n \binom{n}{k} a_{n-k} b_k. \tag{7}$$

By the ultrametric triangle inequality,

$$||F \cdot G|| = \sup_{n \ge 0} \left| \sum_{k=0}^{n} {n \choose k} a_{n-k} b_k \right|_p \le \sup_{m \ge 0} |a_m|_p \cdot \sup_{n \ge 0} |b_n|_p = ||F|| \, ||G||.$$

The product operation on  $\mathbb{C}_p[X]$  is continuous for the norm topology, and therefore extends uniquely to a product on its completion V. We have shown  $(V, \|\cdot\|)$  is a Banach algebra.

**Definition 3.5.** Let  $m_1, m_2, \ldots \in \mathbb{Q}_p$ . The zeta function associated to  $m = (m_1, m_2, \ldots)$  is the formal power series defined by

$$\zeta_m(X) := \exp\left(\sum_{k=1}^{\infty} \frac{m_k}{k} X^k\right). \tag{8}$$

**Theorem 3.6.** Let  $m_1, m_2, \ldots \in \mathbb{Q}_p$  and suppose  $m_r \in \mathbb{Z}_p$  for all  $r \geq 2$ . Then  $\zeta_m$  is in V if and only if  $m_1 \equiv m_p \pmod{p}$ .

To prove the theorem we establish two lemmas. For the proofs, we will make repeated use of a well-known formula of Legendre for the p-adic valuation of n!,

$$v_p(n!) = \frac{n - s_p(n)}{p - 1}$$

where  $s_p(n)$  is the sum of the digits of n in base p.

**Lemma 3.7.** For  $m \in \mathbb{C}_p$  and  $r \in \mathbb{N}^{>0}$  the power series  $\exp(mX^r/r)$  is in V if and only if  $v_p(m) > v_p(r) - \frac{r-1}{p-1}$ .

*Proof.* For  $n \ge 0$  the nth EGF-coefficient  $c_n$  of  $\exp(mX^r/r) = \sum_{n\ge 0} c_n X^n/n!$  is given by

$$c_n = \begin{cases} m^k \frac{(rk)!}{r^k k!} & \text{if } n = rk \text{ for } k \in \mathbb{N}, \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$\liminf_{n\to\infty} \frac{1}{n} v_p(c_n) = \liminf_{k\to\infty} \frac{1}{rk} v_p\left(m^k \frac{(rk)!}{r^k k!}\right) = \frac{1}{r} \left(v_p(m) - v_p(r) + \frac{r-1}{p-1}\right).$$

If  $v_p(m) - v_p(r) + \frac{r-1}{p-1} > 0$ , then  $v_p(c_n) \to \infty$  as  $n \to \infty$  so  $\exp(mX^r/r)$  is in V by Theorem 2.2.

To finish we show  $\exp(mX^r/r)$  is not in V if  $v_p(m) - v_p(r) + \frac{r-1}{p-1} \leq 0$ . If  $v_p(m) - v_p(r) + \frac{r-1}{p-1} < 0$ , then  $v_p(c_n) \not\to \infty$  so  $\exp(mX^r/r)$  is not in V. If  $v_p(m) - v_p(r) + \frac{r-1}{p-1} = 0$  then  $v_p(c_n) = (p-1)^{-1}(s_p(rn) - s_p(n))$  where  $s_p(n)$  is the sum of the base-p digits of n. This shows  $v_p(c_n) < \infty$  is constant for  $n \in \{1, p, p^2, \ldots\}$  so  $\exp(mX^r/r)$  is not in V.

**Lemma 3.8.** Let  $m_1, m_2, \ldots \in \mathbb{Z}_p$ . If  $m_1 = m_p = 0$  then the zeta function  $\zeta_m$  (cf. (8)) is in V.

Proof. We claim that  $v_p(r) - \frac{r-1}{p-1} < 0$  if  $r \in \mathbb{N}^{>0} \setminus \{1, p\}$ . The claim is clear if  $v_p(r) = 0$ . If  $v_p(r) = 1$  then r > p, so  $v_p(r) - \frac{r-1}{p-1} < 1 - \frac{p-1}{p-1} = 0$ . If  $v_p(r) = v \ge 2$  then  $r \ge p^v$ , and  $v - \frac{r-1}{p-1} \le v - \frac{p^v-1}{p-1} \le v - 1 - 2^{v-1}$ . The quantity  $v - 1 - 2^{v-1}$  is always negative, which establishes the claim.

By the claim,  $v_p(m_r) \geq 0 > v_p(r) - \frac{r-1}{p-1}$  whenever  $r \in \mathbb{N}^{>0} \setminus \{1, p\}$ , and by Lemma 3.7 we conclude  $\exp(m_r X^r/r)$  is in V for any  $r \in \mathbb{N}^{>0}$ . Let  $G_N = \prod_{r=1}^N \exp(m_r X^r/r)$  denote the partial product, which is contained in V by Proposition 3.4. To prove the lemma it suffices to show that  $(G_N)_{N\geq 1}$  is a Cauchy sequence by completeness of V.

For any positive integer r we have that

$$\left\| \exp\left(\frac{m_r X^r}{r}\right) \right\| = \sup_{n \ge 0} \left| m_r^n \frac{(rn)!}{r^n n!} \right|_p \le \sup_{n \ge 0} \left| \prod_{\substack{1 \le k \le rn \\ k \ne r \mathbb{Z}}} k \right|_p = 1.$$

By submultiplicativity this implies that  $||G_N|| \leq 1$ . Moreover,

$$\left\| \exp\left(\frac{m_r X^r}{r}\right) - 1 \right\| = \sup_{n \ge 1} \left| m_r^n \frac{(rn)!}{r^n n!} \right|_p \le \sup_{n \ge 1} \left| \prod_{\substack{1 \le k \le rn \\ k \ne r, \mathbb{Z}}} k \right|_p = \left| (r-1)! \right|_p.$$

Putting these together,

$$||G_N - G_{N-1}|| \le ||G_{N-1}|| \left| \exp\left(\frac{m_N X^N}{N}\right) - 1 \right| \le |(N-1)!|_p.$$

By the ultrametric triangle inequality, whenever  $N \leq M$  we have

$$||G_M - G_N|| \le \max_{N+1 \le i \le M} ||G_i - G_{i-1}|| \le \max_{N+1 \le i \le M} |(i-1)!|_p \le |N!|_p.$$

The upper bound goes to zero as  $N \to \infty$ . This verifies that  $(G_N)_{N \ge 1}$  is a Cauchy sequence and therefore converges to  $\zeta_m \in V$ .

Proof of Theorem 3.6. We can factor  $\zeta_m$  as

$$\zeta_m = \exp(m_1 X) \exp\left(\frac{m_p X^p}{p}\right) \prod_{r \neq 1, p} \exp\left(\frac{m_r X^r}{r}\right).$$

Let  $m^{\underline{n}} = m(m-1)(m-2)\cdots(m-n+1)$  denote the *n*th falling factorial of m. Note that  $|m^{\underline{n}}|_p \to 0$  as  $n \to \infty$  for any  $m \in \mathbb{Z}_p$ ; hence

$$(1-X)^m = \sum_{n\geq 0} {m \choose n} (-1)^n X^n = \sum_{n\geq 0} (-1)^n m^n \frac{X^n}{n!}$$

is in V for any  $m \in \mathbb{Z}_p$ . We also have the power series identity  $\prod_{r \geq 1} \exp\left(-\frac{X^r}{r}\right) = 1 - X$ .

We see that V contains  $\zeta_m$  if and only if V contains

$$\zeta_m \cdot (1 - X)^{m_p} = \exp(m_1 X) \exp\left(\frac{m_p X^p}{p}\right) \prod_{r \neq 1, p} \exp\left(\frac{m_r X^r}{r}\right) \prod_{r > 1} \exp\left(\frac{-m_p X^r}{r}\right)$$

$$= \exp((m_1 - m_p)X) \prod_{r \neq 1, p} \exp\left(\frac{(m_r - m_p)X^r}{r}\right).$$

By Lemma 3.8,  $\prod_{r\neq 1,p} \exp\left(\frac{(m_r-m_p)X^r}{r}\right)$  is an invertible element of V, so V contains  $\zeta_m$  if and only if it contains  $\exp((m_1-m_p)X)$ . By Lemma 3.7,  $\exp((m_1-m_p)X)$  is in V if and only if  $|m_1-m_p|_p < 1$ .

Proof of Theorem 3.2. It is a well-known symmetric function identity that

$$\sum_{n=0}^{\infty} h_n X^n = \exp\left(\sum_{k=1}^{\infty} \frac{p_k}{k} X^k\right)$$

where  $h_n = \sum_{1 \leq i_1 \leq \cdots \leq i_n} x_{i_1} \cdots x_{i_n}$  and  $p_k = \sum_{i \geq 1} x_i^k$ . The set  $\{h_n\}_n$  is algebraically independent in  $\mathbb{Q}[\{x_i\}_i]$ , and by specializing  $h_n \mapsto f(n)/n!$  we obtain

$$\sum_{n=0}^{\infty} \frac{f(n)}{n!} X^n = \exp\left(\sum_{k=1}^{\infty} \frac{m_k}{k} X^k\right) = \zeta_m.$$

Let  $c_n = \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} f(k)$  be the *n*th finite difference of f. Then by (7)

$$1 + c_1 X + c_2 \frac{X^2}{2!} + \dots = e^{-X} \zeta_m,$$

so f is p-adically continuous if and only if  $e^{-X}\zeta_m$  is in V. Applying Theorem 3.6 to  $m_1 - 1, m_2, m_3, \ldots$  shows that  $e^{-X}\zeta_m$  is in V if and only if

$$f(1) - 1 = m_1 - 1 \equiv m_p \pmod{p}.$$

3.3. Cycle-restricted permutations. A derangement is defined as a permutation with no cycle of length one. In this section we consider a generalization by placing an arbitrary restriction on the cycle lengths of a permutation. Let  $S_n$  denote the symmetric group on n letters.

Given a set L of positive integers, let

$$d_n^L = \#\{\sigma \in S_n : \text{cycle lengths of } \sigma \text{ are in } L\}.$$

We show that the functions  $n \mapsto d_n^L$  for different choices of L are precisely the functions whose m-values (cf. (1)) are all in  $\{0,1\}$ .

**Theorem 3.9** (Theorem 1.4). Let  $f: \mathbb{N} \to \mathbb{Q}$  be a function whose m-values (cf. (1)) are all in  $\{0,1\}$ . Then f(n) is equal to the number  $d_n^L$  of permutations in  $S_n$  whose cycle lengths are in  $L = \{r : m_r = 1\}$ .

Combining the above theorem with Theorem 3.2 results in the following p-adic interpolation result for counts of cycle-restricted permutations. (Note that  $n \mapsto (-1)^n$  is 2-adically continuous, so part (c) holds equally for  $n \mapsto (-1)^n d_n^L$ .)

**Theorem 3.10** (Theorem 1.5). Let L be a set of positive integers, and let  $p \geq 3$  be prime. Let  $d_n^L$  denote the number of permutations in  $S_n$  whose cycle lengths are in L.

- (a)  $n \mapsto d_n^L$  is p-adically continuous if and only if  $1 \in L$  and  $p \notin L$ .
- (b)  $n \mapsto (-1)^n d_n^L$  is p-adically continuous if and only if  $1 \notin L$  and  $p \in L$ .
- (c)  $n \mapsto d_n^L$  is 2-adically continuous if and only if one of the following hold:
  - $1 \in L$  and  $2 \notin L$ , or
  - $1 \notin L$  and  $2 \in L$ .

This theorem encompasses a large class of sequences. In the following examples, we write  $P_L$  for the set of primes for which the displayed sequence extends to a continuous function on  $\mathbb{Z}_p$ .

**Example 2** (Squares). If  $L = \{1, 4, 9, 16, \ldots\}$  consists of the perfect squares, then

$$(d_n^L) = (1, 1, 1, 1, 7, 31, 91, 211, 1681, 52417, \ldots)$$

and  $P_L = \{\text{all primes}\}.$ 

**Example 3** (Non-squares). If  $L = \mathbb{N}^{>0} \setminus \{1, 4, 9, 16, \ldots\}$ , then

$$((-1)^n d_n^L) = (1, 0, 1, -2, 3, -44, 175, -1434, 12313, -59912, \ldots)$$

and  $P_L = \{\text{all primes}\}.$ 

**Example 4** (Powers of 3). If  $L := \{1, 3, 9, 27, \ldots\}$ , then

$$(d_n^L) = (1, 1, 1, 3, 9, 21, 81, 351, 1233, 46089, \ldots)$$

and  $P_L = \{\text{all primes except } 3\}.$ 

**Example 5** (Complement of powers of 3). If  $L := \mathbb{N}^{>0} \setminus \{1, 3, 9, 27, \ldots\}$ , then

$$((-1)^n d_n^L) = (1, 0, 1, 0, 9, -24, 225, -1224, 11025, -53136, \ldots)$$

and  $P_L = \{\text{all primes except } 3\}.$ 

**Example 6** (Classical derangements). If  $L := \{2, 3, 4, 5, \ldots\}$ , then

$$((-1)^n d_n^L) = (1, 0, 1, -2, 9, -44, 265, -1854, 14833, -133496, \ldots)$$

are the classical derangement numbers (up to sign) and  $P_L = \{\text{all primes}\}.$ 

**Example 7** (Prime numbers and complement). If  $L := \{2, 3, 5, 7, \ldots\}$  consists of the prime numbers, then

$$((-1)^n d_n^L) = (1, 0, 1, -2, 3, -44, 55, -1434, 3913, -39752, \dots)$$

and  $P_L = \{\text{all primes}\}.$  If  $L' := \mathbb{N}^{>0} \setminus L$ , then

$$(d_n^{L'}) = (1, 1, 1, 1, 7, 31, 211, 1051, 10081, 107857, \ldots)$$

and  $P_{L'} = \{\text{all primes}\}\ \text{also.}$ 

**Example 8** (Artin–Hasse complement). If  $L_{(p)} = \{p, p^2, p^3, \ldots\}$ , then  $(-1)^n d_n^{L_{(p)}}$  is the *n*th EGF-coefficient of  $e^X E_p(-X)$  and  $P_{L_{(p)}} = \{p\}$ . For instance, if p = 3 we have

$$e^{X}E_{3}(-X) = 1 - \frac{1}{3}X^{3} + \frac{1}{18}X^{6} - \frac{19}{162}X^{9} + \frac{73}{1944}X^{12} - \frac{181}{29160}X^{15} + O(X^{18})$$

and

$$((-1)^n d_n^{L_{(3)}}) = (1, 0, 0, -2, 0, 0, 40, 0, 0, -42560, 0, 0, 17987200, 0, 0, \ldots).$$

Theorem 3.9 and Theorem 3.10 are proven using the following formal power series identity.

**Proposition 3.11.** Let  $d_n^L$  denote the number of permutations in  $S_n$  whose cycle lengths are in L. Then the exponential generating function of  $d_n^L$  factors as

$$\sum_{n>0} d_n^L \frac{X^n}{n!} = \prod_{r \in L} \exp\left(\frac{X^r}{r}\right). \tag{9}$$

For the proof it is convenient to use combinatorial species (cf. [20, §1] for a quick introduction or [5] for a comprehensive treatment). In the next proof, "generating function of a species" always means the exponential generating function.

*Proof.* For a positive integer r, consider the subset of permutations in  $S_n$  equal to an r-cycle (empty if  $n \neq r$ ). We have

$$\#\{\sigma \in S_n : \sigma \text{ is an } r\text{-cycle}\} = \begin{cases} (r-1)! & \text{if } n=r, \\ 0 & \text{otherwise.} \end{cases}$$

The generating function of this sequence is  $\frac{1}{r}X^r$ . By the theory of species,  $\exp(X^r/r)$  is the generating function of the sequence

$$n \mapsto \#\{\sigma \in S_n : \sigma \text{ decomposes into disjoint } r\text{-cycles}\}.$$

A standard result from the theory of species says that the product of the generating functions of two species A and B is the generating function of the species where we partition [n] into disjoint subsets  $S \sqcup T$ , and impose an A-species on S and a B-species on T. This generalizes by induction to finite products.

Consider the species

$$[n] \mapsto \begin{cases} \{ \sigma \in S_n : \text{ every cycle length of } \sigma \text{ is in } L \} & \text{if } n \leq N, \\ \emptyset & \text{otherwise.} \end{cases}$$

The generating function of this species is  $\sum_{n=0}^{N} d_n^L \frac{X^n}{n!}$ . By the previous paragraphs, this generating function factors as

$$\sum_{n=0}^{N} d_n^L \frac{X^n}{n!} = \prod_{\substack{r \in L \\ r < N}} \exp\left(\frac{X^r}{r}\right) = \prod_{r \in L} \exp\left(\frac{X^r}{r}\right) \pmod{X^{N+1}}.$$

Taking the limit as  $N \to \infty$  in the X-adic topology verifies (9).

Proof of Theorem 3.10. We apply the criterion of Theorem 3.2, that a function on  $\mathbb{N}$  with integral  $m_k$ -values for  $k \geq 2$  is p-adically continuous if and only if its m-values satisfy  $m_p \equiv m_1 - 1 \pmod{p}$ . Let  $\alpha \in \mathbb{Z}$  and consider the function  $n \mapsto \alpha^n d_n^L$ . This function has m-values

$$m_k = \begin{cases} \alpha^k & \text{if } k \in L, \\ 0 & \text{otherwise.} \end{cases}$$

Let  $\chi_L$  denote the characteristic function of L, so that  $m_k = \alpha^k \chi_L(k)$ . Then  $n \mapsto \alpha^n d_n^L$  is p-adically continuous if and only if  $m_1 - m_p \equiv 1 \pmod{p}$ , i.e.

$$\alpha \chi_L(1) - \alpha^p \chi_L(p) \equiv \alpha(\chi_L(1) - \chi_L(p)) \equiv 1 \pmod{p}.$$

The theorem now follows easily from this congruence for  $\alpha \in \{\pm 1\}$ .

## 4. Cyclic arrangements and derangements

In this section we study the p-adic properties of a(n,r) and demonstrate a formula for a(n,r) involving the floor function (Theorem 4.6). We prove that a(n,r) is p-adically continuous as a function of two variables (Theorem 4.5), and give a precise description of the p-adic regularity of  $n \mapsto a(n,r)$  in terms of  $|r|_p$  (Theorem 4.3). The results of this section will be used to construct our p-adic analogue of the incomplete gamma function in §5.

4.1. Wreath products. Let r and n be positive integers. Let  $C_r$  denote the cyclic group of order r. The symmetric group  $S_n$  acts by precomposition on the set  $\operatorname{Fun}([n], C_r) = \{f : [n] \to C_r\}$  of functions, which is itself a group under pointwise operations. The wreath product is defined by

$$C_r \wr S_n := \operatorname{Fun}([n], C_r) \rtimes S_n.$$

This group has a natural permutation action on the set  $C_r \times [n]$  given by

$$(f,\sigma)(s,m) = (s + f(\sigma(m)), \sigma(m)). \tag{10}$$

The group  $C_r \wr S_n$  can also be represented as the group of complex  $n \times n$  matrices generated by diagonal matrices of order r and permutation matrices. In this representation, the action on the set  $C_r \times [n]$  defined above is isomorphic to matrix multiplication on the set  $\{\zeta^i e_j : i = 0, 1, \ldots, r - 1, j = 1, 2, \ldots, n\}$  where  $\zeta$  is a primitive rth root of unity and  $e_i$  is a standard basis vector of  $\mathbb{C}^n$ .

**Definition 4.1.** An r-cyclic derangement of degree n is an element of  $C_r \wr S_n$  whose action on  $C_r \times [n]$  has no fixed point. An r-cyclic arrangement of degree n is a choice of subset  $A \subset [n]$  and an element of  $C_r \wr S_{|A|}$ .

**Example 9.** There are five 2-cyclic derangements of degree 2. They are given by the matrices

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}.$$

There are three 2-cyclic arrangements of degree 1. In the notation (A, f) with  $f \in C_r \wr S_{|A|}$  in its matrix form, they are given by  $(\varnothing, ()), ([1], (1)), ([1], (-1))$ .

Recall that we have defined the combinatorial function (cf. (3))

$$a(n,r) = \begin{cases} \# \text{ of } r\text{-cyclic arrangements of degree } n & \text{if } r > 0, \\ (-1)^n (\# \text{ of } |r|\text{-cyclic derangements of degree } n) & \text{if } r < 0. \end{cases}$$

We begin by deriving a formula for a(n,r) which makes sense for nonintegral values of r.

**Lemma 4.2.** For any nonzero integer r and positive integer n,

$$a(n,r) = \sum_{k=0}^{n} \binom{n}{k} k! r^k. \tag{11}$$

*Proof.* We use a standard inclusion-exclusion argument. When r is positive the formula follows by summing  $\#(C_r \wr S_k) = k! r^k$  over the possible cardinalities k of the  $\binom{n}{k}$  choices of subset  $A \subset [n]$ . Suppose r is negative. Observe from (10) that the set of fixed points of  $(f, \sigma)$  is equal to  $C_{|r|} \times A$  where  $A = \{m \in [n] : \sigma(m) = m, f(m) = 0\}$ . Then

$$\#(|r|\text{-cyclic derangements}) = \#(x \in C_{|r|} \wr S_n) - \sum_{i \in [n]} \#(x \text{ which fix } C_{|r|} \times \{i\})$$

$$+ \sum_{\{i,j\} \subset [n]} \#(x \text{ which fix } C_{|r|} \times \{i,j\}) - \cdots$$

$$= \sum_{k=0}^{n} (-1)^k \sum_{\substack{A \subset [n] \\ |A| = k}} \#(x \in C_{|r|} \wr S_n \text{ which fix } C_{|r|} \times A).$$

For any subset  $A \subset [n]$ , the subgroup of  $C_{|r|} \wr S_n$  fixing every element of  $C_{|r|} \times A$  is  $C_{|r|} \wr S_{n-\#A}$  which has cardinality  $(-r)^{n-\#A}(n-\#A)!$ . Thus  $(-1)^n a(n,r)$  is equal to

$$\#(|r|\text{-cyclic derangements}) = \sum_{k=0}^{n} (-1)^k \binom{n}{k} (-r)^{n-k} (n-k)!$$
$$= (-1)^n \sum_{k=0}^{n} \binom{n}{k} r^k k!.$$

We henceforth use (11) to define a(n,r) for all  $r \in \mathbb{C}_p$  and  $n \in \mathbb{N}$ .

4.2. p-adic interpolation of a(n,r). In this section we prove that a(n,r) is interpolated by a two-variable p-adically continuous function. First we interpolate  $n \mapsto a(n,r)$  for fixed  $r \in \mathbb{C}_p^{\times}$  (Theorem 4.3), and afterwards we interpolate a(n,r) in both variables (Theorem 4.5).

**Theorem 4.3.** Let  $r \in \mathbb{C}_p^{\times}$  and let  $a(\cdot, r) \colon \mathbb{N} \to \mathbb{C}_p$  be defined by (11). Then  $n \mapsto a(n, r)$  extends to an analytic function (resp. metric map, locally analytic function) on the closed unit disk if and only if  $|r|_p < 1$  (resp.  $|r|_p \le 1$ ,  $|r|_p < p^{(p-1)^{-1}}$ ). If  $|r|_p \ge p^{(p-1)^{-1}}$  then  $n \mapsto a(n, r)$  does not extend to a continuous function on  $\mathbb{Z}_p$ .

**Corollary 4.4.** For any prime p and integer r, the function  $n \mapsto a(n,r)$  extends to a locally analytic metric map on  $\mathbb{Z}_p$ . If p divides r, then  $n \mapsto a(n,r)$  extends to an analytic metric map on  $\mathbb{Z}_p$ .

**Remark 1.** We note that  $n \mapsto a(n,r)$  has integral m-values if  $r \in \mathbb{Z}_p$  (cf. (1)), and is therefore encompassed by Theorem 3.2. Indeed from (7) we have

$$\sum_{n=0}^{\infty} a(n,r) \frac{X^n}{n!} = e^X \cdot \sum_{n=0}^{\infty} n! r^n \frac{X^n}{n!} = e^X (1 - rX)^{-1}$$
$$= \exp\left( (1 + r) \frac{X}{1} + r^2 \frac{X^2}{2} + r^3 \frac{X^3}{3} + \cdots \right)$$

from which we can read off the *m*-values. The required congruence (2) for continuity is satisfied:  $m_p = r^p \equiv r = (r+1) - 1 \pmod{p\mathbb{Z}_p}$ .

Proof of Theorem 4.3. The kth finite difference of  $n \mapsto a(n,r)$  is  $c(k) = r^k k!$ . By Theorem 2.2, a(n,r) extends to a p-adic analytic function on the closed unit disk if and only if  $|c(n)/n!|_p = |r|_p^n \to 0$  as  $n \to \infty$ , and this occurs if and only if  $|r|_p < 1$ . This proves the first equivalence.

Now observe that  $p^{\lfloor \frac{\log n}{\log p} \rfloor}$  is the largest power of p that is less than or equal to n. It follows that

$$\sup_{n \ge 1} |n!|_p \, p^{\lfloor \frac{\log n}{\log p} \rfloor} \le 1.$$

Clearly, if  $|r|_p \leq 1$  then we have

$$\sup_{n\geq 1} |c(n)|_p p^{\lfloor \frac{\log n}{\log p} \rfloor} = \sup_{n\geq 1} |r^n n!|_p p^{\lfloor \frac{\log n}{\log p} \rfloor} \leq 1,$$

in which case a(n,r) extends to a metric map on  $\mathbb{Z}_p$  by Theorem 2.2. On the other hand, if  $|r|_p > 1$  then

$$\sup_{n>1} |r^n n!|_p p^{\lfloor \frac{\log n}{\log p} \rfloor} \ge |r|_p > 1,$$

showing that a(n,r) does not extend to a metric map on  $\mathbb{Z}_p$ .

Now we prove the third equivalence. Using Legendre's formula for  $v_p(n!)$  we obtain  $v_p(c(n)) = n ((p-1)^{-1} + v_p(r)) - s_p(n)(p-1)^{-1}$  where  $s_p(n)$  is the sum of the digits of n in base-p. In particular, we have

$$v_p(c(n)) = n\left((p-1)^{-1} + v_p(r)\right) - O(\log n).$$
(12)

Thus

$$\liminf_{n \to \infty} \left( \frac{v_p(c(n))}{n} \right) = (p-1)^{-1} + v_p(r).$$

By Amice's theorem (Theorem 2.2), a(n,r) extends to a p-adic locally analytic function on the closed unit disk if and only if  $(p-1)^{-1} + v_p(r) > 0$ , or equivalently,  $|r|_p < p^{(p-1)^{-1}}$ .

For the last claim, from (12) it is clear that  $v_p(c(n)) \to -\infty$  if  $(p-1)^{-1} + v_p(r) < 0$ , or equivalently, if  $|r|_p > p^{(p-1)^{-1}}$ . On the other hand, if  $(p-1)^{-1} + v_p(r) = 0$  then

$$v_p(c(n)) = -s_p(n)(p-1)^{-1}$$
.

Since  $s_p(p^N) = 1$  for any positive integer N, we see that in this case  $v_p(c(n))$  does not go to  $\infty$  as  $n \to \infty$ . We conclude that a(n,r) does not admit an extension to a continuous function on  $\mathbb{Z}_p$  when  $|r|_p \geq p^{(p-1)^{-1}}$ .

Recall a two-variable function  $f: \mathbb{Z}_p \times \mathbb{Z}_p \to \mathbb{Z}_p$  is a metric map if for all x, y, r, s,

$$|f(x,r) - f(y,s)|_p \le \max\{|x - y|_p, |r - s|_p\}.$$

**Theorem 4.5.** The function  $\mathbb{N} \times \mathbb{N}^{>0} \to \mathbb{Z}_p$ :  $(n,r) \mapsto a(n,r)$  extends to a metric map  $\tilde{a}: \mathbb{Z}_p \times \mathbb{Z}_p \to \mathbb{Z}_p$ . Explicitly, it is given by the convergent series

$$a_p(y,r) = 1 + \sum_{k=1}^{\infty} r^k y(y-1) \cdots (y-k+1).$$

*Proof.* Let  $(x,r), (y,s) \in \mathbb{Z}_p \times \mathbb{Z}_p$ . The function  $n \mapsto a(n,r)$  admits a continuous extension in the first variable (denoted  $a_p(\cdot,r)$ ) by Theorem 4.3. From (11), the Mahler series of  $a_p(\cdot,r)$  is

$$a_p(y,r) = \sum_{k=0}^{\infty} r^k k! \binom{y}{k} = 1 + \sum_{k=1}^{\infty} r^k y^{\underline{k}}$$
 (13)

where  $y^{\underline{k}} := y(y-1)\cdots(y-k+1)$ . This sum is absolutely convergent since  $a_p(\cdot,r)$  is continuous. This shows that the assignment  $(y,r)\mapsto a_p(y,r)$  is well-defined.

By absolute convergence we can rearrange (13) to obtain

$$|a_p(y,r) - a_p(y,s)|_p = \Big| \sum_{k=1}^{\infty} (r^k - s^k) y^{\underline{k}} \Big|_p$$

$$\leq \max_{k \geq 1} \left( |r^k - s^k|_p |y^{\underline{k}}|_p \right)$$

$$\leq |r - s|_p.$$
(14)

In the last inequality, we have used that  $|r^k - s^k|_p \le |r - s|_p$  and  $|y^k|_p \le 1$  for  $k \ge 1$  and  $y, r, s \in \mathbb{Z}_p$ .

We have the inequality  $|a_p(x,r) - a_p(y,r)|_p \le |x-y|_p$  since  $a_p(\cdot,r)$  is a metric map by Theorem 4.3. Combining this inequality with (14) we obtain

$$|a_p(x,r) - a_p(y,s)|_p \le \max\{|a_p(x,r) - a(y,r)|_p, |a_p(y,r) - a_p(y,s)|_p\} \le \max\{|x-y|_p, |r-s|_p\}.$$

This inequality completes the proof.

4.3. An archimedean formula for a(n,r). The conclusions of Theorem 4.3 and Theorem 4.5 are rather surprising in view of the following beautiful formula for a(n,r) in terms of the floor function.

**Theorem 4.6.** Let r be a nonzero integer and let n be a positive integer. Then

$$a(n,r) = \begin{cases} \lfloor e^{1/r} r^n n! + \frac{1}{2} \rfloor & \text{if } r < 0, \\ \lfloor e^{1/r} r^n n! \rfloor & \text{if } r > 0. \end{cases}$$
 (15)

The cases  $r \in \{\pm 1\}$  are well-known (e.g. [12]). In view of its p-adic properties, the function a(n,r) thus exhibits a surprising harmony between the p-adic and real topologies, the latter of which is used in (15) to define the ordering on  $\mathbb{R}$ .

**Proposition 4.7.** Let r be a positive integer and let a(n,r) denote the number of r-cyclic arrangements of degree n. Then

$$a(n,r) = \lfloor e^{\frac{1}{r}} r^n n! \rfloor \quad (n \in \mathbb{N}^{>0}).$$

*Proof.* Observe that

$$e^{\frac{1}{r}}r^{n}n! = r^{n}n! \left(1 + r^{-1} + \frac{r^{-2}}{2!} + \frac{r^{-3}}{3!} + \cdots\right)$$

$$= r^{n}n! + r^{n-1}(n-1)! \binom{n}{1} + r^{n-2}(n-2)! \binom{n}{2} + \cdots + r\binom{n}{n-1} + 1$$

$$+ \frac{r^{-1}}{n+1} + \frac{r^{-2}}{(n+1)(n+2)} + \cdots$$

Let  $R_{r,n}$  denote the 'remainder term'

$$R_{r,n} = \sum_{k>1} \frac{n!}{(n+k)!} r^{-k} = \frac{r^{-1}}{n+1} + \frac{r^{-2}}{(n+1)(n+2)} + \cdots$$
 (16)

We have  $e^{1/r}r^n n! = a(n,r) + R_{r,n}$ . Each summand of  $R_{r,n}$  is a decreasing function of r and a decreasing function of n. Since  $r, n \ge 1$ , we have  $R_{r,n} \le R_{1,1} = e - 2 < 1$ . It follows that

$$\lfloor e^{\frac{1}{r}}r^n n! \rfloor = \lfloor a(n,r) + R_{r,n} \rfloor = a(n,r).$$

**Proposition 4.8.** Let r be a negative integer and let a(n,r) denote the quantity  $(-1)^n$  (number of |r|-cyclic derangements of degree n). Then

$$a(n,r) = \lfloor e^{\frac{1}{r}} r^n n! + \frac{1}{2} \rfloor \quad (n \in \mathbb{N}^{>0}).$$

*Proof.* Continue letting  $R_{r,n}$  be as in (16) so that  $e^{1/r}r^n n! = a(n,r) + R_{r,n}$ . For all positive integers n we have

$$-R_{r,n} = \frac{|r|^{-1}}{n+1} - \frac{|r|^{-2}}{(n+1)(n+2)} + \frac{|r|^{-3}}{(n+1)(n+2)(n+3)} - \cdots$$

$$= \sum_{\ell=1}^{\infty} \frac{|r|^{-2\ell+1}}{(n+1)(n+2)\cdots(n+2\ell)} \Big( (n+2\ell) - |r|^{-1} \Big). \tag{17}$$

Clearly each summand of (17) is positive, so  $R_{r,n}$  is negative. We will show that  $|R_{r,n}|$  is a decreasing function of n by considering each summand individually.

Regarding the summand of (17) as a function of a continuous variable n, the expression has log-derivative

$$\frac{d}{dn} \left( \log(|r|^{-2\ell+1}) + \log(n+2\ell - |r|^{-1}) - \sum_{k=1}^{2\ell} \log(n+k) \right)$$

$$= \frac{1}{n+2\ell - |r|^{-1}} - \sum_{k=1}^{2\ell} \frac{1}{n+k}. \quad (18)$$

Since  $|r| \ge 1$ , we have the bounds

$$\frac{1}{n+2\ell-|r|^{-1}} \le \frac{1}{n+2\ell-1} < \sum_{k=1}^{2\ell} \frac{1}{n+k},$$

which imply that (18) is negative. Therefore each (positive) summand in (17) is decreasing as a function of n.

This shows in particular that  $|R_{r,n}| \leq |R_{r,1}|$ . We have

$$|R_{r,1}| = 1 + r - re^{\frac{1}{r}} \in (0, \frac{1}{2})$$
 for  $r \le -1$ .

Thus we have  $e^{\frac{1}{r}}r^n n! = a(n,r) + R_{r,n}$  with  $R_{r,n} \in (-\frac{1}{2},0)$ . This shows that for all  $n \geq 1$ ,

$$\left\lfloor e^{\frac{1}{r}}r^n n! + \frac{1}{2} \right\rfloor = \left\lfloor a(n,r) + R_{r,n} + \frac{1}{2} \right\rfloor = a(n,r).$$

## 5. The p-adic incomplete gamma function

In 1975, Morita [17] discovered a p-adic counterpart to the gamma function. In this section, we show that the incomplete gamma function also has a p-adic counterpart.

5.1. The classical incomplete gamma function. We recall the classical construction of the incomplete gamma function. For a complex number s let  $\omega_s$  denote the differential given by

$$\omega_s(t) := t^s e^{-t} \frac{dt}{t}$$

where we take the principal value for  $t^s = \exp(s \log t)$ . We choose the branch cut  $(-\infty, 0]$  so that  $\omega_s$  is holomorphic and single-valued on the domain  $U = \mathbb{C} \setminus (-\infty, 0]$ . For any oriented half-infinite curve  $\gamma$  in U tending towards positive

infinity, the integral  $\int_{\gamma} \omega_s$  converges and defines a holomorphic function of s. We thus obtain, for any complex number  $z \in U$ , the entire function of s given by

$$\Gamma(s,z) := \int_{\gamma} \omega_s$$

where  $\gamma$  is any curve in U beginning at z and tending to positive infinity. This defines the *incomplete gamma function*  $\Gamma \colon \mathbb{C} \times U \to \mathbb{C}$ . By Hartog's theorem on separate analyticity,  $\Gamma(s,z)$  is an analytic function of two variables.

The incomplete gamma function recovers the gamma function in the limit as z goes to zero:

$$\lim_{z \to 0^+} \Gamma(s, z) = \Gamma(s).$$

In general, the incomplete gamma function takes different values as z approaches the branch cut. For  $s \in \mathbb{C}$  and z on the negative real axis we have

$$\Gamma(s, z + i0^{+}) = e^{2\pi i s} \Gamma(s, z - i0^{+}) + (1 - e^{2\pi i s}) \Gamma(s)$$
(19)

where  $f(0^+) := \lim_{\epsilon \to 0^+} f(\epsilon)$ ; see [9, Eq. 8.2.9]. If s is a positive integer this shows  $\Gamma(s,z)$  extends continuously in the second variable across the branch cut, and  $\Gamma$  extends to the domain  $\Gamma : (\mathbb{C} \times U) \cup (\mathbb{N}^{>0} \times \mathbb{C}) \to \mathbb{C}$ .

The incomplete gamma function is closely related to the combinatorial function a(n,r) counting cyclic derangements and arrangements. Just as the count of all permutations is interpolated by the gamma function, the counts of cyclic derangements and cyclic arrangements are interpolated by the incomplete gamma function (up to an explicit entire function). Recall from §4 that we extended the combinatorial definition of a(n,r) to  $a(n,r) = \sum_{k=0}^{n} {n \choose k} k! r^k$  for all  $r \in \mathbb{C}_p$ .

**Proposition 5.1.** Let  $z \in \mathbb{C}^{\times}$  be fixed. The function  $s \mapsto \Gamma(1+s,z)z^{-s}e^{z}$  is holomorphic for all  $s \in \mathbb{C}$ . For  $n \in \mathbb{N}$  we have  $\Gamma(1+n,z)z^{-n}e^{z} = a(n,1/z)$ .

$$(\Gamma(1+n,z))$$
 is defined for  $(n,z) \in \mathbb{N} \times \mathbb{C}$  by (19).)

*Proof.* The function  $z^{-s} = \exp(-s \log z)$  is entire in the argument s, where we take the principal value for the logarithm to determine the constant  $\log z$ , and  $\Gamma(1+s,z)$  is entire in s. This proves the first part of the proposition.

By [9, Eq. 8.7.3], for any  $(s,z) \in \mathbb{N}^{>0} \times \mathbb{C}$  we have

$$\Gamma(s,z) = \Gamma(s) \left( 1 - z^s e^{-z} \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(s+k+1)} \right).$$

Using that  $\Gamma(s+k+1)\Gamma(s)^{-1}=(s+k)\cdots(s+1)s$  and writing s=1+n we have

$$\Gamma(1+n,z) = n! \left( 1 - z^{1+n} e^{-z} \sum_{k=0}^{\infty} \frac{z^k}{(n+k+1)!} \right)$$

$$= n! \left( 1 - e^{-z} \sum_{k=n+1}^{\infty} \frac{z^k}{k!} \right)$$

$$= n! e^{-z} \left( e^z - \sum_{k=n+1}^{\infty} \frac{z^k}{k!} \right)$$

$$= n! e^{-z} \sum_{k=0}^{n} \frac{z^k}{k!}$$

$$= e^{-z} \sum_{k=0}^{n} \frac{n!}{(n-k)!} z^{n-k} = a(n, 1/z) z^n e^{-z}.$$

Rearranging terms proves the proposition.

Corollary 5.2. For any  $z \in \mathbb{C}^{\times}$ ,  $\mathbb{Q}(\{\Gamma(n,z) : n \in \mathbb{N}^{>0}\}) = \mathbb{Q}(z,e^z)$ .

*Proof.* From Proposition 5.1 it is easily seen that for any positive integer n,  $\Gamma(n,z)$  is a polynomial function in z and  $e^z$ . This shows that  $\mathbb{Q}(\{\Gamma(n,z):n\in\mathbb{N}^{>0}\})\subset\mathbb{Q}(z,e^z)$ . Moreover,  $e^{-z}=\Gamma(1,z)$  and  $z=\frac{\Gamma(2,z)}{\Gamma(1,z)}-1$ , proving that  $\mathbb{Q}(z,e^z)\subset\mathbb{Q}(\{\Gamma(n,z):n\in\mathbb{N}^{>0}\})$ .

5.2. The p-adic incomplete gamma function. We will interpolate the incomplete gamma function on its integral values  $(n,r) \in \mathbb{N}^{>0} \times \mathbb{Z}$ . Before doing so, it is necessary to realize the integral values of the incomplete gamma function (which are transcendental elements of  $\mathbb{C}$ ) as p-adic numbers. As we shall see, there is a reasonably natural way of doing this. The field generated by the integral values of the incomplete gamma function is

$$\mathbb{Q}(\{\Gamma(n,r):(n,r)\in\mathbb{N}^{>0}\times\mathbb{Z}\})=\mathbb{Q}(e)$$

by Corollary 5.2. Therefore, to obtain a homomorphism from the field generated by  $\{\Gamma(n,r):(n,r)\in\mathbb{N}^{>0}\times\mathbb{Z}\}$  to the field of p-adic numbers, it suffices to specify a transcendental element of  $\mathbb{Q}_p$  corresponding to e. Let  $\tau_p:\mathbb{Q}(e)\to\mathbb{Q}_p$  be the unique field homomorphism satisfying

$$\tau_p(e^{-1}) = \exp_p(p) = 1 + p + p^2/2! + p^3/3! + \cdots$$

The element  $\exp_p(p) \in \mathbb{Q}_p$  is transcendental over  $\mathbb{Q}$  (cf. [1, §3]).

**Theorem 5.3** (Theorem 1.1). There exists a continuous function

$$\Gamma_p \colon \mathbb{Z}_p \times (1 + p\mathbb{Z}_p) \to \mathbb{Z}_p$$

satisfying

$$\Gamma_p(n,r) = \tau_p \Gamma(n,r)$$

for any positive integers n and r satisfying  $r \equiv 1 \pmod{p}$ .

*Proof.* We have that

$$\tau_p \Gamma(n,r) = \tau_p \left( a(n-1,1/r)r^{n-1}e^{-r} \right) = a(n-1,1/r)r^{n-1}\tau_p(e^{-r})$$

by Proposition 5.1. We will show that each of the three functions  $(n,r) \mapsto a(n-1,1/r)$ ,  $(n,r) \mapsto r^{n-1}$ , and  $r \mapsto \tau_p(e^{-r})$  is interpolated by a uniformly continuous function in the *p*-adic topology.

We have already proven in Theorem 4.5 that  $(n,r) \mapsto a(n,r)$  extends to the metric map  $\mathbb{Z}_p \times \mathbb{Z}_p \to \mathbb{Z}_p$  given by the Mahler series (13)

$$a(y,s) = \sum_{k=0}^{\infty} s^k k! {y \choose k} \text{ for } (y,s) \in \mathbb{Z}_p \times \mathbb{Z}_p.$$

This suffices to prove that  $(n,r) \mapsto a(n-1,1/r)$  is interpolated by a metric map, hence interpolated by a uniformly continuous map.

Let  $\exp_p: \mathbb{Z}_p \to 1 + p\mathbb{Z}_p$  (resp.  $\log_p: 1 + p\mathbb{Z}_p \to \mathbb{Z}_p$ ) denote the p-adic exponential (resp. p-adic logarithm). Then, for any positive integers n and r satisfying  $r \equiv 1 \pmod{p}$ , we have that  $r^n = \exp_p(n \log_p r)$ . Therefore the function  $\mathbb{Z}_p \times (1 + p\mathbb{Z}_p) \to \mathbb{Z}_p: (x, s) \mapsto \exp_p(x \log_p s)$  is a continuous extension of  $\mathbb{N}^{>0} \times (1 + p\mathbb{N}) \to \mathbb{Z}_p: (n, r) \mapsto r^n$  which is also uniformly continuous since  $\mathbb{Z}_p \times (1 + p\mathbb{Z}_p)$  is compact.

 $\mathbb{Z}_p \times (1+p\mathbb{Z}_p)$  is compact. Finally,  $\mathbb{N}^{>0} \to \mathbb{Z}_p \colon r \mapsto \tau_p(e^{-r}) = \exp_p(p)^r$  is interpolated by the *p*-adic locally analytic function  $\mathbb{Z}_p \to \mathbb{Z}_p \colon r \mapsto \exp_p(pr)$  which is also uniformly continuous since  $\mathbb{Z}_p$  is compact.

Any uniformly continuous function  $f: U \to Y$  mapping to a complete metric space Y, and defined on a dense subset  $U \subset X$  of a topological space X, extends uniquely to a uniformly continuous function  $X \to Y$ . We conclude that  $\tau_p\Gamma(n,r)$  is interpolated by the uniformly continuous function

$$\Gamma_p(y,r) = r^{y-1} \exp_p(pr)a(y-1,1/r)$$
 (20)

for 
$$(y,r) \in \mathbb{Z}_p \times (1+p\mathbb{Z}_p)$$
.

**Remark 2** (*m*-values of  $\Gamma_p$ ). We can corroborate Theorem 1.1 using Theorem 3.2. For fixed  $r \in \mathbb{Z}$  the *m*-values of  $n \mapsto \exp_p(-pr)\tau_p\Gamma(n,r)$  are found to be

$$m_k = \begin{cases} r+1 & \text{if } k = 1, \\ 1 & \text{if } k \ge 2, \end{cases}$$

and the required congruence (2) for continuity is that  $r \equiv 1 \pmod{p}$ .

We collect a few facts about  $\Gamma_p$ .

**Theorem 5.4.** Let  $\Gamma_p \colon \mathbb{Z}_p \times (1+p\mathbb{Z}_p) \to \mathbb{Z}_p$  denote the p-adic incomplete gamma function of Theorem 1.1.

(1) For any  $y \in \mathbb{Z}_p$  and  $r \in 1 + p\mathbb{Z}_p$  we have the series expression

$$\Gamma_p(y,r) = \sum_{k=0}^{\infty} r^{y-k-1} \exp_p(pr)k! {y-1 \choose k}.$$

(2) For any  $n \in \mathbb{N}$  and  $r \in \mathbb{Z}$  satisfying  $r \equiv 1 \pmod{p}$ , we have

$$\Gamma_p(n+1,1/r) = r^{-n} \exp_p(p/r) \cdot \begin{cases} \lfloor e^{1/r} r^n n! + \frac{1}{2} \rfloor & \text{if } r < 0, \\ \lfloor e^{1/r} r^n n! \rfloor & \text{if } r > 0. \end{cases}$$

(3) For any  $n \in \mathbb{N}$  and  $r \in 1 + p\mathbb{Z}_p$ , we have

$$\Gamma_p(n+1,r) = \exp_p(pr)n!f_n(r)$$

where  $f_n(X)$  is the nth truncated exponential polynomial

$$f_n(X) := 1 + X + \frac{1}{2!}X^2 + \dots + \frac{1}{n!}X^n \quad (n \ge 1).$$

*Proof.* The first formula follows at once from (20) and (13). The second formula follows from (20) and (15). The third formula follows from (20) and the formula

$$a(n,r) = \sum_{k=0}^{n} \frac{n!}{(n-k)!} r^k = r^n n! \sum_{k=0}^{n} \frac{1}{k!} r^{-k} = r^n n! f_n(1/r).$$

Using a classical result of Schur on the irreducibility of truncated exponential polynomials, we can make the following observation.

**Theorem 5.5.**  $\Gamma_p$  has finitely many zeros in  $\mathbb{Z}_p \times (1 + p\mathbb{Z}_p)$ .

Observe that if  $(y,r) \in \mathbb{Z}_p \times (1+p\mathbb{Z}_p)$  is a zero of  $\Gamma_p$ , then the formula  $\Gamma_p(y,r) = r^{y-1} \exp_p(pr) a(y-1,1/r)$  shows that (y-1,1/r) is a zero of a (the formulas  $1 = r^y r^{-y} = \exp_p(x) \exp_p(-x)$  show that the other terms never vanish). The result now follows from the next proposition.

**Proposition 5.6.** Let  $r \in \mathbb{C}_p^{\times}$  and suppose  $|r|_p < p^{(p-1)^{-1}}$ . Let  $a_p(\cdot,r)$  denote the unique continuous function  $\mathbb{Z}_p \to \mathbb{C}_p$  which extends (11) (this function exists by Theorem 4.3). Then the set  $S = \{s \in \mathbb{Z}_p : a_p(s,r) = 0\}$  is finite.

A nonzero analytic function on a closed region has only finitely many zeros, in which case the result follows immediately from Theorem 4.3 when  $|r|_p < 1$ . However, a nonzero p-adic locally analytic function may have infinitely many zeros on a closed region. Thus, for  $1 \leq |r|_p < p^{(p-1)^{-1}}$ , the function  $s \mapsto a_p(s,r)$  may have infinitely many zeros in  $\mathbb{Z}_p$  a priori. The proposition shows that this does not occur.

*Proof.* By Theorem 4.3,  $a_p(\cdot, r)$  is locally analytic on the closed unit disk. This means that there is a covering of  $\mathbb{Z}_p$  by open disks such that the restriction of  $a_p(\cdot, r)$  to any one of them is given by a convergent power series with coefficients in  $\mathbb{C}_p$ . As  $\mathbb{Z}_p$  is compact, we may take this open covering to be finite.

Suppose that S is infinite for a contradiction. Then at least one disk D of the open covering must contain infinitely many zeros of  $a_p(\cdot, r)$ . By Weierstrass's Preparation Theorem, a nonzero analytic function on an open disk has only finitely many zeros (cf. e.g. [18, §9]). Thus the restriction of  $a_p(\cdot, r)$  to D is identically zero.

Let m and n be distinct integers chosen from  $D \cap \mathbb{N}$  (this is possible because  $\mathbb{N}$  is dense in  $\mathbb{Z}_p$ ). By assumption  $a_p(m,r) = a_p(n,r) = 0$ . We have

$$a_p(n,r) = \sum_{k=0}^n \frac{n!}{(n-k)!} r^k = r^n n! \sum_{k=0}^n \frac{1}{k!} r^{-k} = r^n n! f_n(1/r).$$

This implies that 1/r is a root of  $f_m(X)$  as well as  $f_n(X)$ . These polynomials are both irreducible over  $\mathbb{Q}$  by a classical theorem of Schur [25], implying that the degree of 1/r as an algebraic number is simultaneously equal to m and n, a contradiction. This implies S is finite.

5.3. Some questions. The classical gamma function  $\Gamma(s)$  is recovered from the incomplete gamma function  $\Gamma(s,z)$  in the archimedean limit as  $z \to 0^+$ . Can Morita's p-adic gamma function [17] or Diamond's p-adic log gamma function [8] be related to the p-adic incomplete gamma function? In 1984, Schikhof [24, p. 17] asked (in our notation) whether  $a_p(-1,-1) = (-1)^{p-1}d_{-1,p} = 1 + 1 + 2! + 3! + 4! + 5! + \cdots$  is in  $\mathbb{Q} \cap \mathbb{Z}_p$ . R. Murty–Sumner [19] conjecture a negative outcome to Schikhof's question. More generally, we ask the same question of  $a_p(-1,r)$ . Is there a nonzero integer r such that  $a_p(-1,r) \in \mathbb{Q} \cap \mathbb{Z}_p$ ?

There are interesting examples of functions  $f: \mathbb{N} \to \mathbb{C}_p$  whose m-values are not in  $\mathbb{Z}_p$ . Does Theorem 1.2 admit a generalization for arbitrary functions  $f: \mathbb{N} \to \mathbb{C}_p$ ? Can stronger p-adic regularity (Lipschitz continuity, differentiability, integrality of divided differences, analyticity) be detected with congruences on m-values?

### ACKNOWLEDGEMENTS

We thank the reviewer for their useful suggestions. We are also grateful to Daniel Barsky and Julian Rosen for pointing out an error on an earlier version of this manuscript. We thank Jeffrey Lagarias for some helpful comments. We also thank A. Suki Dasher for a valuable suggestion which led to the connection with the incomplete gamma function.

## References

- [1] W. W. Adams. Transcendental numbers in the p-adic domain. American Journal of Mathematics, 88(2):279–308, 1966.
- [2] Y. Amice. Interpolation p-adique. Bull. Soc. Math. France, 92:117–180, 1964.
- [3] S. H. Assaf. Cyclic derangements. Electron. J. Combin., 17(1):Research Paper 163, 14, 2010.

- [4] D. Barsky. Analyse p-adique et suites classiques de nombres. Séminaire Lotharingien de combinatoire, 1981.
- [5] F. Bergeron, G. Labelle, and P. Leroux. Combinatorial species and tree-like structures, volume 67 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 1998.
- [6] S. Chowla, I. N. Herstein, and W. R. Scott. The solutions of  $x^d = 1$  in symmetric groups. Norske Vid. Selsk. Forh., Trondheim, 25:29–31 (1953), 1952.
- [7] K. Conrad. Artin–Hasse-type series and roots of unity, 1994.
- [8] J. Diamond. The p-adic log gamma function and p-adic Euler constants. Trans. Amer. Math. Soc., 233:321–337, 1977.
- [9] NIST Digital Library of Mathematical Functions. http://dlmf.nist.gov/, Release 1.0.28 of 2020-09-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [10] D. Eric and R. Tanguy. On primary pseudo-polynomials (Around Ruzsa's Conjecture), 2021.
- [11] R. R. Hall. On pseudo-polynomials. Mathematika, 18:71–77, 1971.
- [12] M. Hassani. Derangements and applications. J. Integer Seq., 6(1):Article 03.1.2, 8, 2003.
- [13] H. Ishihara, H. Ochiai, Y. Takegahara, and T. Yoshida. p-divisibility of the number of solutions of  $x^p = 1$  in a symmetric group. Ann. Comb., 5(2):197-210, 2001.
- [14] V. Kuperberg. On pseudo-polynomials divisible only by a sparse set of primes and  $\alpha$ -primary pseudo-polynomials, 2021.
- [15] K. Mahler. An interpolation series for continuous functions of a p-adic variable. J. Reine Angew. Math., 199:23–34, 1958.
- [16] P. Miska. Arithmetic properties of the sequence of derangements. J. Number Theory, 163:114–145, 2016.
- [17] Y. Morita. A p-adic analogue of the  $\Gamma$ -function. J. Fac. Sci. Univ. Tokyo Sect. IA Math., 22(2):255-266, 1975.
- [18] M. R. Murty. Introduction to p-adic analytic number theory, volume 27 of AMS/IP Studies in Advanced Mathematics. American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2002.
- [19] M. R. Murty and S. Sumner. On the *p*-adic series  $\sum_{n=1}^{\infty} n^k \cdot n!$ . In *Number theory*, volume 36 of *CRM Proc. Lecture Notes*, pages 219–227. Amer. Math. Soc., Providence, RI, 2004.
- [20] A. M. Nelson. A generalized cyclotomic identity. Adv. Math., 83(1):1–29, 1990.
- [21] A. O'Desky. Functions with integer-valued divided differences. Journal of Number Theory, 2021.
- [22] V. Paşol and A. Zaharescu. Sondow's conjecture, convergents to e, and p-adic analytic functions. Math. Z., 292(1-2):499–511, 2019.
- [23] A. M. Robert. A course in p-adic analysis, volume 198 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2000.
- [24] W. H. Schikhof. Ultrametric calculus, volume 4 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2006.
- [25] I. Schur. Einige Sätze über Primzahlen mit Anwendungen auf Irreduzibilitätsfragen I, II. Sitzungsberichte Preuss. Akad. Wiss. Phys.-Math. Klasse, pages 125–136, 152–173, 1929.
- [26] G. Serafin. Identities behind some congruences for r-Bell and derangement polynomials. Res. Number Theory, 6(4):Paper No. 39, 8, 2020.
- [27] Y. Sun, X. Wu, and J. Zhuang. Congruences on the Bell polynomials and the derangement polynomials. *J. Number Theory*, 133(5):1564–1571, 2013.
- [28] Z.-W. Sun and D. Zagier. On a curious property of Bell numbers. Bull. Aust. Math. Soc., 84(1):153–158, 2011.

DEPARTMENT OF MATHEMATICS PRINCETON UNIVERSITY PRINCETON, NJ 08544-1000, USA

 $URL: \ \ \, \texttt{HTTPS://www.ANDREWODESKY.COM/} \\ Email \ \, address: \ \ \, \texttt{andy.odesky@gmail.com} \\$ 

DEPARTMENT OF MATHEMATICS UNIVERSITY OF WASHINGTON SEATTLE, WA 98195-4350, USA Email address: hrichman@uw.edu