ZACHARY STIER

UC Berkeley

ABSTRACT. Parzanchevski–Sarnak recently adapted an algorithm of Ross–Selinger for factorization of PU(2)-diagonal elements to within distance  $\varepsilon$  into an efficient probabilistic algorithm for any PU(2)-element, using at most  $3 \log_p \frac{1}{\varepsilon^3}$  factors from certain well-chosen sets. The Clifford+T gates are one such set arising from p = 2. In that setting, we leverage recent work of Carvalho Pinto–Petit to improve this to  $\frac{7}{3} \log_2 \frac{1}{\varepsilon^3}$ , and implement the algorithm in Haskell.

## 1. INTRODUCTION

Factoring in a matrix group given a set of topological generators is a problem of fundamental and practical significance. Because individual gates in quantum circuits acting on n qubits exist as (projective) elements in  $U(\mathbb{C}^{2^n})$ , a question of importance is to fabricate only a specific (finite) set which may then approximate any given gate to arbitrary precision; that this is possible is the Solovay–Kitaev theorem (cf. Sarnak [Sar15]). A golden gate set [Sar15, PS18] is a certain choice of these topological generators for single-qubit gates, and their construction is associated to a particular prime p. [PS18] shows that most elements of PU(2) will have a factorization to within  $\varepsilon$ (in the bi-invariant metric d on PU(2)) of length up to  $k = (1 + o(1)) \log_p \frac{1}{\varepsilon^3}$ , but that computing such factorizations for general elements is NP-complete. Instead, [PS18, §2.3] extends [RS16]'s  $(3 + o(1)) \log_2 \frac{1}{\varepsilon^3}$  efficient heuristic factorization algorithm for Clifford+T gates, to any golden gate set. The bulk of this factorization is accomplished by thrice applying an algorithm adapted from [RS16] (initially written for Clifford+T gates, defined below), each time factoring a diagonal into up to length  $(1 + o(1)) \log_p \frac{1}{\varepsilon^3}$  in topological generators.

[CP18] point out that their factorization for the discrete case of the LPS Ramanujan graphs  $X^{p,q}$ , attaining factorization length  $7 \log_p q$ , should analogize in the continuous setting with  $q \approx \frac{1}{\varepsilon}$ , and indeed it does, as we now show for the setting of Clifford+T gates. We understand that Kliuchnikov–Lauter–Minko–Paetznick–Petit have a similar extension underway [KLM<sup>+</sup>20].

In PU(2), let

$$H = \frac{i}{\sqrt{2}} \begin{pmatrix} 1 & 1\\ 1 & -1 \end{pmatrix}, \qquad \qquad S = \begin{pmatrix} 1 & \\ & i \end{pmatrix}, \qquad \qquad T = \begin{pmatrix} e^{\frac{i\pi}{8}} & \\ & e^{-\frac{i\pi}{8}} \end{pmatrix}.$$

Associated to the prime p = 2 is a golden gate set S consisting of the Clifford group  $C = \langle H, S \rangle$ along with T. Define  $\Gamma = \langle S \rangle = \langle C \sqcup \{T\} \rangle = \langle H, T \rangle$ ; for this reason we often think of S as just

E-mail address: zstier@berkeley.edu.

Date: December 2020.

<sup>\*</sup>Which is optimal due to the set of such gates having size  $\approx p^k$ , and the ball of radius  $\varepsilon$  in PU(2) having volume (in Haar measure) some constant multiple of  $\varepsilon^3$ .

 $\{H, T\}$ .  $\Gamma$  is dense in PU(2), and we are interested in tracking the *T*-count of  $\Gamma$ -approximants to elements of PU(2), defined as follows: for  $\gamma \in \Gamma$ , factor  $\gamma = \prod_{i=1}^{n} s_i$  where  $s_i \in S$ ; the *T*-count is the minimum number of  $s_i$  equalling *T* among all factorizations  $\{s_i\}$  of  $\gamma$ . The *T*-count is of interest because the Clifford+*T* gates are commonly used for modern quantum circuitry, and elements of *C* are assumed to be easy to fabricate, while *T* is much more costly, hence why we ignore the number of *C*-elements required.

Each element of  $\Gamma$  has entries in the standard quaternion realization [Sar15] in the ring of integers  $\mathcal{O} = \mathbb{Z}[\sqrt{2}]$ , a unique factorization domain, with fraction field  $K = \mathbb{Q}[\sqrt{2}]$ . We assume that there exists a  $O(\text{poly} \log N_{K/\mathbb{Q}}(n))$ -time algorithm to factor  $n \in \mathcal{O}$  into primes.

**Theorem** ([RS16]). There is a  $O(\text{poly} \log \frac{1}{\varepsilon})$ -time algorithm to factor almost any element of PU(2) using the gates S to within  $\varepsilon$  in d using T-count at most  $(3 + o(1)) \log_2 \frac{1}{\varepsilon^3}$ .

(This result, accomplished with Euler angles, generalizes to when S is any golden gate set [PS18, §2.3].) We shall leverage the two-dimensional lattice structure of  $\mathcal{O}$  to apply Lenstra's algorithm [Len83, Paz83]. The main result is the following:

**Theorem 1.** There is a  $O\left(\operatorname{poly}\log\frac{1}{\varepsilon}\right)$ -time algorithm to factor almost any element of PU(2) using the gates S to within  $\varepsilon$  in d using T-count at most  $\left(\frac{7}{3} + o(1)\right)\log_2\frac{1}{\varepsilon^3}$ .

The heart of this improvement lies with replacing the middle diagonal factor of a typical element of PU(2) with a well-chosen element of  $\Gamma$ , selected to have particularly small *T*-count.

In §2, we present some technical lemmas that enable this factorization. In §3, we describe the algorithm which accomplishes Theorem 1, analyzing it is §4. In §5, we give some information about the Haskell implementation.

# 2. Nearby elements in PU(2)

The contents of this section are entirely independent of the choice of topological generators. Instead, the goal is Lemma 4 by which we establish sufficient conditions for two elements of PU(2) to be nearby with respect to the following metric.

**Definition** (PU(2)'s bi-invariant metric, cf. [Sar15, PS18]). PU(2) has the bi-invariant metric

$$d(x,y) = 1 - \frac{1}{2} |\operatorname{tr} x^* y|.$$

Note that this is also the bi-invariant metric on SU(2), and that  $PU(2) \cong PSU(2)$ , so quotienting by (norm-one) scalars does not change this metric due to the presence of the absolute value. Therefore, for concreteness we shall work work elements of SU(2).

We introduce the following convenient notation for elements of the group.

**Notation.** Suppose  $\alpha, \beta \in \mathbb{C}$  satisfy  $|\alpha|^2 + |\beta|^2 = 1$ . Then we let  $u(\alpha, \beta)$  denote the canonical element corresponding to  $\alpha$  and  $\beta$ , i.e.

$$u(\alpha,\beta) = \begin{pmatrix} \alpha & \beta \\ -\overline{\beta} & \overline{\alpha} \end{pmatrix}$$

We use the additional shorthand of  $u(\theta)$  to denote the canonical diagonal element corresponding to rotation by  $\theta \in \mathbb{R}$ , i.e.

$$u(\theta) = u(e^{i\theta}, 0) = \begin{pmatrix} e^{i\theta} & 0\\ 0 & e^{-i\theta} \end{pmatrix}.$$

Now, we are equipped to begin discussing the elements of SU(2), beginning with a fact about real numbers, representing the norms of the complex numbers comprising the matrix. There is a catch, namely that these results only hold for matrices outside of a small neighborhood of the identity matrix parameterized by  $\varepsilon_0$ ; however,  $\varepsilon_0$  can be made as small as one likes, with the only cost being (approximately inversely) to multiples of  $\varepsilon$  obtained in subsequent bounds. Therefore it is ideal to have  $\varepsilon \ll \varepsilon_0$ .

**Lemma 2.** Select absolute constants  $\tilde{\varepsilon}, \varepsilon_0 > 0$ . Suppose  $a_1, a_2, b_1, b_2 \ge 0$  satisfy the following conditions, for some  $\varepsilon < \tilde{\varepsilon}$ :

$$a_1^2 + b_1^2 = 1,$$
  $a_2^2 + b_2^2 = 1,$   $|a_1 - a_2| < \varepsilon,$ 

and either  $a_1 < \sqrt{1 - \varepsilon_0^2}$  or  $a_2 < \sqrt{1 - \varepsilon_0^2}$ . Then there exists  $c = c(\tilde{\varepsilon}, \varepsilon_0)$  for which  $|b_1 - b_2| < c\varepsilon$ .

A pictorial view of this result is as follows. Consider points in the upper half-plane and on the unit circle, at least some fixed distance above the x-axis. Then, if they have nearby x-coordinates, it follows that they have comparably nearby y-coordinates.

*Proof.* For convenience, write  $a_2 = a_1 + \delta$  for  $|\delta| < \varepsilon$ . Then

$$b_2^2 = 1 - a_2^2$$
  
=  $1 - a_1^2 - 2\delta a_1 - \delta^2$   
=  $b_1^2 - \delta(2a_1 + \delta)$   
 $|b_1 - b_2| = \frac{|\delta| |2a_1 + \delta|}{b_1 + b_2}$   
 $< \frac{2 + \tilde{\varepsilon}}{\varepsilon_0} \varepsilon$ 

where  $c(\tilde{\varepsilon}, \varepsilon_0) = \frac{2+\tilde{\varepsilon}}{\varepsilon_0}$ .

We leverage this result and apply it to the absolute values of entries of elements of SU(2), to show the following about such matrices having nearby entrywise absolute values and arguments.

**Lemma 3.** Select absolute constants  $\tilde{\varepsilon}, \varepsilon_0 > 0$ . Suppose  $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{C}$  satisfy the following conditions, for some  $\varepsilon < \tilde{\varepsilon}$ :

$$\begin{split} \arg \alpha_1 &= \arg \alpha_2, \\ \arg \beta_1 &= \arg \beta_2, \\ & ||\alpha_1|^2 + |\beta_1|^2 = 1, \\ & |\alpha_2|^2 + |\beta_2|^2 = 1, \\ & ||\alpha_1| - |\alpha_2|| < \varepsilon, \end{split}$$

and either  $|\alpha_1| < \sqrt{1 - \varepsilon_0^2}$  or  $|\alpha_2| < \sqrt{1 - \varepsilon_0^2}$ . Write  $\gamma_1 = u(\alpha_1, \beta_1)$  and  $\gamma_2 = u(\alpha_2, \beta_2)$ . Then there exists  $\tilde{c} = \tilde{c}(\tilde{\varepsilon}, \varepsilon_0)$  for which  $d(\gamma_1, \gamma_2) < \tilde{c}\varepsilon$ .

*Proof.* We first fix  $c = \frac{2+\tilde{\varepsilon}}{\varepsilon_0}$  and immediately we apply Lemma 2 to find  $||\beta_1| - |\beta_2|| < c\varepsilon$ . We have that

$$\gamma_1^* \gamma_2 = \begin{pmatrix} \overline{\alpha_1} \alpha_2 + \beta_1 \overline{\beta_2} & \star \\ \star & \alpha_1 \overline{\alpha_2} + \overline{\beta_1} \beta_2 \end{pmatrix}$$

where the  $\star$ 's indicate irrelevant quantities. Then,

$$\operatorname{tr} \gamma_1^* \gamma_2 = \overline{\alpha_1} \alpha_2 + \alpha_1 \overline{\alpha_2} + \beta_1 \beta_2 + \beta_1 \beta_2$$
  
= 2 |\alpha\_1| |\alpha\_2| + 2 |\beta\_1| |\beta\_2|  
> 2 |\alpha\_1| (|\alpha\_1| - \varepsilon) + 2 |\beta\_1| (|\beta\_1| - c\varepsilon)  
= 2 - 2 (|\alpha\_1| + c |\beta\_1|) \varepsilon.

Elementarily,\*  $|\alpha_1| + c |\beta_1| \leq \sqrt{1 + c^2}$  for all choices of  $\alpha_1, \beta_1$  satisfying  $|\alpha_1|^2 + |\beta_1|^2 = 1$ . Therefore, tr $\gamma_1^* \gamma_2 > 2 - 2\sqrt{1 + c^2}\varepsilon$ . We conclude with

$$d(\gamma_1, \gamma_2) = 1 - \frac{1}{2} |\operatorname{tr} \gamma_1^* \gamma_2| < \sqrt{1 + c^2} \varepsilon$$

where  $\tilde{c}(\tilde{\varepsilon}, \varepsilon_0) = \sqrt{1 + c^2}$ .

Finally, we use this guarantee about the bi-invariant metric to explicitly describe rotations which send one matrix to have close entries in absolute value to another while also ensuring closeness in the metric.

**Lemma 4.** Select absolute constants  $\tilde{\varepsilon}, \varepsilon_0 > 0$ . Take  $\gamma_1, \gamma_2 \in SU(2)$  and write them as  $\gamma_1 = u(\alpha_1, \beta_1)$  and  $\gamma_2 = u(\alpha_2, \beta_2)$ . If  $||\alpha_1| - |\alpha_2|| < \varepsilon$  for some  $\varepsilon < \tilde{\varepsilon}$  and either  $|\alpha_1| < \sqrt{1 - \varepsilon_0^2}$  or  $|\alpha_2| < \sqrt{1 - \varepsilon_0^2}$  then there exist  $\theta_1, \theta_2 \in \mathbb{R}$  and  $\tilde{c} = \tilde{c}(\tilde{\varepsilon}, \varepsilon_0)^{\dagger}$  for which, writing  $\delta_1 = u(\theta_1)$  and  $\delta_2 = u(\theta_2)$ , we have

$$d(\gamma_1, \delta_1 \gamma_2 \delta_2) < \tilde{c}\varepsilon.$$

Proof. First, let

$$\theta_1 + \theta_2 = \arg \alpha_1 - \arg \alpha_2$$
  
$$\theta_1 - \theta_2 = \arg \beta_1 - \arg \beta_2$$

which reduces to

$$\theta_1 = \frac{1}{2} \left( \arg \alpha_1 - \arg \alpha_2 + \arg \beta_1 - \arg \beta_2 \right)$$
  
$$\theta_2 = \frac{1}{2} \left( \arg \alpha_1 - \arg \alpha_2 - \arg \beta_1 + \arg \beta_2 \right).$$

We then multiply out  $\delta_1 \gamma_2 \delta_2$  as<sup>‡</sup>

$$\delta_1 \gamma_2 \delta_2 = \begin{pmatrix} e^{i(\theta_1 + \theta_2)} \alpha_2 & e^{i(\theta_1 - \theta_2)} \beta_2 \\ -e^{i(\theta_1 - \theta_2)} \beta_2 & e^{i(\theta_1 + \theta_2)} \alpha_2 \end{pmatrix} = \begin{pmatrix} \alpha_3 & \beta_3 \\ -\overline{\beta_3} & \overline{\alpha_3} \end{pmatrix} \in \mathrm{PU}(2)$$

where  $|\alpha_3| = |\alpha_2|$ ,  $|\beta_3| = |\beta_2|$ ,  $\arg \alpha_3 = \arg \alpha_1$ , and  $\arg \beta_3 = \arg \beta_1$ . Therefore we apply Lemma 3 and immediately conclude the result.

<sup>\*</sup>e.g. using Lagrange multipliers.

 $<sup>^{\</sup>dagger}\tilde{c}$  as in Lemma 3.

<sup>&</sup>lt;sup>‡</sup>The bars represent complex conjugates, not fractions.

## 3. Algorithm for short paths

Select absolute constants  $\tilde{\varepsilon}, \varepsilon_0 > 0$  where  $\tilde{\varepsilon} < \frac{1}{2}$ . Take any  $g = u(\alpha, \beta) \in PU(2)$  where  $|\alpha| < \sqrt{1 - \varepsilon_0^2}$ , and pick  $\varepsilon < \tilde{\varepsilon}$ .\* We wish to approximate g using  $\gamma \in \Gamma$  of the form

(5) 
$$\gamma = \frac{1}{2^{\frac{1}{2}k}} \begin{pmatrix} x_0 + x_1 i & x_2 + x_3 i \\ -x_2 + x_3 i & x_0 - x_1 i \end{pmatrix}$$

having k, the factorization length, minimized, and so we begin with k = 0. (We also have  $x_0, x_1, x_2, x_3 \in \mathcal{O}$ .) In particular, the objective is to approximate g as  $\gamma_1 \gamma \gamma_2$  where  $\gamma_1, \gamma_2 \in \Gamma$  approximate well-chosen diagonals, and  $\gamma \in \Gamma$  has factorization computable by Kliuchnikov–Maslov–Mosca [KMM13]. (We will compute  $\gamma_1$  and  $\gamma_2$  using [RS16].) We will see that  $\gamma$  is designed to have factorization typically shorter than that of  $\gamma_1$  and  $\gamma_2$ , giving rise to the desired improvement.

factorization typically shorter than that of  $\gamma_1$  and  $\gamma_2$ , giving rise to the desired improvement. In order to apply Lemma 4 we need to have  $\left|\frac{x_0+x_1i}{2^{\frac{1}{2}k}}\right| = \sqrt{\frac{x_0^2+x_1^2}{2^k}}$  near  $|\alpha|$  (that is, within  $\varepsilon$ ). Because  $\left|\frac{x_0+x_1i}{2^{\frac{1}{2}k}}\right| + |\alpha| \ge |\alpha|$  which is fixed, it suffices to first find candidate values for  $x_0, x_1 \in \mathcal{O}$  with  $\left|\left|\frac{x_0+x_1i}{2^{\frac{1}{2}k}}\right|^2 - |\alpha|^2\right| < \varepsilon |\alpha|$ , rewritten to

(6) 
$$\left|x_0^2 + x_1^2 - \left|\alpha\right|^2 2^k\right| < \varepsilon \left|\alpha\right| 2^k.$$

Viewing  $\gamma$  as an element of SU(2), we also have det  $\gamma = 1$ , i.e.  $x_0^2 + x_1^2 + x_2^2 + x_3^2 = 2^k$ .  $[K : \mathbb{Q}] = 2$ , so we explicitly work with the Galois group elements

$$\sigma_{+}: 1 \longmapsto 1$$

$$\sqrt{2} \longmapsto \sqrt{2},$$

$$\sigma_{-}: 1 \longmapsto 1$$

$$\sqrt{2} \longmapsto -\sqrt{2},$$

both of which are real embeddings, so that as  $x_i \in \mathcal{O} \subset K \subset \mathbb{R}$ , it follows that  $\sigma_{\pm}(x_0^2 + x_1^2) + \sigma_{\pm}(x_2^2 + x_3^2) = \sigma_{\pm}2^k = 2^k$ , and so

(7) 
$$\sigma_{\pm}(x_0^2 + x_1^2) \leqslant 2^k.$$

Now, let  $m = x_0^2 + x_1^2 \in \mathcal{O}$ . Considering  $\mathcal{O}$  as an integer lattice, we adapt (6) and (7) and seek to solve

$$\left| m - |\alpha|^2 2^k \right| < \varepsilon |\alpha| 2^k$$
$$|\sigma_{\pm} m| \leqslant 2^k$$

which are convex constraints on m when written in its lattice components. Since this is an integer programming problem in two dimensions, we apply Lenstra's algorithm to efficiently list all such lattice points m. For each m, using efficient factorization in  $\mathcal{O}$ , we attempt to write m as a sum of two squares; if possible, say  $m = x_0^2 + x_1^2$ , and so we attempt to write  $\tilde{m} = 2^k - m$  as a sum of two squares. If possible, say  $\tilde{m} = x_2^2 + x_3^2$ , so we simply halt and return  $\gamma$  corresponding to (5). However, if  $\tilde{m}$  may not be represented as a sum of two squares, we simply move on to the next value of m and try this process again. If this fails for all m arising from k, we increment k and run Lenstra's algorithm for the new inequalities.

<sup>\*</sup>By choice of sufficiently small  $\varepsilon_0$ , we exclude at this stage an arbitrarily small ball in Haar measure about the identity matrix.

Supposing we have halted and constructed  $\gamma$ , we compute  $\delta_1$  and  $\delta_2$  guaranteed by Lemma 4. These are efficiently approximable by [RS16, Algorithm 7.3] to  $\gamma_1$  and  $\gamma_2$ , respectively. Chaining together the three approximations  $\gamma_1 \gamma \gamma_2$  gives the final desired approximation to g.

**Remark.** We assume to begin that g is far from a diagonal, i.e. that  $|\alpha| < \sqrt{1 - \varepsilon_0^2}$ . However, we also note that if  $\gamma = u(\arg \alpha)$ , then  $d(\gamma, g) = 1 - |\alpha| = \frac{\varepsilon_0^2}{2} + O(\varepsilon_0^4)$  (by considering the Taylor series of  $\sqrt{1 - x^2}$ ). Therefore, if we have the additional assumption that  $\varepsilon_0^2 \approx \varepsilon$  (which is sensible, since if  $\varepsilon$  is "small" then  $\sqrt{\varepsilon}$  will be "large", relatively speaking, and this is not incompatible with  $\varepsilon \ll \varepsilon_0$ ), we get a bifurcated approach for any  $g = u(\alpha, \beta) \in PU(2)!$  Namely, in the case that  $|\alpha| < \sqrt{1 - \varepsilon_0^2}$  we run our algorithm as specified above, and otherwise we give  $u(\arg \alpha)$  to [RS16]'s algorithm.

## 4. Analysis of the algorithm

We begin the analysis by establishing the *T*-count and tightness of the approximation. In particular,  $d(\gamma_1, \delta_1) < \varepsilon$  and  $d(\gamma_2, \delta_2) < \varepsilon$  with factorization lengths each  $(1 + o(1)) \log_2 \frac{1}{\varepsilon^3}$ . By Lemma 4,  $d(g, \delta_1 \gamma \delta_2) < \tilde{c}\varepsilon$ . Therefore,  $d(g, \gamma_1 \gamma \gamma_2) < (\tilde{c} + 2)\varepsilon$  (by the triangle inequality) and since  $\gamma$  has a factorization of *T*-count approximately  $\frac{1}{3} \log_2 \frac{1}{\varepsilon^3}$ , this constitutes a factorization of an element in *g*'s neighborhood of *T*-count  $\frac{7}{3} \log_2 \frac{1}{\varepsilon^3}$ .

The efficiency of this algorithm—that is, that it runs in time  $O(\text{poly} \log \frac{1}{\varepsilon})$ —is because we expect to halt when  $2^k \varepsilon^3 \in O(1)$  (so only  $k \approx \frac{1}{3} \log_2 \frac{1}{\varepsilon^3}$  calls are expected), and only call polynomiallymany polynomial-time subroutines. The dominant subroutines are calls to Lenstra's algorithm, which as shown in [Len83] which runs in time polynomial in the size of the constraints for any fixed dimension n. Indeed, here we have only m = 6 linear constraints (two per absolute value), and the largest value a in the constraints is  $p^k$ , so the runtime is polynomial in  $nm \log a \in \Theta(k)$ .

The reason we expect to halt when  $2^k \varepsilon^3 \in O(1)$  is that  $\Gamma$  is viewed as an infinite 3-regular tree with each edge taking one further from the identity matrix's vertex represents an increase in the weight. The number of vertices of *T*-count exactly k is  $3 \cdot 2^{k-1}$ , so there are  $\Theta(2^k)$  elements of *T*-count up to k. Expecting them to cover PU(2) (of total volume 1) well for large k, with a ball of volume some constant multiple c of  $\varepsilon^3$ , we therefore have  $c2^k\varepsilon^3 \approx 1$ , i.e.  $k \approx \frac{1}{3}\log_2 \frac{1}{\varepsilon^3}$ . As *T*-count is increasing in the quantity k as in (5), we expect to only have about  $\frac{1}{3}\log_2 \frac{1}{\varepsilon^3}$ -many iterations to achieve *T*-count  $\frac{1}{3}\log_2 \frac{1}{\varepsilon^3}$ .

When attempting to write elements of  $\mathcal{O}$  as a sum of two squares, we primarily rest on a belief, in the style of Cramér's conjecture and a conjecture of Sardari [Sar19, (\*)], that sums of squares are dense in  $\mathbb{N}$ . Seeking to analogize [Sar19, (\*)] in particular, we note that the operative aspect is that a dense cluster of lattice points will represent a sum of two squares, and that a point accomplishing this will be found quickly through Lenstra's algorithm.

The significance of this result is to accomplish a factorization in PU(2) shorter than the  $3 \log_2 \frac{1}{\varepsilon^3}$ -factorization first demonstrated in [PS18], precisely the desired outcome.

## 5. Implementation and an example

Visit https://math.berkeley.edu/~zstier/ugthesis/HTLenstra.hs for a Haskell implementation of code for the algorithm described in §3, using many functions implemented in [RS18], and visit https://math.berkeley.edu/~zstier/ugthesis/HTLenstra.nb for a partial Mathematica implementation. Let us now attempt to approximate

$$g = \frac{1}{3} \begin{pmatrix} 1 & 2+2i \\ -2+2i & 1 \end{pmatrix}$$

in d to within  $\varepsilon = 10^{-10}$  (times a small positive constant) using the Haskell implementation. Our heuristics provide an expectation of T-count near  $\frac{7}{3}\log_2 10^{30} \approx 232.5$ .

The algorithm described in §3 halts at k = 17 and returns the approximation

$$\gamma = \frac{1}{2^9} \begin{pmatrix} -121 + 145\sqrt{2} + (123 - 192\sqrt{2})i & 103 + 78\sqrt{2} - (211 + 157\sqrt{2})i \\ -103 - 78\sqrt{2} - (211 + 157\sqrt{2})i & -121 + 145\sqrt{2} - (123 - 192\sqrt{2})i \end{pmatrix}.$$

This factors as

# 

with T-count 32—actually shorter than the expectation of  $\frac{1}{3}\log_2 10^{30} \approx 33.2$ . We readily compute

$$\theta_1 \approx 1.477137$$
  
 $\theta_2 \approx -0.421352$ 

and approximate the corresponding matrices  $\delta_1 = u(\theta_1)$  and  $\delta_2 = u(\theta_2)$  as

both of *T*-counts 102, very near to  $\log_2 10^{30} \approx 99.6$ . In total, this factorization gives *T*-count 236 (fewer than four more than the predicted value of  $\frac{7}{3}\log_2 10^{30}$ ), and explicitly multiplying out the matrices gives precision with respect to *d* within  $\frac{4}{10^{21}}$ , well within the guarantees of Lemma 4 with  $\tilde{\varepsilon} = \frac{1}{10^9}$  and  $\varepsilon_0 = \frac{5}{6}$  (any  $\tilde{\varepsilon} > \varepsilon$  and  $\varepsilon_0 < \frac{2\sqrt{2}}{3}$  suffice), which predicts an approximation to within  $\left(2 + \frac{2+10^{-9}}{\frac{5}{6}}\right) \frac{1}{10^{10}} \approx 4.4 \cdot \frac{1}{10^{10}}$ .

### 6. Other gate sets

Consider the golden gate set of "V-gates,"\* with their notation preserved from [Sar15]:

$$s_1 = \frac{1}{\sqrt{5}} \begin{pmatrix} 1+2i \\ 1-2i \end{pmatrix}, \qquad s_2 = \frac{1}{\sqrt{5}} \begin{pmatrix} 1 & 2i \\ 2i & 1 \end{pmatrix}, \qquad s_3 = \frac{1}{\sqrt{5}} \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix},$$

with  $S = \{s_1^{\pm 1}, s_2^{\pm 1}, s_3^{\pm 1}\}$  and  $\Gamma = \langle s_1, s_2, s_3 \rangle$ . This case is actually even simpler than that of the Clifford+*T* gates, requires just a 1-dimensional variant of Lenstra's algorithm, and is described in full, and implemented, in [Sti20, §2.2].

The case of general golden gate sets will be the subject of a future paper.

### Acknowledgements

Part of this work appears in my senior thesis at Princeton University. I am grateful to my advisor Peter Sarnak for his support and guidance throughout. I am supported by a National Science Foundation Graduate Research Fellowship, grant number DGE-1752814.

#### References

- [CP18] E. Carvalho Pinto and C. Petit. "Better path-finding algorithms in LPS Ramanujan graphs." In: Journal of Mathematical Cryptology 12(4) (2018).
- [KLM<sup>+</sup>20] V. Kliuchnikov, K. Lauter, R. Minko, A. Paetznick, and C. Petit. Private communication.
- [KMM13] V. Kliuchnikov, D. Maslov, and M. Mosca. "Fast and efficient exact synthesis of single qubit unitaries generated by Clifford and T gates." In: *Quantum Information & Computation* (2013).
- [Len83] H. W. Lenstra, Jr. "Integer Programming with a Fixed Number of Variables." In: Mathematics of Operations Research 8(4) (1983).
- [Paz83] A. Paz. "A simplified version of H.W. Lenstra's integer programming algorithm and some applications" (1983).
- [PS18] O. Parzanchevski and P. Sarnak. "Super-Golden-Gates for PU(2)." In: Advances in Mathematics 327 (2018), pp. 869–901.
- [RS16] N. Ross and P. Selinger. "Optimal ancilla-free Clifford+T approximation of z-rotations." In: Quantum Information & Computation (2016).
- [RS18] N. Ross and P. Selinger. "newsynth: Exact and approximate synthesis of quantum circuits." In: Hackage: The Haskell Package Repository (2018).
- [Sar15] P. Sarnak. "Letter to Scott Aaronson and Andy Pollington on the Solovay–Kitaev Theorem and Golden Gates." (2015).
- [Sar19] N. Sardari. "Complexity of Strong Approximation on the Sphere." In: International Mathematics Research Notices (2019).
- [Sti20] Z. Stier. "Optimal topological generators of U(1) and short paths in  $X^{p,q}$  and SU(2)." Princeton University senior thesis (2020).

<sup>\*</sup>So named due to the connection with the Roman numeral V.