On the compactness threshold in the critical Kirchhoff equation^{*}

Erisa Hasani and Kanishka Perera

Department of Mathematical Sciences Florida Institute of Technology Melbourne, FL 32901, USA ehasani2016@my.fit.edu & kperera@fit.edu

Abstract

We study a class of critical Kirchhoff problems with a general nonlocal term. The main difficulty here is the absence of a closed-form formula for the compactness threshold. First we obtain a variational characterization of this threshold level. Then we prove a series of existence and multiplicity results based on this variational characterization.

1 Introduction

The purpose of this paper is to study the critical Kirchhoff problem

$$\begin{cases} -h\left(\int_{\Omega} |\nabla u|^2 \, dx\right) \Delta u = f(x, u) + |u|^{2^* - 2} \, u \quad \text{in } \Omega\\ u = 0 \qquad \qquad \text{on } \partial\Omega, \end{cases}$$
(1.1)

where Ω is a bounded domain in \mathbb{R}^N , $N \geq 3$, $h: [0, \infty) \to [0, \infty)$ is a continuous and nondecreasing function, $2^* = 2N/(N-2)$ is the critical Sobolev exponent, and f is a Carathéodory function on $\Omega \times \mathbb{R}$ satisfying the subcritical growth condition

 $|f(x,t)| \le c_1 |t|^{p-1} + c_2 \quad \text{for a.a. } x \in \Omega \text{ and all } t \in \mathbb{R}$ (1.2)

for some constants $c_1, c_2 > 0$ and 1 . The class of nonlocal terms considered here includes sums of powers

$$h(t) = \sum_{i=1}^{n} a_i t^{\gamma_i - 1}, \quad t \ge 0,$$

^{*}*MSC2010:* Primary 35J60, Secondary 35B33, 35J20

Key Words and Phrases: critical Kirchhoff equation, general nonlocal term, compactness threshold, existence, multiplicity

where $a_1, \ldots, a_n > 0$ and $1 \le \gamma_1 < \cdots < \gamma_n < +\infty$. A model case is

$$h(t) = a + bt^{\gamma - 1},$$

where $a, b \ge 0$ with a + b > 0 and $1 < \gamma < +\infty$. The classical case h(t) = a + bt corresponds to $\gamma = 2$.

As is usually the case with problems of critical growth, problem (1.1) lacks compactness. The standard approach to such problems is to determine a threshold level below which there is compactness and construct minimax critical levels below this threshold. This approach has been used in the classical case

$$h(t) = a + bt, \quad a, b \ge 0, \quad a + b > 0$$

in dimensions N = 3 and 4 to obtain nontrivial solutions in the recent literature (see, e.g., Huang et al. [6], Liao et al. [8], Naimen [9, 10], Xie et al. [13], Yao and Mu [14], Zhang and Liu [15], and the references therein). However, in the general case considered in the present paper such a threshold level cannot be found in closed form. Our first contribution here is a variational characterization of this threshold level (see Theorem 2.3). Then we give a series of existence and multiplicity results based on this variational characterization (see Section 3). This requires novel arguments due to the absence of a closed-form compactness threshold.

We will state and prove our compactness, existence, and multiplicity results for problem (1.1) with a general nonlocal term h in the next two sections. To illustrate our results while keeping the presentation simple, we state them here for the model problem

$$\begin{cases} -\left[a+b\left(\int_{\Omega}|\nabla u|^{2} dx\right)^{\gamma-1}\right]\Delta u = \lambda u + |u|^{2^{*}-2} u \text{ in } \Omega \\ u = 0 \text{ on } \partial\Omega, \end{cases}$$
(1.3)

where $1 < \gamma < +\infty$, $a \ge 0$, b > 0, and $\lambda > 0$.

Weak solutions of this problem coincide with critical points of the functional

$$J(u) = \frac{a}{2} \int_{\Omega} |\nabla u|^2 dx + \frac{b}{2\gamma} \left(\int_{\Omega} |\nabla u|^2 dx \right)^{\gamma} - \frac{\lambda}{2} \int_{\Omega} u^2 dx - \frac{1}{2^*} \int_{\Omega} |u|^{2^*} dx, \quad u \in H^1_0(\Omega).$$

Recall that J satisfies the Palais-Smale compactness condition at the level $c \in \mathbb{R}$, or the (PS)_c condition for short, if every sequence (u_j) in $H_0^1(\Omega)$ such that $J(u_j) \to c$ and $J'(u_j) \to 0$ has a strongly convergent subsequence. Let S be the best Sobolev constant (see (2.1)) and let $\lambda_1 > 0$ be the first Dirichlet eigenvalue of the Laplacian in Ω . We have the following compactness results for the cases $\gamma < 2^*/2$, $\gamma = 2^*/2$, and $\gamma > 2^*/2$ (see Corollary 2.5, Corollary 2.9, and Corollary 2.11).

Theorem 1.1. Let $1 < \gamma < 2^*/2$, a, b > 0, and $0 < \lambda \le a\lambda_1$. Let t_0 be the unique positive solution of the equation $a + bt^{\gamma-1} = S^{-2^*/2} t^{2^*/2-1}$ and set

$$c^* = \frac{1}{N}at_0 + \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right)bt_0^{\gamma}.$$

Then J satisfies the $(PS)_c$ condition for all $c < c^*$.

Theorem 1.2. Let $\gamma = 2^*/2$.

(i) Let
$$a > 0, 0 < b < S^{-2^*/2}$$
, and $0 < \lambda < a\lambda_1$. Set
$$c^* = \frac{1}{N} \left(\frac{a^{2^*/2}}{S^{-2^*/2} - b}\right)^{2/(2^* - 2)}.$$

Then J satisfies the $(PS)_c$ condition for all $c < c^*$.

(ii) If $a \ge 0$ and $b > S^{-2^*/2}$, then J satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$ for any $\lambda > 0$.

Theorem 1.3. If $\gamma > 2^*/2$ and

$$a^{\gamma-2^*/2} b^{2^*/2-1} > \frac{(\gamma-2^*/2)^{\gamma-2^*/2} (2^*/2-1)^{2^*/2-1}}{(\gamma-1)^{\gamma-1}} S^{-(2^*/2)(\gamma-1)},$$

then J satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$ for any $\lambda > 0$.

Theorems 1.1–1.3 have the following corollary for the classical case $\gamma = 2$, where $c^* = +\infty$ means that J satisfies the (PS)_c condition for all $c \in \mathbb{R}$.

Corollary 1.4. Let $\gamma = 2$.

(i) If
$$N = 3$$
, $a, b > 0$, and $0 < \lambda \le a\lambda_1$, then $c^* = \frac{1}{4}abS^3 + \frac{1}{24}b^3S^6 + \frac{1}{24}(4aS + b^2S^4)^{3/2}$.

(ii) If
$$N = 4$$
, $a > 0$, $0 < b < S^{-2}$, and $0 < \lambda < a\lambda_1$, then $c^* = \frac{a^2}{4(S^{-2} - b)}$.

(iii) If N = 4, $a \ge 0$, and $b > S^{-2}$, then $c^* = +\infty$ for any $\lambda > 0$.

(iv) If
$$N \ge 5$$
 and $a^{N-4}b^2 > \frac{4(N-4)^{N-4}}{(N-2)^{N-2}}S^{-N}$, then $c^* = +\infty$ for any $\lambda > 0$.

Remark 1.5. The threshold levels in Corollary 1.4 (i)-(iii) were also obtained using different arguments in Naimen [10], Naimen [9], and Liao et al. [8], respectively.

We have the following existence and multiplicity results for problem (1.3) (see Corollary 3.6, Theorem 3.10, and Theorem 3.11).

Theorem 1.6. If $1 < \gamma < 2^*/2$, a, b > 0, and $N \ge 4$, then problem (1.3) has a nontrivial solution for $0 < \lambda < a\lambda_1$.

Theorem 1.7. Let $\gamma = 2^*/2$.

- (i) If a > 0, $0 < b < S^{-2^*/2}$, and $N \ge 4$, then problem (1.3) has a nontrivial solution for $0 < \lambda < a\lambda_1$.
- (ii) If a = 0 and $b > S^{-2^*/2}$, then problem (1.3) has a nontrivial solution for all $\lambda > 0$.
- (iii) If a > 0 and $b > S^{-2^*/2}$, then problem (1.3) has two nontrivial solutions for $\lambda > a\lambda_1$.

Theorem 1.8. If $\gamma > 2^*/2$ and

$$a^{\gamma-2^*/2} b^{2^*/2-1} > \frac{(\gamma-2^*/2)^{\gamma-2^*/2} (2^*/2-1)^{2^*/2-1}}{(\gamma-1)^{\gamma-1}} S^{-(2^*/2)(\gamma-1)},$$

then problem (1.3) has two nontrivial solutions for $\lambda \geq a\lambda_1$.

Theorems 1.6–1.8 have the following corollaries for the classical case $\gamma = 2$.

Corollary 1.9. Let $\gamma = 2$ and N = 4.

- (i) If a > 0 and $0 < b < S^{-2}$, then problem (1.3) has a nontrivial solution for $0 < \lambda < a\lambda_1$.
- (ii) If a = 0 and $b > S^{-2}$, then problem (1.3) has a nontrivial solution for all $\lambda > 0$.
- (iii) If a > 0 and $b > S^{-2}$, then problem (1.3) has two nontrivial solutions for $\lambda > a\lambda_1$.

Corollary 1.10. If $\gamma = 2$, $N \ge 5$, and

$$a^{N-4}b^2 > \frac{4(N-4)^{N-4}}{(N-2)^{N-2}}S^{-N},$$

then problem (1.3) has two nontrivial solutions for $\lambda \geq a\lambda_1$.

Remark 1.11. The result in Corollary 1.9 (i) was also obtained in Naimen [9] using a different method. Liao et al. [8] obtained one nontrivial solution when $a \ge 0$, $b > S^{-2}$, and $\lambda > a\lambda_1$. See also Perera and Zhang [12] for a related result in the subcritical case in dimensions $N \le 3$.

Remark 1.12. Corollary 1.10 complements the results in Naimen and Shibata [11], where two positive solutions were obtained when a = 1, b > 0 is sufficiently small, and $0 < \lambda < \lambda_1$.

In the borderline case where $\gamma = 2^*/2$ and $b = S^{-2^*/2}$, lower-order terms come into play. Consider the problem

$$\begin{cases} -\left[a + S^{-2^{*}/2} \left(\int_{\Omega} |\nabla u|^{2} dx\right)^{2^{*}/2-1} + \eta \left(\int_{\Omega} |\nabla u|^{2} dx\right)^{\sigma-1}\right] = \lambda u + |u|^{2^{*}-2} u \quad \text{in } \Omega \\ u = 0 \qquad \qquad \text{on } \partial\Omega, \end{cases}$$
(1.4)

where $a \ge 0$, $\eta > 0$, $1 < \sigma < 2^*/2$, and $\lambda > 0$. We have the following existence and multiplicity result (see Theorem 3.13).

Theorem 1.13. Let $\eta > 0$ and $1 < \sigma < 2^*/2$.

- (i) If a = 0, then problem (1.4) has a nontrivial solution for all $\lambda > 0$.
- (ii) If a > 0, then problem (1.4) has two nontrivial solutions for $\lambda > a\lambda_1$.

2 Compactness threshold

A weak solution of problem (1.1) is a function u that belongs to the Sobolev space $H_0^1(\Omega)$ and satisfies

$$h\left(\int_{\Omega} |\nabla u|^2 \, dx\right) \int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} f(x, u) \, v \, dx + \int_{\Omega} |u|^{2^* - 2} \, uv \, dx \quad \forall v \in H^1_0(\Omega).$$

Weak solutions coincide with critical points of the C^1 -functional

$$J(u) = \frac{1}{2} H\left(\int_{\Omega} |\nabla u|^2 \, dx\right) - \int_{\Omega} F(x, u) \, dx - \frac{1}{2^*} \int_{\Omega} |u|^{2^*} dx, \quad u \in H_0^1(\Omega),$$

where $F(x,t) = \int_0^t f(x,s) \, ds$ is the primitive of f.

Definition 2.1. The functional J satisfies the Palais-Smale compactness condition at the level $c \in \mathbb{R}$, or the (PS)_c condition for short, if every sequence (u_j) in $H_0^1(\Omega)$ such that

$$J(u_j) \to c \qquad J'(u_j) \to 0,$$

called a $(PS)_c$ sequence, has a strongly convergent subsequence.

Let

$$S = \inf_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\int_{\Omega} |\nabla u|^2 \, dx}{\left(\int_{\Omega} |u|^{2^*} dx\right)^{2/2^*}}$$
(2.1)

be the best Sobolev constant. The set

$$I = \left\{ t > 0 : h(t) \le S^{-2^*/2} t^{2^*/2 - 1} \right\}$$

will play an important role in our compactness results. We begin with a simple but useful proposition.

Proposition 2.2. If (u_i) is a sequence in $H_0^1(\Omega)$ such that

$$J'(u_j) \to 0, \qquad u_j \rightharpoonup u, \qquad ||u_j - u||^2 \to t,$$

then either t = 0 or $t \in I$. In particular, if $I = \emptyset$, then every bounded sequence (u_j) in $H_0^1(\Omega)$ such that $J'(u_j) \to 0$ has a strongly convergent subsequence.

Proof. Since $J'(u_j) \to 0$,

$$h\left(\int_{\Omega} |\nabla u_j|^2 \, dx\right) \int_{\Omega} \nabla u_j \cdot \nabla v \, dx - \int_{\Omega} f(x, u_j) \, v \, dx - \int_{\Omega} |u_j|^{2^* - 2} \, u_j \, v \, dx = \mathrm{o}(\|v\|) \quad (2.2)$$

for all $v \in H_0^1(\Omega)$, and since $u_j \rightharpoonup u$ and $||u_j - u||^2 \rightarrow t$,

$$\int_{\Omega} |\nabla u_j|^2 \, dx \to \int_{\Omega} |\nabla u|^2 \, dx + t =: s.$$

Passing to a renamed subsequence, we may assume that $u_j \to u$ strongly in $L^p(\Omega)$ and a.e. in Ω . So taking $v = u_j$ in (2.2) gives

$$h(s)\left(\int_{\Omega} |\nabla u|^2 \, dx + t\right) - \int_{\Omega} u f(x, u) \, dx - \int_{\Omega} |u_j|^{2^*} \, dx = o(1), \tag{2.3}$$

while taking v = u and passing to the limit gives

$$h(s) \int_{\Omega} |\nabla u|^2 \, dx - \int_{\Omega} u f(x, u) \, dx - \int_{\Omega} |u|^{2^*} \, dx = 0.$$
(2.4)

Since

$$\int_{\Omega} |u_j|^{2^*} dx - \int_{\Omega} |u|^{2^*} dx = \int_{\Omega} |u_j - u|^{2^*} dx + o(1)$$

by the Brézis-Lieb lemma (see [2]), subtracting (2.4) from (2.3) and using (2.1) gives

$$th(s) = \int_{\Omega} |u_j - u|^{2^*} dx + o(1) \le S^{-2^*/2} \left(\int_{\Omega} |\nabla(u_j - u)|^2 dx \right)^{2^*/2} + o(1).$$

If t > 0, then passing to the limit and noting that $h(s) \ge h(t)$ since $s \ge t$ and h is nondecreasing gives $h(t) \le S^{-2^*/2} t^{2^*/2-1}$, so $t \in I$.

First we consider the case where I is nonempty. Let

$$H(t) = \int_0^t h(s) \, ds, \quad t \ge 0$$

be the primitive of h, and set

$$K(t) = \frac{1}{2}H(t) - \frac{1}{2^*}th(t), \quad t \ge 0.$$

For $1 \leq \gamma \leq 2^*/2$, let

$$\lambda_1(\gamma) = \inf_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\left(\int_{\Omega} |\nabla u|^2 \, dx\right)^{\gamma}}{\int_{\Omega} |u|^{2\gamma} \, dx}$$
(2.5)

be the first eigenvalue of the nonlinear eigenvalue problem

$$\begin{cases} -\left(\int_{\Omega} |\nabla u|^2 dx\right)^{\gamma-1} \Delta u = \lambda \, |u|^{2\gamma-2} \, u & \text{in } \Omega\\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

which is positive by the Sobolev embedding theorem. We note that $\lambda_1(1) = \lambda_1$, the first Dirichlet eigenvalue of the Laplacian in Ω , and $\lambda_1(2^*/2) = S^{2^*/2}$. We assume that

(A₁) for some constants $\alpha_1, \ldots, \alpha_n > 0$, $1 \le \gamma_1 < \cdots < \gamma_n < 2^*/2$, and $\mu_1 \le \lambda_1(\gamma_1), \ldots, \mu_n \le \lambda_1(\gamma_n)$ with at least one of the inequalities strict,

$$K(t) \ge \sum_{i=1}^{n} \alpha_i t^{\gamma_i} \quad \forall t \ge 0$$

and

$$F(x,t) - \frac{1}{2^*} t f(x,t) \le \sum_{i=1}^n \mu_i \,\alpha_i \, |t|^{2\gamma_i} \quad \text{for a.a. } x \in \Omega \text{ and all } t \in \mathbb{R};$$

 (A_2) K is superadditive, i.e.,

$$K(t_1 + t_2) \ge K(t_1) + K(t_2) \quad \forall t_1, t_2 \ge 0;$$

 $(A_3) h(t)/t^{2^*/2-1}$ is strictly decreasing for t > 0 and

$$0 \le b := \lim_{t \to +\infty} \frac{h(t)}{t^{2^*/2 - 1}} < S^{-2^*/2} < \lim_{t \to 0} \frac{h(t)}{t^{2^*/2 - 1}} \le +\infty.$$

We note that K is nonnegative by (A_1) and hence nondecreasing by (A_2) . We have the following theorem.

Theorem 2.3. Assume that $I \neq \emptyset$ and (1.2), (A₁), and (A₂) hold. Set

$$c^* = \inf_{t \in I} K(t).$$
 (2.6)

Then J satisfies the $(PS)_c$ condition for all $c < c^*$. If, in addition, (A_3) holds, then the equation

$$h(t) = S^{-2^*/2} t^{2^*/2 - 1}$$
(2.7)

has a unique positive solution t_0 and

$$c^* = K(t_0), \tag{2.8}$$

in particular, $c^* > 0$.

Proof. First we note that for all $u \in H_0^1(\Omega)$,

$$K\left(\int_{\Omega} |\nabla u|^2 \, dx\right) - \int_{\Omega} \left[F(x,u) - \frac{1}{2^*} \, uf(x,u)\right] dx \ge \sum_{i=1}^n \alpha_i \left(1 - \frac{\mu_i}{\lambda_1(\gamma_i)}\right) \left(\int_{\Omega} |\nabla u|^2 \, dx\right)^{\gamma_i} \tag{2.9}$$

by (A_1) and (2.5).

Let $c < c^*$ and let (u_j) be a $(PS)_c$ sequence. Then

$$\frac{1}{2}H\left(\int_{\Omega}|\nabla u_j|^2\,dx\right) - \int_{\Omega}F(x,u_j)\,dx - \frac{1}{2^*}\int_{\Omega}|u_j|^{2^*}dx = c + o(1) \tag{2.10}$$

and

$$h\left(\int_{\Omega} |\nabla u_j|^2 \, dx\right) \int_{\Omega} \nabla u_j \cdot \nabla v \, dx - \int_{\Omega} f(x, u_j) \, v \, dx - \int_{\Omega} |u_j|^{2^* - 2} \, u_j \, v \, dx = \mathrm{o}(\|v\|) \quad (2.11)$$

for all $v \in H_0^1(\Omega)$. Taking $v = u_j$ in (2.11), dividing by 2^{*}, and subtracting from (2.10) gives

$$K\left(\int_{\Omega} |\nabla u_j|^2 \, dx\right) - \int_{\Omega} \left[F(x, u_j) - \frac{1}{2^*} \, u_j \, f(x, u_j)\right] dx = c + o(1 + ||u_j||),$$

which together with (2.9) implies that (u_j) is bounded in $H_0^1(\Omega)$. So a renamed subsequence of (u_j) converges to some u weakly in $H_0^1(\Omega)$, strongly in $L^p(\Omega)$, and a.e. in Ω . For a further subsequence, $||u_j - u||^2$ converges to some $t \ge 0$. We will show that t = 0. Suppose $t \ne 0$. Then $t \in I$ by Proposition 2.2 and hence

$$K(t) \ge c^* \tag{2.12}$$

by (2.6). As in the proof of Proposition 2.2,

$$\int_{\Omega} |\nabla u_j|^2 \, dx \to \int_{\Omega} |\nabla u|^2 \, dx + t =: s$$

and

$$sh(s) - \int_{\Omega} uf(x,u) \, dx - \int_{\Omega} |u_j|^{2^*} dx = o(1).$$
 (2.13)

Moreover, passing to the limit in (2.10) gives

$$\frac{1}{2}H(s) - \int_{\Omega} F(x,u) \, dx - \frac{1}{2^*} \int_{\Omega} |u_j|^{2^*} dx = c + o(1),$$

and combining this with (2.13) gives

$$c = K(s) - \int_{\Omega} \left[F(x, u) - \frac{1}{2^*} u f(x, u) \right] dx.$$

Since

$$K(s) \ge K(t) + K\left(\int_{\Omega} |\nabla u|^2 dx\right)$$

by (A_2) and

$$K\left(\int_{\Omega} |\nabla u|^2 \, dx\right) - \int_{\Omega} \left[F(x, u) - \frac{1}{2^*} \, uf(x, u)\right] \, dx \ge 0$$

by (2.9), then

 $c \ge K(t).$

This together with (2.12) gives $c \ge c^*$, contrary to assumption. So t = 0.

If (A_3) holds, then the equation $h(t)/t^{2^*/2-1} = S^{-2^*/2}$ has a unique positive solution t_0 and $I = [t_0, \infty)$. Since K is nondecreasing, then $c^* = K(t_0)$, in particular, (A_1) implies that $c^* > 0$.

Next we consider the case where I is empty. We assume that

- (A_4) h satisfies one of the following conditions:
 - (i) for some constants $\eta > 0$ and $p/2 < \gamma < 2^*/2$,

$$h(t) \ge S^{-2^*/2} t^{2^*/2-1} + \eta t^{\gamma-1} \quad \forall t \ge 0;$$

(*ii*) for some constant $b > S^{-2^*/2}$,

$$h(t) \ge bt^{2^*/2-1} \quad \forall t \ge 0;$$

(*iii*) for some constants b > 0 and $\gamma > 2^*/2$,

$$h(t) > \max \left\{ S^{-2^*/2} t^{2^*/2-1}, bt^{\gamma-1} \right\} \quad \forall t > 0.$$

We have the following theorem.

Theorem 2.4. Assume that (1.2) and (A_4) hold. Then J is bounded from below, coercive, and satisfies the (PS)_c condition for all $c \in \mathbb{R}$. In particular, J has a global minimizer.

Proof. By (1.2) and (2.1),

$$J(u) \ge \frac{1}{2} H\left(\int_{\Omega} |\nabla u|^2 \, dx\right) - c_3 \int_{\Omega} |u|^p \, dx - c_4 - \frac{1}{2^*} S^{-2^*/2} \left(\int_{\Omega} |\nabla u|^2 \, dx\right)^{2^*/2}$$

for some constants $c_3, c_4 > 0$. By $(A_4), H$ satisfies one of the following:

(i)
$$H(t) \ge \frac{2}{2^*} S^{-2^*/2} t^{2^*/2} + \frac{\eta}{\gamma} t^{\gamma}$$
 for all $t \ge 0$, where $\eta > 0$ and $p/2 < \gamma < 2^*/2$;

- (*ii*) $H(t) \ge \frac{2b}{2^*} t^{2^*/2}$ for all $t \ge 0$, where $b > S^{-2^*/2}$;
- (*iii*) $H(t) \ge \frac{b}{\gamma} t^{\gamma}$ for all $t \ge 0$, where b > 0 and $\gamma > 2^*/2$.

It follows that J is bounded from below and coercive.

Let $c \in \mathbb{R}$ and let (u_j) be a $(PS)_c$ sequence. By coercivity, (u_j) is bounded. By (A_4) , $h(t) > S^{-2^{*/2}} t^{2^{*/2}-1}$ for all t > 0, so $I = \emptyset$. So (u_j) has a strongly convergent subsequence by Proposition 2.2.

Finally we apply our results to the model problem

$$\begin{cases} -\left[a+b\left(\int_{\Omega}|\nabla u|^{2} dx\right)^{\gamma-1}\right]\Delta u = f(x,u)+|u|^{2^{*}-2} u \quad \text{in } \Omega\\ u=0 \qquad \qquad \text{on } \partial\Omega, \end{cases}$$
(2.14)

where $a, b \ge 0$ with a + b > 0 and $1 < \gamma < +\infty$. Here

$$h(t) = a + bt^{\gamma - 1}, \qquad H(t) = at + \frac{b}{\gamma}t^{\gamma}, \qquad K(t) = \frac{1}{N}at + \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right)bt^{\gamma}$$

and

$$J(u) = \frac{a}{2} \int_{\Omega} |\nabla u|^2 dx + \frac{b}{2\gamma} \left(\int_{\Omega} |\nabla u|^2 dx \right)^{\gamma} - \int_{\Omega} F(x, u) dx - \frac{1}{2^*} \int_{\Omega} |u|^{2^*} dx, \quad u \in H_0^1(\Omega).$$

Theorem 2.3 has the following corollary for the case $\gamma < 2^*/2$.

Corollary 2.5. Let $1 < \gamma < 2^*/2$ and $a, b \ge 0$. Assume that f satisfies (1.2) and

$$F(x,t) - \frac{1}{2^*} tf(x,t) \le \frac{1}{N} \lambda a t^2 + \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right) \mu b |t|^{2\gamma} \quad for \ a.a. \ x \in \Omega \ and \ all \ t \in \mathbb{R}$$

for some constants $\lambda \leq \lambda_1$ and $\mu \leq \lambda_1(\gamma)$ with either a > 0 and $\lambda < \lambda_1$, or b > 0 and $\mu < \lambda_1(\gamma)$. Let t_0 be the unique positive solution of the equation

 $a + bt^{\gamma - 1} = S^{-2^{*/2}} t^{2^{*/2} - 1}$

and set

$$c^* = \frac{1}{N} a t_0 + \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right) b t_0^{\gamma}.$$

Then J satisfies the $(PS)_c$ condition for all $c < c^*$.

Remark 2.6. For a = 1 and b = 0, Corollary 2.5 gives the well-known compactness threshold

$$c^* = \frac{1}{N} \, S^{N/2}$$

in the Brézis-Nirenberg problem (see [3]).

Remark 2.7. An interesting special case of problem (2.14) is

$$\begin{cases} -\left(\int_{\Omega} |\nabla u|^2 dx\right)^{\gamma-1} \Delta u = \mu |u|^{2\gamma-2} u + |u|^{2^{*-2}} u & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\gamma > 1$ and $\mu > 0$. For $\gamma < 2^*/2$ and $\mu < \lambda_1(\gamma)$, Corollary 2.5 gives the compactness threshold

$$c^* = \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right) S^{2^*\gamma/(2^* - 2\gamma)}$$

for the associated variational functional

$$J(u) = \frac{1}{2\gamma} \left(\int_{\Omega} |\nabla u|^2 \, dx \right)^{\gamma} - \frac{\mu}{2\gamma} \int_{\Omega} |u|^{2\gamma} \, dx - \frac{1}{2^*} \int_{\Omega} |u|^{2^*} \, dx, \quad u \in H_0^1(\Omega).$$

Remark 2.8. The classical case h(t) = a + bt when N = 3 is one of the few cases with both a and b positive and $\gamma < 2^*/2$ where c^* can be found in closed form. Here Corollary 2.5 gives

$$c^* = \frac{1}{4} abS^3 + \frac{1}{24} b^3 S^6 + \frac{1}{24} \left(4aS + b^2 S^4\right)^{3/2}.$$

This threshold level was also obtained in Naimen [10, Lemma 2.5] using concentration compactness arguments. Our approach here is simpler.

Theorem 2.3 also has the following corollary for the case $\gamma = 2^*/2$.

Corollary 2.9. Let $\gamma = 2^*/2$, a > 0, and $0 < b < S^{-2^*/2}$. Assume that f satisfies (1.2) and

$$F(x,t) - \frac{1}{2^*} tf(x,t) \le \frac{1}{N} \lambda a t^2$$
 for a.a. $x \in \Omega$ and all $t \in \mathbb{R}$

for some constant $\lambda < \lambda_1$. Set

$$c^* = \frac{1}{N} \left(\frac{a^{2^*/2}}{S^{-2^*/2} - b} \right)^{2/(2^* - 2)}$$

Then J satisfies the $(PS)_c$ condition for all $c < c^*$.

Remark 2.10. For the classical case h(t) = a + bt with N = 4, a > 0, and $0 < b < S^{-2}$, Corollary 2.9 gives

$$c^* = \frac{a^2}{4\left(S^{-2} - b\right)}$$

This threshold level was also obtained in Naimen [9, Lemma 2.1] by analyzing the behavior of Palais-Smale sequences. Our approach here is simpler again.

Theorem 2.4 has the following corollary for $\gamma \geq 2^*/2$.

Corollary 2.11. Assume that f satisfies (1.2). Then J is bounded from below, satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$, and has a global minimizer in each of the following cases:

- (i) $\gamma = 2^*/2, a \ge 0, and b > S^{-2^*/2};$
- (ii) $\gamma > 2^*/2$ and

$$a^{\gamma-2^*/2} b^{2^*/2-1} > \frac{(\gamma-2^*/2)^{\gamma-2^*/2} (2^*/2-1)^{2^*/2-1}}{(\gamma-1)^{\gamma-1}} S^{-(2^*/2)(\gamma-1)}.$$

Proof. The minimum of $a + bt^{\gamma-1} - S^{-2^*/2} t^{2^*/2-1}$, t > 0 is positive if and only if the last inequality holds.

Remark 2.12. For the classical case h(t) = a + bt with N = 4, $a \ge 0$, and $b > S^{-2}$, Corollary 2.11 (i) implies that J satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$. This was also observed in Liao et al. [8, Proposition 2.1].

Remark 2.13. For the classical case h(t) = a + bt with $N \ge 5$, Corollary 2.11 (*ii*) implies that J satisfies the (PS)_c condition for all $c \in \mathbb{R}$ if

$$a^{N-4}b^2 > \frac{4(N-4)^{N-4}}{(N-2)^{N-2}}S^{-N}.$$

3 Existence and multiplicity results

In the case where I is nonempty, our main existence result for problem (1.1) is the following theorem.

Theorem 3.1. Assume (1.2) and (A_1) – (A_3) . Assume further that

$$H(t) \ge b_0 t^{\gamma_0} \quad for \ 0 \le t \le \delta \tag{3.1}$$

for some constants $\delta, b_0 > 0$ and $1 \leq \gamma_0 < 2^*/2$,

$$F(x,t) \le \frac{1}{2} \mu_0 b_0 |t|^{2\gamma_0}$$
 for a.a. $x \in \Omega$ and $|t| \le \delta$ (3.2)

for some $\mu_0 < \lambda_1(\gamma_0)$, and

$$F(x,t) \ge \frac{1}{q} \nu t^q \quad \text{for a.a. } x \in B_r(x_0) \text{ and all } t \ge 0$$
(3.3)

for some ball $B_r(x_0) \subset \Omega$, $\nu > 0$, and $2\gamma_0 \leq q \leq 2\gamma_n$. Then problem (1.1) has a nontrivial solution in each of the following cases:

- (i) N = 3 and q > 4,
- (ii) $N \ge 4$ and $q \ge N/(N-2)$.

We will show that the functional J has the mountain pass geometry and the mountain pass level is below the compactness threshold c^* in (2.8).

Lemma 3.2. If (1.2), (3.1), and (3.2) hold, then $\exists \rho > 0$ such that

$$\inf_{\|u\|=\rho} J(u) > 0.$$
(3.4)

Proof. By (1.2) and (3.2),

$$F(x,t) \le \frac{1}{2} \mu_0 b_0 |t|^{2\gamma_0} + c_5 |t|^{2^*}$$
 for a.a. $x \in \Omega$ and all $t \in \mathbb{R}$

for some constant $c_5 > 0$. This together with (3.1) implies that for $||u|| \leq \sqrt{\delta}$,

$$J(u) \ge \frac{1}{2} b_0 \left[\left(\int_{\Omega} |\nabla u|^2 \, dx \right)^{\gamma_0} - \mu_0 \int_{\Omega} |u|^{2\gamma_0} \, dx \right] - \left(c_5 + \frac{1}{2^*} \right) \int_{\Omega} |u|^{2^*} dx$$
$$\ge \frac{1}{2} b_0 \left(1 - \frac{\mu_0}{\lambda_1(\gamma_0)} + o(1) \right) \|u\|^{2\gamma_0} \quad \text{as } \|u\| \to 0$$

since $2^* > 2\gamma_0$. Since $\mu_0 < \lambda_1(\gamma_0)$, the desired conclusion follows from this.

Next we show that for a suitably chosen $v \in H_0^1(\Omega) \setminus \{0\}$, $J(sv) \to -\infty$ as $s \to +\infty$ and the maximum of J on the ray $sv, s \ge 0$ is strictly less than c^* . Take a function $\psi \in C_0^{\infty}(B_r(x_0))$ such that $0 \le \psi \le 1$ on $B_r(x_0)$ and $\psi = 1$ on $B_{r/2}(x_0)$, and set

$$u_{\varepsilon}(x) = \frac{\psi(x)}{\left(\varepsilon + |x - x_0|^2\right)^{(N-2)/2}}$$

and

$$v_{\varepsilon} = \frac{u_{\varepsilon}}{|u_{\varepsilon}|_2}$$

for $\varepsilon > 0$. Then

$$\int_{\Omega} |\nabla v_{\varepsilon}|^2 dx = S + \mathcal{O}\left(\varepsilon^{(N-2)/2}\right)$$
(3.5)

and

$$\int_{\Omega} v_{\varepsilon}^{q} dx = \begin{cases} \kappa \varepsilon^{(2N-(N-2)q)/4} + \mathcal{O}(\varepsilon^{(N-2)q/4}) & \text{if } q > N/(N-2) \\ \\ \kappa \varepsilon^{N/4} |\log \varepsilon| + \mathcal{O}(\varepsilon^{N/4}) & \text{if } q = N/(N-2) \end{cases}$$
(3.6)

for some constant $\kappa > 0$ (see, e.g., Drábek and Huang [5]).

Lemma 3.3. For all sufficiently small $\varepsilon > 0$,

$$J(sv_{\varepsilon}) \to -\infty \quad as \ s \to +\infty. \tag{3.7}$$

Proof. Since $|v_{\varepsilon}|_{2^*} = 1$, $|v_{\varepsilon}|_p$ is bounded and (1.2) gives

$$J(sv_{\varepsilon}) \leq \frac{1}{2} H\left(s^2 \int_{\Omega} |\nabla v_{\varepsilon}|^2 dx\right) + c_6 s^p + c_7 - \frac{s^{2^*}}{2^*}, \quad s \geq 0$$

$$(3.8)$$

for some constants $c_6, c_7 > 0$. Set

$$t = s^2 \int_{\Omega} |\nabla v_{\varepsilon}|^2 \, dx.$$

Then $t \to +\infty$ as $s \to +\infty$ and (3.8) gives

$$J(sv_{\varepsilon}) \leq \frac{1}{2} H(t) + c_6 t^{p/2} \left(\int_{\Omega} |\nabla v_{\varepsilon}|^2 \, dx \right)^{-p/2} + c_7 - \frac{t^{2^*/2}}{2^*} \left(\int_{\Omega} |\nabla v_{\varepsilon}|^2 \, dx \right)^{-2^*/2}.$$
 (3.9)

By (A_3) ,

$$\lim_{t \to +\infty} \frac{H(t)}{t^{2^*/2}} = \frac{2b}{2^*}$$

so (3.9) gives

$$J(sv_{\varepsilon}) \le c_6 t^{p/2} \left(\int_{\Omega} |\nabla v_{\varepsilon}|^2 \, dx \right)^{-p/2} + c_7 - \frac{t^{2^*/2}}{2^*} \left[\left(\int_{\Omega} |\nabla v_{\varepsilon}|^2 \, dx \right)^{-2^*/2} - b + \mathrm{o}(1) \right]$$

as $t \to +\infty$. Since $\int_{\Omega} |\nabla v_{\varepsilon}|^2 dx \to S$ as $\varepsilon \to 0$ by (3.5), $b < S^{-2^*/2}$, and $p < 2^*$, the desired conclusion follows.

Lemma 3.4. In each of the two cases in Theorem 3.1,

$$\max_{s \ge 0} J(sv_{\varepsilon}) < c^* \tag{3.10}$$

for all sufficiently small $\varepsilon > 0$.

Proof. Since $v_{\varepsilon} = 0$ outside $B_r(x_0)$, (3.3) gives

$$J(sv_{\varepsilon}) \leq \frac{1}{2} H\left(s^2 \int_{\Omega} |\nabla v_{\varepsilon}|^2 dx\right) - \frac{1}{q} \nu s^q \int_{\Omega} v_{\varepsilon}^q dx - \frac{s^{2^*}}{2^*} =: z_{\varepsilon}(s),$$

so it suffices to show that

$$\max_{s \ge 0} z_{\varepsilon}(s) < c^*$$

for sufficiently small $\varepsilon > 0$. Suppose this is false. Then there are sequences (ε_j) and (s_j) , with $\varepsilon_j, s_j > 0$ and $\varepsilon_j \to 0$, such that

$$z_{\varepsilon_j}(s_j) = \frac{1}{2} H\left(s_j^2 \int_{\Omega} |\nabla v_{\varepsilon_j}|^2 dx\right) - \frac{1}{q} \nu s_j^q \int_{\Omega} v_{\varepsilon_j}^q dx - \frac{s_j^{2^*}}{2^*} \ge c^*$$
(3.11)

and

$$s_j z_{\varepsilon_j}'(s_j) = h\left(s_j^2 \int_{\Omega} |\nabla v_{\varepsilon_j}|^2 dx\right) s_j^2 \int_{\Omega} |\nabla v_{\varepsilon_j}|^2 dx - \nu s_j^q \int_{\Omega} v_{\varepsilon_j}^q dx - s_j^{2^*} = 0.$$
(3.12)

Set

$$t_j = s_j^2 \int_{\Omega} |\nabla v_{\varepsilon_j}|^2 \, dx.$$

Then (3.12) gives

$$\frac{h(t_j)}{t_j^{2^*/2-1}} = \frac{1}{\left(\int_{\Omega} |\nabla v_{\varepsilon_j}|^2 \, dx\right)^{2^*/2}} + \nu t_j^{-(2^*-q)/2} \frac{\int_{\Omega} v_{\varepsilon_j}^q \, dx}{\left(\int_{\Omega} |\nabla v_{\varepsilon_j}|^2 \, dx\right)^{q/2}}.$$
(3.13)

If $t_j \to +\infty$ for a renamed subsequence, then the left-hand side goes to b by (A_3) , while the right-hand side goes to $S^{-2^*/2}$ since $\int_{\Omega} |\nabla v_{\varepsilon_j}|^2 dx \to S$ by (3.5) and $\int_{\Omega} v_{\varepsilon_j}^q dx \to 0$ by (3.6), contradicting our assumption that $b < S^{-2^*/2}$. So (t_j) is bounded, and hence converges to some $t \ge 0$ for a renamed subsequence. Then $s_j^2 \to S^{-1} t$ and hence passing to the limit in (3.11) gives

$$\frac{1}{2} H(t) - \frac{1}{2^*} S^{-2^*/2} t^{2^*/2} > 0$$

since $c^* > 0$, so t > 0. On the other hand, passing to the limit in (3.12) shows that t satisfies (2.7). Since t_0 is the unique positive solution of this equation, it follows that $t = t_0$.

Now combining (3.13) with (3.5) and (3.6) gives

$$\frac{h(t_j)}{t_j^{2^*/2-1}} = S^{-2^*/2} + \begin{cases} \sigma_j \, \varepsilon_j^{(2N-(N-2)q)/4} + \mathcal{O}\big(\varepsilon_j^{(N-2)/2}\big) & \text{if } q > N/(N-2) \\ \sigma_j \, \varepsilon_j^{N/4} \, |\log \varepsilon_j| + \mathcal{O}\big(\varepsilon_j^{\min\{(N-2)/2, N/4\}}\big) & \text{if } q = N/(N-2), \end{cases}$$

where $\sigma_j \to \kappa \nu S^{-q/2} t_0^{-(2^*-q)/2} > 0$. It follows from this that in each of the two cases in the lemma,

$$\frac{h(t_j)}{t_j^{2^*/2-1}} \ge S^{-2^*/2} = \frac{h(t_0)}{t_0^{2^*/2-1}}$$

for all sufficiently large j. Then $t_j \leq t_0$ by (A_3) . Since K is nondecreasing, then

$$K(t_j) \le K(t_0) = c^*.$$

However, dividing (3.12) by 2^* and subtracting from (3.11) gives

$$K(t_j) - \left(\frac{1}{q} - \frac{1}{2^*}\right) \nu s_j^q \int_{\Omega} v_{\varepsilon_j}^q \, dx \ge c^*,$$

so $K(t_j) > c^*$. This contradiction completes the proof.

We are now ready to prove Theorem 3.1.

Proof of Theorem 3.1. Let ρ be as in Lemma 3.2 and fix $\varepsilon > 0$ such that (3.7) and (3.10) hold. Then $\exists R > \rho$ such that $J(Rv_{\varepsilon}) \leq 0$. Let

$$\Gamma = \left\{ \varphi \in C([0,1], H_0^1(\Omega)) : \varphi(0) = 0, \, \varphi(1) = Rv_{\varepsilon} \right\}$$

be the class of paths in $H_0^1(\Omega)$ joining the origin to Rv_{ε} , and set

$$c := \inf_{\varphi \in \Gamma} \max_{u \in \varphi([0,1])} J(u).$$

By (3.4), c > 0. Since the path $\varphi_0(s) = sRv_{\varepsilon}, s \in [0, 1]$ is in Γ ,

$$c \le \max_{u \in \varphi_0([0,1])} J(u) \le \max_{s \ge 0} J(sv_{\varepsilon}) < c^*,$$

so J satisfies the $(PS)_c$ condition by Theorem 2.3. Hence J has a critical point u with J(u) = c by the mountain pass theorem (see Ambrosetti and Rabinowitz [1]). Then u is a weak solution of problem (1.1) and u is nontrivial since c > 0.

Theorem 3.1 has many interesting consequences, some of which we now present. First we consider the problem

$$\begin{cases} -h\left(\int_{\Omega} |\nabla u|^2 \, dx\right) \Delta u = \lambda u + |u|^{2^* - 2} \, u \quad \text{in } \Omega \\ u = 0 \qquad \text{on } \partial\Omega, \end{cases}$$
(3.14)

where $\lambda > 0$. Assume that

 $K(t) \ge \alpha t \quad \forall t \ge 0 \tag{3.15}$

for some constant $\alpha > 0$ and

$$H(t) \ge a_0 t \quad \text{for } 0 \le t \le \delta \tag{3.16}$$

for some constants $\delta, a_0 > 0$. We have $f(x, t) = \lambda t$ and

$$F(x,t) = \frac{1}{2} \lambda t^2, \qquad F(x,t) - \frac{1}{2^*} t f(x,t) = \frac{1}{N} \lambda t^2,$$

so (A_1) holds with $\mu_1 = \lambda/N\alpha$ if $\lambda < N\alpha\lambda_1$, (3.2) holds with $\gamma_0 = 1$ and $\mu_0 = \lambda/a_0$ if $\lambda < a_0\lambda_1$, and (3.3) holds with q = 2 if $\lambda > 0$. So Theorem 3.1 has the following corollary for problem (3.14).

Corollary 3.5. Assume (3.15), (A_2) , (A_3) , and (3.16). If

 $0 < \lambda < \min\{a_0, N\alpha\}\lambda_1$

and $N \geq 4$, then problem (3.14) has a nontrivial solution.

In particular, we have the following corollary in the model case $h(t) = a + bt^{\gamma-1}$.

Corollary 3.6. The problem

$$\begin{cases} -\left[a+b\left(\int_{\Omega}|\nabla u|^{2} dx\right)^{\gamma-1}\right]\Delta u = \lambda u + |u|^{2^{*}-2} u \quad in \ \Omega \\ u = 0 \qquad on \ \partial\Omega, \end{cases}$$

where either $1 < \gamma < 2^*/2$ and $b \ge 0$, or $\gamma = 2^*/2$ and $0 \le b < S^{-2^*/2}$, has a nontrivial solution if $0 < \lambda < a\lambda_1$ and $N \ge 4$.

Next we consider the problem

$$\begin{cases} -h\left(\int_{\Omega} |\nabla u|^2 \, dx\right) \Delta u = \mu \, |u|^{2\gamma - 2} \, u + |u|^{2^* - 2} \, u & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$
(3.17)

where $\mu > 0$ and $1 < \gamma < 2^*/2$. Assume that

 $K(t) \ge \beta t^{\gamma} \quad \forall t \ge 0 \tag{3.18}$

for some constant $\beta > 0$ and

 $H(t) \ge b_0 t^{\gamma} \quad \text{for } 0 \le t \le \delta \tag{3.19}$

for some constants $\delta, b_0 > 0$. We have $f(x, t) = \mu |t|^{2\gamma - 2} t$ and

$$F(x,t) = \frac{1}{2\gamma} \,\mu \,|t|^{2\gamma}, \qquad F(x,t) - \frac{1}{2^*} \,tf(x,t) = \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right) \mu \,|t|^{2\gamma},$$

so (A₁) holds with $\mu_1 = (1/2\gamma - 1/2^*) \mu/\beta$ if

$$\mu < \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right)^{-1} \beta \lambda_1(\gamma),$$

(3.2) holds with $\gamma_0 = \gamma$ and $\mu_0 = \mu/\gamma b_0$ if $\mu < \gamma b_0 \lambda_1(\gamma)$, and (3.3) holds with $q = 2\gamma$ if $\mu > 0$. So Theorem 3.1 has the following corollary for problem (3.17).

Corollary 3.7. Assume (3.18), (A_2) , (A_3) , and (3.19). If

$$0 < \mu < \min\left\{\gamma b_0, \left(\frac{1}{2\gamma} - \frac{1}{2^*}\right)^{-1}\beta\right\}\lambda_1(\gamma)$$

and $N \ge 4$, or N = 3 and $\gamma > 2$, then problem (3.17) has a nontrivial solution.

In particular, we have the following corollary in the model case $h(t) = a + bt^{\gamma-1}$.

Corollary 3.8. The problem

$$\begin{cases} -\left[a+b\left(\int_{\Omega}|\nabla u|^{2} dx\right)^{\gamma-1}\right]\Delta u=\mu |u|^{2\gamma-2} u+|u|^{2^{\ast}-2} u \quad in \ \Omega \\ u=0 \qquad \qquad on \ \partial\Omega, \end{cases}$$

where $a \ge 0$ and $1 < \gamma < 2^*/2$, has a nontrivial solution if $0 < \mu < b\lambda_1(\gamma)$ and $N \ge 4$, or N = 3 and $\gamma > 2$.

Finally we consider the problem

$$\begin{cases} -h\left(\int_{\Omega} |\nabla u|^2 \, dx\right) \Delta u = \nu \, |u|^{q-2} \, u + |u|^{2^*-2} \, u \quad \text{in } \Omega \\ u = 0 \qquad \qquad \text{on } \partial\Omega, \end{cases}$$
(3.20)

where $\nu > 0$ and $2 < q < 2^*$. Assume that for some constants $\alpha, \beta > 0$ and $q/2 < \gamma < 2^*/2$,

$$K(t) \ge \alpha t + \beta t^{\gamma} \quad \forall t \ge 0. \tag{3.21}$$

Since h is nonnegative, $H(t) \ge 2K(t) \ge 2\alpha t$, so (3.1) holds with $b_0 = 2\alpha$ and $\gamma_0 = 1$. We have $f(x,t) = \nu |t|^{q-2} t$ and

$$F(x,t) = \frac{1}{q}\nu|t|^{q}, \qquad F(x,t) - \frac{1}{2^{*}}tf(x,t) = \left(\frac{1}{q} - \frac{1}{2^{*}}\right)\nu|t|^{q}.$$

Since q > 2, (3.2) holds for any $\mu_0 > 0$ if $\delta > 0$ is sufficiently small. Theorem 3.1 has the following corollary for problem (3.20).

Corollary 3.9. Assume (3.21), (A_2) , and (A_3) . If

$$0 < \nu < (2\gamma - 2) \left(\frac{1}{q} - \frac{1}{2^*}\right)^{-1} \left(\frac{\alpha\lambda_1}{2\gamma - q}\right)^{(2\gamma - q)/(2\gamma - 2)} \left(\frac{\beta\lambda_1(\gamma)}{q - 2}\right)^{(q-2)/(2\gamma - 2)}$$

and $N \ge 4$, or N = 3 and q > 4, then problem (3.20) has a nontrivial solution. Proof. To see that (A_1) holds, note that the minimum of

$$\lambda \alpha t^2 + \mu \beta |t|^{2\gamma} - \left(\frac{1}{q} - \frac{1}{2^*}\right) \nu |t|^q, \quad t \in \mathbb{R}$$

is nonnegative if and only if

$$\nu \le (2\gamma - 2) \left(\frac{1}{q} - \frac{1}{2^*}\right)^{-1} \left(\frac{\alpha\lambda}{2\gamma - q}\right)^{(2\gamma - q)/(2\gamma - 2)} \left(\frac{\beta\mu}{q - 2}\right)^{(q - 2)/(2\gamma - 2)}.$$

In the case where I is empty, first we consider the model problem

$$\begin{cases} -\left[a+b\left(\int_{\Omega}|\nabla u|^{2} dx\right)^{\gamma-1}\right]\Delta u = \lambda u + g(x,u) + |u|^{2^{*}-2} u \quad \text{in } \Omega \\ u = 0 \qquad \qquad \text{on } \partial\Omega, \end{cases}$$
(3.22)

where $a, b \ge 0$ and $2^*/2 \le \gamma < +\infty$ satisfy one of the two conditions in Corollary 2.11, $\lambda > 0$, and g is a Carathéodory function on $\Omega \times \mathbb{R}$ satisfying

$$g(x,t) = o(t)$$
 as $t \to 0$, uniformly a.e. in Ω (3.23)

and

$$|g(x,t)| \le c_8 |t|^{p-1} + c_9 \quad \text{for a.a. } x \in \Omega \text{ and all } t \in \mathbb{R}$$
(3.24)

for some constants $c_8, c_9 > 0$ and 2 . The associated variational functional is

$$J(u) = \frac{a}{2} \int_{\Omega} |\nabla u|^2 dx + \frac{b}{2\gamma} \left(\int_{\Omega} |\nabla u|^2 dx \right)^{\gamma} - \frac{\lambda}{2} \int_{\Omega} u^2 dx - \int_{\Omega} G(x, u) dx - \frac{1}{2^*} \int_{\Omega} |u|^{2^*} dx,$$
$$u \in H_0^1(\Omega),$$

where $G(x,t) = \int_0^t g(x,s) \, ds$ is the primitive of g. We note that

$$\int_{\Omega} G(x, u) \, dx = \mathrm{o}(\|u\|^2) \quad \text{as } \|u\| \to 0 \tag{3.25}$$

by (3.23) and (3.24).

When a = 0, we have the following existence result.

Theorem 3.10. Assume that g satisfies (3.23) and (3.24). If $\gamma = 2^*/2$, a = 0, and $b > S^{-2^*/2}$, then problem (3.22) has a nontrivial solution for all $\lambda > 0$.

Proof. By Corollary 2.11, J has a global minimizer u_0 . For any $u \in H_0^1(\Omega) \setminus \{0\}$,

$$J(su) = -\frac{\lambda s^2}{2} \int_{\Omega} u^2 dx + o(s^2) \quad \text{as } s \to 0$$

by (3.25), so J(su) < 0 if s > 0 is sufficiently small. So $J(u_0) = \inf_{H_0^1(\Omega)} J < 0$ and hence u_0 is nontrivial.

When a > 0, we prove a multiplicity result. Let $0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots$ be the Dirichlet eigenvalues of $-\Delta$ on Ω , repeated according to multiplicity.

Theorem 3.11. Assume that g satisfies (3.23) and (3.24).

- (i) If $\gamma = 2^*/2$, a > 0, and $b > S^{-2^*/2}$, then problem (3.22) has at least two nontrivial solutions in each of the following cases:
 - (a) $a\lambda_k < \lambda < a\lambda_{k+1}$ for some $k \ge 1$;
 - (b) $a\lambda_k < \lambda = a\lambda_{k+1}$ for some $k \ge 1$ and $G(x,t) \le 0$ for a.a. $x \in \Omega$ and $|t| \le \delta$ for some $\delta > 0$.

(*ii*) If $\gamma > 2^*/2$ and

$$a^{\gamma-2^*/2} b^{2^*/2-1} > \frac{(\gamma-2^*/2)^{\gamma-2^*/2} (2^*/2-1)^{2^*/2-1}}{(\gamma-1)^{\gamma-1}} S^{-(2^*/2)(\gamma-1)},$$

then problem (3.22) has at least two nontrivial solutions in each of the following cases:

- (a) $a\lambda_k < \lambda < a\lambda_{k+1}$ for some $k \ge 1$;
- (b) $a\lambda_k = \lambda < a\lambda_{k+1}$ for some $k \ge 1$ and $G(x,t) \ge 0$ for a.a. $x \in \Omega$ and $|t| \le \delta$ for some $\delta > 0$.

We will prove this theorem using the following result of Brezis and Nirenberg [4, Theorem 4].

Proposition 3.12. Let J be a C^1 -functional on a Banach space X. Assume that J is bounded from below, $\inf_X J < 0$, and J satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$. Assume further that X has a direct sum decomposition $X = V \oplus W$, u = v + w with $\dim V < +\infty$ and

$$\begin{cases} J(v) \le 0 & \text{for } v \in V \cap B_r(0) \\ J(w) \ge 0 & \text{for } w \in W \cap B_r(0) \end{cases}$$

for some r > 0. Then J has at least two nontrivial critical points.

Proof of Theorem 3.11. By Corollary 2.11, J is bounded from below and satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$. We have the direct sum decomposition $H_0^1(\Omega) = V \oplus W$, u = v + w,

where V is the span of the eigenfunctions associated with $\lambda_1, \ldots, \lambda_k$ and W is the orthogonal complement of V. For $v \in V$,

$$J(v) \leq -\frac{1}{2} \left(\frac{\lambda}{\lambda_k} - a\right) \int_{\Omega} |\nabla v|^2 dx + \frac{b}{2\gamma} \left(\int_{\Omega} |\nabla v|^2 dx\right)^{\gamma} - \int_{\Omega} G(x, v) dx - \frac{1}{2^*} \int_{\Omega} |v|^{2^*} dx$$
$$= -\frac{1}{2} \left(\frac{\lambda}{\lambda_k} - a\right) \int_{\Omega} |\nabla v|^2 dx + o(||v||^2) \quad \text{as } ||v|| \to 0 \quad (3.26)$$

by (3.25), so J(v) < 0 if $\lambda > a\lambda_k$ and ||v|| > 0 is sufficiently small. For $w \in W$,

$$\begin{aligned} J(w) &\geq \frac{1}{2} \left(a - \frac{\lambda}{\lambda_{k+1}} \right) \int_{\Omega} |\nabla w|^2 \, dx + \frac{b}{2\gamma} \left(\int_{\Omega} |\nabla w|^2 \, dx \right)^{\gamma} - \int_{\Omega} G(x, w) \, dx - \frac{1}{2^*} \int_{\Omega} |w|^{2^*} dx \\ &= \frac{1}{2} \left(a - \frac{\lambda}{\lambda_{k+1}} \right) \int_{\Omega} |\nabla w|^2 \, dx + \mathrm{o}(\|w\|^2) \quad \text{as} \ \|w\| \to 0 \end{aligned}$$

by (3.25), so $J(w) \ge 0$ if $\lambda < a\lambda_{k+1}$ and ||w|| is sufficiently small. So J has at least two nontrivial critical points by Proposition 3.12 in the cases (i)(a) and (ii)(a).

In the case (i)(b), (2.1) gives

$$J(w) \ge \int_{\Omega} \left[\frac{a}{2} \left(|\nabla w|^2 - \lambda_{k+1} w^2 \right) - G(x, w) \right] dx + \frac{b - S^{-2^*/2}}{2^*} \left(\int_{\Omega} |\nabla w|^2 dx \right)^{2^*/2} \quad \forall w \in W.$$

The local sign condition on G in this case implies that the first integral on the right-hand side is nonnegative if ||w|| is sufficiently small (see Li and Willem [7]). Since $b > S^{-2^{*/2}}$, then $J(w) \ge 0$ when ||w|| is small. In the case (ii)(b), (3.26) gives

$$J(v) \le \frac{b}{2\gamma} \left(\int_{\Omega} |\nabla v|^2 \, dx \right)^{\gamma} - \int_{\{|v| > \delta\}} G(x, v) \, dx - \frac{1}{2^*} \int_{\Omega} |v|^{2^*} dx \quad \forall v \in V.$$

Since V is a finite dimensional subspace of $H_0^1(\Omega)$ consisting of L^{∞} -functions and $\gamma > 2^*/2$, it follows from this that J(v) < 0 if ||v|| > 0 is sufficiently small. So J has two nontrivial critical points in these cases also.

In the borderline case where $\gamma = 2^*/2$ and $b = S^{-2^*/2}$, lower-order terms come into play. We consider the problem

$$\begin{cases} -h\left(\int_{\Omega} |\nabla u|^2 \, dx\right) \Delta u = \lambda u + g(x, u) + |u|^{2^* - 2} \, u \quad \text{in } \Omega \\ u = 0 \qquad \qquad \text{on } \partial\Omega, \end{cases}$$
(3.27)

where

$$h(t) = a + S^{-2^*/2} t^{2^*/2 - 1} + \eta t^{\sigma - 1}, \quad t \ge 0,$$

 $a \ge 0, \eta > 0, p/2 < \sigma < 2^*/2, \lambda > 0$, and g is a Carathéodory function on $\Omega \times \mathbb{R}$ satisfying (3.23) and (3.24). The associated functional is

$$\begin{split} J(u) &= \frac{a}{2} \int_{\Omega} |\nabla u|^2 \, dx + \frac{S^{-2^*/2}}{2^*} \left(\int_{\Omega} |\nabla u|^2 \, dx \right)^{2^*/2} + \frac{\eta}{2\sigma} \left(\int_{\Omega} |\nabla u|^2 \, dx \right)^{\sigma} \\ &- \frac{\lambda}{2} \int_{\Omega} u^2 \, dx - \int_{\Omega} G(x, u) \, dx - \frac{1}{2^*} \int_{\Omega} |u|^{2^*} dx, \quad u \in H^1_0(\Omega), \end{split}$$

where $G(x,t) = \int_0^t g(x,s) \, ds$ satisfies (3.25) as before. We have the following existence and multiplicity result.

Theorem 3.13. Let $\eta > 0$ and $p/2 < \sigma < 2^*/2$, and assume that g satisfies (3.23) and (3.24).

- (i) If a = 0, then problem (3.27) has a nontrivial solution for all $\lambda > 0$.
- (ii) If a > 0, then problem (3.27) has at least two nontrivial solutions in each of the following cases:
 - (a) $a\lambda_k < \lambda < a\lambda_{k+1}$ for some $k \ge 1$;
 - (b) $a\lambda_k < \lambda = a\lambda_{k+1}$ for some $k \ge 1$ and $G(x,t) \le 0$ for a.a. $x \in \Omega$ and $|t| \le \delta$ for some $\delta > 0$.

Proof. (i) By Theorem 2.4, J has a global minimizer u_0 . For any $u \in H_0^1(\Omega) \setminus \{0\}$,

$$J(su) = -\frac{\lambda s^2}{2} \int_{\Omega} u^2 dx + o(s^2) \quad \text{as } s \to 0$$

by (3.25), so J(su) < 0 if s > 0 is sufficiently small. So $J(u_0) = \inf_{H_0^1(\Omega)} J < 0$ and hence u_0 is nontrivial.

(*ii*) By Theorem 2.4, J is bounded from below and satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$. We have the direct sum decomposition $H_0^1(\Omega) = V \oplus W$, u = v + w, where V is the span of the eigenfunctions associated with $\lambda_1, \ldots, \lambda_k$ and W is the orthogonal complement of V. For $v \in V$,

$$J(v) \leq -\frac{1}{2} \left(\frac{\lambda}{\lambda_k} - a\right) \int_{\Omega} |\nabla v|^2 dx + \frac{S^{-2^*/2}}{2^*} \left(\int_{\Omega} |\nabla v|^2 dx\right)^{2^*/2} + \frac{\eta}{2\sigma} \left(\int_{\Omega} |\nabla v|^2 dx\right)^{\sigma} - \int_{\Omega} G(x, v) dx - \frac{1}{2^*} \int_{\Omega} |v|^{2^*} dx = -\frac{1}{2} \left(\frac{\lambda}{\lambda_k} - a\right) \int_{\Omega} |\nabla v|^2 dx + o(||v||^2) \quad \text{as } ||v|| \to 0$$

by (3.25), so J(v) < 0 if $\lambda > a\lambda_k$ and ||v|| > 0 is sufficiently small. For $w \in W$,

$$J(w) \ge \frac{1}{2} \left(a - \frac{\lambda}{\lambda_{k+1}} \right) \int_{\Omega} |\nabla w|^2 \, dx + \frac{S^{-2^*/2}}{2^*} \left(\int_{\Omega} |\nabla w|^2 \, dx \right)^{2^*/2} + \frac{\eta}{2\sigma} \left(\int_{\Omega} |\nabla w|^2 \, dx \right)^{\sigma} \\ - \int_{\Omega} G(x, w) \, dx - \frac{1}{2^*} \int_{\Omega} |w|^{2^*} \, dx = \frac{1}{2} \left(a - \frac{\lambda}{\lambda_{k+1}} \right) \int_{\Omega} |\nabla w|^2 \, dx + o(||w||^2) \quad \text{as } ||w|| \to 0$$

by (3.25), so $J(w) \ge 0$ if $\lambda < a\lambda_{k+1}$ and ||w|| is sufficiently small. So J has at least two nontrivial critical points by Proposition 3.12 in the case (a). In the case (b), (2.1) gives

$$J(w) \ge \int_{\Omega} \left[\frac{a}{2} \left(|\nabla w|^2 - \lambda_{k+1} w^2 \right) - G(x, w) \right] dx \quad \forall w \in W.$$

The local sign condition on G implies that the right-hand side is nonnegative when ||w|| is small (see Li and Willem [7]). So J has two nontrivial critical points in this case also.

References

- [1] Antonio Ambrosetti and Paul H. Rabinowitz. Dual variational methods in critical point theory and applications. J. Functional Analysis, 14:349–381, 1973.
- [2] Haïm Brézis and Elliott Lieb. A relation between pointwise convergence of functions and convergence of functionals. Proc. Amer. Math. Soc., 88(3):486–490, 1983.
- [3] Haïm Brézis and Louis Nirenberg. Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. Comm. Pure Appl. Math., 36(4):437–477, 1983.
- [4] Haïm Brezis and Louis Nirenberg. Remarks on finding critical points. Comm. Pure Appl. Math., 44(8-9):939–963, 1991.
- [5] Pavel Drábek and Yin Xi Huang. Multiplicity of positive solutions for some quasilinear elliptic equation in \mathbf{R}^N with critical Sobolev exponent. J. Differential Equations, 140(1):106–132, 1997.
- [6] Yisheng Huang, Zeng Liu, and Yuanze Wu. On Kirchhoff type equations with critical Sobolev exponent. J. Math. Anal. Appl., 462(1):483–504, 2018.
- [7] Shu Jie Li and Michel Willem. Applications of local linking to critical point theory. J. Math. Anal. Appl., 189(1):6–32, 1995.
- [8] Jia-Feng Liao, Xiao-Feng Ke, Jiu Liu, and Chun-Lei Tang. The Brezis-Nirenberg result for the Kirchhoff-type equation in dimension four. Appl. Anal., 97(15):2720–2726, 2018.
- [9] Daisuke Naimen. The critical problem of Kirchhoff type elliptic equations in dimension four. J. Differential Equations, 257(4):1168–1193, 2014.
- [10] Daisuke Naimen. Positive solutions of Kirchhoff type elliptic equations involving a critical Sobolev exponent. NoDEA Nonlinear Differential Equations Appl., 21(6):885– 914, 2014.
- [11] Daisuke Naimen and Masataka Shibata. Two positive solutions for the Kirchhoff type elliptic problem with critical nonlinearity in high dimension. *Nonlinear Anal.*, 186:187– 208, 2019.
- [12] Kanishka Perera and Zhitao Zhang. Nontrivial solutions of Kirchhoff-type problems via the Yang index. J. Differential Equations, 221(1):246–255, 2006.

- [13] Qi-Lin Xie, Xing-Ping Wu, and Chun-Lei Tang. Existence and multiplicity of solutions for Kirchhoff type problem with critical exponent. *Commun. Pure Appl. Anal.*, 12(6):2773–2786, 2013.
- [14] Xianzhong Yao and Chunlai Mu. Multiplicity of solutions for Kirchhoff type equations involving critical Sobolev exponents in high dimension. *Math. Methods Appl. Sci.*, 39(13):3722–3734, 2016.
- [15] Chunhong Zhang and Zhisu Liu. Multiplicity of nontrivial solutions for a critical degenerate Kirchhoff type problem. Appl. Math. Lett., 69:87–93, 2017.