# LINKEDNESS OF CARTESIAN PRODUCTS OF COMPLETE GRAPHS

LEIF K. JØRGENSEN

Department of Mathematical Sciences, Aalborg University, Denmark

GUILLERMO PINEDA-VILLAVICENCIO

Centre for Informatics and Applied Optimisation, Federation University, Australia School of Information Technology, Deakin University, Geelong, Australia

## JULIEN UGON

Centre for Informatics and Applied Optimisation, Federation University, Australia School of Information Technology, Deakin University, Geelong, Australia

ABSTRACT. This paper is concerned with the linkedness of Cartesian products of complete graphs. A graph with at least 2k vertices is *k*-linked if, for every set of 2k distinct vertices organised in arbitrary k pairs of vertices, there are k vertex-disjoint paths joining the vertices in the pairs.

We show that the Cartesian product  $K^{d_1+1} \times K^{d_2+1}$  of complete graphs  $K^{d_1+1}$  and  $K^{d_2+1}$  is  $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for  $d_1, d_2 \ge 2$ , and this is best possible.

This result is connected to graphs of simple polytopes. The Cartesian product  $K^{d_1+1} \times K^{d_2+1}$  is the graph of the Cartesian product  $T(d_1) \times T(d_2)$  of a  $d_1$ -dimensional simplex  $T(d_1)$  and a  $d_2$ -dimensional simplex  $T(d_2)$ . And the polytope  $T(d_1) \times T(d_2)$  is a simple polytope, a  $(d_1 + d_2)$ -dimensional polytope in which every vertex is incident to exactly  $d_1 + d_2$  edges.

While not every *d*-polytope is  $\lfloor d/2 \rfloor$ -linked, it may be conjectured that every simple *d*-polytope is. Our result implies the veracity of the revised conjecture for Cartesian products of two simplices.

*E-mail addresses*: leifkjorgensen@gmail.com, work@guillermo.com.au, julien.ugon@deakin.edu.au.

Date: July 11, 2021.

<sup>2020</sup> Mathematics Subject Classification. Primary 05C40; Secondary 52B05.

 $K\!ey$  words and phrases.  $k\mbox{-linked},$  cyclic polytope, connectivity, separator, dual polytope, linkedness, Cartesian product.

2

#### 1. INTRODUCTION

Denote by V(X) the vertex set of a graph. Given sets A, B of vertices in a graph, a path from A to B, called an A - B path, is a (vertex-edge) path  $L := u_0 \ldots u_n$  in the graph such that  $V(L) \cap A = \{u_0\}$  and  $V(L) \cap B = \{u_n\}$ . We write a - B path instead of  $\{a\} - B$  path, and likewise, write A - b path instead of  $A - \{b\}$ .

Let G be a graph and X a subset of 2k distinct vertices of G. The elements of X are called *terminals*. Let  $Y := \{\{s_1, t_1\}, \ldots, \{s_k, t_k\}\}$  be an arbitrary labelling and (unordered) pairing of all the vertices in X. We say that Y is *linked* in G if we can find disjoint  $s_i - t_i$  paths for  $i \in [1, k]$ , the interval  $1, \ldots, k$ . The set X is *linked* in G if every such pairing of its vertices is linked in G. Throughout this paper, by a set of disjoint paths, we mean a set of vertex-disjoint paths. If G has at least 2k vertices and every set of exactly 2k vertices is linked in G, we say that G is k-linked.

This paper studies the linkedness of Cartesian products of complete graphs. Linkedness of Cartesian products has been studied in the past (Mészáros, 2016). The *Cartesian product*  $G_1 \times G_2$  of two graphs  $G_1$  and  $G_2$  is the graph defined on the pairs  $(v_1, v_2)$  with  $v_i \in G_i$  and with two pairs  $(u_1, u_2)$  and  $(v_1, v_2)$  being adjacent if, for some  $\ell \in \{1, 2\}$ ,  $u_\ell v_\ell \in E(G_\ell)$  and  $u_i = v_i$  for  $i \neq \ell$ . We prove that the Cartesian product  $K^{d_1+1} \times K^{d_2+1}$  of complete graphs  $K^{d_1+1}$  and  $K^{d_2+1}$ is  $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for  $d_1, d_2 \ge 0$ , and that there are products that are not  $\lfloor (d_1 + d_2 + 1)/2 \rfloor$ -linked; hence this result is best possible. Here  $K^t$  denotes the complete graph on t vertices.

Our result is connected to questions on the linkedness of a polytope. A (convex) polytope is the convex hull of a finite set X of points in  $\mathbb{R}^d$ ; the *convex hull* of X is the smallest convex set containing X. The *dimension* of a polytope in  $\mathbb{R}^d$  is one less than the maximum number of affinely independent points in the polytope; a set of points  $\vec{p}_1, \ldots, \vec{p}_k$  in  $\mathbb{R}^d$  is affinely independent if the k-1 vectors  $\vec{p}_1 - \vec{p}_k, \ldots, \vec{p}_{k-1} - \vec{p}_k$  are linearly independent. A polytope of dimension d is referred to as a d-polytope.

The Cartesian product  $P \times P'$  of a *d*-polytope  $P \subset \mathbb{R}^d$  and a *d'*-polytope  $P' \subset \mathbb{R}^{d'}$  is the Cartesian product of the sets P and P':

$$P \times P' = \left\{ \begin{pmatrix} p \\ p' \end{pmatrix} \in \mathbb{R}^{d+d'} \middle| p \in P, p' \in P \right\}.$$

The resulting polytope is (d + d')-dimensional. The graph G(P) of a polytope P is the undirected graph formed by the vertices and edges of the polytope. It follows that the graph  $G(P \times P')$  of the Cartesian product  $P \times P'$  is the Cartesian product  $G(P) \times G(P')$  of the graphs G(P) and G(P').

A *d*-simplex T(d) is the convex hull of d + 1 affinely independent points in  $\mathbb{R}^d$ . The graph of T(d) is the complete graph  $K^{d+1}$ . As a consequence, our result implies that the graph of the Cartesian product  $T(d_1) \times T(d_2)$  is  $\lfloor (d_1 + d_2)/2 \rfloor$ linked for  $d_1, d_2 \ge 0$ . Henceforth, if the graph of a polytope is *k*-linked we say that the polytope is also *k*-linked.

A-B path

terminals linked pairs linked set

k-linked

The first edition of the Handbook of Discrete and Computational Geometry (Goodman & O'Rourke, 1997, Problem 17.2.6) posed the question of whether or not every *d*-polytope is  $\lfloor d/2 \rfloor$ -linked. This question was answered in the negative by Gallivan (1985). None of the known counterexamples are *simple d-polytopes*, *d*-polytopes in which every vertex is incident to exactly *d* edges. Hence, it may be hypothesised that the conjecture holds for such polytopes.

#### **Conjecture 1.** Every simple d-polytope is $\lfloor d/2 \rfloor$ -linked for $d \ge 2$ .

Cartesian products of simplices are simple polytopes, and so our result supports this revised conjecture. Furthermore, Cartesian products of simplices and duals of cyclic polytopes are related; the dual of a cyclic *d*-polytope with d + 2 vertices is the Cartesian product of a  $\lfloor d/2 \rfloor$ -simplex and a  $\lceil d/2 \rceil$ -simplex (Ziegler, 1995, Ex. 0.6). Hence we obtain that the dual of a cyclic *d*-polytope on d + 2 vertices is also  $\lfloor d/2 \rfloor$ -linked for  $d \ge 2$ .

Unless otherwise stated, the graph theoretical notation and terminology follows from Diestel (2017) and the polytope theoretical notation and terminology from Ziegler (1995). Moreover, when referring to graph-theoretical properties of a polytope such as linkedness and connectivity, we mean properties of its graph.

#### 2. LINKEDNESS OF CARTESIAN PRODUCTS OF COMPLEX GRAPHS

The contribution of this section is a sharp theorem (Theorem 2) that tells the story of the linkedness of Cartesian product of two complete graphs.

**Theorem 2.** The Cartesian product of two complete graphs  $K^{d_1+1}$  and  $K^{d_2+1}$  is  $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for every  $d_1, d_2 \ge 0$ .

*Remark* 3. Theorem 2 is best possible. There are products  $K^{d_1+1} \times K^{d_2+1}$  that are not  $\lfloor (d_1 + d_2 + 1)/2 \rfloor$ -linked:

- (i)  $K^2 \times K^{d_2+1}$  for even  $d_2 \ge 1$ , and
- (ii)  $K^3 \times K^{d_2+1}$  for  $d_2 = 1, 3, 5, 7, 9$ .

For each of these cases, Fig. 1 provides a pairing of terminals that cannot be  $\lfloor (d_1 + d_2 + 1)/2 \rfloor$ -linked. We conjecture these are the only such cases.

An immediate corollary of Theorem 2 is the following.

**Corollary 4.** The Cartesian product of two simplices  $T(d_1)$  and  $T(d_2)$  is  $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for every  $d_1, d_2 \ge 0$ .

The notions of linkage, linkage problem, and valid path will simplify our arguments. A *linkage* in a graph is a subgraph in which every component is a path. Let X be a set of vertices in a graph and let  $Y := \{\{s_1, t_1\}, \ldots, \{s_k, t_k\}\}$  be a pairing of all the vertices of X. A Y-linkage  $\{L_1, \ldots, L_k\}$  is a set of disjoint paths with the path  $L_i$  joining the pair  $\{s_i, t_i\}$  for  $i = 1, \ldots, k$ . We may also say that Y represents our *linkage problem*, and if Y is linked in G then our linkage problem is *feasible* and *infeasible* otherwise. A path in the graph is called X-valid if no inner vertex

	${}^{s_1}{\mathbf{o}}$	$\overset{s_2}{\mathbf{O}}$	0	<sup>33</sup> t	$\overset{3}{5}\cdots$	$\mathbf{\hat{b}}$	$\mathbf{o}^{t_k}$		0	0	o t <sub>3</sub>	$s^{s_1}$		
									0	0	$o t_1$ o	о <i>s</i> 2		
	$\mathbf{o}_{t_2}$	$\mathbf{o}_{t_1}$	0	0 0	o… c	0	0		0	0	$ot_2$ o	<b>ว</b> <i>s</i> 3		
$v_2 = v_1$ (a)						(b)								
0	0	0	ot2	<b>o</b> <i>s</i> <sub>4</sub>	$o^{s_1}$		0	0	0	0	$\mathbf{o} t_2$	0	$\mathbf{o}^{s_4}$	$o^{s_1}$
0	0	0	<b>o</b> <i>t</i> <sub>4</sub>	<b>o</b> <i>t</i> <sub>3</sub>	<b>o</b> <i>s</i> <sub>2</sub>		0	0	0	0	$\mathbf{o} t_4$	0 <i>t</i> 3	$\mathbf{o} s_5$	$o s_2$
0	0	0	0	$ot_1$	<b>o</b> <i>s</i> <sub>3</sub>		0	0	0	0	0	$ot_5$	$\mathbf{o} t_1$	o <i>s</i> <sub>3</sub>
(c)									(d	)				
		0	ο	0	0	0	0	ot <sub>6</sub>	ot <sub>5</sub>	<b>0</b> <i>s</i> <sub>4</sub>	<b>o</b> <i>s</i> <sub>1</sub>			
		0	0	0	0	0	ot <sub>3</sub>	0	<b>o</b> <i>t</i> <sub>4</sub>	<b>o</b> s <sub>5</sub>	o s <sub>2</sub>			
		0	0	0	0	0	<b>o</b> <i>t</i> <sub>2</sub>	$\mathbf{o}^{t_1}$	0	<b>o</b> <i>s</i> <sub>6</sub>	o <i>s</i> 3			
(e)														

LINKEDNESS OF CARTESIAN PRODUCTS OF COMPLETE GRAPHS

FIGURE 1. No feasible linkage problems for  $K^{d_1+1} \times K^{d_2+1}$ ,  $k = \lfloor (d_1 + d_2 + 1)/2 \rfloor$ ,  $d_1 \leq 2$  and  $d_2 > d_1$ . (a) The case  $d_1 = 1$  and  $d_2 > d_1$ . (b) The case  $d_1 = 2$  and  $d_2 = 3$ . (c) The case  $d_1 = 2$  and  $d_2 = 5$ . (d) The case  $d_1 = 2$  and  $d_2 = 7$ . (e) The case  $d_1 = 2$  and  $d_2 = 9$ . Each row of each part (a)-(e) is a complete graph whose edges have not been drawn.

of the path is in X. Let X be a set of vertices in a graph G. Denote by G[X] the subgraph of G induced by X, the subgraph of G that contains all the edges of G with vertices in X. Write G - X for  $G[V(G) \setminus X]$ .

Consider a linkage problem  $Y := \{\{s_1, t_1\}, \ldots, \{s_k, t_k\}\}$  on a set X of 2k vertices in a graph G. Consider a linkage  $\mathcal{L}$  from a subset Z of X to some set Z' disjoint from X and label the vertices of Z' such that the path in  $\mathcal{L}$  with end  $z_i \in Z$  has its other end  $z'_i \in Z'$ . Then the linkage  $\mathcal{L}$  in G induces a linkage problem Y' in  $(G - V(\mathcal{L})) \cup Z'$  where the vertices of  $X \setminus Z$  remain and the vertices of Z have been replaced by the vertices of Z'. Slightly abusing terminology, we also call terminals the vertices of Z'. If the problem Y' is feasible in  $(G - V(\mathcal{L})) \cup Z'$ , so is the problem Y in G.

Since we make heavy use of Menger's theorem (Diestel, 2017, Thm. 3.3.1), we next remind the reader of one of its consequences.

**Theorem 5** (Menger's theorem). Let G be a k-connected graph, and let A and B be two subsets of its vertices, each of cardinality at least k. Then there are k disjoint A - B paths in G.

We fix some notation and terminology for the remaining of the section. Let G denote the graph  $K^{d_1+1} \times K^{d_2+1}$ . We think of  $G = K^{d_1+1} \times K^{d_2+1}$  as a grid with

4

0	0	0	0	0	)
Ò	Ò	:	÷	÷	$\alpha$ rows
ò	ò	0	0	Ó	J
0	0	0	0	0	)
$: B_{\alpha}$	÷	$A_{\alpha}$	÷	÷	$d_1 + 1 - \alpha$ rows
0	0	0	0	0	J
$C_{12}$	2	Ē	212		

FIGURE 2. Depiction of the subgraphs  $B_{\alpha+1}$ ,  $A_{\alpha+1}$ ,  $C_{12}$ , and  $\overline{C}_{12}$  of  $K^{d_1+1} \times K^{d_2+1}$ .

 $d_1 + 1$  rows and  $d_2 + 1$  columns. In this way, the entry in Row *i* and Column *j* can be referred to as G[i, j].

When we write about a row r of subgraph G' of G, we think of r as a subgraph of G' and as the number r so that we can write about the rth row of G' or G; this ambiguity should cause no confusion. An entry in the grid  $K^{d_1+1} \times K^{d_2+1}$  with no terminal is said to be *free*, as is a row or a column of a subgraph of G with no terminal. A row or a column of a subgraph of G with every entry being occupied by a terminal is said to be *full*.

We need the following induced subgraphs of G:

 $C_{ab...z}$ , the subgraph formed by the union of Columns  $a, b, \ldots, z$ ;

 $\overline{C}_{ab...z}$ , the subgraph obtained by removing Columns  $a, b, \ldots, z$ ;

 $R_{ab...z}$ , the subgraph formed by the union of Rows  $a, b, \ldots, z$ ;

 $\bar{R}_{ab...z}$ , the subgraph obtained by removing Rows  $a, b, \ldots, z$ ;

 $A_{\alpha}$ , the induced subgraph of  $\bar{C}_{12}$  obtained by removing its first  $\alpha$  rows; and

 $B_{\alpha}$ , the subgraph of  $C_{12}$  obtained by removing its first  $\alpha$  rows.

For instance,  $\overline{C}_1$  denotes the subgraph of G obtained by removing the first column,  $C_{12}$  the subgraph formed by the first two columns of G, and  $\overline{C}_{12}$  denotes the subgraph obtained by removing the first two columns of G; observe  $\overline{C}_{12}$  is isomorphic to  $K^{d_1+1} \times K^{d_2-1}$ . Figure 2 depicts some of the aforementioned subgraphs of  $K^{d_1+1} \times K^{d_2+1}$ .

The connectivity of  $K^{d_1+1} \times K^{d_2+1}$  is stated below.

**Lemma 6** (Špacapan 2008, Thm. 1). The (vertex)connectivity of  $K^{d_1+1} \times K^{d_2+1}$ is precisely  $d_1 + d_2$ .

We continue fixing further notation. Henceforth let  $k := \lfloor (d_1 + d_2)/2 \rfloor$ . And let X be a subset of 2k vertices of G and let  $Y := \{\{s_1, t_1\}, \ldots, \{s_k, t_k\}\}$  be a pairing of all the vertices in X.

We first settle the simple cases of  $(0, d_2)$  and  $(1, d_2)$  for  $d_2 \ge 0$ .

**Proposition 7** (Base cases). For  $d_2 \ge 0$  the Cartesian products  $K^1 \times K^{d_2+1}$  and  $K^2 \times K^{d_2+1}$  are both  $|(1 + d_2)/2|$ -linked. This statement is best possible.

*Proof.* The lemma is true for the pair  $(0, d_2)$  for each  $d_2 \ge 0$ , since  $K^1 \times K^{d_2+1} = K^{d_2+1}$  and  $K^{d_2+1}$  is  $\lfloor (1+d_2)/2 \rfloor$ -linked. This is best possible.

The graph  $K^2 \times K^{d_2+1}$  is  $(1 + d_2)$ -connected by Lemma 6. Use Menger's theorem (5) to bring the  $1 + d_2$  terminals to the subgraph  $\bar{R}_1$  through a linkage  $\{S_1, \ldots, S_k, T_1, \ldots, T_k\}$  with  $S_i := s_i - \bar{R}_1$  and  $T_i := t_i - \bar{R}_1$  for  $i \in [1, k]$ . Letting  $\{\bar{s}_i\} := V(S_i) \cap V(\bar{R}_1)$  and  $\{\bar{t}_i\} := V(T_i) \cap V(\bar{R}_1)$ , we produce a new linkage problem  $Y' := \{(\bar{s}_1, \bar{t}_1\}, \ldots, (\bar{s}_k, \bar{t}_k)\}$  in  $\bar{R}_1$  whose feasibility implies that of Y in G. To solve Y' link the pairs of Y' in the subgraph  $\bar{R}_1$ , which is isomorphic to  $K^{d_2+1}$ , using the  $\lfloor (1 + d_2)/2 \rfloor$ -linkedness of  $K^{d_2+1}$ . Figure 1(a) shows an infeasible linkage problem with  $\lfloor (2 + d_2)/2 \rfloor$  pairs in the graph  $K^2 \times K^{d_2+1}$ .

In what follows we aim to find a Y-linkage  $\{L_1, \ldots, L_k\}$  in G with  $L_i$  joining the pair  $\{s_i, t_i\}$  of Y for  $i \in [1, k]$ . Our proof is by induction on  $(d_1, d_2)$  with the base cases settled in Proposition 7. If there is a pair of Y, say  $\{s_1, t_1\}$ , lying in some column or row of G, say in Column 1, we send every terminal  $s_i \in C_1$  that is different from  $s_1$  and  $t_1$  and that is not adjacent to  $t_i$  to the subgraph  $\overline{C}_1$ , and apply the induction hypothesis on  $\overline{C}_1$ . Otherwise, we may assume every pair of Y lies in two distinct columns or rows, say the pair  $\{s_1, t_1\}$  lies in  $C_{12}$ ; then we send every terminal  $s_i \in C_{12}$  that is different from  $s_1$  and  $t_1$  and that is not adjacent to  $t_i$  to the subgraph  $\overline{C}_{12}$ , and apply the induction hypothesis to  $\overline{C}_{12}$ . We develop these ideas below.

The definition of k-linkedness gives the following lemma at once; we will use it implicitly hereafter.

**Lemma 8.** Let  $\ell \leq k$ . Let X be a set of  $2\ell$  distinct vertices of a k-linked graph K, let Y be a labelling and pairing of the vertices in X, and let Z be a set of  $2k - 2\ell$  vertices in K such that  $X \cap Z = \emptyset$ . Then there exists a Y-linkage in K that avoids every vertex in Z.

Besides, basic algebraic manipulation yields the following inequality.

**Lemma 9.** If  $x \ge 2$  and  $y \ge 2$  then x(y-1) > x + y - 3.

*Proof.* The inequality simplifies to (x-1)(y-2) > -1.

We are now ready to put together all the elements of the proof of Theorem 2.

Proof of Theorem 2. Let  $k := \lfloor (d_1 + d_2)/2 \rfloor$ . Then  $d_1 + d_2 \ge 2k$ .

Proposition 7 gives the result for the pairs  $(d_1, 0)$ ,  $(0, d_2)$ ,  $(d_1, 1)$ , and  $(1, d_2)$  for each  $d_1, d_2 \ge 0$ . Hence, our bidimensional induction on  $(d_1, d_2)$  can start with the assumption of  $d_1, d_2 \ge 2$ .

We first deal with the case where a pair in Y, say  $\{s_1, t_1\}$ , lies in some column or some row of G, say in Column 1. Case 1. A pair in Y, say  $\{s_1, t_1\}$ , lies in Column 1.

The induction hypothesis ensures that the subgraph  $\overline{C}_1$  is (k-1)-linked. Hence it suffices to show that all the terminals in  $C_1$  other than  $s_1, t_1$  can be moved to  $\overline{C}_1$ via a linkage; Menger's theorem (5) guarantees this.

Let U be the set of terminals in  $C_1$  other than  $s_1$  and  $t_1$ , and let W be the set of terminals in  $\overline{C}_1$ . Then  $|U| + |W| = d_1 + d_2 - 2$ . Besides, the subgraph  $G - (W \cup \{s_1, t_1\})$  is |U|-connected, as G is  $(d_1 + d_2)$ -connected (Lemma 6). In the case of  $d_1, d_2 \ge 2$ , Lemma 9 yields that  $\overline{C}_1$  has more than  $|U \cup W|$  vertices:

$$|\bar{C}_1| = (d_1 + 1)d_2 > d_1 + 1 + d_2 + 1 - 3 > d_1 + d_2 - 2 = |U| + |W|.$$

Menger's theorem (5) applies and gives disjoint  $C_1 - \overline{C}_1$  paths from the terminals in U to |U| free entries in  $\overline{C}_1$ . The (k-1)-linkedness of  $\overline{C}_1$  now settles the case.

By symmetry, we can assume that every pair  $\{s_i, t_i\}$  in Y lies in two different columns or rows and that  $s_i, t_i$  are not adjacent. Without loss of generality, assume that

(\*)  $s_1$  is Column 1 and  $t_1$  is in Column 2 of  $C_{12}$ .

The induction hypothesis also ensures that both  $\overline{C}_{12}$  and  $\overline{R}_{12}$  are (k-1)-linked. We consider two further cases based on the number of terminals in  $C_{12}$  or  $R_{12}$ .

Case 2. The subgraph  $C_{12}$  contains precisely  $d_1 + 2 - \alpha$  terminals, including  $\{s_1, t_1\}$ , where  $0 \leq \alpha \leq d_1$ .

Excluding  $\{s_1, t_1\}$ , there are at most  $d_1$  terminals in  $C_{12}$ , and there are  $d_1 + 1$  internally-disjoint  $s_1 - t_1$  paths in  $C_{12}$  of length at most three: two length-two paths and  $d_1 - 1$  length-three paths. One of these  $s_1 - t_1$  paths, say  $L_1$ , avoids every other terminal in  $C_{12}$ .

Without loss of generality, assume that Row 1 in  $C_{12}$  is part of the path  $L_1$ ; that is,

(\*\*) 
$$\{G[1,1], G[1,2]\} \subseteq V(L_1).$$

It is true that  $(V(L_1) \cap V(B_1)) \subseteq \{s_1, t_1\}.$ 

In the subcase  $\alpha = d_1$ , every pair in  $Y \setminus \{s_1, t_1\}$  is in  $\overline{C}_{12}$ , and the induction hypothesis on  $\overline{C}_{12}$  settles the subcase.

Suppose that  $\alpha = d_1 - 1$ , say  $C_{12}$  contains  $\{s_1, t_1, s_2\}$ . Then  $s_2 \in B_1$  and  $t_2 \in \overline{C}_{12}$ . We may assume  $s_1, s_2$  are in Column 1 and  $t_1$  is in Column 2. We show there is an X-valid  $s_2 - A_1$  path  $L'_2$  such that the vertex  $x \in V(L'_2) \cap V(A_1)$  is either  $t_2$  or a nonterminal.

Through each entry of Column 1 of  $B_1$ , there is a  $s_2 - A_1$  path of length at most two: one of length one and  $d_1-1$  of length two. Moreover,  $d_1-1$  of such paths avoid  $s_1$ . To ensure the existence of  $L'_2$ , it suffices to show that  $A_1$  cannot have  $d_1 - 1$ rows that are full of terminals other than  $t_2$ . According to Lemma 9, the inequality

$$(d_1 - 1)(d_2 - 1) > d_1 - 1 + d_2 - 3 = |X \setminus \{s_1, t_1, s_2, t_2\}|$$

holds for  $d_1, d_2 \ge 2$ . Hence we get the existence of  $L'_2$ . As a result, the solution of the new problem  $Y' := \{\{x, t_2\}, \{s_3, t_3\}, \ldots, \{s_k, t_k\}\}$  in  $\overline{C}_{12}$  induces a solution of the problem Y in G. And the solution of Y' follows from the (k-1)-linkedness of  $\overline{C}_{12}$ .

Henceforth assume that  $\alpha \leq d_1 - 2$ . To finalise Case 2, we require a couple of claims.

**Claim 1.** Suppose that there are at most  $d_1 + 2$  terminals in  $B_1 = K^{d_1} \times K^2$ . Then there is an injection from the set of rows of  $B_1$  that contain two terminals  $x_1, x_2$  such that  $\{x_1, x_2\} \cap \{s_1, t_1\} = \emptyset$  to the set of rows of  $B_1$  that contain either no terminal or a terminal in  $\{s_1, t_1\}$  but no other terminal.

*Proof.* This follows from a simple counting argument. The number of rows in  $B_1$  is  $d_1$ . Let m denote the number of rows of  $B_1$  that contain two terminals  $x_1, x_2$  such that  $\{x_1, x_2\} \cap \{s_1, t_1\} = \emptyset$  and let  $n := |(X \cap V(B_1)) \setminus \{s_1, t_1\}|$ . It follows that the number of rows of  $B_1$  that contains precisely one terminal  $x \notin \{s_1, t_1\}$  is n - 2m; either  $s_1$  or  $t_1$  may be in these rows. As a result, the number of rows of  $B_1$  that contain either no terminal or a terminal in  $\{s_1, t_1\}$  but no other terminal is  $d_1 - m - (n - 2m)$ . Combining  $n \leqslant d_1$  with all these numbers, we get that

$$d_1-m-(n-2m)=d_1-n+m\geqslant d_1-d_1+m=m.$$

The claim is proved.

**Claim 2.** Suppose that there are at most  $d_1 + 2$  terminals in  $B_1 = K^{d_1} \times K^2$ . If every row in the subgraph  $A_1 = K^{d_1} \times K^{d_2-1}$  of  $\overline{C}_{12}$  has a free entry, then, for every terminal  $x \notin \{s_1, t_1\}$  in  $B_1$ , there is a  $B_1 - A_1$  path L from x to a free entry in  $A_1$  such that L is X-valid; and all these X-valid paths are disjoint.

*Proof.* If a row of  $B_1$  contains exactly one terminal  $x \notin \{s_1, t_1\}$ , then send x to a free entry in the same row of  $A_1$ . Let  $x_1$  and  $x_2$  be two terminals in  $B_1$  that satisfy  $\{x_1, x_2\} \cap \{s_1, t_1\} = \emptyset$  and occupy a row  $r_f$  of  $B_1$ . From Claim 1 ensues the existence of a row  $r_e$  of  $B_1$  that contains either no terminal or a terminal in  $\{s_1, t_1\}$ but no other terminal; in short, there is at least a free entry in  $r_e$ .

Consider a pair  $(r_f, r_e)$  of rows granted by Claim 1. Send either  $x_1$  or  $x_2$ , say  $x_1$ , to the free entry in the row  $r_e$  of  $A_1$  passing through the corresponding free entry in the row  $r_e$  of  $B_1$ , and send  $x_2$  to a free entry in the row  $r_f$  of  $A_1$ . The proof of the claim is now complete.

Now suppose that  $\alpha = 0$  or  $2 \leq \alpha \leq d_1 - 2$ . In this subcase, the subgraph  $\bar{C}_{12}$  contains at most  $\alpha$  full rows: if  $\alpha + 1$  rows were full in  $\bar{C}_{12}$  then there would be at least  $(\alpha+1)(d_2-1)$  terminals in  $\bar{C}_{12}$  but  $(\alpha+1)(d_2-1) > d_2-2+\alpha$  (Lemma 9). Even when the path  $L_1$  uses the first row of  $C_{12}$  by (\*\*), there is no loss of generality by assuming that the full rows of  $\bar{C}_{12}$  are among the first  $\alpha + 1$  rows of  $\bar{C}_{12}$ . It follows that every row of  $A_{\alpha+1}$  has a free entry.

We send to  $B_{\alpha+1}$  the terminals other than  $s_1$  and  $t_1$  that are in the rows 2 to  $\alpha + 1$  of  $C_{12}$ , the terminals other than  $s_1$  and  $t_1$  that are in  $B_1 \setminus B_{\alpha+1}$ . For  $\alpha = 0$ ,  $B_1 \setminus B_{\alpha+1} = \emptyset$  and there is nothing to do.

We now focus on the subcase  $2 \leq \alpha \leq d_1 - 2$ . Let  $n_1$  and  $n_2$  denote the number of terminals in  $B_1 \setminus B_{\alpha+1}$  and  $B_{\alpha+1}$ , respectively. Then the following inequalities hold

$$n_1 + n_2 \leqslant d_1 + 2 - \alpha \leqslant d_1 \quad (\text{since } 2 \leqslant \alpha),$$
  
$$n_1 + n_2 \leqslant d_1 + 2 - \alpha \leqslant 2d_1 - 2\alpha = |B_{\alpha+1}| \quad (\text{since } \alpha \leqslant d_1 - 2).$$

Since  $B_1$  is  $d_1$ -connected by Lemma 6, Menger's theorem gives  $n_1$  disjoint paths from the terminals in  $B_1 \setminus B_{\alpha+1}$  to  $n_1$  free entries in  $B_{\alpha+1}$ . These free entries in  $B_{\alpha+1}$  will henceforth play the role of the terminals of  $B_1 \setminus B_{\alpha+1}$ , and so we call them terminals as well.

Let  $d'_1 := d_1 - \alpha$  and  $d'_2 := d_2$ . Then,  $d'_1 \ge 2$ , there are at most  $d'_1 + 2$  terminals in  $B_{\alpha+1} = K^{d'_1} \times K^2$ , and every row in  $A_{\alpha+1} = K^{d'_1} \times K^{d'_2} - 1$  has a free entry. Hence, Claim 2 applies, and there is a linkage formed by X-valid paths from the terminals in  $B_{\alpha+1}$ , other than  $t_1$ , to free entries in  $A_{\alpha+1}$ . Now we have a new linkage problem Y' in  $\overline{C}_{12}$  with at most 2(k-1) pairs. The solution of Y' in  $\overline{C}_{12}$ implies a solution of the linkage problem Y in G. To link the pairs of Y' use the (k-1)-linkedness of  $\overline{C}_{12}$ .

Finally assume that  $\alpha = 1$ . In a first scenario suppose that either both entries in  $B_1 \setminus B_2$  are nonterminals or each terminal in  $B_1 \setminus B_2$  is adjacent to a nonterminal in  $B_2$ . Then we can send the terminals in  $B_1 \setminus B_2$  to  $B_2$ . In the second scenario suppose that there is a terminal  $s_i$  in  $B_1 \setminus B_2$  whose neighbours in  $B_2$  are all terminals. Then the column of  $s_1$  in  $B_1$  would contain exactly  $d_1$  terminals, including  $s_i$ . We send  $s_i$  to a free entry in  $A_1$ , in the same row as  $s_i$  (the first row of  $A_1$ ): if this free entry didn't exist, then  $s_i$  would be adjacent to the  $d_2 - 1$  terminals in  $A_1$  and the  $d_1 - 1$  terminals in  $B_2$ . Since there are  $d_1 + d_2$  terminals in total, it would follow that  $s_i$  is adjacent to  $t_i$ . This contradiction shows that we can send  $s_i$  to a free entry in  $A_1$ . This completes the proof of Case 2.

In both scenarios, it remains to send the terminals in  $B_2$  to  $A_2$ . To do so, we reason as in the subcase  $2 \leq \alpha \leq d_1 - 2$ . Let  $d'_1 := d_1 - 1$  and  $d'_2 := d_2$ . It follows that  $d'_1 \geq 2$ , that there are at most  $d_1 + 2'$  terminals in  $B_2$ , and that every row in  $A_2$  has a free entry. Claim 2 applies again and gives a linkage formed by X-valid paths from the terminals in  $B_2$ , other than  $s_1, t_1$ , to free entries in  $A_2$ .

With all the terminals in  $\overline{C}_{12}$ , therein we have a new linkage problem Y' with at most 2(k-1) pairs whose solution in  $\overline{C}_{12}$  implies a solution of the linkage problem Y in G. To solve Y' in  $\overline{C}_{12}$  use the (k-1)-linkedness of  $\overline{C}_{12}$ .

By symmetry, we also have the result if there are at most  $d_2 + 2$  terminals in  $R_{12}$ , including  $\{s_1, t_1\}$ .

Case 3. The subgraph  $C_{12}$  contains at least  $d_1 + 3$  terminals, including  $\{s_1, t_1\}$ .

This case reduces to the previous case. If  $C_{12}$  contains at least  $d_1 + 3$  terminals then  $R_{12}$  contains at most  $d_2 - 3 + 4 = d_2 + 1$  terminals, since there are four entries shared by  $C_{12}$  and  $R_{12}$ . Since we make no distinction between columns and rows, this case is already covered. This completes the proof of the theorem.

#### 3. Duals of cyclic polytopes

There is a close connection between duals of cyclic *d*-polytopes with d+2 vertices and Cartesian products of complete graphs.

The moment curve in  $\mathbb{R}^d$  is defined by  $x(t) := (t, t^2, \ldots, t^d)$  for  $t \in \mathbb{R}$ , and the convex hull of any n > d points on it gives a *cyclic polytope* C(n, d). The *combinatorics* of a cyclic polytope, the face lattice of the polytope faces partially ordered by inclusion, is independent of the points chosen on the moment curve. Hence we talk of the cyclic *d*-polytope on *n* vertices (Ziegler, 1995, Example 0.6).

For a polytope P that contains the origin in its interior, the *dual polytope*  $P^*$  is defined as

$$P^* = \{ y \in \mathbb{R}^d \mid x \cdot y \leq 1 \text{ for all } x \text{ in } P \}.$$

If P does not contain the origin, we translate the polytope so that it does. Translating the polytope P changes the geometry of  $P^*$  but not its face lattice. The face lattice of  $P^*$  is the inclusion reversed face lattice of P. In particular, the vertices of  $P^*$  correspond to the facets of P, and the edges of  $P^*$  correspond to the (d-2)faces of P. The *dual graph* of a polytope P is the graph of the dual polytope, or equivalently, the graph on the set of facets of P where two facets are adjacent in the dual graph if they share a (d-2)-face.

Duals of cyclic *d*-polytopes are simple *d*-polytopes. It is also the case that the dual of a cyclic *d*-polytope with d + 2 vertices can be expressed as  $T(\lfloor d/2 \rfloor) \times T(\lceil d/2 \rceil)$  (Ziegler 1995, Ex. 0.6). From this observation and Theorem 2 the next corollary follows at once.

**Corollary 10.** Duals of cyclic polytopes with d + 2 vertices are  $\lfloor d/2 \rfloor$ -linked for every  $d \ge 2$ .

### 4. Acknowledgments

Guillermo would like to thank the hospitality of Leif Jørgensen, and the Department of Mathematical Sciences at Aalborg University where this research started.

Julien Ugon's research was supported by the ARC discovery project DP180100602.

#### References

- Diestel, R. (2017). *Graph Theory*, volume 173 of *Graduate Texts in Mathematics*. Berlin: Springer-Verlag, 5th edition.
- Gallivan, S. (1985). Disjoint edge paths between given vertices of a convex polytope. J. Combin. Theory Ser. A, 39(1), 112–115.

- Goodman, J. E. & O'Rourke, J., Eds. (1997). Handbook of discrete and computational geometry. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 1st edition.
- Mészáros, G. (2016). On linkedness in the Cartesian product of graphs. Period. Math. Hungar., 72(2), 130–138.
- Špacapan, S. (2008). Connectivity of cartesian products of graphs. Applied Mathematics Letters, 21(7), 682–685.
- Ziegler, G. M. (1995). Lectures on polytopes, volume 152 of Graduate Texts in Mathematics. New York: Springer-Verlag.