

THE PRODUCT ON \mathcal{W} -SPACES OF RATIONAL FORMS

A. ZUEVSKY

ABSTRACT. Let V be a quasi-conformal grading-restricted vertex algebra, \mathcal{W} be its module, and $\mathcal{W}_{z_1, \dots, z_n}$ be the space of rational differential forms with complex parameters (z_1, \dots, z_n) for $n \geq 0$. Using geometric interpretation in terms of two Riemann spheres sewing we define a product of elements of two spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$, and study its properties. A product is introduced also for elements of two spaces $C_m^k(V, \mathcal{W}) \times C_{m'}^n(V, \mathcal{W}) \rightarrow C_{m+m'}^{k+n}(V, \mathcal{W})$ of the corresponding chain complex of rational differential forms invariant with respect to transformations of complex parameters.

1. CONFLICT OF INTEREST

The author states that:

- 1.) The paper does not contain any potential conflicts of interests.

2. DATA AVAILABILITY STATEMENT

The author confirms that:

- 1.) All data generated or analyzed during this study are included in this published article.
- 2.) Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

3. INTRODUCTION

The problem of defining a product on the space $\mathcal{W}_{z_1, \dots, z_n}$ (or \mathcal{W} -spaces) of rational differential forms (and, in particular, on $C_m^n(V, \mathcal{W})$ -spaces of corresponding chain complex) is very important for cohomology theory of vertex algebras, continual Lie algebras, theory of integrable models, as well as for further applications to cohomology of smooth manifolds. A cohomology theory for grading-restricted vertex algebras was introduced in [10]. Vertex algebras, generalizations of ordinary Lie algebras, are essential in conformal field theory [4], and it is a rapidly developing field of studies. Algebraic nature of methods applied in this field helps to understand and compute the structure of vertex algebra characters [1, 3, 4, 11, 24]. On the other hand, the geometric side of a vertex algebra characters is in association of their formal parameters with local coordinates on a complex variety. Depending on geometry of a manifold, one

Key words and phrases. Vertex algebras; Riemann surfaces; product of \mathcal{W} -spaces; chain complexes.

can obtain various consequences for a vertex algebra and its space of characters. Vice-versa, one can study geometric property of a manifold by using algebraic nature of a vertex algebra attached.

For purposes of cohomology invariants construction for vertex algebras, it is important to define products of elements of chain-cochain complex spaces. The natural and well-developed theory of vertex operator algebra correlation functions [7, 8, 12, 15–18, 20, 21] considered on higher genus Riemann surfaces [23] and constructed from lower genus Riemann surfaces proved its convenience for computations, and served as a motivation for our construction of a product of rational forms. In that direction, an extremely difficult question of composability with vertex operators occurs. For the cohomology theory of vertex algebras, one has to assume that the chain-cochains are composable with vertex operators which assumes convergence. Especially, when we want to compute cohomology of a vertex algebra, we have to deal with the convergence problem first. In the case of grading-restricted vertex algebras, the difficulty is that chain-cochains are not represented by vertex or intertwining operators. The techniques for vertex operators or intertwining operators in general do not work. The aim of this paper is to develop such new techniques.

For products of spaces of chain-cochains, we propose to involve the geometric procedure [23] of sewing of Riemann surfaces as auxiliary model spaces in a geometric interpretation of algebraic products of spaces associated to vertex algebras. Similar to various other structures in the theory of vertex operator algebras, this is not a usual associative product. The product that occurs is parametrized by a nonzero complex number ϵ identified to the complex parameter of the sewing procedure involved. More generally, the product is constructed from two Riemann spheres with a collection of marked points with local coordinates vanishing at these points. The same scheme works, for example, for tensor products of modules which are in fact parametrized by such geometric objects. Because of this, the existence of such products involves the convergence. In addition to that, a vertex operator algebra must satisfy some conditions in order for such convergence to hold.

In this paper we introduce the product of elements of \mathcal{W} -spaces of rational differential forms for a grading-restricted vertex algebra. For the construction of complexes (cf. Section 7) we make use of maps from tensor powers of V to the space $\mathcal{W}_{z_1, \dots, z_n}$ to define cochains in vertex algebra cohomology theory. For that purpose, in particular, to define coboundary operators, we have to compose chain-cochains with vertex operators. However, the images of vertex operator maps in general do not belong to algebras or their modules. Such objects belong to corresponding algebraic completions which constitute one of the most subtle features of the theory of vertex algebras. Because of this, we might not be able to compose vertex operators directly. In order to overcome this problem, one first writes a series by projecting an element of the algebraic completion of an algebra or a module to its homogeneous components. Then we compose these homogeneous components with vertex operators, and take formal sums. If such formal sums are absolutely convergent, then these operators can be composed and can be used in constructions.

The plan of the paper is the following. In Section 4 we recall the definition of the space of \mathcal{W} -valued rational forms for grading-restricted vertex algebras, and remind

properties of their elements. In Section 5 we introduce a product for elements of two $\mathcal{W}_{z_1, \dots, z_n}$ -spaces. In Section 6 we study properties of the resulting product. In Section 7 we recall the definition and properties of spaces $C_m^n(V, \mathcal{W})$ of the chain-cochain complex for a grading-restricted vertex algebra. In Section 8 we define the product for elements of $C_m^n(V, \mathcal{W})$ -spaces and study its properties. In Section 9 we consider the particular case of a short exceptional complex associated to certain $C_m^n(V, \mathcal{W})$ subspaces. In Subsection 9.1 we provide a specific example of application of our product to cohomology invariants for vertex algebras. In Appendixes we provide the material needed for the construction of the product for \mathcal{W} -spaces. In Appendix 10 we recall the notion of a quasi-conformal grading-restricted vertex algebra. In Appendix 11 we describe the geometric procedure of forming a Riemann sphere by sewing two initial Riemann spheres. Finally, Appendix 12 contains a proof of Proposition 10.

4. SPACES OF \mathcal{W} -VALUED RATIONAL FORMS

4.1. The space \mathcal{W} of rational forms. Part of notions and notations in this Subsection originates from [10]. We define the configuration spaces:

$$F_n \mathbb{C} = \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid z_i \neq z_j, i \neq j\},$$

for $n \in \mathbb{Z}_+$. Recall definitions and properties given in Appendix 10. Let V be a grading-restricted vertex algebra, and W a grading-restricted generalized V -module. By \overline{W} we denote the algebraic completion of W ,

$$\overline{W} = \prod_{n \in \mathbb{C}} W_{(n)} = (W')^*.$$

Let $w' \in W'$ be an arbitrary element of W' dual to W with respect to the canonical pairing $\langle \cdot, \cdot \rangle$ with the dual space of W .

Definition 1. A \overline{W} -valued rational function f in (z_1, \dots, z_n) with the only possible poles at $z_i = z_j, i \neq j$, is a map

$$\begin{aligned} f : F_n \mathbb{C} &\rightarrow \overline{W}, \\ (z_1, \dots, z_n) &\mapsto f(z_1, \dots, z_n), \end{aligned}$$

such that for any $w' \in W'$,

$$R(z_1, \dots, z_n) = R(\langle w', f(z_1, \dots, z_n) \rangle), \quad (4.1)$$

is a rational function in (z_1, \dots, z_n) with the only possible poles at $z_i = z_j, i \neq j$. In this paper, such a map is called \overline{W} -valued rational function in (z_1, \dots, z_n) with possible other poles. The space of \overline{W} -valued rational functions is denoted by $\overline{W}_{z_1, \dots, z_n}$.

Here $R(\cdot)$ denotes the following. Namely, if a meromorphic function $f(z_1, \dots, z_n)$ on a region in \mathbb{C}^n can be analytically extended to a rational function in (z_1, \dots, z_n) , then the notation $R(f(z_1, \dots, z_n))$ is used to denote such rational function. Note that a set of a grading-restricted vertex algebra elements (v_1, \dots, v_n) associated with the corresponding (z_1, \dots, z_n) plays the role of non-commutative parameters for a function f in (4.1). Recall (Appendix 10) the definition of a quasi-conformal grading-restricted vertex algebra V . Let us introduce the definition of a $\mathcal{W}_{z_1, \dots, z_n}$ -space:

Definition 2. We define the space $\mathcal{W}_{z_1, \dots, z_n}$ of $\overline{W}_{z_1, \dots, z_n}$ -valued rational forms

$$\begin{aligned} \mathcal{F}(v_1, z_1; \dots; v_n, z_n) \\ = \Phi \left(dz_1^{\text{wt}(v_1)} \otimes v_1, z_1; \dots; dz_n^{\text{wt}(v_n)} \otimes v_n, z_n \right), \end{aligned} \quad (4.2)$$

formed by all \overline{W} -valued rational functions Φ with each vertex algebra element entry v_i , $1 \leq i \leq n$ of a grading-restricted quasi-conformal vertex algebra V tensored with power $\text{wt}(v_i)$ -differential of the corresponding formal parameter z_i . We call \mathcal{W} the space of $\mathcal{W}_{z_1, \dots, z_n}$ for $n = 0$.

Recall definitions and properties of Subsection 10.2 of Appendix 10. Let us denote $\mathcal{O}^{(n)}$ the space of formal Taylor series in n variables. In Appendix 12 we give a proof of the following

Proposition 1. *For primary vectors of a quasi-conformal grading-restricted vertex algebra V , the form (4.2) is invariant with respect to elements*

$$(\rho_1(z_1, \dots, z_n), \dots, \rho_n(z_1, \dots, z_n)),$$

of the group $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$, i.e., under the changes

$$z_i \mapsto z'_i = \rho_i(z_1, \dots, z_n),$$

of formal parameters (z_1, \dots, z_n) .

4.2. Properties of rational functions for \mathcal{W} -valued elements. Let V be a grading-restricted vertex algebra and W a grading-restricted generalized V -module (cf. Appendix 10). Let us give here modifications of definitions and facts about matrix elements for a grading-restricted vertex algebra.

Definition 3. For $n \in \mathbb{Z}_+$, a map

$$\mathcal{F}(v_1, z_1; \dots; v_n, z_n) \in V^{\otimes n} \rightarrow \mathcal{W}_{z_1, \dots, z_n},$$

is said to have the $L_V(-1)$ -derivative property if

$$(i) \quad \partial_{z_i} \mathcal{F}(v_1, z_1; \dots; v_n, z_n) = \mathcal{F}(v_1, z_1; \dots; L_V(-1)v_i, z_i; \dots; v_n, z_n), \quad (4.3)$$

for $i = 1, \dots, n$, $(v_1, \dots, v_n) \in V$, $w' \in W'$, and

$$(ii) \quad \sum_{i=1}^n \partial_{z_i} \mathcal{F}(v_1, z_1; \dots; v_n, z_n) = L_W(-1) \cdot \mathcal{F}(v_1, z_1; \dots; v_n, z_n). \quad (4.4)$$

In [10] we find the following

Proposition 2. *Let \mathcal{F} be a map having the $L_W(-1)$ -derivative property. Then for $(v_1, \dots, v_n) \in V$, $(z_1, \dots, z_n) \in F_n \mathbb{C}$, $z \in \mathbb{C}$ such that $(z_1 + z, \dots, z_n + z) \in F_n \mathbb{C}$,*

$$e^{zL_W(-1)} \mathcal{F}(v_1, z_1; \dots; v_n, z_n) = \mathcal{F}(v_1, z_1 + z; \dots; v_n, z_n + z), \quad (4.5)$$

and $1 \leq i \leq n$ such that

$$(z_1, \dots, z_{i-1}, z_i + z, z_{i+1}, \dots, z_n) \in F_n \mathbb{C},$$

the power series expansion of

$$\mathcal{F}(v_1, z_1; \dots; v_{i-1}, z_{i-1}; v_i, z_i + z; v_{i+1}, z_{i+1}; \dots; v_n, z_n), \quad (4.6)$$

in z is equal to the power series

$$\mathcal{F} \left(v_1 z_1; \dots; v_{i-1}, z_{i-1}; e^{zL_V(-1)} v_i, z_i; v_{i+1}, z_{i+1}; \dots; v_n, z_n \right), \quad (4.7)$$

in z . In particular, the power series (4.7) in z is absolutely convergent to (4.6) in the disk $|z| < \min_{i \neq j} \{|z_i - z_j|\}$.

One states

Definition 4. A map

$$\mathcal{F} : V^{\otimes n} \rightarrow \mathcal{W}_{z_1, \dots, z_n}$$

has the $L_W(0)$ -conjugation property if for $(v_1, \dots, v_n) \in V$, $(z_1, \dots, z_n) \in F_n \mathbb{C}$, and $z \in \mathbb{C}^\times$, such that $(zz_1, \dots, zz_n) \in F_n \mathbb{C}$,

$$z^{L_W(0)} \mathcal{F}(v_1, z_1; \dots; v_n, z_n) = \mathcal{F}(z^{L_V(0)} v_1, zz_1; \dots; z^{L_V(0)} v_n, zz_n). \quad (4.8)$$

Let S_n be the permutation group. One defines the action of S_n on the space $\text{Hom}(V^{\otimes n}, \mathcal{W}_{z_1, \dots, z_n})$ of maps from $V^{\otimes n}$ to $\mathcal{W}_{z_1, \dots, z_n}$ by

$$\sigma(\mathcal{F})(v_1, z_1; \dots; v_n, z_n) = \mathcal{F}(v_{\sigma(1)}, z_{\sigma(1)}; \dots; v_{\sigma(n)}, z_{\sigma(n)}), \quad (4.9)$$

for $\sigma \in S_n$, and $(v_1, \dots, v_n) \in V$. We will use the notation $\sigma_{i_1, \dots, i_n} \in S_n$, to denote the permutation given by $\sigma_{i_1, \dots, i_n}(j) = i_j$, for $j = 1, \dots, n$.

Finally, the following result was proved in [3]:

Proposition 3. For $(v_1, \dots, v_n) \in V$, $w \in W$ and $w' \in W'$,

$$\langle w', Y_W(v_1, z_1) \dots Y_W(v_n, z_n) w \rangle,$$

is absolutely convergent in the region $|z_1| > \dots > |z_n| > 0$ to a rational function

$$R(\langle w', Y_W(v_1, z_1) \dots Y_W(v_n, z_n) w \rangle),$$

in (z_1, \dots, z_n) with the only possible poles at $z_i = z_j$, $i \neq j$, and $z_i = 0$. The following commutativity holds: for $\sigma \in S_n$,

$$\begin{aligned} R(\langle w', Y_W(v_1, z_1) \dots Y_W(v_n, z_n) w \rangle) \\ = R(\langle w', Y_W(v_{\sigma(1)}, z_{\sigma(1)}) \dots Y_W(v_{\sigma(n)}, z_{\sigma(n)}) w \rangle). \end{aligned}$$

5. PRODUCT OF SPACES OF \mathcal{W} -VALUED FORMS

5.1. Motivation and geometric interpretation. The structure of $\mathcal{W}_{z_1, \dots, z_n}$ -spaces is quite complicated and it is difficult to introduce algebraically a product of its elements. In order to define an appropriate product of two elements of $\mathcal{W}_{z_1, \dots, z_n}$ -spaces we first have to interpret them geometrically. Basically, a $\mathcal{W}_{z_1, \dots, z_n}$ -space must be associated with a certain model space, the algebraic \mathcal{W} -language should be transferred to a geometric one, two model spaces should be "related" appropriately, and, finally, a product should be defined.

For two $\mathcal{W}_{x_1, \dots, x_k}$ - and $\mathcal{W}_{y_1, \dots, y_n}$ -spaces we first associate formal complex parameters in the sets (x_1, \dots, x_k) and (y_1, \dots, y_n) to parameters of two auxiliary spaces. Then we describe a geometric procedure to form a resulting model space by combining two original model spaces. Formal parameters of $\mathcal{W}_{z_1, \dots, z_{k+n}}$ should be then identified with parameters of the resulting space.

Note that according to our assumption, $(x_1, \dots, x_k) \in F_k\mathbb{C}$, and $(y_1, \dots, y_n) \in F_n\mathbb{C}$. As it follows from definition of the configuration space $F_n\mathbb{C}$ given in Subsection 4.1, in the case of coincidence of two formal parameters they are excluded from $F_n\mathbb{C}$. In general, it may happen that some number r of formal parameters of $\mathcal{W}_{x_1, \dots, x_k}$ coincides with some r formal parameters of $\mathcal{W}_{y_1, \dots, y_n}$. Thus, we require that the set of formal parameters (z_1, \dots, z_{k+n-r}) for the resulting model space would belong to $F_{k+n-r}\mathbb{C}$. This leads to the fall off of the total number of formal parameters for the resulting model space $\mathcal{W}_{z_1, \dots, z_{k+n-r}}$. In what follows we consider the case when all formal parameters (x_1, \dots, x_k) differ from formal parameters of (y_1, \dots, y_n) . This singular case can then be treated similar to the ordinary one in lower dimension.

5.2. Definition of the product for \mathcal{W} -valued rational forms. Recall the definition (10.15) of the intertwining operator Y_{WV}^W given in Appendix 10. We then formulate

Definition 5. For a quasi-conformal module W of a grading-restricted vertex algebra V , a set of quasi-primary V -elements (v_1, \dots, v_n) , $(v'_1, \dots, v'_n) \in V$, $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in \mathcal{W}_{x_1, \dots, x_k}$, $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in \mathcal{W}_{y_1, \dots, y_n}$, introduce the ϵ -product for $\epsilon = \zeta_1 \zeta_2$, for $|\zeta_a| > 0$, $a = 1, 2$,

$$\cdot_\epsilon : \mathcal{W}_{x_1, \dots, x_k} \times \mathcal{W}_{y_1, \dots, y_n} \rightarrow \mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}, \quad (5.1)$$

for $(x_1, \dots, x_k; y_1, \dots, y_n) \in F_{k+n}\mathbb{C}$. For arbitrary $w' \in W'$, the product is associated to the form

$$\begin{aligned} & \mathcal{R}(x_1, \dots, x_k; y_1, \dots, y_n; \epsilon, \zeta_1, \zeta_2) \\ &= \sum_{l \in \mathbb{Z}} \epsilon^l \sum_{u \in V_l} \langle w', Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\ & \quad \langle w', Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle, \end{aligned} \quad (5.2)$$

via (4.1), parametrized by $\zeta_a \in \mathbb{C}$, $|\zeta_a| > 0$, $a = 1, 2$, for all pairs of matrix elements $\langle w', Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle$, $\langle w', Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle$ such that (5.2) converges in ϵ . The sum is taken over any V_l -basis $\{u\}$, where \bar{u} is the dual of u with respect to the canonical pairing $\langle \cdot, \cdot \rangle_\lambda$ (10.28) with the dual space of V , (see Appendix 10).

By the standard reasoning [3, 24], (5.2) does not depend on the choice of a basis of $u \in V_l$, $l \in \mathbb{Z}$. In the case when multiplied forms \mathcal{F} do not contain V -elements, i.e., for $\Phi, \Psi \in \mathcal{W}$, (5.2) defines the product $\Phi \cdot_\epsilon \Psi$ associated to a rational function:

$$\mathcal{R}(\epsilon) = \sum_{l \in \mathbb{Z}} \epsilon^l \sum_{u \in V_l} \langle w', Y_{WV}^W(\Phi, \zeta_1) u \rangle \langle w', Y_{WV}^W(\Psi, \zeta_2) \bar{u} \rangle, \quad (5.3)$$

which defines $\mathcal{F}(\epsilon) \in \mathcal{W}$ via $\mathcal{R}(\epsilon) = \langle w', \mathcal{F}(\epsilon) \rangle$. As we will see in Section 7, Definition 5 is also supported by Proposition (8).

Remark 1. Note that due to (10.15), in Definition 5, and in (5.2) in particular, it is assumed that $\mathcal{F}(v_1, x_1; \dots; v_k, x_k)$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$ are composable with the V -module W vertex operators $Y_W(u, -\zeta_1)$ and $Y_W(\bar{u}, -\zeta_2)$ correspondingly (see Section 7 for the definition of composability). The product (5.2) is actually defined by

sum of products of matrix elements of ordinary V -module W vertex operators acting on $\mathcal{W}_{z_1, \dots, z_n}$ elements. In what follows we will see that, since $u \in V$ and $\bar{u} \in V'$ are connected by (10.29), ζ_1 and ζ_2 appear in a relation to each other. The form of the product defined above is natural in terms of the theory of characters for vertex operator algebras in conformal field theory [4, 15–18, 20–22, 24] and many others.

5.3. Existence of the corresponding rational form for the ϵ -product. In order to describe the ϵ product of elements of two spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$ of rational \mathcal{W} -valued forms, we use a geometric interpretation [11, 23]. Recall that a $\mathcal{W}_{z_1, \dots, z_n}$ -space is defined by means of matrix elements of the form (4.1). For a vertex algebra V , this corresponds [3] to a matrix element of a number of V -vertex operators with formal parameters identified with local coordinates on a Riemann sphere. Geometrically, each space $\mathcal{W}_{z_1, \dots, z_n}$ can be also associated to a Riemann sphere with a few marked points with local coordinates vanishing at these points [11]. An extra point can be associated to a center of an annulus used in order to sew the sphere with another sphere. The product (5.2) has then a geometric interpretation. The resulting model space would also be associated to a Riemann sphere formed as a result of sewing procedure. In Appendix 11 we describe explicitly the geometric procedure of sewing of two spheres [23].

Let us identify (as in [1, 4, 11, 22–24]) two sets (x_1, \dots, x_k) and (y_1, \dots, y_n) of complex formal parameters, with local coordinates of two sets of points on the first and the second Riemann spheres correspondingly. Identify complex parameters ζ_1, ζ_2 of (5.2) with coordinates (11.1) of the annuluses (11.3). After identification of annuluses \mathcal{A}_a and $\mathcal{A}_{\bar{a}}$, r coinciding coordinates may occur. This takes into account case of coinciding formal parameters.

As we will see in the next Subsection, the product is defined by a sum of products of matrix elements [3] associated to each of two spheres. Such sum is supposed to describe a \mathcal{W} -valued rational differential form defined on the sphere formed as a result of geometric sewing [23] of two initial spheres. Since two initial spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$ are defined through rational-valued forms with the conditions of absolute convergence expressed by matrix elements of the form (4.1), it is then showed (Proposition 4), that the resulting product defines a $\mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$ -valued rational form by means of a convergent matrix element on the resulting sphere. In what follows we prove the existence of such rational form. The complex sewing parameter, parameterizing the module space of the sewing spheres, parametrizes also the product of the corresponding \mathcal{W} -spaces.

In this Subsection and the next Section we formulate the results of this paper for the ϵ -product of \mathcal{W} -spaces.

Proposition 4. *The product (5.2) of elements of the spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$ corresponds to a rational form with only possible poles at $x_i = x_j$, $y_{i'} = y_{j'}$, and $x_i = y_{j'}$, $1 \leq i, i' \leq k$, $1 \leq j, j' \leq n$.*

Proof. In order to prove this proposition we use the geometric interpretation of the product (5.2) in terms of the Riemann spheres with marked points (see Appendix 11). We consider two sets of vertex algebra elements (v_1, \dots, v_k) and (v'_1, \dots, v'_k) ,

and two sets of formal complex parameters $(x_1, \dots, x_k), (y_1, \dots, y_n)$. Formal parameters are identified with local coordinates of k points on the Riemann sphere $\widehat{\Sigma}_1^{(0)}$, and n points on $\widehat{\Sigma}_2^{(0)}$, with excised annuluses \mathcal{A}_a (see definitions and notations in Appendix 11). The existence of an appropriate $\mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$ -valued rational form is clear from Definition 5 corresponding to the absolute convergent rational form $\mathcal{R}(x_1, \dots, x_k; y_1, \dots, y_n; \epsilon)$ defining the ϵ -product of elements of the spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$. For all choices of elements of the spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$ there exists an element $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \in \mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$ such that the product (5.2) is $R(x_1, \dots, x_k; y_1, \dots, y_n; \epsilon) = \langle w', \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle$. \square

As we see, the ϵ -product is parametrized by a non-zero complex parameter ϵ , and a collection of points on auxiliary spheres with formal parameters vanishing at these points. We then have

Definition 6. Let W be a quasi-conformal module for a grading restricted vertex algebra V . For fixed sets $(v_1, \dots, v_k), (v'_1, \dots, v'_n) \in V$, $(x_1, \dots, x_k) \in \mathbb{C}$, $(y_1, \dots, y_n) \in \mathbb{C}$, we call the set of all $\mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$ -valued rational forms $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon)$ defined by (5.2) with the parameter ϵ exhausting all possible values, the complete product of the spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$.

6. PROPERTIES OF THE \mathcal{W} -PRODUCT

In this Section we study properties of the product $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon)$ of (5.2). Since we assume that $(x_1, \dots, x_k; y_1, \dots, y_n) \in F_{k+n}\mathbb{C}$, i.e., coincidences of x_i and y_j are excluded by the definition of $F_{k+n}\mathbb{C}$. We have

Definition 7. We define the action of $\partial_l = \partial_{z_l} = \partial/\partial z_l$, $1 \leq l \leq k+n$, the differentiation of $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon)$ with respect to the l -th entry of $(x_1, \dots, x_k; y_1, \dots, y_n)$ as follows

$$\begin{aligned} & \langle w', \partial_l \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\ &= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', \partial_{x_i}^{\delta_{l,i}} Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\ & \quad \langle w', \partial_{y_j}^{\delta_{l,j}} Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle. \end{aligned} \quad (6.1)$$

Proposition 5. *The product (5.2) satisfies the $L_V(-1)$ -derivative (4.3) and $L_V(0)$ -conjugation (4.8) properties.*

Proof. By using (4.3) for $\mathcal{F}(v_1, x_1; \dots; v_k, x_k)$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$, we consider

$$\begin{aligned}
& \langle w', \partial_l \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', \partial_{x_i}^{\delta_{i,i}} Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', \partial_{y_j}^{\delta_{i,j}} Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', \partial_{x_i}^{\delta_{i,i}} Y_W(u, -\zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle u \rangle \\
&\quad \langle w', \partial_{y_j}^{\delta_{i,j}} Y_W(\bar{u}, -\zeta_2) \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', Y_{WV}^W(\partial_{x_i}^{\delta_{i,i}} \mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', Y_{WV}^W(\partial_{y_j}^{\delta_{i,j}} \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; (L_V(-1))^{\delta_{i,i}} v_i, x_i; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; (L_V(-1))^{\delta_{i,j}} v'_j, y_j; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&= \langle w', \mathcal{F}(v_1, x_1; \dots; (L_V(-1))_l; \dots; v'_n, y_n; \epsilon) \rangle, \tag{6.2}
\end{aligned}$$

where $(L_V(-1))_l$ acts on the l -th entry of $(v_1, \dots; v_k; v'_1, \dots; v'_n)$. Summing over l we obtain

$$\begin{aligned}
& \sum_{l=1}^{k+n} \partial_l \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\
&= \sum_{l=1}^{k+n} \langle w', \mathcal{F}(v'_1, x_1; \dots; (L_V(-1))_l; \dots; v'_n, y_n; \epsilon) \rangle \\
&= \langle w', L_W(-1) \cdot \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle. \tag{6.3}
\end{aligned}$$

Due to (4.8), (10.5), (10.29), (10.30), and (10.13), we have

$$\begin{aligned}
& \langle w', \mathcal{F}(z^{L_V(0)} v_1, z x_1; \dots; z^{L_V(0)} v_k, z x_k; z^{L_V(0)} v'_1, z y_1; \dots; z^{L_V(0)} v'_n, z y_n; \epsilon) \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', Y_{WV}^W(\mathcal{F}(z^{L_V(0)} v_1, z x_1; \dots; z^{L_V(0)} v_k, z x_k), \zeta_1) u \rangle \\
&\quad \langle w', Y_{WV}^W(\mathcal{F}(z^{L_V(0)} v'_1, z y_1; \dots; z^{L_V(0)} v'_n, z y_n), \zeta_2) \bar{u} \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', Y_{WV}^W(z^{L_V(0)} \mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', Y_{WV}^W(z^{L_V(0)} \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle
\end{aligned}$$

$$\begin{aligned}
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) z^{L_V(0)} \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
&\quad \langle w', e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) z^{L_V(0)} \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', e^{\zeta_1 L_W(-1)} z^{L_V(0)} Y_W(z^{-L_V(0)} u, -z \zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
&\quad \langle w', e^{\zeta_2 L_W(-1)} z^{L_W(0)} Y_W(z^{-L_V(0)} \bar{u}, -z \zeta_2) \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', e^{\zeta_1 L_W(-1)} z^{L_W(0)} z^{-wtu} Y_W(u, -z \zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
&\quad \langle w', e^{\zeta_2 L_W(-1)} z^{L_W(0)} z^{-wt\bar{u}} Y_W(\bar{u}, -z \zeta_2) \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', z^{L_W(0)} e^{\zeta_1 L_W(-1)} Y_W(u, -z \zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
&\quad \langle w', z^{L_W(0)} e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -z \zeta_2) \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', z^{L_W(0)} Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), z \zeta_1) u \rangle \\
&\quad \langle w', z^{L_W(0)} Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), z \zeta_2) \bar{u} \rangle \\
&= \sum_{m \in \mathbb{Z}} \epsilon^m \sum_{u \in V_m} \langle w', z^{L_W(0)} Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta'_1) u \rangle \\
&\quad \langle w', z^{L_W(0)} Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta'_2) \bar{u} \rangle \\
&= \langle w', (z^{L_W(0)}) \cdot \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle.
\end{aligned}$$

With (11.4), we obtain (4.8) for (5.2). \square

Remark 2. As we see in the last expressions, the $L_V(0)$ -conjugation property (4.8) for the product (5.2) includes the action of $z^{L_V(0)}$ -operator on complex parameters ζ_a , $a = 1, 2$.

We also have

Proposition 6. *For primary elements $v_i, v'_j \in V$, $1 \leq i \leq k$, $1 \leq j \leq n$, of a quasi-conformal grading-restricted vertex algebra V and its grading-restricted generalized module W , the product (5.2) is canonical with respect to the action of the group $\text{Aut}_{x_1, \dots, x_k; y_1, \dots, y_n} \mathcal{O}^{(k+n)}$ of $k+n$ -dimensional changes*

$$\begin{aligned}
&(x_1, \dots, x_k; y_1, \dots, y_n) \mapsto (x'_1, \dots, x'_k; y'_1, \dots, y'_n) \\
&= (\rho_1(x_1, \dots, x_k; y_1, \dots, y_n), \dots, \rho_{k+n}(x_1, \dots, x_k; y_1, \dots, y_n)), \quad (6.4)
\end{aligned}$$

of formal parameters.

Proof. Note that due to Proposition 1

$$\begin{aligned}\mathcal{F}(v_1, x'_1; \dots; v_k, x'_k) &= \mathcal{F}(v_1, x_1; \dots; v_k, x_k), \\ \mathcal{F}(v_1, y'_1; \dots; v_n, y'_n) &= \mathcal{F}(v_1, y_1; \dots; v_n, y_n).\end{aligned}$$

Thus,

$$\begin{aligned}& \langle w', \mathcal{F}(v_1, x'_1; \dots; v_k, x'_k; v'_1, y'_1; \dots; v'_n, y'_n; \epsilon) \rangle \\ &= \sum_{l \in \mathbb{Z}} \epsilon^l \sum_{u \in V_l} \langle w', Y_{WV}^W(\mathcal{F}(v_1, x'_1; \dots; v_k, x'_k), \zeta_1) u \rangle \\ & \quad \langle w', Y_{WV}^W(\mathcal{F}(v'_1, y'_1; \dots; v'_n, y'_n), \zeta_2) \bar{u} \rangle \\ &= \sum_{l \in \mathbb{Z}} \epsilon^l \sum_{u \in V_l} \langle w', Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\ & \quad \langle w', Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\ &= \langle w', \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle.\end{aligned}$$

Thus, the product (5.2) is invariant under (6.4). \square

In the geometric interpretation in terms of auxiliary spaces, the definition (5.2) depends on the choice of insertion points p_i , $1 \leq i \leq k$, with local coordinates x_i on $\widehat{\Sigma}_1^{(0)}$, and p'_j , $1 \leq j \leq k$, with local coordinates y_j on $\widehat{\Sigma}_2^{(0)}$. Suppose we change the distribution of points among two Riemann spheres. We formulate the following

Lemma 1. *For a fixed set $(\tilde{v}_1, \dots, \tilde{v}_n) \in V$, of vertex algebra elements, the ϵ -product $\mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_n, z_n; \epsilon) \in \mathcal{W}_{z_1, \dots, z_n}$,*

$$\cdot_\epsilon : \mathcal{W}_{z_1, \dots, z_k} \times \mathcal{W}_{z_{k+1}, \dots, z_n} \rightarrow \mathcal{W}_{z_1, \dots, z_n}, \quad (6.5)$$

remains the same for elements $\mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_k, z_k) \in \mathcal{W}_{z_1, \dots, z_k}$ and $\mathcal{F}(\tilde{v}_{k+1}, z_{k+1}; \dots; \tilde{v}_n, z_n) \in \mathcal{W}_{z_{k+1}, \dots, z_n}$, for any $0 \leq k \leq n$.

Remark 3. This Lemma is important for the formulation of cohomology invariants associated to grading-restricted vertex algebras on smooth manifolds. In case $k = 0$, we obtain from (6.6),

$$\cdot_\epsilon : \mathcal{W} \times \mathcal{W}_{z_1, \dots, z_n} \rightarrow \mathcal{W}_{z_1, \dots, z_n}. \quad (6.6)$$

Proof. Let $\tilde{v}_i \in V$, $1 \leq i \leq k$, $\tilde{v}_j \in V$, $1 \leq j \leq k$, and z_i, z_j are corresponding formal parameters. We show that the ϵ -product of $\mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_k, z_k)$ and $\mathcal{F}(\tilde{v}_{k+1}, z_{k+1}; \dots; \tilde{v}_n, z_n)$, i.e., the $\mathcal{W}_{z_1, \dots, z_{k+n}}$ -valued differential form

$$\mathcal{F}((\tilde{v}_1, z_1; \dots; \tilde{v}_k, z_k); (\tilde{v}_{k+1}, z_{k+1}; \dots; \tilde{v}_n, z_n); \zeta_1, \zeta_2; \epsilon) \quad (6.7)$$

is independent of the choice of $0 \leq k \leq n$. Consider

$$\begin{aligned}& \langle w', \mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_k, z_k; \tilde{v}_{k+1}, z_{k+1}; \dots; \tilde{v}_n, z_n; \zeta_1, \zeta_2; \epsilon) \rangle \\ &= \sum_{l \in \mathbb{Z}} \epsilon^l \sum_{u \in V_l} \langle w', Y_{WV}^W(\mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_k, z_k), \zeta_1) u \rangle \\ & \quad \langle w', Y_{WV}^W(\mathcal{F}(\tilde{v}_{k+1}, z_{k+1}; \dots; \tilde{v}_n, z_n), \zeta_2) \bar{u} \rangle.\end{aligned} \quad (6.8)$$

On the other hand, for $0 \leq m \leq k$, consider

$$\begin{aligned} & \sum_{l \in \mathbb{Z}} \epsilon^l \sum_{u \in V_l} \langle w', Y_{WV}^W(\mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_m, z_m), \zeta_1) u) \\ & \quad \langle w', Y_{WV}^W(\mathcal{F}(\tilde{v}_{m+1}, z'_{m+1}; \dots; \tilde{v}_k, z'_k; \tilde{v}_{k+1}, z_1; \dots; \tilde{v}_n, z_n), \zeta_2) \bar{u} \rangle \\ & = \langle w', \mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_m, z_m; \tilde{v}_{m+1}, z'_{m+1}; \dots; \tilde{v}_k, z'_k; \tilde{v}_{k+1}, z_{k+1}; \dots; \tilde{v}_n, z_n) \rangle. \end{aligned}$$

The last is the ϵ -product (5.2) of $\mathcal{F}(\tilde{v}_1, z_1; \dots; \tilde{v}_m, z_m) \in \mathcal{W}_{z_1, \dots, z_m}$ and $\mathcal{F}(\tilde{v}_{m+1}, z'_{m+1}; \dots; \tilde{v}_k, z'_k; \tilde{v}_{k+1}, z_1; \dots; \tilde{v}_n, z_n) \in \mathcal{W}_{z'_{m+1}, \dots, z'_k; z_1, \dots, z_n}$. Let us apply the invariance with respect to a subgroup of $\text{Aut}_{z_1, \dots, z_{k+n}} \mathcal{O}^{(n)}$, with (z_1, \dots, z_m) and (z_{k+1}, \dots, z_n) remaining unchanged. Then we obtain the same product (6.8). \square

Next, we formulate

Definition 8. We define the action of an element $\sigma \in S_{k+n}$ on the product of $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in \mathcal{W}_{x_1, \dots, x_k}$, and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in \mathcal{W}_{y_1, \dots, y_n}$, as

$$\begin{aligned} & \langle w', \sigma(\mathcal{F})(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\ & = \langle w', \mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}; v'_{\sigma(1)}, y_{\sigma(1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}; \epsilon) \rangle \\ & = \sum_{u \in V} \langle w', Y_{WV}^W(\mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}), \zeta_1) u) \\ & \quad \langle w', Y_{WV}^W(\mathcal{F}(v'_{\sigma(1)}, y_{\sigma(1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}), \zeta_2) \bar{u} \rangle. \end{aligned} \quad (6.9)$$

7. SPACES $C_m^n(V, \mathcal{W})$ OF A COMPLEX

The spaces $C_m^n(V, \mathcal{W})$, $n \geq 0$, $m \geq 0$, and differentials δ_m^n , for the chain-cochain complex $(C_m^n(V, \mathcal{W}), \delta_m^n)$ were introduced in [10, 19]). In this Section we recal the definition and properties of $C_m^n(V, \mathcal{W})$.

7.1. **E -elements.** For $w \in W$, the \overline{W} -valued function given by

$$E_W^{(n)}(v_1, z_1; \dots; v_n, z_n; w) = E(\omega_W(v_1, z_1) \dots \omega_W(v_n, z_n) w),$$

where an element $E(\phi)$ is a \overline{W} -valued rational function,

$$\omega_W(v_i, z_i) = Y_W(dz_i^{\text{wt}(v_i)} \otimes v_i, z_i), \quad (7.1)$$

$\phi \in \overline{W}$ is given by (see notations for $\omega_W(\cdot, \cdot)$ in Section 7.3)

$$E(\phi) = R(\langle w', \phi \rangle).$$

One defines

$$E_{WV}^{W;(n)}(w; v_1, z_1; \dots; v_n, z_n) = E_W^{(n)}(v_1, z_1; \dots; v_n, z_n; w),$$

where $E_{WV}^{W;(n)}(w; v_1, z_1; \dots; v_n, z_n)$ is an element of $\overline{W}_{z_1, \dots, z_n}$. For $(z_1, \dots, z_n, \zeta) \in F_{n+1}\mathbb{C}$, $(v_1, \dots, v_n) \in V$, and $w \in W$, set

$$E_W^{(n,1)}(v_1, z_1; \dots; v_n, z_n; w, \zeta) = E(Y_W(v_1, z_1) \dots Y_W(v_n, z_n) Y_{WV}^W(w, \zeta) \mathbf{1}_V).$$

One defines

$$\mathcal{F} \circ \left(E_{V; \mathbf{1}}^{(l_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(l_n)} \right) : V^{\otimes m+n} \rightarrow \overline{W}_{z_1, \dots, z_{m+n}},$$

by

$$\begin{aligned} & (\mathcal{F} \circ (E_{V; \mathbf{1}}^{(l_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(l_n)}))(v_1 \otimes \dots \otimes v_{m+n-1}) \\ &= E(\mathcal{F}(E_{V; \mathbf{1}}^{(l_1)}(v_1 \otimes \dots \otimes v_{l_1}) \otimes \dots \\ & \quad \otimes E_{V; \mathbf{1}}^{(l_n)}(v_{l_1+\dots+l_{n-1}+1} \otimes \dots \otimes v_{l_1+\dots+l_{n-1}+l_n}))), \end{aligned}$$

and

$$E_W^{(m)} \circ_0 \mathcal{F} : V^{\otimes m+n} \rightarrow \overline{W}_{z_1, \dots, z_{m+n-1}},$$

is given by

$$\begin{aligned} & (E_W^{(m)} \circ_0 \mathcal{F})(v_1 \otimes \dots \otimes v_{m+n}) \\ &= E(E_W^{(m)}(v_1 \otimes \dots \otimes v_m; \mathcal{F}(v_{m+1} \otimes \dots \otimes v_{m+n}))). \end{aligned}$$

Finally,

$$E_{WV}^{W;(m)} \circ_{m+1} \mathcal{F} : V^{\otimes m+n} \rightarrow \overline{W}_{z_1, \dots, z_{m+n-1}},$$

is defined by

$$(E_{WV}^{W;(m)} \circ_{m+1} \mathcal{F})(v_1 \otimes \dots \otimes v_{m+n}) = E(E_{WV}^{W;(m)}(\mathcal{F}(v_1 \otimes \dots \otimes v_n); v_{n+1} \otimes \dots \otimes v_{n+m})).$$

In the case that $l_1 = \dots = l_{i-1} = l_{i+1} = 1$ and $l_i = m - n - 1$, for some $1 \leq i \leq n$, we will use $\mathcal{F} \circ_i E_{V; \mathbf{1}}^{(l_i)}$ to denote $\mathcal{F} \circ (E_{V; \mathbf{1}}^{(l_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(l_n)})$.

7.2. Maps composable with vertex operators. Let us recall the definition of maps composable with a number of vertex operators [10].

Definition 9. For a V -module

$$W = \coprod_{n \in \mathbb{C}} W_{(n)},$$

and $m \in \mathbb{C}$, let

$$P_m : \overline{W} \rightarrow W_{(m)},$$

be the projection from \overline{W} to $W_{(m)}$. Let

$$\mathcal{F} : V^{\otimes n} \rightarrow \mathcal{W}_{z_1, \dots, z_n},$$

be a map. For $m \in \mathbb{N}$, \mathcal{F} is called composable with m vertex operators if the following conditions are satisfied:

1) Let $l_1, \dots, l_n \in \mathbb{Z}_+$ such that $l_1 + \dots + l_n = m + n$, $v_1, \dots, v_{m+n} \in V$, and $w' \in W'$. Set

$$\Psi_i = E_V^{(l_i)}(v_{k_1}, z_{k_1} - \zeta_i; \dots; v_{k_i}, z_{k_i} - \zeta_i; \mathbf{1}_V), \quad (7.2)$$

where

$$k_1 = l_1 + \dots + l_{i-1} + 1, \quad \dots, \quad k_i = l_1 + \dots + l_{i-1} + l_i, \quad (7.3)$$

for $i = 1, \dots, n$. Then there exist positive integers $N_m^n(v_i, v_j)$ depending only on v_i and v_j for $i, j = 1, \dots, k$, $i \neq j$ such that the series

$$\mathcal{I}_m^n(\mathcal{F}) = \sum_{r_1, \dots, r_n \in \mathbb{Z}} \langle w', \mathcal{F}(P_{r_1} \Psi_1; \zeta_1; \dots; P_{r_n} \Psi_n, \zeta_n) \rangle, \quad (7.4)$$

is absolutely convergent when

$$|z_{l_1+\dots+l_{i-1}+p}-\zeta_i|+|z_{l_1+\dots+l_{j-1}+q}-\zeta_j|<|\zeta_i-\zeta_j|, \quad (7.5)$$

for $i, j = 1, \dots, k, i \neq j$ and for $p = 1, \dots, l_i$ and $q = 1, \dots, l_j$. The sum must be analytically extended to a rational function in (z_1, \dots, z_{m+n}) , independent of $(\zeta_1, \dots, \zeta_n)$, with the only possible poles at $z_i = z_j$, of order less than or equal to $N_m^n(v_i, v_j)$, for $i, j = 1, \dots, k, i \neq j$.

2) For $v_1, \dots, v_{m+n} \in V$, and $(z_1, \dots, z_{n+m}) \in \mathbb{C}$ there exist positive integers $N_m^n(v_i, v_j)$, depending only on v_i and v_j , for $i, j = 1, \dots, k, i \neq j$, such that for arbitrary $w' \in W'$, and such that the series

$$\mathcal{J}_m^n(\mathcal{F}) = \sum_{q \in \mathbb{C}} \langle w', E_W^{(m)}(v_1, z_1; \dots; v_m, z_m; P_q(\mathcal{F}(v_{m+1}, z_{m+1}; \dots; v_{m+n}, z_{m+n})) \rangle, \quad (7.6)$$

is absolutely convergent when

$$\begin{aligned} z_i &\neq z_j, \quad i \neq j, \\ |z_i| &> |z_k| > 0, \end{aligned} \quad (7.7)$$

for $i = 1, \dots, m$, and $k = m+1, \dots, m+n$, and the sum can be analytically extended to a rational function in (z_1, \dots, z_{n+m}) with the only possible poles at $z_i = z_j$, of orders less than or equal to $N_m^n(v_i, v_j)$, for $i, j = 1, \dots, k, i \neq j$.

In [10], we the following useful proposition was proven:

Proposition 7. *Let $\mathcal{F} : V^{\otimes n} \rightarrow \mathcal{W}_{z_1, \dots, z_n}$ be composable with m vertex operators. Then we have:*

- (1) *For $p \leq m$, \mathcal{F} is composable with p vertex operators and for $p, q \in \mathbb{Z}_+$ such that $p+q \leq m$ and $l_1, \dots, l_n \in \mathbb{Z}_+$ such that $l_1 + \dots + l_n = p+n$, $\mathcal{F} \circ (E_{V; \mathbf{1}}^{(l_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(l_n)})$ and $E_W^{(p)} \circ_{p+1} \mathcal{F}$ are composable with q vertex operators.*
- (2) *For $p, q \in \mathbb{Z}_+$ such that $p+q \leq m$, $l_1, \dots, l_n \in \mathbb{Z}_+$ such that $l_1 + \dots + l_n = p+n$ and $k_1, \dots, k_{p+n} \in \mathbb{Z}_+$ such that $k_1 + \dots + k_{p+n} = q+p+n$, we have*

$$\begin{aligned} &(\mathcal{F} \circ (E_{V; \mathbf{1}}^{(l_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(l_n)})) \circ (E_{V; \mathbf{1}}^{(k_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(k_{p+n})}) \\ &= \mathcal{F} \circ (E_{V; \mathbf{1}}^{(k_1+\dots+k_{l_1})} \otimes \dots \otimes E_{V; \mathbf{1}}^{(k_{l_1+\dots+l_{n-1}+1+\dots+k_{p+n})}). \end{aligned}$$

- (3) *For $p, q \in \mathbb{Z}_+$ such that $p+q \leq m$ and $l_1, \dots, l_n \in \mathbb{Z}_+$ such that $l_1 + \dots + l_n = p+n$, we have*

$$E_W^{(q)} \circ_{q+1} (\mathcal{F} \circ (E_{V; \mathbf{1}}^{(l_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(l_n)})) = (E_W^{(q)} \circ_{q+1} \mathcal{F}) \circ (E_{V; \mathbf{1}}^{(l_1)} \otimes \dots \otimes E_{V; \mathbf{1}}^{(l_n)}).$$

- (4) *For $p, q \in \mathbb{Z}_+$ such that $p+q \leq m$, we have*

$$E_W^{(p)} \circ_{p+1} (E_W^{(q)} \circ_{q+1} \mathcal{F}) = E_W^{(p+q)} \circ_{p+q+1} \mathcal{F}.$$

Finally, in [10] we find the proof of the following

Proposition 8. *Let now $P_n : W \rightarrow W_{(n)}$, for $n \in \mathbb{C}$ be the projection from W to $W_{(n)}$. For $k, l_1, \dots, l_{n+1} \in \mathbb{Z}_+$ and $v_1^{(1)}, \dots, v_{l_1}^{(1)}, \dots, v_1^{(n+1)}, \dots, v_{l_{n+1}}^{(n+1)} \in V, w \in W$,*

and $w' \in W'$, the series

$$\begin{aligned} & \sum_{r_1, \dots, r_n \in \mathbb{Z}, r_{n+1} \in \mathbb{C}} \langle w', E_W^{(n,1)}(P_{r_1}(E_V^{(l_1)}(v_1^{(1)}, z_1^{(1)}; \dots; v_{l_1}^{(1)}, z_{l_1}^{(1)}; \mathbf{1}_V, z_1^{(0)})); \dots; \\ & P_{r_n}(E_V^{(l_n)}(v_1^{(n)}, z_1^{(n)}; \dots; v_{l_n}^{(n)}, z_{l_n}^{(n)}; \mathbf{1}_V, z_n^{(0)})) \\ & P_{r_{n+1}}(E_W^{(l_{n+1})}(v_1^{(n+1)}, z_1^{(n+1)}; \dots; v_{l_{n+1}}^{(n+1)}, z_{l_{n+1}}^{(n+1)}; w, z_{n+1}^{(0)})) \rangle, \end{aligned} \quad (7.8)$$

converges absolutely to

$$\begin{aligned} & \langle w', E_W^{(n)}(v_1^{(1)}, z_1^{(1)} + z_1^{(0)}; \dots; v_{l_1}^{(1)}, z_{l_1}^{(1)} + z_1^{(0)}; \dots; \\ & v_1^{(n+1)}, z_1^{(n+1)} + z_{n+1}^{(0)}; v_{l_{n+1}}^{(n+1)}, z_{l_{n+1}}^{(n+1)} + z_{n+1}^{(0)}; w) \rangle, \end{aligned}$$

when $0 < |z_p^{(i)}| + |z_q^{(j)}| < |z_i^{(0)} - z_j^{(0)}|$ for $i, j = 1, \dots, n+1$, $i \neq j$, $p = 1, \dots, l_i$, $q = 1, \dots, l_j$.

7.3. Definition of $C_m^n(V, \mathcal{W})$ -spaces. In this Subsection we recall the definition of spaces $C_m^n(V, \mathcal{W})$ for a grading-restricted vertex algebra V . First, recall the definition of shuffles [10]. For $l \in \mathbb{N}$ and $1 \leq s \leq l-1$, let $J_{l;s}$ be the set of elements of S_l which preserve the order of the first s numbers and the order of the last $l-s$ numbers, that is,

$$J_{l;s} = \{\sigma \in S_l \mid \sigma(1) < \dots < \sigma(s), \sigma(s+1) < \dots < \sigma(l)\}.$$

The elements of $J_{l;s}$ are called shuffles, and we use the notation

$$J_{l;s}^{-1} = \{\sigma^{-1} \mid \sigma \in J_{l;s}\}.$$

For a set of n elements (v_1, \dots, v_n) of a grading-restricted vertex algebra V , we consider maps

$$\mathcal{F}(v_1, z_1; \dots; v_n, z_n) : V^{\otimes n} \rightarrow \mathcal{W}_{z_1, \dots, z_n} \quad (7.9)$$

(see Section 4 for the definition of a $\mathcal{W}_{z_1, \dots, z_n}$ space). Note that similar to considerations of [1], (4.2) can be treated as $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$ -torsor of the product of groups of formal parameter transformations. In what follows, according to definitions of Appendix 4, when we write an element \mathcal{F} of the space $\mathcal{W}_{z_1, \dots, z_n}$, we actually have in mind corresponding matrix element $\langle w', \mathcal{F} \rangle$ that absolutely converges (in a certain domain) to a rational form-valued function $R(\langle w', \mathcal{F} \rangle)$. Quite frequently we will write $\langle w', \mathcal{F} \rangle$ which would denote a rational \mathcal{W} -valued form. In notations, we would keep tensor products of vertex algebra elements with wt -powers of z -differentials when it is inevitable only.

In the next Section we prove, that for arbitrary $v_i \in V$, $1 \leq i \leq n$, with formal parameters z_i an element (4.2) as well as the vertex operators (7.1) are invariant with respect to the action of the group $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$. In (7.1) we mean the ordinary vertex operator (as defined in Appendix 10) not affecting the tensor product with corresponding differential. In [10] one finds:

Proposition 9. *The subspace of $\text{Hom}(V^{\otimes n}, \mathcal{W}_{z_1, \dots, z_n})$ consisting of maps having the $L_V(-1)$ -derivative property, having the $L_V(0)$ -conjugation property or being composable with m vertex operators is invariant under the action of S_n . \square*

We next have

Definition 10. For an arbitrary set of vertex algebra elements $v_i, v_j \in V$, and formal complex parameters $z_i, z_j, 1 \leq i \leq n, 1 \leq j \leq m, n \geq 0, m \geq 0$, we denote by $C_m^n(V, \mathcal{W})$, the space of all maps (7.9)

$$\mathcal{F}(v_1, z_1; \dots; v_n, z_n) : V^{\otimes n} \rightarrow \mathcal{W}_{z_1, \dots, z_n}, \quad (7.10)$$

composable with a m of vertex operators (7.1) with vertex algebra elements v_j , with formal parameters z_j . We assume also that (4.2) satisfy $L_V(-1)$ -derivative (4.3), $L_V(0)$ -conjugation (4.8) properties, and the symmetry property with respect to action of the symmetric group S_n :

$$\sum_{\sigma \in J_{n;1}^{-1}} (-1)^{|\sigma|} \mathcal{F}(v_{\sigma(1)}, z_{\sigma(1)}; \dots; v_{\sigma(n)}, z_{\sigma(n)}) = 0. \quad (7.11)$$

In Appendix 12 we give a proof of the following

Proposition 10. For primary vectors of a quasi-conformal grading-restricted vertex algebra V , Definition 10 is canonical, i.e., invariant with respect to the group of n -dimensional transformations

$$(z_1, \dots, z_n) \mapsto (z'_1, \dots, z'_n) = (\rho_1(z_1, \dots, z_n), \dots, \rho_n(z_1, \dots, z_n)),$$

of formal parameters $z_i, 1 \leq i \leq n$.

In Appendix 12 we recall the proof of Proposition 10.

Remark 4. The condition of quasi-conformality is necessary in the proof of invariance of elements of the space $\mathcal{W}_{z_1, \dots, z_n}$ with respect to a vertex algebraic representation (cf. Appendix 10) of the group $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$. In what follows, we will always assume the quasi-conformality of V -modules when it concerns the spaces $C_m^n(V, \mathcal{W})$.

7.4. Coboundary operators. In this Subsection we recall the definition of the coboundary operator for the spaces $C_m^n(V, \mathcal{W})$,

$$\begin{aligned} \delta_m^n \mathcal{F} &= \sum_{i=1}^n (-1)^i \mathcal{F}(\omega_V(v_i, z_i - z_{i+1}) v_{i+1}) \\ &+ \omega_W(v_1, z_1) \mathcal{F}(v_2, z_2; \dots; v_n, z_n) \\ &+ (-1)^{n+1} \omega_W(v_{n+1}, z_{n+1}) \mathcal{F}(v_1, z_1; \dots; v_n, z_n). \end{aligned} \quad (7.12)$$

Note that it is assumed that the coboundary operator does not affect $dz_i^{\text{wt}(v_i)}$ -tensor multipliers in \mathcal{F} . In [10] the following proposition is proved

Proposition 11. The operator (7.12) obeys

$$\delta_m^n : C_m^n(V, \mathcal{W}) \rightarrow C_{m-1}^{n+1}(V, \mathcal{W}), \quad (7.13)$$

$$\delta_{m-1}^{n+1} \circ \delta_m^n = 0, \quad (7.14)$$

$$0 \longrightarrow C_m^0(V, \mathcal{W}) \xrightarrow{\delta_m^0} C_{m-1}^1(V, \mathcal{W}) \xrightarrow{\delta_{m-1}^1} \dots \xrightarrow{\delta_1^{m-1}} C_0^m(V, \mathcal{W}) \longrightarrow 0, \quad (7.15)$$

i.e., it provides the chain-cochain complex $(C_m^n(V, \mathcal{W}), \delta_m^n)$. \square

8. APPLICATION: THE PRODUCT OF $C_m^n(V, \mathcal{W})$ -SPACES

In this Section we consider an application of the material of Section 5 to the spaces $C_m^n(V, \mathcal{W})$ of a complex (Definition 10) described in the previous Section. We introduce the product of elements of two spaces of the complex with the image in another space of the complex coherent with respect to the original differential (7.12), and the symmetry property (7.11). We prove the canonicity of the product, and derive an analogue of Leibniz formula.

Definition 11. For $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in C_m^k(V, \mathcal{W})$, and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in C_{m'}^n(V, \mathcal{W})$ the product

$$\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \mapsto \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon),$$

is a $\mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$ -valued rational form

$$\begin{aligned} & \langle w', \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\ &= \langle w', \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\ &= \sum_{u \in V} \langle w', Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\ & \qquad \langle w', Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle, \end{aligned} \quad (8.1)$$

defined by (5.2).

Remark 5. Let t be the number of common vertex operators the mappings $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in C_m^k(V, \mathcal{W})$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in C_{m'}^n(V, \mathcal{W})$, are composable with. Similar to the case of common formal parameters, this case is separately treated with a decrease to $m + m' - t$ of the number of composable vertex operators. In what follows, we exclude this case from considerations.

The action of $\sigma \in S_{k+n}$ on the product $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_{k+1}, y_1; \dots; v'_n, y_n; \epsilon)$ (8.1) is given by (4.9). We then have

Proposition 12. For $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in C_m^k(V, \mathcal{W})$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in C_{m'}^n(V, \mathcal{W})$, the product $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon)$ (8.1) belongs to the space $C_{m+m'}^{k+n}(V, \mathcal{W})$, i.e.,

$$\cdot_\epsilon : C_m^k(V, \mathcal{W}) \times C_{m'}^n(V, \mathcal{W}) \rightarrow C_{m+m'}^{k+n}(V, \mathcal{W}).$$

Proof. In Proposition 4 we proved that $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \in \mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$. It is clear that

$$\cdot_\epsilon : C_l^k(V, \mathcal{W}) \times C_l^m(V, \mathcal{W}) \rightarrow C_l^{k+n}(V, \mathcal{W}),$$

for some l . First, we show that (7.11) for $\sigma \in S_{k+n}$,

$$\sum_{\sigma \in J_{k+n; s}^{-1}} (-1)^{|\sigma|} \mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}; v'_{\sigma(k+1)}, y_{\sigma(k+1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}) = 0.$$

For arbitrary $w' \in W'$, we have

$$\begin{aligned}
& \sum_{\sigma \in J_{k+n;s}^{-1}} (-1)^{|\sigma|} \langle w', \mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}; v'_{\sigma(1)}, y_{\sigma(1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}) \rangle \\
&= \sum_{\sigma \in J_{k+n;s}^{-1}} (-1)^{|\sigma|} \sum_{u \in V} \langle w', Y_{WV}^W(\mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}), \zeta_1) u \rangle \\
&\quad \langle w', Y_{WV}^W(\mathcal{F}(v'_{\sigma(1)}, y_{\sigma(1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}), \zeta_2) \bar{u} \rangle \\
&= \sum_{u \in V} \sum_{\sigma \in J_{k+n;s}^{-1}} (-1)^{|\sigma|} \langle w', e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}) \rangle \\
&\quad \langle w', e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) \mathcal{F}(v'_{\sigma(1)}, y_{\sigma(1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}) \rangle \\
&= \sum_{u \in V} \langle w', e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \sum_{\sigma \in J_{k;s}^{-1}} (-1)^{|\sigma|} \mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}) \rangle \\
&\quad \langle w', e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) \mathcal{F}(v'_{\sigma(1)}, y_{\sigma(1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}) \rangle \\
&+ \sum_{u \in V} \langle w', e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \mathcal{F}(v_{\sigma(1)}, x_{\sigma(1)}; \dots; v_{\sigma(k)}, x_{\sigma(k)}) \rangle \\
&\quad \langle w', e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) \sum_{\sigma \in J_{n;s}^{-1}} (-1)^{|\sigma|} \mathcal{F}(v'_{\sigma(1)}, y_{\sigma(1)}; \dots; v'_{\sigma(n)}, y_{\sigma(n)}) \rangle = 0,
\end{aligned}$$

since, $J_{k+n;s}^{-1} = J_{k;s}^{-1} \times J_{n;s}^{-1}$, and due to the fact that $\mathcal{F}(v_1, x_1; \dots; v_k, x_k)$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$ satisfy (4.9).

Next, we show that $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon)$ (8.1) is composable with $m + m'$ vertex operators. Recall that $\mathcal{F}(v_1, x_1; \dots; v_k, x_k)$ is composable with m vertex operators, and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$ is composable with m' vertex operators. For $\mathcal{F}(v_1, x_1; \dots; v_k, x_k)$ we have:

1) Let $l_1, \dots, l_k \in \mathbb{Z}_+$ such that $l_1 + \dots + l_k = k + m$, and $v_1, \dots, v_{k+m} \in V$, and arbitrary $w' \in W'$. Set

$$\Psi_i = E_V^{(l_i)}(v_{k_1}, x_{k_1} - \zeta_i; \dots; v_{k_i}, x_{k_i} - \zeta_i; \mathbf{1}_V), \quad (8.2)$$

where

$$k_1 = l_1 + \dots + l_{i-1} + 1, \quad \dots, \quad k_i = l_1 + \dots + l_{i-1} + l_i, \quad (8.3)$$

for $i = 1, \dots, k$. Then the series

$$\mathcal{I}_m^k(\mathcal{F}) = \sum_{r_1, \dots, r_k \in \mathbb{Z}} \langle w', \mathcal{F}(P_{r_1} \Psi_1; \zeta_1; \dots; P_{r_k} \Psi_k, \zeta_k) \rangle, \quad (8.4)$$

is absolutely convergent when

$$|x_{l_1 + \dots + l_{i-1} + p} - \zeta_i| + |x_{l_1 + \dots + l_{j-1} + q} - \zeta_i| < |\zeta_i - \zeta_j|, \quad (8.5)$$

for $i, j = 1, \dots, k$, $i \neq j$ and for $p = 1, \dots, l_i$ and $q = 1, \dots, l_j$. There exist positive integers $N_m^k(v_i, v_j)$, depending only on v_i and v_j for $i, j = 1, \dots, k$, $i \neq j$, such that the sum is analytically extended to a rational function in (x_1, \dots, x_{k+m}) , independent

of $(\zeta_1, \dots, \zeta_k)$, with the only possible poles at $x_i = x_j$, of order less than or equal to $N_m^k(v_i, v_j)$, for $i, j = 1, \dots, k$, $i \neq j$.

For $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$ we have:

1') Let $l'_1, \dots, l'_n \in \mathbb{Z}_+$ such that $l'_1 + \dots + l'_n = n + m'$, $v'_1, \dots, v_{n+m'} \in V$ and arbitrary $w' \in W'$. Set

$$\Psi'_{i'} = E_V^{(l'_{i'})}(v'_{k'_1}, y_{k'_1} - \zeta'_{i'}; \dots; v'_{k'_{i'}}, y_{k'_{i'}} - \zeta'_{i'}; \mathbf{1}_V), \quad (8.6)$$

where

$$k'_1 = l'_1 + \dots + l'_{i'-1} + 1, \quad \dots, \quad k'_{i'} = l'_1 + \dots + l'_{i'-1} + l'_{i'}, \quad (8.7)$$

for $i' = 1, \dots, n$. Then the series

$$\mathcal{I}_{m'}^{n'}(\mathcal{F}) = \sum_{r'_1, \dots, r'_n \in \mathbb{Z}} \langle w', \mathcal{F}(P_{r'_1} \Psi'_1; \zeta'_1; \dots; P_{r'_n} \Psi'_n, \zeta'_n) \rangle, \quad (8.8)$$

is absolutely convergent when

$$|y_{l'_1 + \dots + l'_{i'-1} + p'} - \zeta'_{i'}| + |y_{l'_1 + \dots + l'_{j'-1} + q'} - \zeta'_{j'}| < |\zeta'_{i'} - \zeta'_{j'}|, \quad (8.9)$$

for $i', j' = 1, \dots, n$, $i' \neq j'$ and for $p' = 1, \dots, l'_{i'}$ and $q' = 1, \dots, l'_{j'}$. There exist positive integers $N_{m'}^n(v'_{i'}, v'_{j'})$, depending only on $v'_{i'}$ and $v'_{j'}$, for $i, j = 1, \dots, n$, $i' \neq j'$, such that the sum is analytically extended to a rational function in $(y_1, \dots, y_{n+m'})$, independent of $(\zeta'_1, \dots, \zeta'_n)$, with the only possible poles at $y_{i'} = y_{j'}$, of order less than or equal to $N_{m'}^n(v'_{i'}, v'_{j'})$, for $i', j' = 1, \dots, n$, $i' \neq j'$.

Now let us consider the first condition of Definition 9 of composability for the product (8.1) of $\mathcal{F}(v_1, x_1; \dots; v_k, x_k)$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$ with a number of vertex operators. Then we obtain for $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon)$ the following. We redefine the notations for the set

$$\begin{aligned} & (v''_1, \dots, v''_k; v''_{k+1}, \dots, v''_{k+m}; v''_{k+m+1}, \dots, v''_{k+n+m+m'}; v_{n+1}, \dots, v'_{n+m'}) \\ & = (v_1, \dots, v_k; v_{k+1}, \dots, v_{k+m}; v'_1, \dots, v'_n; v'_{n+1}, \dots, v'_{n+m'}), \\ & (z_1, \dots, z_k; z_{k+1}, \dots, z_{k+n}) = (x_1, \dots, x_k; y_1, \dots, y_n), \end{aligned}$$

of vertex algebra V elements. Introduce $l''_1, \dots, l''_{k+n} \in \mathbb{Z}_+$, such that $l''_1 + \dots + l''_{k+n} = k + n + m + m'$. Define

$$\Psi''_i = E_V^{(l''_i)}(v''_{k''_1}, z_{k''_1} - \zeta''_{i''}; \dots; v''_{k''_{i''}}, z_{k''_{i''}} - \zeta''_{i''}; \mathbf{1}_V), \quad (8.10)$$

where

$$k''_1 = l''_1 + \dots + l''_{i''-1} + 1, \quad \dots, \quad k''_{i''} = l''_1 + \dots + l''_{i''-1} + l''_{i''}, \quad (8.11)$$

for $i'' = 1, \dots, k + n$, and we take

$$(\zeta''_1, \dots, \zeta''_{k+n}) = (\zeta_1, \dots, \zeta_k; \zeta'_1, \dots, \zeta'_n).$$

Then we consider

$$\mathcal{I}_{m+m'}^{k+n}(\mathcal{F}) = \sum_{r''_1, \dots, r''_{k+n} \in \mathbb{Z}} \langle w', \mathcal{F}(P_{r''_1} \Psi''_1; \zeta''_1; \dots; P_{r''_{k+n}} \Psi''_{k+n}, \zeta''_{k+n}) \rangle, \quad (8.12)$$

and prove it is absolutely convergent with some conditions.

The condition

$$|z_{l_1''+\dots+l_{i-1}''+p''} - \zeta_i''| + |z_{l_1''+\dots+l_{j-1}''+q''} - \zeta_j''| < |\zeta_i'' - \zeta_j''|, \quad (8.13)$$

of absolute convergence for (8.12) for $i'', j'' = 1, \dots, k+n$, $i \neq j$ and for $p'' = 1, \dots, l_i''$ and $q'' = 1, \dots, l_j''$, follows from the conditions (8.5) and (8.19). The action of $e^{\zeta L_W(-1)} Y_W(\cdot, \cdot)$, $a = 1, 2$, in

$$\begin{aligned} & \langle w', e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta) \sum_{r_1, \dots, r_k \in \mathbb{Z}} \mathcal{F}(P_{r_1} \Psi_1; \zeta_1; \dots; P_{r_k} \Psi_k, \zeta_k) \rangle, \\ & \langle w', e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\tilde{\zeta}) \sum_{r'_1, \dots, r'_n \in \mathbb{Z}} \mathcal{F}(P_{r'_1} \Psi'_1; \zeta'_1; \dots; P_{r'_k} \Psi'_n, \zeta'_n) \rangle, \end{aligned}$$

does not affect the absolute convergence of (8.4) and (8.8). We obtain

$$\begin{aligned} |\mathcal{I}_{m+m'}^{k+n}(\mathcal{F})| &= \\ &= \left| \sum_{r''_1, \dots, r''_{k+n} \in \mathbb{Z}} \langle w', \mathcal{F}(P_{r''_1} \Psi''_1; \zeta''_1; \dots; P_{r''_{k+n}} \Psi''_{k+n}, \zeta''_{k+n}) \rangle \right| \\ &= \left| \sum_{u \in V} \langle w', Y_{VW}^W \left(\sum_{r_1, \dots, r_k \in \mathbb{Z}} \mathcal{F}(P_{r_1} \Psi_1; \zeta_1; \dots; P_{r_k} \Psi_k, \zeta_k), \zeta \right) u \right. \\ & \quad \left. \langle w', Y_{VW}^W \left(\sum_{r'_1, \dots, r'_n \in \mathbb{Z}} \mathcal{F}(P_{r'_1} \Psi'_1; \zeta'_1; \dots; P_{r'_n} \Psi'_n, \zeta'_n), \tilde{\zeta} \right) \bar{u} \right| \\ &\leq |\mathcal{I}_m^k(\mathcal{F})| |\mathcal{I}_{m'}^n(\mathcal{F})|. \end{aligned}$$

Thus, we infer that (8.12) is absolutely convergent. Recall that the maximal orders of possible poles of (8.12) are $N_m^k(v_i, v_j)$, $N_{m'}^n(v'_{i'}, v'_{j'})$ at $x_i = x_j$, $y_{i'} = y_{j'}$. From the last expression we infer that there exist positive integers $N_{m+m'}^{k+n}(v''_{i''}, v''_{j''})$ for $i, j = 1, \dots, k$, $i \neq j$, $i', j' = 1, \dots, n$, $i' \neq j'$, depending only on $v''_{i''}$ and $v''_{j''}$ for $i'', j'' = 1, \dots, k+n$, $i'' \neq j''$ such that the series (8.12) can be analytically extended to a rational function in $(x_1, \dots, x_k; y_1, \dots, y_n)$, independent of $(\zeta''_1, \dots, \zeta''_{k+n})$, with extra possible poles at $x_i = y_j$, of order less than or equal to $N_{m+m'}^{k+n}(v''_{i''}, v''_{j''})$, for $i'', j'' = 1, \dots, n$, $i'' \neq j''$.

Let us proceed with the second condition of composability. For $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in C_m^k(V, \mathcal{W})$, and $(v_1, \dots, v_{k+m}) \in V$, $(x_1, \dots, x_{k+m}) \in \mathbb{C}$, we have

2) For arbitrary $w' \in W'$, the series

$$\mathcal{J}_m^k(\mathcal{F}) = \sum_{q \in \mathbb{C}} \langle w', E_W^{(m)} \left(v_1, x_1; \dots; v_m, x_m; P_q(\mathcal{F}(v_{m+1}, x_{m+1}; \dots; v_{m+k}, x_{m+k})) \right) \rangle, \quad (8.14)$$

is absolutely convergent when

$$\begin{aligned} x_i &\neq x_j, \quad i \neq j, \\ |x_i| &> |x_{k'}| > 0, \end{aligned} \quad (8.15)$$

for $i = 1, \dots, m$, and $k' = m + 1, \dots, k + m$, and the sum can be analytically extended to a rational function in (x_1, \dots, x_{k+m}) with the only possible poles at $x_i = x_j$, of orders less than or equal to $N_m^k(v_i, v_j)$, for $i, j = 1, \dots, k$, $i \neq j$.

2') For $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in C_{m'}^n(V, \mathcal{W})$, $(v'_1, \dots, v'_{n+m'}) \in V$, and $(y_1, \dots, y_{n+m'}) \in \mathbb{C}$, the series

$$\begin{aligned} \mathcal{J}_{m'}^n(\mathcal{F}) &= \sum_{q \in \mathbb{C}} \langle w', E_W^{(m')} (v'_1, y_1; \dots; v'_{m'}, y_{m'}; \\ &P_q(\mathcal{F}(v'_{m'+1}, y_{m'+1}; \dots; v'_{m'+n}, y_{m'+n}))) \rangle, \end{aligned} \quad (8.16)$$

is absolutely convergent when

$$\begin{aligned} y_{i'} &\neq y_{j'}, \quad i' \neq j', \\ |y_{i'}| &> |y_{k''}| > 0, \end{aligned} \quad (8.17)$$

for $i' = 1, \dots, m'$, and $k'' = m' + 1, \dots, n + m'$, and the sum can be analytically extended to a rational function in $(y_1, \dots, y_{n+m'})$ with the only possible poles at $y_{i'} = y_{j'}$, of orders less than or equal to $N_{m'}^n(v'_{i'}, v'_{j'})$, for $i', j' = 1, \dots, n$, $i' \neq j'$.

2'') Thus, for the product (8.1) we obtain $(v''_1, \dots, v''_{k+n+m+m'}) \in V$, and $(z_1, \dots, z_{k+n+m+m'}) \in \mathbb{C}$, we find positive integers $N_{m+m'}^{k+n}(v'_i, v'_j)$, depending only on v'_i and v'_j , for $i'', j'' = 1, \dots, k + n$, $i'' \neq j''$, such that for arbitrary $w' \in W'$. First we note

Lemma 2.

$$\begin{aligned} &\sum_{q \in \mathbb{C}} \langle w', E_W^{(m+m')} (v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \\ &P_q(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}))) \rangle \\ &= \sum_{u \in V} \langle w', E_W^{(m)} (v_{k+1}, x_{k+1}; \dots; v_{k+m}, x_{k+m}; \\ &P_q(Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u)) \rangle \\ &\langle w', E_W^{(m')} (v'_{n+1}, y_{n+1}; \dots; v'_{n+m'}, y_{n+m'}) \\ &P_q(Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u}) \rangle. \end{aligned}$$

Proof. Consider

$$\begin{aligned} &\sum_{u \in V} \langle w', E_W^{(m+m')} (v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \\ &P_q(Y_{WV}^W(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}), \zeta_1) u)) \rangle \\ &\langle w', E_W^{(m+m')} (v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}) \\ &P_q(Y_{WV}^W(\mathcal{F}(v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; \\ &v''_{m+m'+k+n}, z_{m+m'+k+n}), \zeta_2) \bar{u}) \rangle \end{aligned}$$

$$\begin{aligned}
&= \sum_{q \in \mathbb{C}} \sum_{u \in V} \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\
&\quad \left. P_q \left(e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}) \right) \right) \rangle \\
&\quad \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\
&\quad \left. P_q \left(e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) \mathcal{F}(v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; \right. \right. \\
&\quad \quad \left. \left. v''_{m+m'+k+n}, z_{m+m'+k+n}) \right) \right) \rangle.
\end{aligned}$$

The action of exponentials $e^{\zeta_a L_W(-1)}$, $a = 1, 2$, of the differential operator $L_W(-1)$, and W -module vertex operators $Y_W(u, -\zeta_1)$, $Y_W(u, -\zeta_2)$ shifts the grading index q of W_q -subspaces by $\alpha \in \mathbb{C}$ which can be later rescaled to q . Thus, we can rewrite the last expression as

$$\begin{aligned}
&= \sum_{q \in \mathbb{C}} \sum_{u \in V} \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\
&\quad \left. e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}) \right) \right) \rangle \\
&\quad \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\
&\quad \left. e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}) \right) \right) \rangle \\
&= \sum_{q \in \mathbb{C}} \sum_{u \in V} \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\
&\quad \left. Y_{WV}^W \left(P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}) \right), \zeta_1 \right) u \right) \rangle \\
&\quad \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\
&\quad \left. Y_{WV}^W \left(P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}), -\zeta_2 \right) \bar{u} \right) \right) \rangle \\
&= \sum_{q \in \mathbb{C}} \sum_{\tilde{w} \in W} \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \tilde{w} \right) \rangle \\
&\quad \sum_{u \in V} \langle w', Y_{WV}^W \left(P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}), -\zeta_1 \right) u \right) \rangle \\
&\quad \langle \tilde{w}', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \tilde{w} \right) \rangle \\
&\quad \langle w', Y_{WV}^W \left(P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}), -\zeta_2 \right) \bar{u} \right) \rangle
\end{aligned}$$

$$= \sum_{q \in \mathbb{C}} \langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\ \left. P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}; \right. \right. \\ \left. \left. v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}) \right) \right) \rangle.$$

Now note that, according to Proposition 6, as an element of $\mathcal{W}_{z_1, \dots, z_{k+n+m+m'}}$

$$\langle w', E_W^{(m+m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\ \left. P_{q+\alpha} \left(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}; \right. \right. \\ \left. \left. v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}) \right) \right) \rangle, \quad (8.18)$$

is invariant with respect to the action of $\sigma \in S_{k+n+m+m'}$. Thus we are able to use this invariance to show that (8.18) is reduced to

$$\langle w', E_W^{(m+m')} \left(v''_{k+1}, z_{k+1}; \dots; v''_{k+1+m}, z_{k+1+m}; v''_{n+1}, z_{n+1}; \dots; v''_{n+1+m'}, z_{n+1+m'}; \right. \\ \left. P_{q+\alpha} \left(\mathcal{F}(v''_1, z_1; \dots; v''_k, z_k; v''_{k+1}, z_{k+1}; \dots; v''_{k+n}, z_{k+n}) \right) \right) \rangle \\ = \langle w', E_W^{(m+m')} \left(v_{k+1}, x_{k+1}; \dots; v_{k+1+m}, x_{k+1+m}; v'_{n+1}, y_{n+1}; \dots; v'_{n+1+m'}, y_{n+1+m'}; \right. \\ \left. P_{q+\alpha} \left(\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n) \right) \right) \rangle.$$

Similarly, since

$$\langle w', E_W^{(m)} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\ \left. P_q \left(Y_{WV}^W \left(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k}, z_{m+m'+k}), \zeta_1 \right) u \right) \right) \rangle, \\ \langle w', E_W^{(m')} \left(v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\ \left. P_q \left(Y_{WV}^W \left(\mathcal{F}(v''_{m+m'+k+1}, z_{m+m'+k+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}), \zeta_2 \right) \bar{u} \right) \right) \rangle.$$

correspond to elements of $\mathcal{W}_{z_1, \dots, z_{m+m'+k}}$ and $\mathcal{W}_{z_{m+m'+k+1}, \dots, z_{m+m'+k+n}}$, we use Proposition 6 again and obtain

$$\langle w', E_W^{(m)} \left(v_{k+1}, x_{k+1}; \dots; v_{k+m}, x_{k+m}; P_q \left(Y_{WV}^W \left(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1 \right) u \right) \right) \rangle \\ \langle w', E_W^{(m')} \left(v'_{n+1}, y_{n+1}; \dots; v'_{n+m'}, y_{n+m'}; P_q \left(Y_{WV}^W \left(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2 \right) \bar{u} \right) \right) \rangle,$$

correspondingly. Thus, the assertion of Lemma follows. \square

Under conditions

$$z_{i''} \neq z_{j''}, \quad i'' \neq j'', \\ |z_{i''}| > |z_{k''}| > 0, \quad (8.19)$$

for $i'' = 1, \dots, m + m'$, and $k''' = m + m' + 1, \dots, m + m' + k + n$, let us introduce

$$\begin{aligned} \mathcal{J}_{m+m'}^{k+n}(\mathcal{F}) &= \sum_{q \in \mathbb{C}} \langle w', E_W^{(m+m')} (v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \\ &P_q(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}); \epsilon) \rangle. \end{aligned} \quad (8.20)$$

Using Lemma 2 we obtain

$$\begin{aligned} &|\mathcal{J}_{m+m'}^{k+n}(\mathcal{F})| \\ &= \left| \sum_{q \in \mathbb{C}} \langle w', E_W^{(m+m')} (v''_1, z_1; \dots; v''_{m+m'}, z_{m+m'}; \right. \\ &\quad \left. P_q(\mathcal{F}(v''_{m+m'+1}, z_{m+m'+1}; \dots; v''_{m+m'+k+n}, z_{m+m'+k+n}); \epsilon) \rangle \right| \\ &= \left| \sum_{q \in \mathbb{C}} \sum_{u \in V} \langle w', E_W^{(m)} (v_{k+1}, x_{k+1}; \dots; v_{k+m}, x_{k+m}; \right. \\ &\quad \left. P_q(Y_{WV}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u) \rangle \right. \\ &\quad \left. \langle w', E_W^{(m')} (v'_{n+1}, y_{n+1}; \dots; v'_{n+m'}, y_{n+m'}; \right. \\ &\quad \left. P_q(Y_{WV}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u}) \rangle \right| \\ &\leq |\mathcal{J}_m^k(\mathcal{F})| |\mathcal{J}_{m'}^n(\mathcal{F})|, \end{aligned}$$

where we have used the invariance of (8.1) with respect to $\sigma \in S_{m+m'+k+n}$. According to Proposition 8 $\mathcal{J}_m^k(\mathcal{F})$ and $\mathcal{J}_{m'}^n(\mathcal{F})$ in the last expression are absolute convergent. Thus, we infer that $\mathcal{J}_{m+m'}^{k+n}(\mathcal{F})$ is absolutely convergent, and the sum (8.12) is analytically extendable to a rational function in $(z_1, \dots, z_{k+n+m+m'})$ with the only possible poles at $x_i = x_j$, $y_{i'} = y_{j'}$, and at $x_i = y_{j'}$, i.e., the only possible poles at $z_{i''} = z_{j''}$, of orders less than or equal to $N_{m+m'}^{k+n}(v''_{i''}, v''_{j''})$, for $i'', j'' = 1, \dots, k''', i'' \neq j''$. This finishes the proof of the proposition. \square

Now we prove the following

Corollary 1. *For $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in C_m^k(V, \mathcal{W})$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in C_{m'}^n(V, \mathcal{W})$, the product*

$$\begin{aligned} &\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \\ &= \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \end{aligned} \quad (8.21)$$

is canonical with respect to the action

$$\begin{aligned} &(x_1, \dots, x_k; y_1, \dots, y_n) \mapsto (x'_1, \dots, x'_k; y'_1, \dots, y'_n) \\ &= (\rho_1(x_1, \dots, x_k; y_1, \dots, y_n), \dots, \rho_{k+n}(x_1, \dots, x_k; y_1, \dots, y_n)), \end{aligned} \quad (8.22)$$

of elements of the group $\text{Aut}_{x_1, \dots, x_k; y_1, \dots, y_n} \mathcal{O}^{(k+n)}$.

Proof. In Subsection 6 we have proved that the product (5.2) belongs to $W_{x_1, \dots, x_k; y_1, \dots, y_n}$, and is invariant with respect to the group $\text{Aut}_{x_1, \dots, x_k; y_1, \dots, y_n} \mathcal{O}^{(k+n)}$. Similar as in the proof of Proposition 10, vertex operators $\omega_V(v_i, x_i)$, $1 \leq i \leq m$, composable with $\mathcal{F}(v_1, x_1; \dots; v_k, x_k)$, and vertex operators $\omega_V(v_j, y_j)$, $1 \leq j \leq m'$, composable with $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$, are also invariant with respect to $(\rho_1(x_1, \dots, x_k; y_1, \dots, y_n), \dots, \rho_{k+n}(x_1, \dots, x_k; y_1, \dots, y_n)) \in \text{Aut}_{x_1, \dots, x_k; y_1, \dots, y_n} \mathcal{O}^{(k+n)}$. \square

Since the product of $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in C_m^k(V, \mathcal{W})$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in C_{m'}^n(V, \mathcal{W})$ results in an element of $C_{m+m'}^{k+n}(V, \mathcal{W})$, then, similar to Proposition 9 [10], the following corollary follows directly from Proposition (12) and Definition 8:

Corollary 2. *For the spaces $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$ with the product (5.2) $\mathcal{F} \in \mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$, the subspace of $\mathcal{W}_{x_1, \dots, x_k; y_1, \dots, y_n}$ consisting of maps having the $L_W(-1)$ -derivative property, having the $L_V(0)$ -conjugation property or being composable with m vertex operators is invariant under the action of S_{k+n} .*

Finally, we have the following

Corollary 3. *For a fixed set $(v_1, \dots, v_k; v_{k+1}, \dots, v_{k+n}) \in V$ of vertex algebra elements, and fixed $k+n$, and $m+m'$, the ϵ -product $\mathcal{F}(v_1, z_1; \dots; v_k, z_k; v_{k+1}, z_{k+1}; \dots; v_{k+n}, z_{k+n}; \epsilon)$,*

$$\cdot_\epsilon : C_m^k(V, \mathcal{W}) \times C_{m'}^n(V, \mathcal{W}) \rightarrow C_{m+m'}^{k+n}(V, \mathcal{W}),$$

of the spaces $C_m^k(V, \mathcal{W})$ and $C_{m'}^n(V, \mathcal{W})$, for all choices of $k, n, m, m' \geq 0$, is the same element of $C_{m+m'}^{k+n}(V, \mathcal{W})$ for all possible $k \geq 0$.

Proof. In Proposition 4 we have proved that the result of the maps belongs to $W_{x_1, \dots, x_k; y_1, \dots, y_n}$, for all $k, n \geq 0$, and fixed $k+n$. As in the proof of Proposition 12, by checking conditions for the forms (8.4) and (8.8), we see that, by Proposition 7, the product $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n)$ is composable with fixed $m+m'$ vertex operators for every $0 \leq l \leq k$. \square

8.1. Coboundary operator acting on the product space. In Proposition 12 we proved that the product (8.1) of elements of spaces $C_m^k(V, \mathcal{W})$ and $C_{m'}^n(V, \mathcal{W})$ belongs to $C_{m+m'}^{k+n}(V, \mathcal{W})$. Thus, the product admits the action of the differential operator $\delta_{m+m'}^{k+n}$ defined in (7.12). The co-boundary operator (7.12) has a variation of Leibniz law with respect to the product (8.1). Indeed, we state here

Proposition 13. *For $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \in C_m^k(V, \mathcal{W})$ and $\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \in C_{m'}^n(V, \mathcal{W})$, the action of $\delta_{m+m'}^{k+n}$ on their product (8.1) is given by*

$$\begin{aligned} & \delta_{m+m'}^{k+n} (\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)) \\ &= (\delta_m^k \mathcal{F}(v_1, x_1; \dots; v_k, x_k)) \cdot_\epsilon \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \\ &+ (-1)^k \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon (\delta_{m'}^n \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)). \end{aligned} \quad (8.23)$$

Remark 6. Checking (7.12) we see that an extra arbitrary vertex algebra element $v_{n+1} \in V$, as well as a corresponding extra arbitrary formal parameter z_{n+1} appear as a result of the action of δ_m^n on $\mathcal{F} \in C_m^n(V, \mathcal{W})$ mapping it to $C_{m-1}^{n+1}(V, \mathcal{W})$. In

application to the ϵ -product (8.1) these extra arbitrary elements are involved in the definition of the action of $\delta_{m+m'}^{k+n}$ on $\mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n)$.

Proof. According to (7.12), the action of $\delta_{m+m'}^{k+n}$ on $\mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon)$ is given by

$$\begin{aligned}
& \langle w', \delta_{m+m'}^{k+n} \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\
&= \langle w', \sum_{i=1}^k (-1)^i \mathcal{F}(v_1, x_1; \dots; v_{i-1}, x_{i-1}; \omega_V(v_i, x_i - x_{i+1})v_{i+1}, x_{i+1}; v_{i+2}, x_{i+2}; \\
&\quad \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\
&\quad + \sum_{i=1}^n (-1)^i \langle w', \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_{i-1}, y_{i-1}; \\
&\quad \quad \omega_V(v'_i, y_i - y_{i+1};) v'_{i+1}, y_{i+1}; v'_{i+2}, y_{i+2}; \dots; v'_n, y_n; \epsilon) \rangle \\
&\quad + \langle w', \omega_W(v_1, x_1) \mathcal{F}(v_2, x_2; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\
&\quad + \langle w, (-1)^{k+n+1} \omega_W(v'_{n+1}, y_{n+1}) \mathcal{F}(v_1, x_1; \dots; v_k, x_k; v'_1, y_1; \dots; v'_n, y_n; \epsilon) \rangle \\
&= \sum_{u \in V} \langle w', \sum_{i=1}^k (-1)^i Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_{i-1}, x_{i-1}; \omega_V(v_i, x_i - x_{i+1})v_{i+1}, x_{i+1}; \\
&\quad v_{i+2}, x_{i+2}; \dots; v_k, x_k), \zeta_1) u \rangle \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&\quad + \sum_{u \in V} \sum_{i=1}^n (-1)^i \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_{i-1}, y_{i-1}; \\
&\quad \quad \omega_V(v'_i, y_i - y_{i+1};) v'_{i+1}, y_{i+1}; v'_{i+2}, y_{i+2}; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&\quad + \sum_{u \in V} \langle w', Y_{VW}^W(\omega_W(v_1, x_1) \mathcal{F}(v_2, x_2; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&\quad + \sum_{u \in V} \langle w', Y_{VW}^W((-1)^{k+1} \omega_W(v_{k+1}, x_{k+1}) \mathcal{F}(v_1, x_1; \dots; v_k), \zeta_1) u \rangle \\
&\quad \quad \langle w', Y_{VW}^W(\mathcal{F}(x_k; v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle
\end{aligned}$$

$$\begin{aligned}
& - \sum_{u \in V} \langle w', (-1)^{k+1} \langle w', Y_{VW}^W(\omega_W(v_{k+1}, x_{k+1}) \mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u) \\
& \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& + \sum_{u \in V} \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u) \\
& \quad \langle w', Y_{VW}^W(\omega_W(v'_1, y_1) \mathcal{F}(v'_2, y_2; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& \sum_{u \in V} \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) \rangle \\
& \quad \langle w', Y_{VW}^W(\omega_W(v'_1, y_1) \mathcal{F}(v'_2, y_2; \dots; v'_n, y_n), \zeta_2) \rangle \\
& = \sum_{u \in V} \langle w', Y_{VW}^W(\delta_m^k \mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u) \\
& \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& + (-1)^k \sum_{u \in V} \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u) \\
& \quad \langle w', Y_{VW}^W(\delta_m^n \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& = \langle w', \delta_m^k \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon \langle w', \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \rangle \\
& \quad + (-1)^k \langle w', \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \cdot_\epsilon \delta_m^n \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle,
\end{aligned}$$

since,

$$\begin{aligned}
& \sum_{u \in V} \langle w', (-1)^{k+1} Y_{VW}^W(\omega_W(v_{k+1}, x_{k+1}) \mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u) \\
& \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& = \sum_{u \in V} \langle w', (-1)^{k+1} e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \omega_W(v_{k+1}, x_{k+1}) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
& \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& = \sum_{u \in V} \langle w', (-1)^{k+1} e^{\zeta_1 L_W(-1)} \omega_W(v_{k+1}, x_{k+1}) Y_W(u, -\zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
& \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& = \sum_{u \in V} \langle w', (-1)^{k+1} \omega_W(v_{k+1}, x_{k+1} + \zeta_1) e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
& \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
& = \sum_{v \in V} \sum_{u \in V} \langle v', (-1)^{k+1} \omega_W(v_{k+1}, x_{k+1} + \zeta_1) w \rangle \\
& \quad \langle w', e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
& \quad \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle
\end{aligned}$$

$$\begin{aligned}
&= \sum_{u \in V} \langle w', e^{\zeta_1 L_W(-1)} Y_W(u, -\zeta_1) \mathcal{F}(v_1, x_1; \dots; v_k, x_k) \rangle \\
&\quad \sum_{v \in V} \langle w', (-1)^{k+1} \omega_W(v_{k+1}, x_{k+1} + \zeta_1) w \rangle \langle w', Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&= \sum_{u \in V} \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', (-1)^{k+1} \omega_W(v_{k+1}, x_{k+1} + \zeta_1) Y_{VW}^W(\mathcal{F}(v'_1, y_1; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle \\
&= \sum_{u \in V} \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', (-1)^{k+1} \omega_W(v_{k+1}, x_{k+1} + \zeta_1) e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\
&= \sum_{u \in V} \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', (-1)^{k+1} e^{\zeta_2 L_W(-1)} Y_W(\bar{u}, -\zeta_2) \omega_W(v_{k+1}, x_{k+1} + \zeta_1 - \zeta_2) \mathcal{F}(v'_1, y_1; \dots; v'_n, y_n) \rangle \\
&= \sum_{u \in V} \langle w', Y_{VW}^W(\mathcal{F}(v_1, x_1; \dots; v_k, x_k), \zeta_1) u \rangle \\
&\quad \langle w', Y_{VW}^W(\omega_W(v'_1, y_1) \mathcal{F}(v'_2, y_2; \dots; v'_n, y_n), \zeta_2) \bar{u} \rangle,
\end{aligned}$$

due to locality (10.7) of vertex operators, and arbitrariness of $v_{k+1} \in V$ and x_{k+1} , we can always put

$$\omega_W(v_{k+1}, x_{k+1} + \zeta_1 - \zeta_2) = \omega_W(v'_1, y_1),$$

for $v_{k+1} = v'_1$, $x_{k+1} = y_1 + \zeta_2 - \zeta_1$. \square

Finally, we have the following

Corollary 4. *The multiplication (8.1) extends the chain-cochain complex (7.13)–(7.15) structure for all products $C_m^k(V, \mathcal{W}) \times C_{m'}^n(V, \mathcal{W})$, $k, n \geq 0$, $m, m' \geq 0$. \square*

Corollary 5. *The product (8.1) and the product operator (7.12) endow the space $C_m^k(V, \mathcal{W}) \times C_{m'}^n(V, \mathcal{W})$, $k, n \geq 0$, $m, m' \geq 0$, with the structure of a bi-graded differential algebra $\mathcal{G}(V, \mathcal{W}, \cdot, \delta_m^n)$. \square*

9. EXAMPLE: EXCEPTIONAL COMPLEX

In addition to the complex $(C_m^n(V, \mathcal{W}), \delta_m^n)$ provided by (7.13)–(7.15), there exists an exceptional short complex $(C_{ex}^2(V, \mathcal{W}), \delta_{ex}^2)$. In [10] we have

Lemma 3. *For $n = 2$, there exists a subspace $C_{ex}^0(V, \mathcal{W})$*

$$C_m^2(V, \mathcal{W}) \subset C_{ex}^2(V, \mathcal{W}) \subset C_0^2(V, \mathcal{W}),$$

for all $m \geq 1$, with the action of coboundary operator δ_m^2 defined. \square

Let us recall some facts about the exceptional complex. Consider the space $C_0^2(V, \mathcal{W})$. It consist of \mathcal{W}_{z_1, z_2} -elements with zero vertex operators composable. The space $C_0^2(V, \mathcal{W})$ contains elements of \mathcal{W}_{z_1, z_2} so that the action of δ_0^2 is zero. Nevertheless, as for $\mathcal{J}_m^n(\Phi)$ in (7.6), Definition 9, let us consider sum of projections

$$P_r : \mathcal{W}_{z_i, z_j} \rightarrow W_r,$$

for $r \in \mathbb{C}$, and $(i, j) = (1, 2), (2, 3)$, so that the condition (7.6) is satisfied for some elements similar to the action (7.6) of δ_0^2 . Separating the first two and the second two summands in (7.12), we find that for a subspace of $C_0^2(V, \mathcal{W})$ (which we denote as $C_{ex}^2(V, \mathcal{W})$), for $v_1, v_2, v_3 \in V$, and arbitrary $w' \in W'$, $\zeta \in \mathbb{C}$, the following elements

$$\begin{aligned} & G_1(z_1, z_2, z_3) \\ &= \sum_{r \in \mathbb{C}} \left(\langle w', E_W^{(1)}(v_1, z_1; P_r(\mathcal{F}(v_2, z_2 - \zeta; v_3, z_3 - \zeta))) \rangle \right. \\ & \quad \left. + \langle w', \mathcal{F}(v_1, z_1; P_r(E_V^{(2)}(v_2, z_2 - \zeta; v_3, z_3 - \zeta; \mathbf{1}_V), \zeta)) \rangle \right) \\ &= \sum_{r \in \mathbb{C}} \left(\langle w', \omega_W(v_1, z_1) P_r(\mathcal{F}(v_2, z_2 - \zeta; v_3, z_3 - \zeta)) \rangle \right. \\ & \quad \left. + \langle w', \mathcal{F}(v_1, z_1; P_r(\omega_V(v_2, z_2 - \zeta) \omega_V(v_3, z_3 - \zeta) \mathbf{1}_V), \zeta) \rangle \right), \end{aligned} \tag{9.1}$$

and

$$\begin{aligned} & G_2(z_1, z_2, z_3) \\ &= \sum_{r \in \mathbb{C}} \left(\langle w', \mathcal{F}(P_r(E_V^{(2)}(v_1, z_1 - \zeta, v_2, z_2 - \zeta; \mathbf{1}_V)), \zeta; v_3, z_3) \rangle \right. \\ & \quad \left. + \langle w', E_{WV}^{W;(1)}(P_r(\mathcal{F}(v_1, z_1 - \zeta; v_2, z_2 - \zeta), \zeta; v_3, z_3)) \rangle \right) \\ &= \sum_{r \in \mathbb{C}} \left(\langle w', \mathcal{F}(P_r(\omega_V(v_1, z_1 - \zeta) \omega_V(v_2, z_2 - \zeta) \mathbf{1}_V), \zeta; v_3, z_3) \rangle \right. \\ & \quad \left. + \langle w', \omega_V(v_3, z_3) P_r(\mathcal{F}(v_1, z_1 - \zeta; v_2, z_2 - \zeta)) \rangle \right), \end{aligned} \tag{9.2}$$

are absolutely convergent in the regions

$$\begin{aligned} & |z_1 - \zeta| > |z_2 - \zeta|, \\ & |z_2 - \zeta| > 0, \\ & |\zeta - z_3| > |z_1 - \zeta|, \\ & |z_2 - \zeta| > 0, \end{aligned}$$

where z_i , $1 \leq i \leq 3$. These functions can be analytically extended to rational form-valued functions in z_1 and z_2 with the only possible poles at $z_1, z_2 = 0$, and $z_1 = z_2$. Note that (9.1) and (9.2) constitute the first two and the last two terms of (7.12) correspondingly. According to Proposition 7 (cf. Appendix 7.2), $C_m^2(V, \mathcal{W})$ is a subspace of $C_{ex}^2(V, \mathcal{W})$, for $m \geq 0$, and $\mathcal{F} \in C_m^2(V, \mathcal{W})$ are composable with m vertex operators. Then we have

Definition 12. The coboundary operator

$$\delta_{ex}^2 : C_{ex}^2(V, \mathcal{W}) \rightarrow C_0^3(V, \mathcal{W}), \quad (9.3)$$

is defined by

$$\begin{aligned} \delta_{ex}^2 \mathcal{F} &= \langle w', \omega_W(v_1, z_1) \mathcal{F}(v_2, z_2; v_3, z_3) \rangle \\ &- \langle w', \mathcal{F}(\omega_V(v_1, z_1) \omega_V(v_2, z_2) \mathbf{1}_V; v_3, z_3) \rangle \\ &+ \langle w', \mathcal{F}(v_1, z_1; \omega_V(v_2, z_2) \omega_V(v_3, z_3) \mathbf{1}_V) \rangle \\ &+ \langle w', \omega_W(v_3, z_3) \mathcal{F}(v_1, z_1; v_2, z_2) \rangle, \end{aligned} \quad (9.4)$$

for arbitrary $w' \in W'$, $\mathcal{F} \in C_{ex}^2(V, \mathcal{W})$, $(v_1, v_2, v_3) \in V$ and $(z_1, z_2, z_3) \in F_3\mathcal{C}$.

In [10] we also find

Proposition 14. *The operator (9.4) provides the chain-cochain complex*

$$\begin{aligned} &\delta_{ex}^2 \circ \delta_2^1 = 0, \\ 0 \longrightarrow C_3^0(V, \mathcal{W}) \xrightarrow{\delta_3^0} C_2^1(V, \mathcal{W}) \xrightarrow{\delta_2^1} C_{ex}^2(V, \mathcal{W}) \xrightarrow{\delta_{ex}^2} C_0^3(V, \mathcal{W}) \longrightarrow 0. \end{aligned} \quad (9.5)$$

□

Since

$$\delta_2^1 C_2^1(V, \mathcal{W}) \subset C_1^2(V, \mathcal{W}) \subset C_{ex}^2(V, \mathcal{W}),$$

the second formula follows from the first one, and

$$\delta_{ex}^2 \circ \delta_2^1 = \delta_1^2 \circ \delta_2^1 = 0.$$

For elements of the spaces $C_{ex}^2(V, \mathcal{W})$ we have the following

Corollary 6. *The product of elements of the spaces $C_{ex}^2(V, \mathcal{W})$ and $C_m^n(V, \mathcal{W})$ is given by (8.1),*

$$\cdot_\epsilon : C_{ex}^2(V, \mathcal{W}) \times C_m^n(V, \mathcal{W}) \rightarrow C_m^{n+2}(V, \mathcal{W}), \quad (9.6)$$

and, in particular,

$$\cdot_\epsilon : C_{ex}^2(V, \mathcal{W}) \times C_{ex}^2(V, \mathcal{W}) \rightarrow C_0^4(V, \mathcal{W}).$$

Proof. The fact that the number of formal parameters is $n+2$ in the product (8.1) follows from Proposition (4). Consider the product (8.1) for $C_{ex}^2(V, \mathcal{W})$ and $C_m^n(V, \mathcal{W})$. It is clear that, similar to considerations of the proof of Proposition 12, the total number m of vertex operators the product \mathcal{F} is composable to remains the same. □

9.1. An application to cohomology invariants of vertex algebras. In this Subsection we give an explicit application of the product constructed in Section 5 to cohomology invariants of vertex algebras by using the results of [14]. Let us define a further product of a pair of elements of the spaces $C_m^k(V, \mathcal{W})$ and $C_{m'}^n(V, \mathcal{W})$, suitable for the formulation of cohomology invariants. Consider the forms $\mathcal{F}(v_1, z_1; \dots; v_n, z_n) \in C_m^k(V, \mathcal{W})$, $\tilde{\mathcal{F}}(v_{k+1}, z_{k+1}; \dots; v_{k+n}, z_{k+n}) \in C_{m'}^n(V, \mathcal{W})$, with r common vertex algebra elements (and, correspondingly, r formal variables), and t common vertex operators mappings Φ and Ψ are composable with. Define an extra product of \mathcal{F} and $\tilde{\mathcal{F}}$, $\mathcal{F} \cdot \tilde{\mathcal{F}} : V^{\otimes(k+n-r)} \rightarrow \mathcal{W}_{z_1, \dots, z_{k+n-r}}$, $\mathcal{F} \cdot \tilde{\mathcal{F}} = [\mathcal{F}, \cdot_\epsilon \tilde{\mathcal{F}}] = \mathcal{F} \cdot_\epsilon \tilde{\mathcal{F}} - \tilde{\mathcal{F}} \cdot_\epsilon \mathcal{F}$, where

brackets denote the ordinary commutator on $\mathcal{W}_{z_1, \dots, z_{k+n-r}}$. Due to the properties of the maps $\mathcal{F} \in C_m^k(V, \mathcal{W})$ and $\tilde{\mathcal{F}} \in C_{m'}^n(V, \mathcal{W})$, the product $\Phi \cdot \Psi$ belongs to the space $C_{m+m'-t}^{k+n-r}(V, \mathcal{W})$. For $k = n$ and $\tilde{\mathcal{F}}(v_{n+1}, z_{n+1}; \dots; v_{2n}, z_{2n}) = \mathcal{F}(v_1, z_1; \dots; v_n, z_n)$, and we obtain from (8.1) that $\mathcal{F}(v_1, z_1; \dots; v_n, z_n) \cdot \mathcal{F}(v_1, z_1; \dots; v_n, z_n) = 0$. Let us now introduce cohomology classes associated to grading-restricted vertex algebras. Usually, (e.g., [2, 6, 14]) cohomology classes are introduced by means of an extra condition (in particular, the orthogonality condition) applied to differential forms, and leading to an integrability condition. One calls a map $\mathcal{F} \in C_k^n(V, \mathcal{W})$, closed if $\delta_k^n \mathcal{F} = 0$. For $k \geq 1$, one calls it exact if there exists $\tilde{\mathcal{F}} \in C_{k-1}^{n+1}(V, \mathcal{W})$, such that $\tilde{\mathcal{F}} = \delta_k^n \mathcal{F}$. For $\mathcal{F} \in C_k^n(V, \mathcal{W})$ we call the cohomology class of mappings $[\mathcal{F}]$ the set of all closed forms that differ from \mathcal{F} by an exact mapping, i.e., for $\Lambda \in C_{k+1}^{n-1}(V, \mathcal{W})$, $[\mathcal{F}] = \mathcal{F} + \delta_{k+1}^{n-1} \Lambda$. In contrast to [14], our cohomology class is a functional of $v \in V$. That means that the actual functional form of $\mathcal{F}(v, z)$ (and, therefore, $\langle w', \mathcal{F} \rangle$, for $w' \in W'$) varies with various choices of $v \in V$. Under a natural extra condition, the double complexes (7.15) and (9.5) allow us to establish relations among elements of $C_m^n(V, \mathcal{W})$ -spaces. By analogy with the notion of the integrability for differential forms [6], we use here the notion of the orthogonality for the spaces of a complex. For the complexes (7.15) and (9.5) let us require that for a pair of complex spaces $C_m^k(V, \mathcal{W})$ and $C_{m'}^n(V, \mathcal{W})$, there exist subspaces $\tilde{C}_m^k(V, \mathcal{W}) \subset C_m^k(V, \mathcal{W})$, $\tilde{C}_{m'}^n(V, \mathcal{W}) \subset C_{m'}^n(V, \mathcal{W})$, such that for all $\mathcal{F} \in \tilde{C}_m^k(V, \mathcal{W})$ and all $\tilde{\mathcal{F}} \in \tilde{C}_{m'}^n(V, \mathcal{W})$, $\mathcal{F} \cdot \delta_m^n \tilde{\mathcal{F}} = 0$. Namely, \mathcal{F} is supposed to be orthogonal to $\delta_m^n \tilde{\mathcal{F}}$ with respect to the new product. We call this the orthogonality condition for mappings of the double complexes (7.15) and (9.5). Note that in the case of differential forms considered on a smooth manifold, the Frobenius theorem for a distribution provides the orthogonality condition [6]. The fact that both sides of the orthogonality condition belong to the same double complex space, applies limitations to possible combinations of (k, m) and (n, m') . Analogously to the proof given in [14], it turns out that the orthogonality condition applied to elements of complexes (7.15) and (9.5) gives non-vanishing cohomology classes $[(\delta_m^n \mathcal{F}) \cdot \mathcal{F}]$ independent on the choice of $\mathcal{F} \in C_m^n(V, \mathcal{W})$, in particular, for $(n, m) = (1, 2), (0, 3), (1, t), 0 \leq t \leq 2$.

ACKNOWLEDGMENT

The author is supported by the Institute of Mathematics of the Academy of Sciences of the Czech Republic (RVO 67985840). I would like to thank two unanimous referees for very valuable comments and corrections.

10. APPENDIX: GRADING-RESTRICTED VERTEX ALGEBRAS AND THEIR MODULES

In this Section, following [10], we recall basic properties of grading-restricted vertex algebras and their grading-restricted generalized modules. We work over the base field \mathbb{C} of complex numbers.

Definition 13. A vertex algebra $(V, Y_V, \mathbf{1}_V)$, (cf. [13]), consists of a \mathbb{Z} -graded complex vector space

$$V = \coprod_{n \in \mathbb{Z}} V_{(n)}, \quad \dim V_{(n)} < \infty,$$

for each $n \in \mathbb{Z}$, and a linear map

$$Y_V : V \rightarrow \text{End}(V)[[z, z^{-1}]],$$

for a formal parameter z , and a distinguished vector $\mathbf{1}_V \in V$. The evaluation of Y_V on $v \in V$ is the vertex operator

$$Y_V(v) \equiv Y_V(v, z) = \sum_{n \in \mathbb{Z}} v(n) z^{-n-1}, \quad (10.1)$$

with components $(Y_V(v))_n = v(n) \in \text{End}(V)$, where $Y_V(v, z)\mathbf{1}_V = v + O(z)$. The vertex operators satisfy the vacuum axiom (the identity property)

$$Y_V(\mathbf{1}_V, z) = \text{Id}_V,$$

and the locality axiom

$$(z - w)^N [Y_V(u_1, z), Y_V(u_2, w)] = 0,$$

for some $N \gg 0$, all $u_1, u_2 \in V$.

Definition 14. A grading-restricted vertex algebra satisfies the following conditions:

- (1) Grading-restriction condition: $V_{(n)}$ is finite dimensional for all $n \in \mathbb{Z}$, and $V_{(n)} = 0$ for $n \ll 0$;
- (2) Lower-truncation condition: For $u, v \in V$, $Y_V(u, z)v$ contains only finitely many negative power terms, that is,

$$Y_V(u, z)v \in V((z)),$$

(the space of formal Laurent series in z with coefficients in V);

- (3) Creation property: For $u \in V$,

$$Y_V(u, z)\mathbf{1}_V \in V[[z]],$$

and

$$\lim_{z \rightarrow 0} Y_V(u, z)\mathbf{1}_V = u;$$

- (4) Duality: For $u_1, u_2, v \in V$,

$$v' \in V' = \prod_{n \in \mathbb{Z}} V'_{(n)},$$

where $V'_{(n)}$ denotes the dual vector space to $V_{(n)}$ and $\langle \cdot, \cdot \rangle$ the evaluation pairing $V' \otimes V \rightarrow \mathbb{C}$, the series

$$\langle v', Y_V(u_1, z_1)Y_V(u_2, z_2)v \rangle, \quad (10.2)$$

$$\langle v', Y_V(u_2, z_2)Y_V(u_1, z_1)v \rangle, \quad (10.3)$$

$$\langle v', Y_V(Y_V(u_1, z_1 - z_2)u_2, z_2)v \rangle, \quad (10.4)$$

are absolutely convergent in the regions

$$|z_1| > |z_2| > 0,$$

$$|z_2| > |z_1| > 0,$$

$$|z_2| > |z_1 - z_2| > 0,$$

respectively, to a common rational function in z_1 and z_2 with the only possible poles at $z_1 = 0 = z_2$ and $z_1 = z_2$;

- (5) $L_V(0)$ -bracket formula: Let $L_V(0) : V \rightarrow V$, be defined by

$$L_V(0)v = nv, \quad n = \text{wt}(v),$$

for $v \in V_{(n)}$. Then

$$[L_V(0), Y_V(v, z)] = Y_V(L_V(0)v, z) + z \frac{d}{dz} Y_V(v, z), \quad (10.5)$$

for $v \in V$.

- (6) $L_V(-1)$ -derivative property: Let

$$L_V(-1) : V \rightarrow V,$$

be the operator given by

$$L_V(-1)v = \text{Res}_z z^{-2} Y_V(v, z) \mathbf{1}_V = Y_{(-2)}(v) \mathbf{1}_V,$$

where $Y_{(-2)}(v)$ is the (-2) -th vertex algebra mode of the vertex operator $Y(v, z)$ for $v \in V$. Then for $v \in V$,

$$\frac{d}{dz} Y_V(u, z) = Y_V(L_V(-1)u, z) = [L_V(-1), Y_V(u, z)]. \quad (10.6)$$

In addition to that, we recall here the following definition (cf. [1]):

Definition 15. A grading-restricted vertex algebra V is called conformal of central charge $c \in \mathbb{C}$, if there exists a non-zero conformal vector (Virasoro vector) $\omega \in V_{(2)}$ such that the corresponding vertex operator

$$Y_V(\omega, z) = \sum_{n \in \mathbb{Z}} L_V(n) z^{-n-2},$$

is determined by modes of Virasoro algebra $L_V(n) : V \rightarrow V$ satisfying

$$[L_V(m), L_V(n)] = (m-n)L_V(m+n) + \frac{c}{12}(m^3 - m)\delta_{m+n,0} \text{Id}_V.$$

10.1. Grading-restricted generalized V -module.

Definition 16. A grading-restricted generalized V -module is a vector space W equipped with a vertex operator map

$$\begin{aligned} Y_W : V \otimes W &\rightarrow W[[z, z^{-1}]], \\ u \otimes w &\mapsto Y_W(u, w) \equiv Y_W(u, z)w = \sum_{n \in \mathbb{Z}} (Y_W)_n(u, w) z^{-n-1}, \end{aligned}$$

and linear operators $L_W(0)$ and $L_W(-1)$ on W satisfying the following conditions:

- (1) Grading-restriction condition: The vector space W is \mathbb{C} -graded, that is,

$$W = \prod_{\alpha \in \mathbb{C}} W_{(\alpha)},$$

such that $W_{(\alpha)} = 0$ when the real part of α is sufficiently negative;

- (2) Lower-truncation condition: For $u \in V$ and $w \in W$, $Y_W(u, z)w$ contains only finitely many negative power terms, that is, $Y_W(u, z)w \in W((z))$;

(3) Identity property: Let Id_W be the identity operator on W . Then

$$Y_W(\mathbf{1}_V, z) = \text{Id}_W;$$

(4) Duality: For $u_1, u_2 \in V$, $w \in W$,

$$w' \in W' = \prod_{n \in \mathbb{Z}} W_{(n)}^*,$$

W' denotes the dual V -module to W and $\langle \cdot, \cdot \rangle$ their evaluation pairing, the series

$$\langle w', Y_W(u_1, z_1) Y_W(u_2, z_2) w \rangle, \quad (10.7)$$

$$\langle w', Y_W(u_2, z_2) Y_W(u_1, z_1) w \rangle, \quad (10.8)$$

$$\langle w', Y_W(Y_V(u_1, z_1 - z_2) u_2, z_2) w \rangle, \quad (10.9)$$

are absolutely convergent in the regions

$$|z_1| > |z_2| > 0,$$

$$|z_2| > |z_1| > 0,$$

$$|z_2| > |z_1 - z_2| > 0,$$

respectively, to a common rational function in z_1 and z_2 with the only possible poles at $z_1 = 0 = z_2$ and $z_1 = z_2$.

(5) $L_W(0)$ -bracket formula: For $v \in V$,

$$[L_W(0), Y_W(v, z)] = Y_W(L_V(0)v, z) + z \frac{d}{dz} Y_W(v, z);$$

(6) $L_W(0)$ -grading property: For $w \in W_{(\alpha)}$, there exists $N \in \mathbb{Z}_+$ such that

$$(L_W(0) - \alpha)^N w = 0; \quad (10.10)$$

(7) $L_W(-1)$ -derivative property: For $v \in V$,

$$\frac{d}{dz} Y_W(u, z) = Y_W(L_V(-1)u, z) = [L_W(-1), Y_W(u, z)]. \quad (10.11)$$

The translation property of vertex operators

$$Y_W(u, z) = e^{-z' L_W(-1)} Y_W(u, z + z') e^{z' L_W(-1)}, \quad (10.12)$$

for $z' \in \mathbb{C}$, follows from (10.11). For $a \in \mathbb{C}$, the conjugation property with respect to the grading operator $L_W(0)$ is given by

$$a^{L_W(0)} Y_W(v, z) a^{-L_W(0)} = Y_W(a^{L_W(0)} v, az). \quad (10.13)$$

For $v \in V$, and $w \in W$, the intertwining operator

$$\begin{aligned} Y_{WV}^W : V &\rightarrow W, \\ v &\mapsto Y_{WV}^W(w, z)v, \end{aligned} \quad (10.14)$$

is defined by

$$Y_{WV}^W(w, z)v = e^{z L_W(-1)} Y_W(v, -z)w. \quad (10.15)$$

10.2. Group of automorphisms of formal parameters. Assume that W is a quasi-conformal grading-restricted vertex algebra V -module. Let us recall some further facts from [1] relating generators of Virasoro algebra with the group of automorphisms in complex dimension one. Let us represent an element of $\text{Aut}_z \mathcal{O}^{(1)}$ by the map

$$z \mapsto \rho = \rho(z), \quad (10.16)$$

given by the power series

$$\rho(z) = \sum_{k \geq 1} a_k z^k, \quad (10.17)$$

$\rho(z)$ can be represented in an exponential form

$$f(z) = \exp \left(\sum_{k > -1} \beta_k z^{k+1} \partial_z \right) (\beta_0)^{z \partial_z} .z, \quad (10.18)$$

where we express $\beta_k \in \mathbb{C}$, $k \geq 0$, through combinations of a_k , $k \geq 1$. A representation of Virasoro algebra modes in terms of differential operators is given by [13]

$$L_W(m) \mapsto -\zeta^{m+1} \partial_\zeta, \quad (10.19)$$

for $m \in \mathbb{Z}$. By expanding (10.18) and comparing to (12.1) we obtain a system of equations which, can be solved recursively for all β_k . In [1], $v \in V$, they derive the formula

$$[L_W(n), Y_W(v, z)] = \sum_{m \geq -1} \frac{1}{(m+1)!} (\partial_z^{m+1} z^{m+1}) Y_W(L_V(m)v, z), \quad (10.20)$$

of a Virasoro generator commutation with a vertex operator. Given a vector field

$$\beta(z) \partial_z = \sum_{n \geq -1} \beta_n z^{n+1} \partial_z, \quad (10.21)$$

which belongs to local Lie algebra of $\text{Aut} \mathcal{O}^{(1)}$, one introduces the operator

$$\beta = - \sum_{n \geq -1} \beta_n L_W(n).$$

We conclude from (10.21) with the following

Lemma 4.

$$[\beta, Y_W(v, z)] = - \sum_{m \geq -1} \frac{1}{(m+1)!} (\partial_z^{m+1} \beta(z)) Y_W(L_V(m)v, z). \quad (10.22)$$

Here we introduce the following definitions.

Definition 17. We call a grading-restricted vertex algebra quasi-conformal if it carries an action of $\text{Der} \mathcal{O}^{(n)}$ such that commutation formula (10.22) holds for any $v \in V$, and $z = z_j$, $1 \leq j \leq n$, the element $L_V(-1) = -\partial_z$ acts as the translation operator

$$L_V(0) = -z \partial_z,$$

acts semi-simply with integral eigenvalues, and the Lie subalgebra $\text{Der}_+ \mathcal{O}^{(n)}$ acts locally nilpotently.

Definition 18. A vector A which belongs to a quasi-conformal grading-restricted vertex algebra V is called primary of conformal dimension $\Delta(A) \in \mathbb{Z}_+$ if

$$\begin{aligned} L_V(k)A &= 0, \quad k > 0, \\ L_W(0)A &= \Delta(A)A. \end{aligned}$$

The formula (10.22) is used in [1] in order to prove invariance of vertex operators multiplied by conformal weight differentials in case of primary states, and in generic case.

Let us give some further definitions:

Definition 19. A conformal grading-restricted vertex algebra is a conformal vertex algebra V , such that it module W is equipped with an action of the Virasoro algebra and hence its Lie subalgebra $\text{Der}_0 \mathcal{O}^{(n)}$ given by the Lie algebra of $\text{Aut } \mathcal{O}^{(n)}$.

Definition 20. A grading-restricted vertex algebra V -module W is called quasi-conformal if it carries an action of local Lie algebra of $\text{Aut } \mathcal{O}$ such that commutation formula (10.22) holds for any $v \in V$, the element $L_W(-1) = -\partial_z$, as the translation operator T ,

$$L_W(0) = -z\partial_z,$$

acts semi-simply with integral eigenvalues, and the Lie subalgebra of the positive part of local Lie algebra of $\text{Aut } \mathcal{O}^{(n)}$ acts locally nilpotently.

Recall [1] the exponential form $f(\zeta)$ (10.18) of the coordinate transformation (10.16) $\rho(z) \in \text{Aut } \mathcal{O}^{(1)}$. A quasi-conformal vertex algebra has the formula (10.22), thus it is possible by using the identification (10.19), to introduce the linear operator representing $f(\zeta)$ (10.18) on $\mathcal{W}_{z_1, \dots, z_n}$,

$$P(f(\zeta)) = \exp \left(\sum_{m>0} (m+1) \beta_m L_V(m) \right) \beta_0^{L_W(0)}, \quad (10.23)$$

(note that we have a different normalization in it). In [1] it was shown that the action of an operator similar to (10.23) on a vertex algebra element $v \in V_n$ contains finitely many terms, and subspaces

$$V_{\leq m} = \bigoplus_{n \geq K}^m V_n,$$

are stable under all operators $P(f)$, $f \in \text{Aut } \mathcal{O}^{(1)}$. In [1] they proved the following

Lemma 5. *The assignment*

$$f \mapsto P(f),$$

defines a representation of $\text{Aut } \mathcal{O}^{(1)}$ on V ,

$$P(f_1 * f_2) = P(f_1) P(f_2),$$

which is the inductive limit of the representations $V_{\leq m}$, $m \geq K$ with some K for some product $$ of f_1 and f_2 .*

Similarly, (10.23) provides a representation operator on $\mathcal{W}_{z_1, \dots, z_n}$.

10.3. Non-degenerate invariant bilinear form on V . In this Subsection we recall [20] the notion of a non-degenerate invariant bilinear form defined on V . The subalgebra

$$\{L_V(-1), L_V(0), L_V(1)\} \cong SL(2, \mathbb{C}),$$

associated with Möbius transformations on z naturally acts on V , (cf., e.g. [13]). In particular, for $\lambda \in \mathbb{C}$,

$$\gamma_\lambda = \begin{pmatrix} 0 & \lambda \\ -\lambda & 0 \end{pmatrix} : z \mapsto w = -\frac{\lambda^2}{z}, \quad (10.24)$$

is generated by

$$T_\lambda = \exp(\lambda L_V(-1)) \exp(\lambda^{-1} L_V(1)) \exp(\lambda L_V(-1)),$$

where

$$T_\lambda Y(u, z) T_\lambda^{-1} = Y\left(\exp\left(-\frac{z}{\lambda^2} L_V(1)\right) \left(-\frac{z}{\lambda}\right)^{-2L_V(0)} u, -\frac{\lambda^2}{z}\right). \quad (10.25)$$

In our considerations (cf. Appendix 11) of the Riemann sphere sewing, we use in particular, the Möbius map

$$z \mapsto z' = \epsilon/z,$$

associated with the sewing condition (11.4) with

$$\lambda = -\xi \epsilon^{\frac{1}{2}}, \quad (10.26)$$

and $\xi \in \{\pm\sqrt{-1}\}$. The adjoint vertex operator [3, 13] is defined by

$$Y^\dagger(u, z) = \sum_{n \in \mathbb{Z}} u^\dagger(n) z^{-n-1} = T_\lambda Y(u, z) T_\lambda^{-1}. \quad (10.27)$$

A bilinear form $\langle \cdot, \cdot \rangle_\lambda$ on V is invariant if for all $a, b, u \in V$,

$$\langle Y(u, z)a, b \rangle_\lambda = \langle a, Y^\dagger(u, z)b \rangle_\lambda, \quad (10.28)$$

i.e.,

$$\langle u(n)a, b \rangle_\lambda = \langle a, u^\dagger(n)b \rangle_\lambda.$$

Thus, it follows that

$$\langle L_V(0)a, b \rangle_\lambda = \langle a, L_V(0)b \rangle_\lambda, \quad (10.29)$$

so that

$$\langle a, b \rangle_\lambda = 0, \quad (10.30)$$

if $wt(a) \neq wt(b)$ for homogeneous a, b . One also finds

$$\langle a, b \rangle_\lambda = \langle b, a \rangle_\lambda,$$

and it is non-degenerate if and only if V is simple. Given any V basis $\{u^\alpha\}$ we define the dual V basis $\{\bar{u}^\beta\}$ where $\langle u^\alpha, \bar{u}^\beta \rangle_\lambda = \delta^{\alpha\beta}$.

11. APPENDIX: A SPHERE FORMED FROM SEWING OF TWO SPHERES

In this appendix we recall some facts from [20]. The matrix element for a number of vertex operators of a vertex algebra is usually associated [3,4,22] with a vertex algebra character on a sphere. We extrapolate this notion to the case of $\mathcal{W}_{z_1, \dots, z_n}$ spaces. In Section 5 we explained that a space $\mathcal{W}_{z_1, \dots, z_n}$ can be associated with a Riemann sphere with marked points. The product of two such spaces is then associated with a sewing of two such spheres with a number of marked points and extra points with local coordinates identified with formal parameters of $\mathcal{W}_{x_1, \dots, x_k}$ and $\mathcal{W}_{y_1, \dots, y_n}$. In order to supply an appropriate geometric construction for the product, we use the ϵ -sewing procedure (described in this Appendix) for two initial spheres to obtain a matrix element associated with (5.1).

Remark 7. In addition to the ϵ -sewing procedure of two initial spheres, one can alternatively use the self-sewing procedure [23] of the sphere to get the torus. Then by sending parameters to an appropriate limit by shrinking the genus to zero. As a result, one obtains again the sphere but with a different parameterization. In this paper we focus on the ϵ -formalism only.

In our particular case of \mathcal{W} -valued rational functions obtained from matrix elements (4.1) we take two Riemann spheres $\Sigma_a^{(0)}$, $a = 1, 2$, as two initial auxiliary spaces. The resulting space is formed by the sphere $\Sigma^{(0)}$ obtained by the sewing procedure of $\Sigma_a^{(0)}$. The formal parameters (x_1, \dots, x_k) and (y_1, \dots, y_n) are identified with local coordinates of k and n points on two initial spheres $\Sigma_a^{(0)}$, $a = 1, 2$ correspondingly. In the ϵ sewing procedure, some r points parameters among (p_1, \dots, p_k) may coincide with points among (p'_1, \dots, p'_n) when we identify the annuluses (11.3) below. This corresponds to the singular case of coincidence of r formal parameters.

Consider the sphere formed by sewing together two initial spheres in the sewing scheme referred to as the ϵ -formalism in [23]. Let $\Sigma_a^{(0)}$, $a = 1, 2$ be two initial spheres. Introduce a complex sewing parameter ϵ where

$$|\epsilon| \leq r_1 r_2,$$

Consider k distinct points on $p_i \in \Sigma_1^{(0)}$, $i = 1, \dots, k$, with local coordinates $(x_1, \dots, x_k) \in F_k \mathbb{C}$, and distinct points $p_j \in \Sigma_2^{(0)}$, $j = 1, \dots, n$, with local coordinates $(y_1, \dots, y_n) \in F_n \mathbb{C}$, with

$$\begin{aligned} |x_i| &\geq |\epsilon|/r_2, \\ |y_j| &\geq |\epsilon|/r_1. \end{aligned}$$

Choose a local coordinate $z_a \in \mathbb{C}$ on $\Sigma_a^{(0)}$ in the neighborhood of points $p_a \in \Sigma_a^{(0)}$, $a = 1, 2$. Consider the closed disks $|\zeta_a| \leq r_a$, and excise the disk

$$\{\zeta_a, |\zeta_a| \leq |\epsilon|/r_a\} \subset \Sigma_a^{(0)}, \quad (11.1)$$

to form a punctured sphere

$$\widehat{\Sigma}_a^{(0)} = \Sigma_a^{(0)} \setminus \{\zeta_a, |\zeta_a| \leq |\epsilon|/r_a\}.$$

We use the convention

$$\overline{1} = 2, \quad \overline{2} = 1. \quad (11.2)$$

Define the annulus

$$\mathcal{A}_a = \{\zeta_a, |\epsilon|/r_a \leq |\zeta_a| \leq r_a\} \subset \widehat{\Sigma}_a^{(0)}, \quad (11.3)$$

and identify \mathcal{A}_1 and \mathcal{A}_2 as a single region $\mathcal{A} = \mathcal{A}_1 \simeq \mathcal{A}_2$ via the sewing relation

$$\zeta_1 \zeta_2 = \epsilon. \quad (11.4)$$

In this way we obtain a genus zero compact Riemann surface

$$\Sigma^{(0)} = \left\{ \widehat{\Sigma}_1^{(0)} \setminus \mathcal{A}_1 \right\} \cup \left\{ \widehat{\Sigma}_2^{(0)} \setminus \mathcal{A}_2 \right\} \cup \mathcal{A}.$$

This the sphere form a suitable geometric model for the construction of a product of \mathcal{W} -valued rational forms in Section 5.

12. APPENDIX: A PROOF OF PROPOSITION 1

In this Appendix we give a proof of Proposition 10, namely, we prove that Definition 3 is independent of the choice of formal parameters. Let us first recall definitions required for that. Let

$$\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)} = \text{Aut}_{\mathbb{C}}[[z_1, \dots, z_n]],$$

be the group of formal automorphisms of n -dimensional formal power series algebra $\mathbb{C}[[z_1, \dots, z_n]]$.

Let W be a quasi-conformal module for a grading-restricted vertex algebra V . The \mathbb{Z} -grading on W is bounded from below,

$$W = \bigoplus_{k > k_0} W_k,$$

for some $k_0 \in \mathbb{Z}$. Since the vector fields $z^{k+1} \partial_z$ with $k \in \mathbb{N}$ acts on W as the operators $-L_W(k)$ of degree $-k$, the action of the Lie subalgebra $\text{Der}_+ \mathcal{O}^{(n)}$ is locally nilpotent. Furthermore, $z \partial_z$ acts as the grading operator $L_W(0)$, which is diagonalizable with integral eigenvalues. Thus, the action of $\text{Der} \mathcal{O}^{(n)}$ on a conformal vertex algebra V can be exponentiated to an action of $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$.

We write an element of $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$ as

$$\begin{aligned} (z_1, \dots, z_n) \rightarrow \rho &= (\rho_1, \dots, \rho_n), \\ \rho_i &= \rho_i(z_1, \dots, z_n), \end{aligned}$$

for $i = 1, \dots, n$, where ρ_i are defined by elements of $\mathfrak{m} \in \mathcal{O}^{(n)}$

$$\rho_i(z_1, \dots, z_n) = \sum_{\substack{i_1 \geq 0, \dots, i_n \geq 0, \\ \sum_{j=1}^n i_j \geq 1}} a_{(i_1, \dots, i_k)} z_1^{i_1} \dots z_k^{i_k}, \quad a_{(i_1, \dots, i_k)} \in \mathbb{C}, \quad (12.1)$$

and the images of ρ_i , $i = 1, \dots, n$, in the finite dimensional \mathbb{C} -vector space $\mathfrak{m}/\mathfrak{m}^2$ are linearly independent. Let us denote

$$\begin{aligned} \mathbf{v} &= (v_1 \otimes \dots \otimes v_n), \\ \mathbf{z} &= (z_1, \dots, z_n), \\ w_i &= \rho_i(z_1, \dots, z_n), \\ \mathbf{w} &= (w_1, \dots, w_n). \end{aligned}$$

The natural object that turns to be invariant with respect to the action of the group $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$ is given by the matrix element of the n -vector

$$\langle w', \overline{\Phi}(\mathbf{v}, \mathbf{z} \, d\mathbf{z}) \rangle = \langle w', [\Phi(v_1, z_1 \, dz_{i(1)}; \dots; v_n, z_n \, dz_{i(n)})] \rangle, \quad (12.2)$$

containing n \mathcal{F} -entries, where $i(j)$ denotes the cyclic permutation of $(1, \dots, n)$ starting with j . In this paper we use (4.2) which is related to (12.4). Due to (4.8), (12.4) can be written in the form

$$\begin{aligned} \langle w', \overline{\Phi}(\mathbf{d}\mathbf{z} \, \mathbf{v})(\mathbf{z}) \rangle &= \langle w', \left[(dz_{i(j)})^{-L(0)w} \Phi \left(\left((dz_{i(j)})^{L_0^{(v)}} \mathbf{v} \right) (\mathbf{z}) \right) \right] \rangle \\ &= \langle w', \left[(dz_{i(j)})^{-L(0)w} \Phi \left((dz_{i(j)})^{\text{wt}(v_j)} \mathbf{v}(\mathbf{z}) \right) \right] \rangle, \end{aligned}$$

coherent with the one-dimensional case of [1] and containing $\text{wt}(v)$ -differentials. The idea to use torsors [1] is to represent the action of ρ (12.1) of the group $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$ on formal parameters \mathbf{z} in vectors (\mathbf{v}, \mathbf{z}) to the action by V -operators on vertex algebra states \mathbf{v} . Recall the standard representation of the Virasoro mode [13]

$$z_j^{m+1} \partial_{z_j} \mapsto -L_m, \quad m \in \mathbb{Z}.$$

In order to represent the action of the group $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$ on the variables (z_1, \dots, z_n) of $\overline{\mathcal{F}}$ (12.4) on (v_1, \dots, v_n) , we have to transfer (as in $n = 1$ case of [1]) to an exponential form of (12.1). The coefficients $\beta_{r_1, \dots, r_n}^{(j)} \in \mathbb{C}$ are recursively found [9] in terms of coefficients $a_{r_1, \dots, r_n}^{(i)}$ of (12.1). We introduce the linear operators

$$R(\rho_1, \dots, \rho_n) : V^{\otimes n} \rightarrow V^{\otimes n},$$

and define the action

$$\overline{\Phi}(\mathbf{v}, \mathbf{w} \, d\mathbf{w}) = R(\rho_1, \dots, \rho_n) \overline{\Phi}(\mathbf{v}, \mathbf{z} \, d\mathbf{w}). \quad (12.3)$$

Proof. Consider the vector

$$\overline{\Phi} = [\Phi(v_1, w_1 \, dw_{i(1)}; \dots; v_n, w_n \, dw_{i(n)})], \quad (12.4)$$

with primary (v_1, \dots, v_n) . Note that

$$dw_j = \sum_{i=1}^n \frac{\partial \rho_j}{\partial z_i} dz_i, \quad \partial_{z_i} \rho_j = \frac{\partial \rho_j}{\partial z_i}, \quad (12.5)$$

(as in [1], we skip the complex conjugated part $d\overline{z}_i$). By definition (12.3) of the action of $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$, and due to (12.5) by rewriting dw_i , we have

$$\begin{aligned} \overline{\Phi}(\mathbf{v}, \mathbf{w} \, d\mathbf{w}) &= R(\rho_1, \dots, \rho_n) [\Phi(v_1, z_1 \, dw_{i(1)}; \dots; v_n, z_n \, dw_{i(n)})] \\ &= R(\rho_1, \dots, \rho_n) \left[\Phi \left(v_1, z_1 \sum_{j=1}^n \partial_j \rho_{i(1)} \, dz_j; \dots; v_n, z_n \sum_{j=1}^n \partial_j \rho_{i(n)} \, dz_j \right) \right]. \end{aligned}$$

By using (4.3) and linearity of the mapping Φ , we obtain from the last equation

$$\overline{\Phi}(\mathbf{v}, \mathbf{w} \, d\mathbf{w}) = [\Phi(v_1, z_1 \, dz_{i(1)}; \dots; v_n, z_n \, dz_{i(n)})], \quad (12.6)$$

with

$$R(\rho_1, \dots, \rho_n) = \begin{bmatrix} \widehat{\partial}_J \rho_{i_1(I)} \\ \widehat{\partial}_J \rho_{i_2(I)} \\ \dots \\ \widehat{\partial}_J \rho_{i_n(I)} \end{bmatrix}. \quad (12.7)$$

The index operator J takes the value of index z_j of arguments in the vector (12.6), while the index operator I takes values of index of differentials dz_i in each entry of the vector $\overline{\Phi}$ (12.4). Thus, the index operator $i(I) = (i_1, \dots, i_n(I))$ is given by consequent cycling permutations of I . Taking into account the property (4.3), we define the operator

$$\widehat{\partial}_J \rho_a = \exp \left(- \sum_{(r_1 \dots r_n), \sum_{i=1}^n r_i \geq 1} r_J \beta_{r_1, \dots, r_n}^{(a)} \zeta_1^{r_1} \dots \zeta_J^{r_J} \dots \zeta_n^{r_n} L_{(W)}(-1) \right), \quad (12.8)$$

which contains index operators J as index of a dummy variable ζ_J turning into z_j , $j = 1, \dots, n$. (12.8) acts on each argument of mappings Φ in the vector $\overline{\Phi}$ (12.4). Due to the definition of a grading-restricted vertex algebra, the action of operators $R(\rho_1, \dots, \rho_n)$ for $i = 1, \dots, n$, on $v \in V$ results in a sum of finitely many terms. Similar to [1], for $n = 1$, one proves

Lemma 6. *The mappings $\rho(z_1, \dots, z_n) \mapsto R(\rho_1, \dots, \rho_n)$, for $i, j = 1, \dots, n$, define a representation of $\text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$ on $V^{\otimes n}$ by*

$$R(\rho \circ \rho') = R(\rho) R(\rho'),$$

for $\rho, \rho' \in \text{Aut}_{z_1, \dots, z_n} \mathcal{O}^{(n)}$.

We then conclude with

$$[\overline{\Phi}(v_1, z_1 dz_1; \dots; v_n, \dots, z_n dz_n)]. \quad (12.9)$$

Thus the vector $\overline{\Phi}$ (12.4) is invariant, i.e.,

$$\overline{\Phi}(\mathbf{v}, \mathbf{w} \, d\mathbf{w}) = \overline{\Phi}(\mathbf{v}, \mathbf{z} \, d\mathbf{z}). \quad (12.10)$$

Recall that the construction of the spaces $C_m^n(V, \mathcal{W})$ of the complex. It assumes that $\Phi \in C_m^n(V, \mathcal{W})$ is composable with m vertex operators. In one-dimensional complex case, [1] they proved that a vertex operator multiplied to the $\text{wt}(v_i)$ -power of the differential $Y_W(v_i, z_i) dz_i^{\text{wt}(v_i)}$ is invariant with respect to the action of the group $\text{Aut}_{z_i} \mathcal{O}^{(1)}$. Here we prove that $Y_W(v_i, z_i) dz_i^{\text{wt}(v_i)}$ is invariant with respect to the change of the local coordinates $z_i \mapsto w_i(z_1, \dots, z_n)$.

Let (z_1, \dots, z_n) be an open ball $D_{\mathbf{z}_0}^{(n)}$ of local formal coordinates around a fixed-value \mathbf{z}_0 of (z_1, \dots, z_n) . Define a $\text{wt}(v_i)$ -differential on $D_{\mathbf{z}_0}^{(n)}$ with values in $\text{End}(W)_{\mathbf{z}_0}$ as follows: identify $\text{End}(W)_{\mathbf{z}_0}$ with $\text{End} W$ using the formal parameters (z_1, \dots, z_n) , and set

$$\omega_{i,x} = Y_W(v_i, z_i) dz_i^{\text{wt}(v_i)}.$$

Let $(w_1, \dots, w_n) = (\rho_1(z_1, \dots, z_n), \dots, \rho_n(z_1, \dots, z_n))$, be another set of formal parameters on an n -dimensional ball $D_{\mathbf{z}_0}^{(n)}$. Let us express the set of $\text{wt}(v_i)$ -differentials on $D_{\mathbf{z}_0}^{(n), \times}$

$$Y_W(v_i, w_i) dw_i^{\text{wt}(v_i)},$$

$i = 1, \dots, n$, in terms of the parameters (z_1, \dots, z_n) . We would like to show that it coincides with the set of $\text{wt}(v_i)$ -differentials $Y_W(v_i, w_i) dz_i^{\text{wt}(v_i)}$.

Recall the notion of torsors (Section 10). Consider a vector $(v_i, z_1, \dots, z_n) \in W_{\mathbf{z}_0}$ with $v_i \in V$. Then the same vector equals

$$(R_i^{-1}(\rho_1, \dots, \rho_n) v_i, w_1, \dots, w_n),$$

i.e., it is identified with

$$R_i^{-1}(\rho_1, \dots, \rho_n) v_i \in V,$$

using the formal parameters (w_1, \dots, w_n) . Here $R_i(\rho_1, \dots, \rho_n)$ is an operator representing transformation of $z_i \rightarrow w_i$, as an action on V . Therefore if we have an operator on $W_{\mathbf{z}_0}$ which is equal to a $\text{Aut } \mathcal{O}^{(n)}$ -torsor S under the identification $\text{End } W_{\mathbf{z}_0} \in \text{End } W$ using the formal parameters (w_1, \dots, w_n) , then this operator equals

$$R_i(\rho_1, \dots, \rho_n) S R_i^{-1}(\rho_1, \dots, \rho_n),$$

under the identification $\text{End } W_{\mathbf{z}_0} \in \text{End } W^{(i)}$ using the combined parameters (v_i, z_1, \dots, z_n) . Thus, in terms of (v_i, z_1, \dots, z_n) , the differential $Y_W(v_i, w_i) dw_i^{\text{wt}(v_i)}$ becomes

$$Y_W(v_i, z_i) dz_i^{\text{wt}(v_i)} = R_i(\rho) Y_W(v_i, \rho(z_1, \dots, z_n)) R_i^{-1}(\rho) dw_i^{\text{wt}(v_i)}.$$

According to Definition (10), elements Φ are composable with m vertex operators. Thus we see that (12.4) is a canonical object of $C_m^n(V, \mathcal{W})$. We have proved that elements of the spaces $C_m^n(V, \mathcal{W})$ are independent on the choice of formal parameters. \square

REFERENCES

- [1] Ben-Zvi D., E. Frenkel E.: *Vertex algebras on algebraic curves*. American Mathematical Society, 2 edition, 2004.
- [2] Crainic, M., Moerdijk, I.: Čech-De Rham theory for leaf spaces of foliations. *Math. Ann.* 328 (2004), no. 1–2, 59–85.
- [3] Frenkel, I., Huang Y., Lepowsky, J.: On axiomatic approaches to vertex operator algebras and modules, *Memoirs American Math. Soc.* **104**, 1993.
- [4] Francesco, Ph., Mathieu, P., Senechal, D.: *Conformal Field Theory*. Graduate Texts in Contemporary Physics. 1997.
- [5] Fuks, D. B. : *Cohomology of infinite-dimensional Lie algebras*, Contemporary Soviet Mathematics, Consultant Bureau, New York, 1986.
- [6] Ghys, E.: L'invariant de Godbillon-Vey. *Seminaire Bourbaki*, 41-eme annee, n 706, S. M. F. Asterisque 177–178 (1989).
- [7] Gilroy, T., Tuite, M. P.: Genus Two Zhu Theory for Vertex Operator Algebras, arXiv:1511.07664.
- [8] Gilroy, T., Tuite, M. P.: Genus Two Virasoro Correlation Functions for Vertex Operator Algebras, arXiv:1612.02060.

- [9] Gradshteyn, I. S.; Ryzhik, I. M. Table of integrals, series, and products. Translated from the Russian. Eighth edition, Elsevier/Academic Press, Amsterdam, 2015. xlvii+1133 pp.
- [10] Huang, Y.-Zh. : A cohomology theory of grading-restricted vertex algebras, *Comm. Math. Phys.* 327 (2014), no. 1, 279–307.
- [11] Huang, Y.-Zh. : *Two-dimensional conformal geometry and vertex operator algebras*, Progress in Mathematics, Vol. 148, Birkhäuser, Boston, 1997.
- [12] Hurley, D. Tuite, M. P.: On the torus degeneration of the genus two partition function. *Internat. J. Math.* 24 (2013), no. 7, 3050056, 17 pp.
- [13] Kac V.: *Vertex Operator Algebras for Beginners*, University Lecture Series **10**, AMS, Providence 1998.
- [14] Kotschick, D.: Godbillon-Vey invariants for families of foliations. Symplectic and contact topology: interactions and perspectives (Toronto, ON/Montreal, QC, 2001), 131–144, Fields Inst. Commun., 35, Amer. Math. Soc., Providence, RI, 2003.
- [15] Mason, G., Tuite, M. P.: On genus two Riemann surfaces formed from sewn tori. *Comm. Math. Phys.* 270 (2007), no. 3, 587–634.
- [16] Mason, G., Tuite, M. P.: Vertex operators and modular forms. A window into zeta and modular physics, 183–278, *Math. Sci. Res. Inst. Publ.*, 57, Cambridge Univ. Press, Cambridge, 2010.
- [17] Mason, G., Tuite, M. P.: Free bosonic vertex operator algebras on genus two Riemann surfaces I. *Comm. Math. Phys.* 300 (2010), no. 3, 673–713.
- [18] Mason, G., Tuite, M. P.: Free bosonic vertex operator algebras on genus two Riemann surfaces II. Conformal field theory, automorphic forms and related topics, 183–225, *Contrib. Math. Comput. Sci.*, 8, Springer, Heidelberg, 2014
- [19] Qi, F.: Representation theory and cohomology theory of meromorphic open string vertex algebras, Ph.D. dissertation, (2018).
- [20] Tuite, M.P. , Zuevsky, A.: Genus two partition and correlation functions for fermionic vertex operator superalgebras I, *Comm. Math. Phys.* 306, 419–447 (2011).
- [21] Tuite, M.P. , Zuevsky, A.: Alexander The Szego kernel on a sewn Riemann surface. *Comm. Math. Phys.* 306 (2011), no. 3, 617–645.
- [22] Tsuchiya, A., Ueno, K., Yamada, Y.: Conformal field theory on universal family of stable curves with gauge symmetries, *Adv. Stud. Pure. Math.* **19** (1989), 459–566.
- [23] Yamada, A.: Precise variational formulas for abelian differentials. *Kodai Math.J.* **3** (1980) 114–143.
- [24] Zhu. Y.: Modular invariance of characters of vertex operator algebras. *J. Amer. Math. Soc.* 9(1), 237-302, (1996).

INSTITUTE OF MATHEMATICS, CZECH ACADEMY OF SCIENCES,, ZITNA 25, 11567, PRAGUE, CZECH REPUBLIC

Email address: zuevsky@yahoo.com