

The existence of homologically fibered links and solutions of some equations

Nozomu Sekino

Abstract

There is one generalization of fibered links in 3-manifolds, called homologically fibered links. It is known that the existence of a homologically fibered link whose fiber surface has a given homeomorphic type is determined by the first homology group and its torsion linking form of the ambient 3-manifold. In this paper, we interpret the existence of homologically fibered links with that of a solution of some equation, in terms of the first homology group and its torsion linking form or a surgery diagram of the ambient manifold. As an application, we compute the invariant $\text{hc}(\cdot)$, defined through homologically fibered knots, for 3-manifolds whose torsion linking forms represent a generator of linkings.

1 Introduction

It is known that every connected orientable closed 3-manifold M has a fibered link [1], i.e., a link L in M such that the complement of a small open neighborhood of L admits a structure of a connected orientable compact surface bundle over S^1 and that each boundary component of a fiber surface is a longitude of a component of L . Moreover, it is known that every connected orientable closed 3-manifold M has a fibered knot [9]. However, finding fibered links in a given 3-manifold is difficult in general. For example $\text{op}(M)$, defined as the minimal genus of fiber surfaces of fibered knots which M has, is a topological invariant which is difficult to calculate. There is one generalization of fibered links, called *homologically fibered links* [4] (defined in Definition 2.2 below). A homologically fibered link requests some Seifert surface such that the result of cut a 3-manifold along the surface is a homological product of a surface and an interval, whereas a fibered link requests some Seifert surface such that the result of cutting a 3-manifold along the surface is the product of a surface and an interval. Clearly, a fibered link is also a homologically fibered link, and $\text{hc}(M)$ [4], defined as the minimal genus of homological fiber surfaces of homologically fibered knots which M has gives a lower bound of $\text{op}(M)$. Homological products of surfaces (of a fixed homeomorphic type) and intervals are not merely the homological constraint for fiber structures, but they form a monoid by stacking, which contains the mapping class group as a submonoid (see [4]). This attracts attention as a generalization of the mapping class groups. Homological products of surfaces and intervals are studied using “clasper theory”. As a consequence of clasper theory, it is known that the existence of a homologically fibered link whose fiber is a given homeomorphic type in M depends on the first homology group of M with its torsion linking form. In this paper, we interpret the existence of a homologically fibered link whose fiber is a given homeomorphic type with the existence of a solution for some equation as follows.

As a convention, for an $m \times m$ -matrix A and an $n \times n$ -matrix B , $A \oplus B$ is an $(m+n) \times (m+n)$ -matrix whose (i, j) -entry is the same as the (i, j) -entry of A for $1 \leq i, j \leq m$, the $(m+i', m+j')$ -entry is the same as the (i', j') -entry of B for $1 \leq i', j' \leq n$, and the other entries are 0.

Let O_r be an $r \times r$ -zero-matrix, I_r an $r \times r$ -identity matrix, and we regard 0×0 -matrix (O_0 and I_0) as the identity element for \oplus .

Let P, Q, F_0^k, F_1^k and G_1^k be matrices as follows, where $a \geq 0$ and $k \geq 0$:

- P is a diagonal $a \times a$ -matrix whose (i, i) -entry is $p_i \in \mathbb{Z}$ for $a > 0$ or I_0 for $a = 0$.

- Q is a diagonal $a \times a$ -matrix whose (i, i) -entry is $-q_i \in \mathbb{Z}$ for $a > 0$ or I_0 for $a = 0$.
- F_0^0, F_1^0 and G_1^0 to be I_0 ,
and $F_0^k = \begin{pmatrix} 0 & 2^k \\ 2^k & 0 \end{pmatrix}$, $F_1^k = \begin{pmatrix} 2^{k+1} & -2^k \\ -3 \cdot 2^k & 2^{k+1} \end{pmatrix}$, $G_1^k = \begin{pmatrix} -1 & 0 \\ 0 & -3 \end{pmatrix}$ for $k > 0$.

Also let S and T be matrices as follows, where $r \geq 0, e_0 \geq 0, e_1 \geq 0, k_0 = k'_0 = 0, k_i \geq 1$ for $i \neq 0$, and $k'_j \geq 2$ for $j \neq 0$:

- $S = O_r \oplus P \oplus (\bigoplus_{0 \leq i \leq e_0} F_0^{k_i}) \oplus (\bigoplus_{0 \leq j \leq e_1} F_1^{k'_j})$.
- $T = I_r \oplus Q \oplus I_{2e_0} \oplus (\bigoplus_{0 \leq j \leq e_1} G_1^{k'_j})$.

As usual notation, $\Sigma_{g,n+1}$ denotes a connected orientable compact surface of genus g with $n + 1$ boundary components.

Theorem 1.1. *Let M be a connected closed oriented 3-manifold whose free part of the first homology group with integer coefficient is isomorphic to \mathbb{Z}^r and whose torsion linking form is equivalent to $(\bigoplus_{0 \leq l \leq a} A^{p_l}(q_l)) \oplus (\bigoplus_{0 \leq i \leq e_0} E_0^{k_i}) \oplus (\bigoplus_{0 \leq j \leq e_1} E_1^{k'_j})$, where a, e_0, e_1 are non-negative integers and notations $A^{p_l}(q_l), E_0^{k_i}, E_1^{k'_j}$ represent linkings, defined in Subsection 3.1.*

Suppose $(g, n) \neq (0, 0)$. Then M has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{g,n+1}$ if and only if there exist $(r + a + 2e_0 + 2e_1) \times (2g + n)$ -matrix of integer coefficients X and $(2g + n) \times (2g + n)$ - symmetric matrix of integer coefficients Y satisfying the following equation:

$$\begin{vmatrix} S & TX \\ X^t & Y + (\mathcal{E} \oplus O_n) \end{vmatrix} = \pm 1, \quad (1)$$

where \mathcal{E} is a $2g \times 2g$ -matrix whose $(2i - 1, 2i)$ -entry is 1 for every $1 \leq i \leq g$ and the others are 0.

Remark 1.1. For the case which is not covered by Theorem 1.1, it is known that a connected closed orientable 3-manifold M has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{0,1}$ if and only if M is an integral homology 3-sphere. See the definitions in the Subsection 2.1.

In general, it is not easy to compute the torsion linking form of a 3-manifold. Sometimes surgery diagrams of a 3-manifold may be easy to handle. A statement in terms of surgery diagrams is given in Proposition 5.1.

The rest of this paper is organized as follows: In Section 2, we recall the definition of homologically fibered links and the fact of their dependence on torsion linking forms. In Section 3, we recall three types of generators of linking forms and give rational homology 3-spheres whose torsion linking forms are the generators. In Section 4, we give a correspondence between surfaces in 3-manifolds and tuples of annuli and bands for Theorem 1.1. In Section 5, we give a proof of Theorem 1.1. In Section 6, we compute $\text{hc}(\cdot)$ for 3-manifolds whose torsion linking forms are two of three types of generators in Section 3.

Acknowledgements

The author would like to thank Takuya Sakasai and Yuta Nozaki for introducing him this subject and giving him many ingredients about it. He also is grateful to referee for his or her kindness, patience and correcting many mistakes in the proofs and the arguments.

2 Homologically fibered links

In this section, we review the definition of homologically fibered links and the fact that the existence of homologically fibered links depends on the torsion linking forms. In this paper, the homology groups are with integer coefficients.

2.1 Definitions

Definition 2.1. (homology cobordism, [3, Section 2.4])

A *homology cobordism* over $\Sigma_{g,n+1}$ is a triad $(X, \partial_+ X, \partial_- X)$, where X is an oriented connected compact 3-manifold and $\partial_+ X \cup \partial_- X$ is a partition of ∂X , and $\partial_\pm X$ are homeomorphic to $\Sigma_{g,n+1}$ satisfying:

- $\partial_+ X \cup \partial_- X = \partial X$.
- $\partial_+ X \cap \partial_- X = \partial(\partial_+ X)$.
- The induced maps $(i_\pm)_* : H_*(\partial_\pm X) \rightarrow H_*(X)$ are isomorphisms, where $i_\pm : \partial_\pm X \rightarrow X$ are the inclusions.

Note that the third condition is equivalent to the condition that i_\pm induce isomorphisms on $H_1(\cdot)$. Moreover by using the Poincaré-Lefschetz duality, we see that it is sufficient to require only one of i_+ and i_- to induce an isomorphism on $H_1(\cdot)$.

Definition 2.2. [4] Let L be a link in a closed orientable 3-manifold M . L is called *homologically fibered link* if there exists a Seifert surface F of L such that $(M \setminus F, F_+, F_-)$ is a homology cobordism over a surface homeomorphic to F , where F_\pm are the cut ends. In this situation, we call F a *homological fiber* of L .

Clearly every fibered link is a homologically fibered link since the former requests that the manifold cut opened by some Seifert surface to be the product of an interval and a surface and the latter requests it to be the homological product of an interval and a surface. By the observations above the Definition 2.2, for a Seifert surface F in a closed 3-manifold M to be a homological fiber, it is enough to check that push-ups (or push-downs) of oriented simple closed curves on F which form a basis of $H_1(F)$ also form a free basis of $H_1(M \setminus \text{Int}(F \times [0, 1]))$.

Remark 2.1. By definition, when a connected closed orientable 3-manifold M has a homologically fibered link with homological fiber F , $H_1(M)$ can be generated by $1 - \chi(F)$ elements, where $\chi(F)$ is the Euler number of F . Therefore the minimal order of generating sets for $H_1(M)$ imposes some constraint on homeomorphic types of homological fibers in M .

2.2 Dependence on the torsion linking forms

Definition 2.3. (Torsion linking form)

Let M be a connected closed oriented 3-manifold, $TH_1(M)$ the torsion part of $H_1(M)$. For every $a \in TH_1(M)$, there is a non-zero integer n such that na vanishes in $H_1(M)$, and we fix such an integer n_a of minimal absolute value. This a has a representative as oriented curves L_a in M such that the n_a -parallel copy of L_a bounds a surface S_a in M . The *torsion linking form* of M , $\psi_M : TH_1(M) \times TH_1(M) \rightarrow \mathbb{Q}/\mathbb{Z}$ maps (a, b) to $\frac{\langle S_a, L_b \rangle}{n_a} \bmod \mathbb{Z}$, where $\langle S_a, L_b \rangle$ represents the algebraic intersection number of S_a and L_b . It is known that ψ_M is well-defined and is non-singular, symmetric bilinear pairing, and that $\psi_{-M} = -\psi_M$, where $-M$ denotes the manifold homeomorphic to M with the opposite orientation.

There is the operation of 3-manifolds preserving the first homology groups and the torsion linking forms, called the ‘‘Borromean surgery’’ introduced in [8].

Definition 2.4. Let V be a standard handlebody of genus three in S^3 which contains 0-framed link as in Figure 1. Consider a compact orientable 3-manifold M and an embedding $f : V \rightarrow M$. Let N be a manifold obtained as a result of the Dehn surgery on M along the framed link in $f(V)$. This N is called the manifold obtained from M (and f) by a *Borromean surgery*.

About Borromean surgeries, the following facts are known:




Figure 1: A standard handlebody of genus three containing a 0-framed link

Fact 2.1. [5], [7] *Let X be a homology cobordism over $\Sigma_{g,n+1}$ and Y a manifold obtained from X by a Borromean surgery. Then Y is also a homology cobordism over $\Sigma_{g,n+1}$.*

Fact 2.2. [8] *Let M and N be closed oriented 3-manifolds. Then the following are equivalent:*

- *There is an isomorphism between $H_1(M)$ and $H_1(N)$ such that ψ_M and ψ_N are equivalent under this isomorphism.*
- *M is obtained from N by a finite sequence of Borromean surgeries.*

By the facts above, we can see the dependence on the first homology groups and the torsion linking forms about the existence of homologically fibered links:

Proposition 2.1. *Suppose that for connected closed oriented 3-manifolds M and N , there exists an isomorphism between their first homology groups such that their torsion linking forms are equivalent under the isomorphism. Then N has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{g,n+1}$ if and only if M has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{g,n+1}$.*

Proof. We prove the if part and note that it is enough. By Fact 2.2, there is a finite sequence of Borromean surgeries starting from M and ending at N . Since it is enough to prove when N is obtained from M by one Borromean surgery, we assume that. Suppose M has a homological fibered link with homological fiber F which is homeomorphic to $\Sigma_{g,n+1}$. Take a spine T of F and assume that F lies in a small neighborhood of T . Isotope F so that it is disjoint from the surgery link of the Borromean surgery. This can be done since T and the surgery link are graphs in a 3-manifold. Let F' be a surface, which is the image of F after the Borromean surgery. Since F is disjoint from the surgery link, a triad $(N \setminus F', F'^+, F'^-)$ is obtained from $(M \setminus F, F^+, F^-)$ by the Borromean surgery. Then by Fact 2.1, $(N \setminus F', F'^+, F'^-)$ is a homology cobordism over $\Sigma_{g,n+1}$. This implies that N has a homologically fibered link with homological fiber F' which is homeomorphic to $\Sigma_{g,n+1}$. \square

Note that if a connected closed oriented 3-manifold M has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{g,n+1}$, then $-M$ also has such a homologically fibered link. By Proposition 2.1, for a connected closed oriented 3-manifold M containing a homologically fibered link with homological fiber of a given homeomorphic type we can replace M with another one whose first homology group with the torsion linking form is the same as arbitrary one of M and $-M$.

3 Representatives with respect to the first homology groups and their torsion linking forms

In this section, at first we recall a part of the result of [12], which gives a generator of the semigroup of the linkings on finite abelian groups, and next we fix representatives of 3-manifolds with respect to

the first homology groups and their torsion linking forms. In fact, representatives are given in [6]. We give another representatives for our use. Note that they may coincide.

3.1 Linking pair

A *linking* is a pair (G, ψ) such that G is a finite abelian group and ψ is a non-singular, symmetric bilinear pairing $G \times G \rightarrow \mathbb{Q}/\mathbb{Z}$. Sometimes ψ is called a linking on G . We call two linkings (G, ψ) and (G', ψ') equivalent if there exists a group isomorphism between G and G' through which ψ and ψ' are equivalent. By fixing a basis of G , ψ is represented as a non-singular, symmetric matrix with coefficients in \mathbb{Q}/\mathbb{Z} , whose (i, j) -entry is the image of the i -th element and the j -th element of the basis under ψ . For two linkings (G_1, ψ_1) and (G_2, ψ_2) , define the product as $(G_1 \oplus G_2, \psi_1 \oplus \psi_2)$, where $\psi_1 \oplus \psi_2$ maps $((g_1, g_2), (h_1, h_2))$ to $\psi_1(g_1, h_1) + \psi_2(g_2, h_2)$ for every $g_1, h_1 \in G_1$ and $g_2, h_2 \in G_2$. Under this product, linkings form an abelian semigroup \mathcal{R} . For a connected closed orientable 3-manifold M , its torsion linking form, $(TH_1(M), \psi_M)$ is an example of linkings. Moreover, for two connected closed oriented 3-manifolds M and N , the torsion linking form of $M \# N$ is $\psi_M \oplus \psi_N$. In [12], a generator of \mathcal{R} is given as following:

Fact 3.1. [12]

- Let $A^p(q)$ be a linking on $\mathbb{Z}/p\mathbb{Z}$ (with a generator $1 \in \mathbb{Z}/p\mathbb{Z}$) which is represented by $\left(\frac{q}{p}\right)$ for coprime integers $p > 1$ and q , and
- let E_0^k be a linking on $\mathbb{Z}/2^k\mathbb{Z} \oplus \mathbb{Z}/2^k\mathbb{Z}$ (with a basis $\{(1, 0), (0, 1)\}$) which is represented by $\begin{pmatrix} 0 & 2^{-k} \\ 2^{-k} & 0 \end{pmatrix}$ for every integer $k > 0$, and
- let E_1^k be a linking on $\mathbb{Z}/2^k\mathbb{Z} \oplus \mathbb{Z}/2^k\mathbb{Z}$ (with a basis $\{(1, 0), (0, 1)\}$) which is represented by $\begin{pmatrix} 2^{1-k} & 2^{-k} \\ 2^{-k} & 2^{1-k} \end{pmatrix}$ for every integer $k > 1$.

Then \mathcal{R} is generated by $A^p(q)$, E_0^k and E_1^k .

Moreover, the presentation of \mathcal{R} is revealed by combining the results in [12] and [6]. As a notation we regard $A^1(q)$, E_0^0 and E_1^0 as ϕ , the identity element of \mathcal{R} .

Thus every linking is represented as $(\bigoplus_{0 \leq l \leq a} A^{p_l}(q_l)) \oplus (\bigoplus_{0 \leq i \leq e_0} E_0^{k_i}) \oplus (\bigoplus_{0 \leq j \leq e_1} E_1^{k'_j})$, where $a \geq 0$, $p_0 = 1$, p_l is an integer greater than 1 for $l \neq 0$, q_l is an integer prime to p_l , and $e_0 \geq 0$, $k_0 = 0$, $k_i > 0$ for $i \neq 0$, and $e_1 \geq 0$, $k'_0 = 0$, $k'_j > 1$ for $j \neq 0$. Note that this representation is not unique.

3.2 Representatives

We give 3-manifolds representing the generators in Fact 3.1. On showing that the torsion linking forms of these are the generators, we adopt a method of calculating torsion linking forms of rational homology 3-spheres by using their Heegaard splittings [2]. At first, we review the method.

3.2.1 Calculating torsion linking forms by using Heegaard splittings [2]

Let M be a rational homology 3-sphere and $M = V \cup W$ a Heegaard splitting i.e. M is obtained from two handlebodies of same genus, say g , V and W by pasting their boundaries by some orientation reversing homeomorphism, say $f : \partial V \rightarrow \partial W$. We give V and W orientations as standard handlebodies in \mathbb{R}^3 , and we give M an orientation coming from V . Take a symplectic basis $\{a_1^V, \dots, a_g^V, b_1^V, \dots, b_g^V\}$ of $H_1(\partial V)$ such that b_i^V is 0 in $H_1(V)$ for every i , where symplectic basis means that the intersection form on ∂V satisfies $\langle a_i^V, a_j^V \rangle = 0$, $\langle b_i^V, b_j^V \rangle = 0$, and $\langle a_i^V, b_j^V \rangle = \delta_{i,j}$ for every $1 \leq i, j \leq g$. Similarly, take a symplectic basis $\{a_1^W, \dots, a_g^W, b_1^W, \dots, b_g^W\}$ of $H_1(\partial W)$ such that b_i^W is 0 in $H_1(W)$ for every i .

Denote by $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ the matrix representing f_* , the map between the first homology group induced by f with respect to these bases, where A, B, C and D are $(g \times g)$ -matrices over \mathbb{Z} .

Fact 3.2. [2] *In the situation above, $\det(B) \neq 0$ and $H_1(M)$ is isomorphic to $\mathbb{Z}^g/B^t\mathbb{Z}^g$. Moreover, the torsion linking form ψ_M is equivalent to $\mathbb{Z}^g/B^t\mathbb{Z}^g \times \mathbb{Z}^g/B^t\mathbb{Z}^g \rightarrow \mathbb{Q}/\mathbb{Z}; (v, w) \mapsto -v^t(B^{-1}A)w$.*

Remark 3.1. We review the procedure for the calculation in terms of Heegaard diagrams rather than gluing maps used in the following. Let $M = V \cup W$ be a genus g Heegaard splitting of a closed oriented 3-manifold (not necessarily a rational homology three sphere), and S the splitting surface. Give S the orientation as ∂V . Take a family of pairwise disjoint oriented simple closed curves $\{b_1^V, \dots, b_g^V\}$ on S such that each b_i^V bounds pairwise disjoint disks in V and that these disks cut V into a ball. Similarly, take a family of pairwise disjoint oriented simple closed curves $\{b_1^W, \dots, b_g^W\}$ on S such that each b_i^W bounds pairwise disjoint disks in W and that these disks cut W into a ball. Note that the triplet $(S, \{b_1^V, \dots, b_g^V\}, \{b_1^W, \dots, b_g^W\})$ determines the homeomorphic type of $M = V \cup W$. It is well-known that we can compute $H_1(M)$ using the curves: $H_1(M) = \mathbb{Z}\langle x_1, \dots, x_g \rangle / (\sum_{i=1}^g \langle b_i^V, b_j^W \rangle_S \cdot x_i \text{ for } 1 \leq j \leq g)$, where $\langle \cdot, \cdot \rangle_S$ is the algebraic intersection form on S . By this, we can see whether M is a rational homology 3-sphere or not. Suppose that M is a rational homology 3-sphere in the following. Choose a family of pairwise disjoint oriented simple closed curves $\{a_1^V, \dots, a_g^V\}$ on S such that $\langle a_i^V, b_j^V \rangle_S = \delta_{i,j}$ for every $1 \leq i, j \leq g$. Similarly, choose a family of pairwise disjoint oriented simple closed curves $\{a_1^W, \dots, a_g^W\}$ on S such that $\langle a_i^W, b_j^W \rangle_S = -\delta_{i,j}$ for every $1 \leq i, j \leq g$. Then the homology class of $\{a_1^V, \dots, a_g^V, b_1^V, \dots, b_g^V\}$ in S forms a symplectic basis of $H_1(\partial V)$ such that each b_i^V vanishes in $H_1(V)$, and the homology class of $\{a_1^W, \dots, a_g^W, b_1^W, \dots, b_g^W\}$ in S forms a symplectic basis of $H_1(\partial W)$ such that each b_i^W vanishes in $H_1(W)$. In this situation, we can compute the matrices A, B using curves: The (i, j) -entry of A is $\langle b_i^W, a_j^V \rangle_S$, and the (i, j) -entry of B is $\langle b_i^W, b_j^V \rangle_S$.

3.2.2 Representatives for $A^p(q)$

Let M be a lens space of type (p, q) for $p > 1$. M has a surgery presentation as in Figure 2 and a Heegaard splitting $V \cup W$ as in Figure 3. In Figure 3, V is the inner handlebody and W is the outer handlebody, and a box containing $\frac{-q}{p}$ represents curves in it, a result of resolving the intersection points of horizontal $|q|$ lines and vertical $|p|$ lines so that they twist in left-hand (or right-hand) side if $\frac{-q}{p} > 0$ (or < 0 , respectively). We take the $(p, -q)$ -curve as b_1^V and the $(0, 1)$ -curve as b_1^W on the splitting torus. Note that the orientation of $L(p, q)$ coming from the standard one of S^3 corresponds to that coming from V . By computation, we see that M is a rational homology 3-sphere. Let a_1^V be the (r, s) -curve for r, s such that $-rq - sp = 1$. Then the matrices in 3.2.1 are $A = (-r)$ and $B = (-p)$ by computing the algebraic intersections $\langle a_1^V, b_1^W \rangle$ and $\langle b_1^V, b_1^W \rangle$ on ∂W . Note that $\partial V = -\partial W$. Thus by Fact 3.2, $H_1(M) = \mathbb{Z}/p\mathbb{Z}$ and the torsion linking form is $(-\frac{r}{p})$. Let x be a generator of $H_1(M) = \mathbb{Z}/p\mathbb{Z}$ corresponding to the 1×1 -matrix $(-\frac{r}{p})$. Note that qx is also a generator of $H_1(M) = \mathbb{Z}/p\mathbb{Z}$ since p and q are coprime. Under this new generator, the matrix representation of the torsion linking form is $(-\frac{rq^2}{p}) = (\frac{q}{p})$ since $rq \equiv -1 \pmod{p}$, the same as $A^p(q)$. Henceforth, $M(A^p(q))$ denotes $L(p, q)$.

3.2.3 Representatives for E_0^k

Let M be a connected closed orientable 3-manifold having a surgery description as in Figure 4. It has a Heegaard splitting $V \cup W$ as in Figure 5, so-called a vertical Heegaard splitting. The convention of boxes containing rational numbers is as in Figure 3. Note that this Heegaard diagram is not minimally intersecting. We regard V as the inner handlebody and W as the outer handlebody. Note that the orientation coming from the standard one of S^3 corresponds to that coming from V . By computation, we see that M is a rational homology 3-sphere. Let a_1^V, a_2^V be curves as in Figure 6, where b_1^V, b_2^V




Figure 2: A surgery description of $L(p, q)$




Figure 3: left: A Heegaard splitting of $L(p, q)$ right: Examples of boxes contain rational numbers

is the same as in Figure 5, which we abbreviate. Then the matrices in 3.2.1 are $A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 2^k & 0 \\ -2^k & 2^k \end{pmatrix}$. Thus by Fact 3.2, $H_1(M) = \mathbb{Z}\langle x, y \rangle / (2^k x - 2^k y, 2^k y)$ and $\psi_M = -B^{-1}A = \begin{pmatrix} 0 & -2^{-k} \\ -2^{-k} & 2^{1-k} \end{pmatrix}$ under the basis $\{x, y\}$. By changing a basis, $H_1(M) = (\mathbb{Z}\langle x + y \rangle / (2^k(x + y))) \oplus (\mathbb{Z}\langle -x \rangle / (2^k(-x)))$ and $\psi_M = \begin{pmatrix} 0 & 2^{-k} \\ 2^{-k} & 0 \end{pmatrix}$ under the basis $\{x + y, -x\}$, the same as E_0^k . Henceforth, $M(E_0^k)$ denotes this M .




Figure 4: Surgery descriptions of M




Figure 5: A Heegaard splitting of M




Figure 6: Curves $a_1^V, a_2^V, b_1^W, b_2^W$

3.2.4 Representatives for E_1^k

Let M be a connected closed orientable 3-manifold having a surgery description as in Figure 7. It has a Heegaard splitting $V \cup W$ as in Figure 8, so-called a vertical Heegaard splitting. The convention of boxes containing rational numbers is as in Figure 3. We regard V as the inner handlebody and W as the outer handlebody. Note that the orientation coming from the standard one of S^3 corresponds to that coming from V . By computation, we see that M is a rational homology 3-sphere. Let a_1^V, a_2^V be curves as in Figure 9, where b_1^V, b_2^V are the same as in Figure 8, which we abbreviate. Then the matrices in 3.2.1 are $A = \begin{pmatrix} 0 & -1 \\ -3 & 3 \end{pmatrix}$ and $B = \begin{pmatrix} 2^k & 2 \cdot 2^k \\ -2^k & -3 \cdot 2^k \end{pmatrix}$. Thus by Fact 3.2, $H_1(M) = \mathbb{Z}\langle x, y \rangle / (2^k x - 2^k y, 2 \cdot 2^k x - 3 \cdot 2^k y)$ and $\psi_M = -B^{-1}A = \begin{pmatrix} 3 \cdot 2^{1-k} & -3 \cdot 2^{-k} \\ -3 \cdot 2^{-k} & 2^{1-k} \end{pmatrix}$ under the basis $\{x, y\}$. By changing a basis, $H_1(M) = (\mathbb{Z}\langle x+y \rangle / (2^k(x+y))) \oplus (\mathbb{Z}\langle -y \rangle / (2^k(-y)))$ and $\psi_M = \begin{pmatrix} 2^{1-k} & 2^{-k} \\ 2^{-k} & 2^{1-k} \end{pmatrix}$ under the basis $\{x+y, -y\}$, the same as E_1^k . Henceforth, $M(E_1^k)$ denotes this M .




Figure 7: Surgery descriptions of M




Figure 8: A Heegaard splitting of M




Figure 9: Curves $a_1^V, a_2^V, b_1^W, b_2^W$

From the above, we have representatives for connected closed oriented 3-manifolds with respect to the first homology groups and their torsion linking forms.

Proposition 3.1. *Suppose that a connected closed oriented 3-manifold M whose free part of the first homology group is isomorphic to \mathbb{Z}^r and whose torsion linking form is equivalent to $(\bigoplus_{0 \leq l \leq a} A^{p_l}(q_l)) \oplus (\bigoplus_{0 \leq i \leq e_0} E_0^{k_i}) \oplus (\bigoplus_{0 \leq j \leq e_1} E_1^{k'_j})$ for integers $a, p_l, q_l, e_0, k_i, e_1, k'_j$ as in the end of Subsection 3.1.*

Then a 3-manifold $(\#^r S^2 \times S^1) \# (\#_{0 \leq l \leq a} M(A^{p_l}(q_l))) \# (\#_{0 \leq i \leq e_0} M(E_0^{k_i})) \# (\#_{0 \leq j \leq e_1} M(E_1^{k'_j}))$ has isomorphic first homology group to that of M and has equivalent torsion linking forms to that of M , where as a notation we regard $\#^0 S^2 \times S^1, M(A^1(q_0)), M(E_0^0)$ and $M(E_1^0)$ as S^3 .

We use this in Section 5.

4 Thickened surface and its spine bands

In this section, we explain how surfaces (with their spines) in a given 3-manifold correspond to pairs of annuli and bands in the manifold. All surfaces and 3-manifolds in this section are oriented, and we can use the terminology “front” and “back” sides of surfaces.

We state the statement as a proposition:

Proposition 4.1. *Let M be an oriented 3-manifold. Fix non-negative integers g and n .*

There is a bijection between the set of isotopy classes of oriented surfaces in M , each of which is homeomorphic to $\Sigma_{g,n+1}$ with the spine as in Figure 10 and the set of oriented annuli A_1, \dots, A_{2g+n} with oriented core curves and the oriented bands (regarded as small rectangles) $B_1, \dots, B_g, C_1, \dots, C_{g+n-1}$ satisfying the following condition (see Figure 11 for example):

- *One side of ∂B_i , which consists of four sides, is on the back side of A_{2i-1} and is a properly embedded essential arc on A_{2i-1} , and the opposite side of ∂B_i is on the front side of A_{2i} and is a part of the core curve of A_{2i} for $1 \leq i \leq g$.*
- *For $1 \leq i \leq g$, near the intersection of B_i with A_{2i-1} , the front side of B_i is on the positive direction with respect to the orientation of the core curve of A_{2i-1} , and near that of with A_{2i} , the front side of B_i is the “right” side of the core curve of A_{2i} , where we look the intersection (this is on the front side of A_{2i}) so that the core curve of A_{2i} runs from the bottom to the top.*
- *For $1 \leq i \leq g-1$, C_i connects the “left” boundary of A_{2i} and the “right” boundary of A_{2i+1} so that the front sides are attached, and for $g \leq j \leq g+n-1$, C_j connects the “left” boundary of*

A_{j+g} and the “left” boundary of A_{j+g+1} so that the front sides are attached, where we look the front side of A_k so that the core curve runs from the bottom to the top.

We call the condition for annuli and bands in Proposition 4.1 the condition \spadesuit . We construct oriented annuli and bands from a surface with its spine in Subsection 4.1, and construct a surface with its spine from oriented annuli and bands in Subsection 4.2. We can see that these operations are the inverses of each other. Thus Proposition 4.1 holds.




Figure 10: A spine of $\Sigma_{g,n+1}$




Figure 11: Annuli and bands

4.1 Annuli and bands obtained from surfaces

Let F be a surface homeomorphic to $\Sigma_{g,n+1}$ in M . Fix an embedding $\iota : \Sigma_{g,n+1} \times [-1, 1] \rightarrow M$ such that $\iota(\Sigma_{g,n+1} \times \{0\}) = F$ and identify $\Sigma_{g,n+1}$ with F via ι . Take a spine $(K_1 \cup K_2) \cup w_1 \cup \dots \cup w_{g-1} \cup (K_{2g-1} \cup K_{2g}) \cup w_g \cup K_{2g+1} \cup w_{g+1} \cup \dots \cup w_{g+n-1} \cup K_{2g+n}$ of F as in Figure 10. We assign an orientation to K_i for each $1 \leq i \leq 2g+n$. Denote the point $K_{2i-1} \cap K_{2i}$ for $1 \leq i \leq g$ by p_i . Let U_i be the square centered at p_i and V_j the rectangle around w_j as in Figure 12. Take charts $\phi_i : [-1, 1]^2 \rightarrow U_i$ such that $\phi_i([-1, 1] \times \{0\})$ is a part of K_{2i-1} , $\phi_i(\{0\} \times [-1, 1])$ is a part of K_{2i} , and $\psi_j : [-1, 1]^2 \rightarrow V_j$ such that $\psi_j([-1, 1] \times \{0\})$ is a part of w_j and $\{-1\} \times [-1, 1]$ is near K_{2j} for $1 \leq j \leq g$, near K_{j+g} for $g+1 \leq j \leq g+n-1$.

Denote the annulus whose core curve is K_i by A'_i for $1 \leq i \leq 2g+n$ as in Figure 13. Consider $F \times [-1, 1]$, see Figure 14. We regard that it is in M via ι . Let A_{2i-1} , A_{2i} and A_j be $A'_{2i-1} \times \{1\}$, $A'_{2i} \times \{-1\}$ and $A'_j \times \{1\}$, respectively for $1 \leq i \leq g$ and $2g+1 \leq j \leq 2g+n$. These annuli can be regarded as framed knots in M . Let B_i , C_i and C_j be $\phi_i(\{0\} \times [-1, 1]) \times [-1, 1]$, $\{\psi_i(\{t\} \times [-1, 1]) \times \{t\} | t \in [-1, 1]\}$ and $\psi_j([-1, 1]^2) \times \{1\}$, respectively for $1 \leq i \leq g$ and $g+1 \leq j \leq g+n-1$. These are bands in M . We get




Figure 12: Charts

an oriented annuli A_k 's and bands B_i 's and C_j 's (these are rectangles) in M as in Figure 11. Observe that these satisfy the condition \spadesuit with appropriate orientations.




Figure 13: Annuli

4.2 Surfaces from annuli and bands

Let A_1, \dots, A_{2g+n} be oriented annuli and $B_1, \dots, B_g, C_1, \dots, C_{g+n-1}$ be bands satisfying the condition \spadesuit in M . These look like some embeddings of the union of annuli and bands as in Figure 11. Thicken these (denoted by N) and consider curves with colors as in Figure 15. In this figure, a part of the core curve of A_{2i} , which is also one edge of the band B_i is replaced with the other three edges of B_i for each $1 \leq i \leq g$. A small neighborhood F of these curves in ∂N is homeomorphic to $\Sigma_{g,n+1}$ with a spine as in Figure 10. Let K_{2i-1} denote the red curve coming from A_{2i-1} and K_{2i} the blue curve coming from A_{2i} and B_i for $1 \leq i \leq g$, and w_j the green curve coming from C_j for $1 \leq j \leq g+n-1$. For each $1 \leq i \leq 2g+n$, fix an essential arc α_i in A_i such that it is disjoint from w_j for $1 \leq j \leq g+n-1$ and K_l for $l \neq i$, and let D_i be an essential disk in N obtained by thickening α_i . Then each ∂D_i intersects ∂F




Figure 14: Thickened F

twice with the opposite orientations and D_i 's cut N into a ball. Therefore, F gives N the structure of $\Sigma_{g,n+1} \times [-1, 1]$ so that F is $\Sigma_{g,n+1} \times \{1\}$. The colored curves play a role of a spine of $\Sigma_{g,n+1}$. We get a surface corresponding to $\Sigma_{g,n+1} \times \{0\}$ with a spine obtained by projecting the curves along the product structure. In other word, the surface $\Sigma_{g,n+1} \times \{0\}$ with its spine is isotopic to F with its spine in M . We can easily show that the operations in Subsections 4.1 and 4.2 are inverses of each other.

Remark 4.1. Let $m_{A_{2i-1}}$ be a meridian curve of A_{2i-1} (i.e. an oriented circle on the boundary of a small neighborhood of A_{2i-1} which bounds a disk in the neighborhood intersecting the core of A_{2i-1} once positively) for $1 \leq i \leq g$. We assume $m_{A_{2i-1}}$ is in a small neighborhood of A_{2i-1} and disjoint from A_{2i-1} . Note that the curve obtained from A_{2i} by the above procedure represents $[A_{2i}] - [m_{A_{2i-1}}]$ in $H_1(M \setminus N)$, where $[A_{2i}]$ and $[A_{2i-1}]$ denote the elements represented by push-ups of the core curves of A_{2i} and A_{2i-1} , respectively.




Figure 15: Thickened annuli and bands

5 A proof of Theorem 1.1

In this section, we give a proof of Theorem 1.1. At first, we consider the condition for a connected closed orientable 3-manifold having a homologically fibered link whose homological fiber is given homeomorphic type using a surgery diagram. In order to define the linking number $lk(\cdot, \cdot)$ among knots in the surgery link, we have to give the link an orientation. We fix any one. However, this choice does not matter for the existence of a homologically fibered link whose fiber is the given homeomorphism type, see Remark 5.1.

Proposition 5.1. *Let M be a connected closed orientable 3-manifold. Suppose that M is obtained from S^3 by the surgery along a link $L = L_1 \sqcup \cdots \sqcup L_m$, where each L_i is an oriented knot and its coefficient is $\frac{p_i}{q_i}$.*

Then M has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{g,n+1}$ if and only if there exist $m \times (2g+n)$ -matrix of integer coefficients X and $(2g+n) \times (2g+n)$ -symmetric matrix of integer coefficients Y satisfying the following equation:

$$\begin{vmatrix} \Phi & \Psi X \\ X^t & Y + (\mathcal{E} \oplus O_n) \end{vmatrix} = \pm 1, \quad (2)$$

where Φ is an $m \times m$ -matrix whose (i, i) -entry is p_i and (i, j) -entry is $q_i lk(L_i, L_j)$ for $i \neq j$, Ψ is an $m \times m$ -diagonal matrix whose (i, i) -entry is q_i , and \mathcal{E} is a $2g \times 2g$ -matrix whose $(2i-1, 2i)$ -entry is 1 for every $1 \leq i \leq g$ and the others are 0.

Proof. Suppose M has a homologically fibered link whose homological fiber F is homeomorphic to $\Sigma_{g,n+1}$. We will construct a solution of (2). By choosing a spine of F , and using the construction in Section 4, we get oriented annuli A_1, \dots, A_{2g+n} and oriented bands $B_1, \dots, B_g, C_1, \dots, C_n$ satisfying the condition \spadesuit . We regard these annuli and bands are in the surgery diagram on S^3 missing the surgery link. Let T_i be the canonical longitude (regarded in S^3) of the core curve of A_i and m_{A_i} the meridian of (the core curve of) A_i such that $lk(T_i, m_{A_i}) = 1$ and t_i the framing number of A_i regarded as a framed knot. Note that $H_1(M \setminus F) \cong H_1\left(M \setminus \left(\bigcup_{1 \leq i \leq 2g+n} A_i\right) \cup \left(\bigcup_{1 \leq j \leq g} B_j\right) \cup \left(\bigcup_{1 \leq l \leq n} C_l\right)\right) \cong H_1\left(M \setminus \left(\bigcup_{1 \leq i \leq 2g+n} A_i\right)\right)$, and that the push-up of K_i , which is in spines of F as in Figure 10 represents

$$\begin{cases} [T_{2i'-1}] + t_{2i'-1}[m_{A_{2i'-1}}] & \text{if } 1 \leq i \leq 2g \text{ and } i = 2i' - 1, \\ [T_{2i'}] + t_{2i'}[m_{A_{2i'}}] - [m_{A_{2i'-1}}] & \text{if } 1 \leq i \leq 2g \text{ and } i = 2i', \\ [T_i] + t_i[m_{A_i}] & \text{if } 2g+1 \leq i \leq 2g+n \end{cases}$$

in $H_1\left(M \setminus \left(\bigcup_{1 \leq i \leq 2g+n} A_i\right)\right)$.

Let m_{L_i} be the meridian of L_i and set $l_{i,j}$ to be $lk(L_i, L_j)$ and $\tilde{l}_{i,j}$ to be $lk(L_i, T_j)$ and $\bar{l}_{i,j}$ to be $lk(T_i, T_j)$.

Note that $H_1\left(S^3 \setminus \left(L \cup \left(\bigcup_{1 \leq i \leq 2g+n} A_i\right)\right)\right) \cong \mathbb{Z}^{m+2g+n}$ is generated by $\{[m_{L_1}], \dots, [m_{L_m}], [m_{A_1}], \dots, [m_{A_{2g+n}}]\}$,

and that $\frac{p_i}{q_i}$ -slope of L_i represents an element $p_i[m_{L_i}] + q_i\{\sum_{1 \leq j \leq m, j \neq i} \tilde{l}_{i,j}[m_{L_j}] + \sum_{1 \leq k \leq 2g+n} \bar{l}_{i,k}[m_{A_k}]\}$ in

$H_1\left(S^3 \setminus \left(L \cup \left(\bigcup_{1 \leq i \leq 2g+n} A_i\right)\right)\right)$, denoted by \mathbb{A}_i . Note also that the push-up of K_i , which is in a spine of F as in Figure 10 represents

$$\begin{cases} t_{2i'-1}[m_{A_{2i'-1}}] + \sum_{1 \leq j \leq m, j \neq i} \tilde{l}_{j,i}[m_{L_j}] + \sum_{1 \leq k \leq 2g+n, k \neq i} \bar{l}_{i,k}[m_{A_k}] & \text{if } 1 \leq i \leq 2g \text{ and } i = 2i' - 1, \\ t_{2i'}[m_{A_{2i'}}] - [m_{A_{2i'-1}}] + \sum_{1 \leq j \leq m, j \neq i} \tilde{l}_{j,i}[m_{L_j}] + \sum_{1 \leq k \leq 2g+n, k \neq i} \bar{l}_{i,k}[m_{A_k}] & \text{if } 1 \leq i \leq 2g \text{ and } i = 2i', \\ t_i[m_{A_i}] + \sum_{1 \leq j \leq m, j \neq i} \tilde{l}_{j,i}[m_{L_j}] + \sum_{1 \leq k \leq 2g+n, k \neq i} \bar{l}_{i,k}[m_{A_k}] & \text{if } 2g+1 \leq i \leq 2g+n \end{cases}$$

under the basis $\{[m_{L_1}], \dots, [m_{L_m}], [m_{A_1}], \dots, [m_{A_{2g+n}}]\}$ of $H_1\left(S^3 \setminus \left(L \cup \left(\bigcup_{1 \leq i \leq 2g+n} A_i\right)\right)\right)$, denoted by \mathbb{B}_i .

Since F is homological fiber, $H_1(M \setminus F)$ must be \mathbb{Z}^{2g+n} and the push-ups of K_i 's form a free basis of $H_1(M \setminus F)$. This implies that $\{[m_{L_1}], \dots, [m_{L_m}], [m_{A_1}], \dots, [m_{A_{2g+n}}]\} / \langle \mathbb{A}_1, \dots, \mathbb{A}_m \rangle$ is isomorphic to \mathbb{Z}^{2g+n} , which also implies that $\mathbb{A}_1, \dots, \mathbb{A}_m$ are linearly independent in $H_1\left(S^3 \setminus \left(L \cup \left(\bigcup_{1 \leq i \leq 2g+n} A_i\right)\right)\right)$,

and \mathbb{B}_j 's modulo $\langle \mathbb{A}_1, \dots, \mathbb{A}_m \rangle$ form a basis of it. Thus $\{\mathbb{A}_1, \dots, \mathbb{A}_m, \mathbb{B}_1, \dots, \mathbb{B}_{2g+n}\}$ is a basis of \mathbb{Z}^{m+2g+n} freely generated by $\{[m_{L_1}], \dots, [m_{L_m}], [m_{A_1}], \dots, [m_{A_{2g+n}}]\}$. Thus considering the change of

basis matrix, we get a solution of (2) by setting X equal to a matrix whose (i, j) -entry is $\tilde{l}_{i,j}$, Y equal to a symmetric matrix whose (k, k) -entry is t_k , $(2i-1, 2i)$ -entry and $(2i, 2i-1)$ -entry for $i \leq g$ are $\bar{l}_{2i-1, 2i} - 1$ and other (i, j) -entry is $\bar{l}_{i,j}$.

Conversely, suppose we have a solution X, Y for (2). Then we can construct oriented framed knots K_1, \dots, K_{2g+n} in the surgery diagram on S^3 , which will be the core curves of annuli A_1, \dots, A_{2g+n} , such that $lk(L_i, K_j)$ is the (i, j) -entry of X , the framing of K_i is the (i, i) -entry of Y , $lk(K_{2i-1}, K_{2i})$ for $i \leq g$ is equal to 1 plus the $(2i-1, 2i)$ -entry of Y , and $lk(K_i, K_j)$ for other pair (i, j) is the (i, j) -entry

of Y . Take bands $B_1, \dots, B_g, C_1, \dots, C_{2g+n-1}$ satisfying the condition \spadesuit . Note that we can choose bands arbitrarily since these do not affect on the homology. We get a surface which is homeomorphic to $\Sigma_{g,n+1}$ by the construction in Section 4. By reversing the argument in the previous paragraph, we see that this surface is a homological fiber. \square

Remark 5.1. Suppose there is a solution X, Y for the equation (2) in Proposition 5.1 for some orientation of the surgery link. If we reverse the orientation of the i -th component of m -component surgery link, then the signs of the i -th row and i -th column of Φ except for the (i, i) -entry and the (i, i) -th entry of Ψ in the left hand side of the equation (2) are switched. Then a pair of the matrix obtained from X by switching the signs of the i -th row and Y is a solution of the new equation. Therefore the existence of a solution of the equation (2) is preserved.

Remark 5.2. For a given surgery diagram and a solution of (2), we can concretely construct a homologically fibered link as in the latter part of the proof of Proposition 5.1. However we have many choices in construction: There are many links with linking each others in the given number and linking the surgery link in the given number, and we can arbitrary take bands. These homologically fibered links corresponding to one solution are not always different, i.e. they may be isotopic. Moreover, two homological fibered links corresponding to two different solution may be isotopic.

Proof of Theorem 1.1. Suppose that M is a connected closed oriented 3-manifold whose free part of the first homology group is isomorphic to \mathbb{Z}^r and whose torsion linking form is equivalent to $(\bigoplus_{0 \leq l \leq a} A^{p_l}(q_l)) \oplus (\bigoplus_{0 \leq i \leq e_0} E_0^{k_i}) \oplus (\bigoplus_{0 \leq j \leq e_1} E_1^{k'_j})$.

Then $\tilde{M} = (\#^r S^2 \times S^1) \# (\#_{0 \leq l \leq a} M(A^{p_l}(q_l))) \# (\#_{0 \leq i \leq e_0} M(E_0^{k_i})) \# (\#_{0 \leq j \leq e_1} M(E_1^{k'_j}))$ has the isomorphic first homology group to M and has a torsion linking form equivalent to that of M by Proposition 3.1. By Proposition 2.1, M has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{g,n+1}$ if and only if \tilde{M} has such homologically fibered link. We have a surgery diagram on S^3 for \tilde{M} which consists of r copies of an unknot with surgery slope 0 (for $\#^r S^2 \times S^1$), a copies of an unknot with surgery slope $\frac{p_l}{q_l}$ for $0 \leq l \leq a$ (for $\#_{0 \leq l \leq a} M(A^{p_l}(q_l))$), e_0 copies of 2-component link as in the right of Figure 4 for $0 \leq i \leq e_0$ (for $\#_{0 \leq i \leq e_0} M(E_0^{k_i})$), and e_1 copies of 2-component link as in the right of Figure 7 for $0 \leq j \leq e_1$ (for $\#_{0 \leq j \leq e_1} M(E_1^{k'_j})$). Applying Proposition 5.1 with this surgery diagram with appropriate order, we get the equality (1). \square

Almost the same statement as the next corollary was given by Nozaki.

Corollary 5.1. *Let M be a connected closed orientable 3-manifold. Suppose that $M \# (S^2 \times S^1)$ has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{0,n+1}$ for $n \geq 1$ (resp. $\Sigma_{1,1}$). Then M has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{0,n}$ (resp. $\Sigma_{0,2}$).*

Proof. At first, for the case where $M \# (S^2 \times S^1)$ has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{0,2}$, we can see that $H_1(M \# (S^2 \times S^1)) \cong H_1(M) \oplus H_1(S^2 \times S^1)$ is generated by one element by Remark 2.1. Thus M is an integral homology 3-sphere and has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{0,1}$. In the following, for the case where $M \# (S^2 \times S^1)$ has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{0,n+1}$, we assume that $n \geq 2$.

Fix a surgery diagram \mathcal{D} of M , and let $m-1$ be the number of knots in \mathcal{D} . A surgery diagram \mathcal{D}' of $M \# (S^2 \times S^1)$ is obtained by adding disjoint unknot U with surgery slope 0 ($= \frac{0}{1}$) to \mathcal{D} . We give an order among the link in \mathcal{D}' so that U is the first component. Then by Proposition 5.1, we have a solution of (2) for $\Sigma_{0,n+1}$ (resp. $\Sigma_{1,1}$) and fix it. Let \mathbb{X} denote the matrix in the left hand side of (2) and let $\mathbb{X}_{i,j}$ denote the (i, j) -entry of \mathbb{X} . Note that $\mathbb{X}_{1,i} = \mathbb{X}_{i,1} = 0$ for $1 \leq i \leq m$, and that $\mathbb{X}_{1,i} = \mathbb{X}_{i,1}$ for $m+1 \leq i \leq m+n$ (resp. $m+1 \leq i \leq m+2$).

Since (2) holds, $\mathbb{X}_{1,m+1}, \mathbb{X}_{1,m+2}, \dots, \mathbb{X}_{1,m+n}$ (resp. $\mathbb{X}_{1,m+1}, \mathbb{X}_{1,m+2}$) are coprime. This implies that the (ordered) set $\{\mathbb{X}_{1,m+1}, \mathbb{X}_{1,m+2}, \dots, \mathbb{X}_{1,m+n}\}$ (resp. $\{\mathbb{X}_{1,m+1}, \mathbb{X}_{1,m+2}\}$) can be changed into n -element set (resp. two-element set) such that the first element is 1 or -1 and the other elements are 0 in

finitely many steps, at each of which the l -times of the i -th element is added to the j -th element for some integer l and $1 \leq i \neq j \leq n$ (resp. $1 \leq i \neq j \leq 2$). Fix one of such sequence of steps. Along this sequence, at each step, say the l -times of the i -th element is added to the j -th element, we change \mathbb{X} as follows: Add l times the $(m+i)$ -th column to the $(m+j)$ -th column and then add l times the $(m+i)$ -th row to the $(m+j)$ -th row. Note that at any time in the sequence, each matrix obtained from \mathbb{X} is also a “solution” of (2). A solution for a homologically fibered link whose homological fiber is homeomorphic to neither $\Sigma_{0,n+1}$ nor $\Sigma_{1,1}$ is not necessarily preserved under this operation. At the end of the sequence we get a matrix \mathbb{X}' obtained from \mathbb{X} whose $(1, i)$ -entry is 0 for $i \neq m+1$ and ± 1 for $i = m+1$. Note that the $(i, 1)$ -entry of \mathbb{X}' is 0 for $i \neq m+1$ and ± 1 for $i = m+1$.

Expand the determinant of \mathbb{X}' along the first row (the only one cofactor survives) and expand the determinant of the cofactor along the first column (the only one cofactor survives). The cofactor finally obtained is a solution for $\Sigma_{0,n}$ (resp. $\Sigma_{0,2}$) in a 3-manifold which has a surgery diagram \mathcal{D} , representing M . □

6 Examples

Though we have Theorem 1.1, it is difficult in general to find a solution of (1) for a given manifold and homeomorphic type of a surface. Nozaki [10] proved that $L(p, q)$ has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{1,1}$ for every pair (p, q) by solving some equations, which is equivalent to (1), using the density theorem. The author [11] proved that $L(p, q)$ has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{0,3}$ for every pair (p, q) by solving (1) following the Nozaki’s argument. Thanks to Nozaki [10], we see that $\text{hc}(M(A^p(q))) = 1$ for every (p, q) . Moreover, he also proved in his thesis that $\text{hc}((\#^{2m}S^2 \times S^1)\#M(A^p(q))) = m + 1$, and $\text{hc}((\#^{2m+1}S^2 \times S^1)\#M(A^p(q))) = m + 1$ if q or $-q$ is quadratic residue modulo p and $\text{hc}((\#^{2m+1}S^2 \times S^1)\#M(A^p(q))) = m + 2$ otherwise for every non-negative integer m . Thus we know $\text{hc}(\cdot)$ for all 3-manifolds whose torsion linking form is $A^p(q)$. In this section, we determine $\text{hc}(\cdot)$ for 3-manifolds whose torsion linking forms are the other generators for linkings, $(\#^r S^2 \times S^1)\#M(E_0^k)$ and $(\#^r S^2 \times S^1)\#M(E_1^k)$ for a non-negative integer r , we state as a proposition, proved at Subsections 6.1 and 6.2:

Proposition 6.1. *In the following, $\lceil n \rceil$ and $\lfloor n \rfloor$ for a real number n represent the minimal integer greater than or equal to n and the maximal integer less than or equal to n , respectively.*

- For $r \geq 0$, $\text{hc}((\#^r S^2 \times S^1)\#M(E_0^k)) = \lceil \frac{r}{2} \rceil + 1$ if $k = 1$ or $k \geq 3$.
- $\text{hc}(M(E_0^2)) = 2$, and for $r \geq 1$ $\text{hc}((\#^r S^2 \times S^1)\#M(E_0^2)) = \lceil \frac{r}{2} \rceil + 1$.
- For $r \geq 0$, $\text{hc}((\#^r S^2 \times S^1)\#M(E_1^2)) = \lceil \frac{r}{2} \rceil + 1$.
- For $r \geq 0$, $\text{hc}((\#^r S^2 \times S^1)\#M(E_1^k)) = \lfloor \frac{r}{2} \rfloor + 2$ if $k \geq 3$.

As a preparation, we review two observations: Firstly, the invariant $\text{hc}(\cdot)$ is subadditive under the connected sum i.e. $\text{hc}(M_1\#M_2) \leq \text{hc}(M_1) + \text{hc}(M_2)$ since we get a homological fiber by the plumbing of two homological fibers, see [11] for example. Secondly, it is known that $S^2 \times S^1$ has a fibered link whose fiber surface is an annulus. By the plumbing it and a Hopf annulus in S^3 , and by the plumbing two fibered annuli of $S^2 \times S^1$, we get genus one fibered knots in $S^2 \times S^1$ and $\#^2 S^2 \times S^1$. Since they are not integral homology 3-spheres, we see that $\text{hc}(S^2 \times S^1) = 1$ and $\text{hc}(\#^2 S^2 \times S^1) = 1$.

6.1 $\text{hc}((\#^r S^2 \times S^1)\#M(E_0^k))$

We divide the argument into three cases, where r is zero, where r is positive even, and where r is odd.

6.1.1 $\text{hc}(M(E_0^k))$

We will compute $\text{hc}(M(E_0^k))$. Since $M(E_0^k)$ is not an integral homology 3-sphere, $\text{hc}(M(E_0^k)) \geq 1$. Moreover, it is known that $M(E_0^k)$ has a fibered link whose fiber surface is homeomorphic to $\Sigma_{0,3}$ as in Figure 16: If we ignore the green curve, the right of Figure 16 represents a fiber surface of a fibered link whose fiber surface is homeomorphic to $\Sigma_{0,3}$ in $L(2^k, 1) \# L(2^k, 1)$, which is obtained by the plumbing of fibered annuli in each prime components in appropriate way. The green curve is on this fiber, and note that the canonical framing of this curve is identical with the surface framing. Then $(-\frac{1}{2^k})$ -surgery (with respect to the canonical framing) along the green curve preserves the fiber structure. By the plumbings of two Hopf annuli, we get a genus two fibered knot in $M(E_0^k)$. This implies $\text{hc}(M(E_0^k)) \leq 2$. Thus it is enough to show whether $M(E_0^k)$ has a genus one homologically fibered knot or not.




Figure 16: Surgery link on a planar fiber surface

By Theorem 1.1, $M(E_0^k)$ has a genus one homologically fibered knot if and only if there exist integers $x, y, z, w, \alpha, \beta$ and γ satisfying

$$\begin{aligned} \pm 1 &= \begin{vmatrix} 0 & 2^k & x & y \\ 2^k & 0 & z & w \\ x & z & \alpha & \beta + 1 \\ y & w & \beta & \gamma \end{vmatrix} \\ &= -2^{2k}\{\alpha\gamma - \beta(\beta + 1)\} + 2^{k+1}\{x(z\gamma - w\beta) - y(z\beta - w\alpha)\} - 2^k(xw + yz) + (xw - yz)^2 \quad (3) \end{aligned}$$

(i) $k = 1$

In this case, it is known that $M(E_0^1)$ has a genus one fibered knot as in Figure 17: If we ignore the green curve, the right of Figure 17 represents a fiber surface of a genus one fibered knot in $L(2, 1) \# L(2, 1)$, which is obtained by the plumbing of fibered annuli in each prime components in appropriate way. The green curve is on this fiber, and note that the surface framing of this curve is the (-1) -slope with the canonical framing. This implies that the $(-\frac{1}{2})$ -slope of the green curve with respect to the canonical framing is the $(-\frac{1}{2})$ -slope with respect to the surface framing. Then the surgery along the green curve preserves the fiber structure. Thus $M(E_0^1)$ has a genus one homologically fibered knot. In fact, $x = 3, y = 1, z = 0, w = 1, \alpha = -1, \beta = 0$ and $\gamma = 0$ is one of the solutions. This solution may correspond to a non-fibered knot.

(ii) $k = 2$

In this case, there exist no solutions. If there was, $(xw - yz)$ must be odd since the other terms in the right hand side of (7) are even. If $(xw - yz)$ is odd, then the right hand side of (7) is congruent to 5 modulo 8 by noting that $xw + yz = xw - yz + 2yz$ and that the square of every odd integer is congruent to 1 modulo 8. This cannot be ± 1 .




Figure 17: Surgery link on a fiber surface of genus one

(iii) $k \geq 3$

In this case, $x = 2^{k-1}, y = 1, z = 1, w = 1, \alpha = 2^{k-3} + 1, \beta = 0$ and $\gamma = 0$ is one of the solutions of (7).

6.1.2 $\text{hc}((\#^{2m} S^2 \times S^1) \# M(E_0^k))$ for $m \geq 1$

Since $H_1((\#^{2m} S^2 \times S^1) \# M(E_0^k)) \cong \mathbb{Z}^{2m} \oplus \mathbb{Z}/2^k \mathbb{Z} \oplus \mathbb{Z}/2^k \mathbb{Z}$ requires at least $2m + 2$ elements for generating, $\text{hc}((\#^{2m} S^2 \times S^1) \# M(E_0^k)) \geq m + 1$ by Remark 2.1. By the plumbing of two fibered annuli of $S^2 \times S^1$ to a fiber surface of $M(E_0^k)$ in the right of Figure 16, we get a genus two fibered knot in $(\#^2 S^2 \times S^1) \# M(E_0^k)$. Thus

$$\begin{aligned} \text{hc}((\#^{2m} S^2 \times S^1) \# M(E_0^k)) &\leq (m-1) \text{hc}(\#^2 S^2 \times S^1) + \text{hc}((\#^2 S^2 \times S^1) \# M(E_0^k)) \\ &\leq m + 1. \end{aligned}$$

Therefore, $\text{hc}((\#^{2m} S^2 \times S^1) \# M(E_0^k)) = m + 1$ for $m \geq 1$.

6.1.3 $\text{hc}((\#^{2m+1} S^2 \times S^1) \# M(E_0^k))$ for $m \geq 0$

Since $H_1((\#^{2m+1} S^2 \times S^1) \# M(E_0^k)) \cong \mathbb{Z}^{2m+1} \oplus \mathbb{Z}/2^k \mathbb{Z} \oplus \mathbb{Z}/2^k \mathbb{Z}$ requires at least $2m + 3$ elements for generating, $\text{hc}((\#^{2m+1} S^2 \times S^1) \# M(E_0^k)) \geq m + 2$ by Remark 2.1. By the plumbing of two fibered annuli, one is in $S^2 \times S^1$ and the other is in S^3 , to a fiber surface of $M(E_0^k)$ in the right of Figure 16, we get a genus two fibered knot in $(S^2 \times S^1) \# M(E_0^k)$. Thus

$$\begin{aligned} \text{hc}((\#^{2m+1} S^2 \times S^1) \# M(E_0^k)) &\leq m \cdot \text{hc}(\#^2 S^2 \times S^1) + \text{hc}((S^2 \times S^1) \# M(E_0^k)) \\ &\leq m + 2. \end{aligned}$$

Therefore, $\text{hc}((\#^{2m+1} S^2 \times S^1) \# M(E_0^k)) = m + 2$ for $m \geq 0$.

6.2 $\text{hc}((\#^r S^2 \times S^1) \# M(E_1^k))$

We divide the argument into four cases, where r is zero, where k is two, where k is not two and r is even, and where k is not two and r is odd.

6.2.1 $\text{hc}(M(E_1^k))$

We will compute $\text{hc}(M(E_1^k))$ for $k \geq 2$. Since $M(E_1^k)$ is not an integral homology 3-sphere, $\text{hc}(M(E_1^k)) \geq 1$. Moreover, $M(E_1^k)$ has a genus two fibered knot as in Figure 18: If we ignore the red curve and the blue curve, two of the bottom of Figure 18 represent fiber surfaces of genus two fibered knots in $L(2^k, 1) \# L(2^k, 1) \# L(-\frac{2}{3}\{2^{k-1} - (-1)^{k-1}\}, 1) \# L(2, (-1)^{k-1})$ for odd k and even k , which are obtained by the plumbings of fibered annuli in each prime components in appropriate way. The red curve

and blue curve are on this fiber. Note that the 0-slope of the red curve with respect to the canonical framing is the (-1) -slope with respect to the surface framing, and that the $((-1)^k \cdot 2)$ -slope of the blue curve with respect to the canonical framing is the 1-slope with respect to the surface framing. Then the surgery along the red curve and the blue curve preserves the fiber structure. Therefore we have $\text{hc}(M(E_1^k)) \leq 2$. Thus it is enough to show whether $M(E_1^k)$ has a genus one homologically fibered knot or not.




Figure 18: Surgery link on a fiber surface of genus two

By Theorem 1.1, $M(E_1^k)$ has a genus one homologically fibered knot if and only if there exist integers $x, y, z, w, \alpha, \beta$ and γ satisfying

$$\begin{aligned} \pm 1 &= \begin{vmatrix} 2^{k+1} & -2^k & -x & -y \\ -3 \cdot 2^k & 2^{k+1} & -3z & -3w \\ x & z & \alpha & \beta + 1 \\ y & w & \beta & \gamma \end{vmatrix} \\ &= 2^{2k} \{ \alpha \gamma - \beta(\beta + 1) \} + 2^{k+1} \{ \gamma(x^2 + 3xz + 3z^2) + \alpha(y^2 + 3yw + 3w^2) - (2\beta + 1)(xy + 3zw) \} \\ &\quad - 3 \cdot 2^k (2\beta + 1)(xw + yz) + 3(xw - yz)^2 \end{aligned} \quad (4)$$

(i) $k = 2$

In this case, we get a genus one fibered knot as in Figure 19: If we ignore the green curve, the right of

Figure 19 represents a fiber surface of a genus one fibered knot in $L(4, 1) \# L(4, 1)$, which is obtained by the plumbing of fibered annuli in each prime components in appropriate way. The green curve is on this fiber, and note that the surface framing of this curve is the (-1) -slope with the canonical framing. This implies that the $(-\frac{3}{4})$ -slope of the green curve with respect to the canonical framing is the (14) -slope with respect to the surface framing. Then the surgery along the green curve preserves the fiber structure. Thus $M(E_1^2)$ has a genus one homologically fibered knot. In Fact, $x = -1, y = 1, z = 1, w = 0, \alpha = 1, \beta = 0$ and $\gamma = 0$ is one of the solutions. This solution may correspond to a non-fibered knot. Thus we have $\text{hc}(M(E_1^2)) = 1$.




Figure 19: Surgery link on a fiber surface of genus one

(ii) $k \geq 3$

In this case, there exist no solutions of (4). In fact, all terms but the last in the right hand side of (4) are divisible by 8 and the last term is congruent to neither 1 nor -1 modulo 8. Thus we have $\text{hc}(M(E_1^k)) \geq 2$. We conclude $\text{hc}(M(E_1^k)) = 2$.

6.2.2 $\text{hc}((\#^r S^2 \times S^1) \# M(E_1^2))$ for $r \geq 0$

Since $H_1((\#^r S^2 \times S^1) \# M(E_1^2)) \cong \mathbb{Z}^r \oplus \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$ requires at least $r+2$ elements for generating, $\text{hc}((\#^r S^2 \times S^1) \# M(E_1^2)) \geq \lceil \frac{r}{2} \rceil + 1$ by Remark 2.1. On the other hand,

$$\begin{aligned} \text{hc}((\#^r S^2 \times S^1) \# M(E_1^2)) &\leq \text{hc}(\#^r S^2 \times S^1) + \text{hc}(M(E_1^2)) \\ &\leq \lceil \frac{r}{2} \rceil + 1. \end{aligned}$$

Therefore, $\text{hc}((\#^r S^2 \times S^1) \# M(E_1^2)) = \lceil \frac{r}{2} \rceil + 1$ for $r \geq 0$.

6.2.3 $\text{hc}((\#^{2m} S^2 \times S^1) \# M(E_1^k))$ for $m \geq 0$ and $k \geq 3$

Note that

$$\begin{aligned} \text{hc}((\#^{2m} S^2 \times S^1) \# M(E_1^k)) &\leq m \cdot \text{hc}(\#^2 S^2 \times S^1) + \text{hc}(M(E_1^k)) \\ &= m + 2. \end{aligned}$$

We will show that $(\#^{2m} S^2 \times S^1) \# M(E_1^k)$ has no homologically fibered links whose homological fibers are homeomorphic to $\Sigma_{m+1,1}$. This implies that $\text{hc}((\#^{2m} S^2 \times S^1) \# M(E_1^k)) = m + 2$. Suppose that $(\#^{2m} S^2 \times S^1) \# M(E_1^k)$ had a homologically fibered links whose homological fibers are homeomorphic to $\Sigma_{m+1,1}$. By applying Theorem 1.1, we have a solution X , $(2m+2) \times (2m+2)$ -integer matrix, and Y , $(2m+2) \times (2m+2)$ -symmetric integer matrix, for an equation below.

$$\left| \begin{array}{c} O_{2m} \oplus \begin{pmatrix} 2^{k+1} & -2^k \\ -3 \cdot 2^k & 2^{k+1} \end{pmatrix} \\ X^t \end{array} \begin{pmatrix} I_{2m} \oplus \begin{pmatrix} -1 & 0 \\ 0 & -3 \end{pmatrix} \\ Y + \mathcal{E} \end{pmatrix} X \right| = \pm 1 \quad (5)$$

Note that the left hand side is congruent to $\left| X^t \begin{pmatrix} I_{2m} \oplus \begin{pmatrix} -1 & 0 \\ 0 & -3 \end{pmatrix} \end{pmatrix} X \right| = 3|X|^2$ modulo 8. This cannot be ± 1 , and this leads a contradiction.

6.2.4 $\text{hc}((\#^{2m+1}S^2 \times S^1)\#M(E_1^k))$ for $m \geq 0$ and $k \geq 3$

First, we show that $M(E_1^k)$ has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{1,2}$. By Theorem 1.1, the existence of such a link is equivalent to that of a solution $a, b, c, d, e, f, v, w, x, y, z$ of the equation below.

$$\left| \begin{array}{ccccc} 2^{k+1} & -2^k & -x & -y & -u \\ -3 \cdot 2^k & 2^{k+1} & -3z & -3w & -3v \\ x & z & a & b+1 & d \\ y & w & b & c & e \\ u & v & d & e & f \end{array} \right| = \pm 1 \quad (6)$$

By substituting $u = 1, v = d = e = f = 0$, the left hand side of (6) becomes equal to $\left| \begin{array}{ccc} 2^{k+1} & -3z & -3w \\ z & a & b+1 \\ w & b & c \end{array} \right|$.

Note that the existence of a, b, c, z, w such that the determinant becomes ± 1 is equivalent to that of a genus one homologically fibered knot in $L(2^{k+1}, 3)$ by Theorem 1.1, and the existence is guaranteed by Nozaki. Thus $M(E_1^k)$ has a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{1,2}$. Moreover, we get a homologically fibered link whose homological fiber is homeomorphic to $\Sigma_{2,1}$ in $(S^2 \times S^1)\#M(E_1^k)$ by the plumbing with a fibered annulus in $S^2 \times S^1$.

Since $H_1((\#^{2m+1}S^2 \times S^1)\#M(E_1^k)) \cong \mathbb{Z}^{2m+1} \oplus \mathbb{Z}/2^k\mathbb{Z} \oplus \mathbb{Z}/2^k\mathbb{Z}$ requires at least $2m+3$ elements for generating, $\text{hc}((\#^{2m+1}S^2 \times S^1)\#M(E_1^k)) \geq m+2$ by Remark 2.1. On the other hand,

$$\begin{aligned} \text{hc}((\#^{2m+1}S^2 \times S^1)\#M(E_1^k)) &\leq m \cdot \text{hc}(\#^2S^2 \times S^1) + \text{hc}((S^2 \times S^1)\#M(E_1^k)) \\ &\leq m + 2. \end{aligned}$$

Therefore, $\text{hc}((\#^{2m+1}S^2 \times S^1)\#M(E_1^k)) = m + 2$ for $m \geq 0$.

References

- [1] J. W. Alexander. A lemma on systems of knotted curves. *Proceedings of the National Academy of Sciences*, 9(3):93–95, 1923.
- [2] A. Conway, S. Friedl and G. Herrmann. Linking forms revisited. *Pure and Applied Mathematics Quarterly*, volume 12, Number 4, 493–515, 2016.
- [3] S. Garoufalidis and J. Levine. Tree-level invariants of three-manifolds, Massey products and the Johnson homomorphism. In *Graphs and patterns in mathematics and theoretical physics*, volume 73 of *Proc. Sympos. Pure Math.*, pages 173–203. Amer. Math. Soc., Providence, RI, 2005.
- [4] H. Goda and T. Sakasai. Homology cylinders and sutured manifolds for homologically fibered knots. *Tokyo J. Math.*, 36(1):85–111, 2013.

- [5] K. Habiro. Claspers and finite type invariants of links. *Geom. Topol.* 4 (2000), 1–83.
- [6] A. Kawauchi and S. Kojima. Algebraic classification of linking pairings on 3-manifolds. *Math. Ann.* 253 (1980), 29–42.
- [7] G. Massuyeau and J.-B. Meilhan. Equivalence relations for homology cylinders and the core of the Casson invariant. *Trans. Amer. Math. Soc.* 365 (2013), 5431–5502.
- [8] S. Matveev. Generalized surgeries of three-dimensional manifolds and representations of homology spheres. *Mat. Zametki* 42 (1987) 268–278, 345 (Russian); English translation: *Math. Notes* 42 (1987), 651–656.
- [9] R. Myers. Open book decompositions of 3-manifolds. *Proc. Amer. Math. Soc.*, 72(2):397–402, 1978.
- [10] Y. Nozaki. Every lens space contains a genus one homologically fibered knot. *Illinois J. Math.* volume 62, Number 1–4 (2018), 99–111.
- [11] N. Sekino. Lens spaces which are realizable as closures of homology cobordisms over planar surfaces. *Illinois J. Math.* volume 64, Number 4 (2020), 481–492.
- [12] Wall, C.T.C.: Quadratic forms on finite groups, and related topics, *Topology* 2, 281–298 (1964).

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO,
 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914, JAPAN
 E-mail address: `sekinonozomu@g.ecc.u-tokyo.ac.jp`