The ternary Goldbach-Vinogradov theorem with almost equal Piatetski-Shapiro primes

Yanbo Song^{a,*}

^aSchool of Mathematics and Statistics, Xi'an Jiaotong University, Xi'an, 710049, China

Abstract

In this paper, we proved a theorem that every large enough odd number can be represented as the sum of three almost equal Piatetski-Shapiro primes. *Keywords:* Goldbach-Vinogradov theorem, Piatetski-Shapiro prime, almost-equal prime 2010 MSC: 11P32, 11P05

1. Introduction

In 1937, it was proved by I.M. Vinogradov that every large enough odd number could be represented as the sum of three primes. Today we usually called this theorem the ternary Goldbach-Vinogradov theorem. Later many scholars studied the problem that large enough odd number can be represented as the sum of three primes. In [9, 10], C. D. Pan and C. B. Pan showed that every large enough odd number n the Diophantine equation

$$n = p_1 + p_2 + p_3, \quad |p_i - \frac{n}{3}| \le y, \quad 1 \le i \le 3,$$
 (1)

with prime variable is solvable for $y = n^{2/3} (\log n)^c$, where c is a positive constant.

The set of Piatetski-Shapiro primes is of the form

$$\mathcal{A}_{\gamma} = \{ prime \ p | p = [n^{1/\gamma}], \ n \ge 1 \}.$$

Preprint submitted to Indagationes mathematicae-new series

December 14, 2020

^{*}Corresponding author

Email address: syb888@stu.xjtu.edu.cn (Yanbo Song)

⁵ For more information about Piatetski-Shapiro primes, we refer to [4, 5, 8, 12, 13, 14].

In 1992, A. Balog and J. Friedlander [1] studied the ternary Goldbach-Vinogradov theorem restricted to the set of Piatetski-Shapiro primes. They proved the following result:

$$\sum_{\substack{p_1+p_2+p_3=n\\p_i\in\mathcal{A}_{\gamma}}} \frac{1}{\gamma} p_1^{1-\gamma} \frac{1}{\gamma} p_2^{1-\gamma} \frac{1}{\gamma} p_3^{1-\gamma} = \mathfrak{S}_3(n) \frac{3n^2}{\log^3 n} + O\left(\frac{n^2}{\log^4 n}\right),\tag{2}$$

where $\mathfrak{S}_3(n) = \prod_{p|n} \left(1 - \frac{1}{(p-1)^2}\right) \prod_{p \nmid n} \left(1 + \frac{1}{(p-1)^3}\right)$ and $20/21 < \gamma < 1$. Later C.H. Jia [6] extended the range of γ to $15/16 < \gamma < 1$, and A.Kumchev [7] improved the result to $50/53 < \gamma < 1$.

In this paper, we combined these two ideas to study whether (1.1) can hold restricted to the set of Piatetski-Shapiro primes. Specifically, we proved the following theorem that gives an affirmative answer.

Theorem 1. For every large enough odd number n the following equation

$$p_1 = p_1 + p_2 + p_3, \quad |p_i - \frac{n}{3}| \le y, \quad p_i \in \mathcal{A}_{\gamma} \quad 1 \le i \le 3,$$
 (3)

with prime variable is solvable for $y > n^{(52-45\gamma)/8}$ and $44/45 < \gamma < 1$.

2. Notations

γ

Throughout the paper, the letters m and n are natural numbers, and palways denotes a prime number. Furthermore, c > 1 is a fixed real number and and p always denotes a prime number. Furthermore, c > 1 is a fixed real number and

we put $\gamma = \frac{1}{c}$. And we denote the \mathcal{A}_c the set of the Piatetski-Shapiro primes.

We shall frequently use ε with a slight abuse of notation to mean a small positive number, possibly different from line to line.

Given a real number x, we write $e(x) = e^{2\pi i x}$, $\{x\}$ for the fractional part of x, [x] for the greatest integer not exceeding x. We write $\mathcal{L} = \log N$.

We recall that for functions F and real nonnegative G the notations $F \ll G$ and F = O(G) are equivalent to the statement that the inequality $|F| \leq \alpha G$ holds for some constant $\alpha > 0$. We also write $F \simeq G$ to indicate that $F \ll G$ and $G \ll F$. 30 For any function f, we put

$$\Delta f(x) = f(-(x+1)^{\gamma}) - f(-x^{\gamma}), \quad (x > 0).$$

3. Some lemmas

Lemma 1. Put $\psi(x) = x - [x] - 1/2$. Then, there are numbers a_h , b_h such that

$$\left|\psi(x) - \sum_{0 < |h| \le H} a_h e(xh)\right| \le \sum_{|h| \le H} b_h e(xh),$$

where $a_h \ll \frac{1}{|h|}$, $b_h \ll \frac{1}{H}$. Moreover, we let $\psi'(x) = \sum_{0 < |h| \le H} a_h e(xh)$, and 35 $\psi''(x) = \psi(x) - \psi'(x)$.

Proof. For the proof of this lemma, you can see [2], Appendix.

Lemma 2. Let $u, v \ge 1$ be real numbers. For any n > u, we have

$$\Lambda(n) = \sum_{\substack{cd=n\\d \le v}} \log c\mu(d) - \sum_{k|n} a(k) - \sum_{\substack{kc=n\\k>1\\c>u}} \Lambda(c)b(k),$$

where

$$a(k) = \sum_{\substack{cd=k\\c\leq u\\d\leq v}} \Lambda(c)\mu(d), \quad b(k) = \sum_{\substack{d|k\\d\leq v}} \mu(d).$$

Proof. This is Prop. 13.4 in [4].

40 Lemma 3. Let

$$g(\alpha) = \sum_{|p - \frac{n}{3}| < y} e(\alpha p),$$

we have

$$\int_0^1 g(\alpha)^3 e(-n\alpha) d\alpha = \mathfrak{S}_3(n) \frac{3y^2}{\log^3 n} + O(\frac{y^2}{\log^4 n}),$$

where $\mathfrak{S}_{3}(n) = \prod_{p|n} (1 - \frac{1}{(p-1)^{2}}) \prod_{p \nmid n} (1 + \frac{1}{(p-1)^{3}}).$ **Proof.** This is Theorem 3 in [11]. Lemma 4. We put

$$f(\alpha) = \sum_{\substack{|p - \frac{n}{3}| < y \\ p \in \mathcal{A}_1/\gamma}} \frac{1}{\gamma} p^{1 - \gamma} e(\alpha p).$$

⁴⁵ And $g(\alpha)$ are defined in the previous lemma. For any α , there are coprime integers a and q with $0 \le a \le q \le \log n$ such that $|q\alpha - a| < \log^{-1} n$. and $y > n^{(52-45\gamma)/8}$, $44/45 < \gamma < 1$, Then we have

$$f(\alpha) = g(\alpha) + O(n^{21/31 - 14\gamma/31 + \varepsilon} y^{23/31}).$$

Proof. Let $\chi(n)$ be the characteristic function of the set \mathcal{A}_c , then we have

$$\chi(n) = [-n^{\gamma}] - [-(n+1)^{\gamma}] = \gamma n^{\gamma-1} + (\psi(-(n+1)^{\gamma}) - \psi(-n^{\gamma})) + O(n^{\gamma-2}).$$

Thus,

$$f(\alpha) = g(\alpha) + \sum_{|p - \frac{n}{3}| < y} cp^{1 - \gamma} e(\alpha p)(\psi(-(p+1)^{\gamma}) - \psi(-p^{\gamma})) + O(1/\log x).$$

50 It is enough to estimate,

$$\sum_{\substack{|p-\frac{n}{3}| < y}} cp^{1-\gamma} e(\alpha p)(\psi(-(p+1)^{\gamma}) - \psi(-p^{\gamma}))$$

=
$$\sum_{\substack{|p-\frac{n}{3}| < y}} cp^{1-\gamma} e(\alpha p) \Delta \psi'(p) + O\left(\sum_{\substack{|p-\frac{n}{3}| < y}} cp^{1-\gamma} e(\alpha p) \Delta \psi''(p)\right)$$

:= $S_1 + S_2.$

By Lemma 3.1, we have

$$S_2 \ll H^{-1}yn^{1-\gamma} + H^{-1}\sum_{1 \le h < H} \left| \sum_{|x-n/3| < y} x^{1-\gamma} e(hx^{\gamma}) \right|.$$

Partial summation yields

$$\sum_{|x-n/3| < y} x^{1-\gamma} e(hx^{\gamma}) \ll n^{1-\gamma} \max_{t < y} \left| \sum_{|x-n/3| < t} e(hx^{\gamma}) \right|.$$

Using the exponent pair (1/2, 1/2) ([2], eqn. 3.3.4), we have

$$\sum_{|x-n/3| < t} e(hx^{\gamma}) \ll h^{1/2} n^{\gamma/2} + h^{-1} n^{1-\gamma}.$$

Concluding the above estimate, we have

$$S_2 \ll H^{1/2} n^{1-\gamma/2} + H^{-1} y n^{1-\gamma}.$$

55 For S_1 , we first observe that

$$S_1 \ll \frac{1}{\log n} \max_{t < y} \left| \sum_{|x - n/3| < t} cx^{1 - \gamma} e(\alpha x) \Lambda(n) \Delta \psi'(x) \right| + O(n^{3/2 - \gamma}).$$

Noting that $a_h \ll |h|^{-1}$, we have

$$\Delta \psi'(x) = \sum_{1 \le |h| \le H} a_h e(hx^{\gamma})(e(h(x+1)^{\gamma} - hx^{\gamma}) - 1).$$

So we by partial summation, we have

$$\sum_{|x-n/3| < t} cx^{1-\gamma} e(\alpha x) \Lambda(n) \Delta \psi'(x) \ll \max_{z < t} \sum_{1 \le |h| \le H} \left| \sum_{|x-n/3| < z} \Lambda(x) e(\alpha n + hx^{\gamma}) \right|.$$

We have shown so far

$$S_1 \ll \frac{1}{\log n} \max_{z < y} \sum_{1 \le |h| \le H} \left| \sum_{|x - n/3| < z} \Lambda(x) e(\alpha n + hx^{\gamma}) \right| + O(n^{3/2 - \gamma}).$$

Because for any α , there are coprime integers a and q with $0 \le a \le q \le \log n$ such that $|q\alpha - a| < \log^{-1} n$. Then we have

$$\sum_{|x-n/3| < z} \Lambda(x) e(\alpha n + hx^{\gamma}) = q^{-1} \sum_{-q/2 < b \le q/2} S(a+b,q) \sum_{|x-n/3| < z} \Lambda(x) e(R_b(x)),$$

where $R_b(t) = \beta t + ht^{\gamma} - bt/q$, $\beta = \alpha - a/q$ and

$$S(a,q) = \sum_{m=1}^{q} e(\frac{am}{q}).$$

Apply Lemma 3.2 with $uv \leq n$ we have

$$\sum_{|x-n/3| < z} \Lambda(x) e(R_b(x)) = \Sigma_1 + \Sigma_2 + \Sigma_3,$$

where

$$\begin{split} \Sigma_1 &= \sum_{r \leq u} \mu(r) \sum_{\substack{|rs-n/3| < z \\ r \leq u}} e(R_b(rs)) \log s - \sum_{\substack{|rs-n/3| < z \\ r \leq u}} (\sum_{\substack{r = wt \\ w \leq u \\ t \leq v}} \mu(w) \Lambda(t)) e(R_b(rs)), \\ \Sigma_2 &= -\sum_{\substack{|rs-n/3| < z \\ r > u}} \Lambda(s) \sum_{\substack{d|r \\ d \leq u}} \mu(d) e(R_b(rs)), \\ \Sigma_3 &= -\sum_{\substack{|rs-n/3| < z \\ u < r \leq uv}} (\sum_{\substack{r = wt \\ w \leq u \\ t \leq v}} \mu(w) \Lambda(t)) e(R_b(rs)). \end{split}$$

By partial summation

$$\Sigma_1 \ll \log x \sum_{r \le u} \max_{z \le y} \left| \sum_{|rs-n/3| < z} e(R_b(rs)) \right|.$$

Let $f(s) = R_b(rs)$, and noting that $rs \sim n$ and $f''(s) \approx r^2 n^{\gamma-2} |h|$. And applying the van der Corput's estimate in [3] (Cor. 8.13) to f(s) in the inner sum, we have

$$\Sigma_1 \ll \log^2 x(u|h|^{1/2}x^{\gamma/2-1}y + x^{1-\gamma/2}|h|^{-1/2}).$$

For Σ_2 , we use dyadic division we can write

$$\Sigma_2 \ll x^{\varepsilon} \sum_{R,S} |T(R,S)|,$$

where

$$T(R,S) = \sum_{r \sim R} a_r \sum_{\substack{s \sim S \\ |rs - n/3| < z}} b_s e(R_b(rs))$$

with R > u, s > v, $RS \asymp x$, and $|b_s|, |a_r| \le 1$. By Lemma 2.5 in [2], we have

$$T(R,S)^2 \ll \frac{(RS)^2}{L} + \frac{RS^2}{L} \sum_{1 \le |l| \le L} \max_{S < s, s+l \le 2S} \sum_{r \in I} e(R_b(r(s+l)) - R_b(rs)),$$

where $1 \le L \le S$ is to be chosen later, and $I \subseteq (R, 2R]$ is an interval determined by the conditions $r \sim R$, |rs - n/3| < z, |r(s + l) - n/3| < z. Let g(r) = $R_b(r(s+l)) - R_b(rs)$, so we have $g''(r) \simeq x^{\gamma-1} |hl| R^{-1}$. By Cor. 8.13 in [3], we have

$$\sum_{r \in I} e(R_b(r(s+l)) - R_b(rs)) \ll R^{1/2} x^{\gamma/2 - 3/2} y |hl|^{1/2} + x^{1/2 - \gamma/2} R^{1/2} |hl|^{-1/2}.$$

 $So \ we \ have$

$$T(R,S) \ll xL^{-1/2} + R^{3/4}SL^{1/4}x^{\gamma/4-3/4}y^{1/2}|h|^{1/4} + R^{3/4}SL^{-1/4}x^{1/4-\gamma/4}|h|^{-1/4}.$$

And using Lemma 2.4 in $\left[2\right]$ to choose L optimally we obtain

$$\begin{split} T(R,S) \ll & R^{-1/6} n^{(\gamma+3)/6} y^{1/3} |h|^{1/6} + n^{1-\gamma/4} |h|^{-1/4} \\ & + R^{1/2} n^{1/2} + R^{-1/4} n^{(\gamma+1)/4} y^{1/2} |h|^{1/4} + n^{3/4} y^{1/4} R^{-1/4}. \end{split}$$

So we have

$$\begin{aligned} x^{-\varepsilon} \Sigma_2 \ll & nv^{-1/2} + n^{1-\gamma/4} |h|^{-1/4} + n^{3/4} y^{1/4} u^{-1/4} \\ &+ u^{-1/6} n^{(\gamma+3)/6} y^{1/3} |h|^{1/6} + u^{-1/4} n^{(\gamma+1)/4} y^{1/2} |h|^{1/4}. \end{aligned}$$

Similarly we have

$$\begin{split} x^{-\varepsilon} \Sigma_3 \ll & n^{1/2} (uv)^{1/2} + + n^{1-\gamma/4} |h|^{-1/4} + n^{3/4} y^{1/4} u^{-1/4} \\ & + u^{-1/6} n^{(\gamma+3)/6} y^{1/3} |h|^{1/6} + u^{-1/4} n^{(\gamma+1)/4} y^{1/2} |h|^{1/4}. \end{split}$$

 $So \ we \ have$

$$\begin{aligned} x^{-\varepsilon}(\Sigma_2+\Sigma_3) \ll & nv^{-1/2} + n^{1/2}(uv)^{1/2} + n^{1-\gamma/4}|h|^{-1/4} + n^{3/4}y^{1/4}u^{-1/4} \\ & + u^{-1/6}n^{(\gamma+3)/6}y^{1/3}|h|^{1/6} + u^{-1/4}n^{(\gamma+1)/4}y^{1/2}|h|^{1/4}. \end{aligned}$$

Choosing $v = (n/u)^{1/2}$, we have

$$\begin{aligned} x^{-\varepsilon}(\Sigma_1 + \Sigma_2 + \Sigma_3) \ll & u^{1/4} x^{3/4} + n^{1-\gamma/4} |h|^{-1/4} + n^{3/4} y^{1/4} u^{-1/4} \\ &+ u^{-1/6} n^{(\gamma+3)/6} y^{1/3} |h|^{1/6} + u^{-1/4} n^{(\gamma+1)/4} y^{1/2} |h|^{1/4} \\ &+ u |h|^{1/2} x^{\gamma/2 - 1} y. \end{aligned}$$

And using Lemma 2.4 in [2] to choose u optimally we obtain

$$\begin{split} x^{-\varepsilon}(\Sigma_1 + \Sigma_2 + \Sigma_3) \ll & n^{1-\gamma/4} |h|^{-1/4} + x^{3/4} y^{1/8} + |h|^{1/2} x^{\gamma/2 - 1} y + n^{(\gamma+2)/6} y^{1/3} |h|^{1/6} \\ &+ n^{\gamma/4} y^{1/2} |h|^{1/4} + n^{\gamma/8 + 1/2} y^{1/4} |h|^{1/8} + n^{(\gamma+6)/10} y^{1/5} |h|^{1/10} \\ &+ n^{\gamma/10 + 2/5} y^{2/5} |h|^{1/10} + n^{3\gamma/10} y^{3/5} |h|^{3/10} + n^{3\gamma/14 + 2/7} y^{3/7} |h|^{3/14} \end{split}$$

Noting the identity

$$\sum_{a=0}^{p-1} e\left(\frac{ma}{p}\right) = \begin{cases} p, & \text{if } p \mid m;\\ 0, & \text{if } p \nmid m \end{cases}$$

and sum over h, we have

$$\begin{split} S_1 \ll & n^{3/2-\gamma} + n^{\varepsilon} (n^{1-\gamma/4} H^{3/4} + n^{(\gamma+4)/8} y^{1/4} H^{9/8} + n^{(3\gamma+4)/14} y^{3/7} H^{17/14} \\ & + n^{3/4} y^{1/8} H + n^{\gamma/2-1} y H^{3/2} + n^{3\gamma/10} y^{3/5} H^{13/10} \\ & + n^{(\gamma+6)/10} y^{1/5} H^{11/10} + n^{(\gamma+2)/6} y^{1/3} H^{7/6} + n^{\gamma} y^{1/2} H^{5/4}. \end{split}$$

Combining with S_2 , and using Lemma 2.4 in [2] to choose H optimally, we have

$$\begin{split} f(\alpha) - g(\alpha) \ll & n^{3/2 - \gamma} + n^{\varepsilon} (n^{1 - \gamma/4} + n^{(\gamma + 2)/6} y^{1/3} + n^{\gamma/4} y^{1/2} + n^{3/4} y^{1/8} + n^{\gamma/8 + 1/2} y^{1/4} \\ &\quad + n^{\gamma/10 + 3/5} y^{1/5} + n^{3\gamma/10} y^{3/5} + n^{3\gamma/14 + 2/7} y^{3/7} + n^{1 - \gamma/3} y^{1/3} \\ &\quad + n^{1 - 4\gamma/7} y^{3/7} + n^{7/8 - \gamma/2} y^{1/2} + n^{1/5 - 2\gamma/5} y + n^{9/13 - 6\gamma/13} y^{9/13} \\ &\quad + n^{5/9 - 4\gamma/9} y^{7/9} + n^{7/8 - \gamma/2} y^{9/16} + n^{13/17 - 8\gamma/17} y^{11/17} \\ &\quad + n^{17/21 - 10\gamma/21} y^{13/21} + n^{13/23 - 10\gamma/23} y^{19/23} + n^{21/31 - 14\gamma/31} y^{23/31}). \end{split}$$

The proof of the lemma is complete by noting that $n^{21/31-14\gamma/31+\varepsilon}y^{23/31}$ dominates the other terms.

Lemma 5. Let $f(\alpha)$, $g(\alpha)$ are defined in the previous lemma, then we have

$$\int_{0}^{1} |f(\alpha)|^2 d\alpha \ll n^{1-\gamma} y, \tag{4}$$

$$\int_0^1 |g(\alpha)|^2 d\alpha \ll y. \tag{5}$$

Proof. We only prove (3.1), since the proof of (3.2) is similar but simpler.

$$\int_{0}^{1} |f(\alpha)|^{2} d\alpha$$

$$= \sum_{p_{1}} \sum_{p_{2}} p_{1}^{1-\gamma} p_{2}^{1-\gamma} \int_{0}^{1} e(\alpha(p_{1}-p_{2})) d\alpha$$

$$= \sum_{\substack{p \in \mathcal{A}_{c} \\ |p-\frac{n}{3}| < y}} p^{2-2\gamma} \ll n^{2-2\gamma} y n^{\gamma-1} = n^{1-\gamma} y.$$

4. Proof of the theorem

$$\begin{split} R(n) &= \sum_{\substack{p_1 + p_2 + p_3 = n \\ p_i \in \mathcal{A}_c \\ |p_i - \frac{n}{3}| < y \\}} 1 = \int_0^1 f^3(\alpha) e(-n\alpha) d\alpha + O\left(\int_0^1 f^2(\alpha)(f(\alpha) - g(\alpha)) d\alpha\right) \\ &+ O\left(\int_0^1 f(\alpha) g(\alpha)(f(\alpha) - g(\alpha)) d\alpha\right) + O\left(\int_0^1 g^2(\alpha)(f(\alpha) - g(\alpha)) d\alpha\right). \end{split}$$

By Cauchy inequality and trivial estimate, we have

$$\begin{aligned} R(n) &= \int_0^1 g^3(\alpha) e(-n\alpha) d\alpha + O\left(\max_{\alpha \in [0,1]} |f(\alpha) - g(\alpha)| \int_0^1 |f^2(\alpha)| d\alpha\right) \\ &+ O\left(\max_{\alpha \in [0,1]} |f(\alpha) - g(\alpha)| \int_0^1 |g^2(\alpha)| d\alpha\right). \end{aligned}$$

 $_{\tt 80}$ $\,$ By Lemma 3.3, Lemma 3.4 and Lemma 3.5, we have

$$R(n) = \mathfrak{S}_3(n) \frac{3y^2}{\log^3 n} + O(\frac{y^2}{\log^4 n}),$$

where $\mathfrak{S}_3(n) = \prod_{p|n} (1 - \frac{1}{(p-1)^2}) \prod_{p \nmid n} (1 + \frac{1}{(p-1)^3})$. This complete the proof of the theorem.

Acknowledgements

The author would thank his supervisor for his help and encouragement.

85 References

- A. Balog and J. Friedlander, 'A hybrid of theorems of Vinogradov and Piatetski- Shapiro', Pacific J. of Math. 156 (1992), 45-62.
- [2] S. W. Graham and G. Kolesnik, 'van der Corput's Method of Exponential Sums', London Math. Soc. Lecture Note Ser., vol.126, (Cambridge University Press, Cambridge, (1991)).

90

95

105

110

- [3] H. Iwaniec and E. Kowalski, 'Analytic Number Theory', Amer. Math. Soc.
- [5] H. Iwaniec and E. Kowalski, Analytic Number Theory, Amer. Math. Soc. Colloq. Publ., vol. 53, (American mathematical Society, Providence, RI, (2004)).
- [4] C. H. Jia, 'On Piatetski-Shapiro's prime number theorem II', Sci. China 36 (1993),913-926.
- [5] C. H. Jia, 'On Piatetski-Shapiro's prime number theorem I', Chinese Ann. Math. Ser. B 15 (1994), 9-22.
- [6] C. H. Jia, 'On the Piatetski-Shapiro-Vinogradov theorem', Acta Arith. 73 (1995), 1-28.
- [7] A. Kumchev, 'the Piatetski-Shapiro-Vinogradov theorem', J. Théor. Nombres Bordeaux, 9 (1997), 11-23.
 - [8] A. Kumchev, 'On the distribution of prime numbers of the form [n^c]', Glasgow Math. J. 41 (1999), 85-102.
 - [9] C. D. Pan, C. B. Pan, 'On estimation of Trigonometric sums over primes in short intervals (II)', Sci.China, 32 (1989), 641-653.
 - [10] C. D. Pan, C. B. Pan, 'On estimation of Trigonometric sums over primes in short intervals (III)', Chin. Ann. Math., Ser B 11 (1990),138-147
 - [11] C. D. Pan, C. B. Pan, 'Goldbach Conjecture', Science Press, Beijing, (1988).
 - [12] H. Q. Liu and Rivat, 'On the Piatetski-Shapiro prime number theorem', Bull. Lon. Math. Soc. 24 (1993), 143-147.

- [13] I. I. Piatetski-Shapiro, 'On the distribution of prime numbers in sequences of the form [f(m)]', Mat. Sb. 33 (1953), 559-566.
- [14] J. Rivat, J. Wu, 'Prime numbers of the form $[n^c]$ '. Glasg. Math. J. 43(2) 237-254 (2001).