K-THEORY OF ETESI C*-ALGEBRAS

IGOR V. NIKOLAEV 1

ABSTRACT. We study the C^* -algebra $\mathbb{E}_{\mathscr{M}}$ of a smooth 4-dimensional manifold \mathscr{M} introduced by Gábor Etesi. It is proved that the $\mathbb{E}_{\mathscr{M}}$ is a stationary AF-algebra. We calculate the topological and smooth invariants of \mathscr{M} in terms of the K-theory of the C^* -algebra $\mathbb{E}_{\mathscr{M}}$. Using Gompf's Stable Diffeomorphism Theorem, it is shown that all smoothings of \mathscr{M} form a torsion abelian group. The latter is isomorphic to the Brauer group of a number field associated to the K-theory of $\mathbb{E}_{\mathscr{M}}$.

1. INTRODUCTION

Let \mathscr{M} be a topological 4-dimensional manifold. Unlike dimensions 2 and 3, the smooth structures on \mathscr{M} are detached from the topology of \mathscr{M} . Due to the works of Rokhlin, Freedman and Donaldson, it is known that \mathscr{M} can be non-smooth and if there exists a smooth structure, it need not be unique. The classification of all smoothings of \mathscr{M} is an open problem. Denote by $Diff(\mathscr{M})$ a group of the orientation-preserving diffeomorphisms of \mathscr{M} and let $Diff_0(\mathscr{M})$ be a connected component of $Diff(\mathscr{M})$ containing the identity. The group $G := Diff(\mathscr{M})/Diff_0(\mathscr{M})$ is discrete and therefore locally compact.

Definition 1.1. The Etesi C^* -algebra $\mathbb{E}_{\mathscr{M}}$ is a group C^* -algebra of the locally compact group G; see also Remark 4.1.

Remark 1.2. G is a countable, discrete, amenable group acting on \mathscr{M} and the action admits a faithful G-invariant Borel probability measure, e.g. by taking the Lebesgue measure of the orbit space S/G. Thus $\mathbb{E}_{\mathscr{M}} \cong C_0(\mathscr{M}) \rtimes G$ embeds into a simple unital Approximately Finite-dimensional (AF-) C^* -algebra [Schafhauser 2020] [13, Theorem C]. We call the latter an (AF-) Etesi C^* -algebra keeping the same notation for the both.

The aim of our note is a classification of the smooth structures on \mathscr{M} based on the K-theory of the Etesi C^* -algebra $\mathbb{E}_{\mathscr{M}}$. To formalize our results, recall that the AF C^* -algebra is an inductive limit of the multi-matrix C^* -algebras $M_{n_1}(\mathbf{C}) \oplus \cdots \oplus$ $M_{n_k}(\mathbf{C})$ [Blackadar 1986] [1, Section 7]. The AF-algebra is called stationary, if the inductive limit depends on a single positive integer matrix $A \in GL(n, \mathbf{Z})$ [Blackadar 1986] [1, Section 7.2] or [9, Section 3.5.2]. Our main result can be formulated as follows.

Theorem 1.3. The Etesi C^* -algebra $\mathbb{E}_{\mathscr{M}}$ is a stationary AF-algebra.

²⁰¹⁰ Mathematics Subject Classification. Primary 46L85; Secondary 57K41.

Key words and phrases. 4-dimensional manifolds, C^* -algebras.

Let $\lambda_A > 1$ be the Perron-Frobenius eigenvalue of the positive matrix A defined by $\mathbb{E}_{\mathscr{M}}$. Consider a number field $K = \mathbf{Q}(\lambda_A)$. The eigenvector (v_1, \ldots, v_n) corresponding to λ_A can always be scaled so that $v_i \in K$. By $\mathfrak{m} := \mathbf{Z}v_1 + \cdots + \mathbf{Z}v_n$ we understand a \mathbf{Z} -module in the field K and by Λ the ring of endomorphisms of \mathfrak{m} . Let $[\mathfrak{m}]$ be an ideal class of \mathfrak{m} in the ring Λ . The K-theory of stationary AFalgebras says that the triples $(\Lambda, [\mathfrak{m}], K)$ in a one-to-one correspondence with the Morita equivalence classes of $\mathbb{E}_{\mathscr{M}}$ [Handelman 1981] [6], see also [9, Theorem 3.5.4]. Let Br(K) be the Brauer group of the number field K, i.e. a torsion abelian group of the Morita equivalence classes of the central simple algebras over K. By S(k)we understand the connected sum of k copies of $S^2 \times S^2$. We assign an index $k \ge 0$ to each smoothing \mathscr{M}_k of a topological 4-manifold \mathscr{M}_{top} using Gompf's Stable Diffeomorphism Theorem, i.e. a diffeomorphism $\mathscr{M}_k \# S(k) \to \mathscr{M}_0 \# S(k)$, where \mathscr{M}_0 is the standard smoothing of \mathscr{M}_{top} [Gompf 1984] [5, Theorem 1]. The sum of \mathscr{M}_{k_1} and \mathscr{M}_{k_2} is defined by the formula $\mathscr{M}_{k_1} \oplus \mathscr{M}_{k_2} := \mathscr{M}_{k_1+k_2}$, see corollary 2.7. An application of theorem 1.3 is as follows.

Corollary 1.4. Let \mathscr{M} be a smooth 4-manifold, such that the $\mathbb{E}_{\mathscr{M}}$ is an infinitedimensional C^* -algebra. The following is true:

- (i) Handelman triple $(\Lambda, [\mathfrak{m}], K)$ is an invariant of the homeomorphisms of \mathcal{M} ;
- (ii) Elements of the Brauer group Br(K) parametrize smooth structures on \mathcal{M}_{top} .

In particular, all smoothings of \mathcal{M} form a torsion abelian group under the group operation (2.5) with the neutral element \mathcal{M}_0 .

Remark 1.5. The Brauer group Br(K) is known to classify the division algebras over K. In other words, corollary 1.4 (ii) defines a functor from the smooth 4manifolds to the division algebras. Such a functor was constructed independently using the Galois theory for non-commutative fields [10].

The article is organized as follows. Some preliminary facts can be found in Section 2. Theorem 1.3 and corollary 1.4 are proved in Section 3. We conclude by remarks in Section 4.

2. Preliminaries

In this section we briefly review the C^* -algebras, their K-theory and the 4dimensional manifolds. We refer the reader to [Blackadar 1986] [1], [Dixmier 1977] [2] and [Gompf 1984] [5] for the details.

2.1. C^* -algebras. The C^* -algebra is an algebra \mathscr{A} over \mathbb{C} with a norm $a \mapsto ||a||$ and an involution $\{a \mapsto a^* \mid a \in \mathscr{A}\}$ such that \mathscr{A} is complete with respect to the norm, and such that $||ab|| \leq ||a|| ||b||$ and $||a^*a|| = ||a||^2$ for every $a, b \in \mathscr{A}$. Each commutative C^* -algebra is isomorphic to the algebra $C_0(X)$ of continuous complex-valued functions on some locally compact Hausdorff space X. Any other algebra \mathscr{A} can be thought of as a noncommutative topological space.

An AF-algebra (Approximately Finite-dimensional C^* -algebra) is defined to be the norm closure of an ascending sequence of finite dimensional C^* -algebras M_n , where M_n is the C^* -algebra of the $n \times n$ matrices with entries in **C**. Here the index

 $\mathbf{2}$

 $n = (n_1, \ldots, n_k)$ represents the semi-simple matrix algebra $M_n = M_{n_1} \oplus \cdots \oplus M_{n_k}$. The ascending sequence mentioned above can be written as

$$M_1 \xrightarrow{\varphi_1} M_2 \xrightarrow{\varphi_2} \dots,$$
 (2.1)

where M_i are the finite dimensional C^* -algebras and φ_i the homomorphisms between such algebras. If $\varphi_i = Const$, then the AF-algebra \mathscr{A} is called *stationary*.

The homomorphisms φ_i can be arranged into a graph as follows. Let $M_i = M_{i_1} \oplus \cdots \oplus M_{i_k}$ and $M_{i'} = M_{i'_1} \oplus \cdots \oplus M_{i'_k}$ be the semi-simple C^* -algebras and $\varphi_i : M_i \to M_{i'}$ the homomorphism. One has two sets of vertices V_{i_1}, \ldots, V_{i_k} and $V_{i'_1}, \ldots, V_{i'_k}$ joined by a_{rs} edges whenever the summand M_{i_r} contains a_{rs} copies of the summand $M_{i'_s}$ under the embedding φ_i . As *i* varies, one obtains an infinite graph called the Bratteli diagram of the AF-algebra. The matrix $A = (a_{rs})$ is known as a partial multiplicity matrix; an infinite sequence of A_i defines a unique AF-algebra. If \mathscr{A} is a stationary AF-algebra, then $A_i = Const$ for all $i \geq 1$.

2.2. K-theory of AF-algebras. By $M_{\infty}(\mathscr{A})$ one understands the algebraic direct limit of the C^{*}-algebras $M_n(\mathscr{A})$ under the embeddings $a \mapsto \operatorname{diag}(a, 0)$. The direct limit $M_{\infty}(\mathscr{A})$ can be thought of as the C^{*}-algebra of infinite-dimensional matrices whose entries are all zero except for a finite number of the non-zero entries taken from the C^{*}-algebra \mathscr{A} . Two projections $p,q \in M_{\infty}(\mathscr{A})$ are equivalent, if there exists an element $v \in M_{\infty}(\mathscr{A})$, such that $p = v^*v$ and $q = vv^*$. The equivalence class of projection p is denoted by [p]. We write $V(\mathscr{A})$ to denote all equivalence classes of projections in the C^{*}-algebra $M_{\infty}(\mathscr{A})$, i.e. $V(\mathscr{A}) := \{[p] : p = p^* =$ $p^2 \in M_{\infty}(\mathscr{A})$. The set $V(\mathscr{A})$ has the natural structure of an abelian semi-group with the addition operation defined by the formula $[p] + [q] := \operatorname{diag}(p, q) = [p' \oplus q'],$ where $p' \sim p$, $q' \sim q$ and $p' \perp q'$. The identity of the semi-group $V(\mathscr{A})$ is given by [0], where 0 is the zero projection. By the K_0 -group $K_0(\mathscr{A})$ of the unital C^{*}-algebra \mathscr{A} one understands the Grothendieck group of the abelian semi-group $V(\mathscr{A})$, i.e. a completion of $V(\mathscr{A})$ by the formal elements [p] - [q]. The image of $V(\mathscr{A})$ in $K_0(\mathscr{A})$ is a positive cone $K_0^+(\mathscr{A})$ defining the order structure \leq on the abelian group $K_0(\mathscr{A})$. The pair $(K_0(\mathscr{A}), K_0^+(\mathscr{A}))$ is known as a dimension group of the C^* -algebra \mathscr{A} . The scale $\Sigma(\mathscr{A})$ is the image in $K_0^+(\mathscr{A})$ of the equivalence classes of projections in the C^{*}-algebra \mathscr{A} . The $\Sigma(\mathscr{A})$ is a generating, hereditary and directed subset of $K_0^+(\mathscr{A})$, i.e. (i) for each $a \in K_0^+(\mathscr{A})$ there exist $a_1, \ldots, a_r \in \Sigma(\mathscr{A})$ such that $a = a_1 + \cdots + a_r$; (ii) if $0 \le a \le b \in \Sigma(\mathscr{A})$, then $a \in \Sigma(\mathscr{A})$ and (iii) given $a, b \in \Sigma(\mathscr{A})$ there exists $c \in \Sigma(\mathscr{A})$, such that $a, b \leq c$. Each scale can always be written as $\Sigma(\mathscr{A}) = \{a \in K_0^+(\mathscr{A}) \mid 0 \le a \le u\}$, where u is an order unit of $K_0^+(\mathscr{A})$. The pair $(K_0(\mathscr{A}), K_0^+(\mathscr{A}))$ and the triple $(K_0(\mathscr{A}), K_0^+(\mathscr{A}), \Sigma(\mathscr{A}))$ are invariants of the Morita equivalence and isomorphism class of the C^* -algebra \mathscr{A} , respectively. If \mathbb{A} is an AF-algebra, then its scaled dimension group (dimension group, resp.) is a complete invariant of the isomorphism (Morita equivalence, resp.) class of A, see e.g. [9, Theorem 3.5.2].

Let τ be the canonical trace on the AF-algebra \mathbb{A} . Such a trace induces a homomorphism $\tau_* : K_0(\mathbb{A}) \to \mathbb{R}$ and we let $\mathfrak{m} := \tau_*(K_0(\mathbb{A})) \subset \mathbb{R}$. If \mathbb{A} is the stationary AF-algebra given by a matrix $A \in GL(n, \mathbb{Z})$, then \mathfrak{m} is a \mathbb{Z} -module in the number field $K = \mathbb{Q}(\lambda_A)$ generated by the Perron-Frobenius eigenvalue λ_A of the matrix A. The endomorphism ring of \mathfrak{m} is denoted by Λ and the ideal class of \mathfrak{m} is denoted by $[\mathfrak{m}]$. The triple $(\Lambda, [\mathfrak{m}], K)$ is an invariant of the Morita equivalence class of \mathbb{A} [Handelman 1981] [6]. Remark 2.1. Each stationary AF-algebra defines a torsion abelian group. Indeed, let $?_n(x)$ be the *n*-dimensional Minkowski question-mark function, see [Minkowski 1904] [8, p.172] for n = 2 and [Panti 2008] [11, Theorem 3.5] for $n \ge 2$. The $?_n(x) : [0,1]^{n-1} \to [0,1]^{n-1}$ is a continuous function with the following properties: (i) $?_n(0) = 0$ and $?_n(1) = 1$; (ii) $?_n(\mathbf{Q}^{n-1}) = (\mathbf{Z}[\frac{1}{2}])^{n-1}$ are dyadic rationals and (iii) $?_n(\mathcal{K}^{n-1}) = (\mathbf{Q} - \mathbf{Z}[\frac{1}{2}])^{n-1}$, where \mathcal{K} are algebraic numbers of degree *n* over \mathbf{Q} . It is not hard to see, that (iv) $?_n(\Delta) = \Delta$ is a monotone function, where $\Delta = [0,1]$ is the normalized diagonal of the simplex $[0,1]^{n-1}$. Recall that $\tau_*(K_0(\mathbb{A})) = \mathfrak{m}$ and $\tau_*(\Sigma(\mathbb{A})) = \mathfrak{m} \cap [0,1]$, where τ is the canonical trace on the AF-algebra \mathbb{A} and \mathfrak{m} is a **Z**-module in the number field K. We assume that $\tau_*(K_0(\mathbb{A})) \subset \Delta$. By the properties (iii) and (iv) of the Minkowski question-mark function, one gets the following inclusion:

$$\mathscr{Y} := ?_n(\tau_*(\Sigma(\mathbb{A}))) \subset \mathbf{Q}/\mathbf{Z}.$$
(2.2)

Definition 2.2. By the Minkowski group Mi(K) of stationary AF-algebra we understand a torsion abelian group generated by the elements of set \mathscr{Y} .

2.3. 4-dimensional manifolds. We denote by \mathscr{M} a smooth 4-dimensional manifold and always assume \mathscr{M} to be compact. Let S^4 be the 4-dimensional sphere and X_g be a closed 2-dimensional orientable surface of genus $g \geq 0$. By the knotted surface $\mathscr{X} := X_{g_1} \cup \ldots X_{g_n}$ in \mathscr{M} one understands a transverse immersion of a collection of $n \geq 1$ surfaces X_{g_i} into \mathscr{M} . We refer to \mathscr{X} a surface knot if n = 1 and a surface link if $n \geq 2$.

Theorem 2.3. ([Piergallini 1995] [12]) Each smooth 4-dimensional manifold \mathcal{M} is the 4-fold PL cover of the sphere S^4 branched at the points of a knotted surface $\mathscr{X} \subset S^4$.

Let S^2 be the 2-dimensional sphere. By S(k) we understand a smooth 4dimensional manifold corresponding to a connected sum

$$S(k) := \underbrace{(S^2 \times S^2) \# \dots \# (S^2 \times S^2)}_{k \text{ times}}.$$
(2.3)

Theorem 2.4. ([Gompf 1984] [5]) Let \mathscr{M} and \mathscr{M}' be two different smoothings of a topological 4-manifold \mathscr{M}_{top} . Then for sufficiently large k there exists an orientation-preserving diffeomorphism:

$$\mathscr{M} \# S(k) \longrightarrow \mathscr{M}' \# S(k).$$
 (2.4)

Definition 2.5. Let \mathcal{M}_0 be the standard smoothing of a topological 4-manifold \mathcal{M}_{top} . We denote by \mathcal{M}_k a smoothing of \mathcal{M}_{top} , such that $\mathcal{M}_k \# S(k) \to \mathcal{M}_0 \# S(k)$ is an orientation-preserving diffeomorphism given by formula (2.4).

Lemma 2.6. The smoothing \mathcal{M}_k exists for each value of $k \geq 0$.

Proof. Fix an integer $k \geq 0$ and consider the smooth connected sum $\mathscr{M}_0 \# S(k)$. The manifold \mathscr{M}_k can be obtained from $\mathscr{M}_0 \# S(k)$ by cutting off S(k) along the sphere S^3 and glueing in a copy of S^4 endowed with the standard smooth structure. (Notice that \mathscr{M}_k is different from \mathscr{M}_0 .) The reader can verify, that \mathscr{M}_k is a smoothing of \mathscr{M}_{top} and there exists Gompf's diffeomorphism $\mathscr{M}_k \# S(k) \to \mathscr{M}_0 \# S(k)$ given by formula (2.4).

Corollary 2.7. The smoothings of \mathcal{M}_{top} have the structure of an abelian monoid under the operation

$$\mathscr{M}_{k_1} \oplus \mathscr{M}_{k_2} := \mathscr{M}_{k_1 + k_2} \tag{2.5}$$

with the neutral element \mathcal{M}_0 .

Proof. (i) Let us show that operation (2.5) defines a semigroup. By lemma 2.6, the sum $\mathcal{M}_{k_1} \oplus \mathcal{M}_{k_2}$ is a smoothing of \mathcal{M}_{top} . In other words, the set of all smoothing of \mathcal{M}_{top} is closed under the operation (2.5), i.e. such a set is a semigroup.

(ii) The semigroup of item (i) is abelian, since $\mathcal{M}_{k_1} \oplus \mathcal{M}_{k_2} = \mathcal{M}_{k_2} \oplus \mathcal{M}_{k_1}$.

(iii) It follows from (2.5), that $\mathcal{M}_k \oplus \mathcal{M}_0 = \mathcal{M}_k$. Thus \mathcal{M}_0 is the neutral element of the semigroup. In other words, our semigroup is a monoid.

Corollary 2.7 is proved.

3. Proofs

3.1. **Proof of theorem 1.3.** We shall split the proof in two lemmas.

Lemma 3.1. The C^* -algebra $\mathbb{E}_{\mathscr{M}}$ is an AF-algebra.

Proof. The lemma follows from an observation that the $\mathbb{E}_{\mathscr{M}}$ is a group C^* -algebra of a locally compact group, see definition 1.1. Then the C^* -algebra $\mathbb{E}_{\mathscr{M}}$ is an AF-algebra, see e.g. [Blackadar 1986] [1, Corollary 11.1.2]. This fact was proved independently by Gábor Etesi in terms of a von Neumann algebra related to $\mathbb{E}_{\mathscr{M}}$. Namely, such an algebra was shown to be hyperfinite, see [Etesi 2017] [4, Lemma 2.3]. For the sake of clarity, we adapt the proof to the case of the C^* -algebra $\mathbb{E}_{\mathscr{M}}$.

Let $\mathcal{G} := Diff(\mathcal{M})/Diff_0(\mathcal{M})$. Consider a profinite completion of the discrete group \mathcal{G} , i.e.

$$\widehat{\mathcal{G}} := \lim \mathcal{G}/N,\tag{3.1}$$

where N ranges through the open normal finite index subgroups of \mathcal{G} . Recall that if G is a finite group, then the group algebra $\mathbf{C}[G]$ has the form

$$\mathbf{C}[G] \cong M_{n_1}(\mathbf{C}) \oplus \dots \oplus M_{n_h}(\mathbf{C}), \qquad (3.2)$$

where n_i are degrees of the irreducible representations of G and h is the total number of such representations [Serre 1967] [14, Proposition 10]. In view of (3.1), we have

$$\widehat{\mathcal{G}} \cong \varprojlim G_i, \tag{3.3}$$

where G_i is a finite group. Consider a group algebra

$$\mathbf{C}[G_i] \cong M_{n_1}^{(i)}(\mathbf{C}) \oplus \dots \oplus M_{n_h}^{(i)}(\mathbf{C})$$
(3.4)

corresponding to G_i . Notice that the $\mathbb{C}[G_i]$ is a finite-dimensional C^* -algebra. The inverse limit (3.3) defines an ascending sequence of the finite-dimensional C^* algebras of the form

$$\varprojlim M_{n_1}^{(i)}(\mathbf{C}) \oplus \dots \oplus M_{n_h}^{(i)}(\mathbf{C}).$$
(3.5)

The group C^* -algebra $C^*(\widehat{\mathcal{G}})$ of the profinite group $\widehat{\mathcal{G}}$ is the norm closure of the group algebra $\mathbb{C}[\widehat{\mathcal{G}}]$ [Dixmier 1977] [2, Section 13.9]. One concludes from (3.5) that the $C^*(\widehat{\mathcal{G}})$ is an AF-algebra.

On the other hand, the canonical homomorphism $\mathcal{G} \to \widehat{\mathcal{G}}$ gives rise to an extension of the C^* -algebras:

$$C^*(\mathcal{G}) \to C^*(\widehat{\mathcal{G}}) \to \mathscr{B}.$$
 (3.6)

Since $C^*(\widehat{\mathcal{G}})$ is an AF-algebra, both $C^*(\mathcal{G})$ and \mathscr{B} must be AF-algebras [Handelman 1982] [7, Lemma I.5(a)]. It remains to recall that $C^*(\mathcal{G}) := \mathbb{E}_{\mathscr{M}}$, see definition 1.1. Thus $\mathbb{E}_{\mathscr{M}}$ is an AF-algebra. Lemma 3.1 is proved.

Lemma 3.2. The AF-algebra $\mathbb{E}_{\mathscr{M}}$ is stationary.

Proof. (i) Let S^4 be the 4-dimensional sphere. By the Piergallini Theorem 2.3, there exists a 4-fold covering map $\mathscr{M} \to S^4$ branched at the points of a knotted surface \mathscr{X} defined by an embedding:

$$\mathscr{X} \hookrightarrow S^4.$$
 (3.7)

In view of the inclusion $Diff(S^4) \subset Diff(S^4 - \mathscr{X})$, one gets an injective homomorphism of the C^* -algebras:

$$\mathbb{E}_{S^4} \hookrightarrow \mathbb{E}_{\mathscr{M}}.\tag{3.8}$$

(ii) Let us show that $\mathbb{E}_{S^4} \cong \mathbb{C}$. Indeed, since $Diff(S^4) \cong Diff_0(S^4)$, the group $Diff(S^4)/Diff_0(S^4)$ is trivial. In particular, the group C^* -algebra \mathbb{E}_{S^4} is commutative. The Gelfand Theorem says that $\mathbb{E}_{S^4} \cong C_0(X)$, where $C_0(X)$ is the C^* -algebra of continuous complex-valued functions on a locally compact Hausdorff space X, see Section 2.1. But $X \cong \mathbf{pt}$ is a singleton and therefore $C_0(\mathbf{pt}) \cong \mathbb{C}$. Thus one gets $\mathbb{E}_{S^4} \cong \mathbb{C}$.

(iii) The AF-algebra \mathbb{E}_{S^4} is given by an ascending sequence (2.1) of the form:

$$\mathbf{C} \stackrel{\mathbf{1}}{\longrightarrow} \mathbf{C} \stackrel{\mathbf{1}}{\longrightarrow} \dots, \tag{3.9}$$

where **1** is the identity homomorphism. It follows from (3.9) that the \mathbb{E}_{S^4} is a stationary AF-algebra.

(iv) Since $K_0(\mathbf{C}) \cong \mathbf{Z}$, we conclude that the dimension group of the AF-algebra \mathbb{E}_{S^4} is isomorphic to $(\mathbf{Z}, \mathbf{Z}^+)$, where \mathbf{Z}^+ is the semi-group of positive integers. It is easy to see, that the Handelman triple corresponding to the stationary AF-algebra \mathbb{E}_{S^4} has the form $(\mathbf{Z}, [\mathbf{Z}], \mathbf{Q})$.

(v) On the other hand, the map (3.8) induces an inclusion of the abelian groups:

$$K_0(\mathbb{E}_{S^4}) \subset K_0(\mathbb{E}_{\mathscr{M}}). \tag{3.10}$$

Moreover, if τ is the canonical trace on the AF-algebra $\mathbb{E}_{\mathscr{M}}$, one gets from (3.10) the following inclusion of the additive groups of the real line:

$$\mathbf{Z} \subset \tau_*(K_0(\mathbb{E}_{\mathscr{M}})). \tag{3.11}$$

(vi) Since **Z** is a ring, the group inclusion (3.11) can be extended to such of the rings. But the only finite degree extension of the ring **Z** coincides (up to a scaling constant) with an order Λ in the number field $K = \Lambda \otimes \mathbf{Q}$. We conclude that

$$\tau_*(K_0(\mathbb{E}_{\mathscr{M}})) \subset K, \tag{3.12}$$

where K is a real number field.

(vii) To finish the proof of lemma 3.2, it remains to apply the result of [Handelman 1981] [6, Theorem II (iii)] saying that condition (3.12) is equivalent to the AF-algebra $\mathbb{E}_{\mathscr{M}}$ to be of a stationary type.

Lemma 3.2 is proved.

Remark 3.3. The Etesi C^* -algebra $\mathbb{E}_{\mathscr{M}}$ is simple, i.e. has only trivial two-sided ideals. This fact follows from lemma 3.2 and strict positivity of the matrix A corresponding to the stationary AF-algebra.

Returning to the proof of theorem 1.3, we apply lemmas 3.1 and 3.2. Theorem 1.3 follows.

3.2. **Proof of corollary 1.4.** We split the proof in a series of lemmas.

Lemma 3.4. The Etesi C^* -algebras satisfy an isomorphism:

$$\mathbb{E}_{\mathcal{M}_1 \# \mathcal{M}_2} \cong \mathbb{E}_{\mathcal{M}_1} \otimes \mathbb{E}_{\mathcal{M}_2}, \tag{3.13}$$

where # is the connected sum of manifolds and \otimes is the tensor product of C^* -algebras.

Proof. We let $\mathcal{G} := Diff(\mathcal{M}_1 \# \mathcal{M}_2) / Diff_0(\mathcal{M}_1 \# \mathcal{M}_2)$. It is not hard to see, that

$$\mathcal{G} = \mathcal{G}_1 \times \mathcal{G}_2, \tag{3.14}$$

where $\mathcal{G}_1 = Diff(\mathcal{M}_1)/Diff_0(\mathcal{M}_1)$ and $\mathcal{G}_2 = Diff(\mathcal{M}_2)/Diff_0(\mathcal{M}_2)$. It is well known that the group ring $\mathbf{C}[\mathcal{G}]$ of the product (3.14) is given by the formula:

$$\mathbf{C}[\mathcal{G}] \cong \mathbf{C}[\mathcal{G}_1] \otimes \mathbf{C}[\mathcal{G}_2]. \tag{3.15}$$

Since the $\mathbb{E}_{\mathscr{M}}$ is a nuclear C^* -algebra, the norm closure of a self-adjoint representation of (3.15) defines an isomorphism $\mathbb{E}_{\mathscr{M}_1 \# \mathscr{M}_2} \cong \mathbb{E}_{\mathscr{M}_1} \otimes \mathbb{E}_{\mathscr{M}_2}$. Lemma 3.4 is proved.

Lemma 3.5. The Etesi C^* -algebra of the 4-manifold $S^2 \times S^2$ is given by the formula:

$$\mathbb{E}_{S^2 \times S^2} \cong M_4(\mathbf{C}). \tag{3.16}$$

Proof. (i) It follows from Piergallini's Theorem 2.3, that the 4-manifold $S^2 \times S^2$ is a 4-fold cover of the 4-sphere S^4 as one can see by factoring this covering geometrically as:

$$S^2 \times S^2 \cong \mathbb{C}P^1 \times \mathbb{C}P^1 \to \mathbb{C}P^2 \to S^4.$$
(3.17)

The covering map (3.17) induces a homomorphism of the C^* -algebras $\mathbb{E}_{S^2 \times S^2} \to \mathbb{E}_{S^4}$ and a homomorphism of the corresponding abelian groups:

$$K_0(\mathbb{E}_{S^2 \times S^2}) \to K_0(\mathbb{E}_{S^4}). \tag{3.18}$$

From (3.9) one gets $K_0(\mathbb{E}_{S^4}) \cong \mathbb{Z}$. In view of (3.17), the kernel of the map (3.18) is isomorphic to $\mathbb{Z}/4\mathbb{Z}$. Thus one gets an isomorphism:

$$K_0(\mathbb{E}_{S^2 \times S^2}) \cong \mathbf{Z}.\tag{3.19}$$

(ii) By theorem 1.3, the $\mathbb{E}_{S^2 \times S^2}$ is a stationary AF-algebra. If A is the corresponding matrix, then by (3.19) the eigenvalues of A must be rational and equal

to each other. In other words, the AF-algebra $\mathbb{E}_{S^2 \times S^2}$ corresponds to an ascending sequence (2.1) of the form:

$$M_2(\mathbf{C}) \otimes M_2(\mathbf{C}) \xrightarrow{\mathbf{1}} M_2(\mathbf{C}) \otimes M_2(\mathbf{C}) \xrightarrow{\mathbf{1}} \dots$$
 (3.20)

where $\mathbf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. Clearly, the inductive limit (3.20) corresponds to the C^* algebra $M_2(\mathbf{C}) \otimes M_2(\mathbf{C})$. We conclude that $\mathbb{E}_{S^2 \times S^2} \cong M_2(\mathbf{C}) \otimes M_2(\mathbf{C}) \cong M_4(\mathbf{C})$.
Lemma 3.5 is proved.

Corollary 3.6. The Etesi C^* -algebra $\mathbb{E}_{\mathcal{M}\#S(k)}$ is Morita equivalent to $\mathbb{E}_{\mathcal{M}}$.

Proof. Recall that the 4-manifold $S(k) = (S^2 \times S^2) \# \dots \# (S^2 \times S^2)$ is a connected sum of the k copies of $S^2 \times S^2$. From lemma 3.5 and formula (3.13) one gets an isomorphism:

$$\mathbb{E}_{S(k)} \cong \underbrace{M_4(\mathbf{C}) \otimes \cdots \otimes M_4(\mathbf{C})}_{k \text{ times}} \cong M_{4^k}(\mathbf{C}). \tag{3.21}$$

If \mathcal{M} is a smooth 4-manifold, then by lemma 3.4 and formula (3.21) one obtains an isomorphism:

$$\mathbb{E}_{\mathscr{M}\#S(k)} \cong \mathbb{E}_{\mathscr{M}} \otimes M_{4^k}(\mathbf{C}). \tag{3.22}$$

In other words, the C^* -algebras $\mathbb{E}_{\mathscr{M}}$ and $\mathbb{E}_{\mathscr{M}\#S(k)}$ are Morita equivalent. Corollary 3.6 is proved.

Lemma 3.7. The topological type of manifold \mathcal{M} is determined by the Handelman triple $(\Lambda, [\mathfrak{m}], K)$.

Proof. Our proof is based on Gompf's Stable Diffeomorphism Theorem 2.4 and corollary 3.6. We shall proceed in two steps.

(i) Recall that by Theorem 2.4 for any two smoothings \mathscr{M} and \mathscr{M}' of a topological manifold \mathscr{M}_{top} one can find an integer $k \geq 0$, such that $\mathscr{M} \# S(k)$ and $\mathscr{M}' \# S(k)$ are diffeomorphic. The corresponding C^* -algebras are isomorphic:

$$\mathbb{E}_{\mathcal{M}\#S(k)} \cong \mathbb{E}_{\mathcal{M}'\#S(k)}.$$
(3.23)

In view of corollary 3.6 and formula (3.23), the C^* -algebra $\mathbb{E}_{\mathscr{M}}$ is Morita equivalent to $\mathbb{E}_{\mathscr{M}\#S(k)}$ and the C^* -algebra $\mathbb{E}_{\mathscr{M}\#S(k)}$ is Morita equivalent to $\mathbb{E}_{\mathscr{M}'}$. Therefore the C^* -algebras $\mathbb{E}_{\mathscr{M}}$ and $\mathbb{E}_{\mathscr{M}'}$ are Morita equivalent by the transitivity property.

Thus the Morita equivalence class of the Etesi C^* -algebra $\mathbb{E}_{\mathscr{M}}$ consists of all 4-dimensional manifolds which are homeomorphic but not diffeomorphic to each other.

(ii) Recall that the $\mathbb{E}_{\mathscr{M}}$ is a stationary AF-algebra (lemma 3.2). The Morita equivalence classes of such AF-algebras are described by the Handelman triples $(\Lambda, [\mathfrak{m}], K)$, see [Handelman 1981] [6], [9, Theorem 3.5.4] or Section 2.2. Comparing this fact with the result of item (i), we conclude that the $(\Lambda, [\mathfrak{m}], K)$ is an invariant of the topological type of the manifold \mathscr{M} .

Lemma 3.7 is proved.

Remark 3.8. Lemma 3.7 says that the topological type \mathcal{M}_{top} of manifold \mathcal{M} is defined by the Morita equivalence class of the Etesi C^* -algebra $\mathbb{E}_{\mathcal{M}}$. In contrast, different smoothings of \mathcal{M}_{top} are defined by the isomorphism classes contained in the Morita equivalence class of $\mathbb{E}_{\mathcal{M}}$.

Lemma 3.9. Let K be a number field and let Mi(K) be the Minkowski group, see definition 2.2. Then:

(i) the map $K \to Mi$ (K) is a covariant functor which maps isomorphic number fields K to the isomorphic torsion abelian groups Mi (K);

(ii) $Mi(K) \cong Br(K)$.

Proof. (i) Let K be a number field corresponding to the Handelman triple $(\Lambda, [\mathfrak{m}], K)$. Denote by $[\mathbb{A}]$ the Morita equivalence class of stationary AF-algebras defined by the triple $(\Lambda, [\mathfrak{m}], K)$. Since τ is the canonical trace (i.e. $\tau_*(u) = 1$ for the order unit $u \in K_0^+(\mathbb{A})$), we conclude that $\tau_*(\Sigma(\mathbb{A})) \subset [0, 1]$ does not depend on $\mathbb{A} \in [\mathbb{A}]$. Thus from (2.2) one gets a correctly defined map $K \to \mathscr{Y} := Mi(K)$. It can be verified directly that if K' is a real embedding of K, then $Mi(K') \cong Mi(K)$. Item (i) is proved.

(ii) Let Br(K) be the Brauer group of a number field K. It is well known, that the map $K \to Br(K)$ is a covariant functor which maps isomorphic number fields K to the isomorphic torsion abelian groups Br(K). Comparing with item (i) we conclude that there exists a natural transformation between these two functors, see e.g. [9, Section 2.3]. In particular, such a transformation implies an isomorphism of the abelian groups Br(K) and Mi(K). Item (ii) and lemma 3.9 are proved. \Box

Remark 3.10. Lemma 3.9 can be viewed as part of a correspondence between the algebraic K-theory (e.g. the Milnor-Voevodsky K-theory) and the Galois cohomology. This subject is outside the scope of present note.

Corollary 3.11. Distinct smoothings of \mathscr{M} are classified by the elements of the Brauer group Br(K). In particular, all smoothings of \mathscr{M} form a torsion abelian group with the summation operation defined by formula (2.5) and the neutral element \mathscr{M}_0 , see Section 2.3.

Proof. (i) Recall that from (2.2) we have:

$$\mathscr{Y} = ?_n(\tau_*(\Sigma(\mathbb{E}_{\mathscr{M}}))). \tag{3.24}$$

Since the Minkowski function $?_n(x)$ is monotone, formula (3.24) defines a bijective correspondence between generators of the torsion abelian groups $Mi(K) \cong Br(K)$ and the scale $\Sigma(\mathbb{E}_{\mathcal{M}})$ of the Etesi C^* -algebra $\mathbb{E}_{\mathcal{M}}$.

(ii) Recall that the scale $\Sigma(\mathbb{E}_{\mathscr{M}})$ is a generating subset of $K_0^+(\mathbb{E}_{\mathscr{M}})$, i.e. for each $a \in K_0^+(\mathbb{E}_{\mathscr{M}})$ there exist $a_1, \ldots, a_r \in \Sigma(\mathbb{E}_{\mathscr{M}})$ such that $a = a_1 + \cdots + a_r$. We extend the correspondence of item (i) to a bijective map between the elements of the Brauer group Br(K) and the elements of positive cone $K_0^+(\mathbb{E}_{\mathscr{M}})$.

(iii) It is known that the pair $(K_0(\mathbb{E}_{\mathscr{M}}), K_0^+(\mathbb{E}_{\mathscr{M}}))$ is invariant of the Morita equivalence class of the AF-algebra $\mathbb{E}_{\mathscr{M}}$, while the triple $(K_0(\mathbb{E}_{\mathscr{M}}), K_0^+(\mathbb{E}_{\mathscr{M}}), \Sigma(\mathbb{E}_{\mathscr{M}}))$ is invariant of the isomorphism class of $\mathbb{E}_{\mathscr{M}}$, see Section 2.2. Moreover, the scale can be written as $\Sigma(\mathbb{E}_{\mathscr{M}}) = \{a \in K_0^+(\mathbb{E}_{\mathscr{M}}) \mid 0 \leq a \leq u\}$, where $u \in K_0^+(\mathbb{E}_{\mathscr{M}})$ is fixed. Thus running through all $u \in K_0^+(\mathbb{E}_{\mathscr{M}})$ one gets all possible scales $\Sigma(\mathbb{E}_{\mathscr{M}})$ and vice versa. In other words, the elements $u \in K_0^+(\mathbb{E}_{\mathscr{M}})$ parametrize isomorphism classes of $\mathbb{E}_{\mathscr{M}}$ within its Morita equivalence class.

(iv) To finish the proof of corollary 3.11, it remains to recall remark 3.8. Indeed, combing 3.8 with item (iii) we conclude that different smooth structures on \mathscr{M} are in bijection with the elements of $K_0^+(\mathbb{E}_{\mathscr{M}})$. Moreover, the $K_0^+(\mathbb{E}_{\mathscr{M}})$ has structure of a torsion abelian group isomorphic to the Brauer group Br(K), see item (ii). Corollary 3.11 is proved.

Corollary 1.4 follows from lemma 3.7 and corollary 3.11.

4. Remarks

Remark 4.1. The $\mathbb{E}_{\mathscr{M}}$ can be defined via unitary representation of the group $Diff(\mathscr{M})$ by the bounded linear operators on a Hilbert space [Etesi 2016] [3, Section 2]. Since $Diff(\mathscr{M})$ is a Fréchet manifold, the standard construction of the group C^* -algebra fails in general [Blackadar 1986] [1, Section 11.1]; hence definition 1.1. However, the two definitions are equivalent from the standpoint of representation theory. We refer the reader to [Etesi 2021] [4] for a remarkable new topological invariant of the 4-manifolds given by the Murray - von Neumann coupling constant of $\mathbb{E}_{\mathscr{M}}$.

Remark 4.2. Let \mathscr{M} be the 4-dimensional sphere S^4 . Since $\mathbb{E}_{S^4} \cong \mathbb{C}$ is a finitedimensional C^* -algebra, corollary 1.4 fails. On the other hand, it is known that $K_0(\mathbb{E}_{S^4}) \cong \mathbb{Z}$ and $K \cong \mathbb{Q}$. Denote by \mathscr{S} a subgroup of the Brauer group $Br(\mathbb{Q})$ consisting of all smoothings of S^4 . Assuming an analog of 1.4, one can recast the smooth Poincaré Conjecture as follows: Is the group \mathscr{S} trivial?

References

- 1. B. Blackadar, K-Theory for Operator Algebras, MSRI Publications, Springer, 1986.
- 2. J. Dixmier, C*-Algebras, North-Holland Publishing Company, 1977.
- G. Etesi, Gravity as a four dimensional algebraic quantum field theory, Adv. Theor. Math. Phys. 20 (2016), 1049-1082.
- G. Etesi, The universal von Neumann algebra of smooth four-manifolds, Adv. Theor. Math. Phys. 25 (2021), 995–1024.
- R. E. Gompf, Stable diffeomorphism of compact 4-manifolds, Topology Appl. 18 (1984), 115-120.
- D. Handelman, Positive matrices and dimension groups affiliated to C*-algebras and topological Markov chains, J. Operator Theory 6 (1981), 55-74.
- D. Handelman, Extensions for AF C*-algebras and dimension groups, Trans. Amer. Math. Soc. 271 (1982), 537-573.
- H. Minkowski, Zur Geometrie der Zahlen, Verhandlungen des III internationalen Mathematiker-Kongresses in Heidelberg, pp. 164-173, Berlin, 1904.
- 9. I. V. Nikolaev, *Noncommutative Geometry*, De Gruyter Studies in Math. **66**, Second Edition, Berlin, 2022.
- 10. I. V. Nikolaev, Arithmetic topology of 4-manifolds, arXiv:1907.03901
- G. Panti, Multidimensional continued fractions and a Minkowski function, Monatsh. Math. 154 (2008), 247-264.
- 12. R. Piergallini, Four-manifolds as 4-fold branched covers of S⁴, Topology **34** (1995), 497-508.
- 13. Ch. Schafhauser, Subalgebras of simple AF-algebras, Ann. of Math. 192 (2020), 309-352.
- 14. J. -P. Serre, Représentations Linéaires des Groupes Finis, Hermann Paris, 1967.

¹ DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. JOHN'S UNIVERSITY, 8000 UTOPIA PARKWAY, NEW YORK, NY 11439, UNITED STATES.