### Hardness of Learning Halfspaces with Massart Noise

Ilias Diakonikolas\* University of Wisconsin Madison ilias@cs.wisc.edu Daniel M. Kane<sup>†</sup> University of California, San Diego dakane@cs.ucsd.edu

June 29, 2022

#### Abstract

We study the complexity of PAC learning halfspaces in the presence of Massart (bounded) noise. Specifically, given labeled examples (x, y) from a distribution D on  $\mathbb{R}^n \times \{\pm 1\}$  such that the marginal distribution on x is arbitrary and the labels are generated by an unknown halfspace corrupted with Massart noise at rate  $\eta < 1/2$ , we want to compute a hypothesis with small misclassification error. Characterizing the efficient learnability of halfspaces in the Massart model has remained a longstanding open problem in learning theory.

Recent work gave a polynomial-time learning algorithm for this problem with error  $\eta + \epsilon$ . This error upper bound can be far from the information-theoretically optimal bound of  $OPT + \epsilon$ . More recent work showed that *exact learning*, i.e., achieving error  $OPT + \epsilon$ , is hard in the Statistical Query (SQ) model. In this work, we show that there is an exponential gap between the information-theoretically optimal error and the best error that can be achieved by a polynomial-time SQ algorithm. In particular, our lower bound implies that no efficient SQ algorithm can approximate the optimal error within any polynomial factor.

<sup>\*</sup>Supported by NSF Award CCF-1652862 (CAREER) and a Sloan Research Fellowship.

<sup>&</sup>lt;sup>†</sup>Supported by NSF Award CCF-1553288 (CAREER) and a Sloan Research Fellowship.

### 1 Introduction

#### 1.1 Background and Motivation

A halfspace, or Linear Threshold Function (LTF), is any function  $f : \mathbb{R}^m \to \{\pm 1\}$  of the form  $f(x) = \operatorname{sign}(w \cdot x - \theta)$ , for some weight vector  $w \in \mathbb{R}^m$  and threshold  $\theta \in \mathbb{R}$ . (The function sign :  $\mathbb{R} \to \{\pm 1\}$  is defined as  $\operatorname{sign}(t) = 1$  if  $t \ge 0$  and  $\operatorname{sign}(t) = -1$  otherwise.) Halfspaces are a fundamental class of Boolean functions that have been extensively studied in computational complexity and learning theory over several decades [MP68, Yao90, GHR92, STC00, O'D14]. The problem of learning an unknown halfspace is as old as the field of machine learning, starting with the Perceptron algorithm [Ros58, Nov62], and has been one of the most influential problems in this field with techniques such as SVMs [Vap98] and AdaBoost [FS97] coming out of its study.

In the realizable PAC model [Val84], i.e., when the labels are consistent with the target function, halfspaces are efficiently learnable via Linear Programming (see, e.g., [MT94]). In the presence of noisy data, the computational complexity of learning halfspaces depends on the underlying noise model. Here we study the complexity of learning halfspaces with Massart noise. In the Massart (or bounded noise) model, the label of each example x is flipped independently with probability  $\eta(x) \leq \eta$ , for some parameter  $\eta < 1/2$ . The flipping probability is strictly bounded above by 1/2, but can depend on the example x in a potentially adversarial manner. The following definition encapsulates the PAC learning problem in this noise model.

**Definition 1.1** (PAC Learning with Massart Noise). Let  $\mathcal{C}$  be a concept class of Boolean-valued functions over  $X = \mathbb{R}^m$ ,  $\mathcal{D}_x$  be a fixed but unknown distribution over X, and  $0 \leq \eta < 1/2$  be the noise parameter. Let  $f : X \to \{\pm 1\}$  be the unknown target concept with  $f \in \mathcal{C}$ . A Massart example oracle,  $\mathrm{EX}^{\mathrm{Mas}}(f, \mathcal{D}_x, \eta)$ , works as follows: Each time  $\mathrm{EX}^{\mathrm{Mas}}(f, \mathcal{D}_x, \eta)$  is invoked, it returns a labeled example (x, y), where  $x \sim \mathcal{D}_x$ , y = f(x) with probability  $1 - \eta(x)$  and y = -f(x) with probability  $\eta(x)$ , for some unknown function  $\eta(x) : X \to [0, 1/2)$  with  $\eta(x) \leq \eta$  for all  $x \in X$ . Let  $\mathcal{D}$  denote the joint distribution on (x, y) generated by the Massart example oracle. A PAC learning algorithm is given i.i.d. samples from  $\mathcal{D}$  and its goal is to output a hypothesis  $h : X \to \{\pm 1\}$ such that with high probability the error  $\mathbf{Pr}_{(x,y)\sim\mathcal{D}}[h(x) \neq y]$  is as small as possible. We will use  $\mathrm{OPT} \stackrel{\text{def}}{=} \inf_{g \in \mathcal{C}} \mathbf{Pr}_{(x,y)\sim\mathcal{D}}[g(x) \neq y]$  to denote the optimal misclassification error.

We emphasize here that, throughout this paper, we focus on *improper learning*, where the learning algorithm is allowed to output any polynomially evaluatable hypothesis.

The Massart noise model is a natural semi-random input model that was formulated in [MN06]. An equivalent noise model had already been defined in the 80s by Sloan and Rivest [Slo88, Slo92, RS94, Slo96] (under the name "malicious misclassification noise") and a very similar definition had been proposed and studied even earlier by Vapnik [Vap82]. The sample complexity of PAC learning halfspaces with Massart noise is well-understood. For example, it is known (see, e.g., [MN06]) that for any concept class C of VC dimension d,  $O(d/\epsilon^2)$  samples information-theoretically suffice to compute a hypothesis with misclassification error OPT +  $\epsilon$ . We note here that, by definition, we have that OPT =  $\mathbf{E}_{x\sim \mathcal{D}_x}[\eta(x)] \leq \eta$ . In particular, this implies that halfspaces on  $\mathbb{R}^m$  are learnable in the Massart model with  $O(m/\epsilon^2)$  samples.

In sharp contrast, our understanding of the *algorithmic aspects* of PAC learning various natural concept classes with Massart noise is startlingly poor and has remained a tantalizing open problem in computational learning theory since the 1980s. In [Slo88] (see also [Slo92]), Sloan defined the malicious misclassification noise model (an equivalent formulation of Massart noise) and asked whether there exists an efficient learning algorithm for Boolean disjunctions, i.e., ORs of Boolean literals — a very special case of halfspaces — in this model. About a decade later, Edith Cohen [Coh97]

asked the same question for the general class of halfspaces. The problem remained open, even for weak learning, and was highlighted in Avrim Blum's FOCS 2003 tutorial [Blu03]. Surprisingly, until fairly recently, it was not even known whether there exists an efficient algorithm that achieves misclassification error 49% for Massart halfspaces with noise rate upper bound of  $\eta = 1\%$ .

Recent work [DGT19] made the first algorithmic progress on this learning problem. Specifically, [DGT19] gave a poly $(m, 1/\epsilon)$  time learning algorithm for Massart halfspaces with error guarantee of  $\eta + \epsilon$ , where  $\eta$  is the upper bound on the Massart noise rate. This is an *absolute* error guarantee which cannot be improved in general — since it may well be the case that OPT =  $\eta$ . Motivated by [DGT19], more recent work [DIK<sup>+</sup>21] gave an efficient boosting algorithm, achieving error  $\eta + \epsilon$ for any concept class, assuming the existence of a weak learner for the class.

The aforementioned error bound of  $\eta + \epsilon$  can be very far from the information-theoretically optimum of OPT +  $\epsilon$ . We remind the reader that OPT =  $\mathbf{E}_{x \sim \mathcal{D}_x}[\eta(x)] \leq \eta$  and it could well be the case that OPT  $\ll \eta$ . Follow-up work [CKMY20] showed that *exact* learning, i.e., obtaining the *optimal* error of OPT +  $\epsilon$ , when OPT is very close to 1/2, requires super-polynomial time in the Statistical Query (SQ) model [Kea98]. Unfortunately, this SQ lower bound is very fragile, in the sense that it does not rule out *any* constant factor approximation C > 1.

The aforementioned progress notwithstanding, a very large gap remains in our understanding of the *approximate efficient learnability* of halfspaces in the presence of Massart noise.

### Is there a polynomial-time learning algorithm for Massart halfspaces achieving a relative (distribution-independent) error guarantee?

Specifically, is there a polynomial-time algorithm with misclassification error  $C \cdot \text{OPT} + \epsilon$ , for some constant C > 1? If not, can we efficiently achieve a polynomial approximation ratio, i.e., misclassification error of  $O(\text{OPT}^c) + \epsilon$ , for some universal constant 0 < c < 1? More generally, what is the best error (as a function of OPT and  $\eta$ ) that can be achieved in polynomial time?

As the main contribution of this paper, we prove a strong hardness result which rules out any polynomial factor (relative) approximation in the Statistical Query (SQ) model. Before we formally state our contributions, we require some background on SQ algorithms.

**Statistical Query Model.** Statistical Query (SQ) algorithms are a class of algorithms that are only allowed to query expectations of bounded functions of the underlying distribution rather than directly access samples. The SQ model was introduced by Kearns [Kea98] in the context of supervised learning as a natural restriction of the PAC model [Val84] and has been extensively studied in learning theory. A recent line of work [FGR<sup>+</sup>13, FPV15, FGV17, Fel17] generalized the SQ framework for search problems over distributions. The reader is referred to [Fel16] for a survey.

One can prove lower bounds on the complexity of SQ algorithms via an appropriate notion of *Statistical Query dimension*. Such a complexity measure was introduced in [BFJ<sup>+</sup>94] for PAC learning of Boolean functions and has been generalized to the unsupervised setting in [FGR<sup>+</sup>13, Fel17]. A lower bound on the SQ dimension of a learning problem provides an unconditional lower bound on the computational complexity of any SQ algorithm for the problem.

The class of SQ algorithms is fairly broad: a wide range of known algorithmic techniques in machine learning are known to be implementable using SQs. These include spectral techniques, moment and tensor methods, local search (e.g., Expectation Maximization), and many others (see, e.g., [CKL+06, FGR+13, FGV17]). In the context of PAC learning classes of Boolean functions (the topic of this paper), with the exception of learning algorithms using Gaussian elimination (in particular for the concept class of parities, see, e.g., [BKW03]), all known algorithms with non-trivial performance guarantees are either SQ or are implementable using SQs. Finally, we acknowledge very recent work [BBH<sup>+</sup>20] which established a near-equivalence between the SQ model and low-degree algorithms under mild assumptions.

### **1.2 Our Contributions**

Our main result shows that any SQ algorithm that achieves a constant or even a polynomial relative approximation for the problem of PAC learning Massart halfspaces (in the distribution-independent setting) requires at least super-polynomial number of statistical queries of inverse super-polynomial accuracy. Formally, we establish the following theorem:

**Theorem 1.2** (SQ Hardness for Halfspaces on  $\mathbb{R}^M$ ). Let OPT > 0 and  $M \in \mathbb{Z}_+$  be such that  $\log(M)/(\log \log(M))^3$  is at least a sufficiently large constant multiple of  $\log(1/\text{OPT})$ . There exists a parameter  $\tau = M^{-\Omega(\log(M)/(\log \log(M))^3 \log(1/\text{OPT}))}$  such that no SQ algorithm can learn a halfspace on  $\mathbb{R}^M$  in the presence of Massart noise with  $\eta = 1/3$  within error better than 1/polylog(M) with  $1/\tau$  queries of accuracy  $\tau$ . This holds even if the optimal classifier has error at most OPT.

As an immediate corollary of Theorem 1.2, by taking  $\log(1/\text{OPT}) = \sqrt{\log(M)}$ , we obtain a super-polynomial SQ lower bound against learning a hypothesis with error better than 1/polylog(M), even when error OPT (which is exponentially smaller) is possible. This a fortiori implies that it is SQ-hard to learn a hypothesis with error O(OPT) or even poly(OPT).

We note that the  $\eta = 1/3$  assumption is not needed in our SQ hardness result. We can replace the upper bound on the noise rate by an arbitrarily small constant  $\eta$ ; see Remark 2.16.

It is worth comparing Theorem 1.2 to the hardness result of Daniely [Dan16] for PAC learning halfspaces in the agnostic model. Daniely's result is qualitatively similar to our Theorem 1.2 with two differences: (1) The lower bound in [Dan16] only applies against the (much more challenging) agnostic model. (2) In the agnostic setting, it is hard to learn halfspaces within error significantly better than 1/2, rather than error 1/polylog(M) in the Massart setting. (We remind the reader that there exists an efficient Massart halfspace learning algorithm that achieves error arbitrarily close to  $\eta < 1/2$  [DGT19].) Theorem 1.2 proves an SQ lower bound for a much more benign noise model at the cost of allowing somewhat better error in polynomial time.

### 1.3 Related and Prior Work

We have already provided some background on the Massart noise model. Here we summarize the most relevant literature on learning halfspaces in related noise models.

**Random Classification Noise.** Random Classification Noise (RCN) [AL88] is the special case of Massart noise where each label is flipped with probability *exactly*  $\eta < 1/2$ . Halfspaces are known to be efficiently learnable to optimal accuracy in the (distribution-independent) PAC model with RCN [BFKV96, BFKV97]. In fact, it is well-known that any SQ learning algorithm [Kea98] can be transformed to an RCN noise tolerant learning algorithm — a fact that no longer holds in the presence of Massart noise. Roughly speaking, the ability of the Massart adversary to choose *whether* to flip a given label and, if so, with what probability, makes the algorithmic problem in this model challenging.

Agnostic Model. The agnostic model [Hau92, KSS94] is the more challenging model where an adversary is allowed to arbitrarily corrupt an arbitrary OPT < 1/2 fraction of the labels. In the distribution-independent setting, even *weak* agnostic learning of halfspaces (i.e., obtaining a hypothesis with any non-trivial accuracy) is known to be intractable. A long line of work (see,

e.g., [GR06, FGKP06]) has established NP-hardness of weak agnostic *proper* learning. (See [Fel15] for a survey on hardness of proper learning results.) More recently, [Dan16] gave super-polynomial lower bounds for *improper* learning, under certain average-case assumptions, and simultaneously established SQ lower bounds for the problem. Concretely, [Dan16] showed that no polynomial-time SQ algorithm for agnostically learning halfspaces on  $\mathbb{R}^m$  can compute a hypothesis with error  $1/2 - 1/m^c$ , for some constant c > 0, even for instances with optimal error OPT =  $2^{-\log^{1-\nu}(m)}$ , for some constant  $\nu \in (0, 1/2)$ .

Finally, it is worth noting that *exact* learning in the agnostic model is known to be hard even in the distribution-specific PAC model, and in particular under the Gaussian distribution [KK14, GGK20, DKZ20, DKPZ21].

**Prior SQ Lower Bound for Massart Model.** [CKMY20] showed an SQ lower bound of  $m^{\Omega(\log(1/\epsilon))}$  for learning halfspaces with Massart to error OPT +  $\epsilon$ , when OPT is close to 1/2. Specifically, [CKMY20] observed a connection between SQ learning with Massart noise and the *Correlational Statistical Query (CSQ)* model, a restriction of the SQ model defined in [BF02] (see also [Fel08, Fel11]). Given this observation, [CKMY20] deduced their SQ lower bound by applying *as a black-box* a previously known CSQ lower bound by Feldman [Fel11]. This approach is inherently limited to exact learning. Establishing lower bounds for approximate learning requires new ideas.

Efficient Algorithms for Distribution-Specific Learning. Finally, we note that  $poly(m, 1/\epsilon)$  time learning algorithms for homogeneous Massart halfspaces with optimal error guarantees have been developed when the marginal distribution on examples is well-behaved [ABHU15, ABHZ16, ZLC17, YZ17, ZSA20, DKTZ20a, DKTZ20b, DKK<sup>+</sup>20, DKK<sup>+</sup>21]. The hardness result obtained in this paper provides additional motivation for such distributional assumptions. As follows from our inapproximability result, without some niceness assumption on the distribution of examples, obtaining even extremely weak relative approximations to the optimal error is hard.

**Broader Context.** This work is part of the broader direction of understanding the computational complexity of robust high-dimensional learning in the distribution-independent setting. A long line of work, see, e.g., [KLS09, ABL17, DKK<sup>+</sup>16, LRV16, DKK<sup>+</sup>17, DKK<sup>+</sup>18, DKS18, KKM18, DKS19, DKK<sup>+</sup>19] and the recent survey [DK19], has given efficient robust learners for a range of high-dimensional estimation tasks (both supervised and unsupervised) in the presence of a small constant fraction of adversarial corruptions. These algorithmic results inherently rely on the assumption that the clean data is drawn from a "well-behaved" distribution.

On the other hand, the recent work [DGT19] established that efficient robust learners with non-trivial error guarantees are achievable even in the distribution-independent setting, under the more "benign" Massart model. This result provided compelling evidence that there are realistic noise models in which efficient algorithms are possible without imposing assumptions on the good data distribution. Conceptually, the result of this paper shows that, even in such benign noise models, there can be strong computational limitations in learnability — in the sense that it is computationally hard to achieve even weak relative approximations to the optimal error.

### 1.4 Overview of Techniques

At a high level, our proof leverages the SQ lower bound framework developed in [DKS17]. We stress that employing this framework in our context requires a number of new conceptual and technical ideas, as we explain in the proceeding discussion. Roughly speaking, [DKS17] established the following generic SQ-hardness result: Let A be a one-dimensional distribution that matches the first k moments with the standard Gaussian G. Suppose we want to distinguish between the standard high-dimensional Gaussian N(0, I) on  $\mathbb{R}^m$ and a distribution that is a copy of A in a random direction and is a standard Gaussian in the orthogonal complement. Then any SQ algorithm for this hypothesis testing task requires superpolynomial time (roughly,  $m^{\Omega(k)}$  time).

In the context of the current paper, we will in fact require a generalization of the latter generic result that holds even if the one-dimensional distribution A nearly matches the first k moments with G. Such a statement (Proposition 2.6) is not hard to derive from the techniques of [DKS17].

To leverage the aforementioned result in our circumstances, we would like to establish the existence of a distribution (X, Y) on  $\mathbb{R} \times \{\pm 1\}$  that corresponds to a halfspace with Massart noise such that both the distributions of X conditioned on Y = 1 (X | Y = 1) and X conditioned on Y = -1 (X | Y = -1) approximately match their first k moments with the standard Gaussian. Naturally, a number of obstacles arise while trying to achieve this. In particular, achieving the above goal directly is impossible for the following reason. Any distribution X that even approximately matches a constant number of low-order moments with the standard Gaussian will have  $\mathbf{E}[f(X)] \approx \mathbf{E}[f(G)]$  for any halfspace f. To see this, we can use the known fact (see, e.g.  $[\mathrm{DGJ}^+10]$ ) that f can be sandwiched between low-degree polynomials  $f_+ \geq f \geq f_-$  with  $\mathbf{E}[f_+(G) - f_-(G)]$  small. This implies that if both conditional distributions X | Y = 1 and X | Y = -1 approximately match their low-degree moments with G, then  $\mathbf{E}[f(X)|Y = 1]$  will necessarily be close to  $\mathbf{E}[f(X)|Y = -1]$ , which cannot hold with Massart noise.

In order to circumvent this obstacle, we will instead prove a super-polynomial SQ lower bound against learning degree-*d* polynomial threshold functions (PTFs) under the Gaussian distribution with Massart noise, for an appropriate (super-constant) value of the degree *d*. Since a degree-*d* PTF on X is equivalent to an LTF on  $X^{\otimes d}$  — a random variable in  $m^d$  dimensions — we will thus obtain an SQ lower bound for the original halfspace Massart learning problem. A similar idea was used in [Dan16] to prove SQ lower bounds in the agnostic model.

The next challenge is, of course, to construct the required moment-matching distributions in one dimension. Even for our reformulated PTF learning problem, it remains unclear whether this is even possible. For example, let  $f(x) = \operatorname{sign}(p(x))$  be a degree-*d* PTF. Then it will be the case that  $\mathbf{E}[p(X)Y] = \mathbf{E}[p(X)f(X)(1-2\eta(X))] = \mathbf{E}[|p(X)|(1-2\eta(X))] > 0$ . This holds despite the fact that  $\mathbf{E}[p(X)|Y = 1] \approx \mathbf{E}[p(X)|Y = -1] \approx \mathbf{E}[p(G)]$ . If  $\mathbf{E}[p(G)] > 0$ , it will be the case that  $\mathbf{E}[p(X)|Y = -1]$  will be positive, despite the fact that the conditional distribution of  $X \mid Y = -1$ is almost entirely supported on the region where p(X) < 0. Our construction will thus need to take advantage of finding points where |p(X)| is very large.

Fortunately for us, something of a miracle occurs here. Consider a discrete univariate Gaussian  $G_{\delta}$  with spacing  $\delta$  between its values. It is not hard to show that  $G_{\delta}$  approximately matches moments with the standard Gaussian G to error  $\exp(-\Omega(1/\delta^2))$  (Lemma 2.9). On the other hand, all but a tiny fraction of the probability mass of  $G_{\delta}$  is supported on  $d = \tilde{O}(1/\delta)$  points. We use this observation to construct one of our conditional distributions. In more detail, we will use a  $\delta$ -spaced discrete Gaussian, widened slightly by convolving it with a narrow Gaussian. We note that the univariate degree-d PTF that we are approximating will be a union of d/2 narrow intervals around the most significant peaks of this distribution. This construction gives us a distribution almost entirely supported on the negative values of a degree-d PTF, while still matching moments with a standard Gaussian to error approximately  $\exp(-\tilde{\Omega}(d^2))$ . Since this error is still small relative to the underlying dimension of the induced LTF problem, i.e.,  $m^d$ , this will provide us with the desired super-polynomial SQ lower bounds.

So far, we have described our construction of one of the conditional distributions (namely, the

distribution of X | Y = -1). Additionally, we require a distribution for the other conditional X | Y = 1 that nearly matches many of its low-degree moments with the Gaussian G and has much larger density when our PTF, f(x) is 1, and much smaller density when f(x) is -1. We achieve this as follows: We start with a standard Gaussian distribution. By making our final joint distribution on (X, Y) output Y = 1 with substantially higher probability than Y = -1, it is not hard to arrange that the conditional probability that Y = 1 is large when f(X) = 1. However, we still need to deal with the case where f(X) = -1 (which happens on a union of short intervals). To deal with this, we show how to move the mass from these short intervals to nearby regions, in such a way as to not affect any of the low-order moments. To achieve this, we make use of an LP duality argument (Lemma 2.13).

The one-dimensional moment matching construction described above and its proof of correctness are given in Proposition 2.8. Given this construction, we can use (an adaptation of) the SQ machinery in [DKS17] to complete the proof of Theorem 1.2.

#### 1.5 Preliminaries

**Notation.** For  $n \in \mathbb{Z}_+$ , we denote  $[n] \stackrel{\text{def}}{=} \{1, \ldots, n\}$ . We will use standard notation for norms of vectors and functions, that will be presented before it is used in subsequent sections. We use  $\mathbf{E}[X]$  for the expectation of random variable X and  $\mathbf{Pr}[\mathcal{E}]$  for the probability of event  $\mathcal{E}$ .

**Basics on Statistical Query Algorithms.** We will use the framework of Statistical Query (SQ) algorithms for problems over distributions introduced in [FGR<sup>+</sup>13]. We start by defining a decision problem over distributions.

**Definition 1.3** (Decision Problem over Distributions). We denote by  $\mathcal{B}(\mathcal{D}, D)$  the decision (or hypothesis testing) problem in which the input distribution D' is promised to satisfy either (a) D' = D or (b)  $D' \in \mathcal{D}$ , and the goal of the algorithm is to distinguish between these two cases.

We define SQ algorithms as algorithms that do not have direct access to samples from the distribution, but instead have access to an SQ oracle. We consider the following standard oracle.

**Definition 1.4** (STAT Oracle). For a tolerance parameter  $\tau > 0$  and any bounded function  $f : \mathbb{R}^n \to [-1, 1]$ , STAT $(\tau)$  returns a value  $v \in [\mathbf{E}_{x \sim D}[f(x)] - \tau, \mathbf{E}_{x \sim D}[f(x)] + \tau]$ .

We note that [FGR<sup>+</sup>13] introduced another related oracle, which is polynomially equivalent to STAT. Since we prove super-polynomial lower bounds here, there is no essential distinction between these oracles. To define the SQ dimension, we need the following definitions.

**Definition 1.5** (Pairwise Correlation). The pairwise correlation of two distributions with probability density functions  $D_1, D_2 : \mathbb{R}^m \to \mathbb{R}_+$  with respect to a distribution with density  $D : \mathbb{R}^m \to \mathbb{R}_+$ , where the support of D contains the supports of  $D_1$  and  $D_2$ , is defined as  $\chi_D(D_1, D_2) \stackrel{\text{def}}{=} \int_{\mathbb{R}^m} D_1(x)D_2(x)/D(x)dx - 1$ .

We remark that when  $D_1 = D_2$  in the above definition, the pairwise correlation is identified with the  $\chi^2$ -divergence between  $D_1$  and D, i.e.,  $\chi^2(D_1, D) \stackrel{\text{def}}{=} \int_{\mathbb{R}^m} D_1(x)^2 / D(x) dx - 1$ .

**Definition 1.6.** We say that a set of s distributions  $\mathcal{D} = \{D_1, \ldots, D_s\}$  over  $\mathbb{R}^m$  is  $(\gamma, \beta)$ -correlated relative to a distribution D if  $|\chi_D(D_i, D_j)| \leq \gamma$  for all  $i \neq j$ , and  $|\chi_D(D_i, D_j)| \leq \beta$  for i = j.

We are now ready to define our notion of dimension.

**Definition 1.7** (Statistical Query Dimension). For  $\beta, \gamma > 0$ , a decision problem  $\mathcal{B}(\mathcal{D}, D)$ , where D is a fixed distribution and  $\mathcal{D}$  is a family of distributions over  $\mathbb{R}^m$ , let s be the maximum integer such that there exists a finite set of distributions  $\mathcal{D}_D \subseteq \mathcal{D}$  such that  $\mathcal{D}_D$  is  $(\gamma, \beta)$ -correlated relative to D and  $|\mathcal{D}_D| \geq s$ . We define the *Statistical Query dimension* with pairwise correlations  $(\gamma, \beta)$  of  $\mathcal{B}$  to be s and denote it by  $\mathrm{SD}(\mathcal{B}, \gamma, \beta)$ .

Our proof bounds below the Statistical Query dimension of the considered learning problem. This implies lower bounds on the complexity of any SQ algorithm for the problem using the following standard result.

**Lemma 1.8** (Corollary 3.12 in [FGR<sup>+</sup>13]). Let  $\mathcal{B}(\mathcal{D}, D)$  be a decision problem, where D is the reference distribution and  $\mathcal{D}$  is a class of distributions. For  $\gamma, \beta > 0$ , let  $s = \text{SD}(\mathcal{B}, \gamma, \beta)$ . For any  $\gamma' > 0$ , any SQ algorithm for  $\mathcal{B}$  requires at least  $s \cdot \gamma'/(\beta - \gamma)$  queries to the  $\text{STAT}(\sqrt{\gamma + \gamma'})$  oracle.

# 2 SQ Hardness of Learning Halfspaces with Massart Noise: Proof of Theorem 1.2

The structure of this section is as follows: In Section 2.1, we review the SQ framework from [DKS17] with the necessary modifications required for our setting. In Section 2.2, we establish the existence of the one-dimensional distributions with the desired moment-matching properties. Finally, in Section 2.3, we put everything together to complete the proof of Theorem 1.2. Section 2.2 is the main technical contribution of this section.

### 2.1 Generic SQ Lower Bound Construction

We start with the following definition:

**Definition 2.1** (High-Dimensional Hidden Direction Distribution). For a distribution A on the real line with probability density function A(x) and a unit vector  $v \in \mathbb{R}^m$ , consider the distribution over  $\mathbb{R}^m$  with probability density function  $\mathbf{P}_v^A(x) = A(v \cdot x) \exp\left(-\|x - (v \cdot x)v\|_2^2/2\right)/(2\pi)^{(m-1)/2}$ . That is,  $\mathbf{P}_v$  is the product distribution whose orthogonal projection onto the direction of v is A, and onto the subspace perpendicular to v is the standard (m-1)-dimensional normal distribution.

We consider the following condition:

**Condition 2.2.** Let  $k \in \mathbb{Z}_+$  and  $\nu > 0$ . The distribution A is such that (i) the first k moments of A agree with the first k moments of N(0,1) up to error at most  $\nu$ , and (ii)  $\chi^2(A, N(0,1))$  is finite.

Note that Condition 2.2-(ii) above implies that the distribution A has a pdf, which we will denote by A(x). We will henceforth blur the distinction between a distribution and its pdf.

Our main result in this subsection makes essential use of the following key lemma:

**Lemma 2.3** (Correlation Lemma). Let  $k \in \mathbb{Z}_+$ . If the univariate distribution A satisfies Condition 2.2, then for all  $v, v' \in \mathbb{R}^m$ , with  $|v \cdot v'|$  less than a sufficiently small constant, we have that

$$|\chi_{N(0,I)}(\mathbf{P}_{v}^{A}, \mathbf{P}_{v'}^{A})| \le |v \cdot v'|^{k+1} \chi^{2}(A, N(0,1)) + \nu^{2} .$$
(1)

This lemma is a technical generalization of Lemma 3.4 from [DKS17], which applied under *exact* moment matching assumptions. The proof is deferred to Appendix A.

We will also use the following standard fact:

**Fact 2.4.** For any constant c > 0 there exists a set S of  $2^{\Omega_c(m)}$  unit vectors in  $\mathbb{R}^m$  such that any pair  $u, v \in S$ , with  $u \neq v$ , satisfies  $|u \cdot v| < c$ .

In fact, an appropriate size set of random unit vectors satisfies the above statement with high probability. We note that [DKS17] made use of a similar statement, albeit with different parameters.

We will establish an SQ lower bound for the following binary classification problem.

**Definition 2.5** (Hidden Direction Binary Classification Problem). Let A and B be distributions on  $\mathbb{R}$  satisfying Condition 2.2 with parameters  $k \in \mathbb{Z}_+$  and  $\nu \in \mathbb{R}_+$ , and let  $p \in (0, 1)$ . For  $m \in \mathbb{Z}_+$ and a unit vector  $v \in \mathbb{R}^m$ , define the distribution  $\mathbf{P}_v^{A,B,p}$  on  $\mathbb{R}^m \times \{\pm 1\}$  that returns a sample from  $(\mathbf{P}_v^A, 1)$  with probability p and a sample from  $(\mathbf{P}_v^B, -1)$  with probability 1 - p. The corresponding binary classification problem is the following: Given access to a distribution on labeled examples of the form  $\mathbf{P}_v^{A,B,p}$ , for a fixed but unknown unit vector v, output a hypothesis  $h : \mathbb{R}^m \to \{\pm 1\}$ such that  $\mathbf{Pr}_{(X|Y)\sim\mathbf{P}_v^{A,B,p}}[h(X) \neq Y]$  is (approximately) minimized.

Note that it is straightforward to obtain misclassification error  $\min\{p, 1 - p\}$  (as one of the identically constant functions achieves this guarantee). We show that obtaining slightly better error is hard in the SQ model. The following result is the basis for our SQ lower bounds:

**Proposition 2.6** (Generic SQ Lower Bound). Consider the classification problem of Definition 2.5. Let  $\tau \stackrel{\text{def}}{=} \nu^2 + 2^{-k} (\chi^2(A, N(0, 1)) + \chi^2(B, N(0, 1)))$ . Then any SQ algorithm that, given access to a distribution  $\mathbf{P}_v^{A,B,p}$  for an unknown  $v \in \mathbb{R}^m$ , outputs a hypothesis  $h : \mathbb{R}^m \to \{\pm 1\}$  such that  $\mathbf{Pr}_{(X,Y)\sim\mathbf{P}_v^{A,B,p}}[h(X) \neq Y] < \min(p, 1-p) - 2\sqrt{\tau}$  must either make queries of accuracy better than  $2\sqrt{\tau}$  or must make at least  $2^{\Omega(m)}\tau/(\chi^2(A, N(0, 1)) + \chi^2(B, N(0, 1)))$  statistical queries.

Proof of Proposition 2.6. The proof proceeds as follows: We start by defining a related hypothesis testing problem  $\mathcal{H}$  and show that  $\mathcal{H}$  efficiently reduces to our learning (search) problem. We then leverage the machinery of this section (specifically, Lemma 2.3 and Fact 2.4) to prove an SQ lower bound for  $\mathcal{H}$ , which in turns implies an SQ lower bound for our learning task.

Let S be a set of  $2^{\Omega(m)}$  unit vectors in  $\mathbb{R}^m$  whose pairwise inner products are at most a sufficiently small universal constant c. (In fact, any constant c < 1/2 suffices.) By Fact 2.4, such a set is guaranteed to exist. Given S, our hypothesis testing problem is defined as follows.

**Definition 2.7** (Hidden Direction Hypothesis Testing Problem). In the context of Definition 2.5, the testing problem  $\mathcal{H}$  is the task of distinguishing between: (i) the distribution  $\mathbf{P}_{v}^{A,B,p}$ , for v randomly chosen from S, and (ii) the distribution G' on  $\mathbb{R}^{m} \times \{\pm 1\}$ , where for  $(X,Y) \sim G'$  we have that X is a standard Gaussian  $G \sim N(0, I)$ , and Y is independently 1 with probability p and -1 with probability 1 - p.

We claim that  $\mathcal{H}$  efficiently reduces to our learning task. In more detail, any SQ algorithm that computes a hypothesis h satisfying  $\mathbf{Pr}_{(X,Y)\sim\mathbf{P}_v^{A,B,p}}[h(X)\neq Y] < \min(p,1-p) - 2\sqrt{\tau}$  can be used as a black-box to distinguish between  $\mathbf{P}_v^{A,B,p}$ , for v randomly chosen from S, and G'. Indeed, suppose we have such a hypothesis h. Then, with one additional query to estimate the  $\mathbf{Pr}[h(X)\neq Y]$ , we can distinguish between  $\mathbf{P}_v^{A,B,p}$ , for v randomly chosen from S, and G' for the following reason: For any function h, we have that  $\mathbf{Pr}_{(X,Y)\sim G'}[h(X)\neq Y] \geq \min(p,1-p)$ .

It remains to prove that solving the hypothesis testing problem  $\mathcal{H}$  is impossible for an SQ algorithm with the desired parameters. We will show this using Lemma 1.8.

More specifically, we need to show that for  $u, v \in S$  we have that  $|\chi_{G'}(\mathbf{P}_v^{A,B,p}, \mathbf{P}_u^{A,B,p})|$  is small. Since  $G', \mathbf{P}_v^{A,B,p}$ , and  $\mathbf{P}_u^{A,B,p}$  all assign Y = 1 with probability p, it is not hard to see that

$$\chi_{G'}(\mathbf{P}_{v}^{A,B,p}, \mathbf{P}_{u}^{A,B,p}) = p \; \chi_{G'|Y=1} \left( (\mathbf{P}_{v}^{A,B,p} \mid Y=1), (\mathbf{P}_{u}^{A,B,p} \mid Y=1) \right) + (1-p) \; \chi_{G'|Y=-1} \left( (\mathbf{P}_{v}^{A,B,p} \mid Y=-1), (\mathbf{P}_{u}^{A,B,p} \mid Y=-1) \right) \\ = p \; \chi_{G}(\mathbf{P}_{v}^{A}, \mathbf{P}_{u}^{A}) + (1-p) \; \chi_{G}(\mathbf{P}_{v}^{B}, \mathbf{P}_{u}^{B}).$$

By Lemma 2.3, it follows that

$$\chi_{G'}(\mathbf{P}_v^{A,B,p}, \mathbf{P}_u^{A,B,p}) \le \nu^2 + 2^{-k} (\chi^2(A, N(0,1)) + \chi^2(B, N(0,1))) = \tau$$

A similar computation shows that

$$\chi_{G'}(\mathbf{P}_v^{A,B,p}, \mathbf{P}_v^{A,B,p}) = \chi^2(\mathbf{P}_v^{A,B,p}, G') \le \chi^2(A, N(0,1)) + \chi^2(B, N(0,1)) .$$

An application of Lemma 1.8 for  $\gamma = \gamma' = \tau$  and  $\beta = \chi^2(A, N(0, 1)) + \chi^2(B, N(0, 1))$  completes the proof.

### 2.2 Construction of Univariate Moment-Matching Distributions

Here we give our univariate approximate moment-matching construction (Proposition 2.8), which is the main technical contribution of this section.

**Notation.** We will be working with moments of distributions that are best described as normalizations of (unnormalized positive) measures. For notational convenience, by slight abuse of notation, we define  $\mathbf{E}[X]$ , for any non-negative measure on  $\mathbb{R}$ , by  $\mathbf{E}[X] = \int t dX(t)$ . Note that this is equivalent to  $\mathbf{E}[X] = ||X||_1 \mathbf{E}[Y]$ , where  $Y = X/||X||_1$  is the normalized version of X. Furthermore, we denote the  $k^{th}$  moment of such an X by  $\mathbf{E}[X^k]$ .

We write  $E \gtrsim F$  for two expressions E and F to denote that  $E \geq cF$ , where c > 0 is a sufficiently large universal constant (independent of the variables or parameters on which E and F depend). Similarly, we write  $E \lesssim F$  to denote that  $E \leq cF$ , where c > 0 is a sufficiently small universal constant.

We view the real line as a measurable space endowed with the  $\sigma$ -algebra of Lebesgue measurable sets. We will construct two (non-negative, finite) measures  $\mathcal{D}_+$  and  $\mathcal{D}_-$  on this space with appropriate properties. Let  $S_c\mu$  be the scaling of measure  $\mu$  by a constant  $c \neq 0$ , i.e.,  $S_c\mu(A) = \mu((1/c)A)$ for Lebesgue measurable sets  $A \subset \mathbb{R}$ . We use f \* g for the convolution between two measures f, g.

We will use G for the measure of the univariate standard Gaussian distribution N(0,1) and  $g(x) = \frac{1}{\sqrt{2\pi}} \exp(-x^2/2)$  for its probability density function.

The main technical result of this section is captured in the following proposition.

**Proposition 2.8.** Let  $d, k \in \mathbb{Z}_+$  and  $\delta, \epsilon, \zeta \in (0, 1/2)$  with  $\delta > \epsilon$  and such that:

- (i)  $\epsilon \sqrt{\log(1/\zeta)}/\delta \lesssim 1/k^2$ ,
- (ii)  $\delta \lesssim d^{-1/2}$ , and
- (iii)  $d \gtrsim \sqrt{\log(1/\zeta)}/\delta$ .

There exist measures  $\mathcal{D}_+$  and  $\mathcal{D}_-$  over  $\mathbb{R}$  and a union J of d intervals on  $\mathbb{R}$  such that:

1. (a)  $\mathcal{D}_+ = 0$  on J, and (b)  $\mathcal{D}_+ > 2\mathcal{D}_-$  on  $\overline{J} = \mathbb{R} \setminus J$ .

- 2. All but  $\zeta$  fraction of the measure of  $\mathcal{D}_{-}$  lies in J.
- 3. The distributions  $\mathcal{D}_+/\|\mathcal{D}_+\|_1$  and  $\mathcal{D}_-/\|\mathcal{D}_-\|_1$  have their first k moments matching those of G within additive error at most  $(k+1)! \exp(-\Omega(1/\delta^2))$ .
- 4. (a)  $\mathcal{D}_+$  is at most  $O(\delta/\epsilon)$  G and (b)  $\|\mathcal{D}_+\|_1 = \Theta(\delta/\epsilon)$ .
- 5. (a)  $\mathcal{D}_{-} = O(\delta/\epsilon)G$ , and (b)  $\|\mathcal{D}_{-}\|_{1} \gtrsim 1$ .

**Discussion.** Essentially, in our final construction,  $\mathcal{D}_+$  will be proportional to the distribution of X conditioned on Y = 1 and  $\mathcal{D}_-$  proportional to X conditioned on Y = -1, with the ratio of the probability of Y = 1 to the probability of Y = -1 being equal to  $\|\mathcal{D}_+\|_1/\|\mathcal{D}_-\|_1$ . The Massart PTF that X is supposed to mimick will be -1 on J and 1 elsewhere (making it a degree-2d PTF). Property 1 says that Y will deterministically be -1 on J, while Properties 4 and 5 together show that Y will be 1 with probability at least 2/3 on  $\overline{J}$ , thus implying that Y is a noisy version of this PTF (with  $\eta \leq 1/3$ ). Property 2 says that Y only disagrees with this PTF with probability roughly  $\zeta$ , and this OPT will be less than  $\zeta$ . Property 3 says that  $\mathcal{D}_+$  and  $\mathcal{D}_-$ , after rescaling, approximately match many moments with the standard Gaussian, which will be necessary in applying our SQ lower bounds, while Properties 4 and 5 show that  $\mathcal{D}_+$  and  $\mathcal{D}_-$  have relatively small chi-squared norm.

The rest of this subsection is devoted to the proof of Proposition 2.8.

Proof of Proposition 2.8. Let  $G_{\delta}$ ,  $\delta > 0$ , denote the measure of the " $\delta$ -spaced discrete Gaussian distribution". In particular, for each  $n \in \mathbb{Z}$ ,  $G_{\delta}$  assigns mass  $\delta g(n\delta)$  to the point  $n\delta$ . Note that  $G_{\delta}$  is not a probability measure, as its total measure is not equal to one. However, it is not difficult to show (see Lemma 2.9 below) that the measure of  $G_{\delta}$  ic close to one for small  $\delta > 0$ , hence can be intuitively thought of as a probability distribution.

The following lemma shows that the moments of  $G_{\delta}$  approximately match the moments of the standard Gaussian measure G.

**Lemma 2.9.** For all  $t \ge 0$ , we have that  $\left| \mathbf{E}[G_{\delta}^t] - \mathbf{E}[G^t] \right| = t! O(\delta)^t \exp(-\Omega(1/\delta^2)).$ 

The proof of Lemma 2.9 proceeds by analyzing the Fourier transform of  $G_{\delta}$  and using the fact that the  $t^{th}$  moment of a measure is proportional to the  $t^{th}$  derivative of its Fourier transform at 0. The proof is deferred to Appendix B.

Note that Lemma 2.9 for t = 0 implies the total measure of  $G_{\delta}$  is  $\exp(-\Omega(1/\delta^2))$  close to 1.

**Definition of the Measure**  $\mathcal{D}_{-}$ **.** We define the measure  $\mathcal{D}_{-}$  as follows:

$$\mathcal{D}_{-} = S_{\sqrt{1-\epsilon^2}} G_{\delta} * S_{\epsilon} G , \qquad (2)$$

i.e.,  $\mathcal{D}_{-}$  is the convolution of a  $\sqrt{1-\epsilon^2}$  rescaled copy of  $G_{\delta}$  and an  $\epsilon$  rescaled independent copy of G. Formally speaking,  $\mathcal{D}_{-}$  is not a probability measure, because  $G_{\delta}$  (and, therefore, its scaling  $S_{\sqrt{1-\epsilon^2}}G_{\delta}$ ) does not have total measure exactly one. Intuitively,  $\mathcal{D}_{-}$  can be thought of as a countable (unnormalized) mixture of Gaussians (see Figure 1 for an illustration).

It will be useful to derive a closed-form formula for the density function of  $\mathcal{D}_-$ . Note that the measure  $S_{\sqrt{1-\epsilon^2}}G_{\delta}$  has mass function supported on the countable set  $\{x_n\}_{n\in\mathbb{Z}}$ , where  $x_n \stackrel{\text{def}}{=} n\delta\sqrt{1-\epsilon^2}$ , and its value at  $x_n$  is  $\delta g(n\delta)$ . Similarly, the probability measure  $S_{\epsilon}G$  is supported on the real line and its pdf at point  $x \in \mathbb{R}$  is  $(1/\epsilon)g(x/\epsilon)$ . By linearity of the convolution and the sifting property of the delta function, it follows that the density function of  $\mathcal{D}_{-}$  at x is

$$\sum_{n \in \mathbb{Z}} \delta g(n/\delta) (1/\epsilon) g((x - x_n)/\epsilon) = \frac{\delta}{\epsilon \sqrt{2\pi}} \sum_{n \in \mathbb{Z}} \exp\left(-(x - x_n)^2/(2\epsilon^2) - (n\delta)^2/2\right) .$$
(3)

Figure 1: Illustration of measure  $\mathcal{D}_{-}$ :  $\mathcal{D}_{-}$  is an unnormalized countable mixture of Gaussians, each of standard deviation  $\epsilon$ , with pairwise mean separation of  $\delta\sqrt{1-\epsilon^2} \approx \delta$ . The measure  $\mathcal{D}_{-}$  is bounded above by  $O(\delta/\epsilon)G$ . Almost all of the mass of  $\mathcal{D}_{-}$  lies in the union of d intervals J.

The following claim is the key ingredient to establish the desired moment matching condition (property 3 of Proposition 2.8) for the measure  $\mathcal{D}_{-}$ :

Claim 2.10. For  $t \leq k$ , we have that  $\left| \mathbf{E}[\mathcal{D}_{-}^{t}] - \mathbf{E}[G^{t}] \right| \leq t! \exp(-\Omega(1/\delta^{2})).$ 

Proof of Claim 2.10. For  $t \leq k$ , we can calculate the  $t^{th}$  moment of  $\mathcal{D}_{-}$  as follows

$$\begin{split} \mathbf{E}[\mathcal{D}_{-}^{t}] &= \sum_{i=0}^{t} \binom{t}{i} (1-\epsilon^{2})^{i/2} \epsilon^{t-i} \mathbf{E}[G_{\delta}^{i}] \mathbf{E}[G^{t-i}] \\ &= \sum_{i=0}^{t} \binom{t}{i} (1-\epsilon^{2})^{i/2} \epsilon^{t-i} \left( \mathbf{E}[G^{i}] \pm i! O(\delta)^{i} \exp\left(-\Omega(1/\delta^{2})\right) \right) \mathbf{E}[G^{t-i}] \\ &= \sum_{i=0}^{t} \binom{t}{i} (1-\epsilon^{2})^{i/2} \epsilon^{t-i} \mathbf{E}[G^{i}] \mathbf{E}[G^{t-i}] \pm t! \exp\left(-\Omega(1/\delta^{2})\right) \sum_{i=0}^{t} \binom{t}{i} O(\delta)^{i} \epsilon^{t-i} O(\sqrt{t})^{t-i} \\ &= \mathbf{E}[G^{t}] \pm t! \exp\left(-\Omega(1/\delta^{2})\right) O(\delta + \epsilon \sqrt{t})^{t} \\ &= \mathbf{E}[G^{t}] \pm t! \exp\left(-\Omega(1/\delta^{2})\right) \ , \end{split}$$

where the first line follows from the binomial theorem using the independence of  $S_{\sqrt{1-\epsilon^2}}G_{\delta}$  and  $S_{\epsilon}G$  (noting that  $\mathbf{E}[(S_c\mu)^j] = c^j \mathbf{E}[\mu^j]$  for any measure  $\mu, c \neq 0$ , and  $j \in \mathbb{Z}_+$ ), and the second line uses Lemma 2.9. The third line uses the fact that  $\mathbf{E}[G^j] = O(\sqrt{j})^j, j \in \mathbb{Z}_+$ . The fourth line follows by observing that if G, G' are independent standard Gaussians, the random variable X + Y, where  $X \sim S_{\sqrt{1-\epsilon^2}}G, Y \sim S_{\epsilon}G'$ , is a standard Gaussian. For the last line, note that  $\epsilon k^{1/2} < \epsilon k^2 < \delta$ , where the second inequality follows by assumption (i). By assumption (ii), it follows that  $\delta$  is at

most a sufficiently small universal constant. Therefore, the term  $O(\delta + \epsilon \sqrt{t})$  is less than 1. This completes the proof of Claim 2.10.

Claim 2.10 essentially directly implies property 3 of Proposition 2.8 for  $\mathcal{D}_-$ . A minor issue is that  $\mathcal{D}_-$  is itself not normalized, while property 3 refers to its normalized version  $\mathcal{D}_-/||\mathcal{D}_-||_1$ . To establish property 3, it suffices to bound from above the quantity  $|\mathbf{E}[\mathcal{D}_-^t] - \mathbf{E}[(\mathcal{D}_-/||\mathcal{D}_-||_1)^t]|$ , for all  $t \leq k$ . Claim 2.10 for t = 0 implies that  $|||\mathcal{D}_-||_1 - 1| \leq \exp(-\Omega(1/\delta^2))$ . We can thus write

$$\begin{aligned} \left| \mathbf{E}[\mathcal{D}_{-}^{t}] - \mathbf{E}[(\mathcal{D}_{-}/\|\mathcal{D}_{-}\|_{1})^{t}] \right| &= \mathbf{E}[\mathcal{D}_{-}^{t}] \left| 1 - 1/\|\mathcal{D}_{-}\|_{1}^{t} \right| \\ &\leq \left( t! + t! \exp(-\Omega(1/\delta^{2})) \right) t \exp(-\Omega(1/\delta^{2})) \\ &\leq (t+1)! \exp(-\Omega(1/\delta^{2})) , \end{aligned}$$

where the second line uses Claim 2.10, the fact that  $\mathbf{E}[G^t] \leq t!$ , and the elementary inequality  $(1+x)^r \geq 1+rx$ , for all  $x \geq -1$  and  $r \in \mathbb{Z}_+$ . This proves property 3 for  $\mathcal{D}_-$ .

We are now ready to establish the desired upper bound for  $\mathcal{D}_{-}$  (property 5(a) of Proposition 2.8):

**Claim 2.11.** For any measurable set  $A \subseteq \mathbb{R}$ , we have that  $\mathcal{D}_{-}(A) = O(\delta/\epsilon)G(A)$ .

Proof of Claim 2.11. We will show the desired inequality for the corresponding density functions. Recall that the density function of  $\mathcal{D}_{-}$  is given by (3). The exponent  $-(x - x_n)^2/(2\epsilon^2) - (n\delta)^2/2$  of the *n*-th term in (3), where  $x_n = n\delta\sqrt{1-\epsilon^2}$ , is a degree-2 polynomial in  $y = n\delta$ , which can be equivalently written as  $-(1/(2\epsilon^2))y^2 + (\sqrt{1-\epsilon^2}x/\epsilon^2)y - x^2/(2\epsilon^2)$ . This quadratic is maximized at  $y_0 = x\sqrt{1-\epsilon^2}$  and its maximum value is  $-x^2/2$ . Rewriting the quadratic in vertex form, we get  $-x^2/2 - (1/(2\epsilon^2))(y-y_0)^2$ . Since the square term is multiplied by a constant less than -1, the infinite sum of (3) can be bounded above (via a geometric series) by  $O((\delta/\epsilon) \exp(-x^2/2)) = O((\delta/\epsilon)g(x))$ . This completes the proof of Claim 2.11.

We also note that property 5(b) follows directly from Lemma 2.9.

**Definition of the Union of Intervals** *J*. We define *J* to be the union of *d* intervals,  $J_n$ , each of length  $L \stackrel{\text{def}}{=} C\epsilon \sqrt{\log(1/\zeta)}$ , for C > 0 a sufficiently large universal constant, centered at the points  $x_n = n\delta\sqrt{1-\epsilon^2}$ , for integers  $-\lfloor d/2 \rfloor \le n \le \lceil d/2 \rceil$ ; that is,  $J_n = [x_n - L/2, x_n + L/2]$ . See Figure 1 for an illustration.

The following claim establishes that at least  $1 - \zeta$  fraction of the measure of  $\mathcal{D}_{-}$  lies in J (property 2 of Proposition 2.8):

Claim 2.12. We have that  $\mathcal{D}_{-}(\overline{J}) \leq \zeta$ .

Proof. By Gaussian concentration, it follows that for each of the components of  $\mathcal{D}_-$  centered at  $x_n = n\delta\sqrt{1-\epsilon^2}$ , for an  $n \in \mathbb{Z}$  such that  $-\lfloor d/2 \rfloor \leq n \leq \lceil d/2 \rceil$ , all but a  $\zeta/2$ -fraction of their probability measure lies in J. It remains to argue that at most  $\zeta/2$  fraction of the total measure comes from the remaining components. This follows from the assumption that  $d \gtrsim \sqrt{\log(1/\zeta)}/\delta$ , which implies that  $\sum_{n \in \mathbb{Z}, |n| > d/2} \exp(-(n\delta)^2/2) \ll \zeta/2$ . This completes the proof of Claim 2.12.  $\Box$ 

**Definition of the Measure**  $\mathcal{D}_+$ . Intuitively, we would like to define  $\mathcal{D}_+$  to be equal to some suitable multiple M > 0 of the standard Gaussian measure G. Such a definition would satisfy the desired moment-matching conditions (property 3 of Proposition 2.8) with zero error and would also guarantee that  $\mathcal{D}_+ > \mathcal{D}_-$  on  $\overline{J}$ , as desired (property 1(b)). However, this candidate definition does not satisfy property 1(a), i.e., that  $\mathcal{D}_+$  be equal to 0 on J. To satisfy the latter property, we will need to carefully modify this the measure MG.



Figure 2: Illustration of the moment-matching construction given in Lemma 2.13. The measure  $\mathcal{D}$  is defined on [-x, x]. The modified measure  $\mathcal{D}'$  (i) is equal to zero in the subinterval [-c x, c x], (ii) matches the first k moments of  $\mathcal{D}$ , and (iii) is pointwise within a small constant factor of  $\mathcal{D}$  in  $[-x, x] \setminus [-c x, c x]$ .

The key technical tool to achieve this is the following lemma:

**Lemma 2.13.** Let  $x, c \in \mathbb{R}_+$ ,  $k \in \mathbb{Z}_+$ , and  $\mathcal{D}$  be a measure on [-x, x] such that the density function of  $\mathcal{D}$  varies by at most a factor of 2. If  $c \leq 1/k^2$ , there exists a measure  $\mathcal{D}'$  on [-x, x] such that:

- (i)  $\mathcal{D}'$  is 0 on [-cx, cx].
- (ii)  $\mathcal{D}'$  and  $\mathcal{D}$  agree on their first k moments.
- (iii)  $\mathcal{D}/2 \leq \mathcal{D}' \leq 3\mathcal{D}/2$  on  $[-x, x] \setminus [-cx, cx]$ .

*Proof.* First observe that we can assume without loss of generality that x = 1. We will prove the existence of  $\mathcal{D}'$  via LP duality. Indeed, we note that the three desired conditions on  $\mathcal{D}'$  define an LP. This LP has a feasible solution unless the dual LP has a solution. In particular, we show:

**Claim 2.14.** There exists a measure  $\mathcal{D}'$  satisfying (i), (ii), and (iii) unless there exists a degree at most k polynomial p such that

$$\left| \int_{-c}^{c} p(x) \mathcal{D}(x) \right| \ge \int_{[-1,1] \setminus [-c,c]} |p(x)| \mathcal{D}(x)/2 .$$

$$\tag{4}$$

*Proof.* We begin by assuming that Equation (4) has no solution. Note that the system in question can be rewritten as:

- (i)  $\int h(x)\mathcal{D}'(x) = 0$  for any h supported on [-c, c].
- (ii)  $\int p(x)\mathcal{D}'(x) = \mathbf{E}[p(\mathcal{D})]$  for any polynomial p of degree at most k.
- (iii)  $(1/2) \int h(x) \mathcal{D}(x) \leq \int h(x) \mathcal{D}'(x) \leq (3/2) \int h(x) \mathcal{D}(x)$  for any non-negative h supported on  $[-1,1] \setminus [-c,c]$ .

In other words, for any such degree at most k polynomial  $p, h_0 \in L^1([-c,c]), h_+, h_- \in L^1([-1,1] \setminus [-c,c])$  with  $h_+, h_-$  non-negative, we have that

$$\int (p(x) + h_0(x) + h_+(x) - h_-(x))\mathcal{D}'(x) \le \int (p(x) + (3/2)h_+(x) - (1/2)h_-(x))\mathcal{D}(x) \ .$$

Or, equivalently,

$$\int F(x)\mathcal{D}'(x) \le t$$

for any  $F = p + h_0 + h_+ - h_-$  and  $t \ge \int (p(x) + (3/2)h_+(x) - (1/2)h_-(x))\mathcal{D}(x)$  for some  $p, h_0, h_+, h_-$ . Given values of p and F, we note that for each  $x \in [-1, 1] \setminus [-c, c]$  that  $F(x) = p(x) + h_+(x) - h_-(x)$  and, therefore,  $(p(x) + (3/2)h_+(x) - (1/2)h_-(x)) = F(x) + (h_+(x) + h_-(x))/2 \ge F(x) + |p(x)|/2$ . Therefore,

$$t \ge \int_{-c}^{c} p(x)\mathcal{D}(x) + \int_{[-1,1]\setminus[-c,c]} (F(x) + |p(x)|/2)\mathcal{D}(x) .$$
(5)

We show that if there is no solution to Equation (4), there is a  $\mathcal{D}'$  with

$$\int F(x)\mathcal{D}'(x) \le t \; ,$$

whenever  $F(x) \in L^1([-1, 1])$ , and for some degree at most k polynomial p that Equation (5) holds.

Since the collection of such (F, t) is a cone, by [Fan68], as long as the set of such (F, t) is closed, there is such a  $\mathcal{D}'$ , unless we have some  $p, h_0, h_+, h_-$  for which F = 0 and t < 0. We note that unless there are solutions to Equation (4), that (0, t) cannot be in our cone for t < 0. It remains to show that this cone is closed.

Assuming there are no solutions to Equation (4), for any such polynomial p with  $||p||_2 = 1$ , we have that

$$\int_{-c}^{c} p(x)\mathcal{D}(x) + \int_{[-1,1]\setminus[-c,c]} |p(x)|\mathcal{D}(x)/2 > a$$

for some a > 0. By homogeneity, this implies that for any such p, we have

$$\int_{-c}^{c} p(x)\mathcal{D}(x) + \int_{[-1,1]\setminus[-c,c]} |p(x)|\mathcal{D}(x)/2 > a ||p||_2.$$

Assume that there is a sequence of points  $(F_n, t_n)$  in our cone converging to a limit  $(F_0, t_0)$ . We assume that the point  $(F_n, t_n)$  comes from a polynomial  $p_n$ . As

$$t_n \ge \int_{-c}^{c} p_n(x)\mathcal{D}(x) + \int_{[-1,1]\setminus[-c,c]} |p_n(x)|\mathcal{D}'(x)/2 + \int_{[-1,1]\setminus[-c,c]} F_n(x)\mathcal{D}(x) \ge a ||p_n||_2 + \int_{[-1,1]\setminus[-c,c]} F_n(x)\mathcal{D}(x)$$

we can see that  $|p_n|$  must be bounded. By compactness, there must be a subsequence of the  $p_n$  converging to some  $p_0$ . It is then easy to see taking the limits that

$$t_0 \ge \int_{-c}^{c} p_0(x)\mathcal{D}(x) + \int_{[-1,1]\setminus[-c,c]} (F_0(x) + |p_0(x)|/2)\mathcal{D}(x) + \int_$$

thus proving that  $(F_0, t_0)$  is in our cone. This shows that the cone is closed, completing our proof.

We are now left with determining whether or not there exists a degree at most k polynomial p(x) so that  $\left|\int_{-c}^{c} p(x)\mathcal{D}(x)\right| \geq \int_{[-1,1]\setminus[-c,c]} |p(x)|\mathcal{D}(x)/2$ . This in particular would imply that

$$c \sup_{[-1,1]} |p(x)| \gg \int_{-1}^{1} |p(x)| dx$$
,

or equivalently that  $c \|p\|_{\infty} \gg \|p\|_1$ . By Hölder's inequality, this implies that  $\sqrt{c} \|p\|_{\infty} \gg \|p\|_2$ .

Note that the measure on [-1,1] is identical to the projection of the measure on the 2-sphere. Therefore, lifting p to the 2-sphere, we get a degree at most k polynomial in the 2-sphere such that the max of |p| is  $\Omega(c^{-1/2})$  times the  $L_2$ -norm of p over the 2-sphere. We write p as a sum of the orthogonal polynomials  $\phi_i$  on the sphere, as  $p(x) = \sum_i a_i \phi_i(x)$ . Then we have that  $||p||_2^2 = \sum_i a_i^2$ . From this it is easy to see that  $|p(x)| \leq \sqrt{\sum_i \phi_i(x)^2} ||p||_2$ . However,  $\sum_i \phi_i(x)^2$  can easily be seen to be a spherically symmetric function with average value equal to the dimension of the space of degree at most k polynomials on the sphere, which is  $O(k^2)$ . Therefore, this function is a constant function, which is bounded by  $O(k^2)$ . Hence, for any such p, we have that  $||p||_{\infty} = O(k)||p||_2$ , which gives a contradiction when c is a sufficiently small constant multiple of  $1/k^2$ . This completes the proof of Lemma 2.13.



Figure 3: Illustration of the measure  $\mathcal{D}_+$ :  $\mathcal{D}_+$  is zero in J (union of red intervals), it is equal to MG at the tails, is within a constant factor of MG in the remaining intervals, and matches the first k moments of MG.

We are now ready to explain how to construct the measure  $\mathcal{D}_+$ . We start with the measure  $\mathcal{D}^0_+ = M \ G$ , for M > 0 a sufficiently large constant multiple of  $\delta/\epsilon$ . Then, for each integer  $-\lfloor d/2 \rfloor \leq n \leq \lceil d/2 \rceil$ , we apply Lemma 2.13 to the measure  $\mathcal{D}^0_+ |_{I_n}$ ,  $\mathcal{D}^0_+$  restricted to the interval

$$I_n = \left[ x_n - \delta \sqrt{1 - \epsilon^2}/2, x_n + \delta \sqrt{1 - \epsilon^2}/2 \right] ,$$

with the parameter c set to  $c = C\epsilon \sqrt{\log(1/\zeta)}/(\delta\sqrt{1-\epsilon^2})$ , recalling that  $x_n = n\delta\sqrt{1-\epsilon^2}$ , to replace  $\mathcal{D}^0_+|_{I_n}$  by another measure  $\mathcal{D}' =: \mathcal{D}_+|_{I_n}$ . See Figure 3 for an illustration.

By Lemma 2.13 (applied for  $\mathcal{D} = \mathcal{D}^0_+ |_{I_n}$  and integer  $-\lfloor d/2 \rfloor \leq n \leq \lceil d/2 \rceil$ ), the restriction of  $\mathcal{D}_+$  to  $I_n$  has matching first k moments with  $\mathcal{D}^0_+ |_{I_n}$  and is equal to zero on  $J_n$ , the corresponding interval of J. Note that our choice of  $I_n$  and c causes the sub-interval of length 2cx in the statement of Lemma 2.13 to correspond exactly to the interval  $J_n$  of J (since  $2x = \delta\sqrt{1-\epsilon^2}$ , hence 2cx = L).

It remains to verify that Lemma 2.13 can in fact be applied in our setting. That is, we want to argue that (1) the density of  $\mathcal{D}^0_+ = M G$  varies by at most a factor of 2 in each interval  $I_n$ , for integer  $-\lfloor d/2 \rfloor \leq n \leq \lfloor d/2 \rfloor$ , and (2)  $c \leq 1/k^2$ .

For the first statement, it suffices to show the desired claim for the pdf of G. Note that  $\max_{x \in I_n} g(x) / \min_{x \in I_n} g(x)$  is  $\exp(O(\delta^2))$  for n = 0. For  $n \neq 0$ , we have that

$$\max_{x \in I_n} g(x) / \min_{x \in I_n} g(x) = \exp(O(\delta^2((n+1/2)^2 - (n-1/2)^2)))) = \exp(O(n\delta^2)) = \exp(O(d\delta^2)) .$$

This quantity is at most 2, since by assumption we have that  $\delta \leq d^{-1/2}$ . For the second statement, note that by assumption we have that  $\epsilon \sqrt{\log(1/\zeta)}/\delta \leq 1/k^2$ . Therefore, we get that  $c \leq 1/k^2$ , as desired. Thus, the hypotheses of Lemma 2.13 are satisfied.

In summary, the final measure  $\mathcal{D}_+$  thus constructed matches its first k moments with those of M G, has no support on the union of intervals J, and elsewhere has magnitude  $\Theta(M G)$ . This gives properties 1(a), 3, and 4.

It only remains to verify property 1(b), i.e., the ratio of  $\mathcal{D}_+$  to  $\mathcal{D}_-$  on  $\overline{J}$ . By Claim 2.11 and since M is a sufficiently large constant multiple of  $\delta/\epsilon$ , the relative sizes of the distributions hold as desired. This completes the proof of Proposition 2.8.

### 2.3 Putting Everything Together: Proof of Theorem 1.2

We now have the necessary ingredients to complete the proof of Theorem 1.2.

*Proof of Theorem 1.2.* The proof leverages the SQ framework of Section 2.1 combined with the one-dimensional construction of Proposition 2.8.

**Parameter Setting.** Let C > 0 be a sufficiently large universal constant. We define positive integers m and d as follows:  $m = \lceil C \log(1/\tau) \rceil$  and  $d = \lceil C \sqrt{\log(1/\text{OPT}) \log(1/\tau) \log\log(1/\tau)} \rceil$ . Observe that

$$\binom{2d+m}{m} \le m^{2d} = \exp(O(C\sqrt{\log(1/\text{OPT})\log(1/\tau)(\log\log(1/\tau))^3})) .$$
(6)

We note that if  $\log(1/\tau)$  is a sufficiently small constant multiple of  $\log^2(M)/(\log \log(M))^3 \log(1/\text{OPT})$ , then the RHS of (6) is less than M. Thus, by decreasing M if necessary, we can assume that  $M = \binom{2d+m}{m}$ . Consider the Veronese mapping, denote by  $V_{2d} : \mathbb{R}^m \to \mathbb{R}^M$ , such that the coordinate functions of  $V_{2d}$  are exactly the monomials in m variables of degree at most 2d.

Hard Distributions. We can now formally construct the family of high-dimensional distributions on labeled examples that (1) corresponds to Massart halfspaces, and (2) is SQ-hard to learn. We define univariate measures  $\mathcal{D}_+$  and  $\mathcal{D}_-$  on  $\mathbb{R}$ , as given by Proposition 2.8 with  $\zeta = \text{OPT}$ ,  $\delta$  a sufficiently large constant multiple of  $\sqrt{\log(1/\zeta)}/d$ , k a small constant multiple of  $(1/\delta)^2/\log(1/\delta)$ , and  $\epsilon$  a small constant multiple of  $\delta/(\sqrt{\log(1/\zeta)}k^2)$ . It is easily verified that these parameters satisfy the assumptions of Proposition 2.8.

For a unit vector  $v \in \mathbb{R}^{m}$ , consider the distribution  $\mathbf{P}_{v}^{\mathcal{D}_{+},\mathcal{D}_{-},p}$ , as in Proposition 2.6, with  $p = \|\mathcal{D}_{+}\|_{1}/(\|\mathcal{D}_{+}\|_{1} + \|\mathcal{D}_{-}\|_{1})$ . Our hard distribution is the distribution (X',Y') on  $\mathbb{R}^{M} \times \{\pm 1\}$  obtained by drawing (X,Y) from  $\mathbf{P}_{v}^{\mathcal{D}_{+},\mathcal{D}_{-},p}$  and letting  $X' = V_{2d}(X)$  and Y' = Y.

**Claim 2.15.** The distribution (X', Y') on  $\mathbb{R}^M \times \{\pm 1\}$  is a Massart LTF distribution with optimal misclassification error OPT and Massart noise rate upper bound of  $\eta = 1/3$ .

Proof. For a unit vector  $v \in \mathbb{R}^m$ , let  $g_v : \mathbb{R}^m \to \{\pm 1\}$  be defined as  $g_v(x) = -1$  if and only if  $v \cdot x \in J$ , where J is the union of intervals in the construction of Proposition 2.8. Note that  $g_v$  is a degree-2d PTF on  $\mathbb{R}^m$ , since  $g_v$  is a (2d+1)-piecewise constant function of  $v \cdot x$ . Therefore, there exists some LTF  $L : \mathbb{R}^M \to \{\pm 1\}$  such that  $g_v(x) = L(V_{2d}(x))$  for all  $x \in \mathbb{R}^m$ .

Note that our hard distribution returns (X', Y') with Y' = L(X'), unless it picked a sample corresponding to a sample of  $\mathcal{D}_-$  coming from  $\overline{J}$ , which happens with probability at most  $\zeta = OPT$ . Additionally, suppose that our distribution returned a sample with  $X' = V_{2d}(X)$ , for some  $X \in \mathbb{R}^m$ . By construction, conditioned on this event, we have that Y' = 1 with probability proportional to  $\mathcal{D}_+(v \cdot X)$ , and Y' = -1 with probability proportional to  $\mathcal{D}_-(v \cdot X)$ . We note that if  $L(V_{2d}(X)) = 1$ , then  $v \cdot X \notin J$ ; so, by Proposition 2.8 property 1(b), this ratio is at least 2 : 1. On the other hand, if  $L(V_{2d}(X)) = -1$ , then  $v \cdot X \in J$ , so  $\mathcal{D}_+(v \cdot X) = 0$ . This implies that the pointwise probability of error  $\eta(X')$  is at most 1/3, completing the proof of the claim.

We are now ready to complete the proof of our SQ lower bound. It is easy to see that finding a hypothesis that predicts Y' given X' is equivalent to finding a hypothesis for Y given X (since Y = Y' and there is a known 1-1 mapping between X and X'). The pointwise bounds on  $\mathcal{D}_+$  and  $\mathcal{D}_-$ , specifically properties 4 and 5 in Proposition 2.8, imply that

$$\chi^{2}(\mathcal{D}_{+}/\|\mathcal{D}_{+}\|_{1}, G) \leq \chi^{2}(O(\delta/\epsilon)G, G)/\|\mathcal{D}_{+}\|_{1} = O(\delta/\epsilon)$$

and

$$\chi^2(\mathcal{D}_-/\|\mathcal{D}_-\|_1, G) \le \chi^2(O(\delta/\epsilon)G, G)/\|\mathcal{D}_-\|_1 = O(\delta/\epsilon)^2.$$

Therefore, we get that

$$\chi^2(\mathcal{D}_+/\|\mathcal{D}_+\|_1, G) + \chi^2(\mathcal{D}_-/\|\mathcal{D}_-\|_1, G) = O(\delta/\epsilon)^2 = \text{polylog}(M)$$
.

The parameter  $\nu$  in Proposition 2.6 is  $k! \exp(-\Omega(1/\delta^2)) = \exp(-\Omega(1/\delta^2))$ . We note that k is a small constant multiple of  $(1/\delta)^2/\log(1/\delta)$ , which is a large constant multiple of

 $d^2/(\log(1/\text{OPT})/\log(1/\delta)) = C \log(1/\tau) \log \log(1/\tau) / \log \log(1/\delta) \gg C \log(1/\tau)$ .

Since  $(1/\delta)^2$  is larger than the above, our value of  $\nu$  is at most a large constant power of  $\tau$ . Thus, the  $\tau$  parameter relevant to our application of Proposition 2.6 is equal to  $\nu^2 + 2^{-k}(\chi^2(\mathcal{D}_+, G) + \chi^2(\mathcal{D}_-, G))$ , which is at most a large constant power of our "real"  $\tau$ . It is furthermore easy to see that  $\min(p, 1-p) = 1 - p = \Theta(\epsilon/\delta) = 1/\text{polylog}(M)$ .

Therefore, Proposition 2.6 implies that it is impossible for an SQ algorithm to learn a hypothesis with error better than 1/polylog(M) without either using queries of accuracy better than  $\tau$  or making at least  $2^{\Omega(m)}\tau/\text{polylog}(M) > 1/\tau$  many queries. This completes the proof of Theorem 1.2.

**Remark 2.16.** Note that if we rescale  $\mathcal{D}_+$  and  $\mathcal{D}_-$ , they will still satisfy Condition 2.2. Therefore, by Proposition 2.6, it will still be hard for an SQ algorithm to find a hypothesis with error better than  $\min(p, 1-p) - 2\sqrt{\tau}$ . There are a couple of settings of interest here.

On the one hand, if we rescale  $\mathcal{D}_+$  so that  $||\mathcal{D}_+||_1 = ||\mathcal{D}_-||_1$ , then  $\min(p, 1-p)$  will equal 1/2. Therefore, in this case, it will be SQ-hard to learn to error substantially better than 1/2 even though OPT will still be close to 0. It should be noted, however, that this rescaling will break Property 1(b). This means that Y will no longer be an LTF of X with Massart noise. Interestingly, this provides another proof of the result of [Dan16] that distribution-independent agnostic learning of halfspaces is SQ-hard, even to get error slightly better than 1/2 and when the optimal error is close to 0. Another useful rescaling is to increase the size of  $\mathcal{D}_+$  by a factor of  $1/\eta$ , for some  $1/2 > \eta > 0$ . It is easy to see that after making this replacement, Y is now an LTF of X with Massart noise with parameter at most  $\eta$  (rather than parameter 1/3). This costs our result a bit in that now  $\min(p, 1-p) = \eta/\operatorname{poly}(\log(M))$ . However, this shows that learning an LTF with Massart noise rate  $\eta$  to error better than  $\eta/\operatorname{poly}(\log(M))$  is SQ-hard, even when OPT is substantially smaller.

### References

- [ABHU15] P. Awasthi, M. F. Balcan, N. Haghtalab, and R. Urner. Efficient learning of linear separators under bounded noise. In *Proceedings of The 28th Conference on Learning Theory, COLT 2015*, pages 167–190, 2015.
- [ABHZ16] P. Awasthi, M. F. Balcan, N. Haghtalab, and H. Zhang. Learning and 1-bit compressed sensing under asymmetric noise. In *Proceedings of the 29th Conference on Learning Theory, COLT 2016*, pages 152–192, 2016.
- [ABL17] P. Awasthi, M. F. Balcan, and P. M. Long. The power of localization for efficiently learning linear separators with noise. J. ACM, 63(6):50:1–50:27, 2017.
- [AL88] D. Angluin and P. Laird. Learning from noisy examples. *Mach. Learn.*, 2(4):343–370, 1988.
- [BBH<sup>+</sup>20] M. Brennan, G. Bresler, S. B. Hopkins, J. Li, and T.Schramm. Statistical query algorithms and low-degree tests are almost equivalent. *CoRR*, abs/2009.06107, 2020.
- [BF02] N. Bshouty and V. Feldman. On using extended statistical queries to avoid membership queries. *Journal of Machine Learning Research*, 2:359–395, 2002.
- [BFJ<sup>+</sup>94] A. Blum, M. Furst, J. Jackson, M. Kearns, Y. Mansour, and S. Rudich. Weakly learning DNF and characterizing statistical query learning using Fourier analysis. In *Proceedings of the Twenty-Sixth Annual Symposium on Theory of Computing*, pages 253–262, 1994.
- [BFKV96] A. Blum, A. M. Frieze, R. Kannan, and S. Vempala. A polynomial-time algorithm for learning noisy linear threshold functions. In 37th Annual Symposium on Foundations of Computer Science, FOCS '96, pages 330–338, 1996.
- [BFKV97] A. Blum, A. Frieze, R. Kannan, and S. Vempala. A polynomial time algorithm for learning noisy linear threshold functions. *Algorithmica*, 22(1/2):35–52, 1997.
- [BKW03] A. Blum, A. Kalai, and H. Wasserman. Noise-tolerant learning, the parity problem, and the statistical query model. J. ACM, 50(4):506–519, 2003.
- [Blu03] A. Blum. Machine learning: My favorite results, directions, and open problems. In 44th Symposium on Foundations of Computer Science (FOCS 2003), pages 11–14, 2003.
- [CKL<sup>+</sup>06] C.-T. Chu, S. K. Kim, Y. A. Lin, Y. Yu, G. Bradski, A. Y. Ng, and K. Olukotun. Map-reduce for machine learning on multicore. In *Proceedings of the 19th International Conference on Neural Information Processing Systems*, NIPS'06, pages 281–288, Cambridge, MA, USA, 2006. MIT Press.
- [CKMY20] S. Chen, F. Koehler, A. Moitra, and M. Yau. Classification under misspecification: Halfspaces, generalized linear models, and connections to evolvability. *CoRR*, abs/2006.04787, 2020.
- [Coh97] E. Cohen. Learning noisy perceptrons by a perceptron in polynomial time. In Proceedings of the Thirty-Eighth Symposium on Foundations of Computer Science, pages 514–521, 1997.

- [Dan16] A. Daniely. Complexity theoretic limitations on learning halfspaces. In Proceedings of the 48th Annual Symposium on Theory of Computing, STOC 2016, pages 105–117, 2016.
- [DGJ<sup>+</sup>10] I. Diakonikolas, P. Gopalan, R. Jaiswal, R. Servedio, and E. Viola. Bounded independence fools halfspaces. SIAM J. on Comput., 39(8):3441–3462, 2010.
- [DGT19] I. Diakonikolas, T. Gouleakis, and C. Tzamos. Distribution-independent PAC learning of halfspaces with massart noise. In Hanna M. Wallach, Hugo Larochelle, Alina Beygelzimer, Florence d'Alché-Buc, Emily B. Fox, and Roman Garnett, editors, Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019, pages 4751–4762, 2019.
- [DIK<sup>+</sup>21] I. Diakonikolas, R. Impagliazzo, D. M. Kane, R. Lei, J. Sorrell, and C. Tzamos. Boosting in the presence of massart noise. In *Conference on Learning Theory, COLT 2021*, volume 134 of *Proceedings of Machine Learning Research*, pages 1585–1644. PMLR, 2021.
- [DK19] I. Diakonikolas and D. M. Kane. Recent advances in algorithmic high-dimensional robust statistics. CoRR, abs/1911.05911, 2019.
- [DKK<sup>+</sup>16] I. Diakonikolas, G. Kamath, D. M. Kane, J. Li, A. Moitra, and A. Stewart. Robust estimators in high dimensions without the computational intractability. In *Proceedings* of FOCS'16, pages 655–664, 2016.
- [DKK<sup>+</sup>17] I. Diakonikolas, G. Kamath, D. M. Kane, J. Li, A. Moitra, and A. Stewart. Being robust (in high dimensions) can be practical. In *Proceedings of the 34th International Conference on Machine Learning, ICML 2017*, pages 999–1008, 2017.
- [DKK<sup>+</sup>18] I. Diakonikolas, G. Kamath, D. M. Kane, J. Li, A. Moitra, and A. Stewart. Robustly learning a gaussian: Getting optimal error, efficiently. In *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2018*, pages 2683–2702, 2018. Full version available at https://arxiv.org/abs/1704.03866.
- [DKK<sup>+</sup>19] I. Diakonikolas, G. Kamath, D. Kane, J. Li, J. Steinhardt, and A. Stewart. Sever: A robust meta-algorithm for stochastic optimization. In *Proceedings of the 36th International Conference on Machine Learning, ICML 2019*, pages 1596–1606, 2019.
- [DKK<sup>+</sup>20] I. Diakonikolas, D. Kane, V. Kontonis, C. Tzamos, and N. Zarifis. A polynomial time algorithm for learning halfspaces with tsybakov noise. *CoRR*, abs/2010.01705, 2020.
- [DKK<sup>+</sup>21] I. Diakonikolas, D. M. Kane, V. Kontonis, C. Tzamos, and N. Zarifis. Efficiently learning halfspaces with tsybakov noise. *STOC*, 2021.
- [DKPZ21] I. Diakonikolas, D. M. Kane, T. Pittas, and N. Zarifis. The optimality of polynomial regression for agnostic learning under gaussian marginals. In *Proceedings of The 34th Conference on Learning Theory, COLT*, 2021.
- [DKS17] I. Diakonikolas, D. M. Kane, and A. Stewart. Statistical query lower bounds for robust estimation of high-dimensional gaussians and gaussian mixtures. In 58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, pages 73–84, 2017. Full version at http://arxiv.org/abs/1611.03473.

- [DKS18] I. Diakonikolas, D. M. Kane, and A. Stewart. Learning geometric concepts with nasty noise. In Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2018, pages 1061–1073, 2018.
- [DKS19] I. Diakonikolas, W. Kong, and A. Stewart. Efficient algorithms and lower bounds for robust linear regression. In *Proceedings of the Thirtieth Annual ACM-SIAM Sympo*sium on Discrete Algorithms, SODA 2019, pages 2745–2754, 2019.
- [DKTZ20a] I. Diakonikolas, V. Kontonis, C. Tzamos, and N. Zarifis. Learning halfspaces with massart noise under structured distributions. In Jacob D. Abernethy and Shivani Agarwal, editors, Conference on Learning Theory, COLT 2020, volume 125 of Proceedings of Machine Learning Research, pages 1486–1513. PMLR, 2020.
- [DKTZ20b] I. Diakonikolas, V. Kontonis, C. Tzamos, and N. Zarifis. Learning halfspaces with tsybakov noise. *CoRR*, abs/2006.06467, 2020.
- [DKZ20] I. Diakonikolas, D. M. Kane, and N. Zarifis. Near-optimal SQ lower bounds for agnostically learning halfspaces and relus under gaussian marginals. *CoRR*, abs/2006.16200, 2020. In NeurIPS'20.
- [Fan68] K. Fan. On infinite systems of linear inequalities. Journal of Mathematical Analysis and Applications, 21(3):475 – 478, 1968.
- [Fel08] V. Feldman. Evolvability from learning algorithms. In Cynthia Dwork, editor, Proceedings of the 40th Annual ACM Symposium on Theory of Computing, 2008, pages 619–628. ACM, 2008.
- [Fel11] V. Feldman. Distribution-independent evolvability of linear threshold functions. In Sham M. Kakade and Ulrike von Luxburg, editors, COLT 2011 - The 24th Annual Conference on Learning Theory, June 9-11, 2011, volume 19 of JMLR Proceedings, pages 253–272. JMLR.org, 2011.
- [Fel15] V. Feldman. Hardness of proper learning (1988; pitt, valiant). In Encyclopedia of Algorithms. 2015.
- [Fel16] V. Feldman. Statistical query learning. In *Encyclopedia of Algorithms*, pages 2090–2095. 2016.
- [Fel17] V. Feldman. A general characterization of the statistical query complexity. In Satyen Kale and Ohad Shamir, editors, Proceedings of the 30th Conference on Learning Theory, COLT 2017, volume 65 of Proceedings of Machine Learning Research, pages 785– 830. PMLR, 2017.
- [FGKP06] V. Feldman, P. Gopalan, S. Khot, and A. Ponnuswami. New results for learning noisy parities and halfspaces. In *Proc. FOCS*, pages 563–576, 2006.
- [FGR<sup>+</sup>13] V. Feldman, E. Grigorescu, L. Reyzin, S. Vempala, and Y. Xiao. Statistical algorithms and a lower bound for detecting planted cliques. In *Proceedings of STOC'13*, pages 655–664, 2013. Full version in Journal of the ACM, 2017.
- [FGV17] V. Feldman, C. Guzman, and S. S. Vempala. Statistical query algorithms for mean vector estimation and stochastic convex optimization. In Philip N. Klein, editor, Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2017, pages 1265–1277. SIAM, 2017.

- [FPV15] V. Feldman, W. Perkins, and S. Vempala. On the complexity of random satisfiability problems with planted solutions. In *Proceedings of the Forty-Seventh Annual ACM on* Symposium on Theory of Computing, STOC, 2015, pages 77–86, 2015.
- [FS97] Y. Freund and R. Schapire. A decision-theoretic generalization of on-line learning and an application to boosting. *Journal of Computer and System Sciences*, 55(1):119–139, 1997.
- [GGK20] S. Goel, A. Gollakota, and A. R. Klivans. Statistical-query lower bounds via functional gradients. *CoRR*, abs/2006.15812, 2020. In NeurIPS'20.
- [GHR92] M. Goldmann, J. Håstad, and A. Razborov. Majority gates vs. general weighted threshold gates. *Computational Complexity*, 2:277–300, 1992.
- [GR06] V. Guruswami and P. Raghavendra. Hardness of learning halfspaces with noise. In Proc. 47th IEEE Symposium on Foundations of Computer Science (FOCS), pages 543–552. IEEE Computer Society, 2006.
- [Hau92] D. Haussler. Decision theoretic generalizations of the PAC model for neural net and other learning applications. *Information and Computation*, 100:78–150, 1992.
- [Kea98] M. J. Kearns. Efficient noise-tolerant learning from statistical queries. Journal of the ACM, 45(6):983–1006, 1998.
- [KK14] A. R. Klivans and P. Kothari. Embedding hard learning problems into gaussian space. In Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX/RANDOM 2014, pages 793–809, 2014.
- [KKM18] A. R. Klivans, P. K. Kothari, and R. Meka. Efficient algorithms for outlier-robust regression. In Conference On Learning Theory, COLT 2018, pages 1420–1430, 2018.
- [KLS09] A. Klivans, P. Long, and R. Servedio. Learning halfspaces with malicious noise. To appear in Proc. 17th Internat. Colloq. on Algorithms, Languages and Programming (ICALP), 2009.
- [KSS94] M. Kearns, R. Schapire, and L. Sellie. Toward Efficient Agnostic Learning. Machine Learning, 17(2/3):115–141, 1994.
- [LRV16] K. A. Lai, A. B. Rao, and S. Vempala. Agnostic estimation of mean and covariance. In *Proceedings of FOCS'16*, 2016.
- [MN06] P. Massart and E. Nedelec. Risk bounds for statistical learning. Ann. Statist., 34(5):2326–2366, 10 2006.
- [MP68] M. Minsky and S. Papert. *Perceptrons: an introduction to computational geometry*. MIT Press, Cambridge, MA, 1968.
- [MT94] W. Maass and G. Turan. How fast can a threshold gate learn? In S. Hanson, G. Drastal, and R. Rivest, editors, *Computational Learning Theory and Natural Learning Systems*, pages 381–414. MIT Press, 1994.
- [Nov62] A. Novikoff. On convergence proofs on perceptrons. In *Proceedings of the Symposium* on Mathematical Theory of Automata, volume XII, pages 615–622, 1962.

- [O'D14] R. O'Donnell. Analysis of Boolean Functions. Cambridge University Press, 2014.
- [Ros58] F. Rosenblatt. The Perceptron: a probabilistic model for information storage and organization in the brain. *Psychological Review*, 65:386–407, 1958.
- [RS94] R. Rivest and R. Sloan. A formal model of hierarchical concept learning. *Information* and Computation, 114(1):88–114, 1994.
- [Slo88] R. H. Sloan. Types of noise in data for concept learning. In Proceedings of the First Annual Workshop on Computational Learning Theory, COLT '88, pages 91–96, San Francisco, CA, USA, 1988. Morgan Kaufmann Publishers Inc.
- [Slo92] R. H. Sloan. Corrigendum to types of noise in data for concept learning. In Proceedings of the Fifth Annual ACM Conference on Computational Learning Theory, COLT 1992, page 450, 1992.
- [Slo96] R. H. Sloan. Pac Learning, Noise, and Geometry, pages 21–41. Birkhäuser Boston, Boston, MA, 1996.
- [STC00] J. Shawe-Taylor and N. Cristianini. An introduction to support vector machines. Cambridge University Press, 2000.
- [Val84] L. G. Valiant. A theory of the learnable. In Proc. 16th Annual ACM Symposium on Theory of Computing (STOC), pages 436–445. ACM Press, 1984.
- [Vap82] V. Vapnik. Estimation of Dependences Based on Empirical Data: Springer Series in Statistics. Springer-Verlag, Berlin, Heidelberg, 1982.
- [Vap98] V. Vapnik. Statistical Learning Theory. Wiley-Interscience, New York, 1998.
- [Yao90] A. Yao. On ACC and threshold circuits. In *Proceedings of the Thirty-First Annual Symposium on Foundations of Computer Science*, pages 619–627, 1990.
- [YZ17] S. Yan and C. Zhang. Revisiting perceptron: Efficient and label-optimal learning of halfspaces. In Advances in Neural Information Processing Systems 30: Annual Conference on Neural Information Processing Systems 2017, pages 1056–1066, 2017.
- [ZLC17] Y. Zhang, P. Liang, and M. Charikar. A hitting time analysis of stochastic gradient langevin dynamics. In *Proceedings of the 30th Conference on Learning Theory, COLT* 2017, pages 1980–2022, 2017.
- [ZSA20] C. Zhang, J. Shen, and P. Awasthi. Efficient active learning of sparse halfspaces with arbitrary bounded noise. *coRR*, abs/2002.04840, 2020. In NeurIPS'20.

# Appendix

### A Proof of Lemma 2.3

Let  $\theta$  be the angle between v and v'. By making an orthogonal change of variables, we can reduce to the case where v = (1, 0, ..., 0) and  $v' = (\cos(\theta), \sin(\theta), 0, 0, ..., 0)$ . Then by definition we have that  $\chi_{N(0,I)}(\mathbf{P}_v, \mathbf{P}_{v'}) + 1$  is

$$\int_{\mathbb{R}^m} \left( \frac{A(x_1)A(\cos(\theta)x_1 + \sin(\theta)x_2)g(x_2)g(\sin(\theta)x_1 - \cos(\theta)x_2)}{g(x_1)g(x_2)} \right) g(x_3)\cdots g(x_m)dx_1\cdots dx_m \ .$$

Noting that the integral over  $x_3, \ldots, x_m$  separates out, we are left with

$$\int_{\mathbb{R}^2} \left( \frac{A(x)A(\cos(\theta)x + \sin(\theta)y)g(y)g(\sin(\theta)x - \cos(\theta)y)}{g(x)g(y)} \right) dxdy \ .$$

Integrating over y gives

$$\int \frac{A(x)}{g(x)} \left( \int A(\cos(\theta)x + \sin(\theta)y)g(\sin(\theta)x - \cos(\theta)y)dy \right) dx = \int \frac{A(x)U_{\cos(\theta)}A(x)}{g(x)}dx ,$$

where  $U_t$  is the Ornstein-Uhlenbeck operator. We will simplify our computations by expressing the various quantities in terms of the eigenbasis for this operator.

In particular, let  $h_n(x) = He_n(x)/\sqrt{n!}$  where  $He_n(x)$  is the probabilist's Hermite polynomial. We note the following basic facts about them:

1.  $\int_{\mathbb{R}} h_i(x) h_j(x) g(x) dx = \delta_{i,j}.$ 

2. 
$$U_t(h_n(x)g(x)) = t^n h_n(x)g(x)$$
.

We can now write A(x) in this basis as

$$A(x) = \sum_{n=0}^{\infty} a_n h_n(x) g(x) \; .$$

From this, we obtain that

$$\chi^{2}(A, N(0, 1)) = \int_{\mathbb{R}} \left( \sum_{n=0}^{\infty} a_{n}h_{n}(x)g(x) \right)^{2} / g(x)dx$$
$$= \int_{\mathbb{R}} \sum_{n,m=0}^{\infty} a_{n}a_{m}h_{n}(x)h_{m}(x)g(x)dx$$
$$= \sum_{n=0}^{\infty} a_{n}^{2}.$$

Furthermore, we have that

$$\int_{\mathbb{R}} h_s(x) A(x) dx = \int_{\mathbb{R}} \sum_{n=0}^{\infty} a_n h_s(x) h_n(x) g(x) dx = a_s \; .$$

For  $1 \leq s \leq k$ , we have that

$$h_s(x) = \sqrt{s!} \sum_{t=0}^{\lfloor s/2 \rfloor} \frac{(-1)^t x^{s-2t}}{2^t t! (n-2t)!} \ .$$

We therefore have that

$$a_s = \sum_{t=0}^{\lfloor s/2 \rfloor} \left( \frac{\sqrt{s!}(-1)^t x^{s-2t}}{2^t t! (s-2t)!} \right) \mathbf{E}[A^{s-2t}] \ .$$

Note that the above is close to

$$\sum_{t=0}^{\lfloor s/2 \rfloor} \left( \frac{\sqrt{s!}(-1)^t x^{s-2t}}{2^t t! (s-2t)!} \right) \mathbf{E}[G^{s-2t}] = \mathbf{E}[h_s(G)] = 0 \; .$$

In particular, the difference between the two quantities is at most

$$\nu \sum_{t=0}^{\lfloor s/2 \rfloor} \left( \frac{\sqrt{s!}}{2^t t! (s-2t)!} \right) \ .$$

It is easy to see that the denominator is minimized when  $t = s/2 - O(\sqrt{s})$ . From this it follows that this sum is  $2^{O(s)}\nu$ . Therefore, we have that  $a_s = 2^{O(s)}\nu$ , for  $1 \le s \le k$ . Furthermore,  $a_0 = \int A(x)dx = 1$ . Thus, we have that

$$\begin{split} \chi_{N(0,I)}(\mathbf{P}_{v},\mathbf{P}_{v'})+1 &= \int_{\mathbb{R}} \frac{A(x)U_{v\cdot v'}A(x)}{g(x)}dx \\ &= \int_{\mathbb{R}} \left(\sum_{n=0}^{\infty} a_{n}h_{n}(x)g(x)\right) \left(\sum_{n'=0}^{\infty} a_{n}'(v\cdot v')^{n'}h_{n}'(x)g(x)\right) / g(x)dx \\ &= \int_{\mathbb{R}} \sum_{n,n'=0}^{\infty} a_{n}a_{n}'(v\cdot v')^{n'}h_{n}(x)h_{n}'(x)g(x)dx \\ &= \sum_{n=0}^{\infty} a_{n}^{2}(v\cdot v')^{n} \\ &= 1 + \sum_{n=1}^{k} a_{n}^{2}(v\cdot v')^{n} + \sum_{n=k+1}^{\infty} a_{n}^{2}(v\cdot v')^{n} . \end{split}$$

Therefore,

$$\begin{aligned} |\chi_{N(0,I)}(\mathbf{P}_{v},\mathbf{P}_{v'})| &\leq O(\nu^{2}) \sum_{n=1}^{k} 2^{O(n)} |v \cdot v'|^{n} + |v \cdot v'|^{k+1} \sum_{n=0}^{\infty} a_{n}^{2} \\ &\leq \nu^{2} + |v \cdot v'|^{k+1} \chi^{2}(A, N(0,1)) . \end{aligned}$$

This completes our proof.

# B Proof of Lemma 2.9

We consider the Fourier transform of  $G_{\delta}$ . Note that  $G_{\delta}$  is the pointwise product of G with a mesh of delta-functions. Therefore, its Fourier transform is the convolution of their Fourier transforms.

The Fourier transform of G is  $\sqrt{2\pi}G$ . The Fourier transform of the net of delta-functions is a new net with spacing  $1/\delta$ . Thus, we have that the Fourier transform of  $G_{\delta}$  at  $\xi$  is

$$\sum_{n\in\mathbb{Z}}\sqrt{2\pi}g(\xi+n/\delta)\;.$$

The  $t^{th}$  moment of a pseudodistribution is proportional to the value of the  $t^{th}$  derivative of its Fourier transform at  $\xi = 0$ . For G, this is  $\sqrt{2\pi}g^{(t)}(0)$ . For  $G_{\delta}$ , it is equal to this term plus

$$\sum_{n \in \mathbb{Z}, n \neq 0} \sqrt{2\pi} g^{(t)}(n/\delta).$$

Computing the derivative of g using Cauchy's integral formula (integrating around a circle of radius  $1/(2\delta)$  centered at  $n/\delta$ ), we find that

$$|g^{(t)}(n/\delta)| = t! O(\delta)^t \exp(-\Omega(n/\delta)^2).$$

Taking a sum over n yields our result.