On the Soundness of Bealer's Logics

Clarence Protin

August 2021

Abstract

Bealer's intensional logics T1 and T2 were proposed and expounded most fully in his book Quality and Concept (1982) [1] as well in [2]. These logics are unique in being extensions of classical first-order associated to a non-nominalist or non-inscriptionalist ontology and theory of meaning. Structurally they are similar to the second-order systems proposed about the same time by Zalta [30]. In the book and article referenced above Bealer presents a detailed sketch of a proof of soundness and completeness for T1 and T2 (something which seems to be lacking for Zalta's systems). However there are key steps to the soundness proofs which are stated without proof and which seem to be non-trivial. In this paper we both simplify the original presentation of systems T1 and T2 and supply the rather complex and involved proofs of Bealer's missing lemmas.

This paper only addresses technical details of Bealer's logic. For a recent survey of the intensional logic from a philosophical perspective which includes a discussion of the merits of Bealer's approach see [27]. The aims of the paper are (i) to give an alternative simplified presentation of Bealer's syntactic decomposition and (ii) to provide full details of the non-trivial lemmas required to the prove the soundness of T1 and T2.

The paper is organised as follows. In section 1 we present a slightly different and hopefully simpler version of the common framework for T1 and T2. In sections 2 and 3 we present systems T1 and T2 together with same basic results. In section 4 we introduce the class of models we will be working with. In section 5 we present the missing proofs of all the lemmas required to prove the soundness of T1 and in section 6 we do the same for T2.

1 Bealer Decomposition

Definition 1.1 The language L^{ω} consists of a countable ordered collection of variables x, y, z, ... and n-ary predicate symbols $(n \ge 1)$ F, G, H, ... with a distinguished binary predicate =, logical connectives $\&, \sim, \exists$ and the intensional abstraction operator $[\quad]_{x_1...x_n}$ where $x_1...x_n$ is a possibly empty sequence of distinct variables.

We use the notation \bar{x} for a (possibly empty) sequence of distinct variables. We use lower-case letters (possibly with subscripts) towards the end of the Latin alphabet to denote variables.

Definition 1.2 Formulas and terms of L^{ω} are defined by simultaneous induction:

- Variables are terms.
- If $t_1, ..., t_n$ are terms and F is n-ary predicate, then $F(t_1, ..., t_n)$ is a formula.
- If A and B are formulas, v a variable, then (A&B), $\neg A$ and $\exists v.A$ are formulas.

• If A is a formula and $v_1, ..., v_m, 0 \le m$, is a sequence of distinct variables, then $[A]_{v_1...v_m}$ is a term.

A term of the form $[A]_{v_1,...,v_n}$ is called an *intensional abstract*. Intensional abstracts of the form $[F(v_1,...,v_n)]_{v_1...v_n}$ are called *elementary*. Intensional abstracts $[A]_{\alpha}$ where A is not an atomic formula are called *complex*.

The intensional abstraction operator generalises the set-theoretic extension operator $\{x : A\}$. The standard notions of bound and free variable, bound-variable renaming (α -equivalence) as well as a term being free for x in A carry over to L^{ω} in the expected way (note that intensional abstraction binds variables like quantifiers). Terms that differ only by bound variable renaming will be called *variants*.

We will define a set of (partially defined) syntactic operations on intensional abstracts and an algorithm to decompose an intensional abstract in a unique way in terms of such syntactic operations, elementary intensional abstracts and variables.

A term $[A]_{\bar{x}}$ is normalized if all the variables in \bar{x} are free in A and they display the order in which these variables first occur free.

If A is atomic then we call $[A]_{\bar{x}}$ a prime term if the variables in \bar{x} are free in A, independently of the order.

Given an atomic formula $F(t_1, ..., t_n)$ if a variable occurs free in more than one of the terms t_i then it is called a *reflected variable*. A prime term $[A]_{\bar{x}}$ is called a *prime reflection term* if \bar{x} contains a reflected variable in A.

Consider a prime term $[F(t_1, ..., [B]_{\bar{y}}, ..., t_j)]_{\bar{x}}$ which is not a prime reflection term. If there is a variable in \bar{x} which is free in $[B]_{\bar{y}}$ and all the previous arguments are variables in \bar{x} , then the prime term is a *prime relativized predication term* and such a variable is called a *relativized variable*.

Consider a prime term $[F(t_1, ..., t_k, ..., t_j)]_{\bar{x}}$ which is not a prime reflection term. If there is term t_k which is a variable not in \bar{x} or which contains no free variables which are in \bar{x} and all the previous arguments are variables in \bar{x} , then it is a *prime absolute predication term*.

These definitions allow us to divide all intensional abstracts into seven disjoint categories. The following is clear:

Lemma 1.3 Given an intensional abstract $[A]_{\bar{x}}$ it falls into one and one only of the following seven categories which depend only on A and \bar{x} independently of order (only on the underlying set). They are also invariant under the renaming of bound variables.

- 1. Complex and \bar{x} contains variables not free in A.
- 2. Complex and all the variables in \bar{x} are free in A.
- 3. Not complex and \bar{x} contains variables not free in A.
- 4. Prime reflection term.
- 5. Prime relativized predication term.
- 6. Prime absolute predication term.
- 7. $[A]_{\bar{x}'}$ is elementary for some permutation \bar{x}' of \bar{x} .

Before defining Bealer decomposition we need the following considerations on permutations. It is clear that any permutation $\sigma^n \in S^n (n \geq 2)$ can be decomposed into permutations of the form $\sigma_c^n = (1 \ 2 \ ... \ n)$ (for $n \geq 3$) and $\sigma_i^n = (n-1 \ n)$. However, we need a uniquely defined decomposition defined as follows. Assume that our permutations are acting on sequences

 $x_1...x_n$ of variables and let $x'_1...x'_n$ be a permutation of $x_1...x_n$. We use the notation $\sigma^n x_1...x_n$ to denote the results of apply σ^n to the sequence. We associate to $x_1...x_n$ and $x'_1...x'_n$ (or equivalently the permutation which transforms the first sequence into the second) a uniquely defined decomposition as follows:

Example 1.4 Consider $x_1x_2x_3x_4$ and $x_2x_4x_3x_1$. The the process yields σ_i^4 and $x_1x_2x_4x_3$, $\sigma_c^4\sigma_i^4$ and $x_3x_1x_2x_4$, $\sigma_c^4\sigma_c^4\sigma_i^4$ and $x_4x_3x_1x_2$ and finally $\sigma_c^4\sigma_c^4\sigma_c^4\sigma_i^4$ and $x_2x_4x_3x_1$.

Definition 1.5 The Bealer operations consists in eight syntactic operations defined on intensional abstracts. These consists in six unary operations C, I, N, U, E, R, a binary operation K and partially-defined binary operations P_n for $n \geq 0$. These operations can be seen as defined on equivalence classes modulo renaming of bound variables. Let $[A]_{\bar{x}}$ be an intensional abstract with \bar{x} of length n. Then we have:

$$C[A]_{\bar{x}} = [A]_{\sigma_c^n \bar{x}} \quad n \ge 3$$

$$I[A]_{\bar{x}} = [A]_{\sigma_i^n \bar{x}} \quad n \ge 2$$

$$N[A]_{\bar{x}} = [\neg A]_{\bar{x}}$$

$$U[A]_{\bar{x}} = [\exists x_n A]_{x_1 \dots x_{n-1}} \quad n \ge 1$$

$$E[A]_{\bar{x}} = [A]_{\bar{x}y} \quad y \text{ the first variable not in } A \text{ or } \bar{x}$$

$$R[A]_{\bar{x}} = [A[x_{n-1}/x_n]]_{x_1 \dots x_{n-1}}$$

$$K[A]_{\bar{x}}[B]_{\bar{y}} = [A\&B']_{\bar{x}} \quad [B']_{\bar{x}} \text{ variant of } [B]_{\bar{y}}$$

$$P_0[A]_{\bar{x}}[B]_{\bar{y}} = [A[[B]_{\bar{y}}/x_n]]_{x_1...x_{n-1}}$$
 where $n \ge 1$ and $[B]_{\bar{y}}$ is free for x_n in $[A]_{x_1...x_{n-1}}$

$$P_n[A]_{\bar{x}}[B]_{\bar{y}} = [A[[B]_{\bar{z}}/x_n]]_{x_1...x_{n-1}\bar{w}} \quad \text{where } \bar{y} = \bar{z}\bar{w}, \bar{w} \text{ length } n \geq 1, [B]_{\bar{z}} \text{ is free for } x_n \text{ in } [A]_{x_1...x_{n-1}\bar{w}} = [A[B]_{\bar{z}}/x_n]_{x_1...x_n}$$

Consider a sequence \bar{x} and a permutation $\bar{x'}$. We associate to $\beta(\bar{x}, \bar{x'})$ a sequence $\Sigma(\bar{x}, \bar{x'})$ of operations I and C obtained by replacing σ_i^n with I and σ_c^n with C in $\beta(\bar{x}, \bar{x'})$. Thus in the previous example we have $\Sigma(x_1x_2x_3x_4, x_2x_4x_3x_1) = CCCI$. This sequence is to be interpreted compositionally. In general we shall use the notation Σ for any (possibly empty) sequence of operations C and I. We use the notation E^n to denote E composed n times. Also we shall use the notation $P_0^n tt_1...t_n$ to mean $P_0...P_0(P_0 tt_1)t_2....t_n$.

We now define Bealer decomposition.

Definition 1.6 Given an intensional abstract $[A]_{\bar{x}}$ or variable x the Bealer decomposition $\mathcal{B}[A]_{\bar{x}}$ is defined inductively as follows:

- Suppose $[A]_{\bar{x}}$ belongs to categories 1 or 3. And let \bar{x}' be a permutation of \bar{x} of the form $\bar{y}\bar{z}$ so that $[A]_{\bar{y}}$ is normalized and \bar{z} (of length m) consists of the variables in \bar{x} that do not occur free in A in order of their occurrence in \bar{x} . Then $\mathcal{B}[A]_{\bar{x}} := \Sigma(\bar{x}, \bar{x}')E^m\mathcal{B}[A]_{\bar{y}}$.
- Suppose $[A]_{\bar{x}}$ belongs to category 2 and is not normalized. And let \bar{x}' be a permutation of \bar{x} so that $[A]_{\bar{x}'}$ is normalized. Then $\mathcal{B}[A]_{\bar{x}} := \Sigma(\bar{x}, \bar{x}')\mathcal{B}[A]_{\bar{x}'}$.
- Suppose the term belongs to category 2 and is normalized. Then we have

$$\mathcal{B}[A\&B]_{\bar{x}} := K\mathcal{B}[A]_{\bar{x}}\mathcal{B}[B]_{\bar{x}}$$

$$\mathcal{B}[\neg A]_{\bar{x}} := N\mathcal{B}[A]_{\bar{x}}$$

 $\mathcal{B}[\exists v A|_{\bar{x}} := U\mathcal{B}[A]_{\bar{x}v}$ where v is the first variable not in A or \bar{x} .

• Let the term belong to category 4. Of the reflected variables in \bar{x} consider the one v with the right-most occurrence and let be t the right-most argument of A in which v has a free occurrence. Let w be the alphabetically earliest variable not occurring in A or \bar{x} . Let \bar{x}' be the permutation of \bar{x} such that $[F(t_1, ..., t_n)]_{\bar{x}'}$ is normalized and consider $\bar{y}v$ where \bar{y} is \bar{x}' with v removed. Then

$$\mathcal{B}[F(t_1,...,t_n)]_{\bar{x}} := \Sigma(\bar{x},\bar{y}v)R\mathcal{B}[F(...t[w/v]...)]_{\bar{y}vw}$$

• Let the term belong to category 6 and let t be the argument which is not a variable in \bar{x} or which contains no free variables which are in \bar{x} and such that all the previous arguments are variables in \bar{x} . Let v be the alphabetically earliest variable not occurring in A or \bar{x} .. Let \bar{x}' be the permutation of \bar{x} such that $[F(t_1, ..., t_n)]_{\bar{x}'}$ is normalized. Then

$$\mathcal{B}[F(...,t,...)]_{\bar{x}} := \Sigma(\bar{x},\bar{x'})P_0\mathcal{B}[F(...,v,...)]_{\bar{x'}v}\mathcal{B}t$$

• Let the term belong to category 5. Then it is a prime term $[F(t_1, ..., [B]_{\bar{y}}, ..., t_j)]_{\bar{x}}$, where there is a variable in \bar{x} which is free in $[B]_{\bar{y}}$ and such that all the previous arguments are variables in \bar{x} . Let \bar{z} be the sequence, as they first occur, of all m free variables in $[B]_{\bar{y}}$ which are in \bar{x} . Let v be the alphabetically earliest variable not occurring in A or \bar{x} . Then if \bar{x}' is the normalizing permutation of \bar{x} it is of the form $\bar{x}''\bar{z}\bar{x}'''$. Let $\bar{w}=\bar{x}''\bar{x}'''\bar{z}$. Then

$$\mathcal{B}[F(...,[B]_{\bar{y}},...)]_{\bar{x}} := \Sigma(\bar{x},\bar{w})P_m\mathcal{B}[F(...,v,...)]_{\bar{x}''\bar{x}'''v},\mathcal{B}[B]_{\bar{y}\bar{z}}$$

• Finally let the term be in category 7 and let \bar{x}' be such that $[F(x_1,...,x_n)]_{\bar{x}'}$ is elementary.

$$\mathcal{B}[F(x_1,...,x_n)]_{\bar{x}} := \Sigma(\bar{x},\bar{x}')[F(x_1,...,x_n)]_{\bar{x}'}$$

• In the case of a variable we have $\mathcal{B}x := x$.

It is easy to check that this decomposition is well-defined and unique modulo renaming of bound variables. Bealer decomposition transforms an intensional abstract into an expression (in Polish notation) involving the eight Bealer operations, elementary abstracts and variables.

Example 1.7 Consider the term $[\neg F(x, [G(x, y)])]_y$. Then

$$\begin{split} \mathcal{B}[\neg F(x,[G(x,y)])]_y &= N\mathcal{B}[F(x,[G(x,y)])]_y = NP_0\mathcal{B}[F(v,[G(x,y)])]_{yv}x \\ &= NP_0I\mathcal{B}[F(v,[G(x,y)])]_{vy}x = NP_0IP_1\mathcal{B}[F(v,w)]_{vw}\mathcal{B}[G(x,y)]_yx \\ &= NP_0IP_1\mathcal{B}[F(v,w)]_{vw}P_0\mathcal{B}[G(u,y)]_{yu}xx = NP_0IP_1\mathcal{B}[F(v,w)]_{vw}P_0I\mathcal{B}[G(u,y)]_{uy}xx \end{split}$$

The following observation will be used further ahead:

Lemma 1.8 Let $[F(t_1,...,t_n)]_{\bar{x}}$ be a prime term. Then its Bealer decomposition has the form

$$\Gamma_1...\Gamma_m[F(v_1,...,v_n)]_{v_1...v_n}\mathcal{B}t'_{i_1}...\mathcal{B}t'_{i_m}$$

where the Γ_i consist in sequences of the form

$$\Sigma_1 R \Sigma_2 R \Sigma_3 \dots \Sigma_{l-1} R \Sigma_l P_{n_i}$$

and $t'_{i_k} = s_{\bar{y}}$ where $s = t_{i_k}$ and if $s = [A]_{\bar{z}}$ then $s_{\bar{y}}$ is $[A]_{\bar{z}\bar{y}}$ where \bar{y} is the sequence of free variables (as they first occur) of s which are in \bar{x} . If s is a variable then \bar{y} is empty and $t'_{i_k} = s$.

2 System T1

We now consider Bealer's axiomatic system T1 over the language L^{ω} . The conception of intensionality behind T1 is that there is only one necessary truth and that the intensional abstracts of necessarily equivalent formulas are equal and interchangeable. We define necessitation in terms of equality of intensional abstracts:

$$\Box A \equiv_{df} [A] = [[A] = [A]]$$

Here [A] = [A] represents the one necessary truth.

A Hilbert system for T1 consists of the following axioms and rules:

- 1. All propositional tautologies
- 2. (Inst) $\forall v.A(v) \rightarrow A(t)$, t free for v in A.
- 3. (QImp) $\forall x.(A \to B) \to (A \to \forall x.B)$, x not free in A
- 4. (Id) x = x
- 5. (L) $x = y \to (A(x, x) \leftrightarrow A(x, y)), y$ free for x in the positions it replaces x.
- 6. $[A]_{\bar{x}} \neq [B]_{\bar{y}}$, for \bar{x} and \bar{y} of different lengths
- 7. $[A]_{\bar{x}} = [A']_{\bar{x}'}$ (equality modulo renaming bound variables)
- 8. (B) $[A]_{\bar{x}} = [B]_{\bar{x}} \leftrightarrow \Box \forall \bar{x}. (A \leftrightarrow B)$
- 9. (T) $\Box A \to A$
- 10. (K) $\Box (A \to B) \to (\Box A \to \Box B)$
- 11. (S5) $\Diamond A \to \Box \Diamond A$
- 1. (MP) From A and $A \to B$ infer B
- 2. (N) From A infer $\Box A$
- 3. (Gen) From A infer $\forall x.A$

Here, as usual, $\Diamond A =_{df} \neg \Box \neg A$.

In order to prove soundness and completeness it is convenient to replace axioms (L),(B) and (S5) with:

1. (L') $x = y \to (A(x, x) \to A(x, y)), y$ free for x in the positions it replaces x, A atomic.

2. (B1)
$$\square(A \leftrightarrow B) \leftrightarrow [A] = [B]$$

3. (B2)
$$\forall v.[A(v)]_{\bar{x}} = [B(v)]_{\bar{x}} \leftrightarrow [A(x)]_{\bar{x}v} = [B(v)]_{\bar{x}v}$$

4. (S5')
$$x \neq y \rightarrow \Box x \neq y$$

The rest of this section is devoted to proving that we obtain equivalent axiomatic systems.

Lemma 2.1 In T1 we have:

$$i) \vdash x = y \rightarrow \Box x = y$$

$$ii) \vdash \neg \Box x = y \leftrightarrow x \neq y$$

Proof. For i), by the identity axiom we have x = x. Hence using the necessitation rule we get

$$\Box x = x$$

which is

$$[x = x] = [[x = x] = [x = x]]$$

We denote this expression by B(x,x). Using B(x,x) in (L)

$$x = y \to A(x, x) \leftrightarrow A(x, y)$$

which is equivalent to $A(x,x) \to (x=y \to A(x,y))$ we obtain as desired.

For ii), take the converse of the (T) instance $\Box x = y \to x = y$ to obtain $x \neq y \to \neg \Box x = y$. For the other direction simply take the converse of i)

Lemma 2.2 We obtain an equivalent system if in T1 we replace (S5) with $(*) x \neq y \rightarrow \Box x \neq y$.

Proof. Assuming (*), since $\Diamond A$ is by definition $\neg([\neg A] = [[\neg A] = [\neg A]])$ we get (S5) immediately.

In the other direction take the (S5) instance

$$\neg \Box \neg \neg x = y \rightarrow \Box \neg \Box \neg \neg x = y$$

that is,

$$\neg \Box x = y \to \Box \neg \Box x = y$$

Now by ii) of the previous lemma we have

$$\neg \Box x = y \rightarrow \neg x = y$$

Applying necessitation and using (K) we get

$$\Box \neg \Box x = y \rightarrow \Box \neg x = y$$

and combining with (**) we get as desired.

We will use the standard properties of S5 modal logic:

Lemma 2.3 In T1 we have

$$i) \vdash A \rightarrow \Diamond A$$

$$ii) \vdash A \rightarrow \Box \Diamond A$$

$$iii) \Diamond \Box A \to A$$

- iv) If $\vdash A \rightarrow B$ then we can derive $\vdash \Diamond A \rightarrow \Diamond B$
- v) We have $\vdash \Diamond A \rightarrow B$ iff $\vdash A \rightarrow \Box B$

Proof. For i) take the converse of the (T) instance $\Box \neg A \to \neg A$. For ii) combine i) and (S5). For iii) take the converse of the (S5) instance $\neg \Box \Diamond \neg A \to \neg \Diamond \neg A$ and use (T). For iv) take the converse of the premise, apply (T) and take the converse again. Finally v) follows easily from (T), iv), iii) and ii).

Lemma 2.4 $\vdash \Box \forall v. A(v) \rightarrow \forall v. \Box A(v)$

Proof. Take the (Ins) instance

$$\forall v. A(v) \rightarrow A(v)$$

Applying necessitation, (K) and modus ponens yields:

$$\Box \forall v. A(v) \to \Box A(v)$$

Applying generalization:

$$\forall v. (\Box \forall v. A(v) \rightarrow \Box A(v))$$

Since v is not free on the left side of the implication we can apply (QImp) and modus ponens to obtain

$$\Box \forall v. A(v) \to \forall v. \Box A(v)$$

Lemma 2.5 We have the Barcan Formula $\vdash \forall v. \Box A(v) \rightarrow \Box \forall v. A(v)$.

Proof. We have by (Ins)

$$\forall v. \Box A \rightarrow \Box A$$

Applying \Diamond yields, by lemma 2.3 iii) and iv):

$$\Diamond \forall v. \Box A \rightarrow \Diamond \Box A \rightarrow A$$

Hence by generalization, (QImp) and modus ponens:

$$\Diamond \forall v. \Box A \rightarrow \forall v. A$$

which again by lemma 2.3 yields

$$\forall v. \Box A \rightarrow \Box \forall v. A$$

We can now prove:

Lemma 2.6 We obtain an equivalent system to T1 if instead of (B) take

$$\Box(A \leftrightarrow B) \leftrightarrow ([A] = [B])$$

(B'2)
$$\forall v.([A(v)]_{\bar{x}} = [B(v)]_{\bar{x}}) \leftrightarrow [A(v)]_{\bar{x}v} = [B(v)]_{\bar{x}v}$$

Proof. Using Barcan's formula and its converse it is easy to see that (B'1) and (B'2) follow from (B).

Assume (B'1) and (B'2). We must show that

$$[A]_{\bar{x}} = [B]_{\bar{x}} \leftrightarrow \Box \forall \bar{x}. (A \leftrightarrow B)$$

Let $\bar{x} = \bar{x'}v$.

Since $[A]_{\bar{x'}v} = [B]_{\bar{x'}v} \leftrightarrow \forall v.([A(v)]_{\bar{x'}} = [B(v)]_{\bar{x'}})$ using (B'2) repeatedly we get

$$[A]_{\bar{x}v} = [B]_{\bar{x}v} \leftrightarrow \forall \bar{x}, v.([A(v)] = [B(v)])$$

Using (B'1) we get

$$[A]_{\bar{x}} = [B]_{\bar{x}} \leftrightarrow \forall \bar{x}. \Box (A \leftrightarrow B)$$

Using repeatedly Barcan's formula and its converse we get as desired.

Finally:

Lemma 2.7 We obtain an equivalent system to T1 when the Leibniz axiom (L) is restricted to atomic predicates and restricted to the form (L') $x = y \rightarrow (A(x, x) \rightarrow A(x, y))$.

Proof. L obviously implies (L') as a particular case. In the other direction we first show that $x = y \to (A(x,x) \leftarrow A(x,y))$. We have that $x = y \to (A(x,x) \to A(x,y))$ and A(x,y) arises from A(x,x) by replacing some occurrences of x by y. So A(x,x) arises from replacing some occurrences of y in A(x,y) by x. Hence we have $y = x \to (A(x,y) \to A(x,x))$. Also from (L') we can derive $x = y \to y = x$ and using this we get as desired. We can now proceed inductively on the structure of the formula A. The case of negation is immediate. For conjunction we use the tautology $(A \to B) \to (C \to D) \to (A\&C \to B\&D)$ and for quantification we use generalisation, (QImp) and modus ponens.

This lemma justifies that "there is only one necessary truth".

Lemma 2.8 We have $\forall x, y . [x = x] = [y = y].$

Proof. We have $x = x \leftrightarrow y = y$. We apply necessitation and (B'1) and then generalisation to get as desired.

3 System T2

Bealer's logic T2 is given the following Hilbert axiomatic system:

- 1. All propositional tautologies
- 2. (Ins) $\forall x. A(x) \rightarrow A(t)$, t free for v in A.
- 3. (QImp) $\forall x.(A \to B) \to (A \to \forall x.B)$, x not free in A
- 4. (Id) x = x
- 5. (L) $x = y \to (A(x, x) \leftrightarrow A(x, y)), y$ free for x in the positions it replaces x.
- 6. $[A]_{\bar{x}} \neq [B]_{\bar{y}}$, \bar{x} and \bar{y} of different lengths
- 7. $[A]_{\bar{x}} = [A']_{\bar{x}'}$ (equality modulo renaming bound variables)
- 8. $[A]_{\bar{x}} = [B]_{\bar{x}} \rightarrow (A \leftrightarrow B)$
- 9. $[A]_{\bar{x}} \neq [B]_{\bar{y}}$, non-elementary terms belonging to different categories.
- 10. Let t' = Bt and r' = Br where B is the first operation in the Bealer decomposition of t' and r'. Then $t = r \leftrightarrow t' = r'$
- 11. Let t = Bt'r' and r = Bt''r'' where B is the first operation in the Bealer decomposition of t and r. The $t = r \leftrightarrow t' = t'' \& r' = r''$.
- 12. If $[F(x_1,...,x_n)]_{x_1...x_n} = s$ and G occurs in s then $[G(y_1,...,y_m)]_{y_1...y_m} \neq t$ whenever F occurs in t (Non-circularity)
- 1. (MP) From A and $A \to B$ infer B
- 2. (TGen) Suppose that F does not occur in A(v). If $\vdash A([F(x_1, ..., x_n)]_{x_1...x_n})$ then we can infer $\vdash A(t)$ whenever t is complex, of arity n and free for v in A.
- 3. (Gen) From A infer $\forall x.A$

4 Model Structures

Let $F = \emptyset$ and $T = \{\emptyset\}$. A model \mathcal{M} consists of a set \mathcal{D} with a decomposition into the union of disjoint sets \mathcal{D}_i for $i \geq -1$, a distinguished element $id \in \mathcal{D}_2$, a set \mathcal{H} of functions H whose domain is \mathcal{D} and such that H is the identify on \mathcal{D}_{-1} and on \mathcal{D}_0 has range $\{T, F\}$ and on \mathcal{D}_i has range $\mathcal{P}(\mathcal{D}^i)$ where $\mathcal{D}^i = \mathcal{D} \times ... \times \mathcal{D}$ (i terms) for $i \geq 1$. All these functions must satisfy $H(id) = \{(x, y) : x = y\}$. There is a distinguished function \mathcal{G} representing the actual extension. Furthermore we have unary (partial) operations n, e, u, c, i, r and binary (partial) operations k, p_n with $n \geq 0$. These operations restrict to the following domains and ranges:

$$n: \mathcal{D}_{i} \to \mathcal{D}_{i}, i \geq 0$$

$$e: \mathcal{D}_{i} \to \mathcal{D}_{i+1}, i \geq 0$$

$$u: \mathcal{D}_{i} \to \mathcal{D}_{i-1}, i \geq 1 \text{ and } u: \mathcal{D}_{0} \to \mathcal{D}_{0}.$$

$$c: \mathcal{D}_{i} \to \mathcal{D}_{i}, i \geq 3$$

$$r: \mathcal{D}_i \to \mathcal{D}_{i-1}, i \geq 2$$

 $i: \mathcal{D}_i \to \mathcal{D}_i, i \geq 2$

$$a: \mathcal{D}_i \times \mathcal{D}_i \to \mathcal{D}_i, i \ge 0$$

$$p_n: \mathcal{D}_i \times \mathcal{D}_j \to \mathcal{D}_{i+n-1}, i \ge 1, j \ge n$$

We define inductively $p_0^n(d, x_1, ..., x_n) = p_0(p_0^{n-1}(d, x_2, ..., x_{n-1}), x_1)$ and $p_0^1(d, x_1) = p_0(d, x_1)$. Furthermore we have the important constraints on how the functions in \mathcal{H} relate to these operations. Here d denotes a suitable member of \mathcal{D}^i . For all $\mathcal{H} \in \mathcal{H}$ we must have:

```
d' \in \mathcal{H}n(d) \text{ iff } d' \notin \mathcal{H}d \\ (x_1, ..., x_i, x_{i+1}) \in \mathcal{H}e(d) \text{ iff } (x_1, ..., x_i) \in \mathcal{H}d \\ (x_1, ..., x_{i-1}) \in \mathcal{H}u(d) \text{ iff there is a } x_i \text{ such that } (x_1, ..., x_i) \in \mathcal{H}d \\ (x_1, ..., x_{i-1}, x_i) \in \mathcal{H}c(d) \text{ iff } (x_i, x_1, ..., x_{i-1}) \in \mathcal{H}d \\ (x_1, ..., x_{i-1}, x_i) \in \mathcal{H}i(d) \text{ iff } (x_1, ..., x_i, x_{i-1}) \in \mathcal{H}d \\ (x_1, ..., x_{i-1}, x_i) \in \mathcal{H}r(d) \text{ iff } (x_1, ..., x_i, x_i) \in \mathcal{H}d \\ d'' \in \mathcal{H}k(d, d') \text{ iff } d'' \in \mathcal{H}d \text{ and } d'' \in \mathcal{H}d' \\ (x_1, ..., x_{i-1}, y_1, ..., y_n) \in \mathcal{H}p_n(d, d') \text{ iff } (x_1, ..., x_{i-1}, p_n^n(d', y_1, ..., y_n)) \in \mathcal{H}d
```

Here we consider also the 0-tuple as being \emptyset . Notice that elements of D^0 are propositions seen as objective entities according to their meaning. The extension functions \mathcal{H} determine their Boolean values which are $T = \{\emptyset\}$ and $F = \emptyset$.

Definition 4.1 Given a model \mathcal{M} , an interpretation \mathcal{I} assigns *i*-ary predicate elements of \mathcal{D}_i and = to id. An assignment \mathcal{A} assigns to variables elements in \mathcal{D} .

Bealer decomposition is crucial to define the denotation of a term relative to a model, interpretation and assignment.

Definition 4.2 Given \mathcal{M}, \mathcal{I} and \mathcal{A} we define the *denotation* $D_{\mathcal{I},\mathcal{A},\mathcal{M}}t$ of a term t of L^{ω} as follows. If t is a variable then it is $\mathcal{A}(t)$. If t is an elementary term $[F(v_1, ..., v_n)]_{v_1...v_n}$ then it is $\mathcal{I}(F)$. Otherwise, consider the Bealer decomposition $\mathcal{B}t$ of t. If $\mathcal{B}t = Bt'$ (respectively Bt't'') where B is a Bealer operation then we define inductively

$$D_{\mathcal{I},\mathcal{A},\mathcal{M}}t = bD_{\mathcal{I},\mathcal{A},\mathcal{M}}t'$$
 (respectively $D_{\mathcal{I},\mathcal{A},\mathcal{M}}t = bD_{\mathcal{I},\mathcal{A},\mathcal{M}}t'D_{\mathcal{I},\mathcal{A},\mathcal{M}}t''$)

where b is i, c, e, u, n, r, k, p_n if B is I, C, E, U, N, R, K, P_n respectively.

Definition 4.3 We say that a formula A is true for \mathcal{M}, \mathcal{I} and \mathcal{A} (we also write $T_{\mathcal{I},\mathcal{A},\mathcal{M}}(A)$) if $\mathcal{G}D_{\mathcal{I},\mathcal{A},\mathcal{M}}[A] = T$.

It is simple to see that the denotation is invariant modulo renaming bound variables. From now one, we will in most cases drop the subscripts in $D_{\mathcal{I},\mathcal{A},\mathcal{M}}$ and assume we a working with a given model, interpretation and assignment.

We are now interested in special classes of models for which T1 and T2 are sound and complete.

Definition 4.4 A model \mathcal{M} is called type 1 if for all $i \geq -1$, $x, y \in \mathcal{D}_i$

(type 1 model condition)
$$\forall \mathcal{H} \in H \quad \mathcal{H}(x) = \mathcal{H}(x) \to x = y$$

A model \mathcal{M} is called type 2 if the operations are:

- 1. one-to-one
- 2. disjoint on their ranges
- 3. and non-cycling

By the first two conditions each element of the model has a unique (possibly infinite) decomposition in terms of the operations. This decomposition is a tree having as leaves indecomposable elements. The non-cycling condition means that the same element cannot appear in more than one place in the tree.

A formula A is T1-valid (we write $\vdash_{T1} A$) iff A is true for all type 1 model \mathcal{M} , interpretation \mathcal{I} and assignment \mathcal{A} . In the same way a formula A is T2-valid ($\vdash_{T2} A$) iff A is true for all type 2 model \mathcal{M} , interpretation \mathcal{I} and assignment \mathcal{A} .

Note that for type 2 models only \mathcal{G} is relevant. The type 1 condition restricts the possible operations on the model.

Example 4.5 A standard model for Kelley-Morse set theory furnishes an example of a type 1 model with a single extensional function. Type 1 models can be seen as generalised set theory models with a plurality of membership predicates $\in_{\mathcal{H}}$ such that the axiom of extensionality is only valid globally. Consider a language with a single unary predicate M. We can construct a type 2 model with no constants, D_0 , D_1 and $D_i = \emptyset$ for $i \geq 2$.

Lemma 4.6 Let v be an externally quantifiable variable in $[B(v)]_{\bar{x}}$ and let t be free for v in $[B(v)]_{\bar{x}}$. Consider any model structure \mathcal{M} and any interpretation \mathcal{I} and assignment \mathcal{A} . Let \mathcal{A}' be an assignment which is just like \mathcal{A} except that $\mathcal{A}'(v) = Dt$. Then

$$D_{\mathcal{A}'}[B(v)]_{\bar{x}} = D[B(t)]_{\bar{x}}$$

Proof. By induction on the Bealer decomposition of $[B(v)]_{\bar{x}}$. The base case is either an elementary abstract (in which case there are no externally quantifiable variables) or else a variable. If it is not v we are done and if it is v the result is trivial.

Suppose that $[B(v)]_{\bar{x}} = Bs(v)$ were B is some unary operation. If B is I, C, N, U or E then t is still free for v in d(v) and hence by the induction hypothesis $D_{\mathcal{A}'}s(v) = Ds(t)$ so that $D_{\mathcal{A}'}[B(v)]_{\bar{x}} = bD_{\mathcal{A}'}s(v) = bDs(t) = DBs(t)$ where b is the corresponding model function. If B is B then there is a problem that B has a new bound variable which may occur in B in such a way that B is no longer free for B in B. But we may rename this bound variable to obtain an B-equivalent term B where this problem does not occur. Note that the substitution does not effect which of the seven syntactic categories an intensional abstract belongs to. Suppose B is B in the previous argument applies. If B is B then B is B then B in the problem of the new bound variable in the first argument possibly occurring in B. We thus need to use B-equivalence so B remains free for B in B in the case of 0-predication we have

$$D_{\mathcal{A}'}[B(v)]_{\bar{x}} = p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, D_{\mathcal{A}'}s'(v))$$

If s' is not a variable or a variable distinct from v, then we may apply the induction hypothesis to obtain:

$$p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, D_{\mathcal{A}'}s'(v)) = p_0(D[B'(t)]_{\bar{x}w}, Ds'(t)) = D[B(t)]_{\bar{x}}$$

If s' is v then we obtain

$$p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, D_{\mathcal{A}'}v)$$

$$= p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, \mathcal{A}'v)$$

$$= p_0(D[B'(t)]_{\bar{x}w}, Dt)$$

$$= D[B(t)]_{\bar{x}}$$

In the k-predication case for k > 0 we have, for $s'(v) = [C(v)]_{\bar{y}\bar{z}}$ where \bar{z} are relativized variables of $[C(v)]_{\bar{y}}$ in $[B(v)]_{\bar{x}}$

$$D_{\mathcal{A}'}[B(v)]_{\bar{x}} = p_k(D_{\mathcal{A}'}[B'(v)]_{\bar{x}'w}, D_{\mathcal{A}'}[C(v)]_{\bar{y}\bar{z}})$$

where \bar{x}' is obtained from \bar{x} by omitting \bar{z} . The result now follows easily from induction.

Lemma 4.7 For all \mathcal{M} and \mathcal{I} , \mathcal{A} we have $D[A]_{x_1...x_k} \in \mathcal{D}_k$

Proof. This follows by easy induction from the definition of D and the model operations.

Given a sequence of Bealer operations of the form Σ we denote its associated n-permutation by σ . This permutation can also be applied to n-tuples $(x_1, ... x_n)$ and we denote such an application by $\sigma(x_1, ..., x_n)$ just as we denote its application to variable sequences by $\sigma x_1 ... x_n$. We will make frequent use of the following:

Lemma 4.8 For any permutation σ of $(x_1,...,x_n)$ we have

$$\sigma(v_1, ..., v_n) \in \mathcal{H}D[F]_{\sigma x_1, ..., x_n} \leftrightarrow (v_1, ..., v_n) \in \mathcal{H}D[F]_{x_1, ...x_n}$$

Proof. Let the Bealer decomposition of $[F]_{x_1...x_n}$ be $\Sigma_1 \mathcal{B}[F]_{x'_1...x'_n}$ where $\mathcal{B}[F]_{x'_1...x'_n}$ does not begin with C or I. Then Bealer decomposition of $[F]_{\sigma x_1,...,x_n}$ will be of the form $\Sigma_2 \mathcal{B}[F]_{x'_1...x'_n}$. By the definition of denotation and model we have for any $(x_1,...,x_n)$:

$$(x_1, ..., x_n) \in \mathcal{H}D[F]_{x_1...x_n} \text{ iff } \sigma_1(x_1, ..., x_n) \in \mathcal{H}D[F]_{x'_1...x'_n}$$

$$(x_1, ..., x_n) \in \mathcal{H}D[F]_{\sigma x_1, ..., x_n} \text{ iff } \sigma_2(x_1, ..., x_n) \in \mathcal{H}D[F]_{x'_1...x'_n}$$

where $x'_1...x'_n = \sigma_1(x_1, ..., x_n)$ and $x'_1...x'_n = \sigma_2\sigma(x_1, ..., x_n)$ and hence $\sigma = \sigma_2^{-1}\sigma_1$ and $\sigma_1 = \sigma_2\sigma$.

Hence given $(x_1,...x_n)$ we have $(x_1,...,x_n) \in \mathcal{H}D[F]_{x_1...x_n}$ iff $\sigma_1(x_1,...,x_n) = \sigma_2\sigma(x_1,...,x_n) \in \mathcal{H}D[F]_{x'_1...x'_n}$ iff $\sigma(x_1,...,x_n) \in \mathcal{H}D[F]_{\sigma x_1,...,x_n}$.

Lemma 4.9 For all $\mathcal{I}, \mathcal{A}, \mathcal{M}, F(t_1, ..., t_n)$ is true iff $(Dt_1, ..., Dt_n) \in \mathcal{I}(F)$.

Proof. By definition $F(t_1,...,t_n)$ is true iff $\mathcal{G}D([F(t_1,...,t_n)]) = T$. But

 $[F(t_1,...,t_n)] = P_0[F(v_1,...,t_n)]_{v_1}t_1 = P_0P_p[F(v_1,v_2,...,t_n)]_{v_1v_2}t_2t_1 = P_0^n[F(v_1,...,v_n)]_{v_1...v_n}t_n...t_2t_1$ hence by definition $\mathcal{G}D([F(t_1,...,t_n)]) = T$ iff

$$Dt_1 \in \mathcal{G}p_0(p_0(...p_0(D[F(v_1,...,v_n)]_{v_1...v_n},Dt_n),...,Dt_3),Dt_2)$$

iff

$$(Dt_1, Dt_2) \in \mathcal{G}p_0(p_0(...p_0(D[F(v_1, ..., v_n)]_{v_1...v_n}, Dt_n)..., Dt_3)$$

and so on until obtaining

$$(Dt_1, ..., Dt_n) \in D[F(v_1, ..., v_n)]_{v_1...v_n} = \mathcal{I}(F)$$

Lemma 4.10 $T_{\mathcal{I}A\mathcal{M}}(A\&B)$ iff $T_{\mathcal{I}A\mathcal{M}}(A)$ and $T_{\mathcal{I}A\mathcal{M}}(B)$. Also $T_{\mathcal{I}A\mathcal{M}}(\neg A)$ iff it is not the case that $T_{\mathcal{I}A\mathcal{M}}(A)$.

This is an immediate consequence of the definition of $T_{\mathcal{IAM}}$.

5 Soundness of T1

In this section we work in a type 1 model \mathcal{M} . We will use Polish notation for the model operations and omit parentheses when possible. We note first that Lemma 4.9 (for the case of equality) and Lemma 4.10 yield immediately:

Lemma 5.1 For all models we have propositional tautologies, (MP), (Id) and (L') are sound.

It is immediate by lemma 4.7 that axiom 6 is sound and the soundness of axiom 7 is obvious. The following is of central importance

Lemma 5.2 (Bealer's lemma) Let v be free in $[A(v)]_{\bar{x}}$. Let \mathcal{M} be type 1 and \mathcal{I} , \mathcal{A} be an interpretation and assignment. Then

$$D[A(v)]_{\bar{x}} = p_0 D[A(v)]_{\bar{x}v} \mathcal{A}(v)$$

Note that when we write $(x_1, ..., x_n)$ we include the case n = 0 in which case the sequence is taken to be the empty set \emptyset .

Proof. We proceed by induction on the Bealer decomposition of $[A(v)]_{\bar{x}}$. Let $\mathcal{B}[A(v)]_{\bar{x}} = K\mathcal{B}[B(v)]_{\bar{x}}\mathcal{B}[C(v)]_{\bar{x}}$. Then

$$D[A(v)]_{\bar{x}} = kD[B(v)]_{\bar{x}}D[C(v)]_{\bar{x}} = kp_0D[B(v)]_{\bar{x}v}\mathcal{A}(v)p_0D[C(v)]_{\bar{x}v}\mathcal{A}(v)$$

by the induction hypothesis. We use the type 1 condition to show that

$$kp_0D[B(v)]_{\bar{x}v}\mathcal{A}(v)p_0D[C(v)]_{\bar{x}v}\mathcal{A}(v) = p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

For any \mathcal{H} let $(x_1,...,x_n) \in \mathcal{H}kp_0D[B(v)]_{\bar{x}v}\mathcal{A}(v)p_0D[C(v)]_{\bar{x}v}\mathcal{A}(v)$. Then this is equivalent to

$$(x_1,...,x_n) \in \mathcal{H}p_0D[B(v)]_{\bar{x}v}\mathcal{A}(v)$$
 and $(x_1,...,x_n) \in \mathcal{H}p_0D[C(v)]_{\bar{x}v}\mathcal{A}(v)$

which is equivalent to

$$(x_1, ..., x_n, \mathcal{A}(v)) \in \mathcal{H}D[B(v)]_{\bar{x}v}$$
 and $(x_1, ..., x_n, \mathcal{A}(v)) \in \mathcal{H}D[C(v)]_{\bar{x}v}$

The Bealer decomposition of $[A(v)]_{\bar{x}v}$ is $\Sigma K[B(v)]_{\bar{x}'}[C(v)]_{\bar{x}'}$ were Σ is such that $\bar{x'} = \sigma \bar{x}v$ is the normalised permutation of $\bar{x}v$ for $[A(v)]_{\bar{x}v}$. But then the above condition is equivalent to

$$\sigma(x_1, ..., x_n, \mathcal{A}(v)) \in \mathcal{H}D[B(v)]_{\bar{x}'}$$
 and $\sigma(x_1, ..., x_n, \mathcal{A}(v)) \in \mathcal{H}D[C(v)]_{\bar{x}'}$

which is equivalent to $\sigma(x_1,...,x_n,\mathcal{A}(v)) \in \mathcal{H}kDB[(v)]_{\bar{x}'}DC[(v)]_{\bar{x}'}$. Let s be the composition of operations corresponding to Σ . Then the previous condition is equivalent to $(x_1,...,x_n,\mathcal{A}(v)) \in \mathcal{H}skDB[(v)]_{\bar{x}'}DC[(v)]_{\bar{x}'} = \mathcal{H}D[A(v)]_{\bar{x}v}$. But this is equivalent to $(x_1,...,x_n) \in \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$. Hence the conclusion follows from the type 1 condition.

Let $\mathcal{B}[A(v)]_{\bar{x}} = N\mathcal{B}[A'(v)]_{\bar{x}}$ where $A(v) = \neg A'(v)$. We have $D[A(v)]_{\bar{x}} = nD[A'(v)]_{\bar{x}} = np_0D[A'(v)]_{\bar{x}v}\mathcal{A}(v)$ by the induction hypothesis. We use the type 1 condition to show that

$$np_0D[A'(v)]_{\bar{x}v}\mathcal{A}(v) = p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

Then for any \mathcal{H} ,

$$(x_1,...,x_n) \in \mathcal{H}np_0D[A'(v)]_{\bar{x}v}\mathcal{A}(v) \text{ iff } (x_1,...,x_n) \notin \mathcal{H}p_0D[A'(v)]_{\bar{x}v}\mathcal{A}(v)$$

iff
$$(x_1, ..., x_n, \mathcal{A}(v)) \notin \mathcal{H}[A'(v)]_{\bar{x}v}$$

Let \bar{x}' be the normalised sequence for $[A'(v)]_{\bar{x}v}$ and σ the associated permutation. Then the above condition holds iff

$$\sigma(x_1,...,x_n,\mathcal{A}(v)) \notin \mathcal{H}[A'(v)]_{\bar{x}'} \text{ iff } \sigma(x_1,...,x_n,\mathcal{A}(v)) \in \mathcal{H}n[A'(v)]_{\bar{x}'}$$

The Bealer decomposition of $[A(v)]_{\bar{x}v}$ is $\Sigma N[A'(v)]_{\bar{x}'}$ where σ corresponds to Σ so $D[A(v)]_{\bar{x}v} = snD[A'(v)]_{\bar{x}'}$ where s is the composition of model operation corresponding to Σ . Hence we have $\sigma(x_1,...,x_n,\mathcal{A}(v)) \in \mathcal{H}n[A'(v)]_{\bar{x}'}$ iff $(x_1,...,x_n,\mathcal{A}(v)) \in \mathcal{H}D[A(v)]_{\bar{x}v}$ iff $(x_1,...,x_n) \in \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$ and our conclusion follows from the type 1 model condition.

Let $\mathcal{B}[A(v)]_{\bar{x}} = U[A'(v)]_{\bar{x}w}$ where $A(v) = \exists w.A'(v)$. Then $D[A(v)]_{\bar{x}} = uD[A'(v)]_{\bar{x}w} = up_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v)$ by induction. We show that $up_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v) = p_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$. Take a \mathcal{H} . Then

$$(x_1,...,x_n) \in \mathcal{H}up_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v)$$
 iff there is y such that $(x_1,...,x_n,y) \in \mathcal{H}p_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v)$

iff there is a y such that
$$(x_1,...,x_n,y,\mathcal{A}(v)) \in \mathcal{H}D[A'(v)]_{\bar{x}wv}$$

As previously let $\bar{x}' = \sigma \bar{x}v$ be the normalized sequence for $[A(v)]_{\bar{x}v}$. The Bealer decomposition of $[A(v)]_{\bar{x}v}$ is $\Sigma U\mathcal{B}[A'(v)]_{\bar{x}'w}$ so $p_0[A(v)]_{\bar{x}v}\mathcal{A}(v) = p_0suD[A'(v)]_{\bar{x}'w}\mathcal{A}(v)$. The above condition is equivalent to

there is a y such that $(x'_1, ..., x'_{n+1}, y) \in \mathcal{H}D[A'(v)]_{\bar{x}'w}$

where $\mathcal{A}(v)$ occupies the position x'_i of v in \bar{x}' . But this is equivalent to

$$(x'_1,...,x'_{n+1}) \in \mathcal{H}uD[A'(v)]_{\bar{x}'w}$$

which is equivalent to

$$(x_1, ..., x_n, \mathcal{A}(v)) \in \mathcal{H}suD[A'(v)]_{\bar{x}'w} \text{ iff } (x_1, ..., x_n) \in \mathcal{H}p_0suD[A'(v)]_{\bar{x}'w}\mathcal{A}(v) = \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

and the result follows from the type 1 model condition.

Let $\mathcal{B}[A(v)]_{\bar{x}} = E\mathcal{B}[A(v)]_{\bar{x}'}$ with $\bar{x} = \bar{x}'y$ and y does not occur free in A(v). We have $D[A(v)]_{\bar{x}} = eD[A(v)]_{\bar{x}'} = ep_0[A(v)]_{\bar{x}'v}\mathcal{A}(v)$ by induction. As previously take a \mathcal{H} . Then

$$(x_1,...,x_n) \in \mathcal{H}ep_0[A(v)]_{\bar{x}'v}\mathcal{A}(v) \text{ iff } (x_1,...,x_{n-1}) \in \mathcal{H}p_0[A(v)]_{\bar{x}'v}\mathcal{A}(v)$$

iff
$$(x_1, ..., x_{n-1}, \mathcal{A}(v)) \in \mathcal{H}[A(v)]_{\bar{x}'v}$$

The Bealer decomposition of $[A(v)]_{\bar{z}v}$ is $\Sigma \mathcal{B}[A(v)]_{\bar{z}} = \Sigma E \mathcal{B}[A(v)]_{\bar{z}'}$ where $\bar{z} = \sigma \bar{x}v$ is the normalizing permutation which puts v in its proper place and \bar{z}' drops the last element of \bar{z} (not v and the same as x_n). Put $\bar{z}' = \sigma' \bar{x}' v$ and let σ' correspond to composition of model operations s'. Then the above condition is equivalent to

$$(x_1, ..., x_{n-1}, \mathcal{A}(v)) \in \mathcal{H}s'D[A(v)]_{\sigma'\bar{x}'v} = \mathcal{H}s'D[A(v)]_{\bar{z}'}$$

which is equivalent to

$$(x_1, ..., x_{n-1}, x_n, \mathcal{A}(v)) \in \mathcal{H}ies'D[A(v)]_{\bar{z}'} = \mathcal{H}seD[A(v)]_{\bar{z}'}$$

which is turn is equivalent to:

$$(x_1, \ldots, x_n) \in \mathcal{H}p_0 seD[A(v)]_{\bar{z}'} \mathcal{A}(v) = \mathcal{H}p_0 D[A(v)]_{\bar{x}v} \mathcal{A}(v)$$

and the result follows as in the other cases.

Let $\mathcal{B}[A(v)]_{\bar{x}} = B\mathcal{B}[A(v)]_{\bar{x'}}$ with B equal to C or I where $\bar{x'} = \sigma_b \bar{x}$ for b equal to c or i. Then $D[A(v)]_{\bar{x}} = bD[A(v)]_{\bar{x'}} = bp_0[A(v)]_{\bar{x'}} \mathcal{A}(v)$ using the induction hypothesis. We use the type 1 condition to show that

$$bp_0[A(v)]_{\bar{x'}v}\mathcal{A}(v) = Dp_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

For any \mathcal{H} we have

$$(x_1, ..., x_n) \in \mathcal{H}bp_0[A(v)]_{\bar{x'}v}\mathcal{A}(v) \text{ iff } (x'_1, ..., x'_n) \in \mathcal{H}p_0[A(v)]_{\bar{x'}v}\mathcal{A}(v)$$

where $(x_1, ..., x_n) = \sigma_b(x_1, ..., x_n)$. But

$$(x'_1,...,x'_n) \in \mathcal{H}p_0[A(v)]_{\bar{x'}v}\mathcal{A}(v) \text{ iff } (x'_1,...,x'_n,\mathcal{A}(v)) \in \mathcal{H}[A(v)]_{\bar{x'}v}$$

iff
$$(x_1, ..., x_n, \mathcal{A}(v)) \in \mathcal{H}[A(v)]_{\bar{x}v}$$
 iff $(x_1, ..., x_n) \in \mathcal{H}p_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$

Hence by the type 1 model condition we get that $bp_0[A(v)]_{\bar{x}'v}\mathcal{A}(v) = Dp_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$.

Now let $\mathcal{B}[A(v)]_{\bar{x}} = R\mathcal{B}[A'(v)]_{\bar{x}w}$. Then $[A(v)]_{\bar{x}}$ has a reflected variable and hence so does $[A(v)]_{\bar{x}v}$ (it is also belongs to category 4 and is a prime reflection term). We distinguish between two cases. Either v is the right-most reflected variable or it is not. Assume that is is not. Let t be the right-most argument in which t occurs. Then the right-most reflected variable is the last element x_n of \bar{x} . We have $D[A(v)]_{\bar{x}} = rp_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v)$ by induction, were A'(v) is obtained from A(v) by replacing the argument t with $t[w/x_n]$. Fix \mathcal{H} . Then

$$(x_1,...x_n) \in \mathcal{H}rp_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v) \text{ iff } (x_1,...,x_n,x_n,\mathcal{A}(v)) \in \mathcal{H}D['A(v)]_{\bar{x}wv}$$

Now the Bealer decomposition of $[A(v)]_{\bar{x}v}$ in this case will be $\Sigma R\mathcal{B}[A'(v)]_{\bar{x}'w}$. Here $\bar{x}v = \sigma \bar{x}'$. \bar{x}' moves v to its proper place so that \bar{x}' is the normalized permutation of $\bar{x}v$ except that x_n is placed at the end, being the reflected variable with right-most occurrence. We have $D[A(v)]_{\bar{x}v} = srD[A'(v)]_{\bar{x}'w}$. Then the condition above is equivalent to:

$$(x'_1,...,x'_{n+1},x'_{n+1}) \in \mathcal{H}D[A'(v)]_{\bar{x}'w}$$

with $x'_{n+1} = x_n$. But this is equivalent to:

$$(x'_1,...,x'_{n+1}) \in \mathcal{H}rD[A'(v)]_{\bar{x}'w}$$

which is turn is equivalent to

$$(x_1, ..., x_n, \mathcal{A}(v)) \in \mathcal{H}srD[A'(v)]_{\bar{x}'w}$$

which is finally equivalent to

$$(x_1,, x_n) \in \mathcal{H}p_0srD[A'(v)]_{\bar{x}'w}\mathcal{A}(v) = \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

and the result follows as in the other cases. Now for the second case: v is a reflected variable with the right-most occurrence. Let the right-most argument in which it occurs be t. The

Bealer decomposition of $[A(v)]_{\bar{x}v}$ in this case will be $\Sigma R\mathcal{B}[A'(v)]_{\bar{x}'vw}$ where A'(v) is obtained from A(v) by replacing t with t[w/v]. Here $\bar{x}v = \sigma \bar{x}'v$. \bar{x}' moves x_n back to its proper place so that \bar{x}' is a normalized sequence. Now either in $[A'(v)]_{\bar{x}'vw}$ v is still the right-most reflected variable or else it is x_n . If v it is still the right-most reflected variable we continue with the Bealer decomposition until we arrive at term in which the right-most reflected variable is x_n . The decomposition will be of the form:

$$[A(v)]_{\bar{x}v} = \Sigma_1 R \Sigma_2 R \dots \Sigma_k R \mathcal{B}[A^{(k)}(v)]_{\bar{z}w_k}$$

where \bar{z} (normalized except for w_k being placed at the end) contains the elements of \bar{x}, v and new variables $w_1, ..., w_{k-1}$. Here $[A^{(k)}(v)]_{\bar{z}w_k}$ is now as the previous case (modulo a permutation). It is easy to see that

$$(x_1, ...x_n) \in \mathcal{H}p_0 s_1 r s_2 r s_k RD[A^{(k)}(v)]_{\bar{z}w_k} \mathcal{A}(v)$$

iff

$$(x'_1, ..., x'_{n+k+1}) \in \mathcal{H}D[A^{(k)}(v)]_{\bar{z}w_k}$$

Here $(x'_1, ..., x'_{n+k+1})$ follows the sequence of $\bar{z}w_k$, having x_i for x_i (here we are abusing notation) but has $x'_k = \mathcal{A}(v)$ whenever x'_k is a w_i or v.

Now $D[A^{(k)}(v)]_{\bar{z}w_k} = sD[A^{(k)}(v)]_{\bar{u}x_n} = srD[A_0^{(k)}(v)]_{\bar{u}x_nw}$ where \bar{u} is the normalized permutation of $\bar{z}w_k$ but with x_n omitted. Hence

(*)
$$(x'_1, ..., x'_{n+k+1}) \in \mathcal{H}D[A^{(k)}(v)]_{\bar{z}w_k} \text{ iff } (x''_1, ..., x''_{n+k}, x_n, x_n) \in \mathcal{H}D[A_0^{(k)}(v)]_{\bar{u}x_nw}$$

where the x_i'' follows the normalized sequence but with x_n omitted and $A_0^{(k)}(v)$ has the right-most argument t in which x_n occurs replaced with $t[w/x_n]$. Now we have by induction,

$$(x_1, ..., x_n) \in \mathcal{H}D[A(v)]_{\bar{x}} \text{ iff } (x_1, ..., x_n) \in \mathcal{H}rp_0[A_1(v)]_{\bar{x}wv}\mathcal{A}(v)$$

iff
$$(x_1, ..., x_n, x_n, \mathcal{A}(v)) \in \mathcal{H}D[A_1(v)]_{\bar{x}wv}$$

where A_1 has the right-most argument t in which x_n appears replaced by $t[w/x_n]$. Now $[A_1(v)]_{\bar{x}wv}$ has v reflected in the same arguments and positions as $[A(v)]_{\bar{x}v}$. Hence

$$[A_1(v)]_{\bar{x}wv} = \Sigma_1' R \Sigma_2' R \Sigma_k' R \mathcal{B}[A_1^{(k)}(v)]_{\bar{q}w_k}$$

Hence

$$(x_1, ..., x_n, x_n, \mathcal{A}(v)) \in \mathcal{H}D[A_1(v)]_{\bar{x}wv} \text{ iff } (q_1, ..., q_{n+k+2}) \in \mathcal{H}D[A_1^{(k)}(v)]_{\bar{q}w_k}$$

Here \bar{q} is the normalized permutation of $\bar{x}wv$ except that w_k is moved to the end. It has new variables $w_1, ..., w_{k_1}$. $(q_1, ..., q_{n+k+2})$ follows this sequence and for v or w_i it has A and for w it has x_n . It is clear that $A_1^{(k)}(v)$ is the same as $A_0^{(k)}(v)$. Examining (*) and noticing that applying a permutation we have

$$(x_1'', ..., x_{n+k}'', x_n, x_n) \in \mathcal{H}D[A_0^{(k)}(v)]_{\bar{u}x_nw} \text{ iff } (q_1, ..., q_{n+k+2}) \in \mathcal{H}D[A_0^{(k)}(v)]_{\bar{q}w_k}$$

and the result follows.

Finally, consider the case in which $\mathcal{B}[A(v)]_{\bar{x}} = P_n \mathcal{B}[A'(v)]_{\bar{u}w} t_{\bar{y}}$. Here $t_{\bar{y}}$ means that if $t = [B]_{\bar{z}}$ then $t_{\bar{y}} = [B]_{\bar{z}\bar{y}}$ or else t is a variable and \bar{y} is the empty sequence. Let the argument in A(v) which w replaced be t. Here $\bar{x} = \bar{x}'\bar{x}''\bar{y}$ is normalized except that the sequence \bar{y} of variables

in \bar{x} which are free in t are placed at the end. All arguments before t are variables in \bar{x} . So v may occur in t and in arguments after t and v may be reflected. Hence we will have (if v is reflected)

$$[A(v)]_{\bar{x'}\bar{x''}\bar{y}v} = \Sigma_1 R \Sigma R ... R \Sigma_k \Sigma [A^{(k)}(v)]_{\bar{x'}\bar{p}\bar{y'}} = \overline{\Sigma} R \Sigma [A^{(k)}(v)]_{\bar{x'}\bar{p}\bar{y'}}$$

Here $\bar{x}'\bar{q}$ is the normalized permutation of $\bar{x}v$ and the k new variables w_k but with the sequence \bar{y}' of variables of \bar{q} which occur free in t placed at the end. \bar{y}' is either equal to \bar{y} or contains in addition v. Notice how \bar{x}' is not affected by the permutations. For clarity, we wrote the last permutation as a composition. If v is not reflected then we have only Σ which puts v in its proper place (which is either in \bar{y} or in \bar{p}).

We have that

$$(x'_1,...,x'_a,x''_1,...,x''_b,y_1,...,y_c,v) \in \mathcal{H}\overline{sr}sD[A^{(k)}(v)]_{\bar{x}'\bar{p}\bar{p}'}$$

iff

$$(x'_1,...,x'_a,p_1,...,p_{b+k},y'_1,...,y'_{c+j}) \in \mathcal{H}D[A^{(k)}(v)]_{\bar{x'}\bar{p}\bar{y'}}$$

where j is either 0 and v is in \bar{p} or else is 1 and v is in \bar{y}' . Here the $(p_1, ..., p_{b+k})$ is permutation of the x_i'' and w_i (and possibly v) corresponding to the normalized sequence of these variables and which has v for w_i .

W.l.o.g let v be in \bar{p} and t. Fix a \mathcal{H} . Then

$$(x'_1, ..., x'_a, x''_1, ..., x''_b, y_1, ..., y_c) \in \mathcal{H}D[A(v)]_{\bar{x}'\bar{x}''\bar{y}} = \mathcal{H}P_nD[A'_1(v)]_{\bar{x}'\bar{x}''w}Dt_{\bar{y}} \text{ iff}$$

$$(x'_1, ..., x'_a, x''_1, ..., x''_b, p^c(Dt_{\bar{y}}, y_1, ...y_1)) \in \mathcal{H}D[A'_1(v)]_{\bar{x}\bar{x}'w} = \mathcal{H}p_0D[A'_1(v)]_{\bar{x}'\bar{x}''wv}\mathcal{A}(v)$$

by induction. This is equivalent to

$$(x'_1,...,x'_a,x''_1,...,x''_b,p^c(Dt_{\bar{y}},y_1,...y_1),\mathcal{A}(v))) \in \mathcal{H}D[A'_1(v)]_{\bar{x}'\bar{x}''wv}$$

Let the Bealer decomposition of $[A'_1(v)]_{\bar{x}'\bar{x}''wv}$ be

$$\Sigma_1'R\Sigma_2'R...R\Sigma_{k-1}'\Sigma'[A'^{(k-1)}(v)]_{\bar{x}'\bar{q}} = \overline{\Sigma'R}\Sigma'[A'^{(k)}(v)]_{\bar{x}'\bar{q}}$$

where \bar{q} consists of variables of \bar{x}'' , w, v and k new variables w_k . The previous condition is equivalent to

$$(x'_1,...,x'_a,x''_1,...,x''_b,p^c(Dt_{\bar{y}},y_1,...y_1),\mathcal{A}(v))) \in \mathcal{H}\overline{s'R}s'D[A'^{(k-1)}(v)]_{\bar{x}'\bar{q}}$$

which is equivalent to

$$(x_1', ..., x_a', r_1, ..., r_s) \in \mathcal{H}D[A'^{(k-1)}(v)]_{\bar{x}'\bar{q}}$$

where $(r_1, ..., r_s)$ corresponds to the sequence \bar{q} with v and w_k corresponding to $\mathcal{A}(v)$ and w corresponding to $p^c(Dt_{\bar{y}}, y_1, ...y_1)$.

Now consider

$$(x'_1,...,x'_a,x''_1,...,x''_b,y_1,...,y_c) \in \mathcal{H}p_0D[A(v)]_{\bar{x}\bar{x}'\bar{y}v}\mathcal{A}(v)$$

this is equivalent to

$$(x'_1, ..., x'_a, x''_1, ..., x''_b, y_1, ..., y_c, \mathcal{A}(v)) \in \mathcal{H}D[A(v)]_{\bar{x}\bar{x}'\bar{y}v} = \mathcal{H}\bar{sr}sD[A^{(k)}(v)]_{\bar{x}'\bar{p}\bar{y}'}$$

where $\bar{y'} = \bar{y''}v\bar{y'''}$. This is turn is equivalent to

$$(x'_1,...,x'_n,x''_1,...,x''_h,y_1,...,y_c,\mathcal{A}(v)) \in \mathcal{H}\overline{sr}sP_{n+1}D[A^{(k)'}(v)]_{\bar{x'}\bar{p}w}t_{\bar{v'}}$$

that is,

$$(x'_1,...,x'_a,e_1,...,e_{b+k},y'_1,...,y'_{c+1}) \in \mathcal{H}P_{n+1}D[A^{(k)'}(v)]_{\bar{x'}\bar{p}w}t_{\bar{y'}}$$

where $(e_1, ..., e_{b+k+1})$ corresponds to the sequence \bar{p} and has $\mathcal{A}(v)$ corresponding to w_k and $(y'_1, ..., y'_{c+1})$ corresponds to $\bar{y'} = \bar{y''}v\bar{y'''}$ with $\mathcal{A}(v)$ corresponding to v. This in turn is the same

$$(x'_1,...,x'_a,e_1,...,e_{b+k},p_0^{c+1}(Dt_{\bar{v'}},y'_1,...,y'_{c+1})) \in \mathcal{H}D[A^{(k)'}(v)]_{\bar{x'}\bar{p}w}$$

It is easy to check that by induction we have

$$p_0^{c+1}(Dt_{\bar{y'}}, y'_1, ..., y'_{c+1})) = p_0 p_0^c(Dt_{\bar{y}v}, y_1, ..., y_c)) \mathcal{A}(v) = p_0^c(Dt_{\bar{y}}, y_1, ..., y_c)$$

Inspecting (**), noticing that $[A^{(k)'}(v)]_{\bar{x'}\bar{p}w} = [A'^{(k)}(v)]_{\bar{x'}\bar{p}w}$ (modulo renaming bound variables) and applying a permutation we get as desired.

Lemma 5.3 $T_{\mathcal{I}A\mathcal{M}}(\exists v.A)$ iff there is an assignment \mathcal{A}' like \mathcal{A} except perhaps for what it assigns to v and such that $T_{\mathcal{I}A'\mathcal{M}}(A(v))$.

Proof. Assume w.l.o.g that v occurs free in A. We have $T_{\mathcal{I}A\mathcal{M}}(\exists v.A)$ iff $\mathcal{G}D([\exists v.A]) = T$ iff $\mathcal{G}uD[A]_v = T$. The last condition is equivalent to the existence of a $x \in \mathcal{D}$ such that

$$x \in \mathcal{G}D[A(v)]_v$$

Let \mathcal{A}' be like \mathcal{A} but with $\mathcal{A}'(v) = x$. We must show that $\mathcal{G}D_{\mathcal{A}'}[A(v)] = T$. Notice that $D_{\mathcal{A}'}[A(v)]_v = D[A(v)]_v$. By the proof of Bealer's lemma (note that we are not using the type 1 condition) we have that:

$$\mathcal{G}D_{\mathcal{A}'}[A(v)] = T \text{ iff } \mathcal{G}p_0D[A(v)]_v\mathcal{A}'(v) = T$$

Hence we must show that $\mathcal{G}p_0D[A(v)]_v\mathcal{A}'(v) = T$. But this is equivalent to $\mathcal{A}'(v) \in \mathcal{G}D[A(v)]_v$ and the result follows since $\mathcal{A}'(v) = x$. The other direction is similar.

Hence if $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\forall v.A(v))$ then for any assignment \mathcal{A}' which only differs at most on v we have that $T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(A(v))$. Hence lemma 5.3 yields:

Lemma 5.4 For any models we have $(Inst) \models \forall v.A(v) \rightarrow A(t)$ where t is free for v in A.

Likewise we have

Lemma 5.5 For any models we have $(QImp) \models \forall v.(A \rightarrow B(v)) \rightarrow (A \rightarrow \forall v.B(v))$ where v does not occur free in A.

Proof. Assume $T_{\mathcal{IAM}}(\forall v.A(\rightarrow B(v)))$. Then for all assignment \mathcal{A}' differing from \mathcal{A} at most on v we have that

$$(*) T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(A \to B(v))$$

We must show that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A \to \forall v.B(v))$. Assume then that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A)$. We must show that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A \to \forall v.B(v))$. Take any assignment \mathcal{A}' differing from \mathcal{A} at most on v. Since v does not occur in A we have that $T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(A)$. Hence by (*) we get that $T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(\forall v.A(v))$. Since this holds for any \mathcal{A}' we get the conclusion by lemma 5.3.

Similarly we get the soundness of the (Gen) rule:

Lemma 5.6 For any models if $\vDash A(v)$ then $\vDash \forall v. A(v)$.

Lemma 5.7 For all \mathcal{M} of type 1 and \mathcal{I} , \mathcal{A} and terms t, t' we have

$$D([t = t]) = D([t' = t'])$$

Proof. Suppose that

$$D([t=t]) \neq D([t'=t'])$$

Then the type 1 condition implies that there is a \mathcal{H} such that

$$\mathcal{H}D([t=t]) \neq \mathcal{H}D([t'=t'])$$

But

$$[t = t] = P_0(P_0([v = w]_{vw}, t), t)$$

Hence applying the definition of \mathcal{H} yields

$$\mathcal{H}D([t=t]) = T \leftrightarrow D(t) = D(t)$$

hence we have $\mathcal{H}D([t=t])=T$ and also analogously $\mathcal{H}D([t'=t'])=T$ and so we obtain a contradiction.

Lemma 5.8 ([1]A8*(a)) For type 1 models we have $(B1) \models \Box(A \leftrightarrow B) \leftrightarrow [A] = [B]$.

Proof. We must show that $\mathcal{G}D_{[\mathcal{IAM}}(\Box(A \leftrightarrow B) \leftrightarrow [A] = [B]]) = T$, which is equivalent of showing that

$$\mathcal{G}D_{\mathcal{IAM}}([\Box(A \leftrightarrow B)]) = T \text{ iff } \mathcal{G}D_{\mathcal{IAM}}([[A] = [B]]) = T$$

that is,

$$\mathcal{G}D_{\mathcal{IAM}}([[A \leftrightarrow B] = [[A = B] = [A = B]]]) = T \text{ iff } \mathcal{G}D_{\mathcal{IAM}}([[A] = [B]]) = T$$

which is equivalent to

$$D_{\mathcal{IAM}}([A \leftrightarrow B]) = D_{\mathcal{IAM}}([[A = B] = [A = B]]) \text{ iff } D_{\mathcal{IAM}}([[A]) = D_{\mathcal{IAM}}([B])$$

Lemma 5.9 ([1] A8*(b)) For type 1 models we have $(B2) \models \forall v. [A(v)]_{\alpha} = [B(v)]_{\alpha} \leftrightarrow [A(v)]_{\alpha v} = [B(v)]_{\alpha v}$.

Proof. We have

$$D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha} = p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v}, \mathcal{A}(v))$$

and

$$D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha} = p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}, \mathcal{A}(v))$$

Consider all assignments \mathcal{A}' like \mathcal{A} except for the assignment on v. Then $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_{\alpha v}$ and $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_{\alpha v}$. So if $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$ the for any \mathcal{A}' we have $D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_{\alpha} = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_{\alpha}$ and one direction follows.

We now need to show that if for all such A' we have

$$D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_{\alpha} = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_{\alpha}$$

then $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$. But if $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} \neq D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$ then by the type 1 condition there would be a \mathcal{H} and a $(x_1, ..., x_m)$ such that $(x_1, ..., x_m) \in \mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v}$ but $(x_1, ..., x_m) \notin \mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$. But taking $\mathcal{A}'(v) = x_n$ this means that

$$(x_1, ..., x_{m-1}) \in \mathcal{H}p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v}, \mathcal{A}'(v)) = \mathcal{H}D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_{\alpha v}$$

but

$$(x_1, ..., x_{m-1}) \notin \mathcal{H}p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}, \mathcal{A}'(v)) = \mathcal{H}D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_{\alpha v}$$

and we obtain a contradiction.

Lemma 5.10 ([1] A11*) In the same conditions we have $(S5') \vDash v_i \neq v_j \rightarrow \Box v_i \neq v_j$.

Proof. We must show that if $T_{\mathcal{IAM}}(v_i \neq v_j)$ then $T_{\mathcal{IAM}}(\square v_i \neq v_j)$.

But if $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(v_i \neq v_j)$ then $D_{\mathcal{I}\mathcal{A}\mathcal{M}}(v_i) \neq D_{\mathcal{I}\mathcal{A}\mathcal{M}}(v_k)$. But this is equivalent to $\mathcal{A}(v_i) \neq \mathcal{A}(v_k)$. This means that for all $\mathcal{H} \in \mathcal{H}$ we have $\mathcal{H}([v_i \neq v_j]) = T$ and the conclusion follows.

Lemma 5.11 ([1] A9) In the same conditions we have $(T) \vDash \Box A \rightarrow A$.

Proof. If $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box A)$ then $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$ and so in particular $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A)$

Lemma 5.12 ([1] A10) In the same conditions we have $(K) \vDash \Box(A \to B) \to (\Box A \to \Box B)$.

Proof. Assume $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box(A \to B))$. Then $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A \to B] = T$ which means that if $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$ then $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([B])$. To show that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box A \to \Box B)$ consider any $\mathcal{K} \in H$. Assume that $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$. Then $\mathcal{K}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$ and hence $\mathcal{K}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([B]) = T$. Hence we have $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([B]) = T$ and thus $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box A \to \Box B)$.

6 Soundness of T2

We have already proven the soundness of the standard first-order logical axioms and rules for any model. Also the soundess of axioms 6 and 7. Axioms 9 to 11 are clearly sound by the type 2 model condition.

Lemma 6.1 TGen is sound: suppose that F does not occur in A(v). If $\vdash A([F(x_1, ..., x_n)]_{x_1...x_n})$ then we can infer $\vdash A(t)$ whenever t complex, of arity n and is free for v in A.

Proof. By (L) we have

$$\vdash [F(x_1, ..., x_n)]_{x_1...x_n} = t \to A([F(x_1, ..., x_n)]_{x_1...x_n}) \to A(t)$$

from which it follows that

$$\vdash A([F(x_1,...,x_n)]_{x_1...x_n}) \to [F(x_1,...,x_n)]_{x_1...x_n} = t \to A(t)$$

Assume $\vDash A([F(x_1,...,x_n)]_{x_1...x_n})$. Then by soundness of (L), (MP) and tautologies we get

$$\vDash [F(x_1,...,x_n)]_{x_1...x_n} = t \to A(t)$$

By the non-cycling condition for T2 models F does not occur in t and hence the interpretation of F does not affect the denotation of Dt. But choose an interpretation \mathcal{I} assigning to $[F(x_1,...,x_n)]_{x_1...x_n}$ the value Dt. In this case we have $\models [F(x_1,...,x_n)]_{x_1...x_n} = t$ and hence $T_{\mathcal{I}}(A(t))$. But since D[A(t)] cannot depend on the interpretation is we must have $\models A(t)$.

Lemma 6.2 We have that $[A]_{\bar{x}} = [B]_{\bar{x}} \to A \leftrightarrow B$ is sound.

Proof. For any T2 model, interpretation and assignment we must show that $T([A]_{\bar{x}} = [B]_{\bar{x}} \to A \leftrightarrow B)$. We must show that if $T([A]_{\bar{x}} = [B]_{\bar{x}})$ then $T(A \leftrightarrow B)$. Now if $T([A]_{\bar{x}} = [B]_{\bar{x}})$ then we have $D[A]_{\bar{x}} = D[B]_{\bar{x}}$. We now show that $T(A \leftrightarrow B)$. We show only one implication, the other is similar. We need only show that $T(\forall \bar{x}.A \to B)$, that is, $\mathcal{G}D[\forall \bar{x}.A \to B] = T$. Now

$$D[\forall \bar{x}.A \to B] = D[\neg \exists \bar{x}. \neg (A \to B)] = D[\neg \exists \bar{x}. A \& \neg B]$$

$$= ne....eD[A\& \neg B]_{\bar{x}} = ue...ekD[A]_{\bar{x}}nD[B]_{\bar{x}} = ue...ekD[A]_{\bar{x}}nD[A]_{\bar{x}}$$

So $\emptyset \in \mathcal{H}ue...ekD[A]_{\bar{x}}nD[A]_{\bar{x}}$ iff there is no $(x_1,...,x_n)$ such that $(x_1,...,x_n) \in \mathcal{H}D[A]_{\bar{x}}$ and $(x_1,...,x_n) \notin \mathcal{H}D[A]_{\bar{x}}$. Hence $\mathcal{G}D[\forall \bar{x}.A \to B] = T$.

Note that this rule is sound for any model.

7 Conclusion

References

- [1] Bealer, G. (1982). Quality and Concept, Oxford University Press.
- [2] Bealer, G. (1983). Completeness in the theory of properties, relations, and propositions, Journal of Symbolic Logic 48 (2): 415-426.
- [3] Bealer, G. (1994). Property theory: the type-free approach v. the Church approach, Journal of Philosophical Logic 23: 139-171.
- [4] Bealer, G. (1993). A solution to Frege's puzzle. In J. E. Tomberlin (Ed.), Philosophical perspectives 7: Language and logic (pp. 17–60). Atascadero, CA: Ridgeview Publishing Company.
- [5] Bealer, G. (1998). Propositions. Mind, 107(425), 1–32.
- [6] Bealer, G. and Mönnich, U (1989). Property Theories, In Dov Gabbay & Franz Guenthner (eds.), Handbook of Philosophical Logic, Volume IV. Kluwer Academic Publishers. pp. 133-251.
- [7] Bealer, G. (1989). Fine-Grained Type-Free Intensionality, n Gennero Chierchia, Barbara H. Partee & Raymond Turner (eds.), Properties, Types, and Meaning, Volume 1. Kluwer Academic Publishers. pp. 177-230.
- [8] Bressan, A. (1972). A General Interpreted Modal Calculus, New Haven: Yale University Press.
- [9] Carnap, R. (1947). Meaning and Necessity, Chicago: University of Chicago Press. Enlarged edition 1956.
- [10] Church, A. (1951). A formulation of the logic of sense and denotation, in P. Henle (Ed.), Structure, Method and Meaning, New York: The Liberal Arts Press. pp 3–24.
- [11] Church, A. (1973). Outline of a revised formulation of the logic of sense and denotation (part I), Noûs, 7: 24–33.

- [12] Church, A. (1974). Outline of a revised formulation of the logic of sense and denotation (part II), Noûs, 8: 135–156.
- [13] Cresswell, M. J. (1973). Logic and languages. London: Meuthen.
- [14] Frege, G. (1892). Über Sinn und Bedutung. Zeitschrift für Philosophie und philosophische Kritik, 100: 25–50. English translation as 'On Sinn and Bedeutung,' in (Beaney 1997).
- [15] Gallin, D. (1975). Intensional and Higher-Order Modal Logic, Amsterdam: North-Holland.
- [16] Kripke, S. (1963). "Semantical considerations on modal logics," in Acta Philosophica Fennica, 16: 83–94.
- [17] Kripke, S. (1980). Naming and Necessity (Second edition), Cambridge, MA: Harvard University Press.
- [18] Kripke, S. (2008). Frege's Theory of Sense and Reference: Some Exegetical Notes, Theoria, 74: 181–218.
- [19] Marcus, R. (1946). "A Functional Calculus of First Order Based on Strict Implication," The Journal of Symbolic Logic, 11: 1–16.
- [20] Marcus, R. (1961) "Modalities and intensional Languages," Synthese, XIII: 303–322. Reprinted in Modalities, Philosophical Essays, Ruth Barcan Marcus, Oxford University Press, 1993.
- [21] Montague, R. (1960). "On the nature of certain philosophical entities," The Monist, 53: 159–194. Reprinted in (Thomason 1974), 148–187.
- [22] Montague, R. (1970). "Pragmatics and intensional logic," Synthèse, 22: 68–94. Reprinted in (Thomason 1974), 119–147.
- [23] Moschovakis, Y. N. (1989). The formal language of recursion. 54, 1216–1252.
- [24] Moschovakis, Y. N. (1994). Sense and denotation as algorithm and value. In J. Oikkonen and J. Vaananen (Eds.), Lecture Notes in Logic, Volume 2, pp. 210–249. Springer. [Full version of proof of Theorem 4.1 from that paper is available online.]
- [25] Moschovakis, Y. (2006). "A logical calculus of meaning and synonymy," Linguistics and Philosophy, 29: 27–89.
- [26] Quine, W. V. (1963). "Reference and modality," in From a Logical Point of View (second ed.), Chapter VIII, pp. 139–159. New York: Harper Torchbooks.
- [27] Parsons, D. (2016). Theories of Intensionality, Springer, Singapore.
- [28] Tichý, P. (1971). "An Approach to Intensional Analysis," Noûs, 5: 273–297.
- [29] Tichý, P. (1988). The foundations of Frege's logic, Berlin and New York: De Gruyter.
- [30] Zalta, E. (1988). Intensional Logic and the Metaphysics of Intentionality, Cambridge, MA: MIT Press.