

Bealer's Intensional Logic

Clarence Lewis Protin^{1*}

¹Independent Researcher

June 15, 2022

Abstract

Many intuitively valid arguments involving intensionality cannot be captured by first-order logic, even when extended by modal and epistemic operators. Indeed, previous attempts at providing an adequate treatment of the phenomenon of intensionality in logic and language, such as those of Frege, Church, Russell, Carnap, Quine, Montague and others are fraught with numerous philosophical and technical difficulties and shortcomings. We present Bealer's solution to this problem which hinges on an ontological commitment to theory of Properties, Propositions and Relations (PRP). At the most basic level we can distinguish two conceptions in the theory of PRPs. An objective one tied to modality and necessary equivalence, and a mental (intentional) one tied to concepts and the requirement of non-circularity in definitions. Building on the work of Russell, Church and Quine, Bealer proposes two distinct intensional logics T1 and T2 (presented in Hilbert form) corresponding to these two conceptions, both based on the language of first-order logic extended with an intensional abstraction operator. In T1 necessitation can be directly defined and the axioms entail that we obtain standard S5 modal logic. These logics have a series of striking features and desirable aspects which set them apart from higher-order approaches. Bealer constructs a non-Tarskian algebraic semantic framework, distinct from possible worlds semantics and its problematic ontological commitments, yielding two classes of models for which T1 and T2 are both sound and complete. Other features include being able to deal with quantifying-in, and the various substitution puzzles, being free from artificial type restrictions, having a Russellian semantics, satisfying Davidson's learnability requirement, etc. Bealer proposes his logic as the basis of a larger philosophical project in the tradition of logicism (or logical realism) concerning which we refer to his book *Quality and Concept* (1982). This includes a neo-Fregean logicist foundation of arithmetic and set-theory in which various (according to him) purely logical predication axioms (and intensional analogues of ZF, NGB, or Kelley-Morse axioms) are adjoined to T2, thereby explaining incompleteness as a property of pure logic rather than of mathematics. Surprisingly, and rather ironically, Bealer's logic also fulfills Carnap's thesis of extensionality due precisely to its ontological commitment to the reality of PRPs. The proof of these results consists either in lemmas which are merely stated or which are given but brief sketches of a proof. We aim to give detailed proofs of all the mathematical logical results that appear in Bealer's *Quality and Concept* and in [2] and to clarify and simplify some of the concepts and techniques so as to bring Bealer's work to a larger audience of philosophers, logicians, linguists and mathematicians and to be better equipped to address some of the unsolved problems and challenges. We also include a brief introduction to other approaches to intensionality in natural language and discuss how Bealer's approach compares favourably to some of them and is likely to benefit from the insights offered by others.

*The author declares that there is no conflict of interest.

Keywords: *Intensional Logic; Modal Logic; Non-classical Logics; Semantics of Natural Language; Logicism; First-order Logic.*

1 Approaches to Intensionality

Natural language abounds with syntactic constructions which *prima facie* have no analogue in the restricted domain of mathematical discourse. Our major example will be propositional attitudes (for example, an agent knowing or believing something), modalities and that-clauses. These constructions are all examples of *intensionality*: they seem to involve the meaning of the embedded sentence rather than its denotation. This is the cause of many problems. If two terms are merely co-extensive then we cannot substitute one for another according to the standard Leibniz rule. Suppose John believes that Scott is not the author of *Waverley*. Suppose "the author of *Waverley*" and Scott designate the same individual. Then if we substitute one for the other in the sentence above we obtain \therefore John believes that Scott is not Scott. Following [27] we call this type of problem (A), a particular case of a wide class of *substitutivity problems* such as Mates' puzzle and the paradox of analysis(see [1][ch.1 and 3] for an introduction to these problems). Another problem involves terms which have no denotation when we attempt to apply the standard \exists -introduction rule. From John desires a unicorn we get $\therefore \exists x$. John desires x . How are we to interpret this existential quantifier ? What precisely does it quantify ? We refer to this type of problem as (B).

Historically[19, 20] modalities were some of the first of these intentional constructs to be studied formally, but only in the context of adjoining modal operators to already known systems of formal logic such as propositional or classical first-order logic. Epistemic and temporal constructs are dealt with in an *ad hoc* fashion by introducing the corresponding operators. With this approach it is clear that if we wish to have a general logic capable of dealing with all these constructs then such a logic would have to be rather eclectic and uneconomical in form, abounding in primitive symbols. On the other hand a new kind of semantics was proposed, the *possible worlds* semantics associated with Carnap, Kripke and Hintikka. This is the semantics that served as the basis for Montague's approach, which was later axiomatised by Gallin[15]. The key features are that meaning is a *function* from states to extensions involving individuals and a type hierarchy and thus a limitation on self-applicability of predicates. For instance, the expression *number of the planets* denoted different numbers at different historical epochs. The many-worlds approach has served as the basis for the majority of general formal approaches to intensionality in language until today. For example, the work of Cresswell[13], Bressan[8] and Tichý[30, 31].

Another approach is the *sententialist* (or *inscriptionalist*) one which is associated to Quine, Davidson and Sheffler. The sentences which occur within intentional constructs are taken as *concrete inscriptions*. A recent formulation of Sheffler's approach by David Parsons[27] is first-order and employs standard Tarskian semantics. In this system belief is taken as a binary relation between an agent and an inscription. Sentences are the truth-bearers.

The third class of approaches are those which are neither based on many-worlds semantics or sententialism. Within this class are the approach of Moschovakis[23, 24, 25] based on the "sense as algorithm" idea, the Meinongian approaches of Zalta[32] and Priest[32][p.129-145] and the approach of George Bealer, based on the theory of Properties, Relations and Propositions (PRP). Bealer's logic is a classical first-order logic with equality augmented with a term-forming abstraction bracket operator $[A]_{x_1 \dots x_n}$ in the style of Quine's corner-bracket. This operator is syntactically very similar to the extensional operator $\{x : A\}$ of set theory. Bealer starts out with two axiomatic systems T1 and T2. The axiomatic systems are quite simple and natural. In T1 the box operator can be defined to obtain an embedding of S5 logic. The semantics are directly inspired by the ontology of PRPs, propositions and n -ary relations appear as irreducible

entities alongside individuals. Formulas within the bracket operator are interpreted by means of such entities. In the class of models adequate for T1 each entity is assigned a set of possible extensions ranging over the entire model. In T2 only one extension is considered.

We now consider very briefly some reasons for the preferability of Bealer's approach relative to the many-worlds and sententialist/inscriptionalist approaches.

Let us consider the sententialist approach of Scheffler which has been formalised by D. Parsons in [27][185-208] as a first-order system ISM_D^{L*} . In order to deal with that-clauses [27][p.204] an infinite number of unanalysable one-place predicates "That-P" need to be introduced, corresponding to the infinite number of sentence-inscriptions of the object language L . It seems plausible that Bealer would have argued that this violates Davidson's learnability requirement. Although system ISM_D^{L*} can deal with problems (A) and (B) it fails in elegance and economy as it requires numerous axioms and additional recursive axiom schemes (rules) introduced *ad hoc* to deal with the combinatorics of sentence-inscriptions and that-clauses. We note also that the question of completeness is not addressed unlike for Bealer's logics. Leaving aside the violation of Davidson's learnability requirement, all things being equal, a simpler system which solves the same problems is clearly preferable to the Scheffler-Parsons approach. A serious problem with the sententialist approach lies in the nature of sentence-inscriptions. What exactly is a "sentence-inscription"? The burden of the sententialist is to give a precise account of such sentence-inscriptions and to justify the counter-intuitive stance that propositional attitudes relate to sentence-inscriptions conceived as physical objects or sense perceptions (or classes of such). As Bealer observes in [6][p.174], how can the prelinguistic cognitive states of a child be described as relating to sentence-inscriptions? The PRP theory on the other hand complies with our intuition: in knowing something we relate to propositions or states of affairs, not sentence-inscriptions. There are many arguments against a sententialist and nominalist account of propositional attitudes. See for instance [28] for an analysis of nominalist accounts of mathematical knowledge and [6][174-185] for a very detailed discussion of Scheffler's brand of sententialism. Also, Bealer has shown in [7][p.216-223] how Quine's and Davidson's scepticism regarding the need for entities such as PRPs can be fully met.

We now consider the many-worlds approach in which intensional constructs are interpreted as functions from worlds to extensions. It is a desideratum that an intensional logic should give a non-circular account for what is it for a proposition to be necessary, rather than leave the box as an undefined symbol. The Kripkean explanation in terms of truth in all possible worlds only sweeps the problem under the rug as it leaves open the question of nature of the accessibility relation R between worlds. Bealer's logic T1 makes the box a defined notion in terms of more primitive concepts. Bealer makes a convincing case for a non-circular definition of necessity in [1][p.204] based on the theory of PRPs. A serious problem in the many-worlds approach is the fine-grain distinction between rigid concepts, between predicates which have the same extension in all possible worlds yet clearly differ in sense. This is not a problem in Bealer's PRP approach as sense is taken as an ontological primitive. Another similar problem concerns the intensional interpretation of fictional names, predicates which arguably have empty extension in all possible worlds. What exactly are these objects which allow us in particular to make an application of rule (B)? To believe in unicorns is not the same as believing in fairies but both concepts correspond to the same function yielding the empty set in all worlds, a function which is difficult to attach any interpretation to. It is clear that the PRP approach is much more promising for it takes properties as primitives and unicorns and fairies correspond to different concepts. There are many more problems and objections to the many-worlds approach, such as regards to its higher-order presentation which imposes an artificial type-restriction on certain "transcendental" predicates. Higher-order systems also have the disadvantage of being incomplete.

The main problem which confronts Bealer's system (in particular in the form T2) is dealing

with the wide class of substitutivity problems. While several of them are completely solved, others demand a further fine-grained distinction between PRPs. Bealer made further progress on these problems since [1] and in [7] he deploys the distinction between platonic and non-platonic modes of presentation in order to deal with the nature of proper names and the associated substitutivity puzzles. Although in [1] a pragmatic approach was proposed to solve Mates' puzzle, in [3] he develops the idea of "logical form" to allow a much finer distinction between intensional entities than is possible in T2. Although these theories are not entirely worked out they seem promising. We believe that a comparison or combination with the "sense as algorithm" approach of Moschovakis or the emphasis on sentences as *constructions* in Tichý may lead to the adequate concept of fine-grained distinction between intensional entities needed to solve the substitutivity puzzles.

2 The Nature of Bealer's Logic

This paper aims to reformulate, clarify and simplify Bealer's intensional logics, in particular by furnishing detailed proofs, so as to bring his work to a larger audience of logicians, linguists and philosophers. In this section we give our own motivation for Bealer's logic and in section 3 we give a brief outline of the Bealer's own logico-linguistic motivation for his logic, the details of which can be found in [1][Intr. and Ch.1]. Most of the lemmas in [1, 2] are either merely stated or given a sketchy outline of a proof. The rest of the sections are dedicated to proving in detail the soundness and completeness of Bealer's systems T1 and T2 as well as reformulating and simplifying certain concepts and details. We then conclude with a discussion of the work that remains to be done.

In certain versions of set theory (such as Kelley-Morse Set theory as presented in the classic book *General Topology*) one starts from standard first-order logic with equality endowed with an *extension operator*, a term-forming operator which in general takes a formula ϕ and forms the class $\{x : \phi(x)\}$ where ϕ can have other free variables besides x . Here the variable x functions like a quantified variable and the concepts of bound and free variable are extended in the expected way, including the usual proviso for performing substitution of terms for variables. The other fundamental feature is the presence of the distinguished binary membership predicate ϵ . The two fundamental axioms are the classification axiom scheme and the axiom of extensionality which express how ϵ determines equality.

The basic idea of George Bealer's first-order intensional logic is to consider an intensional version of a similar term-forming operator $[x : \phi]$ (where we use square braces instead of curly ones). We call these "classes" intensional abstracts. Semantically, we can think of this logic as being a set theory with a plurality of membership predicates ϵ_w (with some analogy to "possible worlds" semantics) and in which equality is intensional equality, a stronger property than mere membership equivalence (in this case, membership equivalence for all membership predicates). Such a membership equality for all membership predicates is called *necessary extensional equality* and is weaker than strict intensional equality. We also consider a special distinguished membership predicate representing the "actual world".

Consider a version of Bealer's logic without the intensional analogue of the membership predicate and consider two primitive notions of equality, ordinary equality = corresponding to strict intensional equality and \approx expressing necessary extensional equality. In this version we do not have the analogue of the Kuratowski construction of ordered-pairs and so we must take as primitives abstracts over (possibly empty) sequences of distinct variables $[x_1 \dots x_n : \phi]$. We can remedy this by introducing an ordered-pair operation as a primitive with the expected axiom $(x, y) = (x', y')$ iff $x = x' \ \& \ y = y'$. From this we can form arbitrary n-tuples (x_1, \dots, x_n) in the expected way. Thus we can define $[(x_1, \dots, x_n) : \phi(x_1, \dots, x_n)]$ as

$\{x : \exists x_1, \dots, x_n. x = (x_1, \dots, x_n) \ \& \ \phi(x_1, \dots, x_n)\}$. We also consider the 0-tuple which we denote by \emptyset , an additional primitive constant. Then we have that $[\emptyset : \phi]$ is $[x : x = \emptyset \ \& \ \phi]$. In Bealer's logic this is interpreted as *the proposition that ϕ* . In extensional set theory, when ϕ is closed, these classes can either be the ordinary empty set \emptyset or equal to the set $\{\emptyset\}$ depending on whether ϕ holds or not. Thus, in this case these kinds of abstract correspond to truth values. In general they correspond to taking the proposition ϕ as a term. Consider the classical set-theoretic operations (functions) that can be performed on classes such as intersection, complement, projection (of a binary relation). These are the term analogues of the logical operations conjunction, negation and existential quantification. We can furthermore consider an expansion which involves transforming an n -ary relation R into a $n + 1$ -ary one R' in a trivial way $(x_1, \dots, x_n, x_{n+1}) \in R'$ iff $(x_1, \dots, x_n) \in R$. There is also the inversion or, in general, a permutation of an n -ary relation as well as the identification of pairs of variables (generalised diagonals). In our situation these must be considered in principle as merely meta-syntactic operations. The central idea of first-order logic intensional logic, which allows us to define an adequate semantics, is the decomposing the structure of intensional abstracts in terms of such meta-syntactic operations, the most notable aspect being the use of a generalised form of substitution in which intensionally bound variables of the term to be inserted are added to the bound variables of the term they are substituted into.

3 Brief Motivation

3.1 Quantifying-in

Consider the following intuitively valid inference:

$$\frac{\begin{array}{l} \text{Whatever } x \text{ believes is necessary.} \\ \text{Whatever is necessary is true.} \end{array}}{\text{Whatever } x \text{ believes is true.}}$$

In standard quantifier logic this would be formalized as follows:

$$\frac{\begin{array}{l} \forall y. B(x, y) \rightarrow N(y) \\ \forall y. N(y) \rightarrow T(y) \end{array}}{\forall y. B(x, y) \rightarrow T(y)}$$

Now consider:

$$\frac{\begin{array}{l} \text{Whatever } x \text{ believes is true.} \\ x \text{ believes that } A. \end{array}}{\text{It is true that } A.}$$

If we treat that-clauses as singular terms, and represent for a formula A the term 'the proposition that A ' by $[A]$ then we can formalize this inference as follows:

$$\frac{\begin{array}{l} \forall y. B(x, y) \rightarrow T(y) \\ B(x, [A]) \end{array}}{T([A])}$$

But now consider:

$$\frac{\begin{array}{l} \text{Whatever } x \text{ believes is true.} \\ x \text{ believes that } v \text{ believes something.} \end{array}}{\text{It is true that } v \text{ believes something.}}$$

By analogy we would formalize this as follows:

$$\frac{\forall y.B(x, y) \rightarrow T(y)}{\frac{B(x, [\exists v.B(x, v)])}{T([\exists v.B(x, v)])}}$$

But now consider the inference:

$$\frac{x \text{ believes that he believes something.}}{\text{There is someone } v \text{ such that } x \text{ believes that } v \text{ believes something.}}$$

A candidate for formalization would be:

$$\frac{B(x, [\exists y.B(x, y)])}{\exists v.B(x, [\exists y.B(v, y)])}$$

In conclusion: *that-clauses should be treated as singular terms which can contain externally quantifiable variables.*

3.2 Semantic Problems

We need to find a way of dealing with substitutivity failures for materially equivalent and co-referential formulas. For instance:

$$\frac{\begin{array}{l} x \text{ believes that everything runs.} \\ \text{Everything runs if and only if everything walks.} \end{array}}{x \text{ believes that everything walks.}}$$

$$\frac{\begin{array}{l} B(x, [\forall y.R(y)]) \\ \forall y.R(y) \leftrightarrow \forall y.W(y) \end{array}}{B(x, [\forall y.W(y)])}$$

$$\frac{\begin{array}{l} x \text{ wonders whether } y \text{ is the author of } Waverley \\ y \text{ is the author of } Waverley \end{array}}{x \text{ wonders whether } y = y}$$

$$\frac{\begin{array}{l} W(x, [y = \iota z.A(z)]) \\ y = \iota z.A(z) \end{array}}{W(x, [y = y])}$$

Intensional entities can thus be different even if their *extension* be the same. What kind of entities do the $[A]$ correspond to ? Some approaches:

- Linguistic entities (Carnap, Quine): structural descriptions of a formula or inscription of a formula. But there is a difficulty with the Langford-Church translation test.
- Nothing at all: the above metalinguistics names and descriptions can be analysed away completely (Scheffler). According to Bealer these approaches seem to violate Davidson's learnability requirement.
- Extra-linguistic entities such as propositions (Bealer's neologicism).

According to Bealer there are two kinds of intensional entities. According to the first (traced back to Leibniz and one of Church's formulations of Frege's theory of *sense*) two intensional entities are equal *if and if they are necessarily equivalent*. This is related to possible-worlds semantics and deals with necessity and possibility. A definition need only be a necessary equivalence

According to the second conception each intensional entity when defined completely *must possess a unique non-circular definition*. According to Bealer this is found in Leibniz's distinction between simple and complex properties and Russell's doctrine of logical atomism. This conception is used to treat intentionality: belief, desire, perception, etc.

Example of a deduction valid in the first conception but not in the second:

$$\frac{\begin{array}{l} x \text{ wonders whether there are trilaterals which are not triangles.} \\ \text{Necessarily all and only triangles are trilaterals.} \end{array}}{x \text{ wonders whether there are triangles which are not triangles.}}$$

Other nominative formations in language such as the gerundive and infinitive are formalized by *intensional abstraction* over variables $[A]_{x_1, \dots, x_n}$. Intensional abstracts in general thus correspond to *propositions, relations and properties*.

3.3 Bealer's Intensional Logic

What is Bealer's intensional logic ? The language of Bealer's intensional logic is that of first-order logic with equality with a finite number of predicates symbols extended with term-forming intensional abstraction operators $[]_{x_1 \dots x_n}$. There are two kinds of intensional entities and thus two distinct axiomatic systems of intensional logic, $T1$ and $T2$. Bealer's formulates two new classes of model for which $T1$ and $T2$ are both sound and complete. $T1$ contains standard S5 modal logic. Using a single primitive binary predication predicate Δ and additional so-called "predication" axiom-schemes we obtain a neo-Fregean formulation of arithmetic. In [1][ch.7] Bealer argues that if we accept the ontology of intensional entities then we can translate intensional logic into standard extensional logic using standard Tarskian semantics (the thesis of extensionality).

The multi-modal approach originating with C. I. Lewis, Carnap, Hintikka and Kripke has some disadvantages. Operators apply only to formulas and not to singular terms so many intuitively valid arguments cannot be captured. It is eclectic and incomplete rather than based on a single intensional operator. According to Bealer, the higher-order approach has the following disadvantages:

- Does not have a sound and complete semantics.
- Predicates can be linguistic subjects which leads us to consider two sorts of meaning relation to avoid Frege's 'a=a'/'a=b' puzzle.
- Needs type theory to avoid paradoxes but this imposes implausible existence restrictions on intensional entities.
- Division of variables into sorts makes it impossible to deal with "transcendental" predicates without violating Davidson's learnability requirement.
- Requires possible-world semantics and commitment to non-actual possibilia.

We recommend the paper [3] for an interesting further discussion and defense of Bealer's Intensional Logic.

4 Bealer Decomposition

Definition 4.1 The language L^ω consists of a countable ordered collection of variables x, y, z, \dots and n -ary predicate symbols ($n \geq 1$) F, G, H, \dots , with a distinguished binary predicate $=$, logical connectives $\&, \sim, \exists$ and the intensional abstraction operator $[\]_{x_1 \dots x_n}$ where $x_1 \dots x_n$ is a possibly empty sequence of distinct variables.

We use the notation \bar{x} for a (possibly empty) sequence of distinct variables. We use lower-case letters (possibly with subscripts) towards the end of the Latin alphabet to denote variables.

Definition 4.2 Formulas and terms of L^ω are defined by simultaneous induction:

- Variables are terms.
- If t_1, \dots, t_n are terms and F is n -ary predicate, then $F(t_1, \dots, t_n)$ is a formula.
- If A and B are formulas, v a variable, then $(A \& B), \neg A$ and $\exists v.A$ are formulas.
- If A is a formula and $v_1, \dots, v_m, 0 \leq m$, is a sequence of *distinct* variables, then $[A]_{v_1 \dots v_m}$ is a term.

A term of the form $[A]_{v_1, \dots, v_n}$ is called an *intensional abstract*. Intensional abstracts of the form $[F(v_1, \dots, v_n)]_{v_1 \dots v_n}$ are called *elementary*. Intensional abstracts $[A]_\alpha$ where A is not an atomic formula are called *complex*.

The intensional abstraction operator generalises the set-theoretic extension operator $\{x : A\}$. The standard notions of bound and free variable, bound-variable renaming α -equivalence) as well as a term being free for x in A carry over to L^ω in the expected way (note that intensional abstraction binds variables like quantifiers). Terms that differ only by bound variable renaming will be called *variants*.

We will define a set of (partially defined) syntactic operations on intensional abstracts and an algorithm to decompose an intensional abstract in a unique way in terms of such syntactic operations, elementary intensional abstracts and variables.

A term $[A]_{\bar{x}}$ is *normalized* if all the variables in \bar{x} are free in A and they display the order in which these variables first occur free.

If A is atomic then we call $[A]_{\bar{x}}$ a *prime term* if the variables in \bar{x} are free in A , independently of the order.

Given an atomic formula $F(t_1, \dots, t_n)$ if a variable occurs free in more than one of the terms t_i then it is called a *reflected variable*. A prime term $[A]_{\bar{x}}$ is called a *prime reflection term* if \bar{x} contains a reflected variable in A .

Consider a prime term $[F(t_1, \dots, [B]_{\bar{y}}, \dots, t_j)]_{\bar{x}}$ which is not a prime reflection term. If there is a variable in \bar{x} which is free in $[B]_{\bar{y}}$ and all the previous arguments are variables in \bar{x} , then the prime term is a *prime relativized predication term* and such a variable is called a *relativized variable*.

Consider a prime term $[F(t_1, \dots, t_k, \dots, t_j)]_{\bar{x}}$ which is not a prime reflection term. If there is term t_k which is a variable not in \bar{x} or which contains no free variables which are in \bar{x} and all the previous arguments are variables in \bar{x} , then it is a *prime absolute predication term*.

These definitions allow us to divide all intensional abstracts into seven disjoint categories. The following is clear:

Lemma 4.3 *Given an intensional abstract $[A]_{\bar{x}}$ it falls into one and one only of the following seven categories which depend only on A and \bar{x} independently of order (only on the underlying set). They are also invariant under the renaming of bound variables.*

1. Complex and \bar{x} contains variables not free in A .
2. Complex and all the variables in \bar{x} are free in A .
3. Not complex and \bar{x} contains variables not free in A .
4. Prime reflection term.
5. Prime relativized predication term.
6. Prime absolute predication term.
7. $[A]_{\bar{x}'}$ is elementary for some permutation \bar{x}' of \bar{x} .

Before defining Bealer decomposition we need the following considerations on permutations. It is clear that any permutation $\sigma^n \in S^n (n \geq 2)$ can be decomposed into permutations of the form $\sigma_c^n = (1 \ 2 \ \dots \ n)$ (for $n \geq 3$) and $\sigma_i^n = (n-1 \ n)$. However, we need a uniquely defined decomposition defined as follows. Assume that our permutations are acting on sequences $x_1 \dots x_n$ of variables and let $x'_1 \dots x'_n$ be a permutation of $x_1 \dots x_n$. We use the notation $\sigma^n x_1 \dots x_n$ to denote the results of apply σ^n to the sequence. We associate to $x_1 \dots x_n$ and $x'_1 \dots x'_n$ (or equivalently the permutation which transforms the first sequence into the second) a uniquely defined decomposition as follows:

Suppose $\pi' = \pi_n \dots \pi_2 \pi_1$ has already been defined, where π_i is either σ_c^n or σ_i^n . Then let s be the maximal subsegment of $\pi' x_1 \dots x_n$ equal to an initial segment $b = x'_1 \dots x'_k$ of $x'_1 \dots x'_n$. If b is the whole $x'_1 \dots x'_n$ we are done. Otherwise x'_{k+1} will correspond to a certain x in $\pi' x_1 \dots x_n$. If x is the last element of this sequence then we choose σ_i^n , otherwise we chose σ_c^n . This process terminates and the decomposition is clearly unique. Given a permutation σ (or two sequences \bar{x}, \bar{x}' of the same length), we denote its decomposition by $\beta(\sigma)$ (or $\beta(\bar{x}, \bar{x}')$).

Example 4.4 Consider $x_1 x_2 x_3 x_4$ and $x_2 x_4 x_3 x_1$. The the process yields σ_i^4 and $x_1 x_2 x_4 x_3$, $\sigma_c^4 \sigma_i^4$ and $x_3 x_1 x_2 x_4$, $\sigma_c^4 \sigma_c^4 \sigma_i^4$ and $x_4 x_3 x_1 x_2$ and finally $\sigma_c^4 \sigma_c^4 \sigma_c^4 \sigma_i^4$ and $x_2 x_4 x_3 x_1$.

Definition 4.5 The Bealer operations consists in eight syntactic operations defined on intensional abstracts. These consists in six unary operations C, I, N, U, E, R , a binary operation K and partially-defined binary operations P_n for $n \geq 0$. These operations can be seen as defined on equivalence classes modulo renaming of bound variables. Let $[A]_{\bar{x}}$ be an intensional abstract with \bar{x} of length n . Then we have:

$$C[A]_{\bar{x}} = [A]_{\sigma_c^n \bar{x}} \quad n \geq 3$$

$$I[A]_{\bar{x}} = [A]_{\sigma_i^n \bar{x}} \quad n \geq 2$$

$$N[A]_{\bar{x}} = [\neg A]_{\bar{x}}$$

$$U[A]_{\bar{x}} = [\exists x_n A]_{x_1 \dots x_{n-1}} \quad n \geq 1$$

$$E[A]_{\bar{x}} = [A]_{\bar{x}y} \quad y \text{ the first variable not in } A \text{ or } \bar{x}$$

$$R[A]_{\bar{x}} = [A[x_{n-1}/x_n]]_{x_1 \dots x_{n-1}}$$

$$K[A]_{\bar{x}}[B]_{\bar{y}} = [A \& B']_{\bar{x}} \quad [B']_{\bar{x}} \text{ variant of } [B]_{\bar{y}}$$

$$P_0[A]_{\bar{x}}[B]_{\bar{y}} = [A[[B]_{\bar{y}}/x_n]]_{x_1 \dots x_{n-1}} \quad \text{where } n \geq 1 \text{ and } [B]_{\bar{y}} \text{ is free for } x_n \text{ in } [A]_{x_1 \dots x_{n-1}}$$

$$P_n[A]_{\bar{x}}[B]_{\bar{y}} = [A[[B]_{\bar{z}}/x_n]]_{x_1 \dots x_{n-1}\bar{w}} \quad \text{where } \bar{y} = \bar{z}\bar{w}, \bar{w} \text{ length } n \geq 1, [B]_{\bar{z}} \text{ is free for } x_n \text{ in } [A]_{x_1 \dots x_{n-1}}$$

Consider a sequence \bar{x} and a permutation \bar{x}' . We associate to $\beta(\bar{x}, \bar{x}')$ a sequence $\Sigma(\bar{x}, \bar{x}')$ of operations I and C obtained by replacing σ_i^n with I and σ_c^n with C in $\beta(\bar{x}, \bar{x}')$. Thus in the previous example we have $\Sigma(x_1x_2x_3x_4, x_2x_4x_3x_1) = CCCI$. This sequence is to interpreted compositionally. In general we shall use the notation Σ for any (possibly empty) sequence of operations C and I . We use the notation E^n to denote E composed n times. Also we shall use the notation $P_0^n tt_1 \dots t_n$ to mean $P_0 \dots P_0(P_0 tt_1)t_2 \dots t_n$.

We now define Bealer decomposition.

Definition 4.6 Given an intensional abstract $[A]_{\bar{x}}$ or variable x the Bealer decomposition $\mathcal{B}[A]_{\bar{x}}$ is defined inductively as follows:

- Suppose $[A]_{\bar{x}}$ belongs to categories 1 or 3. And let \bar{x}' be a permutation of \bar{x} of the form $\bar{y}\bar{z}$ so that $[A]_{\bar{y}}$ is normalized and \bar{z} (of length m) consists of the variables in \bar{x} that do not occur free in A in order of their occurrence in \bar{x} . Then $\mathcal{B}[A]_{\bar{x}} := \Sigma(\bar{x}, \bar{x}')E^m\mathcal{B}[A]_{\bar{y}}$.
- Suppose $[A]_{\bar{x}}$ belongs to category 2 and is not normalized. And let \bar{x}' be a permutation of \bar{x} so that $[A]_{\bar{x}'}$ is normalized. Then $\mathcal{B}[A]_{\bar{x}} := \Sigma(\bar{x}, \bar{x}')\mathcal{B}[A]_{\bar{x}'}$.
- Suppose the term belongs to category 2 and is normalized. Then we have

$$\mathcal{B}[A \& B]_{\bar{x}} := K\mathcal{B}[A]_{\bar{x}}\mathcal{B}[B]_{\bar{x}}$$

$$\mathcal{B}[\neg A]_{\bar{x}} := N\mathcal{B}[A]_{\bar{x}}$$

$$\mathcal{B}[\exists v A]_{\bar{x}} := U\mathcal{B}[A]_{\bar{x}v} \text{ where } v \text{ is the first variable not in } A \text{ or } \bar{x}.$$

- Let the term belong to category 4. Of the reflected variables in \bar{x} consider the one v with the right-most occurrence and let be t the right-most argument of A in which v has a free occurrence. Let w be the alphabetically earliest variable not occurring in A or \bar{x} . Let \bar{x}' be the permutation of \bar{x} such that $[F(t_1, \dots, t_n)]_{\bar{x}'}$ is normalized and consider $\bar{y}v$ where \bar{y} is \bar{x}' with v removed. Then

$$\mathcal{B}[F(t_1, \dots, t_n)]_{\bar{x}} := \Sigma(\bar{x}, \bar{y}v)R\mathcal{B}[F(\dots t[w/v] \dots)]_{\bar{y}vw}$$

- Let the term belong to category 6 and let t be the argument which is not a variable in \bar{x} or which contains no free variables which are in \bar{x} and such that all the previous arguments are variables in \bar{x} . Let v be the alphabetically earliest variable not occurring in A or \bar{x} . Let \bar{x}' be the permutation of \bar{x} such that $[F(t_1, \dots, t, \dots, t_n)]_{\bar{x}'}$ is normalized. Then

$$\mathcal{B}[F(\dots, t, \dots)]_{\bar{x}} := \Sigma(\bar{x}, \bar{x}')P_0\mathcal{B}[F(\dots, v, \dots)]_{\bar{x}'v}\mathcal{B}t$$

- Let the term belong to category 5. Then it is a prime term $[F(t_1, \dots, [B]_{\bar{y}}, \dots, t_j)]_{\bar{x}}$, where there is a variable in \bar{x} which is free in $[B]_{\bar{y}}$ and such that all the previous arguments are variables in \bar{x} . Let \bar{z} be the sequence, as they first occur, of all m free variables in $[B]_{\bar{y}}$ which are in \bar{x} . Let v be the alphabetically earliest variable not occurring in A or \bar{x} . Then if \bar{x}' is the normalizing permutation of \bar{x} it is of the form $\bar{x}''\bar{z}\bar{x}'''$. Let $\bar{w} = \bar{x}''\bar{x}'''\bar{z}$. Then

$$\mathcal{B}[F(\dots, [B]_{\bar{y}}, \dots)]_{\bar{x}} := \Sigma(\bar{x}, \bar{w})P_m\mathcal{B}[F(\dots, v, \dots)]_{\bar{x}''\bar{w}v}\mathcal{B}[B]_{\bar{y}\bar{z}}$$

- Finally let the term be in category 7 and let \bar{x}' be such that $[F(x_1, \dots, x_n)]_{\bar{x}'}$ is elementary.

$$\mathcal{B}[F(x_1, \dots, x_n)]_{\bar{x}} := \Sigma(\bar{x}, \bar{x}')[F(x_1, \dots, x_n)]_{\bar{x}'}$$

- In the case of a variable we have $\mathcal{B}x := x$.

It is easy to check that this decomposition is well-defined and unique modulo renaming of bound variables. Bealer decomposition transforms an intensional abstract into an expression (in Polish notation) involving the eight Bealer operations, elementary abstracts and variables.

Example 4.7 Consider the term $[\neg F(x, [G(x, y)])]_y$. Then

$$\begin{aligned} \mathcal{B}[\neg F(x, [G(x, y)])]_y &= N\mathcal{B}[F(x, [G(x, y)])]_y = NP_0\mathcal{B}[F(v, [G(x, y)])]_{yv}x \\ &= NP_0I\mathcal{B}[F(v, [G(x, y)])]_{vy}x = NP_0IP_1\mathcal{B}[F(v, w)]_{vw}\mathcal{B}[G(x, y)]_yx \\ &= NP_0IP_1\mathcal{B}[F(v, w)]_{vw}P_0\mathcal{B}[G(u, y)]_{yu}xx = NP_0IP_1\mathcal{B}[F(v, w)]_{vw}P_0I\mathcal{B}[G(u, y)]_{uy}xx \end{aligned}$$

The following observation will be used further ahead:

Lemma 4.8 *Let $[F(t_1, \dots, t_n)]_{\bar{x}}$ be a prime term. Then its Bealer decomposition has the form*

$$\Gamma_1 \dots \Gamma_m [F(v_1, \dots, v_n)]_{v_1 \dots v_n} \mathcal{B}t'_{i_1} \dots \mathcal{B}t'_{i_m}$$

where the Γ_i consist in sequences of the form

$$\Sigma_1 R \Sigma_2 R \Sigma_3 \dots \Sigma_{l-1} R \Sigma_l P_{n_i}$$

and $t'_{i_k} = s_{\bar{y}}$ where $s = t_{i_k}$ and if $s = [A]_{\bar{z}}$ then $s_{\bar{y}}$ is $[A]_{\bar{z}\bar{y}}$ where \bar{y} is the sequence of free variables (as they first occur) of s which are in \bar{x} . If s is a variable then \bar{y} is empty and $t'_{i_k} = s$.

5 The Logic of Necessity

We now consider Bealer's axiomatic system T1 over the language L^ω . The conception of intensionality behind T1 is that there is only one necessary truth and that the intensional abstracts of necessarily equivalent formulas are equal and interchangeable. We define necessitation in terms of equality of intensional abstracts:

$$\Box A \equiv_{df} [A] = [[A] = [A]]$$

Here $[[A] = [A]]$ represents the one necessary truth.

A Hilbert system for T1 consists of the following axioms and rules:

1. All propositional tautologies
2. (Inst) $\forall v. A(v) \rightarrow A(t)$, t free for v in A .
3. (QImp) $\forall x. (A \rightarrow B) \rightarrow (A \rightarrow \forall x. B)$, x not free in A
4. (Id) $x = x$
5. (L) $x = y \rightarrow (A(x, x) \leftrightarrow A(x, y))$, y free for x in the positions it replaces x .
6. $[A]_{\bar{x}} \neq [B]_{\bar{y}}$, for \bar{x} and \bar{y} of different lengths

7. $[A]_{\bar{x}} = [A']_{\bar{x}'}$ (equality modulo renaming bound variables)
8. (B) $[A]_{\bar{x}} = [B]_{\bar{x}} \leftrightarrow \Box \forall \bar{x}. (A \leftrightarrow B)$
9. (T) $\Box A \rightarrow A$
10. (K) $\Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)$
11. (S5) $\Diamond A \rightarrow \Box \Diamond A$
1. (MP) From A and $A \rightarrow B$ infer B
2. (N) From A infer $\Box A$
3. (Gen) From A infer $\forall x.A$

Here, as usual, $\Diamond A =_{df} \neg \Box \neg A$.

In order to prove soundness and completeness it is convenient to replace axioms (L),(B) and (S5) with:

1. (L') $x = y \rightarrow (A(x, x) \rightarrow A(x, y))$, y free for x in the positions it replaces x , A atomic.
2. (B1) $\Box(A \leftrightarrow B) \leftrightarrow [A] = [B]$
3. (B2) $\forall v. [A(v)]_{\bar{x}} = [B(v)]_{\bar{x}} \leftrightarrow [A(x)]_{\bar{x}v} = [B(v)]_{\bar{x}v}$
4. (S5') $x \neq y \rightarrow \Box x \neq y$

The rest of this section is devoted to proving that we obtain equivalent axiomatic systems.

Lemma 5.1 *In T1 we have:*

- i) $\vdash x = y \rightarrow \Box x = y$
- ii) $\vdash \neg \Box x = y \leftrightarrow x \neq y$

Proof. For i), by the identity axiom we have $x = x$. Hence using the necessitation rule we get

$$\Box x = x$$

which is

$$[x = x] = [[x = x] = [x = x]]$$

We denote this expression by $B(x, x)$. Using $B(x, x)$ in (L)

$$x = y \rightarrow A(x, x) \leftrightarrow A(x, y)$$

which is equivalent to $A(x, x) \rightarrow (x = y \rightarrow A(x, y))$ we obtain as desired.

For ii), take the converse of the (T) instance $\Box x = y \rightarrow x = y$ to obtain $x \neq y \rightarrow \neg \Box x = y$. For the other direction simply take the converse of i) ■

Lemma 5.2 *We obtain an equivalent system if in T1 we replace (S5) with*

$$(*) \ x \neq y \rightarrow \Box x \neq y.$$

Proof. Assuming (*), since $\Diamond A$ is by definition $\neg([\neg A] = [[\neg A] = [\neg A]])$ we get (S5) immediately.

In the other direction take the (S5) instance

$$\neg\Box\neg\neg x = y \rightarrow \Box\neg\Box\neg\neg x = y$$

that is,

$$(**) \quad \neg\Box x = y \rightarrow \Box\neg\Box x = y$$

Now by ii) of the previous lemma we have

$$\neg\Box x = y \rightarrow \neg x = y$$

Applying necessitation and using (K) we get

$$\Box\neg\Box x = y \rightarrow \Box\neg x = y$$

and combining with (**) we get as desired. ■

We will use the standard properties of S5 modal logic:

Lemma 5.3 *In T1 we have*

- i) $\vdash A \rightarrow \Diamond A$
- ii) $\vdash A \rightarrow \Box\Diamond A$
- iii) $\Diamond\Box A \rightarrow A$
- iv) *If $\vdash A \rightarrow B$ then we can derive $\vdash \Diamond A \rightarrow \Diamond B$*
- v) *We have $\vdash \Diamond A \rightarrow B$ iff $\vdash A \rightarrow \Box B$*

Proof. For i) take the converse of the (T) instance $\Box\neg A \rightarrow \neg A$. For ii) combine i) and (S5). For iii) take the converse of the (S5) instance $\neg\Box\Diamond\neg A \rightarrow \neg\Diamond\neg A$ and use (T). For iv) take the converse of the premise, apply (T) and take the converse again. Finally v) follows easily from (T), iv), iii) and ii). ■

Lemma 5.4 $\vdash \Box\forall v.A(v) \rightarrow \forall v.\Box A(v)$

Proof. Take the (Ins) instance

$$\forall v.A(v) \rightarrow A(v)$$

Applying necessitation, (K) and modus ponens yields:

$$\Box\forall v.A(v) \rightarrow \Box A(v)$$

Applying generalization:

$$\forall v.(\Box\forall v.A(v) \rightarrow \Box A(v))$$

Since v is not free on the left side of the implication we can apply (QImp) and modus ponens to obtain

$$\Box\forall v.A(v) \rightarrow \forall v.\Box A(v)$$

■

Lemma 5.5 *We have the Barcan Formula $\vdash \forall v. \Box A(v) \rightarrow \Box \forall v. A(v)$.*

Proof. We have by (Ins)

$$\forall v. \Box A \rightarrow \Box A$$

Applying \Diamond yields, by lemma 5.3 iii) and iv):

$$\Diamond \forall v. \Box A \rightarrow \Diamond \Box A \rightarrow A$$

Hence by generalization, (QImp) and modus ponens:

$$\Diamond \forall v. \Box A \rightarrow \forall v. A$$

which again by lemma 5.3 yields

$$\forall v. \Box A \rightarrow \Box \forall v. A$$

■

We can now prove:

Lemma 5.6 *We obtain an equivalent system to T1 if instead of (B) take*

$$(B'1) \quad \Box(A \leftrightarrow B) \leftrightarrow ([A] = [B])$$

$$(B'2) \quad \forall v. ([A(v)]_{\bar{x}} = [B(v)]_{\bar{x}}) \leftrightarrow [A(v)]_{\bar{x}v} = [B(v)]_{\bar{x}v}$$

Proof. Using Barcan's formula and its converse it is easy to see that (B'1) and (B'2) follow from (B).

Assume (B'1) and (B'2). We must show that

$$[A]_{\bar{x}} = [B]_{\bar{x}} \leftrightarrow \Box \forall \bar{x}. (A \leftrightarrow B)$$

Let $\bar{x} = \bar{x}'v$.

Since $[A]_{\bar{x}'v} = [B]_{\bar{x}'v} \leftrightarrow \forall v. ([A(v)]_{\bar{x}'} = [B(v)]_{\bar{x}'})$ using (B'2) repeatedly we get

$$[A]_{\bar{x}v} = [B]_{\bar{x}v} \leftrightarrow \forall \bar{x}, v. ([A(v)] = [B(v)])$$

Using (B'1) we get

$$[A]_{\bar{x}} = [B]_{\bar{x}} \leftrightarrow \forall \bar{x}. \Box(A \leftrightarrow B)$$

Using repeatedly Barcan's formula and its converse we get as desired.

■

Finally:

Lemma 5.7 *We obtain an equivalent system to T1 when the Leibniz axiom (L) is restricted to atomic predicates and restricted to the form (L') $x = y \rightarrow (A(x, x) \rightarrow A(x, y))$.*

Proof. L obviously implies (L') as a particular case. In the other direction we first show that $x = y \rightarrow (A(x, x) \leftarrow A(x, y))$. We have that $x = y \rightarrow (A(x, x) \rightarrow A(x, y))$ and $A(x, y)$ arises from $A(x, x)$ by replacing some occurrences of x by y . So $A(x, x)$ arises from replacing some occurrences of y in $A(x, y)$ by x . Hence we have $y = x \rightarrow (A(x, y) \rightarrow A(x, x))$. Also from (L') we can derive $x = y \rightarrow y = x$ and using this we get as desired. We can now proceed inductively on the structure of the formula A . The case of negation is immediate. For conjunction we use the tautology $(A \rightarrow B) \rightarrow (C \rightarrow D) \rightarrow (A \& C \rightarrow B \& D)$ and for quantification we use generalisation, (QImp) and modus ponens. ■

This lemma justifies that "there is only one necessary truth".

Lemma 5.8 *We have $\forall x, y. [x = x] = [y = y]$.*

Proof. We have $x = x \leftrightarrow y = y$. We apply necessitation and (B'1) and then generalisation to get as desired. ■

6 The Logic of Definition

The logic T1 is insufficient in many instances.

Lemma 6.1 *In T1 if $\vdash W(x, [\exists y(T(y) \& \neg S(y))])$ and $\vdash \Box \forall y(S(y) \leftrightarrow T(y))$ then*

$$\vdash W(x, [\exists y(T(y) \& \neg T(y))]).$$

Proof. We only need to show that $[\exists y(T(y) \& \neg S(y))] = [\exists y(T(y) \& \neg T(y))]$, that is, by (B), $\Box(\exists y(T(y) \& \neg S(y)) \leftrightarrow \exists y(T(y) \& \neg T(y)))$. From the second hypothesis we get $\vdash S(y) \leftrightarrow T(y)$ and using the tautology:

$$(T(y) \leftrightarrow S(y)) \rightarrow ((T(y) \& \neg S(y)) \leftrightarrow (T(y) \& \neg T(y)))$$

we get easily

$$\forall y(T(y) \leftrightarrow S(y)) \rightarrow \exists y((T(y) \& \neg S(y)) \leftrightarrow (T(y) \& \neg T(y)))$$

and the result follows from necessitation, (K), modus ponens and (B).

To avoid this in our intensional logic we accept only one implication in axiom (B).

Bealer's logic T2 is given the following Hilbert axiomatic system:

1. All propositional tautologies
2. (Ins) $\forall x.A(x) \rightarrow A(t)$, t free for v in A .
3. (QImp) $\forall x.(A \rightarrow B) \rightarrow (A \rightarrow \forall x.B)$, x not free in A
4. (Id) $x = x$
5. (L) $x = y \rightarrow (A(x, x) \leftrightarrow A(x, y))$, y free for x in the positions it replaces x .
6. $[A]_{\bar{x}} \neq [B]_{\bar{y}}$, \bar{x} and \bar{y} of different lengths
7. $[A]_{\bar{x}} = [A']_{\bar{x}'}$ (equality modulo renaming bound variables)
8. $[A]_{\bar{x}} = [B]_{\bar{x}} \rightarrow (A \leftrightarrow B)$

9. $[A]_{\bar{x}} \neq [B]_{\bar{y}}$, non-elementary terms belonging to different categories.
10. Let $t' = Bt$ and $r' = Br$ where B is the first operation in the Bealer decomposition of t' and r' . Then $t = r \leftrightarrow t' = r'$
11. Let $t = Bt'r'$ and $r = Bt''r''$ where B is the first operation in the Bealer decomposition of t and r . The $t = r \leftrightarrow t' = t'' \& r' = r''$.
12. If $[F(x_1, \dots, x_n)]_{x_1 \dots x_n} = s$ and G occurs in s then $[G(y_1, \dots, y_m)]_{y_1 \dots y_m} \neq t$ whenever F occurs in t (Non-circularity)
1. (MP) From A and $A \rightarrow B$ infer B
2. (TGen) Suppose that F does not occur in $A(v)$. If $\vdash A([F(x_1, \dots, x_n)]_{x_1 \dots x_n})$ then we can infer $\vdash A(t)$ whenever t is complex, of arity n and free for v in A .
3. (Gen) From A infer $\forall x.A$

7 Model Structures

Let $F = \emptyset$ and $T = \{\emptyset\}$. A model \mathcal{M} consists of a set \mathcal{D} with a decomposition into the union of *disjoint* sets \mathcal{D}_i for $i \geq -1$, a distinguished element $id \in \mathcal{D}_2$, a set \mathcal{H} of functions H whose domain is \mathcal{D} and such that H is the identify on \mathcal{D}_{-1} and on \mathcal{D}_0 has range $\{T, F\}$ and on \mathcal{D}_i has range $\mathcal{P}(\mathcal{D}^i)$ where $\mathcal{D}^i = \mathcal{D} \times \dots \times \mathcal{D}$ (i terms) for $i \geq 1$. All these functions must satisfy $H(id) = \{(x, y) : x = y\}$. There is a distinguished function \mathcal{G} representing the *actual extension*. Furthermore we have unary (partial) operations n, e, u, c, i, r and binary (partial) operations k, p_n with $n \geq 0$. These operations restrict to the following domains and ranges:

$$\begin{aligned}
n &: \mathcal{D}_i \rightarrow \mathcal{D}_i, i \geq 0 \\
e &: \mathcal{D}_i \rightarrow \mathcal{D}_{i+1}, i \geq 0 \\
u &: \mathcal{D}_i \rightarrow \mathcal{D}_{i-1}, i \geq 1 \text{ and } u : \mathcal{D}_0 \rightarrow \mathcal{D}_0. \\
c &: \mathcal{D}_i \rightarrow \mathcal{D}_i, i \geq 3 \\
i &: \mathcal{D}_i \rightarrow \mathcal{D}_i, i \geq 2 \\
r &: \mathcal{D}_i \rightarrow \mathcal{D}_{i-1}, i \geq 2 \\
a &: \mathcal{D}_i \times \mathcal{D}_i \rightarrow \mathcal{D}_i, i \geq 0 \\
p_n &: \mathcal{D}_i \times \mathcal{D}_j \rightarrow \mathcal{D}_{i+n-1}, i \geq 1, j \geq n
\end{aligned}$$

We define inductively $p_0^n(d, x_1, \dots, x_n) = p_0(p_0^{n-1}(d, x_2, \dots, x_{n-1}), x_1)$ and $p_0^1(d, x_1) = p_0(d, x_1)$.

Furthermore we have the important constraints on how the functions in \mathcal{H} relate to these operations. Here d denotes a suitable member of \mathcal{D}^i . For all $\mathcal{H} \in H$ we must have:

$$\begin{aligned}
d' \in \mathcal{H}n(d) &\text{ iff } d' \notin \mathcal{H}d \\
(x_1, \dots, x_i, x_{i+1}) \in \mathcal{H}e(d) &\text{ iff } (x_1, \dots, x_i) \in \mathcal{H}d \\
(x_1, \dots, x_{i-1}) \in \mathcal{H}u(d) &\text{ iff there is a } x_i \text{ such that } (x_1, \dots, x_i) \in \mathcal{H}d \\
(x_1, \dots, x_{i-1}, x_i) \in \mathcal{H}c(d) &\text{ iff } (x_i, x_1, \dots, x_{i-1}) \in \mathcal{H}d \\
(x_1, \dots, x_{i-1}, x_i) \in \mathcal{H}i(d) &\text{ iff } (x_1, \dots, x_i, x_{i-1}) \in \mathcal{H}d \\
(x_1, \dots, x_{i-1}, x_i) \in \mathcal{H}r(d) &\text{ iff } (x_1, \dots, x_i, x_i) \in \mathcal{H}d \\
d'' \in \mathcal{H}k(d, d') &\text{ iff } d'' \in \mathcal{H}d \text{ and } d' \in \mathcal{H}d' \\
(x_1, \dots, x_n) \in \mathcal{H}p_0(d, d') &\text{ iff } (x_1, \dots, x_n, d') \in \mathcal{H}d \\
(x_1, \dots, x_{i-1}, y_1, \dots, y_n) \in \mathcal{H}p_n(d, d') &\text{ iff } (x_1, \dots, x_{i-1}, p_0^n(d', y_1, \dots, y_n)) \in \mathcal{H}d
\end{aligned}$$

Here we consider also the 0-tuple as being \emptyset . Notice that elements of D^0 are propositions seen as objective entities according to their meaning. The extension functions \mathcal{H} determine their Boolean values which are $T = \{\emptyset\}$ and $F = \emptyset$.

Definition 7.1 Given a model \mathcal{M} , an *interpretation* \mathcal{I} assigns i -ary predicate elements of \mathcal{D}_i and $=$ to id . An *assignment* \mathcal{A} assigns to variables elements in \mathcal{D} .

Bealer decomposition is crucial to define the denotation of a term relative to a model, interpretation and assignment.

Definition 7.2 Given \mathcal{M}, \mathcal{I} and \mathcal{A} we define the *denotation* $D_{\mathcal{I}, \mathcal{A}, \mathcal{M}}t$ of a term t of L^ω as follows. If t is a variable then it is $\mathcal{A}(t)$. If t is an elementary term $[F(v_1, \dots, v_n)]_{v_1 \dots v_n}$ then it is $\mathcal{I}(F)$. Otherwise, consider the Bealer decomposition $\mathcal{B}t$ of t . If $\mathcal{B}t = Bt'$ (respectively $Bt't''$) where B is a Bealer operation then we define inductively

$$D_{\mathcal{I}, \mathcal{A}, \mathcal{M}}t = bD_{\mathcal{I}, \mathcal{A}, \mathcal{M}}t' \text{ (respectively } D_{\mathcal{I}, \mathcal{A}, \mathcal{M}}t = bD_{\mathcal{I}, \mathcal{A}, \mathcal{M}}t'D_{\mathcal{I}, \mathcal{A}, \mathcal{M}}t'')$$

where b is i, c, e, u, n, r, k, p_n if B is I, C, E, U, N, R, K, P_n respectively.

Definition 7.3 We say that a formula A is *true* for \mathcal{M}, \mathcal{I} and \mathcal{A} (we also write $T_{\mathcal{I}, \mathcal{A}, \mathcal{M}}(A)$) if $\mathcal{G}D_{\mathcal{I}, \mathcal{A}, \mathcal{M}}[A] = T$.

It is simple to see that the denotation is invariant modulo renaming bound variables. From now on, we will in most cases drop the subscripts in $D_{\mathcal{I}, \mathcal{A}, \mathcal{M}}$ and assume we are working with a given model, interpretation and assignment.

We are now interested in special classes of models for which T1 and T2 are sound and complete.

Definition 7.4 A model \mathcal{M} is called type 1 if for all $i \geq -1$, $x, y \in \mathcal{D}_i$

$$(\text{type 1 model condition}) \quad \forall \mathcal{H} \in H \quad \mathcal{H}(x) = \mathcal{H}(y) \rightarrow x = y$$

A model \mathcal{M} is called type 2 if the operations are:

1. one-to-one
2. disjoint on their ranges
3. and non-cycling

By the first two conditions each element of the model has a unique (possibly infinite) decomposition in terms of the operations. This decomposition is a tree having leaves indecomposable elements. The non-cycling condition means that the same element cannot appear in more than one place in the tree.

A formula A is T1-valid (we write $\vdash_{T1} A$) iff A is true for all type 1 model \mathcal{M} , interpretation \mathcal{I} and assignment \mathcal{A} . In the same way a formula A is T2-valid ($\vdash_{T2} A$) iff A is true for all type 2 model \mathcal{M} , interpretation \mathcal{I} and assignment \mathcal{A} .

Note that for type 2 models only \mathcal{G} is relevant. The type 1 condition restricts the possible operations on the model.

Example 7.5 A standard model for Kelley-Morse set theory furnishes an example of a type 1 model with a single extensional function. Type 1 models can be seen as generalised set theory models with a plurality of membership predicates $\in_{\mathcal{H}}$ such that the axiom of extensionality is only valid globally. Consider a language with a single unary predicate M . We can construct a type 2 model with no constants, D_0 , D_1 and $D_i = \emptyset$ for $i \geq 2$.

Lemma 7.6 *Let v be an externally quantifiable variable in $[B(v)]_{\bar{x}}$ and let t be free for v in $[B(v)]_{\bar{x}}$. Consider any model structure \mathcal{M} and any interpretation \mathcal{I} and assignment \mathcal{A} . Let \mathcal{A}' be an assignment which is just like \mathcal{A} except that $\mathcal{A}'(v) = Dt$. Then*

$$D_{\mathcal{A}'}[B(v)]_{\bar{x}} = D[B(t)]_{\bar{x}}$$

Proof. By induction on the Bealer decomposition of $[B(v)]_{\bar{x}}$. The base case is either an elementary abstract (in which case there are no externally quantifiable variables) or else a variable. If it is not v we are done and if it is v the result is trivial.

Suppose that $[B(v)]_{\bar{x}} = Bs(v)$ where B is some unary operation. If B is I , C , N , U or E then t is still free for v in $d(v)$ and hence by the induction hypothesis $D_{\mathcal{A}'}s(v) = Ds(t)$ so that $D_{\mathcal{A}'}[B(v)]_{\bar{x}} = bD_{\mathcal{A}'}s(v) = bDs(t) = DBs(t)$ where b is the corresponding model function. If B is R then there is a problem that s has a new bound variable which may occur in t in such a way that t is no longer free for v in s . But we may rename this bound variable to obtain an α -equivalent term s' where this problem does not occur. Note that the substitution *does not effect which of the seven syntactic categories an intensional abstract belongs to*. Suppose $[B(v)]_{\bar{x}} = Bs(v)s'(v)$ for a binary operation B . If B is A then t remains free for v in $s(v)$ and $s(v')$ and the previous argument applies. If B is P_k then we have the problem of the new bound variable in the first argument possibly occurring in t . We thus need to use α -equivalence so t remains free for v in $s(v)$. In the case of 0-predication we have

$$D_{\mathcal{A}'}[B(v)]_{\bar{x}} = p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, D_{\mathcal{A}'}s'(v))$$

If s' is not a variable or a variable distinct from v , then we may apply the induction hypothesis to obtain:

$$p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, D_{\mathcal{A}'}s'(v)) = p_0(D[B'(t)]_{\bar{x}w}, Ds'(t)) = D[B(t)]_{\bar{x}}$$

If s' is v then we obtain

$$\begin{aligned} & p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, D_{\mathcal{A}'}v) \\ &= p_0(D_{\mathcal{A}'}[B'(v)]_{\bar{x}w}, \mathcal{A}'v) \\ &= p_0(D[B'(t)]_{\bar{x}w}, Dt) \\ &= D[B(t)]_{\bar{x}} \end{aligned}$$

In the k -predication case for $k > 0$ we have, for $s'(v) = [C(v)]_{\bar{y}\bar{z}}$ where \bar{z} are relativized variables of $[C(v)]_{\bar{y}}$ in $[B(v)]_{\bar{x}}$

$$D_{\mathcal{A}'}[B(v)]_{\bar{x}} = p_k(D_{\mathcal{A}'}[B'(v)]_{\bar{x}'w}, D_{\mathcal{A}'}[C(v)]_{\bar{y}\bar{z}})$$

where \bar{x}' is obtained from \bar{x} by omitting \bar{z} . The result now follows easily from induction.

Lemma 7.7 *For all \mathcal{M} and \mathcal{I} , \mathcal{A} we have $D[A]_{x_1 \dots x_k} \in \mathcal{D}_k$*

Proof. This follows by easy induction from the definition of D and the model operations.

Given a sequence of Bealer operations of the form Σ we denote its associated n -permutation by σ . This permutation can also be applied to n -tuples (x_1, \dots, x_n) and we denote such an application by $\sigma(x_1, \dots, x_n)$ just as we denote its application to variable sequences by $\sigma x_1 \dots x_n$.

We will make frequent use of the following:

Lemma 7.8 *For any permutation σ of (x_1, \dots, x_n) we have*

$$\sigma(v_1, \dots, v_n) \in \mathcal{HD}[F]_{\sigma x_1, \dots, \sigma x_n} \leftrightarrow (v_1, \dots, v_n) \in \mathcal{HD}[F]_{x_1 \dots x_n}$$

Proof. Let the Bealer decomposition of $[F]_{x_1 \dots x_n}$ be $\Sigma_1 \mathcal{B}[F]_{x'_1 \dots x'_n}$ where $\mathcal{B}[F]_{x'_1 \dots x'_n}$ does not begin with C or I . Then Bealer decomposition of $[F]_{\sigma x_1, \dots, \sigma x_n}$ will be of the form $\Sigma_2 \mathcal{B}[F]_{x'_1 \dots x'_n}$. By the definition of denotation and model we have for any (x_1, \dots, x_n) :

$$(x_1, \dots, x_n) \in \mathcal{HD}[F]_{x_1 \dots x_n} \text{ iff } \sigma_1(x_1, \dots, x_n) \in \mathcal{HD}[F]_{x'_1 \dots x'_n}$$

$$(x_1, \dots, x_n) \in \mathcal{HD}[F]_{\sigma x_1, \dots, \sigma x_n} \text{ iff } \sigma_2(x_1, \dots, x_n) \in \mathcal{HD}[F]_{x'_1 \dots x'_n}$$

where $x'_1 \dots x'_n = \sigma_1(x_1, \dots, x_n)$ and $x'_1 \dots x'_n = \sigma_2 \sigma(x_1, \dots, x_n)$ and hence $\sigma = \sigma_2^{-1} \sigma_1$ and $\sigma_1 = \sigma_2 \sigma$.

Hence given (x_1, \dots, x_n) we have $(x_1, \dots, x_n) \in \mathcal{HD}[F]_{x_1 \dots x_n}$ iff $\sigma_1(x_1, \dots, x_n) = \sigma_2 \sigma(x_1, \dots, x_n) \in \mathcal{HD}[F]_{x'_1 \dots x'_n}$ iff $\sigma(x_1, \dots, x_n) \in \mathcal{HD}[F]_{\sigma x_1, \dots, \sigma x_n}$.

Lemma 7.9 *For all $\mathcal{I}, \mathcal{A}, \mathcal{M}$, $F(t_1, \dots, t_n)$ is true iff $(Dt_1, \dots, Dt_n) \in \mathcal{I}(F)$.*

Proof. By definition $F(t_1, \dots, t_n)$ is true iff $\mathcal{GD}([F(t_1, \dots, t_n)]) = T$. But

$$[F(t_1, \dots, t_n)] = P_0[F(v_1, \dots, v_n)]_{v_1} t_1 = P_0 P_p[F(v_1, v_2, \dots, v_n)]_{v_1 v_2} t_2 t_1 = P_0^n[F(v_1, \dots, v_n)]_{v_1 \dots v_n} t_n \dots t_2 t_1$$

hence by definition $\mathcal{GD}([F(t_1, \dots, t_n)]) = T$ iff

$$Dt_1 \in \mathcal{G}p_0(p_0(\dots p_0(D[F(v_1, \dots, v_n)]_{v_1 \dots v_n}, Dt_n), \dots, Dt_3), Dt_2)$$

iff

$$(Dt_1, Dt_2) \in \mathcal{G}p_0(p_0(\dots p_0(D[F(v_1, \dots, v_n)]_{v_1 \dots v_n}, Dt_n), \dots, Dt_3)$$

and so on until obtaining

$$(Dt_1, \dots, Dt_n) \in D[F(v_1, \dots, v_n)]_{v_1 \dots v_n} = \mathcal{I}(F)$$

Lemma 7.10 *$T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A \& B)$ iff $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A)$ and $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(B)$. Also $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\neg A)$ iff it is not the case that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A)$.*

This is an immediate consequence of the definition of $T_{\mathcal{I}\mathcal{A}\mathcal{M}}$.

8 Soundness of T1

In this section we work in a type 1 model \mathcal{M} . We will use Polish notation for the model operations and omit parenthesis when possible. We note first that Lemma 7.9 (for the case of equality) and Lemma 7.10 yields immediately:

Lemma 8.1 *For all models we have propositional tautologies, (MP), (Id) and (L') are sound.*

It is immediate by lemma 7.7 that axiom 6 is sound and the soundness of axiom 7 is obvious. The following is of central importance

Lemma 8.2 (Bealer's lemma) *Let v be free in $[A(v)]_{\bar{x}}$. Then for all \mathcal{M} of type 1 and \mathcal{I} , \mathcal{A} be an interpretation and assignment. Then*

$$D[A(v)]_{\bar{x}} = p_0 D[A(v)]_{\bar{x}v} \mathcal{A}(v)$$

Note that when we write (x_1, \dots, x_n) we include the case $n = 0$ in which case the sequence is taken to be the empty set \emptyset .

Proof. We proceed by induction on the Bealer decomposition of $[A(v)]_{\bar{x}}$.

Let $\mathcal{B}[A(v)]_{\bar{x}} = K\mathcal{B}[B(v)]_{\bar{x}}\mathcal{B}[C(v)]_{\bar{x}}$. Then

$$D[A(v)]_{\bar{x}} = kD[B(v)]_{\bar{x}}D[C(v)]_{\bar{x}} = kp_0 D[B(v)]_{\bar{x}v} \mathcal{A}(v) p_0 D[C(v)]_{\bar{x}v} \mathcal{A}(v)$$

by the induction hypothesis. We use the type 1 condition to show that

$$kp_0 D[B(v)]_{\bar{x}v} \mathcal{A}(v) p_0 D[C(v)]_{\bar{x}v} \mathcal{A}(v) = p_0 D[A(v)]_{\bar{x}v} \mathcal{A}(v)$$

For any \mathcal{H} let $(x_1, \dots, x_n) \in \mathcal{H}kp_0 D[B(v)]_{\bar{x}v} \mathcal{A}(v) p_0 D[C(v)]_{\bar{x}v} \mathcal{A}(v)$. Then this is equivalent to

$$(x_1, \dots, x_n) \in \mathcal{H}p_0 D[B(v)]_{\bar{x}v} \mathcal{A}(v) \text{ and } (x_1, \dots, x_n) \in \mathcal{H}p_0 D[C(v)]_{\bar{x}v} \mathcal{A}(v)$$

which is equivalent to

$$(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}D[B(v)]_{\bar{x}v} \text{ and } (x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}D[C(v)]_{\bar{x}v}$$

The Bealer decomposition of $[A(v)]_{\bar{x}v}$ is $\Sigma K[B(v)]_{\bar{x}'}[C(v)]_{\bar{x}'}$ where Σ is such that $\bar{x}' = \sigma \bar{x}v$ is the normalised permutation of $\bar{x}v$ for $[A(v)]_{\bar{x}v}$. But then the above condition is equivalent to

$$\sigma(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}D[B(v)]_{\bar{x}'} \text{ and } \sigma(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}D[C(v)]_{\bar{x}'}$$

which is equivalent to $\sigma(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}skDB[(v)]_{\bar{x}'}DC[(v)]_{\bar{x}'}$. Let s be the composition of operations corresponding to Σ . Then the previous condition is equivalent to $(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}skDB[(v)]_{\bar{x}'}DC[(v)]_{\bar{x}'} = \mathcal{H}D[A(v)]_{\bar{x}v}$. But this is equivalent to $(x_1, \dots, x_n) \in \mathcal{H}p_0 D[A(v)]_{\bar{x}v} \mathcal{A}(v)$. Hence the conclusion follows from the type 1 condition.

Let $\mathcal{B}[A(v)]_{\bar{x}} = N\mathcal{B}[A'(v)]_{\bar{x}}$ where $A(v) = \neg A'(v)$. We have $D[A(v)]_{\bar{x}} = nD[A'(v)]_{\bar{x}} = np_0 D[A'(v)]_{\bar{x}v} \mathcal{A}(v)$ by the induction hypothesis. We use the type 1 condition to show that

$$np_0 D[A'(v)]_{\bar{x}v} \mathcal{A}(v) = p_0 D[A(v)]_{\bar{x}v} \mathcal{A}(v)$$

Then for any \mathcal{H} ,

$$(x_1, \dots, x_n) \in \mathcal{H}np_0 D[A'(v)]_{\bar{x}v} \mathcal{A}(v) \text{ iff } (x_1, \dots, x_n) \notin \mathcal{H}p_0 D[A'(v)]_{\bar{x}v} \mathcal{A}(v)$$

$$\text{iff } (x_1, \dots, x_n, \mathcal{A}(v)) \notin \mathcal{H}[A'(v)]_{\bar{x}v}$$

Let \bar{x}' be the normalised sequence for $[A'(v)]_{\bar{x}v}$ and σ the associated permutation. Then the above condition holds iff

$$\sigma(x_1, \dots, x_n, \mathcal{A}(v)) \notin \mathcal{H}[A'(v)]_{\bar{x}'} \text{ iff } \sigma(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}n[A'(v)]_{\bar{x}'}$$

The Bealer decomposition of $[A(v)]_{\bar{x}v}$ is $\Sigma N[A'(v)]_{\bar{x}'}$ where σ corresponds to Σ so $D[A(v)]_{\bar{x}v} = snD[A'(v)]_{\bar{x}'}$ where s is the composition of model operation corresponding to Σ . Hence we have $\sigma(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}n[A'(v)]_{\bar{x}'}$ iff $(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}D[A(v)]_{\bar{x}v}$ iff $(x_1, \dots, x_n) \in \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$ and our conclusion follows from the type 1 model condition.

Let $\mathcal{B}[A(v)]_{\bar{x}} = U[A'(v)]_{\bar{x}w}$ where $A(v) = \exists w.A'(v)$. Then $D[A(v)]_{\bar{x}} = uD[A'(v)]_{\bar{x}w} = up_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v)$ by induction. We show that $up_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v) = p_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$. Take a \mathcal{H} . Then

$$(x_1, \dots, x_n) \in \mathcal{H}up_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v) \text{ iff there is } y \text{ such that } (x_1, \dots, x_n, y) \in \mathcal{H}p_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v)$$

$$\text{iff there is a } y \text{ such that } (x_1, \dots, x_n, y, \mathcal{A}(v)) \in \mathcal{H}D[A'(v)]_{\bar{x}wv}$$

As previously let $\bar{x}' = \sigma\bar{x}v$ be the normalized sequence for $[A(v)]_{\bar{x}v}$. The Bealer decomposition of $[A(v)]_{\bar{x}v}$ is $\Sigma U\mathcal{B}[A'(v)]_{\bar{x}'w}$ so $p_0[A(v)]_{\bar{x}v}\mathcal{A}(v) = p_0suD[A'(v)]_{\bar{x}'w}\mathcal{A}(v)$. The above condition is equivalent to

$$\text{there is a } y \text{ such that } (x'_1, \dots, x'_{n+1}, y) \in \mathcal{H}D[A'(v)]_{\bar{x}'w}$$

where $\mathcal{A}(v)$ occupies the position x'_i of v in \bar{x}' . But this is equivalent to

$$(x'_1, \dots, x'_{n+1}) \in \mathcal{H}uD[A'(v)]_{\bar{x}'w}$$

which is equivalent to

$$(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}suD[A'(v)]_{\bar{x}'w} \text{ iff } (x_1, \dots, x_n) \in \mathcal{H}p_0suD[A'(v)]_{\bar{x}'w}\mathcal{A}(v) = \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

and the result follows from the type 1 model condition.

Let $\mathcal{B}[A(v)]_{\bar{x}} = E\mathcal{B}[A(v)]_{\bar{x}'}$ with $\bar{x} = \bar{x}'y$ and y does not occur free in $A(v)$. We have $D[A(v)]_{\bar{x}} = eD[A(v)]_{\bar{x}'} = ep_0[A(v)]_{\bar{x}'v}\mathcal{A}(v)$ by induction. As previously take a \mathcal{H} . Then

$$(x_1, \dots, x_n) \in \mathcal{H}ep_0[A(v)]_{\bar{x}'v}\mathcal{A}(v) \text{ iff } (x_1, \dots, x_{n-1}) \in \mathcal{H}p_0[A(v)]_{\bar{x}'v}\mathcal{A}(v)$$

$$\text{iff } (x_1, \dots, x_{n-1}, \mathcal{A}(v)) \in \mathcal{H}[A(v)]_{\bar{x}'v}$$

The Bealer decomposition of $[A(v)]_{\bar{x}v}$ is $\Sigma\mathcal{B}[A(v)]_{\bar{z}} = \Sigma E\mathcal{B}[A(v)]_{\bar{z}'}$ where $\bar{z} = \sigma\bar{x}v$ is the normalizing permutation which puts v in its proper place and \bar{z}' drops the last element of \bar{z} (not v and the same as x_n). Put $\bar{z}' = \sigma'\bar{x}'v$ and let σ' correspond to composition of model operations s' . Then the above condition is equivalent to

$$(x_1, \dots, x_{n-1}, \mathcal{A}(v)) \in \mathcal{H}s'D[A(v)]_{\sigma'\bar{x}'v} = \mathcal{H}s'D[A(v)]_{\bar{z}'}$$

which is equivalent to

$$(x_1, \dots, x_{n-1}, x_n, \mathcal{A}(v)) \in \mathcal{H}ies'D[A(v)]_{\bar{z}'} = \mathcal{H}seD[A(v)]_{\bar{z}'}$$

which is turn is equivalent to:

$$(x_1, \dots, x_n) \in \mathcal{H}p_0seD[A(v)]_{\bar{x}'}\mathcal{A}(v) = \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

and the result follows as in the other cases.

Let $\mathcal{B}[A(v)]_{\bar{x}} = B\mathcal{B}[A(v)]_{\bar{x}'}$ with B equal to C or I where $\bar{x}' = \sigma_b\bar{x}$ for b equal to c or i . Then $D[A(v)]_{\bar{x}} = bD[A(v)]_{\bar{x}'} = bp_0[A(v)]_{\bar{x}'v}\mathcal{A}(v)$ using the induction hypothesis. We use the type 1 condition to show that

$$bp_0[A(v)]_{\bar{x}'v}\mathcal{A}(v) = Dp_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

For any \mathcal{H} we have

$$(x_1, \dots, x_n) \in \mathcal{H}bp_0[A(v)]_{\bar{x}'v}\mathcal{A}(v) \text{ iff } (x'_1, \dots, x'_n) \in \mathcal{H}p_0[A(v)]_{\bar{x}'v}\mathcal{A}(v)$$

where $(x_1, \dots, x_n) = \sigma_b(x_1, \dots, x_n)$. But

$$(x'_1, \dots, x'_n) \in \mathcal{H}p_0[A(v)]_{\bar{x}'v}\mathcal{A}(v) \text{ iff } (x'_1, \dots, x'_n, \mathcal{A}(v)) \in \mathcal{H}[A(v)]_{\bar{x}'v}$$

$$\text{iff } (x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}[A(v)]_{\bar{x}v} \text{ iff } (x_1, \dots, x_n) \in \mathcal{H}p_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

Hence by the type 1 model condition we get that $bp_0[A(v)]_{\bar{x}'v}\mathcal{A}(v) = Dp_0[A(v)]_{\bar{x}v}\mathcal{A}(v)$.

Now let $\mathcal{B}[A(v)]_{\bar{x}} = R\mathcal{B}[A'(v)]_{\bar{x}w}$. Then $[A(v)]_{\bar{x}}$ has a reflected variable and hence so does $[A(v)]_{\bar{x}v}$ (it is also belongs to category 4 and is a prime reflection term). We distinguish between two cases. Either v is the right-most reflected variable or it is not. Assume that it is not. Let t be the right-most argument in which t occurs. Then the right-most reflected variable is the last element x_n of \bar{x} . We have $D[A(v)]_{\bar{x}} = rp_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v)$ by induction, where $A'(v)$ is obtained from $A(v)$ by replacing the argument t with $t[w/x_n]$. Fix \mathcal{H} . Then

$$(x_1, \dots, x_n) \in \mathcal{H}rp_0D[A'(v)]_{\bar{x}wv}\mathcal{A}(v) \text{ iff } (x_1, \dots, x_n, x_n, \mathcal{A}(v)) \in \mathcal{H}D[A'(v)]_{\bar{x}wv}$$

Now the Bealer decomposition of $[A(v)]_{\bar{x}v}$ in this case will be $\Sigma R\mathcal{B}[A'(v)]_{\bar{x}'w}$. Here $\bar{x}v = \sigma\bar{x}'$. \bar{x}' moves v to its proper place so that \bar{x}' is the normalized permutation of $\bar{x}v$ except that x_n is placed at the end, being the reflected variable with right-most occurrence. We have $D[A(v)]_{\bar{x}v} = srD[A'(v)]_{\bar{x}'w}$. Then the condition above is equivalent to:

$$(x'_1, \dots, x'_{n+1}, x'_{n+1}) \in \mathcal{H}D[A'(v)]_{\bar{x}'w}$$

with $x'_{n+1} = x_n$. But this is equivalent to:

$$(x'_1, \dots, x'_{n+1}) \in \mathcal{H}rD[A'(v)]_{\bar{x}'w}$$

which in turn is equivalent to

$$(x_1, \dots, x_n, \mathcal{A}(v)) \in \mathcal{H}srD[A'(v)]_{\bar{x}'w}$$

which is finally equivalent to

$$(x_1, \dots, x_n) \in \mathcal{H}p_0srD[A'(v)]_{\bar{x}'w}\mathcal{A}(v) = \mathcal{H}p_0D[A(v)]_{\bar{x}v}\mathcal{A}(v)$$

and the result follows as in the other cases. Now for the second case: v is a reflected variable with the right-most occurrence. Let the right-most argument in which it occurs be t . The Bealer decomposition of $[A(v)]_{\bar{x}v}$ in this case will be $\Sigma R\mathcal{B}[A'(v)]_{\bar{x}'vw}$ where $A'(v)$ is obtained from $A(v)$ by replacing t with $t[w/v]$. Here $\bar{x}v = \sigma\bar{x}'v$. \bar{x}' moves x_n back to its proper place so that \bar{x}' is a normalized sequence. Now either in $[A'(v)]_{\bar{x}'vw}$ v is still the right-most reflected variable or else it is x_n . If v it is still the right-most reflected variable we continue with the

Bealer decomposition until we arrive at term in which the right-most reflected variable is x_n . The decomposition will be of the form:

$$[A(v)]_{\bar{x}v} = \Sigma_1 R \Sigma_2 R \dots \Sigma_k R \mathcal{B}[A^{(k)}(v)]_{\bar{z}w_k}$$

where \bar{z} (normalized except for w_k being placed at the end) contains the elements of \bar{x}, v and new variables w_1, \dots, w_{k-1} . Here $[A^{(k)}(v)]_{\bar{z}w_k}$ is now as the previous case (modulo a permutation).

It is easy to see that

$$(x_1, \dots, x_n) \in \mathcal{H}p_0 s_1 r s_2 r \dots s_k R D[A^{(k)}(v)]_{\bar{z}w_k} \mathcal{A}(v)$$

iff

$$(x'_1, \dots, x'_{n+k+1}) \in \mathcal{H}D[A^{(k)}(v)]_{\bar{z}w_k}$$

Here $(x'_1, \dots, x'_{n+k+1})$ follows the sequence of $\bar{z}w_k$, having x_i for x_i (here we are abusing notation) but has $x'_k = \mathcal{A}(v)$ whenever x'_k is a w_i or v .

Now $D[A^{(k)}(v)]_{\bar{z}w_k} = sD[A^{(k)}(v)]_{\bar{u}x_n} = srD[A_0^{(k)}(v)]_{\bar{u}x_n w}$ where \bar{u} is the normalized permutation of $\bar{z}w_k$ but with x_n omitted. Hence

$$(*) \quad (x'_1, \dots, x'_{n+k+1}) \in \mathcal{H}D[A^{(k)}(v)]_{\bar{z}w_k} \text{ iff } (x''_1, \dots, x''_{n+k}, x_n, x_n) \in \mathcal{H}D[A_0^{(k)}(v)]_{\bar{u}x_n w}$$

where the x''_i follows the normalized sequence but with x_n omitted and $A_0^{(k)}(v)$ has the right-most argument t in which x_n occurs replaced with $t[w/x_n]$. Now we have by induction,

$$(x_1, \dots, x_n) \in \mathcal{H}D[A(v)]_{\bar{x}} \text{ iff } (x_1, \dots, x_n) \in \mathcal{H}rp_0[A_1(v)]_{\bar{x}wv} \mathcal{A}(v)$$

$$\text{iff } (x_1, \dots, x_n, x_n, \mathcal{A}(v)) \in \mathcal{H}D[A_1(v)]_{\bar{x}wv}$$

where A_1 has the right-most argument t in which x_n appears replaced by $t[w/x_n]$. Now $[A_1(v)]_{\bar{x}wv}$ has v reflected in the same arguments and positions as $[A(v)]_{\bar{x}v}$. Hence

$$[A_1(v)]_{\bar{x}wv} = \Sigma'_1 R \Sigma'_2 R \dots \Sigma'_k R \mathcal{B}[A_1^{(k)}(v)]_{\bar{q}w_k}$$

Hence

$$(x_1, \dots, x_n, x_n, \mathcal{A}(v)) \in \mathcal{H}D[A_1(v)]_{\bar{x}wv} \text{ iff } (q_1, \dots, q_{n+k+2}) \in \mathcal{H}D[A_1^{(k)}(v)]_{\bar{q}w_k}$$

Here \bar{q} is the normalized permutation of $\bar{x}wv$ except that w_k is moved to the end. It has new variables w_1, \dots, w_{k-1} . (q_1, \dots, q_{n+k+2}) follows this sequence and for v or w_i it has \mathcal{A} and for w it has x_n . It is clear that $A_1^{(k)}(v)$ is the same as $A_0^{(k)}(v)$. Examining (*) and noticing that applying a permutation we have

$$(x''_1, \dots, x''_{n+k}, x_n, x_n) \in \mathcal{H}D[A_0^{(k)}(v)]_{\bar{u}x_n w} \text{ iff } (q_1, \dots, q_{n+k+2}) \in \mathcal{H}D[A_0^{(k)}(v)]_{\bar{q}w_k}$$

and the result follows.

Finally, consider the case in which $\mathcal{B}[A(v)]_{\bar{x}} = P_n \mathcal{B}[A'(v)]_{\bar{u}w} t_{\bar{y}}$. Here $t_{\bar{y}}$ means that if $t = [B]_{\bar{z}}$ then $t_{\bar{y}} = [B]_{\bar{z}\bar{y}}$ or else t is a variable and \bar{y} is the empty sequence. Let the argument in $A(v)$ which w replaced be t . Here $\bar{x} = \bar{x}' \bar{x}'' \bar{y}$ is normalized except that the sequence \bar{y} of variables in \bar{x} which are free in t are placed at the end. All arguments before t are variables in \bar{x} . So v may occur in t and in arguments after t and v may be reflected. Hence we will have (if v is reflected)

$$[A(v)]_{\bar{x}'\bar{x}''\bar{y}v} = \Sigma_1 R \Sigma R \dots R \Sigma_k \Sigma [A^{(k)}(v)]_{\bar{x}'\bar{p}\bar{y}'} = \overline{\Sigma R} \Sigma [A^{(k)}(v)]_{\bar{x}'\bar{p}\bar{y}'}$$

Here $\bar{x}'\bar{q}$ is the normalized permutation of $\bar{x}v$ and the k new variables w_k but with the sequence \bar{y}' of variables of \bar{q} which occur free in t placed at the end. \bar{y}' is either equal to \bar{y} or contains in addition v . Notice how \bar{x}' is not affected by the permutations. For clarity, we wrote the last permutation as a composition. If v is not reflected then we have only Σ which puts v in its proper place (which is either in \bar{y} or in \bar{p}).

We have that

$$(x'_1, \dots, x'_a, x''_1, \dots, x''_b, y_1, \dots, y_c, v) \in \mathcal{H}\overline{s}r s D[A^{(k)}(v)]_{\bar{x}'\bar{p}\bar{y}'}$$

iff

$$(x'_1, \dots, x'_a, p_1, \dots, p_{b+k}, y'_1, \dots, y'_{c+j}) \in \mathcal{H}D[A^{(k)}(v)]_{\bar{x}'\bar{p}\bar{y}'}$$

where j is either 0 and v is in \bar{p} or else is 1 and v is in \bar{y}' . Here the (p_1, \dots, p_{b+k}) is permutation of the x''_i and w_i (and possibly v) corresponding to the normalized sequence of these variables and which has v for w_i .

W.l.o.g let v be in \bar{p} and t . Fix a \mathcal{H} . Then

$$(x'_1, \dots, x'_a, x''_1, \dots, x''_b, y_1, \dots, y_c) \in \mathcal{H}D[A(v)]_{\bar{x}'\bar{x}''\bar{y}} = \mathcal{H}P_n D[A'_1(v)]_{\bar{x}'\bar{x}''w} D t_{\bar{y}} \text{ iff } \\ (x'_1, \dots, x'_a, x''_1, \dots, x''_b, p^c(D t_{\bar{y}}, y_1, \dots, y_1)) \in \mathcal{H}D[A'_1(v)]_{\bar{x}\bar{x}'w} = \mathcal{H}p_0 D[A'_1(v)]_{\bar{x}\bar{x}''wv} \mathcal{A}(v)$$

by induction. This is equivalent to

$$(x'_1, \dots, x'_a, x''_1, \dots, x''_b, p^c(D t_{\bar{y}}, y_1, \dots, y_1), \mathcal{A}(v))) \in \mathcal{H}D[A'_1(v)]_{\bar{x}'\bar{x}''wv}$$

Let the Bealer decomposition of $[A'_1(v)]_{\bar{x}'\bar{x}''wv}$ be

$$\Sigma'_1 R \Sigma'_2 R \dots R \Sigma'_{k-1} \Sigma' [A'^{(k-1)}(v)]_{\bar{x}'\bar{q}} = \overline{\Sigma' R} \Sigma' [A'^{(k)}(v)]_{\bar{x}'\bar{q}}$$

where \bar{q} consists of variables of \bar{x}'' , w , v and k new variables w_k .

The previous condition is equivalent to

$$(x'_1, \dots, x'_a, x''_1, \dots, x''_b, p^c(D t_{\bar{y}}, y_1, \dots, y_1), \mathcal{A}(v))) \in \mathcal{H}\overline{s'}\overline{R}s' D[A'^{(k-1)}(v)]_{\bar{x}'\bar{q}}$$

which is equivalent to

$$(**) \quad (x'_1, \dots, x'_a, r_1, \dots, r_s) \in \mathcal{H}D[A'^{(k-1)}(v)]_{\bar{x}'\bar{q}}$$

where (r_1, \dots, r_s) corresponds to the sequence \bar{q} with v and w_k corresponding to $\mathcal{A}(v)$ and w corresponding to $p^c(D t_{\bar{y}}, y_1, \dots, y_1)$.

Now consider

$$(x'_1, \dots, x'_a, x''_1, \dots, x''_b, y_1, \dots, y_c) \in \mathcal{H}p_0 D[A(v)]_{\bar{x}\bar{x}'\bar{y}v} \mathcal{A}(v)$$

this is equivalent to

$$(x'_1, \dots, x'_a, x''_1, \dots, x''_b, y_1, \dots, y_c, \mathcal{A}(v)) \in \mathcal{H}D[A(v)]_{\bar{x}\bar{x}'\bar{y}v} = \mathcal{H}\overline{s}r s D[A^{(k)}(v)]_{\bar{x}'\bar{p}\bar{y}'}$$

where $\bar{y}' = \bar{y}''v\bar{y}'''$. This in turn is equivalent to

$$(x'_1, \dots, x'_a, x''_1, \dots, x''_b, y_1, \dots, y_c, \mathcal{A}(v)) \in \mathcal{H}\overline{s}r s P_{n+1} D[A^{(k)'}(v)]_{\bar{x}'\bar{p}w\bar{t}_{\bar{y}'}}$$

that is,

$$(x'_1, \dots, x'_a, e_1, \dots, e_{b+k}, y'_1, \dots, y'_{c+1}) \in \mathcal{H}P_{n+1}D[A^{(k)'}(v)]_{\bar{x}'\bar{p}wt_{\bar{y}'}}$$

where (e_1, \dots, e_{b+k+1}) corresponds to the sequence \bar{p} and has $\mathcal{A}(v)$ corresponding to w_k and (y'_1, \dots, y'_{c+1}) corresponds to $\bar{y}' = y''vy'''$ with $\mathcal{A}(v)$ corresponding to v . This in turn is the same as

$$(x'_1, \dots, x'_a, e_1, \dots, e_{b+k}, p_0^{c+1}(Dt_{\bar{y}'}, y'_1, \dots, y'_{c+1})) \in \mathcal{H}D[A^{(k)'}(v)]_{\bar{x}'\bar{p}w}$$

It is easy to check that by induction we have

$$p_0^{c+1}(Dt_{\bar{y}'}, y'_1, \dots, y'_{c+1})) = p_0 p_0^c(Dt_{\bar{y}'}, y_1, \dots, y_c))\mathcal{A}(v) = p_0^c(Dt_{\bar{y}'}, y_1, \dots, y_c)$$

Inspecting (**), noticing that $[A^{(k)'}(v)]_{\bar{x}'\bar{p}w} = [A^{(k)}(v)]_{\bar{x}'\bar{p}w}$ (modulo renaming bound variables) and applying a permutation we get as desired.

Lemma 8.3 $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\exists v.A)$ iff there is an assignment \mathcal{A}' like \mathcal{A} except perhaps for what it assigns to v and such that $T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(A(v))$.

Proof. Assume w.l.o.g that v occurs free in A . We have $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\exists v.A)$ iff $\mathcal{G}D([\exists v.A]) = T$ iff $\mathcal{G}uD[A]_v = T$. The last condition is equivalent to the existence of a $x \in \mathcal{D}$ such that

$$x \in \mathcal{G}D[A(v)]_v$$

Let \mathcal{A}' be like \mathcal{A} but with $\mathcal{A}'(v) = x$. We must show that $\mathcal{G}D_{\mathcal{A}'}[A(v)] = T$. Notice that $D_{\mathcal{A}'}[A(v)]_v = D[A(v)]_v$. By the proof of Bealer's lemma (note that we are not using the type 1 condition) we have that:

$$\mathcal{G}D_{\mathcal{A}'}[A(v)] = T \text{ iff } \mathcal{G}p_0D[A(v)]_v\mathcal{A}'(v) = T$$

Hence we must show that $\mathcal{G}p_0D[A(v)]_v\mathcal{A}'(v) = T$. But this is equivalent to $\mathcal{A}'(v) \in \mathcal{G}D[A(v)]_v$ and the result follows since $\mathcal{A}'(v) = x$. The other direction is similar.

Hence if $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\forall v.A(v))$ then for any assignment \mathcal{A}' which only differs at most on v we have that $T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(A(v))$. Hence lemma 8.3 yields:

Lemma 8.4 For any models we have $(Inst) \models \forall v.A(v) \rightarrow A(t)$ where t is free for v in A .

Likewise we have

Lemma 8.5 For any models we have $(QImp) \models \forall v.(A \rightarrow B(v)) \rightarrow (A \rightarrow \forall v.B(v))$ where v does not occur free in A .

Proof. Assume $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\forall v.A \rightarrow B(v))$. Then for all assignment \mathcal{A}' differing from \mathcal{A} at most on v we have that

$$(*) \quad T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(A \rightarrow B(v))$$

We must show that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A \rightarrow \forall v.B(v))$. Assume then that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A)$. We must show that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A \rightarrow \forall v.B(v))$. Take any assignment \mathcal{A}' differing from \mathcal{A} at most on v . Since v does not occur in A we have that $T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(A)$. Hence by (*) we get that $T_{\mathcal{I}\mathcal{A}'\mathcal{M}}(\forall v.A(v))$. Since this holds for any \mathcal{A}' we get the conclusion by lemma 8.3.

Similarly we get the soundness of the (Gen) rule:

Lemma 8.6 For any models if $\models A(v)$ then $\models \forall v.A(v)$.

Lemma 8.7 For all \mathcal{M} of type 1 and \mathcal{I} , \mathcal{A} and terms t, t' we have

$$D([t = t]) = D([t' = t'])$$

Proof. Suppose that

$$D([t = t]) \neq D([t' = t'])$$

Then the type 1 condition implies that there is a \mathcal{H} such that

$$\mathcal{H}D([t = t]) \neq \mathcal{H}D([t' = t'])$$

But

$$[t = t] = P_0(P_0([v = w]_{vw}, t), t)$$

Hence applying the definition of \mathcal{H} yields

$$\mathcal{H}D([t = t]) = T \leftrightarrow D(t) = D(t)$$

hence we have $\mathcal{H}D([t = t]) = T$ and also analogously $\mathcal{H}D([t' = t']) = T$ and so we obtain a contradiction.

Lemma 8.8 ([1]A8*(a)) For type 1 models we have $(B1) \models \Box(A \leftrightarrow B) \leftrightarrow [A] = [B]$.

Proof. We must show that $\mathcal{G}D_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box(A \leftrightarrow B) \leftrightarrow [A] = [B]) = T$, which is equivalent to showing that

$$\mathcal{G}D_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box(A \leftrightarrow B)) = T \text{ iff } \mathcal{G}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A] = [B]) = T$$

that is,

$$\mathcal{G}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A \leftrightarrow B] = [[A = B] = [A = B]]) = T \text{ iff } \mathcal{G}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A] = [B]) = T$$

which is equivalent to

$$D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A \leftrightarrow B]) = D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A = B] = [A = B]) \text{ iff } D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = D_{\mathcal{I}\mathcal{A}\mathcal{M}}([B])$$

Lemma 8.9 ([1] A8*(b)) For type 1 models we have $(B2) \models \forall v.[A(v)]_\alpha = [B(v)]_\alpha \leftrightarrow [A(v)]_{\alpha v} = [B(v)]_{\alpha v}$.

Proof. We have

$$D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_\alpha = p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v}, \mathcal{A}(v))$$

and

$$D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_\alpha = p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}, \mathcal{A}(v))$$

Consider all assignments \mathcal{A}' like \mathcal{A} except for the assignment on v . Then $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_{\alpha v}$ and $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_{\alpha v}$. So if $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$ then for any \mathcal{A}' we have $D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_\alpha = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_\alpha$ and one direction follows.

We now need to show that if for all such \mathcal{A}' we have

$$D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_\alpha = D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_\alpha$$

then $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} = D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$. But if $D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v} \neq D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$ then by the type 1 condition there would be a \mathcal{H} and a (x_1, \dots, x_m) such that $(x_1, \dots, x_m) \in \mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v}$ but $(x_1, \dots, x_m) \notin \mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}$. But taking $\mathcal{A}'(v) = x_n$ this means that

$$(x_1, \dots, x_{m-1}) \in \mathcal{H}p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A(v)]_{\alpha v}, \mathcal{A}'(v)) = \mathcal{H}D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[A(v)]_\alpha$$

but

$$(x_1, \dots, x_{m-1}) \notin \mathcal{H}p_0(D_{\mathcal{I}\mathcal{A}\mathcal{M}}[B(v)]_{\alpha v}, \mathcal{A}'(v)) = \mathcal{H}D_{\mathcal{I}\mathcal{A}'\mathcal{M}}[B(v)]_{\alpha}$$

and we obtain a contradiction.

Lemma 8.10 ([1] A11*) *In the same conditions we have $(S5') \models v_i \neq v_j \rightarrow \Box v_i \neq v_j$.*

Proof. We must show that if $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(v_i \neq v_j)$ then $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box v_i \neq v_j)$.

But if $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(v_i \neq v_j)$ then $D_{\mathcal{I}\mathcal{A}\mathcal{M}}(v_i) \neq D_{\mathcal{I}\mathcal{A}\mathcal{M}}(v_k)$. But this is equivalent to $\mathcal{A}(v_i) \neq \mathcal{A}(v_k)$. This means that for all $\mathcal{H} \in H$ we have $\mathcal{H}([v_i \neq v_j]) = T$ and the conclusion follows.

Lemma 8.11 ([1] A9) *In the same conditions we have $(T) \models \Box A \rightarrow A$.*

Proof. If $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box A)$ then $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$ and so in particular $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(A)$

Lemma 8.12 ([1] A10) *In the same conditions we have $(K) \models \Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)$.*

Proof. Assume $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box(A \rightarrow B))$. Then $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}[A \rightarrow B] = T$ which means that if $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$ then $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([B]) = T$. To show that $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box A \rightarrow \Box B)$ consider any $\mathcal{K} \in H$. Assume that $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$. Then $\mathcal{K}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([A]) = T$ and hence $\mathcal{K}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([B]) = T$. Hence we have $\forall \mathcal{H} \in H$ we have $\mathcal{H}D_{\mathcal{I}\mathcal{A}\mathcal{M}}([B]) = T$ and thus $T_{\mathcal{I}\mathcal{A}\mathcal{M}}(\Box A \rightarrow \Box B)$.

9 Completeness of T1

In this section we show completeness:

Lemma 9.1 *If a sentence A is T1-valid then $\vdash A$.*

It is enough to show that:

Lemma 9.2 *A set Γ of sentences of T1 is consistent iff Γ has a model.*

By soundness an inconsistent set of formulas cannot have a model so we only need prove one implication. This lemma is enough to prove completeness for is a formula A is T1-valid then $\neg A$ cannot have a model, otherwise there would be a model for which both A and $\neg A$ held. Since $\neg A$ does not have a model by the lemma above $\neg A$ is inconsistent and hence we have $\neg A \vdash \neg x = x$, that is $\vdash \neg A \rightarrow \neg x = x$ or $\vdash x = x \rightarrow A$ from which it follows that $\vdash A$.

Given a consistent set Γ of sentences we form an extension $L^{\omega*}$ of L^{ω} consisting of countably infinite constants and n -ary predicate symbols F_i (including 0-ary propositional constants) and consider any enumeration S_1, S_2, \dots of all sentences. Consider also any enumeration of all closed terms $[B]_{\bar{x}}$. We apply the following process to the sequence S_1, S_2, \dots . Consider the first closed term $[B]_{\bar{x}}$ and chose a F and let $[F(x_1, \dots, x_n)]_{\bar{x}} = [B]_{\bar{x}}$ be S_k . If F does not occur in a sentence before S_k or in B or in Γ then we proceed to the next closed term. Otherwise chose any F' which satisfies this condition and let S_m be $[F'(x_1, \dots, x_n)]_{\bar{x}} = [B]_{\bar{x}}$. Then switch the positions of S_m and S_k in the enumeration S_1, S_2, \dots and proceed to the next closed term. Any finite initial segments of the resulting transformations of the resulting sequence S'_1, S'_2, \dots will stabilise after a finite number of iterations (we do not move an element of a sequence twice).

From now on we assume that we have a consistent set Γ and that we have an enumeration S_1, S_2, \dots of sentences of L^{ω} satisfying the property that for any closed term t there is a predicate symbol F and a S_k of the form $[F(x_1, \dots, x_n)]_{\bar{x}} = t$ such that F does not occur in Γ , t or in any S_j before S_k .

We construct an array $\Delta_{ij}(i, j \geq 1)$ of sets of sentences is the following form:

- $\Delta_{11} = \Gamma$.
- $\Delta_{ij} = \Delta_{ij-1} \cup C_{ij} \cup \{S_j\}$ if $j > 1$ and $\{S_j\} \cup \Delta_{ij-1} \cup E_{ij}$ is consistent.
- $\Delta_{ij} = \Delta_{ij-1} \cup C_{ij}$ if $j > 1$ and $\{S_j\} \cup \Delta_{ij-1} \cup E_{ij}$ is inconsistent.
- $\Delta_{i1} = \{\exists \bar{x}. \neg A \leftrightarrow B\}$ if S_{i-1} is $[A]_{\bar{x}} \neq [B]_{\bar{x}}$ and $S_{i-1} \in \Delta_{i-1i-1}$
- $\Delta_{i1} = \Delta_{i-1i-1}$ otherwise.

here

$$E_{ij} = \{[A]_{\bar{x}} = [B]_{\bar{y}} : \exists k, m. k \leq i, m \leq j, (k, m) \neq (i, j) \& \Delta_{km} \vdash [A]_{\bar{x}} = [B]_{\bar{y}}\}$$

for $j > 1$ and

$$E_{ij} = \{[A]_{\bar{x}} = [B]_{\bar{y}} : \exists k, m. k \leq i-1, m \leq i-1 \& \Delta_{km} \vdash [A]_{\bar{x}} = [B]_{\bar{y}}\}$$

for $j = 1$. Consider the set (in the order of S_1, S_2, \dots) of all existential sentences $\exists x.A(x), \exists x.A'(x), \dots$ in Δ_{ij-1} which occur before S_j . We proceed as follows. We start by picking a constant a_1 such that $A(a_1)$ is the first instantiation occurring after S_j such that a_i does not occur before S_j or in Δ_{ij-1} (or Δ_{i-1i-1} for $j = 1$). We then proceed to chose an constant a_2 such that $A'(a_2)$ is the first instantiation after $A(a_1)$ such that a_2 does not occur before or in $A(a_1)$ and so forth. C_{ij} consists of the set of such instantiations $\{A(a_1), A'(a_2), \dots\}$.

Lemma 9.3 $\Delta_{ij} \cup E_{ij}$ is consistent.

Proof. By induction on the construction of Δ_{ij} . The case Δ_{11} is trivial since Γ is consistent. Let $j \geq 1$ and assume by induction that $\Delta_{ij} \cup E_{ij}$ is consistent.

Suppose that $\Delta_{ij+1} \cup E_{ij+1}$ were inconsistent. Then $\Delta_{ij+1} \cup E_{ij+1} \vdash \neg v = v$ which means that for some finite set of formulas F_1, \dots, F_k of $\Delta_{ij+1} \cup E_{ij+1}$ we have

$$(*) \quad \vdash (F_1 \& F_2 \& \dots \& F_k) \rightarrow \neg v = v$$

Suppose that $\Delta_{ij+1} = \Delta_{ij} \cup C_{ij+1}$. If all the F_i 's belong Δ_{ij} then this contradicts the consistency of $\Delta_{ij} \cup E_{ij}$. Hence there is at least one F_i belonging to $C_{ij+1} \cup E_{ij+1}$.

If F_i is in C_{ij+1} then it is a $C(a)$ where $\exists v.C(v)$ is in Δ_{ij} . We may reorganise (*) into the form

$$\vdash C_1(a_1) \rightarrow (C_2(a_2) \rightarrow \dots \rightarrow (F_k \rightarrow \neg v = v) \dots$$

so that each instantiation constant a_i in the expression does not occur further ahead. Since a_1 does not occur further ahead we may substitute it for a fresh variable and then apply the \exists -introduction rule

$$\vdash \exists x.C_1(x) \rightarrow (C_2(a_2) \rightarrow \dots \rightarrow (F_k \rightarrow \neg v = v) \dots$$

and so on until we have replaced all formulas in C_{ij+1} by formulas in Δ_{ij} .

If an F_i belongs to E_{ij+1} then it is an equality $t = s$ such that we have $\Delta_{km} \vdash t = s$ for $k \leq i, m \leq j+1, (k, m) \neq (i, j+1)$, that is $\vdash (G_1 \& \dots \& G_n) \rightarrow t = s$ for $G_i \in \Delta_{km}$. We replace $t = s$ by $G_1 \& \dots \& G_n$. Thus we may eliminate all such F_i and we obtain a contradiction.

Now suppose that $\Delta_{ij+1} = \Delta_{ij} \cup C_{ij+1} \cup \{S_{j+1}\}$.

Assume that $\Delta_{ij+1} \cup E_{ij+1}$ is inconsistent. Then $\Delta_{ij} \cup C_{ij+1} \cup \{S_{j+1}\} \cup E_{ij+1}$ is inconsistent and using the previous argument we find $F_1 \dots F_k$ in $\Delta_{ij} \cup \{S_{j+1}\}$ such that

$$\vdash (F_1 \ \& \ F_2 \ \& \ \dots \ \& \ F_k) \rightarrow \neg v = v$$

which contradicts the way Δ_{ij+1} is formed in this case.

Finally we must consider the case of Δ_{i1} . We assume by induction that $\Delta_{i-1i-1} \cup E_{i-1i-1}$ is consistent. If $\Delta_{i1} = \Delta_{i-1i-1}$ then assume that $\Delta_{i-1i-1} \cup E_{1i}$ were inconsistent and we have

$$\vdash (F_1 \ \& \ F_2 \ \& \ \dots \ \& \ F_k) \rightarrow \neg v = v$$

where the F_i are in $\Delta_{i-1i-1} \cup E_{1i}$. Now by definition $E_{1i} = E_{i-1i-1} \cup \{[A]_{\bar{x}} = [B]_{\bar{y}} : \Delta_{i-1i-1} \vdash [A]_{\bar{x}} = [B]_{\bar{y}}\}$ so an argument similar to the one above shows that we obtain a contradiction to the consistency of $\Delta_{i-1i-1} \cup E_{i-1i-1}$.

If $\Delta_{i1} = \{\exists \bar{x}. \neg A \leftrightarrow B\}$ then $[A]_{\bar{x}} \neq [B]_{\bar{x}} \in \Delta_{i-1i-1}$. Assume $\{\exists \bar{x}. \neg A \leftrightarrow B\} \cup E_{i1}$ were inconsistent. Then there are equality sentences E_1, \dots, E_k in E_{i1} such that $\vdash E_1 \rightarrow \dots \rightarrow E_k \rightarrow \forall \bar{x}. A \leftrightarrow B$. We then apply necessitation, (T) and (B) to obtain $E_{i1} \vdash [A]_{\bar{x}} = [B]_{\bar{x}}$, a contradiction. ■

We now define $\Delta^k = \bigcup_j \Delta_{kj}$ and consider the set \mathcal{A} consisting of all such Δ^k for $k \geq 1$.

Definition 9.4 Let L_ω^* be an extension of L_ω . A set \mathcal{A} of sets of sentences in L_ω^* is *T1-perfect* if

- Every set in \mathcal{A} is maximal, consistent and ω -complete.
- If an identity sentence $t = t'$ is in one of the sets of \mathcal{A} it is in all of them.
- If a sentence $[A]_{\bar{x}} = [B]_{\bar{x}}$ belongs to one set of \mathcal{A} there is a set in \mathcal{A} which contains the sentence $\exists \bar{x}. \sim A \leftrightarrow B$.
- For every closed term $[B]_{x_1 \dots x_n}$ there is an atomic predicate F such that $[B]_{x_1 \dots x_n} = [F(x_1, \dots, x_n)]_{x_1 \dots x_n}$ belongs to a set in \mathcal{A} .

Lemma 9.5 \mathcal{A} is *T1-perfect*.

Proof. Let $\Delta^k \in \mathcal{A}$. We show first that Δ^k is maximal, consistent and ω -complete. It is obviously consistent because each Δ_{kj} is consistent by the above lemma. To show that it is maximal we need to show that for any sentence S_i either S_i or $\neg S_i = S_j$ is in Δ^k . Suppose then that neither formula was in Δ^k . Then we must have that $S_i \cup \Delta_{kj-1} \cup E_{kj}$ and $S_j \cup \Delta_{kj-1} \cup E_{kj}$ are inconsistent. Suppose w.l.o.g. that $i < j$. Then since $\Delta_{ki} \subseteq \Delta_{kj}$ we get a contradiction. For if both a sentence and it's negation are inconsistent with a set of sentences, that set is inconsistent.

To show that Δ^k is ω -complete we need to show that if $\Delta^k \vdash \exists v. F(v)$ then $\Delta^k \vdash F(t)$ for some term t . If $\Delta^k \vdash \exists v. F(v)$ then by maximality $\exists v. F(v) \in \Delta^k$ and let it have been introduced in stage Δ_{ki} . Then in a subsequent stage j , by construction it is clear that C_{ij} and hence Δ^k will contain a formula of the form $F(c)$ for some constant c .

We now show that if an equality $t = s$ is in one $\Delta^k \in \mathcal{A}$ then it is in them all. Let then $t = s$ be in a certain Δ^k and consider any other Δ^j . If Δ^j is in a previous row then obviously we cannot have $\neg t = s \in \Delta^j$ because $t = s$ must belong to a certain Δ_{km} and $\Delta_{km} \cup E_{km}$ is consistent by the previous lemma. A similar argument show that $\neg t = s$ cannot belong to a Δ^j which is on a subsequent row.

We now show that if $[A]_{\bar{x}} \neq [B]_{\bar{x}}$ is in a Δ^k then there is a Δ^j containing $\exists \bar{x}. \neg A \leftrightarrow B$. Suppose $[A]_{\bar{x}} \neq [B]_{\bar{x}}$ is in Δ_{ki} (minimal i for that condition) which is under or on the diagonal

($i \leq k$). Then the result follows directly from the construction of the transition between rows. Suppose then that it is above the diagonal. Notice that in all subsequent Δ_{ki} 's we cannot have $\Delta_{ki} \vdash [A]_{\bar{x}} = [B]_{\bar{x}}$ and we just "slide" down the column until reaching below the diagonal and get as desired.

Finally we show that for any closed term $[G]_{\bar{x}}$ there is a primitive predicate F such that $[G]_{\bar{x}} = [F(x_1, \dots, x_k)]_{x_1, \dots, x_k}$. We can choose such a F (let $[G]_{\bar{x}} = [F(x_1, \dots, x_k)]_{x_1, \dots, x_k}$ be S_i) so that F does not occur in any of the previous S_j , in G or in Γ . Hence in the stage where S_i may be introduced this formula is consistent with the previous Δ_{kj} in the same row (including for any previous column) and thus belongs to Δ^k .

Definition 9.6 Consider a T1-perfect set \mathcal{A} and a $\Delta \in \mathcal{A}$. We define an order $<$ on predicate symbols (such that the first element is $=$) and variables. We define the *canonical model* as follows. Consider the set D_Δ consisting of those symbols F such that there are no $G < F$ for which $[F(x_1, \dots, x_n)]_{x_1 \dots x_n} = [G(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$ and those constants a such that there is no $[F(x_1, \dots, x_n)]_{x_1 \dots x_n} = a \in \Delta$ and there are no constants b such that $b < a$ and $a = b \in \Delta$. We let D_Δ^i divide the symbols according to arity for $i \geq 0$, the case $i = -1$ corresponding to constants. The distinguished equality element is simply $=$. The extension functions $\mathcal{H} \in H$ are determined by the Δ 's in \mathcal{A} and the actual extension function by Δ . We have that $\mathcal{H}_\Delta(F)$ consists of all (F_1, \dots, F_n) such that $[F_i(v_1, \dots, v_n)]_{v_1 \dots v_n} = t_i \in \Delta$ or F_i is a constant c_i and $F(t_1, \dots, t_n) \in \Delta$. We now define the eight model operations.

$$k(F, G) = H \text{ iff } [H(v_1, \dots, v_k, u_1, \dots, u_m)]_{v_1 \dots v_k u_1 \dots u_m} = [F'(v_1, \dots, v_k) \& G'(u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} \in \Delta$$

$$\text{where } [F(v_1, \dots, v_k, u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} = [F'(v_1, \dots, v_k)]_{v_1 \dots v_k u_1 \dots u_k} \in \Delta \text{ and } [G(v_1, \dots, v_k, u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} = [G'(u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} \in \Delta$$

$$n(F) = H \text{ iff } [H(v_1, \dots, v_k)]_{v_1 \dots v_k} = [\sim F(v_1, \dots, v_k)]_{v_1 \dots v_k} \in \Delta$$

$$u(F) = H \text{ iff } [H(v_1, \dots, v_k)]_{v_1 \dots v_k} = [\exists v. F(v_1, \dots, v_k, v)]_{v_1 \dots v_k} \in \Delta$$

$$e(F) = H \text{ iff } [H(v_1, \dots, v_k)]_{v_1 \dots v_k} = [F(v_1, \dots, v_{k-1})]_{v_1 \dots v_k} \in \Delta$$

$$i(F) = H \text{ iff } [H(v_1, \dots, v_k)]_{v_1 \dots v_k} = [F(v_1, \dots, v_k, v_{k-1})]_{v_1 \dots v_k} \in \Delta$$

$$c(F) = H \text{ iff } [H(v_1, \dots, v_k)]_{v_1 \dots v_k} = [F(v_k, v_1, v_2, \dots, v_{k-1})]_{v_1 \dots v_k} \in \Delta$$

$$r(F) = H \text{ iff } [H(v_1, \dots, v_k)]_{v_1 \dots v_k} = [F(v_1, \dots, v_k, v_k)]_{v_1 \dots v_k} \in \Delta$$

$$p_i(F, G) = H \text{ iff } [H(v_1, \dots, v_k, w_1, \dots, w_i)]_{v_1 \dots v_k w_1 \dots w_i} = [F(v_1, \dots, v_k, [G(x_1, \dots, x_j, w_1, \dots, w_i)]_{x_1 \dots x_j})]_{v_1 \dots v_k w_1 \dots w_i} \in \Delta$$

Note how the condition on $<$ guarantees that these operations are well-defined. Finally there is a *canonical interpretation* \mathcal{I}_Δ which associates to a F the G in D_Δ such that $[G(x_1, \dots, x_n)]_{x_1 \dots x_n} = [F(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$ and to a constant c the constant $d \in D_\Delta$ such that $c = d \in \Delta$.

Lemma 9.7 *The canonical model defined above with the canonical interpretation is a T1-model and a model for Δ .*

Proof. The conditions of the extension functions are verified as follows.

Suppose that $(F_1, \dots, F_k, G_1, \dots, G_m) \in \mathcal{H}_\Delta H$ for $H = k(F, G)$. Then we have closed terms t_i, t'_i such that $F_i[x_1 w, \dots, x_{s_i}]_{x_1 \dots x_{s_j}} = t_i$, $G_i[x_1, \dots, x_{p_i}]_{x_1 \dots x_{p_i}} = t'_i$, and $H(t_1, \dots, t_k, t'_1, \dots, t'_m) \in \Delta$. But since $[H(v_1, \dots, v_k, u_1, \dots, u_m)]_{v_1 \dots v_k u_1 \dots u_m} = [F'(v_1, \dots, v_k) \& G'(u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} \in \Delta$ where

$[F(v_1, \dots, v_k, u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} = [F'(v_1, \dots, v_k)]_{v_1 \dots v_k u_1 \dots u_k} \in \Delta$ and
 $[G(v_1, \dots, v_k, u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} = [G'(u_1, \dots, u_k)]_{v_1 \dots v_k u_1 \dots u_k} \in \Delta$
 one direction of axiom (B) and (T) yield $\forall v_1, \dots, v_k, u_1, \dots, u_m. F'(v_1, \dots, v_k) \& G'(u_1, \dots, u_k) \leftrightarrow$
 $H(v_1, \dots, v_k, u_1, \dots, u_m)$ and
 $\forall v_1, \dots, v_k, u_1, \dots, u_m. F(v_1, \dots, v_k, u_1, \dots, u_k) \leftrightarrow F'(v_1, \dots, v_k) \in \Delta$ and
 $\forall v_1, \dots, v_k, u_1, \dots, u_m. G(v_1, \dots, v_k, u_1, \dots, u_k) \leftrightarrow G'(u_1, \dots, u_k) \in \Delta$.
 whence we deduce that $F'(t_1, \dots, t_k) \& G'(t'_1, \dots, t'_k) \in \Delta$ and hence $F(t_1, \dots, t_k, t'_1, \dots, t'_k)$ and
 $G(t_1, \dots, t_k, t'_1, \dots, t'_k)$ are in Δ which is equivalent to $(F_1, \dots, F_k) \in \mathcal{H}_\Delta F$ and $(G_1, \dots, G_m) \in \mathcal{H}_\Delta G$. The other direction is similar. The conditions for u, e, i, c, r are proven in the same way. We now show the condition for p_i . Suppose we had $(F_1, \dots, F_k, G_1, \dots, G_{i-1}) \in \mathcal{H}_\Delta p_i(F, G)$ where $p_i(F, G) = H$. Then we have closed terms t_i, t'_i such that $F_i[x_1 w, \dots, x_{s_i}]_{x_1 \dots x_{s_i}} = t_i$, $G_i[x_1, \dots, x_{p_i}]_{x_1 \dots x_{p_i}} = t'_i$, and $H(t_1, \dots, t_k, t'_1, \dots, t'_i) \in \Delta$. But since $[H(v_1, \dots, v_k, w_1, \dots, w_i)]_{v_1 \dots v_k w_1 \dots w_i} = [F(v_1, \dots, v_k, [G(x_1, \dots, x_j, w_1, \dots, w_i)]_{x_1 \dots x_j})]_{v_1 \dots v_k w_1 \dots w_i} \in \Delta$ using as previously (B) and (T) we get that

$$F(t_1, \dots, t_k, [G(x_1, \dots, x_j, t'_1, \dots, t'_i)]_{x_1 \dots x_j}) \in \Delta$$

We need to prove that $(F_1, \dots, F_k, p_o^j(G, G_1, \dots, G_j)) \in \mathcal{H}_\Delta F$. This is equivalent to proving that, for $H = p_o^j(G, G_1, \dots, G_j)$ there is a $t = [H(y_1, \dots, y_j)]_{y_1 \dots y_j}$ such that $F(t_1, \dots, t_k, t) \in \Delta$. But this follows from taking the $H \in D_\Delta$ such that $[H(y_1, \dots, y_j)]_{y_1 \dots y_j} = [G(x_1, \dots, x_j, t'_1, \dots, t'_i)]_{x_1 \dots x_j} = [G(x_1, \dots, x_j, G_i[x_1, \dots, x_{p_i}]_{x_1 \dots x_{p_i}}, \dots, G_i[x_1, \dots, x_{p_i}]_{x_1 \dots x_{p_i}})]_{x_1 \dots x_j}$.

We now need prove the T1 condition. Let $F \neq G$ for $F, G \in D_\Delta$. Then $[F(x_1, \dots, x_n)]_{x_1 \dots x_n} \neq [G(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$. Since \mathcal{A} is T1 perfect there exists a Δ' such that $\exists \bar{x}. \neg F(x_1, \dots, x_n) \leftrightarrow G(x_1, \dots, x_n) \in \Delta'$. Hence by ω -completeness there are terms t_1, \dots, t_n such that $\neg F(t_1, \dots, t_n) \leftrightarrow G(t_1, \dots, t_n) \in \Delta'$ and the result follows.

We now prove by induction on complexity that we obtain a model of the sentences in Δ . For the atomic case note that if $t = [F(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$ and $F \in D_\Delta$ then $Dt = F$. Thus if $A(t_1, \dots, t_n) \in \Delta$ consider $t_i = [F_i(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$ with $F_i \in D_\Delta$. Then $(Dt_1, \dots, Dt_n) = (F_1, \dots, F_n) \in \mathcal{GI}(F)$ so $\mathcal{GD}[A(t_1, \dots, t_n)] = T$. Consider $A \& B \in \Delta$. Then by completeness A and B belong to Δ and thus by induction $\mathcal{G}[A] = T$ and $\mathcal{G}[B] = T$ and thus $\mathcal{GD}[A \& B] = T$. The case of $\neg A$ is similar. Finally if $\exists x. A(x) \in \Delta$ then ω -completeness implies that there is a closed term t such that $A(t) \in \Delta$ and so by induction $\mathcal{GD}[A(t)] = T$. Now use lemma 7.6 and 8.3 with $\mathcal{A}(v) = F$ for $t = [F(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$.

10 Soundness and Completeness of T2

We have already proven the soundness of the standard first-order logical axioms and rules for any model. Also the soundness of axioms 6 and 7. Axioms 9 to 11 are clearly sound by the type 2 model condition.

Lemma 10.1 *TGen is sound: suppose that F does not occur in $A(v)$. If $\vdash A([F(x_1, \dots, x_n)]_{x_1 \dots x_n})$ then we can infer $\vdash A(t)$ whenever t complex, of arity n and is free for v in A .*

Proof. By (L) we have

$$\vdash [F(x_1, \dots, x_n)]_{x_1 \dots x_n} = t \rightarrow A([F(x_1, \dots, x_n)]_{x_1 \dots x_n}) \rightarrow A(t)$$

from which it follows that

$$\vdash A([F(x_1, \dots, x_n)]_{x_1 \dots x_n}) \rightarrow [F(x_1, \dots, x_n)]_{x_1 \dots x_n} = t \rightarrow A(t)$$

Assume $\vdash A([F(x_1, \dots, x_n)]_{x_1 \dots x_n})$. Then by soundness of (L), (MP) and tautologies we get

$$\models [F(x_1, \dots, x_n)]_{x_1 \dots x_n} = t \rightarrow A(t)$$

By the non-cycling condition for T2 models F does not occur in t and hence the interpretation of F does not affect the denotation of Dt . But choose an interpretation \mathcal{I} assigning to $[F(x_1, \dots, x_n)]_{x_1 \dots x_n}$ the value Dt . In this case we have $\models [F(x_1, \dots, x_n)]_{x_1 \dots x_n} = t$ and hence $T_{\mathcal{I}}(A(t))$. But since $D[A(t)]$ cannot depend on the interpretation is we must have $\models A(t)$.

Lemma 10.2 *We have that $[A]_{\bar{x}} = [B]_{\bar{x}} \rightarrow A \leftrightarrow B$ is sound.*

Proof. For any T2 model, interpretation and assignment we must show that $T([A]_{\bar{x}} = [B]_{\bar{x}} \rightarrow A \leftrightarrow B)$. We must show that if $T([A]_{\bar{x}} = [B]_{\bar{x}})$ then $T(A \leftrightarrow B)$. Now if $T([A]_{\bar{x}} = [B]_{\bar{x}})$ then we have $D[A]_{\bar{x}} = D[B]_{\bar{x}}$. We now show that $T(A \leftrightarrow B)$. We show only one implication, the other is similar. We need only show that $T(\forall \bar{x}. A \rightarrow B)$, that is, $\mathcal{G}D[\forall \bar{x}. A \rightarrow B] = T$. Now

$$\begin{aligned} D[\forall \bar{x}. A \rightarrow B] &= D[\neg \exists \bar{x}. \neg(A \rightarrow B)] = D[\neg \exists \bar{x}. A \& \neg B] \\ &= ne \dots e D[A \& \neg B]_{\bar{x}} = ue \dots ek D[A]_{\bar{x}} n D[B]_{\bar{x}} = ue \dots ek D[A]_{\bar{x}} n D[A]_{\bar{x}} \end{aligned}$$

So $\emptyset \in \mathcal{H}ue \dots ek D[A]_{\bar{x}} n D[A]_{\bar{x}}$ iff there is no (x_1, \dots, x_n) such that $(x_1, \dots, x_n) \in \mathcal{H}D[A]_{\bar{x}}$ and $(x_1, \dots, x_n) \notin \mathcal{H}D[A]_{\bar{x}}$. Hence $\mathcal{G}D[\forall \bar{x}. A \rightarrow B] = T$.

Note that this rule is sound for any model.

Lemma 10.3 *Let Γ be a consistent set of sentences in T2. Then it is included in a maximal, consistent, ω -complete set of sentences Δ such that for each closed term $[A]_{x_1, \dots, x_n}$ the sentence $[A]_{x_1, \dots, x_n} = [F(x_1, \dots, x_n)]_{x_1, \dots, x_n}$ belongs to Δ , for some symbol F .*

Proof. We assume we have an enumeration S_1, S_2, \dots of the sentences of L^ω as in the proof of completeness of T1. Let $\Delta_0 = \Gamma$. Then given Δ_i , consider the sentence S_{i+1} . If it is consistent with Δ_i then $\Delta_{i+1} = \Delta_i \cup \{S_{i+1}\} \cup C_i$ otherwise $\Delta_{i+1} = \Delta_i \cup C_i$. Consider the set (in the order of S_1, S_2, \dots) of all existential sentences $\exists x. A(x), \exists x. A'(x), \dots$ in Δ_i which occur before S_{i+1} . We proceed as follows. We start by picking a constant a_1 such that $A(a_1)$ is the first instantiation occurring after S_{i+1} such that a_1 does not occur before S_{i+1} or in Δ_i . We then proceed to chose an constant a_2 such that $A'(a_2)$ is the first instantiation after $A(a_1)$ such that a_2 does not occur before or in $A(a_1)$ and so forth. C_i consists of the set of such instantiations $\{A(a_1), A'(a_2), \dots\}$.

Each Δ_i is consistent, by a similar argument as in the T1 case.

We show first that $\Delta = \bigcup \Delta_i$ is maximal, consistent and ω -complete. It is obviously consistent because each Δ_i is consistent. To show that it is maximal we need to show that for any sentence S_i either S_i or $\neg S_i = S_j$ is in Δ . Suppose then that neither formula was in Δ . Then we must have that $S_i \cup \Delta_{i-1}$ and $S_j \cup \Delta_{j-1}$ are inconsistent. Suppose w.l.o.g. that $i < j$. Then since $\Delta_i \subseteq \Delta_j$ we get a contradiction. For if both a sentence and it's negation are inconsistent with a set of sentences, that set is inconsistent.

To show that Δ is ω -complete we need to show that if $\Delta \vdash \exists v. F(v)$ then $\Delta \vdash F(t)$ for some term t . If $\Delta \vdash \exists v. F(v)$ then by maximality $\exists v. F(v) \in \Delta$ and let it have been introduced in stage Δ_i . Then in a subsequent stage j , by construction it is clear that C_j and hence Δ will contain a formula of the form $F(c)$ for some constant c .

Finally we show that for any closed term $[G]_{\bar{x}}$ there is a primitive predicate F such that $[G]_{\bar{x}} = [F(x_1, \dots, x_k)]_{x_1, \dots, x_k}$. We can choose such a F (let $[G]_{\bar{x}} = [F(x_1, \dots, x_k)]_{x_1, \dots, x_k}$ be S_i) so that F does not occur in any of the previous S_j , in G or in Γ . Hence in the stage where S_i may be introduced this formula is consistent with the previous Δ_j and thus belongs to Δ .

We form the canonical model and canonical interpretation as in the T1 case, using Δ . Verification that we obtain in fact a model is the same. Note that were in the T1 case we used (B) and (T) here we can make use of axiom 8. The only difference is that we have only one extension function $\mathcal{G} \in H$ corresponding to Δ itself.

Lemma 10.4 *The canonical model defined above with the canonical interpretation is a T2-model and a model for Δ .*

Proof. We check the type 2 condition. The condition of being one-to-one clearly follows from axioms 10 and 11. For instance if we had $F = n(G) = n(G')$ for $F, G, G' \in D_\Delta$ then $[F(x_1, \dots, x_n)]_{x_1 \dots x_n} = [\neg G(x_1, \dots, x_n)]_{x_1 \dots x_n}$ and $[F(x_1, \dots, x_n)]_{x_1 \dots x_n} = [\neg G'(x_1, \dots, x_n)]_{x_1 \dots x_n}$ are in Δ . But since Δ is maximal we have then $[\neg G(x_1, \dots, x_n)]_{x_1 \dots x_n} = [\neg G'(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$ and hence by axiom 10 that $[G(x_1, \dots, x_n)]_{x_1 \dots x_n} = [G'(x_1, \dots, x_n)]_{x_1 \dots x_n} \in \Delta$. But the conditions on elements of D_Δ imply then that $G = G'$. The condition of having disjoint ranges similarly follows from axiom 9. The non-cycling condition follows from axiom 12.

We prove by induction on complexity that we obtain a model of the sentences in Δ in the same way as in lemma 9.7.

11 Fregean Arithmetic in T2

Let Δ be a marked binary predicate in T2, where we write $x\Delta y$ for $\Delta(x, y)$.

Lemma 11.1 *Bealer, Quality and Concept, p122.*

Proof. We must show that NN0, where

$$0 = [\neg \exists v(v\Delta y)]_y$$

$$NNx \equiv_{df} \forall z(I(z) \rightarrow x\Delta z)$$

where

$$I(z) \equiv_{df} 0\Delta z \ \& \ (\forall y(y\Delta z \rightarrow S(y)\Delta z))$$

and

$$S(x) \equiv_{df} [\exists u(u\Delta x \ \& \ \exists v(\neg v\Delta u \ \& \ y \cong [w = v \vee w\Delta u])_w)]_y$$

$$x \cong y \equiv_{df} \forall w(w\Delta x \leftrightarrow w\Delta y)$$

We have the tautology $0\Delta z \ \& \ (\forall y(y\Delta z \rightarrow S(y)\Delta z)) \rightarrow 0\Delta z$ to which applying \forall -generalization we get immediately the formula NN0.

We now show that $NNx \rightarrow NN(S(x))$. We must show that

$$\forall z(I(z) \rightarrow x\Delta z) \rightarrow \forall z(I(z) \rightarrow S(x)\Delta z)$$

By tautologies and a quantifier axiom we have

$$I(z) \rightarrow \forall y(y\Delta z \rightarrow S(y)\Delta z) \rightarrow (x\Delta z \rightarrow S(x)\Delta z)$$

If we consider the tautology $A \rightarrow (B \rightarrow C) \rightarrow ((A \rightarrow B) \rightarrow (A \rightarrow C))$ then we get

$$\vdash (I(z) \rightarrow x\Delta z) \rightarrow (I(z) \rightarrow S(x)\Delta z)$$

and hence using generalization and $\forall x(A \rightarrow B) \rightarrow (\forall x A \rightarrow \forall x B)$:

$$\vdash \forall z(I(z) \rightarrow x\Delta z) \rightarrow \forall z(I(z) \rightarrow S(x)\Delta z))$$

which is $NN(x) \rightarrow N(S(x))$.

We now show that $\neg\exists x : 0 = S(x)$. But this equivalent to

$$\forall x \neg([\neg\exists v(v\Delta y)]_y = [\exists u(u\Delta x \ \& \ \exists v(\neg v\Delta u \ \& \ y \cong [w = v \vee w\Delta u])_w)]_y)$$

which is an immediate consequence of the T2 axiom for these intensional abstracts belong to different domains.

We now show that $x = y \rightarrow S(x) = S(y)$.

Consider $[u\Delta v]_v$. Then $[u\Delta x]$ is the predication of $[(u\Delta v)]_v$ of x and $[u\Delta y]$ is the predication of $[(u\Delta v)]_v$ of y . The axiom (A11) yields immediately that $x = y \rightarrow [u\Delta x] = [u\Delta y]$. Denoting $[(\exists v(\neg v\Delta u \ \& \ z \cong [(w = v \vee w\Delta u)]_w))]_z$ by s we get by the same axiom that $x = y \rightarrow [(u\Delta x \ \& \ s)]_z = [(u\Delta y \ \& \ s)]_z$ where we first apply (A10) to obtain the expansions $[u\Delta x]_z$ and $[u\Delta y]_z$. Finally we apply the same axiom for existential generalization to obtain as desired.

Finally we must show that $\forall z(I(z) \rightarrow \forall x(NNx \rightarrow x\Delta z))$. The result follows easily from quantifier rules and axioms from $I(z) \rightarrow (NNx \rightarrow x\Delta z)$ which is just the tautology

$$A \rightarrow ((A \rightarrow B) \rightarrow B)$$

Let $A(x) \equiv_{df} NNx \ \& \ \exists y.(NNy \ \& \ y = S(x))$. Consider the following *predication axiom*

$$x\Delta[A(u)]_u \leftrightarrow A(x)$$

Then according to Bealer we have:

Theorem 11.2 *Quality and Concept, p127. If we add the above predication axiom the complete theory for the structure $\langle NN, =, S, 0 \rangle$ is derivable in T2.*

Frege's Law V is generally blamed for the paradoxes. But the paradoxes can be avoided if the law is retained but higher-order variables and quantifiers are dropped, for then we obtain the sound and complete system of first-order logic with extensional abstraction (T1 with modal collapse), law V corresponding to axiom (B).

12 Conclusion and Future Work

We have given detailed proofs of the soundness and completeness of Bealer's systems T1 and T2 as well as simplifying the original presentation of Bealer Decomposition in [1]. There are many other results which are stated (with or without a sketch of a proof) in *Quality and Concept* as well as open problems. There is a whole avenue to explore regarding the addition of "predication" axioms to T2. In particular, we should give a detailed proof of Theorem 11.2 of the previous section. In [1][pp.67 - 68] Bealer proposes a way of combining T1 and T2 into a single logic. Here equality means T2 equality (which is finer) and T1 equality is represented by an additional equivalence relation \approx . One problem with proving soundness and completeness seems to be that we need to use Bealer's lemma. But in this case Bealer lemma will be an identity involving \approx rather than $=$. Thus it would seem that we need to add an extra condition on the extension functions: $x \in \mathcal{H}y$ and $z \approx x$ implies that $z \in \mathcal{H}y$. To prove the completeness of system T2' seems to be the most logical follow-up to the present work. Another interesting aspect would be to abandon the rather cumbersome Hilbert-style axiomatic system and formulate a natural deduction version of T1 and T2 instead. The major challenge in *Quality and Concept* is to find an axiomatization and prove soundness and completeness for

Bealer's System T3. Syntactically T3 is like T1 and T2 except that there are now two distinct term-forming operators $[A]_{\bar{x}}$ and $|A|_{\bar{x}}$. Hence we will have two version of the Bealer operators and the semantics must have two sets of the corresponding functions. We now give a brief descriptions of the structure of the models envisioned by Bealer.

According to Bealer the building blocks of reality consist in primitive qualities, connections and conditions. Bealer, in his realist, anti-representationalist approach in *Quality and Concept*, ch.8, argues that we should consider two sets of syntactic operators, one for forming relations, properties and conditions (pertaining to objective phenomenality) out of the primitives C, I, N, E, U, R, P_n, K and $C', I', N', E', U', R', P'_n, K'$ for forming thoughts and ideas (pertaining to the subjective or conceptual), which includes Cambridge properties. Our thoughts are ultimately grounded in reality itself: they should possess a decomposition tree into objective phenomenal components.

We now consider the model-theoretic counterpart in which to the two classes of syntactic operations C, I, N, E, U, R, P_n, K and $C', I', N', E', U', R', P'_n, K'$ correspond to operations c, i, n, e, u, r, p_n, k and $c', i', n', e', u', r', p'_n, k'$ on a set D partitioned into disjoint sets D_i corresponding to the different arities of intensional terms. We will call the first set of operations primary entity operations o_2 and the second set secondary entity operations o_1 . Secondary-entity operations must satisfy injectivity, disjoint range and non-cycling conditions (Postulate I). In D we consider two fundamental domains: secondary entities \mathcal{E}_2 which are defined as the range of o_2 and primary entities \mathcal{E}_1 defined as the range of o_1 . Given a $d \in D$ we can seek for various possible decompositions through the inverse of the operations o_1 and o_2 . Note that such decompositions do not need to follow the order of the (in this case generalised) Bealer decomposition. We denote a decomposition by $o_i(d_1, \dots, d_k)$ for $d_i \in D$ and we can distinguish those d_i which are the second or further argument of a P_n or P'_n and those that are not (Bealer calls these *subject nodes*). Our decompositions may involve the intertwining of both o_1 and o_2 .

Then there are two kinds of "primes", those that cannot be decomposed in terms of o_1 , which we call simple primary entities \mathfrak{P}_1 and those that cannot be decomposed in terms of o_2 which we call secondary primary entities \mathfrak{P}_2 . For each of these sets we can consider their restrictions to the various D_i . We have obviously

$$D = \mathcal{E}_1 \cup \mathcal{E}_2 \cup \mathfrak{P}_1 \cup \mathfrak{P}_2$$

Clearly $\mathfrak{P}_1 \cap \mathcal{E}_1 = \emptyset$ and $\mathfrak{P}_2 \cap \mathcal{E}_2 = \emptyset$. We put $\mathcal{C} = \mathcal{E}_2 \cap \mathcal{E}_1$ (Cambridge entities). What is the relationship of \mathfrak{P}_2 to \mathcal{E}_1 ? We consider the \mathcal{E}_1 which are not Cambridge entities, that is, $\mathcal{P} = \mathcal{E}_1 \cap \mathfrak{P}_2$, to be primitive entities which cannot be further decomposed via an abstract thought-operation.

The other postulate (Postulate II) is that all secondary entities must admit a decomposition $o_2(d_1, \dots, d_2)$ in which the d_i are either in \mathcal{P} or subject nodes. It would be interesting to understand better the significance of this postulate of Bealer's.

We note that \mathcal{P} need not be closed under the operations o_1 . If we consider its intersection with the various D_i we get primitive beings for $i = -1$, propositions for $i = 0$, qualities for $i = 1$ and connections for $i > 1$. A part $\mathcal{M} \subset \mathcal{P} \cap D_{-1}$ consists of minds, those beings capable of entering into transcendental connections [1][245-248].

After the publication of [1] in 1982 Bealer continued to develop his theory of PRPs for over two decades, in particular in [3, 7] where the distinction between platonic and non-platonic modes of presentation is made together with a further definition of "logical form". It is clear that an analysis of different forms of syntactic decomposition of formulas and terms play a central role. Hopefully a combination with approaches which focus (in a non-nominalistic way) on the constructive or algorithmic meaning of the syntactic structure of expressions, such as those of Tichý[30, 31] or Moschovakis[24], are likely to prove fruitful in bringing the theory of PRPs to a conclusive and satisfactory form.

References

- [1] Bealer, G. (1982). *Quality and Concept*, Oxford University Press.
- [2] Bealer, G. (1983). "Completeness in the theory of properties, relations, and propositions", *Journal of Symbolic Logic* 48 (2): 415-426.
- [3] Bealer, G. (1994). "Property theory: the type-free approach v. the Church approach", *Journal of Philosophical Logic* 23: 139-171.
- [4] Bealer, G. (1993). "A solution to Frege's puzzle". In J. E. Tomberlin (Ed.), *Philosophical perspectives 7: Language and logic* (pp. 17-60). Atascadero, CA: Ridgeview Publishing Company.
- [5] Bealer, G. (1998). "Propositions". *Mind*, 107(425), 1-32.
- [6] Bealer, G. and Mönnich, U (1989). ""Property Theories, In Dov Gabbay & Franz Guenther (eds.), *Handbook of Philosophical Logic, Volume IV*. Kluwer Academic Publishers. pp. 133-251.
- [7] Bealer, G. (1989). "Fine-Grained Type-Free Intensionality", in Gennero Chierchia, Barbara H. Partee & Raymond Turner (eds.), *Properties, Types, and Meaning, Volume 1*. Kluwer Academic Publishers. pp. 177-230.
- [8] Bressan, A. (1972). *A General Interpreted Modal Calculus*, New Haven: Yale University Press.
- [9] Carnap, R. (1947). *Meaning and Necessity*, Chicago: University of Chicago Press. Enlarged edition 1956.
- [10] Church, A. (1951). "A formulation of the logic of sense and denotation", in P. Henle (Ed.), *Structure, Method and Meaning*, New York: The Liberal Arts Press. pp 3-24.
- [11] Church, A. (1973). *Outline of a revised formulation of the logic of sense and denotation (part I)*", *Noûs*, 7: 24-33.
- [12] Church, A. (1974). "Outline of a revised formulation of the logic of sense and denotation (part II)", *Noûs*, 8: 135-156.
- [13] Cresswell, M. J. (1973). *Logic and languages*. London: Meuthen.
- [14] Frege, G. (1892). *Über Sinn und Bedeutung*. *Zeitschrift für Philosophie und philosophische Kritik*, 100: 25-50. English translation as 'On Sinn and Bedeutung,' in (Beaney 1997).
- [15] Gallin, D. (1975). *Intensional and Higher-Order Modal Logic*, Amsterdam: North-Holland.
- [16] Kripke, S. (1963). "Semantical considerations on modal logics," in *Acta Philosophica Fennica*, 16: 83-94.
- [17] Kripke, S. (1980). *Naming and Necessity* (Second edition), Cambridge, MA: Harvard University Press.
- [18] Kripke, S. (2008). *Frege's Theory of Sense and Reference: Some Exegetical Notes*, *Theoria*, 74: 181-218.

- [19] Marcus, R. (1946). “A Functional Calculus of First Order Based on Strict Implication,” *The Journal of Symbolic Logic*, 11: 1–16.
- [20] Marcus, R. (1961) “Modalities and intensional Languages,” *Synthese*, XIII: 303–322. Reprinted in *Modalities, Philosophical Essays*, Ruth Barcan Marcus, Oxford University Press, 1993.
- [21] Montague, R. (1960). “On the nature of certain philosophical entities,” *The Monist*, 53: 159–194. Reprinted in (Thomason 1974), 148–187.
- [22] Montague, R. (1970). “Pragmatics and intensional logic,” *Synthèse*, 22: 68–94. Reprinted in (Thomason 1974), 119–147.
- [23] Moschovakis, Y. N. (1989). The formal language of recursion. 54, 1216–1252.
- [24] Moschovakis, Y. N. (1994). Sense and denotation as algorithm and value. In J. Oikkonen and J. Vaananen (Eds.), *Lecture Notes in Logic, Volume 2*, pp. 210–249. Springer. [Full version of proof of Theorem 4.1 from that paper is available online.]
- [25] Moschovakis, Y. (2006). “A logical calculus of meaning and synonymy,” *Linguistics and Philosophy*, 29: 27–89.
- [26] Quine, W. V. (1963). “Reference and modality,” in *From a Logical Point of View* (second ed.), Chapter VIII, pp. 139–159. New York: Harper Torchbooks.
- [27] Parsons, D. (2016). *Theories of Intensionality*, Springer, Singapore.
- [28] Steiner, M. (1975). *Mathematical Knowledge*, Cornell University Press.
- [29] Thomason, R. H. (Ed.) (1974). *Formal Philosophy, Selected Papers of Richard Montague*, New Haven and London: Yale University Press.
- [30] Tichý, P. (1971). “An Approach to Intensional Analysis,” *Noûs*, 5: 273–297.
- [31] Tichý, P. (1988). *The foundations of Frege’s logic*, Berlin and New York: De Gruyter.
- [32] Zalta, E. (1988). *Intensional Logic and the Metaphysics of Intentionality*, Cambridge, MA: MIT Press.