

Around spin Hurwitz numbers

A. D. Mironov^{a,b,c*}, A. Morozov^{d,b,c†}, S. M. Natanzon^{e,b‡}, A. Yu. Orlov^{f§}

^a *Lebedev Physics Institute, Moscow 119991, Russia*

^b *ITEP, Moscow 117218, Russia*

^c *Institute for Information Transmission Problems, Moscow 127994, Russia*

^d *MIPT, Dolgoprudny, 141701, Russia*

^e *HSE University, Moscow, Russian Federation*

^f *Shirshov Institute of Oceanology, RAS, Nahimovskii Prospekt 36, Moscow 117997, Russia*

Abstract

We present a review of the spin Hurwitz numbers, which count the ramified coverings with spin structures. They are related to peculiar Q Schur functions, which are actually related to characters of the Sergeev group. This allows one to put the whole story into the modern context of matrix models and integrable hierarchies. Hurwitz partition functions are actually broader than the conventional τ -functions, but reduce to them in particular circumstances. We explain how special d -soliton τ -functions of the KdV and Veselov-Novikov hierarchies generate the spin Hurwitz numbers $H^\pm(\Gamma_d^b)$ and $H^\pm(\Gamma_d^b, \Delta)$. The generating functions of the spin Hurwitz numbers are hypergeometric τ -functions of the BKP integrable hierarchy, and we present their fermionic realization. We also explain how one can construct τ -functions of this type entirely in terms of the Q Schur functions. An important role in this approach is played by factorization formulas for the Q Schur functions on special loci.

Boris Dubrovin was one of the brightest minds in modern mathematics, in fact, his well-known achievements cover only a small part of all what he thought about and planned to do. Long ago, as young students, we were inspired by the brilliant volume [1], to which Boris also contributed a big part of his knowledge and vision. During his entire life in science, he tried to make new fields as clear and transparent, as happened to the classical subjects in that unique textbook.

This paper was supposed to have a single author, Sergey Natanzon, and was planned as his personal tribute to his close friend, Boris. Unfortunately, Sergey was not given a chance to fulfill this duty. The present note is a brief review of what he taught us about the increasingly important subject of *spin* Hurwitz numbers, and it certainly lacks the vision which Sergey had on it and which we, survivors, will still need to rediscover in our future work.

*mironov@lpi.ru; mironov@itep.ru

†morozov@itep.ru

‡Sergey Natanzon passed away on December, 7, 2020

§orlovs55@mail.ru

Contents

1	Introduction	2
2	Hurwitz numbers	4
2.1	Classical Hurwitz numbers	4
2.2	Spin Hurwitz numbers	5
3	Q Schur functions	5
3.1	Projective Schur functions	6
3.2	Neutral fermions and projective Schur functions	6
3.2.1	From KP to BKP	7
3.2.2	Q Schur function as fermionic average	8
3.2.3	Sergeev characters	9
4	BKP τ-functions	10
4.1	Hypergeometric BKP vs KP τ -functions	10
4.2	Bosonization	12
4.3	W_∞^B algebra	13
4.4	Cut-and-join equation	16
5	KdV soliton solution as the BKP solutions	17
5.1	Solitons of the KdV and BKP hierarchies which generate Hurwitz numbers	17
5.2	Different fermionic expressions generating Hurwitz numbers	19
5.3	Matrix model τ -functions as fermion averages	21
5.4	On character expansion for KdV τ -functions and BKP	21
6	Factorization on special loci	22
6.1	Specialization at $p_k = \delta_{k,r}$	22
6.2	Basic factorization formula	23
7	Hypergeometric τ-functions entirely made from characters	24
8	Conclusion	25
A	Derivation of (6.6) from (6.9)	26
B	Factorization formula (6.9) from fermion calculus	27
C	Multicomponent KP and BKP τ-functions	29

1 Introduction

Hurwitz numbers [2–5] count ramified coverings, and their significance in physics is dictated by the role that complex curves play in string theory. Applications range from the free field calculus on ramified Riemann surfaces [6] to genus expansion in matrix models and topological recursion [7, 8]. The spin Hurwitz numbers [9, 10] do just the same things to Riemann surfaces with spin structures labeled by theta-characteristics. The matrix model counterparts in this case appear to be the cubic Kontsevich [11, 12] and Brezin-Gross-Witten (BGW) [8, 13] models, which are non-obvious generalizations of the Hermitian matrix model [14] equivalent to a quadratic counterpart of Kontsevich theory [15, 16]. As usual, the most efficient technique to develop spin Hurwitz calculus relies on algebraic approach, which finally puts the problem into the framework of integrable systems. The basic of this approach includes the following ingredients [17]:

- A relevant substitute of the Schur functions, which in the case of cubic Kontsevich and BGW models is provided by the Q Schur functions
- Their relation to the Hall-Littlewood polynomials [18]
- Their relation to characters of the Sergeev algebra [19], which provides a relevant generalization of symmetric group characters
- The Fröbenius formula, expressing the spin Hurwitz numbers through the Sergeev characters and the Q Schur functions
- Commuting system of cut-and-join \hat{W} operators [17, 20], which have Q Schur functions as their common eigenfunctions and the Sergeev characters as eigenvalues
- Free fermion representations [21–24], which allow one to represent Q Schur functions as Pfaffians
- Integrability properties of the spin Hurwitz numbers. As usual [20, 25, 26], Hurwitz τ -functions form a broader and still uncomprehensible variety, but, in special cases, we get soliton and other solutions to the BKP hierarchy [21, 22]
- Hypergeometric τ -functions of the BKP hierarchy [24, 27]. By definition, they are bilinear in the Q Schur functions with coefficients of a very special product form (DP denotes the strict partitions)

$$\tau_{BKP}\{p_k, p_k^*\} = \sum_{\alpha \in \text{DP}} Q_\alpha\{p_k\} Q_\alpha\{p_k^*\} \cdot \prod_{i=1}^{\ell(\alpha)} f(\alpha_i) \quad (1.1)$$

Important examples are provided by the ratios

$$\frac{Q_{N\alpha}\{\delta_{k,r}\}}{Q_\alpha\{\delta_{k,r}\}} = \prod_{i=1}^{\ell(\alpha)} f(\alpha_i) \quad \forall \text{ coprime } N, r \quad (1.2)$$

where $\ell(\alpha)$ denotes the number of lines in the Young diagram α , and $N\alpha$ denotes the Young diagram with lengths $N\alpha_i$. In particular, for the cubic Kontsevich model [28],

$$\tau_{K_3}\{p_k\} = \sum_{\alpha \in \text{DP}} Q_\alpha\{p_k\} Q_\alpha\{\delta_{k,3}\} \cdot \frac{Q_\alpha\{\delta_{k,1}\}}{Q_{2\alpha}\{\delta_{k,1}\}} \frac{Q_{2\alpha}\{\delta_{k,3}\}}{Q_\alpha\{\delta_{k,3}\}} \cdot \frac{1}{2^{\ell(\alpha)}} \quad (1.3)$$

while, for the BGW model [29],

$$\tau_{BGW}\{p_k\} = \sum_{\alpha \in \text{DP}} Q_\alpha\{p_k\} Q_\alpha\{\delta_{k,1}\} \cdot \left(\frac{Q_\alpha\{\delta_{k,1}\}}{Q_{2\alpha}\{\delta_{k,1}\}} \right)^2 \cdot \frac{1}{2^{\ell(\alpha)}} \quad (1.4)$$

We refer to the very recent paper [29] for some complimentary details.

As to the ordinary integrability, its intimate relation to characters is well known. The Schur functions are themselves solutions to the Hirota bilinear equations, and general KP/Toda τ -functions are their linear combinations with the coefficients satisfying the Plücker relations, which have determinants and their free fermion realizations as natural solutions. The Q Schur functions play the same role for the BKP hierarchies, only solutions are now Pfaffians, which can be described in terms of “neutral” fermions. These simple facts are, however, not fully trivial: for example, the KdV hierarchy, which can be considered as a reduction of the both KP and BKP hierarchies [21, 22, 30–32], possesses as solutions only the very special Q Schur functions $Q_{[1]}$, $Q_{[2,1]}$, $Q_{[3,2,1]}$, \dots , $Q_{[\dots,4,3,2,1]} \sim \text{Schur}_{[\dots,4,3,2,1]}$, which are the only Schur functions independent of even time-variables p_{2k} and proportional to the

corresponding Q Schur functions. All other Q Schur functions, which are all independent of p_{2k} , do not solve the KdV hierarchy.

In the present paper, we review some auxiliary aspects, related to the free-fermion description of the Q Schur functions, the BKP and KdV hierarchies. In particular, as in the case of ordinary Hurwitz numbers, the lowest “cut-and-join” \hat{W} operators commute with the BKP Hirota equations and generates an especially simple hypergeometric “tau”-function, which can be also considered as an infinite-soliton τ -function of the KdV hierarchy. Technically, the BKP τ -function can be defined as a direct counterpart of the Toda lattice τ -function, but for the matrix elements of $SL(\infty)$ generated by a restricted set of “neutral” fermions and depending only on odd sets of times. Proper weights made of exponentials of power sums of Young diagram lengths in (1.1) (“completed cycles”) in this formalism provides the hypergeometric τ -functions, which are easy and very straightforward to work with. The full (spin) Hurwitz τ -functions involve far more complicated weights made out of all (Sergeev) symmetric characters, and they provide an important generalization beyond (B)KP theory, which still awaits an efficient language and deep investigation.

Notation. Throughout the paper, we denote through $[x]$ the integer part of a number, through $\{x\}$ its fractional part. For an integer k , $(k)_r = r\{k/r\}$ denotes the value mod r . For the strict partition α , $\ell(\alpha)$ is the number of parts, and $\bar{\ell}(\alpha) := 2 \cdot \left\lceil \frac{\ell(\alpha)+1}{2} \right\rceil$.

2 Hurwitz numbers

2.1 Classical Hurwitz numbers

Consider a compact Riemann surface S of genus g with a finite number of points $x_1, \dots, x_n \in S$. Consider a set of Young diagrams $(\Delta^1, \dots, \Delta^n)$ of the same degree $d = |\Delta_i|$. The lengths of the rows $\Delta_1^i, \dots, \Delta_{\ell_i}^i$ of the Young diagram Δ^i give the partition of the number d .

Denote by $\widetilde{M}(\Delta^1, \dots, \Delta^n)$ the set of holomorphic mappings of compact Riemann surfaces $\varphi : P \rightarrow S$, whose critical values lie in $\{x^1, \dots, x^n\}$, and the pre-images $\varphi^{-1}(x^i)$ consist of points, where φ has degrees $\Delta_1^i, \dots, \Delta_{\ell_i}^i$. We call the mappings $\varphi : P \rightarrow S$ and $\varphi' : P' \rightarrow S$ as equivalent if there exists a biholomorphic mapping ϕ such that $\varphi = \varphi' \phi$. Let $M(\Delta^1, \dots, \Delta^n)$ denote the set of equivalence classes in the set $\widetilde{M}(\Delta^1, \dots, \Delta^n)$.

The classical Hurwitz number [5] is the number

$$H_d(g|\Delta^1, \dots, \Delta^n) = \sum_{\varphi \in M(\Delta^1, \dots, \Delta^n)} \frac{1}{|Aut(\varphi)|} \quad (2.1)$$

There is the Fröbenius formula that gives a combinatorial expression for the Hurwitz numbers [4],

$$H_d(g|\Delta^1, \dots, \Delta^n) = \frac{[\Delta_1] \dots [\Delta_k]}{(d!)^2} \sum_R \frac{\psi_R(\Delta_1) \dots \psi_R(\Delta_k)}{\psi_R(1)^{(k-2)}} \quad (2.2)$$

where $[\Delta]$ is the number of permutations of the cyclic type Δ , i.e. the number of elements in the conjugacy class of the symmetric group \mathfrak{S}_d given by the Young diagram Δ , $|\Delta| = d$; $\psi_R(\Delta)$ is value of the character ψ_R of the representation R of the symmetric group \mathfrak{S}_d on the permutation of cyclic type Δ , $\psi_R(1)$ is the value on the permutation with all unit cycles, $\Delta = \underbrace{[1, \dots, 1]}_{d \text{ times}}$, and the sum is

taken over all characters of irreducible representations of \mathfrak{S}_d .

Among the classical Hurwitz numbers, we will be interested only in the so-called double Hurwitz numbers

$$H_d(\Delta, \Delta^*, b) = H_d(0|\Delta, \Delta^*, \Gamma_2^b) \quad (2.3)$$

Here Γ_2 is a Young diagram with one row of length 2 and the rest of rows of length 1. The Γ_2^b means a set from b diagrams Γ_2 .

Consider two infinite sets of variables $p = (p_1, p_2, \dots)$ and $p^* = (p_1^*, p_2^*, \dots)$. Associate monomial $p_\Delta = p_{\Delta_1} \cdots p_{\Delta_\ell}$ to the Young diagram $\Delta = [\Delta_1, \dots, \Delta_\ell]$ and monomial $p_{\Delta^*}^* = p_{\Delta_1^*}^* \cdots p_{\Delta_\ell^*}^*$ to Young diagram $\Delta^* = [\Delta_1^*, \dots, \Delta_\ell^*]$. To the double Hurwitz number $H_d(\Delta, \Delta^*, b)$ we associate the monomial $H_d(\Delta, \Delta^*, b)p_\Delta p_{\Delta^*}^*$. As a generating function for the double Hurwitz numbers, one usually considers the function proposed in [33]

$$\tau(\Delta, \Delta^*, \beta, q) = \sum_{d>0} \sum_{|\Delta|=|\Delta^*|=0} \sum_{b, \Delta, \Delta^*} q^d \frac{\beta^b}{b!} H_d(\Delta, \Delta^*, b) p_\Delta p_{\Delta^*}^* \quad (2.4)$$

According to [34], [35], [33], [36] this function is a τ -function of the 2D Toda lattice (it was called hypergeometric in [35]).

Moreover, according to [20, 37], it satisfies the *cut-and-join* equation

$$\frac{\partial \tau(\Delta, \Delta^*, \beta, q)}{\partial \beta} = W \tau(\Delta, \Delta^*, \beta, q) \quad (2.5)$$

where

$$W = \frac{1}{2} \sum_{a, b > 0} \left((a+b) p_a p_b \frac{\partial}{\partial p_{a+b}} + a b p_{a+b} \frac{\partial^2}{\partial p_a \partial p_b} \right) \quad (2.6)$$

2.2 Spin Hurwitz numbers

A line bundle L on a Riemann surface is called a spin bundle if the tensor square of L is isomorphic to the cotangent bundle. The parity of the space of holomorphic sections of the bundle L is called the parity of the bundle (see [38, 39]) and is denoted by $Arf(L) \in \{0, 1\}$. The surface of genus 0 has exactly one spin bundle, and it is even.

Consider a holomorphic mapping $\varphi : P \rightarrow S$ whose critical point orders are odd. Such a mapping associates the spin bundle L onto S with the spin bundle $\varphi^*(L)$ onto P . Its parity $Arf(\varphi) = Arf(\varphi^*(L))$ depends only on φ and $Arf(L)$.

The spin Hurwitz number was defined in [9] as

$$H_d^{Arf(L)}(g|\Delta_1, \dots, \Delta_n) = \sum_{\varphi \in M(\Delta_1, \dots, \Delta_n)} \frac{(-1)^{Arf(\varphi)}}{|\text{Aut}(\varphi)|} \quad (2.7)$$

Depending on whether the parity of the bundle is even or odd, later on, we use the superscripts $+$ and $-$ respectively.

We are only interested in the double spin Hurwitz numbers, which, in this case, have the form

$$H_d^{sp}(\Delta, \Delta^*, b) = H_d^0(0|\Delta, \Delta^*, \Gamma_3^b) \quad (2.8)$$

where Γ_3 is a Young diagram with one row of length 3 and other rows of length 1, and Γ_3^b means b of such diagrams. The generating functions for such numbers can be written in different ways. The simplest generating functions are proposed and investigated in [17, 40]. They satisfy both the 2BKP hierarchy and modified cut-and-join equations. We discuss these issues in the present paper.

3 Q Schur functions

In this paper, we define the Q Schur functions in a peculiar normalization, which is conventional for fermionic representations, but different from the natural one for the Cauchy identities and matrix models used in [28] and [29].

3.1 Projective Schur functions

To define the Q Schur functions Q_α , we begin following [18] by defining an infinite skew symmetric matrix $(Q_{ij})_{i,j \in \mathbf{N}}$, whose entries are symmetric functions of the infinite sequence of indeterminates $\mathbf{x} = (x_1, x_2, \dots)$, via the following formula:

$$Q_{ij}(\mathbf{x}) := \begin{cases} q_i(\mathbf{x})q_j(\mathbf{x}) + 2 \sum_{k=1}^j (-1)^k q_{i+k}(\mathbf{x})q_{j-k}(\mathbf{x}) & \text{if } (i, j) \neq (0, 0), \\ 0 & \text{if } (i, j) = (0, 0), \end{cases} \quad (3.1)$$

where the $q_i(\mathbf{x})$'s are defined by the generating function:

$$\prod_{i=1}^{\infty} \frac{1 + zx_i}{1 - zx_i} = \sum_{i=0}^{\infty} z^i q_i(\mathbf{x}) \quad (3.2)$$

For instance, $q_1(\mathbf{x}) = 2 \sum_i x_i$. In particular,

$$Q_{(j,0)}(\mathbf{x}) = -Q_{(0,j)}(\mathbf{x}) = q_j(\mathbf{x}) \quad \text{for } j \geq 1 \quad (3.3)$$

For a strict partition α of even cardinality $2n$ (including a possible zero part $\alpha_{2n} = 0$), let $\mathbf{M}_\alpha(\mathbf{x})$ denote the $2n \times 2n$ skew symmetric matrix with entries

$$(\mathbf{M}_\alpha(\mathbf{x}))_{ij} := Q_{\alpha_i \alpha_j}(\mathbf{x}), \quad 1 \leq i, j \leq 2n. \quad (3.4)$$

The Q Schur function is defined as its Pfaffian [18]

$$\boxed{Q_\alpha(\mathbf{x}) := \text{Pf}(\mathbf{M}_\alpha(\mathbf{x}))} \quad (3.5)$$

and, for completeness,

$$Q_\emptyset := 1. \quad (3.6)$$

Equivalently, these may be viewed as functions $q_j(\mathbf{p})$, $Q_{ij}(\mathbf{p})$ of the odd-indexed power sum symmetric functions $\mathbf{p} = (p_1, p_3, \dots)$

$$p_{2i-1} = p_{2i-1}(\mathbf{x}) = \sum_{a=1}^{\infty} x_a^{2i-1}, \quad a = 1, 2, \dots \quad (3.7)$$

Following [18], we use the agreement

$$Q_\alpha\{p_k(\mathbf{x})\} := Q_\alpha(\mathbf{x}). \quad (3.8)$$

In particular, $Q_{[1]}\{p_k\} = 2p_1 = 2 \sum_i x_i$, $Q_{[2]}\{p_k\} = 2p_1^2 = 2(\sum_i x_i)^2$.

There is the following relation:

$$Q_\alpha\{\delta_{k,1}\} = \frac{2^{|\alpha|}}{\prod_{i=1}^{\ell(\alpha)} \alpha_i!} \Delta^*(\alpha), \quad \Delta^*(\alpha) := \prod_{i < j \leq \ell(\alpha)} \frac{\alpha_i - \alpha_j}{\alpha_i + \alpha_j} \quad (3.9)$$

Remark 3.1. In the literature on integrable systems, the variables often used and called times of the BKP hierarchy (see [21], [30], [23], [24], [27], [41]) are $\frac{2p_m}{m}$ with m odd.

In the present paper, we use the BKP hierarchy, however we re-write known BKP formulas in the power sum variables.

3.2 Neutral fermions and projective Schur functions

In this section, we very briefly recall the known facts (details can be found either in the original papers [21], [22], [23], or in [24], the results of which we will use). Anyway, we need to fix the notation.

3.2.1 From KP to BKP

A natural way to construct the BKP hierarchy is to start with the KP hierarchy. A standard way to describe this latter is to realize the τ -function of the hierarchy as a fermionic average

$$\tau(\mathbf{p}_f, \mathbf{p}_f^*) = \langle 0 | \gamma(\mathbf{p}_f) g \gamma^\dagger(\mathbf{p}_f^*) | 0 \rangle \quad (3.10)$$

where $\mathbf{p}_f := (p_1, p_2, p_3, \dots)$ and analogously \mathbf{p}_f^* are sets of KP time variables,

$$\begin{aligned} \gamma(\mathbf{p}_f) &:= e^{\sum_{m>0} \frac{1}{m} \mathcal{J}_m p_m}, & \gamma^\dagger(\mathbf{p}_f^*) &:= e^{\sum_{m>0} \frac{1}{m} \mathcal{J}_{-m} p_m^*} \\ \mathcal{J}_m &:= \sum_{i \in \mathbb{Z}} \psi_i \psi_{i+m}^*, & g &= \exp \left(\sum_{i,j} A_{ij} \psi_i \psi_j^* \right) \end{aligned} \quad (3.11)$$

and ψ_i, ψ_i^* are charged fermions,

$$[\psi_i, \psi_j]_+ = 0, \quad [\psi_i, \psi_j^*]_+ = \delta_{ij}, \quad [\psi_i^*, \psi_j^*]_+ = 0 \quad (3.12)$$

with the vacuum defined as

$$\psi_i |0\rangle = 0 = \langle 0 | \psi_i^* \quad \forall i < 0, \quad \langle 0 | \psi_i = 0 = \psi_i^* |0\rangle \quad \forall i \geq 0 \quad (3.13)$$

The pairwise expectation values are:

$$\langle 0 | \psi_j^\dagger \psi_k | 0 \rangle = \begin{cases} \delta_{j,k} & \text{if } k > 0, \\ 0 & \text{if } k < 0, \end{cases} \quad (3.14)$$

Now one can notice that an embedding into the KP hierarchy of the τ -function that depends on only odd time variables can be naturally achieved by introducing the two sets of neutral fermions $\{\phi_i, i \in \mathbb{Z}\}$

$$\phi_j = \frac{\psi_j + (-1)^j \psi_{-j}^*}{\sqrt{2}}, \quad \hat{\phi}_j = i \frac{\psi_j - (-1)^j \psi_{-j}^*}{\sqrt{2}} \quad (3.15)$$

with the canonical anticommutation relations:

$$[\phi_j, \phi_k]_+ = (-1)^j \delta_{j+k,0}, \quad [\hat{\phi}_j, \hat{\phi}_k]_+ = (-1)^j \delta_{j+k,0}, \quad [\phi_j, \hat{\phi}_k]_+ = 0 \quad (3.16)$$

In particular, $(\phi_0)^2 = \frac{1}{2}$. Acting on the left and right vacua $|0\rangle, \langle 0|$, one obtains

$$\phi_{-j} |0\rangle = 0 = \langle 0 | \phi_j, \quad \forall j > 0. \quad (3.17)$$

and similarly for the second set of fermions. The pairwise expectation values are:

$$\langle 0 | \phi_j \phi_k | 0 \rangle = \begin{cases} (-1)^k \delta_{j,-k} & \text{if } k > 0, \\ \frac{1}{2} \delta_{j,0} & \text{if } k = 0, \\ 0 & \text{if } k < 0, \end{cases} \quad (3.18)$$

For (Euclidean) Fermi fields $\phi(\zeta) := \sum_{n \in \mathbb{Z}} \zeta^n \phi_n$, where $\zeta \in \mathbb{CP}^1$, the variables $T = -\log |\zeta|$ and $\varphi = \arg \zeta$ are treated as time and space variables respectively. The multipoint correlation functions uses the chronological ordering convention that states that the field assigned to a larger time variable is put to the right of the field assigned to a smaller one, and the permutation sign should be taken into account. For example, if $|\zeta_1| > |\zeta_2|$, one writes $\langle 0 | \phi(\zeta_2) \phi(\zeta_1) | 0 \rangle$ as $-\langle 0 | \phi(\zeta_1) \phi(\zeta_2) | 0 \rangle$, which by (3.18) is

equal to $-\frac{1}{2} + \zeta_2 \zeta_1^{-1} - \dots = -\frac{1}{2} \frac{1 - \zeta_2 \zeta_1^{-1}}{1 + \zeta_2 \zeta_1^{-1}}$. For a given strict partition $\alpha = (\alpha_1, \dots, \alpha_{2r})$ where $\alpha_{2r} \geq 0$, we use the notation $\Delta^*(\zeta_\alpha)$ defined by the relation

$$\langle 0 | \phi(\zeta_{\alpha_1}) \cdots \phi(\zeta_{\alpha_{2r}}) | 0 \rangle = 2^{-r} \prod_{i < j \leq 2r} \frac{\zeta_{\alpha_i} - \zeta_{\alpha_j}}{\zeta_{\alpha_i} + \zeta_{\alpha_j}} =: 2^{-r} \Delta^*(\zeta_\alpha) \quad (3.19)$$

The first equality is obtained by the Wick theorem and by a simple analytical reasoning about poles and zeroes.

An important fact is that the factor $\gamma(\mathbf{p}_f)$ becomes a product of two factors at all even times vanishing:

$$\gamma(\mathbf{p}_f) \Big|_{p_{2k}=0} = e^{\sum_{m \in \mathbb{Z}^+_{\text{odd}}} (\frac{2}{m} p_m J_m + \frac{2}{m} p_m \hat{J}_m)} \quad (3.20)$$

where

$$J_m = \frac{1}{2} \sum_{i \in \mathbb{Z}} (-1)^i : \phi_{-i-m} \phi_i : \quad (3.21)$$

and similarly for $\gamma^\dagger(\mathbf{p}^*)$. Here $: X :$ denotes the normal ordering (which is $: X := X - \langle 0 | X | 0 \rangle$ for X quadratic in fermions).

Now we can consider only ‘‘half’’ of this system leaving only one set of the neutral fermions. In this system, there are two mutually commuting Abelian groups of the BKP flows

$$\gamma(\mathbf{p}) := e^{\sum_{m \in \mathbb{Z}^+_{\text{odd}}} \frac{2}{m} J_m p_m}, \quad \gamma^\dagger(\mathbf{p}) := e^{\sum_{m \in \mathbb{Z}^+_{\text{odd}}} \frac{2}{m} J_{-m} p_m} \quad (3.22)$$

One has

$$J_n J_m - J_m J_n = \frac{n}{2} \delta_{n+m, 0} \quad (3.23)$$

and

$$J_m |0\rangle = 0 = \langle 0 | J_{-m}, \quad \forall m > 0 \quad (3.24)$$

which results in

$$\gamma(\mathbf{p}) |0\rangle = |0\rangle, \quad \langle 0 | \gamma^\dagger(\mathbf{p}) = \langle 0 | \quad (3.25)$$

and in

$$\gamma(\mathbf{p}) \gamma^\dagger(\mathbf{p}^*) = e^{\sum_{m \in \mathbb{Z}^+_{\text{odd}}} \frac{2}{m} p_m p_m^*} \gamma^\dagger(\mathbf{p}^*) \gamma(\mathbf{p}) \quad (3.26)$$

3.2.2 Q Schur function as fermionic average

One can construct the Q Schur functions as fermionic averages much similar to how the ordinary Schur functions are realized as fermionic averages of charged fermions [22].

Any nonzero partition with distinct parts (also known as strict partition), say α , can be written as $\alpha = (\alpha_1, \dots, \alpha_r)$, where r is an even number, and $\alpha_1 > \dots > \alpha_r \geq 0$. We call r the completed length of α and denote $\bar{\ell}(\alpha)$. As usual, the length of a partition is the number of non-vanishing parts of α , and it is denoted $\ell(\alpha)$; thus $\ell(\alpha)$ is an odd number if and only if $\alpha_r = 0$, while $\bar{\ell}(\alpha)$ is always even. The length of the zero partition is 0. Following [18], we denote by DP the set of all partitions with distinct parts (or the same: the set of all strict partitions).

Let us introduce the notation

$$\begin{aligned}\Phi_\alpha &= 2^n \phi_{\alpha_1} \cdots \phi_{\alpha_{2n}}, & \Phi_\alpha^\dagger &= (-1)^{\sum_{i=1}^{2n} \alpha_i} 2^n \phi_{\alpha_{-2n}} \cdots \phi_{-\alpha_1} \\ J_\Delta^\dagger &= J_{\Delta_1} \cdots J_{\Delta_m}, & J_\Delta &= J_{-\Delta_m} \cdots J_{-\Delta_1}\end{aligned}\quad (3.27)$$

where $\Delta = (\Delta_1, \dots, \Delta_m)$ is a partition with odd parts (we denote OP the total set of such partitions). We have

$$\begin{aligned}\langle 0 | \Phi_\beta^\dagger \Phi_\alpha | 0 \rangle &= 2^{2n} \delta_{\alpha, \beta} = \langle Q_\alpha, Q_\beta \rangle \\ \langle 0 | J_\Delta^\dagger J_\Delta | 0 \rangle &= 2^{-m} z_\Delta \delta_{\bar{\Delta}, \Delta} = \langle \mathbf{p}_{\bar{\Delta}}, \mathbf{p}_\Delta \rangle\end{aligned}\quad (3.28)$$

where \langle, \rangle denotes the scalar product in the space of symmetric functions, see (8.12) in [18]. Here z_Δ is the standard symmetric factor of the Young diagram (order of the automorphism), and $\mathbf{p}_\alpha := \prod_i^{\ell(\alpha)} p_{\alpha_i}$.

The key relation we need was found in [23] and, in our notations, is

$$\boxed{Q_\alpha \{p_k\} = \langle 0 | \gamma(\mathbf{p}) \Phi_\alpha | 0 \rangle = \langle 0 | \Phi_\alpha^\dagger \gamma^\dagger(\mathbf{p}) | 0 \rangle}\quad (3.29)$$

which results in (3.5) according to the Pfaffian form of the Wick theorem with the choice $p_m = \sum_i x_i^m$.

As a result, one has

$$\begin{aligned}\gamma^\dagger(\mathbf{p}) | 0 \rangle &= \sum_{\alpha \in \text{DP}} 2^{-2n} \Phi_\alpha | 0 \rangle Q_\alpha \{p_k\} = \sum_{m \geq 0} \sum_{\substack{\Delta \in \text{OP} \\ \ell(\Delta) = m}} 2^m J_\Delta | 0 \rangle \frac{\mathbf{p}_\Delta}{z_\Delta} \\ \langle 0 | \gamma(\mathbf{p}) &= \sum_{\alpha \in \text{DP}} 2^{-2n} Q_\alpha \{p_k\} \langle 0 | \Phi_\alpha^\dagger = \sum_{k \geq 0} \sum_{\substack{\Delta \in \text{OP} \\ \ell(\Delta) = k}} 2^m \frac{\mathbf{p}_\Delta}{z_\Delta} \langle 0 | J_\Delta^\dagger\end{aligned}\quad (3.30)$$

thus, one gets

$$e^{\sum_{m \in \mathbb{Z}_{\text{odd}}^+} \frac{2}{m} p_m p_m^*} = \langle 0 | \gamma(\mathbf{p}) \gamma^\dagger(\mathbf{p}) | 0 \rangle = \sum_{\alpha \in \text{DP}} 2^{-\ell(\alpha)} Q_\alpha \{p_k\} Q_\alpha \{p_k^*\} = \sum_{\Delta \in \text{OP}} 2^{\ell(\Delta)} \frac{\mathbf{p}_\Delta \mathbf{p}_\Delta^*}{z_\Delta}\quad (3.31)$$

where the first equality follows from (3.24 and (3.25)), see also (8.13) in [18].

Let us write down the unity operator which acts in the Fock space spanned by $\Phi_\alpha | 0 \rangle$:

$$\mathbf{1} = \sum_{\alpha \in \text{DP}} \Phi_\alpha | 0 \rangle 2^{-\ell(\alpha)} \langle 0 | \Phi_\alpha^\dagger = \sum_{\Delta \in \text{OP}} J_\Delta | 0 \rangle \frac{2^{\ell(\Delta)}}{z_\Delta} \langle 0 | J_\Delta^\dagger\quad (3.32)$$

The sets $\{\Phi_\alpha | 0 \rangle, \alpha \in \text{DP}\}$ and $\{J_\Delta | 0 \rangle, \Delta \in \text{OP}\}$ form two different bases in the fermionic Fock space.

3.2.3 Sergeev characters

Now note that the quantity

$$\boxed{\chi_\alpha(\Delta) := 2^{-\ell(\alpha)} \langle 0 | J_\Delta^\dagger \Phi_\alpha | 0 \rangle = 2^{-\ell(\alpha)} \langle 0 | \Phi_\alpha^\dagger J_\Delta | 0 \rangle}\quad (3.33)$$

is nothing but the character of the Sergeev group [19]. To see this, we notice that

$$\begin{aligned}J_\Delta | 0 \rangle &= \sum_\alpha \chi_\alpha(\Delta) \Phi_\alpha | 0 \rangle, & \langle 0 | J_\Delta^\dagger &= \sum_\alpha \chi_\alpha(\Delta) \langle 0 | \Phi_\alpha^\dagger \\ \Phi_\alpha | 0 \rangle &= \sum_\Delta \frac{2^{\ell(\alpha) + \ell(\Delta)}}{z_\Delta} \chi_\alpha(\Delta) J_\Delta | 0 \rangle, & \langle 0 | \Phi_\alpha^\dagger &= \sum_\Delta \frac{2^{\ell(\alpha) + \ell(\Delta)}}{z_\Delta} \chi_\alpha(\Delta) \langle 0 | J_\Delta^\dagger\end{aligned}\quad (3.34)$$

Thus, one has

$$\mathbf{p}_\Delta = \sum_{\alpha} \chi_{\alpha}(\Delta) Q_{\alpha}\{p_k\}, \quad Q_{\alpha}\{p_k\} = \sum_{\Delta} \frac{2^{\ell(\alpha)+\ell(\Delta)}}{z_{\Delta}} \chi_{\alpha}(\Delta) \mathbf{p}_{\Delta} \quad (3.35)$$

This is a counterpart of the Fröbenius formula for the Q Schur functions (see [18, Sec.I.7]), and, hence, $\chi_{\alpha}(\Delta)$ are, indeed, characters of the Sergeev group, [10, 40].

Introduce

$$\mathbf{f}_{\alpha}(\Delta) := 2^{-\ell(\Delta)} \langle 0 | J_{\Delta}^{\dagger} \Phi_{\alpha} | 0 \rangle \frac{1}{z_{\Delta}} \frac{1}{Q_{\alpha}\{\delta_{k,1}\}} = \frac{2^{\ell(\alpha)+\ell(\Delta)}}{z_{\Delta} Q_{\alpha}\{\delta_{k,1}\}} \chi_{\alpha}(\Delta) \quad (3.36)$$

In [17], we used the notation $\Phi_{\alpha}(\Delta)$ for this quantity, it is a basic building block for a spin counterpart of the Fröbenius formula for the Hurwitz numbers, see formula (4.42).

With this quantity,

$$Q_{\alpha}\{p_k\} = Q_{\alpha}\{\delta_{k,1}\} \sum_{\Delta \in \text{OP}} \mathbf{f}_{\alpha}(\Delta) \mathbf{p}_{\Delta} \quad (3.37)$$

4 BKP τ -functions

Any vacuum expectation value of the form

$$\tau(\mathbf{p}, \mathbf{p}^*) = \langle 0 | \gamma(\mathbf{p}) g \gamma^{\dagger}(\mathbf{p}^*) | 0 \rangle \quad (4.1)$$

where $\mathbf{p} = (p_1, p_3, \dots)$ and $\mathbf{p}^* = (p_1^*, p_3^*, \dots)$ are two independent sets of parameters, and

$$g = e^{\sum_{i,k \in \mathbf{Z}} A_{ik} : \phi_i \phi_k :} \quad (4.2)$$

is a BKP τ -function with respect to the set of times \mathbf{p} , and, at the same time, it is a BKP τ -function with respect to the set of times \mathbf{p}^* , hence, it can be called a two-sided BKP, we call it either BKP τ -function or 2BKP τ -function. Here $\{A_{ik}\}_{i,k \in \mathbf{Z}}$ are the elements of a doubly infinite skew symmetric matrix A . The choice of matrix A or, what is the same, the choice of g defines a common solution to all equations of the 2BKP integrable hierarchy.

4.1 Hypergeometric BKP vs KP τ -functions

The term hypergeometric τ -function was introduced in [35] where it was emphasized that the series

$$\sum_R \frac{\prod_{l=1}^p (a_l)_R}{\prod_{m=1}^s (b_m)_R} \cdot \text{Schur}_R(\mathbf{x}) \cdot \text{Schur}_R(\mathbf{y}), \quad (z)_R := \prod_{(i,j) \in R} (z + j - i) \quad (4.3)$$

is a generalized hypergeometric series of two sets of variables \mathbf{x} and \mathbf{y} . This series can be presented in the form

$$\sum_R \text{Schur}_R\{p_k\} \text{Schur}_R\{p_k^*\} e^{\sum_k \frac{1}{k} t_k C_k(R)}, \quad C_k(R) := \sum_i (R_i - i + 1/2)^k - (1/2 - i)^k \quad (4.4)$$

(see, e.g., [42]), where $C_k(R)$ are the eigenvalues of the peculiarly chosen Casimir operators of $GL(\infty)$ in representation R of $SL(N)$ at large enough $N > \ell(R)$. This is a KP τ -function with g of the form [33–35]

$$g = e^{\sum_{n \in \mathbf{Z}} : \psi_n \psi_n^* : \sum_k \frac{1}{k} n^k t_k} \quad (4.5)$$

Quite similarly, the special choice of g ,

$$g^\pm = g^\pm(\mathbf{t}) = e^{\sum_{n>0} (-1)^n \phi_n \phi_{-n} \left(\sum_{k \in \mathbb{Z}_{odd}^+} \frac{1}{k} n^k t_k - \delta^\pm \right)}, \quad \delta^\pm = \frac{i\pi(\pm 1 - 1)}{2}, \quad (4.6)$$

where $\mathbf{t} = (t_1, t_3, \dots)$ is a set of parameters, describes the family of τ -functions called hypergeometric BKP τ -functions [24], [27]. We will denote the related τ -function through $\tau^\pm(\mathbf{p}, \mathbf{p}^* | \mathbf{t})$.

One can verify that

$$g^\pm \phi_i (g^\pm)^{-1} = \pm e^{\sum_{k \in \mathbb{Z}_{odd}^+} \frac{1}{k} i^k t_k} \phi_i, \quad i \neq 0 \quad (4.7)$$

Let us note that $g^\pm |0\rangle = |0\rangle$. From (3.16), (3.30), it follows that

$$\begin{aligned} \tau^\pm(\mathbf{p}, \mathbf{p}^* | \mathbf{t}) &:= \langle 0 | \gamma(\mathbf{p}) g^\pm(\mathbf{t}) \gamma^\dagger(\mathbf{p}^*) | 0 \rangle = \\ &= \sum_{\substack{\alpha \in \text{DP} \\ \ell(\alpha) \text{ even}}} e^{\sum_{k \in \mathbb{Z}_{odd}^+} \frac{1}{k} t_k \omega_k(\alpha)} 2^{-\ell(\alpha)} Q_\alpha(\mathbf{p}) Q_\alpha(\mathbf{p}^*) \pm \sum_{\substack{\alpha \in \text{DP} \\ \ell(\alpha) \text{ odd}}} e^{\sum_{k \in \mathbb{Z}_{odd}^+} \frac{1}{k} t_k \omega_k(\alpha)} 2^{-\ell(\alpha)} Q_\alpha(\mathbf{p}) Q_\alpha(\mathbf{p}^*) : \end{aligned} \quad (4.8)$$

i.e.

$$\boxed{\tau^\pm(\mathbf{p}, \mathbf{p}^* | \mathbf{t}) = R_\pm \cdot \sum_{\alpha \in \text{DP}} e^{\sum_{k \in \mathbb{Z}_{odd}^+} \frac{1}{k} t_k \omega_k(\alpha)} 2^{-\ell(\alpha)} Q_\alpha(\mathbf{p}) Q_\alpha(\mathbf{p}^*)} \quad (4.9)$$

where R_\pm is the projection operator, and

$$\omega_k(\alpha) = \sum_{i=1}^{\ell(\alpha)} \alpha_i^k \quad (4.10)$$

are spin counterparts of the completed cycles in the case of ordinary Hurwitz numbers. Notice the difference between these quantities and (4.4): in the spin case, the main quantities involve the integers α_i , while the same quantities in the non-spin case, $\alpha_i - i$. This is related to the fact that, in the former case, all the formulas involve the strict partitions, and, in the latter one, the formulas involve all partitions.

Remark 4.1. For every hypergeometric BKP τ -function (4.9), its square is a hypergeometric KP τ -function, with all even times put to zero, details see in [24, sec.3, Prop.2].

The issue of the hypergeometric τ -functions can be explained in a different way: the KP τ -function can be generally expanded into the Schur functions,

$$\tau(\mathbf{p}) = \sum_{\alpha} c_{\alpha} \text{Schur}_{\alpha} \{p_k\} \quad (4.11)$$

This linear combination solves the KP hierarchy iff the coefficients satisfy the Plücker relations for the infinite-dimensional Grassmannian [22]

$$c(\vec{x} | \vec{y}) [x_i, x_j; y_i, y_j] \cdot c(\vec{x} | \vec{y}) - c(\vec{x} | \vec{y}) [x_i; y_i] \cdot c(\vec{x} | \vec{y}) [x_j; y_j] + c(\vec{x} | \vec{y}) [x_i; y_j] \cdot c(\vec{x} | \vec{y}) [x_j; y_i] = 0 \quad (4.12)$$

where we used the Fröbenius (hook) parametrization of the Young diagrams $\alpha = (x_1, \dots, x_h | y_1, \dots, y_h) = (\vec{x} | \vec{y})$, where x_i, y_i are the hook arms and legs correspondingly [18], and denoted through $[\{x_i\}; \{y_j\}]$ removing a subset $\{x_i\}; \{y_j\}$ from the set of hook legs and arms.

Now we note that if c_{α} 's solve these relation, so do $c_{\alpha} \prod_{i,j \in \alpha} f(i - j)$ with an arbitrary function $f(x)$. Since a particular solution to the Plücker relations is given by the Schur functions of arbitrary

set of times (parameters), and since the exponential in (4.4) is of the type $\prod_{i,j \in \alpha} f(i-j)$ [42], we finally come to the hypergeometric τ -function (4.4).

Similarly, a linear combination of the Q Schur functions,

$$\tau(\mathbf{p}) = \sum_{\alpha \in \text{DP}} c_{\alpha}^{BKP} Q_{\alpha}\{p_k\} \quad (4.13)$$

solves the BKP hierarchy iff the coefficients satisfy the Plücker relations for the isotropic Grassmannian [22]

$$\begin{aligned} & c_{[\alpha_1, \dots, \alpha_k]}^{BKP} c_{[\alpha_1, \dots, \alpha_k, \beta_1, \beta_2, \beta_3, \beta_4]}^{BKP} - c_{[\alpha_1, \dots, \alpha_k, \beta_1, \beta_2]}^{BKP} c_{[\alpha_1, \dots, \alpha_k, \beta_3, \beta_4]}^{BKP} + \\ & + c_{[\alpha_1, \dots, \alpha_k, \beta_1, \beta_3]}^{BKP} c_{[\alpha_1, \dots, \alpha_k, \beta_2, \beta_4]}^{BKP} - c_{[\alpha_1, \dots, \alpha_k, \beta_1, \beta_4]}^{BKP} c_{[\alpha_1, \dots, \alpha_k, \beta_2, \beta_3]}^{BKP} = 0 \end{aligned} \quad (4.14)$$

Now we again note that if c_{α}^{BKP} 's solve these relation, so do $(\pm 1)^{\ell(\alpha)} e^{\sum_{m \in \mathbb{Z}^+_{\text{odd}}} t_m \omega_m(\alpha)} c_{\alpha}^{BKP}$. Since a particular solution to the Plücker relations is given by the Q Schur functions of arbitrary set of times (parameters), we finally come to the hypergeometric τ -function (4.9).

4.2 Bosonization

Now we need to bosonize the operators that act on the Fock vectors[¶]. The BKP hierarchy and related objects that we will review in this section were introduced in a series of papers by Kyoto school [21], [30], [22]. However, we shall use here the approach due to [43], [44].

Let $z \in S^1$

$$V(z, \hat{\mathbf{p}}) = \frac{1}{\sqrt{2}} D_{\eta} e^{\sum_{m \in \mathbb{Z}^+_{\text{odd}}} \frac{2}{m} z^m p_m} e^{-\sum_{m \in \mathbb{Z}^+_{\text{odd}}} z^{-m} \frac{\partial}{\partial p_m}} \quad (4.15)$$

be the vertex operator as it was introduced in [44]. Here $D_{\eta} = \eta + \frac{\partial}{\partial \eta}$, where η is an auxiliary odd Grassmannian variable: $\eta^2 = 0$, $D_{\eta}^2 = 1$. The symbol $\hat{\mathbf{p}}$ denotes the set of two collection p_1, p_3, p_5, \dots and $\frac{\partial}{\partial p_1}, \frac{\partial}{\partial p_3}, \dots$

Introducing

$$2\theta(z, \mathbf{p}) := \sum_{m \in \mathbb{Z}^+_{\text{odd}}} \frac{2}{m} z^m p_m - \sum_{m \in \mathbb{Z}^+_{\text{odd}}} z^{-m} \frac{\partial}{\partial p_m} = \sum_{m \in \mathbb{Z}_{\text{odd}}} \frac{2}{m} J_m^b z^m \quad (4.16)$$

where

$$J_m^b(\hat{\mathbf{p}}) = \begin{cases} p_m & \text{if } m > 0 \text{ odd} \\ -\frac{m}{2} \frac{\partial}{\partial p_{-m}} & \text{if } m < 0 \text{ odd} \\ 0 & \text{if } m \text{ even,} \end{cases} \quad (4.17)$$

one can rewrite (4.15) as

$$V(z, \mathbf{p}) = \frac{D_{\eta}}{\sqrt{2}} \vdots e^{2\theta(z, \mathbf{p})} \vdots \quad (4.18)$$

where \vdots denotes the so-called bosonic normal ordering which means that all derivatives are moved to the right.

Using

$$[J_m^b, J_n^b] = -\frac{m}{2} \delta_{m+n, 0}, \quad (4.19)$$

[¶]Note that, for the first time, such a bosonization relation was obtained in an unpublished preprint preceding the article [45], the article was not accepted for publication because the result was rather unusual.

one can verify that

$$V(z_1, \mathbf{p})V(z_2, \mathbf{p}) = \frac{1 - z_2 z_1^{-1}}{1 + z_2 z_1^{-1}} : V(z_1, \mathbf{p})V(z_2, \mathbf{p}) : \quad (4.20)$$

which results in the commutation relations

$$V(z_1, \mathbf{p})V(z_2, \mathbf{p}) + V(z_2, \mathbf{p})V(z_1, \mathbf{p}) = \sum_{n \in \mathbb{Z}} \left(-\frac{z_1}{z_2} \right)^n = \delta(\varphi_1 - \varphi_2 - \pi) \quad (4.21)$$

where the right hand side is the Dirac delta function, $z_i = e^{\varphi_i}$, $i = 1, 2$. For the Fourier modes $V(z, \mathbf{p}) = \sum_{i \in \mathbb{Z}} V_i(\mathbf{p})z^i$, one obtains

$$V_n(\mathbf{p})V_m(\mathbf{p}) + V_m(\mathbf{p})V_n(\mathbf{p}) = (-1)^n \delta_{n+m} \quad (4.22)$$

Now one can note that up to the sign factor the fermionic and bosonic currents have the same commutation relations:

$$J_n^b J_m^b - J_m^b J_n^b = J_m J_n - J_n J_m = \frac{m}{2} \delta_{n+m, 0} \quad (4.23)$$

and that the operator

$$\phi(z) := \phi_0 \gamma([z]) \gamma^\dagger(-[z^{-1}]) \quad (4.24)$$

where $[z] := (z, z^3, z^5, \dots)$, has the same commutation relations as the vertex operator (4.15). This describes a correspondence between bosonic and fermionic operators.

4.3 W_∞^B algebra

Following the standard procedure, one can expand the product of the two vertex operators in the generators of W_∞^B algebra:

$$\begin{aligned} & \frac{1}{2} V(ze^{\frac{y}{2}}, \mathbf{p}) V(-ze^{-\frac{y}{2}}, \mathbf{p}) - \frac{1}{2} \frac{1 + e^{-y}}{1 - e^{-y}} = \frac{1}{4} \frac{e^y + 1}{e^y - 1} \left(: e^{\theta(ze^{\frac{y}{2}}) + \theta(-ze^{-\frac{y}{2}})} : - 1 \right) = \\ & = \frac{1}{4} \frac{e^y + 1}{e^y - 1} \sum_{k > 0} \frac{1}{k!} : \left(\sum_{m \in \mathbb{Z}_{\text{odd}}} \theta_m z^m \left(e^{\frac{my}{2}} - e^{-\frac{my}{2}} \right) \right)^k : =: \sum_{m \in \mathbb{Z}, n \geq 0} \frac{1}{n!} y^n z^m \Omega_{mn}(\mathbf{p}) \end{aligned} \quad (4.25)$$

As follows from the left hand side of this formula, Ω_{mn} vanish when n and m have the same parity.

In particular, the operators $\Omega_{m,0} = J_m^b$ with odd m form the bosonic current algebra, $\Omega_{m,1}$ with even m form the Virasoro algebra, etc. Our main interest is the commutative algebra of operators $\Omega_n := \Omega_{0,n}$ with odd n . In particular,

$$\begin{aligned} \Omega_1 &= \sum_{n > 0} n p_n \partial_n \\ \Omega_3 &= \frac{1}{2} \sum_{n > 0} n^3 p_n \partial_n + \frac{1}{2} \sum_{n > 0} n p_n \partial_n + 4 \sum_{n_1, n_2, n_3 \text{ odd}} p_{n_1} p_{n_2} p_{n_3} (n_1 + n_2 + n_3) \partial_{n_1 + n_2 + n_3} + \\ & \quad + 3 \sum_{n_1 + n_2 = n_3 + n_4 \text{ odd}} p_{n_1} p_{n_2} n_3 n_4 \partial_{n_3} \partial_{n_4} + \sum_{n_1, n_2, n_3 \text{ odd}} p_{n_1 + n_2 + n_3} \partial_{n_1} \partial_{n_2} \partial_{n_3} \end{aligned} \quad (4.26)$$

The fermionic counterpart of (4.25) is much simpler:

$$\frac{1}{2} : \phi(ze^{\frac{y}{2}}) \phi(-ze^{-\frac{y}{2}}) : = \frac{1}{2} \sum_{m, j \in \mathbb{Z}} z^m e^{\frac{y}{2}(m+2j)} (-1)^j : \phi_{m+j} \phi_{-j} : = \sum_{\substack{m \in \mathbb{Z} \\ n \geq 0}} \frac{1}{n!} y^n z^m \Omega_{mn}^F \quad (4.27)$$

Again, as follows from the left hand side of this formula, $\Omega_{mn} = 0$ when n and m have the same parity. One gets

$$\Omega_{mn}^F = \frac{1}{2} \sum_{j \in \mathbb{Z}} \left(\frac{m}{2} + j\right)^n (-1)^j : \phi_{m+j} \phi_{-j} : \quad (4.28)$$

In particular, the current algebra:

$$J_m := \Omega_{-m,0}^F = \begin{cases} \frac{1}{2} \sum_{j \in \mathbb{Z}} (-1)^j \phi_{j-m} \phi_{-j} & \text{if } m \text{ odd} \\ 0 & \text{if } m \text{ even,} \end{cases} \quad (4.29)$$

the Virasoro algebra

$$L_m^F := \Omega_{-m,1}^F = \begin{cases} \frac{1}{2} \sum_{j \in \mathbb{Z}} \left(\frac{m}{2} + j\right) (-1)^j (-1)^j \phi_{j-m} \phi_{-j} & \text{if } m \text{ even} \\ 0 & \text{if } m \text{ odd,} \end{cases} \quad (4.30)$$

and

$$\Omega_n^F := \Omega_{0,n}^F = \frac{1}{2} \sum_{j \in \mathbb{Z}} (-1)^j j^n : \phi_j \phi_{-j} : = \sum_{j=1,3,\dots} (-1)^j j^n \phi_j \phi_{-j}, \quad n \text{ odd.} \quad (4.31)$$

One can easily see that

$$[\Omega_n^F, \Omega_m^F] = 0$$

for each pair of n, m .

The boson-fermion correspondence gives rise to the following relation

$$e^{t_{mn} \Omega_{m,n}(\mathbf{p} + t_{m'n'}) \Omega_{m',n'}(\mathbf{p})} \cdot \tau(\mathbf{p}, \mathbf{p}^*) = \langle 0 | \gamma(\mathbf{p}) e^{t_{mn} \Omega_{m,n}^F} g e^{t_{m'n'} \Omega_{m',n'}^F} \gamma^\dagger(\mathbf{p}^*) | 0 \rangle = \tau(\mathbf{p}, \mathbf{p}^* | t_{mn}, t_{m'n'})$$

The flows with respect to the parameters t_{mn} and $t_{m'n'}$ are called additional symmetries, see [46] and [43].

Lemma 4.1. For $r \in \mathbb{Z}_{\geq 0}$ introduce

$$d_{2k,n} := z^{-k} \cdot \left(z \frac{\partial}{\partial z} \right)^n \cdot z^{-k} \quad (4.32)$$

$$\Omega_{2k,n}^F(r) := \operatorname{res}_z (d_{2k,n}^r \cdot \phi(z)) \phi(-z) \frac{dz}{z} = \sum_j \phi_{2rk+j} \phi_{-j} (-1)^j (j+k)^n (j+3k)^n \dots (j+(2r+1)k)^n$$

Then:

- $\Omega_{2k,1}(1) = \Omega_{2k,1}$ and $\Omega_{2k,1}(r) = 0$ if r is even.
- $[\Omega_{2k,n}(r), \Omega_{2k,n}(r')] = 0$.

We define

$$\gamma_{m,n}(\mathbf{p}) := e^{\sum_{r>0, \text{odd}} pr \Omega_{m,n}^F(r)} \quad (4.33)$$

where $\mathbf{p} = (p_1, p_3, \dots)$ is a set of parameters.

It can be shown that the vacuum expectation value

$$\langle 0 | \gamma_{m,n}(\mathbf{p}) g \gamma_{m',n'}(\mathbf{p}^*) | 0 \rangle$$

where $m < 0$ and $m' > 0$ is a BKP τ -function with respect to \mathbf{p} variables and a BKP τ -function with respect to \mathbf{p}^* variables.

Consider the Virasoro element

$$\Omega_{2,1} = L_{-1} = \frac{1}{2} \sum_{j \in \mathbb{Z}} \phi_{2+j} \phi_{-j} (-1)^j (1+j)$$

and the set of related commuting operators

$$\Omega_{2,1}(r) =: L_{-1}(r) = \frac{1}{2} \operatorname{res}_z \left(\underbrace{\left(\frac{\partial}{\partial z} \cdot \frac{1}{z} \right)^r}_{d_{2,1}^r} \phi(z) \right) \phi(-z) \frac{dz}{z} = \quad (4.34)$$

$$= \frac{1}{2} \sum_{j \in \mathbb{Z}} \phi_{2r+j} \phi_{-j} (-1)^j (1+j)(3+j) \cdots (2r+j-1), \quad r = 1, 3, 5, \dots \quad (4.35)$$

We have the following counterpart of (3.29)

Proposition 4.2.

$$\langle 0 | \Phi_\alpha^\dagger \gamma_{2,1}(\mathbf{p}) | 0 \rangle = \begin{cases} Q_\alpha(\mathbf{p}) \prod_i \alpha_i!! & \text{each } \alpha_i \text{ is even} \\ 0 & \text{otherwise} \end{cases} \quad (4.36)$$

The proof is achieved with the observation that $\phi_{-j}(\mathbf{p}) := (\gamma_{2,1}(\mathbf{p}))^{-1} \phi_{-j} \gamma_{2,1}(\mathbf{p})$ is a finite linear combination of the modes ϕ_{-i} where $1 \leq i \leq j$ and where i is odd in case j is an odd positive number. Then, it follows that the presence of the odd parts leads to zero vacuum expectation value. We take into account that $\langle 0 | \gamma_{2,1} = \langle 0 |$ and apply the Wick theorem for $\phi_{\alpha_i}(\mathbf{p})$ and use the definition (3.5) of the projective Schur functions to get the right hand side of (4.36).

Corollary 4.3. Let $g = \exp \sum_{n>0} \phi_i \phi_{-i} \log F(i)$ where F is a function on the lattice $\mathbb{Z}_{>0}$. We have

$$\langle 0 | \gamma(\mathbf{p}) g \gamma_{2,1}(\mathbf{p}^*) | 0 \rangle = \sum_{\alpha \in \text{DP}} Q_{2\alpha}(\mathbf{p}) Q_\alpha(\mathbf{p}^*) \prod_{i=1}^{\ell(\alpha)} F(2\alpha_i) \alpha_i!! \quad (4.37)$$

Corollary 4.4. Let $\gamma_{2,1}^{\text{B}}(\mathbf{p}^*; \hat{\mathbf{p}})$ be the bosonic version of $\gamma_{2,1}(\mathbf{p}^*)$. Then

$$\gamma_{2,1}^{\text{B}}(\mathbf{p}^*; \hat{\mathbf{p}}) \cdot 1 = \sum_{\alpha \in \text{DP}} Q_{2\alpha}(\mathbf{p}) Q_\alpha(\mathbf{p}^*) \prod_{i=1}^{\ell(\alpha)} \alpha_i!! \quad (4.38)$$

Let us note that the right hand side of (4.37) has the form of the right hand side of (1.3) and (1.4).

Remark 4.2. The perturbation series for the partition functions of the well-known $N \times N$ two-matrix model (and therefore also one-matrix model) are obtained from the similar construction [35]. Let us consider the set of operators commuting with the KP Virasoro generator L_{-1} :

$$L_{-1}^{\text{KP}}(r) := \operatorname{res}_z \left(\frac{d^r \psi(z)}{dz^r} \right) \psi^\dagger(z), \quad r = 1, 2, 3, \dots \quad (4.39)$$

Then

$$e^{\sum_{r>0} \frac{1}{r} p_r^* L_{-1}^{\text{Bos}}(r)} \cdot 1 = \sum_{\lambda} (N)_\lambda s_\lambda(\mathbf{p}) s_\lambda(\mathbf{p}^*) \quad (4.40)$$

which is a counterpart of (4.33) for $\gamma_{2,1}(\mathbf{p})$, (4.34), and (4.38).

At the end of this section we note that the action of the bosonic version of $\gamma_{m,n}$ with even n (and odd m) on 1 always gives a version of the BKP hypergeometric tau function.

4.4 Cut-and-join equation

A specific form of the exponential means that the BKP τ -function (4.9) is a generating function of the Hurwitz numbers corresponding to the completed cycles. In the case of the ordinary Hurwitz numbers, one could consider the generating function of the simplest double Hurwitz numbers with two branching profiles fixed and all other ramifications being just double ramification points, $H_d(0|\Delta, \Delta^*, \Gamma_2^b)$. This generating function was a (KP) τ -function [33]. On the contrary, in the spin case, even the simplest generating function of the spin Hurwitz numbers $H^\pm(\Gamma_d, \Delta^1, \Delta^2)$ where

$$\Gamma_d = (3, 1^{d-3}) \quad (4.41)$$

is not a τ -function.

Indeed, the Hurwitz numbers can be represented as [10, 17, 40]

$$H^\pm(\Delta^1, \dots, \Delta^k) = R_\pm \cdot \sum_{\alpha \in \text{DP}} (Q_\alpha \{\delta_{k,1}\})^2 \mathbf{f}_\alpha(\Delta^1) \cdots \mathbf{f}_\alpha(\Delta^k) \quad (4.42)$$

and (see [17, Eq.(102) and derivation in sec.6])

$$\mathbf{f}_\alpha(\Gamma_d) = \frac{1}{3}\omega_3(\alpha) - (\omega_1(\alpha))^2 + \frac{2}{3}\omega_1(\alpha), \quad |\alpha| = d \geq 3 \quad (4.43)$$

One can see that this expression is not a linear combination of the completed cycles $\omega_k(\alpha)$. However, we can still derive an equation for $\tau^\pm(\mathbf{p}, \mathbf{p}^*|\mathbf{t})$, which is a counterpart of the celebrated cut-and-join equation [37].

To this end, with the help of (4.43), we re-write (4.9) as follows

$$\begin{aligned} \tau^\pm(\mathbf{p}, \mathbf{p}^*|\mathbf{t}) &= R_\pm \cdot \sum_{\alpha \in \text{DP}} 2^{-\ell(\alpha)} e^{t_1 d + t_3 \left(\mathbf{f}_\alpha(\Gamma) + d^2 - \frac{2}{3}d \right)} e^{\sum_{n>3, \text{odd}} \frac{1}{n} t_n \omega_n(\alpha)} Q_\alpha \{p_k\} Q_\alpha \{p_k^*\} = \\ &= c \sum_{d \geq 0} e^{t_1(d - \frac{1}{3}) + t_3(d - \frac{1}{3})^2} \Phi_d^\pm(\mathbf{p}, \mathbf{p}^*|t_3, t_5, \dots) \end{aligned} \quad (4.44)$$

where $c = e^{\frac{1}{3}t_1 - \frac{1}{9}t_3}$ and where

$$\Phi_d^\pm(\mathbf{p}, \mathbf{p}^*|t_3, t_5, \dots) = R_\pm \cdot \sum_{\substack{\alpha \in \text{DP} \\ |\alpha|=d}} e^{t_3 \mathbf{f}_\alpha(\Gamma)} 2^{-\ell(\alpha)} e^{\sum_{n>3, \text{odd}} \frac{1}{n} t_n \omega_n(\alpha)} Q_\alpha \{p_k\} Q_\alpha \{p_k^*\} \quad (4.45)$$

Let us put $t_i = 0, i > 3$. Then, we get

$$\Phi_d^\pm(\mathbf{p}, \mathbf{p}^*|t_3) = \sum_{\substack{\Delta^1, \Delta^2 \\ |\Delta^1|=|\Delta^2|}} \sum_{b \geq 0} \frac{t_3^b}{b!} H^\pm(\Gamma_d^b, \Delta^1, \Delta^2) \mathbf{p}_{\Delta^1} \mathbf{p}_{\Delta^2}^* \quad (4.46)$$

where

$$H^\pm(\Gamma_d^b, \Delta^1, \Delta^2) = R_\pm \cdot \sum_{\substack{\alpha \in \text{DP} \\ |\alpha|=d}} (Q_\alpha \{\delta_{k,1}\})^2 (\mathbf{f}_\alpha(\Gamma))^b \mathbf{f}_\alpha(\Delta^1) \mathbf{f}_\alpha(\Delta^2) \quad (4.47)$$

Since the BKP τ -function is defined up to a constant factor c , we obtain

Theorem 4.5. Multiply the BKP τ -function (4.9) with the factor $\frac{1}{c} = e^{-\frac{1}{3}t_1 + \frac{1}{9}t_3}$. Then, one gets the cut-and-join equation in form

$$\left(\frac{\partial}{\partial t_3} - \left(\frac{\partial}{\partial t_1} \right)^2 \right) \cdot \tau^\pm(\mathbf{p}, \mathbf{p}^*|\mathbf{t}) = \mathcal{W} \cdot \tau^\pm(\mathbf{p}, \mathbf{p}^*|\mathbf{t}) \quad (4.48)$$

or, which is the same

$$\frac{\partial \Phi_d^\pm(\mathbf{p}, \mathbf{p}^* | t_3)}{\partial t_3} = \mathcal{W} \cdot \Phi_d^\pm(\mathbf{p}, \mathbf{p}^* | t_3) \quad (4.49)$$

where the cut-and-join operator

$$\begin{aligned} \mathcal{W} = & \frac{1}{3}\Omega_3(\mathbf{p}) - \Omega_1(\mathbf{p})\Omega_1(\mathbf{p}) + \frac{2}{3}\Omega_1(\mathbf{p}) = - \left(\sum_{n>0} n p_n \partial_n \right)^2 + \left(\frac{2}{3} + \frac{1}{6} \right) \sum_{n>0} n p_n \partial_n + \\ & + \frac{1}{6} \sum_{n>0} n^3 p_n \partial_n + \frac{4}{3} \sum_{n_1, n_2, n_3 \text{ odd}} p_{n_1} p_{n_2} p_{n_3} (n_1 + n_2 + n_3) \partial_{n_1+n_2+n_3} + \\ & + \sum_{n_1+n_2=n_3+n_4 \text{ odd}} p_{n_1} p_{n_2} n_3 n_4 \partial_{n_3} \partial_{n_4} + \frac{1}{3} \sum_{n_1, n_2, n_3 \text{ odd}} p_{n_1+n_2+n_3} \partial_{n_1} \partial_{n_2} \partial_{n_3} \end{aligned} \quad (4.50)$$

Thus, the cut-and-join equation has a form of the quantum heat equation with the potential \mathcal{W} .

By analogy with the case of the ordinary Hurwitz numbers analyzed by Boris Dubrovin in [47], we have

Remark 4.3. Each $\Omega_n(\mathbf{p})$ can be treated as a Hamiltonian of the quantum dispersionless modified KdV equation on the circle with the eigenstates given by $Q_\alpha(\mathbf{p})$ and the eigenvalues given by $\omega_n(\alpha)$. If one introduces the Plank constant \hbar and puts $v = \hbar^{-\frac{1}{2}} \sum_n e^{in\varphi} J_n^b$, then the first nontrivial Hamiltonian $\hbar^2 \Omega_3$ is

$$\mathcal{H}_3 = \int_0^{2\pi} d\varphi \dot{ : } \left(v^4 + \hbar (v_\varphi)^2 \right) \dot{ : } d\varphi \quad (4.51)$$

5 KdV soliton solution as the BKP solutions

5.1 Solitons of the KdV and BKP hierarchies which generate Hurwitz numbers

So far we discussed integrable properties with respect to time variables \mathbf{p} that are the variables of the Q Schur functions. In this subsection, we discuss a relation to the soliton solution in t_n . That is, we want to show that if we fix both sets \mathbf{p}, \mathbf{p}^* so that they are equal to $\mathbf{p}_1 := (1, 0, 0, \dots)$, then $\tau^\pm(\mathbf{p}_1, \mathbf{p}_1 | \mathbf{t})$ turns into a recognizable soliton τ -function of the KdV equation, where the role of times is played by the set $\mathbf{t} = (t_1, t_3, \dots)$.

Theorem 5.1. Consider the τ -function (4.44) where we restrict the times to be $\mathbf{p} = \mathbf{p}^* = \mathbf{p}_1$ and introduce

$$u^\pm(\mathbf{t}) = 2 \frac{\partial^2}{\partial t_1^2} \log \tau^\pm(\mathbf{p}_1, \mathbf{p}_1 | \mathbf{t}) \quad (5.1)$$

Then each u^\pm is the (∞) -soliton solution of the KdV hierarchy with respect to the times $\mathbf{t} = (t_1, t_3, \dots)$. In particular,

$$12u_{t_3}^\pm = u_{t_1 t_1 t_1}^\pm + 6u_{t_1}^\pm u_{t_1}^\pm \quad (5.2)$$

If one puts all $t_i = 0$ for $i > 3$ and denote $(t_1, t_3) = (x, t)$, so that (x, t) are the space-time coordinates in the standard KdV theory [48], then

$$\tau^\pm(\mathbf{p}_1, \mathbf{p}_1 | x + \log q, t, 0, 0, \dots) = c \sum_{d \geq 3} q^d e^{xd+t(d^2 - \frac{2}{3}d)} \sum_{b \geq 0} \frac{t^b}{b!} H^\pm \left(\Gamma_d^b \right) \quad (5.3)$$

Proof. This solution is well-known: it is a multi-soliton τ -function. Indeed, as it follows from (4.9), (3.9) we have

$$\tau^\pm(\mathbf{p}_1, \mathbf{p}_1 | \mathbf{t}) = 1 + \sum_{k=1}^{\infty} \sum_{1 \leq \alpha_1 < \dots < \alpha_k} 2^{2|\alpha| - k} \prod_{i < j \leq k} \left(\frac{\alpha_i - \alpha_j}{\alpha_i + \alpha_j} \right)^2 \prod_{i=1}^k \frac{e^{\eta(\alpha_i, \mathbf{t}) + \delta^\pm}}{(\alpha_i!)^2} \quad (5.4)$$

where δ^\pm is defined in (4.6) and where

$$\eta(x, \mathbf{t}) := \sum_{m=1,3,5,\dots} \frac{1}{m} t_m x^m \quad (5.5)$$

which is a special case of the general soliton solution of KdV hierarchy given by

$$\tau_{\text{KdV}}^{\text{sol}\pm}(\mathbf{t}) = 1 + \sum_{k>0} \sum_{\substack{\alpha \in \text{DP} \\ \ell(\alpha)=k}} (\Delta^*(\zeta_\alpha))^2 \prod_{i=1}^k e^{\eta(\alpha_i, \mathbf{t}) + a_i - \delta^\pm} \quad (5.6)$$

$$= 1 + \sum_i e^{\eta_i^\pm} + \sum_{i < j} \frac{(\zeta_i - \zeta_j)^2}{(\zeta_i + \zeta_j)^2} e^{\eta_i^\pm + \eta_j^\pm} + \sum_{i < j < k} \frac{(\zeta_i - \zeta_j)^2 (\zeta_i - \zeta_k)^2 (\zeta_j - \zeta_k)^2}{(\zeta_i + \zeta_j)^2 (\zeta_i + \zeta_k)^2 (\zeta_j + \zeta_k)^2} e^{\eta_i^\pm + \eta_j^\pm + \eta_k^\pm} + \dots \quad (5.7)$$

where $\eta_j^\pm := \eta(\zeta_j, \mathbf{t}) + a_j - \delta^\pm$ and

$$\Delta^*(\zeta_\alpha) := \prod_{i < j \leq k} \frac{\zeta_{\alpha_i} - \zeta_{\alpha_j}}{\zeta_{\alpha_i} + \zeta_{\alpha_j}} \quad (5.8)$$

We recall that the parameters ζ_j play the role of the soliton momenta, the parameters a_j define initial positions of solitons, these are the solitons $u^+ \sim \frac{1}{2}\zeta_j^2 \text{ch}^{-2}(\frac{1}{2}\zeta_j x + \frac{1}{6}\zeta_j^3 t + \frac{1}{2}a_j)$ for the δ^+ case, and the solitons $u^- \sim \frac{1}{2}\zeta_i^2 \sinh^{-2}(\frac{1}{2}\zeta_i x + \frac{1}{6}\zeta_i^3 t + \frac{1}{2}a_i)$ for the δ^- case. Factors (5.8) describe interactions of solitons.

Choosing $\zeta_j = j$, $j = 1, 2, 3, \dots$ and $a_j = -\log 2^{1-2j} (j!)^2$, and taking into account (3.9), (4.9) one obtains (5.4). Formula (5.3) translates the sum over the number of solitons in the formula (5.7) into a sum over momenta $d = \sum_k \alpha_k$, d playing the role of the number of sheets in the covering problem related to $H(\Gamma_d^b)$.

Remark 5.1. To generate the Hurwitz numbers for the d -sheeted coverings, it is enough to consider the N -soliton KdV τ -function with the parameters ζ_i filling the string $0, 1, \dots, d$ where d does not exceed the number of solitons N . Such a τ -function is holomorphic in the (x, t) -variables, and one can write

$$H^\pm(\Gamma_d^b) = (2\pi)^{-1} \text{res}_{t=0} t^{-b-1} e^{-t(d^2 + \frac{2}{3}d)} \int_0^{2\pi} dx e^{-ixd} \tau_N^{\text{KdV}, \pm}(ix, t) \quad (5.9)$$

A similar solitonic τ -function can be obtained at $\mathbf{p}^* = \delta_{k,1} := \mathbf{p}_1$: $\tau(\mathbf{p}, \mathbf{p}_1 | \mathbf{t})$ is the τ -function of the two component BKP hierarchy [44] (or of the Veselov-Novikov hierarchy) with respect to the times $\mathbf{p} = (p_1, p_3, \dots)$ and $\mathbf{t} = (t_1, t_3, \dots)$.

Remark 5.2. Two component τ -functions are constructed with the two-component fermions. We recall that any two-component BKP (2-BKP) τ -function is a BKP τ -function in $\mathbf{p} = (p_1, p_3, \dots)$ variables, and it is also a BKP τ -function in $\mathbf{t} = (t_1, t_3, \dots)$ variables, see [22], [44].

Theorem 5.2. The τ -function (4.44) $\tau^\pm(\mathbf{p}, \mathbf{p}_1 | \mathbf{t})$ that generates the Hurwitz numbers is a two-component BKP multisoliton τ -function with respect to the time sets \mathbf{t} and \mathbf{p} .

For the proof, it suffices to represent $\tau^\pm(\mathbf{p}, \mathbf{p}_1|\mathbf{t})$ in the form of an appropriate vacuum expectation value. Indeed, this 2-BKP (Veselov-Novikov) τ -function $\tilde{\tau}^{\text{VN}\pm}(\mathbf{p}, \mathbf{t})$ is of the following solitonic type

$$\tilde{\tau}^{\text{VN}\pm}(\mathbf{p}, \mathbf{t}) = \langle 0|\gamma^{(1)}(\mathbf{t})\gamma^{(2)}(\mathbf{p})g^\pm(0)e^{\sqrt{-1}\sum_{i\geq 0}C_{ij}\phi^{(1)}(\zeta_i)\phi_j^{(2)}}|0\rangle \quad (5.10)$$

$$= \sum_{\alpha, \beta \in \text{DP}} 2^{-\frac{1}{2}\bar{\ell}(\alpha) - \frac{1}{2}\bar{\ell}(\beta)} C_{\alpha, \beta} e^{\eta(\zeta_{\alpha_i}, \mathbf{t}) - \delta^\pm} \Delta^*(\zeta_\alpha) Q_\beta\{\mathbf{p}\} \quad (5.11)$$

where the symbol $g^\pm(0)$ is given by (4.6) where each ϕ_i is replaced by $\phi_i^{(2)}$, and where $\eta(\zeta_{\alpha_i}, \mathbf{t})$ and $\Delta^*(\zeta_\alpha)$ are given respectively by (5.5) and by (5.8), the set $\{\zeta_i\}$ is a set of free parameters (soliton momenta), and a free chosen matrix C defines an interaction between solitons (the net of lines in the (t_1, t_3) -plane in the tropical approximation of the τ -function). The series (5.11) where $C_{\alpha, \beta} = \det(C)_{\alpha_i, \beta_j}$ is obtained by direct calculation of the vacuum expectation value in (5.10) taking into account (3.19) and (3.29). Then, choosing $C_{j,0} = 2^j \frac{1}{j!} \delta_{j,i}$, $i > 0$, $C_{j,0} = 2\delta_{j,0}$ and putting $\zeta_j = j$, one gets $\tilde{\tau}^{\text{VN}\pm}(\mathbf{p}, \mathbf{t}) = \tau^\pm(\mathbf{p}, \mathbf{p}_1|\mathbf{t})$ as it follows from (4.9) and (3.9).

When all $t_i = 0$ for $i > 3$, this τ -function generates the Hurwitz numbers $H^\pm(\Gamma_d^r, \Delta)$:

$$\tau^\pm(\mathbf{p}, \mathbf{p}_1|\mathbf{t}) = \sum_{d \geq 0} e^{t_1 d + t_3 \left(d^2 - \frac{2}{3}d\right)} \sum_{\substack{\Delta \\ |\Delta|=d}} \sum_{b \geq 0} \frac{t_3^b}{b!} H^\pm(\Gamma_d^b, \Delta) \mathbf{p}_\Delta \quad (5.12)$$

Such a τ -function describes “the net of resonant solitons” (it is either the net of regular solitons or the net of singular solitons).[†]

Similarly to the previous case, one gets

Remark 5.3. To generate the Hurwitz numbers $H^\pm(\Gamma_d^b, \Delta)$, it is enough to consider the N -soliton τ -function with the parameters ζ_i which fill the string $0, 1, \dots, d$ where d does not exceed the number of solitons N . Such a τ -function is holomorphic in the (x, t) -variables, and one can write

$$\sum_{\substack{\Delta \\ |\Delta|=d}} H^\pm(\Gamma_d^b, \Delta) \mathbf{p}_\Delta = (2\pi)^{-1} \operatorname{res}_{t=0} t^{-b-1} e^{-t\left(d^2 + \frac{2}{3}d\right)} \int_0^{2\pi} dx e^{-ixd} \tau_N^{\text{VN}, \pm}(ix, t) \quad (5.13)$$

The superscript “-” denotes the singular soliton solution, while “+” is related to the regular one.

5.2 Different fermionic expressions generating Hurwitz numbers

Besides formula (4.8) and the example discussed in (5.10), there are many others representations of the series (4.9) in terms of the neutral fermions. These are different embeddings of a given function of many variables into various families of τ -functions. Here are some more examples. In these examples, $\eta(\alpha_i, \mathbf{t})$ and $\Delta^*(\zeta_\alpha)$ are given respectively by (5.5) and by (5.8).

Lemma 5.3. *The multisoliton BKP τ -function can be written as the following vacuum expectation value:*

$$\tau_1(\mathbf{t}) = \langle 0|\gamma(\mathbf{t})e^{\sum_{i>j\geq 0}A_{ij}\phi(\zeta_i)\phi(\zeta_j)}|0\rangle = \sum_{\alpha \in \text{DP}} 2^{-\frac{1}{2}\bar{\ell}(\alpha)} A_\alpha \Delta^*(\zeta_\alpha) \prod_{i=1}^{\ell(\alpha)} e^{\eta(\zeta_{\alpha_i}, \mathbf{t})} \quad (5.14)$$

where $A_\alpha = \text{Pf} \bar{A}$, and \bar{A} is the skew symmetric $\bar{\ell}(\alpha) \times \bar{\ell}(\alpha)$ submatrix of the matrix A whose entries are selected by the parts of $\alpha = (\alpha_1, \dots, \alpha_k)$ as

$$\bar{A}_{i,j} := A_{\alpha_i, \alpha_j}, \quad i, j \leq k \quad (5.15)$$

[†]The best presentation of such nets of regular solitons in the KP case is given in [49]. Singular resonant solitons were considered in [50]. Networks of singular solitons will be described elsewhere.

when k is even, and

$$\bar{A}_{i,j} := \begin{cases} A_{\alpha_i, \alpha_j}, & i, j \leq k \\ A_{\alpha_i, 0}, & i \leq k, j = k+1 \end{cases} \quad (5.16)$$

when k is odd. For any choice of A , one gets the multisoliton BKP τ -function (which describes the net of resonant solitons with the momenta given by the set $\{\zeta_i\}$). For $\zeta_i = i$, one obtains the BKP multisoliton τ -function with integer momenta.

As an example, one can choose $A_{ij} = \frac{1}{2}Q_{Ni, Nj}\{\mathbf{p}\}F(i)F(j)$, $j > 0$ and $A_{i,0} = Q_{Ni,0}\{\mathbf{p}\}F(i)$. Then, the τ -function (5.14) takes the form

$$\tau_1(\mathbf{t}, \mathbf{p}) = \sum_{\alpha \in \text{DP}} 2^{-\ell(\alpha)} \Delta^*(\zeta_\alpha) Q_{N\alpha}\{\mathbf{p}\} \prod_{i=1}^{\ell(\alpha)} e^{\eta(\zeta_{\alpha_i}, \mathbf{t})} F(\alpha_i) \quad (5.17)$$

The right hand side of (5.17) where $F(i) = \pm 1$ can be equated to the τ -function (5.10) if we choose $C_{i,j} \sim \delta_{Ni,j}$ in (5.11).

If we put $p_k = \delta_{k,1}$ and $F(i) = \pm \frac{(Ni)!}{i!}$, we get KdV τ -function (5.4).

Lemma 5.4. *Another example of the BKP τ -function (A_α is the same as in the previous Lemma) is given by*

$$\tau_2(\mathbf{p}) = \langle 0 | \gamma(\mathbf{p}) e^{\sum_{i>j \geq 0} A_{ij} \phi_i \phi_j} | 0 \rangle = \sum_{\alpha} 2^{-\frac{1}{2}\bar{\ell}(\alpha)} A_\alpha Q_\alpha\{\mathbf{p}\} \quad (5.18)$$

If we choose A to be the same as in example (5.17), the τ -function (5.18) takes the form

$$\tau_2(\mathbf{p}, \mathbf{p}^*) = \sum_{\alpha \in \text{DP}} 2^{-\ell(\alpha)} Q_\alpha\{\mathbf{p}\} Q_{N\alpha}\{\mathbf{p}^*\} \prod_{i=1}^{\ell(\alpha)} F(\alpha_i) \quad (5.19)$$

which is of the form considered in sec.7 below. For $N = 1$, (5.19) is the hypergeometric τ -function given by (4.9) if we choose $F(j) = \pm \exp \sum_m j^m t_m$.

Lemma 5.5. *An example of the 2-component BKP (2-BKP) τ -function is given by*

$$\tau_3(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}) = \langle 0 | \gamma^{(1)}(\mathbf{p}^{(1)}) \gamma^{(2)}(\mathbf{p}^{(2)}) e^{\sqrt{-1} \sum_{i,j \geq 0} C_{ij} \phi_i^{(1)} \phi_j^{(2)}} | 0 \rangle = \sum_{\alpha, \beta \in \text{DP}} 2^{-\frac{1}{2}\bar{\ell}(\alpha) - \frac{1}{2}\bar{\ell}(\beta)} C_{\alpha, \beta} Q_\alpha\{\mathbf{p}^{(1)}\} Q_\beta\{\mathbf{p}^{(2)}\} \quad (5.20)$$

where $C_{\alpha, \beta} = \det(C)_{\alpha_i, \beta_j}$.

For instance, one can choose $C_{ij} = F(j)\delta_{i, Nj}$, $j > 0$, $C_{i,0} = 2\delta_{i,0}$ in order to get once again the series at the right hand side of (5.19).

Lemma 5.6. *Another example of the 2-BKP τ -function is given by*

$$\begin{aligned} \tau_4(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}) &= \langle 0 | \gamma^{(1)}(\mathbf{p}^{(1)}) \gamma^{(2)}(\mathbf{p}^{(2)}) e^{\sqrt{-1} \sum_{i,j \geq 0} C_{ij} \phi^{(1)}(\zeta_i) \phi^{(2)}(\zeta_j)} | 0 \rangle \\ &= \sum_{\alpha, \beta \in \text{DP}} 2^{-\frac{1}{2}\bar{\ell}(\alpha) - \frac{1}{2}\bar{\ell}(\beta)} C_{\alpha, \beta} \prod_{i=1}^{\ell(\alpha)} e^{\eta(\zeta_{\alpha_i}, \mathbf{p}^{(1)})} \prod_{i=1}^{\ell(\beta)} e^{\eta(\zeta_{\beta_i}, \mathbf{p}^{(2)})} \Delta_\alpha^*(\zeta^{(1)}) \Delta_\beta^*(\zeta^{(2)}) \end{aligned} \quad (5.21)$$

$$= \sum_{\alpha, \beta \in \text{DP}} 2^{-\frac{1}{2}\bar{\ell}(\alpha) - \frac{1}{2}\bar{\ell}(\beta)} C_{\alpha, \beta} \prod_{i=1}^{\ell(\alpha)} e^{\eta(\zeta_{\alpha_i}, \mathbf{p}^{(1)})} \prod_{i=1}^{\ell(\beta)} e^{\eta(\zeta_{\beta_i}, \mathbf{p}^{(2)})} \Delta_\alpha^*(\zeta^{(1)}) \Delta_\beta^*(\zeta^{(2)}) \quad (5.22)$$

To get the KdV multisoliton τ -function, it is enough to take $\zeta_i^{(1)} = \zeta_i^{(2)}$, $i \geq 0$. To get the τ -function (5.4), one takes $\zeta_i = i$ and $C_{ij} = 2^{2j}(j!)^{-2}\delta_{i,j}$, $j > 0$, $C_{i,0} = C_{0,i} = 2\delta_{i,0}$.

The examples of the KdV and Veselov-Novikov soliton τ -functions considered above can be presented as a specification of τ_1, \dots, τ_4 .

Remark 5.4. In all cases, the sums in the exponentials can be replaced by integrals with appropriate measures (to get specializations of the hypergeometric τ -functions discussed in sec.5.1, one needs to choose the Γ -function and the contour going around the real semi-axis in the complex plane).

Any particular fermionic representation is convenient for writing down the string equations, which are stationary points for special combinations of W_∞^B symmetries discussed in sec.4.3.

5.3 Matrix model τ -functions as fermion averages

The partition functions of the matrix models in external field considered in the Introduction, (1.3) and (1.4), which are hypergeometric BKP τ -functions can be considered as particular cases of described fermion averages.

The τ -function of the BGW model in (1.4) can be presented as examples considered in sec.5.2 by a proper specification of parameters:

Proposition 5.7.

$$\tau_{BGW}(\mathbf{p}) = \sum_{\alpha \in \text{DP}} 2^{-\ell(\alpha)} Q_\alpha\{\mathbf{p}\} Q_\alpha\{\delta_{k,1}\} \prod_{i=1}^{\ell(\alpha)} \left(\frac{(2\alpha_i)!}{\alpha_i!} \right)^2 \quad (5.23)$$

$$\begin{aligned} &= \langle 0 | \gamma(\mathbf{p}) e^{\sum_{n>0} \log\left(\frac{(2n)!}{n!}\right)^2 (-1)^n \phi_n \phi_{-n}} \gamma\{\delta_{k,1}\} | 0 \rangle = \\ &\langle 0 | \gamma(\mathbf{p}) e^{\sum_{i>j \geq 0} \frac{1}{2} \left(\frac{(2i)!(2j)!}{i!j!}\right)^2 \tilde{Q}_{i,j} \phi_i \phi_j} | 0 \rangle = \langle 0 | \gamma^{(1)}(\mathbf{p}) \gamma^{(2)}\{\delta_{k,1}\} e^{\sqrt{-1} \sum_{i \geq 0} C_i \left(\frac{(2i)!}{i!}\right)^2 \phi_i^{(1)} \phi_i^{(2)}} | 0 \rangle \end{aligned}$$

where we used (3.9). Here $\tilde{Q}_{i,j} = Q_{i,j}\{\delta_{k,1}\} = 2^{i+j} \frac{1}{i!j!} \frac{i-j}{i+j}$; $\tilde{Q}_{i,0} = 2Q_{i,0}\{\delta_{k,1}\} = 2^{i+1} \frac{1}{i!}$ and $C_i = 1$, $i > 0$; $C_0 = 2$

Similarly, the τ -function of the Kontsevich model (1.3) can be presented as as examples considered in sec.5.2:

Proposition 5.8. The series (1.3) can be treated both as the BKP and as the 2-BKP τ -function presented as a fermion average

$$\tau_{K_3}\{\mathbf{p}\} = \sum_{\alpha \in \text{DP}} 2^{-\ell(\alpha)} Q_\alpha\{\mathbf{p}\} Q_{2\alpha}\{\delta_{k,3}\} \prod_{i=1}^{\ell(\alpha)} \frac{(2\alpha_i)!}{\alpha_i!} = \quad (5.24)$$

$$= \langle 0 | \gamma(\mathbf{p}) e^{\sum_{i>j \geq 0} \frac{1}{2} \frac{(2i)!(2j)!}{i!j!} \tilde{Q}_{2i,2j} \{\delta_{k,3}\} \phi_i \phi_j} | 0 \rangle = \langle 0 | \gamma^{(1)}(\mathbf{p}) \gamma^{(2)}\{\delta_{k,3}\} e^{\sqrt{-1} \sum_{i \geq 0} C_i \frac{(2i)!}{i!} \phi_i^{(1)} \phi_{2i}^{(2)}} | 0 \rangle$$

where $\tilde{Q}_{2i,2j} = Q_{2i,2j}\{\delta_{k,3}\}$ and $\tilde{Q}_{2i,0} = 2Q_{2i,0}\{\delta_{k,3}\}$ and where $C_i = 1$, $i > 0$ and $C_0 = 2$.

5.4 On character expansion for KdV τ -functions and BKP

Since the bilinear Plücker relations (4.12) does not contain terms c_α^2 for any diagram α , one can put all the coefficients $c_\alpha = 0$ but one. Hence, any individual Schur function solves the Plücker relations, and is a KP τ -function. For a similar reason, any individual Q Schur function is a τ -function of the BKP hierarchy.

If one applies the Schur function expansion, (4.11) to the KdV hierarchy, which is a KP reduction, one immediately realizes that an individual Schur function (i.e. a single non-vanishing c_α) solves the hierarchy only for the diagrams $[\dots, 4, 3, 2, 1]$. This is not surprising, since the KdV τ -function

does not depend on even time variables p_{2k} , and so do only these Schur functions: $\text{Schur}_{[1]} = p_1$ or $\text{Schur}_{[2,1]} = \frac{1}{3}(-p_3 + p_1^3)$, etc.

On the other hand, the KdV solutions simultaneously solve the BKP hierarchy, and one could expect that a natural character expansion of the KdV hierarchy is in terms of the Q Schur functions (4.13). Of course, this is formally true, because the Q Schur functions form a complete basis in the space of functions of odd times p_{2k+1} , but the restrictions on the coefficients c_α substituting the Plücker relations in KP and BKP cases, is more involved. In particular, most individual Q -functions are not the KdV τ -functions: exceptions are provided just by $Q_{\dots,4,3,2,1}$, which coincide with peculiar ordinary Schur functions, depending on odd times only as discussed above.

Note that the reduction BKP \rightarrow KdV equation has the simplest form when one uses the 2-BKP approach and imposes the condition

$$\left(\frac{\partial}{\partial p_1^{(1)}} - \frac{\partial}{\partial p_1^{(2)}} \right) \tau(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}) = 0 \quad (5.25)$$

and the KdV higher times can be identified with $\mathbf{p}^{(1)} + \mathbf{p}^{(2)}$. It can be also formulated in terms of the Lax operator for the Veselov-Novikov equation, which is a 2D Schrödinger operator (with a reduction 2D Schrödinger \rightarrow 1D Schrödinger operator the latter being the Lax operator for the KdV equation). Another way is to present the KdV hierarchy as the so-called rational reduction of the BKP hierarchy. For some other aspects of relation between the KdV and BKP hierarchies, see [21, 22, 30–32].

6 Factorization on special loci

6.1 Specialization at $p_k = \delta_{k,r}$

The locus $p_k = \delta_{k,1}$ is well known to play a big role in representation theory and character calculus. For instance, in the case of Schur polynomial S_R , this locus is associated with the dimension $|R|! S_R\{\delta_{k,1}\}$ of representation R of the permutation group $S_{|R|}$ and reflects the Schur-Weyl duality [51]. In this case, this quantity has a special notation, $d_R := S_R\{\delta_{k,1}\}$. Somewhat unexpectedly superintegrability relations in the Gaussian matrix models are sensitive also to the values at other loci $p_k = \delta_{k,r}$ [28]. This gets a natural explanation in the study of monomial matrix models, where the Cauchy identity states

$$e^{-\frac{1}{r} \text{Tr} X^r} = \exp \left(- \sum_k \frac{1}{k} \text{Tr} X^k \cdot \delta_{k,r} \right) = \sum_R (-1)^{|R|} S_{R^\vee} \{ \text{Tr} X^k \} \cdot S_R \{ \delta_{k,r} \} \quad (6.1)$$

This makes the study of character values at these delta-loci very important. These values are known to be distinguishably factorizable, and very recently A.Alexandrov [29] conjectured an explicit formula for the ratio of two Q Schur functions, Q_R/Q_{2R} at $p_k = \delta_{k,3}$. In this section, we describe a general and explicit factorization formula for Q_R/Q_{NR} at any N and $p_k = \delta_{k,r}$ (hence, r is odd), and even more general factorization formula for Q_R at only $p_{k,rj}$ non-vanishing (r and j odd). We prove these formulas in Appendices A and B.

The relation important for us is

$$Q_\alpha \{ \delta_{k,r} \} = Q_{N\alpha} \{ \delta_{k,r} \} \prod_{i=1}^{\ell(\alpha)} F(\alpha_i) \quad (6.2)$$

with coprime N and r and with some function $F(\alpha_i)$ that we describe below.

In order to explain (6.2), we list a set of facts about specializations of the symmetric functions at $p_k = \delta_{k,r}$.

- For the Schur functions, the values at $p_k^{(r)} = \delta_{k,r}$ are equal to

$$S_R\{\delta_{k,r}\} = \delta_r(R) \prod_{x \in R} \frac{1}{[h_x]_{0,r}} \quad (6.3)$$

where h_x is the hook length and $[n]_{p,r}$ is defined to be n when n is equal to $p \bmod r$, and to be 1 otherwise. $\delta_r(R)$ is defined in [52, Eq.(3.26)]:

$$\delta_r(R) = \begin{cases} (-1)^{|R|/r} \prod_{x \in R} (-1)^{[c_x/r] + [h_x/r]} & \text{if the } r\text{-core of } R \text{ is trivial} \\ 0 & \text{otherwise} \end{cases} \quad (6.4)$$

where c_x is the content of the box in R .

- Similarly, for the Q Schur functions, the values at $p_k^* = \delta_{k,r}/2$ are equal to

$$Q_\alpha\{\delta_{k,r}/2\} = \tilde{\delta}_r(\alpha) \prod_{x \in S(\alpha)} \frac{1}{[h_x^{(d)}]_{0,r}} \quad (6.5)$$

where $h_x^{(d)}$ is the hook length in the doubled Young diagram, which is defined to be $d(\alpha) := (\alpha_1, \alpha_2, \dots | \alpha_1 - 1, \alpha_2 - 1, \dots)$ in the Fröbenius notation [18], while $S(\alpha)$ denotes here the shifted Young diagram [18], which is the part of $d(\alpha)$ lying above the main diagonal. The sign factor $\tilde{\delta}_r(\alpha)$ vanishes if the r -core of $d(\alpha)$ is non-trivial.

- Suppose we are given a strict partition α such that the size of α is divisible by r , and $\tilde{\delta}_r(\alpha) \neq 0$ (this simultaneously implies that $\tilde{\delta}_r(N\alpha) \neq 0$), i.e. the r -core of $d(\alpha)$ is trivial. Suppose also that N and r are coprime (r is certainly odd). Then, the following formula is correct:

$$\boxed{\frac{Q_\alpha\{\frac{r}{2} \cdot \delta_{k,r}\}}{Q_{N\alpha}\{\frac{r}{2} \cdot \delta_{k,r}\}} = \prod_{i=1}^{\ell(\alpha)} (-1)^{\rho_{N,r}(\alpha_i)} \cdot N^{\{\alpha_i/r\}} \cdot \frac{[N\alpha_i/r]!}{[\alpha_i/r]!}} \quad (6.6)$$

where $\{\dots\}$ at the r.h.s. denotes the fractional part of a number, and $[\dots]$ denotes the integer part.

We derive formula (6.6) in Appendix A, here just point out that the integer-valued function $\rho_{N,r}(x)$ depends only on $(x)_r$ (the value of $x \bmod r$), and on $(Nx)_r$ (the value of $Nx \bmod r$). Manifestly, $\rho_{N,r}(0) = 0$, and all other x_r enters the product in pairs $\rho_{N,r}(k) + \rho_{N,r}(r - k)$ so that there is no difference which sign to choose for an individual $(-1)^{\rho_{N,r}(x)}$ in the pair. For the sake of definiteness, let us choose $(-1)^{\rho_{N,r}(k)} = 1$ with $r \geq k > r/2$. Then one has

$$\rho_{N,r}(x) = \begin{cases} (Nx)_r - (x)_r & \text{for } 0 < (x)_r < r/2 \\ 0 & \text{otherwise} \end{cases} \quad (6.7)$$

In fact, there is a more fundamental factorization formula for p_k non-vanishing only at k divisible by r , (6.9), and formula (6.6) is its straightforward corollary in a particular case. We discuss this in the next subsection.

6.2 Basic factorization formula

Now we consider a more general factorization formula in the case, when an infinite set of times t_{kr} is non-vanishing. That is, we introduce a set of times $\mathbf{p}[r] := (p_1[r], p_3[r], \dots)$ such that

$$\frac{1}{k} p_k[r] = \frac{1}{j} p_j \delta_{k,jr}, \quad \text{both } r, j \text{ odd} \quad (6.8)$$

Let us consider a strict partition α , and produce r new strict partitions μ, a^c, b^c made of $[\alpha_i/r]$. Parting into these $r - 1$ partitions depends on the value $(\alpha_i)_r = x$: the parts with $x = 0$ get to partition μ , the parts with $0 < x \leq (r - 1)/2$ get to partitions $a^c, c = x$, and those with $(r - 1)/2 < x < r$ get to partitions $b^c, c = r - x$. Thus, the partition a^c at each color c has an associated partition b^c .

Thus, the parts of α are parted into three groups:

- parts $r\mu$ that are divisible by r
- parts presented as $ra^c + c$ where $c = 1, \dots, \frac{1}{2}(r - 1)$
- parts presented as $r(b^c + 1) - c$ where $c = 1, \dots, \frac{1}{2}(r - 1)$

Suppose that $|\alpha|$ is divisible by r and that the length of partitions a^c coincides with those of b^c (otherwise, $Q_\alpha\{p_k[r]\} = 0$). We denote this length through κ^c . Then, there is a beautiful factorization formula (which we derive in Appendix B)

$$Q_\alpha\{p_k[r]\} = (-1)^\omega 2^{-\frac{1}{2}\bar{\ell}(\mu)} Q_\mu\{p_k\} \prod_{c=1}^{\frac{1}{2}(r-1)} S_{(a^c|b^c)}\{2p'_k\} \cdot \prod_{i=1}^{\kappa^c} (-1)^{a_i^c + b_i^c + c} \quad (6.9)$$

where $p'_k := (p_1, 0, p_3, 0, p_5, \dots)$, $S_{(a^c|b^c)}$ is the ordinary Schur function in the Fröbenius (hook) notation, and ω depends on the order of embedded parts which belong to one of the three groups. Basically, it is not important for our purposes because we will be interested in rescaling of lengths of the parts $\alpha_i \rightarrow N\alpha_i$ which keeps the order, and we get the same ω .

For example, for $r = 3$ and a partition, say, $\alpha = [6, 5, 4, 3, 2, 1]$, $\mu = [2, 1]$, $a^1 = [1, 0]$, $b^1 = [1, 0]$ so that eq.(6.9) states:

$$Q_{[6,5,4,3,2,1]}\{0, 0, 3p_1, 0, 0, 0, 0, 0, 3p_3, \dots\} \sim Q_{[2,1]}\{p_1, p_3\} \cdot \underbrace{S_{(1,0|1,0)}}_{S_{[2,2]}}\{2p_1, 0, 2p_3, 0\} \quad (6.10)$$

Similarly, for $r = 3$ and a partition $\alpha = [15, 7, 6, 5, 2, 1]$, $\mu = [5, 2]$, $a^1 = [2, 0]$, $b^1 = [1, 0]$ so that eq.(6.9) states:

$$Q_{[15,7,6,5,2,1]}\{0, 0, 3p_1, 0, 0, 0, 0, 0, 3p_3, \dots\} \sim Q_{[5,2]}\{p_1, p_3, p_5, p_7\} \cdot \underbrace{S_{(2,0|1,0)}}_{S_{[3,2]}}\{2p_1, 0, 2p_3, 0, p_5\} \quad (6.11)$$

The basic factorization formula (6.9) immediately leads to formula (6.6), see Appendix A for details.

7 Hypergeometric τ -functions entirely made from characters

In this section, we describe an important construction of particular hypergeometric τ -functions from the ratios of characters at the special loci $p_k = \delta_{k,r}$ described in the previous section. It is actually applicable in a more general context, but we present it in the case of the Q Schur functions and the BKP hierarchy.

The relations (6.2), (6.6) implies that the bilinear combination (1.1), which is a BKP τ -function can be specified in the form

$$\tau_{BKP}\{p, \bar{p}\} = \sum_{\alpha \in \text{DP}} Q_\alpha\{p\} Q_{N\alpha}\{\delta_{k,r}\} \cdot \prod_{i=1}^{\ell(\alpha)} f(\alpha_i) \quad (7.1)$$

Indeed, the restrictions for α in (6.6) implies that $Q_{N\alpha} \neq 0$ and, hence, are satisfied, since otherwise α does not contribute to the sum (7.1).

In its turn, this means that not only the partition function of the Kontsevich model, [28]

$$\tau_{K_3}\{p_k\} = \sum_{\alpha \in \text{DP}} \frac{1}{2^{\ell(\alpha)}} \cdot \frac{Q_\alpha\{p_k\} Q_\alpha\{\delta_{k,1}\} Q_{2\alpha}\{\delta_{k,3}\}}{Q_{2\alpha}\{\delta_{k,1}\}} \quad (7.2)$$

which is of form (7.1) with $N = 2$, $r = 3$ [29], is a BKP τ -function with the weight function given by

$$\prod_{i=1}^{\ell(\alpha)} f(\alpha_i) = \frac{Q_\alpha\{\delta_{k,3}\}}{Q_{2\alpha}\{\delta_{k,3}\}} \frac{Q_\alpha\{\delta_{k,1}\}}{Q_{2\alpha}\{\delta_{k,1}\}} \quad (7.3)$$

but a more general combination with arbitrary coprime N and r is still a BKP τ -function.

From (6.6), it also follows that one can choose it as a product of various ratios $\frac{Q_\alpha\{\delta_{k,r}\}}{Q_{N\alpha}\{\delta_{k,r}\}}$ still preserving the BKP τ -function:

$$\tau_{BKP}\{p, \bar{p}\} = \sum_{\alpha \in \text{DP}} Q_\alpha\{p\} Q_\alpha\{\delta_{k,r}\} \prod_i \frac{Q_\alpha\{\delta_{k,r_i}\}}{Q_{N_i\alpha}\{\delta_{k,r_i}\}} \prod_j \frac{Q_{N_j\alpha}\{\delta_{k,r_j}\}}{Q_\alpha\{\delta_{k,r_j}\}} \quad (7.4)$$

with arbitrary sets of (coprime) N_i , r_i , and N_j , r_j , however, one has to deal carefully with vanishing of $Q_\alpha\{\delta_{k,r_i}\}$ (either choosing proper sets of r_i 's, or continuing formula (6.6) to all sets of α_i by definition).

This means that one can construct non-trivial hypergeometric BKP τ -functions completely in terms of characters. The cubic Kontsevich and BGW partition functions belong to this class, which once again emphasizes **the intimate connection between the hypergeometric and matrix model partition functions**, i.e. between hypergeometricity and string equations.

Since the Virasoro algebra acts on the Q Schur functions in a very simple way [53]:

$$\begin{aligned} \hat{L}_n Q_\alpha \left\{ \frac{p_k}{\sqrt{2}} \right\} &= \sum_{i=1}^{\ell(\alpha)} (-)^{\nu_i} (\alpha_i - n) Q_{\alpha - 2n\epsilon_i} \left\{ \frac{p_k}{\sqrt{2}} \right\}, \\ \hat{L}_n &:= \sum_{k \in \mathbb{Z}_{\text{odd}}^+} (k+n)p_k \frac{\partial}{\partial p_{k+n}} + \frac{1}{2} \sum_{a,b \in \mathbb{Z}_{\text{odd}}^+}^{a+b=2n} ab \frac{\partial^2}{\partial p_a \partial p_b} \end{aligned} \quad (7.5)$$

where $\alpha - 2k\epsilon_i$ denotes the shift of $\alpha_i \rightarrow \alpha_i - 2k$. This can make it shorter than some other lines and thus imply reordering of lines in the diagram to put them back into decreasing order, then $\nu_i(\alpha, b)$ is the number of lines, which the i -th one needs to jump over. It is possible to check that (1.3) and (1.4) satisfy the Virasoro constraints (after an appropriate rescaling of time variables). This illustrates the general claim of [54] that superintegrability of matrix models, which underlies these expressions implies both the ordinary integrability and the Virasoro constraints: the two basic properties of matrix model partition functions [14]. From the very beginning, it was clear that they are intimately related [55], but a nature of this relation remained obscure. Now we understand that they are just two different corollaries of a more fundamental superintegrability feature $\langle \text{character} \rangle = \text{character}$.

8 Conclusion

In this paper, we made a brief review of the details behind the formalism of Q Schur functions, which can be relevant for deeper investigation of the spin Hurwitz partition functions. We put a special emphasize on the ‘‘hypergeometric’’ τ -functions. They are made from Casimir exponentials and give rise to peculiarly-factorized coefficients in the character expansions, which is typically associated with (generalized) hypergeometric series, hence the name. There is a mounting evidence that matrix model τ -functions (i.e. those which satisfy additional string equations) belong to this class. Until recently, this was not so obvious, because important matrix models were not brought to this form, but recently

the reason has been found: the τ -functions for the Kontsevich and BGW models are expanded in the Q Schur functions rather than in the ordinary Schur functions. A posteriori, this is rather obvious because they satisfy KdV rather than the generic KP/Toda hierarchy. Moreover, now it is clear that the generalized Kontsevich model is likewise expressed through appropriately generalized Q Schur functions [56]. In this paper, we, however, concentrated on the standard Q Schur functions and perspectives of spin Hurwitz studies.

Acknowledgements. We are grateful to A. Alexandrov for various discussions and for sharing his results prior to publication. A. O. thanks J. van de Leur and J. Harnad for numerous discussions on topics related to the BKP hierarchy. The publication was partly prepared by Sergey Natanzon within the framework of the Academic Fund Program at the National Research University Higher School of Economics (HSE) in 2020-2021 (grant 20-01-009) and by the Russian Academic Excellence Project “5-100”. A. O. was supported by RFBR grant 18-01-00273a. Our work is also supported in part by the grant of the Foundation for the Advancement of Theoretical Physics “BASIS” (A.Mir.), by RFBR grants 19-01-00680 (A.Mir.) and 19-02-00815 (A.Mor.), by joint grants 19-51-50008-YaF-a (A.Mir.), 19-51-53014-GFEN-a (A.Mir., A.Mor.), 21-52-52004-MNT-a (A.Mir., A.Mor.). The work was also partly funded by RFBR and NSFB according to the research project 19-51-18006 (A.Mir., A.Mor.).

A Derivation of (6.6) from (6.9)

The basic factorization formula (6.9) allows us to obtain formula (6.6) immediately with the following logic. First of all, the basic factorization formula (6.9) expresses the Q Schur functions, Q_R at $p_k[r]$ through those at p_k , these two being related by (6.8). In particular, under the choice $p_k[r] = r\delta_{k,r}$, it expresses the Q Schur functions at $\delta_{k,r}$ through those at $\delta_{k,1}$. Hence, it remains to evaluate the Q Schur functions at $p_k = \delta_{k,1}$, and then insert the obtained results into (6.9) in order to obtain finally (A.9). This is what is done in this Appendix in three steps. The factorization formula itself is proved in Appendix B.

Step I. (6.9) explicitly at $p_k = r\delta_{k,r}/2$. In order to obtain formula (6.6), we evaluate (6.9) at $p_k = r\delta_{k,r}/2$. Using (3.9), we get for individual factors in (6.9) in this case:

$$Q_\mu \left\{ \frac{1}{2} \cdot \delta_{k,1} \right\} = \prod_{i=1}^{\kappa^0} \frac{1}{\mu_i!} \prod_{i < j} \frac{\mu_i - \mu_j}{\mu_i + \mu_j} \quad (\text{A.1})$$

$$S_{(a^c|b^c)}\{\delta_{k,1}\} = \frac{1}{\prod_{i=1}^{\kappa^c} a_i^c! b_i^c!} \frac{\prod_{i < j} (a_i^c - a_j^c)(b_i^c - b_j^c)}{\prod_{i,j} (a_i^c + b_j^c + 1)} \quad (\text{A.2})$$

Step II. (6.9) for $N\alpha$. Now we evaluate (6.9) for $N\alpha$. To this end, we notice that $N\alpha$ consists of the following parts:

- parts $Nr\mu$ that are still divisible by r
- parts presented as $Nra^c + Nc$ where $Nc = N, \dots, \frac{1}{2}N(r-1)$
- parts presented as $Nr(b^c + 1) - Nc$ where $Nc = N, \dots, \frac{1}{2}N(r-1)$

For each c there exists p_c and c_N

$$Nc = rp_c + c_N, \quad c_N < r \quad (\text{A.3})$$

Let

$$N < r, \quad N \text{ and } r \text{ are coprime} \quad (\text{A.4})$$

Equation (A.3) maps each c to a certain c_N . If c and c' are different, then $c_N(c)$ can not coincide with either $r - c_N(c')$ or $c_N(c')$. Indeed, suppose it is not correct, and one has the same c_N for two different c (c and c'):

$$N(c + c') = r(p_c + p'_c + r) \Rightarrow c + c' = \frac{r}{N}(p_c + p'_c + 1)$$

However, this is impossible because of (A.4) and because both c and c' are less than $\frac{1}{2}r$. Similarly impossible is

$$N(c - c') = r(p_c - p'_c) \Rightarrow c - c' = \frac{r}{N}(p_c - p'_c)$$

Notice that the partition associated by c is also associated by c_N :

$$\begin{aligned} (a^c, b^c) &= (ra_i^c + c, r(b_j^c + 1) - c) \rightarrow (Nra_i^c + Nc, Nr(b_j^c + 1) - Nc) = \\ &= (r(Na_i^c + p_c) + c_N, r(Nb_j^c - p_c) - c_N) \end{aligned} \quad (\text{A.5})$$

Thus, one finally obtains

$$Na = r(Na_i + p_i) + c_N, \quad Nb = r(N(b_i + 1) - p_i) - c_N \quad (\text{A.6})$$

Therefore, one gets

$$S_{(Na^c|Nb^c)}\{\delta_{k,1}\} = \frac{N^{-\kappa^c}}{\prod_{i=1}^{\kappa^c} (Na_i^c + p_i)!(Nb_i^c - p_i - 1)!} \frac{\prod_{i < j} (a_i^c - a_j^c)(b_i^c - b_j^c)}{\prod_{i,j=1}^{\kappa^c} (a_i^c + b_j^c + 1)} \quad (\text{A.7})$$

and

$$Q_{N\mu} \left\{ \frac{1}{2} \cdot \delta_{k,1} \right\} = \frac{1}{\prod_{i=1}^k (N\mu_i)!} \prod_{i < j} \frac{\mu_i - \mu_j}{\mu_i + \mu_j} \quad (\text{A.8})$$

Step III. Evaluating the ratio (6.6). Now, using (A.1)-(A.2) and (A.7)-(A.8), we are ready to evaluate the ratio of the Q Schur functions in (6.6):

$$\frac{Q_\alpha \left\{ \frac{r}{2} \cdot \delta_{k,r} \right\}}{Q_{N\alpha} \left\{ \frac{r}{2} \cdot \delta_{k,r} \right\}} = (-1)^g \left(\prod_{i=1}^{\ell(\mu)} \frac{(N\mu_i)!}{\mu_i!} \right)^{\frac{1}{2}(r-1)} \prod_{c=1}^{\frac{1}{2}(r-1)} N^{\kappa^c} \prod_{i=1}^{\kappa^c} \frac{(Na_i^c + p_c)!(Nb_i^c - p_c - 1)!}{a_i^c! b_i^c!} = (-1)^g N^{\frac{1}{2}v} \prod_{i=1}^{\ell(\alpha)} \frac{[N\alpha_i/r]!}{[\alpha_i/r]!} \quad (\text{A.9})$$

where v is the number of parts of α that are not divisible by r , and g originates from the sign factors at the r.h.s. of (6.9) and is equal to

$$g = \sum_{c=1}^{\frac{1}{2}(r-1)} \kappa^c (c - c_N(c)) \quad (\text{A.10})$$

where $c_N(c)$ is given by (A.3). Since this sign factor coincides with (6.7) and $v = 2 \sum_{i=1}^{\ell(\alpha)} \{\alpha_i/r\}$, we finally come to formula (6.6).

B Factorization formula (6.9) from fermion calculus

Here we prove (6.9) using the fermion average representation for the Q Schur functions (3.29). To this end, we consider the fermionic representation for the Q Schur function (3.29)

$$Q_\alpha \{p_k[r]\} = 2^{\bar{\ell}(\alpha)} \langle 0 | \gamma(\mathbf{p}[r]) \phi_{\alpha_1} \cdots \phi_{\alpha_\ell} | 0 \rangle = 2^{\bar{\ell}(\alpha)} \langle 0 | \phi_{\alpha_1}(\mathbf{p}[r]) \cdots \phi_{\alpha_\ell}(\mathbf{p}[r]) | 0 \rangle \quad (\text{B.1})$$

$$\phi_k(\mathbf{p}[r]) := \gamma(\mathbf{p}[r]) \cdot \phi_{\alpha_1} \cdot \gamma(\mathbf{p}[r])^{-1}$$

Using the canonical anticommutation relation (3.16), we have

$$[J_m, \phi_i] = \phi_{i-m}, \quad m \text{ odd} \quad (\text{B.2})$$

and

$$\phi_j(\mathbf{p}[r]) := e^{\sum_{m>0, \text{odd}} \frac{2}{nr} J_{mr} p_{mr}[r]} \phi_j e^{-\sum_{m>0, \text{odd}} \frac{2}{nr} J_{mr} t_{mr}[r]} = \sum_{m \geq 0} \phi_{j-mr} h_m \{2p'_k\} \quad (\text{B.3})$$

where h_i are complete symmetric functions restricted on the set of odd labeled times:

$$e^{\sum_{n>0, \text{odd}} \frac{2}{n} p_n z^n} = \sum_{n \geq 0} z^n h_n \{2p'_k\}, \quad h_n \{2p'_k\} := S_{(n)} \{2p'_k\} \quad (\text{B.4})$$

Let us note that the exponential (B.4) is also a generating function for the elementary projective Schur functions $Q_{n,0}$:

$$h_n \{2p'_k\} = Q_{(n,0)} \{p_k\} \quad (\text{B.5})$$

For evaluation of the averages at the r.h.s of (B.1), we use the Wick theorem. To this end, we consider the pairwise average:

$$-\langle 0 | \phi_a(\mathbf{p}[r]) \phi_b(\mathbf{p}[r]) | 0 \rangle = \langle 0 | \phi_b(\mathbf{p}[r]) \phi_a(\mathbf{p}[r]) | 0 \rangle = \langle 0 | \sum_{m \geq 0} \phi_{b-mr} h_m \{2p'_k\} \sum_{n \geq 0} \phi_{a-nr} h_n \{2p'_k\} | 0 \rangle \quad (\text{B.6})$$

where, according to the canonical pairing (3.18), contribute only the terms with

$$a - nr + b - mr = 0 \quad (\text{B.7})$$

This implies distributing the parts of the strict partition into three groups μ , a^c and b^c as above.

Remark B.1. One may say that there exist a part α_x such that $a_i^c + p_c = [\alpha_x/r]$ and $c = (\alpha_x)_r$, and a part α_y , with $[\alpha_y/r] = b_j^c - p_c - 1$ and $(\alpha_y)_r = -c$. In this case, $\langle 0 | \phi_{\alpha_x}(\mathbf{p}[r]) \phi_{\alpha_y}(\mathbf{p}[r]) | 0 \rangle \neq 0$.

Similarly, one may say that there exist a part α_x such that $N a_i^c + p_c = [N \alpha_x/r]$ and $c_N = (N \alpha_x)_r$, and a part α_y , with $[N \alpha_y/r] = N b_j^c - p_c - 1$ and $(N \alpha_y)_r = -c_N$. In this case, $\langle 0 | \phi_{N \alpha_x}(\mathbf{p}[r]) \phi_{N \alpha_y}(\mathbf{p}[r]) | 0 \rangle \neq 0$.

One obtains for a pair of a^c and b^c of the same color c (we will denote here $a = r a^c + c$, $b = r b^c + r - c$) that (B.6) is equal to

$$\begin{aligned} & \langle 0 | \phi_{-c} \phi_c | 0 \rangle h_{b^c+1} \{2p'_k\} h_{a^c} \{2p'_k\} + \cdots + \langle 0 | \phi_{-a} \phi_a | 0 \rangle h_{a^c+b^c+1} \{2p'_k\} h_0 \{2p'_k\} \\ & = (-1)^c h_{b^c+1} \{2p'_k\} h_{a^c} \{2p'_k\} + \cdots + (-1)^{a+1} h_{a^c+b^c} \{2p'_k\} h_1 \{2p'_k\} + (-1)^a h_{a^c+b^c+1} \{2p'_k\} h_0 \{2p'_k\} \end{aligned} \quad (\text{B.8})$$

For instance, for $r = 3$ (i.e. there is the single color $c = 1 = \frac{1}{2}(r - 1)$), if taking $a^1 = a^2 = 0$, i.e. $a = 1$, $b = 2$, one obtains

$$\langle 0 | \phi_2(\mathbf{p}[3]) \phi_1(\mathbf{p}[3]) | 0 \rangle = \langle 0 | \phi_{-1} \phi_1 | 0 \rangle h_1 \{2p'_k\} h_0 \{2p'_k\} = (-1)^1 h_1 \{2p'_k\} h_0 \{2p'_k\} = -2p_1 \quad (\text{B.9})$$

Now notice that, since r, j are odd, one has from (B.4)

$$h_i \{2p'_k\} = (-1)^i h_i \{-2p'_k\} \quad (\text{B.10})$$

therefore (B.8) is written as

$$(-1)^{c+a^c} h_{b^c+1} \{2p'_k\} h_{a^c} \{-2p'_k\} + \cdots + (-1)^a h_{a^c+b^c} \{2p'_k\} h_1 \{-2p'_k\} + (-1)^a h_{a^c+b^c+1} \{2p'_k\} h_0 \{-2p'_k\} \quad (\text{B.11})$$

(the parity of $a = r a^c + c$ is equal to that of $a^c + c$ because r is odd).

We compare (B.11) with the one-hook Schur function [18]

$$S_{(j|k)} \{2p'_k\} = (-1)^k \sum_{i=0}^k h_{j+i+1} \{2p'_k\} h_{k-i} \{-2p'_k\} \quad (\text{B.12})$$

and obtain that

$$\langle 0 | \gamma(\mathbf{p}[r]) \phi_{r b^c + c^*} \phi_{r a^c + c} | 0 \rangle = (-1)^{b^c + r a^c + c} S_{(a^c | b^c)} \{2p'_k\} = (-1)^{b^c + a^c + c} S_{(a^c | b^c)} \{2p'_k\} \quad (\text{B.13})$$

The case of sub-partition μ can be considered in a similar way. For a pairwise averages, using (B.3) in the same way as before, and using (B.5) and (3.1), one gets

$$\langle 0 | \phi_{r\mu}(\mathbf{p}[r]) \phi_{r\nu}(\mathbf{p}[r]) | 0 \rangle = \langle 0 | \sum_{m \geq 0} \phi_{\mu-mr} h_m \{2p'_k\} \sum_{n \geq 0} \phi_{\nu-nr} h_n \{2p'_k\} | 0 \rangle = 2^{-1} Q_{(\mu, \nu)} \{p_k\} \quad (\text{B.14})$$

(here μ, ν are numbers). The Pfaffian of the Wick theorem yields the projective Schur function labeled by the partition μ .

As one can see, after re-numbering, the neutral fermions from complementary groups a^c and b^c are quite similar to the charged fermions (3.14), while the neutral fermions in the group μ up to re-numbering the Fourier modes remain to be neutral (3.18) inside average (B.1).

Finally, applying the Wick theorem to evaluate averages with all three groups of fermions, one obtains (6.9).

C Multicomponent KP and BKP τ -functions

Following [22], one can define two-component charged fermions by $\psi_j^{(i)} = \psi_{2j+i}$, $\psi_j^{\dagger(i)} = \psi_{2j+i}^\dagger$, where $i = 1, 2$ is the component number. Due to (3.12), we see that the fermions with different components anticommute. For each given component, one can construct the neutral fermions labeled with the same component number via (3.15). The neutral fermions related to different components anticommute. We construct currents labeled with the component number and the corresponding evolution operators,

$$\gamma^{(i)}(\mathbf{p}^i) = e^{\sum_{k>0, \text{ odd}} \frac{1}{k} p_k^{(i)} J_k^{(i)}}, \quad J_k^{(i)} = \sum_{n \in \mathbb{Z}} (-1)^n \phi_{-k-n}^{(i)} \phi_n^{(i)}, \quad i = 1, 2 \quad (\text{C.1})$$

Then any vacuum expectation value

$$\tau(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}) = \langle 0 | \gamma^{(1)}(\mathbf{p}^{(1)}) \gamma^{(2)}(\mathbf{p}^{(2)}) e^{\mathcal{A}} | 0 \rangle \quad (\text{C.2})$$

where \mathcal{A} is quadratic in the neutral fermions is a two-component BKP τ -function, and the sets $\mathbf{p}^{(1)}$ and $\mathbf{p}^{(2)}$ are called the sets of 2-BKP higher times. There is an important

Remark C.1. If one fixes the set $\mathbf{p}^{(2)}$ (the set $\mathbf{p}^{(1)}$), then $\tau(\mathbf{p}^{(1)}, \mathbf{p}^{(2)})$ can be considered as a one-component BKP τ -function with respect to the set $\mathbf{p}^{(1)}$ (to the set $\mathbf{p}^{(2)}$).

Details may be found in [22] or in [44].

References

- [1] B.A. Dubrovin, A.T. Fomenko, S.P. Novikov, *Modern Geometry— Methods and Applications: Part I. The Geometry of Surfaces, Transformation Groups, and Fields* (Springer-Verlag New York, 1984); Part II: The Geometry and Topology of Manifolds (Springer Science and Business Media, 2012)
- [2] A. Hurwitz, Math. Ann. **38** (1891) 452–45
- [3] G. Frobenius, Berl. Ber. **1896** (1896) 985–1021
- [4] W. Burnside, *Theory of groups of finite order*, 2nd edition, Cambridge Univ. Press (1911)
- [5] R.Dijkgraaf, In: *The moduli spaces of curves*, Progress in Math., 129 (1995), 149-163
- [6] V.Knizhnik, Sov.Phys.Usp. **32** (1989) 945 (in Russian Edition: vol.159, p.451)
- [7] A.Alexandrov, A.Mironov and A.Morozov, Physica **D235** (2007) 126-167, hep-th/0608228
B. Eynard and N. Orantin, Commun. Num. Theor. Phys. **1** (2007) 347-452, math-ph/0702045
- [8] A.Alexandrov, A.Mironov and A.Morozov, JHEP **12** (2009) 053, arXiv:0906.3305
- [9] A. Eskin, A. Okounkov, R. Pandharipande, Adv. Math. **217** (2008) 873-888
- [10] S. Gunningham, Geom. Topol. **20** (2016) 1859-1907, arXiv:1201.1273
- [11] M. Kontsevich, Commun. Math. Phys. **147** (1992) 1
- [12] S. Kharchev, A. Marshakov, A. Mironov, A. Morozov, A. Zabrodin, Phys. Lett. **B275** (1992) 311, hep-th/9111037
S. Kharchev, A. Marshakov, A. Mironov, A. Morozov, A. Zabrodin, Nucl. Phys. **B380** (1992) 181, hep-th/9201013
- [13] A.Mironov, A.Morozov and G.Semenoff, Int.J.Mod.Phys., **A10** (1995) 2015, hep-th/9404005
- [14] A. Morozov, Phys.Usp.(UFN) **37** (1994) 1; hep-th/9502091; hep-th/0502010
A. Mironov, Int.J.Mod.Phys. **A9** (1994) 4355; Phys.Part.Nucl. **33** (2002) 537; hep-th/9409190
- [15] L.Chekhov and Yu.Makeenko, Phys.Lett. **B278** (1992) 271-278, hep-th/9202006; Mod.Phys.Lett. **A7** (1992) 1223-1236, hep-th/9201033
- [16] S. Kharchev, A. Marshakov, A. Mironov, A. Morozov, Nucl.Phys. **B397** (1993) 339-378, hep-th/9203043
- [17] A.D. Mironov, A. Yu. Morozov, S. Natanzon, Eur. Phys. J. C **80** (2020) 97, arXiv:1904.11458
- [18] I.G. Macdonald, *Symmetric functions and Hall polynomials*, Second Edition, Oxford University Press, 1995

- [19] A. Sergeev, Math. Sb. USSR, **51** (1985) 419–427
- [20] A.Mironov, A.Morosov, S.Natanzon, J. Geom. Phys. **62** (2012) 148–155
- [21] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, Physica **4D** (1982) 343-365
- [22] M. Jimbo and T. Miwa, Publ. Res. Inst. Math. Sci. **19** (1983) 943-1001
- [23] Y. You, in: *Infinite-Dimensional Lie Algebras and Groups*, Adv. Ser. Math. Phys. **7** (1989). World Sci. Publ., Teaneck, NJ
- [24] A. Yu. Orlov, Theor. Math. Phys. **137** (2003) 1574-1589
- [25] A. Mironov, A. Morozov, S. Natanzon, Theor.Math.Phys.**166** (2011) 1-22, arXiv:0904.4227
- [26] A. Mironov, A. Morozov and A. Sleptsov, Eur. Phys. J. C **73** (2013) 2492, arXiv:1304.7499
- [27] J. J. C. Nimmo, A. Yu. Orlov Glasgow Mathematical Journal **47 (A)** (2005) 149-168
- [28] A. Mironov and A. Morozov, arXiv:2011.12917
- [29] A. Alexandrov, arXiv:2012.07573
- [30] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, “Transformation groups for soliton equations”, In: *Non-linear integrable systems - classical theory and quantum theory*, 39-120. World Scientific (Singapore), eds. M. Jimbo and T. Miwa (1983)
- [31] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, Publ. RIMS, Kyoto Univ. **18** (1982) 1077-1110
- [32] A. Alexandrov, arXiv:2012.10448
- [33] A.Okounkov, Math. Res. Lett. **7** (2000) 447–45
- [34] S. Kharchev, A. Marshakov, A. Mironov, A. Morozov, Int.J.Mod.Phys. **A10** (1995) 2015, hep-th/9312210
- [35] A. Orlov, D.M. Shcherbin, Theor.Math.Phys. **128** (2001) 906-926
- [36] A. Alexandrov, A. Mironov, A. Morozov and S. Natanzon, J. Phys. **A45** (2012) 045209, arXiv:1103.410
- [37] D.Goulden , D.M.Jackson, A.Vainshtein A., Ann. of Comb. **4** (2000) 27-46
- [38] D.Mumford, Ann. Sci. Ecole Norm. Sup. (4) **4** (1971) 181–192
- [39] M. Atiyah, Ann. Sci. Ecole Norm.Sup. (4) **4** (1971) 47–6
- [40] Junho Lee, arXiv:1407.0055
- [41] J. Harnad, J. W. van de Leur and A Yu. Orlov, Theor. Math. Phys. **168** (2011) 951-962
- [42] A. Alexandrov, A. Mironov, A. Morozov and S. Natanzon, JHEP **11** (2014) 080, arXiv:1405.1395
- [43] J. W. van de Leur, arxiv:9411159
- [44] V. Kac and J. van de Leur, CRM Proceedings and Lecture Notes **14** (1998) 159–202
- [45] A. K. Pogrebkov and V. N. Sushko, *Quantization of the $(\sin\psi)_2$ interaction in terms of fermion variables*, Translated from Teoreticheskaya i Matematicheskaya Fizika, **24** (1975) 425-429 (September, 1975). The original paper was submitted on May 15, 1975
- [46] A. Yu. Orlov, “Vertex operator, $\bar{\partial}$ -problem, symmetries, variational identities and Hamiltonian formalism for 2+1 D integrable systems”. Nonlinear and Turbulent Processes in Physics 1987, ed. V. Baryakhtar., V.E.Zakharov —Singapore (1988)
- [47] B.A. Dubrovin, *Symplectic field theory of a disk, quantum integrable systems, and Schur polynomials*, arxiv:1407.5824
- [48] S. V. Manakov, S. P. Novikov, L. P. Pitaevskii and V. E. Zakharov, *Theory of Solitons*, ed. S. P. Novikov, Nauka 1979
- [49] Yuuji Kodama, J. Phys. **A37** (2004) 11169-11190
- [50] A. Yu. Orlov, *Soliton collapse in integrable models*, preprint EAiE N 221 Novosibirsk (1983)
- [51] W. Fulton, J. Harris, *Representation theory. A first course*, Springer-Verlag, 1991

- [52] C. Cordova, B. Heidenreich, A. Papolitov, S. Shakirov, Commun. Math. Phys. **361** (2018) 1235, arXiv:1611.03142
- [53] K. Aokage, E. Shinkawa, HF. Yamada, Lett.Math.Phys. **110**(2020) 1381–1389
- [54] A. Mironov, A. Morozov, Phys. Lett. **B771** (2017) 503, arXiv:1705.00976
A. Mironov, A. Morozov, JHEP **1808** (2018) 163, arXiv:1807.02409
- [55] M. Fukuma, H. Kawai and R. Nakayama, Int. J. Mod. Phys. **A6** (1991) 1385-1406
- [56] A. Mironov, A. Morozov, to appear