

CLASSIFICATION AND A PRIORI ESTIMATES FOR THE SINGULAR PRESCRIBING Q -CURVATURE EQUATION ON 4-MANIFOLD

MOHAMEDEN AHMEDOU, LINA WU, AND LEI ZHANG

ABSTRACT. On (M, g) a compact riemannian 4-manifold we consider the prescribed Q -curvature equation defined on M with finite singular sources. We first prove a classification theorem for singular Liouville equations defined on \mathbb{R}^4 and perform a concentration compactness analysis. Then we derive a quantization result for bubbling solutions and establish a priori estimate under the assumption that certain conformal invariant does not take some quantized values. Furthermore we prove a spherical Harnack inequality around singular sources provided their strength is not an integer. Such an inequality implies that in this case singular sources are *isolated simple blow up points*.

Key Words: Blow-up analysis, Conical metrics, Moving plane method, Q -curvature, Spherical Harnack inequality.

AMS subject classification: 53C21, 35C60, 58J60.

CONTENTS

1. Introduction	1
2. Entire solutions of fourth order singular Liouville type equations in \mathbb{R}^4	7
2.1. Asymptotic behavior of entire solutions	7
2.2. Classification of entire solutions for the case $-1 < \gamma < 0$	18
3. Preliminaries for blowup analysis	22
4. Blow-up analysis near the singularity	24
5. Concentration-Compactness Result and A Priori Estimate	32
6. A spherical Harnack inequality	34
Appendix: Comparision between $d_g(x, q)$ and $ x - q $	36
References	37

1. INTRODUCTION

The question of existence of conformal metrics of constant or more generally prescribed curvature on riemannian manifolds is a recurrent problem in differential geometry and geometric analysis. Indeed a positive or a negative answer to such

Lina Wu is partially supported by the China Scholarship Council (No.201806210165).

Lei Zhang is partially supported by a Simons Foundation Collaboration Grant

Date: December 22, 2024.

a question has far reaching consequences on the geometry and topology of the underlying manifold. The Poincaré uniformization's theorem on closed surfaces, the Yamabe problem on riemannian manifolds of dimension $n \geq 3$ and the Nirenberg's problem on standard spheres \mathbb{S}^n , just to name a few, are well known and well studied mathematical problems.

A similar question, which goes back to Picard [37, 38] deals with the existence of conformal metrics of constant or prescribed curvature on surfaces with conical singularities. After the pioneering work of Picard at the beginning of the last century, such a problem has been systematically investigated by Berger [8], McOwen[41, 42] and Troyanov[46, 45]. More recently Bartolucci-deMarchis-Malchiodi[5] used a Morse theoretical approach to prove further existence and multiplicity results.

In this paper, a first part of a series of papers, we address the problem of existence of conformal conical metrics of constant, or more generally prescribed Q -curvature on four dimensional riemannian manifolds. In the following we will explain in some details the geometric context of such a problem:

Given (M, g) a compact four-dimensional Riemannian manifold, the Q -curvature and the Paneitz operator are defined respectively, by

$$(1.1) \quad Q_g = -\frac{1}{12}(\Delta_g R_g - R_g^2 + 3|\text{Ric}_g|^2),$$

$$(1.2) \quad P_g \varphi = \Delta_g^2 \varphi + \text{div}_g \left(\left(\frac{2}{3} R_g g - 2 \text{Ric}_g \right) \nabla u \right),$$

where Ric_g is the Ricci tensor and R_g is the scalar curvature of (M, g) .

Similar to second order equations a natural question is the following uniformization statement: *given a four-dimensional Riemannian manifold (M, g) , is there a metric $\tilde{g} = e^{2u}g$ in the conformal class of g with constant Q -curvature?*

Under the conformal change of metric above, the Paneitz operator is conformally covariant:

$$(1.3) \quad P_{\tilde{g}} \varphi = e^{-4u} P_g \varphi,$$

and the Q curvature of \tilde{g} is given by

$$(1.4) \quad P_g u + 2Q_g = 2\tilde{Q}e^{4u}.$$

From (1.3) and (1.4), the question above is equivalent to the existence of solution to this fourth order equation:

$$(1.5) \quad P_g u + 2Q_g = 2\tilde{Q}e^{4u},$$

where \tilde{Q} is a real constant.

Integrating with respect to the volume element dV_g , we can see that

$$\kappa_P = \int_M Q_g dV_g$$

is a constant in the conformal class of g , here we also point out the Gauss-Bonnet-Chern formula that links the local curvature to the global topology of M is:

$$\int_M \left(Q_g + \frac{1}{8} |W_g|^2 \right) dV_g = 4\pi^2 \chi_M,$$

where W_g denotes the Weyl's tensor of (M, g) and χ_M is the Euler characteristic of M . From this equality and the aforementioned conformal covariance property it is not hard to imagine that P_g and Q_g are related to a number of studies such as Moser-Trudinger type inequatlities, log-determinant formulas and the compactification of locally flat manifolds, see [7, 9, 10, 16, 17, 19]. In many of these studies the kernel of P_g is usually assumed to consist only of constants:

$$(P) \quad \text{Ker}(P_g) = \{\text{constants}\}.$$

In this paper consider the following prescribed Q -curvature equation involving singular sources

$$(1.6) \quad P_g u + 2Q_g = 2he^{4u} - 8\pi^2 \sum_{j=1}^N \gamma_j \left(\delta_{q_j} - \frac{1}{\text{vol}_g(M)} \right),$$

where h is a smooth positive function, $N \in \mathbb{N}$ is a positive integer, q_1, \dots, q_N are distinct points on M and where Dirac measures δ_{q_j} are located, $\gamma_j > -1$ are constants.

Solutions to (1.6) have the following geometric interpretation: Setting $\tilde{g} := e^{2u}g$ we obtain a metric conformal to g on $M \setminus \{q_1, \dots, q_N\}$ which has a conical singularity at each q_i . One says that \tilde{g} is represented by the divisor $D := \sum_{i=1}^N \gamma_i q_i$. See Fang-Ma [23]. Furthermore due to Gauss-Bonnet-Chern formula for conic four manifolds, see [17], [16], [14], we have that

$$\tilde{\kappa}_P := \int_M Q_{\tilde{g}} dV_{\tilde{g}} = \int_M Q_g dV_g + 8\pi^2 \sum_{i=1}^N \gamma_i$$

is a conformal invariant.

Considerable progress has been made for the regular case of (1.6), that is $N = 0$ in (1.6). Under the assumption that the Kernel of the Paneitz operator contains only constants, Chang-Yang [19] proved existence for $\kappa_P < 8\pi^2$, Djadli-Malchiodi [21] settled the case that $\kappa_P \neq 8\pi^2 n$ for any $n \in \mathbb{N}$. Li-Li-Liu [30] gave a necessary condition for existence in the case $\kappa_P = 8\pi^2$, Ahmedou-Ndiaye [2] developed a Morse theory the case of $\kappa_P = 8\pi^2 n$ and Ndiaye [36] combined the celebrated topological argument of Bahri-Coron [3] with the *critical point theory at infinity* in [2] to derive some existence results. We point out that an essential estimate related to the proof in [21] is a priori estimate when κ_P is away from $8\pi^2 \mathbb{N}$ proved by Malchiodi in [33]. Later in [22], Druet and Robert extended such an a priori estimate to the following more general equation in the same class:

$$(1.7) \quad P_g u + 2b = 2he^{4u},$$

where b is a smooth function. If $b = Q_g$ is the Q -curvature of the conformal metric $e^{2u}g$. More specially, assuming $h_k \rightarrow h_0$, $h_k \geq c_0 > 0$ and $b_k \rightarrow b_0$, then any sequence

of solutions $\{u_k\}$ of (1.7) with $h = h_k$ and $b = b_k$ is uniformly bounded under the condition $\int_M b_0 dV_g \neq 8\pi^2 n$, see also Malchiodi [33].

However, bubbling can occur when $\int_M b_0 dV_g = 8\pi^2 n$ for some positive integer n . The understanding of this bubbling phenomenon is vital for the existence problem. The study of the blow-up profile and other blow-up phenomena for the Paneitz operator and other elliptic equations has attracted much interest recently and the reference is too numerous to be mentioned, we just list a few closely related to our article in our humble opinion: [1, 11, 18, 20, 25, 26, 27, 30, 34, 40, 35, 39, 48, 49, 52]. Particularly, in [52], the third named author and Weinstein have obtained sharp estimates on the difference near the blow-up points between a bubbling sequence of solutions to (1.7) with $h = h_k$ and $b = b_k$ and standard bubbles, and obtained the vanishing rate under the assumption that (M, g) may not be locally conformally flat.

When taking the singularities into the account as in (1.6), we consider the following more general singular equation:

$$(1.8) \quad P_g u + 2b = 2he^{4u} - 8\pi^2 \sum_{j=1}^N \gamma_j \left(\delta_{q_j} - \frac{1}{\text{vol}_g(M)} \right),$$

where h is a positive smooth function on M and $b \in C^1(M)$. Before stating our first main result, we define a *critical set* Γ as follows:

$$\Gamma = \left\{ 16\pi^2 n + 16\pi^2 \sum_{j \in J} (1 + \gamma_j) : n \in \mathbb{N} \cup \{0\} \text{ and } J \subset \{1, \dots, N\} \right\}.$$

In order to obtain the a priori estimates and existence results, we mainly study the blow-up phenomena for (1.8). Let us consider the following equations:

$$(1.9) \quad P_g u_k + 2b_k = 2h_k e^{4u_k} - 8\pi^2 \sum_{j=1}^N \gamma_j \left(\delta_{q_j} - \frac{1}{\text{vol}_g(M)} \right) \quad \text{in } M,$$

with normalized total integration:

$$(1.10) \quad \int_M e^{4u_k} dV_g = 1.$$

Let $\{u_k\}$ be a sequence of solutions of (1.9). We say p is a blowup point of u_k if there exists a sequence $p_k \rightarrow p$ such that

$$u_k(p_k) + 8\pi^2 \sum_{j=1}^N \gamma_j G(p_k, q_j) \rightarrow \infty.$$

u_k is called a sequence of blowup solutions if it has a blowup point. Here $G(x, p)$ is the Green's function of P_g defined in (3.3). For blowup solutions we assume that coefficient functions are regular enough to have limits:

$$(1.11) \quad \|b_k - b_0\|_{C^1(M)} \rightarrow 0, \quad \|h_k - h_0\|_{C^1(M)} \rightarrow 0, \quad 0 < c_0 < h_0 < 1/c_0.$$

Without loss of generality, we assume the integration of $h_k e^{u_k}$ is 1:

Our first main result asserts that a priori estimate holds for u_k , as long as $2 \int_M b_k$ does not tend to the following critical set:

$$\Gamma = \left\{ 16\pi^2 n + 16\pi^2 \sum_{j \in J} (1 + \gamma_j) : n \in \mathbb{N} \cap \{0\} \text{ and } J \subset \{1, \dots, N\} \right\}.$$

Theorem 1.1 (A Priori Estimate). *Suppose (P) holds, b and h satisfy (1.11). If $\{u_k\}$ is a sequence of solutions of (1.9) under restriction (1.10) and $\int_M 2b_0 dV_g \in \mathbb{R}^+ \setminus \Gamma$,*

$$\left| u_k(x) + 8\pi^2 \sum_{j=1}^N \gamma_j G(x, q_j) \right| \leq C, \quad \forall x \in M$$

for some $C > 0$ independent of k .

In particular, the a priori estimate holds for the singular prescribing Q -curvature equations. Indeed Theorem 1.1 is an extension of previous results of Malchiodi [33], Druet-Robert [22] and Fardoun-Regbaoui [24] for the regular prescribed Q -curvature equation. We point out that the argument in the regular case uses in a crucial way the explicit form of the bubble, while our argument uses only the asymptotic behavior of the bubble and is based on a Pohozaev identity for equations under conformal normal coordinates (see [52]).

If $\int_M 2b_0 dV_g \in \Gamma$ and u_k does blowup around a singular source $q \in M$, our next result says that if γ_q is not a positive integer, the Spherical Harnack inequality holds for u_k around q :

Theorem 1.2. *Suppose $\{u_k\}$ be a sequence of solutions of (1.9) that also satisfies (1.10), (P) holds and b, h satisfy (1.11). If q_j is a blowup point of u_k and $\gamma_{q_j} \notin \mathbb{N}$, there exist $C, \delta > 0$ independent of k such that*

$$\max_{B(q_j, \delta)} (u_k(x) + \log|x - q_j|) \leq C, \quad |x - q_j| \leq \delta.$$

Establishing Spherical Harnack inequality for bubbling solutions is critical for applications like a priori estimate, degree counting program and existence results. The readers may look at recent breakthroughs of the third author and Wei [51] for Liouville equations.

One indispensable part of the blowup analysis for the Q curvature equation is the classification of global solutions on \mathbb{R}^4 . For this purpose we consider the limiting equation used to describe the profile of bubbling solutions:

$$(1.12) \quad \Delta^2 u(x) = 6|x|^{4\gamma} e^{4u(x)}, \quad \text{in } \mathbb{R}^4, \quad |x|^{4\gamma} e^{4u(x)} \in L^1(\mathbb{R}^4).$$

Clearly if u is a solution of (1.12), so is u_λ defined by

$$(1.13) \quad u_\lambda(x) = u(\lambda x) + (1 + \gamma) \log \lambda$$

for any given $\lambda > 0$. Our next main result is

Theorem 1.3. *Suppose that u is a solution of (1.12) with $\gamma > -1$ and $|u(x)| = o(|x|^2)$ at infinity. Then*

$$(i) \quad \int_{\mathbb{R}^4} 6|y|^{4\gamma} e^{4u(y)} dy = 16\pi^2(1 + \gamma).$$

- (ii) $u(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|y|}{|x-y|}\right) |y|^{4\gamma} e^{4u(y)} dy + C_0$ for some $C_0 \in \mathbb{R}$,
- (iii) Furthermore, if $-1 < \gamma < 0$, u is radially symmetric about the origin and is unique up to scaling in (1.13).

Note that (i) is a quantization result and is true for all $\gamma > -1$. (iii) completes the classification for all $\gamma \in (-1, 0]$. (ii) is also proved as a part of the main theorem in [27]. As a corollary of the classification in (iii) we now derive a more specific description of the asymptotic behavior of u : For simplicity we use $\mu = 1 + \gamma$.

Corollary 1.4. *Let u be a solution of (1.12) with $\gamma \in (-1, 0)$ and suppose $|u(x)| = o(|x|^2)$ at ∞ . Then for $M = \lfloor \frac{1}{4\mu} \rfloor$, there exists $c_0 \in \mathbb{R}$ such that*

$$(1.14) \quad u(x) = -2\mu \log |x| + c_0 + \sum_{l=1}^M \frac{c_l}{|x|^{4l\mu}} + O\left(\frac{1}{|x|}\right), \quad |x| > 1$$

where $c_l = \frac{3e^4 \sum_{s=0}^{l-1} c_s}{32l^2\mu^2(1-2l\mu)(1+2l\mu)}$, $l = 1, \dots, M$;

$$(1.15) \quad -\Delta u(x) = \frac{4\mu}{|x|^2} + \sum_{l=1}^M \frac{4c_l l \mu (2-4l\mu)}{|x|^{2+4l\mu}} + O\left(\frac{1}{|x|^3}\right), \quad |x| > 1$$

Note that $\lfloor \frac{1}{4\mu} \rfloor$ stands for the largest integer no greater than $\frac{1}{4\mu}$. If $\gamma > -\frac{3}{4}$, $M = 0$, the third term in (1.15) does not exist in this case. If the $o(|x|^2)$ assumption in Theorem 1.3 is removed, it is established in [27] by Hyder et al that if u is a solution of (1.12) with $\gamma > -1$, after an orthogonal transformation, $u(x)$ can be represented by

$$(1.16) \quad u(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|y|}{|x-y|}\right) |y|^{4\gamma} e^{4u(y)} dy - \sum_{j=1}^4 a_j (x_j - x_j^0)^2 + c_0$$

for some $c_0 \in \mathbb{R}$. In this case we can prove a symmetry result: If $\gamma \in (-1, 0)$ and $a_i x_i^0 = 0$ for all i , u is a radial function. See Theorem 2.10 for more detail.

Remark 1.5. *Our quantization and classification results directly point to the following important open questions: 1. Theorem 1.3 does not cover the case that $\gamma > 0$, except for the quantization result. The symmetry and profile for global solutions of (1.12) in this case are largely unknown.*

2. If the sub-quadratic growth assumption ($u(x) = o(|x|^2)$) is removed, it is not clear to us how the total integration of global solutions will change. It is proved in Lin [31] that for $\gamma = 0$ solutions with sub-quadratic growth give the largest integration. It is also proved by Hyder et al [27] that for $\gamma < 0$ there are non-radial solutions whose integration is greater than that of radial ones. It can be easily derived from our Theorem 2.10 that if $\gamma < 0$ and $a_i x_i^0 = 0$ for all i , the one with the sub-quadratic growth gives the largest integration. However, no information is known for other situations.

When we were about to submit this article we heard that Jevnikar-Sire-Yang [28] are working on a similar project independently and their results are to be posted soon.

Here we briefly outline the strategy of the proofs in our paper. For the proof of the classification result for globally defined singular equation, we follow the argument of Lin [31] but we need to take care of all the complications caused by the singular source. In particular the method of moving planes relies crucially on the integral form of the global solutions. We are able to prove the complete classification for $\gamma \in (-1, 0]$ and a quantization result for all $\gamma > -1$. For blowup solutions we first use a small-energy lemma (Lemma 4.1) to prove that there are at most finite blowup points. Then we take advantage of a Pohozaev identity established by Weinstein-Zhang [52] to describe a precise asymptotic behavior of blowup solutions around a blowup point. Then the total integration as well as precise asymptotic behavior of solutions can be further determined. With this information the critical set Γ can be identified and if the total integral of $2b$ does not tend to Γ we obtain a priori estimate. The idea to prove Theorem 1.2 is as follows: If the spherical harnack inequality is violated, there should be finite small bubbling circles around the singular source all tending to the singular source. Around each tiny bubbling disk there is a Pohozaev identity, and a “big” ball that includes all these tiny balls also has a Pohozaev identity. The comparison of these Pohozaev identities implies that the strength of the singular source has to be an integer.

The organization of this paper is as follows. In section 2 we analyze the globally defined solutions and proved the quantization and the classification results stated in Theorem 1.3 and Corollary 1.4. Then in Section 3, we list some useful facts about the conformal normal coordinates and Pohozaev identity and in section 4, we perform a blow-up analysis near singular points. Section 5 is devoted to a concentration-compactness theorem and a priori estimate for the singular prescribing Q -curvature equation on 4-manifolds while in section 6 we prove our spherical Harnack inequality. Finally we provide in the appendix an useful estimate of the difference between the geodesic distance and the Euclidean one for nearby points on the manifold.

2. ENTIRE SOLUTIONS OF FOURTH ORDER SINGULAR LIOUVILLE TYPE EQUATIONS IN \mathbb{R}^4

In this section, we will follow the argument of Lin [31] to analyze solutions of (1.12) and prove Theorem 1.3 and Theorem 2.10.

2.1. Asymptotic behavior of entire solutions.

Our argument is progressive in nature and we shall obtain a rough estimate of u at infinity. For this purpose we set

$$(2.1) \quad v(x) := \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|x-y|}{|y|}\right) |y|^{4\gamma} e^{4u(y)} dy,$$

which is obviously a solution of

$$(2.2) \quad \Delta^2 v(x) = -6|x|^{4\gamma} e^{4u(x)}, \quad \text{in } \mathbb{R}^4.$$

The asymptotic behavior of u has a large to do with that of v , so in the first lemma we derive a rough upper bound of v . For convenience we set

$$(2.3) \quad \alpha = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} |y|^{4\gamma} e^{4u} dy.$$

Lemma 2.1. *Suppose u is a solution of (1.12) and let α be given as in (2.3). Then*

$$(2.4) \quad v(x) \leq \alpha \log |x| + C$$

for some constant C .

Proof. Since the goal is to describe asymptotic behavior it is natural to assume $|x| > 4$. For such x , we decompose $\mathbb{R}^4 = A_1 \cup A_2$, where

$$A_1 = \left\{ y : |y - x| \leq \frac{|x|}{2} \right\}, \quad A_2 = \left\{ y : |y - x| \geq \frac{|x|}{2} \right\}.$$

For $y \in A_1$, $\log \frac{|x-y|}{|y|} \leq 0$ because $|y| \geq |x| - |x-y| \geq \frac{|x|}{2} \geq |x-y|$, Thus

$$\int_{A_1} \log \left(\frac{|x-y|}{|y|} \right) |y|^{4\gamma} e^{4u(y)} dy \leq 0$$

and

$$v(x) \leq \frac{3}{4\pi^2} \int_{A_2} \log \left(\frac{|x-y|}{|y|} \right) |y|^{4\gamma} e^{4u(y)} dy \leq 0.$$

To evaluate the integral over A_2 , we first make two trivial observations:

$$\begin{aligned} |x-y| &\leq |x| + |y| \leq |x||y|, & \text{if } |y| \geq 2, \\ \log |x-y| &\leq \log |x| + C, & \text{if } |y| \leq 2 \end{aligned}$$

where $|x| > 4$ is used. Consequently

$$\begin{aligned} v(x) &\leq \frac{3}{4\pi^2} \int_{A_2} \log \left(\frac{|x-y|}{|y|} \right) |y|^{4\gamma} e^{4u(y)} dy \\ &\leq \frac{3}{4\pi^2} \left\{ \log |x| \int_{A_2 \cap \{|y| \geq 2\}} |y|^{4\gamma} e^{4u} dy + \int_{A_2 \cap \{|y| \leq 2\}} \log \left(\frac{|x-y|}{|y|} \right) |y|^{4\gamma} e^{4u} dy \right\} \\ &\leq \frac{3}{4\pi^2} \left\{ \log |x| \int_{A_2} |y|^{4\gamma} e^{4u} dy + C \int_{A_2 \cap \{|y| \leq 2\}} |y|^{4\gamma} e^{4u} dy \right. \\ &\quad \left. - \int_{A_2 \cap \{|y| \leq 2\}} (\log |y|) |y|^{4\gamma} e^{4u} dy \right\} \\ &\leq \frac{3}{4\pi^2} \log |x| \int_{\mathbb{R}^4} |y|^{4\gamma} e^{4u} dy + C. \end{aligned}$$

Lemma 2.1 is established. \square

Before proving a lower bound of $v(x)$ we derive an expression of $\Delta u(x)$ in integral form.

Lemma 2.2. *Suppose u is a solution of (1.12). Then there exists a constant $C_1 \geq 0$ such that*

$$(2.5) \quad \Delta u(x) = -\frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x-y|^2} |y|^{4\gamma} e^{4u(y)} dy - C_1.$$

Proof. Let $w(x) = u(x) + v(x)$. Then from the equations of u and v in (1.12) and (2.2), we have $\Delta^2 w = 0$ in \mathbb{R}^4 . Hence, Δw is a harmonic function in \mathbb{R}^4 . By the mean value property of harmonic functions, we have, for any $x_0 \in \mathbb{R}^4$ and $r > 0$,

$$\Delta w(x_0) = \frac{2}{\pi^2 r^4} \int_{B(x_0, r)} \Delta w(y) dy = \frac{2}{\pi^2 r^4} \int_{\partial B(x_0, r)} \Delta w(y) d\sigma,$$

where $\frac{\pi^2}{2}$ is the volume of the unit ball. That is

$$(2.6) \quad \frac{r}{4} \Delta w(x_0) = \oint_{|y-x_0|=r} \frac{\partial w}{\partial r}(y) d\sigma,$$

where $\oint_{|y-x_0|=r} f(y) d\sigma = \frac{1}{2\pi^2 r^3} \int_{|y-x_0|=r} f(y) d\sigma$ denotes the integral average of f over $\partial B(x_0, r)$. Then integrating the identity above along r , we get

$$\frac{r^2}{8} \Delta w(x_0) = \oint_{|y-x_0|=r} w d\sigma - w(x_0).$$

Therefore, the Jensen inequality implies

$$\begin{aligned} \exp\left(\frac{r^2}{2} \Delta w(x_0)\right) &\leq e^{-4w(x_0)} \exp\left(4 \oint_{|y-x_0|=r} w d\sigma\right) \\ &\leq e^{-4w(x_0)} \oint_{|y-x_0|=r} e^{4w} d\sigma \end{aligned}$$

From Lemma 2.1, we have $w(x) = u(x) + v(x) \leq u(x) + \alpha \log|x| + C$, and as a consequence

$$\begin{aligned} \int_0^\infty r^{3-4\alpha+4\gamma} \exp\left(\frac{\Delta w(x_0)}{2} r^2\right) dr &\leq \int_{\mathbb{R}^4} |x|^{-4\alpha+4\gamma} e^{-4w(x_0)} e^{4w(x)} dx \\ &\leq C \int_{\mathbb{R}^4} |x|^{-4\alpha+4\gamma} e^{4u(x)} |x|^{4\alpha} dx = C \int_{\mathbb{R}^4} |x|^{4\alpha} e^{4u} dx < +\infty, \end{aligned}$$

which means

$$r^{3-4\alpha+4\gamma} \exp\left(\frac{\Delta w(x_0)}{2} r^2\right) \in L^1([1, +\infty)).$$

From here we see $\Delta w(x_0) \leq 0$ for all $x_0 \in \mathbb{R}^4$. Liouville's Theorem implies that there exists some constant $C_1 \geq 0$ such that $\Delta w(x) \equiv -C_1$ in \mathbb{R}^4 . Lemma 2.2 is established based on this and

$$\Delta v(x) = \frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x-y|^2} |y|^{4\gamma} e^{4u(y)} dy.$$

□

With the help of the representation for Δu , we can estimate $v(x)$ from below for $|x|$ large. We will use following result in Lemma 2.3 of [31].

Let $h(x)$ be the solution of

$$\begin{cases} \Delta^2 h(x) = f(x), & \text{in } \Omega, \\ \Delta h(x) = h(x) = 0, & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded domain of \mathbb{R}^4 .

Lemma 1. [31]

Suppose $f \in L^1(\bar{\Omega})$. Then for any $\delta \in (0, 32\pi^2)$, there exists a constant $C_\delta > 0$ such that

$$\int_{\Omega} \exp\left(\frac{\delta|h|}{\|f\|_{L^1}}\right) dx \leq C_\delta (\text{diam } \Omega)^4,$$

where $\text{diam } \Omega$ denotes the diameter of Ω .

Lemma 2.3. Let u be a solution of (1.12) and v be in (2.1). Then for given $\varepsilon > 0$, there exists a constant $R = R(\varepsilon)$ only depending on ε such that

$$(2.7) \quad v(x) \geq (\alpha - \varepsilon) \log |x|, \quad |x| > R(\varepsilon).$$

Proof. We first prove a claim slightly weaker than (2.7): for any $\varepsilon > 0$, there exists $R = R(\varepsilon) > 0$ such that

$$(2.8) \quad v(x) \geq (\alpha - \frac{\varepsilon}{2}) \log |x| + \frac{3}{4\pi^2} \int_{B(x,1)} (\log |x-y|) |y|^{4\gamma} e^{4u(y)} dy.$$

To prove (2.8) we consider \mathbb{R}^4 as a disjoint union of three sets: $\mathbb{R}^4 = A_1 \cup A_2 \cup A_3$, where

$$\begin{aligned} A_1 &= \{y : |y| < R_0\}, \\ A_2 &= \{y : |x-y| \leq |x|/2, |y| \geq R_0\}, \\ A_3 &= \{y : |x-y| \geq |x|/2, |y| \geq R_0\}. \end{aligned}$$

Then we choose $R_0 = R_0(\varepsilon)$ sufficiently large so that

$$\begin{aligned} & \frac{3}{4\pi^2} \int_{A_1} \log\left(\frac{|x-y|}{|y|}\right) |y|^{4\gamma} e^{4u(y)} dy - \alpha \log |x| \\ &= \frac{3}{4\pi^2} \log |x| \int_{A_1} \frac{\log |x-y| - \log |x| - \log |y|}{\log |x|} |y|^{4\gamma} e^{4u(y)} dy - \frac{\varepsilon}{8} \log |x| \\ &\geq -\frac{\varepsilon}{4} \log |x| \end{aligned}$$

for large $|x|$. Thus we have

$$(2.9) \quad \frac{3}{4\pi^2} \int_{A_1} \log\left(\frac{|x-y|}{|y|}\right) |y|^{4\gamma} e^{4u(y)} dy \geq (\alpha - \frac{\varepsilon}{4}) \log |x|.$$

For $x \in A_2$ and $|x|$ large, we have $\frac{|x|}{2} \leq |x| \leq \frac{3}{2}|x|$. Then

$$\begin{aligned} & \int_{A_2} \log\left(\frac{|x-y|}{|y|}\right) |y|^{4\gamma} e^{4u(y)} dy \\ (2.10) \quad &= \int_{A_2} (\log |x-y|) |y|^{4\gamma} e^{4u(y)} dy - \int_{A_2} (\log |y|) |y|^{4\gamma} e^{4u(y)} dy \\ &\geq \int_{B(x,1)} (\log |x-y|) |y|^{4\gamma} e^{4u(y)} dy - \log(2|x|) \int_{A_2} |y|^{4\gamma} e^{4u(y)} dy. \end{aligned}$$

For $y \in A_3$, we use two trivial inequalities: $|x - y| \geq \frac{|x|}{2} \geq \frac{|y|}{4}$ if $|y| \leq 2|x|$ and $|x - y| \geq |y| - |x| \geq \frac{|y|}{2}$ if $|y| \geq 2|x|$. Clearly in both cases, we have

$$\frac{|x - y|}{|y|} \geq \frac{1}{4}, \quad y \in A_3.$$

Therefore,

$$(2.11) \quad \frac{3}{4\pi^2} \int_{A_3} \log\left(\frac{|x - y|}{|y|}\right) |y|^{4\gamma} e^{4u(y)} dy \geq \log \frac{1}{4} \int_{A_3} |y|^{4\gamma} e^{4u(y)} dy.$$

From (2.9), (2.10), (2.11) and $|y|^{4\gamma} e^{4u(y)} \in L^1(\mathbb{R}^4)$, we obtain (2.8).

Next, we show that

$$(2.12) \quad \int_{B(x,1)} (\log |x - y|) |y|^{4\gamma} e^{4u(y)} dy \geq -C$$

for some positive constant C . For this purpose we set

$$(2.13) \quad \tilde{u}(x) = u(x) - \gamma \log |x|$$

Then \tilde{u} satisfies

$$(2.14) \quad \begin{cases} \Delta^2 \tilde{u}(x) = 6e^{4\tilde{u}(x)} - 8\pi^2 \gamma \delta_0, & \text{in } \mathbb{R}^4, \\ e^{4\tilde{u}} \in L^1(\mathbb{R}^4). \end{cases}$$

Let $0 < \varepsilon_0 < \pi^2$ and $R_0 = R_0(\varepsilon_0)$ be sufficiently large such that

$$(2.15) \quad 6 \int_{B(x,4)} |y|^{4\gamma} e^{4u(y)} dy = 6 \int_{B(x,4)} e^{4\tilde{u}(y)} dy \leq \varepsilon_0, \quad \text{for } |x| \geq R_0,$$

then we let h be the solution of

$$\begin{cases} \Delta^2 h(x) = 6e^{4\tilde{u}(y)}, & \text{in } B(x,4), \\ h(x) = \Delta h(x) = 0, & \text{on } \partial B(x,4). \end{cases}$$

From Lemma 1, we can see that for $\varepsilon_0 > 0$ small,

$$(2.16) \quad \int_{B(x,4)} e^{24|h|} dy \leq c_1$$

for some constant c_1 independent of x .

Next we set $q(y) = \tilde{u}(y) - h(y)$ for $y \in B(x,4)$, which clearly satisfies

$$\begin{cases} \Delta^2 q(y) = 0, & \text{in } B(x,4), \\ q(y) = \tilde{u}(y), \quad \Delta q(y) = \Delta \tilde{u}(y), & \text{on } \partial B(x,4). \end{cases}$$

Let $\tilde{q}(y) = -\Delta q(y)$. Then by Lemma 2.2, we see that for $|x|$ large enough and $y \in \partial B(x,4)$

$$\tilde{q}(y) = -\Delta \tilde{u}(y) = -\Delta u(y) + \gamma \Delta(\log |y|) = -\Delta u(y) - \frac{2\gamma}{|y|^2}.$$

By setting

$$\hat{q}(y) = \tilde{q}(y) + \frac{2\gamma}{|y|^2} = -\Delta u(y) = \frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|z - y|^2} |z|^{4\gamma} e^{4u(z)} dz + C_1,$$

we obviously have $\hat{q}(y) > 0$ on $\partial B(x, 4)$, and hence $\tilde{q}(y) > -2\gamma/|y|^2$ on $\partial B(x, 4)$. Observing that $1/|y|^2$ is the fundamental solution of Δ . In other words, $\hat{q}(y)$ is harmonic in $B(x, 4)$ with positive boundary value on $\partial B(x, 4)$. The maximum principle implies $\hat{q} > 0$ in $B(x, 4)$. Thus, by the Harnack inequality and mean value property of harmonic functions, we have

$$(2.17) \quad \begin{aligned} \tilde{q}(y) &= \hat{q}(y) - \frac{2\gamma}{|y|^2} \\ &\leq c_2 \hat{q}(x) - \frac{2\gamma}{|y|^2} = -\int_{\partial B(x, 4)} \Delta u d\sigma - \frac{2\gamma}{|y|^2} = -\int_{\partial B(x, 4)} \Delta \tilde{u} d\sigma, \quad y \in \overline{B(x, 2)}, \end{aligned}$$

with a constant c_2 .

Integrating (2.14) along r , we have

$$\begin{aligned} \int_{\partial B(x, 4)} \Delta \tilde{u} d\sigma - \Delta \tilde{u}(x) &= \int_0^r \frac{3}{\pi^2 s^3} \int_0^s \int_{\partial B(x, t)} e^{4\tilde{u}} d\sigma dt ds \\ &= \int_0^r \frac{3}{\pi^2 s^3} \int_0^s t^3 \int_{\partial B(x, 1)} e^{4\tilde{u}} d\sigma dt ds \\ &= \int_0^r t^3 \int_{\partial B(x, 1)} e^{4\tilde{u}} \left(\frac{3}{2\pi^2 t^2} - \frac{3}{2\pi^2 r^2} \right) d\sigma dt. \end{aligned}$$

That is

$$(2.18) \quad \int_{\partial B(x, 4)} \Delta \tilde{u} d\sigma - \Delta \tilde{u}(x) = \frac{3}{2\pi^2} \int_{B(x, r)} \left(\frac{1}{|x-y|^2} - \frac{1}{r^2} \right) e^{4\tilde{u}} dy.$$

Next by Lemma 2.2 and (2.18), we can see

$$\begin{aligned} -\int_{\partial B(x, 4)} \Delta \tilde{u} d\sigma &= -\Delta \tilde{u}(x) - \frac{3}{2\pi^2} \int_{B(x, r)} \frac{1}{|x-y|^2} e^{4\tilde{u}} dy + \frac{3}{2\pi^2 r^2} \int_{B(x, r)} e^{4\tilde{u}} dy \\ &= -\Delta u(x) - \frac{3}{2\pi^2} \int_{B(x, r)} \frac{1}{|x-y|^2} e^{4\tilde{u}} dy + \frac{3}{2\pi^2 r^2} \int_{B(x, r)} e^{4\tilde{u}} dy - \frac{2\gamma}{|x|^2} \\ &= \frac{3}{2\pi^2 r^2} \int_{|x-y| \geq r} \frac{1}{|x-y|^2} e^{4\tilde{u}} dy + \frac{3}{2\pi^2 r^2} \int_{B(x, r)} e^{4\tilde{u}} dy - \frac{2\gamma}{|x|^2} + C_1. \end{aligned}$$

In particular, for $r = 4$ and $|x|$ large,

$$(2.19) \quad -\int_{\partial B(x, 4)} \Delta \tilde{u} d\sigma \leq c_3.$$

Hence, from (2.17), we get

$$(2.20) \quad \tilde{q}(y) \leq c_4, \quad y \in \overline{B(x, 2)},$$

and immediately

$$|\tilde{q}(y)| \leq c_5, \quad y \in \overline{B(x, 2)}.$$

Since q satisfies

$$\begin{cases} \Delta q(y) = -\tilde{q}(y), & \text{in } B(x, 4), \\ q(y) = \tilde{u}(y), & \text{on } \partial B(x, 4), \end{cases}$$

by estimates for linear elliptic equations, we have for any $p > 1$ and $\sigma > 2$,

$$(2.21) \quad \sup_{B(x,1)} q \leq c \left(\|q^+\|_{L^p(B(x,2))} + \|\tilde{q}\|_{L^\sigma(B(x,2))} \right),$$

where $c = c(p, \sigma)$.

On the other hand, we observe that $q^+(y) \leq \tilde{u}^+(y) + |h(y)|$ for $y \in B(x, 4)$. Then by (2.16), we get

$$\int_{B(x,2)} (q^+)^p \leq c_6 \int_{B(x,2)} e^{2q^+} \leq c_5 \left(\int_{B(x,2)} e^{4\tilde{u}^+} \right)^{\frac{1}{2}} \left(\int_{B(x,2)} e^{4|h|} \right)^{\frac{1}{2}} \leq c_7 \left(\int_{B(x,2)} e^{4\tilde{u}^+} \right)^{\frac{1}{2}}$$

Since $e^{4\tilde{u}^+} \leq 1 + e^{4\tilde{u}}$, we have $\|q^+\|_{L^p(B(x,2))} \leq c_7$, which together with (2.20) and (2.21) implies

$$(2.22) \quad \sup_{B(x,1)} q \leq c_8.$$

In view of $\tilde{u} = h + q$, we now obtain

$$(2.23) \quad \tilde{u}(y) \leq h(y) + q(y) \leq c_8 + |h(y)|, \quad y \in \overline{B(x, 2)}.$$

Therefore,

$$(2.24) \quad \int_{B(x,1)} e^{24\tilde{u}} \leq c_9 \int_{B(x,1)} e^{24|h|} dy \leq c_{10},$$

Then

$$\left| \int_{B(x,1)} (\log|x-y|) e^{4\tilde{u}} dy \right| \leq \left(\int_{B(x,1)} (\log|x-y|)^2 dy \right)^{\frac{1}{2}} \left(\int_{B(x,1)} e^{8\tilde{u}(y)} dy \right)^{\frac{1}{2}} \leq c_{11},$$

which means

$$(2.25) \quad \left| \int_{B(x,1)} (\log|x-y|) |y|^{4\gamma} e^{4u(y)} dy \right| \leq c_{11},$$

where c_{11} is a constant independent of x ($|x|$ large). As a consequence, (2.8) and (2.25) lead to

$$v(x) \geq \left(\alpha - \frac{\varepsilon}{2}\right) \log|x| - c_{11} \geq (\alpha - \varepsilon) \log|x|$$

for $|x|$ large, which is (2.7).

To estimate Δv , we observe that

$$\Delta v(x) = -\frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x-y|^2} |y|^{4\gamma} e^{4u(y)} dy = -\frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x-y|^2} e^{4\tilde{u}(y)} dy.$$

We decompose $\mathbb{R}^4 = B_1 \cup B_2 \cup B_3$ for $|x|$ large, where

$$\begin{aligned} B_1 &= \{y : |x-y| \geq |x|/2\}, \\ B_2 &= \{y : 1 \leq |x-y| \leq |x|/2, |y| \geq R_0\}, \\ B_3 &= \{y : |x-y| \leq 1\}. \end{aligned}$$

Then

$$\int_{B_1} \frac{1}{|x-y|^2} e^{4\tilde{u}(y)} dy \leq \frac{C}{|x|^2} \rightarrow 0, \quad \text{as } |x| \rightarrow \infty,$$

$$\int_{B_2} \frac{1}{|x-y|^2} e^{4\tilde{u}(y)} dy \leq \int_{B_2} e^{4\tilde{u}(y)} dy \rightarrow 0, \quad \text{as } |x| \rightarrow \infty,$$

and

$$\begin{aligned} & \int_{B_3} \frac{1}{|x-y|^2} e^{4\tilde{u}(y)} dy \\ & \leq \left(\int_{|x-y| \leq 1} \left(\frac{1}{|x-y|} \right)^{\frac{8}{5}} dy \right)^{\frac{5}{8}} \left(\int_{|x-y| \leq 1} e^{4\tilde{u}} dy \right)^{\frac{1}{4}} \left(\int_{|x-y| \leq 1} e^{24\tilde{u}} dy \right)^{\frac{1}{8}} \\ & \leq c \left(\int_{|x-y| \leq 1} e^{4\tilde{u}} dy \right)^{\frac{1}{4}} \rightarrow 0, \quad \text{as } |x| \rightarrow \infty, \end{aligned}$$

where we have used (2.24). Lemma 2.3 is established. \square

With the estimates of $v(x)$ near infinity and the expression of Δu , we can show the expression of u in integral form under the condition $|u(x)| = o(|x|^2)$ at ∞ :

Lemma 2.4. *Suppose $|u(x)| = o(|x|^2)$ at ∞ . Then there exists a constant C_0 such that*

$$(2.26) \quad u(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log \left(\frac{|y|}{|x-y|} \right) |y|^{4\gamma} e^{4u(y)} dy + C_0.$$

Furthermore, for any given $\varepsilon > 0$,

$$(2.27) \quad -\alpha \log |x| - C \leq u(x) \leq (-\alpha + \varepsilon) \log |x|, \quad |x| \geq R(\varepsilon),$$

where $R(\varepsilon)$ comes from Lemma 2.3.

Proof. We start from the integral expression of Δu in Lemma 2.2:

$$\Delta u(x) = -\frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x-y|^2} |y|^{4\gamma} e^{4u(y)} dy - C_1, \quad C_1 \geq 0,$$

and we first prove $C_1 = 0$ by contradiction. If $C_1 > 0$ we have

$$\Delta u(x) \leq -C_1 < 0, \quad |x| \geq R_0,$$

where R_0 is large. Let

$$(2.28) \quad h(y) = u(y) + \varepsilon |y|^2 + A(|y|^{-2} - R_0^{-2}).$$

Under the assumption of $|u(y)| = o(|y|^2)$ at ∞ , we have $\lim_{|y| \rightarrow +\infty} h(y) = +\infty$ for any fixed $\varepsilon > 0$ and $A > 0$. So we choose $\varepsilon > 0$ small to make

$$(2.29) \quad \Delta h(y) = \Delta u(y) + 8\varepsilon < -\frac{C_1}{2} < 0, \quad |y| \geq R_0,$$

and A sufficiently large such that $\inf_{|y| \geq R_0} h(y)$ is achieved by some $y_0 \in \mathbb{R}^4$ and $|y_0| > R_0$. Clearly we have obtained a contradiction to the maximum principle. Hence, $C_1 = 0$ and $u + v$ is harmonic in \mathbb{R}^4 .

From Lemma 2.1 and Lemma 2.3, we know for $|x|$ large enough

$$(\alpha - \varepsilon) \log |x| \leq v(x) \leq \alpha \log |x| + C,$$

which together with the assumption $|u(x)| = o(|x|^2)$ at ∞ indicates

$$(2.30) \quad |u(x) + v(x)| = o(|x|^2) \quad \text{at } \infty.$$

Since $u + v$ is harmonic, by the gradient estimates for harmonic functions, we have

$$u(x) + v(x) = \sum_{j=1}^4 a_j x_j + a_0$$

with some constants $a_j \in \mathbb{R}$, $j = 0, \dots, 4$. Therefore, for $|x|$ large enough, we get

$$e^{4u(x)} = e^{a_0} e^{-4v(x)} e^{\sum_{j=1}^4 a_j x_j} \geq C|x|^{-4\alpha} e^{\sum_{j=1}^4 a_j x_j}.$$

Since $|y|^{4\gamma} e^{4u(y)} \in L^1(\mathbb{R}^4)$, we have $a_j = 0$ for $1 \leq j \leq 4$. Therefore,

$$u(x) = -v(x) + a_0 = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|y|}{|x-y|}\right) |y|^{4\gamma} e^{4u(y)} dy + C_0,$$

and then

$$-\alpha \log|x| - C \leq u(x) \leq (-\alpha + \varepsilon) \log|x|,$$

for $|x|$ large. Lemma 2.4 is established. \square

Next we need a Pohozaev identity for u of

$$(2.31) \quad \Delta u = Q(x)e^{4u} \text{ in } \mathbb{R}^4.$$

Lemma 2.5. *Suppose u is an entire smooth solution of (2.31). Then for any bounded domain, we have*

$$(2.32) \quad \begin{aligned} & \int_{\Omega} \left(Q + \frac{1}{4} \langle x, \nabla Q \rangle e^{4u} \right) dx \\ &= \frac{1}{4} \int_{\partial\Omega} \langle x, \nu \rangle Q(x) e^{4u} d\sigma + \int_{\partial\Omega} \left\{ \frac{1}{2} |\Delta u|^2 \langle x, \nu \rangle - 2 \frac{\partial u}{\partial \nu} \Delta u \right. \\ & \quad \left. - \langle x, \nabla u \rangle \frac{\partial \Delta u}{\partial \nu} - \langle x, \nabla \Delta u \rangle \frac{\partial u}{\partial \nu} + \langle x, \nu \rangle \langle \nabla u, \nabla \Delta u \rangle \right\} d\sigma. \end{aligned}$$

In particular, taking $\Omega = B_R$, we have

$$(2.33) \quad \begin{aligned} & \int_{B_R} Q(x) e^{4u} dx + \frac{1}{4} \int_{B_R} \langle x, \nabla Q \rangle e^{4u} dx \\ &= \frac{1}{4} \int_{\partial B_R} |x| Q e^{4u} d\sigma + \frac{1}{2} \int_{\partial B_R} |x| |\Delta u|^2 d\sigma - 2 \int_{\partial B_R} \frac{\partial u}{\partial r} \Delta u d\sigma - \int_{\partial B_R} |x| \frac{\partial u}{\partial r} \frac{\partial \Delta u}{\partial r} d\sigma. \end{aligned}$$

Proof. Multiplying (2.31) by $x \cdot \nabla u$, we have

$$(2.34) \quad \int_{\Omega} \nabla^2(x \cdot \nabla u) = \int_{\Omega} Q(x) e^{4u} (x \cdot \nabla u).$$

After integrating by parts and direct computation, we get

$$(\text{RHS of (2.34)}) = \frac{1}{4} \int_{\partial\Omega} \langle x, \nu \rangle Q(x) e^{4u} d\sigma - \int_{\Omega} \left(Q + \frac{1}{4} \langle x, \nabla Q \rangle e^{4u} \right) dx,$$

and

$$\begin{aligned} \text{(LHS) of (2.34)} &= -\frac{1}{2} \int_{\partial\Omega} |\Delta u|^2 \langle x, \nu \rangle + 2 \int_{\partial\Omega} \frac{\partial u}{\partial \nu} \Delta u + \int_{\partial\Omega} \langle x, \nabla u \rangle \frac{\partial \Delta u}{\partial \nu} \\ &\quad + \int_{\partial\Omega} \langle x, \nabla \Delta u \rangle \frac{\partial u}{\partial \nu} - \int_{\partial\Omega} \langle x, \nu \rangle \langle \nabla u, \nabla \Delta u \rangle. \end{aligned}$$

Thus we establish (2.32). Taking $\Omega = B_R$, we immediately obtain (2.33) from (2.32). Note that the singularity at the origin is insignificant that contributes nothing to the final form of the Pohozaev identity. \square

From the Pohozaev identity we shall determine the exact value of α .

Lemma 2.6. *Let u be a solution of (1.12). Assume $|u(x)| = o(|x|^2)$ at ∞ , then $\alpha = 2(1 + \gamma)$.*

Proof. Taking $Q(x) = 6|x|^{4\gamma}$ in (2.33), we have

$$\begin{aligned} (2.35) \quad & 6(1 + \gamma) \int_{B_R} |x|^{4\gamma} e^{4u} dx \\ &= \frac{1}{4} \int_{\partial B_R} 6r^{4\gamma+1} e^{4u} d\sigma + \frac{1}{2} \int_{\partial B_R} r |\Delta u|^2 - 2 \int_{\partial B_R} \frac{\partial u}{\partial r} \Delta u - \int_{\partial B_R} r \frac{\partial u}{\partial r} \frac{\partial \Delta u}{\partial r}. \end{aligned}$$

In view of Lemma 2.4, we have obtained

$$u(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|y|}{|x-y|}\right) |y|^{4\gamma} e^{4u(y)} dy + C_0,$$

and $e^{u(y)} \geq |y|^{-4\alpha}$ for $|y|$ large enough. Then $|y|^{4\gamma} e^{4u(y)} \geq |y|^{4(\gamma-\alpha)}$. Hence $|y|^{4\gamma} e^{4u(y)} \in L^1(\mathbb{R}^4)$ implies $\alpha > 1 + \gamma$. On the other hand, by the representation of u and direct calculations, there hold

$$\begin{aligned} \frac{\partial u}{\partial r}(x) &= -\frac{3}{4\pi^2} \int_{\mathbb{R}^4} \frac{x \cdot (x-y)}{|x||x-y|^2} |y|^{4\gamma} e^{4u(y)} dy, \\ \Delta u(x) &= -\frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x-y|^2} |y|^{4\gamma} e^{4u(y)} dy, \end{aligned}$$

and

$$\frac{\partial}{\partial r} \Delta u(x) = \frac{3}{\pi^2} \int_{\mathbb{R}^4} \frac{x \cdot (x-y)}{|x||x-y|^4} |y|^{4\gamma} e^{4u(y)} dy,$$

Recall the definition of u , Lemma 2.4 and $\alpha > 1 + \gamma$, then we have

$$\begin{aligned} (2.36) \quad & \lim_{r \rightarrow +\infty} \frac{\partial u}{\partial r} = 0, \quad \lim_{r \rightarrow +\infty} r \frac{\partial u}{\partial r} = -\alpha, \\ & \lim_{r \rightarrow +\infty} r^2 \Delta u = -2\alpha, \quad \lim_{r \rightarrow +\infty} r^3 \frac{\partial}{\partial r} \Delta u = 4\alpha, \end{aligned}$$

where $r = |x|$. Therefore, applying the Pohozaev identity (2.35) and (2.36), we have

$$8\pi^2(1 + \gamma)\alpha = 4\pi^2\alpha^2,$$

which leads to $\alpha = 2(1 + \gamma)$. \square

Now we can determine the asymptotic behavior of u at infinity using the exact value of α .

Lemma 2.7. *Let u be a solution of (1.12) and suppose $|u(x)| = o(|x|^2)$ at ∞ . Then there exist $c_0 \in \mathbb{R}$, $\tau > 0$ such that for $|x| > 1$,*

$$(2.37) \quad u(x) = -2(1 + \gamma) \log |x| + c_0 + O(|x|^{-\tau}), \quad |x| > 1$$

and

$$(2.38) \quad \nabla^j(u(x) + 2(1 + \gamma)) = O(|x|^{-\tau-j}), \quad j = 1, 2, 3, 4, \quad |x| > 1.$$

In particular

$$(2.39) \quad -\Delta u(x) = \frac{4(1 + \gamma)}{|x|^2} + O(|x|^{-2-\tau}), \quad |x| > 1.$$

Proof. Let $w(x) = u(\frac{x}{|x|^2}) - 2(1 + \gamma) \log |x|$, then from the equation of u and the assumption, we see w satisfies

$$(2.40) \quad \begin{cases} \Delta^2 w(x) = 6|x|^{4\gamma} e^{4w(x)}, & \text{in } \mathbb{R}^4 \setminus \{0\}, \\ |w(x)| = o(\log \frac{1}{|x|}), \quad |\Delta w(x)| = o(\frac{1}{|x|^2}), & \text{as } |x| \rightarrow 0. \end{cases}$$

Let $h(x)$ be a solution of

$$(2.41) \quad \begin{cases} \Delta^2 h(x) = 6|x|^{4\gamma} e^{4w(x)}, & \text{in } B_1 \setminus \{0\}, \\ h(x) = w(x), \quad \Delta h(x) = \Delta w(x), & \text{on } \partial B_1. \end{cases}$$

and $q(x) = w(x) - h(x)$. Then $q(x)$ satisfies

$$(2.42) \quad \begin{cases} \Delta^2 q(x) = 0, & \text{in } B_1, \\ q(x) = \Delta q = 0, & \text{on } \partial B_1, \\ |q(x)| = o(\log \frac{1}{|x|}), \quad |\Delta q(x)| = o(\frac{1}{|x|^2}), & \text{as } |x| \rightarrow 0. \end{cases}$$

First for Δq , since its growth near the singular source is weaker than fundamental solutions, the singularity is removable, thus $\Delta q = 0$ in B_1 . By exactly the same reason we further conclude that $q \equiv 0$ in B_1 . That means $w(x) = h(x) \in C^{0,\tau}(\bar{B}_1)$. It suffices to consider the regularity of h in B_1 .

Note that

$$\begin{aligned} |x|^{4\gamma} e^{4w(x)} &= |x|^{4\gamma} e^{4u(\frac{x}{|x|^2}) - 8(1+\gamma) \log |x|} \\ &\sim |x|^{4\gamma} |x|^{-8(1+\gamma)} |x|^{4\alpha} \sim |x|^{4\gamma}, \quad \text{near } 0, \end{aligned}$$

where we used Lemma 2.4 and Lemma 2.6.

By standard elliptic estimate, if $\gamma > -\frac{3}{4}$, $w \in C^{1,\tau}$ for some $\tau > 0$, if $-1 < \gamma \leq -\frac{3}{4}$ we have $w \in C^{0,\tau_1}$ for $\tau < 4(1 + \gamma)$. So in either case we use $\tau \in (0, 1)$ to have the following expansion of u :

$$u(x) = -2(1 + \gamma) \log |x| + c_0 + O(|x|^{-\tau}), \quad |x| > 1.$$

Based on this we can use standard elliptic estimate to obtain corresponding gradient estimates: now $|x|^{4\gamma} e^{4u}$ can be written as

$$(2.43) \quad |x|^{4\gamma} e^{4u} = e^{c_0} r^{-8-4\gamma} + O(r^{-8-4\gamma-\tau}), \quad r = |x| > 1$$

Using this in the expression of $u(x)$ in (2.26) and $-\Delta u$ in (2.5) it is easy to obtain (2.39) for Δu . Then the asymptotic behavior for other derivatives in (2.38) is a direct consequence of standard elliptic estimates. \square

Remark 2.8. *It is important to observe from (2.43) that the function $|x|^{4\gamma} e^{4u(x)}$ is strictly decreasing in $r = |x|$ for r large.*

2.2. Classification of entire solutions for the case $-1 < \gamma < 0$.

In this subsection, we will show the solution of (2.14) or (1.12) has radial symmetry and uniqueness property up to scaling if $-1 < \gamma < 0$. Similar to [31], we will use the method of moving planes. But the situation for singular equation is a lot harder since it is difficult to obtain precise asymptotic behavior of u first without knowing its radial symmetry. In this proof the integral expressions of u and Δu play a crucial role.

Suppose that u is a smooth entire solution of (1.12) with $|u(x)| = o(|x|^2)$ at ∞ . Recall $-\Delta u > 0$ in \mathbb{R}^4 and (2.37)~(2.38), so we will apply the method of moving planes to $-\Delta u$. Let $v(x) = -\Delta u(x)$. Then by Lemma 2.7,

$$(2.44) \quad v(x) = \frac{4(1+\gamma)}{|x|^2} + O(|x|^{-2-\tau}), \quad |x| > 1.$$

First, we state some conventional notations for moving planes. For any $\lambda \in \mathbb{R}$, let $T_\lambda = \{x \in \mathbb{R}^4 : x_1 = \lambda\}$, $\Sigma_\lambda = \{x : x_1 > \lambda\}$ and $x^\lambda = (2\lambda - x_1, x_2, x_3, x_4)$ be the reflection point of x with respect to T_λ . In addition, when applying the method of moving plane, we need two lemmas in [15], also can be seen in [31].

Proof of Theorem 1.3. Lemma 2.6 establishes (i) in Theorem 1.3. Next we aim to prove the radial symmetry of solutions by the method of moving planes.

Step 1: We start moving planes along x_1 -direction. For any λ , we consider $w_\lambda(x) = u(x) - u(x^\lambda)$ in Σ_λ . Then $w_\lambda(x)$ satisfies

$$(2.45) \quad \begin{cases} \Delta^2 w_\lambda(x) = b_\lambda(x) w_\lambda(x), & \text{in } \Sigma_\lambda, \\ w_\lambda(x) = \Delta w_\lambda(x) = 0, & \text{on } \partial T_\lambda, \end{cases}$$

where

$$b_\lambda(x) = 6 \frac{|x|^{4\gamma} e^{4u(x)} - |x^\lambda|^{4\gamma} e^{4u(x^\lambda)}}{u(x) - u(x^\lambda)}.$$

First we claim that there exists $\lambda_0 < 0$ such that $w_{\lambda_0} > 0$ and $v(x) - v_{\lambda_0}(x) > 0$ in Σ_{λ_0} . Using the expression of u in (2.26) we can obviously write $u(x) - u_\lambda(x)$ as

$$u(x) - u_\lambda(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log \frac{|x^\lambda - y|}{|x - y|} |y|^{4\gamma} e^{4u(y)} dy = \int_{\Sigma_\lambda} + \int_{\mathbb{R}^4 \setminus \Sigma_\lambda},$$

where $u_\lambda(x) = u(x^\lambda)$. Changing the integration over $\mathbb{R}^4 \setminus \Sigma_\lambda$ to Σ_λ by a change of variable: $z = (z_1, z_2, z_3, z_4) = (2\lambda - y_1, y_2, y_3, y_4)$ we see that

$$(2.46) \quad u(x) - u_\lambda(x) = \frac{3}{4\pi^2} \int_{\Sigma_\lambda} \log \frac{|x^\lambda - y|}{|x - y|} (|y|^{4\gamma} e^{4u(y)} - |y^\lambda|^{4\gamma} e^{4u(y^\lambda)}) dy.$$

Here we remark that writing w_λ in the form of (2.46) is crucial for our argument, it prevents us from very delicate asymptotic analysis for u and its derivatives. From

(2.46) we claim that when λ is very negative and when $x \in \Sigma_\lambda$, $u - u_\lambda > 0$ in Σ_λ . Indeed, first it is obvious that $\log(|x^\lambda - y|/|x - y|) > 0$. Next we observe the integration of y over B_R and outside B_R , respectively. For integration over B_R , we clearly have $|y|^{4\gamma} e^{4u} > c_0 > 0$ for some $c_0 > 0$. Based on the asymptotic behavior of u we can easily make $|y^\lambda|^{4\gamma} e^{4u(y^\lambda)} < c_0$ for λ very negative. For $y \in \Sigma_\lambda \setminus B_R$, by Remark 2.8 we have

$$|y|^{4\gamma} e^{4u(y)} > |y^\lambda|^{4\gamma} e^{4u(y^\lambda)} \text{ because } |y^\lambda| > |y|.$$

We also observe that for $v = -\Delta u$,

$$v(x) - v_\lambda(x) = \frac{3}{2\pi^2} \int_{\mathbb{R}^4} \left(\frac{1}{|x - y|^2} - \frac{1}{|x^\lambda - y|^2} \right) |y|^{4\gamma} e^{4u(y)} dy.$$

After a similar transformation we have

$$v(x) - v_\lambda(x) = \frac{3}{2\pi^2} \int_{\Sigma_\lambda} \left(\frac{1}{|x - y|^2} - \frac{1}{|x^\lambda - y|^2} \right) (|y|^{4\gamma} e^{4u(y)} - |y^\lambda|^{4\gamma} e^{4u(y^\lambda)}) dy.$$

By exactly the same reasoning we see that for λ very negative, $v(x) > v_\lambda(x)$ in Σ_λ .

Let λ_0 be the starting position for the moving plane process and let $\bar{\lambda} \leq 0$ be the upper limit:

$$(2.47) \quad \bar{\lambda} := \sup \{ \lambda \leq 0 : v(x^\mu) < v(x) \text{ for all } x \in \Sigma_\lambda \text{ and } \mu \leq \lambda \}$$

Next we claim that $\bar{\lambda} = 0$. If this is not the case, we have $\bar{\lambda} < 0$. Looking at the equation for $w_{\bar{\lambda}}$:

$$(2.48) \quad \Delta^2 w_{\bar{\lambda}}(x) = 6|x^{\bar{\lambda}}|^{4\gamma} (e^{4u(x)} - e^{4u_{\bar{\lambda}}(x)}) + 6(|x|^{4\gamma} - |x^{\bar{\lambda}}|^{4\gamma}) e^{4u(x)}, \quad x \in \Sigma_{\bar{\lambda}}.$$

On one hand the continuity gives $\Delta w_{\bar{\lambda}}(x) \leq 0$, since $w_{\bar{\lambda}} \rightarrow 0$ as $|x| \rightarrow +\infty$ and $w_{\bar{\lambda}}|_{T_{\bar{\lambda}}} = 0$. On the other hand we have $|x^{\bar{\lambda}}| > |x|$. This strict inequality and the strong maximum principle combined gives

$$w_{\bar{\lambda}}(x) > 0, \quad \Delta w_{\bar{\lambda}}(x) < 0, \quad \text{in } \Sigma_{\bar{\lambda}}.$$

Then standard procedure can be applied to move the plane a little further to the right of $\bar{\lambda}$. A contradiction to the definition of $\bar{\lambda}$. The fact that $\bar{\lambda} = 0$ means $u(x) \geq u(x^\lambda)$ in the x_1 direction. Applying the moving plane method to all directions we obtain the radial symmetry of u .

Step 2: Now we prove the uniqueness of the solution of (1.12) modulus scaling in (1.13).

Let w_1 and w_2 be two radial solutions of (1.12). Then w_i satisfies $w'_i(0) = w'_i(0) = 0$. Without loss of generality, we may assume $w_1(0) = w_2(0)$, on account of the rescale invariance. By the uniqueness of ODE, we only need to prove $w''_1(0) = w''_1(0)$.

If $w''_1(0) < w''_2(0)$, then $w_1(r) < w_2(r)$ for small $r > 0$. We will prove $w_1(0) < w_2(0)$ for all $r > 0$. Suppose there exists $r_0 > 0$ such that $w_1(r_0) = w_2(r_0)$ and $w_1(r) < w_2(r)$ for $0 < r < r_0$. Then by (1.12), we have

$$\frac{\partial}{\partial r} \Delta(w_1(r) - w_2(r)) = 6r^{4\gamma} (e^{4w_1(r)} - e^{4w_2(r)}) < 0, \quad 1 < r \leq r_0,$$

which together with the assumption implies

$$\Delta(w_1 - w_2) < 0, \quad \text{in } B(0, r_0).$$

Since $w_1(r_0) - w_2(r_0) = 0$, from the maximum principle, we have $w_1(r) - w_2(r) > 0$ for $0 < r < r_0$, which contradicts with $w_1(0) = w_2(0)$. Thus, $w_1(r) < w_2(r)$ for all $r > 0$. Hence $\frac{\partial}{\partial r} \Delta(w_1(r) - w_2(r)) < 0$ for all $r > 0$, which means $\Delta(w_1(r) - w_2(r))$ is decreasing in r . Thus $w_1(r) - w_2(r) \leq -cr^2$ as $r \rightarrow +\infty$ for some constant $c > 0$, which yields a contradiction to the assumption $w_i(r) = o(r^2)$ at ∞ .

Similarly, it is impossible for $w_1''(0) > w_2''(0)$. Thus, the radial solution of (1.12) is unique under the scaling $u_\lambda(x) = u(\lambda x) + (1 + \gamma) \log \lambda$ for some $\lambda > 0$, and it is valid for (2.14) after scaling. \square

Proof of Corollary 1.4: First we use the rough expansion of u in (2.37) to rewrite $|x|^{4\gamma} e^{4u}$ as

$$r^{4\gamma} e^{4u} = e^{4c_0} r^{-4-4\mu} + O(r^{-4-4\mu-\tau}), \quad r > 1, \quad \text{for some } \tau > 0.$$

Here we use $\mu = 1 + \gamma > 0$ for convenience. Then for $v = -\Delta u$, we have (see (2.39))

$$\lim_{r \rightarrow \infty} v'(r) r^3 = -8\mu.$$

The equation for v can be written as

$$v''(r) + \frac{3}{r} v'(r) = 6e^{4c_0} r^{-4-4\mu} + O(r^{-4-4\mu-\tau}), \quad r > 1.$$

Multiplying r^3 to both sides and integrating from r to ∞ , we have

$$r^3 v'(r) + 8\mu = -\frac{6e^{4c_0}}{4\mu} r^{-4\mu} + O(r^{-4\mu-\tau}), \quad r > 1.$$

Thus

$$v'(r) = -\frac{8\mu}{r^3} - \frac{3e^{4c_0}}{2\mu} r^{-3-4\mu} + O(r^{-3-4\mu-\tau}), \quad r > 1$$

and

$$v(r) = \frac{4\mu}{r^2} + \frac{3e^{4c_0}}{2\mu(2+4\mu)} r^{-2-4\mu} + O(r^{-2-4\mu-\tau}), \quad r > 1,$$

where $\lim_{r \rightarrow \infty} v(r) = 0$ is used. This expression of v , which is $-u'' - \frac{3}{r} u'$, gives

$$(2.49) \quad u(r) = -2\mu \log r + c_0 + c_1 r^{-4\mu} + O(r^{-4\mu-\tau}) + O(r^{-1}), \quad r > 1$$

where

$$c_1 = \frac{3e^{4c_0}}{32\mu^2(1+2\mu)(1-2\mu)}.$$

Now (2.49) has improved the estimate of $r^{4\gamma} e^{4u}$ to

$$(2.50) \quad r^{4\gamma} e^{4u} = e^{4c_0} r^{-4-4\mu} + e^{4(c_0+c_1)} r^{-4-8\mu} + O(r^{-4-8\mu-\tau}), \quad r > 1.$$

Using (2.50) in computation we obtain

$$u(r) = -2\mu \log r + c_0 + c_1 r^{-4\mu} + c_2 r^{-8\mu}, \quad r > 1$$

where

$$c_2 = \frac{3e^{4(c_0+c_1)}}{128\mu^2(1+4\mu)(1-4\mu)}.$$

In general

$$c_l = \frac{3e^{4(c_0+\dots+c_{l-1})}}{32l^2\mu^2(1-2l\mu)(1+2l\mu)}.$$

Finally the expansions of Δu as well as other derivatives can be obtained in standard argument. Corollary 1.4 is established. \square

Next, we consider the case without the assumption $|u(x)| = o(|x|^2)$ at ∞ .

The following lemma is similar to Lemma 3.3 in [31].

Lemma 2.9. *Suppose that $\Delta u = a$ in \mathbb{R}^n for a constant $a \in \mathbb{R}$ such that $\exp(u - c|x|^2) \in L^1(\mathbb{R}^n)$ for some $c > 0$. Then u is a polynomial of order at most 2.*

Proof. Let P be parabola that satisfies $\Delta P = a$. Then $u - P$ is a harmonic function, which also satisfies $\exp(u - P - c|x|^2) \in L^1(\mathbb{R}^n)$ for some $c > 0$ obviously. Thus Lemma 3.3 in [31] asserts that $u - P$ is a parabola, so is u . \square

If the $o(|x|^2)$ assumption is removed, we have the following result:

Theorem 2.10. *Let u be a solution of (1.12) with $\gamma > -1$. Then after an orthogonal transformation, $u(x)$ can be represented by (1.16), which has an asymptotic expansion of*

$$u(x) = -2(1+\gamma)\log|x| - \sum_{j=1}^4 a_j(x_j - x_j^0)^2 + c_0 + O(|x|^{-\tau}), \quad |x| > 1$$

for some $\tau > 0$. The function Δu satisfies

$$(2.51) \quad \Delta u(x) = -\frac{3}{2\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x-y|^2} |y|^{4\gamma} e^{4u(y)} dy - 2 \sum_{j=1}^4 a_j, \quad |x| > 1$$

where a_j are nonnegative constants and $x^0 = (x_1^0, \dots, x_4^0) \in \mathbb{R}^4$. Moreover, if $-1 < \gamma < 0$ and $a_i x_i^0 = 0$ for all $i = 1, \dots, 4$, u is radially symmetric with respect to the origin.

Proof of Theorem 2.10. Suppose that u is a solution of (1.12). Let v be (2.1) and $w(x) = u(x) + v(x)$. By Lemma 2.2, we have $\Delta w(x) \equiv -C_1 \leq 0$ in \mathbb{R}^4 . Since v has only logarithmic growth at infinity, the integrability of u guarantees that the assumption of Lemma 2.9 is satisfied. Thus there exist constants c_0 and a_{ij} ($i, j = 1, \dots, 4$) such that $a_{ij} = a_{ji}$ and

$$w(x) = \sum_{i,j,k=1}^4 (a_{ij}x_i x_j + b_k x_k) + c_0.$$

After an orthogonal transformation, we may assume

$$u(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|y|}{|x-y|}\right) |y|^{4\gamma} e^{4u(y)} dy - \sum_{j=1}^4 (a_j x_j^2 + b_j x_j) + c_0.$$

Since $|x|^{4\gamma} e^{4u} \in L^1(\mathbb{R}^4)$, we have $a_j \geq 0$ for all $j = 1, \dots, 4$, and $b_j = 0$ if $a_j = 0$. Hence, we can rewrite $u(x)$ as follows:

$$u(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|y|}{|x-y|}\right) |y|^{4\gamma} e^{4u(y)} dy - \sum_{j=1}^4 a_j (x_j - x_j^0)^2 + c_0$$

and (2.51) holds from the previous argument.

Next, we show the radial symmetry under the assumption $a_i x_i^0 = 0$ for all i . Clearly in this case

$$u(x) = \frac{3}{4\pi^2} \int_{\mathbb{R}^4} \log\left(\frac{|y|}{|x-y|}\right) |y|^{4\gamma} e^{4u(y)} dy - \sum_{j=1}^4 a_j x_j^2 + c_0$$

Let $\hat{u}(x) = u(x) + \sum_{j=1}^4 a_j x_j^2$. Then

$$(2.52) \quad \Delta^2 \hat{u}(x) = 6|x|^{4\gamma} e^{-4\sum_{j=1}^4 a_j x_j^2} e^{4\hat{u}(x)}, \quad \text{in } \mathbb{R}^4.$$

As in Lemma 2.7, we set $\hat{w}(x) = \hat{u}(\frac{x}{|x|^2}) - \alpha \log|x|$, then $|\hat{w}(x)| = o(\log \frac{1}{|x|})$ near 0 from Lemma 2.1 and Lemma 2.3, and \hat{w} satisfies

$$(2.53) \quad \Delta^2 \hat{w}(x) = 6|x|^{4\gamma} e^{-4\sum_{j=1}^4 a_j (\frac{x_j}{|x|^2})^2} e^{4\hat{w}(x)}, \quad \text{in } \mathbb{R}^4 \setminus \{0\}.$$

Note that $a_j \geq 0$, then we can follow the argument in the proof of Lemma 2.7 to obtain for $|x|$ large and a $\tau \in (0, 1)$ such that

$$(2.54) \quad \hat{u}(x) = -\alpha \log|x| + c_0 + O(|x|^{-\tau}),$$

and

$$(2.55) \quad \begin{cases} -\Delta \hat{u}(x) = \frac{2\alpha}{|x|^2} + O(|x|^{-2-\tau}), \\ -\frac{\partial}{\partial x_i} \Delta \hat{u}(x) = -4\alpha \frac{x_i}{|x|^4} + O(|x|^{-3-\tau}). \end{cases}$$

At this point, we establish (1.16). Finally, as in the proof of Theorem 1.3, we can apply the method of moving planes to show that $\hat{u}(x)$ is symmetric with respect to the origin. □

3. PRELIMINARIES FOR BLOWUP ANALYSIS

Let $G : M \times M \setminus \text{diag}$ denote the Green's function for the Paneitz operator

$$(3.1) \quad f(x) - \bar{f} = \int_M G(x, y) P_g f(y) dV_g(y), \quad \int_M G(x, y) dV_g(y) = 0,$$

where $\bar{f} = \frac{1}{\text{vol}_g(M)} \int_M f dV_g$ is the average of f over M . Then the weak form of (3.1) is

$$(3.2) \quad P_{g,y} G(x, y) = \delta_x - \frac{1}{\text{vol}_g(M)}.$$

Set R be the regular part of the Green function. Then by the Appendix A in [52], for y in a neighborhood of x ,

$$(3.3) \quad G(x, y) = -\frac{1}{8\pi^2} \log d_g(x, y) \chi + R(x, y).$$

where χ is a cut-off function to avoid cut locus. Using G we can decompose u_k as the sum of its regular part and singular part

$$(3.4) \quad u_k(x) = \tilde{u}_k(x) - 8\pi^2 \sum_{j=1}^N \gamma_j G(x, q_j).$$

Then \tilde{u}_k satisfies

$$(3.5) \quad P_g \tilde{u}_k + 2b_k = 2H_k e^{4\tilde{u}_k} \quad \text{in } M,$$

where

$$(3.6) \quad H_k(x) = h_k(x) \prod_{j=1}^N e^{-32\pi^2 \gamma_j G(x, q_j)}.$$

Clearly, (1.9) and (1.11) imply that

$$(3.7) \quad \int_M H_k e^{4\tilde{u}_k} dV_g \leq C$$

We will work with \tilde{u}_k in the later blow-up analysis.

Similar with [52], since the metric g may not be locally conformally flat, we will apply the *conformal normal coordinates*, whose existence has been proved in [29]. More specially, for $q \in M$, there exists a normal coordinate around q such that g can be deformed to \mathfrak{g} which satisfies $\det(\mathfrak{g}) = 1$. We use R_{ijkl} to denote the curvature tensor under \mathfrak{g} .

We will apply the expansions of the metric \mathfrak{g} and its derivatives in the conformal normal coordinates (seen in the Appendix B of [52]), which are

$$(3.8) \quad \begin{aligned} \mathfrak{g}_{ab}(x) &= \delta_{ab} + \frac{1}{3} R_{aijb} x^i x^j + O(r^3), \\ \mathfrak{g}^{ab}(x) &= \delta^{ab} - \frac{1}{3} R^{aijb} x_i x_j + O(r^3), \\ \partial_c \mathfrak{g}^{ab}(x) &= -\frac{1}{3} (R_{acib} + R_{aicb}) x^i + O(r^2), \\ \partial_{ab} \mathfrak{g}^{ab}(x) &= \frac{1}{3} R_{ia,a} x^i + O(r^2), \end{aligned}$$

In addition, the following Pohozaev identity from the Appendix D in [52] will play an important role when the blow-up analysis is carried out in the conformal normal coordinates.

Lemma 2. [52]

For equation $P_g u + 2b = 2he^{4u}$ in M and $\Omega = B(0, r)$, there holds

$$\begin{aligned}
 (3.9) \quad & \int_{\Omega} \left(2he^{4u} + \frac{1}{2} x^i \partial_i h e^{4u} \right) \\
 &= \int_{\partial\Omega} \left(\frac{1}{2} x^i \nu_i h e^{4u} - x^k \nu_j g^{ij} \partial_i (\Delta_g u) \partial_k u + \nu_j g^{ij} \Delta_g u \partial_i u + x^k \nu_j g^{ij} \Delta_g u \partial_{ik} u - \frac{1}{2} x^i \nu_i (\Delta_g u)^2 \right) \\
 &+ \int_{\Omega} \left(\Delta_g u \partial_i g^{ij} \partial_j u + x^k \Delta_g u \partial_{ik} g^{ij} \partial_j u + x^k \Delta_g u \partial_k g^{ij} \partial_{ij} u - 2b x^i \partial_i u \right) \\
 &+ 2 \int_{\partial\Omega} \left(R_{ij,l}(q) x^l x^k \nu_i \partial_j u \partial_k u + O(r^3) |\nabla u|^2 \right) \\
 &- \int_{\Omega} \left(2R_{ij,l}(q) (x^l \partial_j u \partial_i u + x^k x^l \partial_j u \partial_{ik} u) + 0(r^2) |\nabla u|^2 + O(r^4) |\nabla^2 u| \right)
 \end{aligned}$$

Note that we use $B(p, r)$ to denote a ball centered at p with radius r . Sometimes if the center is the origin, we use B_r instead of $B(0, r)$.

4. BLOW-UP ANALYSIS NEAR THE SINGULARITY

In this section, we focus on the blow-up analysis near q_j , and to simplify the notation, we will omit the superscript j . Similar to the argument in [52], we will work in the conformal normal coordinates near q from [29]. To be specific, we can find some function w defined on M , such that in a small neighborhood $B(q, \delta)$ of q , $\delta > 0$, we have

$$(4.1) \quad \det(\hat{g}) = 1$$

in the normal coordinates of the conformal metric $\hat{g} = e^{2w} g$. For convenience we just use g instead of \hat{g} . Note that in a neighborhood of q ,

$$(4.2) \quad w(x) = O(d_g(x, q)^2).$$

where $d_g(x, q)$ stands for the distance between x and q under metric g .

Using the conformal covariance property of P_g the function $u_k = \tilde{u}_k - w$ satisfies

$$(4.3) \quad P_g u_k + 2b_k = 2H_k e^{4u_k},$$

and

$$(4.4) \quad \int_M H_k e^{4u_k} dV_g \leq C,$$

where $2b_k = P_g w + 2b_k$ and $\tilde{H}_k = H_k e^{4w}$. For simplicity, we still denote \tilde{H}_k by H_k .

We still use G to denote the Green's function for P_g . Then we have the following Green's representation formula

$$(4.5) \quad u_k(x) = \bar{u}_k + 2 \int_M G(x, y) H_k(y) e^{4u_k(y)} dV_g(y) - 2 \int_M G(x, y) b_k(y) dV_g(y),$$

where \bar{u}_k is the average of u_k over (M, g) . Using the expression of G in (3.3), we have

$$(4.6) \quad u_k(x) = \bar{u}_k + 2 \int_M \left(-\frac{1}{8\pi^2} \log d_g(x, y) \chi \right) H_k(y) e^{4u_k(y)} dV_g(y) + \phi_k(x),$$

where

$$(4.7) \quad \phi_k(x) = 2 \int_M R(x, y) H_k(y) e^{4u_k(y)} dV_g(y) - 2 \int_M G(x, y) h_k(y) dV_g(y).$$

Note that $\det(g) = 1$ in $B(q, \delta)$, we have $dV_g(y) = dy$ in $B(q, \delta)$. Taking the difference of (4.6) evaluated at x and q , we get

$$(4.8) \quad \begin{aligned} & u_k(x) - u_k(q) \\ &= \frac{1}{4\pi^2} \int_M (\chi(r_q) \log |y - q| - \chi(r_x) \log d_g(x, y)) H_k(y) e^{4u_k(y)} dV_g(y) + \phi_k(x) - \phi_k(q), \end{aligned}$$

where $r_q = |y - q|$ and $r_x = d_g(x, y)$. Here since the coordinates are normal, we have $d_g(y, q) = |y - q|$.

Thanks to the cut-off function χ , we can replace the integral over M by an integral over $B(q, 2\delta)$:

$$(4.9) \quad \begin{aligned} & u_k(x) - u_k(q) \\ &= \frac{1}{4\pi^2} \int_{B(q, 4\delta)} (\chi(r_q) \log |y - q| - \chi(r_x) \log d_g(x, y)) H_k(y) e^{4u_k(y)} dV_g(y) \\ & \quad + \phi_k(x) - \phi_k(q), \quad x \in B(q, 2\delta). \end{aligned}$$

We next give an upper bound of the mass near q when u_k cannot blow up at q . Before stating such a small energy Lemma we point out that the function H_k can be written in the neighborhood of a singular source q as

$$(4.10) \quad H_k(x) = h_k(x) d_g(x, q)^{4\gamma},$$

with $h_k(q) \neq 0$. Using this notation, our result states as follows:

Lemma 4.1. *Let q be a singular source with index γ . If*

$$(4.11) \quad \int_{B(q, 2\delta)} 2h_k d_g(x, q)^{4\gamma} e^{4u_k(x)} dV_g < \min\{8\pi^2, 8\pi^2(1 + \gamma)\},$$

h_k is defined in (4.10). Then $u_k \leq C$ in $B(q, \delta)$.

Proof. Note that

$$P_g u_k = 2h_k d_g(x, q)^{4\gamma} e^{4u_k(x)} - 2h_k.$$

We write $u_k = u_{1k} + u_{2k}$ where u_{1k} is the solution of

$$(4.12) \quad \begin{cases} \Delta^2 u_{1k} = 2h_k d_g(x, q)^{4\gamma} e^{4u_k(x)} & \text{in } B(q, 2\delta) \\ u_{1k}(x) = \Delta u_{1k}(x) = 0 & \text{on } \partial B(q, 2\delta). \end{cases}$$

By Lemma 1 from [31] we have

$$(4.13) \quad \int_{B(q, 2\delta)} \exp \left\{ \frac{\tilde{\delta} |u_{1k}|}{\|2H_k e^{4u_k}\|_{L^1(B(q, 2\delta), g)}} \right\} dV_g \leq C,$$

with any $\tilde{\delta} \in (0, 32\pi^2)$ and some constant $C = C(\tilde{\delta}, \delta)$. On one hand in $B(q, 2\delta)$,

$$(4.14) \quad u_{1k}(x) = \int_{B(q, \delta)} G_\delta(x, y) 2h_k d_g(\eta, q)^{4\gamma} e^{4u_k} d\eta, \quad x \in B(q, 2\delta)$$

where $G_\delta(x, y)$ is the Green's function of Δ^2 on $B(q, 2\delta)$:

$$G_\delta(x, y) = -\frac{1}{8\pi^2} \log |x - y| + R_\delta(x, y),$$

with

$$G_\delta(x, y) = \Delta_y G_\delta(x, y) = 0, \quad \text{for } x \in B(q, 2\delta), y \in \partial B(q, 2\delta).$$

In particular for $x \in B(q, \frac{3}{2}\delta)$,

$$u_{1k}(x) = -\frac{1}{8\pi^2} \int_{B(q, 2\delta)} \log |x - q| 2\mathfrak{h}_k |\eta - q|^{4\gamma} e^{4u_k} d\eta + O(1), \quad x \in B(q, \frac{3}{2}\delta).$$

On the other hand the Green's representation formula of u_k gives

$$u_k(x) = u_{1k}(x) + u_{2k}(x) = \overline{u_k} + \int_M G(x, \eta) 2\mathfrak{h}_k d_g(x, q)^{4\gamma} e^{4u_k} d\eta.$$

Since the leading term of G and G_δ are both $-\frac{1}{8\pi^2} \log d_g(x, q)$, we have

$$u_{2k}(x) = \overline{u_k} + O(1).$$

From $\int_M e^{4u_k} dV_g \leq C$ and Jensen's inequality

$$e^{4\overline{u_k}} \leq \int_M e^{4u_k} dV_g \leq C.$$

Therefore, $u_{2k} \leq C$ in $B(q, \frac{3}{2}\delta)$. Now we focus on u_{1k} .

If $\gamma > 0$, (4.11) is

$$\int_{B(q, 2\delta)} 2\mathfrak{h}_k d_g(x, q)^{4\gamma} e^{4u_k(x)} dV_g < 8\pi^2.$$

Since u_{2k} is bounded from above in $B(q, \frac{3}{2}\delta)$, we see from (4.13) that there exists some $p > 1$ to make $e^{4u_{1k}} \in L^p(B(q, 2\delta))$:

$$(4.15) \quad \|2\mathfrak{h}_k d_g(x, q)^{4\gamma} e^{4u_k(x)}\|_{L^p(B(q, \frac{3}{2}\delta))} \leq C.$$

The estimate (4.15) leads to a L^∞ of u_{1k} in $B(q, \delta)$ based on two reasons. First the integration of (4.14) gives a L^1 bound of u_{1k} in $B(q, \frac{3}{2}\delta)$:

$$(4.16) \quad \|u_{1k}\|_{L^1(B(q, \frac{3}{2}\delta))} \leq C.$$

Second, the standard interior regularity results in [13] (Theorem 1 in Section 3 of [13]) gives

$$\|u_{1k}\|_{W^{4,p}(B(q, \delta))} \leq \|2\mathfrak{h}_k d_g(x, q)^{4\gamma} e^{4u_k(x)}\|_{L^p(B(q, \frac{3}{2}\delta))} + \|u_{1k}\|_{L^1(B(q, \frac{3}{2}\delta))} \leq C.$$

Thus we have obtained the L^∞ bound of u_{1k} in $B(q, \delta)$ by standard Sobolev embedding theorem.

If $-1 < \gamma < 0$, (4.11) is

$$\int_{B(q, 2\delta)} 2\mathfrak{h}_k d_g(x, q)^{-4|\gamma|} e^{4u_k(x)} dV_g < 8\pi^2(1 - |\gamma|).$$

Since $\frac{\tilde{\delta}}{8\pi^2(1-|\gamma|)} < \frac{4}{1-|\gamma|}$, this strict inequality makes it possible to choose $\tilde{\delta} \in (0, 32\pi^2)$ and $p > 1$ with

$$p < \frac{1}{|\gamma|} \quad p < \frac{\tilde{\delta}}{32\pi^2(1-|\gamma|)}.$$

Then Hölder inequality tells us that (4.15) is also true in this case and the L^∞ bound of u_{1k} over $B(q, \delta)$ follows immediately. The combination of the L^∞ bound of u_{1k} and the upper bound of u_{2k} implies that $u_k \leq C$ in $B(q, \delta)$. \square

Remark 4.2. *If q is a regular point the same proof obviously leads to a similar conclusion: Suppose $B(q, 2\delta)$ has no singular source and $\int_{B(q, 2\delta)} 2h_k e^{4u_k} < 8\pi^2$, there is an upper bound of u_k in $B(q, \delta)$.*

Another immediate consequence of Lemma 4.1 is that blowup sequence only converges to point measures. The following theorem takes one step further to assert that the point measure is quantized and the bubbling solutions tend to $-\infty$ away from blowup points.

Theorem 4.3 (Concentration and Quantization). *Let $\{u_k\}$ be a sequence of solution to (4.3) with (4.4). Assume that q is the only blow-up point of u_k in $B(q, \delta)$, then as $k \rightarrow +\infty$, along a subsequence, there hold*

$$(4.17) \quad u_k \rightarrow -\infty, \quad \text{uniformly on any compact set of } B(q, \delta) \setminus \{q\},$$

$$(4.18) \quad 2h_k d_g(x, q)^{4\gamma} e^{4u_k} \rightarrow \beta \delta_q, \quad \text{weakly in the sense of measure on } M,$$

with $\beta = 16\pi^2(1 + \gamma)$ and h_k is defined in (4.10).

In particular, along a subsequence,

$$(4.19) \quad 2 \int_{B(q, \delta)} H_k(x) e^{4u_k(x)} dV_g \rightarrow 16\pi^2(1 + \gamma), \quad \text{as } k \rightarrow +\infty.$$

Proof. Suppose q is the only blowup pint in $B(q, 2\delta)$, then we observe that for any given $K \subset\subset B(q, \delta)$

$$(4.20) \quad |u_k(x) - u_k(y)| \leq C(K)$$

because the $|x - q|$ is comparable to $|y - q|$ and the total integration of $h_k e^{4u_k}$ is bounded. It is easy to obtain (4.20) from the Green's representation formula. Then it is also immediate to observe that

$$(4.21) \quad |\nabla_g^j u_k(x)|_g \leq C(K), \quad \text{in } K, \quad j = 1, 2, 3$$

because when $x \in K$,

$$|\nabla^j G(x, y)| \leq C|x - y|^{-j}, \quad x, y \in B(q, 2\delta),$$

and u_k has a upper bound. These two facts imply (4.21) by trivial estimates. Then the equation for u_k further provides the estimate for the fourth order derivatives of u_k :

$$|\nabla^\alpha u(x)| \leq C(k), \quad x \in K, \quad |\alpha| = 4.$$

Now we prove (4.17) by contradiction. Suppose that there exists a point $x_0 \in B(q, \delta) \setminus \{q\}$ such that $\{u_k(x_0)\}_{k \in \mathbb{N}}$ is also bounded from below. By (4.20) we see

that u_k is bounded in L^∞ norm in any compact subset of $B(q, \delta) \setminus \{q\}$. This fact and the gradients estimates of u_k guarantee that along a subsequence

$$u_k \rightarrow u_0 \quad \text{in } C_{loc}^{3,\sigma}(B(q, 2\delta) \setminus \{q\}),$$

with some constant $\sigma \in (0, 1)$ and the limit function u_0 solves

$$P_g u_0(x) + 2b_0(x) = 2h_0(x) d_g(x, q)^{4\gamma} e^{4u_0(x)} \quad \text{in } B(q, 2\delta) \setminus \{q\}.$$

Around q we use $\beta(r)$ and its limit to describe the concentration of energy:

$$\begin{aligned} \beta_k(r) &= \int_{B(q,r)} 2h_k(x) d_g(x, q)^{4\gamma} e^{4u_k(x)} dV_g \\ \beta(r) &= \lim_{k \rightarrow +\infty} \beta_k(r), \quad \beta = \lim_{r \rightarrow 0} \beta(r). \end{aligned}$$

From Lemma 4.1 and Remark 4.2 we see that if q is singular point $\beta \geq \min\{8\pi^2(1 + \gamma), 8\pi^2\}$, if q is a regular point $\beta \geq 8\pi^2$. Fixing any $r > 0$ small, we integrate the equation of u_k in $B(q, r)$ to obtain

$$\int_{B(q,r)} (P_g u_k + 2b_k) dV_g = \int_{B(q,r)} H_k e^{4u_k} dV_g = \beta_k(r),$$

which implies

$$\lim_{r \rightarrow 0} \int_{B(q,r)} (P_g u_0 + 2b_0) dV_g = \beta.$$

Therefore, u_0 satisfies, in the distribution sense,

$$P_g u_0(x) + 2\hat{b}_0(x) = 2h_0(x) d_g(x, q)^{4\gamma} e^{4u_0(x)} + \beta \delta_q \quad \text{in } B(q, 2\delta)$$

Using the Green's representation formula for u_k , we have

$$(4.22) \quad u_0(x) = -\frac{\beta}{8\pi^2} \log d_g(x, q) + v(x) + w(x)$$

where the first term comes from the convolution of $-\frac{1}{8\pi^2} \log d_g(x, y) \chi$ with $\beta \delta_q$, the second term v comes from the convolution of $-\frac{1}{8\pi^2} \log d_g(x, y) \chi$ with $2h_0 d_g(x, q)^{4\gamma} e^{4u_0}$:

$$(4.23) \quad v(x) = -\frac{1}{4\pi^2} \int_{B(q,\delta)} \log d_g(x, y) h_0(y) d_g(y, q)^{4\gamma} e^{4u_0(y)} dV_g(y),$$

and w is the collection of insignificant other terms:

$$(4.24) \quad w \in C^4(M).$$

For v we use (4.23) to denote $v(x)$ in $B(q, \delta)$ and we extend v to make $v \equiv 0$ on $M \setminus B(q, 2\delta)$. Based on the definition of v we now show that $v \in L^\infty(B(q, \delta))$. In fact, from (4.23) we have this lower bound of v in $B(q, \delta)$:

$$(4.25) \quad v(x) \geq \frac{1}{4\pi^2} \log \frac{1}{\delta} \|V\|_{L^1(M)} \geq C \quad \text{in } B(q, \delta)$$

where

$$V(x) = h_0(x) d_g(x, q)^{4\gamma} e^{4u_0(x)}.$$

The lower bounds of \mathfrak{h}_0 and v lead to a lower bound for $V(x)$:

$$V(x) = \mathfrak{h}_0(x)d_g(x, q)^{4\gamma} e^{4\mathfrak{u}_0(x)} \geq C d_g(x, q)^{4\gamma - \frac{\beta}{2\pi^2}} e^{4v(x)+w(x)} \geq \frac{c}{d_g(x, q)^s}$$

with $s = \frac{\beta}{2\pi^2} - 4\gamma$ and suitable $c > 0$. Since $\|V\|_{L^1} < \infty$ we see immediately that $s < 4$, which is

$$(4.26) \quad \beta < 8\pi^2(1 + \gamma).$$

Thus there is no way for \mathfrak{u}_k to be bounded from below away from singular source unless $\gamma > 0$. We have proved (4.17) for $\gamma \leq 0$. For $\gamma > 0$ we have an upper bound for $V(x)$:

$$(4.27) \quad V(x) \leq \frac{c}{d_g(x, q)^s} e^{4v(x)} \quad \text{in } B(q, \delta), \quad \text{if } \gamma > 0.$$

To proceed with the proof of $v \in L^\infty$ we observe from (4.23) and direct computation that

$$\begin{cases} \Delta^2 v(x) = V(x) + \eta(x) & \text{in } B(q, 2\delta) \\ v(x) = \Delta v(x) = 0 & \text{on } \partial B(q, 2\delta), \end{cases}$$

where η is smooth in $B(q, 2\delta)$. Note that the boundary condition of v on $\partial B(q, 2\delta)$ is based on the smooth extension of v mentioned before. Now we employ a standard argument of Brezis-Merle [12] to obtain $e^{\kappa|v|} \in L^1(B(q, 2\delta), g)$ for any constant $\kappa > 0$. Indeed, let $0 < \varepsilon < 1/\kappa$ and $V = V_1 + V_2$ with $\|V_1\|_{L^1(B(q, 2\delta))} < \varepsilon$ and $V_2 \in L^\infty(B(q, 2\delta))$. Correspondingly we write $v = v_1 + v_2$, where v_1 solves

$$\begin{cases} \Delta^2 v_1(x) = V_1(x) & \text{in } B(q, 2\delta) \\ v_1(x) = \Delta v_1(x) = 0 & \text{on } \partial B(q, 2\delta), \end{cases}$$

and v_2 solves

$$\begin{cases} \Delta^2 v_2(x) = V_2(x) + \eta(x) & \text{in } B(q, 2\delta) \\ v_2(x) = \Delta v_2(x) = 0 & \text{on } \partial B(q, 2\delta), \end{cases}$$

Choosing $\tilde{\delta} = 32\pi^2 - 1$ in (4.13), we find

$$\int_{B(q, 2\delta)} e^{\kappa|v_1|} \leq \int_{B(q, 2\delta)} e^{\frac{|v_1|}{\|V_1\|_{L^1(B(q, 2\delta))}}} \leq C$$

based on the smallness of ε . Then by standard elliptic regularity theory, we have $v_2 \in L^\infty(B(q, 2\delta))$. Consequently, $e^{\kappa|v|} \in L^1(B(q, 2\delta), g)$. Since κ is large it is possible to use Hölder inequality to obtain $V \in L^{p'}(B(q, 2\delta), \hat{g})$ for some $p \in (1, \frac{4}{s})$ if $s \geq 0$ and $p \in (1, +\infty)$ if $s \leq 0$. Thus we have proved $v \leq C$ in $B(q, 2\delta)$.

From the L^∞ bound of v we can use two positive constants c_1 and c_2 to bound V from above and below

$$(4.28) \quad \frac{c_1}{d_g(x, q)^s} \leq V(x) = \mathfrak{h}_0(x)d_g(x, q)^{4\gamma} e^{4\mathfrak{u}_0(x)} \leq \frac{c_2}{d_g(x, q)^s}, \quad \text{if } \gamma > 0.$$

Next, we aim to derive a contradiction by taking advantage of the Pohozaev identity (3.9) in Section 3. Set $h(x) = \mathfrak{h}_0(x)d_g(x, q)^{4\gamma}$ and $\Omega = B(q, r)$ in (3.9), then direct computation and (A.1) give rise to

$$\int_{\Omega} x^i \partial_i (\mathfrak{h}_0(x)d_g(x, q)^{4\gamma}) e^{4\mathfrak{u}_0} = \int_{B(q, r)} (x^i \partial_i \mathfrak{h}_0(x)d_g(x, q)^{4\gamma} e^{4\mathfrak{u}_0} + x^i \mathfrak{h}_0 \partial_i d_g(x, q)^{4\gamma} e^{4\mathfrak{u}_0})$$

and

$$\begin{aligned}\partial_i d_g(x, q)^{4\gamma} &= 4\gamma d_g(x, q)^{4\gamma-1} \partial_i d_g(x, q) = 4\gamma d_g(x, q)^{4\gamma-1} (\partial_i |x - q| + C) \\ &= 4\gamma d_g(x, q)^{4\gamma-2} ((x - q)^i + C)\end{aligned}$$

Immediately, together with (A.1) we get

$$\begin{aligned}& \int_{\Omega} \frac{1}{2} (x - q)^i \partial_i (\mathfrak{h}_0(x) d_g(x, q)^{4\gamma}) e^{4u_0} \\ &= \int_{B(q, r)} 2\gamma \mathfrak{h}_0(x) d_g(x, q)^{4\gamma} e^{4u_0} + \int_{B(q, r)} \frac{1}{2} \partial_\nu \mathfrak{h}_0 d_g(x, q)^{4\gamma+1} e^{4u_0}\end{aligned}$$

Therefore, for $r \rightarrow 0$

$$(4.29) \quad (\text{LHS}) \text{ of } (3.9) = 2(1 + \gamma) \int_{B(q, r)} \mathfrak{h}_0 d_g(x, q)^{4\gamma} e^{4u_0} + O(r) = (1 + \gamma)\beta + o_r(1),$$

where $\lim_{r \rightarrow 0} o_r(1) = 0$. Denote the four integrals on the right hand side of (3.9) by I_1, I_2, I_3 and I_4 , respectively. Thanks to the expansions of g , which are in (3.8), we obtain that

$$\begin{aligned}|I_2| &\leq C \int_{B(q, r)} \left(r |\nabla^2 u_0| |\nabla u_0| + |x - q| |\nabla^2 u_0| |\nabla u_0| + r |x - q| |\nabla^2 u_0| + |x - q| |\nabla u_0| \right), \\ |I_3| &\leq C \int_{\partial B(q, r)} \left(|x - q|^2 |\nabla u_0|^2 + O(r^3) |\nabla u_0|^2 \right), \\ |I_4| &\leq C \int_{B(q, r)} \left(|x - q| |\nabla u_0|^2 + |x - q|^2 |\nabla^2 u_0| |\nabla u_0| + O(r^2) |\nabla u_0|^2 + O(r^4) |\nabla^2 u_0| \right),\end{aligned}$$

Next, we shall estimate $|\nabla^j u_0|$ in $B(q, r)$ for $j = 1, 2, 3$. Recalling (4.22)~(4.24), it is important to consider the $\nabla^j v$ in $B(q, r)$ for $j = 1, 2, 3$. By means of the Green's representation formula, we observe that

$$|\nabla^j v(x)| \leq C \int_{B(q, 2\delta)} \frac{1}{d_g(x, y)^j} V(y) + O(1)$$

In order to estimate the integral in the inequality above, we decompose $B(q, 2\delta)$ into two parts

$$\Omega_1 = B(q, 2\delta) \cap \left\{ d_g(x, y) \leq \frac{d_g(x, q)}{2} \right\}, \quad \Omega_2 = B(q, 2\delta) \setminus \Omega_1.$$

In this estimate we use $V(y) = O(1) d_g(y, q)^{-s}$ in (4.28). Hence

$$(4.30) \quad \int_{\Omega_1} \frac{1}{d_g(x, y)^j} V(y) dV_g(y) \leq \frac{C}{d_g(x, q)^s} \int_{B(x, \frac{d_g(x, q)}{2})} \frac{1}{d_g(x, y)^j} dV_g(y) \leq C d_g(x, q)^{4-s-j}.$$

Using (4.28) again, we obtain that

$$\tilde{I} := \int_{\Omega_2} \frac{1}{d_g(x, y)^j} V(y) dV_g(y) \leq C \int_{\Omega_2} \frac{1}{d_g(x, y)^j} \frac{1}{d_g(y, q)^s} dV_g(y).$$

Fixing some t as follows

$$(4.31) \quad t \in \begin{cases} (0, \frac{4}{s}), & s > 0, \\ (1, +\infty), & s \leq 0, \end{cases}$$

we have $-ts > -4$. It follows from the Hölder inequality that

$$\tilde{I} \leq C \left(\int_{\Omega_2} \frac{1}{d_g(x, y)^{jt^*}} dV_g(y) \right)^{\frac{1}{t^*}} \left(\int_{\Omega_1} \frac{1}{d_g(y, q)^{st}} dV_g(y) \right)^{\frac{1}{t}} \leq C \left(\int_{\frac{d_g(x, q)}{2}}^{\tilde{c}} \frac{1}{\rho^{jt^*-3}} d\rho \right)^{\frac{1}{t^*}}$$

where $t^* = \frac{t}{t-1}$ denotes the conjugate of t and \tilde{c} is some positive constant. Then direct computation and the fact $-ts > -4$ imply that

$$(4.32) \quad \tilde{I} = \int_{\Omega_2} \frac{1}{d_g(x, y)^j} V(y) dV_g(y) \leq \begin{cases} C |\log d_g(x, q)|^{\frac{1}{t^*}}, & \text{if } jt^* = 4, \\ C d_g(x, q)^{\frac{4}{t^*}-j} + C, & \text{if } jt^* \neq 4. \end{cases}$$

In view of (4.31), we get that

$$t^* \in \begin{cases} (\frac{4}{4-s}, +\infty), & s > 0, \\ (1, +\infty), & s \leq 0. \end{cases}$$

Hence, there holds $\frac{4}{t^*} - j < 4 - s - j$. Consequently, from (4.30) and (4.32) there exists some $\tau > 0$ such that for any $r \in (0, \delta)$

$$(4.33) \quad |\nabla^j v(x)| \leq C d_g(x, q)^{\tau-j} + C, \quad j = 1, 2, 3, \quad x \in B(q, r).$$

In fact, we may choose $\tau \in (0, 1)$ if $jt^* = 4$, and otherwise $\tau = \frac{4}{t^*}$. At this point, we obtain that

$$(4.34) \quad |\nabla^j u_0(x)| \leq C d_g(x, q)^{-j}, \quad j = 1, 2, 3 \quad x \in B(q, r).$$

Thus by virtue of (A.1) and (4.33), we may adjust $\tau > 0$ such that on $\partial B(q, r)$

$$\begin{aligned} \frac{\partial u_0}{\partial r} &= -\frac{\beta}{8\pi^2} \frac{1}{|x-q|} + O(r^\tau) \frac{1}{|x-q|} + O(1), \\ \Delta u_0 &= -\frac{\beta}{4\pi^2} \frac{1}{|x-q|^2} + O(r^\tau) \frac{1}{|x-q|^2} + O(1), \\ \frac{\partial \Delta u_0}{\partial r} &= \frac{\beta}{2\pi^2} \frac{1}{|x-q|^3} + O(r^\tau) \frac{1}{|x-q|^3} + O(1). \end{aligned}$$

Therefore, the estimates of I_2 , I_3 and I_4 can be improved:

$$\begin{aligned} |I_2| &\leq C \int_{B(q, r)} (rd^{-3} + d^{-2} + 1) dV_g \leq Cr^2, \\ |I_3| &\leq C \int_{\partial B(q, r)} r^2 d_g(x, q)^{-2} dV_g \leq Cr^3, \\ |I_4| &\leq C \int_{B(q, r)} d_g(x, q)^{-1} dV_g \leq Cr^3. \end{aligned}$$

Finally for I_1 and we use the expansions of g^{ij} to obtain

$$\begin{aligned} I_1 &= \int_{\partial B(q, r)} \left(-rv_i \partial_i (\Delta_g u_0) \partial_\nu u_0 + \Delta_g u_0 \partial_\nu u_0 + (x-q)^k v_i \Delta_g u_0 \partial_{ik} u_0 - \frac{1}{2} r (\Delta_g u_0)^2 \right) \\ &\quad + O(r) \\ &= \int_{\partial B(q, r)} \left(-rv_i \partial_i (\Delta u_0) \partial_\nu < x-q, u_0 >_{flat} + \Delta u_0 \partial_\nu u_0 - \frac{1}{2} r (\Delta u_0)^2 \right) + O(r), \end{aligned}$$

where we have used $\det(g) = 1$ in $B(q, \delta)$ and $\Delta_g u = \partial_i(g^{ij}\partial_j u)$. Consequently,

$$(4.35) \quad (\text{RHS of (3.9)}) = I_1 + o_r(1) = \frac{\beta^2}{16\pi^2} + o_r(1).$$

Combining (4.29) and (4.35), we derive that $\beta = 16\pi^2(1 + \gamma)$, which yields a contradiction to (4.26) in the case $\gamma > 0$. Therefore $u_k \rightarrow -\infty$ uniformly on any compact subset of $B(q, 2\delta) \setminus \{q\}$, $b_k(x)d_g(x, q)e^{4u_k(x)} \rightarrow 0$ uniformly on any compact subset of $B(q, 2\delta) \setminus \{q\}$ and

$$b_k(x)d_g(x, q)e^{4u_k(x)} \rightarrow \beta\delta_q \text{ weakly in the sense of measure on } M.$$

In the end, we show the quantization β is exactly $16\pi^2(1 + \gamma)$. To see this, set $c_k = \int u_k d\sigma$ and $\check{u}_k(x) = u_k(x) - c_k$. Then, we have $c_k \rightarrow -\infty$ and $\check{u}_k \rightarrow \check{u}$ in $C^4(B(q, \delta))$ as $k \rightarrow +\infty$. Moreover, there exists a smooth function \check{v} such that $\check{u}(x) = \frac{\beta}{8\pi^2} \log d_g(x, q) + \check{v}$. Taking advantage of the Pohozaev identity as before, we obtain $\beta = 16\pi^2(1 + \gamma)$ and

$$2b_k(x)d_g(x, q)^{4\gamma}e^{4u_k(x)} \rightarrow 16\pi^2(1 + \gamma)\delta_q.$$

□

Based on Theorem 4.3 and its proof, we immediately obtain the following corollaries.

Corollary 4.4. *Suppose that u_k satisfies the assumptions in Theorem 4.3, then along a subsequence, there holds*

$$u_k - c_k \rightarrow -2(1 + \gamma) \log d_g(x, q) + \hat{v}, \quad \text{in } C_{loc}^4(B(q, 2\delta) \setminus \{q\}),$$

where $c_k = \int_{\partial B(q, \delta)} u_k d\sigma_g \rightarrow -\infty$ and \hat{v} is a smooth function in $B(q, 2\delta)$.

Corollary 4.5. *Suppose that u_k satisfy*

$$P_g u_k(x) + 2b_k = 2H_k e^{4u_k} \quad \text{in } M$$

with $\int_{B(q_j, 2\delta)} 2H_k e^{4u_k} dV_g \rightarrow \rho_j < 16\pi^2(1 + \gamma_j)$ for some $j \in \{1, \dots, N\}$. Then $\{u_k\}$ is uniformly bounded from above on any subset of $B(q_j, 2\delta)$. In particular, $\{u_k\}$ can not blow up in $B(q_j, 2\delta)$.

5. CONCENTRATION-COMPACTNESS RESULT AND A PRIORI ESTIMATE

In this section, we aim to establish the concentration-compactness principle and a priori estimate basing on the result in the section 4. Indeed, we will derive the following concentration-compactness type result for the regular part of $\{u_k\}$.

Set

$$\rho_k = \int_M 2H_k e^{4u_k} dV_{u_k}.$$

Theorem 5.1 (Concentration-Compactness). *Let $\{\tilde{u}_k\}$ be a sequence of solution to (3.5) and (3.7) with $\rho_k \rightarrow \rho$. Then there exists a subsequence, still denoted $\{\tilde{u}_k\}$, for which one of the following alternative holds:*

- (i) $\sup_{\Sigma} |\tilde{u}_k| \leq C_{\Sigma}$, for any $\Sigma \subset\subset M$.
- (ii) $\sup_{\Sigma} \tilde{u}_k \rightarrow -\infty$, for any $\Sigma \subset\subset M$.

(iii) *There exist a finite set $S = \{p^1, \dots, p^m\} \subset M$ with $m \in \mathbb{N}$, and sequences of points $\{x_k^1\}_{k \in \mathbb{N}}, \dots, \{x_k^m\}_{k \in \mathbb{N}} \subset M$, such that for all $i = 1, \dots, m$*

$$x_k^i \rightarrow p^i, \quad \sup_{\Sigma} \tilde{u}_k \rightarrow -\infty \text{ for any } \Sigma \subset M \setminus S$$

and

$$2H_k e^{4\tilde{u}_k} \rightarrow \sum_{i=1}^m \beta_i \delta_{p^i} \text{ weakly in the sense of measures in } M.$$

Futhermore, $\beta_i \in 16\pi^2\mathbb{N}$ if $p^i \notin \{q_1, \dots, q_N\}$, and $\beta_i = 16\pi^2(1 + \gamma)$ if $p^i = q_j$ for some $j \in \{1, \dots, N\}$.

Proof of Theorem 5.1. We define S to be the set of blow-up points of u_k in M , that is,

$$S = \{x \in M : \exists x_k \in M, \text{ s.t. } x_k \rightarrow x \text{ and } \tilde{u}_k(x_k) \rightarrow +\infty \text{ as } k \rightarrow +\infty\}.$$

We distinguish two cases.

Case 1: $S \neq \emptyset$.

For $p \in S$, Lemma 4.1 say that the mass of $\{\tilde{u}_k\}$ near p is no less than $8\pi^2(1 + \gamma)$. Then finite integral assumption $\int_M H_k e^{4\tilde{u}_k} dV_g \leq C$ implies $\text{card}(S) \leq C$. We may denote $S = \{p_1, \dots, p_m\}$ with some $n \in \mathbb{N}$. Therefore, there exists $r_0 \in (0, 1)$ such that for any $p^i \in S$, p^i is the only blow-up point of \tilde{u}_k in $B(p^i, r_0)$. Therefore, from the results in [22] and Theorem 4.3, we obtain the alternative (iii).

Case 2: $S = \emptyset$.

In this case, we have $\sup_M \tilde{u}_k \leq C$, which implies $H_k e^{4\tilde{u}_k}$ is uniformly bounded in M . Taking into account of the Green's representation formula,

$$\tilde{u}_k(x) - \tilde{u}_k = \int_M G(x, y) H_k(y) e^{4\tilde{u}_k} dV_g(y) = O(1).$$

Hence, after taking a subsequence, the alternative (i) occurs if $\limsup_{k \rightarrow +\infty} \int_M \tilde{u}_k dV_g > -\infty$, the alternative (ii) holds if $\limsup_{k \rightarrow +\infty} \int_M \tilde{u}_k dV_g \rightarrow -\infty$,

□

Immediately, we derive the following two corollaries from Theorem 5.1.

Corollary 5.2. *Suppose that \tilde{u}_k satisfies the assumption in Theorem 5.1 and alternative (iii) occurs, then $\rho \in \Gamma$.*

Corollary 5.3. *Suppose that \tilde{u}_k satisfies the assumption in Theorem 5.1. Then for every $\rho \in \mathbb{R}^+ \setminus \Gamma$, there exists a constant C only depending on ρ , such that $\tilde{u}_k \leq C$. In particular, if we additionally assume $\int_M \tilde{u}_k dV_g = 0$ or $\int_M H_k e^{4\tilde{u}_k} dV_g \geq c$ for some constant $c > 0$, then*

$$\|\tilde{u}_k\|_{L^\infty(M)} \leq C.$$

The following result explains that Γ is some critical set to (1.8) or (1.9).

Proposition 5.4 (Critical Set). *Suppose $\text{Ker}(P_g) = \{\text{constants}\}$ and that $\{u_k\}$ is a sequence of solutions to (1.9)~(1.10) with the coefficients satisfying (1.11). If the blow-up phenomena occur, then $\int_M 2bdV_g \in \Gamma$.*

Proof of Propertion 5.4 and Theorem 1.1. Since

$$\int_M H_k e^{4u_k} dV_g = \int_M h_k e^{4u_k} dV_g = \int_M b_k dV_g,$$

Proposition 5.4 and Theorem 1.1 obviously follow from the two corollaries above. \square

6. A SPHERICAL HARNACK INEQUALITY

As a corollary of the Concentration-Compactness theorem, we derive a spherical Harnack inequality near a singular source q_j when its strength γ_j is not an integer. Recall that if q_j is a blow-up point of u_k , then there exist $\delta > 0$ and a sequence $\{x_k^j\}_{k \in \mathbb{N}}$ such that $u_k(x_k) = \max_{B(q_j, \delta)} u_k \rightarrow +\infty$ and $S \cap B(q_j, \delta) = \{q_j\}$. Thus, we have

$$\varepsilon_k^j = \exp\left(-\frac{u_k(x_k^j)}{1 + \gamma_j}\right) \rightarrow 0, \quad \text{as } k \rightarrow +\infty.$$

Corollary 6.1. *Suppose that \tilde{u}_k satisfies the assumption in Theorem 5.1 and q_j is a blow-up point of \tilde{u}_k . If $\gamma_j \notin \mathbb{N}$, then $d_g(x_k^j, q_j) = O(\varepsilon_k^j)$, that is*

$$y_k^j := \frac{x_k^j - q_j}{\varepsilon_k^j} \rightarrow y^0 \in \mathbb{R}^4$$

and \tilde{u}_k has only one bubble at q_j .

Proof. Otherwise, we assume that

$$\frac{x_k^j - q_j}{\varepsilon_k^j} \rightarrow +\infty \quad \text{as } k \rightarrow +\infty.$$

For simplicity, we omit the superscript j and work on the u_k under the conformal normal coordinate. Let $r_k = d_g(x_k, q)$ and define the map $\varphi_k : B(0, \delta r_k^{-1}) \rightarrow B(q, \delta)$ by

$$\varphi_k : y \mapsto r_k y + q,$$

where on the right hand side we are using the conformal normal coordinates on $B(q, \delta)$. We use the notation $\check{f} = \varphi_{k*} f = f \circ \varphi_k$ to denote the pull-back of a function f defined on $B(q, \delta)$, and we let $\check{g}_k = r_k^{-2} \varphi_{k*} g$ be the blow-up metric, i.e., a rescaling of the pull-back metric.

We define

$$v_k(y) = \check{u}_k + (1 + \gamma) \log r_k, \quad y \in B(0, \delta/r_k).$$

Then v_k satisfies

$$\begin{cases} P_{g_k} v_k + 2\check{b}_k = 2\check{h}_k(y) d_{g_k}(y, 0)^{4\gamma} e^{4v_k}, & \text{in } B(0, \delta/r_k), \\ \int_{B(0, \delta/r_k)} \check{h}_k(y) d_{g_k}(y, 0)^{4\gamma} e^{4v_k} dV_{g_k} \leq C, \end{cases}$$

where $\check{b}_k(y) = b_k(r_k y + q)$, $\check{h}_k(y) = h_k(r_k y + q)$.

Denote $e_k = \frac{x_k - q}{r_k}$, then $e_k \rightarrow e_0 \in S^1 \subset \mathbb{R}^4$ and e_0 is a blow-up point of \tilde{v}_k . Applying Theorem 5.1, there exists a finite blow-up set $\tilde{S} = \{y^1, \dots, y^l\}$ such that $v_k \rightarrow -\infty$ uniformly on any compact subset of $\mathbb{R}^4 \setminus \tilde{S}$ and

$$2\check{h}_k(y)d_{g_k}(y, 0)^{4\gamma}e^{4v_k} \rightarrow \sum_{i=1}^l \alpha_i \delta_{y^i}, \quad \text{weakly in the sense of measure,}$$

where

$$\alpha_i = \lim_{k \rightarrow +\infty} \int_{B(y^i, r_0)} 2\check{h}_k(y)d_{g_k}(y, 0)^{4\gamma}e^{4v_k(y)}dV_{g_k} \quad \text{and} \quad B(y^i, r_0) \cap \tilde{S} = \{y^i\}.$$

First, we claim that 0 is not a blow-up point of v_k . Suppose $0 \in \tilde{S}$, then by means of Theorem 5.1 we have

$$\lim_{k \rightarrow +\infty} \int_{B(0, r_0)} \check{h}_k(y)d_{g_k}(y, 0)^{4\gamma}e^{4v_k(y)}dV_{g_k} = 16\pi^2(1 + \gamma).$$

On the other hand, since $e_0 \in S^1$ is also a blow-up point of v_k in $B(0, R)$ for R large. Then

$$\begin{aligned} 16\pi^2(1 + \gamma) &= \lim_{k \rightarrow +\infty} \int_{B(q, \delta)} 2\hat{h}_k(x)d_{g_k}(x, q)^{4\gamma}e^{4u_k(x)}dV_{g_k} \\ &= \lim_{k \rightarrow +\infty} \int_{B(0, \delta/r_k)} 2\check{h}_k(y)d_{g_k}(y, 0)^{4\gamma}e^{4v_k(y)}dV_{g_k} \\ &\geq \lim_{k \rightarrow +\infty} \int_{B(0, r_0) \cup B(e_0, r_0)} 2\check{h}_k(y)d_{g_k}(y, 0)^{4\gamma}e^{4v_k(y)}dV_{g_k} \\ &= 16\pi^2(1 + \gamma) + 16\pi^2, \end{aligned}$$

which yields a contradiction. Thus, $0 \notin \tilde{S}$.

As a consequence, we have

$$16\pi^2(1 + \gamma) = 16\pi^2 l$$

which is impossible, since $\gamma \notin \mathbb{N}$ while l is an integer. In other words, if γ is not an integer, then $d_g(x_k, q) = O(\varepsilon_k)$ and \hat{u}_k has only one bubble at q . \square

Theorem 1.2 is obviously equivalent to the following form:

Theorem 6.2 (A Spherical Harnack Inequality ($\gamma \notin \mathbb{N}$)). *Suppose that \tilde{u}_k satisfies the assumption in Theorem 5.1 and q_j is a blow-up point of \tilde{u}_k . If $\gamma_j \notin \mathbb{N}$, then near q_j there holds the following spherical Harnack inequality:*

$$(6.1) \quad \max_{x \in B(q_j, \delta)} \{\tilde{u}_k(x) + (1 + \gamma_j) \log |x - q_j|\} \leq C$$

with some constant C .

Proof. Suppose that (6.1) fails, then there exists a sequence $\{x_k\} \subset B(q_j, \delta)$ such that $\max_{x \in B(q_j, \delta)} \{\tilde{u}_k(x) + (1 + \gamma_j) \log |x - q_j|\} \rightarrow +\infty$. By means of the selection process and quantization property in [33, 22], we obtain that $\int_{B(q, \delta)} H_k(x)e^{4\tilde{u}_k(x)}dV_g \rightarrow$

$16\pi^2 n$ for some positive integer n . However, from Theorem 5.1, we know that

$$(6.2) \quad \int_{B(q,\delta)} H_k(x) e^{4\tilde{u}_k(x)} dV_g \rightarrow 16\pi^2(1+\gamma), \quad \text{as } k \rightarrow +\infty,$$

which is impossible. □

APPENDIX: COMPARISON BETWEEN $d_g(x, q)$ AND $|x - q|$

In this appendix, we will establish the comparison between the distance $d_g(x, q)$ and its derivatives and their Euclidean counterparts as in the Appendix B in [52]. We will follow the argument in [52] and give the detail for completeness. We claim that for $j = 0, 1, 2, 3$, there holds

$$(A.1) \quad \nabla^j (\log |x - y| - \log d_g(x, y)) = O(r^{2-j}), \quad x \in B(q, 2r) \setminus B(q, r/2).$$

We recall that g is the conformal normal metric centered at q , and we identify $x, y \in T_q M$ with $\exp_q x$ and $\exp_q y$ respectively, where \exp_q is the exponential map at q with respect to the metric g . Thus,

$$d_g(x, y) = d(\exp_q x, \exp_q y), \quad x, y \in B(q, \delta).$$

Also, we denote ∇_g by ∇ for convenience. First, let us note that the following simple estimates on d hold:

$$(A.2) \quad |\nabla^j (\log d_g(x, y))| \leq C|x - y|^{-j}, \quad j = 1, 2, 3, 4.$$

Set

$$f(x) = \log |x - y| - \log d_g(x, y), \quad x \in B(q, 2r) \setminus B(q, r/2).$$

We aim to show that

$$|\nabla^j f(x)| \leq C|x - q|^{2-j}, \quad j = 0, 1, 2, 3, \quad x \in B(q, 2r) \setminus B(q, r/2).$$

Let R , R_{ij} and R_{ijkl} respectively denote the scalar, Ricci and Riemann curvature of g . From the definitions of g and R_{ijkl}^i , we obtain that

$$\nabla^j R_{ijkl}^i(x) = O(1), \quad j = 1, 2.$$

In conformal normal coordinates, there holds $R(q) = R_{ij}(q) = |\nabla R(q)| = 0$. As a consequence, we have further

$$R(x) = O(r^2), \quad R_{ij}(x) = O(r).$$

We shall derive an estimate on $\Delta_g^2 f(x)$. By the definition of g and (A.4) in [52], that is

$$P_{g,y} \left(-\frac{1}{8\pi^2} \chi(r) \log d_g(x, y) \right) = \delta_x + E(x, y), \quad \text{with } E \text{ bounded,}$$

we have that

$$(A.3) \quad P_g \log d_g(x, q) = O(r^4), \quad x \in B(q, 2r) \setminus B(q, r/2).$$

In view of the rough estimates (A.2), we can estimate the term:

$$(P_g - \Delta_g^2) \log d_g(x, q) = \partial_m \left(g^{mi} \left(\frac{2}{3} R(x) g_{ij} - 2R_{ij}(x) \right) g^{lj} \partial_j (\log d_g(x, q)) \right) = O(r^{-1}).$$

Therefore, we can get

$$\Delta_g^2(\log d_g(x, q)) = O(r^{-1}), \quad x \in B(q, 2r) \setminus B(q, r/2).$$

Finally, we consider the term $\Delta_g^2(\log|x - q|)$. Since $\Delta^2(\log|x - q|) = 0$, it suffices to estimate $\Delta_g^2 - \Delta^2$. For any function u , the direct computation leads to

$$\begin{aligned} \Delta_g^2 u = & g^{ab} g^{ij} \partial_{ijab} u + 2 \partial_{ija} u (\partial_b g^{ab} g^{ij} + g^{ab} \partial_b g^{ij}) \\ (A.4) \quad & + \partial_{ij} u (\partial_a g^{ab} \partial_b g^{ij} + 2 g^{ai} \partial_{ab} g^{bj} + g^{ab} \partial_{ab} g^{ij} + \partial_a g^{ia} \partial_b g^{bj}) \\ & + \partial_j u (\partial_a g^{ab} \partial_{ib} g^{ij} + g^{ab} \partial_{iab} g^{ij}), \end{aligned}$$

where we have used $\det(g) = 1$. Using the expansion of g^{ab} :

$$g^{ab}(x) = \delta_{ab} - \frac{1}{3} R_{mab}(q) x^a x^b + O(|x - q|^3),$$

and replacing u by $\log|x - q|$ in (A.4) above, we obtain that

$$\Delta_g^2(\log|x - q|) = O(r^{-2}).$$

Consequently,

$$(A.5) \quad \Delta_g^2 f(x) = O(r^{-2}), \quad x \in B(q, 2r) \setminus B(q, r/2).$$

An estimate on the L^∞ -norm of $f(x)$ can easily be seen as follows:

$$(A.6) \quad d_g(x, q) = \int_0^{x(t)} \sqrt{x'_i(t) g_{ij}(t) x'_j(t)} dt = |x - q| (1 + O(r^2)),$$

which implies

$$(A.7) \quad f(x) = O(r^2), \quad x \in B(q, 2r) \setminus B(q, r/2).$$

Applying the elliptic theory to (A.5) and (A.7), we obtain the claim (A.1) and

$$(A.8) \quad d_g(x, q) = |x - q| (1 + O(r^2)).$$

REFERENCES

- [1] R. F. Adimurthi, M. Struwe, and F. Robert, Concentration phenomena for Liouville's equation in dimension four. *J. Eur. Math. Soc.*, **8** (2006), 171-180.
- [2] M. Ahmedou and C. B. Ndiaye, Morse theory and the resonant Q -curvature problem, Preprint arXiv:1409.7919.
- [3] A. Bahri and J.-M. Coron, On a nonlinear elliptic equation involving the critical Sobolev exponent: the effect of the topology of the domain. *Comm. Pure Appl. Math.* **41** (1988), no. 3, 253-294.
- [4] M. Ben Ayed, Mohamed, M. Ould Ahmedou, Existence and multiplicity results for a fourth order mean field equation. *J. Funct. Anal.*, **258** (2010), no. 9, 3165-3194.
- [5] Bartolucci, D; De Marchis, F.; Malchiodi, A. *Supercritical conformal metrics on surfaces with conical singularities*, Int. Math. Res. Not. IMRN 2011, no. 24, 5625-5643.
- [6] D. Bartolucci, G. Tarantello, Liouville type equations with singular data and their application to periodic multivortices for the electroweak theory. *Comm. Math. Phys.*, **229** (2002), 3-47.
- [7] W. Beckner, Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality. *Ann. of Math.*, **138-1** (1993), 213-242.
- [8] M.S. Berger, *Riemannian structure of prescribed Gaussian curvature for compact 2-manifolds*, J. Differential Geom. **5** (1971), 325-332.

- [9] T.P. Branson, S.A. Chang, P. Yang, Estimates and extremal problems for the log-determinant on 4-manifolds. *Comm. Math. Phys.*, **149** (1992), 241-262.
- [10] T.P. Branson, B.Oersted. Explicit functional determinants in four dimensions. *Proc. Amer. Math. Soc.*, **113** (1991), 669-682.
- [11] S. Brendle, Convergence of the Q-curvature flow on S^4 . *Adv. Math.*, **205**(1) (2006), 1-32.
- [12] H. Brezis, F. Merle, Uniform estimates and blow-up behaviour for solutions of $\Delta u = V(x)e^u$ in two dimensions. *Comm. Partial Differential Equations*, **16** (1991), 1223-1253.
- [13] F. E. Browder, On the regularity properties of solutions of elliptic differential equations. *Comm. Pure Appl. Math.*, **9** (1956), 351-361.
- [14] R. Buzano, H.-T. Nguyen, The Chern-Gauss-Bonnet formula for singular non compact four dimensional manifolds. *Communications in Analysis and Geometry* **27**(2019), 1697-1736.
- [15] L. Caffarelli, B. Gidas and J. Spruck, Asymptotic Symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth. *Comm. Pure Appl. Math.*, **42** (1989), 271-297.
- [16] S.A. Chang, J. Qing, P. Yang, Compactification of a class of conformally flat 4-manifold. *Invent. Math.*, **142** (2000), 65-93.
- [17] S.A. Chang, J. Qing, P. Yang, On the Chern-Gauss-Bonnet integral for conformal metrics on \mathbb{R}^4 . *Duke Math. J.*, **103** (2000), 523-544.
- [18] S.A. Chang, J. Qing, P. Yang, Some progress in conformal geometry. *SIGMA Symmetry Integrability Geom. Methods Appl.*, **3** (2007), 122.
- [19] S.A. Chang, P. Yang, Extremal metrics of zeta function determinants on 4-manifolds. *Ann. of Math.*, **142** (1995) 171-212.
- [20] Z. Djadli, A. Malchiodi, A fourth order uniformization theorem on some four manifolds with large total Q-curvature. *C. R. Math. Acad. Sci. Paris*, **340** (5) (2005) 341-346.
- [21] Z. Djadli, A. Malchiodi, Existence of conformal metrics with constant Q-curvature. *Ann. of Math.*, **168** (3) (2008) 813-858.
- [22] O. Druet, F. Robert, Bubbling phenomena for fourth-order four-dimensional PDEs with exponential growth. *Proc. Amer. Math. Soc.*, **134** (2006), 897-908.
- [23] H. Fang and B. Ma, Constant Q-curvature metrics on conic 4-manifolds, *Advances in Calculus of Variations*, 2020.
- [24] Fardoun, Ali; Regbaoui, Rachid Compactness properties for geometric fourth order elliptic equations with application to the Q-curvature flow. *J. Reine Angew. Math.* **734** (2018), 229-264.
- [25] C. Fefferman, K. Hirachi, Ambient metric construction of Q-curvature in conformal and CR-geometries. *Math. Res. Lett.*, **10**(5-6) (2003) 819-831.
- [26] M. Gursky, J. Viaclovsky, A fully nonlinear equation on four-manifolds with positive scalar curvature. *J. Differential Geom.*, **63**(1) (2003) 131-154.
- [27] Ali Hyder, Gabriele Mancini, Luca Martinazzi, Local and nonlocal singular Liouville equations in Euclidean spaces, *International Mathematics Research Notices*, Vol 00. No.0, pp 1-33.
- [28] Aleks Jevnikar, Yannick Sire, Wen Yang, Conformal Q-curvature metrics on supercritical conic 4-manifolds, in preparation.
- [29] J.M. Lee, T.H. Parker, The Yamabe problem. *Bull. Amer. Math. Soc. (N.S.)*, **17**(1) (1987) 37-91.
- [30] J. Li, Y. Li, P. Liu, The Q-curvature on a 4-dimensional Riemannian manifold (M, g) with $\int_M Q dV_g = 8\pi^2$. *Adv. Math.*, **231** (2012) 2194-2223.
- [31] C. S. Lin, A classification of solutions of conformally invariant fourth order equation in \mathbb{R}^n , *Comm. Math. Helv.*, **73** (1998), 206-231.
- [32] C.S. Lin, J.C. Wei, Sharp estimates for bubbling solutions of a fourth order mean field equation. *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, (5) **6** (4) (2007) 599-630.
- [33] A. Malchiodi, Compactness of solutions to some geometric fourth-order equations. *J. Reine Angew. Math.*, **594** (2006) 137-174.
- [34] A. Malchiodi, Conformal metrics with constant Q-curvature. *SIGMA Symmetry Integrability Geom. Methods Appl.*, **3** (2007) paper 120.
- [35] C.B. Ndiaye, Constant Q-curvature metrics in arbitrary dimension. *J. Funct. Anal.*, **251** (1) (2007) 1-58.

- [36] C. B. Ndiaye, Algebraic topological methods for the supercritical Q -curvature problem. *Adv. Math.*, **277** (2015), 56–99.
- [37] E. Picard, *De l'équation $\Delta u = ke^u$ sur une surface de Riemann fermée*, J. Math. Pure Appl. **9** (1893), 273–291.
- [38] E. Picard, *De l'intégration de l'équation $\Delta u = e^u$ sur une surface de Riemann fermée*, Crelle's J. **130** (1905), 243–258.
- [39] J. Qing, D. Raske, Compactness for conformal metrics with constant Q -curvature on locally conformally flat manifolds. *Calc. Var. Partial Differential Equations*, **26** (3) (2006) 343–356.
- [40] A. Malchiodi, M. Struwe, Q -curvature flow on S^4 . *J. Differential Geom.*, **73** (1) (2006) 1–44.
- [41] R.C. McOwen *Point Singularities and Conformal Metrics on Riemann Surfaces*, Proceedings of the American Mathematical Society **103** (1988), 222–224.
- [42] R.C. McOwen, *Conformal metrics in \mathbb{R}^2 with prescribed Gaussian curvature and positive total curvature*, Indiana Univ. Math. J. **34** (1985), 97–104.
- [43] F. Robert, Quantization effects for a fourth-order equation of exponential growth in dimension 4, *Proc. Roy. Soc. Edinburgh Sect. A*, **137** (2007), 531–553.
- [44] M. Struwe, Quantization for a fourth order equation with critical exponential growth. *Math. Z.*, **256** (2) (2007) 397–424.
- [45] M. Troyanov, “Metrics of constant curvature on a sphere with two conical singularities.” In Proceedings of the Third International Symposium on Differential Geometry (Peniscola 1988), 296–308. Lecture Notes in Mathematics 1410. Springer.
- [46] M. Troyanov, *Prescribing curvature on compact surfaces with conical singularities*, Trans. Amer. Math. Soc. **324** (1991), 793–821.
- [47] K. Uhlenbeck, J. Viaclovsky, Regularity of weak solutions to critical exponent variational equations. *Math. Res. Lett.*, **7** (2000) 651–656.
- [48] J. Wei, Asymptotic behavior of a nonlinear fourth order eigenvalue problem. *Comm. Partial Differential Equations*, **21** (1996) 1451–1467.
- [49] J. Wei, X. Xu, On conformal deformations of metrics on S^n . *J. Funct. Anal.*, **157**(1) (1998) 292–325.
- [50] J. Wei, D. Ye, Nonradial solutions for a conformally invariant fourth order equation in \mathbb{R}^4 . *Calc. Var. Partial Differential Equations*, **32** (2008), no. 3, 373–386.
- [51] J. Wei, L. Zhang, Estimates for Liouville equation with quantized singularities. *Adv. Math.* **380** (2021), 107606.
- [52] G. Weinstein, L. Zhang, The profile of bubbling solutions of a class of fourth order geometric equations on 4-manifolds. *J. Funct. Anal.*, **257** (2009) 3895–3929.
- [53] X. Xu, Classification of solutions of certain fourth-order nonlinear elliptic equations in \mathbb{R}^4 . *Pacific J. Math.*, **225** (2006), no. 2, 361–378.

MOHAMEDEN AHMEDOU, DEPARTMENT OF MATHEMATICS, GIESSEN UNIVERSITY, ARNDTSTRASSE 2, 35392, GIESSEN, GERMANY

Email address: Mohameden.Ahmedou@math.uni-giessen.de

LINA WU, DEPARTMENT OF MATHEMATICS, BEIJING JIAOTONG UNIVERSITY, BEIJING, 100044, CHINA

Email address: lnwu@bjtu.edu.cn

LEI ZHANG, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, 1400 STADIUM RD, GAINESVILLE FL 32611

Email address: leizhang@ufl.edu