# GLOBAL WELL-POSEDNESS AND DECAY FOR VISCOUS WATER WAVE MODELS

#### RAFAEL GRANERO-BELINCHÓN AND STEFANO SCROBOGNA

ABSTRACT. The motion of the free surface of an incompressible fluid is a very active research area. Most of these works examine the case of an inviscid fluid. However, in several practical applications, there are instances where the viscous damping needs to be considered. In this paper we derive and study a new asymptotic model for the motion of unidirectional viscous water waves. In particular, we establish the global well-posedness in Sobolev spaces. Furthermore, we also establish the global well-posedness and decay of a fourth order PDE modelling bidirectional water waves with viscosity moving in deep water with or without surface tension effects.

#### CONTENTS

1. Introduction	1
1.1. Main results	4
2. Proof of Theorem 1	6
2.1. The linear semigroup	6
2.2. Decay in the low regularity space	7
2.3. Boundedness in the high regularity space	10
2.4. Finishing the proof of Theorem 1	12
3. Derivation of $(5)$	13
4. Proof of Theorem 2	15
4.1. Local well-posedness	15
4.2. The linear semigroup	18
4.3. Decay in the low regularity space	18
4.4. Boundedness in the high regularity space	19
4.5. Finishing the proof of Theorem $2$	20
Acknowledgments	20
References	21

### 1. INTRODUCTION

The motion of waves in fluids has been a hot research topic since the XVIIIth century with the works of Laplace and Lagrange. On the one hand there is a large number of papers dealing with the free boundary Euler and Navier-Stokes equations [6, 30]. These are free boundary problems and as a consequence the domain of definition  $\Omega(t)$  of the functions (the bulk of the fluid) is an unknown of the system that has to be determined from the dynamics (see Figure 1).

 $Key\ words\ and\ phrases.$  Water waves, damping, moving interfaces, free-boundary problems.



FIGURE 1. Scheme of the problem

On the other hand the literature on asymptotic models of such free boundary problems is even larger (cf. [31]). These asymptotic models allow to have very good approximate description of the actual dynamics while simplifying the equations under study. In this direction there are many papers dealing with the case of inviscid fluids and, in particular dealing with asymptotic models for shallow water waves (see for instance [31] and the references therein) and models of water waves with small steepness (we refer to [1,3,33-35] for example). Similar small steepness asymptotics models can be derived for other free boundary problems, such as the Muskat problem [8, 15], see [18, 20, 38]. Many of such asymptotic models are used in different applications in Coastal Engineering and Physics.

Although it is a classical topic, the works studying the case of a viscous fluid are more scarce. The first works studying the case of a viscous water wave date back to Boussinesq [7], Basset [5] and Lamb [29]. Since then there are many other papers studying damped water waves. For instance, we refer to the manuscripts of Kakutani & Matsuuchi [26], Ruvinsky & Freidman [37], Longuet-Higgins [32], Jiang, Ting, Perlin & Schultz [23], Joseph & Wang [24], Wang & Joseph [39] and Wu, Liu & Yue [40].

According to the work by Dias, Dyachenko & Zakharov [10], the viscous damping of gravity water waves can be described by the following free boundary problem:

$$\Delta \phi = 0 \qquad \qquad \text{in } \Omega(t), \qquad (1a)$$

$$\rho\left(\phi_t + \frac{1}{2}|\nabla\phi|^2 + Gh\right) = -2\mu\partial_2^2\phi \qquad \text{on } \Gamma(t), \qquad (1b)$$

$$h_t = \nabla \phi \cdot (-\partial_1 h, 1) + 2\frac{\mu}{\rho} \partial_1^2 h$$
 on  $\Gamma(t)$ , (1c)

where h denotes the height of the wave,  $\phi$  is the velocity potential and  $G, \rho$  and  $\mu$  are the gravity acceleration, density and viscosity of the fluid.

Since its appearance, this system was considered by several other authors (see [11-14]). The need for simplified asymptotic models for damped waterwaves systems was highlighted at first by Longuet-Higgins, which in [32] stated that

> For certain applications, however, viscous damping of the waves is important, and it would be highly convenient to have equations and boundary conditions of comparable simplicity as for undamped waves.

In this spirit, Kakleas & Nicholls [25] derived a quadratic asymptotic model while Bae, Lin & Shin [4] derived a cubic asymptotic model of (1). The well-posedness of this quadratic model was studied by Ambrose, Bona & Nicholls [2] while the well-posedness of the full Dias-Dyachenko-Zakharov was proved by Ngom & Nicholls in [36] in the case of a nonzero surface tension and by Granero-Belinchón & Scrobogna [21] in the case in which the surface tension can be zero.

In a series of works [19,22], the authors, starting with the Dias-Dyachenko-Zakharov (1) free boundary problem, derived and studied the following bidirectional models of viscous water waves

$$\begin{cases} f_{tt} + 2\delta\Lambda^2 f_t + \Lambda f + \beta\Lambda^3 f + \delta^2\Lambda^4 f = \varepsilon \bigg\{ -\Lambda \left( (\mathcal{H}f_t)^2 \right) \\ + \partial_x \llbracket \mathcal{H}, f \rrbracket \Lambda f + \beta \partial_x \llbracket \mathcal{H}, f \rrbracket \Lambda^3 f + \delta \partial_x \llbracket \mathcal{H}, \mathcal{H}f_t \rrbracket \mathcal{H}\partial_x^2 f \\ + \delta\Lambda \left( \mathcal{H}f_t \mathcal{H}\partial_x^2 f \right) - \delta\partial_x \llbracket \partial_x^2, f \rrbracket \mathcal{H}f_t + \delta^2 \partial_x \llbracket \partial_x^2, f \rrbracket \Lambda \partial_x f \\ - \delta^2 \partial_x \llbracket \mathcal{H}, \partial_x^2 f \rrbracket \partial_x^2 f \bigg\}, \end{cases}$$
(2)  
$$f(x, 0) = f_0(x), \\ f_t(x, 0) = f_1(x), \end{cases}$$

and

$$\begin{cases} f_{tt} + 2\delta\Lambda^{2}f_{t} + \Lambda f + \beta\Lambda^{3}f + \delta^{2}\Lambda^{4}f = \varepsilon \left\{ -\Lambda \left( (\mathcal{H}f_{t})^{2} \right) \\ + \partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda f + \beta\partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda^{3}f + \delta\partial_{x} \llbracket \mathcal{H}, \mathcal{H}f_{t} \rrbracket \mathcal{H}\partial_{x}^{2}f \\ + \delta\Lambda \left( \mathcal{H}f_{t}\mathcal{H}\partial_{x}^{2}f \right) - \delta\partial_{x} \llbracket \partial_{x}^{2}, f \rrbracket \mathcal{H}f_{t} \right\}, \qquad (3)$$
$$f(x,0) = f_{0}(x), \\ f_{t}(x,0) = f_{1}(x), \end{cases}$$

where  $\varepsilon$  is the steepness parameters that measures the ratio between the anplitude and the wavelength,  $\delta > 0$  is a dimensionless parameter reflecting the viscous effects,  $\beta \geq 0$  is the Bond number measuring the ratio between capillary and gravity forces. The operators  $\mathcal{H}$  and  $\Lambda$  denote the Hilbert transform and the square root of the Laplacian

$$\widehat{\mathcal{H}f}(k) = -i\mathrm{sgn}(k)\widehat{f}(k), \quad \widehat{\Lambda f}(k) = |k|\widehat{f}(k), \quad (4)$$

and

$$\llbracket A, B \rrbracket f = A(Bf) - B(Af),$$

is the commutator between two operators acting on the function f. In what follows we consider  $(x,t) \in \mathbb{S}^1 \times [0,T]$  where  $\mathbb{S}^1$  denotes the interval  $[-\pi,\pi]$  with periodic boundary conditions. Furthermore, we will consider zero-mean initial data  $f_0$  and  $f_1$ .

The purpose of this work is twofold. First we prove that the system (3) is globally well posed for initial data which are sufficiently small.

Second we derive a new asymptotic model of unidirectional viscous water waves. In particular, we obtain the following nonlocal and nonlinear equation

$$2\varepsilon u_{t} = \mathcal{N}u_{x} + 2\delta \mathcal{N}u_{xx} + \mathcal{N}\mathcal{H}u - \beta \mathcal{N}\mathcal{H}\partial_{x}^{2}u + \delta^{2}\mathcal{N}\partial_{x}^{3}u - \varepsilon \mathcal{N}\bigg\{2uu_{x} + \Lambda [\![\mathcal{H}, \Lambda^{-1}u]\!]u + \beta \Lambda [\![\mathcal{H}, \Lambda^{-1}u]\!]\Lambda^{2}u - \delta \Lambda [\![\mathcal{H}, u]\!]u_{x} + \delta \partial_{x} (uu_{x}) + \delta \Lambda [\![\partial_{x}^{2}, \Lambda^{-1}u]\!]u\bigg\}.$$
(5)

where the operator

$$\mathcal{N} = (1 - \delta^2 \partial_x^2)^{-1} (1 - \delta \partial_x)$$

is defined in Fourier variables as

$$\widehat{\mathcal{N}} = \frac{1 - \delta i k}{1 + \delta^2 |k|^2}.$$

1.1. Main results. We start this section introducing some notation that we will use along the paper. We denote with C any positive constant independent of any physical parameter of the problem. The explicit value of C may vary from line to line.

We recall the definition of the homogeneous Sobolev spaces of fractional order

$$\dot{H}^{s} = \dot{H}^{s}\left(\mathbb{S}^{1}\right) = \left\{f \in L^{1} \mid \Lambda^{s} f \in L^{2}\right\},\$$

for any  $s \in \mathbb{R}$ . It is well known that for zero mean function we have that  $H^s = \dot{H}^s$ . As both equations preserve the zero mean property from now on we will always use the non-homogeneous notation in order to indicate a Sobolev space of regularity s. Similarly, we define the homogeneous Wiener spaces

$$\dot{A}^{s} = \dot{A}^{s} \left( \mathbb{S}^{1} \right) = \left\{ f \in L^{1} \mid \widehat{\Lambda^{s} f} \in \ell^{1} \right\},\$$

where  $\hat{f}$  denotes the Fourier series of f.

The first main result of this work is the following theorem:

**Theorem 1.** Let  $\delta > 0$  and  $\beta \ge 0$ . There exists a  $c_0 > 0$  such that for any  $(f_0, f_1) \in H^6 \times H^4$  such that

$$\|f_0\|_{H^6} + \|f_1\|_{H^4} \le c_0,$$

then, there exist a unique global solution  $(f, f_t)$  of (3) stemming from the initial data  $(f_0, f_1)$  which belongs to the energy space

$$f \in C\left(\mathbb{R}_+; H^6\right),$$
  
$$f_t \in C\left(\mathbb{R}_+; H^4\right) \cap L^2(\mathbb{R}_+; H^5).$$

Furthermore,

$$\begin{split} \|f\|_{A^0} + \|f_t\|_{A^0} &\leq C e^{-t\delta}, \\ \|f\|_{H^r} + \|f_t\|_{H^s} &\leq C e^{-C(\delta,r,s)t}, \qquad \forall \ (r,s) \in [0,6) \times [0,4) \,. \end{split}$$

4

Once the local existence and uniqueness was obtained in [22], we only need to provide with appropriate energy estimates. In order to do that we will consider a space of low regularity X and a space of high regularity Y, which will be explicitly defined below. Next we are going to define an energy having the form

$$\|\|(f, f_t)\|\|_T = \sup_{t \in [0,T]} \left\{ e^{\alpha t} \|(f(\bullet, t), f_t(\bullet, t))\|_X \right\} + \|(f, f_t)\|_Y \quad \text{for } \alpha > 0.$$

Equipped with this definition of energy, the rest of the proof is focused on obtaining an inequality of the form

$$|||(f, f_t)|||_T \leq C_0 (f_0, f_1) + P (|||(f, f_t)|||_T),$$

for certain polynomial P and constant  $C_0(f_0, f_1)$  that depends on the initial data. The previous inequality implies that for small enough  $C_0(f_0, f_1)$ , the solution satisfies

$$|||(f_0, f_t)|||_T \le 2\mathcal{C}_0(f_0, f_1),$$

for all T > 0, then a standard continuation argument allow us to extend the solution to arbitrary long time intervals.

Next, we derive a new asymptotic model of unidirectional viscous water waves. This new model takes the form (5). Our second main result is

**Theorem 2.** Let  $\delta > 0$  and  $\beta \ge 0$ . Then given and arbitrary zero mean  $u_0 \in H^2$ , there exists a unique local strong solution to (5)

$$u \in C([0, T^*], H^2) \cap L^2([0, T^*]; H^3),$$

for a small enough  $T^*$  depending only on  $||u_0||_{H^2}$  and the physical parameters of the problem. Furthermore, there exists a  $c_0 > 0$  such that for any  $u_0 \in H^2$ satisfying

$$||u_0||_{H^2} \le c_0,$$

then, there exist a unique global solution u of (5) stemming from the initial data  $u_0$  which belongs to the energy space

$$u \in C\left(\mathbb{R}_+; H^2\right) \cap L^2(\mathbb{R}_+; H^3).$$

Moreover,

$$\begin{split} \|u\|_{A^0} &\leq C e^{-\frac{o}{2}t}, \\ \|u\|_{H^r} &\leq C e^{-C(\delta,r)t}, \end{split} \qquad \forall \ r \in [0,2). \end{split}$$

In order to prove the local existence part of this theorem we use Picard's theorem together with energy estimates in  $H^2$  and the commutator structure of part of the nonlinearity. Once the local existence and uniqueness has been obtained, to ensure the global existence and decay we only need to provide with appropriate energy estimates. To do that we are going to define a modified energy  $|||f|||_T$  that has two different contributions. On the one hand we consider the low regularity space X where the solution will decay while on the other hand we will also regard a high regularity space Y where the solution will only remain bounded. The particular choice of X and Y will be clear below. Then the energy will take the form

$$|||f|||_T = \sup_{t \in [0,T]} \left\{ e^{\alpha t} \, ||f(\bullet,t)||_X \right\} + ||f||_Y \quad \text{for } \alpha > 0.$$

Equipped with this definition of energy, the rest of the proof will be devoted to obtain an inequality of the form

$$||u|||_T \le C_0 (f_0) + P(|||u|||_T),$$

for certain polynomial P of degree larger than 1, constant  $C_0(u_0)$  that depends on the initial data. The previous inequality implies that for small enough  $C_0(u_0)$ , the solution satisfies

$$|||u|||_T \leq 2\mathcal{C}_0(u_0),$$

for all T > 0, then a standard continuation argument allow us to extend the solution to arbitrary long time intervals.

### 2. Proof of Theorem 1

Without loss of generality, we consider  $\varepsilon = 1$  in (3). According to the result in [22], there is a local in time solution  $(f, f_t)$  for the problem (3). Let us define the modified energy

$$\begin{split} \| (f, f_t) \|_T &= e^{\delta T} \max_{t' \in [0, T]} \left\{ \left\| \left( f\left(t'\right), f_t\left(t'\right) \right) \right\|_{A^0} \right\} \\ &+ \max_{t' \in [0, T]} \left\{ \left\| f_t\left(t'\right) \right\|_{H^4} + \left\| f\left(t'\right) \right\|_{H^6} \right\}. \end{split}$$

The estimates of [22] assures us moreover that the solution exists at least in a time interval  $[0, T_{\text{max}}]$  where  $T_{\text{max}} = T_{\text{max}}(f_0, f_1)$  is the maximal lifespan of the solution.

2.1. The linear semigroup. We consider the linear nonhomogeneous problem

$$f_{tt} + 2\delta\Lambda^2 f_t + \Lambda f + \beta\Lambda^3 f + \delta^2\Lambda^4 f = F,$$
(6)

where F is a zero mean forcing. Let us denote with

$$u(x,t) = \begin{pmatrix} f(x,t) \\ f_t(x,t) \end{pmatrix}, \qquad u_0(x) = \begin{pmatrix} f_0(x) \\ f_1(x) \end{pmatrix},$$

so that (6) becomes

$$u_t + \mathcal{L}u = \begin{pmatrix} 0 \\ F \end{pmatrix}, \qquad \mathcal{L} = \begin{pmatrix} 0 & -1 \\ \Lambda + \beta \Lambda^3 + \delta^2 \Lambda^4 & 2\delta \Lambda^2 \end{pmatrix}.$$

Applying Duhamel principle we write  $u = u_{\rm L} + u_{\rm NL}$  where

$$u_{\rm L}(t) = e^{-t\mathcal{L}}u_0, \qquad \qquad u_{\rm NL}(t) = \int_0^t e^{-(t-t')\mathcal{L}} \begin{pmatrix} 0\\ F(t') \end{pmatrix} \mathrm{d}t'.$$

The eigenvalues of L are the Fourier multipliers

$$\lambda_{\pm}(n) = \delta |n|^2 \pm i \sqrt{|n| \left(1 + \beta |n|^2\right)},$$

so we see that the linear operator L induces both parabolic smoothing effects and oscillating behavior of the solution. Since the solution has zero mean, we have that  $\lambda_{\pm}(n) \neq 0$ . The two orthonormal eigenvectors associated to  $\lambda_{\pm}(n)$  are

$$\mathbf{e}_{\pm}\left(n\right) = \frac{1}{\sqrt{1 + \left|\lambda_{\pm}\left(n\right)\right|^{2}}} \left(\begin{array}{c} 1\\ -\lambda_{\pm}\left(n\right) \end{array}\right),$$

so that, if we denote

$$D = \begin{pmatrix} \lambda_{-} & 0 \\ 0 & \lambda_{+} \end{pmatrix},$$
  

$$S = \begin{pmatrix} 1 & 1 \\ -\lambda_{-} & -\lambda_{+} \end{pmatrix},$$
  

$$S^{-1} = \frac{1}{\lambda_{-} - \lambda_{+}} \begin{pmatrix} -\lambda_{+} & -1 \\ \lambda_{-} & 1 \end{pmatrix},$$

we have that

$$e^{-t\mathcal{L}} = S^{-1}e^{-tD}S.$$

With the above considerations we write  $u_{\rm L}$  and  $u_{\rm NL}$  in terms of  $f_0, f_1$  and F as

$$\hat{u}_{\mathrm{L}}(t) = \frac{1}{\lambda_{-} - \lambda_{+}} \begin{pmatrix} \lambda_{-}e^{-t\lambda_{+}} - \lambda_{+}e^{-t\lambda_{-}} \\ \lambda_{-} \left(e^{-t\lambda_{-}} - e^{-t\lambda_{+}}\right) \end{pmatrix} \hat{f}_{0} \\ + \frac{1}{\lambda_{-} - \lambda_{+}} \begin{pmatrix} \lambda_{+} \left(e^{-t\lambda_{+}} - e^{-t\lambda_{-}}\right) \\ \lambda_{-}e^{-t\lambda_{-}} - \lambda_{+}e^{-t\lambda_{+}} \end{pmatrix} \hat{f}_{1}, \\ \hat{u}_{\mathrm{NL}}(t) = \int_{0}^{t} \frac{1}{\lambda_{-} - \lambda_{+}} \begin{pmatrix} \lambda_{+} \left(e^{-(t-t')\lambda_{+}} - e^{-(t-t')\lambda_{-}}\right) \\ \lambda_{-}e^{-(t-t')\lambda_{-}} - \lambda_{+}e^{-(t-t')\lambda_{+}} \end{pmatrix} \hat{F}(t') \, \mathrm{d}t'.$$

We want to obtain now the decay rates of the linear semigroup. Let us at first check the time-decay of  $u_{\rm L}$ . We can compute that

$$\left|\frac{\lambda_{\pm}}{\lambda_{-} - \lambda_{+}}\right| \leq \frac{1}{2} \left(1 + \delta \sqrt{\frac{|n|}{1 + \beta}}\right),\tag{7}$$

since  $|n| \ge 1$  due to conservation of average. We deduce that, for j = 0, 1

$$\left|\frac{\lambda_{\pm}\left(n\right)}{\lambda_{-}\left(n\right)-\lambda_{+}\left(n\right)}e^{-t\lambda_{\pm}\left(n\right)}\hat{f}_{j}\left(n\right)\right| \leq \frac{e^{-\delta t}}{2}\left(1+\delta\sqrt{\frac{|n|}{1+\beta}}\right)\left|\hat{f}_{j}\left(n\right)\right|,$$

which in turn implies that

$$e^{\delta T} \max_{t' \in [0,T]} \left\| u_{\mathcal{L}}(t') \right\|_{A^0} \le C \left\| (f_0, f_1) \right\|_{A^{1/2}}.$$
(8)

Equivalently, we have that

$$\|e^{-t\mathcal{L}}\|_{A^{1/2}\mapsto A^0} \le Ce^{-\delta t}.$$
(9)

2.2. Decay in the low regularity space. If we write the equation in its mild formulation using Duhamel's principle, we have that the nonlinear forcing is given by  $F = \sum_{j=1}^{6} F_j,$ 

where

$$F_{1} = -\Lambda \left( \left( \mathcal{H}f_{t} \right)^{2} \right),$$
  

$$F_{2} = \partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda f,$$
  

$$F_{3} = \beta \partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda^{3} f,$$
  

$$F_{4} = \delta \partial_{x} \llbracket \mathcal{H}, \mathcal{H}f_{t} \rrbracket \mathcal{H} \partial_{x}^{2} f,$$

$$F_5 = \delta \Lambda \left( \mathcal{H} f_t \mathcal{H} \partial_x^2 f \right),$$
  
$$F_6 = \delta \partial_x \left[ \left[ \partial_x^2, f \right] \right] \mathcal{H} f_t.$$

The goal of the present computations is to provide a control of the form

 $\|F\left(t\right)\|_{A^{1/2}} \leq C e^{-\delta t(1+q)} \|\|(f,f_t)\|_T^2, \qquad t \in [0,T]\,, \ q > 0.$ 

We are going to use the Sobolev embedding

 $||a||_{A^s} \le C_{\delta} ||a||_{H^{s+1/2+\delta}} \le C ||a||_{H^{s+1}},$ 

together with interpolation between Sobolev spaces and the fractional product rule

$$|ab||_{A^s} \le C_s(||a||_{A^0} ||b||_{A^s} + ||a||_{A^s} ||b||_{A^0}) \le C_s ||a||_{A^s} ||b||_{A^s},$$

to estimate  $F_j$ . We compute

$$\begin{split} \|F_1\|_{A^{1/2}} &\leq C \|f_t\|_{A^0} \|f_t\|_{A^{3/2}} \\ &\leq C \|f_t\|_{A^0} \|f_t\|_{H^{5/2}} \\ &\leq C \|f_t\|_{A^0}^{1+3/8} \|f_t\|_{H^4}^{5/8} \end{split}$$

Using linear interpolation in Wiener spaces

$$||a||_{A^s} \le C ||a||_{A^r}^{s/r} ||a||_{A^0}^{1-s/r},$$

we find that

$$\begin{split} \|F_2\|_{A^{1/2}} &\leq \|\llbracket \mathcal{H}, f \rrbracket \Lambda f \|_{A^{3/2}} \\ &\leq C(\|f\|_{A^{3/2}} \|f\|_{A^1} + \|f\|_{A^{5/2}} \|f\|_{A^0}) \\ &\leq C \|f\|_{A^{5/2}} \|f\|_{A^0} \\ &\leq C \|f\|_{H^{7/2}} \|f\|_{A^0} \\ &\leq C \|f\|_{H^{6}}^{7/12} \|f\|_{A^0}. \end{split}$$

Similarly,

$$\begin{split} \|F_3\|_{A^{1/2}} &\leq C \|\llbracket \mathcal{H}, f \rrbracket \Lambda^3 f \|_{A^{3/2}} \\ &\leq C (\|f\|_{A^{3/2}} \|f\|_{A^3} + \|f\|_{A^{3+3/2}} \|f\|_{A^0}) \\ &\leq C \|f\|_{A^{9/2}} \|f\|_{A^0} \\ &\leq C \|f\|_{H^{11/2}} \|f\|_{A^0} \\ &\leq C \|f\|_{H^{6}}^{11/12} \|f\|_{A^0}^{1+1/12}. \end{split}$$

Similarly, we have that

$$\begin{split} \|F_4\|_{A^{1/2}} &\leq C \|\llbracket \mathcal{H}, \mathcal{H}f_t \rrbracket \mathcal{H}\partial_x^2 f \|_{A^{3/2}} \\ &\leq C \left( \|f_t\|_{A^0} \|f\|_{A^{7/2}} + \|f_t\|_{A^{3/2}} \|f\|_{A^2} \right) \\ &\leq C \left( \|f_t\|_{A^0} \|f\|_{H^6} + \|f_t\|_{A^0}^{1/2} \|f_t\|_{A^3}^{1/2} \|f\|_{A^0}^{3/5} \|f\|_{A^5}^{2/5} \right) \\ &\leq C \left( \|f_t\|_{A^0} \|f\|_{H^6} + \|f_t\|_{A^0}^{1/2} \|f_t\|_{H^4}^{1/2} \|f\|_{A^0}^{3/5} \|f\|_{H^6}^{2/5} \right), \\ \|F_5\|_{A^{1/2}} &\leq C \left( \|f_t\|_{A^0} \|f\|_{H^6} + \|f_t\|_{A^0}^{1/2} \|f_t\|_{H^4}^{1/2} \|f\|_{A^0}^{3/5} \|f\|_{H^6}^{2/5} \right), \end{split}$$

Using the commutator structure together with the product rule in Wiener spaces, we estimate

$$\begin{aligned} \|F_6\|_{A^{1/2}} &\leq C \|\partial_x^3 f \mathcal{H} f_t + 3\partial_x^2 f \partial_x \mathcal{H} f_t + \partial_x f \partial_x^2 \mathcal{H} f_t \|_{A^{1/2}} \\ &\leq C \bigg( \|f\|_{A^{3+1/2}} \|f_t\|_{A^0} + \|f\|_{A^3} \|f_t\|_{A^{1/2}} \\ &+ \|f\|_{A^{2+1/2}} \|f_t\|_{A^1} + \|f\|_{A^2} \|f_t\|_{A^{1+1/2}} \\ &+ \|f\|_{A^{1+1/2}} \|f_t\|_{A^2} + \|f\|_{A^1} \|f_t\|_{A^{2+1/2}} \bigg). \end{aligned}$$

Using interpolation in Wiener spaces and then the Sobolev embedding

 $\|f\|_{A^{5+2/5}} \leq C \|f\|_{H^6} \qquad \text{and} \qquad \|f_t\|_{A^{3+2/5}} \leq C \|f_t\|_{H^4},$  we compute that

$$\begin{aligned} \|F_6\|_{A^{1/2}} &\leq C \bigg( \|f\|_{A^0}^{19/54} \|f\|_{H^6}^{35/54} \|f_t\|_{A^0} + \|f\|_{A^0}^{4/9} \|f\|_{H^6}^{5/9} \|f_t\|_{A^0}^{29/34} \|f_t\|_{H^4}^{5/34} \\ &+ \|f\|_{A^0}^{29/54} \|f\|_{H^6}^{25/54} \|f_t\|_{A^0}^{12/17} \|f_t\|_{H^4}^{5/17} \\ &+ \|f\|_{A^0}^{17/27} \|f\|_{H^6}^{10/27} \|f_t\|_{A^0}^{19/34} \|f_t\|_{H^4}^{15/34} \\ &+ \|f\|_{A^0}^{13/18} \|f\|_{H^6}^{5/18} \|f_t\|_{A^0}^{7/10} \|f_t\|_{H^4}^{10/17} \\ &+ \|f\|_{A^0}^{22/27} \|f\|_{H^6}^{5/27} \|f_t\|_{A^0}^{9/34} \|f_t\|_{H^4}^{25/34} \bigg). \end{aligned}$$

Let us recall that using Duhamel formulation the solution then can be written as

$$u(x,t) = e^{-tL}u_0 + \int_0^t e^{-(t-t')L} \begin{pmatrix} 0 \\ F(t') \end{pmatrix} dt'$$

and satifies,

$$|\hat{u}(n,t)| \le Ce^{-\delta t} (1+\sqrt{|n|}) \left| \hat{f}_j(n) \right| + C \int_0^t e^{-(t-t')\delta n^2} \sqrt{|n|} \left| \hat{F}(n,t') \right| \mathrm{d}t'.$$

Using

$$-(t-t')\,\delta n^2 \le -(t-t')\,\delta \le 0,$$

we can estimate

$$\|u(t)\|_{A^0} \le Ce^{-\delta t} \|(f_0, f_1)\|_{A^{1/2}} + Ce^{-\delta t} \int_0^t e^{\delta t'} \sum_{j=1}^6 \|F_j(t')\|_{A^{1/2}} \mathrm{d}t'.$$

Recalling the previous estimates for  $||F_j(t')||_{A^{1/2}}$  and the definition of the norm  $|||(f, f_t)||_T$ , we have that

$$\|F(t')\|_{A^{1/2}} \le Ce^{-\delta t'(1+73/918)} \|\|(f, f_t)\|_T^2, \qquad 0 \le t' \le t \le T.$$

We conclude that

$$e^{\delta t} \max_{t' \in [0,t]} \left\{ \left\| \left( f\left(t'\right), f_t\left(t'\right) \right) \right\|_{A^0} \right\} \\ \leq C \| (f_0, f_1) \|_{A^{1/2}} + C \| (f, f_t) \|_T^2 \int_0^t e^{-(73/918)\delta t'} dt' \\ \leq C \| (f_0, f_1) \|_{A^{1/2}} + C \| (f, f_t) \|_T^2$$

$$(10)$$

2.3. Boundedness in the high regularity space. Similarly as in [22], we test the equation against  $\Lambda^8 f_t$ , integrate in  $\mathbb{S}^1$  and integrate by parts obtaining the energy balance

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\mathfrak{E}(t) + \mathfrak{D}\left(t'\right) = \sum_{i=1}^{6} I_{i}\left(t\right),\tag{11}$$

with

$$\begin{aligned} \mathfrak{E}(t) &= \left\| f_t\left(t'\right) \right\|_{H^4}^2 + \beta \left\| f\left(t'\right) \right\|_{H^{4+3/2}}^2 + \delta^2 \left\| f\left(t'\right) \right\|_{H^6}^2 + \left\| f\left(t'\right) \right\|_{H^{4+1/2}}^2, \\ \mathfrak{D}(t) &= 2\delta \left\| f_t(t) \right\|_{H^5}^2, \end{aligned}$$

and

$$I_{1}(t) = -\int_{\mathbb{S}^{1}} \Lambda\left((\mathcal{H}f_{t})^{2}\right) \Lambda^{7} f_{t} \, \mathrm{d}x$$
$$I_{2}(t) = \int_{\mathbb{S}^{1}} \partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda f \Lambda^{8} f_{t} \, \mathrm{d}x$$
$$I_{3}(t) = \beta \int_{\mathbb{S}^{1}} \partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda^{3} f \Lambda^{8} f_{t} \, \mathrm{d}x$$
$$I_{4}(t) = \delta \int_{\mathbb{S}^{1}} \partial_{x} \llbracket \mathcal{H}, \mathcal{H}f_{t} \rrbracket \mathcal{H}\partial_{x}^{2} f \Lambda^{8} f_{t} \, \mathrm{d}x$$
$$I_{5}(t) = \delta \int_{\mathbb{S}^{1}} \Lambda \left(\mathcal{H}f_{t}\mathcal{H}\partial_{x}^{2} f\right) \Lambda^{8} f_{t} \, \mathrm{d}x$$
$$I_{6}(t) = -\delta \int_{\mathbb{S}^{1}} \partial_{x} \llbracket \partial_{x} \llbracket \partial_{x}^{2}, f \rrbracket \mathcal{H}f_{t} \Lambda^{8} f_{t} \, \mathrm{d}x.$$

Using the self-adjointness of the operator  $\Lambda$  together with Hölder's inequality and the Sobolev embedding

$$\|g\|_{L^4} \le C \|g\|_{H^{0.25}},$$

we find that

$$\begin{split} I_{1}(t) &= -\int_{\mathbb{S}^{1}} \left( (\mathcal{H}f_{t})^{2} \right) \Lambda^{9} f_{t} \, \mathrm{d}x \\ &= -\int_{\mathbb{S}^{1}} \left( (\mathcal{H}f_{t})^{2} \right) \partial_{x}^{4} \Lambda^{5} f_{t} \, \mathrm{d}x \\ &= -\int_{\mathbb{S}^{1}} \partial_{x}^{4} \left( (\mathcal{H}f_{t})^{2} \right) \Lambda^{5} f_{t} \, \mathrm{d}x \\ &= -\int_{\mathbb{S}^{1}} \left( 2\mathcal{H}f_{t} \Lambda \partial_{x}^{3} f_{t} + 6(\Lambda \partial_{x}f_{t})^{2} + 8\Lambda f_{t} \partial_{x}^{2} \Lambda f_{t} \right) \Lambda^{5} f_{t} \, \mathrm{d}x \\ &\leq C \|f_{t}\|_{H^{5}} \left( \|f_{t}\|_{H^{4}} \|\mathcal{H}f_{t}\|_{L^{\infty}} + \|f_{t}\|_{H^{2.25}}^{2} + \|f_{t}\|_{H^{3}} \|\Lambda f_{t}\|_{L^{\infty}} \right) \\ &\leq C \|f_{t}\|_{H^{5}} \|f_{t}\|_{H^{4}} \|f_{t}\|_{H^{2.25}} \\ &\leq \sigma \|f_{t}\|_{H^{5}}^{2} + C \|f_{t}\|_{H^{4}}^{2} \|f_{t}\|_{H^{2.25}}^{2} , \end{split}$$
(12)

for  $\sigma > 0$  to be fixed below.

Furthermore, using interpolation between Sobolev spaces, the embedding

$$H^s \subset H^r, r \leq s,$$

we obtain the estimate

$$I_1(t) \le \sigma \|f_t\|_{H^5}^2 + C \|(f, f_t)\|_T^4 e^{-(\delta/2)t}.$$
(13)

Equipped with (29), we can estimate  $I_2$  as follows

$$I_{2}(t) = \int_{\mathbb{S}^{1}} \Lambda^{4} \partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda f \Lambda^{4} f_{t} \, \mathrm{d}x$$
  
$$\leq \left\| \partial_{x}^{5} \llbracket \mathcal{H}, f \rrbracket \Lambda f \right\|_{L^{2}} \left\| \Lambda^{4} f_{t} \right\|_{L^{2}}$$
  
$$\leq \left\| \partial_{x}^{5} f \right\|_{L^{\infty}} \left\| \Lambda f \right\|_{L^{2}} \left\| \Lambda^{4} f_{t} \right\|_{L^{2}}$$

As a consequence, by interpolation in Wiener and Sobolev spaces, we have that

$$I_2(t) \le C |||(f, f_t)|||_T^3 e^{-(\delta/2)t}.$$
(14)

Analogously, we find that

$$I_{3}(t) \leq \|\partial_{x}^{4}f\|_{L^{\infty}} \|\Lambda^{3}f\|_{L^{2}} \|\Lambda^{5}f_{t}\|_{L^{2}}$$
$$\leq \|\partial_{x}^{4}f\|_{L^{\infty}}^{2} \|\Lambda^{3}f\|_{L^{2}}^{2} + \sigma \|\Lambda^{5}f_{t}\|_{L^{2}}^{2}$$
$$\leq C \|\|(f,f_{t})\|_{T}^{4}e^{-(\delta/2)t} + \sigma \|\Lambda^{5}f_{t}\|_{L^{2}}^{2}.$$
(15)

We can decompose  $I_4(t)$  as follows

$$I_4(t) = \delta \int_{\mathbb{S}^1} \left[ \Lambda(\mathcal{H}f_t \Lambda \partial_x f) + \partial_x (\mathcal{H}f_t \partial_x^2 f) \right] \Lambda^8 f_t \, \mathrm{d}x$$
$$= \delta \int_{\mathbb{S}^1} \left[ \Lambda(\mathcal{H}f_t \Lambda \partial_x f) - \partial_x (\mathcal{H}f_t \Lambda^2 f) \right] \Lambda^8 f_t \, \mathrm{d}x$$
$$= J_1^4 + J_2^4,$$

with

$$J_1^4 = \delta \int_0^t \int_{\mathbb{S}^1} \Lambda(\mathcal{H}f_t \Lambda \partial_x f) \Lambda^8 f_t \, \mathrm{d}x \, \mathrm{d}t'$$
$$J_2^4 = -\delta \int_0^t \int_{\mathbb{S}^1} \partial_x (\mathcal{H}f_t \Lambda^2 f) \Lambda^8 f_t \, \mathrm{d}x \, \mathrm{d}t'.$$

We will use the fractional Leibniz rule (see [16, 27, 28]):

.

 $\|\Lambda^{s}(uv)\|_{L^{p}} \leq C\left(\|\Lambda^{s}u\|_{L^{p_{1}}}\|v\|_{L^{p_{2}}} + \|\Lambda^{s}v\|_{L^{p_{3}}}\|u\|_{L^{p_{4}}}\right),$ 

which holds whenever

$$\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_3} + \frac{1}{p_4} \qquad \text{where } 1/2$$

and  $s > \max\{0, 1/p - 1\}$ . Using the fractional Leibniz rule and the self-adjointness of the operator  $\Lambda$ , we compute

$$J_{1}^{4}(t) = \delta \int_{\mathbb{S}^{1}} \Lambda^{4} (\mathcal{H}f_{t}\Lambda\partial_{x}f)\Lambda^{5}f_{t} \, \mathrm{d}x$$
  

$$\leq \delta \|\Lambda^{4} (\mathcal{H}f_{t}\Lambda\partial_{x}f)\|_{L^{2}} \|\Lambda^{5}f_{t}\|_{L^{2}}$$
  

$$\leq \delta C (\|f_{t}\|_{H^{1}}\|f\|_{H^{6}} + \|f_{t}\|_{H^{4}}\|f\|_{H^{3}})\|f_{t}\|_{H^{5}}$$
  

$$\leq \delta C (\|f_{t}\|_{H^{1}}^{2}\|f\|_{H^{6}}^{2} + \|f_{t}\|_{H^{4}}^{2}\|f\|_{H^{3}}^{2}) + \sigma \|f_{t}\|_{H^{5}}^{2}$$
  

$$\leq C \|\|f\|_{T}^{4} e^{-(\delta/2)t} + \sigma \|f_{t}\|_{H^{5}}^{2}.$$

The terms  $J_2^4$  and  $I_5 = J_1^4$  can be estimated in a similar way and we find that

$$I_4(t) + I_5(t) \le C |||f|||_T^4 e^{-(\delta/2)t} + \sigma ||f_t||_{H^5}^2.$$
(16)

Now we are left with  $I_6$ . We remark that

$$I_6(t) = -\delta \int_{\mathbb{S}^1} \partial_x \left[ \partial_x^2 f \mathcal{H} f_t + 2\partial_x f \Lambda f_t \right] \partial_x^4 \Lambda^4 f_t \, \mathrm{d}x.$$

Integrating by parts, we find that

$$I_6(t) = \delta \int_{\mathbb{S}^1} \partial_x^4 \left[ \partial_x^2 f \mathcal{H} f_t + 2 \partial_x f \Lambda f_t \right] \partial_x \Lambda^4 f_t \, \mathrm{d}x.$$

Hence, using the same ideas as before, we have that

$$I_{6}(t) \leq |||(f, f_{t})|||_{T}^{4} e^{-(\delta/2)t} + \sigma ||f_{t}||_{H^{5}}^{2} + 2 \int_{\mathbb{S}^{1}} \partial_{x} f \Lambda^{5} f_{t} \partial_{x} \Lambda^{4} f_{t} \, \mathrm{d}x.$$

The term

$$J_1^6 = 2 \int_{\mathbb{S}^1} \partial_x f \Lambda^5 f_t \partial_x \Lambda^4 f_t \, \mathrm{d}x$$

is the highest order term. However, it has an inner commutator structure that we can exploit as follows:

$$J_1^6(t) = \int_{\mathbb{S}^1} \mathcal{H}(\partial_x f \Lambda^5 f_t) \Lambda^5 f_t \, \mathrm{d}x - \int_{\mathbb{S}^1} \partial_x f \Lambda^5 f_t \mathcal{H} \Lambda^5 f_t \, \mathrm{d}x$$
$$= \int_{\mathbb{S}^1} [\![\mathcal{H}, \partial_x f]\!] \Lambda^5 f_t \Lambda^5 f_t \, \mathrm{d}x.$$

Then, recalling (29), we conclude that

$$I_6(t) \le |||(f, f_t)|||_T^4 e^{-(\delta/2)t} + \sigma ||f_t||_{H^5}^2.$$
(17)

2.4. Finishing the proof of Theorem 1. Collecting (13), (14), (15), (16) and (17) and taking  $\sigma$  small enough, we conclude

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathfrak{E}(t) + \mathfrak{D}(t') \le C |||(f, f_t)|||_T^2 e^{-(\delta/2)t} + C |||(f, f_t)|||_T^4 e^{-(\delta/2)t}.$$
(18)

Integrating in time and using (10), we conclude the polynomial bound

$$\|\|(f, f_t)\|\|_T + \int_0^T \mathfrak{D}(t') dt' \le C \|(f_0, f_1)\|_{A^{1/2}} + \mathfrak{E}(0) + C(\|\|(f, f_t)\|\|_T^2 + \|\|(f, f_t)\|\|_T^4),$$

thus, there exists a (fixed, positive) constant  $1 < C^+$  such that

$$\|\|(f,f_t)\|\|_T \le C^+ \left[ (\|f_0\|_{H^6} + \|f_1\|_{H^4}) + \left( \|\|(f,f_t)\|_T^2 + \|\|(f,f_t)\|_T^4 \right) \right].$$
(19)

We observe that, in the previous estimates, we have not used any hypothesis on the size of the initial data and the previous bound is valid for every solution and  $T \in (0, T_{\text{max}})$ .

We want to prove that, there exists a  $c_0 > 0$  such that for any solution of (3) stemming from an initial data

$$||f_0||_{H^6} + ||f_1||_{H^4} \le c_0, \tag{20}$$

the inequality

$$|||(f, f_t)|||_T \le C \left( ||f_0||_{H^6} + ||f_1||_{H^4} \right),$$

holds true for any T > 0 and thus the solution is global by a standard continuation argument.

Let us assume that the solution does not stay bounded for all times, the contrary being true would imply that the solution is global by a continuation argument. If the initial data is small enough, we can find T such that

$$|||(f, f_t)|||_T = \frac{3}{4} < 1$$

The above inequality allow us to deduce the polynomial bound

$$|||(f, f_t)|||_T \le 2C^+ \left[ (||f_0||_{H^6} + ||f_1||_{H^4}) + |||(f, f_t)|||_T^2 \right],$$
(21)

or equivalently,

$$2C^{+} |||(f, f_{t})|||_{T} \leq (2C^{+})^{2} (||f_{0}||_{H^{6}} + ||f_{1}||_{H^{4}}) + [2C^{+}|||(f, f_{t})|||_{T}]^{2}.$$
(22)

Then, without loss of generality we can restrict our analysis to a polynomial of the form

$$|||(f, f_t)|||_T \le C_0 (f_0, f_1) + |||(f, f_t)|||_T^2.$$
(23)

Now we observe that if

$$C_0 \ll 1$$

is small enough, the polynomial

$$\mathcal{Q}(y) = \mathcal{C}_0 - y + y^2$$

has two positive real roots

$$y_{\pm} = \frac{1 \pm \sqrt{1 - 4\mathcal{C}_0}}{2},$$

moreover if  $0 < C_0 \ll 1$ 

$$y_{-} = \min\{y_{+}, y_{-}\} = \frac{1 - \sqrt{1 - 4C_0}}{2} \le 2C_0.$$

Furthermore, analogously as in in [22], we know that the application  $t \mapsto |||(f, f_t)|||_t$  is continuous for  $t \in [0, T_{\max})$ . This, together with the smallness in the initial data, implies that

$$||(f, f_t)||_T \in [0, y_-].$$

We combine the above deduction with the estimate  $y_{-} \leq 2C_0$  and we deduce that

$$|||(f, f_t)|||_T \le 2\mathcal{C}_0 < \frac{3}{4},$$

if we take  $C_0$  small enough. This is a contradiction with the definition of T and implies that the solution is global.

## 3. Derivation of (5)

Our starting point in this section is (2):

$$f_{tt} = -2\delta\Lambda^{2}f_{t} - \Lambda f - \beta\Lambda^{3}f - \delta^{2}\Lambda^{4}f + \varepsilon \left\{ -\Lambda \left( \left(\mathcal{H}f_{t}\right)^{2} \right) + \partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda f + \beta\partial_{x} \llbracket \mathcal{H}, f \rrbracket \Lambda^{3}f + \delta\partial_{x} \llbracket \mathcal{H}, \mathcal{H}f_{t} \rrbracket \mathcal{H}\partial_{x}^{2}f + \delta\Lambda \left( \mathcal{H}f_{t} \mathcal{H}\partial_{x}^{2}f \right) + \delta^{2}\partial_{x} \llbracket \partial_{x}^{2}, f \rrbracket \Lambda \partial_{x}f - \delta\partial_{x} \llbracket \partial_{x}^{2}, f \rrbracket \mathcal{H}f_{t} - \delta^{2}\partial_{x} \llbracket \mathcal{H}, \partial_{x}^{2}f \rrbracket \partial_{x}^{2}f \right\}.$$
(24)

Let us introduce the 'far-field' variables,

$$\chi = x - t, \quad \tau = \varepsilon t.$$

Then, we have that

$$\frac{\partial}{\partial t}f(\chi(x,t),\tau(t)) = -f_{\chi} + \varepsilon f_{\tau},$$

and

$$\frac{\partial^2}{\partial t^2} f(\chi(x,t),\tau(t)) = f_{\chi\chi} - 2\varepsilon f_{\tau\chi} + \varepsilon^2 f_{\tau\tau}.$$

After neglecting terms of  $O(\varepsilon^2)$ , (24) reads

$$\begin{split} (f_{\chi} - 2\varepsilon f_{\tau})_{\chi} &= -2\delta\Lambda^{2}(-f_{\chi} + \varepsilon f_{\tau}) - \Lambda f - \beta\Lambda^{3}f - \delta^{2}\Lambda^{4}f \\ &+ \varepsilon \bigg\{ -\Lambda \left( \left(\mathcal{H}f_{\chi}\right)^{2} \right) + \partial_{\chi} \llbracket \mathcal{H}, f \rrbracket \Lambda f + \beta\partial_{\chi} \llbracket \mathcal{H}, f \rrbracket \Lambda^{3}f \\ &- \delta\partial_{\chi} \llbracket \mathcal{H}, \mathcal{H}f_{\chi} \rrbracket \mathcal{H}\partial_{\chi}^{2}f - \delta\Lambda \left(\mathcal{H}f_{\chi}\mathcal{H}\partial_{\chi}^{2}f\right) + \delta^{2}\partial_{\chi} \llbracket \partial_{x}^{2}, f \rrbracket \Lambda \partial_{\chi}f \\ &+ \delta\partial_{\chi} \llbracket \partial_{x}^{2}, f \rrbracket \mathcal{H}f_{\chi} - \delta^{2}\partial_{\chi} \llbracket \mathcal{H}, \partial_{x}^{2}f \Biggr] \partial_{\chi}^{2}f \bigg\}. \end{split}$$

Integrating in  $\chi$  and using our previous notation for the space and time variables we find the equation

$$\begin{aligned} f_x - 2\varepsilon f_t &= 2\delta\partial_x (-f_x + \varepsilon f_\tau) - \mathcal{H}f + \beta \mathcal{H}\partial_x^2 f - \delta^2 \partial_x^3 f \\ &+ \varepsilon \bigg\{ - \mathcal{H}\left( (\mathcal{H}f_x)^2 \right) + \llbracket \mathcal{H}, f \rrbracket \Lambda f + \beta \llbracket \mathcal{H}, f \rrbracket \Lambda^3 f \\ &- \delta \llbracket \mathcal{H}, \Lambda f \rrbracket \mathcal{H}\partial_x^2 f - \delta \mathcal{H} \left( \Lambda f \mathcal{H}\partial_x^2 f \right) + \delta^2 \llbracket \partial_x^2, f \rrbracket \Lambda \partial_x f \\ &+ \delta \llbracket \partial_x^2, f \rrbracket \mathcal{H}f_x - \delta^2 \llbracket \mathcal{H}, \partial_x^2 f \rrbracket \partial_x^2 f \bigg\}. \end{aligned}$$

Regrouping terms we can equivalently write

$$(1+\delta\partial_x)2\varepsilon f_t = f_x + 2\delta f_{xx} + \mathcal{H}f - \beta\mathcal{H}\partial_x^2 f + \delta^2\partial_x^3 f$$
$$-\varepsilon \bigg\{ -\mathcal{H}\left((\mathcal{H}f_x)^2\right) + \llbracket\mathcal{H}, f\rrbracket\Lambda f + \beta\llbracket\mathcal{H}, f\rrbracket\Lambda^3 f$$
$$-\delta\llbracket\mathcal{H}, \Lambda f\rrbracket\mathcal{H}\partial_x^2 f - \delta\mathcal{H}\left(\Lambda f\mathcal{H}\partial_x^2 f\right) + \delta^2\llbracket\partial_x^2, f\rrbracket\Lambda\partial_x f$$
$$+\delta\llbracket\partial_x^2, f\rrbracket\mathcal{H}f_x - \delta^2\llbracket\mathcal{H}, \partial_x^2 f\rrbracket\partial_x^2 f\bigg\}.$$
(25)

We observe that, taking the operator

$$\mathcal{N} = (1 - \delta^2 \partial_x^2)^{-1} (1 - \delta \partial_x),$$

we find that

$$2\varepsilon f_{t} = \mathcal{N}f_{x} + 2\delta\mathcal{N}f_{xx} + \mathcal{N}\mathcal{H}f - \beta\mathcal{N}\mathcal{H}\partial_{x}^{2}f + \delta^{2}\mathcal{N}\partial_{x}^{3}f - \varepsilon\mathcal{N}\bigg\{-\mathcal{H}\left((\mathcal{H}f_{x})^{2}\right) + \llbracket\mathcal{H}, f\rrbracket\Lambda f + \beta\llbracket\mathcal{H}, f\rrbracket\Lambda^{3}f - \delta\llbracket\mathcal{H}, \Lambda f\rrbracket\mathcal{H}\partial_{x}^{2}f - \delta\mathcal{H}\left(\Lambda f\mathcal{H}\partial_{x}^{2}f\right) + \delta^{2}\llbracket\partial_{x}^{2}, f\rrbracket\Lambda\partial_{x}f + \delta\llbracket\partial_{x}^{2}, f\rrbracket\mathcal{H}f_{x} - \delta^{2}\llbracket\mathcal{H}, \partial_{x}^{2}f\rrbracket\partial_{x}^{2}f\bigg\}.$$
 (26)

As in [17], we now define

$$u = \Lambda f.$$

This new unknown solves the following equation

$$\begin{split} 2\varepsilon u_t &= \mathcal{N}u_x + 2\delta \mathcal{N}u_{xx} + \mathcal{N}\mathcal{H}u - \beta \mathcal{N}\mathcal{H}\partial_x^2 u + \delta^2 \mathcal{N}\partial_x^3 u \\ &- \varepsilon \Lambda \mathcal{N} \bigg\{ -\mathcal{H} \left( u^2 \right) + \big[\!\big[\mathcal{H}, \Lambda^{-1}u\big]\!\big] u + \beta \big[\!\big[\mathcal{H}, \Lambda^{-1}u\big]\!\big] \Lambda^2 u \\ &+ \delta \big[\!\big[\mathcal{H}, u\big]\!\big] \mathcal{H}\Lambda u + \delta \mathcal{H} \left( u \mathcal{H}\Lambda u \right) + \delta^2 \big[\!\big[\partial_x^2, \Lambda^{-1}u\big]\!\big] u_x \\ &+ \delta \big[\!\big[\partial_x^2, \Lambda^{-1}u\big]\!\big] u - \delta^2 \big[\!\big[\mathcal{H}, \Lambda u\big]\!\big] \Lambda u \bigg\}. \end{split}$$

The previous equation can be written equivalently as

$$2\varepsilon u_{t} = \mathcal{N}u_{x} + 2\delta \mathcal{N}u_{xx} + \mathcal{N}\mathcal{H}u - \beta \mathcal{N}\mathcal{H}\partial_{x}^{2}u + \delta^{2}\mathcal{N}\partial_{x}^{3}u - \varepsilon \mathcal{N}\bigg\{2uu_{x} + \Lambda [\![\mathcal{H}, \Lambda^{-1}u]\!]u + \beta \Lambda [\![\mathcal{H}, \Lambda^{-1}u]\!]\Lambda^{2}u - \delta \Lambda [\![\mathcal{H}, u]\!]u_{x} + \delta \partial_{x} (uu_{x}) + \delta^{2}\Lambda [\![\partial_{x}^{2}, \Lambda^{-1}u]\!]u_{x} + \delta \Lambda [\![\partial_{x}^{2}, \Lambda^{-1}u]\!]u - \delta^{2}\Lambda [\![\mathcal{H}, \Lambda u]\!]\Lambda u\bigg\}.$$
(27)

Neglecting now the nonlinear terms that are  $O(\varepsilon \delta^2)$  we conclude (5).

## 4. Proof of Theorem 2

4.1. Local well-posedness. First, we observe that (5) can be written as

$$2\varepsilon u_{t} = \mathcal{N}u_{x} + 2\delta \mathcal{N}u_{xx} + \mathcal{N}\mathcal{H}u - \beta \mathcal{P}\mathcal{H}\partial_{x}^{2}u + \beta \delta \mathcal{P}\Lambda\partial_{x}^{2}u + \delta^{2}\mathcal{P}\partial_{x}^{3}u - \delta^{3}\mathcal{P}\partial_{x}^{4}u - \varepsilon \mathcal{N}\bigg\{2uu_{x} + \Lambda \big[\![\mathcal{H}, \Lambda^{-1}u\big]\!]u + \beta \Lambda \big[\![\mathcal{H}, \Lambda^{-1}u\big]\!]\Lambda^{2}u - \delta \Lambda \big[\![\mathcal{H}, u\big]\!]u_{x} + \delta \partial_{x} (uu_{x}) + \delta \Lambda \big[\![\partial_{x}^{2}, \Lambda^{-1}u\big]\!]u\bigg\}, \quad (28)$$

where the operator

$$\mathcal{P} = (1 - \delta^2 \partial_x^2)^{-1}$$

is defined in Fourier variables as

$$\widehat{\mathcal{P}} = \frac{1}{1 + \delta^2 |k|^2}.$$

Then, using

$$\mathcal{P} = Id + \mathcal{P}\delta^2 \partial_x^2$$

we can observe that the terms

\_

$$\beta \delta \mathcal{P} \Lambda \partial_x^2 u = -\frac{\beta}{\delta} \Lambda u + \frac{\beta}{\delta} \mathcal{P} \Lambda u$$

and

$$-\delta^3 \mathcal{P} \partial_x^4 u = \delta \partial_x^2 u - \delta \mathcal{P} \partial_x^2 u$$

are of parabolic type.

To simplify the notation, in the course of this proof we take  $\varepsilon = 1$ . Now we obtain the *a priori* estimates in the  $H^2$  Sobolev space. These estimates implies the local existence of solution after a standard regularization approach using the periodic heat kernel as mollifier.

We start noticing that

$$\int_{\mathbb{S}^1} u(x,t) \mathrm{d}x = \int_{\mathbb{S}^1} u(x,0) \mathrm{d}x = 0$$

Now we test (28) against  $\Lambda^4 u$ . Then we obtain that

$$\frac{\mathrm{d}}{\mathrm{d}t} \|u\|_{H^2}^2 = L + NL_1 + NL_2 + NL_3 + NL_4 + NL_5 + NL_6,$$

where

$$\begin{split} L &= \int_{\mathbb{S}^1} \{ \mathcal{N}u_x + 2\delta \mathcal{N}u_{xx} + \mathcal{N}\mathcal{H}u - \beta \mathcal{P}\mathcal{H}\partial_x^2 u \\ &+ \beta \delta \mathcal{P}\Lambda \partial_x^2 u + \delta^2 \mathcal{P}\partial_x^3 u - \delta^3 \mathcal{P}\partial_x^4 u \} \Lambda^4 u dx, \\ NL_1 &= -2 \int_{\mathbb{S}^1} \mathcal{N}(uu_x) \Lambda^4 u dx, \\ NL_2 &= - \int_{\mathbb{S}^1} \mathcal{N}\Lambda \llbracket \mathcal{H}, \Lambda^{-1} u \rrbracket u \Lambda^4 u dx, \\ NL_3 &= -\beta \int_{\mathbb{S}^1} \mathcal{N}\Lambda \llbracket \mathcal{H}, \Lambda^{-1} u \rrbracket \Lambda^2 u \Lambda^4 u dx, \\ NL_4 &= \delta \int_{\mathbb{S}^1} \mathcal{N}\Lambda \llbracket \mathcal{H}, u \rrbracket \partial_x u \Lambda^4 u dx, \\ NL_5 &= -\delta \int_{\mathbb{S}^1} \mathcal{N}\partial_x (uu_x) \Lambda^4 u dx, \\ NL_6 &= -\delta \int_{\mathbb{S}^1} \mathcal{N}\Lambda \llbracket \partial_x^2, \Lambda^{-1} u \rrbracket u \Lambda^4 u dx. \end{split}$$

After a number of integrations by parts, we find that

$$L = \int_{\mathbb{S}^1} \left( \delta \mathcal{P} u_{xx} - \delta \mathcal{P} \Lambda u + \beta \delta \mathcal{P} \Lambda u_{xx} - \delta^3 \mathcal{P} \partial_x^4 u \right) \Lambda^4 u dx$$
  
=  $-\delta \| \mathcal{P}^{1/2} u_{xxx} \|_{L^2}^2 - \delta \| \mathcal{P}^{1/2} \Lambda^{5/2} u \|_{L^2}^2 - \delta \beta \| \mathcal{P}^{1/2} \Lambda^{1/2} u_{xxx} \|_{L^2}^2 - \delta^3 \| \mathcal{P}^{1/2} \Lambda^4 u \|_{L^2}^2$ 

Furthermore, using the parabolic character of some of the terms in L, we find that

$$L \le -\frac{1}{\delta} \|u_{xx}\|_{L^2}^2 - \delta \|\Lambda^3 u\|_{L^2}^2 + C \|u\|_{H^2}^2.$$

For the first nonlinear term  $NL_1$ , we integrate by parts and use that  $\mathcal{N}$  can absorb one derivative to find the estimate

$$NL_{1} \leq \|u\|_{H^{3}} \|u^{2}\|_{H^{1}} \leq C \|u\|_{H^{3}} \|u\|_{H^{1}}^{2} \leq \sigma \|u\|_{H^{3}}^{2} + C \|u\|_{H^{1}}^{4},$$

for  $\sigma > 0$  that will be fixed later.

We recall the following commutator estimate (see equation (1.13) in [9])

$$\left\|\partial_x^{\ell} \llbracket \mathcal{H}, U \rrbracket \partial_x^m V\right\|_{L^p} \le C \left\|\partial_x^{\ell+m} U\right\|_{L^{\infty}} \|V\|_{L^p}, \quad p \in (1,\infty), \quad \ell, m \in \mathbb{N}.$$
(29)

Using (29) and the Sobolev embedding

$$\|\partial_x \Lambda^{-1} u\|_{L^{\infty}} \le C \|\Lambda^{-1} u\|_{H^2} \le C \|u\|_{H^1},$$

we find that

$$NL_2 \le C \|u\|_{H^3} \|u\|_{H^1}^2 \le \sigma \|u\|_{H^3}^2 + C \|u\|_{H^1}^4.$$

Similarly,

$$NL_3 \le C \|u\|_{H^3} \|u\|_{H^2} \|u\|_{H^1} \le \sigma \|u\|_{H^3}^2 + C \|u\|_{H^2}^4,$$

$$NL_4 \le C \|u\|_{H^3} \|u\|_{H^2} \|u\|_{H^1} \le \sigma \|u\|_{H^3}^2 + C \|u\|_{H^2}^4.$$

Integrating by parts in  $NL_5$  and using the regularizing effect from  $\mathcal{N}$ , we can obtain that

$$NL_5 \le C \|u\|_{H^3} \left( \|\partial_x u\|_{L^4}^2 + \|u\|_{L^\infty} \|u\|_{H^2} \right).$$

Using the Sobolev embeddings

$$\|g\|_{L^4} \le C \|g\|_{H^{0.25}},$$

and

$$||g||_{L^{\infty}} \leq C ||g||_{H^1}$$

we find that

$$NL_5 \le \sigma \|u\|_{H^3}^2 + C \|u\|_{H^2}^4.$$

Integrating by parts and using the previous ideas we can estimate the last nonlinear contribution as

$$NL_6 \le C \|u\|_{H^3} \|u\|_{H^2}^2 \le \sigma \|u\|_{H^3}^2 + C \|u\|_{H^2}^4.$$

Taking now  $0<\sigma\ll$  small enough we can ensure that

$$\frac{\mathrm{d}}{\mathrm{d}t} \|u\|_{H^2}^2 + \frac{\delta}{2} \|\Lambda^3 u\|_{L^2}^2 \le C \|u\|_{H^2}^2 + C \|u\|_{H^2}^4, \tag{30}$$

which ensures the existence of a uniform time  $T^*$  such that

$$u \in C([0, T^*), H^2) \cap L^2(0, T^*; H^3).$$

The local existence of solution will follow now from a standard application of Picard's theorem to a sequence of approximate problems. At this level of regularity, the uniqueness of such local strong solution can be easily obtained from a standard contradiction argument that we skip for the sake of brevity.

The rest of this section is devoted to the global existence of solution for small initial data. In order to do that, we define the modified energy

$$|||u|||_{T} = e^{\delta/2T} \max_{t' \in [0,T]} \{ ||u(t')||_{A^{0}} \} + ||u(t')||_{H^{2}}.$$

Then, our goal is to conclude the polynomial inequality

$$|||u|||_T \le C_0 (f_0) + P(|||u|||_T).$$

4.2. The linear semigroup. We consider the linear nonhomogeneous problem

 $2f_t - \mathcal{N}(f_x + 2\delta\partial_x^2 f + \mathcal{H}f) + \beta \mathcal{P}\mathcal{H}f_{xx} - \beta\delta \mathcal{P}\Lambda f_{xx} - \delta^2 \mathcal{P}\partial_x^3 f + \delta^3 \mathcal{P}\partial_x^4 f = F, \quad (31)$ where F is the forcing. This linear equation can then be written as

 $2u_t + \mathscr{L}u = F,$  $u = \Lambda f$ ,

with

$$\mathscr{L}u = \lambda(k)\hat{u}(k)$$

and

$$\begin{split} \lambda(k) &= -\frac{1 - \delta i k}{1 + \delta^2 k^2} (ik - 2\delta |k|^2 - i \text{sgn}(k)) \\ &+ \frac{\beta i k |k|}{1 + \delta^2 k^2} + \frac{\beta \delta |k|^3}{1 + \delta^2 k^2} + \frac{i \delta^2 k^3}{1 + \delta^2 k^2} + \frac{\delta^3 |k|^4}{1 + \delta^2 k^2} \end{split}$$

Then, we have that the homogeneous problem satisfies

$$\hat{u}(k,t) = \hat{u}_0(k)e^{-\lambda(k)t}, \\ \left| \hat{u}_0(k)e^{-\lambda(k)t} \right| \le e^{-\delta t} \left| \hat{u}_0(k) \right|,$$

which in turn implies that

$$e^{\delta T} \max_{t' \in [0,T]} \left\| u(t') \right\|_{A^0} \le \| u_0 \|_{A^0} \,. \tag{32}$$

Equivalently, we have that

$$\|e^{-t\mathscr{L}}\|_{A^0 \mapsto A^0} \le e^{-\delta t}.$$
(33)

4.3. Decay in the low regularity space. Using Duhamel's principle, we can write the mild formulation of our problem as

$$\hat{u}(k,t) = e^{-\lambda(k)t/2}\hat{u}_0(k) + e^{-\lambda(k)t/2}\int_0^t e^{\lambda(k)s/2}\hat{F}(k,s)ds$$

with

$$F = -\frac{\mathcal{N}}{2} \left\{ 2uu_x + \Lambda \llbracket \mathcal{H}, \Lambda^{-1}u \rrbracket u + \beta \Lambda \llbracket \mathcal{H}, \Lambda^{-1}u \rrbracket \Lambda^2 u - \delta \Lambda \llbracket \mathcal{H}, u \rrbracket u_x + \delta \partial_x (uu_x) + \delta \Lambda \llbracket \partial_x^2, \Lambda^{-1}u \rrbracket u \right\}.$$

We observe that

$$(\widehat{\llbracket \mathcal{H}, a \rrbracket} b)(n) = \sum_{n} -i \left( \operatorname{sgn} k - \operatorname{sgn} (k - n) \right) \hat{a}(n) \hat{b}(k - n), \qquad (34)$$

from where

$$0 \le |k| \le |n|,$$

so that this commutator does not vanish. A consequence of the above monotonicity relation is that

$$|n-k| \le |n|.$$

This implies that the above bilinear form presents a nontrivial commutation which allows to commute any derivative acting on the entire bilinear form as a differential operator acting onto a only. In a similar fashion, we find that

$$\left| \left[ \mathcal{H}, \widehat{\Lambda^{-1}u} \right] \Lambda^2 u(k) \right| = \left| \sum_n -i \frac{|k-n|^2}{|n|} \, \hat{u}(n) \hat{u}(k-n) \left( \operatorname{sgn} k - \operatorname{sgn} \left( k - n \right) \right) \right|$$
$$\leqslant C \sum_n |n| \left| \hat{u}(n) \right| \, \left| \hat{u}(k-n) \right|. \tag{35}$$

Using that  $A^0$  is an algebra, the fact that  $\mathcal{N}$  gains one derivative, (35) and Sobolev embedding, we find the estimate

$$\|F\|_{A^{0}} \leq C\left(\|u\|_{A^{0}}^{2} + \|\Lambda^{-1}u\|_{A^{0}}\|u\|_{A^{0}} + \|u\|_{A^{0}}\|\Lambda u\|_{A^{0}}\right).$$

In particular using

$$||u||_{A^1} \le C ||u||_{A^{1,25}}^{4/5} ||u||_{A^0}^{1/5} \le C ||u||_{H^2}^{4/5} ||u||_{A^0}^{1/5},$$

we find that

$$||F||_{A^0} \le C ||u||_{A^0}^{6/5} \left( ||u||_{A^0}^{4/5} + ||u||_{H^2}^{4/5} \right) \le C e^{-3\delta/5t} |||u|||_T^2.$$

As a consequence, we conclude that

$$e^{(\delta/2)t} \max_{t' \in [0,t]} \left\{ \left\| u\left(t'\right) \right\|_{A^{0}} \right\} \leq C \left\| u\left(t'\right) \right\|_{A^{0}} + C \left\| \left\| u\left(t'\right) \right\| \right\|_{T}^{2} \int_{0}^{t} e^{-(\delta/10)t'} dt' \\ \leq C \left\| u\left(t'\right) \right\|_{A^{0}} + C \left\| \left\| u\left(t'\right) \right\| \right\|_{T}^{2}, \quad (36)$$

this concludes the low-regularity estimates.

4.4. Boundedness in the high regularity space. To achieve the required estimate, we have to perform a finer analysis of the nonlinearity. In particular, we need to remove the term

$$||u||_{H^2}^2$$

from the right hand side of (30). In order to do that, we compute

$$NL_{1} = -2 \int_{\mathbb{S}^{1}} \mathcal{N}(uu_{x})\partial_{x}^{4}udx,$$
  
$$= 2 \int_{\mathbb{S}^{1}} \partial_{x}^{2} \mathcal{N}(uu_{x})\partial_{x}^{3}udx,$$
  
$$= 2 \int_{\mathbb{S}^{1}} \partial_{x}(1 - \delta\partial_{x})\mathcal{P}^{1/2}(uu_{x})\mathcal{P}^{1/2}\partial_{x}^{3}udx,$$
  
$$\leq C \|u\|_{H^{1}} \|u\|_{H^{2}} \|\mathcal{P}^{1/2}\partial_{x}^{3}u\|_{L^{2}}.$$

Similarly, invoking (29), we find that

$$NL_{2} = \int_{\mathbb{S}^{1}} \partial_{x} (1 - \delta \partial_{x}) \mathcal{P}^{1/2} \Lambda \llbracket \mathcal{H}, f \rrbracket u \mathcal{P}^{1/2} \partial_{x}^{3} u dx,$$
  

$$\leq C \lVert u \rVert_{L^{2}} \lVert u \rVert_{H^{1.75}} \lVert \mathcal{P}^{1/2} \partial_{x}^{3} u \rVert_{L^{2}},$$
  

$$NL_{3} = \beta \int_{\mathbb{S}^{1}} \partial_{x} (1 - \delta \partial_{x}) \mathcal{P}^{1/2} \Lambda \llbracket \mathcal{H}, f \rrbracket \Lambda^{2} u \mathcal{P}^{1/2} \partial_{x}^{3} u dx,$$
  

$$\leq C \lVert \Lambda^{2} u \rVert_{L^{2}} \lVert u \rVert_{H^{1.75}} \lVert \mathcal{P}^{1/2} \partial_{x}^{3} u \rVert_{L^{2}}.$$

For the term  $NL_4$  we compute as follows

$$NL_4 = \delta \int_{\mathbb{S}^1} (1 - \delta \partial_x) \mathcal{P}^{1/2} \Lambda \llbracket \mathcal{H}, u \rrbracket \partial_x u \mathcal{P}^{1/2} \Lambda^4 u \mathrm{d}x,$$
$$\leq C \lVert \partial_x u \rVert_{L^2} \lVert u \rVert_{H^{1.75}} \lVert \mathcal{P}^{1/2} \Lambda^4 u \rVert_{L^2}.$$

Similarly, since  $\mathcal{P}^{1/2}\partial_x(uu_x) = m_0(D)\partial_x(u^2)$  with  $m_0$  a Fourier multiplier of order zero and using the classical fact that the space  $H^s \cap L^\infty$ ,  $s \in \mathbb{R}$  is a Banach algebra we can argue that

$$NL_{5} = -\delta \int_{\mathbb{S}^{1}} (1 - \delta \partial_{x}) \mathcal{P}^{1/2} \partial_{x} (uu_{x}) \mathcal{P}^{1/2} \Lambda^{4} u dx,$$
  
$$\leq C(\|u\|_{H^{2}} \|u\|_{A^{0}} + \|\partial_{x}u\|_{L^{4}}^{2}) \|\mathcal{P}^{1/2} \Lambda^{4}u\|_{L^{2}},$$
  
$$\leq C\|u\|_{H^{2}} \|u\|_{A^{0}} \|\mathcal{P}^{1/2} \Lambda^{4}u\|_{L^{2}},$$

and

$$NL_{6} = -\delta \int_{\mathbb{S}^{1}} (1 - \delta \partial_{x}) \mathcal{P}^{1/2} \Lambda \llbracket \partial_{x}^{2}, f \rrbracket u \mathcal{P}^{1/2} \Lambda^{4} u \mathrm{d}x,$$
  
$$\leq C \|u\|_{H^{2}} \|u\|_{A^{0}} \|\mathcal{P}^{1/2} \Lambda^{4} u\|_{L^{2}},$$

where we have used the inequality

$$\|\partial_x u\|_{L^4}^2 \le C \|u\|_{H^2} \|u\|_{A^0}.$$

As a consequence, using Young's inequality, we can find the inequality

$$\frac{\mathrm{d}}{\mathrm{d}t} \|u\|_{H^2}^2 + \frac{\delta}{2} \|\mathcal{P}^{1/2} \Lambda^4 u\|_{L^2}^2 \le C \|\|u(t')\|\|_T^3 \|u\|_{H^2} e^{-\delta/16t}.$$
(37)

4.5. Finishing the proof of Theorem 2. Collecting (36), (37) we conclude the polynomial bound

$$|||u|||_T \le \mathcal{C}_0(u_0) + |||u|||_T^2.$$

From here we can finish the argument as in the proof of Theorem 1 and we obtain that the solution is global.

#### Acknowledgments

The research of S.S. is supported by the European Research Council through the Starting Grant project H2020-EU.1.1.-639227. R.G-B was supported by the project "Mathematical Analysis of Fluids and Applications" with reference PID2019-109348GA-I00/AEI/ 10.13039/501100011033 and acronym "MAFyA" funded by Agencia Estatal de Investigación and the Ministerio de Ciencia, Innovacion y Universidades (MICIU). Project supported by a 2021 Leonardo Grant for Researchers and Cultural Creators, BBVA Foundation. The BBVA Foundation accepts no responsability for the opinions, statements and contents included in the project and/or the results thereof, which are entirely the responsability of the authors.

#### References

- Benjamin Akers and Paul A Milewski, Dynamics of three-dimensional gravitycapillary solitary waves in deep water, SIAM Journal on Applied Mathematics 70 (2010), no. 7, 2390–2408.
- [2] David Ambrose, Jerry Bona, and David Nicholls, Well-posedness of a model for water waves with viscosity, (2012).
- [3] CH Aurther, Rafael Granero-Belinchón, Steve Shkoller, and Jon Wilkening, Rigorous asymptotic models of water waves, Water Waves 1 (2019), no. 1, 71–130.
- [4] H. Bae, W. Lee, and J. Shin, Global existence and decay rates of solutions to the viscous water-waves system, Preprint (2020).
- [5] Alfred Barnard Basset, A treatise on hydrodynamics: with numerous examples, vol. 2, Deighton, Bell and Company, 1888.
- [6] J. Thomas Beale, The initial value problem for the Navier-Stokes equations with a free surface, Comm. Pure Appl. Math. 34 (1981), no. 3, 359–392. MR 611750
- [7] J Boussinesq, Lois de l'extinction de la houle en haute mer, CR Acad. Sci. Paris 121 (1895), no. 15-20, 2.
- [8] Peter Constantin, Diego Córdoba, Francisco Gancedo, and Robert M. Strain, On the global existence for the Muskat problem, J. Eur. Math. Soc. (JEMS) 15 (2013), no. 1, 201–227.
- [9] L Dawson, H McGahagan, and G Ponce, On the decay properties of solutions to a class of schrödinger equations, Proceedings of the American Mathematical Society 136 (2008), no. 6, 2081–2090.
- [10] Frederic Dias, Alexander I Dyachenko, and Vladimir E Zakharov, Theory of weakly damped free-surface flows: a new formulation based on potential flow solutions, Physics Letters A 372 (2008), no. 8, 1297–1302.
- [11] Denys Dutykh, Visco-potential free-surface flows and long wave modelling, European Journal of Mechanics-B/Fluids 28 (2009), no. 3, 430–443.
- [12] Denys Dutykh and Frédéric Dias, Dissipative boussinesq equations, Comptes Rendus Mecanique 335 (2007), no. 9-10, 559–583.
- [13] \_\_\_\_\_, Viscous potential free-surface flows in a fluid layer of finite depth, Comptes Rendus Mathematique 345 (2007), no. 2, 113–118.
- [14] Denys Dutykh and Olivier Goubet, Derivation of dissipative boussinesq equations using the dirichlet-to-neumann operator approach, Mathematics and Computers in Simulation 127 (2016), 80–93.
- [15] Francisco Gancedo, Rafael Granero-Belinchón, and Stefano Scrobogna, Surface tension stabilization of the Rayleigh-Taylor instability for a fluid layer in a porous medium, Ann. Inst. H. Poincaré Anal. Non Linéaire **37** (2020), no. 6, 1299–1343.
- [16] Loukas Grafakos and Seungly Oh, The kato-ponce inequality, Communications in Partial Differential Equations 39 (2014), no. 6, 1128–1157.
- [17] Rafael Granero-Belinchón and Alejandro Ortega, On the motion of gravity-capillary waves with odd viscosity, arXiv preprint arXiv:2103.01062 (2021).
- [18] Rafael Granero-Belinchón and Stefano Scrobogna, Asymptotic models for free boundary flow in porous media, Physica D: Nonlinear Phenomena **392** (2019), 1–16.
- [19] \_\_\_\_\_, Models for damped water waves, SIAM Journal on Applied Mathematics 79 (2019), no. 6, 2530–2550.
- [20] \_\_\_\_\_, On an asymptotic model for free boundary Darcy flow in porous media, SIAM
   J. Math. Anal. 52 (2020), no. 5, 4937–4970. MR 4161753
- [21] \_\_\_\_\_, Well-posedness of the water-wave with viscosity problem, to appear in J. Differential Equations, arXiv preprint arXiv:2003.11454 (2020).
- [22] \_\_\_\_\_, Well-posedness of water wave model with viscous effects, Proceedings of the American Mathematical Society 148 (2020), no. 12, 5181–5191.
- [23] Lei Jiang, Chao-Lung Ting, Marc Perlin, and William W Schultz, Moderate and steep faraday waves: instabilities, modulation and temporal asymmetries, Journal of Fluid Mechanics **329** (1996), 275–307.
- [24] Daniel D Joseph and Jing Wang, The dissipation approximation and viscous potential flow, Journal of Fluid Mechanics 505 (2004), 365–377.

- [25] Maria Kakleas and David P Nicholls, Numerical simulation of a weakly nonlinear model for water waves with viscosity, Journal of Scientific Computing 42 (2010), no. 2, 274–290.
- [26] Tsunehiko Kakutani and Kazuo Matsuuchi, Effect of viscosity on long gravity waves, Journal of the physical society of Japan 39 (1975), no. 1, 237–246.
- [27] Tosio Kato and Gustavo Ponce, Commutator estimates and the euler and navierstokes equations, Communications on Pure and Applied Mathematics 41 (1988), no. 7, 891–907.
- [28] Carlos E Kenig, Gustavo Ponce, and Luis Vega, Well-posedness and scattering results for the generalized korteweg-de vries equation via the contraction principle, Communications on Pure and Applied Mathematics 46 (1993), no. 4, 527–620.
- [29] H Lamb, Hydrodynamics, Cambridge Univ Press,, 1932.
- [30] David Lannes, Well-posedness of the water-waves equations, J. Amer. Math. Soc. 18 (2005), no. 3, 605–654. MR 2138139
- [31] \_\_\_\_\_, The water waves problem, Mathematical Surveys and Monographs, vol. 188, American Mathematical Society, Providence, RI, 2013, Mathematical analysis and asymptotics.
- [32] Michael S Longuet-Higgins, Theory of weakly damped stokes waves: a new formulation and its physical interpretation, Journal of Fluid Mechanics 235 (1992), 319–324.
- [33] Y Matsuno, Nonlinear evolutions of surface gravity waves on fluid of finite depth, Physical review letters 69 (1992), no. 4, 609.
- [34] Yoshimasa Matsuno, Nonlinear evolution of surface gravity waves over an uneven bottom, Journal of fluid mechanics 249 (1993), 121–133.
- [35] \_\_\_\_\_, Two-dimensional evolution of surface gravity waves on a fluid of arbitrary depth, Physical Review E 47 (1993), no. 6, 4593.
- [36] Marième Ngom and David P Nicholls, Well-posedness and analyticity of solutions to a water wave problem with viscosity, Journal of Differential Equations 265 (2018), no. 10, 5031–5065.
- [37] KD Ruvinsky and GI Freidman, The fine structure of strong gravity-capillary waves, Nonlinear waves: Structures and Bifurcations, AV Gaponov-Grekhov and MI Rabinovich, eds. Moscow: Nauka (1987), 304–326.
- [38] Stefano Scrobogna, Well-posedness of an asymptotic model for capillarity-driven free boundary Darcy flow in porous media in the critical Sobolev space, Nonlinear Anal. Real World Appl. 60 (2021).
- [39] Jing Wang and Daniel D Joseph, Purely irrotational theories of the effect of the viscosity on the decay of free gravity waves, Journal of Fluid Mechanics 559 (2006), 461–472.
- [40] Guangyu Wu, Yuming Liu, and Dick KP Yue, A note on stabilizing the benjamin-feir instability, Journal of Fluid Mechanics 556 (2006), 45–54.

DEPARTAMENTO DE MATEMÁTICAS, ESTADÍSTICA Y COMPUTACIÓN, UNIVERSIDAD DE CANTABRIA. AVDA. LOS CASTROS S/N, SANTANDER, SPAIN. *Email address*: rafael.granero@unican.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO & IMUS, AV. DE LA REINA MERCEDES, S/N, 41012 SEVILLA, SPAIN

Email address: scrobogna@us.es

22