

---

# ON THE CONDITIONS OF ABSORPTION PROPERTY FOR MORPHOLOGICAL OPENING AND CLOSING

---

A PREPRINT

**Chuan-Shen Hu**

Department of Mathematics  
National Taiwan Normal University  
Taipei, Taiwan  
peterbill126@hotmail.com

**Yu-Min Chung**

Department of Mathematics and Statistics  
University of North Carolina at Greensboro  
Greensboro, USA  
y\_chung2@uncg.edu

November 27, 2024

## ABSTRACT

This paper aims to establish the theoretical foundation for *shift inclusion* in mathematical morphology. In this paper, we prove that the morphological opening and closing concerning structuring elements of shift inclusion property would preserve the ordering of images, while this property is important in granulometric analysis and related image processing tasks. Furthermore, we proposed a systematic way, called the *decomposition theorem for shift inclusion*, to construct sequences of structuring elements with shift inclusion property. Moreover, the influences of the image domain are discussed and the condition named *weak shift inclusion* is defined, which is proved as an equivalent condition for ensuring the order-preserving property.

**Keywords** Shift inclusion · Decomposition theorem · Absorption property · Mathematical morphology

## 1 Introduction

Morphological opening and closing are well-known operations in mathematical morphology. Opening and closing are used for erasing small white and black objects in a binary image, which are applied frequently in many areas, such as image and signal processing [1, 2, 3, 4, 5], script segmentation [6, 7], computer vision [8, 9, 10, 11, 12] and granulometric analysis [13, 14, 15, 16]. In mathematical morphology, a digital image can be viewed as a non-negative real-valued functional defined on a fixed subset of an abelian group. For a structuring element  $B$  in an abelian group, opening  $O_B$  and closing  $C_B$  operations are mappings that map each image  $g$  into new images  $O_B(g)$ ,  $C_B(g)$  respectively. In granulometric analysis, for structuring elements  $B_1 \subseteq B_2$ , the following property is crucial: for every image  $g$ ,

$$O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0) \text{ and } C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0) \quad (1)$$

where  $O_{B_i}(g)^{-1}(0)$ 's and  $C_{B_i}(g)^{-1}(0)$ 's are sets of black pixels of images  $O_{B_i}(g)$ 's and  $C_{B_i}(g)$ 's (cf. Figure 1). For opening operations, the property is a weaker form of the *absorption property* [14, 17, 18]<sup>1</sup> (cf. Corollary 1).

However, Example 1 in Section 3 shows that (1) may not hold for arbitrary  $B_1 \subseteq B_2$ . To tackle this problem, in [14], Soille constructed structuring elements  $B_1 \subseteq B_2 \subseteq \dots \subseteq B_n$  as Minkowski sums of discrete line segments in

---

<sup>1</sup>In [14], the absorption property is defined as  $O_{B_j}(g) \leq O_{B_i}(g)$  whenever  $i \leq j$ .




Figure 1: An example of  $B_1 \subseteq B_2 \subseteq B_3 \subseteq B_4$  with property (1). The subsets of black pixels form a filtration of sets. In the process, small white objects would be erased, while larger holes are still alive in the final step.

2-dimensional lattice (Fig. 11.8 [14]). Based on our investigation, an equivalent criterion for constructing filtered structuring elements with property (1) is still unknown in presented literature.

Structuring elements are soul of opening and closing operations, they describe the neighbourhoods of points in the space. The first consideration of structuring elements were *morphological filters* defined on topological spaces [1, 19, 20, 21, 22]. Recently, such morphological setting were also referred as weighted functions (*e.g.* non-flat structuring elements) and were combined with measure theory and differential geometry for further applications in image analysis [23, 24]. Operations of continuous aspect could be adapted in more general cases because of the support of fruitful foundations in geometry and topology. On the other hand, discrete objects are more suitable for implementing and computing in computers. Contrary to continuous objects, images defined on lattices or  $\mathbb{Z}^m$  may face different combinatorial problems, they were also researched in previous works [2, 14, 25, 26, 27]. In the paper we follow the second approach and consider the combinatorial properties of structuring elements  $B_1 \subseteq B_2$  that satisfy (1).


Contributions of the paper are summarized as the following three-fold:

- We propose the *shift inclusion* property  $B_1 \subseteq_{S,P} B_2$  for structuring elements  $B_1$  and  $B_2$  in an arbitrary abelian group. Theorem 1 is the first contribution, which states that  $B_1 \subseteq_{S,P} B_2$  is a sufficient condition for obtaining (1).
- The second contribution is Theorem 3, it suggests that if  $P = M$  is the based abelian group and  $B_1 \subseteq_{S,M} B_2$ , then  $B_2$  can be represented by  $B_1 \cup (\bigcup_{i=1}^n B_1 + v_i)$  where  $v_1, \dots, v_n$  are some vectors in  $M$ .
- We clarify the relation between property (1) and  $\subseteq_{S,P}$ . Example 9 shows that  $\subseteq_{S,P}$  is not necessary for obtaining property (1). Therefore, we propose the *weak shift inclusion*  $B_1 \subseteq_{WS,P} B_2$ , it is equivalent to property (1) (Theorem 6).

## 1.1 Outline

In Section 2, we introduce the notations and definitions used in the paper, every gray-scale image is referred as a non-negative real-valued functional on a subset  $P$  of an abelian group  $M$ . Hence morphological erosion, dilation, opening, and closing are functions from the space of all images on  $P$  to itself [14, 22, 26]. In Section 3, we introduce the formal definition of shift inclusion and prove that it is a sufficient condition to (1) for images defined on arbitrary  $P$ , and equivalent to (1) for images defined on the whole abelian groups. For images defined on abelian groups, we prove the decomposition theorem and exhibit some feasible examples in Section 3.3. In Section 5, the difference between  $\subseteq_{S,M}$  and  $\subseteq_{S,P}$  are compared and we conclude that there are no implication relations between them. The weak shift inclusion is introduced in Section 5, and we prove that the weak shift inclusion is equivalent to the property (1). The

relations between  $\subseteq_{S,P}$ ,  $\subseteq_{S,M}$ ,  $\subseteq_{WS,P}$  and  $\subseteq_{WS,M}$  can be summarized in the following diagram.



## 2 Background and Notations

We review notations that will be used in this article. Most of our notations and definitions follow the work [26, 14]. We denote integers, natural numbers, and real numbers by the standard notation  $\mathbb{Z}$ ,  $\mathbb{N}$ , and  $\mathbb{R}$ , respectively. We use  $\mathbb{Z}_{\geq 0}$  to denote  $\mathbb{N} \cup \{0\}$ , the set of all non-negative integers. The symbol  $\mathbb{R}_{\geq 0}$  represents the set of all non-negative real numbers. Traditionally, digital images are often defined as real-valued functions on subsets of a fixed free abelian groups [26, 14]. Because the morphological operators concerned in the paper only involving the addition operation of based spaces, instead of considering  $(\mathbb{Z}^m, +)$ , we consider arbitrary abelian groups and their subsets as a more general setting, usually denoted by  $M$  and  $P$  respectively. The group operation and the additional identity element of  $M$  are canonically denoted by  $+$  and  $0$ . When  $M = \mathbb{Z}^m$  for some  $m \in \mathbb{N}$ , elements in  $\mathbb{Z}^m$  are denoted by boldface letters e.g.  $\mathbf{u} \in \mathbb{Z}^m$  to distinguish vectors and scalars. We also use  $\{\mathbf{e}_1, \dots, \mathbf{e}_m\}$  to denote the standard basis for  $\mathbb{R}^m$ , it is also a  $\mathbb{Z}$ -module basis for  $\mathbb{Z}^m$ . When  $m = 2$ , we also use the *dot diagrams* to denote subsets of  $\mathbb{Z}^2$ . For example, if  $P = \{(0, 0), (0, 1), (1, 0), (1, 1)\}$ , then we may represent  $P$  by

$$P = \begin{array}{cc} \bullet & \bullet \\ \circ & \bullet \end{array}$$

where  $\circ$  denotes the origin  $(0, 0)$  in  $\mathbb{Z}^2$ . When we only focus on the shape of  $P$ , we may also replace  $\circ$  by  $\bullet$  directly.

**Definition.** Let  $M$  be an abelian group, a **(digital) image on pixel set**  $P \subseteq M$  is a non-negative function  $g : P \rightarrow \mathbb{R}_{\geq 0}$ . If the range of the function  $g$  is  $\{0, 1\}$ , then  $g$  is called a **binary image**. Otherwise, we refer to  $g$  as a **gray-scale image**. The notation  $\mathcal{I}_P$  denotes the set of all images on  $P$ .

For example, if  $P = \{(2, 2), (2, 3), (3, 2), (3, 3)\}$  is a subset of  $\mathbb{Z}^2$  and  $g$  is an image on  $P$ , then the dot diagrams of  $P$  and  $g$  are represented by

$$P = \begin{array}{cc} \bullet & \bullet \\ \bullet & \bullet \end{array} \quad \text{and} \quad g = \begin{array}{cc} a & b \\ c & d \end{array}$$

where  $a, b, c, d$  are non-negative real numbers. Given two images  $f, g : P \rightarrow \mathbb{R}$ , we establish the following partial order of images by

$$f \leq g \text{ if and only if } f(p) \leq g(p) \text{ for all } p \in P. \quad (3)$$

For functions  $f : S \rightarrow T$ ,  $A \subseteq S$  and  $B \subseteq T$ , the sets  $f(A) := \{f(a) \mid a \in A\}$  and  $f^{-1}(B) := \{s \in S \mid f(s) \in B\}$  are the *image of*  $A$  and the *preimage of*  $B$  under  $f$ . As an abbreviation, if  $B = \{b\}$  is a singleton set, we write  $f^{-1}(b)$  instead of  $f^{-1}(\{b\})$  to denote  $f^{-1}(B)$ .

**Definition.** For an abelian group  $M$ , a **structuring element** is a specified finite set  $B$  satisfying  $0 \in B \subseteq M$ . A structuring element  $B$  is **symmetric** if  $B = -B := \{-x \mid x \in B\}$ .

Here is an example about structuring elements  $B_1 \subseteq B_2 \subseteq \dots \subseteq B_n$  in  $\mathbb{Z}^2$  that are represented dot diagrams

$$B_1 = \circ, \quad B_2 = \begin{array}{cc} \bullet & \bullet \\ \circ & \bullet \end{array}, \quad B_3 = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet \\ \bullet & \bullet & \bullet \end{array}, \quad B_4 = \begin{array}{cccc} \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet \end{array},$$

One may consider structuring elements with infinitely many elements. In [22], Serra defined morphological operations on subsets of a topological space. In the framework, the structuring elements were usually compact subsets. While

the assumption of compact sets is more general than finite sets, additional requirements for opening and closing of gray-scale images  $f : P \rightarrow \mathbb{R}_{\geq 0}$  are necessary.

For example, if  $M = \mathbb{R}^n$  and  $B$  is a compact subset of  $M$ , then any continuous functional  $f : M \rightarrow \mathbb{R}_{\geq 0}$  takes its maximum and minimum on  $B$ . However,  $f$  may not take extrema on  $B$  if  $f$  is not continuous. Erosion, dilation, opening and closing on discrete images strongly depend on the local maximum/minimum of the function  $f$ . Therefore, instead of equipping additional requirements for functional  $f$ , we only concern finite structuring elements in this paper.

The *Minkowski sum* and *difference* for subsets  $A, B \subseteq M$  are defined by

$$\begin{aligned} A + B &:= \{a + b \mid a \in A, b \in B\}, \\ A - B &:= \{a - b \mid a \in A, b \in B\}. \end{aligned}$$

If either  $A = \{a\}$  or  $B = \{b\}$  are sets of singleton points, we would simply use  $a + B$ ,  $a - B$ ,  $A + b$  or  $A - b$  rather than  $\{a\} + B$ ,  $\{a\} - B$ ,  $A + \{b\}$  or  $A - \{b\}$  to denote the Minkowski addition or difference of  $A$  and  $B$ . The following four fundamental morphological operations: *erosion*, *dilation*, *opening*, and *closing* are considered in the work.

**Definition.** For  $g \in \mathcal{I}_P$  and structuring element  $B \subseteq M$ , the **erosion** of  $g$  via  $B$  is an image  $\epsilon_B(g) \in \mathcal{I}_P$  defined by

$$\epsilon_B(g)(x) = \min g \left( (x + B) \cap P \right) = \min \{g(x + b) \mid b \in B, x + b \in P\}. \quad (4)$$

Similarly, the **dilation** of  $g$  via  $B$  is an image  $\delta_B(g) \in \mathcal{I}_P$  defined by

$$\delta_B(g)(x) = \max g \left( (x - B) \cap P \right) = \max \{g(x - b) \mid b \in B, x - b \in P\}. \quad (5)$$

Since  $0 \in B$  and  $B$  is finite,  $(x + B) \cap P$  and  $(x - B) \cap P$  are non-empty, finite sets whenever  $x \in P$ . Therefore,  $\epsilon_B(g)$  and  $\delta_B(g)$  are well-defined.

**Remark.** Observe that if  $B \subseteq M$  is symmetric, then (5) is equivalent to

$$\delta_B(g)(x) = \max g \left( (x + B) \cap P \right) = \max \{g(x + b) \mid b \in B, x + b \in P\}. \quad (6)$$

Since the conditions  $x + b \in P$  and  $x - b \in P$  of elements  $x \in P$ ,  $b \in B$  are crucial for defining morphological operations, for  $P \subseteq M$ ,  $x \in P$ , and structuring element  $B$ , we define

$$B(x; P, +) = \{b \in B : x + b \in P\} \text{ and } B(x; P, -) = \{b \in B : x - b \in P\}. \quad (7)$$

We use dot diagrams to explain  $B(x; P, +)$  and  $B(x; P, -)$ . For example, if

$$P = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet \\ \bullet & \bullet & \bullet \end{array} \text{ and } B = \begin{array}{cc} \bullet & \bullet \\ \circ & \bullet \end{array},$$

then  $B((0, 0); P, +) = B((0, 0); P, -) = B$ . Similarly,

- $B(\mathbf{x}; P, +) = B(-\mathbf{x}; P, -)$  for all  $\mathbf{x} \in P$  (since  $P$  is symmetric);
- $B((1, 1); P, +) = \{(0, 0)\}$  and  $B((1, 1); P, -) = B$ ;
- $B((0, 1); P, +) = \{(0, 0), (1, 0)\}$  and  $B((0, 1); P, -) = B, \dots$  etc.

In particular, if  $P = M$ , then  $B(x; P, +) = B = B(x; P, -)$ .

Therefore, the equations (4) and (5) can also be written as

$$\begin{aligned} \epsilon_B(g)(x) &= \min \{g(x + b) \mid b \in B(x; P, +)\}, \\ \delta_B(g)(x) &= \max \{g(x - b) \mid b \in B(x; P, -)\}. \end{aligned} \quad (8)$$

**Definition** (Section 1.2.1 [26], p. 12). Let  $P, B \subseteq M$  be a pixel set and structuring element respectively. Then **opening** and **closing** operations via  $B$ , denoted by  $O_B$  and  $C_B$ , respectively, are functions  $O_B, C_B : \mathcal{I}_P \rightarrow \mathcal{I}_P$  defined as

$$O_B = \delta_B \circ \epsilon_B \text{ and } C_B = \epsilon_B \circ \delta_B. \quad (9)$$

It is clear that if  $B = \{0\}$ , then  $\epsilon_B = \delta_B = \text{id}_{\mathcal{I}_P}$  where  $\text{id}_{\mathcal{I}_P} : \mathcal{I}_P \rightarrow \mathcal{I}_P$  denotes the identity function and  $O_B = C_B = \text{id}_{\mathcal{I}_P}$ .

In the end of the section, we introduce two thresholding techniques for transforming each gray-scale image to a binary one. These techniques are crucial for obtaining Corollary 1.

**Proposition 1.** For abelian group  $M$  and  $P \subseteq M$ , consider the image  $g \in \mathcal{I}_P$ . For each threshold  $t \in \mathbb{R}_{\geq 0}$  we define  $\tau_t : \mathcal{I}_P \rightarrow \mathcal{I}_P$ ,  $g \mapsto g_t$  by

$$\tau_t(g)(x) = g_t := \begin{cases} 0 & \text{if } g(x) \leq t, \\ 1 & \text{otherwise.} \end{cases} \quad (10)$$

For any structuring element  $B \subseteq M$ , the following diagrams are commutative:

$$\begin{array}{ccc} \mathcal{I}_P & \xrightarrow{\tau_t} & \mathcal{I}_P \\ \epsilon_B \downarrow & & \downarrow \epsilon_B \\ \mathcal{I}_P & \xrightarrow{\tau_t} & \mathcal{I}_P \end{array} \quad \begin{array}{ccc} \mathcal{I}_P & \xrightarrow{\tau_t} & \mathcal{I}_P \\ \delta_B \downarrow & & \downarrow \delta_B \\ \mathcal{I}_P & \xrightarrow{\tau_t} & \mathcal{I}_P \end{array} \quad (11)$$

i.e.,  $\epsilon_B \circ \tau_t = \tau_t \circ \epsilon_B$  and  $\delta_B \circ \tau_t = \tau_t \circ \delta_B$ . Therefore,  $O_B \circ \tau_t = \tau_t \circ O_B$  and  $C_B \circ \tau_t = \tau_t \circ C_B$ .

*Proof.* By definition of erosion, for  $x \in P$  we have

$$\epsilon_B(g)(x) = \min g((x + B) \cap P) \quad \text{and} \quad \epsilon_B(g_t)(x) = \min g_t((x + B) \cap P).$$

If  $b \in B(x; P, +)$  and  $g(x + b) \leq t$ , then  $g_t(x + b) = 0$  and  $\epsilon_B(g)(x) \leq t$ . Hence  $\epsilon_B(g_t)(x) = 0 = \epsilon_B(g)_t(x)$ . On the other hand, if  $g(x + b) > t$  for all  $b \in B(x; P, +)$ , then  $\epsilon_B(g)(x) > t$  and  $g_t(x + b) = 1$  for all  $b \in B(x; P, +)$ . Hence  $\epsilon_B(g)_t(x) = 1 = \epsilon_B(g_t)(x)$ .

Similarly, by definition of dilation, for  $x \in P$  we have

$$\delta_B(g)(x) = \max g((x - B) \cap P) \quad \text{and} \quad \delta_B(g_t)(x) = \max g_t((x - B) \cap P).$$

If  $g(x - b) \leq t$  for all  $b \in B(x; P, -)$ , then  $\delta_B(g_t)(x) = 0$  and  $\delta_B(g)(x) \leq t$ , hence  $\delta_B(g)_t(x) = 0$ . On the other hand, if  $g(x - b) > t$  for some  $b \in B(x; P, -)$ , then  $\delta_B(g)(x) > t$  and  $g_t(x - b) = 1$ , hence  $\delta_B(g)_t(x) = 1 = \delta_B(g_t)(x)$ .  $\square$

**Lemma 1.** For images  $f, g \in \mathcal{I}_P$ ,  $f \leq g$  if and only if  $f_t \leq g_t$  for every  $t \in \mathbb{R}_{\geq 0}$ .

*Proof.* It is clear that  $f \leq g$  implies  $f_t \leq g_t$  for every  $t \in \mathbb{R}_{\geq 0}$ . Conversely, suppose  $0 \leq g(x) < f(x)$  for some  $x \in P$ . Set  $t = g(x)$ , then  $t \in \mathbb{R}_{\geq 0}$  and  $f_t(x) = 1$ ,  $g_t(x) = 0$ .  $\square$

### 3 Shift Inclusion

In this section, we introduce the definition of shift inclusion of two structuring elements and related properties. The main result in the section is Theorem 4, it states shift inclusion of structuring elements is sufficient for ensuring (1). More precisely, consider structuring elements  $B_1 \subseteq B_2$  in an abelian group  $M$ . We hope the following property holds: for every image  $g \in \mathcal{I}_P$ ,

$$O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0) \quad \text{and} \quad C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0).$$

To investigate this, we first observe the following example suggests that (1) may not always hold.

**Example 1.** Consider image  $g : P \rightarrow \{0, 1\}$  with  $P \subseteq \mathbb{Z}^2$  defined by

$$g = \begin{array}{cccccc} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \quad \text{with } P = \begin{array}{cccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

and symmetric structuring elements

$$B_1 = \begin{array}{ccc} & \bullet & \\ \bullet & \circ & \bullet \\ & \bullet & \end{array} \quad \text{and} \quad B_2 = \begin{array}{ccc} & \bullet & \bullet & \bullet \\ \bullet & \bullet & \circ & \bullet \\ & \bullet & \bullet & \bullet \end{array}$$

in  $\mathbb{Z}^2$ . Note that  $B_1 \subseteq B_2$ . By direct calculation,

$$\epsilon_{B_1}(g) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } O_{B_1}(g) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad (12)$$

Similarly,

$$\epsilon_{B_2}(g) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } O_{B_2}(g) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (13)$$

By equations (12) and (13), the set  $O_{B_1}(g)^{-1}(0)$  is not a subset of  $O_{B_2}(g)^{-1}(0)$ .

### 3.1 Definition and Basic Properties

To tackle this problem, we propose an additional property denoted  $B_1 \subseteq_{S,P} B_2$  of structuring elements  $B_1 \subseteq B_2$  in an abelian group  $M$ , called *shift inclusion* with respect to  $P \subseteq M$ . In this paper, we prove that the property (1) holds for arbitrary image  $g \in \mathcal{I}_P$  with  $P \subseteq M$  and  $B_1 \subseteq_{S,P} B_2$ . The definition of  $B_1 \subseteq_{S,P} B_2$  can be separated into two parts: *positive* and *negative* properties, denoted by  $B_1 \subseteq_{S,P,+} B_2$  and  $B_1 \subseteq_{S,P,-} B_2$  respectively.

**Definition.** Given pixel set  $P \subseteq M$  and two structuring elements  $B_1, B_2 \subseteq M$  we say that  $B_1$  is **shift included with respect to  $P$**  in  $B_2$  denoted by  $B_1 \subseteq_{S,P} B_2$  if  $B_1 \subseteq B_2$  and the following properties hold:

- (a) (**Positive property**, denoted by  $B_1 \subseteq_{S,P,+} B_2$ ) For any  $x \in P$  and  $b_2 \in B_2(x; P, +) \subseteq B_2$ , there exists a  $b_1 \in B_1(x; P, +) \subseteq B_1$  such that  $B_1 + (b_2 - b_1) \subseteq B_2$ ;
- (b) (**Negative property**, denoted by  $B_1 \subseteq_{S,P,-} B_2$ ) For any  $x \in P$  and  $b_2 \in B_2(x; P, -) \subseteq B_2$ , there exists a  $b_1 \in B_1(x; P, -) \subseteq B_1$  such that  $B_1 + (b_2 - b_1) \subseteq B_2$ .

A sequence  $B_1, B_2, \dots, B_n$  of structuring elements in  $M$  is called a **shift inclusion sequence** if  $B_i \subseteq_{S,P} B_{i+1}$  for all  $i \in \{1, 2, \dots, n-1\}$ .

The following example shows that the requirement  $B_1 \subseteq B_2$  is not redundant for shift inclusion:

**Example 2.** Consider subset  $P \subseteq \mathbb{Z}^2$  defined by

$$P = \circ \quad \bullet = \{(0, 0), (1, 0)\}$$

and structuring elements

$$B_1 = \begin{smallmatrix} \bullet \\ \circ \end{smallmatrix} = \{(0, 0), (0, 1)\} \text{ and } B_2 = \begin{smallmatrix} \bullet \\ \circ \quad \bullet \end{smallmatrix} = \{(0, 0), (1, 0), (1, 1)\}$$

in  $\mathbb{Z}^2$ . Then  $B_1 \not\subseteq B_2$  while  $B_1$  and  $B_2$  satisfy (a) and (b).

*Proof.* It is sufficient to show that  $B_1$  and  $B_2$  satisfy (a) and (b). Note that

- $B_1((0, 0); P, +) = \{(0, 0)\} = B_1((1, 0); P, +)$ ;
- $B_2((0, 0); P, +) = \{(0, 0), (1, 0)\}$  and  $B_2((1, 0); P, +) = \{(0, 0)\}$ ;
- $B_1((0, 0); P, -) = \{(0, 0)\} = B_1((1, 0); P, -)$ ;
- $B_2((0, 0); P, -) = \{(0, 0)\}$  and  $B_2((1, 0); P, -) = \{(0, 0), (1, 0)\}$ .

Because  $B_1 + (0, 0) \subseteq B_2$  and  $B_1 + (1, 0) \subseteq B_2$ ,  $B_1$  and  $B_2$  satisfy (a) and (b) but  $B_1 \not\subseteq B_2$ . □

We exhibit an example to show that conditions (a) and (b) in Definition 3.1 are not equivalent in general.

**Example 3.** Consider subset  $P \subseteq \mathbb{Z}^2$  defined by

$$P = \begin{array}{c} \bullet \\ \bullet \\ \circ \end{array} = \{(0, 0), (0, 1), (0, 3)\}$$

and structuring elements

$$B_1 = \begin{array}{c} \circ \\ \bullet \end{array} = \{(0, 0), (0, -1)\} \text{ and } B_2 = \begin{array}{c} \bullet \\ \bullet \\ \circ \\ \bullet \end{array} = \{(0, 0), (0, -1), (0, 3), (0, 2)\}$$

in  $\mathbb{Z}^2$ . Then  $B_2 = B_1 \cup (B_1 + 3\mathbf{e}_2)$  and the pair of  $B_1$  and  $B_2$  satisfies (a) but not (b) in Definition 3.1.

*Proof.* We first check that the pair of  $B_1$  and  $B_2$  satisfies (a) by considering all cases:

1. If  $\mathbf{x} = (0, 0)$ , then  $B_1(\mathbf{x}; P, +) = \{(0, 0)\}$  and  $B_2(\mathbf{x}; P, +) = \{(0, 0), (0, 3)\}$ . We have the following possibilities:
  - If  $\mathbf{b}_2 = (0, 0)$ , then  $\mathbf{b}_2 \in B_1$ , take  $\mathbf{b}_1 = \mathbf{b}_2$ ;
  - If  $\mathbf{b}_2 = (0, 3)$ , define  $\mathbf{b}_1 = (0, 0) \in B_1(\mathbf{x}; P, +)$  and  $B_1 + (\mathbf{b}_2 - \mathbf{b}_1) \subseteq B_2$ ;
2.  $\mathbf{x} = (0, 1)$ , then  $B_1(\mathbf{x}; P, +) = \{(0, 0), (0, -1)\}$  and  $B_2(\mathbf{x}; P, +) = \{(0, 0), (0, -1), (0, 2)\}$ . We have the following possibilities:
  - If  $\mathbf{b}_2 = (0, 0)$ , then  $\mathbf{b}_2 \in B_1$ , take  $\mathbf{b}_1 = \mathbf{b}_2$ ;
  - If  $\mathbf{b}_2 = (0, -1)$ , then  $\mathbf{b}_2 \in B_1$ , take  $\mathbf{b}_1 = \mathbf{b}_2$ ;
  - If  $\mathbf{b}_2 = (0, 2)$ , define  $\mathbf{b}_1 = (0, -1) \in B_1$ , then  $\mathbf{x} + \mathbf{b}_1 = (0, 0) \in P$  and  $B_1 + (\mathbf{b}_2 - \mathbf{b}_1) = B_1 + 3\mathbf{e}_2 \subseteq B_2$ .
3. If  $\mathbf{x} = (0, 3)$ , then  $B_1(\mathbf{x}; P, +) = \{(0, 0)\}$  and  $B_2(\mathbf{x}; P, +) = \{(0, 0)\}$ . For  $\mathbf{b}_2 = (0, 0)$ , take  $\mathbf{b}_1 = (0, 0)$ , then  $B_1 + (\mathbf{b}_2 - \mathbf{b}_1) = B_1 + (0, 0) \subseteq B_2$ .

Second, we check that the pair of  $B_1$  and  $B_2$  does not satisfy (b). To prove this, we consider  $\mathbf{x} = (0, 3) \in P$ , then  $\mathbf{b}_2 = (0, 2) \in B_2(\mathbf{x}; P, -) = \{(0, 0), (0, 3), (0, 2)\}$ . Because  $B_1(\mathbf{x}; P, -) = \{(0, 0)\}$ , for  $\mathbf{b}_1 = (0, 0)$ , then  $\mathbf{b}_2 - \mathbf{b}_1 = 2\mathbf{e}_2$ . However,  $B_1 + 2\mathbf{e}_2 = \{(0, 2), (0, 1)\} \not\subseteq B_2$ .

This shows that conditions (a) and (b) are not equivalent.  $\square$

If  $P = M$ , the following proposition shows that positive and negative properties are equivalent.

**Proposition 2.** If  $P = M$  and  $B_1 \subseteq B_2$  are structuring elements, then  $B_1 \subseteq_{S, P, +} B_2$  if and only if  $B_1 \subseteq_{S, P, -} B_2$ .

*Proof.* Because  $P = M$ , we have  $B_1(x; P, +) = B_1 = B_1(x; P, -)$  and  $B_2(x; P, +) = B_2 = B_2(x; P, -)$  whenever  $x \in P$ .

Suppose  $B_1 \subseteq_{S, P, +} B_2$ . Then for any  $x \in P$  and  $b_2 \in B_2(x; P, -) = B_2 = B_2(x; P, +)$ , there exists a  $b_1 \in B_1(x; P, +) = B_1 = B_1(x; P, -)$  such that  $B_1 + (b_2 - b_1) \subseteq B_2$ . This shows that  $B_1 \subseteq_{S, P, -} B_2$ . By symmetric arguments,  $B_1 \subseteq_{S, P, -} B_2$  also implies  $B_1 \subseteq_{S, P, +} B_2$ .  $\square$

Same result follows if  $B_1$  and  $B_2$  are symmetric:

**Proposition 3.** Let  $M$  be an abelian group,  $P \subseteq M$  and  $B_1 \subseteq B_2 \subseteq M$  be structuring elements. If  $B_1, B_2$  are symmetric,  $v \in M$ , and  $B_1 - v \subseteq B_2$ , then  $B_1 + v \subseteq B_2$ . Moreover, (a) and (b) in Definition 3.1 are equivalent.

*Proof.* Suppose  $v \in M$  satisfies  $B_1 - v \subseteq B_2$ . Because  $B_1$  and  $B_2$  are symmetric, we obtain

$$B_1 + v = -B_1 + v = -(B_1 - v) \subseteq -B_2 = B_2.$$

Moreover, if (a) holds, then for  $x \in P$  and  $b_2 \in B_2$  satisfying  $x - b_2 = x + (-b_2) \in P$ , the symmetry of  $B_2$  leads  $-b_2 \in B_2$ . By (a), there is a  $b'_1 \in B_1(x; P, +)$  such that  $B_1 + (-b_2 - b'_1) \subseteq B_2$ . Because  $B_1$  is symmetric,  $b_1 := -b'_1 \in B_1$  and  $x - b_1 = x + b'_1 \in P$ . Because  $B_1 - (b_2 + b'_1) \subseteq B_2$ , preceding discussion shows that  $B_1 + (b_2 - b_1) = B_1 + (b_2 + b'_1) \subseteq B_2$ , this proves (b). By symmetric arguments, (b) implies (a).  $\square$

Similar to the relation  $\subseteq$  on sets, relations  $\subseteq_{S,P,+}$  and  $\subseteq_{S,P,-}$  are a partial orders on the family of all structuring elements in  $M$ . Hence,  $\subseteq_{S,P}$  is also a partial order on the same family. By definition,  $B_1 \subseteq B_2$  is necessary for  $B_1 \subseteq_{S,P,\pm} B_2$ , hence the symmetry and anti-symmetry hold naturally. It remains to show transitivity.

**Proposition 4.** *For pixel set  $P \subseteq M$  and structuring elements  $B_1, B_2, B_3 \subseteq M$ , if  $B_1 \subseteq_{S,P,+} B_2$  (resp.  $B_1 \subseteq_{S,P,-} B_2$ ) and  $B_2 \subseteq_{S,P,+} B_3$  (resp.  $B_2 \subseteq_{S,P,-} B_3$ ), then  $B_1 \subseteq_{S,P,+} B_3$  (resp.  $B_1 \subseteq_{S,P,-} B_3$ ). Therefore,  $\subseteq_{S,P}$  satisfies transitivity.*

*Proof.* Assume  $x \in P$  and  $b_3 \in B_3(x; P, +)$  (resp.  $b_3 \in B_3(x; P, -)$ ). Since  $B_2 \subseteq_{S,P,+} B_3$  (resp.  $B_2 \subseteq_{S,P,-} B_3$ ), there exists  $v = b_3 - b_2 \in b_3 - B_2$  such that  $x + b_2 \in P$  (resp.  $x - b_2 \in P$ ) and  $B_2 + v \subseteq B_3$ . Also, for  $b_2$ , since  $B_1 \subseteq_{S,P,+} B_2$  (resp.  $B_1 \subseteq_{S,P,-} B_2$ ), there exists  $u = b_2 - b_1 \in b_2 - B_1$  such that  $x + b_1 \in P$  (resp.  $x - b_1 \in P$ ) and  $B_1 + u \subseteq B_2$ . Therefore,  $B_1 + (u + v) = (B_1 + u) + v \subseteq B_2 + v \subseteq B_3$  with  $u + v = (b_2 - b_1) + (b_3 - b_2) = b_3 - b_1 \in b_3 - B_1$ , hence  $B_1 \subseteq_{S,P,+} B_3$  (resp.  $B_1 \subseteq_{S,P,-} B_3$ ).  $\square$

### 3.2 Examples

**Lemma 2.** *Let  $P, R_1 \subseteq \mathbb{Z}^m$  be rectangles with  $\mathbf{0} \in R_1$ . Then  $R_1 \subseteq_{S,P} R_1 \cup (R_1 + \mathbf{e}_i)$  and  $R_1 \subseteq_{S,P} R_1 \cup (R_1 - \mathbf{e}_i)$  for every  $i \in \{1, 2, \dots, m\}$ , where  $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_m\} \subseteq \mathbb{Z}^m$  is the standard basis of  $\mathbb{R}^m$ .*

*Proof.* By symmetry, we may assume without loss of generality that  $i = 1$  and prove  $R_1 \subseteq_{S,P} R_1 \cup (R_1 + \mathbf{e}_1)$ . Since  $P, R_1 \subseteq \mathbb{Z}^m$  are rectangles with  $\mathbf{0} \in R_1$ ,

$$P = \mathbb{Z}^m \cap \left( \prod_{i=1}^m [a_i, b_i] \right) \quad \text{and} \quad R_1 = \mathbb{Z}^m \cap \left( \prod_{i=1}^m [r_i, s_i] \right)$$

with integers  $a_i \leq b_i$ , and  $r_i \leq 0 \leq s_i$ . If  $R_2 = R_1 \cup (R_1 + \mathbf{e}_1)$ , then

$$R_2 = \mathbb{Z}^m \cap \left( [r_1, s_1 + 1] \times \prod_{i=2}^m [r_i, s_i] \right) \quad (14)$$

is also a rectangle. Suppose  $\mathbf{x} = (x_1, x_2, \dots, x_m) \in P$  and  $\mathbf{r}_2 = (\lambda_1, \dots, \lambda_m) \in R_2$  satisfy  $\mathbf{x} + \mathbf{r}_2 \in P$ . If  $\mathbf{r}_2 \in R_1$ , then there are nothing need to prove. If  $\mathbf{r}_2 \notin R_1$ , then  $\mathbf{r}_2 = (s_1 + 1, \lambda_2, \dots, \lambda_m)$ . Define  $\mathbf{r}_1 := (s_1, \lambda_2, \dots, \lambda_m) \in R_1$ . Hence  $a_1 \leq x_1 \leq x_1 + s_1 < x_1 + s_1 + 1 = x_1 + \lambda_1 \leq b_1$ . This shows that  $\mathbf{x} + \mathbf{r}_1 \in P$ . Similarly, if  $\mathbf{x} - \mathbf{r}_2 \in P$  and  $\mathbf{r}_2 \notin R_1$ , then  $\mathbf{r}_2 = (s_1 + 1, \lambda_2, \dots, \lambda_m)$  with  $\mathbf{r}_1 := (s_1, \lambda_2, \dots, \lambda_m) \in R_1$ . Hence  $a_1 \leq x_1 - \lambda_1 = x_1 - s_1 - 1 < x_1 - s_1 \leq x_1 \leq b_1$ . This shows that  $\mathbf{x} - \mathbf{r}_1 \in P$ .  $\square$

By applying Lemma 2 inductively, we have the following theorem:

**Theorem 1.** *Let pixel set  $P \subseteq \mathbb{Z}^m$  be a rectangle. If  $\mathbf{0} \in R_1 \subseteq R_2$  where  $R_1, R_2$  are rectangles in  $\mathbb{Z}^m$ , then  $R_1 \subseteq_{S,P} R_2$ .*

*Proof.* We first assume that

$$R_1 = \mathbb{Z}^m \cap \left( \prod_{i=1}^m [r_i, s_i] \right) \quad \text{and} \quad R_2 = \mathbb{Z}^m \cap \left( \prod_{i=1}^m [r_i - c_i, s_i + d_i] \right)$$

with  $r_i \leq 0 \leq s_i$  being integers and  $c_i, d_i \in \mathbb{Z}_{\geq 0}$ . Then we have filtration

$$R_1 = R_1^0 \subseteq R_1^1 \subseteq R_1^2 \cdots \subseteq R_1^m = R_2 \quad (15)$$

where  $R_1^j$  is inductively defined by

$$R_1^j = R_1^{j-1,*} \cup (R_1^{j-1,*} - c_j \cdot \mathbf{e}_j),$$

where

$$R_1^{j-1,*} = R_1^{j-1} \cup (R_1^{j-1} + d_j \cdot \mathbf{e}_j)$$

Because of equation (14), by applying Lemma 2 and Proposition 4 inductively, we deduce that  $R_1^{j-1} \subseteq_{S,P} R_1^j$  for each  $j$ , hence  $R_1 \subseteq_{S,P} R_2$  by Proposition 4.  $\square$



For instance, the following sequence of squares

$$C_0 = \circ, \quad C_1 = \begin{array}{cc} \bullet & \bullet \\ \circ & \bullet \end{array}, \quad C_2 = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet \\ \bullet & \bullet & \bullet \end{array}, \quad C_3 = \begin{array}{cccc} \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet \end{array}, \quad \text{and so on,} \quad (16)$$

satisfies shift inclusion with respect to  $P$ . Squares in (16) can be formulated as:

$$C_0 = \{(0, 0)\},$$

$$C_n = \begin{cases} C_{n-1} \cup (C_{n-1} + \mathbf{e}_1) \cup (C_{n-1} + \mathbf{e}_2) \cup (C_{n-1} + \mathbf{e}_1 + \mathbf{e}_2) & \text{if } n \text{ is odd} \\ C_{n-1} \cup (C_{n-1} - \mathbf{e}_1) \cup (C_{n-1} - \mathbf{e}_2) \cup (C_{n-1} - \mathbf{e}_1 - \mathbf{e}_2) & \text{if } n \text{ is even, } n \geq 2 \end{cases} \quad (17)$$

In addition to rectangular structuring elements, we also provide another family of structuring elements that may be nested by shift inclusion. Recall that for  $\mathbf{x} = (x_1, x_2, \dots, x_m) \in \mathbb{R}^m$ , the  $L^1$ -norm of  $\mathbf{x}$  is defined by

$$\|\mathbf{x}\|_1 = |x_1| + |x_2| + \dots + |x_m|.$$

The same definition can be adapted to  $\mathbb{Z}^m$ , and giving rise to structuring elements of the form

$$Q_\omega = \{\mathbf{x} \in \mathbb{Z}^m : \|\mathbf{x}\|_1 \leq \omega\}$$

for some non-negative integer  $\omega$ . It is clear that  $\mathbf{0} \in Q_\omega$ ,  $Q_\omega \subseteq Q_{\omega+1}$  and  $Q_\omega$  is symmetric.

**Theorem 2.** *Let pixel set  $P \subseteq \mathbb{Z}^m$  be a rectangle. If  $\omega \in \mathbb{Z}_{\geq 0}$  is a non-negative integer, then  $Q_\omega \subseteq_{S,P} Q_{\omega+1}$ .*

*Proof.* Because  $Q_\omega, Q_{\omega+1}$  are symmetric, it is sufficient to show the positive property of shift inclusion. Clearly,  $Q_\omega + \mathbf{e}_i \subseteq Q_{\omega+1}$  and  $Q_\omega - \mathbf{e}_i \subseteq Q_{\omega+1}$  for every  $i \in \{1, 2, \dots, m\}$  by triangle inequality. Assume  $\mathbf{x} = (x_1, \dots, x_m) \in P$  and  $\mathbf{q}_2 = (\lambda_1, \dots, \lambda_m) \in Q_{\omega+1}$  satisfy  $\mathbf{x} + \mathbf{q}_2 \in P$  with

$$P = \mathbb{Z}^m \cap \left( \prod_{i=1}^m [a_i, b_i] \right).$$

Then  $a_i \leq x_i + \lambda_i \leq b_i$  for each  $i \in \{1, 2, \dots, m\}$ . If  $\mathbf{q}_2 \in Q_\omega$ , then there are nothing need to prove. Now we assume  $\mathbf{q}_2 \notin Q_\omega$ , then  $\|\mathbf{q}_2\|_1 = \omega + 1$  and there is a  $j \in \{1, 2, \dots, m\}$  such that  $\lambda_j \neq 0$ . Here we consider two conditions:

- If  $\lambda_j > 0$ , set  $\mathbf{q}_1 = \mathbf{q}_2 - \mathbf{e}_j$ , then

$$\begin{aligned} \|\mathbf{q}_1\|_1 &= |\lambda_1| + \dots + |\lambda_j - 1| + \dots + |\lambda_m| \\ &= |\lambda_1| + \dots + |\lambda_j| - 1 + \dots + |\lambda_m| = \|\mathbf{q}_2\|_1 - 1 = \omega. \end{aligned}$$

This shows that  $\mathbf{q}_1 \in Q_\omega$ . In this case, we also have

$$a_j \leq x_j \leq x_j + \lambda_j - 1 < x_j + \lambda_j \leq b_j.$$

- If  $\lambda_j < 0$ , set  $\mathbf{q}_1 = \mathbf{q}_2 + \mathbf{e}_j$ , then

$$\begin{aligned} \|\mathbf{q}_1\|_1 &= |\lambda_1| + \dots + |\lambda_j + 1| + \dots + |\lambda_m| \\ &= |\lambda_1| + \dots + |\lambda_j| + 1 + \dots + |\lambda_m| = \|\mathbf{q}_2\|_1 + 1 = \omega + 2. \end{aligned}$$

This shows that  $\mathbf{q}_1 \in Q_\omega$ . In this case, we also have

$$a_j \leq x_j + \lambda_j < x_j + \lambda_j + 1 \leq x_j \leq b_j.$$

Both of them show that there is a  $\mathbf{q}_1 \in Q_\omega$  such that  $\mathbf{x} + \mathbf{q}_1 \in P$  and  $Q_\omega + (\mathbf{q}_2 - \mathbf{q}_1) \subseteq Q_{\omega+1}$ .  $\square$

### 3.3 Special case: Shift Inclusion on Whole Space

In Section 3, Theorem 4, we concluded that the shift inclusion is a sufficient condition for obtaining property (1). However, the definition of shift inclusion is quite complicated for verifying. Fortunately, in Theorem 1, we proved that when  $P$  is a rectangle  $\prod_{i=1}^m [a_i, b_i]$  ( $a_i \leq b_i$ ) in  $\mathbb{Z}^m$ , all filtered rectangular structuring elements are shift included, which are widely used in mathematical morphology. Actually, when  $P = M$ , the following decomposition theorem provides a systematical way to construct shift inclusion sequences.

If  $P = M$ , Proposition 2 shows that the positive and negative conditions are equivalent. It can be expressed by the following form:  $B_1 \subseteq_{S,M} B_2$  if and only if for every  $b_2 \in B_2$  there is a  $b_1$  (depends on  $b_2$ ) such that  $B_1 + (b_2 - b_1) \subseteq B_2$ . That is, there is a function  $\phi : B_2 \rightarrow B_1$  (may not unique) such that  $B_1 + (b - \phi(b)) \subseteq B_2$  for every  $b \in B_2$ .

**Theorem 3** (Decomposition Theorem for Shift Inclusion). *If  $M$  is an abelian group and  $B_1 \subseteq B_2 \subseteq M$  are structuring elements, then  $B_1 \subseteq_{S,M} B_2$  if and only if there are finitely many  $v_1, \dots, v_n \in M$  such that*

$$B_2 = B_1 \cup \left( \bigcup_{i=1}^n B_1 + v_i \right). \quad (18)$$

*Proof.* The converse direction is trivial so we prove that  $B_1 \subseteq_{S,M} B_2$  implies the existence of  $v_1, v_2, \dots, v_n \in M$  satisfying (18). If  $B_1 \subseteq_{S,M} B_2$ , then there is a function  $\phi : B_2 \rightarrow B_1$  such that  $B_1 + (b - \phi(b)) \subseteq B_2$  whenever  $b \in B_2$ . Hence

$$B_1 \cup \left( \bigcup_{b \in B_2} B_1 + (b - \phi(b)) \right) \subseteq B_2.$$

Conversely, if  $b_2 \in B_2$ , then  $b_2 = \phi(b_2) + (b_2 - \phi(b_2)) \in B_1 + (b_2 - \phi(b_2))$  and this proves

$$B_2 \subseteq B_1 \cup \left( \bigcup_{b \in B_2} B_1 + (b - \phi(b)) \right).$$

Because  $B_2$  is a finite set, the existence of such  $v_1, v_2, \dots, v_n \in M$  holds.  $\square$

By applying Theorem 3, structuring elements can be constructed inductively. Here we exhibit some examples in  $\mathbb{Z}^2$ .

**Example 4.** *In  $\mathbb{Z}^2$ , the following are shift inclusion sequences:*

$$B_1 = \circ, \quad B_2 = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet \\ \bullet & \bullet & \bullet \end{array}, \quad B_3 = \begin{array}{ccccc} \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \circ & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array},$$

$$B_1 = \circ, \quad B_2 = \begin{array}{ccc} & \bullet & \\ \bullet & \circ & \bullet \\ & \bullet & \end{array}, \quad B_3 = \begin{array}{ccccc} & & \bullet & & \\ & \bullet & \bullet & \bullet & \\ \bullet & \bullet & \circ & \bullet & \bullet \\ & \bullet & \bullet & \bullet & \\ & & \bullet & & \end{array},$$

and

$$B_1 = \circ, \quad B_2 = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet \\ \bullet & \bullet & \bullet \end{array}, \quad B_3 = \begin{array}{ccccc} \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \circ & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}.$$

Sequences of structuring elements shown in Example 4 are symmetric. As we studied in Euclidean spaces, a symmetric structuring elements in  $\mathbb{Z}^m$  are analogue to neighborhood of the origin in  $\mathbb{R}^m$ . However, non-symmetric structuring elements are also used for proposing digital images. Here are some examples:

**Example 5.** *In  $\mathbb{Z}^2$ , the following are shift inclusion sequences:*

$$B_1 = \circ, \quad B_2 = \begin{array}{cc} \bullet & \bullet \\ \circ & \bullet \end{array}, \quad B_3 = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet \\ \bullet & \bullet & \bullet \end{array}, \quad B_4 = \begin{array}{ccccc} \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \circ & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array},$$

$$B_1 = \circ, \quad B_2 = \begin{array}{ccccc} & & \bullet & & \\ \bullet & \circ & \bullet & \bullet & \bullet \\ & \bullet & & & \end{array}, \quad B_3 = \begin{array}{ccccc} & & \bullet & \bullet & \bullet & \bullet \\ & \bullet & \circ & \bullet & \bullet & \bullet \\ & & \bullet & & & \end{array},$$

$$B_1 = \circ, \quad B_2 = \begin{array}{ccc} \bullet & & \\ \circ & \bullet & \end{array}, \quad B_3 = \begin{array}{ccc} \bullet & & \\ \bullet & \bullet & \\ \circ & \bullet & \bullet \end{array},$$

and

$$B_1 = \circ, \quad B_2 = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet \\ \bullet & \bullet & \bullet \end{array}, \quad B_3 = \begin{array}{ccccc} \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \circ & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}.$$

Although images defined on the whole space (e.g.  $\mathbb{Z}^2$ ) is not considered in real applications, the structuring elements obtained from Theorem 3 can still be applied in many real applications. Especially, if  $\{0\} \subseteq_{S,M} B_1 \subseteq_{S,M} B_2 \subseteq_{S,M} \dots \subseteq_{S,M} B_n$  are given and  $g$  is defined on a finite set  $P \subseteq M$ , then  $\tilde{P} := (P + B_n) \cup (P - B_n)$  is still finite and we can define

$$\tilde{g}(x) = \begin{cases} g(x) & \text{if } x \in P, \\ 0 & \text{otherwise.} \end{cases} \quad (19)$$

Then  $\{0\} \subseteq_{S,\tilde{P}} B_1 \subseteq_{S,\tilde{P}} B_2 \subseteq_{S,\tilde{P}} \dots \subseteq_{S,\tilde{P}} B_n$ . Therefore, the property (1) still holds for images  $\tilde{g}, O_{B_1}(\tilde{g}), O_{B_2}(\tilde{g}), \dots, O_{B_n}(\tilde{g})$ .

## 4 Order Preserving Property

To prove  $B_1 \subseteq_{S,P} B_2$  implies (1), the following observation is crucial.

**Lemma 3.** *Let  $M$  be an abelian group,  $x \in P \subseteq M$  and  $g \in \mathcal{I}_P$ . If  $B \subseteq M$  is a structuring element, then the following hold:*

- (a)  $O_B(g)(x) = 0$  if and only if  $b \in B(x; P, -)$  implies there is a  $b' \in B(x - b; P, +)$  such that  $g(x - b + b') = 0$ .
- (b)  $C_B(g)(x) = 0$  if and only if there is a  $b \in B(x; P, +)$  such that  $g(x + b - b') = 0$  whenever  $b' \in B(x + b; P, -)$ .

*Proof.* (a) By definition of dilation, for arbitrary structuring element  $B$  in  $M$ ,  $x \in P$  satisfies  $(\delta_B \circ \epsilon_B)(g)(x) = 0$  if and only if  $\epsilon_B(g)((x - B) \cap P) = \{0\}$  if and only if  $\epsilon_B(g)(x - b) = 0$  whenever  $b \in B(x; P, -)$  if and only if  $\min g((x - b) + B) \cap P = 0$  whenever  $b \in B(x; P, -)$ . It is equivalent to the statement:  $b \in B(x; P, -)$  implies that there is a  $b' \in B(x - b; P, +)$  such that  $g(x - b + b') = 0$ .

(b) By definition of erosion, for arbitrary structuring element  $B$  in  $M$ ,  $x \in P$  satisfies  $(\epsilon_B \circ \delta_B)(g)(x) = 0$  if and only if there is a  $b \in B(x; P, +)$  such that  $\delta_B(g)(x + b) = 0$  if and only if there is a  $b \in B(x; P, +)$  such that  $g((x + b) - B) \cap P = \{0\}$ . It is equivalent to the statement: there is a  $b \in B(x; P, +)$  such that  $g(x + b - b') = 0$  whenever  $b' \in B(x + b; P, -)$ .  $\square$

Next, we show that requiring that structuring elements are nested using shift inclusion with respect to  $P$  will allow us to obtain (1).

**Theorem 4** (Main Theorem, Weak Form). *Let  $P \subseteq M$  be a pixel set and  $B_1, B_2 \subseteq M$  be structuring elements. Then the following statements hold:*

- (a) If  $B_1 \subseteq_{S,P,-} B_2$ , then  $O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_P$ ;
- (b) If  $B_1 \subseteq_{S,P,+} B_2$ , then  $C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_P$ .

In particular, if  $B_1 \subseteq_{S,P} B_2$ , then for  $g \in \mathcal{I}_P$ ,  $O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0)$ .

*Proof.* (a) Suppose  $(\delta_{B_1} \circ \epsilon_{B_1})(g)(x) = 0$  and  $b_2 \in B_2(x; P, -)$ . By assumption  $B_1 \subseteq_{S,P,-} B_2$ , there is a  $b_1 \in B_1(x; P, -)$  such that  $B_1 + (b_2 - b_1) \subseteq B_2$ . By Lemma 3-(a), there is a  $b'_1 \in B_1(x - b_1; P, +)$  such that  $g(x - b_1 + b'_1) = 0$ . Observe that  $x - b_2 + (b'_1 + b_2 - b_1) = x - b_1 + b'_1 \in P$  with  $b'_1 + b_2 - b_1 \in B_1 + (b_2 - b_1) \subseteq B_2$ , this shows that  $O_{B_2}(g)(x) = (\delta_{B_2} \circ \epsilon_{B_2})(g)(x) = 0$  by Lemma 3-(a).

(b) If  $x \in P$  and  $(\epsilon_{B_2} \circ \delta_{B_2})(g)(x) = 0$ , by Lemma 3-(b), there is a  $b_2 \in B_2(x; P, +)$  such that  $g(x + b_2 - b'_2) = 0$  whenever  $b'_2 \in B_2(x + b_2; P, -)$ . By assumption  $B_1 \subseteq_{S,P,+} B_2$ , there is a  $b_1 \in B_1(x; P, +)$  such that  $B_1 + (b_2 - b_1) \subseteq B_2$ . If  $b'_1 \in B_1(x + b_1; P, -)$ , then  $x + b_2 - (b'_1 + b_2 - b_1) = x + b_1 - b'_1 \in P$ . Because  $b'_1 + (b_2 - b_1) \in B_1 + (b_2 - b_1) \subseteq B_2$ , we deduce that  $g(x + b_1 - b'_1) = 0$ , this shows that  $(\epsilon_{B_1} \circ \delta_{B_1})(g)(x) = 0$  by Lemma 3-(b).  $\square$

Actually, by applying the following thresholding technique, the strong form of Theorem 4 follows:

**Corollary 1** (Strong Form). *Let  $P \subseteq M$  be a pixel set and  $B_1, B_2 \subseteq M$  be structuring elements. Then the following hold:*

- (a) If  $B_1 \subseteq_{S,P,-} B_2$ , then  $O_{B_2}(g) \leq O_{B_1}(g)$  whenever  $g \in \mathcal{I}_P$ ;
- (b) If  $B_1 \subseteq_{S,P,+} B_2$ , then  $C_{B_1}(g) \leq C_{B_2}(g)$  whenever  $g \in \mathcal{I}_P$ .

In particular, if  $B_1 \subseteq_{S,P} B_2$ , then for  $g \in \mathcal{I}_P$ ,  $O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0)$ .

By combining Theorem 1, Proposition 1 and Lemma 1, now we can prove Theorem 1.

*Proof for Theorem 1.* For every  $t \in \mathbb{R}_{\geq 0}$ ,  $g_t$  is a binary image. By Theorem 4, we have  $O_{B_2}(g_t) \leq O_{B_1}(g_t)$  and  $C_{B_1}(g_t) \leq C_{B_2}(g_t)$ . By Proposition 1,  $O_{B_2}(g)_t \leq O_{B_1}(g)_t$  and  $C_{B_1}(g)_t \leq C_{B_2}(g)_t$ . By Lemma 1, we have  $O_{B_2}(g) \leq O_{B_1}(g)$  and  $C_{B_1}(g) \leq C_{B_2}(g)$ . The remained part follows from the fact  $\{0\} \subseteq_{S,P} B$ .  $\square$

Properties (a) and (b) in Corollary 1 are stronger than property (1), where (a) is the absorption property in mathematical morphology [14].

#### 4.1 On the whole space

Except the decomposition theorem for structuring elements with respect to  $M$ . Next, we show that the shift inclusion  $\subseteq_{S,M}$  condition, actually, is equivalent to property (1).

**Theorem 5.** *Let  $M$  be an abelian group and  $B_1 \subseteq B_2 \subseteq M$  be structuring elements. Then the following holds:*

- (a)  $O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_M$  if and only if  $B_1 \subseteq_{S,M} B_2$ ;
- (b)  $C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_M$  if and only if  $B_1 \subseteq_{S,M} B_2$ .

*Proof.* The "if" parts of (a) and (b) are just the results of Theorem 4. Conversely, we assume that  $B_1 \not\subseteq_{S,M} B_2$ , then there is a  $b_2 \in B_2$  and a function  $\phi : B_1 \rightarrow B_1$  such that  $\phi(b_1) + (b_2 - b_1) \notin B_2$  for every  $b_1 \in B_1$ . For (a), consider the characteristic function  $\chi_{B_2}$ . Then  $\chi_{B_2}(b_2 - b_1 + \phi(b_1)) = 0$  for every  $b_1 \in B_1$ . This shows that  $O_{B_1}(\chi_{B_2})(b_2) = 0$  by Lemma 3-(a). On the other hand, because  $\chi_{B_2}(b_2 - b_2 + b'_2) = \chi_{B_2}(b'_2) = 1 \neq 0$  for every  $b'_2 \in B_2$ , we obtain  $O_{B_2}(\chi_{B_2})(b_2) \neq 0$  by Lemma 3-(a).

For (b), consider the anti-characteristic function  $\widetilde{\chi_{-B_2}} = 1 - \chi_{-B_2}$ . For every  $b_1 \in B_1$ ,  $-b_2 + b_1 - \phi(b_1) = -(\phi(b_1) + b_2 - b_1) \notin -B_2$ , hence  $\widetilde{\chi_{-B_2}}(-b_2 + b_1 - \phi(b_1)) = 1 \neq 0$ . This shows that  $C_{B_1}(\widetilde{\chi_{-B_2}})(-b_2) \neq 0$  by Lemma 3-(b). On the other hand,  $\widetilde{\chi_{-B_2}}(-b_2 + b_2 - b'_2) = \widetilde{\chi_{-B_2}}(-b'_2) = 0$  whenever  $b'_2 \in B_2$ , hence  $C_{B_2}(\widetilde{\chi_{-B_2}})(-b_2) = 0$  by Lemma 3-(b).  $\square$

## 5 Discussion and Generalization

In this section, we introduce the *weak shift inclusions*  $\subseteq_{WS,P}$  as a generalization of shift inclusions. Theorem 6 shows that the generalization is sufficient and necessary for obtaining the property (1).

We also complete the proofs of true/false of all directions in diagram (2) in the section. The criterion (a) in (2) means that  $\subseteq_{S,M}$  and  $\subseteq_{WS,M}$  are equivalent. Proposition 5 shows that  $\subseteq_{S,M}$  implies  $\subseteq_{WS,M}$ . Conversely, Theorem 6 shows that  $\subseteq_{WS,M}$  implies the property (1). Because  $\subseteq_{S,M}$  is equivalent to (1) (Theorem 5), it shows that  $\subseteq_{WS,M}$  implies  $\subseteq_{S,M}$ . Example 7 and Example 8 prove (b) and (c) respectively. Example 9 shows that  $\subseteq_{S,P}$  is not necessary for (1), hence  $\subseteq_{WS,P}$  doesn't imply  $\subseteq_{S,P}$  i.e., (d). Because (a) shows that  $\subseteq_{S,M}$  and  $\subseteq_{WS,M}$  are equivalent, Example 8 shows that  $\subseteq_{WS,M}$  doesn't imply  $\subseteq_{WS,P}$ , which is (e). The (f) is proved by Proposition 5. Finally, (g) is proved by Theorem 6 and Example 9.

First, we discuss images defined on different domains. The behaviours of shift inclusion defined on different domain are different. For example, in general, for every  $B_1 \subseteq B_2 \subseteq M$  and  $P \subseteq M$ , the relations  $B_1 \subseteq_{S,M} B_2$  and  $B_1 \subseteq_{S,P} B_2$  do not have any directional implications. The following example shows that  $B_1 \subseteq_{S,M} B_2$  may not imply  $B_1 \subseteq_{S,P} B_2$ .

**Example 6.** Consider image  $g : P \rightarrow \{0, 1\}$  with  $P \subseteq \mathbb{Z}^2$  defined by

$$g = \begin{array}{cccccc} & 1 & 0 & 0 & 0 & 0 \\ 1 & & & & & \end{array} \quad \text{with } P = \begin{array}{c} \bullet \bullet \bullet \bullet \bullet \\ \bullet \end{array}$$

and structuring elements

$$B_1 = \begin{array}{cc} \circ & \bullet \end{array} \quad \text{and} \quad B_2 = \begin{array}{cc} \bullet & \bullet \\ \circ & \bullet \end{array}$$

in  $\mathbb{Z}^2$ . Then  $B_1 \subseteq B_2$  and

$$\epsilon_{B_1}(g) = \begin{array}{cccccc} & 0 & 0 & 0 & 0 & 0 \\ 1 & & & & & \end{array} \quad \text{and} \quad O_{B_1}(g) = \begin{array}{cccccc} & 0 & 0 & 0 & 0 & 0 \\ 1 & & & & & \end{array} = \epsilon_{B_1}(g) \quad (20)$$

by applying operator  $\delta_{B_1}$  on  $\epsilon_{B_1}(g)$ . Similarly,

$$\epsilon_{B_2}(g) = \begin{matrix} & 0 & 0 & 0 & 0 & 0 \\ 1 & & & & & \end{matrix} \text{ and } O_{B_2}(g) = \begin{matrix} & 1 & 0 & 0 & 0 & 0 \\ 1 & & & & & \end{matrix}. \quad (21)$$

Observe pixel values in equations (20) and (21),  $O_{B_1}(g)^{-1}(0) \not\subseteq O_{B_2}(g)^{-1}(0)$ , this shows that  $B_1 \not\subseteq_{S,P} B_2$  by Theorem 4. However, Theorem 3 shows that  $B_1 \subseteq_{S,\mathbb{Z}^2} B_2$  since  $B_2 = B_1 + \mathbf{e}_2$ .

On the other hand, the following trivial example shows that  $B_1 \subseteq_{S,P} B_2$  do not implies  $B_1 \subseteq_{S,M} B_2$ .

**Example 7.** Consider image domain  $P \subseteq \mathbb{Z}^2$  defined by

$$P = \bullet \bullet \bullet \bullet \bullet \bullet$$

and consider structuring elements

$$B_1 = \begin{matrix} \bullet \\ \circ \end{matrix} \text{ and } B_2 = \begin{matrix} \bullet & \bullet \\ \circ & \end{matrix}$$

in  $\mathbb{Z}^2$  where  $\circ$  represents the origin  $(0,0)$ . Clearly,  $B_1 \subseteq B_2$  but  $B_1 \not\subseteq_{S,\mathbb{Z}^2} B_2$  by Theorem 3. However,  $B_1 \subseteq_{S,P} B_2$  by directly checking.

Because rectangles in  $\mathbb{Z}^m$  has similar form of  $\mathbb{Z}^m$  and is more regular than domain as in Example 6, a natural question occurs: If  $P$  is a rectangle, dose  $B_1 \subseteq_{S,\mathbb{Z}^m} B_2$  implies  $B_1 \subseteq_{S,P} B_2$ . Unfortunately, here is an counter-example:

**Example 8.** Consider image  $g : P \rightarrow \{0,1\}$  with  $P \subseteq \mathbb{Z}^2$  defined by

$$g = \begin{matrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{matrix} \text{ with } P = \begin{matrix} \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{matrix}$$

and consider structuring elements

$$\begin{aligned} B_1 &= \begin{matrix} \bullet \\ \circ \end{matrix} = \{(0,0), (2,2)\}, \\ B_2 &= \begin{matrix} \bullet \\ \circ \end{matrix} \bullet = \{(0,0), (2,2), (0,-2), (2,0)\}. \end{aligned}$$

in  $\mathbb{Z}^2$  where  $\circ$  represents the origin  $(0,0)$ . Theorem 3 shows that  $B_1 \subseteq_{S,\mathbb{Z}^2} B_2$  since  $B_2 = B_1 + (0,-2)$  and

$$\epsilon_{B_1}(g) = \begin{matrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 \end{matrix}.$$

By applying operator  $\delta_{B_1}$  on  $\epsilon_{B_1}(g)$ , we have

$$O_{B_1}(g) = \begin{matrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 \end{matrix} = \epsilon_{B_1}(g).$$

Similarly,

$$\epsilon_{B_2}(g) = \begin{matrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 \end{matrix} \text{ and } O_{B_2}(g) = (\delta_{B_2} \circ \epsilon_{B_2})(g) = \begin{matrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{matrix} = g.$$

By comparing pixel values, the set  $O_{B_1}(g)^{-1}(0)$  would not be a subset of  $O_{B_2}(g)^{-1}(0)$ . Hence  $B_1 \not\subseteq_{S,P} B_2$ .

However, the following example shows that, for arbitrary pixel domain  $P$ , the condition  $B_1 \subseteq_{S,P} B_2$  is not equivalent to property (1).

**Example 9.** Consider image image domain  $P \subseteq \mathbb{Z}^2$  and structuring elements  $B_1, B_2 \subseteq \mathbb{Z}^2$  defined by

$$P = \begin{smallmatrix} \circ & \bullet \\ \bullet & \end{smallmatrix} = \{(0,0), (1,0), (0,-1)\}, B_1 = \begin{smallmatrix} \circ \\ \bullet \end{smallmatrix} \text{ and } B_2 = \begin{smallmatrix} \circ & \bullet \\ \bullet & \end{smallmatrix}.$$

Then  $B_1 \not\subseteq_{S,P} B_2$  since  $(1,0) \in B_2((0,0); P, +)$ ,  $B_1 + (1,0) \not\subseteq B_2$ , and  $B_1 + (1,1) \not\subseteq B_2$ . However,  $O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_P$ .

*Proof.* We prove the example by exhibiting all images on  $P$  as follow: consider  $a, b, c > 0$ , any image must be one of the following forms:

- If

$$g = \begin{smallmatrix} a & b \\ c & \end{smallmatrix},$$

then the preimages of  $O_{B_1}(g), O_{B_2}(g), C_{B_1}(g)$ , and  $C_{B_2}(g)$  on  $\{0\}$  are  $\emptyset$ ;

- If

$$g = \begin{smallmatrix} 0 & b \\ c & \end{smallmatrix},$$

then  $O_{B_1}(g)^{-1}(0) = \{(0,0)\} = O_{B_2}(g)^{-1}(0)$  and  $C_{B_1}(g)^{-1}(0) = \{(0,0)\} = C_{B_2}(g)^{-1}(0)$ .

- If

$$g = \begin{smallmatrix} a & 0 \\ c & \end{smallmatrix},$$

then  $O_{B_1}(g)^{-1}(0) = \{(1,0)\} \subseteq \{(1,0), (0,0)\} = O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) = \emptyset \subseteq C_{B_1}(g)^{-1}(0)$ .

- If

$$g = \begin{smallmatrix} a & b \\ 0 & \end{smallmatrix},$$

then  $O_{B_1}(g)^{-1}(0) = \{(0,0), (0,-1)\} = O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) = \emptyset = C_{B_1}(g)^{-1}(0)$ .

- If

$$g = \begin{smallmatrix} 0 & 0 \\ c & \end{smallmatrix},$$

then  $O_{B_1}(g)^{-1}(0) = \{(0,0), (1,0)\} = O_{B_2}(g)^{-1}(0)$  and the preimages of closing are  $C_{B_2}(g)^{-1}(0) = \{(0,0), (1,0)\} = C_{B_1}(g)^{-1}(0)$ .

- If

$$g = \begin{smallmatrix} a & 0 \\ 0 & \end{smallmatrix},$$

then  $O_{B_1}(g)^{-1}(0) = P = O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) = \emptyset \subseteq C_{B_1}(g)^{-1}(0)$ .

- If

$$g = \begin{smallmatrix} 0 & b \\ 0 & \end{smallmatrix},$$

then  $O_{B_1}(g)^{-1}(0) = \{(0,0), (0,-1)\} = O_{B_2}(g)^{-1}(0)$  and the preimages of closing are  $C_{B_2}(g)^{-1}(0) = \{(0,0), (0,-1)\} = C_{B_1}(g)^{-1}(0)$ .

- If

$$g = \begin{smallmatrix} 0 & 0 \\ 0 & \end{smallmatrix},$$

then  $O_{B_1}(g)^{-1}(0) = P = O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) = P = C_{B_1}(g)^{-1}(0)$ .

By the discussions,  $O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0)$  and  $C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_P$ .  $\square$

In this section, we introduce the definition of weak shift inclusion of structuring elements, and we prove this condition is equivalent to desired property (1) in Theorem 6.

**Definition.** Given pixel set  $P \subseteq M$  and two structuring elements  $B_1, B_2 \subseteq M$  we say that  $B_1$  is **weak shift included with respect to  $P$**  in  $B_2$ , denoted  $B_1 \subseteq_{WS,P} B_2$ , if  $B_1 \subseteq B_2$  and the following properties hold:

- (a) (**Weak Positive property**, denoted by  $B_1 \subseteq_{WS,P,+} B_2$ ) For any  $x \in P$  and  $b_2 \in B_2(x; P, +)$ , there exists a  $b_1 \in B_1(x; P, +)$  such that if  $b'_1 \in B_1(x+b_1; P, -)$ , then there is a  $b'_2 \in B_2$  such that  $x+b_2-b'_2 = x+b_1-b'_1$ ;
- (b) (**Weak Negative property**, denoted by  $B_1 \subseteq_{WS,P,-} B_2$ ) For any  $x \in P$  and  $b_2 \in B_2(x; P, -)$ , there exists a  $b_1 \in B_1(x; P, -)$  such that if  $b'_1 \in B_1(x-b_1; P, +)$ , then there is a  $b'_2 \in B_2$  such that  $x-b_2+b'_2 = x-b_1+b'_1$ .

The following proposition shows that shift inclusion must implies weak shift inclusion.

**Proposition 5.** Given  $P \subseteq M$  and two structuring elements  $B_1, B_2 \subseteq M$ . Then the following properties hold:

- (a) If  $B_1 \subseteq_{S,P,+} B_2$ , then  $B_1 \subseteq_{WS,P,+} B_2$ ;
- (b) If  $B_1 \subseteq_{S,P,-} B_2$ , then  $B_1 \subseteq_{WS,P,-} B_2$ .

*Proof.* (a) Suppose  $B_1 \subseteq_{S,P,+} B_2$ , then for  $b_2 \in B_2(x; P, +)$ , there is a  $b_1 \in B_1(x; P, +)$  such that  $B_1 + (b_2 - b_1) \subseteq B_2$ . In particular, if  $b'_1 \in B_1(x+b_1; P, -)$ , then  $x+b_1-b'_1 = x+b_2-(b'_1+(b_2-b_1))$  where  $b'_1+(b_2-b_1) \in B_1+(b_2-b_1) \subseteq B_2$ .

(b) Suppose  $B_1 \subseteq_{S,P,-} B_2$ , then for  $b_2 \in B_2(x; P, -)$ , there is a  $b_1 \in B_1(x; P, -)$  such that  $B_1 + (b_2 - b_1) \subseteq B_2$ . In particular, if  $b'_1 \in B_1(x-b_1; P, +)$ , then  $x-b_1+b'_1 = x-b_2+(b'_1+(b_2-b_1))$  where  $b'_1+(b_2-b_1) \in B_1+(b_2-b_1) \subseteq B_2$ .  $\square$

**Theorem 6.** Let  $M$  be an abelian group,  $P \subseteq M$  and  $B_1 \subseteq B_2 \subseteq M$  be structuring elements. Then

- (a)  $O_{B_1}(g)^{-1}(0) \subseteq O_{B_2}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_P$  if and only if  $B_1 \subseteq_{WS,P,+} B_2$ ;
- (b)  $C_{B_2}(g)^{-1}(0) \subseteq C_{B_1}(g)^{-1}(0)$  whenever  $g \in \mathcal{I}_P$  if and only if  $B_1 \subseteq_{WS,P,-} B_2$ .

*Proof.* (a) Suppose  $x \in P$  and  $b_2 \in B_2(x; P, -)$ . If  $B_1 \subseteq_{WS,P,-} B_2$ , there is a  $b_1 \in B_1(x; P, -)$  such that  $b'_1 \in B_1(x-b_1; P, +)$  implies there is a  $b'_2 \in B_2(x-)$  such that  $x-b_1+b'_1 = x-b_2+b'_2$ . By Lemma 3-(a), if  $(\delta_{B_1} \circ \epsilon_{B_1})(g)(x) = 0$ , there is a  $b''_1 \in B_1(x-b_1; P, +)$  such that  $g(x-b_1+b''_1) = 0$ . By assumption, there is a  $b''_2 \in B_2$  such that  $x-b_1+b''_1 = x-b_2+b''_2$ , hence  $g(x-b_2+b''_2) = g(x-b_1+b''_1) = 0$ . This shows that  $O_{B_2}(g)(x) = (\delta_{B_2} \circ \epsilon_{B_2})(g)(x) = 0$  by Lemma 3-(a).

Conversely, suppose  $B_1 \not\subseteq_{WS,P,-} B_2$ . Then there are  $x_0 \in P$  and  $b_2 \in B_2(x_0; P, -)$  satisfy the following property: for every  $b_1 \in B_1(x_0; P, -)$  there is a  $b'_1 \in B_1(x_0-b_1; P, +)$  such that  $x_0-b_2+b'_2 \neq x_0-b_1+b'_1$  for every  $b'_2 \in B_2$ . Consider  $g : P \rightarrow \{0, 1\}$  defined by

$$g(x) = \begin{cases} 1 & \text{if } x \in ((x_0 - b_2) + B_2) \cap P, \\ 0 & \text{otherwise.} \end{cases} \quad (22)$$

Note that  $((x_0 - b_2) + B_2) \cap P$  is non-empty because  $x_0 \in ((x_0 - b_2) + B_2) \cap P$ . This shows that  $g(x_0 - b_1 + b'_1) = 0$  and hence  $O_{B_1}(g)(x_0) = 0$  by Lemma 3-(a). On the other hand, because  $g(((x_0 - b_2) + B_2) \cap P) = \{1\}$ ,  $\epsilon_{B_2}(g)(x_0 - b_2) = 1$  and we obtain

$$O_{B_2}(g)(x_0) = (\delta_{B_2} \circ \epsilon_{B_2})(g)(x_0) = \max \epsilon_{B_2}(g) \left( (x_0 - B_2) \cap P \right) = 1 \neq 0$$

because  $x_0 - b_2 \in (x_0 - B_2) \cap P$  and  $\epsilon_{B_2}(g)(x_0 - b_2) = 1$ .

(b) If  $x \in P$  and  $(\epsilon_{B_2} \circ \delta_{B_2})(g)(x) = 0$ , by Lemma 3-(b), there is a  $b_2 \in B_2(x; P, +)$  such that  $g(x+b_2-b'_2) = 0$  whenever  $b'_2 \in B_2(x+b_2; P, -)$ . By assumption  $B_1 \subseteq_{WS,P,+} B_2$ , there is a  $b_1 \in B_1(x; P, +)$  such that  $b'_1 \in B_1(x+b_1; P, -)$  implies there is a  $b'_2 \in B_2$  such that  $x+b_2-b'_2 = x+b_1-b'_1 \in P$ . For such  $b'_1$ , we deduce that  $g(x+b_1-b'_1) = g(x+b_2-b'_2) = 0$ , this shows that  $(\epsilon_{B_1} \circ \delta_{B_1})(g)(x) = 0$  by Lemma 3-(b).

Conversely, suppose  $B_1 \not\subseteq_{WS,P,+} B_2$ . Then there are  $x_0 \in P$  and  $b_2 \in B_2(x_0; P, +)$  such that for every  $b_1 \in B_1(x_0; P, +)$  there is a  $b'_1 \in B_1(x_0+b_1; P, b'_1)$  such that  $x_0+b_2-b'_2 \neq x_0+b_1-b'_1$  for every  $b'_2 \in B_2$ . Consider  $h : P \rightarrow \{0, 1\}$  defined by

$$h(x) = \begin{cases} 0 & \text{if } x \in ((x_0 + b_2) - B_2) \cap P, \\ 1 & \text{otherwise.} \end{cases} \quad (23)$$

In particular, for such  $b_1$  and  $b'_1$ , we must have  $x_0+b_1-b'_1 \notin (x_0+b_2) - B_2$ , hence  $h(x_0+b_1-b'_1) = 1$ . By Lemma 3-(a), we must have  $C_{B_1}(h)(x_0) \neq 0$ . On the other hand, if  $b'_2 \in B_2$  satisfies  $x_0+b_2-b'_2 \in P$  then  $h(x_0+b_2-b'_2) = 0$ , this shows that  $C_{B_2}(h)(x_0) = 0$  by Lemma 3-(a) again.  $\square$

In this paper, we introduced the definition of shift inclusion of structuring elements as a sufficient condition for the desired absorption property (1). When the image domain is the whole abelian group, the shift inclusion condition is actually equivalent to (1) and can be systemically described by the decomposition theorem (Theorem 3). For an arbitrary image domain in an abelian group, we proved that the shift inclusion condition is strictly stronger than (1). Therefore, the weak shift inclusion was introduced, and we proved it is equivalent to (1) on every subset of a fixed abelian group. All theories introduced here is purely based on the addition operation of the based group, and can be adapted for analyzing real digital images. These properties can be verified and implemented easily, and also provide a solid foundation for constructing filtrations of images via opening and closing operations.

## Acknowledgments

Chuan-Shen Hu is supported by the projects MOST 108-2119-M-002-031 and MOST 108-2115-M-003-005-MY2 hosted by the Ministry of Science and Technology in Taiwan.

## References

- [1] Luc Vincent. Morphological area openings and closings for grey-scale images. In *Shape in Picture*, pages 197–208, Berlin, Heidelberg, 1994. Springer Berlin Heidelberg.
- [2] H.J.A.M. Heijmans and J. Roerdink. *Mathematical Morphology and its Applications to Image and Signal Processing*. Computational Imaging and Vision. Springer Netherlands, 1998.
- [3] Jean Cousty. *Segmentation, hierarchy, mathematical morphology filtering, and application to image analysis*. Habilitation à diriger des recherches, Université Paris-Est, September 2018.
- [4] Huijian Li, Runqiu Wang, Siyuan Cao, Yangkang Chen, Nan Tian, and Xiaoqing Chen. Weak signal detection using multiscale morphology in microseismic monitoring. *Journal of Applied Geophysics*, 133:39 – 49, 2016.
- [5] R. M. Haralick, S. R. Sternberg, and X. Zhuang. Image analysis using mathematical morphology. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, PAMI-9(4):532–550, July 1987.
- [6] George Kour and Raid Saabne. Real-time segmentation of on-line handwritten arabic script. In *Frontiers in Handwriting Recognition (ICFHR), 2014 14th International Conference on*, pages 417–422. IEEE, 2014.
- [7] George Kour and Raid Saabne. Fast classification of handwritten on-line arabic characters. In *Soft Computing and Pattern Recognition (SoCPaR), 2014 6th International Conference of*, pages 312–318. IEEE, 2014.
- [8] R. M. Haralick. Mathematical morphology and computer vision. In *Twenty-Second Asilomar Conference on Signals, Systems and Computers*, volume 1, pages 468–479, 1988.
- [9] M. Sonka, V. Hlavac, and R. Boyle. *Image Processing, Analysis, and Machine Vision*. Cengage Learning, 2014.
- [10] B. Pawar, V. T. Humbe, and L. Kundnani. Morphology based moving vehicle detection. In *2017 International Conference on Big Data Analytics and Computational Intelligence (ICBDAC)*, pages 217–223, 2017.
- [11] Gianni Franchi, Amin Fehri, and Angela Yao. Deep morphological networks. *Pattern Recognition*, 102:107246, 2020.
- [12] Anthony Iheonye, Yvan Garipey, and Vijaya Raghavan. Computer vision for real-time monitoring of shrinkage for peas dried in a fluidized bed dryer. *Drying Technology*, 38(1-2):130–146, 2020.
- [13] J. C. Klein and J. Serra. The texture analyser. *Journal of Microscopy*, 95(2):349–356, 1972.
- [14] Pierre Soille. *Morphological Image Analysis: Principles and Applications*. Springer-Verlag New York, Inc., Secaucus, NJ, USA, 2 edition, 2003.
- [15] A. Colomer, J. Igual, and V. Naranjo. Detection of early signs of diabetic retinopathy based on textural and morphological information in fundus images. *Sensors*, 20(4):1005, 2020.
- [16] Victor Gager, David Legland, Alain Bourmaud, Antoine Le Duigou, Floran Pierre, Karim Behloul, and Christophe Baley. Oriented granulometry to quantify fibre orientation distributions in synthetic and plant fibre composite preforms. *Industrial Crops and Products*, 152:112548, 2020.
- [17] Pritom Jyoti Bora, Irthasa Azeem, K. Vinoy, Praveen Ramamurthy, and Giridhar Madras. Morphology controllable microwave absorption property of polyvinylbutyral (pvp)-mno 2 nanocomposites. *Composites Part B: Engineering*, 132, 09 2017.



- [18] Prakash Krishnaiah, Chantara Thevy Ratnam, and Sivakumar Manickam. Morphology water absorption and biodegradable properties of polylactide biocomposites reinforced with sisal fibers. *Materials Today: Proceedings*, 5(10, Part 3):22506 – 22516, 2018. International Conference on Advances in Science & Engineering ICASE -2017, 19th – 22nd January 2017.
- [19] P. Maragos and R. Schafer. Morphological filters—part i: Their set-theoretic analysis and relations to linear shift-invariant filters. *IEEE Transactions on Acoustics, Speech, and Signal Processing*, 35(8):1153–1169, 1987.
- [20] P. Maragos and R. Schafer. Morphological filters—part ii: Their relations to median, order-statistic, and stack filters. *IEEE Transactions on Acoustics, Speech, and Signal Processing*, 35(8):1170–1184, 1987.
- [21] Lasse Koskinen, Jaakko T. Astola, and Yrjö A. Neuvo. Soft morphological filters. In Paul D. Gader and Edward R. Dougherty, editors, *Image Algebra and Morphological Image Processing II*, volume 1568, pages 262 – 270. International Society for Optics and Photonics, SPIE, 1991.
- [22] J. Serra. *Image Analysis and Mathematical Morphology*. Number 1 in Image Analysis and Mathematical Morphology. Academic Press, 1984.
- [23] Álvaro Ginés Legaz-Aparicio, Rafael Verdú-Monedero, and Jesús Angulo. Adaptive morphological filters based on a multiple orientation vector field dependent on image local features. *Journal of Computational and Applied Mathematics*, 330:965 – 981, 2018.
- [24] Ryan Armstrong, James McClure, Vanessa Robins, Zhishang Liu, Christoph Arns, Steffen Schlüter, and Steffen Berg. Porous media characterization using minkowski functionals: Theories, applications and future directions. *Transport in Porous Media*, 11 2018.
- [25] Cris Luengo Hendriks and Lucas Van Vliet. Discrete morphology with line structuring elements. *10th International Conference, CAIP 2003, Groningen, The Netherlands, Proceedings*, 2003.
- [26] Laurent Najman and Hugues Talbot. *Mathematical Morphology*. Wiley-ISTE, 1d edition, 2010.
- [27] Anders Landström. *Elliptical Adaptive Structuring Elements for Mathematical Morphology*. PhD thesis, Luleå University of Technology, Signals and Systems, 2014.