Benign overfitting without concentration

January 5, 2021

Zong Shang¹

Abstract

We obtain a sufficient condition for benign overfitting of linear regression problem. Our result does not rely on concentration argument but on small-ball assumption and thus can holds in heavy-tailed case. The basic idea is to establish a coordinate small-ball estimate in terms of effective rank so that we can calibrate the balance of epsilon-Net and exponential probability. Our result indicates that benign overfitting is not depending on concentration property of the input vector. Finally, we discuss potential difficulties for benign overfitting beyond linear model and a benign overfitting result without truncated effective rank.

1 Introduction

In recent years, there are tremendous interest in studying generalization property of statistical model when it interpolates the input data. The classical learning theory suggests that when the predictor fits input data perfectly, it will suffer from noise so that it will not generalize well. To overcome this problem, regularization and penalized learning procedures like LASSO are studied to weaken the effect of noise to avoid overfitting. However, some empirical experiments indicate that overfitting may perform well. Why can overfitting perform well? In what cases can overfitting perform well? These questions motivated a series work on this field.

The original motivation is from the Deep Learning community, who empirically revealed that overfitting Deep Neural Network can still generalize well, see [31]. This counter-intuitive phenomenon still appear for linear regression and kernel ridge regression, see [3] and [28]. They believe that investigating benign overfitting phenomenon in linear regression case will benefit to the more complex Deep Neural Networks case.

The cornerstone work [2] presented a minimax bound of generalization error of overfitting linear regression. Their result is in terms of effective ranks, which measures the tail behavior of eigenvalues of covariance matrix and will be defined

 $^{^1{\}rm shang zong 2117}@$ mails.jlu.edu.cn, College of Computer Science and Technology, Jilin University, China.

later in our paper. Recently, [8] improved their results to the large deviation regime. Both their results rely on the assumption that the input vector is a gaussian vector. This assumption is relaxed in [28] to sub-gaussian vector. However, all these work use concentration-argument and thus only adapt to a number of well-behaved distributions. In heavy-tailed case, small-ball method is often employed to study generalization property of statistical models, see [13]. However, the original coordinate small ball estimate cannot be directly used when the input vector is anisotropic, but anisotropicity is a necessary condition for benign overfitting. Fortunately, this issue can be solved by a simple modification. In this paper, we derive a sufficient condition for benign overfitting when the input is heavy-tailed by using small-ball method.

Benign overfitting phenomenon was firstly discovered by [31], and a great deal of effort in it has been devoted to the investigating its reason. [17], [10] studied asymptotic generalization error in random feature setting. In fact, benign overfitting in linear regression is not equivalent to that of random feature because parameters in random features cannot be controlled to minimize empirical loss like that in linear regression(only parameters in second layer can be used to minimize empirical loss while others in first layer are randomized). However, their empirical results illustrates that linear regression and random features of shallow neural network share similar double-descent risk curve. [2] obtained a two-sides non-asymptotic generalization bound of prediction error in gaussian linear regression setting. Their result is in terms of effective ranks, which is a truncated version of stable rank in Asymptotic Geometric Analysis. [28] generalized it into sub-gaussian linear regression. Their result is also in terms of effective ranks. [14], [16], and [15] studied benign overfitting in Reproduced Kernel Hilbert Space. [4] showed that interpolant maybe the optimal predictor in some cases.

The closest to our work is [8], who derived a sufficient condition for benign overfitting for gaussian linear regression in terms of effective rank. We obtain similar results but we only assume the input satisfies small ball assumption, in stead of gaussian distribution. Our result can also be partially compared to [28], where they assumed both sub-gaussian and a marginal small ball assumption.

1.1 Background and notation

In this paper, we consider linear regression problems in \mathbb{R}^p . Given a dataset $D_N = \{(X_i, Y_i)\}_{i=1}^N$ and $Y_i = \langle X_i, \alpha^* \rangle + \xi_i$, where $\alpha^* \in \mathbb{R}^p$ is an unknown vector and $(X_i)_{i=1}^N$ are i.i.d. copies of X, ξ_i are unpredictable i.i.d. centered sub-gaussian noise, which is independent with X. Because we are going to compare linear regression with more general functions class later, we also often use $f_\alpha(\cdot)$ to denote $\langle \cdot, \alpha \rangle$ in the following, and \mathcal{F}_A as a set of f_α such that $\alpha \in A$. Assume the random vector $X \in \mathbb{R}^p$ satisfies weak small ball assumption with parameter (\mathcal{L}, θ) , which will be defined in Definition 2.1 ,and denote its covariance matrix as Σ . Define the design matrix \mathbf{X} with N lines X_i^T . Denote response vector $\mathbf{Y} = (Y_1, \cdots, Y_N)$.

When p > n, the least-square estimator can interpolate D_N . Denote the one

that has the smallest ℓ_2 norm as $\hat{\alpha}$. That is to say,

$$\hat{\alpha} = \mathbf{X}^{\dagger} \mathbf{Y}.$$

where \mathbf{X}^{\dagger} is the Moore-Penrose pseudo inverse of \mathbf{X} . Denote $H_N \subset \mathbb{R}^p$ as $H_N = \{\alpha \in \mathbb{R}^p : \mathbf{X}\alpha = \mathbf{Y}\}$, we call H_N as interpolation space. We have

$$\hat{\alpha} = \operatorname*{argmin}_{\alpha \in H_N} \|\alpha\|_{\ell_2} \,.$$

We assume that $\operatorname{rank}(\Sigma) > N$, then $\hat{\alpha}$ exists almost surely.

Our loss function is squared loss, that is $\ell(t) = t^2$, and the loss of α is denoted by $\ell_{\alpha} = \ell(\langle \alpha - \alpha^*, \mathbf{X} \rangle)$. So the empirical excess risk is defined as

$$\mathbf{P}_{N}\mathcal{L}_{\alpha} = \mathbf{P}_{N}\left(\ell_{\alpha} - \ell_{\alpha^{*}}\right)$$

Benign overfitting depends on effective ranks of Σ . If $A \in \mathbb{R}^{p \times p}$ is a symmetric matrix, denote $\lambda_1(A) \geq \cdots \geq \lambda_p(A)$ as eigenvalues of A and $s_1(A) \geq \cdots \geq s_p(A)$ be its singular values. If there is no ambiguity, we will write s_i in stead of $s_i(A)$.

[2] defined two effective ranks:

$$r_k(\Sigma) = \frac{\sum_{i>k} \lambda_i(\Sigma)}{\lambda_{k+1}}, \quad R_k(\Sigma) = \frac{\left(\sum_{i>k} \lambda_i(\Sigma)\right)^2}{\sum_{i>k} \lambda_i^2(\Sigma)}.$$
 (1.1)

 $R_k(\Sigma)$ is a truncated version of stable rank that occurred in Asymptotic Geometric Analysis, see [30], [25] and [22] for a comprehensive review. Stable rank, denoted as srank_q(A), defined by

$$\operatorname{srank}_{q}(A) = \left(\frac{\|A\|_{S_{2}}}{\|A\|_{S_{q}}}\right)^{\frac{2q}{q-2}},$$

where $\|\cdot\|_{S_q}$ is the q-Schatten norm of A, that is to say, $\|A\|_{S_q} = \left(\sum_{i=1}^p s_i^q(A)\right)^{1/q}$. When q = 4, and $A = \Sigma^{1/2}$, then

$$\operatorname{srank}_{4}(\Sigma^{1/2}) = \left(\frac{\left\|\Sigma^{1/2}\right\|_{S_{2}}}{\left\|\Sigma^{1/2}\right\|_{S_{4}}}\right)^{4} = \frac{\left(\sum_{i=1}^{p} s_{i}^{2}(\Sigma^{1/2})\right)^{2}}{\sum_{i=1}^{p} s_{i}^{4}(\Sigma^{1/2})} = \frac{\left(\sum_{i=1}^{p} \lambda_{i}(\Sigma)\right)^{2}}{\sum_{i=1}^{p} \lambda_{i}^{2}(\Sigma)}.$$

It can be seen that $R_k(\Sigma)$ is the truncated version of srank₄($\Sigma^{1/2}$).

Apart from this, $r_k(\Sigma)$ is also a truncated version of the usual "effective rank" which is actually $\operatorname{tr}(\Sigma)/\lambda_1(\Sigma)$ in statistical literature, see [12],[26].

In fact, our result will be in terms of $R_k(\Sigma)$ instead of $r_k(\Sigma)$, which is the usual choice in most past work. However, this does not matter, because the two effective ranks are closely related, we refer the reader to Appendix A.6 in [2] for a comprehensive review.

For sake of simplicity, we define some extra notations. They have no special meanings, but will make our formula more clear. Denote

$$R_{k,2}(\Sigma) := \left(1 - \sqrt{\frac{k}{\operatorname{srank}_4(\Sigma)}}\right) R_k(\Sigma).$$
(1.2)

When k = 0, we have $R_{k,2}(\Sigma) = \operatorname{srank}_4(\Sigma)$. Denote

$$\mathfrak{R}_k(\Sigma) := \frac{(4p-k)^2}{8p} \frac{c^{\frac{16p^2}{(4p-k)^2}R_{k,2}(\Sigma)}}{R_{k,2}(\Sigma)},$$

where c < 1 is a constant.

Denote the operator norm of a matrix A as ||A||. Denote $||\alpha||_A$ as $\sqrt{\alpha^T A \alpha}$. We use S(r) to denote the sphere in \mathbb{R}^p with radius r with respect to $||\cdot||_{\ell_2}$, B(r) as ball analogously. Denote $S_A(r)$ and $B_A(r)$ as the corresponding sphere and ball with respect to $||\cdot||_A$. Denote $(\varepsilon_i)_{i=1}^N$ are i.i.d. Bernoulli random variables. Denote D as unit ball with respect to L_2 distance. If $F \subset L_2(\mu)$, let $\{G_f : f \in \mathcal{F}\}$ be the canonical gaussian process indexed by F, denote $\mathbb{E} ||G||_F$ as

$$\mathbb{E} \|G\|_F := \sup \left\{ \mathbb{E} \sup_{f \in F'} G_f : F' \subset F, F' \text{ is finite} \right\}.$$

Denote

$$\Lambda_{s_0,u}(\mathcal{F}) = \inf \sup_{f \in \mathcal{F}} \sum_{s \ge s_0} 2^{s/2} \, \|f - \pi_s f\|_{(u^2 2^s)}, \quad u \ge 1, \, s_0 \ge 0.$$

where the infimum is taken with respect to all admissible sequences $(F_s)_{s \ge 0}$ and $\pi_s f$ is the nearest point in F_s to f with respect to $\|\cdot\|_{(u^2 2^s)}$. Here $\|\cdot\|_{(p)} = \sup_{1 \le q \le p} \|\cdot\|_{L_q} / \sqrt{q}$. An admissible sequence is a sequence of partitions on \mathcal{F} such that $|F_s| \le 2^{2^s}$, and $|F_0| = 1$, cf. [21]. Denote $d_q(F)$ as diameter of Fwith respect to $\|\cdot\|_{L_q}$. Denote $\|\cdot\|_{\psi_2}$ as sub-gaussian norm. Denote [N] as set $\{1, 2, \dots, N\}$. Denote $C, c, c_0, c_1, c_2, \cdots$ as absolute constants.

1.2 Structure of this paper

Section 2 contains some preliminaries knowledge. Section 3 contains our main result, Theorem 3.1 and its proof are decomposed into two parts, which will be post-posed to section 4 and section 5. These two sections contains estimation error and prediction error of interpolation procedure in linear regression case. In section 6, we will discuss why it is difficult to obtain oracle inequality beyond linear regression, and give a benign overfitting result without truncated effective rank by using Dvoretzky-Milman Theorem in Asymptotic Geometric Analysis.

2 Preliminaries

In this section, we introduce some preliminary techniques which will be used to formulate and prove our main results. More precisely, we will introduce localization method to yield oracle inequality and small ball method to provide a lower bound of smallest singular value of design matrix.

2.1 Localization Method

To get an oracle inequality, there are two approaches in general. The first one is called Isomorphism Method, which uses the isomorphy between between empirical and actual structures to derive an oracle inequality. The other is called localization method. In this work, we will use localization method to derive an oracle inequality.

For a statistical model \mathcal{F} and $f^* \in \mathcal{F}$ is an oracle. Localization Method uses a L_2 -ball centered at f^* with radius r to localize model \mathcal{F} . This allows us to study statistical properties of a learning procedure \hat{f} on this small ball². More precisely, the radius r captures upper bound estimation error $||f - f^*||_{L_2}$ for all $f \in \mathcal{F} \cap rD$. Therefore, if we can find an upper bound of r, we find the estimation error of learning procedure \hat{f} . Analog to that in [8], our localized set in this paper is

$$\mathcal{F}_{H_{r,\rho}} = \{\langle \cdot, \alpha \rangle : \alpha \in H_{r,\rho}\}, \text{ where } H_{r,\rho} := B(\rho) \cap B_{\Sigma}(r),$$

where ρ is upper bound of estimation error, which will be studied in section 4, and r is upper bound of prediction error, which will be studied in section 5. Obtaining prediction risk is based on estimation risk. In this paper, we obtain estimation risk by studying minimum ℓ_2 interpolation procedure and obtain prediction error by localization method based on it.

Optimal level of r, denoted as r^* is carefully chosen by fixed points called complexity parameters.

2.1.1 Complexity parameters

In classical statistical learning theory, there are two common-used complexity parameters called multiplier complexity r_M and quadratic complexity r_Q , we refer the reader to [20] for a comprehensive view. Quadratic complexity r_Q is defined as follows:

$$r_{Q,1}(\mathcal{F},\zeta) = \operatorname*{arginf}_{r>0} \mathbb{E} \|G\|_{(\mathcal{F}-\mathcal{F})\cap rD} \leqslant \zeta r \sqrt{N},$$

and

$$r_{Q,2}(\mathcal{F},\zeta) = \operatorname*{arginf}_{r>0} \mathbb{E} \sup_{w \in (\mathcal{F}-\mathcal{F})\cap rD} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^{N} \varepsilon_i w(X_i) \right| \leqslant \zeta r \sqrt{N},$$

²This diameter is not necessarily to be of L_2 distance in localization method, though our choice is L_2 distance. We refer the interested reader to [7]

where ζ is an absolute constant.

 $r_Q(\zeta_1, \zeta_2) = \max \{ r_{Q,1}(\mathcal{F}, \zeta_1), r_{Q,2}(\mathcal{F}, \zeta_2) \}$ is an intrinsic parameter. That is to say, r_Q does not rely on noise ξ , but only on \mathcal{F} . This parameter measures the ability of \mathcal{F} to estimate target function f^* .

While multiplier complexity r_M is defined as follows:

$$\phi_N(r) := \sup_{w \in (\mathcal{F} - \mathcal{F}) \cap rD} \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i \xi_i w(X_i),$$
$$r_{M,1}(\kappa, \delta) = \inf_{r>0} \left\{ \mathbb{P} \left\{ \phi_N(r) \leqslant r^2 \kappa \sqrt{N} \right\} \ge 1 - \delta \right\}$$

and

$$r_0(\kappa) = \inf_{r>0} \sup_{w \in (\mathcal{F} - \mathcal{F}) \cap rD} \left\{ \|\xi w(X)\|_{L_2} \leqslant \frac{1}{2} \sqrt{N} \kappa r^2 \right\},$$

where κ is an absolute constant.

Then $r_M(\kappa, \delta) = r_{M,1} + r_0$ is called multiplier complexity, which measures the interplay between noise ξ and function class \mathcal{F} .

Classical learning theory employs r_M to measure the ability of \mathcal{F} to absorb noise ξ . However, this parameter does not make sense in interpolation case. This is because interpolant \hat{f} causes no loss on r_M by interpolating $(X_i, Y_i)_{i=1}^N$ perfectly. Therefore, $r_M = 0$ in this case. However, since \hat{f} interpolates $(X_i, Y_i)_{i=1}^N$, it bears influence from noise ξ so that r_Q is no longer an intrinsic parameter. That is to say, r_Q relies on ξ implicitly because \hat{f} has to estimate both signal and noise. It is this that causes the biggest difference from interpolation case and classical learning theory. Therefore, our complexity parameter is a variant of quadratic complexity, which will be defined in Equation 3.5.

Localization method employed complexity parameters to provide radius of localized set. However, we need to illustrate that interpolant \hat{f} lies in it with high probability. This step is guaranteed by an exclusion argument.

2.1.2 Exclusion

For all $f \in \mathcal{F}$, if f wants to be an interpolant, its empirical excess risk must be lower than a fixed level with high probability. To see this, we first decompose $\inf_{f \in \mathcal{F}} P_N \mathcal{L}_f$ to its lower bound.

There are two decompositions of empirical excess risk into quadratic and multiplier components. The first one is as follows:

$$\inf_{f \in \mathcal{F}} P_N \mathcal{L}_f = \inf_{f \in \mathcal{F}} \frac{1}{N} \sum_{i=1}^N (f(X_i) - Y_i)^2 - (f^*(X_i) - Y_i)^2 \\
\geqslant \inf_{f \in \mathcal{F}} \left\{ \frac{1}{N} \sum_{i=1}^N (f - f^*)^2 (X_i) \right\} - 2 \sup_{f \in \mathcal{F}} \left\{ \frac{1}{N} \sum_{i=1}^N \xi_i (f - f^*) (X_i) \right\} \\
:= \inf_{f \in \mathcal{F}} P_N \mathcal{Q}_{f - f^*} - 2 \sup_{f \in \mathcal{F}} P_N \mathcal{M}_{f - f^*}.$$
(2.1)

This kind of decomposition needs lower bound of quadratic component $\mathbb{P}_N \mathcal{Q}_{f-f^*}$. This lower bound is provided by small ball method. We will use this approach in subsection 6.2 to acquire a sufficient condition for benign overfitting without truncated effective rank. We turn to the second kind of decomposition. Our localized statistical model $\mathcal{F} = \mathcal{F}_{H_{r,\rho}}$ is a class of linear functionals on \mathbb{R}^p . Recall that the optimal choice of r is denoted as r^* , so denote $\theta = r^*/r \in (0, 1)$, and $\alpha_0 = \alpha^* + \theta(\alpha - \alpha^*)$, so $\left\| \Sigma^{1/2}(\alpha_0 - \alpha^*) \right\|_{\ell_2} = r^*$ and $\|\alpha_0 - \alpha^*\|_{\ell_2} \leq \theta \rho$. Denote

$$Q_{r,\rho} = \sup_{\alpha - \alpha^* \in H_{r,\rho}} \left| \frac{1}{N} \sum_{i=1}^N \langle X_i, \alpha - \alpha^* \rangle^2 - \mathbb{E} \langle X_i, \alpha - \alpha^* \rangle^2 \right|$$

and

$$M_{r,\rho} = \sup_{\alpha - \alpha^* \in H_{r,\rho}} \left| \frac{2}{N} \sum_{i=1}^N \xi_i \left\langle X_i, \alpha - \alpha^* \right\rangle \right|$$

Then

$$\inf_{f \in \mathcal{F}} \mathbf{P}_N \mathcal{L}_f = \inf_{f \in \mathcal{F}} \frac{1}{N} \sum_{i=1}^N (f(X_i) - Y_i)^2 - (f^*(X_i) - Y_i)^2$$

$$\geqslant \quad \theta^{-2} \left((r^*)^2 - Q_{r^*,\rho} \right) - 2M_{r^*,\rho}.$$
(2.2)

Suppose $(\xi_i)_{i=1}^N$ are i.i.d. sub-gaussian random variables, then by Bernstein's inequality, with probability at least $1 - \exp(-N/2)$ we have

$$\frac{1}{N}\sum_{i=1}^{N}\xi_{i}^{2} \leqslant \frac{1}{2}\left\|\xi\right\|_{\psi_{2}}^{2}$$

Because interpolation procedure \hat{f} interpolates all these inputs $(X_i, Y_i)_{i=1}^N$, the excess risk of \hat{f} can be obtained by the noises,

$$\mathbf{P}_N \mathcal{L}_{\hat{f}} = \mathbf{P}_N \left(\ell_{\hat{f}} - \ell_{f^*} \right) = -\mathbf{P}_N \ell_{f^*} = -\mathbf{P}_N \xi^2,$$

then with probability at least $1 - \exp(-N/2)$, we have

$$\mathbf{P}_{N}\mathcal{L}_{\hat{f}} \leqslant -\frac{1}{2} \left\|\xi\right\|_{\psi_{2}}^{2}.$$

That is to say, if f wants to be an interpolant, it must satisfy this upper bound. Otherwise, f will be excluded because it has little probability to be an interpolant.

This upper bound is different from the case of non-interpolation setting, where it is 0. It is smaller than 0 because the interpolation procedure is more restrict (in the sense of the interpolation space is smaller than version space) than non-interpolation procedure like ERM. The smaller upper bound can exclude more functions than non-interpolation procedure. Therefore, we just need to upper bound multiplier and quadratic processes in Equation 2.2, such that the lower bound of empirical excess risk over all $f \in \mathcal{F}_{H_{r,\rho}}$ is greater than $- \|\xi\|_{\psi_2}^2 / 2$ when r is greater than a fixed level r^* . So that functions in $\mathcal{F}_{H_{r,\rho}}$ will be excluded from being an interpolant with high probability. Therefore, with high probability, interpolant will lie in $H_{r^*,\rho}$ and we can upper bound its prediction risk by r^* .

2.1.3 Multiplier process and Quadratic process

We employ upper bounds of multiplier process and quadratic process in [21]:

Lemma 2.1 (Theorem 4.4 in [21]:Upper bound of multiplier process). There exist absolute constants c_1 and c_2 for which the following holds. If $\xi \in L_{\psi_2}$ then for every $u, w \ge 8$, with probability at least $1 - 2\exp(-c_1u^22^{s_0}) - 2\exp(-c_1Nw^2)$,

$$\sup_{f\in\mathcal{F}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^{N} \left(\xi_i f(X_i) - \mathbb{E}\xi f \right) \right| \leq cuw \left\| \xi \right\|_{\psi_2} \tilde{\Lambda}_{s_0, u}(\mathcal{F}),$$

where c is an absolute constant.

Lemma 2.2 (Theorem 1.13 in [21]: Upper bound of empirical process). There exists a constant c(q) that depends only on q for which the following holds. Then with probability at least $1 - 2\exp(-c_0 u^2 2^{s_0})$,

$$\sup_{f\in\mathcal{F}} \left| \frac{1}{N} \sum_{i=1}^{N} \left(f^2(X_i) - \mathbb{E}f^2 \right) \right| \leq \frac{c(q)}{N} \left(u^2 \tilde{\Lambda}_{s_0, u}^2(\mathcal{F}) + u\sqrt{N} d_q(\mathcal{F}) \tilde{\Lambda}_{s_0, u}(\mathcal{F}) \right),$$

where c(q) is a constant depending on q. Particularly, if \mathcal{F} is a sub-gaussian class, then with probability at least $1 - 2\exp(-c_0 N)$,

$$\sup_{f \in \mathcal{F}} \left| \frac{1}{N} \sum_{i=1}^{N} \left(f^2(X_i) - \mathbb{E}f^2 \right) \right| \leq cL^2 \mathbb{E} \left\| G \right\|_{\mathcal{F}}^2$$

Set $2^{s_0} = k_{\mathcal{F}}$, where $k_{\mathcal{F}} = (\mathbb{E} ||G||_{\mathcal{F}} / d_2(\mathcal{F}))^2$ is the Dvoretzky-Milman Dimension of \mathcal{F} , we refer the reader to [1] for a comprehensive view. For ℓ_2^p , the Dvoretzky-Milman dimension $k \sim p$, see e.g. Theorem 5.4.1 in [1].And $\tilde{\Lambda}(\mathcal{F})$ is called Λ -complexity of \mathcal{F} , which is a generalization of Gaussian complexity so that Λ -complexity just needs \mathcal{F} has finite order of moments, instead of infinite order of moments. Particularly, when \mathcal{F} happens to be a sub-gaussian class, $\tilde{\Lambda}(\mathcal{F})$ is equivalent to $\mathbb{E} ||G||_{\mathcal{F}}$. We refer the reader to [21] for a comprehensive review.

In this paper, our function class is of linear functionals on \mathbb{R}^p , especially ellipses since we assume the random vector X is not isotropic. Given the covariance matrix Σ of random vector X. By Lemma 5 in [8], we obtain

$$\mathbb{E} \|G\|_{\mathcal{F}_{H_{r,\rho}}} \leqslant \sqrt{2} \sqrt{\sum_{i=1}^{p} \min\left\{\lambda_i(\Sigma)\rho^2, r^2\right\}}.$$
(2.3)

However, estimating $\tilde{\Lambda}(H_{r,\rho})$ is non-trivial unless $H_{r,\rho}$ is a sub-gaussian class. Note that the deviation in Lemma 2.2 is neither optimal nor user-friendly(in the sense of the deviation parameter u is coupled with complexity parameter $\mathbb{E} \|G\|_{\mathcal{F}}$). In fact, upper bound of quadratic process given by [9] is in optimal deviation when \mathcal{F} is a sub-gaussian class. This is not fit to our heavy-tailed setup. It is non-trivial to obtain upper bound of quadratic process in heavytailed case. However, when \mathcal{F} is sub-gaussian, we can omit parameter r_2^* which will be defined in section 3 and get a better bound of $\|\Gamma\|$ in subsection 4.2, which will generate a preciser result, whereas the proof is omitted. To make our result uniform to both heavy-tailed and sub-gaussian case, we employ the one from [21] though it will not generate an optimal bound.

2.2 Small Ball Method

To deal with heavy-tailed case, we employ small ball method, which is a crucial argument in Asymptotic Geometric Analysis, see [1]. Small Ball Method in statistical learning theory is first developed in [13]. It can be viewed as a kind of Paley-Zygmund method, which assumes the random vector is sufficiently spread, so that it will have many large coordinates. We refer the reader to [20] and [22] for a comprehensive view.

Classical small ball assumption is a lower bound on tail of random function, that is

$$\mathbb{P}\left\{ |X_i| \ge \kappa \|X\|_{L_2} \right\} \ge \theta,$$

which can be verified by Paley-Zygmund inequality, see Lemma 3.1 in [11], under $L_4 - L_2$ norm equivalence condition. In this paper, small ball method is used to obtain lower bound of smallest singular value of design matrix. In statistical learning theory, small ball assumption is used to lead to lower bound of quadratic component in Equation 2.1, so that it can provide a lower bound of smallest singular value. As we do not need coordinates of input vector X are independent, we need a small ball method without independent. Fortunately, independence assumption is relaxed in [22], and the corresponding definition of small-ball assumption is as follows:

Definition 2.1 (Small Ball Assumption:[22]). The random vector $X \in \mathbb{R}^p$ satisfies a weak small ball assumption(denoted as wSBA) with constants \mathcal{L}, κ if for every $1 \leq k \leq p-1$, every k dimensional subspace F, every $z \in \mathbb{R}^p$,

$$\mathbb{P}\left\{\left\|P_{F}X-z\right\|_{\ell_{2}}\leqslant\kappa\sqrt{k}\right\}\leqslant\left(\mathcal{L}\kappa\right)^{k},$$

where P_F is the orthogonal projection onto the subspace F.

There are many cases when random vector X satisfying wSBA, we refer the reader to Appendix A in [22] for a comprehensive view.

3 Main Result

In this section, we will formulate our main result, Theorem 3.1. Before this, we have to assume our final assumption and define some parameters.

Firstly, We need to following assumption: There are constants $\delta_1>0$ and $\delta_2\geqslant 1$ such that

$$\left(\frac{1}{p}\sum_{i=1}^{p}\left\|\Sigma^{1/2}e_{i}\right\|_{\ell_{2}}^{2+\delta_{1}}\right)^{\frac{1}{2+\delta_{1}}} \leqslant \delta_{2}\sqrt{\frac{\operatorname{tr}(\Sigma)}{p}},\tag{3.1}$$

where $(e_i)_{i=1}^p$ are ONB of \mathbb{R}^p .

This assumption is used to select a proper (in sense of a uniform lower bound of inner product) subset $\sigma_0 \subset [p]$ such that $|\sigma_0| \ge c_0 p$, where c_0 depends on δ_1 and δ_2 . This assumption is not restrictive. See subsection 3.1 for an example.

Secondly, we define three parameters. Define k^\ast as the smallest integer such that

$$p\log\left(1+\frac{\sqrt{d_q(D)\frac{\tilde{\Lambda}(D)}{\sqrt{p}}+\frac{\tilde{\Lambda}^2(D)}{p}+\lambda_1(\Sigma)}\sqrt{\frac{p}{\operatorname{tr}(\Sigma)}}}{\sqrt{3c_0(1-\Re_k(\Sigma))-1}}\right) \leqslant N\frac{c_0\Re_k(\Sigma)+1-c_0}{2} + \log p(3.2)$$

where the minimum of empty set is defined as ∞ . Denote ν as follows:

$$\nu := N \frac{c_0 \Re_{k^*}(\Sigma) + 1 - c_0}{2} + \log p - p \log \left(1 + \frac{\sqrt{d_q(D) \frac{\tilde{\Lambda}(D)}{\sqrt{p}} + \frac{\tilde{\Lambda}^2(D)}{p} + \lambda_1(\Sigma)} \sqrt{\frac{p}{\operatorname{tr}(\Sigma)}}}{\sqrt{3c_0(1 - \Re_{k^*}(\Sigma)) - 1}} \right).(3.3)$$

Parameter k^* is a level which can balance the two sides in Equation 3.2.

Denote

$$\rho = \|\alpha^*\|_{\ell_2} + \sqrt{\frac{2}{3(1-c_0)c_0 - 1}} \sqrt{\frac{p}{\operatorname{tr}(\Sigma)}} \frac{\|\xi\|_{\psi_2}}{\varepsilon}, \qquad (3.4)$$

where ε is a constant. ρ will be upper bound of estimation error. Denote

$$r_{1}^{*} := \underset{r>0}{\operatorname{arginf}} \left\{ \tilde{\Lambda}(H_{r,\rho}) \leqslant \sqrt{\zeta_{1}pr} \right\}, \quad r_{2}^{*} := \underset{r>0}{\operatorname{arginf}} \left\{ d_{q}(\mathcal{F}_{r,\rho}) \leqslant \zeta_{2}r \right\}.$$
$$r^{*} := r_{1}^{*} + r_{2}^{*}, \tag{3.5}$$

where ζ_1, ζ_2 are absolute constants. Particularly, when $H_{r,\rho}$ is a sub-gaussian class, this definition reduces to that of [8]. r^* will be upper bound of prediction risk.

Now we can formulate our main result as follows.

Theorem 3.1. Suppose $X = \Sigma^{1/2} Z \in \mathbb{R}^p$ is a random vector, where Z is an isotropic random vector that satisfies wSBA with constants \mathcal{L}, κ , and Σ satisfies Equation 3.1. If $(X_i)_{i=1}^N$ are i.i.d. copies of X, forming rows of a random matrix **X**. Let $\hat{\alpha}$ be an interpolation solution on $(X_i, Y_i)_{i=1}^N$, where $Y_i = \langle X_i, \alpha^* \rangle + \xi_i$, and $(\xi_i)_{i=1}^N$ are i.i.d. sub-gaussian random variables. Then there exists absolute constant c such that: with probability at least $1 - \exp(-\nu) - \exp(-cN)$,

$$\left\|\hat{\alpha} - \alpha^*\right\|_{\ell_2} \leqslant \rho, \quad \left\|\Sigma^{1/2} \left(\hat{\alpha} - \alpha^*\right)\right\|_{\ell_2} \leqslant r^*,$$

where ρ , ν and r^* are defined in Equation 3.4, 3.3 and 3.5

3.1 Example

Consider a simple example considered in [2], [8] and [28]. When X is a subgaussian random vector, we have $\tilde{\Lambda}(\mathcal{F}) \sim \mathbb{E} ||G||_{\mathcal{F}}$ at once. Therefore, with probability at least $1 - \exp(-cN)$, we have

$$\|\Gamma\| \lesssim \sqrt{N\lambda_1(\Sigma)}.$$

Consider a concrete case that $\varepsilon = o(1)$ such that for any k,

$$\lambda_k(\Sigma) = e^{-k} + \varepsilon$$
, with $\log \frac{1}{\varepsilon} < N$, $p = cN \log \frac{1}{\varepsilon}$.

If $p\varepsilon = \omega(1)$, then $\operatorname{tr}(\Sigma) = O(1)$, and $d_q(D)\frac{\tilde{\Lambda}(\mathcal{F})}{\sqrt{p}} + \frac{\tilde{\Lambda}^2(\mathcal{F})}{p} + \lambda_1(\Sigma) = O(1)$. So

$$\sqrt{\frac{p}{\operatorname{tr}(\Sigma)}}\sqrt{d_q(D)\frac{\tilde{\Lambda}(D)}{\sqrt{p}} + \frac{\tilde{\Lambda}^2(D)}{p} + \lambda_1(\Sigma)} = O(1).$$

To choose k such that Equation 3.2 holds, we have to bound $\mathfrak{R}_k(\Sigma)$ from below. Firstly, we need to lower bound $R_k(\Sigma)$.

$$R_k(\Sigma) = \Theta\left(\frac{\left(e^{-k} + p\varepsilon\right)^2}{e^{-2k} + p\varepsilon^2}\right)$$

by setting $k = \log(1/\varepsilon) < \frac{p}{2}$, then $R_k(\Sigma) = \Theta(p)$.

Then we estimate $\operatorname{srank}_4(\Sigma)$. We have $\operatorname{srank}_4(\Sigma) = \Theta(p)$. So $R_{k,2}(\Sigma) = \Theta(p)$. Further, we have $\mathfrak{R}_k(\Sigma) = \Theta(2^{-p})$. Further, $2^{-p} = \Theta(\varepsilon^{cN})$.

LHS/p =
$$\Theta\left(\log\left(1 + \frac{1}{\sqrt{3c_0(1 - 2^{-p}) - 1}}\right)\right)$$

= $\Theta\left(-\log\left(3c_0(1 - \varepsilon^{cN}) - 1\right)\right)$
= $\Theta\left(-\log\left(1 - \varepsilon^{cN}\right)\right).$

For the right hand side of Equation 3.2, we have

$$\frac{N}{p}\frac{c_0\Re_k(\Sigma) + 1 - c_0}{2} = \Theta\left(\frac{1 - c_0(1 - 2^{-p})}{2c\log\left(1/\varepsilon\right)}\right) = \Theta\left(\varepsilon^{cN}\right),$$

and

$$\frac{\log p}{p} = \Theta\left(\frac{\log c + \log N + \log\log\left(1/\varepsilon\right)}{cN\log\left(1/\varepsilon\right)}\right) = \Theta\left(\frac{\log N}{N}\right)$$

Recall that $\varepsilon = o(1)$, so Equation 3.2 holds for N large enough.

Consider δ_2 such that $\delta_2^2 \ge 1 + 1/\varepsilon$, then Equation 3.1 holds. Therefore, we can set

$$\nu = \frac{\log N}{N} - \varepsilon^{cN} + \log\left(1 - \varepsilon^{cN}\right),$$

Next, we estimate r^* . Since $\mathcal{F}_{H_{r,\rho}}$ is a sub-gaussian class, $r_2^* = 0$, and

$$r_1^* = \operatorname*{arginf}_{r>0} \left\{ \sum_{i=1}^p \min\left\{ r^2, \lambda_i(\Sigma)\rho^2 \right\} \leqslant \zeta_1 p r^2 \right\}.$$

So $r^* = r_1^* \leq \frac{2}{\sqrt{\zeta_1}} \|\alpha^*\|_{\ell_2} \sqrt{\frac{\operatorname{tr}(\Sigma)}{p}}$. Therefore, by Theorem 3.1, with probability at least $1 - \exp(-\nu) - \exp(-cN)$, we have

$$\begin{aligned} \|\hat{\alpha} - \alpha^*\|_{\ell_2} &\leqslant \|\alpha^*\|_{\ell_2} + C \,\|\xi\|_{\psi_2} \,\sqrt{\frac{p}{p\varepsilon+1}} \leqslant c \,\|\alpha^*\|_{\ell_2} \,, \\ \left\|\Sigma^{1/2}(\hat{\alpha} - \alpha^*)\right\|_{\ell_2}^2 &\leqslant \frac{4}{\zeta_1} \,\|\alpha^*\|_{\ell_2}^2 \left(\frac{p\varepsilon+1}{p}\right) = \frac{4}{\zeta_1} \,\|\alpha^*\|_{\ell_2}^2 \left(\varepsilon + \frac{1}{c\log(1/\varepsilon)N}\right), \end{aligned}$$

if signal-to-noise ratio $\|\alpha^*\|_{\ell_2} / \|\xi\|_{\psi_2}$ is greater than $\sqrt{\frac{p}{p\varepsilon+1}}$.

4 Estimation Error

In this section, we are going to obtain a upper bound of $\|\hat{\alpha} - \alpha^*\|_{\ell_2}$ in high probability. We have

$$\hat{\alpha} = \mathbf{X}^{\dagger}\mathbf{Y} = \mathbf{X}^{\dagger}\mathbf{X}\alpha^{*} + \mathbf{X}^{\dagger}\xi.$$

Therefore,

$$\|\hat{\alpha} - \alpha^*\|_{\ell_2} = \| \left(\mathbf{X}^{\dagger} \mathbf{X} - I \right) \alpha^* \|_{\ell_2} + \| \mathbf{X}^{\dagger} \xi \|_{\ell_2} \leq \| \alpha^* \|_{\ell_2} + \| \mathbf{X}^{\dagger} \| \| \xi \|_{\ell_2} \,. \tag{4.1}$$

For $\|\xi\|_{\ell_2}$, we can obtain

$$\left\|\xi\right\|_{\ell_{2}} \leqslant \sqrt{N} \left\|\xi\right\|_{\psi_{2}},$$

with probability at least $1 - \exp(-N)$ by Bernstein's inequality.

To upper bound $\|\mathbf{X}^{\dagger}\|$, we need a lower bound of the smallest singular value of \mathbf{X} in high probability.

Lemma 4.1 (Lower bound of the smallest singular value). Suppose $X = \Sigma^{1/2} Z \in \mathbb{R}^p$ is a random vector, where Z is an isotropic random vector that satisfies wSBA with constants \mathcal{L}, κ , and Σ satisfies Equation 3.1. If $(X_i)_{i=1}^N$ are i.i.d. copies of X, forming rows of a random matrix X. Then there exists constant c_0 such that the smallest singular value of X has lower bound

$$s_{\min}(\mathbf{X}) \ge \varepsilon \sqrt{\frac{3c_0(1-c_0)-1}{2}} \cdot \sqrt{N} \sqrt{\frac{\operatorname{tr}(\Sigma)}{p}} \gtrsim \sqrt{N \frac{\operatorname{tr}(\Sigma)}{p}}, \quad \forall \varepsilon \in (0,1)$$

with probability at least $1 - \exp(-\nu) - \exp(-cN)$, where c is an absolute constant.

With the help of Lemma 4.1, we can arrive at the estimation error:

Theorem 4.1 (Estimation Error). Suppose $X = \Sigma^{1/2}Z$, where Z is a random vector satisfying wSBA with parameters \mathcal{L} , κ and Σ satisfies Equation 3.1. Let $(X_i)_{i=1}^N$ are i.i.d. copies of X. Let $Y_i = \langle X_i, \alpha^* \rangle + \xi_i$, where $(\xi_i)_{i=1}^N$ are i.i.d. sub-gaussian random variables, and let $\mathbf{Y} = (Y_i)_{i=1}^N$. Let \mathbf{X} as random matrix with lines X_i^T , and $\hat{\alpha} = \mathbf{X}^{\dagger}\mathbf{Y}$. For ν defined in Equation 3.3, there exists constant c_0 such that: with probability at least $1 - \exp(-cN) - \exp(-\nu)$,

$$\begin{split} \|\hat{\alpha} - \alpha^*\|_{\ell_2} &\leqslant \quad \|\alpha^*\|_{\ell_2} + \sqrt{\frac{2}{3(1-c_0)c_0 - 1}} \sqrt{\frac{p}{\operatorname{tr}(\Sigma)}} \frac{\|\xi\|_{\psi_2}}{\varepsilon}, \quad \forall \varepsilon \in (0,1) \\ &\leqslant \quad \|\alpha^*\|_{\ell_2} + c \, \|\xi\|_{\psi_2} \, \sqrt{\frac{p}{\operatorname{tr}(\Sigma)}}. \end{split}$$

The proof is trivial by using Equation 4.1 and Lemma 4.1.

Theorem 4.1 can be compared with Theorem 3 in [8]. Their estimation error is related to effective rank $r_k(\Sigma)$, while our bound depends only on $\operatorname{tr}(\Sigma)/p$. This is because our lower bound on the smallest singular value is given by average eigenvalue, instead of effective rank. We believe that by choosing c_0 , the smallest singular value can be controlled in terms of effective rank, though we think deriving such a bound in our work is not necessary.

In the following subsection, we are going to prove Lemma 4.1. An outline of the proof of Lemma 4.1 is as follows. Firstly, we establish a coordinate small-ball estimation in terms of effective rank in Theorem 4.2. Secondly, we prove a uniform lower bound of $\|\mathbf{X}t\|_{\ell_2}$ on an epsilon-Net of S^{p-1} . Finally, we can lower bound the smallest singular value by combining its minimal ℓ_2 norm and its maximal operator norm.

4.1 Coordinate small ball estimates in terms of effective rank

In this subsection, we prove the following Theorem.

Theorem 4.2 (Coordinate Small Ball Estimate in terms of Effective ranks). If random vector $X = \Sigma^{1/2} Z \in \mathbb{R}^p$ satisfies wSBA with constants (\mathcal{L}, κ) , and $(e_i)_{i=1}^p$ are ONB of \mathbb{R}^p that satisfy Equation 3.1, then for $\varepsilon \in (0, 1)$,

$$\mathbb{P}\left\{ \left| \left\{ i \leqslant p : \left| \left\langle \Sigma^{1/2} Z, e_i \right\rangle \right| \geqslant \varepsilon \sqrt{\frac{\operatorname{tr}(\Sigma)}{p}} \right\} \right| \leqslant c_0 p \right\} \lesssim \frac{(4p-k)^2}{8p} \frac{c^{\frac{16p^2}{(4p-k)^2}R_{k,2}(\Sigma)}}{R_{k,2}(\Sigma)}, (4.2)$$

where c depends on \mathcal{L}, κ .

This is just a simple modification of that in [22]. We divide the proof of Theorem 4.2 into three steps.

Firstly, we select a proper subset $\sigma_0 \subset [p]$. This step can be done by using a probabilistic combinatorics technique. Let u_i be a random vector uniformly distributed on the given ONB $\{e_i\}_{i=1}^p$. Set indicators $(1_i)_{i=1}^p$. If $\|\Sigma^{1/2}u_i\|_{\ell_2} \ge \sqrt{\operatorname{tr}(\Sigma)/(2p)}$, then $1_i = 1$, otherwise $1_i = 0$. Then

$$\mathbb{E}\left[\sum_{i=1}^{p} 1_{i}\right] = \sum_{i=1}^{p} \mathbb{P}\left\{\left\|\Sigma^{1/2} u_{i}\right\|_{\ell_{2}} \ge \sqrt{\frac{\operatorname{tr}(\Sigma)}{2p}}\right\}.$$

Then by Equation 3.1 and Paley-Zygmund inequality, see e.g. Lemma 3.1 in [11], we can get its lower bound: RHS $\geq c_0(\delta_1, \delta_2)p$. Therefore, there exists a subset $\sigma_0 \subset [p]$, whose cardinality is at least c_0p , such that for all $i \in \sigma_0$, there exists

$$\left\|\Sigma^{1/2}e_i\right\|_{\ell_2} \ge \sqrt{\frac{\operatorname{tr}(\Sigma)}{2p}}.\tag{4.3}$$

Secondly, [22] decompose $[c_0p]$ into ℓ coordinate blocks by using restricted invertibility Theorem. That is to say,

Lemma 4.2 (Lemma 3.1 in [22]). Assume that for every $1 \leq i \leq p$, Equation 3.1 holds. Set $k_4 = \operatorname{srank}_4(\Sigma^{1/2})$. Then for any $\lambda \in (0, 1)$, there are disjoint subsets $(\sigma_i)_{i=1}^{\ell} \subset [c_0p]$ such that

• For $1 \leq j \leq \ell$, there is $|\sigma_j| \geq c_0^4 k_q/2048$ and $\sum_{j=1}^{\ell} |\sigma_j| \geq c_0 p/2$. • $\left\| \left((\Sigma^{1/2})^* P_{\sigma_j}^* \right)^{-1} \right\|_{S_{\infty}} \leq 4.$

Next, we derive a uniform lower bound of $|\sigma_j|$ by lower bounding srank₄($\Sigma^{1/2}$). Lemma 4.3 (Lower bound of stable rank in terms of effective rank). For $0 \leq k \leq p-1$ and $\Sigma \in \mathbb{R}^{p \times p}$, we have

$$\operatorname{srank}_{4}(\Sigma^{1/2}) \geqslant \frac{16p^{2}}{(4p-k)^{2}} \left(1 - \sqrt{\frac{k}{\operatorname{srank}_{4}(\Sigma)}}\right) R_{k}(\Sigma),$$

Proof. The proof is separated into two parts. Firstly, we lower bound $\left\|\Sigma^{1/2}\right\|_{S_2}$.

By Ky Fan's maximal principle, see e.g. Lemma 8.1.8 in [18] or Chapter 3 in [5], we have

$$\sum_{i=1}^{p-r} s_i^2(\Sigma^{1/2}) \ge \operatorname{tr}\left(\Sigma\left(I_p - P\right)\right),$$

where $I_p - P$ is an orthogonal projection of rank (p - r), which provides a lower bound of sum of largest (p - r) eigenvalues of Σ . Set p - r = k, then rank(P) = p - k. We have $\sum_{i>k} s_i^2(\Sigma^{1/2}) = \|\Sigma^{1/2}\|_{S_2}^2 - \sum_{i=1}^k s_i^2(\Sigma^{1/2})$. It follows that

$$\sum_{i>k} s_i^2(\Sigma^{1/2}) \leqslant \left\| \Sigma^{1/2} \right\|_{S_2}^2 - \operatorname{tr}\left(\Sigma \left(I_p - P \right) \right).$$
(4.4)

We just need to lower bound tr $(\Sigma (I_p - P))$ in terms of $\|\Sigma^{1/2}\|_{S_2}^2$. Consider Σ and ΣP separately, we have the following identity:

$$\left\|P\Sigma^{1/2}\right\|_{S_2}^2 = \operatorname{tr}(\Sigma) - \operatorname{tr}\left(\Sigma\left(I_p - P\right)\right),$$

 \mathbf{SO}

$$\operatorname{tr}\left(\Sigma\left(I_{p}-P\right)\right) = \left\|\Sigma^{1/2}\right\|_{S_{2}}^{2} - \left\|P\Sigma^{1/2}\right\|_{S_{2}}^{2}.$$
(4.5)

Substitute Equation 4.5 into Equation 4.4, then we just need to upper bound $\|P\Sigma^{1/2}\|_{S_2}^2$. However, by definition of $\|\cdot\|_{S_2}$ and property of Frobenius norm, we have

$$\left\|P\Sigma^{1/2}\right\|_{S_2}^2 = \left\|\Sigma^{1/2}\right\|_{S_2}^2 - \left\|P^C\Sigma^{1/2}\right\|_{S_2}^2,$$

where P^C is complement of projector P. Recall that $\operatorname{rank}(P) = p - k$, so we can set P picking p - k rows of $\Sigma^{1/2}$, so P^C picks k rows of $\Sigma^{1/2}$ and it has lower bound $\|\Sigma^{1/2}\|_{S_2}/(2\sqrt{p})$ by Equation 4.3. Therefore, we have

$$\sum_{i>k} s_i^2(\Sigma^{1/2}) \leqslant \left(1 - \frac{k}{4p}\right) \left\|\Sigma^{1/2}\right\|_{S_2}^2$$

and immediately,

$$\sum_{i=1}^{p} \lambda_i(\Sigma) \ge \left(1 - \frac{k}{4p}\right)^{-1} \sum_{i>k} s_i^2(\Sigma^{1/2}) = \left(1 - \frac{k}{4p}\right)^{-1} \sum_{i>k} \lambda_i(\Sigma).$$
(4.6)

Secondly, we upper bound $\left\|\Sigma^{1/2}\right\|_{S_4}$.

By Holder's inequality,

$$\|\Sigma\|_{S_2}^2 = \sum_{i=1}^k s_i^2(\Sigma) + \sum_{i>k} s_i^2(\Sigma) \leqslant \sqrt{k} \, \|\Sigma\|_{S_4}^2 + \sum_{i>k} s_i^2(\Sigma)$$

So we have

$$\sum_{i=1}^{p} \lambda_i^2(\Sigma) = \left\|\Sigma\right\|_{S_2}^2 \leqslant \left(1 - \sqrt{\frac{k}{\operatorname{srank}_4(\Sigma)}}\right)^{-1} \sum_{i>k} \lambda_i^2(\Sigma),$$
(4.7)

by definition of stable rank.

Combining Equation 4.7 and Equation 4.6, Lemma 4.3 is proved. \Box

The rest of the proof is based on the following Lemma:

Lemma 4.4 ([22]). If random vector X satisfies the wSBA with constant (\mathcal{L}, κ) , and $(e_i)_{i=1}^p$ are ONB of \mathbb{R}^p , and $\Sigma^{1/2} : \mathbb{R}^p \to \mathbb{R}^p$ satisfies Equation 3.1, then for $\varepsilon \in (0, 1)$, we have

$$\mathbb{P}\left\{ \left| \left\{ i \leqslant p : \left| \left\langle \Sigma^{1/2} Z, e_i \right\rangle \right| \geqslant \varepsilon \sqrt{\frac{\operatorname{tr}(\Sigma)}{p}} \right\} \right| \leqslant c_0 p \right\} \leqslant \sum_{j \leqslant \ell} \left(\frac{e\mathcal{L}\varepsilon}{c_0/2} \right)^{|\sigma_j|/(c_0/2)},$$

where σ_j , c_0 , ℓ are the same as Lemma 4.2.

This Lemma is not explicitly given in [22].

Using this, we can prove Theorem 4.2:

Proof of Theorem 4.2. By Lemma 4.4, Lemma 4.3 and Lemma 4.2, Theorem 4.2 is proved easily.

4.2 Lower bound of smallest singular value

In this subsection, we proceed step 2 and 3. Build an epsilon-Net on S^{p-1} , obtain uniform lower bound of smallest singular value on it and extend it on the whole S^{p-1} .

Proof of Lemma 4.1. Fix a random vector $X_j \in \mathbb{R}^p$. Consider p unit vectors $(e_i)_{i=1}^p$ which forming an ONB of \mathbb{R}^p , then by Theorem 4.2, with probability at least $1 - \mathfrak{R}_k(\Sigma)$, there exists a subset σ_j with cardinality at least $c_0 p$, and the vectors in it satisfy that for all $e_i \in \sigma_j$,

$$\left|\left\langle \Sigma^{1/2} Z_j, e_i \right\rangle\right| \ge \varepsilon \sqrt{\frac{\operatorname{tr}(\Sigma)}{p}}.$$
(4.8)

Each X_j has such a subset $\sigma_j \subset (e_i)_{i=1}^p$ of cardinality at least $c_0 p$ with probability at least $1 - \Re_k(\Sigma)$. Pick a $t \in (e_i)_{i=1}^p$ randomly. If $t \in \sigma_j$, then Equation 4.8 holds, that is to say, we can have a lower bound on the inner product at this time.

Denote 1_j as $1_{\{t \notin \sigma_j\}}$, then $\mathbb{E}1_j \leq 1-c_0(1-\Re_k(\Sigma))$. By Bernstein's inequality, with probability at least $1 - \exp(-\min\{t^2, t\}N)$,

$$\sum_{j=1}^{N} 1_j \leqslant N \mathbb{E} 1_j + t \leqslant \frac{3}{2} N(c_0 \Re_k(\Sigma) + 1 - c_0).$$

by setting $t = (c_0 \Re_k(\Sigma) + 1 - c_0)/2$, then with probability at least

$$1 - \exp(-N(c_0 \Re_k(\Sigma) + 1 - c_0)/2),$$

we have $\sum_{j=1}^{N} 1_j \leq 3N(c_0 \Re_k(\Sigma) + 1 - c_0)/2$, that is to say,

$$\sqrt{\sum_{j=1}^{N} \left\langle \Sigma^{1/2} Z_j, u \right\rangle^2} \ge \varepsilon \sqrt{\frac{3c_0(1 - \Re_k(\Sigma)) - 1}{2}N} \sqrt{\frac{\operatorname{tr}(\Sigma)}{p}}, \quad \forall u \in \sigma_j.$$

Build an η -Net Γ_{ε} on B_2^p .

Set

$$\eta = \frac{\sqrt{\operatorname{tr}(\Sigma)}}{2\sqrt{2}} \varepsilon \sqrt{3c_0(1 - \Re_k(\Sigma)) - 1} \sqrt{\frac{N}{p}}$$

By $\log |\Gamma_{\eta}| \leq p \log (1 + 2/\eta)$, we have

$$\log |\Gamma_{\eta}| \leq p \log \left(1 + \frac{4\sqrt{2}}{\varepsilon} \frac{\|\Gamma\|}{\sqrt{\operatorname{tr}(\Sigma)}} \sqrt{\frac{p}{N}} \cdot \frac{1}{\sqrt{3c_0(1 - \Re_k(\Sigma)) - 1}}\right)$$

We just need to ensure

$$p\log\left(1+\frac{\|\Gamma\|}{\sqrt{\operatorname{tr}(\Sigma)}}\sqrt{\frac{p}{N}}\cdot\frac{c}{\sqrt{3c_0(1-\Re_k(\Sigma))-1}}\right)\leqslant N\frac{c_0\Re_k(\Sigma)+1-c_0}{2}+\log p(4.9)$$

by choosing k wisely. Here we just need to repeat the selecting process no more than $\lceil |\Gamma_{\eta}| / (c_0 p) \rceil$ times, so to make Equation 4.8 holds uniformly for all elements in Γ_{η} , we have to pay a log $(\lceil |\Gamma_{\eta}| / (c_0 p) \rceil)$ in exponential term $\exp(-N(c_0 \Re_k(\Sigma) + 1 - c_0)/2)$ and make sure this probability is no greater than 1.

For upper bound of $\|\Gamma\|$, we use Lemma 2.2. With probability at least $1 - \exp(-c_0 N)$,

$$\|\Gamma\| = \sqrt{\max_{t \in S^{p-1}} \sum_{i=1}^{N} \langle \Gamma_{\cdot,i}, t \rangle_{\ell_2}^2} \leq \sqrt{N} \sqrt{C\left(d\frac{\tilde{\Lambda}(D)}{\sqrt{k_D}} + \frac{\tilde{\Lambda}^2(D)}{k_D}\right) + \lambda_1(\Sigma)}.$$

By choosing $k = k^*$ defined in Equation 3.2, the probability can be lower bounded by $1 - \exp(-\nu)$, where ν is defined in Equation 3.3. In summary, with probability at least $1 - \exp(-c_0 N) - \exp(-\nu)$, we have

$$s_{\min}(\mathbf{X}) \ge \varepsilon \sqrt{\frac{3c_0(1-c_0)-1}{8}N} \sqrt{\frac{\operatorname{tr}(\Sigma)}{p}}$$

5 Prediction error

In this section, we obtain an upper bound of prediction error based on upper bound of estimation risk by using localization method introduced in subsetcion 2.1.

Theorem 5.1 (Prediction error). If random vector $X = \Sigma^{1/2} Z \in \mathbb{R}^p$ satisfies wSBA with constants (\mathcal{L}, κ) , and Σ that satisfy Equation 3.1. Then with probability at least $1 - \exp(-\nu) - 2\exp(-cN)$, prediction error satisfies

$$\left\|\Sigma^{1/2}(\hat{\alpha}-\alpha^*)\right\|_{\ell_2}\leqslant r^*,$$

where c is an absolute constant.

The proof is a kind of localization argument. That is to say, we are going to prove $\hat{\alpha} - \alpha^*$ lies in a localized area with respect to $\|\cdot\|_{\Sigma}$. Firstly, we need a localization Lemma from [8]:

Lemma 5.1 (Localization: Lemma 3 in [8]). With probability at least $1 - \exp(-N/16)$, we have $P_N \mathcal{L}_{\hat{\alpha}} \leq - \|\xi\|_{\psi_2}^2/2$. Moreover, for any r, let $\Omega_{r,\rho}$ denote the following event

$$\Omega_{r,\rho} = \left\{ \alpha \in \mathbb{R}^p : \alpha - \alpha^* \in B(\rho) \setminus B_{\Sigma}(r), \text{ and } \mathbb{P}_N \mathcal{L}_\alpha > -\frac{1}{2} \left\| \xi \right\|_{\psi_2}^2 \right\}.$$

On the event

$$\Omega_{r,\delta} \cap \{\hat{\alpha} - \alpha^* \in B(\rho)\} \cap \left\{ \mathbb{P}_N \mathcal{L}_{\hat{\alpha}} \leqslant -\frac{1}{2} \left\| \xi \right\|_{\psi_2}^2 \right\},\tag{5.1}$$

prediction risk has upper bound r, that is to say,

$$\left\|\Sigma^{1/2}\left(\hat{\alpha}-\alpha^*\right)\right\|_{\ell_2}\leqslant r.$$

Lemma 5.1 reduces upper bound of prediction risk to Equation 5.1. Probability of event $\{\hat{\alpha} - \alpha^* \in B(\rho)\}$ can be lower bounded by estimation error, see Lemma 4.1. Recall that $H_{r,\rho} = B(\rho) \cap B_{\Sigma}(r)$, we just need to prove that event

$$\inf_{\alpha:\alpha-\alpha^*\in H_{r,\rho}} \mathbb{P}_N \mathcal{L}_{\alpha} > -\frac{1}{2} \left\|\xi\right\|_{\psi_2}^2$$

holds with high probability for $r > r^*$.

In this section, we are going to find lower bound of $P_N \mathcal{L}_{\alpha}$ in terms of upper bounds of quadratic and multiplier processes according to Equation 2.2.

Firstly, we find upper bound of quadratic process. By Lemma 2.2, with probability at least $1 - \exp(-cN)$, we have

$$Q_{r,\rho} \lesssim_q \left(d_q(H_{r,\rho}) \frac{\tilde{\Lambda}(H_{r,\rho})}{\sqrt{p}} + \frac{\tilde{\Lambda}^2(H_{r,\rho})}{p} \right).$$
(5.2)

Secondly, we need upper bound of multiplier component. This can be done by using Lemma 2.1. That is to say, with probability at least $1 - 2\exp(-cN)$,

$$M_{r,\rho} \lesssim_q \|\xi\|_{\psi_2} \frac{\hat{\Lambda}(H_{r,\rho})}{\sqrt{p}}.$$
(5.3)

since ξ is centered and independent with X. Therefore, when $r > r^*$, we have

$$d_q(H_{r,\rho}) \leqslant \zeta_1 r^*, \quad \frac{\tilde{\Lambda}(H_{r,\rho})}{\sqrt{p}} \leqslant \zeta_2 r^*.$$

Proof of Theorem 5.1. Let $\alpha \in \alpha^* + H_{r,\rho}$. Recall that $r = \left\| \Sigma^{1/2}(\alpha - \alpha^*) \right\|_{\ell_2}$. If $r > r^*$, by substituting Equation 5.3 and Equation 5.2 into Equation 2.2, it follows that

$$\inf_{\alpha \in \alpha^* + H_{r,\rho}} \mathbb{P}_N \mathcal{L}_{\alpha} > (r^*)^2 \left(\theta^{-2} - \zeta_1 \zeta_2 \theta^{-2} - \zeta_2 \theta^{-2} - \zeta_2 \right) - \frac{1}{2} \left\| \xi \right\|_{\psi_2}^2$$

Set ζ_1, ζ_2 small enough, RHS > $-\frac{1}{2} \|\xi\|_{\psi_2}^2$. By Lemma 5.1, Theorem 5.1 is proved.

6 Discussion

In this section, we discuss two aspects. Firstly, we discuss why it is so difficult to investigate benign overfitting beyond linear model. Secondly, we discuss a benign overfitting case without truncated effective rank.

6.1 Why linear model?

In this subsection, we imagine statistical model \mathcal{F} is the affine hull of sub-classes $(\mathcal{F}_j)_{j=1}^d$, that is to say, for all $f \in \mathcal{F}$, there exists f_j in each \mathcal{F}_j and $\alpha_j \in \mathbb{R}$ such that $f = \sum_{j=1}^d \alpha_j f_j$. Denote $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{R}^d$. Of course this is not the problem that we deal with in this paper, but considering such a general case like this would be benefit to understand the role of α and the difficulty to generalize benign overfitting beyond linear model.

Even in this kind of simple "additive model" case, benign overfitting is much more difficult. Firstly, \hat{f} interpolates $(X_i, Y_i)_{i=1}^N$, but \hat{f}_j need not interpolates them. In fact, they may differs a lot, see $\hat{f}(x) = -x + x = \hat{f}_1 + \hat{f}_2$ can interpolate (10,0) but $\hat{f}_1(10) = -\hat{f}_2(10)$. It is a difficult task to derive oracle inequality by studying \mathcal{F}_j .

If we minimizing $\|\alpha\|_{\ell_2}$ analogs to linear case, the minimization of $\|\alpha\|_{\ell_2}$ given interpolation condition $\sum_{j=1}^{p} \alpha_j f(X_i) = Y_i$ for all $i = 1, 2, \dots, N$ can be solved by Moore-Penrose inverse.

Condition on $(f_j)_{j=1}^p$. Denote matrix Γ as

$$\Gamma = \begin{bmatrix} f_1(X_1) & f_2(X_1) & \cdots & f_p(X_1) \\ f_1(X_2) & f_2(X_2) & \cdots & f_p(X_2) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(X_N) & \cdots & \cdots & f_p(X_N). \end{bmatrix}_{N \times p}$$

Denote \mathbf{f}^* as $(f^*(X_1), f^*(X_2), \dots, f^*(X_N))$ and ξ as $(\xi_1, \xi_2, \dots, \xi_N)$. Then the interpolation condition is equivalent to

minimizing
$$\|\alpha\|_{\ell_2}$$
, s.t. $\Gamma \alpha = Y$.

We assume α satisfying interpolation condition always exists. Using Moore-Penrose inverse, we have

$$\hat{\alpha} = \Gamma^{\dagger} Y = \Gamma^{\dagger} \mathbf{f}^* + \Gamma^{\dagger} \xi.$$

Therefore, to establish upper bound of $\|\hat{\alpha}\|_{\ell_2}$, we need a lower bound of the smallest singular value of Γ .

However, as we see in Lemma 4.1, the smallest singular value increases when N increases, causing $\|\Gamma^{\dagger}\mathbf{f}^{*}\|_{\ell_{2}}$ decreasing. This phenomenon is called "signal blood" in [24], which means that the influence caused by signal \mathbf{f}^{*} will decline so that minimizing $\|\hat{\alpha}\|_{\ell_{2}}$ cannot reflect properties true signal unless there are some unrealistic restrictions.

Therefore, \mathbf{f}^* should balance Γ^{\dagger} when N increase to avoid signal blood. This can be done by linear regression, where $\Gamma = \mathbf{X}$. This illustrates that why we choose linear model.

6.2 Benign overfitting without truncated effective rank

Now, we try to establish benign overfitting without truncated effective rank, but on stable rank $r_0(\Sigma)$, see Equation 1.1. Recall that a linear model on $T \subset \mathbb{R}^p$ is $\mathcal{F}_T = \{\langle \cdot, t \rangle : t \in T\}$. Let $\sigma = (X_1, \cdots, X_N)$, then the projection of \mathcal{F}_T by using σ is indeed a random linear transformation of T. That is to say, $P_{\sigma}\mathcal{F}_T = \mathbf{X}T$, where $P_{\sigma}(f_t) = (\langle X_i, t \rangle)_{i=1}^N$. We need lower bound of smallest singular value of \mathbf{X} to derive an upper bound of estimation error, and a lower bound of quadratic component in Equation 2.1. Fortunately, this can be done by Dvoretzky-Milman Theorem, see [1] or [19]. Dvoretzky-Milman Theorem can hold with rather heavy-tailed random vectors, but for the sake of simplicity, we assume $(g_i)_{i=1}^N$ are i.i.d. gaussian random vectors in \mathbb{R}^p .

Lemma 6.1 (Dovoretzky-Milman(one-side)). There exists absolute constants c_1, c_2 such that: If $0 < \delta < \frac{1}{2}$, and

$$N \leqslant c_1 \frac{\delta^2}{\log\left(1/\delta\right)} r_0(\Sigma),\tag{6.1}$$

and $\Gamma = \sum_{i=1}^{N} \langle g_i, \cdot \rangle e_i$, where $(e_i)_{i=1}^{N}$ are ONB of \mathbb{R}^N . Then with probability at least $1 - 2\exp(-c_2 r_0(\Sigma)\delta^4 / \log(1/\delta))$,

$$(1-\delta)\sqrt{\operatorname{tr}(\Sigma)}B_2^N \subset \Gamma\left(\Sigma^{1/2}B_2^p\right)$$

Take $\delta = 1/4$ for example, we have $4\text{tr}(\Sigma)B_2^N \subset \Gamma(\Sigma^{1/2}B_2^p)$, so

$$s_{\min}(\Gamma) = \sqrt{\min_{t \in S^{p-1}} \langle g_i, t \rangle^2} \ge 4\sqrt{\operatorname{tr}(\Sigma)}$$

holds with probability at least $1 - 2\exp(-cr_0(\Sigma))$. Therefore, with probability at least $1 - 2\exp(-c_1r_0(\Sigma)) - 2\exp(-c_2N)$,

$$\|\hat{\alpha} - \alpha^*\|_{\ell_2} \leq \|\alpha^*\|_{\ell_2} + \|\xi\|_{\psi_2} \sqrt{\frac{N}{\operatorname{tr}(\Sigma)}}.$$

As for the prediction risk, we have: when $r > r_1^*$,

$$\inf_{\alpha \in \alpha^* + H_{r,\rho}} \mathcal{P}_N \mathcal{L}_{\alpha} \ge r^2 \left(\frac{16 \operatorname{tr}(\Sigma)}{N} - \frac{1}{2} \zeta_1 \right) - \frac{1}{2} \left\| \xi \right\|_{\psi_2}^2 > -\frac{1}{2} \left\| \xi \right\|_{\psi_2}^2.$$

From here on, the proof is as the same as that of Theorem 5.1, the details are omitted.

Note that $r_0(\Sigma) \leq p$, and $p = cN \log(1/\varepsilon)$, so we can set c_1, δ wisely to adapt to the example discussed in subsection 3.1.

In summary, although interpolation learning suffers from estimating both noise ξ and sign α^* , it still generalize well if the smallest singular value of **X** is large enough such that it can absorb the level of noise, $\sqrt{N} \|\xi\|_{\psi_2}$, see Equation 4.1. The smallest singular value is used to weaken influence of noise. To make the smallest singular value large enough, the number of samples should satisfy an upper bound that depends on covariance of the input vector. This threshold is used to balance the rate of exponential decay(acquired by concentration or small-ball argument) and metric entropy (given by net argument). Therefore, this threshold depends on dimension p, sample size N and covariance Σ . If we fix relationship between p and N(like the example in subsection 3.1), we need Σ has a large trace(or at least heavy tail of eigenvalues), which is the key to benign overfitting. Note that in this interpretation, there is no restrictions on concentration properties of input vector X, but its small-ball property, that is to say, X should fully spread on its margin. It is its spreading that can absorb noise ξ . It is this that make minimum ℓ_2 linear interpolant fit into heavy-tailed case. Finally, we believe that our result could be easily modified to "Informative-Outlier" framework, cf. [7], to obtain a result in a "robust flavor" both for CS community and statistics community.

References

 Artstein-Avidan, S., Giannopoulos, A. and Milman, V.D. (2015) Asymptotic geometric analysis. Part I. American Mathematical Society, Providence.MR3331351.

- [2] Bartlett, P.L., Long, P. M., Lugosi, G. and Tsigler, A. (2019) Benign Overfitting in Linear Regression. Proceedings of the National Academy of Sciences Apr 2020, 201907378.
- [3] Belkin, M., Ma, S. and Mandal, S. (2018) To understand deep learning we need to understand kernel learning. Proceedings of the the 35th International Conference on Machine Learning (ICML 2018).
- [4] Belkin, M., Rakhlin, A. and Tsybakov, A.B. (2019) Does data interpolation contradict statistical optimality? AISTAT 2019.
- [5] Bhatia, R. (1997) Matrix Analysis. Springer-Verlag, New York.MR1477662
- [6] Boucheron, S.,Lugosi, G. and Massart, P. (2013) Concentration inequalities: A nonasymptotic theory of independence. 1nd ed.Oxford university press.MR3185193
- G. Chinot, G. Lecué and M. Lerasle (2020) Statistical Learning with Lipschitz and convex loss functions, Probability Theory and Related Fields, 176, 897– 940.MR4087486
- [8] Chinot, G., Lerasle, M. (2020) Benign overfitting in the large deviation regime. arXiv preprint arXiv:2003.05838.
- [9] Dirksen, S. (2015) Tail bounds via generic chaining. *Electronic Journal of* Probability 20.MR3354613
- [10] Hastie, T., et al. (2019) Surprises in high-dimensional ridgeless least squares interpolation. arXiv preprint arXiv:1903.08560.
- [11] Kallenberg, O. (2002) Foundations of modern probability. 2nd ed. Springer-Verlag, New York.MR1876169
- [12] Koltchinskii, V. and Lounici, K. (2017) Concentration inequalities and moment bounds for sample covariance operators. *Bernoulli* 23, 110-133.MR3556768
- [13] Koltchinskii, V. and Mendelson, S. (2015) Bounding the smallest singular value of a random matrix without concentration. Int. Math. Res. Not. IMRN 23, 12991–13008.MR3431642
- [14] Liang, T., Rakhlin, A. (2020) Just Interpolate: Kernel "Ridgeless" Regression Can Generalize. Annals of Statistics.MR4124325
- [15] Liang, T., Rakhlin, A. and Zhai, X. (2020) On the Multiple Descent of Minimum-Norm Interpolants and Restricted Lower Isometry of Kernels. *Conference on Learning Theory (COLT), 2020.*
- [16] Rakhlin, A. and Zhai, X. (2019) Consistency of Interpolation with Laplace Kernels is a High-Dimensional Phenomenon. *Conference on Learning Theory* (COLT), 2019.

- [17] Song, M. and Montanari, A. (2019) The generalization error of random features regression: Precise asymptotics and double descent curve. *Submitted* to Communications on Pure and Applied Mathematics.
- [18] Størmer, E. (2013) Positive Linear Maps of Operator Algebras. Springer Monographs in Mathematics.MR3012443
- [19] Mendelson, S. (2016a) Dvoretzky type Theorems for subgaussian coordinate projections. J. Theoret. Probab., 29, 1644–1660.MR3571258
- [20] Mendelson, S. (2016b) Learning without concentration for general loss functions. Probability Theory and Related Fields, 171, 459–502.MR3800838
- [21] Mendelson, S. (2016c) Upper bounds on product and multiplier empirical processes. Stochastic Processes and their Applications, 126, 3652– 3680.MR3565471
- [22] Mendelson, S. and Paouris, G. (2019) Stable recovery and the coordinate small-ball behaviour of random vectors. arXiv preprint arXiv:1904.08532.
- [23] Meyer Carl D. (2000) Matrix analysis and applied linear algebra. Society for Industrial and Applied Mathematics (SIAM), 71.MR1777382
- [24] Muthukumar, V., Vodrahalli, K., Subramanian, V. and Sahai, A. (2020) Harmless interpolation of noisy data in regression. *IEEE Journal on Selected Areas in Information Theory.*
- [25] Naor, A. and Youssef, P. (2017) Restricted invertibility revisited. Springer Cham. A journey through discrete mathematics, 657–691.MR3726618
- [26] Rudelson, M. and Vershynin, R. (2007) Sampling from large matrices: an approach through geometric functional analysis. *Journal of the ACM (2007)*, Art. 21, 19 pp.MR2351844
- [27] Talagrand, M. (2014) Upper and lower bounds for stochastic processes: modern methods and classical problems. Springer Science & Business Media.MR3184689
- [28] Tsigler, A. and Bartlett, P. (2020) Benign overfitting in ridge regression.arXiv preprint arXiv:2009.14286.
- [29] Vaart, Aad W and Wellner, Jon A (1996) Weak convergence and empirical processes: with applications to statistics. Springer Series in Statistics.MR1385671
- [30] Vershynin, R. (2018) High-Dimensional Probability: An Introduction with Applications in Data Science. Cambridge University Press:New York.MR3837109
- [31] Zhang, C. et al. (2016) Understanding deep learning requires rethinking generalization. arXiv preprint arXiv:1611.03530.