

Sequential single pion production explaining the dibaryon “ $d^*(2380)$ ” peak

R. Molina,^{1,*} Natsumi Ikeno,^{1,2,†} and Eulogio Oset^{1,‡}

¹*Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC, Institutos de Investigación de Paterna, Aptdo. 22085, 46071 Valencia, Spain*

²*Department of Agricultural, Life and Environmental Sciences, Tottori University, Tottori 680-8551, Japan*

We study the two step sequential one pion production mechanism, $np(I=0) \rightarrow \pi^- pp$, followed by the fusion reaction $pp \rightarrow \pi^+ d$, in order to describe the $np \rightarrow \pi^+ \pi^- d$ reaction with $\pi^+ \pi^-$ in $I=0$, where a narrow peak, so far identified with a “ $d(2380)$ ” dibaryon, has been observed. We find that the second step $pp \rightarrow \pi^+ d$ is driven by a triangle singularity that determines the position of the peak of the reaction and the large strength of the cross section. The combined cross section of these two mechanisms produce a narrow peak with the position, width and strength compatible with the experimental observation within the approximations done. This novel interpretation of the peak without invoking a dibaryon explains why the peak is not observed in other reactions where it has been searched for.

PACS numbers:

The $np \rightarrow \pi^0 \pi^0 d$ reaction exhibits a sharp peak around 2370 MeV with a narrow width of about 70 MeV, which is also seen in the $pp \rightarrow \pi^+ \pi^- d$ reaction with approximately double strength [1–3]. In the absence of a conventional reaction mechanism that can explain these peaks, they have been interpreted as a signal of a dibaryon that has been named $d^*(2380)$. On the base of this hypothesis several other features observed in π production experiments and NN phase shifts have been interpreted (see [4] for a recent review). Actually, the narrow peak in $np \rightarrow \pi \pi d$ affects the inelasticity of the NN phase shifts and should have repercussion in NN phase shifts as emphasized in [5, 6]. Several mechanisms of two pion production leading to $\pi \pi d$ have been studied in [1–3] based upon the model of [7] for $NN \rightarrow NN\pi\pi$, which contain double Δ production, with subsequent $\Delta \rightarrow \pi N$ decay or $N^*(1440)$ production with decay of N^* to $N\pi\pi$, or $N^* \rightarrow \pi \Delta(N\pi)$. In all these cases the resulting np particles are fused into the deuteron. The results of these calculations give rise to cross sections with small strength compared to the peak of the $np \rightarrow \pi^0 \pi^0 d$ reaction and no peak at the energy of the observed one. Such conclusions were already drawn in an early paper [8] and we have explicitly recalculated the cross sections from these mechanisms reconfirming all these earlier findings. Interestingly, in the same work [8] a peak with poor statistics, already visible for the $np \rightarrow \pi^+ \pi^- d$ reaction was explained from a different mechanism, two step sequential π production, $np \rightarrow pp\pi^-$ followed by $pp \rightarrow \pi^+ d$. The cross section for $np \rightarrow \pi^+ \pi^- d$ was evaluated factorizing cross sections for the two latter reactions in an “on-shell” approach that called for further checks concerning its accuracy. Such mechanism has no further been invoked concerning the new improved data on the $np \rightarrow \pi^+ \pi^- d$,




FIG. 1: Two step mechanism for $np \rightarrow \pi^+ \pi^- d$ suggested in [8] with explicit Δ excitation in the $pp \rightarrow \pi^+ d$ last step as found in [9–11]. The mechanism with the nn intermediate state is considered in addition.

$np \rightarrow \pi^0 \pi^0 d$ reactions [1–3].

On the other hand, the time reversal reaction of $pp \rightarrow \pi^+ d$, π^+ absorption in the deuteron, $\pi^+ d \rightarrow pp$, was the subject of study in the past [9–11] and it was shown to have a neat peak corresponding to the Δ excitation. Combining the work of [9–11] with the idea of [8] on the $np \rightarrow \pi^+ \pi^- d$ reaction, the mechanism for $np \rightarrow \pi^+ \pi^- d$ can be expressed diagrammatically as in Fig. 1.

After many years, more refined data and new theoretical developments make most opportune to revise this issue along the same idea. We can quote:

- 1) The data on $np \rightarrow \pi^+ \pi^- d$ and $np \rightarrow \pi^0 \pi^0 d$ have nowadays excellent precision [1–4].
- 2) The $np \rightarrow \pi^0 \pi^0 d$ reaction has $\pi^0 \pi^0$ in isospin $I=0$, and hence the initial np state must also be in $I=0$.

The work of [3] splits the $np \rightarrow \pi^+ \pi^- d$ reaction into $I=0$ and $I=1$, and, as expected, the same peak visible in the $np \rightarrow \pi^0 \pi^0 d$ reaction is seen in the $np(I=0) \rightarrow \pi^+ \pi^- d$ reaction with about double strength. This means that in the $np \rightarrow \pi^- pp$ reaction, the first step of the sequential single pion production mechanism, the initial np state is also in $I=0$. It has only been recently that the first step in Fig. 1, $np \rightarrow \pi^- pp$ with np in $I=0$ has been singled out with relatively good precision [12] (see revision about normalization in [12, 13])

*Electronic address: Raquel.Molina@ific.uv.es

†Electronic address: ikeno@tottori-u.ac.jp

‡Electronic address: Eulogio.Oset@ific.uv.es

3) New developments about triangle singularities [14] allow us to identify the large strength of the $pp \rightarrow \pi^+d$ reaction with the presence of a triangle singularity in the triangle diagram shown in the last part of Fig. 1. This corresponds to having simultaneously the Δ and the two nucleons on shell and collinear. The simplification of the formalism on the triangle singularities done in [15] allows us to see immediately where the peak of the $pp \rightarrow \pi^+d$ cross section should appear, using Eq. (18) of [15] with the d mass slightly unbound to find a solution of that equation. One predicts that a peak of the cross section should appear around $M_{\text{inv}}(pp) \sim 2179$ MeV, very close to where the peak appears in the experiment [16]. The Coleman-Norton theorem [17] is enlightening in this case to visualize the process. It states that a triangle singularity appears when the process visualized in the triangle diagram can occur at the classical level. In our case this would be: the pp system produces ΔN back to back in the pp rest frame; the Δ decays into a π^+ in the direction of the Δ and N' in opposite direction, which is the direction of N . The N' moves faster than N (encoded in Eq. (18) of [15]) and catches up with N to fuse into the deuteron. The fusion of the two nucleons into the deuteron comes out naturally when the mechanism discussed has a triangle singularity, giving rise to a neat peak and a cross section rather large compared with typical fusion reactions [18]. We have done a recalculation of the $pp \rightarrow \pi^+d$ reaction from this new perspective [19], but the details are unnecessary in the derivation done here for the $np \rightarrow \pi^+\pi^-d$ cross section which, as in [8], relies on experimental cross sections, using the new $np(I=0) \rightarrow \pi^-pp$ cross section [12, 13] and the data for $pp \rightarrow \pi^+d$ [16]. We also improve on the on shell approach used in [8]. It is also easy to see that the mechanisms of [7] for $NN \rightarrow NN\pi\pi$ (see Fig. 1 of [7]), which do not contain the sequential pion production and reproduce well the data, do not develop a triangle singularity upon fusion of a final np pair. It is also worth mentioning that while the mechanism for $pp \rightarrow \pi^+d$ in [9–11] was not identified as a triangle singularity, it was shown in [10] that the cross section was blowing up when the Δ width was set to zero, a characteristic of the triangle singularity.

The derivation of the $np \rightarrow \pi^+\pi^-d$ cross section that we do follows the steps of the derivation of the optical theorem [20]. We call t the amplitude for isoscalar $np(I=0) \rightarrow \pi^-pp$, t' for $pp \rightarrow \pi^+d$ and t'' for $np(I=0) \rightarrow \pi^+\pi^-d$. The differential cross section for the isoscalar $np(I=0) \rightarrow \pi^-pp$ reaction is given by

$$\frac{d\sigma_{np \rightarrow \pi^-pp}^I}{dM_{\text{inv}}(p_1p'_1)} = \frac{1}{4ps} (2M_N)^4 \frac{1}{16\pi^3} p_\pi \tilde{p}_1 |\bar{t}|^2 \frac{1}{2} \quad (1)$$

where σ^I stands for the isoscalar cross section, \sqrt{s} is

the center-of-mass (CM) energy of the initial np state, $M_{\text{inv}}(p_1p'_1)$ the invariant mass of the final two protons in this reaction, p the CM momentum of the initial n or p particles, p_π the pion momentum in the np rest frame and \tilde{p}_1 the momentum of the final protons in the pp rest frame. We use the $(2M_N)^4$ factor of fermion field normalization for the nucleons following the formalism of Mandl and Shaw [21]. The magnitude $|\bar{t}|^2$ stands for the angle averaged $|t|^2$ and the factor $\frac{1}{2}$ takes into account the identity of the two final protons.

Similarly, the cross section for $pp \rightarrow \pi^+d$ in the second part of the diagram of Fig. 1 is given by

$$\sigma_{pp \rightarrow \pi^+d} = \frac{1}{16\pi M_{\text{inv}}^2(p_1p'_1)} \frac{p'_\pi}{\tilde{p}_1} |\bar{t}'|^2 (2M_N)^2 (2M_d) \quad (2)$$

where p'_π is the π^+ momentum in the pp rest frame and $|\bar{t}'|^2$ stands for the angle averaged $|t'|^2$. We choose to normalize the deuteron field as the nucleons and add the factor $2M_d$ (it disappears from the final formulas). On the other hand the amplitude for the $np \rightarrow \pi^-\pi^+d$ process in Fig. 1 is given by

$$\begin{aligned} -it'' &= \frac{1}{2} \int \frac{d^4p_1}{(2\pi)^4} \frac{(2M_N)^2}{2E_N(p_1)2E_N(p'_1)} \frac{i}{p_1^0 - E_N(p_1) + i\epsilon} \\ &\times \frac{i}{\sqrt{s} - p_1^0 - \omega_\pi - E_N(p'_1) + i\epsilon} (-i)t(-i)t' \quad (3) \end{aligned}$$

The factor $\frac{1}{2}$ is to account for the intermediate propagator of two identical particles. In the d^4p_1 integrations t and t' would be off shell. In Ref. [8] the pion and the two protons of the intermediate state were taken on shell and t and t' were used with the on shell variables. Theoretical advances done after [8] allow us to go beyond this approximation. Indeed, the chiral unitary approach of [22] for meson-meson interaction, or [23] for meson-baryon interaction, factorizes the vertices on-shell and performs the loop integral of the two intermediate states. A different justification is given in [24] writing a dispersion relation for the inverse of the hadron-hadron scattering amplitude, and it also finds a justification in [25, 26] showing with chiral lagrangians that off shell parts of the amplitudes appearing in the approach get cancelled with counterterms provided with the same theory. This means that in Eq. (3) we can take tt' outside the dp_1^0 integration with their on-shell values and evaluate the remaining of the integral of Eq. (3).

Performing the p_1^0 integration analytically with Cauchy's residues we get

$$\begin{aligned} t'' &= \frac{1}{2} \int \frac{d^3p_1}{(2\pi)^3} \frac{(2M_N)^2}{2E_N(p_1)2E_N(p'_1)} \\ &\times \frac{tt'}{\sqrt{s} - E_N(p_1) - E_N(p'_1) - \omega_\pi + i\epsilon} \quad (4) \end{aligned}$$

where \vec{p}_1, \vec{p}'_1 are the momenta of the intermediate pp particles in Fig. 1, and ω_π the π^- energy. The t, t' amplitudes are Lorentz invariant and we choose to evaluate the

$\int \frac{d^3 p_1}{2E_1(p_1)}$ integral in the pp rest frame, where $|\vec{p}'_1| = |\vec{p}_1|$ and $\sqrt{s} - \omega_\pi$ becomes the invariant mass of the two protons. This integral is logarithmically divergent and requires regularization. The result depends smoothly on a cut off $p_{1,\max}$ for $|\vec{p}_1|$ that we use to regularize the $d^3 p_1$ integration, and we shall take some values for $p_{1,\max}$ in a reasonable range. Yet, we anticipate that the on shell part given by Eq. (5), below, gives the largest contribution to the t'' amplitude. Since $\tilde{p}_1 = 552$ MeV/c for $M_{inv}(p_1 p'_1) = 2179$ MeV, where the triangle singularity would appear for t' for a Δ with zero width, or a pronounced peak when the width is considered, values of $p_{1,\max}$ around 700 – 800 MeV seem reasonable.

The on-shell approximation used in [8] that allows one to write the cross section for $np \rightarrow \pi^+ \pi^- d$ in terms of the $np(I=0) \rightarrow \pi^- pp$ and $pp \rightarrow \pi^+ d$ ones is obtained in the present formalism by taking the imaginary part of the two nucleon propagator

$$\frac{1}{M_{inv}(p_1 p'_1) - 2E_N(p_1) + i\epsilon} \equiv \mathcal{P} \left[\frac{1}{M_{inv}(p_1 p'_1) - 2E_N(p_1)} \right] - i\pi \delta(M_{inv}(p_1 p'_1) - 2E_N(p_1)) \quad (5)$$

We have then

$$t''_{\text{on}} = -i \frac{1}{2} \frac{\tilde{p}_1}{8\pi} \frac{(2M_N)^2}{M_{inv}(p_1 p'_1)} t\bar{t}' \quad (6)$$

where we have factorized the angle averaged value of tt' , $t\bar{t}'$. Using the analogous equation of Eq. (1) for $d\sigma_{np \rightarrow \pi^+ \pi^- d}/dM_{inv}(\pi^+ \pi^-)$, the on-shell approximation of Eq. (6) and Eq. (1) we can already write

$$\frac{d\sigma_{np \rightarrow \pi^+ \pi^- d}}{dM_{inv}(\pi^+ \pi^-)} = (2M_N)^2 (2M_d) p_d \tilde{p}_\pi \frac{1}{4} \frac{\tilde{p}_1^2}{64\pi^2} \times \frac{1}{M_{inv}^2(p_1 p'_1)} \frac{1}{p_\pi \tilde{p}_1} 2|\vec{t}'|^2 \frac{d\sigma_{np \rightarrow \pi^- pp}}{dM_{inv}(p_1 p'_1)} \quad (7)$$

where p_d is the deuteron momentum in the original np rest frame, $|\vec{t}'|^2$ the angle averaged $|t'|^2$, and \tilde{p}_π the pion momentum in the $\pi^+ \pi^-$ rest frame. In Eq. (7) we have assumed that $|t\bar{t}'|^2 = |\vec{t}|^2 |\vec{t}'|^2$. The amplitudes t, t' in Refs. [12, 16] have some angular structure, but these are smooth enough to make this assumption a sensible approximation.

Next we use physical arguments to write the $np \rightarrow \pi^+ \pi^- d$ cross section with an easy compact formula. We note that $\pi^0 \pi^0$ or $\pi^+ \pi^-$ in $I=0$, as we discussed earlier, imply that the two pions are in s -wave and both $\pi^0 \pi^0$ or the symmetrized ($\pi^+ \pi^- + \pi^- \pi^+$) behave as identical particles, which reverts into a Bose enhancement when the two pions go together. Certainly if they are exactly together we shall also have the phase space factor \tilde{p}_π in $d\sigma/dM_{inv}(\pi^+ \pi^-)$ of Eq. (7) which makes null this distribution in the two pion threshold, but some enhancement

for small invariant masses is expected. Our argumentation is supported by the results of [1, 2] for $\pi^0 \pi^0$ (see Fig. 2 of [1] and Fig. 4 of [2]) and also in [3] for charged pions, although the nature of $I=0$ and $I=1$ in this case distorts a bit the mass distribution compared to the clean $I=0$ $\pi^0 \pi^0$ case.

We could take some $M_{inv}(\pi^+ \pi^-)$ distribution as input, but to make the results as model independent as possible we take the $\bar{M}_{inv}(\pi^+ \pi^-) \sim 2m_\pi + 60$ MeV, not far from threshold but we change it to see how the results depend on \bar{M}_{inv} . The stability of the results that we find by changing the value of $\bar{M}_{inv}(\pi\pi)$ justifies this approximation a posteriori. Then, we can write

$$\frac{d\sigma_{np \rightarrow \pi^+ \pi^- d}}{dM_{inv}(\pi^+ \pi^-)} = \sigma_{np \rightarrow \pi^+ \pi^- d} \delta(M_{inv}(\pi^+ \pi^-) - \bar{M}_{\pi\pi}) \quad (8)$$

The approximation of Eq. (8) is sufficiently good and allows us to get a more transparent picture of what is the reason for the appearance of the peak in the $np \rightarrow \pi^+ \pi^- d$ reaction. Note now that the energy of the two pions is obtained as

$$E_{2\pi} = \frac{s + M_{inv}^2(\pi\pi) - M_d^2}{2\sqrt{s}} \quad (9)$$

and since the two pions go relatively together, we take $E_\pi = E_{2\pi}/2$, which allows to relate $M_{inv}(p_1 p'_1)$ with \sqrt{s} via

$$M_{inv}^2(p_1 p'_1) = (P(np) - p_{\pi^-})^2 = s + m_\pi^2 - 2\sqrt{s} E_\pi \quad (10)$$

and formally

$$2M_{inv}(p_1 p'_1) dM_{inv}(p_1 p'_1) = -2\sqrt{s} dE_\pi = -M_{inv}(\pi\pi) dM_{inv}(\pi\pi) \quad (11)$$

Using this relationship we can integrate Eq. (8) with respect to $M_{inv}(\pi\pi)$ and using Eqs. (2) and (7) we obtain,

$$\sigma_{np \rightarrow \pi^+ \pi^- p} = \frac{M_{inv}(p_1 p'_1)}{4\pi} \frac{\sigma_{np \rightarrow \pi^- pp} \sigma_{pp \rightarrow \pi^+ d}}{M_{inv}(\pi\pi)} \frac{\tilde{p}_1^2}{p_\pi p'_\pi} p_d \tilde{p}_\pi \quad (12)$$

One last detail is needed. We have considered the two step $np(I=0) \rightarrow \pi^- pp$ followed by $pp \rightarrow \pi^+ d$. A properly symmetrized t'' amplitude requires the addition of $np(I=0) \rightarrow \pi^+ nn$ followed by $nn \rightarrow \pi^- d$. It is trivial to see considering isospin that the amplitudes $np(I=0) \rightarrow \pi^- pp$ and $np(I=0) \rightarrow \pi^+ nn$ are identical up to the phase of π^+ (-1 in our formalism) and the same happens for $pp \rightarrow \pi^+ d$ and $nn \rightarrow \pi^- d$ for the same configuration of the particles. Hence, the product of the amplitudes is the same. In the case that the π^+ and π^- go exactly together, the two amplitudes will be identical and add coherently, but we saw that the phase space factor \tilde{p}_π of Eq. (12) kills this contributions. When considering the integration over the five degrees of freedom of the three body phase space the terms are expected to sum mostly incoherently and we must multiply by 2 Eq. (12). Similar arguments can be done with respect to the

spin sums and averages. The study of the $pp \rightarrow \pi^+d$ reaction in [19] indicates that there is a certain angular dependence on the different spin transitions and we should expect an incoherent sum over spins. Then by including also in $|\bar{t}|^2$ the average over initial spins and sum over final spins we would be considering in our formula the average over spins of the initial np and the sum over spins of the deuteron, plus the intermediate sum over the pp and nn spins.

Eq. (12) still relies on the on-shell approximation of Eq. (6). To take into account the off shell effects discussed above we replace

$$\left(\frac{\tilde{p}_1}{2\pi M_{\text{inv}}(p_1 p'_1)}\right)^2 \rightarrow |G(M_{\text{inv}})|^2 \quad (13)$$

with G the loop function with two protons

$$G = \int \frac{d^3 p_1}{(2\pi)^3} \frac{1}{E_N(p_1)E_N(p_1)} \frac{1}{M_{\text{inv}}(p_1 p'_1) - 2E_N(p_1) + i\epsilon}$$

The last step in the evaluation of $\sigma_{np \rightarrow \pi^+ \pi^- d}$ requires to use the experimental data for $np(I=0) \rightarrow \pi^- pp$ and $pp \rightarrow \pi^+ d$. We get $\sigma_{pp \rightarrow \pi^+ d}$ directly from experiment [16]. The $\sigma_{np \rightarrow \pi^- pp}$ in $I=0$ requires some thoughts. In [27, 28] the isoscalar $NN \rightarrow \pi NN$ amplitude is obtained via isospin symmetry from $\sigma_{np \rightarrow pp\pi^-}$ and $\sigma_{pp \rightarrow pp\pi^0}$, and relatively precise results are obtained in [12] from improved measurements of these cross sections. In the erratum of [12] and in [13] it is clarified that the actual $\sigma_{pn(I=0) \rightarrow NN\pi}$ is one half of $\sigma_{NN(I=0) \rightarrow NN\pi}$ of [12]. The cross section that we need is $\sigma_{pn(I=0) \rightarrow pp\pi^-}$. It is trivial to see using isospin symmetry that $\sigma_{pn(I=0) \rightarrow pp\pi^-}$, $\sigma_{pn(I=0) \rightarrow nn\pi^+}$ and $(\sigma_{pn(I=0) \rightarrow pn\pi^0} + \sigma_{pn(I=0) \rightarrow np\pi^0})$ are all equal. Then we write the relationship of [12, 27, 28] as

$$\begin{aligned} \sigma_{np(I=0) \rightarrow pp\pi^-} &= \frac{1}{3} \sigma_{np(I=0) \rightarrow NN\pi} \\ &= \frac{1}{6} \sigma_{NN(I=0) \rightarrow NN\pi} \\ &= \frac{1}{6} 3(2\sigma_{np \rightarrow pp\pi^-} - \sigma_{pp \rightarrow pp\pi^0}) \end{aligned} \quad (14)$$

and we take data for $\sigma_{np(I=0) \rightarrow NN\pi}$ from Fig. 1 of [13]. Statistical and some systematic errors are considered in [12, 13]. With the only purpose of making a realistic fit to the data we include also systematic errors from the uncertainty in Eq. (14) when using isospin symmetry. We assume a typical 5% violation of isospin in each of the last two terms of Eq. (14) and sum the errors in quadrature. The systematic errors obtained are of the order of 0.5 mb in $\sigma_{np(I=0) \rightarrow NN\pi}$, which we also add in quadrature to the former ones of [13]. With those errors we have many good fits with reduced χ^2 , (χ_r^2), smaller than 1. We take two of them, one peaking on the lower side of \sqrt{s} and the other one on the upper side for the $np(I=0) \rightarrow NN\pi$ cross section, parameterized as,

$$\sigma_i = \left| \frac{\alpha_i}{\sqrt{s} - \tilde{M}_i + i\frac{\tilde{\Gamma}_i}{2}} \right|^2 \quad (15)$$

TABLE I: Values of the peak strength (“strength”), peak position (“position”), and width, for intermediate particles on shell (columns with $\delta\tilde{M}_{\pi\pi}$, $\tilde{M}_{\pi\pi} = 2m_\pi + \delta\tilde{M}_{\pi\pi}$), and off-shell (“o.s.”), where we have taken $p_{1,\text{max}} = 700$ and 800 MeV and $\delta\tilde{M}_{\pi\pi} = 60$ MeV.

	$\delta\tilde{M}_{\pi\pi}$ (MeV)			$p_{1,\text{max}}^{\text{o.s.}}$ (MeV)	
Set I	40	60	80	700	800
strength (mb)	0.72	0.76	0.75	0.82	0.95
position (MeV)	2332	2332	2332	2332	2332
width (MeV)	76	76	81	75	75
Set II					
strength (mb)	0.75	0.80	0.80	0.85	0.96
position (MeV)	2342	2345	2345	2343	2342
width (MeV)	86	87	88	87	84

and call set I the one with the parameters: $\tilde{M}_1 = 2326$ MeV, $\tilde{\Gamma}_1 = 70$ MeV, $\alpha_1^2 = 2.6 \left(\frac{\tilde{\Gamma}_1}{2}\right)^2$ mb MeV² ($\chi_r^2 = 0.5$), while set II has $\tilde{M}_2 = 2335$ MeV, $\tilde{\Gamma}_2 = 80$ MeV, $\alpha_2^2 = 2.5 \left(\frac{\tilde{\Gamma}_2}{2}\right)^2$ mb MeV² ($\chi_r^2 = 0.55$). The $pp \rightarrow \pi^+ d$ cross section has accurate data and we parameterize it as

$$\sigma_3 = \left| \frac{\alpha_3}{M_{\text{inv}}(p_1 p'_1) - \tilde{M}_3 + i\frac{\tilde{\Gamma}_3}{2}} \right|^2 \quad (16)$$

with $\tilde{M}_3 = 2165$ MeV, $\tilde{\Gamma}_3 = 123.27$ MeV, $\alpha_3^2 = 3.186 \left(\frac{\tilde{\Gamma}_3}{2}\right)^2$ mb MeV².

With the former discussions our final formula on shell is given by

$$\sigma_{np \rightarrow \pi^+ \pi^- p} = \frac{M_{\text{inv}}(p_1 p'_1)}{6\pi} \frac{\sigma_{np \rightarrow NN\pi}^I \sigma_{pp \rightarrow \pi^+ d}}{M_{\text{inv}}(\pi\pi)} \frac{\tilde{p}_1^2}{p_\pi p'_\pi} p_d \tilde{p}_\pi \quad (17)$$

with $\sigma_{np \rightarrow NN\pi}^I = \sigma_{np(I=0) \rightarrow NN\pi}$ of [12, 13] and we show the results in Fig. 2.

We can see that the cross section of $pp \rightarrow \pi^+ d$ and $np(I=0) \rightarrow NN\pi$ overlap around the middle of their energy distributions such that their product in Eq. (17) gives rise to a narrow peak around $\sqrt{s} = 2340$ MeV, close to the position of the experimental $np \rightarrow \pi^+ \pi^- d$ peak around 2365 MeV.

In table I we show the results obtained with set I and set II for the strength of $\sigma_{np \rightarrow \pi^+ \pi^- d}$ at the peak, the peak position and the width of the peak, varying $\tilde{M}_{\pi\pi}$, and $p_{1,\text{max}}$ for the off shell calculations. What one sees is a stability of the results upon changes of $\tilde{M}_{\pi\pi}$, which justifies the use of Eq. (8). We also find that off shell effects




FIG. 2: Plots of $\sigma_{np \rightarrow \pi^- pp}$ ($I = 0$) and $\sigma_{pp \rightarrow \pi^+ d}$, as a function of \sqrt{s} and $M_{\text{inv}}(p_1 p'_1)$, respectively, where $M_{\text{inv}}(p_1 p'_1)$ is evaluated by means of Eq. (10). The results with $\sigma_{np \rightarrow \pi^+ \pi^- d}$ in $I = 0$ of Eq. (17) are multiplied by 10 for a better comparison. Left: Results with set I; Right: Results for set II. $\bar{M}_{\pi\pi} = 2m_\pi + 60$ MeV. Inset: $\sigma_{pp \rightarrow \pi^+ d}$ as a function of $M_{\text{inv}}(p_1 p'_1)$. Data for $pp \rightarrow \pi^+ d$ from [16]. Data for $np(I = 0) \rightarrow \pi^- pp$ are taken from Dakhno et al. [27], and WASA-at-COSY(*) [12, 13], including systematic errors from isospin violation.

using Eq. (13) are small, justifying the on shell approximation used in [8]. The strength at the peak between 0.72–0.96 mb should be considered quite good compared to the experimental one around 0.5 mb, given the different approximations done (fits to the $np(I = 0) \rightarrow \pi^- pp$ cross section with the systematic errors with 20–30% smaller strength at the peak are still acceptable, hence such uncertainties in the resulting $np \rightarrow \pi^+ \pi^- d$ cross section are expected). The peak position from 2332–2345 MeV should also be considered rather good compared to the about 2365 MeV of the experiment [1–3, 12]. The narrow width observed in the experiment of 70–75 MeV is also well reproduced by our results in the range of [75–88] MeV.

The appearance of the peak about 25 MeV below the experimental one is not significant with the perspective that, as discussed in [12], the authors achieve a resolution in \sqrt{s} of about 20 MeV and the $pp \rightarrow pp\pi^0$ and $pn \rightarrow pp\pi^-$ cross sections, from where $\sigma_{np(I=0) \rightarrow pp\pi^-}$ is obtained via Eq. (14) with large cancellations, are measured using data bins of 50 MeV in T_p .

The derivation done contains the basic dynamical ingredients in a skilled way, making some approximations to rely upon experimental cross sections. We think that it is remarkable that a narrow peak, at about the right position, with strength and width comparable to the experimental peak of $np \rightarrow \pi^+ \pi^- d$, appears in spite of the approximations done, and the stability of the results allows us to conclude that a peak with the properties of the experimental one associated so far to the “ $d^*(2380)$ ” dibaryon is unavoidable from the mechanism that we have studied.

From the perspective of the $np \rightarrow \pi^+ \pi^- d$ reaction being tied to the particular reaction mechanism of Fig. 1, with a two step sequential one pion production, it is easy

to understand why the narrow peak of the $np \rightarrow \pi^+ \pi^- d$ reaction is not seen in $\gamma d \rightarrow \pi^+ \pi^- d$ in spite of having the same final state [29]. The first reaction is a fusion reaction, with the last step tied to a triangle singularity. The $\gamma d \rightarrow \pi^+ \pi^- d$ reaction is a coherent reaction, the d is already present in the initial state and the reaction mechanisms are drastically different.

In summary, we have identified the reaction mechanism that produces a narrow peak in the $np \rightarrow \pi^+ \pi^- d$ cross sections without having to invoke a “dibaryon” resonance. From this perspective it is also easy to understand why the peak is not seen in other reactions where it has been searched for, although the peak contributing to the inelastic channels of $pn \rightarrow \text{all}$ can show traces in NN phase shifts, as anticipated in [5, 6] and discussed in [4].

I. ACKNOWLEDGMENTS

R. M. acknowledges support from the CIDEAGENT program with Ref. CIDEAGENT/2019/015 and from the spanish national grant PID2019-106080GB-C21. The work of N. I. was partly supported by JSPS Overseas Research Fellowships and JSPS KAKENHI Grant Number JP19K14709. This work is also partly supported by the Spanish Ministerio de Economía y Competitividad and European FEDER funds under Contracts No. FIS2017-84038-C2-1-P B and No. FIS2017-84038-C2-2-P B. This project has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No. 824093 for the STRONG-2020 project.

-
- [1] M. Bashkanov et al., Phys. Rev. Lett. **102**, 052301 (2009), 0806.4942.
- [2] P. Adlarson et al. (WASA-at-COSY), Phys. Rev. Lett. **106**, 242302 (2011), 1104.0123.
- [3] P. Adlarson et al. (WASA-at-COSY), Phys. Lett. B **721**, 229 (2013), 1212.2881.
- [4] H. Clement and T. Skorodko, Chin. Phys. C **45**, 022001 (2021), 2008.07200.
- [5] G. Faldt and C. Wilkin, Phys. Lett. B **701**, 619 (2011), 1105.4142.
- [6] M. Albaladejo and E. Oset, Phys. Rev. **C88**, 014006 (2013), 1304.7698.
- [7] L. Alvarez-Ruso, E. Oset, and E. Hernandez, Nucl. Phys. A **633**, 519 (1998), nucl-th/9706046.
- [8] I. Bar-Nir et al., Nucl. Phys. B **54**, 17 (1973).
- [9] D. O. Riska, M. Brack, and W. Weise, Phys. Lett. **61B**, 41 (1976).
- [10] A. M. Green and J. A. Niskanen, Nucl. Phys. A **271**, 503 (1976).
- [11] M. Brack, D. O. Riska, and W. Weise, Nucl. Phys. A **287**, 425 (1977).
- [12] P. Adlarson et al. (WASA-at-COSY), Phys. Lett. B **774**, 599 (2017), [Erratum: Phys.Lett.B 806, 135555 (2020)], 1702.07212.
- [13] H. Clement and T. Skorodko (2020), 2010.09217.
- [14] L. D. Landau, Nucl. Phys. **13**, 181 (1960), [Sov. Phys. JETP10,no.1,45(1959); Zh. Eksp. Teor. Fiz.37,no.1,62(1959)].
- [15] M. Bayar, F. Aceti, F.-K. Guo, and E. Oset, Phys. Rev. D **94**, 074039 (2016), 1609.04133.
- [16] C. Richard-Serre, W. Hirt, D. F. Measday, E. G. Michaelis, M. J. M. Saltmarsh, and P. Skarek, Nucl. Phys. B **20**, 413 (1970).
- [17] S. Coleman and R. E. Norton, Nuovo Cim. **38**, 438 (1965).
- [18] P. W. F. Alons, R. D. Bent, J. S. Conte, and M. Dillig, Nucl. Phys. A **480**, 413 (1988).
- [19] N. Ikeno, R. Molina, and E. Oset, To be published (2021).
- [20] C. Itzykson and J. B. Zuber, Quantum Field Theory, International Series In Pure and Applied Physics (McGraw-Hill, New York, 1980), ISBN 978-0-486-44568-7.
- [21] F. Mandl and G. Shaw, QUANTUM FIELD THEORY (1985).
- [22] J. A. Oller and E. Oset, Nucl. Phys. A **620**, 438 (1997), [Erratum: Nucl.Phys.A 652, 407–409 (1999)], hep-ph/9702314.
- [23] E. Oset and A. Ramos, Nucl. Phys. A **635**, 99 (1998), nucl-th/9711022.
- [24] J. A. Oller and U. G. Meissner, Phys. Lett. B **500**, 263 (2001), hep-ph/0011146.
- [25] A. Martinez Torres, K. P. Khemchandani, and E. Oset, Phys. Rev. **C77**, 042203 (2008), 0706.2330.
- [26] A. Martinez Torres, K. P. Khemchandani, L. S. Geng, M. Napsuciale, and E. Oset, Phys. Rev. D **78**, 074031 (2008), 0801.3635.
- [27] L. G. Dakhno et al., Phys. Lett. **114B**, 409 (1982).
- [28] J. Bystricky, J.Phys. France **48**, 1901 (1987).
- [29] M. S. Guenther (A2), PoS **Hadron2017**, 051 (2018).