

Stochastic variational principles for the collisional Vlasov-Maxwell and Vlasov-Poisson equations

Tomasz M. Tyranowski*^{1,2}

¹Max-Planck-Institut für Plasmaphysik

Boltzmannstraße 2, 85748 Garching, Germany

²Technische Universität München, Zentrum Mathematik

Boltzmannstraße 3, 85748 Garching, Germany

Abstract

In this work we recast the collisional Vlasov-Maxwell and Vlasov-Poisson equations as systems of coupled stochastic and partial differential equations, and we derive stochastic variational principles which underlie such reformulations. We also propose a stochastic particle method for the collisional Vlasov-Maxwell equations and provide a variational characterization of it, which can be used as a basis for a further development of stochastic structure-preserving particle-in-cell integrators.

1 Introduction

The collisional Vlasov equation

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} f + \frac{q}{m} (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_{\mathbf{v}} f = C[f] \quad (1.1)$$

describes the time evolution of the particle density function $f = f(\mathbf{x}, \mathbf{v}, t)$ of plasma consisting of charged particles of charge q and mass m which undergo collisions described by the collision operator $C[f]$, and are subject to the electric $\mathbf{E} = \mathbf{E}(\mathbf{x}, t)$ and magnetic $\mathbf{B} = \mathbf{B}(\mathbf{x}, t)$ fields. The vectors $\mathbf{x} = (x^1, x^2, x^3)$ and $\mathbf{v} = (v^1, v^2, v^3)$ denote positions and velocities, respectively. For simplicity, we restrict ourselves to one-specie plasmas. Usually, the particle density function is normalized, so that the total number of particles is $N_{tot} = \iint f(\mathbf{x}, \mathbf{v}, t) d^3 \mathbf{v} d^3 \mathbf{x}$. However, in this work we would like to treat f as a probability density function, and therefore we will use the normalization $\iint f(\mathbf{x}, \mathbf{v}, t) d^3 \mathbf{v} d^3 \mathbf{x} = 1$ instead. A self-consistent model of plasma is obtained by coupling (1.1) with the Maxwell equations

*tomasz.tyranowski@ipp.mpg.de

$$\nabla_x \cdot \mathbf{E} = \rho, \quad (1.2a)$$

$$\nabla_x \cdot \mathbf{B} = 0, \quad (1.2b)$$

$$\nabla_x \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \quad (1.2c)$$

$$\nabla_x \times \mathbf{B} = \frac{\partial \mathbf{E}}{\partial t} + \mathbf{J}, \quad (1.2d)$$

where

$$\rho(\mathbf{x}, t) = qN_{tot} \int_{\mathbb{R}^3} f(\mathbf{x}, \mathbf{v}, t) d^3\mathbf{v}, \quad \mathbf{J}(\mathbf{x}, t) = qN_{tot} \int_{\mathbb{R}^3} \mathbf{v} f(\mathbf{x}, \mathbf{v}, t) d^3\mathbf{v}, \quad (1.3)$$

denote the charge density and the electric current density, respectively, and the factor N_{tot} is due to our normalization. The system (1.1)-(1.3) is usually referred to as the Vlasov-Maxwell equations. It will also be convenient to express the electric and magnetic fields in terms of the scalar $\varphi(\mathbf{x}, t)$ and vector $\mathbf{A}(\mathbf{x}, t)$ potentials

$$\mathbf{E} = -\nabla_x \varphi - \frac{\partial \mathbf{A}}{\partial t}, \quad (1.4a)$$

$$\mathbf{B} = \nabla_x \times \mathbf{A}, \quad (1.4b)$$

as is typical in electrodynamics. The Vlasov-Poisson equations are an approximation of the Vlasov-Maxwell equations in the nonrelativistic zero-magnetic field limit (see Section 6). The main goal of this work is to provide a variational characterization of the Vlasov-Maxwell and Vlasov-Poisson equations via a stochastic Lagrange-d'Alembert type of a principle.

Variational principles have proved extremely useful in the study of nonlinear evolution partial differential equations (PDEs). For instance, they often provide physical insights into the problem being considered; facilitate discovery of conserved quantities by relating them to symmetries via Noether's theorem; allow one to determine approximate solutions to PDEs by minimizing the action functional over a class of test functions (see, e.g., [22]); and provide a way to construct a class of numerical methods called variational integrators (see [71], [72]). A variational principle for the collisionless Vlasov-Maxwell equations was first proposed in [69]. It has been used to derive various particle discretizations of the Vlasov-Maxwell and Vlasov-Poisson equations (see [33], [67], [68], [88], [92]), including structure-preserving variational particle-in-cell methods ([91], [106], [107]). It has also been applied to gyrokinetic theory (see, e.g., [9], [95]). For other formulations and extensions see also [108].

A structure-preserving description of collisional effects is far less developed. A metriplectic framework for the Vlasov-Maxwell-Landau equations has been presented in [43] and [57]. More recently, a stochastic variational principle has been proposed in [59] to describe collisional effects for the Vlasov equation with a fixed external electric field. To the best of our knowledge, to date no variational principle has been derived for the collisional Vlasov-Maxwell and Vlasov-Poisson equations. In this work we extend the notion of the stochastic Lagrange-d'Alembert principle presented in [59] to plasmas evolving in self-consistent electromagnetic fields. The main idea of our approach is to interpret the Vlasov equation (1.1) as a Fokker-Planck equation and consider the associated stochastic differential equations.

The idea of using stochastic differential equations to model collisions has been pursued by a number of authors over the last few decades; see, e.g., [2], [8], [16], [21], [28], [35], [37], [42], [54], [59], [62], [70], [74], [89], [90], [110].

There has been an ever growing body of literature dedicated to stochastic variational principles in recent years. Stochastic variational principles allow the introduction of noise into systems in such a way that the resulting probabilistic models retain all or some of the geometric properties of their deterministic counterparts. For this reason stochastic variational principles have been considered in the context of Lagrangian and Hamiltonian mechanics ([7], [10], [11], [12], [49], [59], [61], [101]), soliton dynamics ([47], [48]), fluid dynamics ([3], [19], [23], [24], [25], [39], [45], [46]), and kinetic plasma theory ([59]).

Main content The main content of the remainder of this paper is, as follows.

In Section 2 we recast the collisional Vlasov-Maxwell equations as a system of coupled stochastic and partial differential equations.

In Section 3 we discuss the relationship between particle methods and stochastic modeling. We formulate a stochastic particle discretization for the collisional Vlasov-Maxwell equations and cast it in a form that allows the derivation of a variational principle.

In Section 4 we describe the variational structure underlying the stochastic particle discretization of the Vlasov-Maxwell system. The main result of this section is Theorem 4.2, in which a stochastic Lagrange-d'Alembert principle for the particle discretization is proved.

In Section 5 we generalize the ideas from Section 4 to the original undiscretized equations. The main result of this section is Theorem 5.1, in which a stochastic Lagrange-d'Alembert principle is proved for a class of the collisional Vlasov-Maxwell equations.

In Section 6 we prove a stochastic Lagrange-d'Alembert principle applicable to the Vlasov-Poisson equations. The main result of this section is Theorem 6.1.

Section 7 contains the summary of our work.

2 The Vlasov-Maxwell-Fokker-Planck equations

2.1 Stochastic reformulation

Various collision models and various forms of the collision operator $C[f]$ are considered in the plasma physics literature (see, e.g., [17], [73]). A key step towards a stochastic variational principle is a probabilistic interpretation of the Vlasov equation (1.1). Therefore, in this work we will be interested only in those collision operators for which (1.1) takes the form of a linear or strongly nonlinear Fokker-Planck equation (see, e.g., [36], [38], [85]). Namely, we will assume that the collision operator can be expressed as

$$C[f] = \frac{1}{2} \sum_{i,j=1}^3 \frac{\partial^2}{\partial v^i \partial v^j} [D_{ij}(\mathbf{x}, \mathbf{v}; f) f] - \sum_{i=1}^3 \frac{\partial}{\partial v^i} [K_i(\mathbf{x}, \mathbf{v}; f) f], \quad (2.1)$$

for some symmetric positive semi-definite matrix $D_{ij}(\mathbf{x}, \mathbf{v}; f)$ and vector $K_i(\mathbf{x}, \mathbf{v}; f)$ functions, where the dependence of D_{ij} and K_i on f may in general be nonlinear, and may involve differential and integral forms of f . In that case (1.1) is an integro-differential equation, the so-called strongly nonlinear Fokker-Planck equation (see [36]). In case D_{ij} and K_i are independent of f , that is, $D_{ij}(\mathbf{x}, \mathbf{v}; f) = D_{ij}(\mathbf{x}, \mathbf{v})$ and $K_i(\mathbf{x}, \mathbf{v}; f) = K_i(\mathbf{x}, \mathbf{v})$, the Vlasov equation (1.1) reduces to the standard linear Fokker-Planck equation. We will further assume that D_{ij} and K_i can be expressed in the form

$$D_{ij}(\mathbf{x}, \mathbf{v}; f) = \sum_{\nu=1}^M g_\nu^i g_\nu^j, \quad K_i(\mathbf{x}, \mathbf{v}; f) = G^i + \frac{1}{2} \sum_{\nu=1}^M \sum_{j=1}^3 \frac{\partial g_\nu^i}{\partial v^j} g_\nu^j, \quad (2.2)$$

for a vector function $\mathbf{G}(\mathbf{x}, \mathbf{v}; f)$, and a family of vector functions $\mathbf{g}_\nu(\mathbf{x}, \mathbf{v}; f)$ with $\nu = 1, \dots, M$. Note that given a symmetric positive semi-definite matrix D_{ij} , a decomposition (2.2) can always be found, but it may not be unique. For instance, one may take $M = 3$ and assume that $g_\nu^i = g_i^\nu$ for $i, \nu = 1, 2, 3$. Then the first equation in (2.2) implies that the family of functions g_ν^i can be determined by calculating the square root of the matrix D_{ij} , and the second equation in (2.2) can be used to calculate the function \mathbf{G} . If (1.1) has the form of a Fokker-Planck equation, then the particle density function f can be interpreted as the probability density function for a stochastic process $(\mathbf{X}(t), \mathbf{V}(t)) \in \mathbb{R}^3 \times \mathbb{R}^3$. This stochastic process then satisfies the Stratonovich stochastic differential equation (see [36], [38], [55], [85])

$$d\mathbf{X} = \mathbf{V} dt, \quad (2.3a)$$

$$d\mathbf{V} = \left(\frac{q}{m} \mathbf{E}(\mathbf{X}, t) + \frac{q}{m} \mathbf{V} \times \mathbf{B}(\mathbf{X}, t) + \mathbf{G}(\mathbf{X}, \mathbf{V}; f) \right) dt + \sum_{\nu=1}^M \mathbf{g}_\nu(\mathbf{X}, \mathbf{V}; f) \circ dW^\nu(t), \quad (2.3b)$$

where $W^1(t), \dots, W^M(t)$ denote the components of the standard M -dimensional Wiener process, and \circ denotes Stratonovich integration. Note that the terms \mathbf{G} and \mathbf{g}_ν can be interpreted as external forces, and that in their absence the equations (2.3) reduce to the equations of motion of a charged particle in an electromagnetic field. We will therefore refer to \mathbf{G} and \mathbf{g}_ν as forcing terms. The electric and magnetic fields are coupled via the Maxwell equations (1.2). It should also be noted that unless (1.1) is linear, the right-hand side of (2.3) depends on f . In order to obtain a self-consistent system, one can express f in terms of the stochastic processes \mathbf{X} and \mathbf{V} as $f(\mathbf{x}, \mathbf{v}, t) = \mathbb{E}[\delta(\mathbf{x} - \mathbf{X}(t))\delta(\mathbf{v} - \mathbf{V}(t))]$, where \mathbb{E} denotes the expected value, and δ is Dirac's delta. This can be further plugged into (1.3). Together, we get

$$f(\mathbf{x}, \mathbf{v}, t) = \mathbb{E}[\delta(\mathbf{x} - \mathbf{X}(t))\delta(\mathbf{v} - \mathbf{V}(t))], \quad (2.4a)$$

$$\rho(\mathbf{x}, t) = qN_{tot}\mathbb{E}[\delta(\mathbf{x} - \mathbf{X}(t))], \quad (2.4b)$$

$$\mathbf{J}(\mathbf{x}, t) = qN_{tot}\mathbb{E}[\mathbf{V}(t)\delta(\mathbf{x} - \mathbf{X}(t))]. \quad (2.4c)$$

Equations (1.2), (2.3), and (2.4) form a self-consistent system of stochastic and partial differential equations whose solutions are the stochastic processes $\mathbf{X}(t)$, $\mathbf{V}(t)$, and the functions $\mathbf{E}(\mathbf{x}, t)$, $\mathbf{B}(\mathbf{x}, t)$.

Remark. Upon substituting (2.4a), the forcing terms \mathbf{G} and \mathbf{g}_ν become functionals of the processes \mathbf{X} and \mathbf{V} , that is, $\mathbf{G}(\mathbf{x}, \mathbf{v}; f) = \mathbf{G}(\mathbf{x}, \mathbf{v}; \mathbf{X}, \mathbf{V})$ and $\mathbf{g}_\nu(\mathbf{x}, \mathbf{v}; f) = \mathbf{g}_\nu(\mathbf{x}, \mathbf{v}; \mathbf{X}, \mathbf{V})$. However, for convenience and simplicity, throughout this work we will stick to the notation $\mathbf{G}(\mathbf{x}, \mathbf{v}; f)$ and $\mathbf{g}_\nu(\mathbf{x}, \mathbf{v}; f)$, understanding that the probability density is given by (2.4a) (or by (3.2a) for particle discretizations; see Section 3).

2.2 Examples

Below we list a few examples of collision operators that fit the description presented in Section 2.1.

2.2.1 Lenard-Bernstein operator

The Lenard-Bernstein collision operator,

$$C[f] = \nu_c \left(\mu \nabla_v \cdot (\mathbf{v}f) + \frac{\gamma^2}{2} \Delta_v f \right), \quad (2.5)$$

where $\nu_c > 0$, $\mu > 0$, and $\gamma > 0$ are parameters, models small-angle collisions, and was originally used to study longitudinal plasma oscillations (see [17], [63], [73]). It can be easily verified that an example decomposition (2.2) for $M = 3$ is given by the functions

$$\mathbf{G}(\mathbf{x}, \mathbf{v}) = -\nu_c \mu \mathbf{v}, \quad \mathbf{g}_1(\mathbf{x}, \mathbf{v}) = \begin{pmatrix} \sqrt{\nu_c} \gamma \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{g}_2(\mathbf{x}, \mathbf{v}) = \begin{pmatrix} 0 \\ \sqrt{\nu_c} \gamma \\ 0 \end{pmatrix}, \quad \mathbf{g}_3(\mathbf{x}, \mathbf{v}) = \begin{pmatrix} 0 \\ 0 \\ \sqrt{\nu_c} \gamma \end{pmatrix}. \quad (2.6)$$

Note that these functions do not explicitly depend on f , therefore in this case (1.1) is a linear Fokker-Planck equation.

2.2.2 Lorentz operator

The Lorentz collision operator models electron-ion interactions via pitch-angle scattering and is given by the formula

$$C[f] = \frac{\nu_c(|\mathbf{v}|)}{2} \nabla_v \cdot (|\mathbf{v}|^2 \mathbb{I} - \mathbf{v} \otimes \mathbf{v}) \nabla_v f, \quad (2.7)$$

where $\nu_c(|\mathbf{v}|)$ is the collisional frequency as a function of the absolute value of velocity, \mathbb{I} is the 3×3 identity matrix, and \otimes denotes tensor product. The primary effect of this type of scattering is a change of the direction of the electron's velocity with negligible energy loss. More information about the Lorentz collision operator, including the exact form of the collision frequency, can be found in, e.g., [5], [17], [53], [73]. It can be verified by a straightforward calculation that an example decomposition (2.2) for $M = 3$ is given by the functions

$$\mathbf{G}(\mathbf{x}, \mathbf{v}) = 0, \quad \mathbf{g}_1(\mathbf{x}, \mathbf{v}) = \sqrt{\nu_c(|\mathbf{v}|)} \begin{pmatrix} 0 \\ -v^3 \\ v^2 \end{pmatrix}, \quad \mathbf{g}_2(\mathbf{x}, \mathbf{v}) = \sqrt{\nu_c(|\mathbf{v}|)} \begin{pmatrix} v^3 \\ 0 \\ -v^1 \end{pmatrix}, \quad \mathbf{g}_3(\mathbf{x}, \mathbf{v}) = \sqrt{\nu_c(|\mathbf{v}|)} \begin{pmatrix} -v^2 \\ v^1 \\ 0 \end{pmatrix}. \quad (2.8)$$

Note that these functions do not explicitly depend on f , therefore also in this case (1.1) is a linear Fokker-Planck equation.

2.2.3 Coulomb/Landau operator

The more general Coulomb collision operator has the form (2.1) with

$$\begin{aligned} D_{ij}(\mathbf{x}, \mathbf{v}; f) &= N_{tot} \Gamma \int_{\mathbb{R}^3} \frac{|\mathbf{v} - \mathbf{u}|^2 \delta_{ij} - (v^i - u^i)(v^j - u^j)}{|\mathbf{v} - \mathbf{u}|^3} f(\mathbf{x}, \mathbf{u}, t) d^3 \mathbf{u}, \\ K_i(\mathbf{x}, \mathbf{v}; f) &= -2N_{tot} \Gamma \int_{\mathbb{R}^3} \frac{v^i - u^i}{|\mathbf{v} - \mathbf{u}|^3} f(\mathbf{x}, \mathbf{u}, t) d^3 \mathbf{u}, \end{aligned} \quad (2.9)$$

where N_{tot} appears due to our normalization of f , δ_{ij} is Kronecker's delta, and $\Gamma = (4\pi q^4/m^2) \ln \Lambda$, with $\ln \Lambda$ denoting the so-called Coulomb logarithm. The Coulomb operator describes collisions in which the fundamental two-body force obeys an inverse square law, and makes the assumption that small-angle collisions are more important than collisions resulting in large momentum changes (see [17], [73], [86]). A decomposition (2.2) can be found, for example, via the procedure outlined in Section 2.1. However, the expressions for \mathbf{G} and \mathbf{g}_ν are complicated, therefore we are not stating them here explicitly. Note that D_{ij} and K_i explicitly depend on f . Therefore, for the Coulomb operator the Vlasov equation (1.1) is a strongly nonlinear Fokker-Planck equation. Note also that D_{ij} and K_i can be explicitly written as functionals of the stochastic processes \mathbf{X} and \mathbf{V} as

$$\begin{aligned} D_{ij}(\mathbf{x}, \mathbf{v}; \mathbf{X}, \mathbf{V}) &= N_{tot} \Gamma \cdot \mathbb{E} \left[\frac{|\mathbf{v} - \mathbf{V}(t)|^2 \delta_{ij} - (v^i - V^i(t))(v^j - V^j(t))}{|\mathbf{v} - \mathbf{V}(t)|^3} \delta(\mathbf{x} - \mathbf{X}(t)) \right], \\ K_i(\mathbf{x}, \mathbf{v}; \mathbf{X}, \mathbf{V}) &= -2N_{tot} \Gamma \cdot \mathbb{E} \left[\frac{v^i - V^i(t)}{|\mathbf{v} - \mathbf{V}(t)|^3} \delta(\mathbf{x} - \mathbf{X}(t)) \right]. \end{aligned} \quad (2.10)$$

The collision operator (2.1) with D_{ij} and K_i as in (2.9) can also be expressed in an equivalent, although more symmetric form, known as the Landau form of the Coulomb operator, or simply the Landau collision operator (see, e.g., [17]).

3 Stochastic particle discretization of the Vlasov-Maxwell equations

Particle modelling is one of the most popular numerical techniques for solving the Vlasov equation (see, e.g., [6], [44]). In this section we discuss the connections between particle methods and stochastic modelling.

The standard particle method for the collisionless Vlasov equation (1.1) (with $C[f] = 0$) consists of substituting the Ansatz $f(\mathbf{x}, \mathbf{v}, t) = \sum_{a=1}^N w_a \delta(\mathbf{x} - \mathbf{X}_a(t)) \delta(\mathbf{v} - \mathbf{V}_a(t))$ for the particle density function, and deriving the corresponding ordinary differential equations satisfied by the 'particle' positions $\mathbf{X}_a(t)$ and velocities $\mathbf{V}_a(t)$, which turn out to be the characteristic equations. Note that we did a qualitatively similar thing in Section 2.1, where we turned the original collisional Vlasov equation into the system of stochastic differential equations (2.3), which in the absence of the forcing

terms \mathbf{G} and \mathbf{g}_ν have the same form as the characteristic equations, and in fact the ‘particles’ $\mathbf{X}_a(t)$ and $\mathbf{V}_a(t)$ can be interpreted as realizations of the stochastic processes $\mathbf{X}(t)$ and $\mathbf{V}(t)$ for different elementary events $\omega \in \Omega$.

When the right-hand side of (2.3) does not depend on f , then (2.3) can in principle be solved numerically with the help of any standard stochastic numerical method (see, e.g., [55]), and each realization of the stochastic processes can be simulated independently of others. When the right-hand side of (2.3) depends on f , then all realizations of the stochastic processes have to be solved for simultaneously, so that at each time step the probability density function f can be numerically approximated (see, e.g., [36]). Such an approach, however, does not quite lend itself to a geometric formulation. Therefore, in order to be able to introduce a variational principle in Section 4, let us consider $2N$ stochastic processes $\mathbf{X}_1, \mathbf{V}_1, \dots, \mathbf{X}_N, \mathbf{V}_N$, with each pair $(\mathbf{X}_a, \mathbf{V}_a)$ satisfying the stochastic differential system

$$d\mathbf{X}_a = \mathbf{V}_a dt, \quad (3.1a)$$

$$d\mathbf{V}_a = \left(\frac{q}{m} \mathbf{E}(\mathbf{X}_a, t) + \frac{q}{m} \mathbf{V}_a \times \mathbf{B}(\mathbf{X}_a, t) + \mathbf{G}(\mathbf{X}_a, \mathbf{V}_a; f) \right) dt + \sum_{\nu=1}^M \mathbf{g}_\nu(\mathbf{X}_a, \mathbf{V}_a; f) \circ dW_a^\nu(t), \quad (3.1b)$$

for $a = 1, \dots, N$, where $\mathbf{W}_a = (W_a^1, \dots, W_a^M)$ are N independent M -dimensional Wiener processes. Note that the systems (3.1) are decoupled from each other for different values of a , and each system is driven by an independent Wiener process \mathbf{W}_a . Therefore, the pairs $(\mathbf{X}_a, \mathbf{V}_a)$ for $a = 1, \dots, N$ are independent identically distributed (i.i.d.) stochastic processes, each with the probability density function f that satisfies the original Fokker-Planck equation (1.1). In that sense (3.1) is equivalent to (2.3). The advantage is that instead of considering N realizations of the 6-dimensional stochastic process (\mathbf{X}, \mathbf{V}) in (2.3), one can consider one realization of the $6N$ -dimensional process $(\mathbf{X}_1, \mathbf{V}_1, \dots, \mathbf{X}_N, \mathbf{V}_N)$ in (3.1). Such a reformulation will allow us to identify an underlying stochastic variational principle in Section 4. The last step leading to the stochastic particle discretization is approximating the probability density function f in (3.1). This can be done with the help of the law of large numbers, namely, one can approximate (2.4) for large N as

$$f(\mathbf{x}, \mathbf{v}, t) \approx \frac{1}{N} \sum_{a=1}^N \delta(\mathbf{x} - \mathbf{X}_a(t)) \delta(\mathbf{v} - \mathbf{V}_a(t)), \quad (3.2a)$$

$$\rho(\mathbf{x}, t) \approx \frac{qN_{tot}}{N} \sum_{a=1}^N \delta(\mathbf{x} - \mathbf{X}_a(t)), \quad (3.2b)$$

$$\mathbf{J}(\mathbf{x}, t) \approx \frac{qN_{tot}}{N} \sum_{a=1}^N \mathbf{V}_a(t) \delta(\mathbf{x} - \mathbf{X}_a(t)). \quad (3.2c)$$

It is easy to see that (3.2a) coincides with the standard Ansatz used in particle modelling (with the weights $w_a = 1/N$). Therefore, the system of stochastic differential equations (3.1) with the approximation (3.2), and with the electromagnetic field coupled via the Maxwell equations (1.2), can be considered as a stochastic particle discretization of the collisional Vlasov-Maxwell equations.

Remark. Upon substituting (3.2a), the forcing terms \mathbf{G} and \mathbf{g}_ν become functionals of the processes $\mathbf{X}_1, \dots, \mathbf{X}_N$ and $\mathbf{V}_1, \dots, \mathbf{V}_N$. Similar to the discussion in Section 2.1, for convenience and

simplicity, throughout this work we will stick to the notation $\mathbf{G}(\mathbf{x}, \mathbf{v}; f)$ and $\mathbf{g}_\nu(\mathbf{x}, \mathbf{v}; f)$, understanding that the probability density is given by (3.2a) for particle discretizations.

4 Variational principle for the particle discretization

In this section we propose an action functional which can be understood as a stochastic version of the Low action functional (see [69]), and we prove a variational principle underlying the particle discretization introduced in Section 3, akin to the stochastic Lagrange-d'Alembert principle first introduced in [59].

4.1 Function spaces

Before we introduce the action functional, we need to identify suitable function spaces on which it will be defined. For simplicity, let our spatial domain be the whole three-dimensional space \mathbb{R}^3 , and let us consider the time interval $[0, T]$ for some $T > 0$. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the probability space with the filtration $\{\mathcal{F}_t\}_{t \geq 0}$, and let $\mathbf{W}_a = (W_a^1, \dots, W_a^M)$ for $a = 1, \dots, N$ denote N independent M -dimensional Wiener processes on that probability space (such that $W_a^\nu(t)$ is \mathcal{F}_t -measurable for all $t \geq 0$). The stochastic processes $\mathbf{X}_a(t)$ and $\mathbf{V}_a(t)$ satisfy (3.1), so they are in particular \mathcal{F}_t -adapted semimartingales, and have almost surely continuous paths (see [84]). We also notice that there is no diffusion term in (3.1a), therefore we even have that the processes $\mathbf{X}_a(t)$ are almost surely of class C^1 . We introduce the notation

$$C_{\Omega, T}^k = \{\mathbf{X} \in L^2(\Omega \times [0, T], \mathbb{R}^3) \mid \mathbf{X} \text{ is a } \mathcal{F}_t\text{-adapted semimartingale, almost surely of class } C^k\}. \quad (4.1)$$

Note that this set is a vector space (see [84]). The potentials φ and \mathbf{A} satisfy the Maxwell equations (1.2) and (1.4), therefore we require them to be of class C^2 . However, since our spatial domain is unbounded, we further need to assume that the vector fields \mathbf{E} and \mathbf{B} are square integrable. We introduce the notation

$$\begin{aligned} \mathfrak{X}(\mathbb{R}^n) &= \{\mathbf{A} \in C^2(\mathbb{R}^3 \times [0, T], \mathbb{R}^n) \cap L^\infty(\mathbb{R}^3 \times [0, T], \mathbb{R}^n) \mid \forall i, j : \frac{\partial A^i}{\partial x^j}, \frac{\partial A^i}{\partial t} \in L^2(\mathbb{R}^3 \times [0, T])\}, \\ \mathfrak{X}_0(\mathbb{R}^n) &= C_0^2(\mathbb{R}^3 \times [0, T], \mathbb{R}^n), \end{aligned} \quad (4.2)$$

where $\mathfrak{X}_0(\mathbb{R}^n)$ is simply the space of compactly supported elements of $\mathfrak{X}(\mathbb{R}^n)$.

4.2 Action functional

Let us consider the action functional

$$S : \Omega \times (C_{\Omega, T}^1)^N \times (C_{\Omega, T}^0)^N \times (C_{\Omega, T}^0)^N \times \mathfrak{X}(\mathbb{R}) \times \mathfrak{X}(\mathbb{R}^3) \longrightarrow \mathbb{R} \quad (4.3)$$

defined by the formula

$$\begin{aligned}
S[\mathbf{X}_1, \dots, \mathbf{X}_N, \mathbf{V}_1, \dots, \mathbf{V}_N, \mathbf{P}_1, \dots, \mathbf{P}_N, \varphi, \mathbf{A}] = \\
\frac{N_{tot}}{N} \sum_{a=1}^N \left[\int_0^T \left(\frac{m}{2} |\mathbf{V}_a|^2 - q\varphi(\mathbf{X}_a, t) + q\mathbf{V}_a \cdot \mathbf{A}(\mathbf{X}_a, t) + \mathbf{P}_a \cdot (\dot{\mathbf{X}}_a - \mathbf{V}_a) \right) dt \right] + \int_0^T \int_{\mathbb{R}^3} \frac{1}{2} (|\mathbf{E}|^2 - |\mathbf{B}|^2) d^3\mathbf{x} dt,
\end{aligned} \tag{4.4}$$

where $\dot{\mathbf{X}}_a$ denotes the time derivative of \mathbf{X}_a , and the electric and magnetic fields \mathbf{E} and \mathbf{B} are expressed in terms of the partial derivatives of the potentials φ and \mathbf{A} as in (1.4). Following the standard convention in stochastic analysis, we will omit writing elementary events $\omega \in \Omega$ as arguments of stochastic processes unless otherwise needed, i.e., $\mathbf{X}_a(t) \equiv \mathbf{X}_a(\omega, t)$. The action functional (4.4) resembles the Low action functional introduced in [69]. In fact, it can be viewed as a particle discretization of the Low action functional, written in terms of stochastic processes (see [33], [67], [88], [91], [106], [107]). The term $\mathbf{P}_a \cdot (\dot{\mathbf{X}}_a - \mathbf{V}_a)$ is the so-called Hamilton-Pontryagin kinematic constraint (see, e.g., [60], [109]) that enforces that $\dot{\mathbf{X}}_a = \mathbf{V}_a$ using the Lagrange multiplier \mathbf{P}_a , which turns out to be the conjugate momentum. In principle, this constraint is not necessary in our context—we could omit it and replace \mathbf{V}_a with $\dot{\mathbf{X}}_a$ in (4.4). We will, however, keep it in order to make a clear connection with the theory developed in [11]. It also makes the notation in the proof of the stochastic Lagrange-d'Alembert principle in Section 4.3 more convenient and elegant. Note that the action functional S is itself a random variable, as $\omega \in \Omega$ is one of its arguments. The variations of S with respect to its arguments are given by (see Appendix A for the details of the derivations)

$$\begin{aligned}
\delta_{\mathbf{X}_a} S = \frac{N_{tot}}{N} \left(\mathbf{P}_a(T) \cdot \delta \mathbf{X}_a(T) - \mathbf{P}_a(0) \cdot \delta \mathbf{X}_a(0) \right) \\
+ \frac{N_{tot}}{N} \left[- \int_0^T \delta \mathbf{X}_a \circ d\mathbf{P}_a + \int_0^T \left(-q \nabla_x \varphi(\mathbf{X}_a, t) \cdot \delta \mathbf{X}_a + q \sum_{i,j=1}^3 V^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}_a, t) \delta X_a^i \right) dt \right],
\end{aligned} \tag{4.5a}$$

$$\delta_{\mathbf{V}_a} S = \frac{N_{tot}}{N} \int_0^T (m \mathbf{V}_a + q \mathbf{A}(\mathbf{X}_a, t) - \mathbf{P}_a) \cdot \delta \mathbf{V}_a dt, \tag{4.5b}$$

$$\delta_{\mathbf{P}_a} S = \frac{N_{tot}}{N} \int_0^T (\dot{\mathbf{X}}_a - \mathbf{V}_a) \cdot \delta \mathbf{P}_a dt, \tag{4.5c}$$

$$\delta_{\mathbf{A}} S = \int_0^T \int_{\mathbb{R}^3} \left(\mathbf{J} + \frac{\partial \mathbf{E}}{\partial t} - \nabla_x \times \mathbf{B} \right) \cdot \delta \mathbf{A} d^3\mathbf{x} dt - \int_{\mathbb{R}^3} (\mathbf{E}(\mathbf{x}, T) \cdot \delta \mathbf{A}(\mathbf{x}, T) - \mathbf{E}(\mathbf{x}, 0) \cdot \delta \mathbf{A}(\mathbf{x}, 0)) d^3\mathbf{x}, \tag{4.5d}$$

$$\delta_{\varphi} S = \int_0^T \int_{\mathbb{R}^3} (\nabla_x \cdot \mathbf{E} - \rho) \cdot \delta \varphi d^3\mathbf{x} dt, \tag{4.5e}$$

where ρ and \mathbf{J} are defined in (3.2b) and (3.2c), respectively. The total variation of S with respect to the variations of all arguments equals

$$\delta S = \sum_{a=1}^N (\delta_{\mathbf{X}_a} S + \delta_{\mathbf{V}_a} S + \delta_{\mathbf{P}_a} S) + \delta_{\varphi} S + \delta_{\mathbf{A}} S. \tag{4.6}$$

4.3 The stochastic Lagrange-d'Alembert principle

While the standard rules of the calculus of variations apply to the variations (4.5d) and (4.5e), the variations (4.5a), (4.5b), (4.5c) involve stochastic processes and stochastic integrals. Therefore, before we can formulate a stochastic variational principle, we need the following lemma, whose proof is given in Appendix B.

Lemma 4.1. *Let $\mathbf{X} \in C_{\Omega, T}^1$ and $\mathbf{V}, \mathbf{P} \in C_{\Omega, T}^0$, and let $\mathbf{R}, \mathbf{r}_\nu : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be of class C^1 for $\nu = 1, \dots, M$. Then*

$$\forall \mathbf{Z} \in C_{\Omega, T}^1 : \int_0^T \left(\mathbf{Z}(t) \circ d\mathbf{P} - \mathbf{R}(\mathbf{X}, \mathbf{V}) \cdot \mathbf{Z}(t) dt - \sum_{\nu=1}^M \mathbf{r}_\nu(\mathbf{X}, \mathbf{V}) \cdot \mathbf{Z}(t) \circ dW^\nu(t) \right) = 0 \quad a.s. \quad (4.7)$$

if and only if

$$\forall t \in [0, T] : \int_0^t \left(d\mathbf{P}(\tau) - \mathbf{R}(\mathbf{X}(\tau), \mathbf{V}(\tau)) d\tau - \sum_{\nu=1}^M \mathbf{r}_\nu(\mathbf{X}(\tau), \mathbf{V}(\tau)) \circ dW^\nu(\tau) \right) = 0 \quad a.s., \quad (4.8)$$

where ‘a.s.’ means almost surely.

Remark. Equation (4.8) means that $\mathbf{P}(t)$, $\mathbf{X}(t)$, and $\mathbf{V}(t)$ satisfy a stochastic differential equation, which can be written in the differential form as

$$d\mathbf{P}(t) = \mathbf{R}(\mathbf{X}(t), \mathbf{V}(t)) dt + \sum_{\nu=1}^M \mathbf{r}_\nu(\mathbf{X}(t), \mathbf{V}(t)) \circ dW^\nu(t). \quad (4.9)$$

We are now in a position to formulate and prove a stochastic variational principle that generalizes the deterministic Lagrange-d'Alembert principle for forced Lagrangian and Hamiltonian systems, akin to the stochastic variational principle introduced in [59].

Theorem 4.2 (Stochastic Lagrange-d'Alembert principle for particles). *Let $\mathbf{X}_a \in C_{\Omega, T}^1$ and $\mathbf{V}_a, \mathbf{P}_a \in C_{\Omega, T}^0$ for $a = 1, \dots, N$ be stochastic processes, and let $\mathbf{A} \in \mathfrak{X}(\mathbb{R}^3)$, $\varphi \in \mathfrak{X}(\mathbb{R})$ be functions. Assume that $\mathbf{G}(\cdot, \cdot; f)$ and $\mathbf{g}_\nu(\cdot, \cdot; f)$ for $\nu = 1, \dots, M$ are C^1 functions of their arguments, where f is given by (3.2a). Then $\mathbf{X}_a, \mathbf{V}_a, \mathbf{P}_a, \mathbf{A}$, and φ satisfy the system of stochastic differential equations*

$$\dot{\mathbf{X}}_a = \mathbf{V}_a, \quad (4.10a)$$

$$\mathbf{P}_a = m\mathbf{V}_a + q\mathbf{A}(\mathbf{X}_a, t), \quad (4.10b)$$

$$dP_a^i = \left(-q \frac{\partial \varphi}{\partial x^i}(\mathbf{X}_a, t) + q \sum_{j=1}^3 V_a^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}_a, t) + m G^i(\mathbf{X}_a, \mathbf{V}_a; f) \right) dt + m \sum_{\nu=1}^M g_\nu^i(\mathbf{X}_a, \mathbf{V}_a; f) \circ dW_a^\nu(t), \quad (4.10c)$$

for $i = 1, 2, 3$ and $a = 1, \dots, N$, together with the Maxwell equations (1.2), (1.4), (3.2) on the time interval $[0, T]$, if and only if they satisfy the following variational principle

$$\delta S + \frac{mN_{tot}}{N} \sum_{a=1}^N \left[\int_0^T \mathbf{G}(\mathbf{X}_a, \mathbf{V}_a; f) \cdot \delta \mathbf{X}_a dt + \sum_{\nu=1}^M \int_0^T \mathbf{g}_\nu(\mathbf{X}_a, \mathbf{V}_a; f) \cdot \delta \mathbf{X}_a \circ dW_a^\nu(t) \right] = 0 \quad (4.11)$$

for arbitrary variations $\delta \mathbf{X}_a \in C_{\Omega, T}^1$, $\delta \mathbf{V}_a, \delta \mathbf{P}_a \in C_{\Omega, T}^0$, $\delta \mathbf{A} \in \mathfrak{X}_0(\mathbb{R}^3)$, and $\delta \varphi \in \mathfrak{X}_0(\mathbb{R})$, with $\delta \mathbf{X}_a(0) = \delta \mathbf{X}_a(T) = 0$ almost surely, and $\delta \mathbf{A}(\mathbf{x}, 0) = \delta \mathbf{A}(\mathbf{x}, T) = 0$ for all $\mathbf{x} \in \mathbb{R}^3$, where the action functional S is given by (4.4).

Proof. Let us first consider the variations with respect to \mathbf{A} in (4.11). Given the boundary conditions for $\delta \mathbf{A}$, from the standard calculus of variations we have that $\delta_{\mathbf{A}} S = 0$ (see Equation (4.5d)) for all $\delta \mathbf{A}$ if and only if (1.2d) is satisfied. Similarly, $\delta_\varphi S = 0$ (see Equation (4.5e)) holds for all $\delta \varphi$ if and only if (1.2a) holds. Further, for variations with respect to \mathbf{V}_a we have that $\delta_{\mathbf{V}_a} S = 0$ (see Equation (4.5b)) for all $\delta \mathbf{V}_a$ if and only if (4.10b) is satisfied almost surely, which follows from the standard theorem of the calculus of variations, since the integral in (4.5b) is a standard Lebesgue integral, and the integrands are almost surely continuous. Similarly, $\delta_{\mathbf{P}_a} S = 0$ (see Equation (4.5c)) for all $\delta \mathbf{P}_a$ if and only if (4.10a) is satisfied almost surely. Finally, for variations with respect to \mathbf{X}_a , Equations (4.5a) and (4.11) give

$$\int_0^T \left(-\delta \mathbf{X}_a \circ d\mathbf{P}_a + \left(-q \nabla_x \varphi(\mathbf{X}_a, t) \cdot \delta \mathbf{X}_a + q \sum_{i,j=1}^3 V_a^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}_a, t) \delta X_a^i + m \mathbf{G}(\mathbf{X}_a, \mathbf{V}_a; f) \cdot \delta \mathbf{X}_a \right) dt + m \sum_{\nu=1}^M \mathbf{g}_\nu(\mathbf{X}_a, \mathbf{V}_a; f) \cdot \delta \mathbf{X}_a \circ dW_a^\nu(t) \right) = 0, \quad (4.12)$$

which, by Lemma 4.1, holds for all $\delta \mathbf{X}_a$ if and only if (4.10c) is satisfied. □

Remark. Equation (4.10) is expressed in terms of the Lagrange multipliers \mathbf{P}_a , which, as can be seen in (4.10b), turn out to be the conjugate momenta. The conjugate momenta can be eliminated, and Equation (4.10) can be recast as Equation (3.1b), which is shown in the following theorem.

Theorem 4.3. *Equations (3.1) and (4.10) are equivalent.*

Proof. By calculating the stochastic differential on both sides of (4.10b) and substituting (4.10a), we obtain

$$dP_a^i = m dV_a^i + q \sum_{j=1}^3 V_a^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}_a, t) dt + q \frac{\partial A^i}{\partial t}(\mathbf{X}_a, t) dt \quad (4.13)$$

for each $i = 1, 2, 3$ and $a = 1, \dots, N$. Comparing this with (4.10c), and using (1.4), one eliminates the conjugate momenta and obtains Equation (3.1b). □

Remark. Theorems 4.2 and 4.3 provide a variational formulation of the stochastic particle method from Section 3. One can further perform a variational discretization of the electromagnetic fields \mathbf{A} and φ , for instance along the lines of [91], [93] or [58], thus obtaining a stochastic particle-in-cell (PIC) discretization of the collisional Vlasov-Maxwell equations. The resulting structure-preserving numerical methods will be investigated in a follow-up work.

5 Variational principle for the Vlasov-Maxwell equations

The form of the action functional (4.4) and of the Lagrange-d'Alembert principle (4.11) suggests that it should be possible to formulate a similar variational principle for the stochastic reformulation of the Vlasov-Maxwell system discussed in Section 2.1. In this section we provide such a variational principle for a class of collision operators.

5.1 Action functional

Let us consider the action functional defined by the formula

$$\bar{S}[\mathbf{X}, \mathbf{V}, \mathbf{P}, \varphi, \mathbf{A}] = N_{tot} \cdot \mathbb{E} \left[\int_0^T \left(\frac{m}{2} |\mathbf{V}|^2 - q\varphi(\mathbf{X}, t) + q\mathbf{V} \cdot \mathbf{A}(\mathbf{X}, t) + \mathbf{P} \cdot (\dot{\mathbf{X}} - \mathbf{V}) \right) dt \right] + \int_0^T \int_{\mathbb{R}^3} \frac{1}{2} (|\mathbf{E}|^2 - |\mathbf{B}|^2) d^3\mathbf{x} dt, \quad (5.1)$$

where $\dot{\mathbf{X}}$ denotes the time derivative of \mathbf{X} , the electric and magnetic fields \mathbf{E} and \mathbf{B} are expressed in terms of the partial derivatives of the potentials φ and \mathbf{A} as in (1.4), and $\mathbb{E}[Y] \equiv \int_{\Omega} Y d\mathbb{P}$ denotes the expected value of the random variable Y . Note that unlike S in (4.4), the action functional \bar{S} is not a random variable, as the dependence on $\omega \in \Omega$ is integrated out with respect to the probability measure by calculating the expected value. In fact, S could be regarded as a Monte Carlo approximation of \bar{S} when the processes $\mathbf{X}_1, \dots, \mathbf{X}_N$ are independent and identically distributed as \mathbf{X} , and similarly for \mathbf{V} and \mathbf{P} . An important issue to consider is the domain of this action functional. In a fashion similar to (4.3), one may want to take as the domain the set

$$C_{\Omega, T}^1 \times C_{\Omega, T}^0 \times C_{\Omega, T}^0 \times \mathfrak{X}(\mathbb{R}) \times \mathfrak{X}(\mathbb{R}^3), \quad (5.2)$$

on which the formula (5.1) is well-defined. This domain, however, turns out to be too big, in the sense that, as will be discussed below, due to the presence of the expected value the variations of \bar{S} do not uniquely determine the set of stochastic evolution equations (2.3). It is therefore necessary to restrict (5.2) to a smaller subspace or submanifold which is compatible with the considered collision operator. Below we will demonstrate how this can be done for a class of collision operators (2.1) for which $D_{ij}(\mathbf{x}, \mathbf{v}; f) = \text{const}$, that is, we have

$$\mathbf{g}_{\nu}(\mathbf{x}, \mathbf{v}; f) = \chi_{\nu} = \text{const}. \quad (5.3)$$

This class encompasses, for instance, the Lenard-Bernstein operator (2.6), or the more general nonlinear energy and momentum preserving Dougherty collision operator and its modifications (see [20], [29], [30], [34], [40], [56], [77], [79]). For a given collision operator of the form (5.3), we define a compatible subset of $C_{\Omega, T}^0$, namely,

$$C_{\text{col}} = \left\{ \mathbf{P} \in C_{\Omega, T}^0 \mid \exists \mathbf{Z} \in C_{\Omega, T}^0 : d\mathbf{P} = \mathbf{Z} dt + m \sum_{\nu=1}^M \boldsymbol{\chi}_\nu dW^\nu(t) \right\}. \quad (5.4)$$

Note that for any $\mathbf{P}_1, \mathbf{P}_2 \in C_{\text{col}}$ we have that $d(\mathbf{P}_1 - \mathbf{P}_2) = (\mathbf{Z}_1 - \mathbf{Z}_2) dt$, that is, $\mathbf{P}_1 - \mathbf{P}_2 \in C_{\Omega, T}^1$. Therefore, the pair $(C_{\text{col}}, C_{\Omega, T}^1)$ is an affine subspace of $C_{\Omega, T}^0$. The action functional \bar{S} can now be defined as

$$\bar{S} : C_{\Omega, T}^1 \times C_{\Omega, T}^0 \times C_{\text{col}} \times \mathfrak{X}(\mathbb{R}) \times \mathfrak{X}(\mathbb{R}^3) \longrightarrow \mathbb{R}. \quad (5.5)$$

Similar to the calculations in Section 4.2, the variations of \bar{S} with respect to \mathbf{V} and \mathbf{P} are given by, respectively,

$$\delta_{\mathbf{V}} \bar{S} = N_{\text{tot}} \cdot \mathbb{E} \left[\int_0^T (m\mathbf{V} + q\mathbf{A}(\mathbf{X}, t) - \mathbf{P}) \cdot \delta\mathbf{V} dt \right], \quad (5.6)$$

$$\delta_{\mathbf{P}} \bar{S} = N_{\text{tot}} \cdot \mathbb{E} \left[\int_0^T (\dot{\mathbf{X}} - \mathbf{V}) \cdot \delta\mathbf{P} dt \right], \quad (5.7)$$

except that here $\delta\mathbf{P} \in C_{\Omega, T}^1$, so that $\mathbf{P} + \epsilon\delta\mathbf{P} \in C_{\text{col}}$. For the variation of \bar{S} with respect to \mathbf{X} we have

$$\begin{aligned} \delta_{\mathbf{X}} \bar{S} = & N_{\text{tot}} \cdot \mathbb{E} \left(\mathbf{P}(T) \cdot \delta\mathbf{X}(T) - \mathbf{P}(0) \cdot \delta\mathbf{X}(0) \right) \\ & + N_{\text{tot}} \cdot \mathbb{E} \left[- \int_0^T \delta\mathbf{X} \circ d\mathbf{P} + \int_0^T \left(-q\nabla_x \varphi(\mathbf{X}, t) \cdot \delta\mathbf{X} + q \sum_{i,j=1}^3 V^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}, t) \delta X^i \right) dt \right]. \end{aligned} \quad (5.8)$$

Since $\mathbf{P} \in C_{\text{col}}$, we have that $d\mathbf{P} = \mathbf{Z} dt + m \sum_{\nu=1}^M \boldsymbol{\chi}_\nu dW^\nu(t)$. Furthermore, the variations $\delta\mathbf{X}$ are almost surely of class C^1 , and therefore have sample paths of almost surely finite variation. Consequently, the quadratic covariation $[\boldsymbol{\chi}_\nu \cdot \delta\mathbf{X}, W^\nu]_0^T = 0$ almost surely (see [84]). Since the expected value of the Itô integral with respect to the Wiener process is zero, we altogether have that

$$\mathbb{E} \left[\int_0^T \boldsymbol{\chi}_\nu \cdot \delta\mathbf{X} \circ dW^\nu(t) \right] = 0, \quad \text{for all } \nu = 1, \dots, M. \quad (5.9)$$

By plugging this in (5.8), we finally obtain

$$\begin{aligned} \delta_{\mathbf{X}} \bar{S} = & N_{\text{tot}} \cdot \mathbb{E} \left(\mathbf{P}(T) \cdot \delta\mathbf{X}(T) - \mathbf{P}(0) \cdot \delta\mathbf{X}(0) \right) \\ & + N_{\text{tot}} \cdot \mathbb{E} \left[\int_0^T \left(-\mathbf{Z} \cdot \delta\mathbf{X} - q\nabla_x \varphi(\mathbf{X}, t) \cdot \delta\mathbf{X} + q \sum_{i,j=1}^3 V^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}, t) \delta X^i \right) dt \right]. \end{aligned} \quad (5.10)$$

The variations with respect to \mathbf{A} and φ are the same as in (4.5d) and (4.5e), respectively, only with the charge and electric current densities given by (2.4) rather than (3.2). The total variation of \bar{S} with respect to the variations of all arguments is given by

$$\delta\bar{S} = \delta_{\mathbf{X}}\bar{S} + \delta_{\mathbf{V}}\bar{S} + \delta_{\mathbf{P}}\bar{S} + \delta_{\varphi}\bar{S} + \delta_{\mathbf{A}}\bar{S}. \quad (5.11)$$

5.2 The stochastic Lagrange-d'Alembert principle

In the following theorems we establish a variational principle for the system of equations (1.2), (2.3), and (2.4) for a class of collision operators with $\mathbf{g}_{\nu}(\mathbf{x}, \mathbf{v}; f) = \chi_{\nu} = \text{const}$ for all $\nu = 1, \dots, M$.

Theorem 5.1 (Stochastic Lagrange-d'Alembert principle for the VM equations). *Let $\mathbf{X} \in C_{\Omega, T}^1$, $\mathbf{V} \in C_{\Omega, T}^0$, $\mathbf{P} \in C_{\text{col}}$ be stochastic processes, and let $\mathbf{A} \in \mathfrak{X}(\mathbb{R}^3)$, $\varphi \in \mathfrak{X}(\mathbb{R})$ be functions. Assume that $\mathbf{G}(\cdot, \cdot; f)$ is a C^1 function of its arguments, where f is given by (2.4a). Then \mathbf{X} , \mathbf{V} , \mathbf{P} , \mathbf{A} , and φ satisfy the system of stochastic differential equations*

$$\dot{\mathbf{X}} = \mathbf{V}, \quad (5.12a)$$

$$\mathbf{P} = m\mathbf{V} + q\mathbf{A}(\mathbf{X}, t), \quad (5.12b)$$

$$dP^i = \left(-q \frac{\partial \varphi}{\partial x^i}(\mathbf{X}, t) + q \sum_{j=1}^3 V^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}, t) + m G^i(\mathbf{X}, \mathbf{V}; f) \right) dt + m \sum_{\nu=1}^M \chi_{\nu}^i dW^{\nu}(t), \quad (5.12c)$$

for $i = 1, 2, 3$, together with the Maxwell equations (1.2), (1.4), (2.4) on the time interval $[0, T]$, if and only if they satisfy the following variational principle

$$\delta\bar{S} + mN_{\text{tot}} \cdot \mathbb{E} \left[\int_0^T \mathbf{G}(\mathbf{X}, \mathbf{V}; f) \cdot \delta\mathbf{X} dt \right] = 0 \quad (5.13)$$

for arbitrary variations $\delta\mathbf{X}, \delta\mathbf{P} \in C_{\Omega, T}^1$, $\delta\mathbf{V} \in C_{\Omega, T}^0$, $\delta\mathbf{A} \in \mathfrak{X}_0(\mathbb{R}^3)$, and $\delta\varphi \in \mathfrak{X}_0(\mathbb{R})$, with $\delta\mathbf{X}(0) = \delta\mathbf{X}(T) = 0$ almost surely, and $\delta\mathbf{A}(\mathbf{x}, 0) = \delta\mathbf{A}(\mathbf{x}, T) = 0$ for all $\mathbf{x} \in \mathbb{R}^3$, where the action functional \bar{S} is given by (5.1) and (5.5).

Proof. Similar to the proof of Theorem 4.2, the equations $\delta_{\varphi}\bar{S} = 0$ and $\delta_{\mathbf{A}}\bar{S} = 0$ are equivalent to (1.2a) and (1.2d), respectively. Note that $C_{\Omega, T}^0$ is a subspace of $L^2(\Omega \times [0, T], \mathbb{R}^3)$, and $\langle \mathbf{Y}_1, \mathbf{Y}_2 \rangle = \mathbb{E}[\int_0^T \mathbf{Y}_1 \cdot \mathbf{Y}_2 dt]$ is an inner product on that space. Therefore, by substituting Equations (5.6), (5.7), and (5.10) in Equation (5.13), and using the fact that the variations are arbitrary, we establish equivalence with Equations (5.12a)-(5.12b), as well as with the equation

$$Z^i = -q \frac{\partial \varphi}{\partial x^i}(\mathbf{X}, t) + q \sum_{j=1}^3 V^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}, t) + m G^i(\mathbf{X}, \mathbf{V}; f), \quad (5.14)$$

for $i = 1, 2, 3$, which in turn is equivalent to Equation (5.12c), given the assumption $\mathbf{P} \in C_{\text{col}}$. \square

Theorem 5.2. *Equation (2.3) with $\mathbf{g}_{\nu}(\mathbf{x}, \mathbf{v}; f) = \chi_{\nu} = \text{const}$ for $\nu = 1, \dots, M$ and Equation (5.12) are equivalent.*

Proof. Similar to the proof of Theorem 4.3, by calculating the stochastic differential on both sides of Equation (5.12b) and comparing with Equation (5.12c), one eliminates \mathbf{P} and obtains Equation (2.3b). □

Remark. Note that the forcing terms \mathbf{g}_ν do not explicitly appear in the variational equation (5.13). By comparing Theorem 4.2 and Theorem 5.1, one could intuitively expect that the relevant variational principle should read

$$\delta\bar{S} + mN_{tot} \cdot \mathbb{E} \left[\int_0^T \mathbf{G}(\mathbf{X}, \mathbf{V}; f) \cdot \delta\mathbf{X} dt + \sum_{\nu=1}^M \int_0^T \mathbf{g}_\nu(\mathbf{X}, \mathbf{V}; f) \cdot \delta\mathbf{X} \circ dW^\nu(t) \right] = 0. \quad (5.15)$$

However, due to the presence of the expected value in this equation, part or all of the information about the Stratonovich integral term is lost, as we saw in (5.9) for instance. Therefore, if the domain (5.2) is chosen for \bar{S} , then the variational equations (5.13) or (5.15) do not determine a unique set of stochastic differential equations that need to be satisfied by the considered stochastic processes. Consequently, it is necessary to encode the missing information about the forcing terms \mathbf{g}_ν in the definition of the action functional \bar{S} by restricting its domain to a subset compatible with the considered collision operator. For the class of collision operators (5.3) a suitable choice of the domain is proposed in (5.5). For other collision operators appropriate domains will be nonlinear subspaces of (5.2), and they will be investigated in a follow-up work.

6 Variational principle for the Vlasov-Poisson equations

In the full Vlasov-Maxwell system the scalar φ and vector \mathbf{A} potentials are independent dynamic variables, and as such have to appear explicitly in the action functional alongside the stochastic processes \mathbf{X} , \mathbf{V} , and \mathbf{P} . In order to ensure the correct coupling between the stochastic processes and the electromagnetic field, an expected value was necessary in the definition of the action functional (5.1). This created a difficulty in deriving a variational principle, as pointed out in the remark following Theorem 5.2. This difficulty can be circumvented for the Vlasov-Poisson equations because in this case the electrostatic potential φ is uniquely determined by the stochastic process \mathbf{X} , as will be demonstrated below.

6.1 The collisional Vlasov-Poisson equations

The collisional Vlasov-Poisson equations

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_x f + \frac{q}{m} \mathbf{E} \cdot \nabla_v f = C[f], \quad (6.1)$$

where

$$\mathbf{E} = -\nabla_x \varphi, \quad (6.2a)$$

$$\Delta_x \varphi = -\rho, \quad (6.2b)$$

and the charge density ρ is given by (1.3), are an approximation of the Vlasov-Maxwell equations in the nonrelativistic zero-magnetic field limit. The associated stochastic differential equations take the form

$$d\mathbf{X} = \mathbf{V} dt, \quad (6.3a)$$

$$d\mathbf{V} = \left(\frac{q}{m} \mathbf{E}(\mathbf{X}, t) + \mathbf{G}(\mathbf{X}, \mathbf{V}; f) \right) dt + \sum_{\nu=1}^M \mathbf{g}_{\nu}(\mathbf{X}, \mathbf{V}; f) \circ dW^{\nu}(t). \quad (6.3b)$$

The equations (2.4b), (6.2), and (6.3) form a stochastic reformulation of the Vlasov-Poisson equations. A stochastic particle discretization and the corresponding stochastic variational principle can be derived just like in Sections 3 and 4, respectively. Also, a variational principle analogous to the Lagrange-d'Alembert principle presented in Section 5 can be derived in a similar fashion. However, by doing so, one encounters the same difficulty with including the Stratonovich integral. In the case of the Vlasov-Poisson equations a different variational principle can be obtained by observing that the electrostatic potential φ can be expressed as a functional of the stochastic process \mathbf{X} ,

$$\varphi : \mathbb{R}^3 \times \mathbb{R} \times C_{\Omega, T}^1 \longrightarrow \mathbb{R}, \quad (6.4)$$

by solving Poisson's equation (6.2b). Given the charge density function (2.4b) and specific boundary conditions, the solution of Poisson's equation can be written using an appropriate Green's function for the Laplacian. Assuming the spatial domain is unbounded, the standard Green's function yields

$$\varphi(\mathbf{x}, t, \mathbf{X}) = \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{\rho(\mathbf{y}, t)}{|\mathbf{x} - \mathbf{y}|} d^3\mathbf{y} = \frac{qN_{tot}}{4\pi} \mathbb{E} \left[\frac{1}{|\mathbf{x} - \mathbf{X}(t)|} \right]. \quad (6.5)$$

From (6.2a) we have the electric field

$$\mathbf{E}(\mathbf{x}, t, \mathbf{X}) = \frac{qN_{tot}}{4\pi} \mathbb{E} \left[\frac{\mathbf{x} - \mathbf{X}(t)}{|\mathbf{x} - \mathbf{X}(t)|^3} \right]. \quad (6.6)$$

6.2 Action functional

Let us consider the action functional

$$\hat{S} : \Omega \times C_{\Omega, T}^1 \times C_{\Omega, T}^1 \times C_{\Omega, T}^0 \times C_{\Omega, T}^0 \longrightarrow \mathbb{R} \quad (6.7)$$

defined by the formula

$$\hat{S}[\mathbf{X}, \mathbf{Y}, \mathbf{V}, \mathbf{P}] = \int_0^T \left(\frac{m}{2} |\mathbf{V}(t)|^2 - q\varphi(\mathbf{X}(t), t, \mathbf{Y}) + \mathbf{P}(t) \cdot (\dot{\mathbf{X}}(t) - \mathbf{V}(t)) \right) dt, \quad (6.8)$$

where the electrostatic potential φ is given by (6.5). Note that similar to S in (4.4), the functional \hat{S} is itself random, and can be viewed as the action functional of particles represented by the process \mathbf{X} which are moving in the electric field generated by particles represented by the process \mathbf{Y} . Similar

to the calculations in Section 4.2, the variations of \hat{S} with respect to \mathbf{X} , \mathbf{V} , and \mathbf{P} are given by, respectively,

$$\begin{aligned} \delta_{\mathbf{X}}\hat{S}[\mathbf{X}, \mathbf{Y}, \mathbf{V}, \mathbf{P}] &= \mathbf{P}(T) \cdot \delta\mathbf{X}(T) - \mathbf{P}(0) \cdot \delta\mathbf{X}(0) \\ &\quad - \int_0^T \delta\mathbf{X}(t) \circ d\mathbf{P}(t) + \int_0^T q\mathbf{E}(\mathbf{X}(t), t, \mathbf{Y}) \cdot \delta\mathbf{X}(t) dt, \end{aligned} \quad (6.9a)$$

$$\delta_{\mathbf{V}}\hat{S}[\mathbf{X}, \mathbf{Y}, \mathbf{V}, \mathbf{P}] = \int_0^T (m\mathbf{V}(t) - \mathbf{P}(t)) \cdot \delta\mathbf{V}(t) dt, \quad (6.9b)$$

$$\delta_{\mathbf{P}}\hat{S}[\mathbf{X}, \mathbf{Y}, \mathbf{V}, \mathbf{P}] = \int_0^T (\dot{\mathbf{X}}(t) - \mathbf{V}(t)) \cdot \delta\mathbf{P}(t) dt, \quad (6.9c)$$

where the electric field \mathbf{E} is given by (6.6). Note that we are not considering variations with respect to \mathbf{Y} . Let us for convenience define the joint variation of \hat{S} with respect to \mathbf{X} , \mathbf{V} , and \mathbf{P} as

$$\delta_{(\mathbf{X}, \mathbf{V}, \mathbf{P})}\hat{S} = \delta_{\mathbf{X}}\hat{S} + \delta_{\mathbf{V}}\hat{S} + \delta_{\mathbf{P}}\hat{S}. \quad (6.10)$$

6.3 The stochastic Lagrange-d'Alembert principle

In the following theorem we formulate a variational principle for the system of equations (2.4b), (6.2), and (6.3). Note that $\mathbf{E}(\mathbf{X}(t), t, \mathbf{X})$ is the electric field generated by a distribution of charged particles represented by the process \mathbf{X} at time t , and evaluated at the random point $\mathbf{x} = \mathbf{X}(t)$ in space. Furthermore, the notation $\delta_{\mathbf{X}}\hat{S}[\mathbf{X}, \mathbf{X}, \mathbf{V}, \mathbf{P}]$ means that the variation of \hat{S} is evaluated for the arguments $\mathbf{X}, \mathbf{Y}, \mathbf{V}, \mathbf{P}$ with $\mathbf{Y} = \mathbf{X}$.

Theorem 6.1 (Stochastic Lagrange-d'Alembert principle for the VP equations). *Let $\mathbf{X} \in C_{\Omega, T}^1$ and $\mathbf{V}, \mathbf{P} \in C_{\Omega, T}^0$ be stochastic processes, and let $\varphi(\cdot, \cdot, \mathbf{X}) \in \mathfrak{X}(\mathbb{R})$ be given by (6.5). Assume that $\mathbf{G}(\cdot, \cdot; f)$ and $\mathbf{g}_{\nu}(\cdot, \cdot; f)$ for $\nu = 1, \dots, M$ are C^1 functions of their arguments, where f is given by (2.4a). Then \mathbf{X} , \mathbf{V} , and \mathbf{P} satisfy the system of stochastic differential equations*

$$\dot{\mathbf{X}}(t) = \mathbf{V}(t), \quad (6.11a)$$

$$\mathbf{P}(t) = m\mathbf{V}(t), \quad (6.11b)$$

$$d\mathbf{P}(t) = \left(q\mathbf{E}(\mathbf{X}(t), t, \mathbf{X}) + m\mathbf{G}(\mathbf{X}(t), \mathbf{V}(t); f) \right) dt + m \sum_{\nu=1}^M \mathbf{g}_{\nu}(\mathbf{X}(t), \mathbf{V}(t); f) \circ dW^{\nu}(t), \quad (6.11c)$$

on the time interval $[0, T]$, if and only if they satisfy the following variational principle

$$\delta_{(\mathbf{X}, \mathbf{V}, \mathbf{P})}\hat{S}[\mathbf{X}, \mathbf{X}, \mathbf{V}, \mathbf{P}] + m \int_0^T \mathbf{G}(\mathbf{X}, \mathbf{V}; f) \cdot \delta\mathbf{X} dt + m \sum_{\nu=1}^M \int_0^T \mathbf{g}_{\nu}(\mathbf{X}, \mathbf{V}; f) \cdot \delta\mathbf{X} \circ dW^{\nu}(t) = 0 \quad (6.12)$$

for arbitrary variations $\delta\mathbf{X} \in C_{\Omega, T}^1$, and $\delta\mathbf{V}, \delta\mathbf{P} \in C_{\Omega, T}^0$, with $\delta\mathbf{X}(0) = \delta\mathbf{X}(T) = 0$ almost surely, where the action functional \hat{S} is given by (6.8).

Proof. Analogous to the proof of Theorem 4.2. □

Remark. It is straightforward to see that Equations (6.11), together with (6.5) and (6.6), are equivalent to the system of equations (2.4b), (6.2), and (6.3). The Lagrange-d'Alembert principle (6.12) is unusual in that the variations of the action functional \hat{S} with respect to the argument \mathbf{Y} are omitted. Thanks to such a form, however, the action functional does not require an expected value, and the collisional effects can be correctly included. A similar idea to solve Poisson's equation and plug the solution into the action functional was presented in [108], where the authors proposed a variational principle for the collisionless Vlasov-Poisson equations. In that approach the energy of the electric field was also included in the variational principle, and the variations were taken with respect to all arguments of the action functional. This approach could be adapted to the stochastic reformulation of the Vlasov-Poisson equations, but the corresponding action functional would have a form similar to (5.1), that is, it would need to contain an expected value, and therefore we would face a similar difficulty as for the Vlasov-Maxwell equations in Section 5.2.

7 Summary and future work

In this work we have considered novel stochastic formulations of the collisional Vlasov-Maxwell and Vlasov-Poisson equations, and we have identified new stochastic variational principles underlying these formulations. We have also proposed a stochastic particle method for the Vlasov-Maxwell equations and proved the corresponding stochastic variational principle.

Our work can be extended in several ways. The stochastic variational principle introduced in Section 4 can be used to construct stochastic variational particle-in-cell numerical algorithms for the collisional Vlasov-Maxwell and Vlasov-Poisson equations. Variational integrators are an important class of geometric integrators. This type of numerical schemes is based on discrete variational principles and provides a natural framework for the discretization of Lagrangian systems, including forced, dissipative, or constrained ones. These methods have the advantage that they are symplectic when applied to systems without forcing, and in the presence of a symmetry, they satisfy a discrete version of Noether's theorem. For this reason they demonstrate superior performance in long-time simulations; see [41], [51], [52], [64], [65], [72], [75], [76], [87], [97], [100]. Variational integrators were introduced in the context of finite-dimensional mechanical systems, but were later generalized to Lagrangian field theories (see [71]) and applied in many computations, for example in elasticity, electrodynamics, fluid dynamics, or plasma physics; see [56], [66], [80], [91], [93], [96], [106], [107]. Stochastic variational integrators were first introduced in [11] and further studied in [10], [47], [49], [59], [101].

In Section 5 we have proposed a general action functional for the collisional Vlasov-Maxwell equations. However, we have also determined that in order to prove a relevant variational principle, the domain of this action functional has to be restricted in a way compatible with the collision operator of interest. We have shown that for a class of collision operators with constant diffusion terms, a suitable subdomain is an affine subspace (i.e., a submanifold). A natural continuation of our work would be to investigate submanifolds of (5.2) which are suitable for other collision operators.

Another aspect worth a more detailed investigation is the issue of existence and uniqueness of the solutions of the stochastic reformulations presented in this work, which are nontrivial systems of coupled stochastic and partial differential equations. This question is closely connected to the issue of existence and uniqueness of the solutions of the original collisional Vlasov-Maxwell and the Vlasov-Poisson equations. General results are available in the collisionless case (see, e.g., [27], [104],

[105]), but the theory for the collisional equations is less developed (see [26], [31], [74], [78], [94], [102], [103] and the references therein).

Furthermore, our stochastic Lagrange-d'Alembert approach could also be adapted to relativistic plasmas (see [13]), and to variational principles with phase-space Lagrangians appearing in gyrokinetic ([9], [14], [95]) and guiding-center theories ([15], [18], [82], [83]). In particular, considering stochastic extensions of the variational principles proposed in [15] could offer an alternative stochastic description of anomalous transport in magnetically confined plasmas (see [4], [99]).

Finally, as is typical for particle methods in general, the stochastic particle discretization proposed in Section 3 will require a large number of particles for accurate numerical simulations, which is computationally expensive. Structure-preserving model reduction methods (see [1], [81]) have been recently successfully applied to particle discretizations of the collisionless Vlasov equation (see [98]). It would be of great practical interest to combine our results with model reduction techniques in order to develop new efficient structure-preserving data-driven numerical methods for the collisional Vlasov-Maxwell equations.

Acknowledgements

We would like to thank Christopher Albert, Darryl Holm, Michael Kraus, Omar Maj, Houman Owhadi, Eric Sonnendrücker, and Cesare Tronci for useful comments and references. The study is a contribution to the Reduced Complexity Models grant number ZT-I-0010 funded by the Helmholtz Association of German Research Centers.

A The variations of the action functional S

We will define the variation of S with respect to the variation $\delta\mathbf{X}_a \in C_{\Omega, T}^1$ of the argument \mathbf{X}_a as

$$\delta_{\mathbf{X}_a} S = \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} S[\mathbf{X}_1, \dots, \mathbf{X}_a + \epsilon \delta\mathbf{X}_a, \dots, \mathbf{X}_N, \mathbf{V}_1, \dots, \mathbf{V}_N, \mathbf{P}_1, \dots, \mathbf{P}_N, \varphi, \mathbf{A}]. \quad (\text{A.1})$$

Since the potentials φ and \mathbf{A} are C^2 , and the processes \mathbf{X}_b , \mathbf{V}_b , and \mathbf{P}_b are almost surely continuous, we can use a dominated convergence argument to interchange the differentiation with respect to ϵ and integration with respect to t to obtain

$$\delta_{\mathbf{X}_a} S = \frac{N_{tot}}{N} \int_0^T \left(-q \nabla_x \varphi(\mathbf{X}_a, t) \cdot \delta\mathbf{X}_a + q \sum_{i,j=1}^3 V^j \frac{\partial A^j}{\partial x^i}(\mathbf{X}_a, t) \delta X_a^i + \mathbf{P}_a \cdot \delta \dot{\mathbf{X}}_a \right) dt. \quad (\text{A.2})$$

Since $\delta\mathbf{X}_a$ is almost surely differentiable, we have that its stochastic differential is simply $d\delta\mathbf{X}_a = \delta \dot{\mathbf{X}}_a dt$. Furthermore, both $\delta\mathbf{X}_a$ and \mathbf{P}_a are almost surely continuous semimartingales, therefore using the integration by parts formula for semimartingales (see [84]) we can write

$$\int_0^T \mathbf{P}_a \cdot \delta \dot{\mathbf{X}}_a dt = \int_0^T \mathbf{P}_a \circ d\delta\mathbf{X}_a = \mathbf{P}_a(t) \cdot \delta\mathbf{X}_a(t) \Big|_0^T - \int_0^T \delta\mathbf{X}_a \circ d\mathbf{P}_a, \quad (\text{A.3})$$

where the Stratonovich integrals are understood in the sense that $\int \delta\mathbf{X}_a \circ d\mathbf{P}_a = \sum_i \int \delta X_a^i \circ dP_a^i$. By substituting (A.3) in (A.2), we obtain (4.5a). Variations with respect to $\delta\mathbf{V}_a, \delta\mathbf{P}_a \in C_{\Omega, T}^0$ are

defined analogously to (A.1). Similar computations (note that integration by parts is not necessary) yield (4.5b) and (4.5c), respectively.

The variation of S with respect to the variation $\delta\mathbf{A} \in \mathfrak{X}_0(\mathbb{R}^3)$ of the vector potential \mathbf{A} is defined as

$$\delta_{\mathbf{A}}S = \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} S[\mathbf{X}_1, \dots, \mathbf{X}_N, \mathbf{V}_1, \dots, \mathbf{V}_N, \mathbf{P}_1, \dots, \mathbf{P}_N, \varphi, \mathbf{A} + \epsilon\delta\mathbf{A}]. \quad (\text{A.4})$$

Switching the order of differentiation and integration, integrating by parts, and using the fact that $\delta\mathbf{A}$ is compactly supported, one arrives at (4.5d), where in the derivations we have used (3.2c) and

$$\begin{aligned} \frac{N_{tot}}{N} \sum_{b=1}^N \left[\int_0^T q\mathbf{V}_b(t) \cdot \delta\mathbf{A}(\mathbf{X}_b, t) dt \right] &= \int_0^T \int_{\mathbb{R}^3} \frac{qN_{tot}}{N} \sum_{b=1}^N [q\mathbf{V}_b(t)\delta(\mathbf{x} - \mathbf{X}_b(t))] \cdot \delta\mathbf{A}(\mathbf{x}, t) d^3\mathbf{x}dt \\ &= \int_0^T \int_{\mathbb{R}^3} \mathbf{J}(\mathbf{x}, t) \cdot \delta\mathbf{A}(\mathbf{x}, t) d^3\mathbf{x}dt, \end{aligned} \quad (\text{A.5})$$

and the remaining calculations are standard, and can be found in, e.g., [32], [50]. The variation of S with respect to the variation $\delta\varphi \in \mathfrak{X}_0(\mathbb{R})$ of the scalar potential φ is defined in a similar fashion, and after similar calculations one obtains (4.5e).

B Proof of Lemma 4.1

Proof. Suppose that (4.8) holds. Then (4.7) follows from the associativity property of the Stratonovich integral (see, e.g., the proof of Theorem 2.1 in [49]). Conversely, assume that (4.7) is satisfied, and let us prove that (4.8) follows. Our reasoning very closely follows the proof of Theorem 3.3 in [11]. Pick any time $t \in [0, T]$. We will use \mathbf{e}_1 , \mathbf{e}_2 , and \mathbf{e}_3 to denote the standard Cartesian basis vectors for \mathbb{R}^3 . Pick a basis vector \mathbf{e}_i . The condition (4.7) in particular holds for \mathbf{Z} 's which are C^1 functions of time, i.e., non-random. The main idea of the proof is to construct a one-parameter family of C^1 functions \mathbf{Z}_ϵ which converge to $\mathbb{1}_{[0,t]}\mathbf{e}_i$ as $\epsilon \rightarrow 0$, and show that the integral in (4.7) converges almost surely to the integral in (4.8). Let us introduce the notation

$$I(\mathbf{X}, \mathbf{V}, \mathbf{P}, \mathbf{Z}) = \int_0^T \left(\mathbf{Z}(\tau) \circ d\mathbf{P}(\tau) - \mathbf{R}(\mathbf{X}, \mathbf{V}) \cdot \mathbf{Z}(\tau) d\tau - \sum_{\nu=1}^M \mathbf{r}_\nu(\mathbf{X}, \mathbf{V}) \cdot \mathbf{Z}(\tau) \circ dW^\nu(\tau) \right), \quad (\text{B.1})$$

$$\begin{aligned} I^*(\mathbf{X}, \mathbf{V}, \mathbf{P}) &= \int_0^T \left(\mathbb{1}_{[0,t]}\mathbf{e}_i \circ d\mathbf{P}(\tau) - \mathbf{R}(\mathbf{X}, \mathbf{V}) \cdot \mathbb{1}_{[0,t]}\mathbf{e}_i d\tau - \sum_{\nu=1}^M \mathbf{r}_\nu(\mathbf{X}, \mathbf{V}) \cdot \mathbb{1}_{[0,t]}\mathbf{e}_i \circ dW^\nu(\tau) \right) \\ &= \int_0^t \left(dP^i(\tau) - R^i(\mathbf{X}(\tau), \mathbf{V}(\tau)) d\tau - \sum_{\nu=1}^M r_\nu^i(\mathbf{X}(\tau), \mathbf{V}(\tau)) \circ dW^\nu(\tau) \right). \end{aligned} \quad (\text{B.2})$$

Define the functions $h_1 : [0, \epsilon] \rightarrow [0, 1]$ and $h_2 : [t - \epsilon, t] \rightarrow [0, 1]$ by the formulas

$$h_1(\tau) = 2\frac{\tau}{\epsilon} - \frac{\tau^2}{\epsilon^2}, \quad h_2(\tau) = \begin{cases} -\frac{2}{\epsilon^2}(\tau - t + \epsilon)^2 + 1 & \text{if } t - \epsilon \leq \tau \leq t - \frac{\epsilon}{2}, \\ \frac{2}{\epsilon^2}(\tau - t + \epsilon)^2 - \frac{4}{\epsilon}(\tau - t + \epsilon) + 2 & \text{if } t - \frac{\epsilon}{2} < \tau \leq t. \end{cases} \quad (\text{B.3})$$

Note that $h_1(0) = h_2(t) = 0$, $h_1(\epsilon) = h_2(t - \epsilon) = 1$, and $h_1'(\epsilon) = h_2'(t - \epsilon) = h_2'(t) = 0$. Define further the family of functions \mathbf{Z}_ϵ by the formula

$$\mathbf{Z}_\epsilon(\tau) = \begin{cases} h_1(\tau)\mathbf{e}_i & \text{if } 0 \leq \tau \leq \epsilon, \\ \mathbf{e}_i & \text{if } \epsilon < \tau < t - \epsilon, \\ h_2(\tau)\mathbf{e}_i & \text{if } t - \epsilon \leq \tau \leq t, \\ 0 & \text{if } t < \tau \leq T. \end{cases} \quad (\text{B.4})$$

It is easy to see that \mathbf{Z}_ϵ is continuously differentiable¹ on $[0, T]$, and converges to $\mathbb{1}_{[0, t]}\mathbf{e}_i$ in the L^2 norm as ϵ goes to zero. Using (B.1), (B.2), (B.3), and (B.4), we have

$$\begin{aligned} I^*(\mathbf{X}, \mathbf{V}, \mathbf{P}) - I(\mathbf{X}, \mathbf{V}, \mathbf{P}, \mathbf{Z}_\epsilon) = & \\ & \int_0^\epsilon \left((1 - h_1(\tau)) \circ dP^i(\tau) - (1 - h_1(\tau))R^i(\mathbf{X}, \mathbf{V}) d\tau - \sum_{\nu=1}^M (1 - h_1(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}) \circ dW^\nu \right) \\ & + \int_{t-\epsilon}^t \left((1 - h_2(\tau)) \circ dP^i(\tau) - (1 - h_2(\tau))R^i(\mathbf{X}, \mathbf{V}) d\tau - \sum_{\nu=1}^M (1 - h_2(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}) \circ dW^\nu \right). \end{aligned} \quad (\text{B.5})$$

By definition, the Stratonovich integrals in (B.5) can be expressed in terms of the Itô integrals as

$$\begin{aligned} \int_0^\epsilon (1 - h_1(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}) \circ dW^\nu &= \int_0^\epsilon (1 - h_1(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}) dW^\nu + \frac{1}{2} \left[(1 - h_1(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}), W^\nu(\tau) \right]_0^\epsilon, \\ \int_{t-\epsilon}^t (1 - h_2(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}) \circ dW^\nu &= \int_{t-\epsilon}^t (1 - h_2(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}) dW^\nu + \frac{1}{2} \left[(1 - h_2(\tau))r_\nu^i(\mathbf{X}, \mathbf{V}), W^\nu(\tau) \right]_{t-\epsilon}^t, \end{aligned} \quad (\text{B.6})$$

for each $\nu = 1, \dots, M$, where $[\cdot, \cdot]$ denotes the quadratic covariation process. Since the quadratic covariation of almost surely continuous semimartingales is itself a semimartingale with almost surely continuous paths (see Theorem 23 in Chapter II.6 of [84]), we have that

$$\left[(1 - h_1(\tau))r_\nu^i(\mathbf{X}(\tau), \mathbf{V}(\tau)), W^\nu(\tau) \right]_0^\epsilon \longrightarrow (1 - h_1(0))r_\nu^i(\mathbf{X}(0), \mathbf{V}(0))W^\nu(0) = 0 \quad \text{a.s. as } \epsilon \longrightarrow 0, \quad (\text{B.7})$$

since $W^\nu(0) = 0$ almost surely. In a similar fashion we show

$$\left[(1 - h_2(\tau))r_\nu^i(\mathbf{X}(\tau), \mathbf{V}(\tau)), W^\nu(\tau) \right]_{t-\epsilon}^t \longrightarrow 0 \quad \text{a.s. as } \epsilon \longrightarrow 0. \quad (\text{B.8})$$

Using (B.5) and (B.6), we have the estimate

¹Note that our definition (B.4) is slightly different from the corresponding definition in [11], because the test functions used in [11] are in fact not differentiable at $\tau = t$. This, however, is of little consequence for the rest of the proof.

$$\begin{aligned}
& |I^*(\mathbf{X}, \mathbf{V}, \mathbf{P}) - I(\mathbf{X}, \mathbf{V}, \mathbf{P}, \mathbf{Z}_\epsilon)| \leq \\
& \underbrace{\left| \int_0^\epsilon \left((1 - h_1(\tau)) \circ dP^i(\tau) - (1 - h_1(\tau)) R^i(\mathbf{X}, \mathbf{V}) d\tau - \sum_{\nu=1}^M (1 - h_1(\tau)) r_\nu^i(\mathbf{X}, \mathbf{V}) dW^\nu \right) \right|}_{\Gamma_1} \\
& + \underbrace{\left| \int_{t-\epsilon}^t \left((1 - h_2(\tau)) \circ dP^i(\tau) - (1 - h_2(\tau)) R^i(\mathbf{X}, \mathbf{V}) d\tau - \sum_{\nu=1}^M (1 - h_2(\tau)) r_\nu^i(\mathbf{X}, \mathbf{V}) dW^\nu \right) \right|}_{\Gamma_2} \\
& + \frac{1}{2} \sum_{\nu=1}^M \left| \left[(1 - h_1(\tau)) r_\nu^i(\mathbf{X}, \mathbf{V}), W^\nu(\tau) \right]_0^\epsilon + \frac{1}{2} \sum_{\nu=1}^M \left| \left[(1 - h_2(\tau)) r_\nu^i(\mathbf{X}, \mathbf{V}), W^\nu(\tau) \right]_{t-\epsilon}^t \right| \right|. \quad (\text{B.9})
\end{aligned}$$

By bounding the integrands and using the Itô isometry theorem, it is shown in [11] that $\Gamma_1 \rightarrow 0$ and $\Gamma_2 \rightarrow 0$ in mean-square as $\epsilon \rightarrow 0$, and consequently, by invoking the Borel-Cantelli lemma, there exists a subsequence (ϵ_n) such that $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$, for which $\Gamma_1 \rightarrow 0$ and $\Gamma_2 \rightarrow 0$ almost surely. Together with (B.7) and (B.8), this means that $I(\mathbf{X}, \mathbf{V}, \mathbf{P}, \mathbf{Z}_{\epsilon_n}) \rightarrow I^*(\mathbf{X}, \mathbf{V}, \mathbf{P})$ almost surely. Given the assumption (4.7), we have that $I^*(\mathbf{X}, \mathbf{V}, \mathbf{P}) = 0$ almost surely, which completes the proof. \square

References

- [1] B. Afkham and J. Hesthaven. Structure preserving model reduction of parametric Hamiltonian systems. *SIAM Journal on Scientific Computing*, 39(6):A2616–A2644, 2017.
- [2] E. Allen and H. Victory. A computational investigation of the random particle method for numerical solution of the kinetic Vlasov-Poisson-Fokker-Planck equations. *Physica A: Statistical Mechanics and its Applications*, 209(3):318 – 346, 1994.
- [3] M. Arnaudon, X. Chen, and A. B. Cruzeiro. Stochastic Euler-Poincaré reduction. *Journal of Mathematical Physics*, 55(8):081507, 2014.
- [4] R. Balescu, H. Wang, and J. H. Misguich. Langevin equation versus kinetic equation: Subdiffusive behavior of charged particles in a stochastic magnetic field. *Physics of Plasmas*, 1(12):3826–3842, 1994.
- [5] J. W. Banks, S. Brunner, R. L. Berger, and T. M. Tran. Vlasov simulations of electron-ion collision effects on damping of electron plasma waves. *Physics of Plasmas*, 23(3):032108, 2016.
- [6] C. Birdsall and A. Langdon. *Plasma Physics via Computer Simulation*. Series in Plasma Physics and Fluid Dynamics. Taylor & Francis, New York, London, 2004.
- [7] J. Bismut. Mécanique aléatoire. In P. Hennequin, editor, *Ecole d’Eté de Probabilités de Saint-Flour X - 1980*, volume 929 of *Lecture Notes in Mathematics*, pages 1–100. Springer Berlin Heidelberg, 1982.

- [8] A. V. Bobylev and K. Nanbu. Theory of collision algorithms for gases and plasmas based on the Boltzmann equation and the Landau-Fokker-Planck equation. *Phys. Rev. E*, 61:4576–4586, 2000.
- [9] A. Bottino and E. Sonnendrücker. Monte Carlo particle-in-cell methods for the simulation of the Vlasov–Maxwell gyrokinetic equations. *Journal of Plasma Physics*, 81(5):435810501, 2015.
- [10] N. Bou-Rabee and H. Owhadi. Stochastic variational partitioned Runge-Kutta integrators for constrained systems. Unpublished, arXiv:0709.2222, 2007.
- [11] N. Bou-Rabee and H. Owhadi. Stochastic variational integrators. *IMA Journal of Numerical Analysis*, 29(2):421–443, 2009.
- [12] N. Bou-Rabee and H. Owhadi. Long-run accuracy of variational integrators in the stochastic context. *SIAM J. Numer. Anal.*, 48(1):278–297, 2010.
- [13] B. J. Braams and C. F. F. Karney. Conductivity of a relativistic plasma. *Physics of Fluids B: Plasma Physics*, 1(7):1355–1368, 1989.
- [14] A. J. Brizard. Variational principle for nonlinear gyrokinetic Vlasov–Maxwell equations. *Physics of Plasmas*, 7(12):4816–4822, 2000.
- [15] A. J. Brizard and C. Tronci. Variational formulations of guiding-center Vlasov-Maxwell theory. *Physics of Plasmas*, 23(6):062107, 2016.
- [16] M. G. Cadjan and M. F. Ivanov. Langevin approach to plasma kinetics with Coulomb collisions. *Journal of Plasma Physics*, 61(1):89–106, 1999.
- [17] J. Callen, K. Bunkers, and J. Kollasch. Plasma kinetic theory. Lecture notes, <http://homepages.cae.wisc.edu/~callen/PlasmaKineticTheory.pdf>, 2018.
- [18] J. R. Cary and A. J. Brizard. Hamiltonian theory of guiding-center motion. *Rev. Mod. Phys.*, 81:693–738, 2009.
- [19] X. Chen, A. B. Cruzeiro, and T. S. Ratiu. Stochastic variational principles for dissipative equations with advected quantities. Unpublished, arXiv:1506.05024, 2018.
- [20] P. Clemmow and J. Dougherty. *Electrodynamics of Particles and Plasmas*. A-W series in advanced physics. Addison-Wesley Publishing Company, 1969.
- [21] B. I. Cohen, A. M. Dimits, A. Friedman, and R. E. Caflisch. Time-step considerations in particle simulation algorithms for Coulomb collisions in plasmas. *IEEE Transactions on Plasma Science*, 38(9):2394–2406, 2010.
- [22] F. Cooper, C. Lucheroni, H. Shepard, and P. Sodano. Variational method for studying solitons in the Korteweg-de Vries equation. *Physics Letters A*, 173(1):33 – 36, 1993.
- [23] C. J. Cotter, G. A. Gottwald, and D. D. Holm. Stochastic partial differential fluid equations as a diffusive limit of deterministic Lagrangian multi-time dynamics. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 473(2205), 2017.

- [24] D. Crisan and D. D. Holm. Wave breaking for the stochastic Camassa-Holm equation. *Physica D: Nonlinear Phenomena*, 376-377:138 – 143, 2018. Special Issue: Nonlinear Partial Differential Equations in Mathematical Fluid Dynamics.
- [25] A. B. Cruzeiro. Stochastic approaches to deterministic fluid dynamics: A selective review. *Water*, 12(3), 2020.
- [26] P. Degond. Global existence of smooth solutions for the Vlasov-Fokker-Planck equation in 1 and 2 space dimensions. *Annales scientifiques de l'École Normale Supérieure*, Ser. 4, 19(4):519–542, 1986.
- [27] P. Degond and H. Neunzert. Local existence of solutions of the Vlasov-Maxwell equations and convergence to the Vlasov-Poisson equations for infinite light velocity. *Mathematical Methods in the Applied Sciences*, 8(1):533–558, 1986.
- [28] A. Dimits, B. Cohen, R. Caflisch, M. Rosin, and L. Ricketson. Higher-order time integration of Coulomb collisions in a plasma using Langevin equations. *Journal of Computational Physics*, 242:561 – 580, 2013.
- [29] J. P. Dougherty. Model Fokker-Planck equation for a plasma and its solution. *The Physics of Fluids*, 7(11):1788–1799, 1964.
- [30] J. P. Dougherty and S. R. Watson. Model Fokker-Planck equations: Part 2. The equation for a multicomponent plasma. *Journal of Plasma Physics*, 1(3):317–326, 1967.
- [31] R. Duan and R. M. Strain. Optimal large-time behavior of the Vlasov-Maxwell-Boltzmann system in the whole space. *Communications on Pure and Applied Mathematics*, 64(11):1497–1546, 2011.
- [32] L. Evans. *Partial Differential Equations*. Graduate studies in mathematics. American Mathematical Society, 2010.
- [33] E. G. Evstatiev and B. A. Shadwick. Variational formulation of particle algorithms for kinetic plasma simulations. *Journal of Computational Physics*, 245:376–398, 2013.
- [34] F. Filbet and E. Sonnendrücker. Comparison of Eulerian Vlasov solvers. *Computer Physics Communications*, 150(3):247–266, 2003.
- [35] T. Frank. Single particle dynamics of many-body systems described by Vlasov-Fokker-Planck equations. *Physics Letters A*, 319(1):173 – 180, 2003.
- [36] T. Frank. *Nonlinear Fokker-Planck Equations: Fundamentals and Applications*. Springer Series in Synergetics. Springer Berlin Heidelberg, 2006.
- [37] Y. Fu, X. Zhang, and H. Qin. An explicitly solvable energy-conserving algorithm for pitch-angle scattering in magnetized plasmas. Unpublished, arXiv:2010.12920, 2020.
- [38] C. Gardiner. *Stochastic Methods: A Handbook for the Natural and Social Sciences*. Springer Series in Synergetics. Springer Berlin Heidelberg, 2009.
- [39] F. Gay-Balmaz and D. D. Holm. Stochastic geometric models with non-stationary spatial correlations in Lagrangian fluid flows. *J. Nonlinear Sci.*, 28:873–904, 2018.

- [40] A. Hakim, M. Francisquez, J. Juno, and G. W. Hammett. Conservative discontinuous Galerkin schemes for nonlinear Dougherty-Fokker-Planck collision operators. *Journal of Plasma Physics*, 86(4):905860403, 2020.
- [41] J. Hall and M. Leok. Spectral variational integrators. *Numer. Math.*, 130(4):681–740, Aug 2015.
- [42] K. J. Havlak and H. D. Victory. The numerical analysis of random particle methods applied to Vlasov-Poisson-Fokker-Planck kinetic equations. *SIAM Journal on Numerical Analysis*, 33(1):291–317, 1996.
- [43] E. Hirvijoki, M. Kraus, and J. W. Burby. Metriplectic particle-in-cell integrators for the Landau collision operator. Preprint arXiv:1802.05263, 2018.
- [44] R. Hockney and J. Eastwood. *Computer Simulation Using Particles*. CRC Press, 1988.
- [45] D. D. Holm. Variational principles for stochastic fluid dynamics. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 471(2176), 2015.
- [46] D. D. Holm. Uncertainty quantification for fluids versus electromagnetic waves. Preprint arXiv:1705.07645, 2017.
- [47] D. D. Holm and T. M. Tyranowski. Variational principles for stochastic soliton dynamics. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 472(2187), 2016.
- [48] D. D. Holm and T. M. Tyranowski. New variational and multisymplectic formulations of the Euler–Poincaré equation on the Virasoro–Bott group using the inverse map. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 474(2213), 2018.
- [49] D. D. Holm and T. M. Tyranowski. Stochastic discrete Hamiltonian variational integrators. *BIT Numerical Mathematics*, 58(4):1009–1048, 2018.
- [50] J. Jackson. *Classical Electrodynamics, 3rd ed.* John Wiley & Sons, New York, 1999.
- [51] L. O. Jay. Structure preservation for constrained dynamics with super partitioned additive Runge–Kutta methods. *SIAM Journal on Scientific Computing*, 20(2):416–446, 1998.
- [52] C. Kane, J. E. Marsden, M. Ortiz, and M. West. Variational integrators and the Newmark algorithm for conservative and dissipative mechanical systems. *International Journal for Numerical Methods in Engineering*, 49(10):1295–1325, 2000.
- [53] C. F. Karney. Fokker-Planck and quasilinear codes. *Computer Physics Reports*, 4(3):183 – 244, 1986.
- [54] R. Kleiber, R. Hatzky, A. Könies, K. Kauffmann, and P. Helander. An improved control-variate scheme for particle-in-cell simulations with collisions. *Computer Physics Communications*, 182(4):1005 – 1012, 2011.
- [55] P. Kloeden and E. Platen. *Numerical Solution of Stochastic Differential Equations*. Applications of Mathematics : Stochastic Modelling and Applied Probability. Springer, 1995.

- [56] M. Kraus. *Variational integrators in plasma physics*. PhD thesis, Technische Universität München, 2013.
- [57] M. Kraus and E. Hirvijoki. Metriplectic integrators for the Landau collision operator. *Physics of Plasmas*, 24(10):102311, 2017.
- [58] M. Kraus, K. Kormann, P. Morrison, and E. Sonnendrücker. GEMPIC: geometric electromagnetic particle-in-cell methods. *Journal of Plasma Physics*, 83(4):905830401, 2017.
- [59] M. Kraus and T. M. Tyranowski. Variational integrators for stochastic dissipative Hamiltonian systems. *IMA Journal of Numerical Analysis*, 07 2020. draa022.
- [60] S. Lall and M. West. Discrete variational Hamiltonian mechanics. *Journal of Physics A: Mathematical and General*, 39(19):5509–5519, 2006.
- [61] J. A. Lázaro-Camí and J. P. Ortega. Stochastic Hamiltonian Dynamical Systems. *Reports on Mathematical Physics*, 61(1):65 – 122, 2008.
- [62] D. S. Lemons, D. Winske, W. Daughton, and B. Albright. Small-angle Coulomb collision model for particle-in-cell simulations. *Journal of Computational Physics*, 228(5):1391 – 1403, 2009.
- [63] A. Lenard and I. B. Bernstein. Plasma oscillations with diffusion in velocity space. *Phys. Rev.*, 112:1456–1459, Dec 1958.
- [64] M. Leok and T. Shingel. General techniques for constructing variational integrators. *Frontiers of Mathematics in China*, 7(2):273–303, 2012.
- [65] M. Leok and J. Zhang. Discrete Hamiltonian variational integrators. *IMA Journal of Numerical Analysis*, 31(4):1497–1532, 2011.
- [66] A. Lew, J. E. Marsden, M. Ortiz, and M. West. Asynchronous variational integrators. *Archive for Rational Mechanics and Analysis*, 167(2):85–146, 2003.
- [67] H. R. Lewis. Energy-conserving numerical approximations for Vlasov plasmas. *Journal of Computational Physics*, 6(1):136–141, 1970.
- [68] H. R. Lewis. Variational algorithms for numerical simulation of collisionless plasma with point particles including electromagnetic interactions. *Journal of Computational Physics*, 10(3):400 – 419, 1972.
- [69] F. E. Low. A Lagrangian formulation of the Boltzmann-Vlasov equation for plasmas. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 248(1253):282–287, 1958.
- [70] W. M. Manheimer, M. Lampe, and G. Joyce. Langevin representation of Coulomb collisions in PIC simulations. *Journal of Computational Physics*, 138(2):563 – 584, 1997.
- [71] J. E. Marsden, G. W. Patrick, and S. Shkoller. Multisymplectic geometry, variational integrators, and nonlinear PDEs. *Communications in Mathematical Physics*, 199(2):351–395, 1998.

- [72] J. E. Marsden and M. West. Discrete mechanics and variational integrators. *Acta Numerica*, 10(1):357–514, 2001.
- [73] D. Montgomery and D. Tidman. *Plasma Kinetic Theory*. McGraw-Hill Advanced Physics Monograph Series. McGraw-Hill, New York, 1964.
- [74] H. Neunzert, M. Pulvirenti, and L. Triolo. On the Vlasov-Fokker-Planck equation. *Mathematical Methods in the Applied Sciences*, 6(1):527–538, 1984.
- [75] S. Ober-Blöbaum. Galerkin variational integrators and modified symplectic Runge-Kutta methods. *IMA Journal of Numerical Analysis*, 37(1):375–406, 2017.
- [76] S. Ober-Blöbaum and N. Saake. Construction and analysis of higher order Galerkin variational integrators. *Advances in Computational Mathematics*, 41(6):955–986, 2015.
- [77] R. S. B. Ong and M. Y. Yu. The effect of weak collisions on ion-acoustic wave instabilities in a current-carrying plasma. *Journal of Plasma Physics*, 3(3):425–433, 1969.
- [78] K. Ono. Global existence of regular solutions for the Vlasov–Poisson–Fokker–Planck system. *Journal of Mathematical Analysis and Applications*, 263(2):626–636, 2001.
- [79] A. Oppenheim. Wave motion in a plasma based on a Fokker-Planck equation. *The Physics of Fluids*, 8(5):900–911, 1965.
- [80] D. Pavlov, P. Mullen, Y. Tong, E. Kanso, J. E. Marsden, and M. Desbrun. Structure-preserving discretization of incompressible fluids. *Physica D: Nonlinear Phenomena*, 240(6):443–458, 2011.
- [81] L. Peng and K. Mohseni. Symplectic model reduction of Hamiltonian systems. *SIAM Journal on Scientific Computing*, 38(1):A1–A27, 2016.
- [82] D. Pfirsch. New variational formulation of Maxwell-Vlasov and guiding center theories local charge and energy conservation laws. *Zeitschrift für Naturforschung A*, 39(1):1–8, 1984.
- [83] D. Pfirsch and P. J. Morrison. Local conservation laws for the Maxwell-Vlasov and collisionless kinetic guiding-center theories. *Phys. Rev. A*, 32:1714–1721, Sep 1985.
- [84] P. Protter. *Stochastic Integration and Differential Equations*. Stochastic Modelling and Applied Probability. Springer Berlin Heidelberg, 2005.
- [85] H. Risken and T. Frank. *The Fokker-Planck Equation: Methods of Solution and Applications*. Springer Series in Synergetics. Springer Berlin Heidelberg, 1996.
- [86] M. N. Rosenbluth, W. M. MacDonald, and D. L. Judd. Fokker-Planck equation for an inverse-square force. *Physical Review*, 107(1):1–6, 1957.
- [87] C. W. Rowley and J. E. Marsden. Variational integrators for degenerate Lagrangians, with application to point vortices. In *Decision and Control, 2002, Proceedings of the 41st IEEE Conference on*, volume 2, pages 1521–1527. IEEE, 2002.
- [88] B. A. Shadwick, A. B. Stamm, and E. G. Evstatiev. Variational formulation of macro-particle plasma simulation algorithms. *Physics of Plasmas*, 21(5):055708, 2014.

- [89] M. Sherlock. A Monte-Carlo method for Coulomb collisions in hybrid plasma models. *J. Comput. Phys.*, 227(4):2286–2292, 2008.
- [90] E. Sonnendrücker, A. Wachter, R. Hatzky, and R. Kleiber. A split control variate scheme for PIC simulations with collisions. *Journal of Computational Physics*, 295:402 – 419, 2015.
- [91] J. Squire, H. Qin, and W. M. Tang. Geometric integration of the Vlasov–Maxwell system with a variational particle-in-cell scheme. *Physics of Plasmas*, 19:084501, 2012.
- [92] A. B. Stamm, B. A. Shadwick, and E. G. Evstatiev. Variational formulation of macroparticle models for electromagnetic plasma simulations. *IEEE Transactions on Plasma Science*, 42(6):1747–1758, 2014.
- [93] A. Stern, Y. Tong, M. Desbrun, and J. E. Marsden. Variational integrators for Maxwell’s equations with sources. *PIERS Online*, 4(7):711–715, 2008.
- [94] R. M. Strain. The Vlasov–Maxwell–Boltzmann system in the whole space. *Communications in Mathematical Physics*, 268:543–567, 2006.
- [95] H. Sugama. Gyrokinetic field theory. *Physics of Plasmas*, 7(2):466–480, 2000.
- [96] T. M. Tyranowski and M. Desbrun. R-adaptive multisymplectic and variational integrators. *Mathematics*, 7(7), 2019.
- [97] T. M. Tyranowski and M. Desbrun. Variational partitioned Runge–Kutta methods for Lagrangians linear in velocities. *Mathematics*, 7(9), 2019.
- [98] T. M. Tyranowski and M. Kraus. Symplectic model reduction methods for the Vlasov equation. Submitted. Preprint arXiv:1910.06026, 2021.
- [99] E. vanden Eijnden and A. Grecos. Stochastic modelling of turbulence and anomalous transport in plasmas. *Journal of Plasma Physics*, 59(4):683–694, June 1998.
- [100] J. Vankerschaver and M. Leok. A novel formulation of point vortex dynamics on the sphere: geometrical and numerical aspects. *J. Nonlin. Sci.*, 24(1):1–37, 2014.
- [101] L. Wang. *Variational Integrators and Generating Functions for Stochastic Hamiltonian Systems*. PhD thesis, Karlsruhe Institute of Technology, 2007.
- [102] X. Wang. The Vlasov–Maxwell–Fokker–Planck system near Maxwellians in \mathbb{R}^3 . *Applicable Analysis*, 0(0):1–28, 2019.
- [103] X. Wang. Global existence and long-time behavior of solutions to the Vlasov-Poisson-Fokker-Planck system. *Acta Applicandae Mathematicae*, 170:853–881, 2020.
- [104] S. Wollman. An existence and uniqueness theorem for the Vlasov-Maxwell system. *Communications on Pure and Applied Mathematics*, 37(4):457–462, 1984.
- [105] S. Wollman. Local existence and uniqueness theory of the Vlasov-Maxwell system. *Journal of Mathematical Analysis and Applications*, 127(1):103–121, 1987.

- [106] J. Xiao, J. Liu, H. Qin, and Z. Yu. A variational multi-symplectic particle-in-cell algorithm with smoothing functions for the Vlasov-Maxwell system. *Physics of Plasmas*, 20(10):102517, 2013.
- [107] J. Xiao, H. Qin, and J. Liu. Structure-preserving geometric particle-in-cell methods for Vlasov-Maxwell systems. *Plasma Science and Technology*, 20(11):110501, sep 2018.
- [108] H. Ye and P. J. Morrison. Action principles for the Vlasov equation. *Physics of Fluids B: Plasma Physics*, 4(4):771–777, 1992.
- [109] H. Yoshimura and J. E. Marsden. Dirac structures in Lagrangian mechanics Part II: Variational structures. *Journal of Geometry and Physics*, 57(1):209 – 250, 2006.
- [110] X. Zhang, Y. Fu, and H. Qin. Simulating pitch angle scattering using an explicitly solvable energy-conserving algorithm. *Phys. Rev. E*, 102:033302, Sep 2020.