TIME-DECAY ESTIMATES FOR LINEARIZED TWO-PHASE NAVIER-STOKES EQUATIONS WITH SURFACE TENSION AND GRAVITY

HIROKAZU SAITO

ABSTRACT. The aim of this paper is to show time-decay estimates of solutions to linearized two-phase Navier-Stokes equations with surface tension and gravity. The original two-phase Navier-Stokes equations describe the two-phase incompressible viscous flow with a sharp interface that is close to the hyperplane $x_N=0$ in the N-dimensional Euclidean space, $N\geq 2$. It is well-known that the Rayleigh-Taylor instability occurs when the upper fluid is heavier than the lower one, while this paper assumes that the lower fluid is heavier than the upper one and proves time-decay estimates of L_P-L_q type for the linearized equations. Our approach is based on solution formulas, given by Shibata and Shimizu (2011), for a resolvent problem associated with the linearized equations.

1. Introduction

Let us consider the motion of two immiscible, viscous, incompressible capillary fluids, $fluid_+$ and $fluid_-$, in the N-dimensional Euclidean space \mathbf{R}^N for $N \geq 2$. Here fluid₊ and fluid₋ occupy $\Omega_+(t)$ and $\Omega_-(t)$, respectively, which are given by

$$\Omega_{\pm}(t) = \{(x', x_N) : x' = (x_1, \dots, x_{N-1}) \in \mathbf{R}^{N-1}, \pm (x_N - H(x', t)) > 0\}$$

for time t > 0 and the so-called height function H = H(x', t). The fluids are thus separated by the interface

$$\Gamma(t) = \{(x', x_N) : x' = (x_1, \dots, x_{N-1}) \in \mathbf{R}^{N-1}, x_N = H(x', t)\}.$$

We denote the density of fluid $_{\pm}$ by ρ_{\pm} , while the viscosity coefficient of fluid $_{\pm}$ by μ_{\pm} . Suppose that ρ_{\pm} and μ_{\pm} are positive constants throughout this paper. That motion of two fluids is governed by the two-phase Navier-Stokes equations where surface tension is included on the interface. In addition, we allow for gravity to act on the fluids. The two-phase Navier-Stokes equations was studied by Prüss and Simonett [4], and they proved that the Rayleigh-Taylor instability occurs in an L_p -setting when the upper fluid is heavier than the lower one, i.e., $\rho_+ > \rho_-$. In the present paper, we assume that the lower fluid is heavier than the upper one, i.e., $\rho_- > \rho_+$, and show time-decay estimates of $L_p - L_q$ type for some linearized system as the first step in proving global existence results for the two-phase Navier-Stokes equations when $\rho_- > \rho_+$.

 $^{2010\ \}textit{Mathematics Subject Classification}.\ \text{Primary: 35Q30; Secondary: 76D07}.$

Key words and phrases. Two-phase flow, semigroup, L_p-L_q decay estimate, surface tension, gravity.

This research was partly supported by JSPS KAKENHI Grant Number JP17K14224.

¹The height function is unknown. This needs to be determined as part of the problem.

Let us define $\dot{\mathbf{R}}^N = \mathbf{R}_+^N \cup \mathbf{R}_-^N$ for

$$\mathbf{R}_{\pm}^{N} = \{(x', x_N) : x' = (x_1, \dots, x_{N-1}) \in \mathbf{R}^{N-1}, \pm x_N > 0\}.$$

This paper is concerned with the following linearized system of the two-phase Navier-Stokes equations:

$$\begin{cases}
\partial_t H - U_N|_{x_N=0} = 0 & \text{on } \mathbf{R}^{N-1} \times (0, \infty), \\
\rho \partial_t \mathbf{U} - \text{Div}(\mu \mathbf{D}(\mathbf{U}) - P\mathbf{I}) = 0 & \text{in } \dot{\mathbf{R}}^N \times (0, \infty), \\
\text{div } \mathbf{U} = 0 & \text{in } \dot{\mathbf{R}}^N \times (0, \infty), \\
-\llbracket (\mu \mathbf{D}(\mathbf{U}) - P\mathbf{I}) \mathbf{e}_N \rrbracket + (\omega - \sigma \Delta') H \mathbf{e}_N = 0 & \text{on } \mathbf{R}^{N-1} \times (0, \infty), \\
\llbracket \mathbf{U} \rrbracket = 0 & \text{on } \mathbf{R}^{N-1} \times (0, \infty), \\
H(x', 0) = d(x') (x' \in \mathbf{R}^{N-1}), \quad \mathbf{U}(x, 0) = \mathbf{f}(x) (x \in \dot{\mathbf{R}}^N),
\end{cases} \tag{1.1}$$

where σ is a positive constant called the surface tension coefficient and one has set for the indicator function $\mathbb{1}_A$ of $A \subset \mathbf{R}^N$

$$\rho = \rho_+ \mathbb{1}_{\mathbf{R}^N_+} + \rho_- \mathbb{1}_{\mathbf{R}^N}, \quad \mu = \mu_+ \mathbb{1}_{\mathbf{R}^N_+} + \mu_- \mathbb{1}_{\mathbf{R}^N}.$$

Here $\mathbf{U} = \mathbf{U}(x,t) = (U_1(x,t),\ldots,U_N(x,t))^{\mathsf{T}2}$ and P = P(x,t) respectively denote the velocity field of the fluid and the pressure field of the fluid at position $x \in \dot{\mathbf{R}}^N$ and time t > 0, while d = d(x') and $\mathbf{f} = \mathbf{f}(x) = (f_1(x),\ldots,f_N(x))^{\mathsf{T}}$ are given initial data. Note that $\mathbf{e}_N = (0,\ldots,0,1)^{\mathsf{T}}$ and \mathbf{I} is the $N \times N$ identity matrix. Let $\partial_j = \partial/\partial x_j$ for $j = 1,\ldots,N$. Then

$$\operatorname{div} \mathbf{U} = \sum_{j=1}^{N} \partial_{j} U_{j}, \quad \Delta' H = \sum_{j=1}^{N-1} \partial_{j}^{2} H,$$

while $\mathbf{D}(\mathbf{U})$ is an $N \times N$ matrix whose (i,j) element is given by $\partial_i U_j + \partial_j U_i$. In addition, for matrix-valued functions $\mathbf{M} = (M_{ij}(x))$,

Div
$$\mathbf{M} = \left(\sum_{j=1}^{N} \partial_{j} M_{1j}, \dots, \sum_{j=1}^{N} \partial_{j} M_{Nj}\right)^{\mathsf{T}}.$$

Let f = f(x) be a function defined on $\dot{\mathbf{R}}^N$. Then $[\![f]\!] = [\![f]\!](x')$ denotes the jump of the quantity f across the interface $\mathbf{R}_0^N = \{(x', x_N) : x' \in \mathbf{R}^{N-1}, x_N = 0\}$, that is,

$$[\![f]\!] = [\![f]\!](x') = f(x', 0+) - f(x', 0-),$$

where $f(x', 0\pm) = \lim_{x_N \to 0, \pm x_N > 0} f(x', x_N)$. Note that $[\![\mathbf{U}]\!] = 0$ on \mathbf{R}^{N-1} implies $U_N|_{x_N=0} = U_N(x', 0+) = U_N(x', 0-)$. For the acceleration of gravity $\gamma_a > 0$, the constant ω is given by

$$\omega = -\llbracket \rho \rrbracket \gamma_a = (\rho_- - \rho_+) \gamma_a,$$

which is positive when $\rho_{-} > \rho_{+}$.

The local well-posedness for the two-phase Navier-Stokes equations with $\Gamma(t)$ as above was proved in Prüss and Simonett [5, 6]. Note that the local well-posedness holds for any positive constants ρ_{\pm} , that is, the condition $\rho_{-} > \rho_{+}$ is not required. Those results were extended to a class of non-Newtonian fluids in [1]. In addition, [8] considered the two-phase inhomogeneous incompressible viscous flow without

 $^{{}^{2}\}mathbf{M}^{\mathsf{T}}$ denotes the transpose of \mathbf{M} .

surface tension when gravity is not taken into account, and proved the local well-posedness in general domains including the above-mentioned $\Omega_{\pm}(t)$. If $\Omega_{\pm}(t)$ are assumed to be layer-like domains, then it is known that the global well-posedness holds when $\rho_{-} > \rho_{+}$. In fact, it was shown in [11] in a horizontally periodic setting, and also we refer to [12].

Let us recall Shibata and Shimizu [10]. They considered the following two resolvent problems:

$$\begin{cases} \lambda \eta - u_N|_{x_N=0} = d & \text{on } \mathbf{R}^{N-1}, \\ \rho \lambda \mathbf{u} - \text{Div}(\mu \mathbf{D}(\mathbf{u}) - \mathfrak{p} \mathbf{I}) = \rho \mathbf{f} & \text{in } \dot{\mathbf{R}}^N, \\ \text{div } \mathbf{u} = 0 & \text{in } \dot{\mathbf{R}}^N, \\ -\llbracket (\mu \mathbf{D}(\mathbf{u}) - \mathfrak{p} \mathbf{I}) \mathbf{e}_N \rrbracket + (\omega - \sigma \Delta') \eta \mathbf{e}_N = 0 & \text{on } \mathbf{R}^{N-1}, \\ \llbracket \mathbf{u} \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}, \end{cases}$$
(1.2)

and also

$$\begin{cases}
\rho \lambda \mathbf{v} - \operatorname{Div}(\mu \mathbf{D}(\mathbf{v}) - \mathfrak{q} \mathbf{I}) = \rho \mathbf{f} & \text{in } \dot{\mathbf{R}}^{N}, \\
\operatorname{div} \mathbf{v} = 0 & \text{in } \dot{\mathbf{R}}^{N}, \\
\llbracket (\mu \mathbf{D}(\mathbf{v}) - \mathfrak{q} \mathbf{I}) \mathbf{e}_{N} \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}, \\
\llbracket \mathbf{v} \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}.
\end{cases} \tag{1.3}$$

We define the sector

$$\Sigma_{\varepsilon} = \{ \lambda \in \mathbf{C} \setminus \{0\} : |\arg \lambda| \le \pi - \varepsilon \} \quad (0 < \varepsilon < \pi/2),$$

where \mathbf{C} is the set of all complex numbers. Let $q \in (1,\infty)$ and $(d,\mathbf{f}) \in X_q := W_q^{2-1/q}(\mathbf{R}^{N-1}) \times L_q(\dot{\mathbf{R}}^N)^N$. In [10], they obtained the following results: there exists a constant $\lambda_0(\varepsilon) \geq 1$ such that, for any $\lambda \in \Sigma_\varepsilon$ with $|\lambda| \geq \lambda_0(\varepsilon)$, (1.2) admits a unique solution $(\eta, \mathbf{u}, \mathfrak{p}) \in W_q^{3-1/q}(\mathbf{R}^{N-1}) \times H_q^2(\dot{\mathbf{R}}^N)^N \times \hat{H}_q^1(\dot{\mathbf{R}}^N)$, which satisfies

$$\|(\lambda \eta, \nabla' \eta)\|_{W_q^{2-1/q}(\mathbf{R}^{N-1})} + \|(\lambda \mathbf{u}, \lambda^{1/2} \nabla \mathbf{u}, \nabla^2 \mathbf{u}, \nabla \mathfrak{p})\|_{L_q(\dot{\mathbf{R}}^N)} \le C_{N,q,\varepsilon,\lambda_0(\varepsilon)} \|(d, \mathbf{f})\|_{X_q}, \tag{1.4}$$

where

$$\nabla' \eta = (\partial_1 \eta, \dots, \partial_{N-1} \eta)^\mathsf{T}, \qquad \nabla \mathfrak{p} = (\partial_1 \mathfrak{p}, \dots, \partial_{N-1} \mathfrak{p}, \partial_N \mathfrak{p})^\mathsf{T},$$
$$\nabla \mathbf{u} = \{\partial_j u_k : j, k = 1, \dots, N\}, \quad \nabla^2 \mathbf{u} = \{\partial_j \partial_k u_l : j, k, l = 1, \dots, N\};$$

for any $\lambda \in \Sigma_{\varepsilon}$, (1.3) admits a unique solution $(\mathbf{v}, \mathfrak{q}) \in H_q^2(\dot{\mathbf{R}}^N)^N \times \widehat{H}_q^1(\dot{\mathbf{R}}^N)$, which satisfies

$$\|(\lambda \mathbf{v}, \lambda^{1/2} \nabla \mathbf{v}, \nabla^2 \mathbf{v}, \nabla \mathfrak{q})\|_{L_q(\dot{\mathbf{R}}^N)} \le C_{N,q,\varepsilon} \|\mathbf{f}\|_{L_q(\dot{\mathbf{R}}^N)}. \tag{1.5}$$

These results hold for any $\rho_{\pm} > 0$ and play a key role in proving time-decay estimates of solutions for (1.1) in the present paper.

This paper is organized as follows: The next section introduces the notation used throughout this paper and states the main results of this paper, that is, time-decay estimates of solutions for (1.1). Section 3 gives the representation formulas of solutions for (1.1) by means of the partial Fourier transform with respect to $x' = (x_1, \ldots, x_{N-1})$ and its inverse transform. Section 4 analyzes the boundary symbol appearing in the representation formulas given in Section 3. Section 5 shows our main result stated in Section 2 for a low frequency part. Section 6 shows our main result stated in Section 2 for a high frequency part.

2. Notation and main results

2.1. **Notation.** First, we introduce function spaces. Let X be a Banach space. Then X^m , $m \geq 2$, stands for the m-product space of X, while the norm of X^m is usually denoted by $\|\cdot\|_X$ instead of $\|\cdot\|_{X^m}$ for the sake of simplicity. For another Banach space Y, we set $\|u\|_{X\cap Y} = \|u\|_X + \|u\|_Y$. Let \mathbf{N} be the set of all natural numbers and $\mathbf{N}_0 = \mathbf{N} \cup \{0\}$. Let $p \geq 1$ or $p = \infty$. For an open set $G \subset \mathbf{R}^M$, $M \geq 1$, the Lebesgue spaces on G are denoted by $L_p(G)$ with norm $\|\cdot\|_{L_p(G)}$, while the Sobolev spaces on G are denoted by $H_p^n(G)$, $n \in \mathbf{N}$, with norm $\|\cdot\|_{H_p^n(G)}$. Let $H_p^0(G) = L_p(G)$. In addition, $C_0^\infty(G)$ is the set of all functions in $C^\infty(G)$ whose supports are compact and contained in G, and $C^\infty(I,X)$ is the set of all C^∞ functions on an interval $I \subset \mathbf{R}$ with value X. For any multi-index $\alpha = (\alpha_1, \ldots, \alpha_M) \in \mathbf{N}_0^M$,

$$\partial^{\alpha} u = \partial_{x}^{\alpha} u = \frac{\partial^{|\alpha|} u(x)}{\partial x_{1}^{\alpha_{1}} \dots \partial x_{M}^{\alpha_{M}}} \quad \text{with } |\alpha| = \alpha_{1} + \dots + \alpha_{M}.$$

Let $q \in (1, \infty)$ and $\widehat{H}_q^1(G) = \{u \in L_{q, loc}(\overline{G}) : \partial^{\alpha} u \in L_q(G) \text{ for } |\alpha| = 1\}$. Let $s \in [0, \infty) \setminus \mathbf{N}$ and [s] be the largest integer less than s. The Sobolev-Slobodeckij spaces on \mathbf{R}^{N-1} are defined by

$$\begin{split} W_q^s(\mathbf{R}^{N-1}) &= \{ u \in L_q(\mathbf{R}^{N-1}) : \|u\|_{W_q^s(\mathbf{R}^{N-1})} < \infty \}, \\ \|u\|_{W_q^s(\mathbf{R}^{N-1})} &= \|u\|_{H_q^{[s]}(\mathbf{R}^{N-1})} \\ &+ \sum_{|\alpha| = [s]} \left(\int_{\mathbf{R}^{N-1}} \int_{\mathbf{R}^{N-1}} \frac{|\partial^\alpha u(x) - \partial^\alpha u(y)|^q}{|x - y|^{N + (s - [s])q}} \, dx dy \right)^{\frac{1}{q}}. \end{split}$$

Let us define a solenoidal space by

$$J_q(\dot{\mathbf{R}}^N)=\{\mathbf{u}\in L_q(\dot{\mathbf{R}}^N)^N: (\mathbf{u},\nabla\varphi)_{\dot{\mathbf{R}}^N}=0 \text{ for any } \varphi\in \widehat{H}^1_{q'}(\mathbf{R}^N)\},$$
 where $q'=q/(q-1)$ and

$$(\mathbf{a}, \mathbf{b})_{\dot{\mathbf{R}}^N} = \int_{\dot{\mathbf{R}}^N} \mathbf{a}(x) \cdot \mathbf{b}(x) \, dx = \sum_{j=1}^N \int_{\dot{\mathbf{R}}^N} a_j(x) b_j(x) \, dx.$$

Furthermore, we set

$$X_{q} = W_{q}^{2-1/q}(\mathbf{R}^{N-1}) \times L_{q}(\dot{\mathbf{R}}^{N})^{N}, \quad JX_{q} = W_{q}^{2-1/q}(\mathbf{R}^{N-1}) \times J_{q}(\dot{\mathbf{R}}^{N}),$$

$$Y_{q} = L_{q}(\mathbf{R}^{N-1}) \times L_{q}(\dot{\mathbf{R}}^{N})^{N}, \qquad JY_{q} = L_{q}(\mathbf{R}^{N-1}) \times J_{q}(\dot{\mathbf{R}}^{N}),$$

and also $Y_1 = L_1(\mathbf{R}^{N-1}) \times L_1(\dot{\mathbf{R}}^N)^N$.

Next, we define the partial Fourier transform with respect to $x'=(x_1,\dots,x_{N-1})$ by

$$\widehat{u}(\xi', x_N) = \int_{\mathbf{R}^{N-1}} e^{-ix' \cdot \xi'} u(x', x_N) \, dx',$$

where $i = \sqrt{-1}$. Its inverse transform is also defined by

$$\mathcal{F}_{\xi'}^{-1}[v(\xi',x_N)](x') = \frac{1}{(2\pi)^{N-1}} \int_{\mathbf{R}^{N-1}} e^{ix'\cdot\xi'} v(\xi',x_N) \, d\xi'.$$

Finally, we introduce some constants and symbols. Let

$$\theta_j = \operatorname{Tan}^{-1}\left(\frac{j}{16}\right) \quad \text{for } j = 1, 2, \quad \lambda_1 = \lambda_0(\theta_1),$$
 (2.1)

where $\lambda_0(\varepsilon)$ is given in (1.4). In addition, we set

$$\alpha = \frac{\omega}{\rho_{+} + \rho_{-}} = \frac{(\rho_{-} - \rho_{+})\gamma_{a}}{\rho_{+} + \rho_{-}}, \quad \beta = \frac{\sqrt{\rho_{+}\mu_{+}}\sqrt{\rho_{-}\mu_{-}}}{(\rho_{+} + \rho_{-})(\sqrt{\rho_{+}\mu_{+}} + \sqrt{\rho_{-}\mu_{-}})}.$$
 (2.2)

The integral path Γ_0 is defined by

$$\Gamma_0 = \Gamma_0^+ \cup \Gamma_0^-, \quad \Gamma_0^+ = \{ \lambda \in \mathbf{C} : \lambda = \frac{2\lambda_1}{\sin \theta_1} + se^{i(\pi - \theta_1)}, s : 0 \to \infty \},$$

$$\Gamma_0^- = \{ \lambda \in \mathbf{C} : \lambda = \frac{2\lambda_1}{\sin \theta_1} + se^{-i(\pi - \theta_1)}, s : \infty \to 0 \}.$$
(2.3)

2.2. Main results. We first introduce the existence of solution operators for (1.1). This immediately follows from the resolvent estimate (1.4) and the standard theory of operator semigroups.

Theorem 2.1. Let $q \in (1, \infty)$ and ρ_{\pm} be any positive constants. Let

$$T(t)(d, \mathbf{f}) = \frac{1}{2\pi i} \int_{\Gamma_0} e^{\lambda t} (\eta(x', \lambda), \mathbf{u}(x, \lambda)) d\lambda,$$

with the solution $(\eta, \mathbf{u}) = (\eta(x', \lambda), \mathbf{u}(x, \lambda))$ to (1.2). Then the following assertions hold.

- (1) For any $(d, \mathbf{f}) \in X_q$, $T(t)(d, \mathbf{f}) \in C^{\infty}((0, \infty), W_q^{3-1/q}(\mathbf{R}^{N-1}) \times H_q^2(\dot{\mathbf{R}}^N)^N)$.
- (2) $\{T(t)\}_{t\geq 0}$ is an analytic C_0 -semigroup on JX_q .
- (3) For any $(d, \mathbf{f}) \in JX_q$, $(H, \mathbf{U}) = T(t)(d, \mathbf{f})$ is a unique solution to (1.1) with some pressure $P \in \widehat{H}^1_q(\dot{\mathbf{R}}^N)$.

Let us define projections P_1 , P_2 by

$$P_1: X_q\ni (a,\mathbf{b})\mapsto a\in W_q^{2-1/q}(\mathbf{R}^{N-1}), \quad P_2: X_q\ni (a,\mathbf{b})\mapsto \mathbf{b}\in L_q(\dot{\mathbf{R}}^N)^N.$$

We set for $(d, \mathbf{f}) \in X_q$

$$\mathcal{H}(t)(d, \mathbf{f}) = P_1 T(t)(d, \mathbf{f}) = \frac{1}{2\pi i} \int_{\Gamma_0} e^{\lambda t} \eta(x', \lambda) \, d\lambda,$$

$$\mathcal{U}(t)(d, \mathbf{f}) = P_2 T(t)(d, \mathbf{f}) = \frac{1}{2\pi i} \int_{\Gamma_0} e^{\lambda t} \mathbf{u}(x, \lambda) \, d\lambda.$$
(2.4)

One now decomposes the solution (\mathbf{u}, \mathbf{p}) of (1.2) into a solution of parabolic system and a solution of hyperbolic-parabolic coupled system as follows:

$$\begin{cases} \rho \lambda \mathbf{u}^{P} - \operatorname{Div}(\mu \mathbf{D}(\mathbf{u}^{P}) - \mathbf{p}^{P} \mathbf{I}) = \rho \mathbf{f} & \text{in } \dot{\mathbf{R}}^{N}, \\ \operatorname{div} \mathbf{u}^{P} = 0 & \text{in } \dot{\mathbf{R}}^{N}, \\ \llbracket (\mu \mathbf{D}(\mathbf{u}^{P}) - \mathbf{p}^{P} \mathbf{I}) \mathbf{e}_{N} \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}, \\ \llbracket \mathbf{u}^{P} \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}, \\ \lambda \eta - u_{N}^{H}|_{x_{N}=0} = d + u_{N}^{P}|_{x_{N}=0} & \text{on } \mathbf{R}^{N-1}, \\ \rho \lambda \mathbf{u}^{H} - \operatorname{Div}(\mu \mathbf{D}(\mathbf{u}^{H}) - \mathbf{p}^{H} \mathbf{I}) = 0 & \text{in } \dot{\mathbf{R}}^{N}, \\ \operatorname{div} \mathbf{u}^{H} = 0 & \text{in } \dot{\mathbf{R}}^{N}, \\ - \llbracket (\mu \mathbf{D}(\mathbf{u}^{H}) - \mathbf{p}^{H} \mathbf{I}) \mathbf{e}_{N} \rrbracket + (\omega - \sigma \Delta') \eta \mathbf{e}_{N} = 0 & \text{on } \mathbf{R}^{N-1}, \\ \llbracket \mathbf{u}^{H} \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}. \end{cases}$$

$$(2.5)$$

It then holds that $\mathbf{u} = \mathbf{u}^P + \mathbf{u}^H$ and $\mathfrak{p} = \mathfrak{p}^P + \mathfrak{p}^H$. In addition, we set

$$\mathcal{U}^P(t)\mathbf{f} = \frac{1}{2\pi i} \int_{\Gamma_0} e^{\lambda t} \mathbf{u}^P(x,\lambda) \, d\lambda, \quad \mathcal{U}^H(t)(d,\mathbf{f}) = \frac{1}{2\pi i} \int_{\Gamma_0} e^{\lambda t} \mathbf{u}^H(x,\lambda) \, d\lambda,$$

which implies

$$\mathcal{U}(t)(d,\mathbf{f}) = \mathcal{U}^{P}(t)\mathbf{f} + \mathcal{U}^{H}(t)(d,\mathbf{f}) \quad \text{for } (d,\mathbf{f}) \in X_{q}.$$
 (2.7)

For the parabolic part, we have the following theorem by the resolvent estimate (1.5) and the standard theory of operator semigroups.

Theorem 2.2. Let $r \in (1, \infty)$ and ρ_{\pm} be any positive constants. Then the following assertions hold.

- (1) For any $\mathbf{f} \in L_r(\dot{\mathbf{R}}^N)^N$, $\mathcal{U}^P(t)\mathbf{f} \in C^{\infty}((0,\infty), H_r^2(\dot{\mathbf{R}}^N)^N)$.
- (2) $\{\mathcal{U}^P(t)\}_{t\geq 0}$ is an analytic C_0 -semigroup on $J_r(\dot{\mathbf{R}}^N)$.
- (3) Let $j \in \mathbf{N}_0$ and k = 0, 1, 2. For any t > 0 and $\mathbf{f} \in J_r(\dot{\mathbf{R}}^N)$,

$$\|\partial_t^j \nabla^k \mathcal{U}^P(t) \mathbf{f}\|_{L_s(\dot{\mathbf{R}}^N)} \le C t^{-j-\frac{k}{2}-\frac{N}{2}\left(\frac{1}{r}-\frac{1}{s}\right)} \|\mathbf{f}\|_{L_r(\dot{\mathbf{R}}^N)},$$

where $r \leq s \leq \infty$ when k = 0, 1 and $r \leq s < \infty$ when k = 2. Here C is a positive constant independent of t and \mathbf{f} .

To complete time-decay estimates for $\mathcal{H}(t)$ and $\mathcal{U}(t)$, we further decompose $(\eta, \mathbf{u}^H, \mathfrak{p}^H)$ satisfying (2.6) as follows: for $z^1 = d$ and $z^2 = u_N^P|_{x_N=0}$, let (η^k, \mathbf{u}^k) be the solution to

$$\begin{cases} \lambda \eta^k - u_N^k|_{x_N = 0} = z^k & \text{on } \mathbf{R}^{N-1}, \\ \rho \lambda \mathbf{u}^k - \text{Div}(\mu \mathbf{D}(\mathbf{u}^k) - \mathfrak{p}^k \mathbf{I}) = 0 & \text{in } \dot{\mathbf{R}}^N, \\ \text{div } \mathbf{u}^k = 0 & \text{in } \dot{\mathbf{R}}^N, \\ -\llbracket (\mu \mathbf{D}(\mathbf{u}^k) - \mathfrak{p}^k \mathbf{I}) \mathbf{e}_N \rrbracket + (\omega - \sigma \Delta') \eta^k \mathbf{e}_N = 0 & \text{on } \mathbf{R}^{N-1}, \\ \llbracket \mathbf{u}^k \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}. \end{cases}$$
(2.8)

It then holds that $\eta = \eta^1 + \eta^2$, $\mathbf{u}^H = \mathbf{u}^1 + \mathbf{u}^2$, and $\mathfrak{p}^H = \mathfrak{p}^1 + \mathfrak{p}^2$. Let $\varphi = \varphi(\xi')$ be a function in $C^{\infty}(\mathbf{R}^{N-1})$ and satisfy $0 \le \varphi \le 1$ with

$$\varphi(\xi') = \begin{cases} 1 & (|\xi'| \le 1), \\ 0 & (|\xi'| \ge 2). \end{cases}$$

In addition, we set $\varphi_{A_0}(\xi') = \varphi(\xi'/A_0)$ and $\varphi_{A_\infty}(\xi') = 1 - \varphi(\xi'/A_\infty)$ for positive constants $A_0 \in (0,1)$ and $A_\infty \geq 2$. Let $\varphi_{[A_0,A_\infty]}(\xi') = 1 - \varphi_{A_0}(\xi') - \varphi_{A_\infty}(\xi')$. Together with these cut-off functions, we define for $Z \in \{A_0,A_\infty,[A_0,A_\infty]\}$ and for an integral path Γ

$$\widehat{\mathcal{H}}_{Z}^{1}(t;\Gamma)d = \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \widehat{\eta}^{1}(\xi',\lambda) d\lambda,
\widehat{\mathcal{H}}_{Z}^{2}(t;\Gamma)\mathbf{f} = \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \widehat{\eta}^{2}(\xi',\lambda) d\lambda,
\widehat{\mathcal{U}}_{Z}^{1}(t;\Gamma)d = \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \widehat{\mathbf{u}}^{1}(\xi',x_{N},\lambda) d\lambda,
\widehat{\mathcal{U}}_{Z}^{2}(t;\Gamma)\mathbf{f} = \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \widehat{\mathbf{u}}^{2}(\xi',x_{N},\lambda) d\lambda.$$
(2.9)

Furthermore, we set for $S \in \{\mathcal{H}, \mathcal{U}\}$ and Γ_0 given by (2.3)

$$\mathcal{S}_Z^1(t)d = \mathcal{F}_{\xi'}^{-1}[\widehat{\mathcal{S}}_Z^1(t;\Gamma_0)d](x'), \quad \mathcal{S}_Z^2(t)\mathbf{f} = \mathcal{F}_{\xi'}^{-1}[\widehat{\mathcal{S}}_Z^2(t;\Gamma_0)\mathbf{f}](x'). \tag{2.10}$$

Noting (2.7), we see that the formulas in (2.4) satisfy

$$\mathcal{H}(t)(d, \mathbf{f}) = \sum_{Z \in \{A_0, A_\infty, [A_0, A_\infty]\}} \left(\mathcal{H}_Z^1(t)d + \mathcal{H}_Z^2(t)\mathbf{f} \right),$$

$$\mathcal{U}(t)(d, \mathbf{f}) = \sum_{Z \in \{A_0, A_\infty, [A_0, A_\infty]\}} \left(\mathcal{U}_Z^1(t)d + \mathcal{U}_Z^2(t)\mathbf{f} \right) + \mathcal{U}^P(t)\mathbf{f}.$$
(2.11)

The following two theorems are our main results of this paper. The first one is time-decay estimates for the low frequency part.

Theorem 2.3. Let $1 \le p < 2 \le q \le \infty$ and suppose that $\rho_- > \rho_+ > 0$. Then there exists a constant $A_0 \in (0,1)$ such that the following assertions hold.

(1) For $\xi' \in \mathbf{R}^{N-1}$ with $|\xi'| \in (0, A_0)$, let $\zeta_{\pm} = \pm i\alpha^{1/2} |\xi'|^{1/2} - \sqrt{2}\alpha^{1/4}\beta(1\pm i)|\xi'|^{5/4}$ and

$$\widehat{\Gamma}_{\mathrm{Res}}^{\pm} = \{\lambda \in \mathbf{C} : \lambda = \zeta_{\pm} + |\xi'|^{6/4} e^{is}, s : 0 \to 2\pi\}.$$

In addition, for $S \in \{\mathcal{H}, \mathcal{U}\}$ and $(d, \mathbf{f}) \in Y_p$, set

$$\mathcal{S}_{A_0}^{1\pm}(t)d = \mathcal{F}_{\xi'}^{-1}[\widehat{\mathcal{S}}_{A_0}^1(t;\widehat{\boldsymbol{\Gamma}}_{\mathrm{Res}}^\pm)d](x'), \quad \mathcal{S}_{A_0}^{2\pm}(t)\mathbf{f} = \mathcal{F}_{\xi'}^{-1}[\widehat{\mathcal{S}}_{A_0}^2(t;\widehat{\boldsymbol{\Gamma}}_{\mathrm{Res}}^\pm)\mathbf{f}](x').$$

Then for any $t \geq 1$

$$\begin{split} &\|(\mathcal{H}_{A_0}^{1\pm}(t)d,\mathcal{H}_{A_0}^{2\pm}(t)\mathbf{f})\|_{L_q(\mathbf{R}^{N-1})} \leq Ct^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)}\|(d,\mathbf{f})\|_{Y_p}, \\ &\|(\mathcal{U}_{A_0}^{1\pm}(t)d,\mathcal{U}_{A_0}^{2\pm}(t)\mathbf{f})\|_{L_q(\dot{\mathbf{R}}^N)} \leq Ct^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{4}{5}\left(\frac{1}{2}-\frac{1}{q}\right)}\|(d,\mathbf{f})\|_{Y_p}, \end{split}$$

with some positive constant C independent of t, d, and \mathbf{f} .

(2) For $S \in \{\mathcal{H}, \mathcal{U}\}$ and $(d, \mathbf{f}) \in Y_p$, set

$$\begin{split} \widetilde{\mathcal{S}}_{A_0}^1(t)d &= \mathcal{S}_{A_0}^1(t)d - \mathcal{S}_{A_0}^{1+}(t)d - \mathcal{S}_{A_0}^{1-}(t)d, \\ \widetilde{\mathcal{S}}_{A_0}^2(t)\mathbf{f} &= \mathcal{S}_{A_0}^2(t)\mathbf{f} - \mathcal{S}_{A_0}^{2+}(t)\mathbf{f} - \mathcal{S}_{A_0}^{2-}(t)\mathbf{f}. \end{split}$$

Let γ_1 be a constant satisfying

$$0 < \gamma_1 < \min\left\{1, 2(N-1)\left(\frac{1}{p} - \frac{1}{2}\right)\right\}.$$

Then for any $t \geq 1$

$$\begin{split} &\|(\widetilde{\mathcal{H}}_{A_0}^1(t)d,\widetilde{\mathcal{H}}_{A_0}^2(t)\mathbf{f})\|_{L_q(\mathbf{R}^{N-1})} \leq Ct^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{3}{4}\gamma_1}\|(d,\mathbf{f})\|_{Y_p},\\ &\|(\widetilde{\mathcal{U}}_{A_0}^1(t)d,\widetilde{\mathcal{U}}_{A_0}^2(t)\mathbf{f})\|_{L_q(\dot{\mathbf{R}}^N)} \leq Ct^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|(d,\mathbf{f})\|_{Y_p}, \end{split}$$

with some positive constant C independent of t, d, and \mathbf{f} .

Remark 2.4. Time-decay estimates for higher-order derivatives of the low frequency part will be discussed in a forthcoming paper.

The second one is time-decay estimates for the high frequency part.

Theorem 2.5. Let $q \in (1, \infty)$ and $j \in \mathbb{N}_0$. Suppose that $\rho_- > \rho_+ > 0$. Then there exist constants $A_{\infty} \geq 2$ and $\gamma_0 > 0$ such that for any $t \geq 1$ and $(d, \mathbf{f}) \in X_q$

$$\|(\partial_t^j \mathcal{H}_Z^1(t)d, \partial_t^j \mathcal{H}_Z^2(t)\mathbf{f})\|_{W_q^{3-1/q}(\mathbf{R}^{N-1})} \le Ce^{-\gamma_0 t} \|(d, \mathbf{f})\|_{X_q},$$
$$\|(\partial_t^j \mathcal{U}_Z^1(t)d, \partial_t^j \mathcal{U}_Z^2(t)\mathbf{f})\|_{H_q^2(\dot{\mathbf{R}}^N)} \le Ce^{-\gamma_0 t} \|(d, \mathbf{f})\|_{X_q},$$

where $Z \in \{A_{\infty}, [A_0, A_{\infty}]\}$ and C is a positive constant independent of t, d, and f. Here A_0 is the positive constant given in Theorem 2.2

Recalling (2.11), we have time-decay estimates of solutions for (1.1) from Theorems 2.2, 2.3, and 2.5 immediately.

Theorem 2.6. Let $1 and <math>(d, \mathbf{f}) \in X_q \cap JY_p$. Then for any $t \ge 1$

$$\|\mathcal{H}(t)(d,\mathbf{f})\|_{L_q(\mathbf{R}^{N-1})} \le Ct^{-\min\left\{\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right),\frac{(N-1)}{2}\left(\frac{1}{p}-\frac{1}{q}\right)+\frac{3}{4}\gamma_1\right\}} \|(d,\mathbf{f})\|_{X_q\cap Y_p},$$

$$\|\mathcal{U}(t)(d,\mathbf{f})\|_{L_q(\dot{\mathbf{R}}^N)} \leq C t^{-\min\left\{\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)+\frac{4}{5}\left(\frac{1}{2}-\frac{1}{q}\right),\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)\right\}} \|(d,\mathbf{f})\|_{X_q\cap Y_p},$$

with a positive constant C independent of t, d, and f, where γ_1 is the positive constant given in Theorem 2.3.

3. Representation formulas for solutions

This section introduces the representation formulas for solutions of (1.1). In this section, we assume that ρ_{\pm} are any positive constants except for Lemma 3.4 (2) below. Here we collect several symbols appearing in the representation formulas. Let $z_0 = \min\{\mu_+/\rho_+, \mu_-/\rho_-\}$. For $\xi' = (\xi_1, \dots, \xi_{N-1}) \in \mathbf{R}^{N-1}$ and $\lambda \in \mathbf{C} \setminus (-\infty, -z_0|\xi'|^2]$, we set

$$A = |\xi'|, \quad B_{\pm} = \sqrt{\frac{\rho_{\pm}}{\mu_{\pm}}\lambda + |\xi'|^2},$$
 (3.1)

where we have chosen a branch cut along the negative real axis and a branch of the square root so that $\Re\sqrt{z} > 0$ for $z \in \mathbb{C} \setminus (-\infty, 0]$. In addition,

$$D_{\pm} = \mu_{\pm}B_{\pm} + \mu_{\mp}A, \quad E = \mu_{+}B_{+} + \mu_{-}B_{-}, \quad \mathcal{M}_{\pm}(a) = \frac{e^{-Aa} - e^{-B_{\pm}a}}{A - B_{\pm}} \quad (a \ge 0),$$
(3.2)

and also

$$F(A,\lambda) = -(\mu_{+} - \mu_{-})^{2} A^{3} + \{(3\mu_{+} - \mu_{-})\mu_{+} B_{+} + (3\mu_{-} - \mu_{+})\mu_{-} B_{-}\} A^{2}$$

$$+ \{(\mu_{+} B_{+} + \mu_{-} B_{-})^{2} + \mu_{+} \mu_{-} (B_{+} + B_{-})^{2}\} A$$

$$+ (\mu_{+} B_{+} + \mu_{-} B_{-})(\mu_{+} B_{+}^{2} + \mu_{-} B_{-}^{2}).$$

$$(3.3)$$

Except for D_{\pm} and E, the above symbols are introduced in [10, (3.3), (3.8), (3.15)]. Furthermore, the following properties are proved in [10, Lemmas 4.7 and 4.8].

Lemma 3.1. Let $\varepsilon \in (0, \pi/2)$ and ρ_{\pm} be any positive constants. Then the following assertions hold.

(1) For any
$$\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$$
 and $\lambda \in \Sigma_{\varepsilon} \cup \{0\}$,

$$C_1(|\lambda|^{1/2} + A) \le \Re B_{\pm} \le |B_{\pm}| \le C_2(|\lambda|^{1/2} + A),$$

$$C_1(|\lambda|^{1/2} + A)^3 < |F(A, \lambda)| < C_2(|\lambda|^{1/2} + A)^3.$$

with positive constants C_1 and C_2 depending on ε , but independent of ξ' and λ . (2) Let $s \in \mathbf{R}$ and $\alpha' \in \mathbf{N}_0^{N-1}$. Then for any $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and $\lambda \in \Sigma_{\varepsilon} \cup \{0\}$

$$\begin{split} |\partial_{\xi'}^{\alpha'}A^s| &\leq CA^{s-|\alpha'|}, \quad |\partial_{\xi'}^{\alpha'}B_{\pm}^s| \leq C(|\lambda|^{1/2} + A)^{s-|\alpha'|}, \\ |\partial_{\xi'}^{\alpha'}E^s| &\leq C(|\lambda|^{1/2} + A)^{s-|\alpha'|}, \end{split}$$

$$|\partial_{\xi'}^{\alpha'}(A+B_{\pm})^s| \le C(|\lambda|^{1/2}+A)^s A^{-|\alpha'|},$$

$$|\partial_{\xi'}^{\alpha'} F(A,\lambda)^s| \le C(|\lambda|^{1/2} + A)^{3s} A^{-|\alpha'|},$$

where C is a positive constant depending on ε , s, and α' , but independent of ξ' and λ .

From Lemma 3.1 and the Bell formula of derivatives of composite functions, we have

Lemma 3.2. Let $\varepsilon \in (0, \pi/2)$ and ρ_{\pm} be any positive constants.

- (1) Let $s \in \mathbf{R}$ and $\alpha' \in \mathbf{N}_0^{N-1}$. Then for any $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and $\lambda \in \Sigma_{\varepsilon} \cup \{0\}$ $|\partial_{\xi'}^{\alpha'} D_{\pm}^s| \leq C(|\lambda|^{1/2} + A)^s A^{-|\alpha'|}, \quad |\partial_{\xi'}^{\alpha'} (D_+ + D_-)^s| \leq C(|\lambda|^{1/2} + A)^s A^{-|\alpha'|},$ where C is a positive constant depending on ε , s, and α' , but independent of ξ' and λ .
- (2) Let $\theta, \nu, \tau > 0$ and $\alpha' \in \mathbf{N}_0^{N-1}$. Then there exists a positive constant $C_{\alpha',\theta}$, independent of ν and τ , such that for any $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$

$$|\partial_{\xi'}^{\alpha'} e^{-\nu\tau A^{\theta}}| \le C_{\alpha',\theta} A^{-|\alpha'|} e^{-(\nu\tau A^{\theta})/2}.$$

3.1. A representation formula for the parabolic part. In this subsection, we introduce a representation formula for solutions of (2.5). Note that (2.5) admits a unique solution $(\mathbf{u}^P, \mathbf{p}^P)$ for $\lambda \in \Sigma_{\varepsilon}$ and $\mathbf{f} \in L_q(\dot{\mathbf{R}}^N)^N$, $\varepsilon \in (0, \pi/2)$ and $q \in (1, \infty)$, as discussed in Section 1. Since $C_0^{\infty}(\dot{\mathbf{R}}^N)$ is dense in $L_p(\dot{\mathbf{R}}^N)$ for $1 \leq p < \infty$, it suffices to consider $\mathbf{f} \in C_0^{\infty}(\dot{\mathbf{R}}^N)^N$ in what follows. Our aim is to prove

Proposition 3.3. Let $\varepsilon \in (0, \pi/2)$ and ρ_{\pm} be any positive constants. Suppose that $(\mathbf{u}^P, \mathbf{p}^P)$ is a solution to (2.5) for some $\lambda \in \Sigma_{\varepsilon}$ and $\mathbf{f} = (f_1, \dots, f_N)^{\mathsf{T}} \in C_0^{\infty}(\dot{\mathbf{R}}^N)^N$. Then there holds

$$\begin{split} &\widehat{u}_{N}^{P}(\xi',x_{N},\lambda)|_{x_{N}=0} \\ &= \sum_{\mathfrak{a},\mathfrak{b}\in\{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)}{A(A+B_{\mathfrak{b}})F(A,\lambda)} \mathcal{M}_{\mathfrak{b}}(y_{N}) \widehat{f_{j}}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &+ \sum_{\mathfrak{a},\mathfrak{b}\in\{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)}{AB_{\mathfrak{b}}(A+B_{\mathfrak{b}})F(A,\lambda)} e^{-B_{\mathfrak{b}}y_{N}} \widehat{f_{j}}(\xi',\mathfrak{a}y_{N}) \, dy_{N}. \end{split}$$

Here $\widehat{u}_N^P(\xi', x_N, \lambda)$ stands for the partial Fourier transform of the Nth component of \mathbf{u}^P and

$$\begin{split} &\Phi^{\mathfrak{a},\mathfrak{b}}_{j}(\xi',\lambda) = \sum_{\substack{|\alpha'|+k+l+m=5\\|\alpha'|+k\geq 2}} C^{\mathfrak{a},\mathfrak{b}}_{j,\alpha',k,l,m}(\xi')^{\alpha'} A^{k} B^{l}_{+} B^{m}_{-}, \\ &\Psi^{\mathfrak{a},\mathfrak{b}}_{j}(\xi',\lambda) = \sum_{\substack{|\alpha'|+k+l+m=5\\|\alpha'|+k\geq 2}} \widetilde{C}^{\mathfrak{a},\mathfrak{b}}_{j,\alpha',k,l,m}(\xi')^{\alpha'} A^{k} B^{l}_{+} B^{m}_{-}, \end{split}$$

with constants $C^{\mathfrak{a},\mathfrak{b}}_{j,\alpha',k,l,m}$ and $\widetilde{C}^{\mathfrak{a},\mathfrak{b}}_{j,\alpha',k,l,m}$ independent of ξ' and λ .

Proof. Let $j=1,\ldots,N-1$ and $J=1,\ldots,N$ in this proof. Although we follow calculations of [10, Section 3], we will achieve a set of equations simpler than [10, (3.6)] in what follows, see (3.24) below.

Step 1. We compute $\widehat{w}_N(\xi', 0-, \lambda)$ for the following resolvent problem:

$$\begin{cases} \rho \lambda \mathbf{w} - \operatorname{Div}(\mu \mathbf{D}(\mathbf{w}) - \mathfrak{r} \mathbf{I}) = 0 & \text{in } \dot{\mathbf{R}}^{N}, \\ \operatorname{div} \mathbf{w} = 0 & \text{in } \dot{\mathbf{R}}^{N}, \\ \llbracket (\mu \mathbf{D}(\mathbf{w}) - \mathfrak{r} \mathbf{I}) \mathbf{e}_{N} \rrbracket = \mathbf{g} & \text{on } \mathbf{R}^{N-1}, \\ \llbracket \mathbf{w} \rrbracket = \mathbf{h} & \text{on } \mathbf{R}^{N-1}, \end{cases}$$
(3.4)

where $\mathbf{g} = (g_1, \dots, g_N)^\mathsf{T}$ and $\mathbf{h} = (h_1, \dots, h_N)^\mathsf{T}$ are suitable functions on \mathbf{R}^{N-1} specified in Step 2 below. The restriction of \mathbf{w} and \mathfrak{r} to \mathbf{R}_{\pm}^N are denoted by \mathbf{w}_{\pm} and \mathfrak{r}_{\pm} , respectively. Let us write the Jth component of \mathbf{w}_{\pm} by $w_{J\pm}$ and observe $\mathrm{Div}(\mu \mathbf{D}(\mathbf{w}_{\pm})) = \mu_{\pm} \Delta \mathbf{w}_{\pm}$ by $\mathrm{div} \mathbf{w}_{\pm} = 0$ in \mathbf{R}_{\pm}^N . Then (3.4) can be written as

$$\begin{cases} \rho_{\pm}\lambda w_{J\pm} - \mu_{\pm}\Delta w_{\pm} \text{ by div } \mathbf{w}_{\pm} = 0 \text{ in } \mathbf{R}_{\pm}^{N}. \text{ Then (5.4) can be written as} \\ \rho_{\pm}\lambda w_{J\pm} - \mu_{\pm}\Delta w_{J\pm} + \partial_{J}\mathbf{r}_{\pm} = 0 & \text{ in } \mathbf{R}_{\pm}^{N}, \\ \sum_{j=1}^{N-1} \partial_{j}w_{j\pm} + \partial_{N}w_{N\pm} = 0 & \text{ in } \mathbf{R}_{\pm}^{N}, \\ \mu_{+}(\partial_{N}w_{j+} + \partial_{j}w_{N+})|_{x_{N}=0} - \mu_{-}(\partial_{N}w_{j-} + \partial_{j}w_{N-})|_{x_{N}=0} = g_{j} & \text{ on } \mathbf{R}^{N-1}, \\ (2\mu_{+}\partial_{N}w_{N+} - \mathbf{r}_{+})|_{x_{N}=0} - (2\mu_{-}\partial_{N}w_{N-} - \mathbf{r}_{-})|_{x_{N}=0} = g_{N} & \text{ on } \mathbf{R}^{N-1}, \\ w_{J+}|_{x_{N}=0} - w_{J-}|_{x_{N}=0} = h_{J} & \text{ on } \mathbf{R}^{N-1}. \end{cases}$$

Let $\widehat{w}_{J\pm}(x_N) = \widehat{w}_{J\pm}(\xi', x_N, \lambda)$ and $\widehat{\mathfrak{r}}_{\pm}(x_N) = \widehat{\mathfrak{r}}_{\pm}(\xi', x_N, \lambda)$. Applying the partial Fourier transform to the last system yields

$$\rho_{\pm}\lambda \widehat{w}_{j\pm}(x_N) - \mu_{\pm}(\partial_N^2 - |\xi'|^2)\widehat{w}_{j\pm}(x_N) + i\xi_j \widehat{\mathfrak{r}}_{\pm}(x_N) = 0, \ \pm x_N > 0, \ (3.5)$$

$$\rho_{\pm}\lambda \widehat{w}_{N\pm}(x_N) - \mu_{\pm}(\partial_N^2 - |\xi'|^2)\widehat{w}_{N\pm}(x_N) + \partial_N \widehat{\mathfrak{r}}_{\pm}(x_N) = 0, \ \pm x_N > 0, \ (3.6)$$

$$\sum_{j=1}^{N-1} i\xi_j \widehat{w}_{j\pm}(x_N) + \partial_N \widehat{w}_{N\pm}(x_N) = 0, \ \pm x_N > 0, \ (3.7)$$

$$\mu_{+}(\partial_{N}\widehat{w}_{j+}(0) + i\xi_{j}\widehat{w}_{N+}(0)) - \mu_{-}(\partial_{N}\widehat{w}_{j-}(0) + i\xi_{j}\widehat{w}_{N-}(0)) = \widehat{g}_{j}, \tag{3.8}$$

$$(2\mu_{+}\partial_{N}\widehat{w}_{N+}(0) - \widehat{\mathfrak{r}}_{+}(0)) - (2\mu_{-}\partial_{N}\widehat{w}_{N-}(0) - \widehat{\mathfrak{r}}_{-}(0)) = \widehat{g}_{N}, \tag{3.9}$$

$$\widehat{w}_{J+}(0) - \widehat{w}_{J-}(0) = \widehat{h}_{J}, \tag{3.10}$$

where $\hat{g}_J = \hat{g}_J(\xi')$ and $\hat{h}_J = \hat{h}_J(\xi')$. Note that (3.5) and (3.6) are respectively equivalent to

$$-\mu_{\pm}(\partial_N^2 - B_{\pm}^2)\widehat{w}_{j\pm}(x_N) + i\xi_j\widehat{\mathfrak{r}}_{\pm}(x_N) = 0, \ \pm x_N > 0, \tag{3.11}$$

$$-\mu_{\pm}(\partial_N^2 - B_{\pm}^2)\widehat{w}_{N\pm}(x_N) + \partial_N \widehat{\mathfrak{r}}_{\pm}(x_N) = 0, \ \pm x_N > 0.$$
 (3.12)

From now on, we look for $\widehat{w}_{J\pm}(x_N)$ and $\widehat{\mathfrak{r}}_{\pm}(x_N)$ of the forms: for $\pm x_N > 0$,

$$\widehat{w}_{J\pm}(x_N) = \alpha_{J\pm}(e^{\mp Ax_N} - e^{\mp B_{\pm}x_N}) + \beta_{J\pm}e^{\mp B_{\pm}x_N}, \quad \widehat{\mathfrak{r}}_{\pm}(x_N) = \gamma_{\pm}e^{\mp Ax_N}. \quad (3.13)$$

Inserting these formulas into (3.11), (3.12), and (3.7)-(3.10) furnishes

$$-\mu_{\pm}\alpha_{i\pm}(A^2 - B_{+}^2) + i\xi_{i}\gamma_{\pm} = 0, \quad -\mu_{\pm}\alpha_{N\pm}(A^2 - B_{+}^2) \mp A\gamma_{\pm} = 0, \quad (3.14)$$

$$i\xi' \cdot \alpha'_{\pm} \mp A\alpha_{N\pm} = 0, \quad -i\xi' \cdot \alpha'_{\pm} + i\xi' \cdot \beta'_{\pm} \pm B_{\pm}\alpha_{N\pm} \mp B_{\pm}\beta_{N\pm} = 0,$$
 (3.15)

$$\mu_{+}\{\alpha_{j+}(-A+B_{+})-\beta_{j+}B_{+}+i\xi_{j}\beta_{N+}\}$$

$$-\mu_{-}\{\alpha_{j-}(A-B_{-}) + \beta_{j-}B_{-} + i\xi_{j}\beta_{N-}\} = \widehat{g}_{j}, \quad (3.16)$$

$$[2\mu_{+}\{\alpha_{N+}(-A+B_{+})-\beta_{N+}B_{+}\}-\gamma_{+}]$$

$$-[2\mu_{-}\{\alpha_{N-}(A-B_{-})+\beta_{N-}B_{-}\}-\gamma_{-}]=\widehat{g}_{N},\quad(3.17)$$

$$\beta_{J+} - \beta_{J-} = \hat{h}_J, \quad (3.18)$$

where $i\xi' \cdot \alpha'_{\pm} = \sum_{j=1}^{N-1} i\xi_j \alpha_{j\pm}$ and $i\xi' \cdot \beta' = \sum_{j=1}^{N-1} i\xi_j \beta_{j\pm}$. Let us solve the equations (3.14)-(3.18). By (3.15), we have

$$\alpha_{N\pm} = \pm \frac{(i\xi' \cdot \beta'_{\pm} \mp B_{\pm}\beta_{N\pm})}{A - B_{+}}, \quad i\xi' \cdot \alpha'_{\pm} = \frac{A(i\xi' \cdot \beta'_{\pm} \mp B_{\pm}\beta_{N\pm})}{A - B_{+}}.$$
 (3.19)

By the first equation of (3.19) and the second equation of (3.14),

$$\gamma_{\pm} = -\frac{\mu_{\pm}(A + B_{\pm})}{A} (i\xi' \cdot \beta_{\pm}' \mp B_{\pm}\beta_{N\pm}). \tag{3.20}$$

Multiplying (3.16) by $i\xi_i$ and summing the resultant formulas yield

$$\mu_{+}\{i\xi'\cdot\alpha'_{+}(-A+B_{+})-i\xi'\cdot\beta'_{+}B_{+}-A^{2}\beta_{N+}\}$$
$$-\mu_{-}\{i\xi'\cdot\alpha'_{-}(A-B_{-})+i\xi'\cdot\beta'_{-}B_{-}-A^{2}\beta_{N-}\}=i\xi'\cdot\widehat{g}',$$

where $i\xi' \cdot \hat{g}' = \sum_{j=1}^{N-1} i\xi_j \hat{g}_j$. Combining this equation with the second one of (3.19),

$$\mu_{+}\{-(A+B_{+})i\xi'\cdot\beta'_{+} + A(B_{+}-A)\beta_{N+}\} -\mu_{-}\{(A+B_{-})i\xi'\cdot\beta'_{-} + A(B_{-}-A)\beta_{N-}\} = i\xi'\cdot\widehat{g}'. \quad (3.21)$$

In addition, by (3.17) and (3.20) together with the first equation of (3.19),

$$\mu_{+}\{(-A+B_{+})i\xi'\cdot\beta'_{+} - B_{+}(A+B_{+})\beta_{N+}\}$$
$$-\mu_{-}\{(-A+B_{-})i\xi'\cdot\beta'_{-} + B_{-}(B_{-}+A)\beta_{N-}\} = A\widehat{g}_{N}. \quad (3.22)$$

Since it follows from (3.18) that $i\xi' \cdot \beta'_{+} = i\xi' \cdot \beta'_{-} + i\xi' \cdot \hat{h}'$ and $\beta_{N+} = \beta_{N-} + \hat{h}_{N+}$ it holds by (3.21) and (3.22) that

$$\{\mu_{+}(A+B_{+}) + \mu_{-}(A+B_{-})\}i\xi' \cdot \beta'_{-} - \{\mu_{+}A(B_{+}-A) - \mu_{-}A(B_{-}-A)\}\beta_{N-}$$

$$= -i\xi' \cdot \hat{g}' - \mu_{+}(A+B_{+})i\xi' \cdot \hat{h}' + \mu_{+}A(B_{+}-A)\hat{h}_{N} =: G(\mathbf{g}, \mathbf{h}),$$

$$- \{\mu_{+}(-A+B_{+}) - \mu_{-}(-A+B_{-})\}i\xi' \cdot \beta'_{-}$$

$$+ \{\mu_{+}B_{+}(A+B_{+}) + \mu_{-}B_{-}(B_{-}+A)\}\beta_{N-}$$

$$= -A\hat{q}_{N} + \mu_{+}(-A+B_{+})i\xi' \cdot \hat{h}' - \mu_{+}B_{+}(A+B_{+})\hat{h}_{N} =: H(\mathbf{g}, \mathbf{h}).$$
(3.23)

We have thus achieved

$$\mathbf{L} \begin{pmatrix} i\xi' \cdot \beta'_{-} \\ \beta_{N-} \end{pmatrix} = \begin{pmatrix} G(\mathbf{g}, \mathbf{h}) \\ H(\mathbf{g}, \mathbf{h}) \end{pmatrix}, \tag{3.24}$$

$$\mathbf{L} = \begin{pmatrix} \mu_{+}(A+B_{+}) + \mu_{-}(A+B_{-}) & -\mu_{+}A(B_{+}-A) + \mu_{-}A(B_{-}-A) \\ -\mu_{+}(-A+B_{+}) + \mu_{-}(-A+B_{-}) & \mu_{+}B_{+}(A+B_{+}) + \mu_{-}B_{-}(B_{-}+A) \end{pmatrix}.$$

Then the inverse matrix L^{-1} of L is given b

$$\mathbf{L}^{-1} = \frac{1}{F(A,\lambda)} \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

where $F(A, \lambda)$ is defined in (3.3) and

$$L_{11} = \mu_{+}B_{+}(A+B_{+}) + \mu_{-}B_{-}(B_{-}+A),$$

$$L_{12} = \mu_{+} A(B_{+} - A) - \mu_{-} A(B_{-} - A),$$

$$L_{21} = \mu_{+} (-A + B_{+}) - \mu_{-} (-A + B_{-}),$$

$$L_{22} = \mu_{+} (A + B_{+}) + \mu_{-} (A + B_{-}).$$
(3.25)

Solving (3.24), we have

$$i\xi' \cdot \beta'_{-} = \frac{L_{11}G(\mathbf{g}, \mathbf{h}) + L_{12}H(\mathbf{g}, \mathbf{h})}{F(A, \lambda)}, \quad \beta_{N-} = \frac{L_{21}G(\mathbf{g}, \mathbf{h}) + L_{22}H(\mathbf{g}, \mathbf{h})}{F(A, \lambda)}.$$
 (3.26)

Since $\widehat{w}_N(\xi',0-,\lambda) = \widehat{w}_{N-}(\xi',0,\lambda) = \beta_{N-}$, there holds

$$\widehat{w}_N(\xi', 0 -, \lambda) = \frac{L_{21}G(\mathbf{g}, \mathbf{h}) + L_{22}H(\mathbf{g}, \mathbf{h})}{F(A, \lambda)}.$$
(3.27)

Step 2. We compute the formula of $\widehat{u}_N^P(\xi',0,\lambda)$. Let

$$(\psi_{\pm}, \phi_{\pm}) = (\psi_{1\pm}, \dots, \psi_{N\pm}, \phi_{\pm})$$

be solutions to the following whole space problems without interface:

$$\begin{cases}
\rho_{+}\lambda\psi_{+} - \operatorname{Div}(\mu_{+}\mathbf{D}(\psi_{+}) - \phi_{+}\mathbf{I}) = \mathbf{F}, & \operatorname{div}\psi_{+} = 0 & \operatorname{in} \mathbf{R}^{N}, \\
\rho_{-}\lambda\psi_{-} - \operatorname{Div}(\mu_{-}\mathbf{D}(\psi_{-}) - \phi_{-}\mathbf{I}) = \mathbf{F}, & \operatorname{div}\psi_{-} = 0 & \operatorname{in} \mathbf{R}^{N},
\end{cases}$$
(3.28)

where $\mathbf{F} = \rho \mathbf{f}$. Set $\psi = \psi_+ \mathbb{1}_{\mathbf{R}_+^N} + \psi_- \mathbb{1}_{\mathbf{R}_-^N}$ and $\phi = \phi_+ \mathbb{1}_{\mathbf{R}_+^N} + \phi_- \mathbb{1}_{\mathbf{R}_-^N}$, and then (ψ, ϕ) satisfies

$$\rho \lambda - \text{Div}(\mu \mathbf{D}(\psi) - \phi \mathbf{I}) = \rho \mathbf{f}, \quad \text{div } \psi = 0 \quad \text{in } \dot{\mathbf{R}}^N.$$

In addition, $\llbracket \phi \rrbracket = 0$ as discussed in the appendix below. Thus $(\mathbf{u}^P, \mathfrak{p}^P)$ are given by $\mathbf{u}^P = \psi + \mathbf{v}$ and $\mathfrak{p}^P = \phi + \mathfrak{q}$ for a solution $(\mathbf{v}, \mathfrak{q})$ to

$$\begin{cases} \rho \lambda \mathbf{v} - \mathrm{Div}(\mu \mathbf{D}(\mathbf{v}) - \mathfrak{q} \mathbf{I}) = 0 & \text{in } \dot{\mathbf{R}}^N, \\ \operatorname{div} \mathbf{v} = 0 & \text{in } \dot{\mathbf{R}}^N, \\ \llbracket (\mu \mathbf{D}(\mathbf{v}) - \mathfrak{q} \mathbf{I}) \mathbf{e}_N \rrbracket = -\llbracket \mu \mathbf{D}(\psi) \mathbf{e}_N \rrbracket & \text{on } \mathbf{R}^{N-1}, \\ \llbracket \mathbf{v} \rrbracket = -\llbracket \psi \rrbracket & \text{on } \mathbf{R}^{N-1}. \end{cases}$$

We now see that by (3.27) with $\mathbf{g} = -\llbracket \mu \mathbf{D}(\psi) \mathbf{e}_N \rrbracket$ and $\mathbf{h} = -\llbracket \psi \rrbracket$

$$\widehat{v}_{N}(\xi', 0 -, \lambda) = \frac{L_{21}}{F(A, \lambda)} \Big\{ \mu_{+}(i\xi' \cdot \partial_{N}\widehat{\psi}'_{+}(\xi', 0, \lambda) - A^{2}\widehat{\psi}_{N+}(\xi', 0, \lambda)) \\
- \mu_{-}(i\xi' \cdot \partial_{N}\widehat{\psi}'_{-}(\xi', 0, \lambda) - A^{2}\widehat{\psi}_{N-}(\xi', 0, \lambda)) \\
+ \mu_{+}(A + B_{+})i\xi' \cdot (\widehat{\psi}'_{+}(\xi', 0, \lambda) - \widehat{\psi}'_{-}(\xi', 0, \lambda)) \\
- \mu_{+}A(B_{+} - A)(\widehat{\psi}_{N+}(\xi', 0, \lambda) - \widehat{\psi}_{N-}(\xi', 0, \lambda)) \Big\} \\
+ \frac{L_{22}}{F(A, \lambda)} \Big\{ A(2\mu_{+}\partial_{N}\widehat{\psi}_{N+} - 2\mu_{-}\partial_{N}\widehat{\psi}_{N-}) \\
- \mu_{+}(-A + B_{+})i\xi' \cdot (\widehat{\psi}'_{+}(\xi', 0, \lambda) - \widehat{\psi}'_{-}(\xi', 0, \lambda)) \\
+ \mu_{+}B_{+}(A + B_{+})(\widehat{\psi}_{N+}(\xi', 0, \lambda) - \widehat{\psi}_{N-}(\xi', 0, \lambda)) \Big\}.$$

Since
$$\widehat{u}_N^P(\xi',0,\lambda) = \widehat{u}_N^P(\xi',0-,\lambda)$$
 by $[\![\mathbf{u}^P]\!] = 0$, there holds
$$\widehat{u}_N^P(\xi',0,\lambda) = \widehat{\psi}_{N-}(\xi',0,\lambda) + \widehat{v}_N(\xi',0-,\lambda).$$

Combing this property with the above formula of $\widehat{v}_N(\xi', 0-, \lambda)$ and (A.1) in the appendix below yields the desired formula of $\widehat{u}_N^P(\xi', 0, \lambda)$. This completes the proof of Proposition 3.3.

3.2. Solution formulas for the hyperbolic-parabolic part. In this subsection, we introduce solution formulas of $(\eta^k, \mathbf{u}^k, \mathfrak{p}^k)$, k = 1, 2, for (2.8). System (2.8) can be written as

$$\begin{cases} \rho \lambda \mathbf{u}^k - \mathrm{Div}(\mu \mathbf{D}(\mathbf{u}^k) - \mathfrak{p}^k \mathbf{I}) = 0 & \text{in } \dot{\mathbf{R}}^N, \\ \operatorname{div} \mathbf{u}^k = 0 & \text{in } \dot{\mathbf{R}}^N, \\ \llbracket (\mu \mathbf{D}(\mathbf{u}^k) - \mathfrak{p}^k \mathbf{I}) \mathbf{e}_N \rrbracket = (\omega - \sigma \Delta') \eta^k \mathbf{e}_N & \text{on } \mathbf{R}^{N-1}, \\ \llbracket \mathbf{u}^k \rrbracket = 0 & \text{on } \mathbf{R}^{N-1}, \end{cases}$$

coupled with

$$\lambda \eta^k - u_N^k|_{x_N = 0} = z^k \quad \text{on } \mathbf{R}^{N-1}.$$
 (3.29)

In what follows, we apply the argumentation in Step 1 for the proof of Proposition 3.3 in the previous subsection. To this end, we set $\mathbf{g} = (0, \dots, 0, (\omega - \sigma \Delta') \eta^k)^\mathsf{T}$ and $\mathbf{h} = 0$ in (3.4). Then $G(\mathbf{g}, \mathbf{h})$ and $H(\mathbf{g}, \mathbf{h})$ in (3.23) are given by

$$G(\mathbf{g}, \mathbf{h}) = 0, \quad H(\mathbf{g}, \mathbf{h}) = -A(\omega + \sigma A^2)\hat{\eta}^k.$$
 (3.30)

Since $\mathbf{h} = 0$, we have by (3.18)

$$\beta_{J+} = \beta_{J-} \quad (J = 1, \dots, N).$$
 (3.31)

Combining this relation with (3.26) and (3.30) furnishes

$$\beta_{N+} = \beta_{N-} = -\frac{L_{22}}{F(A,\lambda)} A(\omega + \sigma A^2) \widehat{\eta}^k, \qquad (3.32)$$

and thus (3.13) gives

$$\widehat{u}_N^k(0) = -\frac{L_{22}}{F(A,\lambda)}A(\omega + \sigma A^2)\widehat{\eta}^k.$$

Inserting this formula into (3.29) yields

$$\lambda \widehat{\eta}^k + \frac{L_{22}}{F(A,\lambda)} A(\omega + \sigma A^2) \widehat{\eta}^k = \widehat{z}^k.$$

Solving this equation, we obtain

$$\widehat{\eta}^k = \frac{F(A,\lambda)}{L(A,\lambda)} \widehat{z}^k, \quad L(A,\lambda) = \lambda F(A,\lambda) + A(\omega + \sigma A^2)(D_+ + D_-), \tag{3.33}$$

where we have used $L_{22} = D_+ + D_-$ with $D_{\pm} = \mu_{\pm} B_{\pm} + \mu_{\mp} A$ given in (3.2). At this point, we note the following lemma.

Lemma 3.4. (1) Let $\varepsilon \in (0, \pi/2)$ and ρ_{\pm} be any positive constants. Then there exists a constant $\delta_0 \geq 1$ such that for any $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and $\lambda \in \Sigma_{\varepsilon}$ with $|\lambda| \geq \delta_0$

$$|L(A,\lambda)| \ge C_{\varepsilon}(|\lambda|^{1/2} + A)\{|\lambda|(|\lambda|^{1/2} + A)^2 + \sigma A^3\},$$
$$|\partial_{\varepsilon'}^{\alpha'} L(A,\lambda)^{-1}| \le C_{\alpha',\varepsilon}[(|\lambda|^{1/2} + A)\{|\lambda|(|\lambda|^{1/2} + A)^2 + \sigma A^3\}]^{-1}A^{-|\alpha'|},$$

where $\alpha' \in \mathbf{N}_0^{N-1}$ and C_{ε} , $C_{\alpha',\varepsilon}$ are positive constants independent of ξ' and λ . (2) Suppose that $\rho_- > \rho_+ > 0$. Let $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and $\lambda \in \mathbf{C}$ with $\Re \lambda \geq 0$. Then $L(A, \lambda) \neq 0$. *Proof.* (1) See [10, Lemma 6.1].

(2) The proof is similar to [7, Lemma 3.2], so that the detailed proof may be omitted. $\hfill\Box$

We continue to calculate the solution formulas for (2.8). By (3.32) and (3.33)

$$\beta_{N+} = \beta_{N-} = -\frac{L_{22}}{L(A,\lambda)} A(\omega + \sigma A^2) \hat{z}^k, \tag{3.34}$$

while by (3.26), (3.30), and (3.31)

$$i\xi' \cdot \beta'_{+} = i\xi' \cdot \beta'_{-} = -\frac{L_{12}}{L(A,\lambda)} A(\omega + \sigma A^{2}) \hat{z}^{k}. \tag{3.35}$$

It thus holds by (3.34) and (3.35) that

$$i\xi' \cdot \beta'_{\pm} \mp B_{\pm}\beta_{N\pm} = -\frac{A(\omega + \sigma A^2)}{L(A,\lambda)} (L_{12} \mp B_{\pm}L_{22}) \hat{z}^k,$$

which, combined with (3.19) and (3.20), furnishes

$$\alpha_{N\pm} = \mp \frac{1}{A - B_{\pm}} \cdot \frac{A(\omega + \sigma A^{2})}{L(A, \lambda)} (L_{12} \mp B_{\pm} L_{22}) \hat{z}^{k},$$

$$\gamma_{\pm} = \frac{\mu_{\pm} (A + B_{\pm})}{A} \cdot \frac{A(\omega + \sigma A^{2})}{L(A, \lambda)} (L_{12} \mp B_{\pm} L_{22}) \hat{z}^{k}.$$
(3.36)

By the first equation of (3.14) and the above formula of γ_{\pm} , we have

$$\alpha_{j\pm} = \frac{i\xi_j}{(A - B_{\pm})A} \cdot \frac{A(\omega + \sigma A^2)}{L(A, \lambda)} (L_{12} \mp B_{\pm} L_{22}) \hat{z}^k \quad (j = 1, \dots, N - 1). \quad (3.37)$$

Noting $g_j = 0$ for j = 1, ..., N - 1, we have by (3.16), (3.31), (3.34), and (3.37)

$$\beta_{j+} = \beta_{j-} =$$

$$-\frac{i\xi_{j}(\omega+\sigma A^{2})}{EL(A,\lambda)}\left[(\mu_{+}+\mu_{-})L_{12}+\left\{\mu_{+}(A-B_{+})-\mu_{-}(A-B_{-})\right\}L_{22}\right]\hat{z}^{k},\quad(3.38)$$

together with $E = \mu_+ B_+ + \mu_- B_-$ given in (3.2).

Let us define for $j = 1, \ldots, N-1$

$$\mathcal{I}_{j\pm}(\xi',\lambda) = i\xi_j(\omega + \sigma A^2)(L_{12} \mp B_{\pm}L_{22}),$$

$$\mathcal{I}_{N\pm}(\xi',\lambda) = \mp A(\omega + \sigma A^2)(L_{12} \mp B_{\pm}L_{22}),$$

$$\mathcal{J}_{j}(\xi',\lambda) = -i\xi_{j}(\omega + \sigma A^{2}) \left[(\mu_{+} + \mu_{-})L_{12} + \{\mu_{+}(A - B_{+}) - \mu_{-}(A - B_{-})\} L_{22} \right],$$

$$\mathcal{J}_{N}(\xi',\lambda) = -EL_{22}A(\omega + \sigma A^{2}).$$

where L_{11} , L_{12} , L_{21} , and L_{22} are given in (3.25). Recall $\mathcal{M}_{\pm}(a)$ in (3.2). Then, in view of (3.13), we have achieved by (3.34), (3.36), (3.37), and (3.38)

$$\widehat{u}_{m\pm}^{k}(x_{N}) = \frac{\mathcal{I}_{m\pm}(\xi',\lambda)}{L(A,\lambda)} \mathcal{M}_{\pm}(\pm x_{N}) \widehat{z}^{k} + \frac{\mathcal{J}_{m}(\xi',\lambda)}{EL(A,\lambda)} e^{\mp B_{\pm}x_{N}} \widehat{z}^{k}$$
(3.39)

for $\pm x_N > 0$ and m = 1, ..., N, with the pressure $\widehat{\mathfrak{p}}_{\pm}^k(x_N) = \gamma_{\pm} e^{\mp x_N} \ (\pm x_N > 0)$. For the above $\widehat{\eta}^k$, $\widehat{u}_{m+}^k(x_N)$, and $\widehat{\mathfrak{p}}_{+}^k(x_N)$, we define

$$\eta^{k} = \mathcal{F}_{\xi'}^{-1} [\widehat{\eta}^{k}(\xi', \lambda)](x'), \quad u_{m\pm}^{k} = \mathcal{F}_{\xi'}^{-1} [\widehat{u}_{m\pm}^{k}(\xi', x_{N}, \lambda)](x'),$$

$$\mathfrak{p}_{\pm}^{k} = \mathcal{F}_{\xi'}^{-1} [\widehat{\mathfrak{p}}_{\pm}^{k}(\xi', x_{N}, \lambda)](x').$$

Then setting

$$u_m^k = u_{m+}^k \mathbb{1}_{\mathbf{R}_+^N} + u_{m-}^k \mathbb{1}_{\mathbf{R}_-^N} \quad \text{and} \quad \mathfrak{p}^k = \mathfrak{p}_+^k \mathbb{1}_{\mathbf{R}_+^N} + \mathfrak{p}_-^k \mathbb{1}_{\mathbf{R}_-^N},$$

we observe that η^k , $\mathbf{u}^k = (u_1^k, \dots, u_N^k)^\mathsf{T}$, and \mathfrak{p}^k become a solution to (2.8). This completes the calculation of solution formulas for (2.8).

3.3. Representation formulas for (1.1). In this subsection, we give the representation formulas of solutions for (1.1). To this end, we first consider $\mathcal{H}_Z^1(t)d$, $\mathcal{H}_Z^2(t)\mathbf{f}$, $\mathcal{U}_Z^1(t)d$, and $\mathcal{U}_Z^2(t)\mathbf{f}$ given in (2.10). Together with Proposition 3.3, inserting $\hat{\eta}^1$ and $\hat{\eta}^2$ of (3.33) into $\hat{\mathcal{H}}_Z^1(t;\Gamma)d$ and $\hat{\mathcal{H}}_Z^2(t;\Gamma)\mathbf{f}$ in (2.9), respectively, yields

$$\begin{split} \widehat{\mathcal{H}}_{Z}^{1}(t;\Gamma)d &= \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{F(A,\lambda)}{L(A,\lambda)} \, d\lambda \, \widehat{d}(\xi'), \\ \widehat{\mathcal{H}}_{Z}^{2}(t;\Gamma)\mathbf{f} &= \sum_{\mathfrak{a},\mathfrak{b} \in \{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \left(\frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda) \mathcal{M}_{\mathfrak{b}}(y_{N})}{A(B_{\mathfrak{b}} + A)L(A,\lambda)} \, d\lambda \right) \widehat{f}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &+ \sum_{\mathfrak{a},\mathfrak{b} \in \{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \left(\frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda) e^{-B_{\mathfrak{b}}y_{N}}}{AB_{\mathfrak{b}}(B_{\mathfrak{b}} + A)L(A,\lambda)} \, d\lambda \right) \widehat{f}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N}. \end{split}$$

Let us define for $\pm x_N > 0$ and m = 1, ..., N

$$\widehat{\mathcal{U}}_{Z,m\pm}^{1}(t;\Gamma)d = \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \widehat{u}_{m\pm}^{1}(\xi', x_{N}, \lambda) d\lambda,$$

$$\widehat{\mathcal{U}}_{Z,m\pm}^{2}(t;\Gamma)\mathbf{f} = \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \widehat{u}_{m\pm}^{2}(\xi', x_{N}, \lambda) d\lambda.$$

Together with Proposition 3.3, inserting $\widehat{u}_{m\pm}^1$ and $\widehat{u}_{m\pm}^2$ of (3.39) into $\widehat{\mathcal{U}}_{Z,m\pm}^1(t;\Gamma)d$ and $\widehat{\mathcal{U}}_{Z,m\pm}^2(t;\Gamma)\mathbf{f}$, respectively, yields the following formulas: for $\pm x_N > 0$ and $m = 1, \ldots, N$

$$\widehat{\mathcal{U}}_{Z,m\pm}^{1}(t;\Gamma)d = \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\mathcal{I}_{m\pm}(\xi',\lambda)}{L(A,\lambda)} \mathcal{M}_{\pm}(\pm x_{N}) d\lambda \, \widehat{d}(\xi')
+ \frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\mathcal{J}_{m}(\xi',\lambda)}{EL(A,\lambda)} e^{\mp B_{\pm}x_{N}} d\lambda \, \widehat{d}(\xi'),$$

and furthermore,

$$\begin{split} \widehat{\mathcal{U}}_{Z,m\pm}^{2}(t;\Gamma)\mathbf{f} &= \sum_{\mathfrak{a},\mathfrak{b}\in\{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \left(\frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda) \mathcal{I}_{m\pm}(\xi',\lambda) \mathcal{M}_{\pm}(\pm x_{N}) \mathcal{M}_{\mathfrak{b}}(y_{N})}{A(B_{\mathfrak{b}} + A)F(A,\lambda)L(A,\lambda)} d\lambda \right) \\ &\times \widehat{f}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &+ \sum_{\mathfrak{a},\mathfrak{b}\in\{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \left(\frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda) \mathcal{I}_{m\pm}(\xi',\lambda) \mathcal{M}_{\pm}(\pm x_{N}) e^{-B_{\mathfrak{b}}y_{N}}}{AB_{\mathfrak{b}}(B_{\mathfrak{b}} + A)F(A,\lambda)L(A,\lambda)} d\lambda \right) \\ &\times \widehat{f}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \end{split}$$

$$+ \sum_{\mathfrak{a},\mathfrak{b}\in\{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \left(\frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda) \mathcal{J}_{m}(\xi',\lambda) e^{\mp B_{\pm}x_{N}} \mathcal{M}_{\mathfrak{b}}(y_{N})}{A(B_{\mathfrak{b}}+A)F(A,\lambda)EL(A,\lambda)} d\lambda \right) \times \hat{f}_{j}(\xi',\mathfrak{a}y_{N}) dy_{N}$$

$$+ \sum_{\mathfrak{a},\mathfrak{b}\in\{+,-\}} \sum_{j=1}^{N} \int_{0}^{\infty} \left(\frac{\varphi_{Z}(\xi')}{2\pi i} \int_{\Gamma} e^{\lambda t} \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda) \mathcal{J}_{m}(\xi',\lambda) e^{\mp B_{\pm}x_{N}} e^{-B_{\mathfrak{b}}y_{N}}}{AB_{\mathfrak{b}}(B_{\mathfrak{b}}+A)F(A,\lambda)EL(A,\lambda)} d\lambda \right) \times \hat{f}_{j}(\xi',\mathfrak{a}y_{N}) dy_{N}.$$

One defines

$$\begin{split} \widehat{\mathcal{U}}_{Z,m}^1(t;\Gamma)d &= (\widehat{\mathcal{U}}_{Z,m+}^1(t;\Gamma)d)\mathbb{1}_{\mathbf{R}_+^N} + (\widehat{\mathcal{U}}_{Z,m-}^1(t;\Gamma)d)\mathbb{1}_{\mathbf{R}_-^N}, \\ \widehat{\mathcal{U}}_{Z,m}^2(t;\Gamma)\mathbf{f} &= (\widehat{\mathcal{U}}_{Z,m+}^2(t;\Gamma)\mathbf{f})\mathbb{1}_{\mathbf{R}_-^N} + (\widehat{\mathcal{U}}_{Z,m-}^2(t;\Gamma)\mathbf{f})\mathbb{1}_{\mathbf{R}_-^N}, \end{split}$$

and then there holds the following relation between these formulas and $\widehat{\mathcal{U}}_Z^1(t;\Gamma)d$, $\widehat{\mathcal{U}}_Z^2(t;\Gamma)\mathbf{f}$ given in (2.9):

$$\widehat{\mathcal{U}}_{Z}^{1}(t;\Gamma)d = (\widehat{\mathcal{U}}_{Z,1}^{1}(t;\Gamma)d, \dots, \widehat{\mathcal{U}}_{Z,N}^{1}(t;\Gamma)d)^{\mathsf{T}},$$

$$\widehat{\mathcal{U}}_{Z}^{2}(t;\Gamma)\mathbf{f} = (\widehat{\mathcal{U}}_{Z,1}^{2}(t;\Gamma)\mathbf{f}, \dots, \widehat{\mathcal{U}}_{Z,N}^{2}(t;\Gamma)\mathbf{f})^{\mathsf{T}}.$$

Summing up the above argumentation, we have obtained the representation formulas of $\mathcal{H}_{Z}^{1}(t)d$, $\mathcal{H}_{Z}^{2}(t)\mathbf{f}$, $\mathcal{U}_{Z}^{1}(t)d$, and $\mathcal{U}_{Z}^{2}(t)\mathbf{f}$ given in (2.10) from the formulas of $\widehat{\mathcal{H}}_{Z}^{1}(t;\Gamma)d$, $\widehat{\mathcal{H}}_{Z}^{2}(t;\Gamma)\mathbf{f}$, $\widehat{\mathcal{U}}_{Z}^{1}(t;\Gamma)d$, and $\widehat{\mathcal{U}}_{Z}^{2}(t;\Gamma)\mathbf{f}$ as above. Furthermore, those representation formulas of $\mathcal{H}_{Z}^{1}(t)d$, $\mathcal{H}_{Z}^{2}(t)\mathbf{f}$, $\mathcal{U}_{Z}^{1}(t)d$, and $\mathcal{U}_{Z}^{2}(t)\mathbf{f}$ give the representation formulas of solutions for (1.1) by the relation (2.11).

Finally, we introduce another useful formula of $L(A, \lambda)$.

Lemma 3.5. Let $L(A, \lambda)$ be given in (3.33) and set

$$\mathcal{L}_A(\lambda) = \lambda^2 + \frac{4AD_+D_-}{(\rho_+ + \rho_-)(D_+ + D_-)}\lambda + \alpha A + \widetilde{\sigma}A^3, \quad \widetilde{\sigma} = \frac{\sigma}{\rho_+ + \rho_-},$$

where D_{\pm} and α are defined in (3.2) and (2.2), respectively. Then

$$L(A, \lambda) = (\rho_+ + \rho_-)(D_+ + D_-)\mathcal{L}_A(\lambda).$$

Proof. The desired relation follows from an elementary calculation, so that the detailed proof may be omitted. \Box

4. Analysis of boundary symbol

We assume $\rho_- > \rho_+ > 0$ throughout this section and analyze mainly the boundary symbol $\mathcal{L}_A(\lambda)$ introduced in the last part of the previous section. Note that α in (2.2) is positive by the assumption $\rho_- > \rho_+ > 0$.

4.1. Analysis of low frequency part. Recalling θ_2 , λ_1 given in (2.1) and $z_0 = \min\{\mu_+/\rho_+, \mu_-/\rho_-\}$, we define for $A = |\xi'|$

$$z_1^{\pm} = -\frac{z_0}{2}A^2 \pm i\frac{z_0}{4}A^2, \quad z_2^{\pm} = \lambda_1 e^{\pm i(\pi - \theta_2)},$$
 (4.1)

and also

$$\widehat{\Gamma}_{1}^{\pm} = \{ \lambda \in \mathbf{C} : \lambda = -\frac{z_{0}}{2} A^{2} + \frac{z_{0}}{4} A^{2} e^{\pm is}, \ 0 \le s \le \frac{\pi}{2} \},$$

$$\widehat{\Gamma}_{2}^{\pm} = \{ \lambda \in \mathbf{C} : \lambda = z_{1}^{\pm} (1 - s) + z_{2}^{\pm} s, \ 0 \le s \le 1 \},$$

$$\widehat{\Gamma}_3 = \{ \lambda \in \mathbf{C} : \lambda = \lambda_1 e^{is}, -(\pi - \theta_2) \le s \le \pi - \theta_2 \}. \tag{4.2}$$

In addition, we set

$$\mathcal{F}_A(\lambda) = (\lambda - \zeta_+)(\lambda - \zeta_-), \quad \mathcal{G}_A(\lambda) = \mathcal{L}_A(\lambda) - \mathcal{F}_A(\lambda),$$

where ζ_{\pm} and $\mathcal{L}_A(A)$ are given in Theorem 2.3 and Lemma 3.5, respectively. Then

$$\mathcal{G}_A(\lambda) = \frac{4AD_+D_-}{(\rho_+ + \rho_-)(D_+ + D_-)} \lambda + \widetilde{\sigma}A^3$$
$$-2\sqrt{2}\alpha^{1/4}\beta A^{5/4}\lambda + 2\sqrt{2}\alpha^{3/4}\beta A^{7/4} - 4\alpha^{1/2}\beta^2 A^{10/4}$$
(4.3)

and the following lemma holds.

Lemma 4.1. There exists a constant $A_1 \in (0,1)$ such that $|\mathcal{F}_A(\lambda)| > |\mathcal{G}_A(\lambda)|$ for $A \in (0,A_1)$ and $\lambda \in \widehat{\Gamma}_1^+ \cup \widehat{\Gamma}_1^- \cup \widehat{\Gamma}_2^+ \cup \widehat{\Gamma}_2^- \cup \widehat{\Gamma}_3$.

Proof. Case 1: $\lambda \in \widehat{\Gamma}_1^+ \cup \widehat{\Gamma}_1^-$. Let $\lambda = -(z_0/2)A^2 + (z_0/4)A^2e^{is}$ $(-\pi/2 \le s \le \pi/2)$. It is clear that

$$|\mathcal{F}_A(\lambda)| \ge CA$$
 for $A \in (0, A_1)$,

with a sufficiently small A_1 and a positive constant C independent of A and λ . Next, we estimate $|\mathcal{G}_A(\lambda)|$ from above. Since

$$D_{\pm} = A \left(\mu_{\pm} \sqrt{\frac{\rho_{\pm}}{\mu_{\pm}} \left(-\frac{z_0}{2} + \frac{z_0}{4} e^{is} \right) + 1} + \mu_{\mp} \right),$$

$$\Re \left(\sqrt{\frac{\rho_{\pm}}{\mu_{\pm}} \left(-\frac{z_0}{2} + \frac{z_0}{4} e^{is} \right) + 1} \right) > 0,$$

there holds

$$\left| \frac{D_{+}D_{-}}{D_{+} + D_{-}} \right| \leq \frac{|D_{+}||D_{-}|}{(\Re D_{+}) + (\Re D_{-})} \leq CA \quad \text{for } A > 0.$$

One thus sees that

$$|\mathcal{G}_A(\lambda)| \le CA^{7/4}$$
 for $A \in (0,1)$,

which implies $|\mathcal{F}_A(\lambda)| > |\mathcal{G}_A(\lambda)|$ for $A \in (0, A_1)$ when A_1 is sufficiently small.

Case 2: $\lambda \in \widehat{\Gamma}_2^+ \cup \widehat{\Gamma}_2^-$. We consider $\lambda \in \widehat{\Gamma}_2^+$ only. Let $\lambda = z_1^+(1-s) + z_2^+s$ $(0 \le s \le 1)$. We write $\lambda = -a + bi$ $(a, b \ge 0)$, that is,

$$a = \frac{z_0}{2}A^2(1-s) + \lambda_1(\cos\theta_2)s, \quad b = \frac{z_0}{4}A^2(1-s) + \lambda_1(\sin\theta_2)s.$$

We then observe that

$$|\lambda - \zeta_{\pm}|^2 = (-a + \sqrt{2}\alpha^{1/4}\beta A^{5/4})^2 + (b \mp \alpha^{1/2}A^{1/2} \pm \sqrt{2}\alpha^{1/4}\beta A^{5/4})^2$$
$$= a^2 + b^2 + \alpha A \mp 2\alpha^{1/2}A^{1/2}b + O(A^{5/4}) \quad \text{as } A \to 0.$$
(4.4)

From this, we immediately see that there exists a constant $A_1 \in (0,1)$ such that

$$|\lambda - \zeta_-| \ge C(|\lambda| + A^{1/2})$$
 for $A \in (0, A_1)$.

Since $2\alpha^{1/2}A^{1/2}b \le \varepsilon \alpha A + b^2/\varepsilon$ ($\varepsilon > 0$), choosing $\varepsilon = 2/3$ furnishes

$$a^2 + b^2 + \alpha A - 2\alpha^{1/2}A^{1/2}b \ge a^2 - \frac{1}{2}b^2 + \frac{1}{3}\alpha A. \tag{4.5}$$

In addition,

$$b = \frac{z_0}{4}A^2(1-s) + \tan\theta_2 \cdot \lambda_1(\cos\theta_2)s = \frac{1}{2}\left\{\frac{z_0}{2}A^2(1-s) + \frac{1}{4}\lambda_1(\cos\theta_2)s\right\} \le \frac{1}{2}a,$$

which implies

$$a^{2} - \frac{1}{2}b^{2} = \frac{1}{4}(a^{2} + b^{2}) + \frac{3}{4}(a^{2} - b^{2}) \ge \frac{1}{4}(a^{2} + b^{2}).$$

It thus holds by (4.4) and (4.5) that $|\lambda - \zeta_+| \ge C(|\lambda| + A^{1/2})$ when A is small enough, and therefore

$$|\mathcal{F}_A(\lambda)| \ge C(|\lambda| + A^{1/2})^2 \quad \text{for } A \in (0, A_1).$$
 (4.6)

In particular,

$$|\mathcal{F}_A(\lambda)| \ge CA^{1/2}(|\lambda| + A^{1/2}) \quad \text{for } A \in (0, A_1).$$
 (4.7)

Next, we estimate $|\mathcal{G}_A(\lambda)|$ from above. By Lemma 3.2 and (4.3), we have for $A \in (0,1)$

$$|\mathcal{G}_A(\lambda)| \le C(A|\lambda|(|\lambda|^{1/2} + A) + A^{5/4}|\lambda| + A^{7/4})$$

$$= CA^{3/4}(A^{1/4}|\lambda|(|\lambda|^{1/2} + A) + A^{1/2}|\lambda| + A). \tag{4.8}$$

Since $|\lambda| \leq \lambda_1$, one sees for $A \in (0,1)$ that

$$A^{1/4}|\lambda|(|\lambda|^{1/2} + A) \le |\lambda|(|\lambda|^{1/2} + A^{1/2}) \le \lambda_1^{1/2}(|\lambda| + \lambda_1^{1/2}A) \le C(|\lambda| + A^{1/2})$$

and that $A^{1/2}|\lambda|+A\leq |\lambda|+A^{1/2}$. Combining these inequalities with (4.8) furnishes

$$|\mathcal{G}_A(\lambda)| \le CA^{3/4}(|\lambda| + A^{1/2})$$
 for $A \in (0, 1)$,

which implies $|\mathcal{F}_A(\lambda)| > |\mathcal{G}_A(\lambda)|$ for $A \in (0, A_1)$ when A_1 is sufficiently small.

Case 3: $\lambda \in \widehat{\Gamma}_3$. Let $\lambda = \lambda_1 e^{is} (-(\pi - \theta_2) \le s \le \pi - \theta_2)$. Then

$$|\mathcal{F}_A(\lambda)| \geq C$$
 for $A \in (0, A_1)$,

with a sufficiently small A_1 and a positive constant C independent of A and λ . Since (4.8) is valid for $\lambda \in \widehat{\Gamma}_3$ when $A \in (0,1)$, there holds

$$|\mathcal{G}_A(\lambda)| < CA^{3/4}$$
 for $A \in (0,1)$.

It therefore holds that $|\mathcal{F}_A(\lambda)| > |\mathcal{G}_A(\lambda)|$ for $A \in (0, A_1)$ when A_1 is sufficiently small. This completes the proof of Lemma 4.1.

By Lemma 4.1 and Rouché's theorem, we immediately have

Proposition 4.2. Let A_1 be the positive constant given in Lemma 4.1 and $A \in (0, A_1)$. Let K be the region enclosed by $\widehat{\Gamma}_1^+ \cup \widehat{\Gamma}_1^- \cup \widehat{\Gamma}_2^+ \cup \widehat{\Gamma}_2^- \cup \widehat{\Gamma}_3$. Then $\mathcal{L}_A(\lambda)$ has two zeros in K.

Recalling $\widehat{\Gamma}_{res}^{\pm}$ given in Theorem 2.3, we prove

Lemma 4.3. There exists a constant $A_2 \in (0,1)$ such that $|\mathcal{F}_A(\lambda)| > |\mathcal{G}_A(\lambda)|$ for $A \in (0,A_2)$ and $\lambda \in \widehat{\Gamma}^+_{Res} \cup \widehat{\Gamma}^-_{Res}$.

Proof. We consider $\lambda \in \widehat{\Gamma}_{Res}^+$ only. Let $\lambda = \zeta_+ + A^{6/4}e^{is}$ $(0 \le s \le 2\pi)$. It is clear that

$$|\mathcal{F}_A(\lambda)| \ge CA^{8/4}$$
 for $A \in (0, A_2)$,

with a sufficiently small A_2 and a positive constant C independent of A and λ . Next, we estimate $|\mathcal{G}_A(\lambda)|$ from above. It holds that

$$B_{\pm} = \sqrt{\frac{\rho_{\pm}}{\mu_{\pm}}} e^{i(\pi/4)} \alpha^{1/4} A^{1/4} (1 + O(A^{3/4}))$$
 as $A \to 0$,

which implies

$$D_{\pm} = \sqrt{\rho_{\pm}\mu_{\pm}}e^{i(\pi/4)}\alpha^{1/4}A^{1/4} + O(A)$$
 as $A \to 0$.

Therefore,

$$\frac{D_{+}D_{-}}{D_{+} + D_{-}} = \frac{\sqrt{\rho_{+}\mu_{+}}\sqrt{\rho_{-}\mu_{-}}}{\sqrt{\rho_{+}\mu_{+}} + \sqrt{\rho_{-}\mu_{-}}} e^{i(\pi/4)} \alpha^{1/4} A^{1/4} + O(A) \quad \text{as } A \to 0.$$

From this, recalling the definition of β given in (2.2), we have

$$\frac{4AD_+D_-}{(\rho_+ + \rho_-)(D_+ + D_-)}\lambda = (-1+i) \cdot 2\sqrt{2}\alpha^{3/4}\beta A^{7/4} + O(A^{10/4}) \quad \text{as } A \to 0.$$

In addition,

$$-2\sqrt{2}\alpha^{1/4}\beta A^{5/4}\lambda = -i \cdot 2\sqrt{2}\alpha^{3/4}\beta A^{7/4} + O(A^{10/4}) \quad \text{as } A \to 0.$$

Combining these two formulas with (4.3) shows that

$$\mathcal{G}_A(\lambda) = O(A^{10/4})$$
 as $A \to 0$,

which implies $|\mathcal{F}_A(\lambda)| > |\mathcal{G}_A(\lambda)|$ for $A \in (0, A_2)$ when A_2 is sufficiently small. This completes the proof of Lemma 4.3.

We now obtain

Proposition 4.4. Let A_1 and A_2 be the positive constants given in Lemma 4.1 and Lemma 4.3, respectively. Then there exists a constant $A_3 \in (0, \min\{A_1, A_2\})$ such that the following assertions hold.

(1) Let $A \in (0, A_3)$. Then for the K in Proposition 4.2

$$\widehat{\Gamma}_{\mathrm{Res}}^{+} \subset K \cap \{\lambda \in \mathbf{C} \setminus \{0\} : \frac{\pi}{2} < \arg \lambda < \frac{3}{4}\pi\},$$

$$\widehat{\Gamma}_{\mathrm{Res}}^{-} \subset K \cap \{\lambda \in \mathbf{C} \setminus \{0\} : -\frac{3}{4}\pi < \arg \lambda < -\frac{\pi}{2}\}.$$

- (2) Let $A \in (0, A_3)$ and K_{\pm} be the regions enclosed by $\widehat{\Gamma}_{Res}^{\pm}$, respectively. Then $\mathcal{L}_A(\lambda)$ has a simple zero denoted by λ_+ in K_+ and another simple zero denoted by λ_- in K_- .
- (3) Let $\mathcal{L}'_A(\lambda)$ be the derivative of $\mathcal{L}_A(\lambda)$ with respect to λ . Then the inequalities

$$|\mathcal{L}'_{A}(\lambda_{+})| \ge CA^{1/2}, \quad |\mathcal{L}'_{A}(\lambda_{-})| \ge CA^{1/2}$$

hold for $A \in (0, A_3)$ and a positive constant C independent of A.

Remark 4.5. The zeros λ_{\pm} of $\mathcal{L}_A(\lambda)$ satisfy

$$\lambda_{\pm} = \zeta_{\pm} + O(A^{6/4}) = \pm i\alpha^{1/2}A^{1/2} - \sqrt{2}\alpha^{1/4}\beta(1\pm i)A^{5/4} + O(A^{6/4})$$
 as $A \to 0$.

When gravity is not taken into account, i.e. $\alpha = 0$, the asymptotics of the zeros of $\mathcal{L}_A(\lambda)$ are obtained in [3].

Proof of Proposition 4.4. (1) The desired properties can be proved by an elementary calculation, so that the detailed proof may be omitted.

- (2) The result follows from Lemma 4.3 and Rouché's theorem immediately.
- (3) Since $\mathcal{L}'_A(\lambda_{\pm}) = \mathcal{F}'_A(\lambda_{\pm}) + \mathcal{G}'_A(\lambda_{\pm})$, one has the desired inequalities immediately by direct calculations together with (4.3). This completes the proof of Proposition 4.4.

4.2. Analysis of high frequency part. Let us define

$$\Lambda(a,b) = \{ \lambda \in \mathbf{C} : \lambda = x + yi, -a \le x \le 0, -b \le y \le b \} \quad (a,b \ge 0). \tag{4.9}$$

We then have

Proposition 4.6. Let a, b > 0. Then there exists a sufficiently large positive number $A_{\text{high}} = A_{\text{high}}(a, b)$ such that the following assertions hold.

(1) For any $A \geq A_{\text{high}}$ and $\lambda \in \Lambda(a, b)$,

$$C_1 A \leq \Re B_+ \leq |B_+| \leq C_2 A$$
,

where C_1 and C_2 are positive constants independent of A and λ .

(2) For any $A \geq A_{\text{high}}$ and $\lambda \in \Lambda(a, b)$,

$$|F(A,\lambda)| > CA^3$$
, $|L(A,\lambda)| > C\sigma A^4$,

where C is a positive constant independent of A, λ , and σ .

Proof. (1) See [7, Lemma 5.3].

(2) First, we estimate $|F(A,\lambda)|$ from below. Since

$$B_{\pm} = A + O(\frac{1}{A}) \quad \text{as } A \to \infty, \tag{4.10}$$

it holds that

$$F(A, \lambda) = 4(\mu_+ + \mu_-)^2 A^3 + O(A)$$
 as $A \to \infty$.

This implies the desired inequality for $|F(A, \lambda)|$.

Next, we estimate $|L(A,\lambda)|$ from below. Recall the formula of $\mathcal{L}_A(\lambda)$ in Lemma 3.5. The asymptotics (4.10) gives

$$D_{\pm} = (\mu_{+} + \mu_{-})A + O(\frac{1}{A})$$
 as $A \to \infty$,

which implies

$$\mathcal{L}_A(\lambda) = \widetilde{\sigma}A^3 + O(A^2)$$
 as $A \to \infty$.

Thus, from the formula of $L(A, \lambda)$ in Lemma 3.5, we see that

$$L(A, \lambda) = 2(\mu_{+} + \mu_{-})\sigma A^{4} + O(A^{3})$$
 as $A \to \infty$.

This yields the desired inequality of $|L(A, \lambda)|$, which completes the proof of Proposition 4.6.

Next, we consider $A \in [M_1, M_2]$ for $M_2 > M_1 > 0$.

Proposition 4.7. Let b > 0 and $M_2 > M_1 > 0$. Then there exist $a_0 \in (0,1)$ such that the following assertions hold.

(1) For any $A \in [M_1, M_2]$ and $\lambda \in \Lambda(a_0, b)$,

$$C_1 \leq \Re B_+ \leq |B_+| \leq C_2$$
,

where C_1 and C_2 are positive constants independent of A and λ .

(2) For any $A \in [M_1, M_2]$ and $\lambda \in \Lambda(a_0, b)$,

$$|F(A,\lambda)| \ge C, \quad |L(A,\lambda)| \ge C,$$

where C is a positive constant independent of A and λ .

Proof. (1) The desired inequalities can be proved by an elementary calculation, so that the detailed proof may be omitted.

(2) By Lemma 3.1, we see that $F(A,\lambda) \neq 0$ for A > 0 and $\Re \lambda \geq 0$. Then the continuity of $F(A,\lambda)$ and the compactness of $[M_1,M_2] \times \Lambda(0,b)$ implies there exists an $m := \min\{|F(A,\lambda)| : A \in [M_1,M_2], \lambda \in \Lambda(0,b)\} > 0$. Choosing a sufficiently small $a_1 \in (0,1)$, we see that $F(A,\lambda)$ is uniform continuous on $[M_1,M_2] \times \Lambda(a_1,b)$. Thus there exists an $a_0 \in (0,a_1]$ such that

$$|F(A,\lambda)| \ge \frac{m}{2}$$
 for $(A,\lambda) \in [M_1, M_2] \times \Lambda(a_0,b)$,

which implies the desired inequality of $|F(A,\lambda)|$ holds. Analogously, the inequality for $|L(A,\lambda)|$ follows from Lemma 3.4 (2). This completes the proof of Proposition 4.7.

5. Time-decay estimates for low frequency part

This section proves Theorem 2.3. Suppose $\rho_- > \rho_+ > 0$ throughout this section. Let us denote the points of intersection between $\lambda = se^{\pm i(3\pi/4)}$ ($s \ge 0$) and Γ_0^{\pm} given in (2.3) by z_3^{\pm} . Then we define

$$\begin{split} \widehat{\Gamma}_4^{\pm} &= \{\lambda \in \mathbf{C} : \lambda = z_1^{\pm}(1-s) + z_3^{\pm}s, 0 \leq s \leq 1\}, \\ \widehat{\Gamma}_5^{\pm} &= \{\lambda \in \mathbf{C} : \lambda = z_3^{\pm} + se^{\pm i(\pi-\theta_1)}, s \geq 0\}, \end{split}$$

where z_1^{\pm} are given in (4.1). Let A_3 be the positive constant given in Proposition 4.4 and let $A_0 \in (0, A_3)$. By Propositions 4.2 and 4.4 together with Cauchy's integral theorem, we see for the operators of (2.10) that

$$\mathcal{S}^1_{A_0}(t)d = \sum_{\mathfrak{a} \in \{+,-\}} \sum_{j \in \{\text{Res},1,4,5\}} \mathcal{S}^1_{A_0}(t;\Gamma_j^{\mathfrak{a}})d, \quad \mathcal{S}^1_{A_0}(t;\Gamma_j^{\mathfrak{a}})d := \mathcal{F}^{-1}_{\xi'}[\widehat{\mathcal{S}}^1_{A_0}(t;\widehat{\Gamma}_j^{\mathfrak{a}})d](x'),$$

$$\mathcal{S}_{A_0}^2(t)\mathbf{f} = \sum_{\mathfrak{a}\in\{+,-\}} \sum_{j\in\{\text{Res},1,4,5\}} \mathcal{S}_{A_0}^2(t;\Gamma_j^{\mathfrak{a}})\mathbf{f}, \quad \mathcal{S}_{A_0}^2(t;\Gamma_j^{\mathfrak{a}})\mathbf{f} := \mathcal{F}_{\xi'}^{-1}[\widehat{\mathcal{S}}_{A_0}^2(t;\widehat{\Gamma}_j^{\mathfrak{a}})\mathbf{f}](x'),$$

$$(5.1)$$

where $S \in \{\mathcal{H}, \mathcal{U}\}$. Here we have used $\widehat{\Gamma}_1^{\pm}$ in (4.2) and the symbols $\widehat{S}_{A_0}^1(t; \widehat{\Gamma}_j^{\mathfrak{a}})d$, $\widehat{S}_{A_0}^2(t; \widehat{\Gamma}_j^{\mathfrak{a}})\mathbf{f}$ introduced in Subsection 3.3.

At this point, we introduce several lemmas used in the following argumentation. From [7, Lemmas 4.3 and 4.4], we have the following two lemmas.

Lemma 5.1. Let $c_1, c_2, d \ge 0$ and $\nu_1, \nu_2 > 0$. Then there exists a positive constant C such that for any $\tau \ge 0$, $a \ge 0$, and $Z \ge 0$

$$e^{-c_1(Z^{\nu_1})\tau}Z^de^{-c_2(Z^{\nu_2})a} \le C(\tau^{d/\nu_1} + a^{d/\nu_2})^{-1}$$

Lemma 5.2. Let $1 \le p, q \le \infty$ and r be the dual exponent of p. Suppose that a > 0, $b_1 > 0$, and $b_2 > 0$.

(1) For $f \in L_p(0, \infty)$, $x_N > 0$, and $\tau > 0$, set

$$I(x_N, \tau) = \int_0^\infty \frac{f(y_N)}{\tau^a + (x_N)^{b_1} + (y_N)^{b_2}} \, dy_N.$$

Then there exists a positive constant C, independent of f, such that for any $\tau > 0$

$$||I(\cdot,\tau)||_{L_q(0,\infty)} \le C\tau^{-a\left(1-\frac{1}{b_1q}-\frac{1}{b_2r}\right)}||f||_{L_p(0,\infty)},$$

provided that $b_1q > 1$ and $b_2r(1 - 1/(b_1q)) > 1$.

(2) For $f \in L_p(0,\infty)$ and $\tau > 0$, set

$$J(\tau) = \int_0^\infty \frac{f(y_N)}{\tau^a + (y_N)^{b_2}} dy_N.$$

Then there exists a positive constant C, independent of f, such that for any $\tau > 0$

$$|J(\tau)| \le C\tau^{-a\left(1-\frac{1}{b_2r}\right)} ||f||_{L_n(0,\infty)},$$

provided that $b_2r > 1$.

Next, we introduce time-decay estimates arising in the study of an evolution equation with the fractional Laplacian.

Lemma 5.3. Let $1 \le p \le q \le \infty$, $\theta > 0$, and $\nu > 0$. Then the following assertions hold.

(1) For any $\tau > 0$ and $\varphi \in L_p(\mathbf{R}^{N-1})$

$$\|\mathcal{F}_{\xi'}^{-1}[e^{-\nu\tau|\xi'|^{\theta}}\widehat{\varphi}(\xi')]\|_{L_{q}(\mathbf{R}^{N-1})} \le C\tau^{-\frac{N-1}{\theta}(\frac{1}{p}-\frac{1}{q})}\|\varphi\|_{L_{p}(\mathbf{R}^{N-1})},$$

with a positive constant C independent of τ and φ .

(2) If it is assumed that $1 \le p \le 2$ additionally, then for any $\tau > 0$ and $\varphi \in L_p(\mathbf{R}^{N-1})$

$$\|e^{-\nu\tau|\xi'|^{\theta}}\widehat{\varphi}\|_{L_2(\mathbf{R}^{N-1})} \le C\tau^{-\frac{N-1}{\theta}\left(\frac{1}{p}-\frac{1}{2}\right)},$$

with a positive constant C independent of τ and φ .

Proof. (1) See e.g. [2, Lemma 3.1] and [13, Lemma 2.5].

(2) The desired estimate follows from (1) and Parseval's identity immediately, so that the detailed proof may be omitted. \Box

Let $L_p(\mathbf{R}^n, X)$ be the X-valued Lebesgue spaces on \mathbf{R}^n , $n \in \mathbf{N}$, for $1 \le p \le \infty$. The following lemma is proved in [9, Theorem 2.3].

Lemma 5.4. Let X be a Banach space and $\|\cdot\|_X$ its norm. Suppose that L and n be a non-negative integer and positive integer, respectively. Let $0 < \sigma \le 1$ and $s = L + \sigma - n$. Let $f(\xi)$ be a C^{∞} -function on $\mathbf{R}^n \setminus \{0\}$ with value X and satisfy the following two conditions:

- (1) $\partial_{\xi}^{\gamma} f \in L_1(\mathbf{R}^n, X)$ for any multi-index $\gamma \in \mathbf{N}_0^n$ with $|\gamma| \leq L$.
- (2) For any multi-index $\gamma \in \mathbf{N}_0^n$, there exists a positive constant M_{γ} such that

$$\|\partial_{\xi}^{\gamma} f(\xi)\|_{X} \le M_{\gamma} |\xi|^{s-|\delta|} \quad (\xi \in \mathbf{R}^{n} \setminus \{0\}).$$

Then there exists a positive constant $C_{n,s}$ such that

$$\|\mathcal{F}_{\xi}^{-1}[f](x)\|_{X} \le C_{n,s} \left(\max_{|\gamma| \le L+2} M_{\gamma} \right) |x|^{-(n+s)} \quad (x \in \mathbf{R}^{n} \setminus \{0\}).$$

5.1. Analysis for $\Gamma_{\rm Res}^{\pm}$. In this subsection, we prove

Theorem 5.5. Let $1 \le p < 2 \le q \le \infty$ and $\langle t \rangle = t + 1$. Then there exists a constant $A_0 \in (0, A_3)$ such that for any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$\begin{split} &\|\mathcal{H}_{A_0}^1(t;\Gamma_{\mathrm{Res}}^{\pm})d\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)} \|d\|_{L_p(\mathbf{R}^{N-1})}, \\ &\|\mathcal{H}_{A_0}^2(t;\Gamma_{\mathrm{Res}}^{\pm})\mathbf{f}\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{4}{5}\left(\frac{1}{p}-\frac{1}{2}\right)} \|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)}, \end{split}$$

$$\begin{split} &\|\mathcal{U}_{A_0}^1(t;\Gamma_{\mathrm{Res}}^{\pm})d\|_{L_q(\dot{\mathbf{R}}^N)} \leq C\langle t \rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{4}{5}\left(\frac{1}{2}-\frac{1}{q}\right)}\|d\|_{L_p(\mathbf{R}^{N-1})}, \\ &\|\mathcal{U}_{A_0}^2(t;\Gamma_{\mathrm{Res}}^{\pm})\mathbf{f}\|_{L_q(\dot{\mathbf{R}}^N)} \leq C\langle t \rangle^{-\frac{4N}{5}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)}, \end{split}$$

where C is a positive constant independent of t, d, and f.

Recalling $A = |\xi'|$ and λ_{\pm} given in Proposition 4.4, we define

$$\mathcal{B}_{\pm} = \sqrt{\frac{\rho_{\pm}}{\mu_{\pm}}\lambda_{\pm} + A^2}, \quad \mathcal{D}_{\pm} = \mu_{\pm}\mathcal{B}_{\pm} + \mu_{\mp}A, \quad \mathcal{E} = \mu_{+}\mathcal{B}_{+} + \mu_{-}\mathcal{B}_{-}.$$

Then we immediately obtain

Lemma 5.6. There exists a constant $A_4 \in (0, A_3)$ such that for any $A \in (0, A_4)$

$$C_1 A^{1/4} \le \Re \mathcal{B}_{\pm} \le |\mathcal{B}_{\pm}| \le C_2 A^{1/4}, \quad C_1 A^{3/4} \le |F(A, \lambda_{\pm})| \le C_2 A^{3/4},$$

with positive constants C_1 and C_2 independent of ξ' , and also

$$|\Phi_j^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda_{\pm})| \le CA^{2+(3/4)}, \quad |\Psi_j^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda_{\pm})| \le CA^{2+(3/4)},$$

 $|\mathcal{I}_{m\pm}(\xi',\lambda_{\pm})| \le CA^{1+(2/4)}, \quad |\mathcal{J}_m(\xi',\lambda_{\pm})| \le CA^{1+(2/4)},$

with a positive constant C independent of ξ' , where $\mathfrak{a}, \mathfrak{b} \in \{+, -\}$ and $j, m = 1, \ldots, N$.

Let $A \in (0, A_4)$. Then, by Lemma 5.6, we have the following estimates for the symbols of the representation formulas given in Subsection 3.3: for the height function,

$$\left| \frac{F(A, \lambda_{\pm})}{\mathcal{D}_{+} + \mathcal{D}_{-}} \right| \leq CA^{1/2}, \quad \left| \frac{\Phi_{j}^{\mathfrak{a}, \mathfrak{b}}(\xi', \lambda_{\pm})}{A(\mathcal{B}_{\mathfrak{b}} + A)(\mathcal{D}_{+} + \mathcal{D}_{-})} \right| \leq CA^{5/4},
\left| \frac{\Psi_{j}^{\mathfrak{a}, \mathfrak{b}}(\xi', \lambda_{\pm})}{A\mathcal{B}_{\mathfrak{b}}(\mathcal{B}_{\mathfrak{b}} + A)(\mathcal{D}_{+} + \mathcal{D}_{-})} \right| \leq CA;$$
(5.2)

for the fluid velocity,

$$\left| \frac{\mathcal{I}_{m\pm}(\xi',\lambda_{\pm})}{\mathcal{D}_{+} + \mathcal{D}_{-}} \right| \leq CA^{5/4}, \quad \left| \frac{\mathcal{J}_{m}(\xi',\lambda_{\pm})}{\mathcal{E}(\mathcal{D}_{+} + \mathcal{D}_{-})} \right| \leq CA,
\left| \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda_{\pm})\mathcal{I}_{m\pm}(\xi',\lambda_{\pm})}{A(\mathcal{B}_{\mathfrak{b}} + A)F(A,\lambda_{\pm})(\mathcal{D}_{+} + \mathcal{D}_{-})} \right| \leq CA^{2},
\left| \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda_{\pm})\mathcal{I}_{m\pm}(\xi',\lambda_{\pm})}{A\mathcal{B}_{\mathfrak{b}}(\mathcal{B}_{\mathfrak{b}} + A)F(A,\lambda_{\pm})(\mathcal{D}_{+} + \mathcal{D}_{-})} \right| \leq CA^{7/4},
\left| \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda_{\pm})\mathcal{J}_{m}(\xi',\lambda_{\pm})}{A(\mathcal{B}_{\mathfrak{b}} + A)F(A,\lambda_{\pm})\mathcal{E}(\mathcal{D}_{+} + \mathcal{D}_{-})} \right| \leq CA^{7/4},
\left| \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda_{\pm})\mathcal{J}_{m}(\xi',\lambda_{\pm})}{A\mathcal{B}_{\mathfrak{b}}(\mathcal{B}_{\mathfrak{b}} + A)F(A,\lambda_{\pm})\mathcal{E}(\mathcal{D}_{+} + \mathcal{D}_{-})} \right| \leq CA^{6/4}.$$
(5.3)

To prove Theorem 5.5, we introduce some technical lemma. Let us define the following operators:

$$[K_{A_0}(t;\Gamma)d](x) = \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_0}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} k(\xi',\lambda) \, d\lambda \, \widehat{d}(\xi') \right] (x'),$$

$$[K_{A_{0},\mathcal{M}}^{\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}](x)$$

$$= \int_{0}^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} k_{\mathcal{M}}(\xi',\lambda) \mathcal{M}_{\mathfrak{b}}(y_{N}) d\lambda \, \widehat{\mathbf{f}}(\xi',\mathfrak{a}y_{N}) \right] (x') \, dy_{N},$$

$$[K_{A_{0},B}^{\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}](x)$$

$$= \int_{0}^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} k_{B}(\xi',\lambda) e^{-B_{\mathfrak{b}}y_{N}} \, d\lambda \, \widehat{\mathbf{f}}(\xi',\mathfrak{a}y_{N}) \right] (x') \, dy_{N}, \qquad (5.4)$$

and also for $\pm x_N > 0$

$$[L_{A_{0},\mathcal{M}}^{\pm}(t;\Gamma)d](x) = \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} l_{\mathcal{M}}(\xi',\lambda) \mathcal{M}_{\pm}(\pm x_{N}) d\lambda \, \widehat{d}(\xi') \right] (x'),$$

$$[L_{A_{0},B}^{\pm}(t;\Gamma)d](x) = \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} l_{B}(\xi',\lambda) e^{\mp B_{\pm}y_{N}} \, d\lambda \, \widehat{d}(\xi') \right] (x'),$$

$$[L_{A_{0},\mathcal{M},\mathcal{M}}^{\pm}(t;\Gamma)\mathbf{f}](x)$$

$$= \int_{0}^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} l_{\mathcal{M},\mathcal{M}}(\xi',\lambda) \mathcal{M}_{\pm}(\pm x_{N}) \mathcal{M}_{\mathfrak{b}}(y_{N}) \, d\lambda \, \widehat{\mathbf{f}}(\xi',\mathfrak{a}y_{N}) \right] (x') \, dy_{N},$$

$$[L_{A_{0},\mathcal{M},B}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}](x)$$

$$= \int_{0}^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} l_{\mathcal{M},B}(\xi',\lambda) \mathcal{M}_{\pm}(\pm x_{N}) e^{-B_{\mathfrak{b}}y_{N}} \, d\lambda \, \widehat{\mathbf{f}}(\xi',\mathfrak{a}y_{N}) \right] (x') \, dy_{N},$$

$$[L_{A_{0},B,\mathcal{M}}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}](x)$$

$$= \int_{0}^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} l_{\mathcal{B},M}(\xi',\lambda) e^{\mp B_{\pm}x_{N}} \mathcal{M}_{\mathfrak{b}}(y_{N}) \, d\lambda \, \widehat{\mathbf{f}}(\xi',\mathfrak{a}y_{N}) \right] (x') \, dy_{N},$$

$$[L_{A_{0},B,B}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}](x)$$

$$= \int_{0}^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi_{A_{0}}(\xi')}{2\pi i} \int_{\widehat{\Gamma}} e^{\lambda t} l_{\mathcal{B},B}(\xi',\lambda) e^{\mp B_{\pm}x_{N}} e^{-B_{\mathfrak{b}}y_{N}} \, d\lambda \, \widehat{\mathbf{f}}(\xi',\mathfrak{a}y_{N}) \right] (x') \, dy_{N}.$$

$$(5.5)$$

Here it is assumed that the symbols

$$k(\xi',\lambda), k_{\mathcal{M}}(\xi',\lambda), k_{B}(\xi',\lambda), l_{\mathcal{M}}(\xi',\lambda), l_{B}(\xi',\lambda), l_{\mathcal{M}\mathcal{M}}(\xi',\lambda), l_{\mathcal{M}\mathcal{M}}(\xi',\lambda), l_{\mathcal{M}\mathcal{M}}(\xi',\lambda)$$

are infinitely many times differentiable with respect to $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and holomorphic with respect to $\lambda \in \mathbf{C} \setminus (-\infty, -z_0|\xi'|^2]$. Then we have

Lemma 5.7. Let $1 \le p < 2 \le q \le \infty$, $\langle t \rangle = t+1$, and $\mathfrak{a}, \mathfrak{b} \in \{+, -\}$. Suppose that

$$Z(\xi',\lambda) = \frac{\widetilde{Z}(\xi',\lambda)}{\mathcal{L}_A(\lambda)} \quad \text{for } Z \in \{k, k_{\mathcal{M}}, k_B, l_{\mathcal{M}}, l_B, l_{\mathcal{MM}}, l_{\mathcal{MB}}, l_{\mathcal{BM}}, l_{\mathcal{BB}}\}$$

and that there exists a constant $A_5 \in (0, A_3)$ such that for any $A \in (0, A_5)$

$$\begin{aligned} |\widetilde{k}(\xi',\lambda_{\pm})| &\leq CA^{1/2}, \quad |\widetilde{k}_{\mathcal{M}}(\xi',\lambda_{\pm})| \leq CA^{5/4}, \quad |\widetilde{k}_{B}(\xi',\lambda_{\pm})| \leq CA, \\ |l_{\mathcal{M}}(\xi',\lambda_{\pm})| &\leq CA^{5/4}, \quad |\widetilde{l}_{B}(\xi',\lambda_{\pm})| \leq CA, \quad |\widetilde{l}_{\mathcal{M}\mathcal{M}}(\xi',\lambda_{\pm})| \leq CA^{2}, \\ |l_{\mathcal{M}B}(\xi',\lambda_{\pm})| &\leq CA^{7/4}, \quad |\widetilde{l}_{B\mathcal{M}}(\xi',\lambda_{\pm})| \leq CA^{7/4}, \quad |\widetilde{l}_{BB}(\xi',\lambda_{\pm})| \leq CA^{6/4}, \end{aligned}$$

with some positive constant C independent of ξ' . Then there exists a constant $A_0 \in (0, A_5)$ such that the following assertions hold.

(1) For any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$\begin{aligned} & \|K_{A_0}(t;\Gamma_{\mathrm{Res}}^{\pm})d\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)} \|d\|_{L_p(\mathbf{R}^{N-1})}, \\ & \|K_{A_0,\mathcal{M}}^{\mathfrak{a},\mathfrak{b}}(t;\Gamma_{\mathrm{Res}}^{\pm})\mathbf{f}\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{4}{5}\left(\frac{1}{p}-\frac{1}{2}\right)} \|\mathbf{f}\|_{L_p(\mathbf{R}_{\mathfrak{a}}^{N})}, \\ & \|K_{A_0,B}^{\mathfrak{a},\mathfrak{b}}(t;\Gamma_{\mathrm{Res}}^{\pm})\mathbf{f}\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{4}{5}\left(\frac{1}{p}-\frac{1}{2}\right)} \|\mathbf{f}\|_{L_p(\mathbf{R}_{\mathfrak{a}}^{N})}, \end{aligned}$$

with some positive constant C independent of t, d, and f.

(2) Let $\Gamma = \Gamma_{\text{Res}}^+$ or $\Gamma = \Gamma_{\text{Res}}^-$. Then for any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$\begin{split} \|L_{A_{0},\mathcal{M}}^{\pm}(t;\Gamma)d\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{4}{5}\left(\frac{1}{2}-\frac{1}{q}\right)}\|d\|_{L_{p}(\mathbf{R}^{N-1})},\\ \|L_{A_{0},B}^{\pm}(t;\Gamma)d\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{4}{5}\left(\frac{1}{2}-\frac{1}{q}\right)}\|d\|_{L_{p}(\mathbf{R}^{N-1})},\\ \|L_{A_{0},\mathcal{M}\mathcal{M}}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{4N}{5}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})},\\ \|L_{A_{0},\mathcal{M}B}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{4N}{5}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})},\\ \|L_{A_{0},B\mathcal{M}}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{4N}{5}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})},\\ \|L_{A_{0},BB}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{4N}{5}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})}, \end{split}$$

with some positive constant C independent of t, d, and \mathbf{f} .

Proof. We here consider

$$K_{A_0}(t;\Gamma_{\mathrm{Res}}^+), \quad K_{A_0,\mathcal{M}}^{+,+}(t;\Gamma_{\mathrm{Res}}^+), \quad L_{A_0,\mathcal{M}}^+(t;\Gamma_{\mathrm{Res}}^+), \quad L_{A_0,\mathcal{M}\mathcal{M}}^{+,+,+}(t;\Gamma_{\mathrm{Res}}^+),$$

only. The other cases can be proved analogously (cf. also [7, Subsection 4.1]).

Case 1: $K_{A_0}(t; \Gamma_{\text{Res}}^+)$. By the residue theorem, we have

$$[K_{A_0}(t; \Gamma_{\mathrm{Res}}^+) d](x') = \mathcal{F}_{\xi'}^{-1} \Big[\varphi_{A_0}(\xi') e^{\lambda_+ t} \frac{\widetilde{k}(\xi', \lambda_+)}{L'_A(\lambda_+)} d\lambda \, \widehat{d}(\xi') \Big](x').$$

Recalling $\Re \zeta_{\pm} = -\sqrt{2}\alpha^{1/4}\beta A^{5/4}$, we write this formula as

$$[K_{A_0}(t;\Gamma_{\mathrm{Res}}^+)d](x') = \mathcal{F}_{\xi'}^{-1} \Big[\varphi_{A_0}(\xi') e^{\frac{\Re \zeta_+}{3} \langle t \rangle} e^{-\frac{\Re \zeta_+}{3} \langle t \rangle} e^{\lambda_+ t} \frac{\widetilde{k}(\xi',\lambda_+)}{L_A'(\lambda_+)} \, \widehat{d}(\xi') \Big](x').$$

Combining this formula with Lemma 5.3 yields

$$||K_{A_0}(t; \Gamma_{\text{Res}}^+) d||_{L_a(\mathbf{R}^{N-1})}$$

$$\leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{2}-\frac{1}{q}\right)} \left\|\mathcal{F}_{\xi'}^{-1}\left[\varphi_{A_0}(\xi')e^{-\frac{\Re\zeta_+}{3}\langle t\rangle}e^{\lambda_+t}\frac{\widetilde{k}(\xi',\lambda_+)}{L_A'(\lambda_+)}\widehat{d}\right]\right\|_{L_2(\mathbf{R}^{N-1})} =: I_1(t).$$

We choose a sufficiently small A_0 so that

$$|e^{-\frac{\Re\zeta_{+}}{3}\langle t\rangle}e^{\lambda_{+}t}| \leq Ce^{\frac{\Re\zeta_{+}}{3}\langle t\rangle}$$
 on $\operatorname{supp}\varphi_{A_{0}}$,

and thus we have by Parseval's identity, Proposition 4.4, and the assumption for $\widetilde{k}(\xi',\lambda_+)$

$$I_1(t) \le C\langle t \rangle^{-\frac{4(N-1)}{5} \left(\frac{1}{2} - \frac{1}{q}\right)} \|\varphi_{A_0}(\xi') e^{\frac{\Re \zeta_+}{3} \langle t \rangle} \widehat{d} \|_{L_2(\mathbf{R}^{N-1})}.$$
 (5.6)

Since $0 \le \varphi_{A_0} \le 1$, this implies

$$I_1(t) \le C\langle t \rangle^{-\frac{4(N-1)}{5} \left(\frac{1}{2} - \frac{1}{q}\right)} \|e^{(\Re \zeta_+/3)\langle t \rangle} \widehat{d}\|_{L_2(\mathbf{R}^{N-1})}.$$

Applying Lemma 5.3 to the right-hand side of the last inequality furnishes the desired estimate for $K_{A_0}(t; \Gamma_{\text{Res}}^+)$. This completes the proof of Case 1.

Case 2: $K_{A_0,\mathcal{M}}^{+,+}(t;\Gamma_{\text{Res}}^+)$. In the same way as we have obtained (5.6), we obtain

$$\begin{aligned} & \|K_{A_0,\mathcal{M}}^{+,+}(t;\Gamma_{\mathrm{Res}}^+)\mathbf{f}\|_{L_q(\mathbf{R}^{N-1})} \\ & \leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{2}-\frac{1}{q}\right)} \int_0^\infty \|\varphi_{A_0}(\xi')e^{\frac{\Re\zeta_+}{3}\langle t\rangle}A^{3/4}\mathcal{M}_+(y_N)\widehat{\mathbf{f}}(\xi',y_N)\|_{L_2(\mathbf{R}^{N-1})}dy_N \\ & =: I_2(t). \end{aligned}$$

We choose a sufficiently small A_0 so that

$$|\mathcal{M}_{+}(y_N)| \le CA^{-1/4}e^{-cAy_N} \quad \text{on supp } \varphi_{A_0} \tag{5.7}$$

for positive constant C and c. Then

$$I_2(t) \le C\langle t \rangle^{-\frac{4(N-1)}{5} \left(\frac{1}{2} - \frac{1}{q}\right)} \int_0^\infty \|e^{\frac{\Re \zeta_+}{3} \langle t \rangle} A^{1/2} e^{-cAy_N} \widehat{\mathbf{f}}(\xi', y_N)\|_{L_2(\mathbf{R}^{N-1})} dy_N,$$

which, combined with Lemmas 5.1 and 5.3, implies

$$I_2(t) \le C\langle t \rangle^{-\frac{4(N-1)}{5} \left(\frac{1}{p} - \frac{1}{q}\right)} \int_0^\infty \frac{\|\mathbf{f}(\xi', y_N)\|_{L_p(\mathbf{R}^{N-1})}}{\langle t \rangle^{2/5} + y_N^{1/2}} \, dy_N.$$

Since $1 \leq p < 2$, applying Lemma 5.2 to the right-hand side of the last inequality shows that the desired estimate for $K_{A_0,\mathcal{M}}^{+,+}(t;\Gamma_{\mathrm{Res}}^+)$ holds. This completes the proof of Case 2.

Case 3: $L_{A_0,\mathcal{M}}^+(t;\Gamma_{\mathrm{Res}}^+)$. In the same way as we have obtained (5.6), we obtain

$$||L_{A_0,\mathcal{M}}^+(t;\Gamma_{\mathrm{Res}}^+)d||_{L_q(\mathbf{R}^{N-1})}$$

$$\leq C\langle t \rangle^{-\frac{4(N-1)}{5} \left(\frac{1}{2} - \frac{1}{q}\right)} \|\varphi_{A_0}(\xi')e^{\frac{\Re \zeta_+}{3}\langle t \rangle} A^{3/4} \mathcal{M}_+(x_N) \widehat{d}(\xi') \|_{L_2(\mathbf{R}^{N-1})} =: I_3(x_N, t).$$

By (5.7), we see that

$$I_3(x_N, t) \le C\langle t \rangle^{-\frac{4(N-1)}{5} \left(\frac{1}{2} - \frac{1}{q}\right)} \|e^{\frac{\Re \zeta_+}{3} \langle t \rangle} A^{1/2} e^{-cAx_N} \widehat{d}(\xi')\|_{L_2(\mathbf{R}^{N-1})}.$$
 (5.8)

When q = 2, it follows from (5.8) that

$$\int_{0}^{\infty} I_{3}(x_{N}, t)^{2} dx_{N} \leq C \int_{0}^{\infty} \|e^{\frac{\Re \zeta_{+}}{3} \langle t \rangle} A^{1/2} e^{-cAx_{N}} \widehat{d}(\xi')\|_{L_{2}(\mathbf{R}^{N-1})}^{2} dx_{N} dx$$

Combining this with Lemma 5.3 yields the desired estimate of $L_{A_0,\mathcal{M}}^+(t;\Gamma_{\mathrm{Res}}^+)$ for q=2. When q>2, it follows from (5.8) and Lemmas 5.1 and 5.3 that

$$I(x_N, t) \le C\langle t \rangle^{-\frac{4(N-1)}{5} (\frac{1}{p} - \frac{1}{q})} \frac{\|d\|_{L_p(\mathbf{R}^{N-1})}}{\langle t \rangle^{2/5} + x_N^{1/2}}.$$

In this inequality, taking L_q norm of both sides with respect to $x_N \in (0, \infty)$ furnishes the desired estimate of $L_{A_0,\mathcal{M}}^+(t;\Gamma_{\mathrm{Res}}^+)$ for q>2. This completes the proof of Case 3.

Case 4: $L_{A_0,\mathcal{M}}^{+,+,+}(t;\Gamma_{\text{Res}}^+)$. In the same way as we have obtained (5.6), we obtain

$$\|L_{A_0,\mathcal{MM}}^{+,+,+}(t;\Gamma_{\mathrm{Res}}^+)d\|_{L_q(\mathbf{R}^{N-1})}$$

$$\leq C\langle t\rangle^{-\frac{4}{5}\left(\frac{1}{2}-\frac{1}{q}\right)} \int_0^\infty \|\varphi_{A_0}(\xi')e^{\frac{\Re\zeta_+}{3}\langle t\rangle}A^{\frac{6}{4}}\mathcal{M}_+(x_N)\mathcal{M}_+(y_N)\widehat{\mathbf{f}}(\xi',y_N)\|_{L_2(\mathbf{R}^{N-1})}dy_N$$

$$=: I_4(x_N, t).$$

Combining this with (5.7) and Lemmas 5.1 and 5.3 yields

$$I_{4}(x_{N},t)$$

$$\leq C\langle t\rangle^{-\frac{4}{5}\left(\frac{1}{2}-\frac{1}{q}\right)} \int_{0}^{\infty} \|e^{\frac{\Re\zeta_{+}}{3}\langle t\rangle} A e^{-cA(x_{N}+y_{N})} \widehat{\mathbf{f}}(\xi',y_{N})\|_{L_{2}(\mathbf{R}^{N-1})} dy_{N}$$

$$\leq C\langle t\rangle^{-\frac{4(N-1)}{5}\left(\frac{1}{p}-\frac{1}{q}\right)} \int_{0}^{\infty} \frac{\|\mathbf{f}(\xi',y_{N})\|_{L_{p}(\mathbf{R}^{N-1})}}{\langle t\rangle^{4/5} + x_{N} + y_{N}} dy_{N}.$$

Lemma 5.2 thus yields the desired estimate for $L_{A_0,\mathcal{MM}}^{+,+,+}(t;\Gamma_{\mathrm{Res}}^+)$. This completes the proof of Lemma 5.7.

Combining Lemma 5.7 with (5.2) and (5.3) yields Theorem 5.5 immediately. This completes the proof of Theorem 5.5.

5.2. Analysis for Γ_1^{\pm} . In this subsection, we prove

Theorem 5.8. Let $1 \le p \le 2 \le q \le \infty$ and $\langle t \rangle = t+1$. Then there exists a constant $A_0 \in (0, A_3)$ such that for any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$\begin{split} &\|\mathcal{H}_{A_0}^1(t;\Gamma_1^{\pm})d\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{3}{2}}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ &\|\mathcal{H}_{A_0}^2(t;\Gamma_1^{\pm})\mathbf{f}\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{p}-\frac{1}{2}\right)-\frac{3}{4}}\|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)},\\ &\|\mathcal{U}_{A_0}^1(t;\Gamma_1^{\pm})d\|_{L_q(\dot{\mathbf{R}}^N)} \leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{3}{4}}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ &\|\mathcal{U}_{A_0}^2(t;\Gamma_1^{\pm})\mathbf{f}\|_{L_q(\dot{\mathbf{R}}^N)} \leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)}, \end{split}$$

where C is a positive constant independent of t, d, and f.

To prove Theorem 5.8, we start with the following lemma.

Lemma 5.9. There exists a constant $A_6 \in (0, A_3)$ such that for any $A \in (0, A_6)$ and $\lambda \in \widehat{\Gamma}_1^+ \cup \widehat{\Gamma}_1^-$

$$C_1 A \le \Re B_{\pm} \le |B_{\pm}| \le C_2 A, \quad C_1 A^3 \le |F(A, \lambda)| \le C_2 A^3,$$

with positive constants C_1 and C_2 independent of ξ' and λ , and also

$$\begin{split} |\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)| &\leq CA^{5}, \quad |\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)| \leq CA^{5}, \\ |\mathcal{I}_{m\pm}(\xi',\lambda)| &\leq CA^{3}, \quad |\mathcal{J}_{m}(\xi',\lambda)| \leq CA^{3}, \end{split}$$

with a positive constant C independent of ξ' and λ , where $\mathfrak{a}, \mathfrak{b} \in \{+, -\}$ and $j, m = 1, \ldots, N$.

Proof. See [7, Lemma 4.9] for B_{\pm} . Then the desired estimates for $\Phi_j^{\mathfrak{a},\mathfrak{b}}$, $\Psi_j^{\mathfrak{a},\mathfrak{b}}$, $\mathcal{I}_{m\pm}$, and \mathcal{J}_m follow from the estimates of B_{\pm} immediately.

We now estimate $|F(A,\lambda)|$. Let $\lambda \in \widehat{\Gamma}_1^+ \cup \widehat{\Gamma}_1^-$. It is clear that $|F(A,\lambda)| \leq CA^3$ by $|B_{\pm}| \leq CA$. In what follows, we prove $|F(A,\lambda)| \geq CA^3$. Since $\lambda = -(z_0/2)A^2 + (z_0/4)A^2e^{is}$ for $s \in [-\pi/2, \pi/2]$, there holds

$$F(A,\lambda) = A^3 F(1,\zeta), \quad \zeta = -\frac{z_0}{2} + \frac{z_0}{4} e^{is}.$$
 (5.9)

It suffices to show that

$$F(1,\zeta) \neq 0 \quad \text{for } s \in [-\pi/2, \pi/2].$$
 (5.10)

Let $s \neq 0$. Then $\zeta \in \Sigma_{\varepsilon}$ for some $\varepsilon = \varepsilon(s)$. Therefore $F(1,\zeta) \neq 0$, since $|F(1,\zeta)| \geq C_{\varepsilon}$ for some positive constant C_{ε} by Lemma 3.1.

Next, we consider s=0. In this case, $\zeta=-z_0/4$ and set $b_{\pm}=\sqrt{(\rho_{\pm}/\mu_{\pm})\zeta+1}$. Then

$$\frac{\sqrt{3}}{2} \le b_{\pm} < 1,\tag{5.11}$$

and $F(1,\zeta)$ can be written as

$$F(1,\zeta) = -(\mu_{+} - \mu_{-})^{2} + (3\mu_{+} - \mu_{-})\mu_{+}b_{+} + (3\mu_{-} - \mu_{+})\mu_{-}b_{-}$$
$$+ (\mu_{+}b_{+} + \mu_{-}b_{-})^{2} + \mu_{+}\mu_{-}(b_{+} + b_{-})^{2} + (\mu_{+}b_{+} + \mu_{-}b_{-})(\mu_{+}b_{+}^{2} + \mu_{-}b_{-}^{2}).$$

Since it follows from (5.11) that

$$-(\mu_{+} - \mu_{-})^{2} + (3\mu_{+} - \mu_{-})\mu_{+}b_{+} + (3\mu_{-} - \mu_{+})\mu_{-}b_{-}$$

$$= -(\mu_{+}^{2} + \mu_{-}^{2}) + 2\mu_{+}\mu_{-} + 3\mu_{+}^{2}b_{+} + 3\mu_{-}^{2}b_{-} - \mu_{+}\mu_{-}(b_{+} + b_{-})$$

$$\geq -(\mu_{+}^{2} + \mu_{-}^{2}) + 2\mu_{+}\mu_{-} + \frac{3\sqrt{3}}{2}(\mu_{+}^{2} + \mu_{-}^{2}) - 2\mu_{+}\mu_{-} > 0,$$

we have $F(1,\zeta) > 0$ for s = 0. Thus (5.10) holds, which implies $|F(1,\zeta)| \geq C$ for any $s \in [-\pi/2, \pi/2]$ and a positive constant C independent of s. Therefore $|F(A,\lambda)| \geq CA^3$ by (5.9), which completes the proof of Lemma 5.9.

Note that $|\mathcal{L}_A(\lambda)| \geq CA$ for $\lambda \in \widehat{\Gamma}_1^+ \cup \widehat{\Gamma}_1^-$ when A is small enough as seen in Case 1 of the proof of Lemma 4.1, and thus it follows from Lemma 5.9 that $|L(A,\lambda)| \geq CA^2$. By this inequality and Lemma 5.9, we have the following estimates for the symbols of the representation formulas given in Subsection 3.3: for the height function,

$$\left| \frac{F(A,\lambda)}{L(A,\lambda)} \right| \le CA, \quad \left| \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)}{A(B_{\mathfrak{b}}+A)L(A,\lambda)} \right| \le CA, \quad \left| \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)}{AB_{\mathfrak{b}}(B_{\mathfrak{b}}+A)L(A,\lambda)} \right| \le C; \tag{5.12}$$

for the velocity

$$\left| \frac{\mathcal{I}_{m\pm}(\xi',\lambda)}{L(A,\lambda)} \right| \leq CA, \qquad \left| \frac{\mathcal{J}_{m}(\xi',\lambda)}{EL(A,\lambda)} \right| \leq C,$$

$$\left| \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{I}_{m\pm}(\xi',\lambda)}{A(B_{\mathfrak{b}}+A)F(A,\lambda)L(A,\lambda)} \right| \leq CA, \qquad \left| \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{I}_{m\pm}(\xi',\lambda)}{AB_{\mathfrak{b}}(B_{\mathfrak{b}}+A)F(A,\lambda)L(A,\lambda)} \right| \leq C,$$

$$\left| \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{J}_{m}(\xi',\lambda)}{A(B_{\mathfrak{b}}+A)F(A,\lambda)EL(A,\lambda)} \right| \leq C, \qquad \left| \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{J}_{m}(\xi',\lambda)}{AB_{\mathfrak{b}}(B_{\mathfrak{b}}+A)F(A,\lambda)EL(A,\lambda)} \right| \leq \frac{C}{A}.$$
(5.13)

Now, recalling the operators difined in (5.4) and (5.5), we introduce the following lemma (cf. [7, Lemma 4.10] for details).

Lemma 5.10. Let $1 \le p \le 2 \le q \le \infty$, $\langle t \rangle = t+1$, and $\mathfrak{a}, \mathfrak{b} \in \{+, -\}$. Suppose that there exists a constant $A_7 \in (0, A_3)$ such that for any $A \in (0, A_7)$ and $\lambda \in \widehat{\Gamma}_1^+ \cup \widehat{\Gamma}_1^-$

$$|k(\xi',\lambda)| \le CA, \quad |k_{\mathcal{M}}(\xi',\lambda)| \le CA, \quad |k_{B}(\xi',\lambda)| \le C,$$

$$|l_{\mathcal{M}}(\xi',\lambda)| \le CA, \quad |l_{B}(\xi',\lambda)| \le C, \quad |l_{\mathcal{M}\mathcal{M}}(\xi',\lambda)| \le CA,$$

$$|l_{\mathcal{M}B}(\xi',\lambda)| \le C, \quad |l_{B\mathcal{M}}(\xi',\lambda)| \le C, \quad |l_{BB}(\xi',\lambda)| \le CA^{-1},$$

with some positive constant C independent of ξ' and λ . Then there exists a constant $A_0 \in (0, A_7)$ such that the following assertions hold.

(1) For any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$||K_{A_0}(t;\Gamma_1^{\pm})d||_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{N-1}{2}(\frac{1}{p}-\frac{1}{q})-\frac{3}{2}}||d||_{L_p(\mathbf{R}^{N-1})},$$

$$||K_{A_0,\mathcal{M}}^{\mathfrak{a},\mathfrak{b}}(t;\Gamma_1^{\pm})\mathbf{f}||_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{N-1}{2}(\frac{1}{p}-\frac{1}{q})-\frac{1}{2}(\frac{1}{p}-\frac{1}{2})-\frac{3}{4}}||\mathbf{f}||_{L_p(\mathbf{R}_a^N)},$$

$$||K_{A_0,B}^{\mathfrak{a},\mathfrak{b}}(t;\Gamma_1^{\pm})\mathbf{f}||_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{N-1}{2}(\frac{1}{p}-\frac{1}{q})-\frac{1}{2}(\frac{1}{p}-\frac{1}{2})-\frac{3}{4}}||\mathbf{f}||_{L_p(\mathbf{R}_a^N)},$$

with some positive constant C independent of t, d, and \mathbf{f} .

(2) Let $\Gamma = \Gamma_1^+$ or $\Gamma = \Gamma_1^-$. Then for any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$\begin{split} \|L_{A_0,\mathcal{M}}^{\pm}(t;\Gamma)d\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{3}{4}}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ \|L_{A_0,B}^{\pm}(t;\Gamma)d\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{3}{4}}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ \|L_{A_0,\mathcal{M}}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_p(\mathbf{R}_{\mathfrak{a}}^N)},\\ \|L_{A_0,\mathcal{M}B}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_p(\mathbf{R}_{\mathfrak{a}}^N)},\\ \|L_{A_0,B}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_p(\mathbf{R}_{\mathfrak{a}}^N)},\\ \|L_{A_0,BB}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)\mathbf{f}\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_p(\mathbf{R}_{\mathfrak{a}}^N)}, \end{split}$$

with some positive constant C independent of t, d, and f.

Combining Lemma 5.10 with (5.12) and (5.13) proves Theorem 5.8 immediately. This completes the proof of Theorem 5.8.

5.3. Analysis for Γ_4^{\pm} . In this subsection, we prove

Theorem 5.11. Let $1 \le p < 2 \le q \le \infty$ and $\langle t \rangle = t+1$ Then there exists a constant $A_0 \in (0, A_3)$ such that for any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$\begin{split} &\|\mathcal{H}_{A_0}^1(t;\Gamma_4^\pm)d\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{3}{4}\gamma_1}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ &\|\mathcal{H}_{A_0}^2(t;\Gamma_4^\pm)\mathbf{f}\|_{L_q(\mathbf{R}^{N-1})} \leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{p}-\frac{1}{2}\right)-\frac{3}{4}\gamma_2}\|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)},\\ &\|\mathcal{U}_{A_0}^1(t;\Gamma_4^\pm)d\|_{L_q(\dot{\mathbf{R}}^N)} \leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{3}{4}\gamma_3}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ &\|\mathcal{U}_{A_0}^2(t;\Gamma_4^\pm)\mathbf{f}\|_{L_q(\dot{\mathbf{R}}^N)} \leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)}, \end{split}$$

where C is a positive constant independent of t, d, and f. Here

$$0 < \gamma_1 < \min\left\{1, 2(N-1)\left(\frac{1}{p} - \frac{1}{2}\right)\right\}, \quad 0 < \gamma_2 < \min\left\{1, 2N\left(\frac{1}{p} - \frac{1}{2}\right)\right\},$$

$$0 < \gamma_3 < \min\left\{1, 2\left((N-1)\left(\frac{1}{p} - \frac{1}{2}\right) + \frac{1}{2} - \frac{1}{q}\right)\right\}.$$

Note that $\widehat{\Gamma}_4^+ \cup \widehat{\Gamma}_4^- \subset \Sigma_{\theta_2}$ for θ_2 given in (2.1). By Lemma 3.1, we have for any $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and $\lambda \in \widehat{\Gamma}_+^4 \cup \widehat{\Gamma}_-^4$

$$|\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)| \le CA^{2}(|\lambda|^{1/2} + A)^{3}, \quad |\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)| \le CA^{2}(|\lambda|^{1/2} + A)^{3},$$

$$|\mathcal{I}_{m\pm}(\xi',\lambda)| \le CA(|\lambda|^{1/2} + A)^{2}, \quad |\mathcal{J}_{m}(\xi',\lambda)| \le CA(|\lambda|^{1/2} + A)^{2}, \quad (5.14)$$

where $\mathfrak{a},\mathfrak{b} \in \{+,-\}$ and $j,m=1,\ldots,N.$ In addition, similarly to (4.6), there holds

$$|\mathcal{L}_A(\lambda)| \ge C(|\lambda| + A^{1/2})^2$$

for a sufficiently small A and $\lambda \in \widehat{\Gamma}_+^4 \cup \widehat{\Gamma}_-^4$, which, combined with $|\lambda| + A^{1/2} \ge (1/2)(|\lambda|^{1/2} + A^{1/4})^2$ and Lemma 3.2, furnishes

$$|L(A,\lambda)| \ge C(|\lambda|^{1/2} + A)(|\lambda|^{1/2} + A^{1/4})^4.$$

By this inequality together with (5.14) and Lemma 3.1, we have the following estimates for the symbols of the representation formulas given in Subsection 3.3: for the height function

$$\left| \frac{F(A,\lambda)}{L(A,\lambda)} \right| \le \frac{C}{(|\lambda|^{1/2} + A^{1/4})^2}, \quad \left| \frac{\Phi_j^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)}{A(B_{\mathfrak{b}} + A)L(A,\lambda)} \right| \le \frac{CA(|\lambda|^{1/2} + A)}{(|\lambda|^{1/2} + A^{1/4})^4}, \\
\left| \frac{\Phi_j^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)}{AB_{\mathfrak{b}}(B_{\mathfrak{b}} + A)L(A,\lambda)} \right| \le \frac{CA}{(|\lambda|^{1/2} + A^{1/4})^4};$$
(5.15)

for the velocity

$$\left| \frac{\mathcal{I}_{m\pm}(\xi',\lambda)}{L(A,\lambda)} \right| \le \frac{CA(|\lambda|^{1/2} + A)}{(|\lambda|^{1/2} + A^{1/4})^4}, \quad \left| \frac{\mathcal{J}_m(\xi',\lambda)}{EL(A,\lambda)} \right| \le \frac{CA}{(|\lambda|^{1/2} + A^{1/4})^4}, \tag{5.16}$$

and also

$$\left| \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{I}_{m\pm}(\xi',\lambda)}{A(B_{\mathfrak{b}}+A)F(A,\lambda)L(A,\lambda)} \right| \leq CA,
\left| \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{I}_{m\pm}(\xi',\lambda)}{AB_{\mathfrak{b}}(B_{\mathfrak{b}}+A)F(A,\lambda)L(A,\lambda)} \right| \leq \frac{CA}{|\lambda|^{1/2}+A},
\left| \frac{\Phi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{J}_{m}(\xi',\lambda)}{A(B_{\mathfrak{b}}+A)F(A,\lambda)EL(A,\lambda)} \right| \leq \frac{CA}{|\lambda|^{1/2}+A},
\left| \frac{\Psi_{j}^{\mathfrak{a},\mathfrak{b}}(\xi',\lambda)\mathcal{J}_{m}(\xi',\lambda)}{AB_{\mathfrak{b}}(B_{\mathfrak{b}}+A)F(A,\lambda)EL(A,\lambda)} \right| \leq \frac{CA}{(|\lambda|^{1/2}+A)^{2}}.$$
(5.17)

We now prove

Lemma 5.12. Let $1 \le p < 2 \le q \le \infty$, $\langle t \rangle = t+1$, and $\mathfrak{a}, \mathfrak{b} \in \{+, -\}$. Suppose that there exists a constant $A_8 \in (0, A_3)$ such that for any $A \in (0, A_8)$ and $\lambda \in \widehat{\Gamma}_4^+ \cup \widehat{\Gamma}_4^-$

$$|k(\xi',\lambda)| \leq \frac{C}{(|\lambda|^{1/2} + A^{1/4})^2}, \quad |k_{\mathcal{M}}(\xi',\lambda)| \leq \frac{CA(|\lambda|^{1/2} + A)}{(|\lambda|^{1/2} + A^{1/4})^4},$$

$$|k_{B}(\xi',\lambda)| \leq \frac{CA}{(|\lambda|^{1/2} + A^{1/4})^4}, \quad |l_{\mathcal{M}}(\xi',\lambda)| \leq \frac{CA(|\lambda|^{1/2} + A)}{(|\lambda|^{1/2} + A^{1/4})^4},$$

$$|l_{B}(\xi',\lambda)| \leq \frac{CA}{(|\lambda|^{1/2} + A^{1/4})^4}, \quad |l_{\mathcal{M}\mathcal{M}}(\xi',\lambda)| \leq CA,$$

$$|l_{\mathcal{M}B}(\xi',\lambda)| \leq \frac{CA}{|\lambda|^{1/2} + A}, \quad |l_{\mathcal{B}\mathcal{M}}(\xi',\lambda)| \leq \frac{CA}{|\lambda|^{1/2} + A},$$

$$|l_{BB}(\xi',\lambda)| \leq \frac{CA}{(|\lambda|^{1/2} + A)^2},$$

with some positive constant of ξ' and λ . Then there exists a constant $A_0 \in (0, A_8)$ such that the following assertions hold.

(1) For any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$||K_{A_0}(t; \Gamma_4^{\pm})d||_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{N-1}{2} \left(\frac{1}{p} - \frac{1}{q}\right) - \frac{3}{4}\gamma_1} ||d||_{L_p(\mathbf{R}^{N-1})},$$

$$||K_{A_0, \mathcal{M}}^{\mathfrak{a}, \mathfrak{b}}(t; \Gamma_4^{\pm})\mathbf{f}||_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{N-1}{2} \left(\frac{1}{p} - \frac{1}{q}\right) - \frac{1}{2} \left(\frac{1}{p} - \frac{1}{2}\right) - \frac{3}{4}\gamma_2} ||\mathbf{f}||_{L_p(\mathbf{R}_a^N)},$$

$$||K_{A_0, B}^{\mathfrak{a}, \mathfrak{b}}(t; \Gamma_4^{\pm})\mathbf{f}||_{L_q(\mathbf{R}^{N-1})} \leq C\langle t \rangle^{-\frac{N-1}{2} \left(\frac{1}{p} - \frac{1}{q}\right) - \frac{1}{2} \left(\frac{1}{p} - \frac{1}{2}\right) - \frac{3}{4}} ||\mathbf{f}||_{L_p(\mathbf{R}_a^N)},$$

with some positive constant C independent of t, d, and \mathbf{f} .

(2) Let $\Gamma = \Gamma_4^+$ or $\Gamma = \Gamma_4^-$. Then for any t > 0 and $(d, \mathbf{f}) \in Y_p$

$$\begin{split} \|L_{A_{0},\mathcal{M}}^{\pm}(t;\Gamma)d\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{3}{4}\gamma_{3}}\|d\|_{L_{p}(\mathbf{R}^{N-1})},\\ \|L_{A_{0},B}^{\pm}(t;\Gamma)d\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{3}{4}}\|d\|_{L_{p}(\mathbf{R}^{N-1})},\\ \|L_{A_{0},\mathcal{M}\mathcal{M}}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})},\\ \|L_{A_{0},\mathcal{M}B}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})},\\ \|L_{A_{0},B\mathcal{M}}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})},\\ \|L_{A_{0},BB}^{\pm}(t;\Gamma)\mathbf{f}\|_{L_{q}(\mathbf{R}_{\pm}^{N})} &\leq C\langle t\rangle^{-\frac{N}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|\mathbf{f}\|_{L_{p}(\mathbf{R}_{a}^{N})}, \end{split}$$

with some positive constant C independent of t, d, and f.

Proof. We here consider $K_{A_0}(t; \Gamma_4^+)$, $K_{A_0,\mathcal{M}}^{+,+}(t; \Gamma_4^+)$, and $L_{A_0,\mathcal{M}}^+(t; \Gamma_4^+)$ only. The desired estimates for

$$L_{A_0,B}^{\pm}(t;\Gamma), \quad L_{A_0,\mathcal{M}\mathcal{M}}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma), \quad L_{A_0,\mathcal{M}B}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma), \quad L_{A_0,B\mathcal{M}}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma), \quad L_{A_0,B\mathcal{B}}^{\pm,\mathfrak{a},\mathfrak{b}}(t;\Gamma)$$

are proved in [7, Lemma 4.13], and $K_{A_0,B}^{\mathfrak{a},\mathfrak{b}}(t;\Gamma_4^{\pm})$ can be proved similarly to the case of $L_{A_0,B}^{\pm}(t;\Gamma)$.

Case 1: $K_{A_0}(t; \Gamma_4^+)$. Since $\lambda = z_1^+(1-s) + z_3^+s$ for $0 \le s \le 1$, there holds $[K_{A_0}(t, \Gamma_4^+)d](x')$

$$= \mathcal{F}_{\xi'}^{-1} \left[e^{-z_0 A^2 \langle t \rangle / 8} \cdot e^{z_0 A^2 \langle t \rangle / 8} \frac{\varphi_{A_0}(\xi')}{2\pi i} \int_0^1 e^{\lambda t} k(\xi', \lambda) (z_3^+ - z_1^+) \, ds \, \widehat{d}(\xi') \right] (x').$$

It thus holds that by Lemma 5.3, Parseval's identity, and the assumption for $k(\xi',\lambda)$

$$||K_{A_0}(t, \Gamma_4^+)d||_{L_q(\mathbf{R}^{N-1})} \le C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{2} - \frac{1}{q}\right)} \int_0^1 ||\varphi_{A_0}(\xi') \frac{e^{z_0 A^2 \langle t \rangle/8} e^{(\Re \lambda)t}}{(|\lambda|^{1/2} + A^{1/4})^2} \widehat{d}(\xi')||_{L_2(\mathbf{R}^{N-1})} ds =: I_1(t).$$

We choose a sufficiently small $A_0 \in (0,1)$ so that

$$e^{z_0A^2\langle t\rangle/8}e^{(\Re\lambda)t} \leq Ce^{-z_0A^2\langle t\rangle/8}e^{-cs\langle t\rangle} \quad \text{on supp}\, \varphi_{A_0}$$

for positive constants C and c. Then

$$I_{1}(t) \leq C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{2} - \frac{1}{q}\right)} \int_{0}^{1} e^{-cs\langle t \rangle} \left\| \varphi_{A_{0}}(\xi') \frac{e^{-z_{0}A^{2}\langle t \rangle/8}}{\left(|\lambda|^{1/2} + A^{1/4}\right)^{2}} \widehat{d}(\xi') \right\|_{L_{2}(\mathbf{R}^{N-1})} ds.$$

$$(5.18)$$

Let $0 < \delta < 2$. Since

$$|\lambda|^{1/2} \ge C(A\sqrt{1-s} + \sqrt{s})$$
 for $s \in [0, 1]$,

we see that

$$(|\lambda|^{1/2} + A^{1/4})^2 \ge (|\lambda|^{1/2})^{2-\delta} A^{\delta/4} \ge C(\sqrt{s})^{2-\delta} A^{\delta/4}. \tag{5.19}$$

Combining this inequality with (5.18) furnishes

$$I_{1}(t) \leq C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{2} - \frac{1}{q}\right)} \int_{0}^{1} \frac{e^{-cs\langle t \rangle}}{\sqrt{s^{2-\delta}}} ds \cdot \left\| e^{-z_{0}A^{2}\langle t \rangle/8} A^{-\delta/4} \widehat{d}(\xi') \right\|_{L_{2}(\mathbf{R}^{N-1})}$$

$$\leq C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{2} - \frac{1}{q}\right) - \frac{\delta}{2}} \left\| e^{-z_{0}A^{2}\langle t \rangle/8} A^{-\delta/4} \widehat{d}(\xi') \right\|_{L_{2}(\mathbf{R}^{N-1})},$$

which, combined with Parseval's identity and Young's inequality, yields

$$\begin{split} I_{1}(t) &\leq C \langle t \rangle^{-\frac{N-1}{2} \left(\frac{1}{2} - \frac{1}{q}\right) - \frac{\delta}{2}} \|\mathcal{F}_{\xi'}^{-1} [e^{-z_{0}A^{2} \langle t \rangle / 8} A^{-\delta/4} \widehat{d}(\xi')] \|_{L_{2}(\mathbf{R}^{N-1})} \\ &\leq C \langle t \rangle^{-\frac{N-1}{2} \left(\frac{1}{2} - \frac{1}{q}\right) - \frac{\delta}{2}} \|\mathcal{F}_{\xi'}^{-1} [e^{-z_{0}A^{2} \langle t \rangle / 8} A^{-\delta/4}] \|_{L_{r}(\mathbf{R}^{N-1})} \|d\|_{L_{p}(\mathbf{R}^{N-1})}, \end{split}$$

where 1 + (1/2) = (1/p) + (1/r).

From now on, we estimate $J_1(t):=\|\mathcal{F}_{\xi'}^{-1}[e^{-z_0A^2\langle t\rangle/8}A^{-\delta/4}]\|_{L_r(\mathbf{R}^{N-1})}$ by Lemma 5.4. By the Leibniz rule and Lemmas 3.1 and 3.2, we have for any multi-index $\alpha'\in\mathbf{N}_0^{N-1}$ and $\xi'\in\mathbf{R}^{N-1}\setminus\{0\}$

$$|\partial_{\varepsilon'}^{\alpha'}(e^{-z_0A^2\langle t\rangle/8}A^{-\delta/4})| \le C_{\alpha'}A^{-(\delta/4)-|\alpha'|}e^{-z_0A^2\langle t\rangle/16},$$

where $C_{\alpha'}$ is a positive constant independent of ξ' and t. Lemma 5.4 with $\sigma = 1 - (\delta/4)$, L = N - 2, and n = N - 1 then furnishes

$$|\mathcal{F}_{\varepsilon'}^{-1}[e^{-z_0 A^2 \langle t \rangle/8} A^{-\delta/4}](x')| \le C|x'|^{-(N-1-(\delta/4))} \quad (x' \in \mathbf{R}^{N-1} \setminus \{0\}).$$

By direct calculations, we also have

$$|\mathcal{F}_{\xi'}^{-1}[e^{-z_0 A^2 \langle t \rangle/8} A^{-\delta/4}](x')| \le C \int_{\mathbf{R}^{N-1}} e^{-z_0 A^2 \langle t \rangle/8} A^{-\delta/4} d\xi' \le C \langle t \rangle^{-(N-1-(\delta/4))/2},$$

and thus we obtain by these two inequalities

$$|\mathcal{F}_{\xi'}^{-1}[e^{-z_0A^2\langle t\rangle/8}A^{-\delta/4}](x')| \le \frac{C}{\langle t\rangle^{(N-1-(\delta/4))/2} + |x'|^{(N-1-(\delta/4))}}.$$

Let us choose the above δ so that

$$0 < \delta < \min \left\{ 2, 4(N-1) \left(\frac{1}{p} - \frac{1}{2} \right) \right\}.$$

Then we have

$$J_1(t) \leq C\langle t \rangle^{-\frac{1}{2}(N-1-\frac{\delta}{4})+\frac{N-1}{2r}} = C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{2}\right)+\frac{\delta}{8}}.$$

Hence

$$I_1(t) \le C\langle t \rangle^{-\frac{N-1}{2}(\frac{1}{p} - \frac{1}{q}) - \frac{3}{8}\delta} \|d\|_{L_p(\mathbf{R}^{N-1})},$$

which implies the desired estimate for $K_{A_0}(t; \Gamma_4^+)$ holds.

Case 2: $K_{A_0,\mathcal{M}}^{+,+}(t;\Gamma_4^+)$. In the same way as we have obtained (5.18), we obtain

$$||K_{A_0,\mathcal{M}}^{+,+}(t;\Gamma_4^+)\mathbf{f}||_{L_q(\mathbf{R}^{N-1})} \le C\langle t \rangle^{-\frac{N-1}{2}(\frac{1}{2}-\frac{1}{q})}$$

$$\times \int_{0}^{\infty} \int_{0}^{1} e^{-cs\langle t \rangle} \left\| \varphi_{A_{0}}(\xi') e^{-z_{0}A^{2}\langle t \rangle/8} k_{\mathcal{M}}(\xi', \lambda) \mathcal{M}_{+}(y_{N}) \widehat{\mathbf{f}}(\xi', y_{N}) \right\|_{L_{2}(\mathbf{R}^{N-1})} ds dy_{N}$$

$$=: I_{2}(t).$$

Since it holds by Lemma 3.1 that $C_1(|\lambda|^{1/2} + A) \leq |B_+ + A| \leq C_2|\lambda|^{1/2}$ for $A \in (0, 1)$ and $\lambda \in \widehat{\Gamma}_4^+$, we see that

$$\left| \frac{1}{B_+ - A} \right| = \left| \frac{B_+ + A}{B_+^2 - A^2} \right| \le \frac{C}{|\lambda|^{1/2}} \le \frac{\widetilde{C}}{|\lambda|^{1/2} + A}.$$

Therefore for $A \in (0,1)$ and $\lambda \in \widehat{\Gamma}_4^+$

$$|\mathcal{M}_+(y_N)| \le \frac{Ce^{-cAy_N}}{|\lambda|^{1/2} + A},$$

with positive constants C and c, which, combined with the assumption for $k_{\mathcal{M}}(\xi',\lambda)$ and (5.19), furnishes

$$|k_{\mathcal{M}}(\xi',\lambda)\mathcal{M}_{+}(y_{N})| \leq C \frac{A^{1/2}e^{-cAy_{N}}}{(|\lambda|^{1/2} + A^{1/4})^{2}} \leq C \frac{A^{1/2}e^{-cAy_{N}}}{(\sqrt{s})^{2-\delta}A^{\delta/4}} \quad (0 < \delta < 2).$$

One now sees that

$$I_{2}(t) \leq C\langle t \rangle^{-\frac{N-1}{2} \left(\frac{1}{2} - \frac{1}{q}\right)} \int_{0}^{1} \frac{e^{-cs\langle t \rangle}}{(\sqrt{s})^{2-\delta}} ds$$

$$\times \int_{0}^{\infty} \|e^{-z_{0}A^{2}\langle t \rangle/8} A^{1/2-\delta/4} e^{-cAy_{N}} \widehat{\mathbf{f}}(\xi', y_{N})\|_{L_{2}(\mathbf{R}^{N-1})} dy_{N}$$

$$\leq C\langle t \rangle^{-\frac{N-1}{2} \left(\frac{1}{2} - \frac{1}{q}\right) - \frac{\delta}{2}} \int_{0}^{\infty} \|e^{-z_{0}A^{2}\langle t \rangle/8} A^{1/2-\delta/4} e^{-cAy_{N}} \widehat{\mathbf{f}}(\xi', y_{N})\|_{L_{2}(\mathbf{R}^{N-1})} dy_{N},$$

$$(5.20)$$

and thus for 1 + (1/2) = (1/p) + (1/r)

$$\begin{split} I_{2}(t) &\leq C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{2} - \frac{1}{q}\right) - \frac{\delta}{2}} \\ &\times \int_{0}^{\infty} \|\mathcal{F}_{\xi'}^{-1} [e^{-z_{0}A^{2}\langle t \rangle/8} A^{1/2 - \delta/4} e^{-cAy_{N}}] \|_{L_{r}(\mathbf{R}^{N-1})} \|\mathbf{f}(\cdot, y_{N})\|_{L_{p}(\mathbf{R}^{N-1})} \, dy_{N} \\ &\leq C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{2} - \frac{1}{q}\right) - \frac{\delta}{2}} J_{2}(t) \|\mathbf{f}\|_{L_{p}(\mathbf{R}_{+}^{N})}, \end{split}$$

where for p' = p/(p-1)

$$J_2(t) =$$

$$\begin{cases} \left(\int_{0}^{\infty} \|\mathcal{F}_{\xi'}^{-1}[e^{-z_{0}A^{2}\langle t \rangle/8}A^{1/2-\delta/4}e^{-cAy_{N}}]\|_{L_{r}(\mathbf{R}^{N-1})}^{p'}dy_{N} \right)^{1/p'} & (1 0} \|\mathcal{F}_{\xi'}^{-1}[e^{-z_{0}A^{2}\langle t \rangle/8}A^{1/2-\delta/4}e^{-cAy_{N}}]\|_{L_{r}(\mathbf{R}^{N-1})} & (p = 1). \end{cases}$$

By the Leibniz rule and Lemmas 3.1 and 3.2, we have for any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$ and $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$

$$|\partial_{\xi'}^{\alpha'}(e^{-z_0A^2\langle t\rangle/8}A^{1/2-\delta/4}e^{-cAy_N})| \le C_{\alpha'}A^{1/2-\delta/4-|\alpha'|}e^{-z_0A^2\langle t\rangle/16}e^{-z_0A^2\langle t\rangle/16}e^{-$$

where $C_{\alpha'}$ is a positive constant independent of ξ' and t. Lemma 5.4 with $\sigma = 1/2 - \delta/4$, L = N - 1, and n = N - 1 then furnishes

$$|\mathcal{F}_{\xi'}^{-1}[e^{-z_0A^2\langle t\rangle/8}A^{1/2-\delta/4}e^{-cAy_N}](x')| \le C|x'|^{-(N-1+\sigma)} \quad (x' \in \mathbf{R}^{N-1} \setminus \{0\}).$$

By direct calculations, we also have

$$|\mathcal{F}_{\xi'}^{-1}[e^{-z_0A^2\langle t\rangle/8}A^{1/2-\delta/4}e^{-cAy_N}](x')|$$

$$\leq C \int_{\mathbf{R}^{N-1}} A^{1/2-\delta/4} e^{-cAy_N} d\xi' \leq C y_N^{-(N-1+\sigma)} \quad (y_N > 0),
|\mathcal{F}_{\xi'}^{-1} [e^{-z_0 A^2 \langle t \rangle/8} A^{1/2-\delta/4} e^{-cAy_N}](x')|
\leq C \int_{\mathbf{R}^{N-1}} e^{-z_0 A^2 \langle t \rangle/8} A^{1/2-\delta/4} d\xi' \leq C \langle t \rangle^{-\frac{1}{2}(N-1+\sigma)} \quad (t > 0).$$

Combining these three inequalities yields

$$|\mathcal{F}_{\xi'}^{-1}[e^{-z_0 A^2 \langle t \rangle/8} A^{1/2 - \delta/4} e^{-cAy_N}](x')| \le \frac{C}{|x'|^{N-1+\sigma} + y_N^{N-1+\sigma} + \langle t \rangle^{(N-1+\sigma)/2}},$$

which implies for any $\delta \in (0,2)$

$$\|\mathcal{F}_{\xi'}^{-1}[e^{-z_0A^2\langle t\rangle/8}A^{1/2-\delta/4}e^{-cAy_N}]\|_{L_r(\mathbf{R}^{N-1})} \le C(y_N + \langle t\rangle^{1/2})^{-(N-1)\left(\frac{1}{p} - \frac{1}{2}\right) - \left(\frac{1}{2} - \frac{\delta}{4}\right)}.$$

Let us choose the above δ so that

$$0<\delta<\min\left\{2,4N\left(\frac{1}{p}-\frac{1}{2}\right)\right\}.$$

Then we have

$$J_2(t) \le C\langle t \rangle^{-\frac{N-1}{2}(\frac{1}{p}-\frac{1}{2})-\frac{1}{2}(\frac{1}{p}-\frac{1}{2})+\frac{\delta}{8}}.$$

Hence

$$I_2(t) \le C\langle t \rangle^{-\frac{N-1}{2}(\frac{1}{p}-\frac{1}{q})-\frac{1}{2}(\frac{1}{p}-\frac{1}{2})-\frac{3}{8}\delta} \|\mathbf{f}\|_{L_p(\mathbf{R}^N)},$$

which implies the desired estimate for $K_{A_0,\mathcal{M}}^{+,+}(t;\Gamma_4^+)$ holds.

Case 3: $L_{A_0,\mathcal{M}}^+(t;\Gamma_4^+)$. In the same way as we have obtained (5.20), we obtain for $\delta \in (0,2)$

$$\begin{split} & \|[L_{A_0,\mathcal{M}}^+(t;\Gamma_4^+)d](\cdot,x_N)\|_{L_q(\mathbf{R}^{N-1})} \\ & \leq C\langle t\rangle^{-\frac{N-1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{\delta}{2}} \|e^{-z_0A^2\langle t\rangle/8}A^{1/2-\delta/4}e^{-cAx_N}\widehat{d}(\xi')\|_{L_2(\mathbf{R}^{N-1})} =: I_3(x_N,t) \end{split}$$

By Parseval's identity and Young's inequality,

 $I_3(x_N,t)$

$$\leq C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{2} - \frac{1}{q}\right) - \frac{\delta}{2}} \|\mathcal{F}_{\xi'}^{-1}[e^{-z_0 A^2 \langle t \rangle / 8} A^{1/2 - \delta/4} e^{-cAx_N}] \|L_{r}(\mathbf{R}^{N-1})\| d\|_{L_p(\mathbf{R}^{N-1})},$$

where 1 + (1/2) = (1/p) + (1/r). Similarly to $J_2(t)$ in Case 2, we observe for any $\delta \in (0,2)$ that

$$\|\mathcal{F}_{\xi'}^{-1}[e^{-z_0A^2\langle t\rangle/8}A^{1/2-\delta/4}e^{-cAx_N}]\|_{L_r(\mathbf{R}^{N-1})} \le C(x_N + \langle t\rangle^{1/2})^{-(N-1)\left(\frac{1}{p} - \frac{1}{2}\right) - \left(\frac{1}{2} - \frac{\delta}{4}\right)}.$$

It thus holds that

$$\left(\int_0^\infty I_3(x_N,t)^q dx_N\right)^{1/q} \le C\langle t \rangle^{-\frac{N-1}{2}\left(\frac{1}{p}-\frac{1}{q}\right)-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{q}\right)-\frac{3}{8}\delta} \|d\|_{L_p(\mathbf{R}^{N-1})}$$

under the assumption

$$0 < \delta < \min \left\{ 2, 4 \left((N-1) \left(\frac{1}{p} - \frac{1}{2} \right) + \frac{1}{2} - \frac{1}{q} \right) \right\}.$$

This implies the desired estimate for $L_{A_0,\mathcal{M}}^+(t;\Gamma_4^+)$, which completes the proof of Lemma 5.12.

Combining Lemma 5.12 with (5.15), (5.16), and (5.17) yields Theorem 5.11 immediately. This completes the proof of Theorem 5.11.

5.4. **Analysis for** Γ_5^{\pm} . Similarly to [7, Subsection 4.4], we can prove by Lemmas 3.1 and 3.4 the following theorem.

Theorem 5.13. Let $1 \le p \le 2 \le q \le \infty$. Then there exist constants $A_0 \in (0, A_3)$ and $c_0 > 0$ such that for any $t \ge 1$ and $(d, \mathbf{f}) \in Y_p$

$$\begin{split} &\|\mathcal{H}_{A_0}^1(t;\Gamma_5^\pm)d\|_{L_q(\mathbf{R}^{N-1})} \leq Ce^{-c_0t}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ &\|\mathcal{H}_{A_0}^2(t;\Gamma_5^\pm)\mathbf{f}\|_{L_q(\mathbf{R}^{N-1})} \leq Ce^{-c_0t}\|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)},\\ &\|\mathcal{U}_{A_0}^1(t;\Gamma_5^\pm)d\|_{L_q(\dot{\mathbf{R}}^N)} \leq Ce^{-c_0t}\|d\|_{L_p(\mathbf{R}^{N-1})},\\ &\|\mathcal{U}_{A_0}^2(t;\Gamma_5^\pm)\mathbf{f}\|_{L_q(\dot{\mathbf{R}}^N)} \leq Ce^{-c_0t}\|\mathbf{f}\|_{L_p(\dot{\mathbf{R}}^N)}, \end{split}$$

where C is a positive constant independent of t, d, and f.

5.5. **Proof of Theorem 2.3.** Recalling (5.1), we observe for $S \in \{H, U\}$ that the operators in Theorem 2.3 are given by

$$\begin{split} \mathcal{S}_{A_0}^{1\pm}(t)d &= \mathcal{S}_{A_0}^1(t;\Gamma_{\mathrm{res}}^\pm)d, & \mathcal{S}_{A_0}^{2\pm}(t)\mathbf{f} = \mathcal{S}_{A_0}^2(t;\Gamma_{\mathrm{res}}^\pm)\mathbf{f}, \\ \widetilde{\mathcal{S}}_{A_0}^1(t)d &= \sum_{\mathfrak{a}\in\{+,-\}} \sum_{j\in\{1,4,5\}} \mathcal{S}_{A_0}^1(t;\Gamma_j^{\mathfrak{a}})d, & \widetilde{\mathcal{S}}_{A_0}^2(t)\mathbf{f} = \sum_{\mathfrak{a}\in\{+,-\}} \sum_{j\in\{1,4,5\}} \mathcal{S}_{A_0}^2(t;\Gamma_j^{\mathfrak{a}})\mathbf{f}. \end{split}$$

Theorems 5.5, 5.8, 5.11, and 5.13 then yields Theorem 2.3 immediately. This completes the proof of Theorem 2.3.

6. Time-decay estimates for high frequency part

This section proves Theorem 2.5. Suppose $\rho_- > \rho_+ > 0$ throughout this section. Let us denote the points of intersection between $\lambda = -1 + is$ $(s \in \mathbf{R})$ and Γ_0^{\pm} given in (2.3) by z_4^{\pm} , and let A_0 be the positive constant given in Theorem 2.3. We define $A_{\infty} = A_{\text{high}}(1, \Im z_4^+)$ for the positive constant A_{high} given in Proposition 4.6. In addition, we set $M_1 = A_0/2$ and $M_2 = 3A_{\infty}$ in Proposition 4.7. Then we have $a_0 \in (0,1)$ from Proposition 4.7 and denote the points of intersection between $\lambda = -a_0 + is$ $(s \in \mathbf{R})$ and Γ_0^{\pm} by z_5^{\pm} . Note that $\Im z_5^- = -\Im z_5^+$.

Now we define integral paths $\widehat{\Gamma}_6$ and $\widehat{\Gamma}_7$ as follows:

$$\widehat{\Gamma}_{6} = \{ \lambda \in \mathbf{C} : \lambda = -a_{0} + si, -\Im z_{5}^{+} \le s \le \Im z_{5}^{+} \},$$

$$\widehat{\Gamma}_{7} = \{ \lambda \in \mathbf{C} : \lambda = -a_{0} + i\Im z_{5}^{+} + se^{i(\pi - \theta_{1})}, s \ge 0 \}$$

$$\cup \{ \lambda \in \mathbf{C} : \lambda = -a_{0} - i\Im z_{5}^{+} + se^{-i(\pi - \theta_{1})}, s \ge 0 \},$$

Then Propositions 4.6 and 4.7 yield the following lemma.

Lemma 6.1. (1) $\Re B_{\pm} > 0$, $F(A,\lambda) \neq 0$, and $L(A,\lambda) \neq 0$ when $A \geq A_0/2$ and $\lambda \in \Lambda(a_0, \Im z_5^+)$, where $\Lambda(a_0, \Im z_5^+)$ is given in (4.9).

(2) Let $s \in \mathbf{R}$ and $\alpha' \in \mathbf{N}_0^{N-1}$. Then for any $\lambda \in \widehat{\Gamma}_6$ it holds that on $\operatorname{supp} \varphi_{A_\infty} \cup \operatorname{supp} \varphi_{[A_0,A_\infty]}$

$$\begin{split} |\partial_{\xi}'B_{\pm}^{s}| &\leq CA^{s-|\alpha'|}, \quad |\partial_{\xi'}^{\alpha'}E^{s}| \leq CA^{s-|\alpha'|}, \quad |\partial_{\xi'}^{\alpha'}(A+B_{\pm})^{s}| \leq CA^{s-|\alpha'|}, \\ |\partial_{\varepsilon'}^{\alpha'}F(A,\lambda)^{s}| &\leq CA^{3s-|\alpha'|}, \quad |\partial_{\varepsilon'}^{\alpha'}L(A,\lambda)^{-1}| \leq CA^{-4-|\alpha'|}. \end{split}$$

Let $S \in \{\mathcal{H}, \mathcal{U}\}$ and $Z \in \{A_{\infty}, [A_0, A_{\infty}]\}$. Recalling (2.10), we have by Lemma 6.1 (1) and Cauchy's integral theorem

$$\begin{split} \mathcal{S}_{Z}^{1}(t)d &= \sum_{j=6}^{7} \mathcal{S}_{Z}^{1}(t;\Gamma_{j})d, \quad \mathcal{S}_{Z}^{1}(t;\Gamma_{j})d := \mathcal{F}_{\xi'}^{-1}[\widehat{\mathcal{S}}_{Z}^{1}(t;\widehat{\Gamma}_{j})d](x'), \\ \mathcal{S}_{Z}^{2}(t)\mathbf{f} &= \sum_{j=6}^{7} \mathcal{S}_{Z}^{2}(t;\Gamma_{j})\mathbf{f}, \quad \mathcal{S}_{Z}^{2}(t;\Gamma_{j})\mathbf{f} := \mathcal{F}_{\xi'}^{-1}[\widehat{\mathcal{S}}_{Z}^{2}(t;\widehat{\Gamma}_{j})\mathbf{f}](x'). \end{split}$$

Similarly to [7, Section 5], we can prove from Lemmas 6.1, 3.1, and 3.4 the following theorem by choosing A_{∞} larger if necessary.

Theorem 6.2. Let $q \in (1, \infty)$ and $Z \in \{A_{\infty}, [A_0, A_{\infty}]\}$. Let j = 6, 7 and $k \in \mathbb{N}_0$. Then for any $t \geq 1$ and $(d, \mathbf{f}) \in X_q$

$$\begin{split} \|(\partial_t^k \mathcal{H}_Z^1(t;\Gamma_j)d,\partial_t^k \mathcal{H}_Z^2(t;\Gamma_j)\mathbf{f})\|_{W_q^{3-1/q}(\mathbf{R}^{N-1})} &\leq Ce^{-ct}\|(d,\mathbf{f})\|_{X_q}, \\ \|(\partial_t^k \mathcal{U}_Z^1(t;\Gamma_j)d,\partial_t^k \mathcal{U}_Z^2(t;\Gamma_j)\mathbf{f})\|_{H^2(\dot{\mathbf{R}}^N)} &\leq Ce^{-ct}\|(d,\mathbf{f})\|_{X_q}, \end{split}$$

where C and c are positive constants independent of t, d, and f.

Theorem 6.2 yields Theorem 2.5 immediately. This completes the proof of Theorem 2.5.

Α

In this appendix, we consider the whole space problems in (3.28) and compute the representation formulas of $\widehat{\psi}_{\pm}|_{x_N=0}$ and $\partial_N \widehat{\psi}_{\pm}|_{x_N=0}$. We define the Fourier transform of f=f(x) and the inverse Fourier transform of $g=g(\xi)$ by

$$\mathcal{F}[f](\xi) = \int_{\mathbf{R}^N} e^{-ix\cdot\xi} f(x) \, dx, \quad \mathcal{F}_{\xi}^{-1}[g](x) = \frac{1}{(2\pi)^N} \int_{\mathbf{R}^N} e^{ix\cdot\xi} g(\xi) \, d\xi.$$

Let us denote the jth component of \mathbf{F} by F_j . In [10, Section 2], we have

$$\psi_{\pm} = \mathcal{F}_{\xi}^{-1} \left[\frac{\mathcal{F}[\mathbf{F}](\xi)}{\rho_{\pm}\lambda + \mu_{\pm}|\xi|^{2}} \right] (x) + \sum_{j=1}^{N} \mathcal{F}_{\xi}^{-1} \left[\frac{(i\xi)(i\xi_{j})\mathcal{F}[F_{j}](\xi)}{|\xi|^{2}(\rho_{\pm}\lambda + \mu_{\pm}|\xi|^{2})} \right] (x),$$

$$\phi_{+} = \phi_{-} = -\sum_{j=1}^{N} \mathcal{F}_{\xi}^{-1} \left[\frac{i\xi_{j}\mathcal{F}[F_{j}](\xi)}{|\xi|^{2}} \right] (x).$$

The formulas of ϕ_{\pm} imply $\llbracket \phi \rrbracket = \phi_{+}(x', 0+) - \phi_{-}(x', 0-) = 0$. For $k = 1, \dots, N-1$, $\psi_{k\pm}$ can be written as

$$\psi_{k\pm} = \mathcal{F}_{\xi}^{-1} \left[\frac{\mathcal{F}[F_k](\xi)}{\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2} \right] (x) - \sum_{j=1}^{N-1} \mathcal{F}_{\xi}^{-1} \left[\frac{\xi_k \xi_j \mathcal{F}[F_j](\xi)}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} \right] (x) + \mathcal{F}_{\xi}^{-1} \left[\frac{(i\xi_k)(i\xi_N)\mathcal{F}[F_N](\xi)}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} \right] (x),$$

and also

$$\psi_{N\pm} = \mathcal{F}_{\xi}^{-1} \left[\frac{A^2 \mathcal{F}[F_N](\xi)}{|\xi|^2 (\rho_{\pm} \lambda + \mu_{\pm} |\xi|^2)} \right] (x) + \sum_{i=1}^{N-1} \mathcal{F}_{\xi}^{-1} \left[\frac{(i\xi_N)(i\xi_j) \mathcal{F}[F_j](\xi)}{|\xi|^2 (\rho_{\pm} \lambda + \mu_{\pm} |\xi|^2)} \right] (x),$$

where A is given in (3.1). It holds for l = 1, ..., N that

$$\begin{split} \mathcal{F}[F_l](\xi) &= \int_0^\infty e^{-iy_N \xi_N} \widehat{F}_l(\xi', y_N) \, dy_N + \int_0^\infty e^{iy_N \xi_N} \widehat{F}_l(\xi', -y_N) \, dy_N \\ &= \sum_{\mathfrak{a} \in \{+, -\}} \int_0^\infty e^{-i\mathfrak{a}y_N \xi_N} \widehat{F}_l(\xi', \mathfrak{a}y_N) \, dy_N \end{split}$$

which, inserted into the above formulas of $\psi_{k\pm}$ and $\psi_{N\pm}$, furnishes

$$\begin{split} &\widehat{\psi}_{k\pm}(\xi',x_{N},\lambda) \\ &= \sum_{\mathfrak{a}\in\{+,-\}} \int_{0}^{\infty} \widehat{F}_{k}(\xi',\mathfrak{a}y_{N}) \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{i(x_{N} - \mathfrak{a}y_{N})\xi_{N}}}{\rho_{\pm}\lambda + \mu_{\pm}|\xi|^{2}} d\xi_{N} \right) dy_{N} \\ &- \sum_{j=1}^{N-1} \sum_{\mathfrak{a}\in\{+,-\}} \int_{0}^{\infty} \xi_{k}\xi_{j}\widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{i(x_{N} - \mathfrak{a}y_{N})\xi_{N}}}{|\xi|^{2}(\rho_{\pm}\lambda + \mu_{\pm}|\xi|^{2})} d\xi_{N} \right) dy_{N} \\ &+ \sum_{\mathfrak{a}\in\{+,-\}} \int_{0}^{\infty} i\xi_{k}\widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{i\xi_{N}e^{i(x_{N} - \mathfrak{a}y_{N})\xi_{N}}}{|\xi|^{2}(\rho_{\pm}\lambda + \mu_{\pm}|\xi|^{2})} d\xi_{N} \right) dy_{N}, \\ &\widehat{\psi}_{N\pm}(\xi',x_{N},\lambda) \\ &= \sum_{\mathfrak{a}\in\{+,-\}} \int_{0}^{\infty} A^{2}\widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{i(x_{N} - \mathfrak{a}y_{N})\xi_{N}}}{|\xi|^{2}(\rho_{\pm}\lambda + \mu_{\pm}|\xi|^{2})} d\xi_{N} \right) dy_{N} \\ &+ \sum_{j=1}^{N-1} \sum_{\mathfrak{a}\in\{+,-\}} \int_{0}^{\infty} i\xi_{j}\widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{i\xi_{N}e^{i(x_{N} - \mathfrak{a}y_{N})\xi_{N}}}{|\xi|^{2}(\rho_{\pm}\lambda + \mu_{\pm}|\xi|^{2})} d\xi_{N} \right) dy_{N}. \end{split}$$

Applying ∂_N to these formulas yields

$$\begin{split} &\partial_N \widehat{\psi}_{k\pm}(\xi',x_N,\lambda) \\ &= \sum_{\mathfrak{a} \in \{+,-\}} \int_0^\infty \widehat{F}_k(\xi',\mathfrak{a}y_N) \left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{i\xi_N e^{i(x_N - \mathfrak{a}y_N)\xi_N}}{\rho_\pm \lambda + \mu_\pm |\xi|^2} \, d\xi_N \right) \, dy_N \\ &- \sum_{j=1}^{N-1} \sum_{\mathfrak{a} \in \{+,-\}} \int_0^\infty \xi_k \xi_j \widehat{F}_j(\xi',\mathfrak{a}y_N) \left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{i\xi_N e^{i(x_N - \mathfrak{a}y_N)\xi_N}}{|\xi|^2 (\rho_\pm \lambda + \mu_\pm |\xi|^2)} \, d\xi_N \right) \, dy_N \\ &+ \sum_{\mathfrak{a} \in \{+,-\}} \int_0^\infty i\xi_k \widehat{F}_N(\xi',\mathfrak{a}y_N) \left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{(i\xi_N)^2 e^{i(x_N - \mathfrak{a}y_N)\xi_N}}{|\xi|^2 (\rho_\pm \lambda + \mu_\pm |\xi|^2)} \, d\xi_N \right) \, dy_N, \\ &\partial_N \widehat{\psi}_{N\pm}(\xi',x_N,\lambda) \\ &= \sum_{\mathfrak{a} \in \{+,-\}} \int_0^\infty A^2 \widehat{F}_N(\xi',\mathfrak{a}y_N) \left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{i\xi_N e^{i(x_N - \mathfrak{a}y_N)\xi_N}}{|\xi|^2 (\rho_\pm \lambda + \mu_\pm |\xi|^2)} \, d\xi_N \right) \, dy_N \\ &+ \sum_{j=1}^{N-1} \sum_{\mathfrak{a} \in \{+,-\}} \int_0^\infty i\xi_j \widehat{F}_j(\xi',\mathfrak{a}y_N) \left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{(i\xi_N)^2 e^{i(x_N - \mathfrak{a}y_N)\xi_N}}{|\xi|^2 (\rho_\pm \lambda + \mu_\pm |\xi|^2)} \, d\xi_N \right) \, dy_N. \end{split}$$

Now we have

Lemma A.1. Let $\xi' = (\xi_1, ..., \xi_{N-1}) \in \mathbf{R}^{N-1} \setminus \{0\}$ and $a \in \mathbf{R} \setminus \{0\}$. Then

$$\begin{split} \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ia\xi_N}}{\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2} \, d\xi_N &= \frac{e^{-B_{\pm}|a|}}{2\mu_{\pm}B_{\pm}}, \\ \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{i\xi_N e^{ia\xi_N}}{\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2} \, d\xi_N &= -\mathrm{sign}(a) \frac{e^{-B_{\pm}|a|}}{2\mu_{\pm}}, \\ \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ia\xi_N}}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} \, d\xi_N &= -\frac{1}{2\mu_{\pm}(A^2 - B_{\pm}^2)} \left(\frac{e^{-A|a|}}{A} - \frac{e^{-B_{\pm}|a|}}{B_{\pm}} \right), \\ \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{i\xi_N e^{ia\xi_N}}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} \, d\xi_N &= -\frac{\mathrm{sign}(a)}{2\mu_{\pm}(A^2 - B_{\pm}^2)} \left(-e^{-A|a|} + e^{-B_{\pm}|a|} \right), \\ \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{(i\xi_N)^2 e^{ia\xi_N}}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} \, d\xi_N &= -\frac{1}{2\mu_{\pm}(A^2 - B_{\pm}^2)} \left(Ae^{-A|a|} - B_{\pm}e^{-B_{\pm}|a|} \right), \end{split}$$

where A and B_{\pm} are defined in (3.1). Here sign(a) = 1 when a > 0 and sign(a) = -1 when a < 0.

Proof. The first and third formulas follow from the residue theorem. Differentiating the first formula with respect to a, we have the second formula. Analogously the fourth and fifth formulas follow from the third formula. This completes the proof of Lemma A.1.

Recall $\mathcal{M}_{\pm}(a)$ given in (3.2), and then

$$e^{-Aa} = (A - B_{\pm})\mathcal{M}_{\pm}(a) + e^{-B_{\pm}a} \quad (a \ge 0).$$

By this relation and Lemma A.1, we have

Lemma A.2. Let $\xi' = (\xi_1, ..., \xi_{N-1}) \in \mathbf{R}^{N-1} \setminus \{0\}$ and $a \in \mathbf{R} \setminus \{0\}$. Then

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ia\xi_N}}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} d\xi_N = -\frac{\mathcal{M}_{\pm}(|a|)}{2\mu_{\pm}A(A + B_{\pm})} + \frac{e^{-B_{\pm}|a|}}{2\mu_{\pm}AB_{\pm}(A + B_{\pm})},$$

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{i\xi_N e^{ia\xi_N}}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} d\xi_N = \frac{\text{sign}(a) \cdot \mathcal{M}_{\pm}(|a|)}{2\mu_{\pm}(A + B_{\pm})},$$

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{(i\xi_N)^2 e^{ia\xi_N}}{|\xi|^2 (\rho_{\pm}\lambda + \mu_{\pm}|\xi|^2)} d\xi_N = -\frac{A\mathcal{M}_{\pm}(|a|)}{2\mu_{\pm}(A + B_{\pm})} - \frac{e^{-B_{\pm}|a|}}{2\mu_{\pm}(A + B_{\pm})}.$$

Together with the above formulas of $\hat{\psi}_{k\pm}$, $\hat{\psi}_{N\pm}$, $\partial_N \hat{\psi}_{k\pm}$, and $\partial_N \hat{\psi}_{N\pm}$, we obtain by Lemmas A.1 and A.2

$$\begin{split} \widehat{\psi}_{k\pm}(\xi',0,\lambda) &= \sum_{\mathfrak{a}\in\{+,-\}} \bigg\{ \int_{0}^{\infty} \frac{e^{-B_{\pm}y_{N}}}{2\mu_{\pm}B_{\pm}} \widehat{F}_{k}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &+ \sum_{j=1}^{N-1} \int_{0}^{\infty} \frac{\xi_{k}\xi_{j}\mathcal{M}_{\pm}(y_{N})}{2\mu_{\pm}A(A+B_{\pm})} \widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &- \sum_{j=1}^{N-1} \int_{0}^{\infty} \frac{\xi_{k}\xi_{j}e^{-B_{\pm}y_{N}}}{2\mu_{\pm}AB_{\pm}(A+B_{\pm})} \widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &- \mathfrak{a} \int_{0}^{\infty} \frac{i\xi_{k}\mathcal{M}_{\pm}(y_{N})}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \bigg\}, \end{split}$$

$$\begin{split} \widehat{\psi}_{N\pm}(\xi',0,\lambda) &= \sum_{\mathfrak{a}\in\{+,-\}} \bigg\{ -\int_{0}^{\infty} \frac{A\mathcal{M}_{\pm}(y_{N})}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &+ \int_{0}^{\infty} \frac{Ae^{-B_{\pm}y_{N}}}{2\mu_{\pm}B_{\pm}(A+B_{\pm})} \widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &- \mathfrak{a} \sum_{j=1}^{N-1} \int_{0}^{\infty} \frac{i\xi_{j}\mathcal{M}_{\pm}(y_{N})}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \bigg\}, \\ \partial_{N}\widehat{\psi}_{k\pm}(\xi',0,\lambda) &= \sum_{\mathfrak{a}\in\{+,-\}} \bigg\{ \mathfrak{a} \int_{0}^{\infty} \frac{e^{-B_{\pm}y_{N}}}{2\mu_{\pm}} \widehat{F}_{k}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &+ \mathfrak{a} \sum_{j=1}^{N-1} \int_{0}^{\infty} \frac{\xi_{j}\xi_{k}\mathcal{M}_{\pm}(y_{N})}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &- \int_{0}^{\infty} \frac{i\xi_{k}A\mathcal{M}_{\pm}(y_{N})}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &- \int_{0}^{\infty} \frac{i\xi_{k}e^{-B_{\pm}y_{N}}}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \bigg\}, \\ \partial_{N}\widehat{\psi}_{N\pm}(\xi',0,\lambda) &= \sum_{\mathfrak{a}\in\{+,-\}} \bigg\{ -\mathfrak{a} \int_{0}^{\infty} \frac{A^{2}\mathcal{M}_{\pm}(y_{N})}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{N}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &- \sum_{j=1}^{N-1} \int_{0}^{\infty} \frac{i\xi_{j}A\mathcal{M}_{\pm}(\mathfrak{a}y_{N})}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \\ &- \sum_{i=1}^{N-1} \int_{0}^{\infty} \frac{i\xi_{j}e^{-B_{\pm}y_{N}}}{2\mu_{\pm}(A+B_{\pm})} \widehat{F}_{j}(\xi',\mathfrak{a}y_{N}) \, dy_{N} \bigg\}. \end{split} \tag{A.1}$$

This completes the proof of the appendix.

References

- [1] M. Hieber and H. Saito. Strong solutions for two-phase free boundary problems for a class of non-Newtonian fluids. J. Evol. Equ., 17(1):335–358, 2017.
- [2] C. Miao, B. Yuan, and B. Zhang. Well-posedness of the Cauchy problem for the fractional power dissipative equations. *Nonlinear Anal.*, 68(3):461–484, 2008.
- [3] J. Prüss and G. Simonett. Analysis of the boundary symbol for the two-phase Navier-Stokes equations with surface tension. *Banach Center Publications*, 86:265–285, 2009.
- [4] J. Prüss and G. Simonett. On the Rayleigh-Taylor instability for the two-phase Navier-Stokes equations. *Indiana Univ. Math. J.*, 59(6):1853–1871, 2010.
- [5] J. Prüss and G. Simonett. On the two-phase Navier-Stokes equations with surface tension. Interfaces Free Bound., 12(3):311–345, 2010.
- [6] J. Prüss and G. Simonett. Analytic solutions for the two-phase Navier-Stokes equations with surface tension and gravity. In *Parabolic Problems*, volume 80 of *Progr. Nonlinear Differential Equations Appl.*, pages 507–540. Birkhäuser/Springer Basel AG, Basel, 2011.
- [7] H. Saito and Y. Shibata. On decay properties of solutions to the Stokes equations with surface tension and gravity in the half space. *J. Math. Soc. Japan*, 68(4):1559–1614, 2016.
- [8] H. Saito, Y. Shibata, and X. Zhang. Some free boundary problem for two-phase inhomogeneous incompressible flows. SIAM J. Math. Anal., 52(4):3397–3443, 2020.
- [9] Y. Shibata and S. Shimizu. A decay property of the Fourier transform and its application to the Stokes problem. J. Math. Fluid Mech., 3(3):213-230, 2001.
- [10] Y. Shibata and S. Shimizu. Maximal L_p-L_q regularity for the two-phase Stokes equations; model problems. J. Differential Equations, 251(2):373–419, 2011.

- [11] Y. Wang, I. Tice, and C. Kim. The viscous surface-internal wave problem: global well-posedness and decay. Arch. Ration. Mech. Anal., 212(1):1–92, 2014.
- [12] L. Xu and Z. Zhang. On the free boundary problem to the two viscous immiscible fluids. J. Differential Equations, 248(5):1044–1111, 2010.
- [13] M. Yamamoto and Y. Sugiyama. Asymptotic expansion of solutions to the drift-diffusion equation with fractional dissipation. *Nonlinear Anal.*, 141:57–87, 2016.

Department of Mathematics, The University of Electro-Communications, 5-1 Chofugaoka 1-chome, Chofu, Tokyo 182-8585, Japan

 $Email\ address: \ {\tt hsaito@uec.ac.jp}$