Strongly connected orientation with minimum lexicographic order of indegrees *

Hongyu Zhou^a, Xinmin Hou^b

^{a,b} Key Laboratory of Wu Wen-Tsun Mathematics
School of Mathematical Sciences
University of Science and Technology of China
Hefei, Anhui 230026, China.

Abstract

Given a simple undirected graph G, an orientation of G is to assign every edge of G a direction. Borradaile et al gave a greedy algorithm SC-Path-Reversal (in polynomial time) which finds a strongly connected orientation that minimizes the maximum indegree, and conjectured that SC-Path-Reversal is indeed optimal for the "minimizing the lexicographic order" objective as well. In this note, we give a positive answer to the conjecture, that is we show that the algorithm SC-PATH-REVERSAL finds a strongly connected orientation that minimizes the lexicographic order of indegrees.

1 Introduction

Graph orientation has long been studied and is a rich field under different conditions. In this note we mainly concern about the strongly-connected orientation with minimum lexicographic order. This objective arises from a telecommunication network design problem [2, 4]. Let G = (V, E) be an undirected simple graph. An orientation Λ of G is an assignment to each edge a direction. By a strongly-connected orientation, we mean the digraph that we obtain is strongly-connected. In a digraph $D = (V, \Lambda)$, the indegree of a vertex v is the number of arcs that are directed to v, denoted by $d_{\Lambda}^-(v)$. The indegree sequence of a digraph (or an orientation) is defined as a non-increasing sequence of the indegrees of all the vertices, that is, we place the indegrees of vertices in a non-increasing order. To compare distinct indegree sequences of two orientations of an undirected graph, we apply the lexicographic order, i.e. let $s = (s_1, s_2, \ldots, s_n)$ and $t = (t_1, t_2, \ldots, t_n)$ be the

^{*}The work was supported by NNSF of China (No. 11671376) and NSF of Anhui Province (No. 1708085MA18) and Anhui Initiative in Quantum Information Technologies (AHY150200).

indegree sequences of two distinct orientations of an undirected graph, respectively, we say s is smaller than t if there exists an integer k with $1 \le k \le n$ such that $s_k < t_k$ and $s_i = t_i$ for all i < k, and vise versa.

Let $D = (V, \Lambda)$ be a digraph and $u, v \in V$, we say u two-reaches to v (or v is two-reachable from u) if there are two arc-disjoint directed paths from u to v in D. A directed path from u to v is called reversible if $d^-(u) < d^-(v) - 1$, and is called strongly reversible if $d^-(u) < d^-(v) - 1$ and u two-reaches v in D.

The following greedy algorithm was first given by de Fraysseix and de Mendez [3]. It has

Algorithm 1.1: PATH-REVERSAL

Input: Undirected simple graph G

Output: Orientation Λ

- 1 Arbitrarily orient every edge of G.
- 2 While there is a reversible path, reverse it.
- з Repeat step 2.

been shown that the algorithm finds an orientation that minimizes the maximum indegree, which is proved by Venkateswaran [4], Asahiro et al. [1], and de Fraysseix and de Mendez [3], respectively. In fact, Path-Reversal can do something more, Borradaile et al. [2] showed that Path-Reversal indeed finds an orientation that minimizes the lexicographic order of the indegrees.

Lemma 1 (Lemma 2 in [2]). Reversing a directed path from u to v maintains the strong connectivity if and only if u two-reaches v. Particularly, v two-reaches v itself.

By Lemma 1, we know that reversing a strongly reversible direct path remains the resulting digraph strongly-connected. Based on PATH-REVERSAL, Borradaile et al. [2] gave a modified version of it as shown in the following.

Algorithm 1.2: SC-PATH-REVERSAL

Input: Undirected simple graph G admitting a strongly connected orientation

Output: Orientation Λ

- 1 Find an arbitrary strongly-connected orientation.
- 2 If there exists a strongly reversible path, reverse it.
- з Repeat step 2.

Borradaile et al [2] showed that the algorithm SC-Path-Reversal finds (in polynomial time) a strongly connected orientation that minimizes the maximum indegree, and conjectured that SC-Path-Reversal is indeed optimal for the "minimizing the lexicographic order" objective as well.

Conjecture 2 (Borradaile et al [2]). The algorithm SC-Path-Reversal finds a strongly connected orientation that minimizes the lexicographic order of indegrees.

In this note, we give a positive answer to Conjecture 2. The following is our main theorem.

Theorem 3. The algorithm SC-PATH-REVERSAL finds an orientation that minimizes the lexicographic order of the indegrees.

In the rest of the note, we give the proof of Theorem 3.

2 Proof of Theorem 3

Before proving that the orientation output has the minimum lexicographic order, we introduce two lemmas given in [2].

Lemma 4 (Lemma 2 in [2]). In a digraph, let s and t be two vertices (can be identical) that 2-reach a vertex v. If there exists a vertex u such that one $u \to s$ and one $u \to t$ paths are arc-disjoint, then u two-reaches v.

The following lemma is a variable version of Lemma 4 in [2], the proof is the same as the one of that lemma, so we omit it here.

Lemma 5. Let v be a vertex in a strongly connected digraph $D = (V, \Lambda)$ and let U be the set of vertices that 2-reach v. Then for any component C of $D[V \setminus U]$, there is exactly one arc from C to U.

Remark 1. By a component we mean a connected component instead of a strongly-connected component.

Now we are ready to give the proof of the main theorem.

Proof of Theorem 3. Let G = (V, E) be the underlying graph. Let $D_{pr} = (V, \Lambda_{pr})$ be the orientation founded by the SC-PATH-REVERSAL, and let $D_{lex} = (V, \Lambda_{lex})$ be a strongly connected orientation that minimizes the lexicographic order of indegrees among all strongly connected orientations of G. Write $d_{pr}^-(v)$ and $d_{lex}^-(v)$ for the indegree of v in D_{pr} and D_{lex} , respectively. Define

$$\Delta := \sum_{v \in V} |d^-_{lex}(v) - d^-_{pr}(v)|$$

and

$$S := \{ v \, | \, d^-_{lex}(v) \neq d^-_{pr}(v) \}.$$

Choose Λ_{lex} such that it minimizes Δ . If $S = \emptyset$, then Λ_{pr} is a strongly-connected orientation having the same lexicographic order of indegree as Λ_{lex} , we are done. So assume $S \neq \emptyset$. Denote $M_1 = \max\{d_{lex}^-(v) | v \in S\}$ and $S_1 = \{u \in S | d_{lex}^-(u) = M_1\}$. Denote $M_2 = \max\{d_{pr}^-(u) | u \in S_1\}$. Choose $v \in S_1$ such that $d_{pr}^-(v) = M_2$.

Case 1: $d_{lex}^-(v) > d_{pr}^-(v)$.

Let U be the set of vertices that two-reach v in D_{lex} . By Lemma 5, there is exactly one arc from each component of $G[V \setminus U]$ to U. Thus, on the one hand,

$$\sum_{u \in U} d^-_{lex}(u) = |E(G[U])| + c(G[V \setminus U]),$$

where c(G) denotes the number of components of a graph G.

While, on the other hand, since D_{pr} is strongly connected, there is at least one arc from each component of $G[V \setminus U]$ to U. Thus

$$\sum_{u \in U} d_{pr}^-(u) \ge |E(G[U])| + c(G[V \setminus U]).$$

So we get $\sum_{u \in U} d^-_{lex}(u) \leq \sum_{u \in U} d^-_{pr}(u)$. Since $d^-_{lex}(v) > d^-_{pr}(v)$, there exists a vertex $w \in U$ such that $d^-_{lex}(w) < d^-_{pr}(w)$. Clearly, $w \in S$. By the choice of v, we have $d^-_{lex}(w) \leq d^-_{lex}(v)$. We claim that $d^-_{lex}(w) < d^-_{lex}(v)$. If not, $w \in S_1$. But $d^-_{pr}(w) > d^-_{lex}(w) = d^-_{lex}(v) > d^-_{pr}(v)$, a contradiction to the choice of v. If $d^-_{lex}(w) < d^-_{lex}(v) - 1$, then reversing a directed path from w to v remains strong connectivity by Lemma 1, but the resulting orientation has a smaller lexicographic order of indegree, a contradiction to the choice of Λ_{lex} . Thus $d^-_{lex}(w) = d^-_{lex}(v) - 1$. Now, reversing a directed path from w to v in D_{lex} , we get another orientation with minimum lexicographic order, and however, with $d^-_{lex}(v) > d^-_{pr}(v)$ and $d^-_{lex}(w) < d^-_{pr}(w)$ before the reverse, we get a smaller Δ , which contradicts to the choice of Λ_{lex} , too.

Case 2: $d_{lex}^-(v) < d_{nr}^-(v)$.

Let U be the set of vertices that 2-reach v in D_{pr} . With a similar discussion as in Case 1, we get $\sum_{u \in U} d^-_{lex}(u) \ge \sum_{u \in U} d^-_{pr}(u)$. Since $d^-_{lex}(v) < d^-_{pr}(v)$, there exists a vertex $w \in U$ such that $d^-_{lex}(w) > d^-_{pr}(w)$. Then we have

$$d^-_{pr}(v) > d^-_{lev}(v) \ge d^-_{lex}(w) > d^-_{pr}(w).$$

This implies that $d_{pr}^-(w) < d_{pr}^-(v) - 1$. So there is a strongly reversible directed path in D_{pr} , which contradicts to the property D_{pr} has no strongly reversible directed path.

We conclude that $S = \emptyset$. Therefore the orientation Λ_{pr} found by the SC-PATH-REVERSAL algorithm has the minimum lexicographic order of indegree.

References

- [1] Y. Asahiro, E. Miyano, H. Ono, and K. Zenmyo. Graph orientation algorithms to minimize the maximum outdegree. International Journal of Foundations of Computer Science, 18(2):197-215, 2007.
- [2] G. Borradaile, J. Iglesias, T. Migler, A. Ochoa, G. Wilfong, L. Zhang. Egalitarian Graph Orientations, J. Graph Algorithms Appl., 21(2017): 687-708.
- [3] H. d. Fraysseix and P. O. de Mendez. Regular orientations, arboricity, and augmentation. In Proceedings of the DIMACS International Workshop on Graph Drawing, GD '94, pages 111-118, London, UK, UK, 1995. Springer-Verlag.
- [4] V. Venkateswaran. Minimizing maximum indegree. Discrete Appl. Math., 143:374-378, 2004.

A Proof of Lemma 5

Let C be a component of $G[V \setminus U]$. Note that there is no arc between C and the other components of $G[V \setminus U]$. Since D is strongly connected, there is at least one arc from C to U. Let v_1, \ldots, v_p be the tails of the arcs from C to U and let W_k be the set of vertices in C that reach v_k for $k = 1, \ldots, p$. Note that there is exactly one arc from v_i to U for $i = 1, \ldots, p$, otherwise v_i 2-reaches v by Lemma 4, which is a contradiction to the choice of U. So in the following it is sufficient to show that p = 1. Denote the head vertex of the arc from v_i to U by u_i , $i = 1, \ldots, p$. We first claim that $W_i \cap W_j \neq \emptyset$ for any pair of different W_i and W_j . If not, suppose there is $x \in W_i \cap W_j$, then there exists a directed path P_i from x to v_i and a directed path P_j from x to v_j in C. Let y be the last common vertex in $V(P_i) \cap V(P_j)$ along the direction of P_i . Then $y \in W_i \cap W_j$ and there are two arc-disjoint directed paths from y to v_i and v_j in C and hence from y to v_i and v_j in C, respectively. By Lemma 4, v_j 2-reaches v_j a contradiction to $v_j \notin U$. Thus $v_j \cap W_j \cap$