

On chromatic symmetric homology and planarity of graphs

Azzurra Ciliberti* and Luca Moci†

Abstract

In [4] the authors defined a categorification of Stanley's chromatic function called the chromatic symmetric homology, given by a suitable family of representations of the symmetric group. In this paper we prove that, as conjectured in [2], if a graph G is non-planar, then its chromatic symmetric homology in bidegree $(1,0)$ contains \mathbb{Z}_2 -torsion. Our proof follows a recursive argument based on Kuratowski's theorem.

1 Introduction

The *chromatic symmetric function* of a graph, defined by Stanley in [5], is a remarkable combinatorial invariant which refines the chromatic polynomial. Recently, in [4], Sazdanovic and Yip categorified this invariant by defining a new homological theory, called the *chromatic symmetric homology* of a graph G . This construction, inspired by Khovanov's categorification of the Jones polynomial [1], is obtained by assigning a graded representation of the symmetric group to every subgraph of G , and a differential to every cover relation in the Boolean poset of subgraphs of G . The chromatic symmetric homology is then defined as the homology of this chain complex; its bigraded Frobenius series $Frob_G(q, t)$, when evaluated at $q = t = 1$, reduces to Stanley's chromatic symmetric function expressed in the Schur basis.

As proved in [2], this categorification produces a truly stronger invariant: in other words, chromatic symmetric homology can distinguish couples of graphs that have the same chromatic symmetric function. Furthermore, in the same paper, the properties of chromatic symmetric homology with integer coefficients have been investigated. The authors of [2] provided examples of graphs whose chromatic symmetric homology has torsion, leaving open the following conjecture:

Conjecture 1. *A graph G is non-planar if and only if its chromatic symmetric homology in bidegree $(1,0)$ contains \mathbb{Z}_2 -torsion.*

In this paper we prove one direction of this conjecture, namely:

Theorem. *Let G be a finite non-planar graph. Then its chromatic symmetric homology in bidegree $(1,0)$ contains \mathbb{Z}_2 -torsion.*

*La Sapienza Università di Roma - azzurra.ciliberti@uniroma1.it

†Università di Bologna - luca.moci2@unibo.it

Our strategy will be based on applying Kuratowsky's theorem: we will show that the torsion elements in the homology of the complete graph K_5 and of the complete bipartite graph $K_{3,3}$ are mapped to torsion elements in the homology of the graphs that are obtained from them by the operations of edge subdivision and graph inclusion, i.e. all the non-planar graphs.

2 Computing q -degree zero homology

The interested reader can find a complete description of the construction of chromatic symmetric homology for a graph in the paper [4]. Here we limit ourselves to briefly recall how to compute homology in q -degree zero. For simplifying the notation, we will denote by C_i the i -th chain module and by H_i the i -th homology module. They correspond respectively to $C_{i,0}$ and $H_{i,0}$ in the notation of [4].

Let G be a graph with n vertices and m edges. We can assume without loss of generality that G is simple: indeed if G has a loop, then its chromatic symmetric homology is zero (Proposition 3.1 of [4]), while if G has two vertices connected by multiple edges, then we can replace them by a single edge without affecting the chromatic symmetric homology (Proposition 3.2 of [4]). Hence we denote by (i, j) the edge incident to the vertices i and j , and we order the set of edges $E(G)$ lexicographically.

Each subset S of $E(G)$ is naturally identified with a subgraph of G , having the same vertices as G and S as set of edges. The authors of [2, 3] call it a "spanning subgraph", while we will call it simply a *subgraph*, because the word "spanning" is sometimes used with a different meaning in graph theory and matroid theory. The set of all subgraphs G has a structure of Boolean lattice $\mathbf{B}(G)$, ordered by reverse inclusion. In the Hasse diagram of $\mathbf{B}(G)$, we direct an edge $\epsilon(F, F')$ from a subgraph F to a subgraph F' if and only if F' can be obtained by removing an edge e from F . The sign of ϵ , $\text{sgn}(\epsilon)$, is defined as $(-1)^k$, where k is the number of edges of F less than e .

Let $F \subseteq E(G)$ a subgraph of G with connected components B_1, \dots, B_r . Then the module associated to it in q -degree zero is the *permutation module*

$$M_F = \text{Ind}_{\mathfrak{S}_{B_1} \times \dots \times \mathfrak{S}_{B_r}}^{\mathfrak{S}_n} (\mathbf{S}_{(b_1)} \otimes \dots \otimes \mathbf{S}_{(b_r)}),$$

where \mathfrak{S}_n is the permutation group on n elements and $\mathbf{S}_{(i)}$ is the Specht module related to the partition (i) .

We define

$$C_i(G) = \bigoplus_{|F|=i} M_F,$$

where the sum is over the subgraphs of G with i edges. Therefore the i -th chain module $C_i(G)$ of the graph is a direct sum of $\binom{m}{i}$ permutation modules of \mathfrak{S}_n . If λ is the partition whose parts are the sizes of the connected components of F , then

$$M_F \cong M_\lambda = \mathbb{C}[\mathfrak{S}_n] \otimes_{\mathbb{C}[\mathfrak{S}_\lambda]} \mathbf{S}_{(n)}.$$

Let F and F' be subgraphs of G where $F' = F - e$. There is an *edge map* $d_{\epsilon(F,F')} : M_F \rightarrow M_{F'}$, defined in our case as the inclusion (for the general definition see [4]).

Finally, the i -th chain map $d_i : C_i(G) \rightarrow C_{i-1}(G)$ is defined as

$$d_i = \sum_{\epsilon} \text{sgn}(\epsilon) d_{\epsilon},$$

where the sum is over all the edges ϵ in $\mathbf{B}(G)$ which join a subgraph of G with i edges to a subgraph with $i - 1$ edges. Sometimes we will use the notation d_i^G , where it may not be clear which graph we are referring to.

We need to recall the following definitions from [2].

Definition 2. Let F be a subgraph of G , and let $\lambda \vdash n$ be the partition whose parts are the sizes of the connected components of F . The numbering $T(F)$ associated to F is the numbering of shape λ such that each row consists of the elements in a connected component of F arranged in increasing order, and rows of $T(F)$ having the same size are ordered so that the minimum element in each row is increasing down the first column.

Let $T = T(F)$. The q -degree zero permutation module M_T associated to the numbering T is cyclically generated by the Young symmetrizer

$$a_T = \sum_{\rho \in R(T)} \rho$$

where $R(T) \leq \mathfrak{S}_n$ is the subgroup of permutations that permute elements within each row of T . We have:

$$M_F \cong M_T = \mathbb{C}[\mathfrak{S}_n] \cdot a_T.$$

Definition 3. For any numberings S and T of shape λ , let

$$v_T^S = \sigma_{T,S} b_T a_T = b_S a_S \sigma_{T,S} \in M_T,$$

where a_T is as above,

$$b_T = \sum_{\zeta \in C(T)} \text{sgn}(\zeta) \zeta,$$

$C(T) \leq \mathfrak{S}_n$ is the subgroup of permutations that permute elements within each column of T , and $\sigma_{T,S} \in \mathfrak{S}_n$ is such that $\sigma_{T,S} \cdot T = S$.

Moreover, the q -degree zero Specht module \mathbf{S}_T associated to the numbering T is cyclically generated by the Young symmetrizer $c_T = b_T a_T$:

$$\mathbf{S}_T = \mathbb{C}[\mathfrak{S}_n] \cdot c_T,$$

and $\mathbf{S}_T \cong \mathbf{S}_\lambda$.

We also recall the following result from [3], Section 7.2, Proposition 2.

Proposition 4. Let F be a subgraph of G with associated numbering T of shape λ . Then

$$\mathbf{S}_T = \text{span}\{b_S a_S \sigma_{T,S} | S \in \text{SYT}(\lambda)\} = \text{span}\{v_T^S | S \in \text{SYT}(\lambda)\},$$

where $\text{SYT}(\lambda)$ is the set of standard Young tableaux of shape λ .

2.1 Computation of $H_1(G)$

We will describe the chromatic homology in terms of Specht modules. Since each Specht module is cyclically generated, then our inclusion maps are completely determined by specifying the image of a cyclic generator for each Specht module. We now show how to achieve these computations systematically.

We restrict ourselves to Specht modules of type $\lambda = (2^k, 1^{n-2k})$ for $k \geq 1$, so we will be computing

$$C_2(G)|_{S_\lambda} \xrightarrow{d_2} C_1(G)|_{S_\lambda} \xrightarrow{d_1} C_0(G)|_{S_\lambda} \rightarrow 0.$$

We order the edges of G in lexicographic order and label these as e_1, \dots, e_m . In homological degree zero, there is only one subgraph without edges. The chain group $C_0(G) = M_{F_\emptyset} \cong M_{(1^n)}$ is the regular representation of \mathfrak{S}_n , where F_\emptyset is the edgeless subgraph. By Corollary 1 in Section 7.3 of [3], the multiplicity of S_λ in $\mathbb{C}[S_n]$ is the number $f^\lambda = K_{\lambda, (1^n)}$ of standard Young tableaux of shape λ . We list the tableaux $Y_1(G), \dots, Y_{f^\lambda}(G) \in SYT(\lambda)$ with respect to the following total order: if T and S are numberings of shape λ such that the i -th row is the lowest row in which the numberings are different, the j -th column is the rightmost column in that row in which the numberings are different and $T(i, j) > S(i, j)$, then we say that $T > S$.

We have

$$C_0(G)|_{S_\lambda} = \bigoplus_{i=1}^{f^\lambda} \mathbb{C}[\mathfrak{S}_n] \cdot v_{Y_i}^{Y_1}.$$

Example 5. Let $G = K_5$. We order the edges of G in lexicographic order; that is,

$$(1, 2), (1, 3), (1, 4), (1, 5), (2, 3), (2, 4), (2, 5), (3, 4), (3, 5), (4, 5),$$

and label these as e_1, \dots, e_{10} . The standard Young tableaux of shape $\lambda = (2^2, 1)$ listed with respect to the ordering defined earlier are

$$Y_1 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & \\ \hline \end{array} Y_2 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & \\ \hline \end{array} Y_3 = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline 5 & \\ \hline \end{array} Y_4 = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 5 \\ \hline 4 & \\ \hline \end{array} Y_5 = \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & \\ \hline \end{array}$$

Then

$$C_0(K_5)|_{S_{(2^2, 1)}} = \bigoplus_{i=1}^5 (\mathbb{C}[S_5] \cdot v_{Y_i}^{Y_1}) \cong \mathbf{s}_{(2^2, 1)}^{\oplus 5}.$$

In homological degree one, there are m subgraphs with exactly one edge, thus

$$C_1(G) = \bigoplus_{i=1}^m M_{F_{e_i}}.$$

If F_{e_i} is the subgraph containing the edge $e_i = (p, q)$, then the permutation module $M_{F_{e_i}}$ has the associated numbering $T(F_{e_i})$ of shape $\mu = (2, 1^{n-2})$, and $M_{F_{e_i}} = \mathbb{C}[\mathfrak{S}_n] \cdot (e + (pq)) \cong M_\mu$.

The multiplicity of S_λ in $M_{F_{e_i}}$ is the number $K_{\lambda, \mu}$ of semistandard Young tableaux of shape λ and weight μ . We next obtain numberings of shape λ that will index these $K_{\lambda, \mu}$ Specht modules \mathbf{S}_λ , by standardizing the set $SSYT(\lambda, \mu)$ of semistandard Young tableaux of shape λ and weight μ with respect to $T(F_{e_i})$ in the following way. For any numbering S , the word $w(S)$ of S is obtained by reading the entries of the rows of S from left to right, and from the top row to the bottom row (note that this is not the usual definition of a reading word for tableaux). So, given $Y \in SSYT(\lambda, \mu)$, let $w(Y) = y_1, \dots, y_n$ be the word of Y , let $w(T) = t_1, \dots, t_n$ be the word of $T = T(F_{e_i})$ and let σ be the permutation that orders y_1, \dots, y_n without exchanging y_i and y_j if $y_i = y_j$. From this we obtain a numbering X of shape λ by replacing the entry in Y that corresponds to y_k by $t_{\sigma(k)}$. We list the numberings $X_i^1(G), \dots, X_i^{K_{\lambda, \mu}}(G)$ obtained using the procedure just described to $SSYT(\lambda, \mu)$ with respect to $T(F_{e_i})$. Observe that since $\mu = (2, 1^{n-2})$ and $\lambda = (2^k, 1^{n-2k})$ where $k \geq 1$, then this procedure guarantees that the first row of each numbering $X_i^j(G)$ is $\boxed{p} \boxed{q}$. So $v_{X_i^j(G)}^{Y_1} \in M_{F_{e_i}}$ and $\mathbb{C}[\mathfrak{S}_n] \cdot v_{X_i^j(G)}^{Y_1} \cong \mathbf{S}_\lambda$ for $j = 1, \dots, K_{\lambda, \mu}$. Thus

$$C_1(G)|_{\mathbf{S}_\lambda} = \bigoplus_{i=1}^m \bigoplus_{j=1}^{K_{\lambda, \mu}} \mathbb{C}[S_n] \cdot v_{X_i^j(G)}^{Y_1}$$

Example 6. Let $G = K_5$. There are 10 subgraphs with exactly one edge. Furthermore, there are two semistandard Young tableaux of shape $\lambda = (2^2, 1)$ and weight $(2, 1^3)$,

$$Z_1 = \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 3 \\ \hline 4 & \\ \hline \end{array} \quad \text{and} \quad Z_2 = \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 4 \\ \hline 3 & \\ \hline \end{array},$$

so the multiplicity of $\mathbf{S}_{(2^2, 1)}$ in each $M_{F_{e_i}} \cong M_{(2, 1^3)}$ is 2. Let $X_i^1(K_5)$ and $X_i^2(K_5)$ denote the numberings which index the two copies of $\mathbf{S}_{(2^2, 1)}$ in each $M_{F_{e_i}}$, again listed with respect to the same ordering. So

$$C_1(K_5)|_{\mathbf{S}_{(2^2, 1)}} = \bigoplus_{i=1}^{10} (\mathbb{C}[S_5] \cdot v_{X_i^1(K_5)}^{Y_1} \oplus \mathbb{C}[S_5] \cdot v_{X_i^2(K_5)}^{Y_1}) \cong \mathbf{S}_{(2^2, 1)}^{\oplus 20}.$$

Consider for instance the subgraph F_{e_1} of K_5 with the edge $e_1 = (1, 2)$ only. The numbering associated to it is

$$T(F_{e_1}) = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline 4 & \\ \hline 5 & \\ \hline \end{array}$$

Let z_1^i, \dots, z_5^i be the word of Z_i , $i = 1, 2$. The permutation that orders z_1^1, \dots, z_5^1 is the identity and the one that orders z_1^2, \dots, z_5^2 is (45). Therefore we have

$$X_1^1 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & \\ \hline \end{array} = Y_1 \quad \text{and} \quad X_1^2 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & \\ \hline \end{array} = Y_2,$$

then

$$v_{X_1^1}^{Y_1} = v_{Y_1}^{Y_1} \quad \text{and} \quad v_{X_1^2}^{Y_1} = v_{Y_2}^{Y_1}.$$

Lastly, we consider the chain module in homological degree two. The subgraphs of G with exactly two edges have connected components of partition type $(2^2, 1^{n-4})$ or $(3, 1^{n-3})$. We are only concerned with Specht modules of type $\lambda = (2^k, 1^{n-2k})$ with $k \geq 2$ necessarily and, since $\lambda \not\vdash (3, 1^{n-3})$, then, by Corollary 1 in Section 7 of [3], S_λ does not appear as a summand in a permutation module isomorphic to $M_{(3, 1^{n-3})}$. Hence, we only need to consider the subgraphs with connected components of partition type $(2^2, 1^{n-4})$.

So suppose G has h subgraphs whose connected components has partition type $\nu = (2^2, 1^{n-4})$. List these subgraphs with respect to the lexicographic order of their edge sets. Suppose F_{e_i, e_j} is the subgraph that contains the edges $e_i = (p, q)$ and $e_j = (r, s)$ with $p < r$. The permutation module $M_{F_{e_i, e_j}}$ has the associated numbering $T(F_{e_i, e_j})$ of shape ν and

$$M_{F_{e_i, e_j}} = \mathbb{C}[S_n] \cdot (e + (pq))(e + (rs)) \cong M_\nu.$$

Similar to the previous case for $C_1(G)$, the multiplicity of S_λ in $M_{F_{e_i, e_j}}$ is $K_{\lambda, \nu}$. We list the numberings $W_{i,j}^1(G), \dots, W_{i,j}^{K_{\lambda, \nu}}(G)$ obtained using the procedure described above to $SSYT(\lambda, \nu)$ with respect to $T(F_{e_i, e_j})$. The procedure guarantees that the top two rows of each numbering W are $\begin{array}{|c|c|} \hline p & q \\ \hline r & s \\ \hline \end{array}$, so $v_W^{Y_1} \in M_{F_{e_i, e_j}}$ and

$$\mathbb{C}[S_n] \cdot v_{W_{i,j}^l(G)}^{Y_1} \cong S_\lambda \text{ for } l = 1, \dots, K_{\lambda, \nu}.$$

Thus

$$C_2(G)|_{S_\lambda} = \bigoplus_{i,j} \bigoplus_{l=1}^{K_{\lambda, \nu}} \mathbb{C}[S_n] \cdot v_{W_{i,j}^l(G)}^{Y_1},$$

where the direct sum is over the values of i and j corresponding to the couples of edges which form subgraphs of type ν , i.e. the non-consecutive edges.

Example 7. Let $G = K_5$. There are 15 subgraphs whose connected components have partition type $(2^2, 1)$. There is only one semistandard Young tableau of shape $\lambda = (2^2, 1)$ and weight $(2^2, 1)$:

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 2 \\ \hline 3 & \\ \hline \end{array}$$

so the multiplicity of $\mathbf{S}_{(2^2,1)}$ in each $M_{F_{e_i e_j}} \cong M_{(2^2,1)}$ is 1, and we let $W_{i,j}$ denote the numbering which indexes the copy of $\mathbf{S}_{(2^2,1)}$ in $M_{F_{e_i e_j}}$. Therefore,

$$C_2(K_5)|_{\mathbf{S}_{(2^2,1)}} = \bigoplus (\mathbb{C}[S_5] \cdot v_{W_{i,j}}^{Y_1}) \cong \mathbf{S}_{(2^2,1)}^{\oplus 15},$$

where the direct sum is over the values of i and j corresponding to the couples of non-consecutive edges.

To compute the edge maps we will need the following theorem (Corollary 2.18 of [2]).

Theorem 8. *For any numberings S of shape λ ,*

$$v_S^T = (-1)^j \sum_{U \in \Xi_{i,j}(S)} v_U^T,$$

where $\Xi_{i,j}(S)$ is the set of all numberings U obtained from S by exchanging the first j entries in the $(i+1)$ -th row of S with j entries in the i -th row of S , preserving the order of each subset of elements.

We let $\pi_{i,j}$ denote the operator on numberings such that

$$\pi_{i,j}(S) = (-1)^j \sum_{U \in \Xi_{i,j}(S)} U.$$

Example 9. Let $G = K_5$. We compute $d_1(v_{X_3^1}^{Y_1})$ and $d_1(v_{X_3^2}^{Y_1})$, so we consider the subgraph F_{e_3} of K_5 with the edge $e_3 = (1, 4)$ only. The numbering associated to it is

$$T(F_{e_3}) = \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & \\ \hline 3 & \\ \hline 5 & \\ \hline \end{array}$$

We have

$$X_3^1 = \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 3 \\ \hline 5 & \\ \hline \end{array} \quad \text{and} \quad X_3^2 = \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & \\ \hline \end{array} = Y_5.$$

We compute

$$\pi_{1,1}(X_3^1) = - \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 4 & 3 \\ \hline 5 & \\ \hline \end{array} - \begin{array}{|c|c|} \hline 2 & 4 \\ \hline 1 & 3 \\ \hline 5 & \\ \hline \end{array}$$

and

$$\pi_{1,2} \begin{bmatrix} 2 & 4 \\ 1 & 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 5 \end{bmatrix} = Y_3.$$

Then, by Theorem 8, we have

$$v_{X_3^1}^{Y_1} = -v_{Y_1}^{Y_1} - v_{Y_3}^{Y_1} \quad \text{and} \quad v_{X_3^2}^{Y_1} = v_{Y_5}^{Y_1},$$

and d_1 sends

$$v_{X_3^1}^{Y_1} \mapsto -v_{Y_1}^{Y_1} - v_{Y_3}^{Y_1} \quad \text{and} \quad v_{X_3^2}^{Y_1} \mapsto v_{Y_5}^{Y_1}.$$

Now we compute $d_2(v_{W_{1,8}}^{Y_1})$, so we have to consider the subgraph F_{e_1, e_8} of K_5 with the edges $e_1 = (1, 2)$ and $e_8 = (3, 4)$. The numbering associated to it is

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 \end{bmatrix}.$$

There are two subgraphs of K_5 with only one edge that can be obtained removing one edge from F_{e_1, e_8} , i.e. F_{e_1} and F_{e_8} . The per-edge map $d_{\epsilon(F_{e_1, e_8}, F_{e_1})}$ appears as a summand in d_2 with a minus sign; instead $d_{\epsilon(F_{e_1, e_8}, F_{e_8})}$ appears in d_2 with a plus sign. Therefore, d_2 sends

$$v_{W_{1,8}}^{Y_1} \mapsto -v_{X_1^1}^{Y_1} + v_{X_8^1}^{Y_1}.$$

3 The case of non-planar graphs

In this section we will prove that if G is a non-planar graph, then the chromatic symmetric homology $H_1(G; \mathbb{Z})$ contains \mathbb{Z}_2 -torsion. We first recall two results from [2]:

Lemma 10. *The chromatic symmetric homology $H_1(K_5; \mathbb{Z})$ contains \mathbb{Z}_2 -torsion.*

Proof. We compute

$$C_2(K_5)|_{\mathbf{S}_{(2^2,1)}} \xrightarrow{d_2} C_1(K_5)|_{\mathbf{S}_{(2^2,1)}} \xrightarrow{d_1} C_0(K_5)|_{\mathbf{S}_{(2^2,1)}} \rightarrow 0,$$

restricted to the $\mathbf{S}_{(2^2,1)}$ modules.

Following the notation introduced in 2.1, let $g = W_{1,8} + W_{1,9} + W_{1,10} + W_{2,6} - W_{2,7} - W_{2,10} + W_{3,5} + W_{3,7} + W_{3,9} + W_{4,5} + W_{4,6} + W_{4,8} - W_{5,10} - W_{6,9} + W_{7,8} \in C_2(K_5)$ and $h = X_9^1 + X_{10}^1 - X_2^1 + X_7^2 + X_9^2 \in C_1(K_5)$. We have that $h \notin \text{im } d_2$, $d_2(g) = 2h$ and $d_1(h) = 0$, so h generates \mathbb{Z}_2 -torsion in $H_1(K_5; \mathbb{Z})$. For more details see [2], Theorem 4.1. \square

Lemma 11. *The chromatic symmetric homology $H_1(K_{3,3}; \mathbb{Z})$ contains \mathbb{Z}_2 -torsion.*

Proof. We compute

$$C_2(K_{3,3})|_{\mathbf{S}_{(2^2,1^2)}} \xrightarrow{d_2} C_1(K_{3,3})|_{\mathbf{S}_{(2^2,1^2)}} \xrightarrow{d_1} C_0(K_{3,3})|_{\mathbf{S}_{(2^2,1^2)}} \rightarrow 0,$$

restricted to the $\mathbf{S}_{(2^2,1^2)}$ modules.

Following the notation introduced in Section 2.1, let $g' = W_{1,6} - W_{1,7} + W_{1,8} + W_{1,9} - W_{2,4} - W_{2,5} + W_{2,7} + W_{2,9} + W_{3,4} - W_{3,5} + W_{3,6} + W_{3,8} + W_{4,8} + W_{4,9} + W_{5,6} + W_{5,7} - W_{6,9} + W_{7,8} \in C_2(K_{3,3})$ and $h' = X_6^3 - X_7^3 + X_8^3 - X_9^2 \in C_1(K_{3,3})$. We have that $h' \notin \text{im } d_2$, $d_2(g') = 2h'$ and $d_1(h') = 0$, so h' generates \mathbb{Z}_2 -torsion in $H_1(K_{3,3}; \mathbb{Z})$. For more details see [2], Theorem 4.2. \square

Proposition 12. *Let G be a graph with n vertices, $n \geq 2$, λ a partition of n of type $(2^k, 1^{n-2k})$ and $h \in C_1(G)|_{\mathbf{S}_\lambda}$ a generator of \mathbb{Z}_p -torsion in $H_1(G; \mathbb{Z})$. Let G' be a subdivision of G with n' vertices, $n' > n$, i.e. a graph obtained from G by inserting $n' - n$ vertices into the edges of G . Then there exists $h' \in C_1(G')|_{\mathbf{S}_{\lambda'}}$, with $\lambda' = (2^k, 1^{n'-2k})$, that generates \mathbb{Z}_p -torsion in $H_1(G'; \mathbb{Z})$.*

Proof. it is enough to show that the statement holds for $n' = n + 1$.

Let G be a graph with n vertices and G' be the graph obtained from G by inserting a vertex into an edge of G . By hypothesis, there exists $h \in C_1(G)|_{\mathbf{S}_\lambda}$, with $\lambda = (2^k, 1^{n-2k})$, that generates \mathbb{Z}_p -torsion in $H_1(G; \mathbb{Z})$.

We number the vertices of G' from 1 to $n + 1$, so that the vertex added is $n + 1$.

We prove that h is mapped in a \mathbb{Z}_p -torsion generator in $H_1(G'; \mathbb{Z})$.

We have that all the edges of G are also edges of G' , except for the edge that has been broken, let it be (i, j) , which is no longer an edge of G' , but it has been replaced by two edges, $(i, n + 1)$ and $(n + 1, j)$. We consider the partition $\lambda' = (2^k, 1^{n+1-2k})$, i.e. the partition λ with an extra box at the bottom. We divide the standard Young tableaux of shape λ' into two groups: those that are obtained simply by adding the box containing $n + 1$ at the bottom of the standard Young tableaux of shape λ and those that don't have $n + 1$ in the last row. We do the same for the semistandard Young tableaux of shape λ' and weight $(2, 1^{n-1})$. If i does not correspond to the two new edges, the $X_i^l(G')$'s are identical to the $X_i^l(G)$'s with the box containing $n + 1$ added at the bottom. Therefore, the differential $d_1^{G'}$ acts on them in exactly the same way as d_1^G , since the $\pi_{i,j}$ operations don't concern the last row. Since $h \in \ker d_1^G$, we also have that $h \in \ker d_1^{G'}$.

We mean h written as 1-chain in G' . This can always be done using the $\pi_{i,j}$ operations and Theorem 8.

For example, consider K_5 and the graph G_1 obtained by adding the vertex 6 into the edge $(1, 5)$. $X_4^1(K_5)$ with a box containing 6 added at the bottom, i.e.

1	5
2	3
4	
6	

a priori is not a 1-chain in G_1 since the edge $(1, 5)$ is not an edge of G_1 ; but

we have

$$\begin{array}{|c|c|} \hline 1 & 5 \\ \hline 2 & 3 \\ \hline 4 & \\ \hline 6 & \\ \hline \end{array} = \pi_{1,1} \begin{array}{|c|c|} \hline 1 & 5 \\ \hline 2 & 3 \\ \hline 4 & \\ \hline 6 & \\ \hline \end{array} = - \begin{array}{|c|c|} \hline 2 & 5 \\ \hline 1 & 3 \\ \hline 4 & \\ \hline 6 & \\ \hline \end{array} - \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & \\ \hline 6 & \\ \hline \end{array} = -X_2^2(G_1) - X_1^2(G_1) \in C_1(G_1).$$

Remark 13. It cannot happen that there is cancellation among 1-cycles in G' that give the cycle h in G . In fact, the $\pi_{i,j}$ operations on numberings correspond, by Theorem 8, to equalities among the elements of $\mathbf{S}_{(2^k, 1^{n-2k})}$ indexed by such numberings. Therefore, if there was cancellation among 1-cycles in G' that give h in G , it would also be among the 1-cycles of G that form h , so h would be the trivial 1-cycle of G , which is not true.

A similar argument applies to the $W_{h,k}^s$'s. Since, in G , there exists a 2-cycle g such that $d_2^G(g) = 2h$ and g can be written as 2-chain in G' with the $\pi_{i,j}$ operations, without becoming trivial as observed for h in Remark 13, we also have that $d_2^{G'}(g) = 2h$.

It remains to prove that $h \notin \text{im } d_2^{G'}$. If h belonged to $\text{im } d_2^{G'}$, since we know that $h \notin \text{im } d_2^G$, it would be linear combination of the columns of $d_2^{G'}$ which are not in d_2^G , but it is not possible because h is a 1-chain in G .

Therefore, we have a \mathbb{Z}_p -torsion generator in $H_1(G', \mathbb{Z})$. \square

Proposition 14. *Let G' be a graph with n' vertices and let G be a subgraph of G' with n vertices, $n \leq n'$. Assume that $h \in C_1(G)_{|\mathbf{S}_\lambda}$ is a generator of \mathbb{Z}_p -torsion in $H_1(G; \mathbb{Z})$. Then there exists $h' \in C_1(G')_{|\mathbf{S}_{\lambda'}}$, with $\lambda' = (2^k, 1^{n'-2k})$, that generates \mathbb{Z}_p -torsion in $H_1(G', \mathbb{Z})$.*

Proof. We have that all the edges of G are also edges of G' . We consider the partition $\lambda' = (2^k, 1^{n'-2k})$, i.e. the partition λ with $n' - n$ extra boxes at the bottom. We number the vertices of G' from 1 to n' , so that the vertices eventually added are $n+1, \dots, n'$. We divide the standard Young tableaux of shape λ' into two groups: those that are obtained simply by adding the boxes containing $n+1, \dots, n'$ at the bottom of the standard Young tableaux of shape λ and those that don't have $n+1, \dots, n'$ in the last rows. We do the same for the semistandard Young tableaux of shape λ' and weight $(2, 1^{n'-2})$. All the $X_i^l(G)$'s with extra boxes containing $n+1, \dots, n'$ at the bottom are among the $X_i^{l'}(G')$'s. Therefore, the differential $d_1^{G'}$ acts on them in exactly the same way as d_1^G , since the $\pi_{i,j}$ operations don't concern the last rows. Since $h \in \ker d_1^G$, we also have that $h \in \ker d_1^{G'}$.

A similar argument applies to the $W_{h,k}^s$'s. Since, in G , there exists a 2-cycle g such that $d_2^G(g) = 2h$, we also have that $d_2^{G'}(g) = 2h$.

It remains to prove that $h \notin \text{im } d_2^{G'}$. If h belonged to $\text{im } d_2^{G'}$, since we know that $h \notin \text{im } d_2^G$, it would be linear combination of the columns of $d_2^{G'}$ which are not in d_2^G , but it is not possible because h is a 1-chain in G .

Therefore, we have a \mathbb{Z}_p -torsion generator in $H_1(G', \mathbb{Z})$. \square

Theorem 15. *Let G be a finite non-planar graph. Then $H_1(G; \mathbb{Z})$ contains \mathbb{Z}_2 -torsion.*

Proof. Since G is non-planar, by Kuratowsky's theorem it contains a subgraph G' which is a subdivision of K_5 or $K_{3,3}$. By Lemma 10 and Lemma 11, both $H_1(K_5; \mathbb{Z})$ and $H_1(K_{3,3}; \mathbb{Z})$ have a generator of \mathbb{Z}_2 -torsion of type $(2^k, 1^{n-2k})$. Hence, by Proposition 12, also $H_1(G; \mathbb{Z})$ contains such a \mathbb{Z}_2 -torsion element; thus by Proposition 14 also $H_1(G; \mathbb{Z})$ does. \square

References

- [1] Dror Bar-Natan. "On Khovanov's categorification of the Jones polynomial". In: *Algebr. Geom. Topol.* 2 (2002), pp. 337–370. ISSN: 1472-2747. DOI: 10.2140/agt.2002.2.337. URL: <https://doi.org/10.2140/agt.2002.2.337>.
- [2] Alex Chandler, Radmila Sazdanovic, Salvatore Stella, and Martha Yip. "On the strength of chromatic symmetric homology for graphs". In: *ArXiv: 1911.13297v1* (2019).
- [3] William Fulton. *Young tableaux*. Vol. 35. London Mathematical Society Student Texts. With applications to representation theory and geometry. Cambridge University Press, Cambridge, 1997, pp. x+260. ISBN: 0-521-56144-2; 0-521-56724-6.
- [4] Radmila Sazdanovic and Martha Yip. "A categorification of the chromatic symmetric function". In: *J. Combin. Theory Ser. A* 154 (2018), pp. 218–246. ISSN: 0097-3165. DOI: 10.1016/j.jcta.2017.08.014. URL: <https://doi.org/10.1016/j.jcta.2017.08>.
- [5] Richard P. Stanley. "A symmetric function generalization of the chromatic polynomial of a graph". In: *Adv. Math.* 111.1 (1995), pp. 166–194. ISSN: 0001-8708. DOI: 10.1006/aima.1995.1020. URL: <https://doi.org/10.1006/aima.1995.1020>.

This figure "Screenshot__308_.png" is available in "png" format from:

<http://arxiv.org/ps/2103.01543v4>

This figure "Screenshot__309_.png" is available in "png" format from:

<http://arxiv.org/ps/2103.01543v4>

This figure "Screenshot__310_.png" is available in "png" format from:

<http://arxiv.org/ps/2103.01543v4>

This figure "Screenshot__311_.png" is available in "png" format from:

<http://arxiv.org/ps/2103.01543v4>