

COLD TRAPS OF HYPERVOLATILES IN THE PROTOSOLAR NEBULA AT THE ORIGIN OF COMET C/2016 R2 (PANSTARRS)'S PECULIAR COMPOSITION

OLIVIER MOUSIS¹, ARTYOM AGUICHINE¹, ALEXIS BOUQUET², JONATHAN I. LUNINE³, GRÉGOIRE DANGER², KATHLEEN E. MANDT⁴, AND ADRIENN LUSPAY-KUTI⁴

Draft version March 3, 2021

ABSTRACT

Recent observations of the long period comet C/2016 R2 (PanSTARRS) indicate an unusually high N₂/CO abundance ratio, typically larger than ~ 0.05 , and at least 2–3 times higher than the one measured in 67P/Churyumov-Gerasimenko. Another striking compositional feature of this comet is its heavy depletion in H₂O (H₂O/CO $\sim 0.32\%$), compared to other comets. Here, we investigate the formation circumstances of a generic comet whose composition reproduces these two key features. We first envisage the possibility that this comet agglomerated from clathrates, but we find that such a scenario does not explain the observed low water abundance. We then alternatively investigate the possibility that the building blocks of the comet C/2016 R2 (PanSTARRS) agglomerated from grains and pebbles made of pure condensates via the use of a disk model describing the radial transport of volatiles. We show that N₂/CO ratios reproducing the value estimated in this comet can be found in grains condensed in the vicinity of the CO and N₂ icelines. Moreover, high CO/H₂O ratios (> 100 times the initial gas phase value) can be found in grains condensed in the vicinity of the CO iceline. If the building blocks of a comet assembled from such grains, they should present N₂/CO and CO/H₂O ratios consistent with the measurements made in comet C/2016 R2 (PanSTARRS)'s coma. Our scenario indicates that comet C/2016 R2 (PanSTARRS) formed in a colder environment than the other comets that share more usual compositions. Our model also explains the unusual composition of the interstellar comet 2I/Borisov.

Subject headings: astrobiology – comets: general – comets: individual (C/2016 R2 (PanSTARRS)) – methods: numerical – solid state: volatile

1. INTRODUCTION

Comets are supposed to be water-rich bodies that are relics of the formation of the solar system. Because they have undergone little alteration during the 4.6 billion years of the solar system evolution, investigating their composition provides indications of the physico-chemical conditions that were at play during their formation in the protosolar nebula (PSN). An outstanding question remaining to be solved is the origin of the apparent N₂ deficiency observed in comets while both Pluto and Triton, also formed in the outer solar system, harbor N₂-rich surfaces and atmospheres (Lellouch et al. 2011; Cruikshank et al. 2015; Mandt et al. 2017). Decades of remote-sensing observations of comets suggest they are depleted in N₂ (Cochran et al. 2000). Only upper limits of $\sim 10^{-4}$ for N₂⁺/CO⁺ were derived in the coma of comets 122P/1995 S1 (deVico) and C/1995 O1 (Hale-Bopp) (Cochran et al. 2000) while very few detections of N₂⁺ emission lines have been reported in other comets from ground-based facilities. This apparent N₂ depletion was interpreted as the result of the selective trapping of CO at the expense of N₂ in the building blocks of comets presumably agglomerated from clathrates (Iro et al. 2003), or as the result of their par-

tial devolatilization due to radiogenic heating (Mousis et al. 2012). To a lesser extent, these important depletions are in agreement with the measurements of the N₂/CO ratio acquired in comet 67P/Churyumov-Gerasimenko (hereafter 67P/C-G) by the ROSINA mass spectrometer aboard the Rosetta spacecraft very early in the mission in October 2014, which suggest a value of $(5.70 \pm 0.66) \times 10^{-3}$ that is depleted by a factor of ~ 25 as compared to the one derived from protosolar N and C abundances (Rubin et al. 2015). However, this N₂/CO ratio was measured at a heliocentric distance beyond 3 au, far beyond perihelion. Rubin et al. (2020) later derived a N₂/CO ratio of $\sim 2.87 \times 10^{-2}$ in 67P/C-G, a value obtained in May 2015 a few months before perihelion passage, and roughly 5 times larger than the previous one.

Contrasting with most of these previous measurements, recent optical spectra of the long period comet C/2016 R2 (PanSTARRS) (hereafter R2) performed at a heliocentric distance of ~ 3 AU, showed that its spectrum was largely dominated by the emission band of CO⁺, but also surprisingly by the presence of the emission band of N₂⁺ (Cochran & McKay 2018). Subsequent observations confirmed an unusually high N₂/CO ratio in R2, a value estimated to range between 0.06 ± 0.01 (Opitom et al. 2019) and 0.08 (Biver et al. 2018), and at least 2–3 times higher than the one measured closer to perihelion in 67P/C-G. Another striking compositional feature of this comet is its heavy depletion in H₂O (H₂O/CO $\sim 0.32\%$; McKay et al. (2019)). These trends are at odds with our understanding of the thermodynamic evolution of cometary nuclei which should be, in principle, depleted

¹ Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France
olivier.mousis@lam.fr

² Aix Marseille Univ, CNRS, PIIM, F-13013 Marseille, France

³ Department of Astronomy, Cornell University, Ithaca, NY 14853, USA

⁴ Applied Physics Laboratory, Johns Hopkins University, 11100 Johns Hopkins Rd., Laurel, MD 20723, USA




FIG. 1.— Comparison of the volatile composition of R2 to the average comet as derived by McKay et al. (2019) [Reprinted from *The Astronomical Journal*, 158:128 (24pp), McKay, A., et al., *The Peculiar Volatile Composition of CO-dominated Comet C/2016 R2* (PanSTARRS), with permission from IOP].

in the most volatile species instead of being enhanced, as R2 shows, after multiple solar passages. Since sublimation of supervolatile ices can occur possibly beyond 40 AU for CO (Womack et al. 2017), one should expect CO outgassing, and then cometary activity over a significant fraction of R2’s orbit. Figure 1 represents a comparison of the volatile composition of R2 to the average comet made by McKay et al. (2019). The figure shows that all oxygen appears to be locked in CO and CO₂, and not H₂O, as is the case for typical comets.

In this paper, we aim at investigating the formation circumstances of a generic comet whose composition reproduces the two key features of R2’s coma, namely its high N₂/CO ratio and the observed significant water depletion, assuming they reflect its primordial composition (McKay et al. 2019). We first investigate the possibility that this comet could have agglomerated from clathrates as it has been suggested in the case of 67P/C-G (Luspay-Kuti et al. 2016; Mousis et al. 2016, 2018), but we find that such a scenario cannot explain the observed low water abundance. We then alternatively investigate the possibility that the building blocks of this comet agglomerated from grains and pebbles made of pure condensates crystallized in the vicinity of the CO iceline in the PSN. This scenario matches both observed features and suggests that this comet formed in a colder environment than the other comets sharing usual compositions.

2. AGGLOMERATION FROM CLATHRATES?

For decades, the agglomeration of comets from clathrates has been regularly invoked to explain their compositional and thermodynamic properties (Lunine & Stevenson 1985; Klinger et al. 1986; Iro et al. 2003; Marboeuf et al. 2010, 2011, 2012). Recent measurements of 67P/C-G’s composition by the Rosina mass spectrometer aboard the *Rosetta* mission have also been interpreted in favor of the presence of clathrates in its interior (Luspay-Kuti et al. 2016; Mousis et al. 2016, 2018). This possibility is now investigated in the case of R2, which presents unusually high CO and N₂ abundances in its coma, compared to typical comets. To do so, we aim at comparing the plausible composition of multiple guest (MG) clathrates formed in the PSN to that of R2, assuming this latter would have agglomerated from these icy structures. These MG clathrates are assumed to be formed from a

gaseous mixture composed of N₂ and CO, assuming these species are the two main C- and N-bearing volatiles in the gas phase of the disk, a hypothesis consistent with the thermochemical models (Lewis & Prinn 1980; Prinn & Fegley 1989; Mousis et al. 2002). Because CO has a much higher propensity for clathration than N₂, the calculations utilized in this work are those performed in the framework of structure I clathrates, namely the structure predicted for a CO-dominated clathrate (Mohammadi et al. 2005). Given the fact that half of protosolar carbon is expected to be used in organic compounds in the PSN (Pollack et al. 1994), only the remaining half is assumed to be in the form of CO. Assuming that 90% of protosolar nitrogen is in N₂ form (Lewis & Prinn 1980), we derive a N₂/CO ratio of 2.66×10^{-1} in the initial gas phase of the PSN, based on the protosolar abundances given by Lodders et al. (2009). In our model, clathrates form in the PSN as long as crystalline water is available. Here we assume that once all the water budget has been used for clathration, the remaining volatiles are not incorporated into solids.

Our model assumes the formation of a multiple guest (MG) clathrate with an equilibrium pressure expressed as (Lunine & Stevenson 1985):

$$P_{eq,MG} = \left[\sum_i \frac{y_i}{P_{eq,i}} \right]^{-1}, \quad (1)$$

where y_i is the mole fraction of the component i in the fluid phase. The equilibrium pressure curves of each species are determined by fitting the available theoretical and laboratory data (Lunine & Stevenson 1985) with equations of the form $\log P_{eq,i} = A/T + B$, where $P_{eq,i}$ and T are the partial equilibrium pressure and temperature of the considered species i , respectively. The relative abundances of guest species incorporated in MG clathrate formed at a given temperature and pressure from the PSN gas phase are calculated following the method described in Lunine & Stevenson (1985) and Mousis et al. (2010). This formalism has been used to interpret 67P/C-G’s ice structure and composition from Rosetta/ROSINA observations of its coma (Mousis et al. 2016, 2018).

Figure 2 shows the influence of the abundance of crystalline water ice in the outer PSN on the N₂/CO ratio in cometary grains at clathration temperatures of 25, 30, 40, and 50 K. These temperatures are all above the formation temperature of N₂ and CO pure condensates. For water abundances below $1.3 \times (O/H_2)_\odot$, the N₂/CO ratios in icy grains correspond to those calculated in clathrates at specified formation temperatures. For example, the N₂/CO ratios measured in Hale-Bopp, deVico and 67P/C-G are closely matched at 25, 30, and 50 K, respectively. At higher water abundances, CO is fully entrapped and significant amounts of N₂ become progressively trapped in clathrates, implying an increase of the global N₂/CO in icy grains. A water abundance of $\sim 1.4 \times (O/H_2)_\odot$ in the PSN allows one to retrieve a N₂/CO ratio in clathrates corresponding to the value measured in R2. Water abundances greater than $\sim 1.7 \times (O/H_2)_\odot$ allows the full trapping of the available N₂ in clathrates crystallized in the PSN, leading to a N₂/CO ratio of $\sim 2.66 \times 10^{-1}$ in cometary grains, corresponding




FIG. 2.— N_2/CO ratio in cometary grains calculated as a function of the water abundance (normalized to the oxygen protosolar abundance (Lodders et al. 2009)) in the PSN for different formation temperatures. The blue and orange horizontal bars represent the N_2/CO ratios measured in comet R2 (Opitom et al. 2019) and 67P/C-G (Rubin et al. 2015, 2020), respectively. The two vertical dashed lines with arrows down correspond to upper limits of the N_2/CO ratio measured in 122P/1995 S1 (deVico) and C/1995 O1 (Hale-Bopp) (Cochran et al. 2000).

to the value calculated from protosolar N and C (with half C in CO and 90% N in N_2). These calculations illustrate the fact that, in principle, a slight increase of the water abundance in the PSN can lead to the full trapping of CO and enough N_2 to reproduce the N_2/CO ratio in R2.

However, the amount of water required by clathrates formed in the PSN to match the N_2/CO ratio measured in R2’s coma is inconsistent with its estimated low abundance. To match the N_2/CO measured in R2, our clathrate model requires a H_2O/CO ratio of 6.1. In contrast, the H_2O/CO mixing ratio measured at 2.8 AU in R2 is $\sim 0.32\%$, a value several orders of magnitude lower than those measured in 67P/C-G ($\sim 200\text{--}1900\%$) at a similar distance range (Gasc et al. 2017) or in many other comets (McKay et al. 2019). As a result, this mechanism fails at challenging to explain the H_2O depletion in R2-like comets via this mechanism.

3. AGGLOMERATION FROM PARTICLES FORMED IN THE VICINITY OF CO AND N_2 ICE LINES

Disk models including radial transport of volatiles can display significant enrichments of both volatile and refractory matter at its condensation location, implying potentially drastic changes of the local metallicity in the disk (Stevenson & Lunine 1988; Cyr et al. 1999; Ali-Dib et al. 2014; Mousis et al. 2019, 2020; Aguichine et al. 2020). Here we investigate this effect by considering the possibility that R2 formed in the vicinity of the condensation locations of CO and N_2 crystalline ices in the PSN.

3.1. Model

The volatile transport and distribution model used in our work is the one described in Aguichine et al. (2020) and Mousis et al. (2020), to which the reader is referred for details. Three energy sources are considered in our model, namely viscous heating, irradiation from the current Sun, and ambient constant irradiation giving a back-

ground temperature of 10 K (see Eq. 6 of Aguichine et al. (2020)). Here, the accretion rate onto the star declines over time, and the disk develops a transition radius between inward and outward gas flows. The location of this transition radius is moving outward with time. In a few words, our time-dependent PSN model is governed by the following differential equation (Lynden-Bell & Pringle 1974):

$$\frac{\partial \Sigma_g}{\partial t} = \frac{3}{r} \frac{\partial}{\partial r} \left[r^{1/2} \frac{\partial}{\partial r} \left(r^{1/2} \Sigma_g \nu \right) \right], \quad (2)$$

This equation describes the time evolution of a viscous accretion disk of surface density Σ_g of viscosity ν , assuming hydrostatic equilibrium in the z direction. The viscosity is calculated in the framework of the α -formalism (Shakura & Sunyaev 1973) using the following method. For each distance r to the Sun, the disk’s properties are calculated by solving the equation of energy balance between viscous heating and radiative transfer at the mid-plane level. This gives us ν , as well as the pressure and temperature profiles of the disk as a function of r . The evolution of the disk starts with an initial profile given by $\Sigma_g \nu \propto \exp(-r^{2-p})$, with $p = \frac{3}{2}$ for an early disk (Lynden-Bell & Pringle 1974). In our computations, the initial disk’s mass is fixed to $0.1 M_\odot$. The computational box is set equal to 500 AU, allowing 99% of the disk mass to be encapsulated within ~ 100 AU. Mass can only be lost by accretion onto the Sun or via outward diffusion, the boundary between these two regimes being defined by location of R_c , the centrifugal radius. R_c is initially located at ~ 6 AU and expands up to ~ 20 and 45 AU after 0.1 and 1 Myr of evolution, respectively. The disk’s initial mass accretion rate onto the Sun is set to $10^{-7.6} M_\odot \text{ yr}^{-1}$ (Hartmann et al. 1998).

The size of dust particles used in our model is determined by a two-populations algorithm derived from Birnstiel et al. (2012). This algorithm computes the representative size of particles through the estimate of the limiting Stokes number in various dynamical regimes. In our model, dust is initially present in the form of particles of sizes $a_0 = 10^{-7}$ m, and grow through mutual collisions. This growth is limited by the maximum sizes imposed by fragmentation or by the drift velocity of the grains (see Aguichine et al. (2020) for details). The dust surface density is the sum over all surface densities of available solids at given time and location, assuming a protosolar ice-to-rock ratio (~ 1.58) (Lodders et al. 2009), and a bulk density of 1000 kg m^{-3} .

We follow the approaches of Desch et al. (2017) and Drażkowska & Alibert (2017) for the dynamics of trace species in terms of motion and thermodynamics, respectively. We assume that the disk is uniformly filled with H_2O , CO, and N_2 . The abundances of CO, and N_2 are those derived in Sec. 2 and the abundance of H_2O is that of the leftover oxygen. No chemistry is assumed to happen between the trace species. In our simulations, grains are considered as homogeneous mixtures of solid species in proportions determined from the aforementioned assumptions (relative abundance ratios, condensation, sublimation). Sublimation of grains occurs during their inward drift when partial pressures of trace species become lower than the corresponding vapor pressures. Once re-

leased, vapors diffuse both inward and outward. Because of the outward diffusion, vapors can recondense back in solid form following the rates defined by Drażkowska & Alibert (2017), and condensation occurs either until thermodynamic equilibrium is reached or until no more gas is available to condense. The position of the iceline of a given species is defined as the location where its gaseous abundance equals that of its coexisting solid (Lodders 2003; Öberg & Wordsworth 2019; Aguichine et al. 2020). This approach results in icelines closer to the Sun than those computed via a simple comparison between the partial and saturation pressures.

The motion of dust and vapor is computed by integrating the 1D radial advection-diffusion equation derived from Birnstiel et al. (2012) and Desch et al. (2017), and detailed in Aguichine et al. (2020). The vapor pressures of trace species are taken from Fray & Schmitt (2009).

3.2. Results

Figure 3 represents the time evolution of the radial profiles of the N_2/CO ratio relative to its assumed initial ratio ($\sim 2.66 \times 10^{-1}$), defined by the enrichment factor f , in both solid and gas phases of the PSN, and for three values of the viscosity parameter α , namely 5×10^{-3} , 10^{-3} , and 10^{-4} . The adopted α values are well within the range of those typically used in models of protoplanetary disks (Hersant et al. 2001; Nelson et al. 2013; Simon et al. 2015). The figure shows that f is flat almost everywhere in the PSN, except in the vicinity of the CO and N_2 icelines, which are located at very close distances from each other, i.e. less than a few tenths of AU, in the 10–15 AU region of the PSN, with the CO iceline situated a bit closer in in the disk.

Depending on the choice of the α parameter, the N_2/CO ratio increases to more than ~ 10 in the gas phase in the area of the two icelines. This peak corresponds to the supply of N_2 vapor when N_2 -rich dust drifts inward the N_2 iceline, which is in excess compared to the CO vapor supplied via backward diffusion beyond the CO iceline. When significant, this peak is preceded by a decrease of the N_2/CO , which corresponds to the the supply of CO vapor when CO-rich dust drifts inward the CO iceline. The N_2/CO ratio in solid phase also experiences important depletions, which correspond to the location where N_2 essentially forms vapor while CO remains in solid phase. The N_2/CO ratio in solid phase also reaches peaks up to $f \sim 10$, depending on the α value, at the location of the N_2 iceline, corresponding to the formation of solid N_2 from N_2 vapor diffusing backward its corresponding iceline. The figure also shows it is possible to form dust in a narrow region of the PSN, i.e. within 10–15 AU, with N_2/CO ratios matching the value estimated in comet R2. A comet agglomerated from these grains would present a N_2/CO ratio consistent with the R2 value, both in its interior and its coma, because the two molecules present similar volatilities, as testified by the proximity of their icelines in the PSN.

Figure 4 represents the time evolution of the radial profiles of the CO/H_2O ratio relative to its assumed initial ratio ($\sim 2.96 \times 10^{-1}$), defined by the enrichment factor f , in both solid and gas phases of the PSN, and for three values of the viscosity parameter α , namely 5×10^{-3} , 10^{-3} , and 10^{-4} . The CO/H_2O ratio in the gas phase in-

creases steeply in the vicinity of the H_2O iceline, which is located at much closer distances from the Sun than the CO and N_2 icelines. This steep increase is due to the freezing of the H_2O vapor when it diffuses outward through its iceline. The CO/H_2O ratio is deeply depleted in the solid phase at distances inward of the CO iceline because this species remains in gaseous form. On the other hand, when approaching the CO iceline, the CO/H_2O ratio presents values exceeding its initial value. A peak forms at this location, and its magnitude depends on the adopted viscosity parameter α of the disk. The CO/H_2O enrichment factor f can typically reach a value of ~ 10 with $\alpha = 10^{-3}$, while it peaks at ~ 200 with $\alpha = 10^{-4}$. If the building blocks of a comet were assembled from grains formed at this location of the PSN, they should present a CO-rich and H_2O -poor composition. Assuming $f = 100$, the CO/H_2O ratio would be ~ 30 , a value exceeding by far those known in typical comets (Bockelée-Morvan & Biver 2017). We note that this value is still about 10 times lower than the one inferred in R2's coma but this latter can be seen as an overestimate of the bulk CO abundance in the comet. Indeed, CO is much more volatile than H_2O and its vapor comes from deeper layers than H_2O vapor in the nucleus, potentially inducing an enriched CO/H_2O ratio in the coma. This effect has been investigated in depth by Marboeuf & Schmitt (2014) who found that the abundance of volatile molecules (relative to H_2O) released from the interiors of nuclei can vary by several orders of magnitude along the comet's orbital evolution.

Interestingly, Figures 3 and 4 show that the locations of the CO and N_2 icelines do not significantly evolve with time and are only weakly affected by the value adopted for the viscosity parameter. The main reason for this is that viscous heating occurs mainly in the dense regions of the PSN. With our parametrization, the gas density quickly decreases after 10 AU, implying that the temperature profile beyond this distance is mostly ruled by solar irradiation, here assumed to be constant.

4. CONCLUSIONS

In this work, we first show that a comet agglomerated from clathrates crystallized in the PSN could, in principle, display N_2/CO ratios corresponding to the value measured in R2's coma. This finding contrasts with the suggestion of Wierzchos & Womack (2018) to rule out the presence of clathrates in R2 only because its N_2/CO ratio is higher than those measured in other comets. This statement is correct in the case of a limited budget of available crystalline water in the PSN. However, if water is abundant enough, as shown in this paper, both molecules are efficient clathrate formers. On the other hand, the amount of water required by clathrates formed in the PSN to match the N_2/CO ratio measured in R2's coma, namely $H_2O/CO \sim 6$, is inconsistent with its estimated extremely low value ($\sim 0.32\%$).

We have then used a disk model describing the radial transport of volatiles in the PSN, evolving both in gaseous and solid (pure condensates) phases. We show that N_2/CO ratios reproducing the value estimated in comet R2 can be found in dust formed in the vicinity of the CO and N_2 icelines, i.e. within the 10–15 AU region of the PSN, depending on the adopted viscosity parameter α parameter of the disk. Meanwhile, very high




FIG. 3.— From left to right: radial profiles of the N_2/CO ratio relative to its initial abundance (defined by the enrichment factor f) calculated as a function of time in the PSN for viscosity parameters $\alpha = 5 \times 10^{-3}$, 10^{-3} , and 10^{-4} . Dashed and solid lines correspond to vapor and solid phases, respectively. The blue bar corresponds to the N_2/CO ratio measured in comet R2.




FIG. 4.— From left to right: radial profiles of the $\text{CO}/\text{H}_2\text{O}$ ratio relative to its initial abundance (defined by the enrichment factor f) calculated as a function of time in the PSN for viscosity parameters $\alpha = 5 \times 10^{-3}$, 10^{-3} , and 10^{-4} . Dashed and solid lines correspond to vapor and solid phases, respectively.

$\text{CO}/\text{H}_2\text{O}$ ratios, i.e. up to more than 200 times the value derived from a protosolar gaseous mixture, can be found in dust formed in the vicinity of the CO iceline (~ 10 AU), the extent of which also depending on the adopted viscosity parameter. If the building blocks of a comet assembled from grains formed close to the CO iceline in the PSN, they should present N_2/CO and $\text{CO}/\text{H}_2\text{O}$ ratios consistent with the measurements made in comet R2’s coma.

The likelihood of finding R2-like comets versus H_2O -rich comets in the PSN can be, to first order, assessed via comparing the masses of their respective reservoirs. To do so, we calculated the mass of H_2O and CO ices contained in a PSN annulus in which CO was found to be more abundant than H_2O (R2-like comets – reservoir 1), and in another annulus where the $\text{CO}/\text{H}_2\text{O}$ ratio was set to 2.96×10^{-1} , i.e. the value derived from protosolar O and C abundances (average comets – reservoir 2). The time evolution of the reservoir 1/reservoir 2 mass ratio is represented in Fig. 5 for $\alpha = 10^{-3}$ and 10^{-4} . Higher α values do not allow for the formation of reservoir 1 in the PSN. This is also shown by Fig. 4 in the case $\alpha = 5 \times 10^{-3}$, where the $\text{CO}/\text{H}_2\text{O}$ ratio in the solid phase never exceeds twice the initial ratio. Figure 5 illustrates the fact that the mass ratio between the two reservoirs strongly depends on the adopted α parameter and the epoch of the PSN evolution. In the case $\alpha = 10^{-3}$, the mass of R2-like comets is about 1% that of the

average comets, if the grains from which they assembled condensed between 0.1 and 1 Myr in the PSN. In the case $\alpha = 10^{-4}$, the mass of R2-like comets varies between less than $\sim 1\%$ and 100% that of the average comets, depending on the particular grain formation epoch in the 0.1–1 Myr timeframe. We conclude that the likelihood of finding R2-like comets versus H_2O -rich comets in the PSN strongly depends on the adopted disk parameters, and thus remains difficult to predict a priori.

The heliocentric distances indicated for the different icelines must be taken with caution and are model-dependent. However, the agglomeration of R2 from dust condensed in the region of the CO iceline indicates that this comet formed at a greater heliocentric distance than a H_2O -rich comet formed from clathrates because the formation temperatures of N_2 - and CO -rich clathrates are always higher than those of the pure condensates at PSN conditions. Our model also applies to other protoplanetary disks and can explain the unusual composition of the interstellar comet 2I/Borisov, which seems to contain substantially more CO than H_2O gas in its coma (Bodewits et al. 2020).

Interestingly, physicochemical models of nitrogen chemistry in protostellar disks show that photodissociation of N_2 leads to production of HCN (Hily-Blant et al. 2017). To overcome this issue, R2 may have formed in a location of the PSN where significant N_2 shielding led to the high N_2 and decreased HCN abundances (Wierz-




FIG. 5.— Time evolution of the mass ratio between reservoir 1 (R2-like comets) and reservoir 2 (average comets) for viscosity parameters $\alpha = 10^{-3}$ and 10^{-4} .

chos & Womack 2018).

We thank the two Referees for their valuable comments on the content of our manuscript and their suggestions for improving the document. O.M. and A.B. acknowledge support from CNES, and J.L. from the JWST project. K.E.M. acknowledges support from the NASA RDAP grant 80NSSC19K1306, and A.L-K. acknowledges support from the NASA RDAP grant 80NSSC18K1620.

REFERENCES

- Aguichine, A., Mousis, O., Devouard, B., et al. 2020, *ApJ*, 901, 97
- Ali-Dib, M., Mousis, O., Petit, J.-M., et al. 2014, *ApJ*, 793, 9
- Birnstiel, T., Klahr, H., & Ercolano, B. 2012, *A&A*, 539, A148
- Biver, N., Bockelée-Morvan, D., Paubert, G., et al. 2018, *A&A*, 619, A127
- Bockelée-Morvan, D. & Biver, N. 2017, *Philosophical Transactions of the Royal Society of London Series A*, 375, 20160252. doi:10.1098/rsta.2016.0252
- Bodewits, D., Noonan, J. W., Feldman, P. D., et al. 2020, *Nature Astronomy*, 4, 867. doi:10.1038/s41550-020-1095-2
- Cochran, A. L., & McKay, A. J. 2018, *ApJ*, 854, L10
- Cochran, A. L., Cochran, W. D., & Barker, E. S. 2000, *Icarus*, 146, 583
- Cruikshank, D. P., Grundy, W. M., DeMeo, F. E., et al. 2015, *Icarus*, 246, 82
- Cyr, K. E., Sharp, C. M., & Lunine, J. I. 1999, *J. Geophys. Res.*, 104, 19003
- Desch, S. J., Estrada, P. R., Kalyaan, A., et al. 2017, *ApJ*, 840, 86
- Drażkowska, J. & Alibert, Y. 2017, *A&A*, 608, A92
- Fray, N. & Schmitt, B. 2009, *Planet. Space Sci.*, 57, 2053
- Gasc, S., Altwegg, K., Balsiger, H., et al. 2017, *MNRAS*, 469, S108. doi:10.1093/mnras/stx1412
- Hartmann, L., Calvet, N., Gullbring, E., et al. 1998, *ApJ*, 495, 385
- Hersant, F., Gautier, D., & Huré, J.-M. 2001, *ApJ*, 554, 391. doi:10.1086/321355
- Hily-Blant, P., Magalhaes, V., Kastner, J., et al. 2017, *A&A*, 603, L6. doi:10.1051/0004-6361/201730524
- Iro, N., Gautier, D., Hersant, F., et al. 2003, *Icarus*, 161, 511
- Klinger, J., Ocampo, J., & Schmitt, B. 1986, *Asteroids, Comets, Meteors II*, 241
- Lellouch, E., de Bergh, C., Sicardy, B., Käufel, H.-U., & Smette, A. 2011, *The Messenger*, 145, 20
- Lewis, J. S., & Prinn, R. G. 1980, *ApJ*, 238, 357
- Lodders, K., Palme, H., & Gail, H.-P. 2009, *Landolt Börnstein*, 712
- Lodders, K. 2003, *ApJ*, 591, 1220. doi:10.1086/375492
- Lunine, J. I., & Stevenson, D. J. 1985, *ApJS*, 58, 493
- Luspay-Kuti, A., Mousis, O., Hässig, M., et al. 2016, *Science Advances*, 2, 1501781
- Lynden-Bell, D. & Pringle, J. E. 1974, *MNRAS*, 168, 603
- Mandt, K., Luspay-Kuti, A., Hamel, M., et al. 2017, *MNRAS*, 472, 118. doi:10.1093/mnras/stx1587
- Marboeuf, U. & Schmitt, B. 2014, *Icarus*, 242, 225. doi:10.1016/j.icarus.2014.07.001
- Marboeuf, U., Schmitt, B., Petit, J.-M., et al. 2012, *A&A*, 542, A82
- Marboeuf, U., Mousis, O., Petit, J.-M., et al. 2011, *A&A*, 525, A144
- Marboeuf, U., Mousis, O., Petit, J.-M., et al. 2010, *ApJ*, 708, 812
- McKay, A. J., DiSanti, M. A., Kelley, M. S. P., et al. 2019, *AJ*, 158, 128
- Mohammadi, A. H., Anderson, R., & Tohidi, B. 2005, *Am. In. Chem. Eng.*, 51, 2825
- Mousis, O., Aguichine, A., Helled, R., et al. 2020, *Phil. Trans. R. Soc. A*, 378: 20200107. http://dx.doi.org/10.1098/rsta.2020.0107
- Mousis, O., Ronnet, T., & Lunine, J. I. 2019, *ApJ*, 875, 9
- Mousis, O., Ronnet, T., Lunine, J. I., et al. 2018, *ApJ*, 865, L11
- Mousis, O., Lunine, J. I., Luspay-Kuti, A., et al. 2016, *ApJ*, 819, L33
- Mousis, O., Guilbert-Lepoutre, A., Lunine, J. I., et al. 2012, *ApJ*, 757, 146
- Mousis, O., Lunine, J. I., Picaud, S., & Cordier, D. 2010, *Faraday Discussions*, 147, 509
- Mousis, O., Gautier, D., & Bockelée-Morvan, D. 2002, *Icarus*, 156, 162
- Nelson, R. P., Gressel, O., & Umurhan, O. M. 2013, *MNRAS*, 435, 2610. doi:10.1093/mnras/stt1475
- Öberg, K. I. & Wordsworth, R. 2019, *AJ*, 158, 194. doi:10.3847/1538-3881/ab46a8
- Opitom, C., Hutsemékers, D., Jehin, E., et al. 2019, *A&A*, 624, A64. doi:10.1051/0004-6361/201834357
- Pollack, J. B., Hollenbach, D., Beckwith, S., et al. 1994, *ApJ*, 421, 615
- Prinn, R. G. P., & Fegley, B., Jr. 1989, *Origin and Evolution of Planetary and Satellite Atmospheres*, 78
- Rubin, M., Engrand, C., Snodgrass, C., et al. 2020, *Space Sci. Rev.*, 216, 102
- Rubin, M., Altwegg, K., Balsiger, H., et al. 2015, *Science*, 348, 232
- Shakura, N. I. & Sunyaev, R. A. 1973, *A&A*, 500, 33
- Simon, J. B., Lesur, G., Kunz, M. W., et al. 2015, *MNRAS*, 454, 1117. doi:10.1093/mnras/stv2070
- Stevenson, D. J. & Lunine, J. I. 1988, *Icarus*, 75, 146
- Wierzchos, K. & Womack, M. 2018, *AJ*, 156, 34. doi:10.3847/1538-3881/aac6bc
- Womack, M., Sarid, G., & Wierzchos, K. 2017, *PASP*, 129, 031001. doi:10.1088/1538-3873/129/973/031001