

A RAMSEY THEOREM FOR GRADED LATTICES

ABHISHEK KHETAN AND AMITAVA BHATTACHARYA

ABSTRACT. We develop a Van der Waerden type theorem in an axiomatic setting of graded lattices and show that this axiomatic formulation can be applied to various lattices, for instance the set partition and the Boolean lattices. We derive the Hales-Jewett theorem as a corollary.

CONTENTS

1. Introduction	1
2. Definitions and Examples	2
3. Main theorem	4
4. Application: The Hales-Jewett Theorem	6
References	9

1. INTRODUCTION

Van der Waerden Theorem is one of the most important and basic results in Ramsey Theory. It states that if k and r are positive integers, then there is a natural number N such that whenever $\chi : [N] \rightarrow [k]$ is a k -coloring of the first N natural numbers, there is contained in $[N]$ an arithmetic progression of length r each of whose elements have the same color. There are many proofs of this result and it has led to many developments in Ramsey theory. See [GRS13] and [GR74].

One can interpret Van der Waerden theorem in the language of posets as follows. Let A_n denote the linear poset $\{1, \dots, n\}$ where the poset relation is the usual comparison relation on the natural numbers. What Van der Waerden theorem says is that if the linear poset A_N is colored with k -colors then there is a monochromatic ‘homothetic’ copy of A_r sitting in A_N , provided N is sufficiently large. It is thus natural to ask if one can prove Van der Waerden like theorems for other posets, for instance, the Boolean and the set partition posets.

The notion of homothety does not make sense for all posets (See [Sta12] for basic notions about posets). For this we need the posets to be graded. Roughly speaking, if P and Q are graded posets then a homothety of P into Q is a ‘translated and scaled’ copy of P in Q (See Definition 2.1). Both the Boolean and the set partition posets are graded posets. Moreover, both of them are lattices. This raises the following natural question. Suppose a large enough Boolean (partition) lattice is colored with k -colors, then can we find a monochromatic homothetic copy of a given small Boolean (partition) lattice in the large Boolean lattice. Further, we would like to insist that this embedding of the smaller lattice into the large lattice preserves the lattice relations. Our main theorem (Theorem 3.1) shows that this is indeed the case.

A key feature of these lattices is ‘self-similarity’. A large Boolean (partition) lattice contains copies of smaller Boolean (partition) lattices. Thus one would like to prove a general theorem

2010 *Mathematics Subject Classification.* 05D10.

Key words and phrases. poset, lattice, coloring, homothety.

about sequences of lattices which satisfy some kind of self similarity condition. To this end, we define a notion of a *homothety system* (See Definition 2.3) on a sequence of graded lattices in Section 2 and prove our main theorem in this abstract setting. Examples 2.7, 2.8, 2.9, 2.10 show that the division lattice, Boolean lattice, chain lattice, and the set partition lattice satisfy these axioms under the *trivial homothety system* (See Definition 2.3).

A substantial generalization of the Van der Waerden type results is the Hales-Jewett theorem proved in [HJ63]. We deduce the Hales-Jewett theorem in Section 4 as a corollary to our main theorem. This is done by considering a non-trivial homothety system on the chain lattices. The Hales-Jewett theorem is precisely the statement that one gets by applying Theorem 3.1 to this system. We hope that more applications can be found using other lattice, especially the set partition lattice. The case of the set partition lattice is especially interesting due to a theorem proved in [PTm80] which states that every lattice can be embedded in the partition lattice.

Various authors have developed Ramsey theorems for posets. See for instance [Tro75], [Tro99], [AW17]. However, to the best of the knowledge of the authors, the homothetic and lattice theoretic aspects have not been considered in literature.

2. DEFINITIONS AND EXAMPLES

We will write \mathbb{N} to denote the set of all the positive integers and \mathbb{N}_0 to denote the set of all the non-negative integers.

Definition 2.1. Let P and Q be graded posets with rank functions r_P and r_Q respectively. We say that a map $f : P \rightarrow Q$ is a **homothety** if f satisfies the following properties

- a) $f(p) \leq f(p')$ if and only if $p \leq p'$.
- b) There is $d \geq 1$, which we call the **scale factor** of f , such that $r_Q(f(p)) - r_Q(f(p')) = d(r_P(p) - r_P(p'))$ for all $p, p' \in P$.

Recall that if P and Q are graded posets with rank functions r_P and r_Q respectively, then $r : P \times Q \rightarrow \mathbb{N}_0$ defined as $r(p, q) = r_P(p) + r_Q(q)$ is a rank function on $P \times Q$, and hence the product is also a graded poset. Thus if $f_1 : P \rightarrow Q_1$ and $f_2 : P \rightarrow Q_2$ are homotheties, then so is the map $f : P \rightarrow Q_1 \times Q_2$ which takes $p \in P$ to $(f_1(p), f_2(p))$. Also note that the composition of two homotheties is again a homothety.

Definition 2.2. Suppose P and Q are graded lattices. A map $f : P \rightarrow Q$ is said to be a **lattice homothety** if f is a homothety and satisfies the condition

$$(2.1) \quad f(p \vee p') = f(p) \vee f(p') \quad \text{and} \quad f(p \wedge p') = f(p) \wedge f(p')$$

for all $p, p' \in P$.

Recall that if Q_1 and Q_2 are lattices then so is $Q_1 \times Q_2$, where

$$(2.2) \quad (q_1, q_2) \vee (q'_1, q'_2) = (q_1 \vee q'_1, q_2 \vee q'_2) \quad \text{and} \quad (q_1, q_2) \wedge (q'_1, q'_2) = (q_1 \wedge q'_1, q_2 \wedge q'_2)$$

Thus if P, Q_1 and Q_2 are graded lattices and $f_1 : P \rightarrow Q_1$ and $f_2 : P \rightarrow Q_2$ are lattice homotheties, then so is the map $f : P \rightarrow Q_1 \times Q_2$ taking p to $(f_1(p), f_2(p))$. A special case is the diagonal map $P \rightarrow P \times P$ which takes p to (p, p) is a lattice homothety whenever P is a graded lattice. Again, the composition of two lattice homotheties is a lattice homothety.

Definition 2.3. Let $A(1), A(2), A(3), \dots$ be a sequence of graded lattices. For each i and j let H_{ij} be a collection of lattice homotheties from $A(i)$ to $A(j)$. We say that $\mathcal{H} = \{H_{ij} : i, j \geq 1\}$ is a **homothety system** on $A(1), A(2), A(3), \dots$ if

- H1. The identity map $A(i) \rightarrow A(i)$ is in H_{ii} for all i .
- H2. Whenever $\varphi \in H_{ij}$ and $\psi \in H_{jk}$, we have $\psi \circ \varphi \in H_{ik}$.

We will refer to the elements of \mathcal{H} as **\mathcal{H} -restricted lattice homotheties**, or simply as **restricted lattice homotheties** when there is no ambiguity about the homothety system under consideration. When each H_{ij} consists of all possible lattice homotheties between $A(i)$ and $A(j)$, we call \mathcal{H} as the **trivial homothety system**. Despite the name, the trivial homothety system is of interest.

Definition 2.4. Let $A(1), A(2), A(3), \dots$ be a sequence of graded lattices equipped with a homothety system \mathcal{H} . Let $m, n, N \geq 1$ and $\varphi : A(m) \times A(n) \rightarrow A(N)$ be a lattice homothety. We say that φ is **compatible** with \mathcal{H} the following two conditions are satisfied.

C1. Whenever $A(i) \rightarrow A(m) \times A(n)$ is a lattice homothety such that each of the compositions

$$(2.3) \quad A(i) \rightarrow A(m) \times A(n) \xrightarrow{\text{proj}_1} A(m) \quad \text{and} \quad A(i) \rightarrow A(m) \times A(n) \xrightarrow{\text{proj}_2} A(n)$$

are restricted lattice homothety, the composition

$$(2.4) \quad A(i) \rightarrow A(m) \times A(n) \xrightarrow{\varphi} A(N)$$

is also a restricted lattice homothety.

C2. For any $p \in A(m)$ and $q \in A(n)$, the maps $\varphi_p : A(n) \rightarrow A(N)$, $\varphi_p(y) = \varphi(p, y)$ and $\varphi_q : A(m) \rightarrow A(N)$, $\varphi_q(x) = \varphi(x, q)$ are both restricted lattice homotheties.

Note that when \mathcal{H} is the trivial homothety system then every lattice homothety $A(m) \times A(n) \rightarrow A(N)$ is compatible with \mathcal{H} .

Definition 2.5. We say that a sequence $A(1), A(2), A(3), \dots$ of graded lattices equipped with a homothety system \mathcal{H} is **good** if for all $N_1, N_2 \geq 1$, there is N such that there is a lattice homothety from $A(N_1) \times A(N_2)$ into $A(N)$ which is compatible with \mathcal{H} .

Definition 2.6. We say that a sequence $A(1), A(2), A(3), \dots$ of graded lattices is **good** if it is good when it is equipped with the trivial homothety system.

Example 2.7. Let F be a finite set. We write $\Pi(F)$ to denote the collection of all partitions of F where we write $\pi \leq \tau$ for two partitions π and τ of F if τ is a refinement of π . Then $r : \Pi(F) \rightarrow \mathbb{N}$ defined as $r(\pi) = |\pi|$ is a grading on $\Pi(F)$. Also, if π and τ are arbitrary partitions of F , then there is a unique coarsest partition which refines both π and τ , and there is a unique finest partition which both π and τ refine. These are defined to be the join and meet of π and τ respectively, thus endowing $\Pi(F)$ with the structure of a graded lattice and we refer to $\Pi(F)$ as the **partition lattice** on F . We write $\Pi(n)$ to denote $\Pi(\{1, \dots, n\})$.

For two disjoint finite sets F and G , there is a natural map $\Pi(F) \times \Pi(G) \rightarrow \Pi(F \sqcup G)$ which sends (π, τ) to $\pi \cup \tau$. It is easy to check that this is a lattice homothety. From this observation one sees that the sequence $\Pi(1), \Pi(2), \Pi(3), \dots$, is a good sequence.

Example 2.8. For any finite set F we write $\text{Bool}(F)$ to denote the set of all the subsets of F . For $S, T \in \text{Bool}(F)$, we write $S \leq T$ if $S \subseteq T$, which endows it with a poset structure. It is easily checked that the function $r : \text{Bool}(F) \rightarrow \mathbb{N}_0$ taking S to $|S|$ is a grading on $\text{Bool}(F)$. Also, the operations of union and intersection play the role of join and meet respectively. Thus $\text{Bool}(F)$ is a graded lattice which we will refer to as the **Boolean lattice** on F . We write $\text{Bool}(n)$ to denote $\text{Bool}(\{1, \dots, n\})$.

When F and G are two disjoint sets, there is a natural map $\text{Bool}(F) \times \text{Bool}(G) \rightarrow \text{Bool}(F \sqcup G)$ taking (S, T) to $S \cup T$. It is clear that this is a lattice homothety and thus the sequence $\text{Bool}(1), \text{Bool}(2), \text{Bool}(3), \dots$ is a good sequence.

Example 2.9. For a positive integer n let D_n denote the set of all the non-negative integers which divide n . For a, b in D_n , we write $a \leq b$ if $a|b$. The function $r : D_n \rightarrow \mathbb{N}_0$ defined as

$$(2.5) \quad a = p_1^{k_1} \cdots p_r^{k_r} \mapsto k_1 + \cdots + k_r$$

where p_i 's are primes, is a rank function on D_n . Also, the least common multiple and the greatest common divisor serve as join and meet of any two elements. Thus D_n is a graded lattice which we refer to as the **division lattice** corresponding to n . Now let $m = p_1^{k_1} \cdots p_r^{k_r}$ and $n = q_1^{\ell_1} \cdots q_s^{\ell_s}$ be two positive integers, where p_i 's and q_j 's are distinct primes. Let $u_1, \dots, u_r, v_1, \dots, v_s$ be pairwise distinct primes and define

$$(2.6) \quad N = u_1^{k_1} \cdots u_r^{k_r} \cdot v_1^{\ell_1} \cdots v_s^{\ell_s}$$

Consider the map $f : D_m \times D_n \rightarrow D_N$ defined as

$$(2.7) \quad f(p_1^{\varepsilon_1} \cdots p_r^{\varepsilon_r}, q_1^{\delta_1} \cdots q_s^{\delta_s}) = u_1^{\varepsilon_1} \cdots u_r^{\varepsilon_r} \cdot v_1^{\delta_1} \cdots v_s^{\delta_s}$$

Then f is a lattice homothety, which shows that the sequence D_1, D_2, D_3, \dots is a good sequence.

Example 2.10. Fix a positive integer k . Define $\text{Chain}_k(F)$ as the set of all maps $F \rightarrow \{0, \dots, k-1\}$. We write $f \leq g$ for two elements f and g in $\text{Chain}_k(F)$ if $f(x) \leq g(x)$ for all $x \in F$. This gives a poset structure to $\text{Chain}_k(F)$. The map $r : \text{Chain}_k(F) \rightarrow \mathbb{N}_0$ defined as $r(f) = \sum_{x \in F} f(x)$ is a grading on $\text{Chain}_k(F)$. Finally, for two elements f and g in $\text{Chain}_k(F)$, we have

$$(2.8) \quad f \vee g = \max(f, g) \quad \text{and} \quad f \wedge g = \min(f, g)$$

serving as meet and join of f and g . Thus $\text{Chain}_k(F)$ is a graded lattice. We will refer to $\text{Chain}_k(F)$ as the k -th **chain lattice** over F .

Now let F and G be two disjoint sets. Consider the map $\text{Chain}_k(F) \times \text{Chain}_k(G) \rightarrow \text{Chain}_k(F \sqcup G)$ which takes (f, g) to the element $h \in \text{Chain}_k(F \sqcup G)$ defined as

$$(2.9) \quad h(x) = \begin{cases} f(x) & \text{if } x \in F \\ g(x) & \text{if } x \in G \end{cases}$$

It can be checked that this is a lattice homothety, which shows that the sequence

$$\text{Chain}_k(1), \text{Chain}_k(2), \text{Chain}_k(3), \dots$$

is a good sequence for any k . Note that $\text{Chain}_2(n)$ is same as $\text{Bool}(n)$.

3. MAIN THEOREM

Throughout this section let $A(1), A(2), A(3), \dots$ be a sequence of graded lattices equipped with a homothety system \mathcal{H} and assume that this system is good.

Theorem 3.1. *Let n and k be positive integers. Then there is a positive integer N such that whenever $\chi : A(N) \rightarrow [k]$ is a k -coloring of $A(N)$, there is a restricted lattice homothety $h : A(n) \rightarrow A(N)$ whose image is χ -monochromatic.*

Before getting into the proof of the above theorem we develop some definitions. Let P be a graded poset. Let $S \subseteq P$ and $p, p' \in P$ be arbitrary. We say that the triple $(S; p, p')$ is **admissible** if

- a) S has a minimum element.
- b) Both p and p' lie in S and are distinct.
- c) p is the minimum element of S .

Let P and Q be graded lattices and $\chi : Q \rightarrow [k]$ be a k -coloring of Q . Let q, q' be two elements in Q such that $q < q'$ and $(S; p, p')$ be an admissible triple in P . A $((S; p \rightarrow q, p' \rightarrow q'), \chi)$ -**lattice homothety** from P to Q is a lattice homothety $f : P \rightarrow Q$ such that

- a) $f(p) = q$ and $f(p') = q'$.
- b) $f(S) \setminus \{q'\}$ is χ -monochromatic.

Let k, ℓ, n and s be positive integers such that $n \geq s$ and $\ell, s \geq 2$. We define $L_n(s, k, \ell)$ as the smallest positive integer (if it exists) N such that whenever $(S; p, p')$ is an admissible triple in $A(n)$ with $|S| = s$, and $\chi : A(N) \rightarrow [k]$ is a k -coloring, then there is a sequence $q_1 < \dots < q_\ell$ in $A(N)$ such that for all $1 \leq i < j \leq \ell$ there is a $((S; p \rightarrow q_i, p' \rightarrow q_j), \chi)$ -restricted lattice homothety from $A(n)$ to $A(N)$.

Lemma 3.2. *If $L_n(s, k, \ell)$ and $L_n(s, k^{B(N_1)}, 2)$ exist then $L_n(s, k, \ell + 1)$ exists, where $N_1 = L_n(s, k, \ell)$ and $B(N_1)$ is the size of $A(N_1)$.*

Proof. Assume $N_1 := L_n(s, k, \ell)$ and $N_2 := L_n(s, k^{B(N_1)}, 2)$ exist. By definition of a good sequence there is N such that there is a lattice homothety $\mathbf{i} : A(N_1) \times A(N_2) \rightarrow A(N)$ compatible with \mathcal{H} . Fix an admissible triple $(S; p, p')$ in $A(n)$ with $|S| = s$ and a k -coloring $\chi : A(N) \rightarrow [k]$. We will show that there is a sequence $q_1 < \dots < q_{\ell+1}$ in $A(N)$ such that for each $1 \leq i < j \leq \ell + 1$, there is a $((S; p \rightarrow q_i, p' \rightarrow q_j), \chi)$ -restricted lattice homothety from $A(n)$ to $A(N)$.

The coloring χ gives an induced $k^{B(N_1)}$ coloring of $A(N_2)$ since the composite $\chi \circ \mathbf{i}$ can be thought of as a map $c : A(N_2) \rightarrow [k]^{A(N_1)}$. For each $y \in A(N_2)$ we thus have a map $c_y : A(N_1) \rightarrow [k]$ which takes $x \in A(N_1)$ to $\chi(x, y)$. Now using the fact that $L_n(s, k^{B(N_1)}, 2)$ exists, we get $b < b'$ in $A(N_2)$ such that there is a $((S; p \rightarrow b, p' \rightarrow b'), c)$ -restricted lattice homothety $h_2 : A(n) \rightarrow A(N_2)$. Thus c_y is same for all $y \in h_2(S) \setminus \{b'\}$. Let $\psi : A(N_1) \rightarrow [k]$ denote c_y for any such y . The existence of $L_n(s, k, \ell)$ now gives a sequence $p_1 < \dots < p_\ell$ in $A(N_1)$ such that for all $1 \leq i < j \leq \ell$ there is a $((S; p \rightarrow p_i, p' \rightarrow p_j), \psi)$ -restricted lattice homothety $h_1^{ij} : A(n) \rightarrow A(N_1)$. Now consider the list

$$(3.1) \quad \mathbf{i}(p_1, b) < \mathbf{i}(p_2, b) < \dots < \mathbf{i}(p_\ell, b) < \mathbf{i}(p_\ell, b')$$

in $A(N)$. We claim that for any two points y and y' with $y < y'$ in this list, there is a $((S; p \rightarrow y, p' \rightarrow y'), \chi)$ -restricted lattice homothety. We consider three cases.

Case 1: Suppose $y = \mathbf{i}(p_i, b)$ and $y' = \mathbf{i}(p_j, b)$ for some $i < j$. Consider the map $h : A(n) \rightarrow A(N)$ defined as $h(x) = \mathbf{i}(h_1^{ij}(x), b)$ for all $x \in S$. By using (C2) it is clear that this map is a $((S; p \rightarrow y, p' \rightarrow y'), \chi)$ -restricted lattice homothety.

Case 2: Suppose $y = \mathbf{i}(p_i, b)$ and $y' = \mathbf{i}(p_\ell, b')$ for some $i < \ell$. Consider the map $h : A(n) \rightarrow A(N)$ defined as

$$(3.2) \quad h(x) = \mathbf{i}(h_1^{i\ell}(x), h_2(x))$$

Now if $x, x' \in S$ with $x' \neq p'$, then we have

$$(3.3) \quad \chi \circ \mathbf{i}(h_1^{i\ell}(x), h_2(x)) = c_{h_2(x)}(h_1^{i\ell}(x)) = \psi(h_1^{i\ell}(x)) = \psi(h_1^{i\ell}(x'))$$

giving

$$(3.4) \quad \chi \circ \mathbf{i}(h_1^{i\ell}(x), h_2(x)) = c_{h_2(x')} (h_1^{i\ell}(x')) = \chi \circ \mathbf{i}(h_1^{i\ell}(x'), h_2(x'))$$

Thus, using (C1), we see that h is a $((S; p \rightarrow y, p' \rightarrow y'), \chi)$ -restricted lattice homothety and we are done.

Case 3: Suppose $y = \mathbf{i}(p_\ell, b)$ and $y' = \mathbf{i}(p_\ell, b')$. Consider the map $h : A(n) \rightarrow A(N)$ defined as $h(x) = \mathbf{i}(p_\ell, h_2(x))$ for each $x \in A(n)$. Then h is a restricted lattice homothety by (C2). We show that $h(S) \setminus \{y'\}$ is χ -monochromatic. So we need to show that if x and x' are in S with $x, x' \neq p'$, then $h(x)$ and $h(x')$ receive the same color under χ . Indeed,

$$(3.5) \quad \chi(h(x)) = \chi \circ \mathbf{i}(p_\ell, h_2(x)) = c_{h_2(x)}(p_\ell) = c_{h_2(x')} (p_\ell) = \chi \circ \mathbf{i}(p_\ell, h_2(x')) = \chi(h(x'))$$

Thus h is a $((S; p \rightarrow y, p' \rightarrow y'), \chi)$ -restricted lattice homothety and we are done. \blacksquare

Lemma 3.3. *If $L_n(s, k, k + 1)$ exists for some $2 \leq s < n$ and some $k \geq 1$ then $L_n(s + 1, k, 2)$ exists.*

Proof. Let $N = L_n(s, k, k+1)$ and $\chi : A(N) \rightarrow [k]$ be a k -coloring. Let $(S'; p, p')$ be an admissible triple in $A(n)$ with $|S'| = s + 1$. We will show that there are $b < b'$ in $A(N)$ such that there is a $((S'; p \rightarrow b, p' \rightarrow b'), \chi)$ -restricted lattice homothety from $A(n)$ to $A(N)$.

Let $S = S' \setminus \{p'\}$ and m' be an arbitrary element in S different from p . Such an m' exists since $|S'| = s + 1 > 2$. By definition of N , there is a sequence $q_1 < \dots < q_{k+1}$ in $A(N)$ such that for each $1 \leq i < j \leq k + 1$ there is a $((S; p \rightarrow q_i, m' \rightarrow q_j), \chi)$ -restricted lattice homothety h^{ij} from $A(n)$ to $A(N)$. By the pigeonhole principle there are q_i and q_j such that $\chi(q_i) = \chi(q_j)$. Thus the image of S under the corresponding restricted lattice homothety h^{ij} is χ -monochromatic in $A(N)$. Write $b = q_i$ and $b' = h^{ij}(p')$. Then we have h_{ij} is a $((S'; p \rightarrow b, p' \rightarrow b'), \chi)$ -restricted lattice homothety from $A(n)$ to $A(N)$ and we are done. \blacksquare

Fix $n \geq s \geq 2$ and let us write $L_n(s, *, *)$ to mean that $L_n(s, k, \ell)$ exists for all k and all $\ell \geq 2$. Similarly, we write $L_n(s, *, \ell)$ and $L_n(s, k, *)$.

Lemma 3.4. $L_n(2, *, *)$ is true.

Proof. First we show that $L_n(2, *, 2)$ is true. Let $k \geq 1$ be arbitrary. We need to show that $L_n(2, k, 2)$ exists. To show this all we need to show is that there is a restricted lattice homothety from $A(n)$ to some $A(N)$. To this end, by definition of a good sequence, we know that there is N such that there is a lattice homothety \mathbf{i} from $A(n) \times A(n)$ into $A(N)$ which is compatible with \mathcal{H} . By using (C1), (H1), and (H2), we see that the composition of \mathbf{i} with the diagonal embedding $A(n) \rightarrow A(n) \times A(n)$ is a restricted lattice homothety $A(n) \rightarrow A(N)$. Now the truth of $L_n(2, *, *)$ follows by repeated applications of Lemma 3.2. \blacksquare

Lemma 3.5. Fix $2 \leq s \leq |A(n)|$. Then $L_n(s, *, *)$ is true.

Proof. We prove this by induction. Lemma 3.4 shows that $L_n(2, *, *)$ is true. Suppose $L_n(i, *, *)$ is true for $i = 2, 3, \dots, s < |A(n)|$. We want to show that $L_n(s + 1, *, \ell)$ is true for all $\ell \geq 2$. By Lemma 3.3, the truth of $L_n(s, *, *)$ implies that $L_n(s + 1, *, 2)$ is true. Suppose we have shown that $L_n(s + 1, *, j)$ is true for $j = 2, \dots, \ell$ for some $\ell \geq 2$. Let $k \geq 1$ be arbitrary. The truth of $L_n(s + 1, *, \ell)$ and $L_n(s + 1, *, 2)$ in particular implies that $N := L_n(s + 1, k, \ell)$ and $L_n(s + 1, k^{B(N)}, 2)$ exist. Thus by Lemma 3.2 we know that $L_n(s + 1, k, \ell + 1)$ exists. Since k was arbitrary, this means that $L_n(s + 1, *, \ell + 1)$ is true, finishing the proof. \blacksquare

Proof of Theorem 3.1. Let $N = L_n(n, k, k + 1)$, which exists by Lemma 3.5. Fix an arbitrary k -coloring $\chi : A(N) \rightarrow [k]$. Let $S = A(n)$ and p and p' be the minimum and maximum elements of $A(n)$ respectively. Thus $(S; p, p')$ is an admissible triple in $A(n)$. By definition of N , there is a sequence $q_1 < \dots < q_{k+1}$ in $A(N)$ such that for each $1 \leq i < j \leq k + 1$ there is a $((S; p \rightarrow q_i, p' \rightarrow q_j), \chi)$ -restricted lattice homothety h from $A(n)$ to $A(N)$. By the pigeonhole principle, there are $i < j$ such that q_i and q_j receive the same color under χ . It is now clear that h^{ij} is the desired restricted lattice homothety. \blacksquare

4. APPLICATION: THE HALES-JEWETT THEOREM

Let t and n be positive integers. Let N be a positive integer and $\mathcal{S} = \{S_1, \dots, S_n\}$ be a collection of pairwise disjoint subsets of $[N]$ with $|S_1| = \dots = |S_n|$. Let $\alpha : [N] \rightarrow \{0, \dots, t - 1\}$ be a function that vanishes on $\bigcup \mathcal{S}$. We write $\varphi_{\mathcal{S}, \alpha}$ to mean the function $\text{Chain}_t(n) \rightarrow \text{Chain}_t(N)$ defined as

$$(4.1) \quad \varphi_{\mathcal{S}, \alpha}(a) = a_1 1_{S_1} + \dots + a_n 1_{S_n} + \alpha$$

for all $a = (a_1, \dots, a_n) \in \text{Chain}_t(n)$. Note that $\varphi_{\mathcal{S}, \alpha}$ is a lattice homothety with scale factor k , where k is the cardinality of any of the S_i 's. We say that a lattice homothety $h : \text{Chain}_t(n) \rightarrow \text{Chain}_t(N)$ is of **type HJ** if there exists a family \mathcal{S} of pairwise disjoint subsets of $[N]$ of equal

size with $|\mathcal{S}| = n$, and a function $\alpha : [N] \rightarrow \{0, \dots, t-1\}$ which vanishes on $\bigcup \mathcal{S}$, such that $h = \varphi_{\mathcal{S}, \alpha}$.

Now suppose $n_1 \leq n_2 \leq n_3$ are positive integers and $h_{12} : \text{Chain}_t(n_1) \rightarrow \text{Chain}_t(n_2)$ and $h_{23} : \text{Chain}_t(n_2) \rightarrow \text{Chain}_t(n_3)$ are homotheties of type HJ. We will show that the composite $h := h_{23} \circ h_{12}$ is also a homothety of type HJ. Let $h_{12} = \varphi_{\mathcal{S}, \alpha}$ and $h_{23} = \varphi_{\mathcal{T}, \beta}$, where $\mathcal{S} = \{S_1, \dots, S_{n_1}\}$ with $|S_1| = \dots = |S_{n_1}|$ and α vanishes on $\bigcup \mathcal{S}$, and $\mathcal{T} = \{T_1, \dots, T_{n_2}\}$ with $|T_1| = \dots = |T_{n_2}|$ and β vanishes on $\bigcup \mathcal{T}$.

Define $\gamma = h(0, \dots, 0)$ and for each $i \in [n_1]$ define U_i as $\bigcup_{p \in S_i} T_p$. It is then readily checked that all the U_i 's are of the same size and are pairwise disjoint, γ vanishes on $\bigcup_i U_i$, and

$$(4.2) \quad h(a_1, \dots, a_{n_1}) = a_1 1_{U_1} + \dots + a_{n_1} 1_{U_{n_1}} + \gamma$$

What we have shown is the following

Lemma 4.1. *Let t be a positive integer and H_{ij} be the set of all the homotheties of type HJ from $\text{Chain}_t(i)$ to $\text{Chain}_t(j)$. Then the family $\mathcal{H} = \{H_{ij} : i, j \geq 1\}$ is a homothety system on the sequence*

$$\text{Chain}_t(1), \text{Chain}_t(2), \text{Chain}_t(3), \dots$$

*We will refer to this homothety system as the **HJ homothety system** on the sequence of chain lattices.*

Lemma 4.2. *The trivial homothety system and the HJ homothety system coincide on the sequence of Boolean lattices.*

Proof. We prove this by induction. The lemma is clearly true for $n = 1, 2$. Suppose the lemma holds for $1, \dots, n$ and let $h : \text{Bool}(n+1) \rightarrow \text{Bool}(N)$ be a lattice homothety. Let k be the scale factor of h . Define

$$(4.3) \quad h_0 = h|\{0, 1\}^n \times \{0\} \quad \text{and} \quad h_1 = h|\{0, 1\}^n \times \{1\}$$

Then both h_1 and h_2 can be thought of as lattice homotheties from $\text{Bool}(n)$ to $\text{Bool}(N)$. By induction, there are families $\mathcal{S} = \{S_1, \dots, S_n\}$ and $\mathcal{T} = \{T_1, \dots, T_n\}$ of pairwise disjoint subsets of $[N]$, and function $\alpha, \beta : [N] \rightarrow \{0, 1\}$ which vanish on $\bigcup \mathcal{S}$ and $\bigcup \mathcal{T}$ respectively such that

$$(4.4) \quad h(a_1, \dots, a_n, 0) = h_0(a_1, \dots, a_n) = a_1 1_{S_1} + \dots + a_n 1_{S_n} + \alpha$$

$$(4.5) \quad h(a_1, \dots, a_n, 1) = h_1(a_1, \dots, a_n) = a_1 1_{T_1} + \dots + a_n 1_{T_n} + \beta$$

for all $(a_1, \dots, a_n) \in \text{Bool}(n)$. Thus

$$(4.6) \quad h(a_1, \dots, a_n, a_{n+1}) = (1 - a_{n+1})h_0 + a_{n+1}h_1$$

Since h has scale factor k , each S_i and each T_j has size k .

We claim that $S_i = T_i$ for all i . To this end, note that the restriction of h to $\{0, 1\} \times \{0\} \times \dots \times \{0\} \times \{0, 1\}$ can be thought of a lattice homothety from $\text{Bool}(2)$ to $\text{Bool}(N)$ having scale factor k . Thus there exist two disjoint sets P and Q of $[N]$ of size k , and function $\gamma : [N] \rightarrow \{0, 1\}$ which vanishes on $P \cup Q$, such that

$$(4.7) \quad h(a_1, 0, \dots, 0, a_{n+1}) = a_1 1_P + a_{n+1} 1_Q + \gamma$$

Using Equation 4.6 we have

$$(4.8) \quad a_1 1_P + \gamma = h(a_1, 0, \dots, 0) = a_1 1_{S_1} + \alpha$$

and

$$(4.9) \quad a_1 1_P + 1_Q + \gamma = h(a_1, 0, \dots, 0, 1) = a_1 1_{T_1} + \beta$$

which give that $S_1 = P = T_1$. It similarly follows that $S_i = T_i$ for all i . The above working also shows that $\beta = 1_Q + \gamma = 1_Q + \alpha$.

Thus we have

$$(4.10) \quad h(a_1, \dots, a_{n+1}) = a_1 1_{S_1} + \dots + a_n 1_{S_n} + a_{n+1} 1_Q + \alpha$$

Thus $S_i \cap Q$ is empty for each i and α vanishes on $Q \cup S_1 \cup \dots \cup S_n$. This shows that $h = \varphi_{\mathcal{F}, \alpha}$ where $\mathcal{F} = \{S_1, \dots, S_n, Q\}$ and we are done. \blacksquare

Example 4.3. We show that the trivial homothety system and HJ homothety system may not be the same for chain lattices. Consider the homothety $h : \text{Chain}_3(2) \rightarrow \text{Chain}_3(4)$ indicated by the following figure.




FIGURE 1. A homothety from $\text{Chain}_3(2)$ to $\text{Chain}_3(4)$.

It is easily checked that this is indeed a lattice homothety but is not of type HJ.

Lemma 4.4. *The sequence of chain lattices equipped with the HJ homothety system is a good sequence.*

Proof. Let $t \geq 1$ be fixed. Let F and G be disjoint non-empty sets and consider the map $\mathbf{i} : \text{Chain}_t(F) \times \text{Chain}_t(G) \rightarrow \text{Chain}_t(F \sqcup G)$ which takes (f, g) in $\text{Chain}_t(F) \times \text{Chain}_t(G)$ to the map $F \sqcup G \rightarrow \{0, \dots, t-1\}$ defined as

$$(4.11) \quad x \mapsto \begin{cases} f(x) & \text{if } x \in F \\ g(x) & \text{if } x \in G \end{cases}$$

For any function $\psi : G \rightarrow \{0, \dots, t-1\}$, let us write $\tilde{\psi}$ to denote the map $F \sqcup G \rightarrow \{0, \dots, t-1\}$ which vanishes on F and takes $y \in G$ to $\psi(y)$. Similarly, for any map $\varphi : F \rightarrow \{0, \dots, t-1\}$ we define $\tilde{\varphi} : F \sqcup G \rightarrow \{0, \dots, t-1\}$.

For $g_0 \in \text{Chain}_t(G)$ fixed, we see that the map $\mathbf{i}_{g_0} : \text{Chain}_t(F) \rightarrow \text{Chain}_t(F \sqcup G)$ takes $f \in \text{Chain}_t(F)$ to $\sum_{i \in F} f(i) 1_{\{i\}} + \tilde{g}_0$. Since \tilde{g}_0 vanishes on F , we see that \mathbf{i}_{g_0} is a homothety of type HJ. Similarly, $\mathbf{i}_{f_0} : \text{Chain}_t(F) \rightarrow \{0, \dots, t-1\}$ is a homothety of type HJ for each $f_0 \in \text{Chain}_t(F)$.

Finally, let S be a finite set and $h_F : \text{Chain}_t(S) \rightarrow \text{Chain}_t(F)$ and $h_G : \text{Chain}_t(S) \rightarrow \text{Chain}_t(G)$ be homotheties of type HJ. Consider the map $h : \text{Chain}_t(S) \rightarrow \text{Chain}_t(F) \times \text{Chain}_t(G)$ defined as $h(s) = (h_F(s), h_G(s))$.

We will show that $\mathbf{i} \circ h$ is a homothety of type HJ. Let $|S| = n$. Since h_F is a homothety of type HJ, there exists a family $\mathcal{F} = \{F_i : i \in S\}$ of pairwise disjoint subsets of F of equal

size, and a function $\alpha : F \rightarrow \{0, \dots, t-1\}$ vanishing on $\bigcup \mathcal{F}$ such that $h_F = \varphi_{\mathcal{F}, \alpha}$. Similarly, there is a family $\mathcal{G} = \{G_i : i \in S\}$ of pairwise disjoint subsets of G of equal size, and a function $\beta : G \rightarrow \{0, \dots, t-1\}$ vanishing on $\bigcup \mathcal{G}$ such that $h_G = \varphi_{\mathcal{G}, \beta}$. It is readily checked that

$$(4.12) \quad \mathbf{i} \circ h(s) = \sum_{i \in S} s(i) 1_{F_i \sqcup G_i} + \tilde{\alpha} + \tilde{\beta}$$

Let $\mathcal{E} = \{F_i \cup G_i : i \in S\}$. It is clear that $\gamma := \tilde{\alpha} + \tilde{\beta}$ vanishes on $\bigcup \mathcal{E}$ and that all elements of \mathcal{E} are of equal size. Thus $\mathbf{i} \circ h = \varphi_{\mathcal{E}, \gamma}$ and we are done. ■

Corollary 4.5. Hales-Jewett Theorem. Let n, t and k be positive integers. Then there is N such that whenever $\text{Chain}_t(N)$ is colored with k colors, there is a lattice homothety $h : \text{Chain}_t(n) \rightarrow \text{Chain}_t(N)$ of type HJ whose image is monochromatic.

Proof. This follows immediately from Lemma 4.4 and Theorem 3.1. ■

REFERENCES

- [AW17] Maria Axenovich and Stefan Walzer. Boolean lattices: Ramsey properties and embeddings. *Order*, 34(2):287–298, 2017.
- [GR74] R. L. Graham and B. L. Rothschild. A short proof of van der Waerden’s theorem on arithmetic progressions. *Proc. Amer. Math. Soc.*, 42:385–386, 1974.
- [GRS13] Ronald L. Graham, Bruce L. Rothschild, and Joel H. Spencer. *Ramsey theory*. Wiley Series in Discrete Mathematics and Optimization. John Wiley & Sons, Inc., Hoboken, NJ, 2013. Paperback edition of the second (1990) edition [MR1044995].
- [HJ63] A. W. Hales and R. I. Jewett. Regularity and positional games. *Trans. Amer. Math. Soc.*, 106:222–229, 1963.
- [PTm80] Pavel Pudlák and Jiří Tma. Every finite lattice can be embedded in a finite partition lattice. *Algebra Universalis*, 10(1):74–95, 1980.
- [Sta12] Richard P. Stanley. *Enumerative combinatorics. Volume 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2012.
- [Tro75] William T. Trotter, Jr. Embedding finite posets in cubes. *Discrete Math.*, 12:165–172, 1975.
- [Tro99] William T. Trotter. Ramsey theory and partially ordered sets. In *Contemporary trends in discrete mathematics (Štířín Castle, 1997)*, volume 49 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 337–347. Amer. Math. Soc., Providence, RI, 1999.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH. 1, HOMI BHABHA ROAD, MUMBAI-400005, INDIA

Email address: khetan@math.tifr.res.in

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH. 1, HOMI BHABHA ROAD, MUMBAI-400005, INDIA

Email address: amitavabhattacharya@protonmail.com