PARISIAN RUIN FOR INSURER AND REINSURER UNDER QUOTA-SHARE TREATY

GRIGORI JASNOVIDOV AND ALEKSANDR SHEMENDYUK

Abstract: In this contribution we study asymptotics of the simultaneous Parisian ruin probability of a two-dimensional fractional Brownian motion risk process. This risk process models the surplus processes of an insurance and a reinsurance companies, where the net loss is distributed between them in given proportions. We also propose an approach for simulation of Pickands and Piterbarg type constants appearing in the asymptotics of the ruin probability.

AMS Classification: Primary 60G15; secondary 60G70

Keywords: Brownian reinsurance risk process; fractional Brownian motion; simultaneous Parisian ruin; exact asymptotics; Piterbarg and Pickands constants

1. Introduction

Consider the risk model defined by

(1)
$$R(t) = u + \rho t - X(t), \quad t \ge 0,$$

where X(t) is a centered Gaussian risk process with a.s. continuous sample paths, $\rho > 0$ is the net profit rate and u > 0 is the initial capital. This model is relevant to insurance and financial applications, see, e.g, [1]. A question of numerous investigations is study of the asymptotics of the classical ruin probability

(2)
$$\lambda(u) := \mathbb{P}\left\{\exists t \ge 0 : R(t) < 0\right\}$$

as $u \to \infty$ under different levels of generality. It turns out, that only for X being a Brownian motion (later on BM) $\lambda(u)$ can be calculated explicitly. Namely, if X is a standard BM, then $\lambda(u) = e^{-2\rho u}$, $u, \rho > 0$, see, e.g., [2]. Since it seems impossible to find the exact value of $\lambda(u)$ in other cases, asymptotics of $\lambda(u)$ as $u \to \infty$ is dealt with. First the problem of a large excursion of a stationary Gaussian process was considered by J. Pickands in 1969, see [3]. We refer to monographs [4–6] for the survey of known results by the recent time. We would like to point out seminal manuscript [7] establishing asymptotics of $\lambda(u)$ under week assumptions on variance and covariance of X. For the discrete-time investigations (i.e., when t in model (1) belongs to a discrete grid $\{0, \delta, 2\delta...\}$ for some $\delta > 0$), we refer to [8–13]. We would like to suggest a reader contributions [14–23] for the related generalizations of the classical ruin problem. Some contributions (see, e.g., [18, 22, 23]), extend the classical ruin problem to the so-called Parisian ruin problem which allows the surplus process to spend a pre-specified time below zero before a ruin is recognized. Formally, the classical Parisian ruin probability is defined by

(3)
$$\mathbb{P}\left\{\exists t \geq 0 : \forall s \in [t, t+T] \ R(s) < 0\right\}, \qquad T \geq 0.$$

Date: January 22, 2024.

As in the classical case, only for X being a BM the probability above can be calculated explicitly (see [24]):

$$\mathbb{P}\left\{\exists t \ge 0 : \forall s \in [t, t+T] \ B(s) - cs > u\right\} = \frac{e^{-c^2T/2} - c\sqrt{2\pi T}\Phi(-c\sqrt{T})}{e^{-c^2T/2} + c\sqrt{2\pi T}\Phi(c\sqrt{T})}e^{-2cu}, \quad T \ge 0$$

where Φ is the distribution function of a standard Gaussian random variable and B is a standard BM. Note in passing, that the asymptotics of the Parisian ruin probability for X being a self-similar Gaussian processes is derived in [23]. We refer to [8, 18] for investigations of some other problems in this field.

Motivated by [25] (see also [10, 26]), we study a model where two companies share the net losses in proportions $\delta_1, \delta_2 > 0$, with $\delta_1 + \delta_2 = 1$, and receive the premiums at rates $\rho_1, \rho_2 > 0$, respectively. Further, the risk process of the *i*th company is defined by

$$R_i(t) = x_i + \rho_i t - \delta_i B(t), \quad t \ge 0, \ i = 1, 2,$$

where $x_i > 0$ is the initial capital of the *i*th company. In this model both claims and net losses are distributed between the companies, which corresponds to the proportional reinsurance dependence of the companies. In this paper we study the asymptotics of the simultaneous Parisian ruin probability defined by

$$\mathbb{P}\left\{\exists t \ge 0 : \forall s \in [t, t+T] \ R_1(s) < 0, R_2(s) < 0\right\}, \quad T \ge 0.$$

Since the probability above does not change under a scaling of (R_1, R_2) , it equals to

$$\mathbb{P}\left\{\exists t \ge 0 : \forall s \in [t, t+T] \ u_1 + c_1 s - B(s) < 0, u_2 + c_2 s - B(s) < 0\right\}, \quad T \ge 0,$$

where $u_i = x_i/\delta_i$ and $c_i = \rho_i/\delta_i$, i = 1, 2. Later on we derive the asymptotics of the probability above as u_1, u_2 tend to infinity at the constant speed (i.e., u_1/u_2 is constant). Therefore, we let $u_i = q_i u$ be fixed constants with $q_i > 0$, i = 1, 2 and deal with asymptotics of

$$\mathcal{P}_T(u) := \mathbb{P}\left\{\exists t > 0 : \forall s \in [t, t+T] \ B(s) > q_1 u + c_1 s, B(s) > q_2 u + c_1 s\right\}, \quad T > 0$$

as $u \to \infty$. Letting the initial capital tends to infinity is not just a mathematical assumption, but also an economic requirement stated by authorities in all developed countries, see [27]. It aims to prevents a company from the bankruptcy because of excessive number of small claims and/or several major claims, before the premium income is able to balance the losses and profits. Observe that $\mathcal{P}_T(u)$ can be rewritten as

$$\mathbb{P}\left\{\exists t \geq 0 : \forall s \in [t, t+T] \ B(s) - \max(c_1 s + q_1 u, c_2 s + q_2 u) > 0\right\}.$$

Thus, the two-dimensional problem may also be considered as a one-dimensional crossing problem over a piece-wise linear barrier. If the two lines $q_1u + c_1t$ and $q_2u + c_2t$ do not intersect over $(0, \infty)$, then the problem reduces to the classical one-dimensional BM risk model, which has been discussed in [18, 23] and thus will not be the focus of this paper. In consideration of that, we assume that

$$(4) c_1 > c_2, \quad q_2 > q_1.$$

Under the assumption above the lines $q_1u + c_1t$ and $q_2u + c_2t$ intersects at point ut_* with

$$(5) t_* = \frac{q_2 - q_1}{c_1 - c_2} > 0$$

that plays a crucial role in the following. The first usual step when dealing with asymptotics of a ruin probability of a Gaussian process is centralizing the process. In our case it can be achieved by the self-similarity of BM:

$$\mathcal{P}_{T}(u) = \mathbb{P}\left\{\exists tu \geq 0 : \inf_{su \in [tu, tu + T]} (B(su) - c_{1}su) > q_{1}u, \inf_{su \in [tu, tu + T]} (B(su) - c_{2}su) > q_{2}u\right\}$$

$$= \mathbb{P}\left\{\exists t \geq 0 : \inf_{s \in [t, t + T/u]} (B(s) - (c_{1}s + q_{1})\sqrt{u}) > 0, \inf_{s \in [t, t + T/u]} (B(s) - (c_{2}s + q_{2})\sqrt{u}) > 0\right\}$$

$$= \mathbb{P}\left\{\exists t \geq 0 : \inf_{s \in [t, t + T/u]} \frac{B(s)}{\max(c_{1}s + q_{1}, c_{2}s + q_{2})} > \sqrt{u}\right\}.$$

The next step is analysis of the variance of the centered process. Note that the variance of $\frac{B(t)}{\max(c_1t+q_1,c_2t+q_2)}$ can achieve its unique maxima only at one of the following points:

$$t_*, \quad \overline{t}_1 := \frac{q_1}{c_1}, \quad \overline{t}_2 := \frac{q_2}{c_2}.$$

From (4) it follows that $\overline{t}_1 < \overline{t}_2$. As we see later, the position of t_* regarding to $(\overline{t}_1, \overline{t}_2)$ determines the asymptotics of $\mathcal{P}_T(u)$. Note, that the variance of $\frac{B(t)}{\max(c_1t+q_1,c_2t+q_2)}$ is not smooth around t_* if (4) is satisfied; this observation does not allow us to obtain the asymptotics of $\mathcal{P}_T(u)$ straightforwardly by [23]. Define for any $L \geq 0$ and some function $h : \mathbb{R} \to \mathbb{R}$ constant

$$\mathcal{F}_{L}^{h} = \mathbb{E} \left\{ \sup_{t \in \mathbb{R}} \inf_{s \in [t, t+L]} e^{\sqrt{2}B(s) - |s| + h(s)} \right\}$$

when the expectation above is finite. For the properties of \mathcal{F}_L^h and related Piterbarg constants we refer to [4, 18, 23, 28]. Notice that \mathcal{F}_0^h is the Piterbarg constant introduced in [25]. Let $\overline{\Phi}$ be the survival function of a standard Gaussian random variable and $\mathbb{I}(\cdot)$ be the indicator function. The next theorem derives the asymptotics of $\mathcal{P}_T(u)$ as $u \to \infty$:

Theorem 1.1. Assume that (4) holds.

1) If $t_* \notin (\overline{t}_1, \overline{t}_2)$, then as $u \to \infty$

(6)
$$\mathcal{P}_{T}(u) \sim \left(\frac{1}{2}\right)^{\mathbb{I}(t_{*}=\overline{t}_{i})} \frac{e^{-c_{i}^{2}T/2} - c_{i}\sqrt{2\pi T}\Phi(-c_{i}\sqrt{T})}{e^{-c_{i}^{2}T/2} + c_{i}\sqrt{2\pi T}\Phi(c_{i}\sqrt{T})} e^{-2c_{i}q_{i}u},$$

where i = 1 if $t_* \leq \overline{t}_1$ and i = 2 if $t_* \geq \overline{t}_2$.

2) If $t_* \in (\overline{t}_1, \overline{t}_2)$, then as $u \to \infty$

$$\mathcal{P}_T(u) \sim \mathcal{F}_{T'}^d \overline{\Phi} \left((c_1 q_2 - c_2 q_1) \sqrt{\frac{q_2 - q_1}{c_1 - c_2}} \sqrt{u} \right),$$

where $\mathcal{F}_{T'}^d \in (0, \infty)$ and

$$(7) T' = T \frac{(c_1 q_2 - q_1 c_2)^2}{2(c_1 - c_2)^2}, d(s) = s \frac{c_1 q_2 + c_2 q_1 - 2c_2 q_2}{c_1 q_2 - q_1 c_2} \mathbb{I}(s < 0) + s \frac{2c_1 q_1 - c_1 q_2 - q_1 c_2}{c_1 q_2 - q_1 c_2} \mathbb{I}(s \ge 0).$$

2. Main Results

In classical risk theory, the surplus process of an insurance company is modeled by the compound Poisson or the general compound renewal risk process, see, e.g., [1]. The calculation of the ruin probabilities is of a particular interest for both theoretical and applied domains. To avoid the technical issues and allow for dependence between claim sizes, these models are often approximated by the risk model (1), driven

by B_H a standard fractional Brownian motion (later on fBm), i.e, Gaussian process with zero-mean and covariance function

$$cov(B_H(t), B_H(s)) = \frac{t^{2H} + s^{2H} - |t - s|^{2H}}{2}, \quad s, t \in \mathbb{R}, \ H \in (0, 1).$$

Since the time spent by the surplus process below zero may depend on u, in the following we allow $T =: T_u$ in (3) to depend on u. As mentioned in [18], for the one-dimensional Parisian ruin probability we need to control the growth of T_u as $u \to \infty$. Namely, we impose the following condition:

(8)
$$\lim_{u \to \infty} T_u u^{1/H-2} = T \in [0, \infty), \ H \in (0, 1).$$

Note that if H > 1/2, then T_u may grow to infinity, while if H < 1/2, then T_u approaches zero as u tends to infinity. As we see later in Proposition 2.2, the condition above is necessary and the result does not hold without it. As for BM, by the self-similarity of fBm we obtain

$$\mathcal{P}_{T_u}(u) := \mathbb{P}\left\{\exists t \ge 0 : \forall s \in [t, t + T_u] \ B_H(s) > q_1 u + c_1 s, B_H(s) > q_2 u + c_1 s\right\}$$
$$= \mathbb{P}\left\{\exists t \ge 0 : \inf_{s \in [t, t + T_u/u]} \frac{B_H(s)}{\max(c_1 s + q_1, c_2 s + q_2)} > u^{1-H}\right\}.$$

The variance of $\frac{B_H(t)}{\max(c_1t+q_1,c_2t+q_2)}$ can achieve its unique maxima only at one of the following points:

(9)
$$t_*, \quad t_1 := \frac{Hq_1}{(1-H)c_1}, \quad t_2 := \frac{Hq_2}{(1-H)c_2}.$$

From (4) it follows that $t_1 < t_2$. Again, the position of t_* regarding to (t_1, t_2) determines the asymptotics of $\mathcal{P}_{T_u}(u)$. Define for $H \in (0, 1)$ and $T \geq 0$ Pickands constants by

$$\mathbb{H}_{2H} = \lim_{S \to \infty} \frac{1}{S} \mathbb{E} \left\{ \sup_{t \in [0,S]} e^{\sqrt{2}B_H(t) - t^{2H}} \right\}, \qquad \mathcal{F}_{2H}(T) = \lim_{S \to \infty} \frac{1}{S} \mathbb{E} \left\{ \sup_{t \in [0,S]} \inf_{s \in [0,T]} e^{\sqrt{2}B_H(t+s) - (t+s)^{2H}} \right\}.$$

It is shown in [23] and [4], respectively, that $\mathcal{F}_{2H}(T)$ and \mathbb{H}_{2H} are finite positive constants. Let

$$(10)\,\mathbb{D}_{H} = \frac{c_{1}t_{*} + q_{1}}{t_{*}^{H}}, \ K_{H} = \frac{2^{\frac{1}{2} - \frac{1}{2H}}\sqrt{\pi}}{\sqrt{H(1 - H)}}, \ \mathbb{C}_{H}^{(i)} = \frac{c_{i}^{H}q_{i}^{1 - H}}{H^{H}(1 - H)^{1 - H}}, \ D_{i} = \frac{c_{i}^{2}(1 - H)^{2 - \frac{1}{H}}}{2^{\frac{1}{2H}}H^{2}}, \ i = 1, 2.$$

Now we are ready to give the asymptotics of $\mathcal{P}_{T_u}(u)$:

Theorem 2.1. Assume that (4) holds and T_u satisfies (8). 1) If $t_* \notin (t_1, t_2)$, then as $u \to \infty$

(11)
$$\mathcal{P}_{T_{u}}(u) \sim \left(\frac{1}{2}\right)^{\mathbb{I}(t_{*}=t_{i})} \times \begin{cases} \frac{e^{-c_{i}^{2}T/2} - c_{i}\sqrt{2\pi T}\Phi(-c_{i}\sqrt{T})}{e^{-c_{i}^{2}T/2} + c_{i}\sqrt{2\pi T}\Phi(c_{i}\sqrt{T})} e^{-2c_{i}q_{i}u}, & H = 1/2, \\ K_{H}\mathcal{F}_{2H}(TD_{i})(\mathbb{C}_{H}^{(i)}u^{1-H})^{\frac{1}{H}-1}\overline{\Phi}(\mathbb{C}_{H}^{(i)}u^{1-H}), & H \neq 1/2, \end{cases}$$

where i = 1 if $t_* \le t_1$ and i = 2 if $t_* \ge t_2$.

2) If $t_* \in (t_1, t_2)$ and $\lim_{u \to \infty} T_u u^{2-1/H} = 0$ for H > 1/2, then as $u \to \infty$

(12)
$$\mathcal{P}_{T_{u}}(u) \sim \overline{\Phi}(\mathbb{D}_{H}u^{1-H}) \times \begin{cases} 1, & H > 1/2, \\ \mathcal{F}_{T'}^{d}, & H = 1/2, \\ \mathcal{F}_{2H}(\overline{D}T)Au^{(1-H)(1/H-2)}, & H < 1/2, \end{cases}$$

where $\mathcal{F}_{T'}^d \in (0, \infty)$ with T' and d defined in (7) and

$$(13) \quad A = \left(|H(c_1 t_* + q_1) - c_1 t_*|^{-1} + |H(c_2 t_* + q_2) - c_2 t_*|^{-1} \right) \frac{t_*^H \mathbb{D}_H^{\frac{1}{H} - 1}}{2^{\frac{1}{2H}}}, \qquad \overline{D} = \frac{(c_1 t_* + q_1)^{\frac{1}{H}}}{2^{\frac{1}{2H}} t_*^2}.$$

The theorem above generalizes Theorem 1.1 and Theorem 3.1 in [25]. Note that if T = 0, then the result above reduces to Theorem 3.1 in [25]. As indicated in [18], it seems extremely difficult to find the exact asymptotics of the one-dimensional Parisian ruin probability if (8) does not hold. The intuitive reason is that the ruin happens over 'too long interval'. To illustrate difficulties arising in approximation of $\mathcal{P}_{T_u}(u)$ in this setup we consider a 'simple' scenario: let $T_u = T > 0$ and H < 1/2. In this case we have

Proposition 2.2. If H < 1/2, $T_u = T > 0$ and $t_* \in (t_1, t_2)$, then

$$(14) \ \overline{C}\overline{\Phi}(\mathbb{D}_{H}u^{1-H})e^{-C_{1,\alpha}u^{2-4H}-C_{2,\alpha}u^{2(1-3H)}} \leq \mathcal{P}_{T_{u}}(u) \leq (2+o(1))\overline{\Phi}(\mathbb{D}_{H}u^{1-H})\overline{\Phi}\left(u^{1-2H}\frac{T^{H}\mathbb{D}_{H}}{2t_{*}^{H}}\right),$$

where $\bar{C} \in (0,1)$ is a fixed constant that does not depend on u and

(15)
$$\alpha = \frac{T^{2H}}{2t_*^{2H}}, \qquad C_{i,\alpha} = \frac{\alpha^i}{i} \mathbb{D}_H^2, \quad i = 1, 2.$$

Note that the result above expands Theorem 3.2 in [18] for fBm case. Comparing the proposition above with Theorem 2.1 we see that right hand part above is exponentially smaller than the corresponding asymptotics in case H < 1/2 in Theorem 2.1 and hence condition (8) indeed is important.

3. Simulation of Piterbarg & Pickands constants

In this section we give algorithms for simulations of Pickands and Piterbarg type constants appearing in Theorems 1.1 and 2.1 and study their properties relevant for simulations. Since the classical Pickands constant \mathbb{H}_{2H} has been investigated in several contributions (see, e.g., [29] and references therein), later on we deal with \mathcal{F}_L^h and $\mathcal{F}_{2H}(L)$. For notation simplicity we denote for any real numbers x < y and $\tau > 0$

$$[x,y]_{\tau} = [x,y] \cap \tau \mathbb{Z}.$$

Simulation of Piterbarg constant. In this subsection we always assume that

$$L \ge 0$$
 and $h(s) = bs \ \mathbb{I}(s < 0) - as \ \mathbb{I}(s \ge 0), \ s \in \mathbb{R}, \ a, b > 0.$

To simulate \mathcal{F}_L^h we use approximation

$$\mathcal{F}_L^h \approx \mathbb{E} \left\{ \sup_{t \in [-M,M]_\tau} \inf_{s \in [t,t+L]_\tau} e^{\sqrt{2}B(s) - |s| + h(s)} \right\},\,$$

where M is sufficiently large and $\tau > 0$ is sufficiently small. The approximation above has several errors: truncation error (i.e., choice of M), discretization error (i.e., choice of τ) and simulation error. It seems difficult to give a precise estimate of the discretization error, we refer to [12, 29] for discussion of such problems. To take an appropriate M and give an upper bound of the truncation error we derive few lemmas. The first lemma provides us bounds for \mathcal{F}_L^h :

Lemma 3.1. It holds that for L > 0

$$2e^{-L\min(a,b)}\overline{\Phi}(\sqrt{2L}) \le \mathcal{F}_L^h < 1 + \frac{1}{a} + \frac{1}{b} - \frac{1}{a+b+1}$$

and

$$\mathcal{F}_0^h = 1 + \frac{1}{a} + \frac{1}{b} - \frac{1}{a+b+1}.$$

Note that the second statement of the lemma gives the explicit expression for the two-sided Piterbarg constant introduced in [25]. In the next lemma we focus on the truncation error:

Lemma 3.2. For $M \geq 0$ it holds that

(16)
$$\mathbb{E}\left\{\sup_{t\in\mathbb{R}\setminus[-M,M]}\inf_{s\in[t,t+L]}e^{\sqrt{2}B(s)-|s|+h(s)}\right\} \leq e^{-aM}\left(1+\frac{1}{a}\right)+e^{-bM}\left(1+\frac{1}{b}\right).$$

Now we are ready to find an appropriate M. We have by Lemma 3.2 that

$$\left| \mathcal{F}_{L}^{h} - \mathbb{E} \left\{ \sup_{t \in [-M,M]} \inf_{s \in [t,t+L]} e^{\sqrt{2}B(s) - |s| + h(s)} \right\} \right| \leq \mathbb{E} \left\{ \sup_{t \in \mathbb{R} \setminus [-M,M]} \inf_{s \in [t,t+L]} e^{\sqrt{2}B(s) - |s| + h(s)} \right\}$$

$$\leq 2 \left(1 + \frac{1}{\min(a,b)} \right) e^{-M\min(a,b)}$$

and on the other hand by Lemma 3.1

$$\mathcal{F}_L^h \ge 2e^{-L\min(a,b)}\overline{\Phi}(\sqrt{2L})$$

hence to obtain a good accuracy we need that

$$\left(1 + \frac{1}{\min(a,b)}\right)e^{-\min(a,b)M} \ll e^{-L\min(a,b)}\overline{\Phi}(\sqrt{2L}).$$

Assume for simulations that $\min(a,b) \geq 1$; otherwise special case $\min(a,b) << 1$ requires a choice of a large M implying very high level of computation capacity. For simulations, we take $M = \frac{7 + L(3 + \min(a,b))}{\min(a,b)}$ providing us truncation error smaller than $3*10^{-3}$; we do not need to have better due to the discretization and simulation errors. Since it seems difficult to estimate these errors, we just take a 'small' τ and a 'big' number of simulation n. The above observations give us the following algorithm:

- 1) take $M = \frac{7 + L(3 + \min(a,b))}{\min(a,b)}, \tau = 0.005$ and $n = 10^4$;
- 2) simulate n times B(t), $t \in [-M, M]_{\tau}$, i.e, obtain $B_i(t)$, $1 \le i \le n$;
- 3) compute

$$\widehat{\mathcal{F}_L^h} := \frac{1}{n} \sum_{i=1}^n \sup_{t \in [-M,M]_\tau} \inf_{s \in [t,t+L]_\tau} e^{\sqrt{2}B_i(s) - |s| + h(s)}.$$

Simulation of Pickands constant. It seems difficult to simulate $\mathcal{F}_{2H}(L)$ relying straightforwardly on its definition. As follows from approach in [29, 30] for any $\eta > 0$ with $W(t) = B_{2H}(t) - |t|^{2H}$

$$\mathcal{F}_{2H}(L) = \mathbb{E}\left\{\frac{\sup_{t \in \mathbb{R}} \inf_{s \in [t, t+L]} e^{W(t)}}{\eta \sum_{k \in \mathbb{Z}} e^{W(k\eta)}}\right\}.$$

The merit of the representation above is that there is no limit as is in the original definition and thus it is much easier to simulate $\mathcal{F}_{2H}(L)$ by the Monte-Carlo method. The second benefit is that there is a sum in the denominator, that can be simulated easily with a good accuracy. The only drawback is that the

sup inf in the nominator is taken on the whole real line. Thus, we approximate $\mathcal{F}_{2H}(L)$ by discrete analog of the formula above:

$$\mathcal{F}_{2H}(L) \approx \mathbb{E} \left\{ \frac{\sup_{t \in [-M,M]_{\tau}} \inf_{s \in [t,t+L]_{\tau}} e^{W(t)}}{\eta \sum_{k \in [-M,M]_{\eta}} e^{W(\eta k)}} \right\},$$

where big M and small τ, η are appropriately chosen positive numbers. In the following lemma we give a lower bound for $\mathcal{F}_{2H}(L)$.

Lemma 3.3. It holds that for any L > 0 and $H \in (0,1)$

$$\mathcal{F}_{2H}(L) \ge \mathbb{E}\left\{ \left(\int_{\mathbb{R}} e^{W(t)} dt \right)^{-1} \right\} e^{-L^{2H}} \sup_{m>0} \left(e^{-\sqrt{2}mL^H} \mathbb{P}\left\{ \sup_{s \in [0,1]} B_H(s) < m \right\} \right)$$
with $\mathbb{E}\left\{ \left(\int_{\mathbb{R}} e^{W(t)} dt \right)^{-1} \right\} \in (0,\infty).$

Taking $m = 1/\sqrt{2}$ in the sup above we obtain a useful for large L estimate

$$\mathcal{F}_{2H}(L) \ge Ce^{-L^{2H} - L^H}, \quad L > 0$$

where C is a some positive number that depends only on H. The following lemma provides us an upper bound for the truncation error:

Lemma 3.4. For some fixed constant c' > 0 and M, L > 0 it holds that

$$\left| \mathcal{F}_{2H}(L) - \mathbb{E} \left\{ \frac{\sup_{t \in [-M,M]} \inf_{s \in [t,t+L]} e^{W(t)}}{\int_{[-M,M]} e^{W(t)} dt} \right\} \right| \le e^{-c'M^{2H}}.$$

Based on 2 lemmas above we propose the following algorithm for simulation of $\mathcal{F}_{2H}(L)$:

- 1) Take $M = \max(10L, 5)$, $\tau = \eta = 0.005$ and $n = 10^4$;
- 2) simulate n times $B_H(t), t \in [-M, M]_{\tau}$, i.e, obtain $B_H^{(i)}(t), 1 \le i \le n$;
- 3) calculate


$$\widehat{\mathcal{F}_{2H}(L)} := \frac{1}{n} \sum_{i=1}^{n} \frac{\sup_{t \in [-M,M]_{\tau}} \inf_{s \in [t,t+L]_{\tau}} e^{\sqrt{2}B_{H}^{(i)}(s) - |s|^{2H}}}{\eta \sum_{k \in [-M,M]_{\eta}} e^{\sqrt{2}B_{H}^{(i)}(k\eta) - |k\eta|^{2H}}}.$$


We give the proofs of all Lemmas above at the end of Section Proofs.


4. Approximate values of Pickands & Piterbarg constants

In this section we apply both algorithms introduced above and obtain approximate numerical values for some particular choices of parameters. To implement our approach we use MATLAB software.

Piterbarg constant. We simulate several graphs of $\widehat{\mathcal{F}_L^h}$ for different choices of a and b:









On each graph above the blue line is simulated value and the dashed lines are theoretical bounds given in Lemma 3.1. We observe that the simulated values are between the theoretical bounds given in Lemma 3.1, $\widehat{\mathcal{F}_L^h}$ is decreasing function and $\widehat{\mathcal{F}_L^h}$ tends to $1 + \frac{1}{a} + \frac{1}{b} - \frac{1}{a+b+1}$ as $L \to 0$.

Pickands constant. We plot several graphs of $\widehat{\mathcal{F}_{2H}(L)}$ for different choices of H. Since the value of $\mathcal{F}_1(L)$ is known, namely, $\mathcal{F}_1(L) = \frac{e^{-L/4} - \sqrt{\pi L}\Phi(-\sqrt{L/2})}{e^{-L/4} + \sqrt{\pi L}\Phi(\sqrt{L/2})}, \ L \geq 0$, (see, e.g., [24]) we consider for simulations only cases H < 0.5 and H > 0.5. To simulate fBm we use Choleski method, (see, e.g., [31]).

Short-range dependence case. We take H equal to 0.1, 0.2, 0.3 and 0.4 and plot $\widehat{\mathcal{F}_{2H}(L)}$ for these values.



Long-range dependence case. Here we take H from $\{0.6, 0.7, 0.8, 0.9\}$ and plot $\widehat{\mathcal{F}_{2H}(L)}$ for these values.



Observe that $\widehat{\mathcal{F}_{2H}(L)}$ is a strictly decreasing function of L for all $H \in (0,1)$. It seems also that $\mathcal{F}_{2H}(L)$ for fixed L is an increasing function of H for $H \in (0,1/2)$ and is not an increasing function of H for $H \in (1/2,1)$.

5. Proofs

Before giving our proofs we formulate a few auxiliary statements. As shown, e.g., in [4]

(17)
$$\overline{\Phi}(x) \sim \frac{e^{-x^2/2}}{\sqrt{2\pi}x}, \quad x \to \infty.$$

Recall that K_H, D_1 and $\mathbb{C}_H^{(1)}$ are defined in (10). The following result immediately follows from [23, 24]:

Proposition 5.1. Assume that T_u satisfies (8). Then as $u \to \infty$

$$\mathbb{P}\left\{\sup_{t\geq 0}\inf_{[t,t+T_u]}(B_H(t)-c_1t)>q_1u\right\}\sim\begin{cases}\frac{e^{-c_1^2T/2}-c_1\sqrt{2\pi T}\Phi(-c_1\sqrt{T})}{e^{-c_1^2T/2}+c_1\sqrt{2\pi T}\Phi(c_1\sqrt{T})}e^{-2c_1q_1u}, & H=1/2,\\K_H\mathcal{F}_{2H}(TD_1)(\mathbb{C}_H^{(1)}u^{1-H})^{\frac{1}{H}-1}\overline{\Phi}(\mathbb{C}_H^{(1)}u^{1-H}), & H\neq 1/2.\end{cases}$$

Now we are ready to present our proofs.

Proof of Theorems 1.1 and 2.1. Since Theorem 1.1 follows immediately from Theorem 2.1 we prove Theorem 2.1 only.

Case (1). Assume that $t_* < t_1$. Let

$$\psi_i(T_u, u) = \mathbb{P}\left\{\sup_{t \ge 0} \inf_{[t, t + T_u]} (B_H(t) - c_i t) > q_i u\right\}, \quad i = 1, 2.$$

For $0 < \varepsilon < t_1 - t_*$ by the self-similarity of fBm we have

$$\psi_1(T_u, u) \ge \mathcal{P}_{T_u}(u) \ge \mathbb{P}\left\{\exists t \in (t_1 - \varepsilon, t_1 + \varepsilon) : \inf_{s \in [t, t + T_u/u]} V_1(t) > u^{1-H}, \inf_{s \in [t, t + T_u/u]} V_2(t) > u^{1-H}\right\}$$

$$= \mathbb{P}\left\{\exists t \in (t_1 - \varepsilon, t_1 + \varepsilon) : \inf_{s \in [t, t + T_u/u]} V_1(t) > u^{1-H}\right\},$$

where

$$V_i(t) = \frac{B_H(t)}{c_i t + q_i}, \quad i = 1, 2.$$

We have by Borel-TIS inequality, see [4] (details are in the Appendix)

(18)
$$\psi_1(T_u, u) \sim \mathbb{P}\left\{\exists t \in (t_1 - \varepsilon, t_1 + \varepsilon) : \inf_{s \in [t, t + T_u/u]} V_1(t) > u^{1-H}\right\}, \quad u \to \infty$$

implying $\mathcal{P}_{T_u}(u) \sim \psi_1(T_u, u)$ as $u \to \infty$. The asymptotics of $\psi_1(T_u, u)$ is given in Proposition 5.1, thus the claim follows.

Assume that $t_* = t_1$. We have

$$\mathbb{P}\left\{\exists t \in [t_1, \infty) : \inf_{s \in [t, t + \frac{T_u}{u}]} V_1(s) > u^{1-H}\right\} \leq \mathcal{P}_{T_u}(u)$$

$$\leq \mathbb{P}\left\{\exists t \in [t_1, \infty) : \inf_{s \in [t, t + \frac{T_u}{u}]} V_1(s) > u^{1-H}\right\}$$

$$+ \mathbb{P}\left\{\exists t \in [0, t_1] : V_2(t) > u^{1-H}\right\}.$$

From the proof of Theorem 3.1, case (4) in [25] it follows that the second term in the last line above is negligible comparing with the final asymptotics of $\mathcal{P}_{T_u}(u)$ given in (11), hence

$$\mathcal{P}_{T_u}(u) \sim \mathbb{P}\left\{\exists t \in [t_1, \infty) : \inf_{s \in [t, t + \frac{T_u}{u}]} V_1(s) > u^{1-H}\right\}, \quad u \to \infty.$$

By the same arguments as in (18) it follows that for $\varepsilon > 0$ the last probability above is equivalent with

$$\mathbb{P}\left\{\exists t \in [t_1, t_1 + \varepsilon] : \inf_{s \in [t, t + T_u/u]} V_1(s) > u^{1-H}\right\}, \quad u \to \infty.$$

Since $\mathcal{F}_1(T) = \frac{e^{-T/4} - \sqrt{\pi T}\Phi(-\sqrt{T/2})}{e^{-T/4} + \sqrt{\pi T}\Phi(\sqrt{T/2})}$, $T \ge 0$ (see [23]) applying Theorem 3.3 in [18] with parameters in the notation therein

$$\widetilde{\sigma} = \frac{t_1^H}{c_1 t + q_1}, \quad \beta_1 = 2, \quad D = \frac{1}{2t_1^{2H}}, \quad \alpha = 2H, \quad A = \frac{q_1^{H-3} H^{H-1} (1 - H)^{4-H}}{2c_1^{H-2}}$$

we obtain

$$\mathbb{P}\left\{\exists t \in [t_1, t_1 + \varepsilon] : \inf_{s \in [t, t + T_u/u]} V_1(s) > u^{1-H}\right\} \sim \frac{1}{2} K_H \mathcal{F}_{2H}(TD_1) (\mathbb{C}_H^{(1)} u^{1-H})^{\frac{1}{H} - 1} \overline{\Phi}(\mathbb{C}_H^{(1)} u^{1-H}), \quad u \to \infty$$

and the claim is established. Case $t_* \geq t_2$ follows by the same arguments.

Case (2). Define

(19)
$$Z_H(t) = \frac{B_H(t)}{\max(c_1 t + q_1, c_2 t + q_2)}, \quad t \ge 0.$$

Similarly to the proof of (18) we have by Borell-TIS inequality for $\varepsilon > 0$ as $u \to \infty$

$$\mathcal{P}_{T_u}(u) = \mathbb{P}\left\{\exists t \ge 0 : \inf_{s \in [t, t + T_u/u]} Z_H(t) > u^{1-H}\right\}$$

$$\sim \mathbb{P}\left\{\exists t \in (t_* - \varepsilon, t_* + \varepsilon) : \inf_{s \in [t, t + T_u/u]} Z_H(t) > u^{1-H}\right\} =: p(u), \quad u \to \infty.$$

Assume that H < 1/2. By "the double-sum" approach, see the proofs of Theorem 3.1, Case (3) H < 1/2 in [25] and Theorem 3.3. case i) in [18] we have as $u \to \infty$

$$(20) \ p(u) \sim \mathbb{P}\left\{\exists t \in (t_*, t_* + \varepsilon) : \inf_{s \in [t, t + \frac{T_u}{u}]} V_1(t) > u^{1-H}\right\} + \mathbb{P}\left\{\exists t \in (t_* - \varepsilon, t_*) : \inf_{s \in [t, t + \frac{T_u}{u}]} V_2(t) > u^{1-H}\right\}.$$

To compute the asymptotics of each probability in the line above we apply Theorem 3.3 in [18]. For the first probability we have in the notation therein

$$\widetilde{\sigma} = \frac{t_*^H}{c_1 t_* + q_1}, \quad \beta_1 = 1, \quad D = \frac{1}{2t_*^{2H}}, \quad \alpha = 2H < 1, \quad A = \frac{t_*^{H-1} |H(c_1 t_* + q_1) - c_1 t_*|}{(c_1 t_* + q_1)^2}$$

implying as $u \to \infty$

$$\mathbb{P}\left\{\exists t \in (t_*, t_* + \varepsilon) : \inf_{s \in [t, t + \frac{T_u}{u}]} V_1(t) > u^{1-H}\right\} \sim \mathcal{F}_{2H}(\frac{(c_1 t_* + q_1)^{\frac{1}{H}}}{2^{\frac{1}{2H}} t_*^2} T) \frac{t_*^H \mathbb{D}_H^{\frac{1}{H} - 1} u^{(1-H)(\frac{1}{H} - 2)}}{|H(c_1 t_* + q_1) - c_1 t_*| 2^{\frac{1}{2H}}} \overline{\Phi}(\mathbb{D}_H u^{1-H}).$$

Applying again Theorem 3.3 in [18] we obtain the asymptotics of the second summand and the claim follows by (20).

Assume that H = 1/2. In order to compute the asymptotics of p(u) applying Theorem 3.3 in [18] with parameters

$$\alpha = \beta_1 = \beta_2 = 1, \ A_{\pm} = \frac{q_1 - c_1 t_*}{q_1 + c_1 t_*}, \ A = \frac{q_2 - c_2 t_*}{q_2 + c_2 t_*}, \ \widetilde{\sigma} = \frac{\sqrt{t_*}}{c_1 t_* + q_1}, \ D = \frac{1}{2t_*}$$

we obtain $(d(\cdot))$ and T' are defined in (7)

$$p(u) \sim \mathcal{F}_{T'}^d \overline{\Phi}(\mathbb{D}_{1/2} \sqrt{u}), \quad u \to \infty.$$

Assume that H > 1/2. Applying Theorem 3.3 in [18] with parameters $\alpha = 2H > 1 = \beta_1 = \beta_2$ we complete the proof since

$$p(u) \sim \overline{\Phi}(\mathbb{D}_H u^{1-H}), \quad u \to \infty.$$

Proof of Proposition 2.2. Lower bound. Take $\kappa = 1 - 3H$ and recall that $\alpha = \frac{T^{2H}}{2t^{2H}}$. We have

$$\mathcal{P}_{T}(u) \geq \mathbb{P}\left\{\forall t \in [t_{*} - T/u, t_{*}]V_{2}(t) > u^{1-H} \text{ and } V_{2}(t_{*}) > u^{1-H} + \alpha u^{\kappa}\right\}$$

$$\geq \bar{C}\mathbb{P}\left\{V_{2}(t_{*}) > u^{1-H} + \alpha u^{\kappa}\right\}$$

$$\sim \bar{C}\overline{\Phi}(\mathbb{D}_{H}u^{1-H})e^{-C_{1,\alpha}u^{1-H+\kappa} - C_{2,\alpha}u^{2\kappa}}, \quad u \to \infty,$$

where \bar{C} is a fixed positive constant that does not depend on u and $C_{1,\alpha}$ and $C_{2,\alpha}$ are defined in (15). Thus, to prove the lower bound we need to show (21). Note that (21) is the same as

$$\mathbb{P}\left\{\exists t \in [t_* - T/u, t_*] : V_2(t) \le u^{1-H} \text{ and } V_2(t_*) > u^{1-H} + \alpha u^{\kappa}\right\} \le \varepsilon' \mathbb{P}\left\{V_2(t_*) > u^{1-H} + \alpha u^{\kappa}\right\},$$

with some $\varepsilon' \in (0,1)$. The last line above is equivalent with

$$\mathbb{P}\left\{\exists t \in [ut_* - T, ut_*] : B_H(t) - c_2 t \le q_2 u \text{ and } B_H(ut_*) - c_2 ut_* > q_2 u + b\alpha u^{\kappa + H}\right\}$$

$$\le \varepsilon' \mathbb{P}\left\{B_H(ut_*) - c_2 ut_* > q_2 u + b\alpha u^{\kappa + H}\right\},$$

where $b = c_2 t_* + q_2$. We have with $\varphi_u(x)$ the density of $B_H(ut_*)$ that the left part of the inequality above does not exceed

$$\mathbb{P}\left\{\exists t \in [ut_* - T, ut_*] : B_H(ut_*) - B_H(t) > b\alpha u^{\kappa + H} \text{ and } B_H(ut_*) > bu\right\}$$

$$= \int_{bu}^{\infty} \mathbb{P}\left\{\exists t \in [ut_* - T, ut_*] : x - B_H(t) > b\alpha u^{\kappa + H} | B_H(ut_*) = x\right\} \varphi_u(x) dx$$

$$\leq \int_{bu}^{bu+1} \mathbb{P}\left\{\exists t \in [ut_* - T, ut_*] : x - B_H(t) > b\alpha u^{\kappa + H} | B_H(ut_*) = x\right\} \varphi_u(x) dx + \int_{bu+1}^{\infty} \varphi_u(x) dx.$$

We also have that

$$\mathbb{P}\left\{B_H(ut_*) - c_2ut_* > q_2u\right\} = \int_{bu}^{\infty} \varphi_u(x)dx \ge \int_{bu}^{bu+1} \varphi_u(x)dx.$$

By (17) we have that $\int_{bu+1}^{\infty} \varphi_u(x)dx$ is negligible comparing with the last integral above. Thus, to prove (21) we need to show

$$\int_{bu}^{bu+1} \mathbb{P}\left\{\exists t \in [ut_* - T, ut_*] : x - B_H(t) > b\alpha u^{\kappa + H} | B_H(ut_*) = x\right\} \varphi_u(x) dx \le \varepsilon' \int_{bu}^{bu+1} \varphi_u(x) dx, \qquad u \to \infty,$$

that follows from the inequality

(22)
$$\sup_{x \in [bu, bu+1]} \mathbb{P}\left\{\exists t \in [ut_* - T, ut_*] : x - B_H(t) > b\alpha u^{\kappa + H} | B_H(ut_*) = x\right\} \le \varepsilon'', \quad u \to \infty,$$

where $\varepsilon'' \in (0,1)$ is some number. We show the line above in the Appendix, thus the lower bound holds.

Upper bound. We have by the self-similarity of fBm

$$\mathcal{P}_T(u) = \mathbb{P}\left\{\sup_{t>0} \inf_{s \in [t, t+T/u]} Z_H(s) > u^{1-H}\right\},\,$$

where Z_H is defined in (19). For $\varepsilon > 0$ by Borell-TIS inequality with $I(t_*) = (-u^{-\varepsilon} + t_*, t_* + u^{-\varepsilon})$ we have

$$\mathbb{P}\left\{\sup_{t\notin I(t_*)}\inf_{s\in[t,t+T/u]}Z_H(s)>u^{1-H}\right\}\leq \mathbb{P}\left\{\sup_{t\notin I(t_*)}Z_H(t)>u^{1-H}\right\}\leq \overline{\Phi}\left(\mathbb{D}_Hu^{1-H}\right)e^{-Cu^{2-2H-2\varepsilon}},\quad u\to\infty,$$

that is asymptotically smaller than the lower bound in (14) for sufficiently small ε . Thus, we focus on estimation of

$$q(u) := \mathbb{P}\left\{ \sup_{t \in I(t_*)} \inf_{s \in [t, t + T/u]} Z_H(s) > u^{1-H} \right\}.$$

Denote $z^2(t) = \text{Var}\{Z_H(t)\}$ and $\overline{Z}_H(t) = Z_H(t)/z(t)$. By Lemma 2.3 in [3] we have with $M = \max(z(t), z(t+T/u))$ (note, $1/M \ge \mathbb{D}_H$)

$$q(u) \leq \mathbb{P}\left\{\exists t \in I(t_*) : Z_H(t) > u^{1-H}, Z_H(t+T/u) > u^{1-H}\right\}$$

$$= \mathbb{P}\left\{\exists t \in I(t_*) : \overline{Z}_H(t) > u^{1-H}/z(t), \overline{Z}_H(t+T/u) > u^{1-H}/z(t+T/u)\right\}$$

$$\leq \mathbb{P}\left\{\exists t \in I(t_*) : \overline{Z}_H(t) > u^{1-H}/M, \overline{Z}_H(t+T/u) > u^{1-H}/M\right\}$$

$$\leq 2(1+o(1))\overline{\Phi}\left(\frac{u^{1-H}}{M}\right)\overline{\Phi}\left(\frac{u^{1-H}}{M}\sqrt{\frac{1-r(t,t+T/u)}{1+r(t,t+T/u)}}\right)$$

$$\leq 2(1+o(1))\overline{\Phi}\left(\frac{u^{1-H}}{M}\right)\overline{\Phi}\left(\mathbb{D}_H u^{1-H}\sqrt{\frac{1-r(t,t+T/u)}{2}}\right),$$

$$(23)$$

where r is the correlation function of Z_H . Since $r(t,s) = corr(B_H(t), B_H(s))$ we have for all $t \in I(t_*)$

$$1 - r(t, t + T/u) = \frac{T^{2H}}{2t_*^{2H}}u^{-2H} + O\left(u^{-2H}(|t - t_*| + |t + T/u - t_*|) + u^{-2}\right), \quad u \to \infty$$

implying

$$\mathbb{D}_H u^{1-H} \sqrt{\frac{1 - r(t, t + T/u)}{2}} = u^{1-2H} \frac{T^H \mathbb{D}_H}{2t_*^H} + O(u^{1-2H} (|t - t_*| + |t + T/u - t_*|) + u^{-1}), \quad u \to \infty.$$

Thus, by (17) we obtain

$$(24) \ \overline{\Phi} \left(\mathbb{D}_H u^{1-H} \sqrt{\frac{1 - r(t, t + T/u)}{2}} \right) \le \overline{\Phi} \left(u^{1-2H} \frac{T^H \mathbb{D}_H}{2t_*^H} \right) e^{Cu^{2-4H}(|t-t_*| + |t+T/u - t_*|)}, \quad u \to \infty.$$

Next we have as $u \to \infty$ for some $C_1 > 0$

$$\overline{\Phi}\left(\frac{u^{1-H}}{M}\right) \sim \overline{\Phi}(\mathbb{D}_H u^{1-H}) e^{-C_1 u^{2-2H}(|t-t_*|+|t+T/u-t_*|)}$$

and by (24) we have for all $t \in I(t_*)$ and large u

$$\overline{\Phi}\left(\frac{u^{1-H}}{M}\right)\overline{\Phi}\left(\mathbb{D}_{H}u^{1-H}\sqrt{\frac{1-r(t,t+T/u)}{2}}\right)\leq \overline{\Phi}\left(\mathbb{D}_{H}u^{1-H}\right)\overline{\Phi}\left(u^{1-2H}\frac{T^{H}\mathbb{D}_{H}}{2t_{*}^{H}}\right)e^{(Cu^{2-4H}-C_{1}u^{2-2H})(|t-t_{*}|+|t+T/u-t_{*}|)}$$

and the claim follows from the line above and (23).

Proof of Lemma 3.1. Lower bound. We have

$$\sup_{t \in \mathbb{R}} \inf_{s \in [t, t+L]} e^{\sqrt{2}B(s) - |s| + h(s)} \ge \inf_{s \in [0, L]} e^{\sqrt{2}B(s) - (1+a)s} \ge e^{-(1+a)L} \inf_{s \in [0, L]} e^{\sqrt{2}B(s)} \stackrel{d}{=} e^{-(1+a)L} e^{-\sup_{s \in [0, L]} \sqrt{2}B(s)},$$

where the symbol $\stackrel{d}{=}$ means equality in distribution between two random variables. Taking expectations of both sides in the line above we obtain

$$\mathcal{F}_L^h \ge e^{-L(1+a)} \mathbb{E} \left\{ e^{-\sup_{s \in [0,L]} \sqrt{2}B(s)} \right\},\,$$

and our next step is to calculate the expectation above. It is known (see, e.g., Chapter 11.1 in [4]) that

$$\mathbb{P}\left\{\sup_{s\in[0,L]}\sqrt{2}B(s)>x\right\}=2\mathbb{P}\left\{\sqrt{2}B(L)>x\right\}=2\overline{\Phi}\left(\frac{x}{\sqrt{2L}}\right),\ x>0$$

hence we obtain that $\frac{e^{-x^2/4L}}{\sqrt{\pi L}}$, x > 0 is the density of $\sup_{s \in [0,L]} \sqrt{2}B(s)$. Thus, we have

$$\mathbb{E}\left\{e^{-\sup_{s\in[0,L]}\sqrt{2}B(s)}\right\} = \int_{0}^{\infty} e^{-x} \frac{e^{-x^{2}/4L}}{\sqrt{\pi L}} dx = \frac{e^{L}}{\sqrt{\pi L}} \int_{0}^{\infty} e^{-(\frac{x}{2\sqrt{L}} + \sqrt{L})^{2}} dx = \frac{2e^{L}}{\sqrt{\pi}} \int_{\sqrt{L}}^{\infty} e^{-z^{2}} dz = 2e^{L} \overline{\Phi}(\sqrt{2L}).$$

and combining all calculations above we obtain

$$\mathcal{F}_L^h \ge 2e^{-La}\overline{\Phi}(\sqrt{2L}), \quad L \ge 0.$$

On the other hand we have

$$\sup_{t \in \mathbb{R}} \inf_{s \in [t, t+L]} e^{\sqrt{2}B(s) - |s| + h(s)} \ge \inf_{s \in [-L, 0]} e^{\sqrt{2}B(s) - (1+b)|s|} \stackrel{d}{=} \inf_{s \in [0, L]} e^{\sqrt{2}B(s) - (1+b)s},$$

and estimating $\inf_{s\in[0,L]}e^{\sqrt{2}B(s)-(1+b)s}$ as above we have $\mathcal{F}_L^h\geq 2e^{-Lb}\overline{\Phi}(\sqrt{2L}),\ L\geq 0$, that completes the proof of the lower bound.

Upper bound. Note that $\mathcal{F}_{2H}^L \leq \mathcal{F}_{2H}^0$ and hence since a BM has independent branches for positive and negative time we have with B_* an independent BM

$$\begin{split} \mathcal{F}^L_{2H} &\leq \mathbb{E}\left\{\sup_{t \in \mathbb{R}} e^{\sqrt{2}B(t) - h(t)}\right\} &= \mathbb{E}\left\{\max\left(\sup_{t \geq 0} e^{\sqrt{2}B(t) - (a+1)t}, \sup_{t \leq 0} e^{\sqrt{2}B(t) - (b+1)|t|}\right)\right\} \\ &= \mathbb{E}\left\{\max\left(\sup_{t \geq 0} e^{\sqrt{2}B(t) - (a+1)t}, \sup_{t \geq 0} e^{\sqrt{2}B^*(t) - (b+1)t}\right)\right\} = \mathbb{E}\left\{e^{\max(\xi_a, \xi_b)}\right\}, \end{split}$$

where ξ_a and ξ_b are exponential random variables with survival functions $e^{-(a+1)x}$ and $e^{-(b+1)x}$, respectively, see [2]. Since ξ_a and ξ_b have exponential distributions the last expectation above is $1 + \frac{1}{a} + \frac{1}{b} - \frac{1}{a+b+1}$ and the claim follows.

Proof of Lemma 3.2. First we have

$$\mathbb{E}\left\{\sup_{t\in\mathbb{R}\backslash[-M,M]}\inf_{s\in[t,t+L]}e^{\sqrt{2}B(s)-|s|+h(s)}\right\}\leq\mathbb{E}\left\{\sup_{s\in[M,\infty)}e^{\sqrt{2}B(s)-(a+1)s}\right\}+\mathbb{E}\left\{\sup_{s\in(-\infty,-M]}e^{\sqrt{2}B(s)-(b+1)|s|}\right\}.$$

Later on we work with the first expectation above. We have

$$\mathbb{E}\left\{\sup_{s\in[M,\infty)}e^{\sqrt{2}B(s)-(1+a)s}\right\}$$

$$=\int_{\mathbb{R}}e^{x}\mathbb{P}\left\{\sup_{s\in[M,\infty)}(\sqrt{2}B(s)-(1+a)s)>x\right\}dx$$

$$=\int_{\mathbb{R}}e^{x}\mathbb{P}\left\{\sup_{s\in[M,\infty)}(\sqrt{2}(B(s)-B(M))-(1+a)(s-M))>x+M(1+a)-\sqrt{2}B(M)\right\}dx.$$

Since a BM has independent increments we have with B^* an independent BM that the last integral above equals

$$\int_{\mathbb{R}} e^{x} \mathbb{P} \left\{ \sup_{s \in [0, \infty)} (\sqrt{2}B(s) - (1+a)s) > x + M(1+a) - \sqrt{2M}B^{*}(1) \right\} dx$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \int_{\mathbb{R}} e^{x} e^{-z^{2}/2} \mathbb{P} \left\{ \sup_{s \in [0, \infty)} (\sqrt{2}B(s) - (1+a)s) > x + M(1+a) - \sqrt{2M}z \right\} dx dz.$$

We know that $\mathbb{P}\left\{\sup_{t\geq 0}(B(t)-ct)>x\right\}=\min(1,e^{-2cx})$ for c>0 and $x\in\mathbb{R}$, thus the expression above equals

$$\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \int_{\mathbb{R}} e^{x-z^2/2} \min(1, e^{-(1+a)(x+M(1+a)-\sqrt{2M}z)}) dx dz$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \int_{\frac{(1+a)M+x}{6M^2}}^{\infty} e^{x-z^2/2} dz dx + \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \int_{-\infty}^{\frac{(1+a)M+x}{\sqrt{2M}}} e^{x-z^2/2-(1+a)(x+M(1+a)-\sqrt{2M}z)} dz dx$$

$$= \int_{\mathbb{R}} e^{x} \overline{\Phi}(\frac{(1+a)M+x}{\sqrt{2M}}) dx + \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-ax} \int_{-\infty}^{\frac{(1+a)M+x}{\sqrt{2M}}} e^{-\frac{(z-\sqrt{2M}(1+a))^{2}}{2}} dz dx$$

$$= \int_{\mathbb{R}} e^{x} \overline{\Phi}(\frac{(1+a)M+x}{\sqrt{2M}}) dx + \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-ax} \int_{-\infty}^{\frac{-(1+a)M+x}{\sqrt{2M}}} e^{-\frac{z^{2}}{2}} dz dx$$

$$= \int_{\mathbb{R}} e^{x} \overline{\Phi}(\frac{(1+a)M+x}{\sqrt{2M}}) dx + \int_{\mathbb{R}} e^{-ax} \Phi(\frac{-(1+a)M+x}{\sqrt{2M}}) dx.$$

Integrating the first integral above by parts we have

$$\int_{\mathbb{R}} e^{x} \overline{\Phi}(\frac{(1+a)M+x}{\sqrt{2M}}) dx = -\int_{\mathbb{R}} (\overline{\Phi}(\frac{(1+a)M+x}{\sqrt{2M}}))' e^{x} dx = \frac{1}{\sqrt{2\pi}\sqrt{2M}} \int_{\mathbb{R}} e^{-\frac{((1+a)M+x)^{2}}{4M}} e^{x} dx \\
= \frac{e^{-aM}}{\sqrt{2\pi}\sqrt{2M}} \int_{\mathbb{R}} e^{-\frac{((a-1)M+x)^{2}}{4M}} dx = e^{-aM}.$$

For the second integral we have similarly

$$\int_{\mathbb{R}} e^{-ax} \Phi(\frac{-(1+a)M+x}{\sqrt{2M}}) dx = -\frac{1}{a} \int_{\mathbb{R}} \Phi(\frac{-(1+a)M+x}{\sqrt{2M}})' e^{-ax} dx
= \frac{1}{a} \frac{1}{\sqrt{2\pi}\sqrt{2M}} \int_{\mathbb{R}} e^{-\frac{(-(1+a)M+x)^2}{4M} - ax} dx = \frac{e^{-aM}}{a\sqrt{2\pi}\sqrt{2M}} \int_{\mathbb{R}} e^{-\frac{((1-a)M+x)^2}{4M}} dx = \frac{e^{-aM}}{a}.$$

Summarizing all calculations above we obtain

$$\mathbb{E}\left\{\sup_{t\in[M,\infty)}e^{\sqrt{2}B(t)-(1+a)t}\right\} = e^{-aM}\left(1+\frac{1}{a}\right).$$

By the same approach and the symmetry of BM around zero we have

$$\mathbb{E}\left\{\sup_{t\in(-\infty,-M]}e^{\sqrt{2}B(t)-(1+b)|t|}\right\} = e^{-bM}\left(1+\frac{1}{b}\right)$$

and hence combining both equations above with the first inequality in the proof we obtain the claim. \Box

Proof of Lemma 3.3. From [29] it follows, that for any $L \geq 0$

(25)
$$\mathcal{F}_{2H}(L) = \mathbb{E}\left\{\frac{\sup_{t \in \mathbb{R}} \inf_{s \in [t, t+L]} e^{W(s)}}{\int_{\mathbb{R}} e^{W(t)} dt}\right\},$$

later on we use this formula in the proof. Observe that $\sup_{t \in \mathbb{R}} \inf_{s \in [t,t+L]} e^{W(s)} \ge \inf_{s \in [0,L]} e^{W(s)}$, hence

$$\mathcal{F}_{2H}(L) \ge \mathbb{E}\left\{\frac{\inf\limits_{s \in [0,L]} e^{W(s)}}{\int\limits_{\mathbb{R}} e^{W(t)} dt}\right\} \ge e^{-L^{2H}} \mathbb{E}\left\{\frac{e^{-\sqrt{2} \sup\limits_{s \in [0,L]} B_H(s)}}{\int\limits_{\mathbb{R}} e^{W(t)} dt}\right\}.$$

Let $\xi = \sup_{s \in [0,L]} B_H(s)$, (Ω, \mathbb{P}) be the general probability space and $\Omega_m = \{\omega \in \Omega : \xi(\omega) < m\}$ for m > 0.

The last expectation above equals

$$\mathbb{E}\left\{\frac{e^{-\sqrt{2}\xi}}{\int\limits_{\mathbb{R}} e^{W(t)}dt}\right\} = \int\limits_{\Omega} \frac{e^{-\sqrt{2}\xi(\omega)}}{\int\limits_{\mathbb{R}} e^{\sqrt{2}B_{H}(t,\omega)-|t|^{2H}}dt}d\mathbb{P}(\omega)$$

$$\geq \int\limits_{\Omega_{m}} \frac{e^{-\sqrt{2}\xi(\omega)}}{\int\limits_{\mathbb{R}} e^{\sqrt{2}B_{H}(t,\omega)-|t|^{2H}}dt}d\mathbb{P}(\omega)$$

$$\geq \mathbb{P}\left\{\Omega_{m}\right\} e^{-\sqrt{2}m}\int\limits_{\Omega_{m}} \frac{1}{\int\limits_{\mathbb{R}} e^{W(t)}dt}d\mathbb{P}(\omega)$$

$$\geq e^{-\sqrt{2}m}\mathbb{P}\left\{\xi < m\right\}\mathbb{E}\left\{\frac{1}{\int\limits_{\mathbb{R}} e^{W(t)}dt}\right\}.$$

Next taking $m = nL^H$ by the self-similarity of fBm we have that

$$e^{-\sqrt{2}m}\mathbb{P}\left\{\xi < m\right\} = e^{-\sqrt{2}nL^H}\mathbb{P}\left\{\sup_{s \in [0,L]} B_H(s) < nL^H\right\} = e^{-\sqrt{2}nL^H}\mathbb{P}\left\{\sup_{s \in [0,1]} B_H(s) < n\right\}.$$

Taking sup with respect to n over $(0, \infty)$ we have

$$\mathcal{F}_{2H}(L) \ge \mathbb{E}\left\{\frac{1}{\int\limits_{\mathbb{R}} e^{W(t)} dt}\right\} e^{-L^{2H}} \sup_{n>0} \left(e^{-\sqrt{2}nL^{H}} \mathbb{P}\left\{\sup_{s \in [0,1]} B_{H}(s) < n\right\}\right)$$

and hence to complete the proof we need to show that the expectation in the expression above is a finite positive constant. Since the classical Pickands constant is finite (see, e.g., [3, 4, 7, 29]) we have

$$0 < \mathbb{E}\left\{\frac{1}{\int\limits_{\mathbb{R}} e^{W(t)} dt}\right\} \le \mathbb{E}\left\{\frac{\sup_{t \in \mathbb{R}} e^{W(t)}}{\int\limits_{\mathbb{R}} e^{W(t)} dt}\right\} = \mathbb{H}_{2H} \in (0, \infty).$$

Proof of Lemma 3.4. By (25) we have that

$$\begin{split} & \left| \mathcal{F}_{2H}(L) - \mathbb{E} \left\{ \frac{\sup_{t \in [-M,M]} \inf_{s \in [t,t+L]} e^{W(s)}}{\int\limits_{[-M,M]} e^{W(t)} dt} \right\} \right| \\ & = \left| \left(\mathbb{E} \left\{ \frac{\sup_{t \in \mathbb{R}} \inf_{s \in [t,t+L]} e^{W(s)}}{\int\limits_{\mathbb{R}} e^{W(t)} dt} \right\} - \mathbb{E} \left\{ \frac{\sup_{t \in [-M,M]} \inf_{s \in [t,t+L]} e^{W(s)}}{\int\limits_{\mathbb{R}} e^{W(t)} dt} \right\} \right) \\ & + \left(\mathbb{E} \left\{ \frac{\sup_{t \in [-M,M]} \inf_{s \in [t,t+L]} e^{W(s)}}{\int\limits_{\mathbb{R}} e^{W(t)} dt} \right\} - \mathbb{E} \left\{ \frac{\sup_{t \in [-M,M]} \inf_{s \in [t,t+L]} e^{W(s)}}{\int\limits_{[-M,M]} e^{W(t)} dt} \right\} \right) \right| \\ & \leq \mathbb{E} \left\{ \frac{\sup_{t \in \mathbb{R} \setminus [-M,M]} e^{W(t)}}{\int\limits_{\mathbb{R}} e^{W(t)} dt} \right\} + \mathbb{E} \left\{ \sup_{t \in [-M,M]} e^{W(t)} \frac{\int\limits_{\mathbb{R} \setminus [-M,M]} e^{W(t)} dt}{\int\limits_{[-M,M]} e^{W(t)} dt} \int\limits_{[-M,M]} e^{W(t)} dt} \right\}. \end{split}$$

As follows from Section 4 in [29], the last line above does not exceed $e^{-c'M^{2H}}$, and the claim holds. \Box

6. Appendix

Proof of (18). To establish the claim we need to show, that

$$\mathbb{P}\left\{\exists t \in \mathbb{R} \setminus [t_1 - \varepsilon, t_1 + \varepsilon] : \inf_{s \in [t, t + T/u]} V_1(s) > u^{1-H}\right\} = o(\psi_1(T_u, u)), \quad u \to \infty.$$

Applying Borell-TIS inequality (see, e.g., [4]) we have as $u \to \infty$

$$\mathbb{P}\left\{\exists t \in \mathbb{R} \backslash [t_1 - \varepsilon, t_1 + \varepsilon] : \inf_{s \in [t, t + T/u]} V_1(s) > u^{1-H}\right\} \leq \mathbb{P}\left\{\exists t \in \mathbb{R} \backslash [t_1 - \varepsilon, t_1 + \varepsilon] : V_1(t) > u^{1-H}\right\} \leq e^{-\frac{(u^{1-H} - \widetilde{M})^2}{2m^2}},$$

where

$$\widetilde{M} = \mathbb{E}\left\{\sup_{\exists t \in \mathbb{R} \setminus [t_1 - \varepsilon, t_1 + \varepsilon]} V_1(t)\right\} < \infty, \quad m^2 = \max_{\exists t \in \mathbb{R} \setminus [t_1 - \varepsilon, t_1 + \varepsilon]} \operatorname{Var}\{V_1(t)\}.$$

Since $Var\{V_1(t)\}$ achieves its unique maxima at t_1 we obtain by (17) that

$$e^{-\frac{(u^{1-H}-\widetilde{M})^2}{2m^2}} = o(\mathbb{P}\{V_1(t_1) < u^{1-H}\}), \quad u \to \infty$$

and the claim follows from the asymptotics of $\psi_1(T_u, u)$ given in Proposition 5.1.

Proof of (22). Define $X_{x,u}(t) = x - B_H(t)|B_H(ut_*) = x$, $t \in [ut_* - T, u]$. To calculate the covariance and expectation of $X_{x,u}$ we use the formulas

$$cov((B,C)|A=x) = cov(B,C) - \frac{cov(A,B)cov(A,C)}{Var\{A\}} \quad and \quad \mathbb{E}\left\{B|A=x\right\} = x \cdot \frac{cov(A,B)}{Var\{A\}}$$

where A,B and C are centered Gaussian random variables and $x\in\mathbb{R}$. We have for $x\in[bu,bu+1]$ and $t,s\in[ut_*-T,ut_*]$ with $v=ut_*,\,y=1-\frac{t}{v}$ and $z=1-\frac{s}{v}$ as $u\to\infty$

$$\begin{aligned}
&= \frac{t^{2H} + s^{2H} - |t - s|^{2H}}{2} - \frac{(t^{2H} + v^{2H} - |t - v|^{2H})(s^{2H} + v^{2H} - |s - v|^{2H})}{4v^{2H}} \\
&= \frac{v^{2H}}{4} \left(2(\frac{t}{v})^{2H} + 2(\frac{s}{v})^{2H} - 2|\frac{t}{v} - \frac{s}{v}|^{2H} - ((\frac{t}{v})^{2H} + 1 - |\frac{t}{v} - 1|^{2H})((\frac{s}{v})^{2H} + 1 - |\frac{s}{v} - 1|^{2H}) \right) \\
&= \frac{v^{2H}}{4} \left(2(1 - y)^{2H} + 2(1 - z)^{2H} - 2|y - z|^{2H} - ((1 - y)^{2H} + 1 - y^{2H})((1 - z)^{2H} + 1 - z^{2H}) \right) \\
&= \frac{v^{2H}}{4} \left(2 - 4Hy + 2 - 4Hz + O(y^2 + z^2) - 2|y - z|^{2H} - (2 - 2Hy - y^{2H} + O(y^2))(2 - 2Hz - z^{2H} + O(z^2)) \right) \\
&= \frac{v^{2H}}{4} \left(2y^{2H} + 2z^{2H} - 2|y - z|^{2H} + O(y^2 + z^2 + z^{2H}y^{2H}) \right) \\
&= \frac{v^{2H}}{4} \left(2y^{2H} + 2z^{2H} - 2|y - z|^{2H} + O(y^2 + z^2 + z^{2H}y^{2H}) \right) \\
&= (1 + o(1)) \frac{(ut_* - t)^{2H} + (ut_* - s)^{2H} - |t - s|^{2H}}{2}.
\end{aligned}$$

For the expectation we have as $u \to \infty$

$$\mathbb{E}\left\{X_{x,u}(t)\right\} = x\left(1 - \frac{v^{2H} + t^{2H} - |v - t|^{2H}}{2v^{2H}}\right) = \frac{x}{2}\left(1 - (t/v)^{2H} + (1 - t/v)^{2H}\right)$$

$$\leq \frac{1}{2}(bu+1)(1-(1-y)^{2H}+y^{2H})$$

$$\leq (bu/2+1)(1-1+2Hy-o(y)+y^{2H})$$

$$\leq Hbuy + \frac{1}{2}buy^{2H} + o(1).$$

From the line above it follows that for some $C_* > 0$, H < 1/2, $x \in [bu, bu + 1]$ and $t \in [ut_* - T, ut_*]$

$$\mathbb{E}\left\{X_{x,u}(t)\right\} \le C_* + \frac{u^{1-2H}b}{2t_*^{2H}}(ut_* - t)^{2H}.$$

We have

$$\sup_{x \in [bu, bu+1]} \mathbb{P} \left\{ \exists t \in [ut_* - T, ut_*] : X_{x,u}(t) > u^{H+\kappa} \alpha b \right\}$$

$$= \sup_{x \in [bu, bu+1]} \mathbb{P} \left\{ \exists t \in [ut_* - T, ut_*] : X_{x,u}(t) - \mathbb{E} \left\{ X_{x,u}(t) \right\} > u^{H+\kappa} \alpha b - \mathbb{E} \left\{ X_{x,u}(t) \right\} \right\}$$

$$\leq \mathbb{P} \left\{ \exists t \in [0, T] : Y_u(t) + f(t) > 0 \right\},$$

where $Y_u(t) = X_{x,u}(ut_* - T + t) - \mathbb{E} \{X_{x,u}(ut_* - T + t)\}$, $t \in [0,T]$ and f(t) is the linear function such that $f(T) = C_1$ and $f(0) = -C_* < 0$. Next we have by (26) for all large u and $t, s \in [0,T]$

$$\mathbb{E}\left\{ (Y_u(t) + f(t) - Y_u(s) - f(s))^2 \right\}$$

$$= \mathbb{E}\left\{ (Y_u(t) - Y_u(s))^2 \right\} + C(t - s)^2$$

$$\leq C_1 \left((ut_* - t)^{2H} + (ut_* - s)^{2H} - (ut_* - t)^{2H} - (ut_* - s)^{2H} + |t - s|^{2H} \right) + C(t - s)^2$$

$$\leq 2|t - s|^{2H}.$$

Thus, by Proposition 9.2.4 in [4] the family $Y_u(t) + f(t)$, u > 0, $t \in [0,T]$ is tight in $\mathcal{B}(C([0,T]))$. As follows from (26), it holds that $\{Y_u(t) + f(t)\}_{t \in [0,T]}$ converges to $\{B_H(t) + f(t)\}_{t \in [0,T]}$ in the sense of convergence of finite-dimensional distributions as $u \to \infty$. Hence by Theorems 4 and 5 in Chapter 5 in [32] the tightness and convergence of finite-dimensional distributions imply weak convergence

$${Y_u(t) + f(t)}_{t \in [0,T]} \Rightarrow {B(t) + f(t)}_{t \in [0,T]}.$$

Since the functional $F(g) = \sup_{t \in [0,T]} g(t)$ is continuous in the uniform metric we obtain

$$\mathbb{P}\left\{\exists t\in[0,T]:Y_u(t)+f(t)>0\right\}\to\mathbb{P}\left\{\exists t\in[0,T]:B_H(t)+f(t)>0\right\},\ \ u\to\infty.$$

Thus, to prove the claim it is enough to show that

(27)
$$\mathbb{P}\left\{\exists t \in [0,T] : B_H(t) + f(t) > 0\right\} < 1.$$

We have for some large m with l(s) the density of $B_H(T)$

$$\mathbb{P}\left\{\sup_{t\in[0,T]}(B_{H}(t)+f(t))<0\right\} \geq \mathbb{P}\left\{\sup_{t\in[0,T]}(B_{H}(t)+f(t))<0 \text{ and } B_{H}(T)<-m\right\} \\
= \int_{-\infty}^{-m}\mathbb{P}\left\{\sup_{t\in[0,T]}(B_{H}(t)+f(t))<0|B_{H}(T)=s\right\}l(s)ds.$$
(28)

Define process $\widetilde{B}_s(t) = B_H(t) + f(t)|B_H(T) = s, \ t \in [0,T]$. We have for s < -m and $t \in [0,T]$

$$\mathbb{E}\left\{\widetilde{B}_s(t)\right\} = f(t) + s \frac{t^{2H} + T^{2H} - |T - t|^{2H}}{2T^{2H}} < -C_1/2, \quad \operatorname{Var}\{\widetilde{B}_s(t)\} = t^{2H} - \frac{(T^{2H} + t^{2H} - |t - s|^{2H})^2}{4T^{2H}} < C_2$$

and thus

$$\mathbb{P}\left\{\sup_{t\in[0,T]}(B_H(t)+f(t))<0|B_H(T)=s\right\}\geq\mathbb{P}\left\{\sup_{t\in[0,T]}\left(\widetilde{B}_s(t)-\mathbb{E}\left\{\widetilde{B}_s(t)\right\}\right)< C_1/2\right\}.$$

The last probability above is positive for any s < -m, see Chapters 10 and 11 in [33] and hence the integral in (28) is positive implying

$$\mathbb{P}\left\{ \sup_{t \in [0,T]} (B_H(t) + f(t)) < 0 \right\} > 0.$$

Consequently (27) holds and the claim is established.

Acknowledgement: Grigori Jasnovidov was supported by Ministry of Science and Higher Education of the Russian Federation grant 075-15-2022-289.

References

- [1] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling extremal events*, volume 33 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1997. For insurance and finance.
- [2] K. Dębicki and M. Mandjes. Queues and Lévy fluctuation theory. Springer, 2015.
- [3] J. Pickands, III. Upcrossing probabilities for stationary Gaussian processes. *Trans. Amer. Math. Soc.*, 145:51–73, 1969.
- [4] V. I. Piterbarg. Twenty Lectures About Gaussian Processes. Atlantic Financial Press London New York, 2015.
- [5] V. I. Piterbarg. Asymptotic methods in the theory of Gaussian processes and fields, volume 148 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 1996. Translated from the Russian by V.V. Piterbarg, revised by the author.
- [6] V. I. Piterbarg and V. R. Fatalov. The Laplace method for probability measures in Banach spaces. *Uspekhi Mat. Nauk*, 50(6(306)):57–150, 1995.
- [7] A. B. Dieker. Extremes of Gaussian processes over an infinite horizon. *Stochastic Process. Appl.*, 115(2):207–248, 2005.
- [8] G. Jasnovidov. Approximation of ruin probability and ruin time in discrete Brownian risk models. Scandinavian Actuarial Journal, 2020.
- [9] I. A. Kozik and V. I. Piterbarg. High excursions of Gaussian nonstationary processes in discrete time. Fundam. Prikl. Mat., 22(2):159–169, 2018.
- [10] G. Jasnovidov. Simultaneous ruin probability for two-dimensional fractional Brownian motion risk process over discrete grid, with supplements. arXiv:2002.04928, 2020.
- [11] Krzysztof Dębicki and Grigori Jasnovidov. Extremes of reflecting Gaussian processes on discrete grid. arXiv:2206.14712, 2022.
- [12] Krzysztof Bisewski and Grigori Jasnovidov. On the speed of convergence of discrete Pickands constants to continuous ones. arXiv:2108.00756, 2021.
- [13] Krzysztof Bisewski and Grigori Jasnovidov. On the speed of convergence of Piterbarg constants. arXiv:2209.13972, 2022.
- [14] K. Dębicki and P. Liu. Extremes of stationary Gaussian storage models. Extremes, 19(2):273–302, 2016.

- [15] Krzysztof Dębicki. Asimptotics of supremum of scaled Brownian motion. 1991.
- [16] K. Dębicki, E. Hashorva, and Peng Liu. Extremes of γ -reflected Gaussian processes with stationary increments.
- [17] K. Dębicki, P. Lui, and Z. Michna. Sojourn times of Gaussian processes with trend. *Journal of Theoretical Probability*, 2019.
- [18] K. Dębicki, E. Hashorva, and L. Ji. Parisian ruin over a finite-time horizon. Science China, 2016.
- [19] L. Ji. On the cumulative Parisian ruin of multi-dimensional Brownian motion models. arXiv:1811.10110, 2018.
- [20] K. Dębicki and G. Sikora. Finite time asymptotics of fluid and ruin models: Multiplexed fractional Brownian motions case. *APPLICATIONES MATHEMATICAE*, 2011.
- [21] K. Dębicki, E. Hashorva, L. Ji, and T. Rolski. Extremal behaviour of hitting a cone by correlated Brownian motion with drift. *Accepted for publication in Stoch. Proc. Appl.*, 2018.
- [22] Long Bai. Asymptotics of Parisian ruin of Brownian motion risk model over an infinite-time horizon. Scandinavian Actuarial Journal, 2018.
- [23] K. Dębicki, E. Hashorva, and L. Ji. Parisian ruin of self-similar Gaussian risk processes. *J. Appl. Probab.*, 52(3):688–702, 2015.
- [24] R. Loeffen, I. Czarna, and Z. Palowski. Parisian ruin probability for spectrally negative Lévy processes. *Bernoulli*, 2013.
- [25] L. Ji and Stephan R. Ruin problem of a two-dimensional fractional Brownian motion risk process. Stoch. Models, 34(1):73–97, 2018.
- [26] Grigori Jasnovidov. Sojourn ruin of a two-dimensional fractional brownian motion risk process. $arXiv:2107.11322,\ 2021.$
- [27] Thomas Mikosch. *Non-life insurance mathematics*. Universitext. Springer-Verlag, Berlin, 2004. An introduction with stochastic processes.
- [28] K. Dębicki and E. Hashorva. Approximation of supremum of max-stable stationary processes and Pickands constants. *Springer Science*, 2019.
- [29] A. B. Dieker and B. Yakir. On asymptotic constants in the theory of extremes for Gaussian processes. Bernoulli, 20(3):1600–1619, 2014.
- [30] Krzysztof Dębicki, Sebastian Engelke, and Enkelejd Hashorva. Generalized Pickands constants and stationary max-stable processes. *Extremes*, 20(3):493–517, 2017.
- [31] A. B. Dieker and M. Mandjes. On spectral simulation of fractional Brownian motion. *Probab. Engrg. Inform. Sci.*, 17(3):417–434, 2003.
- [32] Bylinskii A.V. and Shiryaev A.N. Theory of stochastic processes (in Russian). *M.PHIZMATLIT*, 2005.
- [33] M. Lifshits. Lectures on Gaussian processes (in Russian). Lan, St. Peterburg-Moscow-Krasnodar, 2016.

GRIGORI JASNOVIDOV, ST. PETERSBURG DEPARTMENT OF STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, ST. PETERSBURG, RUSSIA

 $Email\ address: {\tt griga1995@yandex.ru}$

Aleksandr Shemendyuk, Department of Actuarial Science, University of Lausanne,, UNIL-Dorigny, 1015 Lausanne, Switzerland

 $Email\ address: {\tt Aleksandr.Shemendyuk@unil.ch}$