

A NOTE ON \mathcal{F}_n -MULTIPLE ZETA VALUES

MASATAKA ONO, KOSUKE SAKURADA AND SHIN-ICHIRO SEKI

ABSTRACT. For several evaluations of special values and several relations known only in \mathcal{A}_n -multiple zeta values or \mathcal{S}_n -multiple zeta values, we prove that they are uniformly valid in \mathcal{F}_n -multiple zeta values for both the case where $\mathcal{F} = \mathcal{A}$ and $\mathcal{F} = \mathcal{S}$. In particular, the Bowman–Bradley type theorem and sum formulas for \mathcal{S}_2 -multiple zeta values are proved.

1. INTRODUCTION

We call a tuple of positive integers $\mathbf{k} = (k_1, \dots, k_r)$ an *index*. We call $\text{wt}(\mathbf{k}) := k_1 + \dots + k_r$ (resp. $\text{dep}(\mathbf{k}) := r$) the *weight* (resp. *depth*) of \mathbf{k} . If the condition $k_r \geq 2$ is satisfied, then we state that the index $\mathbf{k} = (k_1, \dots, k_r)$ is *admissible*. For an admissible index $\mathbf{k} = (k_1, \dots, k_r)$, the *multiple zeta value* (MZV) $\zeta(\mathbf{k})$ and the *multiple zeta-star value* (MZSV) $\zeta^*(\mathbf{k})$ are defined by

$$\zeta(\mathbf{k}) := \sum_{0 < n_1 < \dots < n_r} \frac{1}{n_1^{k_1} \dots n_r^{k_r}}, \quad \zeta^*(\mathbf{k}) := \sum_{1 \leq n_1 \leq \dots \leq n_r} \frac{1}{n_1^{k_1} \dots n_r^{k_r}}.$$

These series are convergent. We set $\zeta(\emptyset) = \zeta^*(\emptyset) = 1$ for the empty index \emptyset (= the empty tuple).

First we recall the definition of \mathcal{A}_n -multiple zeta(-star) values (\mathcal{A}_n -MZ(S)Vs) introduced by Rosen; see [Ro, Se]. For a positive integer n , set

$$\mathcal{A}_n := \prod_p \mathbb{Z}/p^n \mathbb{Z} \Big/ \bigoplus_p \mathbb{Z}/p^n \mathbb{Z},$$

where p runs over all prime numbers. For an index $\mathbf{k} = (k_1, \dots, k_r)$, the \mathcal{A}_n -MZV $\zeta_{\mathcal{A}_n}(\mathbf{k})$ and the \mathcal{A}_n -MZSV $\zeta_{\mathcal{A}_n}^*(\mathbf{k})$ are defined by

$$\zeta_{\mathcal{A}_n}(\mathbf{k}) := \left(\sum_{0 < n_1 < \dots < n_r < p} \frac{1}{n_1^{k_1} \dots n_r^{k_r}} \text{ mod } p^n \right)_p,$$

$$\zeta_{\mathcal{A}_n}^*(\mathbf{k}) := \left(\sum_{1 \leq n_1 \leq \dots \leq n_r \leq p-1} \frac{1}{n_1^{k_1} \dots n_r^{k_r}} \text{ mod } p^n \right)_p$$

This research was supported in part by JSPS KAKENHI Grant Numbers JP16H06336 and 18J00151.

as elements of \mathcal{A}_n . We also set $\zeta_{\mathcal{A}_n}(\emptyset) = \zeta_{\mathcal{A}_n}^*(\emptyset) = 1$.

Next we recall the definition of *t-adic symmetric multiple zeta values* ($\widehat{\mathcal{S}}$ -MZVs) introduced by Jarossary [J2]. Let t be an indeterminate. For $\bullet \in \{*, \mathfrak{m}\}$ and an index $\mathbf{k} = (k_1, \dots, k_r)$, set

$$\begin{aligned} \zeta_{\widehat{\mathcal{S}}}^\bullet(\mathbf{k}) &= \sum_{i=0}^r (-1)^{k_{i+1} + \dots + k_r} \zeta^\bullet(k_1, \dots, k_i) \\ &\times \sum_{l_{i+1}, \dots, l_r \geq 0} \left[\prod_{j=i+1}^r \binom{k_j + l_j - 1}{l_j} \right] \zeta^\bullet(k_r + l_r, \dots, k_{i+1} + l_{i+1}) t^{l_{i+1} + \dots + l_r} \in \mathcal{Z}[[t]]. \end{aligned}$$

Here, \mathcal{Z} is the \mathbb{Q} -subalgebra of \mathbb{R} generated by all MZVs and $\zeta^*(\mathbf{k}) \in \mathcal{Z}$ (resp. $\zeta^\mathfrak{m}(\mathbf{k}) \in \mathcal{Z}$) is the harmonic (resp. shuffle) regularized MZV. See Subsection 2.1 for details. It is known that $\zeta_{\widehat{\mathcal{S}}}^*(\mathbf{k}) - \zeta_{\widehat{\mathcal{S}}}^\mathfrak{m}(\mathbf{k}) \in (\zeta(2)\mathcal{Z})[[t]]$ for any index \mathbf{k} ([J2, Proposition 3.2.4] and [OSY, Proposition 2.1]). Thus,

$$\zeta_{\widehat{\mathcal{S}}}(\mathbf{k}) := \zeta_{\widehat{\mathcal{S}}}^\bullet(\mathbf{k}) \bmod \zeta(2)$$

is independent of the choice of the regularization $\bullet \in \{*, \mathfrak{m}\}$ and defines a well-defined element of $\overline{\mathcal{Z}}[[t]] := (\mathcal{Z}/\zeta(2)\mathcal{Z})[[t]]$. We call $\zeta_{\widehat{\mathcal{S}}}(\mathbf{k})$ the $\widehat{\mathcal{S}}$ -MZV. We also define the *t-adic symmetric multiple zeta-star value* ($\widehat{\mathcal{S}}$ -MZSV) $\zeta_{\widehat{\mathcal{S}}}^*(\mathbf{k})$ by

$$\zeta_{\widehat{\mathcal{S}}}^*(k_1, \dots, k_r) = \sum_{\substack{\square \text{ is either a comma ','} \\ \text{or a plus '+'}}} \zeta_{\widehat{\mathcal{S}}}(k_1 \square \dots \square k_r).$$

See [HMO, Definition 1.1] for another equivalent definition of the $\widehat{\mathcal{S}}$ -MZSV. For a positive integer n , let $\pi_n: \overline{\mathcal{Z}}[[t]] \rightarrow \overline{\mathcal{Z}}[[t]]/(t^n)$ be the natural projection.

Definition 1.1. For an index $\mathbf{k} = (k_1, \dots, k_r)$, we define the \mathcal{S}_n -multiple zeta(-star) value (\mathcal{S}_n -MZ(S)V) by

$$\zeta_{\mathcal{S}_n}(\mathbf{k}) := \pi_n(\zeta_{\widehat{\mathcal{S}}}(\mathbf{k})), \quad \zeta_{\mathcal{S}_n}^*(\mathbf{k}) := \pi_n(\zeta_{\widehat{\mathcal{S}}}^*(\mathbf{k})) = \sum_{\substack{\square \text{ is either a comma ','} \\ \text{or a plus '+'}}} \zeta_{\mathcal{S}_n}(k_1 \square \dots \square k_r).$$

Note that $\zeta_{\mathcal{S}_1}(\mathbf{k})$ coincides with the usual symmetric multiple zeta value (SMZV) $\zeta_{\mathcal{S}}(\mathbf{k})$ defined by Kaneko and Zagier [KZ].

\mathcal{A}_n -MZ(S)V and \mathcal{S}_n -MZ(S)V are the main objects of this article and together they are called \mathcal{F}_n -MZ(S)V; \mathcal{F} derives from the first letter of the word “finite”. Similar to the conjecture [OSY, Conjecture 4.3], it is conjectured that \mathcal{A}_n -MZVs and \mathcal{S}_n -MZVs satisfy relations of the same form. Hence, a relation among \mathcal{A}_n -MZVs or \mathcal{S}_n -MZVs is always described collectively as a relation of \mathcal{F}_n -MZVs, at least conjecturally. The purpose of

this paper is to confirm that several evaluations of special values and several relations known only in \mathcal{A}_n -MZVs or \mathcal{S}_n -MZVs are uniformly valid in \mathcal{F}_n -MZVs. In some cases, we only deal with $n = 1, 2, 3$.

The remainder of the paper is structured as follows. In Section 2, we prepare relevant tools including Zagier's formula for MZVs, the double shuffle relation for \mathcal{F}_n -MZVs and the relation for \mathcal{F}_n -MZVs derived from the antipode. In Section 3, we put forward some explicit evaluations of \mathcal{F}_n -MZ(S)Vs. In Section 4, we prove the Bowman–Bradley type theorem for \mathcal{F}_2 -MZ(S)Vs. In Section 5, we prove sum formulas for \mathcal{F}_n -MZ(S)Vs with respect to specific n . Some complicated but elementary calculations for binomial coefficients (= the proof of Proposition A.1) are proved in the Appendix.

ACKNOWLEDGEMENT

The authors would like to thank Dr. Kenji Sakugawa for helpful discussion concerning Proposition 3.1 and Theorem 3.2. The authors also would like to thank Hanamichi Kawamura and the anonymous referee for valuable comments on the manuscript.

2. PRELIMINARIES

In this section, we prepare tools which are used in the following sections.

2.1. Algebraic setup. First we recall the notion of the harmonic algebra introduced in [H1]. Let $\mathfrak{H}^1 := \mathbb{Q} + e_1\mathbb{Q}\langle e_0, e_1 \rangle \supset \mathfrak{H}^0 := \mathbb{Q} + e_1\mathbb{Q}\langle e_0, e_1 \rangle e_0$, where $\mathbb{Q}\langle e_0, e_1 \rangle$ is a non-commutative polynomial algebra in two variables e_0 and e_1 . For a positive integer k , we set $e_k := e_1 e_0^{k-1}$. We define the harmonic product $*$ on \mathfrak{H}^1 by $w * 1 = 1 * w = w$, $e_{k_1} w_1 * e_{k_2} w_2 = e_{k_1}(w_1 * e_{k_2} w_2) + e_{k_2}(e_{k_1} w_1 * w_2) + e_{k_1+k_2}(w_1 * w_2)$ (w, w_1, w_2 are words in \mathfrak{H}^1 , $k_1, k_2 \in \mathbb{Z}_{>0}$) with \mathbb{Q} -bilinearity. We also define the shuffle product \mathfrak{m} on $\mathbb{Q}\langle e_0, e_1 \rangle$ by $w \mathfrak{m} 1 = 1 \mathfrak{m} w = w$, $u_1 w_1 \mathfrak{m} u_2 w_2 = u_1(w_1 \mathfrak{m} u_2 w_2) + u_2(u_1 w_1 \mathfrak{m} w_2)$ (w, w_1, w_2 are words in $\mathbb{Q}\langle e_0, e_1 \rangle$, $u_1, u_2 \in \{e_0, e_1\}$) with \mathbb{Q} -bilinearity. Let $\bullet \in \{*, \mathfrak{m}\}$. It is known that \mathfrak{H}^1 becomes a commutative \mathbb{Q} -algebra with respect to the multiplication \bullet , which is denoted by \mathfrak{H}_\bullet^1 . The subspace \mathfrak{H}^0 of \mathfrak{H}^1 is closed under \bullet and becomes a \mathbb{Q} -subalgebra of \mathfrak{H}_\bullet^1 , which is denoted by \mathfrak{H}_\bullet^0 . We define Muneta's shuffle product $\tilde{\mathfrak{m}}$ on \mathfrak{H}^1 ([Mun, §3]) by $w \tilde{\mathfrak{m}} 1 = 1 \tilde{\mathfrak{m}} w = w$, $e_{k_1} w_1 \tilde{\mathfrak{m}} e_{k_2} w_2 = e_{k_1}(w_1 \tilde{\mathfrak{m}} e_{k_2} w_2) + e_{k_2}(e_{k_1} w_1 \tilde{\mathfrak{m}} w_2)$ (w, w_1, w_2 are words in \mathfrak{H}^1 , $k_1, k_2 \in \mathbb{Z}_{>0}$) with \mathbb{Q} -bilinearity.

Next, we recall the harmonic (resp. shuffle) regularized MZV introduced in [IKZ]. It is known that $\mathfrak{H}_\bullet^1 \cong \mathfrak{H}_\bullet^0[e_1]$ as a \mathbb{Q} -algebra (see [H1] for $\bullet = *$ and [Re] for $\bullet = \mathfrak{m}$). Therefore, for $\bullet \in \{*, \mathfrak{m}\}$, any $a \in \mathfrak{H}_\bullet^1$ has a unique expression $a = \sum_{i=0}^n a_i \bullet e_1^{\bullet i}$, where $n \in \mathbb{Z}_{\geq 0}$, $a_i \in \mathfrak{H}_\bullet^0$ ($0 \leq i \leq n$) and $e_1^{\bullet i} := \overbrace{e_1 \bullet \cdots \bullet e_1}^i$. By this expression, we define a

\mathbb{Q} -algebra homomorphism $\text{reg}_\bullet : \mathfrak{H}_\bullet^1 \cong \mathfrak{H}_\bullet^0[e_1] \rightarrow \mathfrak{H}_\bullet^0$ by $\text{reg}_\bullet(\sum_{i=0}^n a_i \bullet e_1^i) := a_0$. We set $e_{\mathbf{k}} := e_{k_1} \cdots e_{k_r}$ for a non-empty index $\mathbf{k} = (k_1, \dots, k_r)$ and $e_\emptyset := 1$. Then we define a \mathbb{Q} -linear map $Z : \mathfrak{H}^0 \rightarrow \mathbb{R}$ by $Z(e_{\mathbf{k}}) := \zeta(\mathbf{k})$ for any admissible index \mathbf{k} . By using this terminology, we define the harmonic (resp. shuffle) regularized MZV $\zeta^*(\mathbf{k})$ (resp. $\zeta^\sharp(\mathbf{k})$) by $\zeta^*(\mathbf{k}) = (Z \circ \text{reg}_*) (e_{\mathbf{k}})$ (resp. $\zeta^\sharp(\mathbf{k}) := (Z \circ \text{reg}_\sharp) (e_{\mathbf{k}})$) for any index \mathbf{k} .

To calculate the shuffle regularized MZV, we use the following fact.

Lemma 2.1 (Regularization formula, [IKZ, Proposition 8]). Let $w = w'e_0$ be an element of \mathfrak{H}^0 with $w' \in \mathfrak{H}^1$. Then, for a non-negative integer m , we have

$$\text{reg}_\sharp(w e_1^m) = (-1)^m (w' \sharp e_1^m) e_0.$$

2.2. Zagier's formulas for MZVs. We quote some results on MZVs. We use these results to evaluate some \mathcal{S}_1 -MZ(S)V_s and \mathcal{S}_2 -MZ(S)V_s.

Theorem 2.2 ([Z, Theorem 1]). For non-negative integers a and b , we have

$$\zeta(\{2\}^a, 3, \{2\}^b) = 2 \sum_{r=1}^{a+b+1} (-1)^r \left\{ \binom{2r}{2a+2} - \left(1 - \frac{1}{2^{2r}}\right) \binom{2r}{2b+1} \right\} \zeta(\{2\}^{a+b-r+1}) \zeta(2r+1),$$

where $\{2\}^a$ denotes a repetitions $\underbrace{2, \dots, 2}_a$. In particular, we have

$$(2.1) \quad \begin{aligned} & \zeta(\{2\}^a, 3, \{2\}^b) \\ & \equiv 2(-1)^{a+b+1} \left\{ \binom{2a+2b+2}{2a+2} - \left(1 - \frac{1}{4^{a+b+1}}\right) \binom{2a+2b+2}{2b+1} \right\} \zeta(2a+2b+3) \pmod{\zeta(2)}. \end{aligned}$$

Theorem 2.3 ([Z, Proposition 7]). Let m and n be positive integers with $n \geq 2$ and $k := m + n$ being odd. Define a positive integer K as $k = 2K + 1$. Then we have

$$\zeta(m, n) = (-1)^m \sum_{s=0}^{K-1} \left\{ \binom{k-2s-1}{m-1} + \binom{k-2s-1}{n-1} - \delta_{n,2s} + (-1)^m \delta_{s,0} \right\} \zeta(2s) \zeta(k-2s).$$

Here $\delta_{x,y}$ is Kronecker's delta, and we understand $\zeta(0) = -\frac{1}{2}$. In particular, we have

$$(2.2) \quad \zeta(m, n) \equiv (-1)^{m+1} \frac{1}{2} \left\{ \binom{k}{m} + (-1)^m \right\} \zeta(k) \pmod{\zeta(2)}.$$

2.3. Double shuffle relation for \mathcal{F}_n -MZVs. The double shuffle relation (DSR) for \mathcal{F}_n -MZVs with $\mathcal{F} \in \{\mathcal{A}, \mathcal{S}\}$ established by Jarossay is a key tool in this paper. We define \mathbb{Q} -linear maps $Z_{\mathcal{A}_n} : \mathfrak{H}^1 \rightarrow \mathcal{A}_n$ and $Z_{\mathcal{S}_n} : \mathfrak{H}^1 \rightarrow \overline{\mathcal{Z}}[t]/(t^n)$ by

$$Z_{\mathcal{A}_n}(e_{\mathbf{k}}) := \zeta_{\mathcal{A}_n}(\mathbf{k}), \quad Z_{\mathcal{S}_n}(e_{\mathbf{k}}) := \zeta_{\mathcal{S}_n}(\mathbf{k})$$

for any index \mathbf{k} .

Theorem 2.4 (DSR for \mathcal{F}_n -MZVs, [J2]. cf. [OSY, Theorems 1.3 and 1.9]). *For indices \mathbf{k} and $\mathbf{l} = (l_1, \dots, l_s)$ and a positive integer n , we have the harmonic relation for \mathcal{F}_n -MZVs*

$$(2.3) \quad Z_{\mathcal{F}_n}(e_{\mathbf{k}} * e_{\mathbf{l}}) = Z_{\mathcal{F}_n}(e_{\mathbf{k}})Z_{\mathcal{F}_n}(e_{\mathbf{l}})$$

and the shuffle relation for \mathcal{F}_n -MZVs

$$(2.4) \quad Z_{\mathcal{F}_n}(e_{\mathbf{k}} \text{III} e_{\mathbf{l}}) = (-1)^{\text{wt}(\mathbf{l})} \sum_{\substack{\mathbf{l}'=(l'_1, \dots, l'_s) \in \mathbb{Z}_{\geq 0}^s \\ \text{wt}(\mathbf{l}') \leq n-1}} \left[\prod_{j=1}^s \binom{l_j + l'_j - 1}{l'_j} \right] Z_{\mathcal{F}_n}(e_{\mathbf{k}} e_{\overline{\mathbf{l}+\mathbf{l}'}}) x_{\mathcal{F}_n}^{\text{wt}(\mathbf{l}')}.$$

Here, we set $\text{wt}(\mathbf{l}') := l'_1 + \dots + l'_s$, $\overline{\mathbf{l}+\mathbf{l}'} := (l_s + l'_s, \dots, l_1 + l'_1)$ and

$$x_{\mathcal{F}_n} := \begin{cases} \mathbf{p}_n := (p \bmod p^n)_p & \text{if } \mathcal{F} = \mathcal{A}, \\ t \bmod t^n & \text{if } \mathcal{F} = \mathcal{S}. \end{cases}$$

We refer to the case $\mathbf{k} = \emptyset$ of the shuffle relation as the *reversal formula*.

We also use the following relation for \mathcal{F}_n -MZVs.

Proposition 2.5. For an index $\mathbf{k} = (k_1, \dots, k_r)$, $n \in \mathbb{Z}_{\geq 1}$ and $\mathcal{F} \in \{\mathcal{A}, \mathcal{S}\}$, we have

$$\sum_{i=0}^r (-1)^i \zeta_{\mathcal{F}_n}(k_1, \dots, k_i) \zeta_{\mathcal{F}_n}^*(k_r, \dots, k_{i+1}) = 0.$$

Proof. This follows from the harmonic relation and [IKOO, Proposition 6] (Note that the sign of [IKOO, Proposition 6] is mistaken). The case $\mathcal{F} = \mathcal{A}$ was first mentioned in [SS, Corollary 3.16 (42)]. \square

3. SPECIAL VALUES

In this section, we explicitly evaluate some \mathcal{F}_n -MZ(S)Vs. For positive integers n and k , set

$$\mathfrak{Z}_{\mathcal{F}_n}(k) := \begin{cases} \left(\frac{B_{p^{n-1}(p-1)-k+1}}{k-1+p^{n-1}} \bmod p^n \right)_p \in \mathcal{A}_n & \text{if } \mathcal{F} = \mathcal{A}, \\ \zeta(k) \bmod \zeta(2) \in \overline{\mathcal{Z}}[t]/(t^n) & \text{if } \mathcal{F} = \mathcal{S}. \end{cases}$$

Here, B_j is the j -th Seki–Bernoulli number and \widehat{B}_j denotes $\frac{B_j}{j}$.

Proposition 3.1. Let n and k be positive integers. For $1 \leq l \leq n-1$,

$$\mathfrak{Z}_{\mathcal{A}_n}(k+l) \mathbf{p}_n^l = \sum_{j=1}^{n-l} (-1)^j \binom{n-l}{j} \left(\widehat{B}_{j(p-1)-k-l+1} \cdot p^l \bmod p^n \right)_p \in \mathcal{A}_n$$

holds. In particular,

$$\mathfrak{Z}_{\mathcal{A}_2}(k+1)\mathbf{p}_2 = \left(\frac{B_{p-k-1}}{k+1} \cdot p \bmod p^2 \right)_p \in \mathcal{A}_2.$$

Proof. Let p be a sufficiently large prime number. Then, by using the Kummer-type congruence proved by Zhi-Hong Sun [Su, Corollary 4.1], we have

$$\begin{aligned} & \frac{B_{p^{n-1}(p-1)-k-l+1}}{k+l-1+p^{n-1}} \cdot p^l \\ & \equiv (-1)^{n+l} \sum_{j=1}^{n-l} (-1)^{j-1} \binom{p^{n-1}-1-j}{n-l-j} \binom{p^{n-1}-1}{j-1} \widehat{B}_{j(p-1)-k-l+1} \cdot p^l \pmod{p^n}. \end{aligned}$$

Since

$$(-1)^{n+l-1} \binom{p^{n-1}-1-j}{n-l-j} \binom{p^{n-1}-1}{j-1} \equiv \binom{n-l}{j} \pmod{p^{n-1}},$$

we have the desired formula. \square

3.1. Depth 1 case.

Theorem 3.2. *For positive integers n and k , we have*

$$\zeta_{\mathcal{F}_n}(k) = (-1)^k \sum_{l=1}^{n-1} \binom{k+l-1}{l} \mathfrak{Z}_{\mathcal{F}_n}(k+l)x_{\mathcal{F}_n}^l.$$

Proof. The case $\mathcal{F} = \mathcal{A}$ is a special case of [W, Theorem 1]. Nevertheless, we can state the direct proof as follows. Let p be a sufficiently large prime number. By Euler's formula and Faulhaber's formula, we have

$$\begin{aligned} \sum_{m=1}^{p-1} \frac{1}{m^k} & \equiv \sum_{m=1}^{p-1} m^{\varphi(p^n)-k} \\ & \equiv \frac{1}{\varphi(p^n)-k+1} \sum_{l=1}^{n-1} \binom{\varphi(p^n)-k+1}{l} B_{\varphi(p^n)-k-l+1} \cdot p^l \\ & = - \sum_{l=1}^{n-1} \binom{\varphi(p^n)-k}{l} \frac{B_{\varphi(p^n)-k-l+1}}{k+l-1+p^{n-1}} \cdot p^l \pmod{p^n}, \end{aligned}$$

where φ is Euler's totient function. By a simple congruence

$$\binom{\varphi(p^n)-k}{l} \equiv (-1)^l \binom{k+l-1}{l} \pmod{p^{n-1}}$$

and the fact that B_j vanishes for odd $j \geq 3$, we have the desired equality in \mathcal{A}_n . Since the case $\mathcal{F} = \mathcal{S}$ is clear by definition, this completes the proof. \square

Remark 3.3. By combining the case $\mathcal{F} = \mathcal{A}$ of Theorem 3.2 and Proposition 3.1, we have

$$\sum_{m=1}^{p-1} \frac{1}{m^k} \equiv (-1)^k \sum_{l=1}^{n-1} \binom{k+l-1}{l} \sum_{j=1}^{n-l} (-1)^j \binom{n-l}{j} \widehat{B}_{j(p-1)-k-l+1} p^l \pmod{p^n}$$

for a sufficiently large prime p . We can check that this holds for $p \geq n+k+1$. This congruence is a generalization of [Su, Theorem 5.1 (a) and Remark 5.1] and [Tau, Theorem 2.1]. However, the proof is identical to that put forward by Sun.

3.2. Depth 2 case. Let $\tau_n: \overline{\mathcal{Z}}[[t]] \rightarrow \overline{\mathcal{Z}}[t]$ be the truncation map defined by $\tau_n(\sum_{l=0}^{\infty} z_l t^l) := \sum_{l=0}^{n-1} z_l t^l$ for a positive integer n . In the following argument, we often identify $\zeta_{\mathcal{S}_n}^{\bullet}(\mathbf{k})$ with $\tau_n(\zeta_{\widehat{\mathcal{S}}}^{\bullet}(\mathbf{k}))$, where $\bullet \in \{\emptyset, \star\}$. Furthermore, we often abbreviate $\zeta(\mathbf{k}) \bmod \zeta(2)$ (resp. $\zeta^{\text{III}}(\mathbf{k}) \bmod \zeta(2)$) to $\zeta(\mathbf{k})$ (resp. $\zeta^{\text{III}}(\mathbf{k})$) in $\overline{\mathcal{Z}}$.

Theorem 3.4. *Let k_1 and k_2 be positive integers. Assume that $k := k_1 + k_2$ is even. Then we have*

$$(3.1) \quad \zeta_{\mathcal{F}_2}(k_1, k_2) = \frac{1}{2} \left\{ (-1)^{k_1} k_2 \binom{k+1}{k_1} - (-1)^{k_2} k_1 \binom{k+1}{k_2} - k \right\} \mathfrak{Z}_{\mathcal{F}_2}(k+1) x_{\mathcal{F}_2},$$

$$(3.2) \quad \zeta_{\mathcal{F}_2}^{\star}(k_1, k_2) = \frac{1}{2} \left\{ (-1)^{k_1} k_2 \binom{k+1}{k_1} - (-1)^{k_2} k_1 \binom{k+1}{k_2} + k \right\} \mathfrak{Z}_{\mathcal{F}_2}(k+1) x_{\mathcal{F}_2}.$$

Proof. The case $\mathcal{F} = \mathcal{A}$ was proved by Zhao, see [Zh, Theorem 3.2]. Hereafter, we consider the case $\mathcal{F} = \mathcal{S}$. First, we prove (3.1) for the case $k_1 \geq 2$. By the definition of $\zeta_{\mathcal{S}_2}(k_1, k_2)$, we have

$$\zeta_{\mathcal{S}_2}(k_1, k_2) = \zeta_{\mathcal{S}_1}(k_1, k_2) + \{k_2 \zeta(k_2+1, k_1) + k_1 \zeta(k_2, k_1+1)\} t.$$

Since $k_1 + k_2$ is even, we have $\zeta_{\mathcal{S}_1}(k_1, k_2) = 0$ by definition. Therefore, by using (2.2), we obtain (3.1) for the case $k_1 \geq 2$. Next, we prove (3.1) for the case $k_1 = 1$ (then k_2 is odd). We have

$$(3.3) \quad \zeta_{\mathcal{S}_2}(1, k_2) = \{k_2 \zeta^{\text{III}}(k_2+1, 1) + \zeta(k_2, 2)\} t.$$

By applying Theorem 2.1 for $w = e_1 e_0^{k_2}$ and the sum formula for MZVs of depth 2, we have

$$(3.4) \quad \zeta^{\text{III}}(k_2+1, 1) = -\zeta(k_2, 2) - \cdots - \zeta(2, k_2) - 2\zeta(1, k_2+1) = -\zeta(k_2+2) - \zeta(1, k_2+1).$$

By (2.2), we have

$$(3.5) \quad \zeta(1, k_2+1) = \frac{k_2+1}{2} \zeta(k_2+2), \quad \zeta(k_2, 2) = \frac{1}{2} \left\{ \frac{(k_2+2)(k_2+1)}{2} - 1 \right\} \zeta(k_2+2).$$

From (3.3), (3.4), (3.5), we obtain (3.1) for the case $k_1 = 1$. The formula (3.2) follows from (3.1), the fact $\zeta_{\mathcal{S}_2}^*(k_1, k_2) = \zeta_{\mathcal{S}_2}(k_1, k_2) + \zeta_{\mathcal{S}_2}(k_1 + k_2)$, and $\zeta_{\mathcal{S}_2}(k) = (-1)^k k \zeta(k+1)t$ (Theorem 3.2 with $\mathcal{F}_n = \mathcal{S}_2$). \square

3.3. Depth 3 case.

Theorem 3.5. *Let k_1, k_2, k_3 be positive integers. Suppose that $k := k_1 + k_2 + k_3$ is odd. Then we have*

$$\zeta_{\mathcal{F}_1}(k_1, k_2, k_3) = -\zeta_{\mathcal{F}_1}^*(k_1, k_2, k_3) = \frac{1}{2} \left\{ (-1)^{k_1} \binom{k}{k_1} - (-1)^{k_3} \binom{k}{k_3} \right\} \mathfrak{Z}_{\mathcal{F}_1}(k).$$

Proof. The case $\mathcal{F} = \mathcal{A}$ was proved by Hoffman and Zhao; see [H2, Theorem 6.2] or [Zh, Theorem 3.5]. Hereafter, we consider the case $\mathcal{F} = \mathcal{S}$. By Proposition 2.5 and the reversal formula for \mathcal{F}_1 -MZVs, we have

$$(3.6) \quad \zeta_{\mathcal{F}_1}^*(k_1, k_2, k_3) = (-1)^{k_1+k_2+k_3} \zeta_{\mathcal{F}_1}(k_1, k_2, k_3) = -\zeta_{\mathcal{F}_1}(k_1, k_2, k_3).$$

From

$$\zeta_{\mathcal{F}_1}^*(k_1, k_2, k_3) = \zeta_{\mathcal{F}_1}(k_1, k_2, k_3) + \zeta_{\mathcal{F}_1}(k_1 + k_2, k_3) + \zeta_{\mathcal{F}_1}(k_1, k_2 + k_3)$$

and the explicit formula for \mathcal{F}_1 -double zeta values [Kan, (7.2), Example 9.4 (2)], we have

$$(3.7) \quad \begin{aligned} \zeta_{\mathcal{F}_1}(k_1, k_2, k_3) &= -\frac{\zeta_{\mathcal{F}_1}(k_1 + k_2, k_3) + \zeta_{\mathcal{F}_1}(k_1, k_2 + k_3)}{2} \\ &= -\frac{1}{2} \left\{ (-1)^{k_3} \binom{k}{k_1 + k_2} + (-1)^{k_2+k_3} \binom{k}{k_1} \right\} \mathfrak{Z}_{\mathcal{F}_1}(k) \\ &= \frac{1}{2} \left\{ (-1)^{k_1} \binom{k}{k_1} - (-1)^{k_3} \binom{k}{k_3} \right\} \mathfrak{Z}_{\mathcal{F}_1}(k). \end{aligned}$$

The formula for $\zeta_{\mathcal{F}_1}^*(k_1, k_2, k_3)$ is obtained by (3.6) and (3.7). \square

3.4. General depth case.

Theorem 3.6. *For positive integers r, k and for $\mathcal{F} \in \{\mathcal{A}, \mathcal{S}\}$, we have*

$$(3.8) \quad \zeta_{\mathcal{F}_2}(\{k\}^r) = (-1)^{r-1} k \mathfrak{Z}_{\mathcal{F}_2}(rk+1) x_{\mathcal{F}_2},$$

$$(3.9) \quad \zeta_{\mathcal{F}_2}^*(\{k\}^r) = k \mathfrak{Z}_{\mathcal{F}_2}(rk+1) x_{\mathcal{F}_2}.$$

Moreover, we have

$$(3.10) \quad \zeta_{\mathcal{F}_3}(\{k\}^r) = (-1)^{r+k+r-1} \left[k \mathfrak{Z}_{\mathcal{F}_3}(rk+1) x_{\mathcal{F}_3} + \left\{ \frac{k(rk+1)}{2} \mathfrak{Z}_{\mathcal{F}_3}(rk+2) - k^2 \sum_{l=1}^{r-1} \mathfrak{Z}_{\mathcal{F}_3}(lk+1) \mathfrak{Z}_{\mathcal{F}_3}((r-l)k+1) \right\} x_{\mathcal{F}_3}^2 \right]$$

and

$$(3.11) \quad \zeta_{\mathcal{F}_3}^*(\{k\}^r) = (-1)^{rk} \left[k \mathfrak{Z}_{\mathcal{F}_3}(rk+1)x_{\mathcal{F}_3} + \left\{ \frac{k(rk+1)}{2} \mathfrak{Z}_{\mathcal{F}_3}(rk+2) + k^2 \sum_{l=1}^{r-1} \mathfrak{Z}_{\mathcal{F}_3}(lk+1) \mathfrak{Z}_{\mathcal{F}_3}((r-l)k+1) \right\} x_{\mathcal{F}_3}^2 \right].$$

Remark 3.7. If rk is odd, then $\mathfrak{Z}_{\mathcal{F}_3}(rk+1)$ and $\mathfrak{Z}_{\mathcal{F}_3}(lk+1)\mathfrak{Z}_{\mathcal{F}_3}((r-l)k+1)$ are 0, and we have

$$\zeta_{\mathcal{F}_3}(\{k\}^r) = (-1)^r \frac{k(rk+1)}{2} \mathfrak{Z}_{\mathcal{F}_3}(rk+2)x_{\mathcal{F}_3}^2, \quad \zeta_{\mathcal{F}_3}^*(\{k\}^r) = -\frac{k(rk+1)}{2} \mathfrak{Z}_{\mathcal{F}_3}(rk+2)x_{\mathcal{F}_3}^2.$$

These formulas for the case $\mathcal{F} = \mathcal{A}$ were first proved by Zhou and Cai in the last remark of [ZC] but our proof differs from theirs.

Proof. Since (3.8) and (3.9) follows from (3.10) and (3.11) by taking modulo $x_{\mathcal{F}_3}^2$, it is sufficient to prove (3.10) and (3.11). Note that $(-1)^{rk} \mathfrak{Z}_{\mathcal{F}_2}(rk+1)x_{\mathcal{F}_2} = \mathfrak{Z}_{\mathcal{F}_2}(rk+1)x_{\mathcal{F}_2}$ holds because if rk is odd, then $\mathfrak{Z}_{\mathcal{F}_2}(rk+1)x_{\mathcal{F}_2} = 0$.

By Theorem 3.2 and the symmetric sum formula (5.1) proved in Section 5 with $\mathbf{k} = (\{k\}^r)$, we have

$$(3.12) \quad \begin{aligned} & r! \zeta_{\mathcal{F}_3}(\{k\}^r) \\ &= (-1)^{rk+r-1} (r-1)! \left\{ rk \mathfrak{Z}_{\mathcal{F}_3}(rk+1)x_{\mathcal{F}_3} + \binom{rk+1}{2} \mathfrak{Z}_{\mathcal{F}_3}(rk+2)x_{\mathcal{F}_3}^2 \right\} \\ & \quad + (-1)^{rk+r-2} \sum_{\substack{B_1 \sqcup B_2 = \{1, \dots, r\} \\ B_1, B_2 \neq \emptyset}} (\#B_1 - 1)! (\#B_2 - 1)! b_1 b_2 \mathfrak{Z}_{\mathcal{F}_3}(b_1+1) \mathfrak{Z}_{\mathcal{F}_3}(b_2+1)x_{\mathcal{F}_3}^2, \end{aligned}$$

where $b_1 = b_1(\{k\}^r)$ and $b_2 = b_2(\{k\}^r)$ are defined as in Theorem 5.1. Set $l := \#B_1$. Then we see that $1 \leq l \leq r-1$, $\#B_2 = r-l$, $b_1 = lk$ and $b_2 = (r-l)k$. Moreover, the number of ways of dividing $\{1, \dots, r\}$ into two non-empty subsets B_1 and B_2 with $\#B_1 = l$ is just $\binom{r}{l}$. Therefore, the summation for the partition in the right-hand side of (3.12) coincides with

$$\begin{aligned} & \sum_{l=1}^{r-1} \binom{r}{l} (l-1)! (r-l-1)! \cdot l(r-l)k^2 \mathfrak{Z}_{\mathcal{F}_3}(lk+1) \mathfrak{Z}_{\mathcal{F}_3}((r-l)k+1)x_{\mathcal{F}_3}^2 \\ &= k^2 \cdot r! \sum_{l=1}^{r-1} \mathfrak{Z}_{\mathcal{F}_3}(lk+1) \mathfrak{Z}_{\mathcal{F}_3}((r-l)k+1)x_{\mathcal{F}_3}^2. \end{aligned}$$

Thus we obtained (3.10). The formula (3.11) is obtained in the same manner. \square

Theorem 3.8. *For non-negative integers a and b , we have*

$$(3.13) \quad \zeta_{\mathcal{F}_1}(\{1\}^a, 2, \{1\}^b) = (-1)^b \binom{a+b+2}{a+1} \mathfrak{Z}_{\mathcal{F}_1}(a+b+2),$$

$$(3.14) \quad \zeta_{\mathcal{F}_1}^*(\{1\}^a, 2, \{1\}^b) = (-1)^b \binom{a+b+2}{a+1} \mathfrak{Z}_{\mathcal{F}_1}(a+b+2).$$

Proof. The case $\mathcal{F} = \mathcal{A}$ was proved by Hessami-Pilehrood–Hessami-Pilehrood–Tauraso; see [HHT, Theorem 4.5]. Hereafter, we consider the case $\mathcal{F} = \mathcal{S}$. By the definition of $\zeta_{\mathcal{S}_1}(\mathbf{k})$ and the fact that $\zeta^{\mathfrak{m}}(\{1\}^k) = 0$ for $k \geq 1$, we have

$$(3.15) \quad \zeta_{\mathcal{S}_1}(\{1\}^a, 2, \{1\}^b) = \zeta^{\mathfrak{m}}(\{1\}^a, 2, \{1\}^b) + (-1)^{a+b} \zeta^{\mathfrak{m}}(\{1\}^b, 2, \{1\}^a).$$

Applying Lemma 2.1 for $w = e_1^{a+1}e_0$ and $m = b$, and using the duality for MZVs, we have

$$(3.16) \quad \zeta^{\mathfrak{m}}(\{1\}^a, 2, \{1\}^b) = (-1)^b \binom{a+b+1}{b} \zeta(a+b+2).$$

From (3.15) and (3.16), we obtain (3.13). The formula (3.14) follows from (3.13), Proposition 2.5 and the fact that $\zeta_{\mathcal{S}_1}(\{1\}^r) = \zeta_{\mathcal{S}_1}^*(\{1\}^r) = 0$ for $r \geq 1$. The last fact is well-known and a special case of Theorem 3.6. \square

Remark 3.9. We can also prove (3.14) using the Hoffman duality ([H2, Theorem 4.6] and [J1, Corollarie 1.12]) and the explicit formula for $\zeta_{\mathcal{F}_1}(a+1, b+1)$.

Theorem 3.10. *For non-negative integers a and b , we have*

$$(3.17) \quad \zeta_{\mathcal{F}_1}(\{2\}^a, 3, \{2\}^b) = \frac{(-1)^{a+b} 2(a-b)}{a+1} \binom{2a+2b+3}{2b+2} \mathfrak{Z}_{\mathcal{F}_1}(2a+2b+3),$$

$$(3.18) \quad \zeta_{\mathcal{F}_1}^*(\{2\}^a, 3, \{2\}^b) = \frac{2(b-a)}{a+1} \binom{2a+2b+3}{2b+2} \mathfrak{Z}_{\mathcal{F}_1}(2a+2b+3).$$

Proof. The case $\mathcal{F} = \mathcal{A}$ was proved by Hessami-Pilehrood–Hessami-Pilehrood–Tauraso; see [HHT, Theorem 4.1]. Hereafter, we consider the case $\mathcal{F} = \mathcal{S}$. By the definition of the \mathcal{S}_1 -MZV and the fact that $\zeta(\{2\}^r) \equiv 0 \pmod{\zeta(2)}$ for $r \geq 1$, we have

$$\zeta_{\mathcal{S}_1}(\{2\}^a, 3, \{2\}^b) = \zeta(\{2\}^a, 3, \{2\}^b) - \zeta(\{2\}^b, 3, \{2\}^a).$$

Thus we obtain (3.17) by the formula (2.1) and straightforward calculation of binomial coefficients. The formula (3.18) is obtained by (3.17), Proposition 2.5 and a special case of Theorem 3.6, that is, the fact that $\zeta_{\mathcal{S}_1}(\{2\}^r) = \zeta_{\mathcal{S}_1}^*(\{2\}^r) = 0$ for $r \geq 1$. \square

Theorem 3.11. *For non-negative integers a and b , we have*

$$(3.19) \quad \zeta_{\mathcal{F}_1}(\{2\}^a, 1, \{2\}^b) = 4(-1)^{a+b} \frac{a-b}{2a+1} \left(1 - \frac{1}{4^{a+b}}\right) \binom{2a+2b+1}{2b+1} \mathfrak{Z}_{\mathcal{F}_1}(2a+2b+1),$$

$$(3.20) \quad \zeta_{\mathcal{F}_1}^*(\{2\}^a, 1, \{2\}^b) = \frac{4(b-a)}{2a+1} \left(1 - \frac{1}{4^{a+b}}\right) \binom{2a+2b+1}{2b+1} \mathfrak{Z}_{\mathcal{F}_1}(2a+2b+1).$$

Proof. The case $\mathcal{F} = \mathcal{A}$ was proved by Hessami-Pilehrood–Hessami-Pilehrood–Tauraso; see [HHT, Theorem 4.2]. Hereafter, we consider the case $\mathcal{F} = \mathcal{S}$. First, we prove (3.19) for the case $a, b \geq 1$. By the definition of the \mathcal{S}_1 -MZV and the duality for MZVs, we obtain

$$\zeta_{\mathcal{S}_1}(\{2\}^a, 1, \{2\}^b) = \zeta(\{2\}^{b-1}, 3, \{2\}^a) - \zeta(\{2\}^{a-1}, 3, \{2\}^b).$$

Then we obtain (3.19) by a similar calculation in Theorem 3.10 using (2.1).

Next we prove (3.19) for the case $a \geq 1$ and $b = 0$. We have

$$(3.21) \quad \zeta_{\mathcal{S}_1}(\{2\}^a, 1) = \zeta^{\text{III}}(\{2\}^a, 1) - \zeta(1, \{2\}^a).$$

Applying Lemma 2.1 for $w = (e_1 e_0)^a$ and $m = 1$, we obtain

$$\zeta^{\text{III}}(\{2\}^a, 1) = -2 \sum_{j=0}^{a-1} \zeta(\{2\}^j, 1, \{2\}^{a-j}).$$

Thus, by the duality for MZVs, we have

$$(3.22) \quad \begin{aligned} \zeta_{\mathcal{S}_1}(\{2\}^a, 1) &= -\zeta(1, \{2\}^a) - 2 \sum_{j=0}^{a-1} \zeta(\{2\}^j, 1, \{2\}^{a-j}) \\ &= -\zeta(\{2\}^{a-1}, 3) - 2 \sum_{j=0}^{a-1} \zeta(\{2\}^{a-j-1}, 3, \{2\}^j). \end{aligned}$$

By (2.1), we obtain

$$(3.23) \quad \zeta(\{2\}^{a-j-1}, 3, \{2\}^j) = 2(-1)^a \left\{ \binom{2a}{2j} - \left(1 - \frac{1}{4^a}\right) \binom{2a}{2j+1} \right\} \zeta(2a+1)$$

for $0 \leq j \leq a-1$. Therefore, from (3.22) and (3.23), we obtain (3.19) for the case $a \geq 1$ and $b = 0$. The case $a = 0$ and $b \geq 1$ of (3.19) follows easily from the reversal formula and (3.22) with $a \geq 1$. This completes the proof of (3.19). The formula (3.20) is obtained by (3.19), Proposition 2.5 and the fact that $\zeta_{\mathcal{S}_1}(\{2\}^r) = \zeta_{\mathcal{S}_1}^*(\{2\}^r) = 0$ for $r \geq 1$. \square

Remark 3.12. Tasaka and Yamamoto proved an analogous formula of Theorem 2.2 for $\zeta^*(\{2\}^a, 1, \{2\}^b)$; see [TY, Theorem 1.6]. We can also obtain Theorem 3.11 by a

similar approach using [TY, Theorem 1.6] and Proposition 2.5 instead of Zagier's formula (Theorem 2.2).

The following theorem is a refinement of the even weight case in Theorem 3.8.

Theorem 3.13. *Let a and b be non-negative integers. Assume that $a + b$ is even. Then we have*

$$(3.24) \quad \zeta_{\mathcal{F}_2}(\{1\}^a, 2, \{1\}^b) = \frac{1}{2} \left\{ 1 + (-1)^a \binom{a+b+3}{b+2} \right\} \mathfrak{Z}_{\mathcal{F}_2}(a+b+3)x_{\mathcal{F}_2},$$

$$(3.25) \quad \zeta_{\mathcal{F}_2}^*(\{1\}^a, 2, \{1\}^b) = \frac{1}{2} \left\{ 1 + (-1)^a \binom{a+b+3}{a+2} \right\} \mathfrak{Z}_{\mathcal{F}_2}(a+b+3)x_{\mathcal{F}_2}.$$

Proof. The case $\mathcal{F} = \mathcal{A}$ was proved by Hessami-Pilehrood–Hessami-Pilehrood–Tauraso [HHT, Theorem 4.5]; there is also another proof by Sakugawa and the third author [SS, Theorem 3.18]. Hereafter, we consider the case $\mathcal{F} = \mathcal{S}$. We first prove the formula (3.24). Set $d := a + b + 1$ and $\mathbf{k} = (k_1, \dots, k_d) := (\{1\}^a, 2, \{1\}^b)$. For $0 \leq i \leq d$, we set

$$P_i(t) := (-1)^{k_{i+1} + \dots + k_d} \zeta^{\mathfrak{m}}(k_1, \dots, k_i) \\ \times \sum_{\substack{l_{i+1}, \dots, l_d \geq 0 \\ l_{i+1} + \dots + l_d \leq 1}} \left[\prod_{j=i+1}^d \binom{k_j + l_j - 1}{l_j} \right] \zeta^{\mathfrak{m}}(k_d + l_d, \dots, k_{i+1} + l_{i+1}) t^{l_{i+1} + \dots + l_d}.$$

Note that this expression for the case $i = d$ means $P_d(t) = \zeta^{\mathfrak{m}}(\mathbf{k})$. Then, by the definition of $\zeta_{\mathcal{S}_2}(\mathbf{k})$, we have

$$\zeta_{\mathcal{S}_2}(\mathbf{k}) = \sum_{i=0}^d P_i(t).$$

We prove that for $1 \leq i \leq a + b + 1$, $P_i(t) = 0$ in $\overline{\mathcal{Z}}[t]$. Since $\zeta^{\mathfrak{m}}(\{1\}^k) = 0$ for a positive integer k , we obtain $P_1(t) = \dots = P_a(t) = 0$. For $0 \leq j \leq b$, we calculate $P_{a+j+1}(t)$. By the definition of $P_i(t)$, we have

$$P_{a+j+1}(t) = (-1)^{b-j} \zeta^{\mathfrak{m}}(\{1\}^a, 2, \{1\}^j) \left(\zeta^{\mathfrak{m}}(\{1\}^{b-j}) + \sum_{i=1}^{b-j} \zeta^{\mathfrak{m}}(\{1\}^{i-1}, 2, \{1\}^{b-j-i}) t \right).$$

By (3.16), we have $\zeta^{\mathfrak{m}}(\{1\}^a, 2, \{1\}^j) = (-1)^j \binom{a+j+1}{j} \zeta(a+j+2)$. Since $\zeta(a+b+2) = 0$ in $\overline{\mathcal{Z}}$, we have $P_{a+b+1}(t) = 0$ in $\overline{\mathcal{Z}}[t]$. Assume that $j < b$. In this case, $\zeta^{\mathfrak{m}}(\{1\}^{b-j}) = 0$ holds. By the shuffle-regularized sum formula [Li, Lemma 3.3] (or [KS, Theorem 1.2]), and the duality for MZVs, we have

$$\sum_{i=1}^{b-j} \zeta^{\mathfrak{m}}(\{1\}^{i-1}, 2, \{1\}^{b-j-i}) = (-1)^{b-j-1} \zeta(b-j+1).$$

Since $a + b$ is even, $a + j + 2 \not\equiv b - j + 1 \pmod{2}$ and thus we have $P_{a+j+1}(t) = 0$ in $\overline{\mathcal{Z}}$.

The calculation of $P_0(t)$ remains. State $P_0(t) = A + Bt$ with $A, B \in \overline{\mathcal{Z}}$. Then from (3.16), we have $A = \zeta^{\boxplus}(\{1\}^b, 2, \{1\}^a) = 0$ in $\overline{\mathcal{Z}}$. By definition, B is expressed as follows:

$$\begin{aligned} B &= \sum_{\substack{l+m=b-1 \\ l, m \geq 0}} \zeta^{\boxplus}(\{1\}^l, 2, \{1\}^m, 2, \{1\}^a) \\ &\quad + 2\zeta^{\boxplus}(\{1\}^b, 3, \{1\}^a) + \sum_{\substack{m+n=a-1 \\ m, n \geq 0}} \zeta^{\boxplus}(\{1\}^b, 2, \{1\}^m, 2, \{1\}^n). \end{aligned}$$

For general non-negative integers l, m and n , by the regularization formula (Lemma 2.1), we have

$$\begin{aligned} \zeta^{\boxplus}(\{1\}^l, 2, \{1\}^m, 2, \{1\}^n) &= (-1)^n \sum_{\substack{r+s=n \\ r, s \geq 0}} \binom{r+l+1}{r} \binom{s+m+1}{s} \zeta(\{1\}^{r+l}, 2, \{1\}^{s+m}, 2), \\ \zeta^{\boxplus}(\{1\}^l, 3, \{1\}^n) &= (-1)^n \sum_{\substack{r+s=n \\ r, s \geq 0}} \binom{r+l+1}{r} \zeta(\{1\}^{r+l}, 2, \{1\}^{s-1}, 2), \end{aligned}$$

where $\zeta(\{1\}^{r+l}, 2, \{1\}^{-1}, 2)$ means $\zeta(\{1\}^{r+l}, 3)$. Thus, with the duality for MZVs, we obtain

$$\begin{aligned} B &= \sum_{\substack{l+m=b-1 \\ l, m \geq 0}} (-1)^a \sum_{\substack{r+s=a \\ r, s \geq 0}} \binom{r+l+1}{r} \binom{s+m+1}{s} \zeta(s+m+2, r+l+2) \\ &\quad + 2(-1)^a \sum_{\substack{r+s=a \\ r, s \geq 0}} \binom{r+b+1}{r} \zeta(s+1, r+b+2) \\ &\quad + \sum_{\substack{m+n=a-1 \\ m, n \geq 0}} (-1)^n \sum_{\substack{r+s=n \\ r, s \geq 0}} \binom{r+b+1}{r} \binom{s+m+1}{s} \zeta(s+m+2, r+b+2). \end{aligned}$$

Since $a + b + 3$ is odd, by using (2.2), we can rewrite B as a rational multiple of the Riemann zeta value $\zeta(a + b + 3)$. Specifically, we have $B = \frac{1}{2}C\zeta(a + b + 3)$ with

$$\begin{aligned}
C &= \sum_{\substack{l+m=b-1 \\ l,m \geq 0}} (-1)^a \sum_{\substack{r+s=a \\ r,s \geq 0}} \binom{r+l+1}{r} \binom{s+m+1}{s} (-1)^{s+m+1} \left\{ \binom{a+b+3}{s+m+2} + (-1)^{s+m} \right\} \\
&\quad + 2(-1)^a \sum_{\substack{r+s=a \\ r,s \geq 0}} \binom{r+b+1}{r} (-1)^s \left\{ \binom{a+b+3}{s+1} + (-1)^{s+1} \right\} \\
&\quad + \sum_{\substack{m+n=a-1 \\ m,n \geq 0}} (-1)^n \sum_{\substack{r+s=n \\ r,s \geq 0}} \binom{r+b+1}{r} \binom{s+m+1}{s} (-1)^{s+m+1} \left\{ \binom{a+b+3}{s+m+2} + (-1)^{s+m} \right\}.
\end{aligned}$$

Therefore, it suffices to prove the following:

$$(3.26) \quad C = 1 + (-1)^a \binom{a+b+3}{b+2}.$$

We prove this in the Appendix. From this, we obtain the desired formula for $\zeta_{\mathcal{S}_2}(\{1\}^a, 2, \{1\}^b)$.

Next, we prove (3.25). By Proposition 2.5, we have

$$\begin{aligned}
\zeta_{\mathcal{S}_2}^*(\{1\}^a, 2, \{1\}^b) - \zeta_{\mathcal{S}_2}(\{1\}^b, 2, \{1\}^a) &= \sum_{j=1}^a (-1)^j \zeta_{\mathcal{S}_2}(\{1\}^b, 2, \{1\}^{a-j}) \zeta_{\mathcal{S}_2}^*(\{1\}^j) \\
&\quad + \sum_{i=1}^b (-1)^{b-i} \zeta_{\mathcal{S}_2}(\{1\}^{b+1-i}) \zeta_{\mathcal{S}_2}^*(\{1\}^a, 2, \{1\}^{i-1}).
\end{aligned}$$

It is sufficient to show that the right-hand side vanishes. If j is odd, then we have $\zeta_{\mathcal{S}_2}^*(\{1\}^j) = 0$ by (3.9). If j is even, then both $\zeta_{\mathcal{S}_2}^*(\{1\}^j)$ and $\zeta_{\mathcal{S}_2}(\{1\}^b, 2, \{1\}^{a-j})$ can be seen as elements of $t\overline{\mathcal{Z}}[t]$ by (3.9) and (3.13). Thus the first summation vanishes in $\overline{\mathcal{Z}}[[t]]/(t^2)$. Similarly, if $b - i$ is even, then we have $\zeta_{\mathcal{S}_2}(\{1\}^{b+1-i}) = 0$ by (3.8). If $b - i$ is odd, then both $\zeta_{\mathcal{S}_2}(\{1\}^{b+1-i})$ and $\zeta_{\mathcal{S}_2}^*(\{1\}^a, 2, \{1\}^{i-1})$ can be seen as elements of $t\overline{\mathcal{Z}}[t]$ by (3.8) and (3.14). Therefore, the second summation also vanishes. \square

Remark 3.14. The proof of the case $\mathcal{F} = \mathcal{A}$ of Theorem 3.13 by Sakugawa and the third author is based on the ‘ \mathcal{A}_2 -duality’ [SS, Remark 3.14 (40)]. If the ‘ \mathcal{S}_2 -duality’ is established, then we can obtain another proof of the case $\mathcal{F} = \mathcal{S}$ of Theorem 3.13. When we were writing this paper, a preprint [TT] by Takeyama and Tasaka appeared on arXiv. Their [TT, Corollary 6.8] contains the \mathcal{S}_2 -duality as a special case.

4. BOWMAN–BRADLEY TYPE THEOREM

Murahara, Onozuka and the third author [MOS] proved the Bowman–Bradley type theorem for \mathcal{A}_2 -MZ(S)V_s (= the case $\mathcal{F} = \mathcal{A}$ of Theorem 4.1). In this section, we prove the \mathcal{S}_2 -counterpart of their theorem. By combining these two theorems, we have the following.

Theorem 4.1 (Bowman–Bradley type theorem for \mathcal{F}_2 -MZ(S)V). *For non-negative integers l and m with $(l, m) \neq (0, 0)$, we have*

$$(4.1) \quad \begin{aligned} & \sum_{\substack{m_0 + \dots + m_{2l} = m \\ m_0, \dots, m_{2l} \geq 0}} \zeta_{\mathcal{F}_2}(\{2\}^{m_0}, 1, \{2\}^{m_1}, 3, \{2\}^{m_2}, \dots, \{2\}^{m_{2l-2}}, 1, \{2\}^{m_{2l-1}}, 3, \{2\}^{m_{2l}}) \\ &= (-1)^m \left\{ (-1)^l 2^{1-2l} \binom{l+m}{l} - 4 \binom{2l+m}{2l} \right\} \mathfrak{Z}_{\mathcal{F}_2}(4l+2m+1)x_{\mathcal{F}_2}, \end{aligned}$$

$$(4.2) \quad \begin{aligned} & \sum_{\substack{m_0 + \dots + m_{2l} = m \\ m_0, \dots, m_{2l} \geq 0}} \zeta_{\mathcal{F}_2}^*(\{2\}^{m_0}, 1, \{2\}^{m_1}, 3, \{2\}^{m_2}, \dots, \{2\}^{m_{2l-2}}, 1, \{2\}^{m_{2l-1}}, 3, \{2\}^{m_{2l}}) \\ &= (-1)^l 2^{1-2l} \binom{l+m}{l} \mathfrak{Z}_{\mathcal{F}_2}(4l+2m+1)x_{\mathcal{F}_2}. \end{aligned}$$

This gives a partial lift of the Bowman–Bradley type theorem for \mathcal{F}_1 -MZ(S)V proved by Saito and Wakabayashi [SW2]. Note that the proof of the $\mathcal{F} = \mathcal{S}$ case in Theorem 4.1 presented here is essentially the same as the proof of the $\mathcal{F} = \mathcal{A}$ case by Murahara, Onozuka and the third author [MOS]. In contrast, proofs of some sum formulas which will be given in the next section are different from those in the previous study.

We prepare two lemmas for the proof.

Lemma 4.2. For non-negative integers l and m with $(l, m) \neq (0, 0)$, we have

$$Z_{\mathcal{S}_2}(e_2^{l+m} \boxplus e_2^l) = (-1)^m 2 \left\{ 1 - 2 \binom{4l+2m}{2l} \right\} \zeta(4l+2m+1)t.$$

Proof. This lemma is the \mathcal{S}_2 -counterpart of [MOS, Lemma 2.5] and is proved from the same argument in [MOS] by using the explicit evaluation of $\zeta_{\mathcal{S}_2}(\{2\}^r)$ ((3.8) with $k = 2$), (3.17) and (2.4) with $\mathcal{F}_n = \mathcal{S}_2$. \square

For a positive integer n , define a \mathbb{Q} -linear map $Z_{\mathcal{S}_n}^* : \mathfrak{H}^1 \rightarrow \overline{\mathcal{Z}}[t]/(t^n)$ by $Z_{\mathcal{S}_n}^*(e_{\mathbf{k}}) := \zeta_{\mathcal{S}_n}^*(\mathbf{k})$ for any index \mathbf{k} .

Lemma 4.3. For non-negative integers l and m , we have

$$\begin{aligned} & Z_{\mathcal{S}_2}^*((e_1 e_3)^l \tilde{\text{III}} e_2^m) \\ &= \sum_{\substack{2i+k+u=2l \\ j+n+v=m}} (-1)^{j+k} \binom{k+n}{k} \binom{u+v}{u} Z_{\mathcal{S}_2}((e_1 e_3)^i \tilde{\text{III}} e_2^j) Z_{\mathcal{S}_2}^*(e_2^{k+n}) Z_{\mathcal{S}_2}^*(e_2^{u+v}). \end{aligned}$$

Proof. This follows immediately from [Y, equation (3.1)] and (2.3) with $\mathcal{F}_n = \mathcal{S}_2$. \square

Proof of the $\mathcal{F} = \mathcal{S}$ case in Theorem 4.1. We prove (4.1) by induction on $l \geq 0$. The case $l = 0$ holds by the explicit evaluation of $\zeta_{\mathcal{S}_2}(\{2\}^r)$ ((3.8) with $k = 2$). Let l be a positive integer and m a non-negative integer. By [MOS, Lemma 2.1], we have

$$\begin{aligned} & Z_{\mathcal{S}_2}((e_1 e_3)^l \tilde{\text{III}} e_2^m) \\ &= 4^{-l} Z_{\mathcal{S}_2}(e_2^{l+m} \text{III} e_2^l) - \sum_{k=0}^{l-1} 4^{k-l} \binom{2l+m-2k}{l-k} Z_{\mathcal{S}_2}((e_1 e_3)^k \tilde{\text{III}} e_2^{2l+m-2k}). \end{aligned}$$

Hence, by Lemma 4.2 and the induction hypothesis, we have

$$\begin{aligned} & Z_{\mathcal{S}_2}((e_1 e_3)^l \tilde{\text{III}} e_2^m) \\ &= (-1)^m 2^{1-2l} \left\{ 1 - 2 \binom{4l+2m}{2l} \right\} \zeta(4l+2m+1)t \\ &\quad - \sum_{k=0}^{l-1} 4^{k-l} \binom{2l+m-2k}{l-k} \\ &\quad \cdot (-1)^m \left\{ (-1)^k 2^{1-2k} \binom{2l+m-k}{k} - 4 \binom{2l+m}{2k} \right\} \zeta(4l+2m+1)t \pmod{\zeta(2)}. \end{aligned}$$

We see that this coincides with the desired formula by using [MOS, Lemma 2.6]. We also obtain (4.2) by the same argument in [MOS] using (4.1), (3.9) with $k = 2$ and Lemma 4.3. \square

5. SUM FORMULAS

In this section, we prove the \mathcal{F}_n -symmetric sum formula (= Theorem 5.1), the \mathcal{F}_n -sum formula over $I_{k,r}$ for $n = 2, 3$ (= Theorem 5.2), and the \mathcal{F}_2 -sum formula over $I_{k,r,i}$ (= Theorem 5.4).

5.1. \mathcal{F}_n -symmetric sum formula. We first state the \mathcal{F}_n -symmetric sum formula.

Theorem 5.1 (\mathcal{F}_n -symmetric sum formula). *Let n and r be positive integers and $\mathbf{k} = (k_1, \dots, k_r)$ an index. Then, we have*

$$(5.1) \quad \sum_{\sigma \in \mathfrak{S}_r} \zeta_{\mathcal{F}_n}(\sigma(\mathbf{k})) = \sum_{\mathcal{B}=\{B_1, \dots, B_l\}} (-1)^{r-l} c(\mathcal{B}) \zeta_{\mathcal{F}_n}(b_1(\mathbf{k})) \cdots \zeta_{\mathcal{F}_n}(b_l(\mathbf{k})),$$

$$(5.2) \quad \sum_{\sigma \in \mathfrak{S}_r} \zeta_{\mathcal{F}_n}^*(\sigma(\mathbf{k})) = \sum_{\mathcal{B}=\{B_1, \dots, B_l\}} c(\mathcal{B}) \zeta_{\mathcal{F}_n}(b_1(\mathbf{k})) \cdots \zeta_{\mathcal{F}_n}(b_l(\mathbf{k})).$$

Here, \mathfrak{S}_r denotes the symmetric group of degree r . For $\sigma \in \mathfrak{S}_r$, set $\sigma(\mathbf{k}) := (k_{\sigma(1)}, \dots, k_{\sigma(r)})$. $\mathcal{B} = \{B_1, \dots, B_l\}$ runs all partitions of $\{1, \dots, r\}$, that is, $\mathcal{B} = \{B_1, \dots, B_l\}$ satisfies that $\{1, \dots, r\} = \bigsqcup_{i=1}^l B_i$ and $B_i \neq \emptyset$ ($1 \leq i \leq l$). Moreover, we set $c(\mathcal{B}) := (\#B_1 - 1)! \cdots (\#B_l - 1)!$ and $b_i(\mathbf{k}) := \sum_{j \in B_i} k_j$.

Proof. Since \mathcal{F}_n -MZVs satisfy the harmonic relation (2.3), we see that the desired formulas hold by the same argument as [H2, Theorem 4.1]. \square

5.2. \mathcal{F}_n -sum formula over $I_{k,r}$ for $n = 2, 3$. Next, we prove the \mathcal{F}_n -sum formula over $I_{k,r}$ for $n = 2, 3$. For positive integers n, r and k with $r \leq k$, $\mathcal{F} \in \{\mathcal{A}, \mathcal{S}\}$ and $\bullet \in \{\emptyset, \star\}$, set

$$S_{\mathcal{F}_n; k, r}^\bullet := \sum_{\mathbf{k} \in I_{k, r}} \zeta_{\mathcal{F}_n}^\bullet(\mathbf{k}),$$

where $I_{k,r}$ denotes the set of all indices \mathbf{k} with $\text{wt}(\mathbf{k}) = k$ and $\text{dep}(\mathbf{k}) = r$.

Theorem 5.2 (\mathcal{F}_n -sum formula over $I_{k,r}$ for $n = 2, 3$). *For positive integers r and k with $r \leq k$, we have*

$$(5.3) \quad S_{\mathcal{F}_2; k, r} = (-1)^{r-1} \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2}, \quad S_{\mathcal{F}_2; k, r}^\star = \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2}.$$

Moreover, we have

$$(5.4) \quad S_{\mathcal{F}_3; k, r} = (-1)^{k+r-1} \left[\binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+1)x_{\mathcal{F}_3} + \left\{ \frac{k+1}{2} \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+2) - \frac{1}{r!} \cdot T_{k,r} \right\} x_{\mathcal{F}_3}^2 \right]$$

and

$$(5.5) \quad S_{\mathcal{F}_3; k, r}^\star = (-1)^k \left[\binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+1)x_{\mathcal{F}_3} + \left\{ \frac{k+1}{2} \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+2) + \frac{1}{r!} \cdot T_{k,r} \right\} x_{\mathcal{F}_3}^2 \right],$$

where

$$T_{k,r} = \sum_{\substack{B_1 \sqcup B_2 = \{1, \dots, r\} \\ B_1, B_2 \neq \emptyset}} \sum_{\substack{b_1 + b_2 = k \\ b_1 \geq \#B_1, b_2 \geq \#B_2}} (b_1)_{\#B_1} (b_2)_{\#B_2} \cdot \mathfrak{Z}_{\mathcal{F}_3}(b_1+1) \mathfrak{Z}_{\mathcal{F}_3}(b_2+1)$$

and the symbol $(n)_m$ denotes $n(n-1) \cdots (n-m+1)$.

Remark 5.3. If $k = b_1 + b_2$ is odd, since $\mathfrak{Z}_{\mathcal{F}_3}(k+1)$ and $\mathfrak{Z}_{\mathcal{F}_3}(b_1+1)\mathfrak{Z}_{\mathcal{F}_3}(b_2+1)$ are 0, we have

$$S_{\mathcal{F}_3;k,r} = (-1)^r \frac{k+1}{2} \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+2)x_{\mathcal{F}_3}^2, \quad S_{\mathcal{F}_3;k,r}^* = -\frac{k+1}{2} \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+2)x_{\mathcal{F}_3}^2,$$

which were first proved by the third author and Yamamoto [SY, Theorem 2.5] for $\mathcal{F} = \mathcal{A}$.

Proof of Theorem 5.2. Since (5.3) is obtained from (5.4) and (5.5) by taking modulo $x_{\mathcal{F}_3}^2$, it is sufficient to prove (5.4) and (5.5). Note that $(-1)^k \mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2} = \mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2}$ holds because if k is odd, then $\mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2} = 0$.

Let us prove (5.4). By Theorem 3.2, we have

$$(5.6) \quad \zeta_{\mathcal{F}_3}(k) = (-1)^k \left\{ k \mathfrak{Z}_{\mathcal{F}_3}(k+1)x_{\mathcal{F}_3} + \binom{k+1}{2} \mathfrak{Z}_{\mathcal{F}_3}(k+2)x_{\mathcal{F}_3}^2 \right\}.$$

Since $x_{\mathcal{F}_3}^l$ with $l \geq 3$ vanishes, by (5.1), we have

$$\begin{aligned} S_{\mathcal{F}_3;k,r} &= \frac{1}{r!} \sum_{\mathbf{k} \in I_{k,r}} \sum_{\sigma \in \mathfrak{S}_r} \zeta_{\mathcal{F}_3}(\sigma(\mathbf{k})) \\ &= \frac{1}{r!} \sum_{\mathbf{k} \in I_{k,r}} \left\{ (-1)^{r-1} (r-1)! \zeta_{\mathcal{F}_3}(k) \right. \\ &\quad \left. + (-1)^{r-2} \sum_{\substack{B_1 \sqcup B_2 = \{1, \dots, r\} \\ B_1, B_2 \neq \emptyset}} (\#B_1 - 1)! (\#B_2 - 1)! \zeta_{\mathcal{F}_3}(b_1(\mathbf{k})) \zeta_{\mathcal{F}_3}(b_2(\mathbf{k})) \right\}. \end{aligned}$$

We calculate the right-hand side. Since $\#I_{k,r} = \binom{k-1}{r-1}$, by (5.6), we have

$$\sum_{\mathbf{k} \in I_{k,r}} \frac{(-1)^{r-1}}{r} \zeta_{\mathcal{F}_3}(k) = (-1)^{k+r-1} \left\{ \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+1)x_{\mathcal{F}_3} + \frac{k+1}{2} \binom{k}{r} \mathfrak{Z}_{\mathcal{F}_3}(k+2)x_{\mathcal{F}_3}^2 \right\}.$$

Furthermore, since $\#\{\mathbf{k} = (k_1, \dots, k_r) \in I_{k,r} \mid \sum_{i \in B_1} k_i = b_1\} = \#I_{b_1, \#B_1} \cdot \#I_{b_2, \#B_2}$ for $B_1, B_2 \neq \emptyset$ with $B_1 \sqcup B_2 = \{1, \dots, r\}$ and b_1, b_2 with $b_1 + b_2 = k$, $b_1 \geq \#B_1$, $b_2 \geq \#B_2$,

we have

$$\begin{aligned}
& \sum_{\mathbf{k} \in I_{k,r}} \sum_{\substack{B_1 \sqcup B_2 = \{1, \dots, r\} \\ B_1, B_2 \neq \emptyset}} (\#B_1 - 1)! (\#B_2 - 1)! \zeta_{\mathcal{F}_3}(b_1(\mathbf{k})) \zeta_{\mathcal{F}_3}(b_2(\mathbf{k})) \\
&= \sum_{\substack{B_1 \sqcup B_2 = \{1, \dots, r\} \\ B_1, B_2 \neq \emptyset}} \sum_{\substack{b_1 + b_2 = k \\ b_1 \geq \#B_1, b_2 \geq \#B_2}} \sum_{\substack{\mathbf{k} = (k_1, \dots, k_r) \in I_{k,r} \\ \sum_{i \in B_1} k_i = b_1}} (\#B_1 - 1)! (\#B_2 - 1)! \zeta_{\mathcal{F}_3}(b_1) \zeta_{\mathcal{F}_3}(b_2) \\
&= (-1)^k \sum_{\substack{B_1 \sqcup B_2 = \{1, \dots, r\} \\ B_1, B_2 \neq \emptyset}} \sum_{\substack{b_1 + b_2 = k \\ b_1 \geq \#B_1, b_2 \geq \#B_2}} (b_1)_{\#B_1} (b_2)_{\#B_2} \cdot \mathfrak{Z}_{\mathcal{F}_3}(b_1 + 1) \mathfrak{Z}_{\mathcal{F}_3}(b_2 + 1) x_{\mathcal{F}_3}^2.
\end{aligned}$$

Note that all terms of $x_{\mathcal{F}_3}^l$ with $l \geq 3$ for \mathcal{F}_3 -MZVs vanish. This completes the calculation for (5.4). The formula (5.5) is obtained by a similar calculation using (5.2). \square

5.3. \mathcal{F}_2 -sum formula over $I_{k,r,i}$. In this subsection, we prove the \mathcal{F}_2 -sum formula over $I_{k,r,i}$. For positive integers k, r, i with $1 \leq i \leq r < k$, let $I_{k,r,i}$ denote the set of indices $\mathbf{k} = (k_1, \dots, k_r)$ with $\text{wt}(\mathbf{k}) = k$, $\text{dep}(\mathbf{k}) = r$ and $k_i \geq 2$. For $\bullet \in \{\emptyset, \star\}$ and a positive integer n , set

$$S_{\mathcal{F}_n; k, r, i}^\bullet := \sum_{\mathbf{k} \in I_{k, r, i}} \zeta_{\mathcal{F}_n}^\bullet(\mathbf{k}).$$

Saito and Wakabayashi [SW1] ($\mathcal{F} = \mathcal{A}$) and Murahara [Mur] ($\mathcal{F} = \mathcal{S}$) proved that

$$\begin{aligned}
S_{\mathcal{F}_1; k, r, i} &= (-1)^i \left\{ \binom{k-1}{i-1} + (-1)^r \binom{k-1}{r-i} \right\} \mathfrak{Z}_{\mathcal{F}_1}(k), \\
S_{\mathcal{F}_1; k, r, i}^\star &= (-1)^i \left\{ \binom{k-1}{r-i} + (-1)^r \binom{k-1}{i-1} \right\} \mathfrak{Z}_{\mathcal{F}_1}(k).
\end{aligned}$$

If k is even, then we have $S_{\mathcal{F}_1; k, r, i} = S_{\mathcal{F}_1; k, r, i}^\star = 0$ by $\mathfrak{Z}_{\mathcal{F}_1}(k) = 0$. Thus it is a natural question what is a lifting of $S_{\mathcal{F}_1; k, r, i}^\bullet$ to \mathcal{F}_2 , that is, $S_{\mathcal{F}_2; k, r, i}^\bullet$. We give the answer in the following form.

Theorem 5.4 (\mathcal{F}_2 -sum formula for $I_{k,r,i}$). *Let k, r, i be positive integers with $1 \leq i \leq r < k$ and suppose that k is even. Then we have*

$$S_{\mathcal{F}_2; k, r, i} = (-1)^{r-1} \frac{b_{k,r,i}}{2} \cdot \mathfrak{Z}_{\mathcal{F}_2}(k+1) x_{\mathcal{F}_2}, \quad S_{\mathcal{F}_2; k, r, i}^\star = \frac{b_{k,r,i}^\star}{2} \cdot \mathfrak{Z}_{\mathcal{F}_2}(k+1) x_{\mathcal{F}_2},$$

where

$$b_{k,r,i} := \binom{k-1}{r} + (-1)^{r-i} \left\{ (k-r) \binom{k}{i-1} + \binom{k-1}{i-1} + (-1)^{r-1} \binom{k-1}{r-i} \right\}$$

and

$$b_{k,r,i}^* := \binom{k-1}{r} + (-1)^{i-1} \left\{ (k-r) \binom{k}{r-i} + \binom{k-1}{r-i} + (-1)^{r-1} \binom{k-1}{i-1} \right\}.$$

The case $\mathcal{F} = \mathcal{A}$ of Theorem 5.4 was proved by the third author and Yamamoto [SY]. In this subsection, we reprove their result and prove the case $\mathcal{F} = \mathcal{S}$ simultaneously by a different method.

Lemma 5.5 (Recurrence relations). For positive integers k, r, i with $2 \leq i+1 \leq r \leq k-1$, we have

$$\begin{aligned} (r-i)S_{\mathcal{F}_2;k,r,i} + iS_{\mathcal{F}_2;k,r,i+1} + (k-r)S_{\mathcal{F}_2;k,r-1,i} &= \sum_{l=1}^{k-r} \zeta_{\mathcal{F}_2}(l)S_{\mathcal{F}_2;k-l,r-1,i}, \\ (r-i)S_{\mathcal{F}_2;k,r,i}^* + iS_{\mathcal{F}_2;k,r,i+1}^* - (k-r)S_{\mathcal{F}_2;k,r-1,i}^* &= \sum_{l=1}^{k-r} \zeta_{\mathcal{F}_2}(l)S_{\mathcal{F}_2;k-l,r-1,i}^*. \end{aligned}$$

Proof. Let $\bullet \in \{\emptyset, \star\}$. From the same argument in [SW1, Lemma 2.1, Proposition 2.2], we see that the sum of the product

$$\sum_{(k_1, \dots, k_{r-1}, l) \in I_{k,r,i}} \zeta_{\mathcal{F}_2}(l) \zeta_{\mathcal{F}_2}^{\bullet}(k_1, \dots, k_{r-1}) = \sum_{l=1}^{k-r} \zeta_{\mathcal{F}_2}(l) S_{\mathcal{F}_2;k-l,r-1,i}^{\bullet}$$

coincides with the left-hand side of the desired recurrence relation by the harmonic relation for \mathcal{F}_2 -MZVs. \square

Corollary 5.6. If k is even, then we have

$$\begin{aligned} (r-i)S_{\mathcal{F}_2;k,r,i} + iS_{\mathcal{F}_2;k,r,i+1} + (k-r)S_{\mathcal{F}_2;k,r-1,i} &= 0, \\ (r-i)S_{\mathcal{F}_2;k,r,i}^* + iS_{\mathcal{F}_2;k,r,i+1}^* - (k-r)S_{\mathcal{F}_2;k,r-1,i}^* &= 0 \end{aligned}$$

for positive integers k, r, i with $2 \leq i+1 \leq r \leq k-1$.

Proof. If l is odd, then $\zeta_{\mathcal{F}_2}(l) = 0$ by Theorem 3.2 or (3.8). If l is even, then $\zeta_{\mathcal{F}_2}(l)$ is a multiple of $x_{\mathcal{F}}$ by Theorem 3.2 or (3.8) and $S_{\mathcal{F}_2;k-l,r-1,i}^{\bullet}$ is also a multiple of $x_{\mathcal{F}}$ by Saito–Wakabayashi and Murahara’s sum formulas. \square

Proof of Theorem 5.4. We prove the non-star case by backward induction on $r \leq k-1$. Since

$$\begin{aligned} b_{k,k-1,i} &= \binom{k-1}{k-1} + (-1)^{k-1-i} \left\{ (k-k+1) \binom{k}{i-1} + \binom{k-1}{i-1} + (-1)^{k-2} \binom{k-1}{k-1-i} \right\} \\ &= 1 + (-1)^{i-1} \binom{k+1}{i}, \end{aligned}$$

we have

$$\begin{aligned} S_{\mathcal{F}_2;k,k-1,i} &= \zeta_{\mathcal{F}_2}(\{1\}^{i-1}, 2, \{1\}^{k-i-1}) = \frac{1}{2} \left\{ 1 + (-1)^{i-1} \binom{k+1}{i} \right\} \mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2} \\ &= \frac{b_{k,k-1,i}}{2} \cdot \mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2}, \end{aligned}$$

by the definition of $S_{\mathcal{F}_2;k,r,i}$ and (3.24). Hence, the case $r = k - 1$ is true. To complete the induction step, by Corollary 5.6, it suffices to prove that

$$(5.7) \quad (r-i)b_{k,r,i} + ib_{k,r,i+1} - (k-r)b_{k,r-1,i} = 0$$

holds for $2 \leq r \leq k - 1$. The left-hand side of (5.7) is

$$\begin{aligned} &(r-i) \left[\binom{k-1}{r} + (-1)^{r-i} \left\{ (k-r) \binom{k}{i-1} + \binom{k-1}{i-1} + (-1)^{r-1} \binom{k-1}{r-i} \right\} \right] \\ &+ i \left[\binom{k-1}{r} + (-1)^{r-i-1} \left\{ (k-r) \binom{k}{i} + \binom{k-1}{i} + (-1)^{r-1} \binom{k-1}{r-i-1} \right\} \right] \\ &- (k-r) \left[\binom{k-1}{r-1} + (-1)^{r-i-1} \left\{ (k-r+1) \binom{k}{i-1} + \binom{k-1}{i-1} + (-1)^{r-2} \binom{k-1}{r-i-1} \right\} \right] \end{aligned}$$

by definition. By $\binom{k-1}{r-1} = \frac{r}{k-r} \binom{k-1}{r}$, we have

$$(5.8) \quad (r-i) \binom{k-1}{r} + i \binom{k-1}{r} - (k-r) \binom{k-1}{r-1} = 0.$$

By $\binom{k}{i} = \frac{k-i+1}{i} \binom{k}{i-1}$, we have

$$(5.9) \quad (r-i)(k-r) \binom{k}{i-1} - i(k-r) \binom{k}{i} + (k-r)(k-r+1) \binom{k}{i-1} = 0.$$

By $\binom{k-1}{i} = \frac{k-i}{i} \binom{k-1}{i-1}$, we have

$$(5.10) \quad (r-i) \binom{k-1}{i-1} - i \binom{k-1}{i} + (k-r) \binom{k-1}{i-1} = 0.$$

By $\binom{k-1}{r-i} = \frac{k-r+i}{r-i} \binom{k-1}{r-i-1}$, we have

$$(5.11) \quad (r-i) \binom{k-1}{r-i} - i \binom{k-1}{r-i-1} - (k-r) \binom{k-1}{r-i-1} = 0.$$

From (5.8), (5.9), (5.10) and (5.11), we obtain (5.7) and we complete the proof of the formula for $S_{\mathcal{F}_2;k,r,i}$. In the star case, we should prove

$$S_{\mathcal{F}_2;k,k-1,i}^* = \frac{b_{k,k-1,i}^*}{2} \cdot \mathfrak{Z}_{\mathcal{F}_2}(k+1)x_{\mathcal{F}_2}$$

and the recurrence relation

$$(r-i)b_{k,r,i}^* + ib_{k,r,i+1}^* - (k-r)b_{k,r-1,i}^* = 0.$$

These are proved similarly to the non-star case. \square

Remark 5.7. We can also prove the star case by connecting $S_{\mathcal{F}_2;k,r,i}$ and $S_{\mathcal{F}_2;k,r,i}^*$ directly using Proposition 2.5. This is the method used in [SY].

APPENDIX A. PROOF OF EQUALITY (3.26)

In this appendix, we prove the following proposition.

Proposition A.1. For non-negative integers a and b , we have

$$C = 1 + (-1)^a \binom{a+b+3}{b+2};$$

see the proof of Theorem 3.13 for the definition of C .

We divide C into six parts. Set

$$\begin{aligned} \text{I} &:= (-1)^{a+1} \sum_{\substack{l+m=b-1 \\ l,m \geq 0}} \sum_{\substack{r+s=a \\ r,s \geq 0}} (-1)^{s+m} \binom{r+l+1}{r} \binom{s+m+1}{s} \binom{a+b+3}{s+m+2}, \\ \text{II} &:= (-1)^{a+1} \sum_{\substack{l+m=b-1 \\ l,m \geq 0}} \sum_{\substack{r+s=a \\ r,s \geq 0}} \binom{r+l+1}{r} \binom{s+m+1}{s}, \\ \text{III} &:= 2(-1)^a \sum_{\substack{r+s=a \\ r,s \geq 0}} (-1)^s \binom{r+b+1}{r} \binom{a+b+3}{s+1}, \\ \text{IV} &:= 2(-1)^{a+1} \sum_{\substack{r+s=a \\ r,s \geq 0}} \binom{r+b+1}{r}, \\ \text{V} &:= \sum_{\substack{m+n=a-1 \\ m,n \geq 0}} (-1)^n \sum_{\substack{r+s=n \\ r,s \geq 0}} (-1)^{s+m+1} \binom{r+b+1}{r} \binom{s+m+1}{s} \binom{a+b+3}{s+m+2}, \\ \text{VI} &:= \sum_{\substack{m+n=a-1 \\ m,n \geq 0}} (-1)^{n+1} \sum_{\substack{r+s=n \\ r,s \geq 0}} \binom{r+b+1}{r} \binom{s+m+1}{s}. \end{aligned}$$

Note that we easily obtain

$$(A.1) \quad \text{IV} = 2(-1)^{a+1} \binom{a+b+2}{a}.$$

By the definition of negative binomial coefficients and the Chu–Vandermonde identity, we also have

$$(A.2) \quad \text{II} = (-1)^{a+1} b \binom{a+b+2}{a}, \quad \text{VI} = (-1)^a \binom{a+b+1}{a-1}.$$

Next, we calculate I, III and V. We use the following equality repeatedly:

$$(A.3) \quad \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{1}{x+k} = \frac{n!}{x(x+1)\cdots(x+n)}.$$

Here, n is a non-negative integer and x is an indeterminate.

Lemma A.2.

$$\text{I} = 0.$$

Proof. By elementary calculations, we have

$$(A.4) \quad \begin{aligned} \text{I} &= (-1)^{a+1} \sum_{m=0}^{b-1} \sum_{s=0}^a (-1)^{s+m} \binom{a+b-m-s}{b-m} \binom{s+m+1}{s} \binom{a+b+3}{s+m+2} \\ &= (-1)^{a+1} (a+1) \binom{a+b+2}{a+1} \sum_{s=0}^a (-1)^s \binom{a}{s} \\ &\quad \times \sum_{m=0}^{b-1} (-1)^m \binom{b+1}{m+1} \left(\frac{1}{a+b+1-s-m} + \frac{1}{s+m+2} \right). \end{aligned}$$

Applying (A.3) with $x = -(a+b+2-s)$ and $x = s+1$, we have

$$(A.5) \quad \begin{aligned} &\sum_{m=0}^{b-1} (-1)^m \binom{b+1}{m+1} \left(\frac{1}{a+b+1-s-m} + \frac{1}{s+m+2} \right) \\ &= \sum_{m=0}^{b+1} (-1)^{m-1} \binom{b+1}{m} \left(\frac{1}{a+b+2-s-m} + \frac{1}{s+m+1} \right) \\ &\quad + \left(\frac{1}{a+b+2-s} + \frac{1}{s+1} \right) + (-1)^{b+1} \left(\frac{1}{a+1-s} + \frac{1}{s+b+2} \right) \\ &= \frac{(-1)^b}{a+b+2-s} \binom{a+b+1-s}{b+1}^{-1} - \frac{1}{s+1} \binom{b+s+2}{b+1}^{-1} \\ &\quad + \frac{1}{a+b+2-s} + \frac{1}{s+1} + \frac{(-1)^{b+1}}{a+1-s} + \frac{(-1)^{b+1}}{s+b+2}. \end{aligned}$$

By substituting (A.5) into (A.4) and then using (A.3) again, we have

$$\begin{aligned}
(-1)^{a+b+1} \sum_{s=0}^a (-1)^s \binom{a+b+2}{s} &= (-1)^{b+1} \binom{a+b+1}{a}, \\
(-1)^a \sum_{s=0}^a (-1)^s \binom{a+b+2}{a-s} &= \sum_{s=0}^a (-1)^s \binom{a+b+2}{s} = (-1)^a \binom{a+b+1}{a}, \\
(-1)^{a+1} (a+1) \binom{a+b+2}{a+1} \sum_{s=0}^a (-1)^s \binom{a}{s} \frac{1}{a+b+2-s} &= -1, \\
(-1)^{a+1} (a+1) \binom{a+b+2}{a+1} \sum_{s=0}^a (-1)^s \binom{a}{s} \frac{1}{s+1} &= (-1)^{a+1} \binom{a+b+2}{a+1}, \\
(-1)^{a+b} (a+1) \binom{a+b+2}{a+1} \sum_{s=0}^a (-1)^s \binom{a}{s} \frac{1}{a+1-s} &= (-1)^b \binom{a+b+2}{a+1}, \\
(-1)^{a+b} (a+1) \binom{a+b+2}{a+1} \sum_{s=0}^a (-1)^s \binom{a}{s} \frac{1}{s+b+2} &= (-1)^{a+b}.
\end{aligned}$$

Since $a+b$ is even, we have the conclusion. \square

Lemma A.3.

$$\text{III} = 2 + 2(-1)^a \binom{a+b+2}{a+1}.$$

Proof. By (A.3), we have

$$\begin{aligned}
\text{III} &= 2(-1)^a \sum_{s=0}^a (-1)^s \binom{a+b+1-s}{a-s} \binom{a+b+3}{s+1} \\
&= 2(-1)^{a-1} (a+b+3) \binom{a+b+2}{a+1} \left\{ \sum_{s=0}^{a+1} (-1)^s \binom{a+1}{s} \frac{1}{a+b+3-s} - \frac{1}{a+b+3} \right\} \\
&= 2(-1)^{a-1} (a+b+3) \binom{a+b+2}{a+1} \left\{ \frac{(-1)^{a-1}}{a+b+3} \binom{a+b+2}{a+1}^{-1} - \frac{1}{a+b+3} \right\} \\
&= 2 + 2(-1)^a \binom{a+b+2}{a+1},
\end{aligned}$$

which completes the proof. \square

Lemma A.4.

$$\text{V} = (-1)^a a \binom{a+b+2}{a+1} + (-1)^a \binom{a+b+1}{a} - 1.$$

Proof. Since

$$\begin{aligned}
V &= (-1)^a \sum_{n=0}^{a-1} \sum_{s=0}^n (-1)^s \binom{b+n+1-s}{b+1} \binom{a+s-n}{s} \binom{a+b+3}{a+s+1-n} \\
&= (-1)^a (a+1) \binom{a+b+2}{a+1} \sum_{n=0}^{a-1} \binom{a}{n} \\
&\quad \times \sum_{s=0}^n (-1)^s \binom{n}{s} \left(\frac{1}{a+s+1-n} + \frac{1}{b+2+n-s} \right)
\end{aligned}$$

and

$$\begin{aligned}
\sum_{s=0}^n (-1)^s \binom{n}{s} \frac{1}{a+s+1-n} &= \frac{1}{a+1-n} \binom{a+1}{n}^{-1}, \\
\sum_{s=0}^n (-1)^s \binom{n}{s} \frac{1}{b+2+n-s} &= \frac{(-1)^n}{b+2+n} \binom{b+n+1}{b+1}^{-1}
\end{aligned}$$

hold by (A.3), we obtain the desired formula. \square

Proof of Proposition 3.26. From (A.1), (A.2), Lemmas A.2, A.3, and A.4, we obtain the desired formula. \square

REFERENCES

- [BB] D. Bowman, D. M. Bradley, *The algebra and combinatorics of shuffles and multiple zeta values*, J. Combin. Theory Ser. A **97** (2002), 43–61.
- [HHT] KH. Hessami Pilehrood, T. Hessami Pilehrood, R. Tauraso, *New properties of multiple harmonic sums modulo p and p -analogues of Leshchiner’s series*, Trans. Amer. Math. Soc. **366** (2014), 3131–3159.
- [HMO] M. Hirose, H. Murahara, M. Ono, *On variants of symmetric multiple zeta-star values and the cyclic sum formula*, to appear in Ramanujan J.
- [H1] M. E. Hoffman, *The algebra of multiple harmonic series*, J. Algebra **194** (1997), 477–495.
- [H2] M. E. Hoffman, *Quasi-symmetric functions and mod p multiple harmonic sums*, Kyushu J. Math. **69** (2015), 345–366.
- [IKOO] K. Ihara, J. Kajikawa, Y. Ohno, J. Okuda, *Multiple zeta values vs. multiple zeta-star values*, J. Algebra **332** (2011), 187–208.
- [IKZ] K. Ihara, M. Kaneko, D. Zagier, *Derivation relation and double shuffle relations for multiple zeta values*, Compositio Math. **142** (2006), 307–338.
- [J1] D. Jarossay, *Double mélange des multizêtas finis et multizêtas symétrisés*, C. R. Acad. Sci. Paris, **352** (2014), 767–771.
- [J2] D. Jarossay, *Adjoint cyclotomic multiple zeta values and cyclotomic multiple harmonic values*, preprint, arXiv:1412.5099v5.

- [Kan] M. Kaneko, *An introduction to classical and finite multiple zeta values*, Publications mathématiques de Besançon, no. 1 (2019), 103–129.
- [KS] M. Kaneko, M. Sakata, *On multiple zeta values of extremal height*, Bull. Aust. Math. Soc. **93** (2016), 186–193.
- [KZ] M. Kaneko, D. Zagier, *Finite multiple zeta values*, in preparation.
- [Li] Z. Li, *Gamma series associated to elements satisfying regularized double shuffle relations*, J. Number Theory **130** (2010), 213–231.
- [Mun] S. Muneta, *A note on evaluations of multiple zeta values*, Proc. Amer. Math. Soc. **137** (2009), 931–935.
- [Mur] H. Murahara, *A note on finite real multiple zeta values*, Kyushu J. Math., **70** (2016), 197–204.
- [MOS] H. Murahara, T. Onozuka, S. Seki, *Bowman-Bradley type theorem for finite multiple zeta values in \mathcal{A}_2* , Osaka Journal of Mathematics **57** (2020), 647–653.
- [OSY] M. Ono, S. Seki, S. Yamamoto, *Truncated t -adic symmetric multiple zeta values and double shuffle relations*, Res. number theory **7**, 15 (2021).
- [Re] C. Reutenauer, *Free Lie Algebras*, Oxford Science Publications (1993).
- [Ro] J. Rosen, *Asymptotic relations for truncated multiple zeta values*, J. Lond. Math. Soc. (2) **91** (2015), 554–572. Int. Math. Res. Notices **23** (2019), 7379–7405.
- [SW1] S. Saito, N. Wakabayashi, *Sum formula for finite multiple zeta values*, J. Math. Soc. Japan **67** (2015), 1069–1076.
- [SW2] S. Saito, N. Wakabayashi, *Bowman-Bradley type theorem for finite multiple zeta values*, Tohoku Math. J. (2) **68** (2016), 241–251.
- [SS] K. Sakugawa, S. Seki, *On functional equations of finite multiple polylogarithms*, J. Algebra **469** (2017), 323–357.
- [Se] S. Seki, *The p -adic duality for the finite star-multiple polylogarithms*, Tohoku Math. J. **71** (2019), 111–122.
- [SY] S. Seki, S. Yamamoto, *Ohno-type identities for multiple harmonic sum*, J. Math. Soc. Japan **72** (2020), 673–686.
- [Su] Z. H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Disc. Appl. Math. **105** (2000), 193–223.
- [TT] Y. Takeyama, K. Tasaka, *Supercongruences of multiple harmonic q -sums and generalized finite/symmetric multiple zeta values*, preprint, arXiv:2012.07067.
- [TY] K. Tasaka, S. Yamamoto, *On some multiple zeta-star values of one-two-three indices*, Int. J. Number Theory **9** (2013), 1171–1184.
- [Tau] R. Tauraso, *More congruences for central binomial coefficients*, J. Number Theory **130** (2010), 2639–2649.
- [W] L. C. Washington, *p -adic L -functions and sums of powers*, J. Number Theory **69** (1998), 50–61.
- [Y] S. Yamamoto, *Explicit evaluation of certain sums of multiple zeta-star values*, Funct. Approx. Comment. Math. **49** (2) (2013), 283–289.
- [Z] D. Zagier, *Evaluation of the multiple zeta values $\zeta(2, \dots, 2, 3, 2, \dots, 2)$* , Ann. of Math. **175** (2012), 977–1000.

- [ZC] X. Zhou, T. Cai, *A generalization of a curious congruence on harmonic sums*, Proc. of Amer. Math. Soc. **135** (2007), no. 5, 1329–1333.
- [Zh] J. Zhao, *Wolstenholme type theorem for multiple harmonic sums*, Int. J. Number Theory **4** (2008), 73–106.

(Masataka Ono) GLOBAL EDUCATION CENTER, WASEDA UNIVERSITY, 1-6-1 NISHI-WASEDA, SHINJUKU-KU, TOKYO, 169-8050, JAPAN

Email address: m-ono@aoni.waseda.jp

(Kosuke Sakurada) SEIWA GAKUEN HIGH SCHOOL, 3-4-1, KINOSHITA, WAKABAYASHI-KU, SENDAI, 984-0047, JAPAN

Email address: sakurada.kosuke@seiwa.ac.jp

(Shin-ichiro Seki) DEPARTMENT OF MATHEMATICAL SCIENCES, AOYAMA GAKUIN UNIVERSITY, 5-10-1 FUCHINOBE, CHUO-KU, SAGAMIHARA-SHI, KANAGAWA, 252-5258, JAPAN

Email address: seki@math.aoyama.ac.jp