# VALUE DISTRIBUTION AND UNIQUENESS FOR Q-DIFFERENCE OF MEROMORPHIC FUNCTIONS SHARING TWO SETS

#### GOUTAM HALDAR

ABSTRACT. In this paper, we investigate the value distribution for linear q-difference polynomials of transcendental meromorphic functions of zero order which improves the results of Xu, Liu and Cao ([26]). We also investigate the uniqueness of zero order meromorphic function with its q-difference operator sharing two sets with finite weight. Some examples have been exhibited which are relevant to the content of the paper.

### 1. Introduction

Let f and g be two non-constant meromorphic functions defined in the open complex plane  $\mathbb{C}$ . If for some  $a \in \mathbb{C} \cup \{\infty\}$ , the zero of f-a and g-a have the same locations as well as same multiplicities, we say that f and g share the value g CM (counting multiplicities). If we do not consider the multiplicities, then g and g are said to share the value g IM (ignoring multiplicities). We adopt the standard notations of the Nevanlinna theory of meromorphic functions explained in ([10]).

A meromorphic function a is said to be a small function of f provided that T(r,a) = S(r,f). i.e., T(r,a) = o(T(r,f)) as  $r \to \infty$ , outside of a possible exceptional set of finite linear measure.

For a set  $S \subset \mathbb{C}$ , we define

$$E_f(S) = \bigcup_{a \in S} \{z | f(z) = a(z)\},$$

where each zero is counted according to its multiplicity and

$$\overline{E}_f(S) = \bigcup_{a \in S} \{z | f(z) = a(z)\}, \text{ where each zero is counted only once.}$$

If  $E_f(S) = E_g(S)$ , we say that f, g share the set S CM and if  $\overline{E}_f(S) = \overline{E}_g(S)$ , we say f, g share the set S IM.

In 2001, Lahiri ([11]) introduced a gradation of sharing of values or sets which is known as weighted sharing. Below we are recalling the notion.

**Definition 1.1.** ([11]) Let k be a non-negative integer or infinity. For  $a \in \mathbb{C} \cup \{\infty\}$  we denote by  $E_k(a, f)$  the set of all a-points of f, where an a point of multiplicity m is counted m times if  $m \leq k$  and k + 1 times if m > k. If  $E_k(a, f) = E_k(a, g)$ , we say that f, g share the value a with weight k.

<sup>2010</sup> Mathematics Subject Classification. 30D35.

Key words and phrases. Meromorphic function, q-difference operator, small function, weighted sharing, zero-order.

We write f, g share (a, k) to mean that f, g share the value a with weight k. Clearly if f, g share (a, k) then f, g share (a, p) for any integer p,  $0 \le p < k$ . Also we note that f, g share a value a IM or CM if and only if f, g share (a, 0) or  $(a, \infty)$  respectively.

**Definition 1.2.** [14] Let p be a positive integer and  $a \in \mathbb{C} \cup \{\infty\}$ .

- (i)  $N(r, a; f \mid \geq p)$   $(\overline{N}(r, a; f \mid \geq p))$  denotes the counting function (reduced counting function) of those a-points of f whose multiplicities are not less than p.
- (ii)  $N(r, a; f \mid \leq p)$  ( $\overline{N}(r, a; f \mid \leq p)$ ) denotes the counting function (reduced counting function) of those a-points of f whose multiplicities are not greater than p.

**Definition 1.3.** [1] Let f and g be two non-constant meromorphic functions such that f and g share the value a IM. Let  $z_0$  be a a-point of f with multiplicity p, a a-point of g with multiplicity q. We denote by  $\overline{N}_L(r,a;f)$  the counting function of those a-points of f and g where p > q, by  $N_E^{(1)}(r,a;f)$  the counting function of those a-points of f and g where p = q = 1 and by  $\overline{N}_E^{(2)}(r,a;f)$  the counting function of those a-points of f and g where  $p = q \ge 2$ , each point in these counting functions is counted only once. Similarly, one can define  $\overline{N}_L(r,a;g)$ ,  $N_E^{(1)}(r,a;g)$ ,  $\overline{N}_E^{(2)}(r,a;g)$ .

**Definition 1.4.** [11, 12] Let f, g share a value a IM. We denote by  $\overline{N}_*(r, a; f, g)$  the reduced counting function of those a-points of f whose multiplicities differ from the multiplicities of the corresponding a-points of g. Clearly  $\overline{N}_*(r, a; f, g) \equiv \overline{N}_*(r, a; g, f)$  and  $\overline{N}_*(r, a; f, g) = \overline{N}_L(r, a; f) + \overline{N}_L(r, a; g)$ .

Recently, many of authors have shown their interest in studying difference equations, the difference product and the q-difference analogues the value distribution theory in the complex plane  $\mathbb{C}$ . A number of remarkable research works (see [5], [6], [7], [8], [9], [16], [17], [18], [19], [22], [25], [31]) have focused on the uniqueness of difference analogues of Nevanlinna theory. In 2006, Halburd and Korhonen ([7]) established a difference analogue of the Logarithmic Derivative Lemma, and then applying it, a lot of results on meromorphic solutions of complex difference equations has been proved. After that Barnett, Halburd, Korhonen and Morgan ([4]) also established a q-difference analogue of the Logarithmic Derivative Lemma.

Let us first recall the notion of the q-shift and q-difference operator of a meromorphic function f.

**Definition 1.5.** For a meromorphic function f and c,  $q \neq 0$   $\in \mathbb{C}$ , let us now denote its q-shift  $E_{q,c}f$  and q-difference operators  $\Delta_{q,c}f$  respectively by  $E_{q,c}f(z) = f(qz+c)$  and  $\Delta_{q,c}f(z) = f(qz+c) - f(z)$ ,  $\Delta_{q,c}^kf(z) := \Delta_{q,c}^{k-1}(\Delta_{q,c}f(z))$ , for all  $k \in \mathbb{N} - \{1\}$ .

We now proceed to define linear q-shift and q-difference operators, denoted respectively by  $L_k(f, E_q)$  and  $L_k(f, \Delta)$  of a meromorphic function f in a more compact and convenient way in the following, which is one of the motivation of writing the paper.

**Definition 1.6.** Let us define

$$L_k(f, E_q) = a_k f(q_k z + c_k) + a_{k-1} f(q_{k-1} z + c_{k-1}) + \dots + a_0 f(q_0 z + c_0)$$
 (1.1)  
and

$$L_k(f,\Delta) = a_k \Delta_{q_k,c_k} f(z) + a_{k-1} \Delta_{q_{k-1},c_{k-1}} f(z) + \dots + a_0 \Delta_{q_0,c_0} f(z), \tag{1.2}$$

where  $a_0, a_1, \ldots, a_k; q_0, q_1, \ldots, q_k; c_0, c_1, \ldots, c_k$  are complex constants. From (1.2), one can easily observe that

$$L_k(f, \Delta) = L_k(f, E_q) - \sum_{j=0}^{k} a_j f(z).$$

If we choose  $q_j = q^j$ ,  $c_j = c$  and  $a_j = (-1)^{k-j} {k \choose j}$  for  $0 \le j \le k$ , then  $L_k(f, \Delta)$  reduces  $\Delta_{q,c}^k f(z)$ .

Let  $P(z) = a_n z^n + a_{n-1} z^{n-1} + \ldots + a_0$  be a nonzero polynomial of degree n, where  $a_n \neq 0$ ,  $a_{n-1}, \ldots, a_0$  are complex constants and m be the number of distinct zeros of P(z). From now on, unless otherwise stated, for  $a \neq 0$ , we denote, by  $S_1 = \{a, aw, aw^2, \ldots, aw^{n-1}\}$ , where  $n \in \mathbb{N}$ , w is a n-th roots of unity and  $S_2 = \{\infty\}$  throughout the paper.

Zhang and Korhonen ([32]) studied the value distribution of q-difference polynomials of meromorphic functions and obtained the following result.

**Theorem A.** [32] Let f be a transcendental meromorphic (resp. entire) function of zero order and q non-zero complex constant. Then for  $n \geq 6$  (resp.  $n \geq 2$ ),  $f(z)^n f(qz)$  assumes every non-zero value  $a \in \mathbb{C}$  infinitely often.

Recently, Liu and Qi [21] firstly investigated value distributions for a q-shift of the meromorphic function and obtained the following result.

**Theorem B.** [21] Let f be a zero-order transcendental meromorphic function,  $n \geq 6$ ;  $q \in \mathbb{C} - \{0\}$ ,  $\eta \in \mathbb{C}$  and R(z) a rational function. Then  $f(z)^n f(qz+\eta) - R(z)$  has infinitely many zeros.

In 2015, Xu, Liu and Cao [26] started investigation about the zeros of  $P(f)f(qz+\eta) = a(z)$  and  $P(f)[f(qz+\eta) - f(z)] = a(z)$ , where a(z) is a small function of f and obtained the following two results.

**Theorem C.** [26] Let f be a zero-order transcendental meromorphic (resp. entire) function,  $q(\neq 0)$ ,  $\eta$  are complex constants. Then for n > m+4 (resp. n > m),  $P(f)f(qz+\eta)=a(z)$  has infinitely many solutions, where a(z) is a non-zero small functions in f.

**Theorem D.** [26] Let f be a zero-order transcendental meromorphic (resp. entire) function,  $q \neq 0$ ,  $\eta$  are complex constants. Then for n > m+6 (resp. n > m+2),  $P(f)\{f(qz+\eta)-f(z)\}=a(z)$  has infinitely many solutions, where a(z) is a non-zero small functions in f.

Regarding uniqueness, in 2011, Qi, Liu and Yang [24] obtained the following result

**Theorem E.** [24] Let f be a zero-order meromorphic function, and  $q \in \mathbb{C} - \{0\}$ ,  $n \geq 4$  be an integer, and let  $F = f^n$ . If F(z) and F(qz) share  $a \in \mathbb{C} - \{0\}$  and  $\infty$  CM, then f(z) = tf(qz) for a constant t that satisfies  $t^n = 1$ .

**Question 1.1.** What would happen if we replace P(f)f(qz+c) in Theorem C by more general q-shift difference polynomial  $P(f)L_k(f, E_k)$  and  $P(f)\{f(qz+c)-f(z)\}$  in Theorem D by  $P(f)L_k(f, \Delta)$ ?

**Question 1.2.** what would happen if we replace  $F = f^n$  in the above theorem by more general polynomial P(f), where P(z) is defined above?

In this paper, we try to find out the possible answer to the above questions, and also investigated the uniqueness of a zero-order meromorphic function f and its linear q-difference operator sharing sets  $S_1$  and  $S_2$  with finite weights. The next section includes our main results.

#### 2. Main Results

**Theorem 2.1.** Let f be a transcendental meromorphic of zero order (resp. entire) function, and  $\alpha(z)$  be a non-zero small function of f(z). Then for n > m + 3k + 4 (resp. n > m + k),  $P(f)L_k(f, E_q) - a(z)$  has infinitely many zeros.

**Theorem 2.2.** Let f be a zero-order transcendental meromorphic (resp. entire) function, and  $\alpha(z)$  be a non-zero small function of f(z). Then for n > m + 5k + 6 (resp. n > m + 2k + 2),  $P(f)L_k(f, \Delta) - \alpha(z)$  has infinitely many zeros.

**Theorem 2.3.** Let f be a meromorphic function of zero-order,  $q \in \mathbb{C} - \{0\}$ , M(z) = P(f(z)). If M(z) and M(qz) share a non-zero complex constant a and  $\infty$  CM, then P(f(z)) = P(f(qz)).

**Theorem 2.4.** Let  $n, k \in \mathbb{N}$ , and f be a non-constant zero order meromorphic function such that  $E_f(S_1, 1) = E_{L_k(f, \Delta_q)}(S_1, 1)$  and  $E_f(S_2, 0) = E_{L_k(f, \Delta_q)}(S_2, 0)$ . If  $n \geq 7$ , then there exists a constant  $t \in \mathbb{C}$  such that  $L_k(f, \Delta_q) \equiv tf$ , where  $t^n = 1$  and  $t \neq -1$ .

**Theorem 2.5.** Let  $n, k \in \mathbb{N}$ , and f be a non-constant zero order meromorphic function such that  $E_f(S_1, 2) = E_{L_k(f, \Delta_q)}(S_1, 2)$  and  $E_f(S_2, 0) = E_{L_k(f, \Delta_q)}(S_2, 0)$ . If  $n \geq 6$ , then there exists a constant  $t \in \mathbb{C}$  such that  $L_k(f, \Delta_q) \equiv tf$ , where  $t^n = 1$  and  $t \neq -1$ .

**Remark 2.1.** Clearly, Theorem 2.1 and Theorem 2.2 are the improvements of Theorem C and Theorem D, respectively.

**Remark 2.2.** The zero order growth restriction in Theorem 2.2 can not be extended to finite order. This can be observed by taking  $P(z) = z^n$ ,  $f(z) = e^z$ , c = 0 and q = -n. Then P(f(z))[f(qz) - f(z)] - 1 have no zeros.

Remark 2.3. Obviously, Theorem 2.3 is an improvement of Theorem E.

**Remark 2.4.** If we closely observe the statement of Theorem E, the we see that if a zero order meromorphic function f(z) and its q-shift operator f(qz) share the sets  $S_1$  and  $S_2$  CM, then f(z) = tf(qz) for a constant t such that  $t^n = 1$ .

The following examples show that Theorem 2.4 and 2.5 hold for n = 7 and n = 6 respectively, for both entire and meromorphic functions.

**Example 2.1.** For a positive integer m, let  $f(z) = z^m$ ,  $L_k(f, \Delta) = f(qz) - f(z)$ , where  $q = (1 + \omega)^{\frac{1}{m}}$ ,  $\omega^7 = 1$  ( $\omega^6 = 1$  for Theorem 2.5). Then one can easily verify that f(z) and  $L_k(f, \Delta)$  share the sets  $S_1$  and  $S_2$  and  $L_k(f, \Delta) = \omega f(z)$  such that  $\omega \neq -1$ ,  $\omega^7 = 1$  ( $\omega^6 = 1$  for Theorem1.5).

**Example 2.2.** For a positive integer m, let  $f(z) = \frac{1}{z^m}$ ,  $L_k(f, \Delta) = f(qz) - f(z)$ , where  $q = \frac{1}{(1+\omega)^{\frac{1}{m}}}$ ,  $\omega^7 = 1$  ( $\omega^6 = 1$  for Theorem 2.5). Then one can easily verify that f(z) and  $L_k(f, \Delta)$  share the sets  $S_1$  and  $S_2$  and  $L_k(f, \Delta) = \omega f(z)$  such that  $\omega \neq -1$ ,  $\omega^7 = 1$  ( $\omega^6 = 1$  for Theorem1.5).

#### 3. Some Lemmas

We now prove several lemmas which will play key roles in proving the main results of the paper. Let  $\mathcal{F}$  and  $\mathcal{G}$  be two non-constant meromorphic functions defined by

$$F = \left(\frac{f(z)}{a}\right)^n \text{ and } G = \left(\frac{L_k(f, \Delta)}{a}\right)^n.$$
 (3.1)

Henceforth we shall denote by H and V in the following.

$$H = \left(\frac{F'''}{F} - \frac{2F'}{F-1}\right) - \left(\frac{G''}{G'} - \frac{2G'}{G-1}\right),\tag{3.2}$$

$$V = \frac{F'}{F(F-1)} - \frac{G'}{G(G-1)}. (3.3)$$

**Lemma 3.1.** [21] Let f be a zero-order meromorphic function, and  $q(\neq 0)$ ,  $c \in \mathbb{C}$ . Then

$$m\left(r, \frac{f(qz+c)}{f(z)}\right) = S(r, f).$$

**Lemma 3.2.** [26] Let f(z) be a transcendental meromorphic function of zero order and q, c two non-zero complex constants. Then

$$\begin{split} T(r,f(qz+c)) &= T(r,f) + S(r,f), \\ N(r,\infty;f(qz+c)) &= N(r,f(z)) + S(r,f), \\ N(r,0;f(qz+c)) &= N(r,f(z)) + S(r,f), \\ \overline{N}(r,\infty;f(qz+c)) &= N(r,f(z)) + S(r,f), \\ \overline{N}(r,0;f(qz+c)) &= N(r,f(z)) + S(r,f) \end{split}$$

on a set of logarithmic density 1.

**Lemma 3.3.** [15] If  $N\left(r, \frac{1}{f^{(k)}} \mid f \neq 0\right)$  denotes the counting function of those zeros of  $f^{(k)}$  which are not the zeros of f, where a zero of  $f^{(k)}$  is counted according to its multiplicity then

$$N\left(r,0;f^{(k)}\mid f\neq 0\right)\leq k\overline{N}(r,\infty;f)+N\left(r,0;f\mid < k\right)+k\overline{N}\left(r,0f\mid \geq k\right)+S(r,f).$$

Lemma 3.4. [23] Let f be a non-constant meromorphic function and let

$$\mathcal{R}(f) = \frac{\sum_{i=0}^{n} a_i f^i}{\sum_{j=0}^{m} b_j f^j}$$

be an irreducible rational function in f with constant coefficients  $\{a_i\}$  and  $\{b_j\}$  where  $a_n \neq 0$  and  $b_m \neq 0$ . Then

$$T(r, \mathcal{R}(f)) = d T(r, f) + S(r, f),$$

where  $d = \max\{n, m\}$ .

**Lemma 3.5.** Let F and G be given by (3.1) satisfying  $E_F(1,m) = E_G(1,m)$ ,  $0 \le m < \infty$  with  $H \not\equiv 0$ , then

$$N_E^{(1)}(r,1;F) \le N(r,H) + S(r,F) + S(r,G).$$

Similar inequality holds for G also.

*Proof.* Let  $z_0$  be a simple 1-point of F. Then by a simple calculation, it can be shown that  $z_0$  is a zero of H. Since m(r,H) = S(r,F) + S(r,G), by First fundamental Theorem of Nevalinna, we obtain

$$N_E^{(1)}(r,1;F) \le N\left(r,\frac{1}{H}\right) \le T(r,H) \le N(r,H) + S(r,F) + S(r,G).$$

**Lemma 3.6.** Let f be a non-constant meromorphic function of zero order, then  $S(r, L_k(f, \Delta))$  can be replaced by S(r, f).

*Proof.* In view of Lemma 3.2, we get

$$T(r, L_k(f, \Delta)) \le \sum_{j=0}^k T(r, f(q_j z)) + T(r, f(z)) + S(r, f) \le (k+2)T(r, f) + S(r, f).$$

Hence, the lemma follows.

**Lemma 3.7.** Let F and G, being defined in (3.1) share (1,m),  $1 \le m < \infty$  and  $(\infty,0)$ . Then

$$\overline{N}(r,1;F,G) \leq \frac{1}{m+1} \left\{ \overline{N}(r,0;f(z)) + \overline{N}(r,0;L_k(f,\Delta)) \right\} + \frac{2}{m+1} \overline{N}(r,\infty;f(z)) + S(r,f).$$

*Proof.* In view of Lemma 3.3 and 3.6, we get

$$\begin{split} \overline{N}_*(r,1;F,G) &= \overline{N}_L(r,1;F) + \overline{N}_L(r,1;G) \\ &\leq \overline{N}(r,1;F) \geq m+2) + \overline{N}(r,1;G) \geq m+2) \\ &\leq \frac{1}{m+1} \Big\{ N(r,0;F' \mid F \neq 0) + N(r,0;G' \mid G \neq 0) \Big\} \\ &\leq \frac{1}{m+1} \Big\{ \overline{N}(r,0;F) + \overline{N}(r,0;G) + 2\overline{N}(r,\infty;F) \Big\} + S(r,f) \\ &\leq \frac{1}{m+1} \Big\{ \overline{N}(r,0;f(z)) + \overline{N}(r,0;L_k(f,\Delta)) \Big\} + \frac{2}{m+1} \overline{N}(r,\infty;f(z)) \\ &+ S(r,f). \end{split}$$

**Lemma 3.8.** Let F, G be given by (3.1) and  $V \not\equiv 0$ . If F, G share (1,m), and f,  $L_k(f,\Delta)$  share  $(\infty,k)$ , where  $0 \leq m,k < \infty$ , then the poles of F and G are zeros of V and

$$(nk+n-1)\overline{N}(r,\infty;f\mid\geq k+1)$$

$$\leq \overline{N}(r,0;f(z)) + \overline{N}(r,0;L_k(f,\Delta)) + \overline{N}_*(r,1;F,G) + S(r,f).$$

*Proof.* Since f(z),  $L_k(f, \Delta)$  share  $(\infty; k)$ , it follows that F, G share  $(\infty; nk)$  and so a pole of F with multiplicity  $p(\geq nk+1)$  is a pole of G with multiplicity  $r(\geq nk+1)$  and vice versa. We note that F and G have no pole of multiplicity q where nk < q < nk+n. Now using the Milloux theorem [[10], p. 55] and Lemma 3.6, we get from the definition of V, m(r, V) = S(r, f(z)). Therefore,

$$(nk + n - 1)\overline{N}(r, \infty; f \mid \geq k + 1) \leq N(r, 0; V) \leq T(r, V) + O(1)$$

$$\leq N(r, \infty; V) + m(r, V) + O(1)$$

$$\leq \overline{N}(r, 0; F) + \overline{N}(r, 0; G) + \overline{N}_*(r, 1; F, G) + S(r, f(z)) + S(r, L(z, f))$$

$$\leq \overline{N}(r, 0; f) + \overline{N}(r, 0; L_k(f, \Delta)) + \overline{N}_*(r, 1; F, G) + S(r, f).$$

**Lemma 3.9.** [27, 31] If F and G share  $(\infty, 0)$  and  $V \equiv 0$ , then  $F \equiv G$ .

**Lemma 3.10.** [28] Let  $H \equiv 0$  and F, G share  $(\infty, 0)$ , then F and G share  $(1, \infty)$ ,  $(\infty, \infty)$ .

**Lemma 3.11.** [3] Let F, G be two non-constant meromorphic functions sharing (1, m), where  $1 \le m < \infty$ . Then

$$\begin{split} & \overline{N}(r,1;F) + \overline{N}(r,1;G) - N_E^{1)}(r,1;F) + \left(t - \frac{1}{2}\right) \overline{N}_*(r,1;F,G) \\ & \leq & \frac{1}{2}[N(r,1;F) + N(r,1;G)]. \end{split}$$

**Lemma 3.12.** [13] Suppose F, G share (1,0),  $(\infty,0)$ . If  $H \not\equiv 0$ , then,

$$N(r,H) \leq \overline{N}(r,0;F \mid \geq 2) + \overline{N}(r,0;G \mid \geq 2) + \overline{N}_*(r,1;F,G) + \overline{N}_*(r,\infty;F,G) + \overline{N}_0(r,0;F') + \overline{N}_0(r,0;G') + S(r,F) + S(r,G).$$

where  $\overline{N}_0(r,0;F')$  is the reduced counting function of those zeros of F' which are not the zeros of F(F-1) and  $\overline{N}_0(r,0;G')$  is similarly defined.

**Lemma 3.13.** Let f be a transcendental meromorphic function function of zero order and  $L_k(f, E_q)$  be a linear q-shift polynomial defined in (1.1) and P(z) be defined as in the introduction. Then we have

$$(n-k-1)T(r,f) + S(r,f) < T(r,P(f)L_k(f,E_q)) < (n+k+1)T(r,f) + S(r,f)$$

If f is a transcendental entire function of zero order, then

$$T(r, P(f))L_k(f, E_q) = (n+1)T(r, f) + S(r, f).$$

*Proof.* We set a function  $F := P(f)L_k(f, E_q)$ , now if f is an entire function of finite order, then

$$\begin{split} T(r,F) &= m(r,F) \leq m(r,P(f)L_k(f,E_q)) + S(r,f) \\ &\leq & m(r,P(f)f) + m\left(r,\frac{L_k(f,E_q)}{f}\right) + S(r,f) \leq T(r,P(f)f) + S(r,f). \end{aligned} (3.4)$$

On the other hand, using Lemma 3.4, we have

$$(n+1)T(r,f) = T(r,P(f)f) + S(r,f) \le m(r,P(f)f) + S(r,f)$$

$$\le m(r,F) + m\left(r,\frac{P(f)f}{F}\right) + S(r,f)$$

$$\le m(r,F) + m\left(r,\frac{f}{L_k(f,E_q)}\right) + S(r,f) \le T(r,F) + S(r,f). \tag{3.5}$$

It follows from (3.4) and (3.4),

$$T(r,F) = (n+1)T(r,f) + S(r,f).$$

If f is a meromorphic function of zero order, then

$$T(r,F) = T(r,P(f)L_k(f,E_q)) \le T(r,P(f)) + T(r,L_k(f,E_q)) + S(r,f)$$
  
 
$$\le (n+k+1)T(r,f) + S(r,f).$$
(3.6)

Also we see that,

$$(n+1)T(r,f) + S(r,f) = T(r,P(f)f) + S(r,f)$$

$$\leq m(r,P(f)f) + N(r,P(f)f) + S(r,f)$$

$$\leq m\left(r,F\frac{f}{L_{k}(f,E_{q})}\right) + N\left(r,F\frac{f}{L_{k}(f,E_{q})}\right) + S(r,f)$$

$$\leq T(r,F) + (k+2)T(r,f) + S(r,f).$$

$$(3.7)$$

Equations (3.5) and (3.7) yield

$$(n-k-1)T(r,f) + S(r,f) \le T(r,F) \le (n+k+1)T(r,f) + S(r,f).$$

**Lemma 3.14.** Let f be a transcendental meromorphic function function of zero order and  $L_k(f, \Delta_q)$  be a linear q-difference polynomial defined in (1.2) and P(z) be defined as in the introduction. Then we have

$$T(r, P(f)L_k(f, E_q)) \ge (n - k - 1)T(r, f) + S(r, f).$$

If f is a transcendental entire function of zero order, we have

$$T(r, P(f)L_k(f, \Delta)) \ge nT(r, f) + S(r, f).$$

*Proof.* In a similar manner as done in Lemma 3.13, we can prove it. So, we omit the details.  $\Box$ 

## 4. Proofs of the theorems

Proof of Theorem 2.1. Case 1. Suppose f be a transcendental meromorphic of zero order function. Let  $P(f)L_k(f, E_q) - a(z)$  has only finitely many zeros.

By the Second fundamental Theorem of Nevalinna, and Lemmas 3.13, 3.2, we obtain

$$T(r, P(f)L_k(f, E_q)) \leq \overline{N}(r, \infty; P(f)L_k(f, E_q)) + \overline{N}(r, 0; P(f)L_k(f, E_q)) + \overline{N}(r, 0; P(f)L_k(f, E_q) - \alpha(z)) + S(r, f)$$

$$\leq (k+2)\overline{N}(r, \infty; f(z)) + (m+k+1)T(r, f(z)) + S(r, f).$$

i.e.,

$$(n-m-3k-4)T(r, f) < S(r, f),$$

which is a contradiction since n > m + 3k + 4.

Case 2. Let f(z) is a transcendental entire function. Then using the similar arguments as done in Case 1, we get

$$(n-m-k)T(r,f) \le S(r,f),$$

which contradicts n > m + k. Hence, the theorem is proved.

Proof of Theorem 2.2. Using Lemma 3.14 and 3.2, the proof of this theorem can be done in the line of proof of the Theorem 2.1. So we the details.  $\Box$ 

Proof of Theorem 2.3. Let  $G_*(z) = \frac{M(z)}{a}$ , then we know  $G_*(z)$  and  $G_*(qz)$  share 1 and  $\infty$  CM. Since the order of f is zero, it follows that

$$\frac{G_*(z)-1}{G_*(qz)-1}=h$$
, where h is a non-zero constant.

Rewriting the above equation, we obtain

$$G_*(z) + \frac{1}{h} - 1 = \frac{G_*(qz)}{h}.$$
 (4.1)

Suppose  $h \neq 1$ . Keeping in view of 3.4 and 4.1, the Second Fundamental theorem of Nevalinna yields

$$nT(r, f(z)) = T(r, G_*(z)) \leq \overline{N}(r, \infty; G_*) + \overline{N}(r, 0; G_*) + \overline{N}(r, 1 - \frac{1}{h}; G_*) + S(r, f)$$

$$\leq N(r, \infty; f(z)) + \overline{N}(r, 0; P(f)) + \overline{N}(r, 0; G_*(qz)) + S(r, f)$$

$$\leq (2m + 1)T(r, f) + S(r, f),$$

which contradicts n > 2m + 1. Hence h = 1, which implies that  $G_*(z) = G_*(qz)$ , i.e., P(f(z)) = tP(f(qz)) for a constant t satisfying  $t^n = 1$ .

Proof of Theorem 2.4. Let F and G be two functions defined in (3.1).

Since  $E_f(S_1, 1) = E_{L_k(f, \Delta)}(S_1, 1)$  and  $E_f(S_2, 0) = E_{L_k(f, \Delta)}(S_2, 0)$ , it follows that F, G share (1, 1) and  $(\infty, 0)$ .

We now consider the following two cases.

**Case 1:** Suppose  $H \not\equiv 0$ . Then  $F \not\equiv G$ . So, by Lemma 3.9, it follows that  $V \not\equiv 0$ . By the Second Fundamental Theorem of Nevalinna, we have

$$T(r,F) + T(r,G)$$

$$\leq \overline{N}(r,1;F) + \overline{N}(r,0;F) + \overline{N}(r,\infty;F) + \overline{N}(r,1;G) + \overline{N}(r,0;G) + \overline{N}(r,\infty;G)$$

$$-\overline{N}_0(r,0;F') - \overline{N}_0(r,0;G') + S(r,F) + S(r,G).$$

$$(4.2)$$

Applying Lemmas 3.5, 3.11 and 3.12 to the 4.2, we obtain,

$$\frac{n}{2} \Big( T(r,F) + T(r,G) \Big)$$

$$\leq N_2(r,0;F) + N_2(r,0;G) + 3\overline{N}(r,\infty;F) - \left( m - \frac{3}{2} \right) \overline{N}_*(r,1;F,G) + S(r,F) + S(r,G)$$

$$\leq 2 \{ \overline{N}(r,0;F) + \overline{N}(r,0;G) \} + 3\overline{N}(r,\infty;F) - \left( m - \frac{3}{2} \right) \overline{N}_*(r,1;F,G) + S(r,F) + S(r,G).$$

Using Lemma 3.7 with m = 1 and Lemma 3.6, the above inequality becomes

$$\frac{n}{2}\Big(T(r,F)+T(r,G)\Big)\leq \frac{9}{4}\Big(\overline{N}(r,0;F)+\overline{N}(r,0;G)\Big)+\frac{7}{2}\overline{N}(r,\infty;F)+S(r,f).$$

Again applying Lemmas 3.4, 3.6, 3.7 and 3.8 with m = 1, k = 0, the above equation turns into

$$\left(\frac{n}{2} - \frac{21}{4(n-2)} - \frac{9}{4}\right) \left(T(r,f) + T(r, L_k(f, \Delta))\right) \le S(r,f),$$

which contradicts  $n \geq 7$ .

Case 2: Suppose  $H \equiv 0$ . After integration we get,

$$F \equiv \frac{\alpha G + \beta}{\gamma G + \delta},\tag{4.3}$$

where  $\alpha, \beta, \gamma, \delta$  are complex constants satisfying  $\alpha \delta - \beta \gamma \neq 0$ .

As F, G share  $(\infty, 0)$ , by Lemma 3.10, it follows that f(z),  $L_k(f, \Delta)$  share  $(1, \infty)$ and  $(\infty, \infty)$ .

Subcase 2.1: Let 
$$\alpha \gamma \neq 0$$
. Then  $F - \frac{\alpha}{\gamma} = \frac{-(\alpha \delta - \beta \gamma)}{\gamma(\gamma G + \delta)} \neq 0$ .

Therefore, by the Second Fundamental Theorem of Nevallina, we get

$$nT(r,f) \le \overline{N}(r,0;F) + \overline{N}(r,\infty;F) + \overline{N}(r,\frac{\alpha}{\gamma};F) + S(r,F)$$
  
  $\le 2T(r,f) + S(r,f)$ 

which is a contradiction since  $n \geq 7$ .

**Subcase 2.2:** Suppose that  $\alpha \gamma = 0$ . Since  $\alpha \delta - \beta \gamma \neq 0$ , both  $\alpha$  and  $\gamma$  are not zero simultaneously.

**Subcase 2.2.1:** Suppose  $\alpha \neq 0$  and  $\gamma = 0$ . Then (4.3) becomes  $F \equiv AG + B$ , where  $A = \frac{\alpha}{\delta}$  and  $B = \frac{\beta}{\delta}$ . Subcase 2.2.1.1: Let F has no 1-point. Then by the Second Fundamental

Theorem, we get

$$T(r,F) \leq \overline{N}(r,0;F) + \overline{N}(r,1;F) + \overline{N}(r,\infty;F) + S(r,F)$$

or,

$$(n-2)T(r,f) < S(r,f),$$

which is a contradiction.

**Subcase 2.2.1.2:** Let F has some 1-point. Then A + B = 1.

**Subcase 2.2.1.2.1:** Suppose  $A \neq 1$ . Then  $F \equiv AG + 1 - A$ .

Therefore, by the Second Fundamental Theorem, we get

$$T(r,F) \leq \overline{N}(r,0;F) + \overline{N}(r,\infty;F) + \overline{N}(r,1-\alpha;F) + S(r,F)$$
  
$$\leq \overline{N}(r,0;F) + \overline{N}(r,\infty;F) + \overline{N}(r,0;G) + +S(r,F)$$
  
$$\leq 3T(r,f) + S(r,f).$$

i.e.,

$$(n-3)T(r,f) \le S(r,f),$$

which is again a contradiction since  $n \geq 7$ .

**Subcase 2.2.1.2.2:** Suppose A=1. Then  $F\equiv G$ . Thus we have  $L_k(f,\Delta)\equiv$ tf(z), where  $t^n = 1$  and  $t \neq -1$ 

**Subcase 2.2.2:** Suppose  $\alpha = 0$  and  $\gamma \neq 0$ .

Then (4.3) becomes

$$F \equiv \frac{1}{CG+D}, \ \ \text{where} \ \ C = \frac{\gamma}{\beta}, \ \ D = \frac{\delta}{\beta}.$$

**Subcase 2.2.2.1:** Let F has no 1-point. Then applying the second fundamental theorem to F, we have

$$nT(r,f) \le \overline{N}(r,\infty;F) + \overline{N}(r,0;F) + \overline{N}(r,1;F) + S(r,F)$$
  
  $\le 2T(r,f) + S(r,f),$ 

which is a contradiction.

**Subcase 2.2.2.2:** Suppose that F has some 1-point. Then C+D=1.

**Subcase 2.2.2.2.1:** Suppose C=1. Then D=0 and thus  $FG\equiv 1$ . *i.e.*,  $f(z)L_k(f,\Delta)\equiv ta^2$ , where  $t^n=1$ . Since F and G share  $(\infty,\infty)$ , so we have  $N\left(r,\frac{L_k(f,\Delta)}{f}\right)=N(r,0;f)$  and so in view of Lemma 3.1 and 3.6, we have

$$2T(r,f) + S(r,f) \le T\left(r, \frac{ta^2}{f^2}\right) \le T\left(r, \frac{L_k(f,\Delta)}{f}\right) + S(r,f)$$

$$\le N\left(r, \frac{L_k(f,\Delta)}{f}\right) + S(r,f) \le N(r,0;f) + S(r,f) \le T(r,f) + S(r,f),$$

which is a contradiction.

Proof of Theorem 2.5. The proof of this theorem can be done in a similar manner as done in Theorem 2.4, so we omit the details.  $\Box$ 

## Acknowledgment.

The authors did not receive support from any organization for the submitted work.

#### References

- T.C. Alzahary, H.X. Yi, Weighted value sharing and a question of I. Lahiri, Complex Var. Theory Appl. 49 (15) (2004) 1063-1078.
- [2] A. Banerjee, Meromorphic functions sharing two sets. Czech. Math. J. 57 (132) (2007) 1199– 1214.
- [3] A. Banerjee, Uniqueness of meromorphic functions sharing two sets with finite weight II. Tamkang J. Math. 41 (4) (2010) 379–392.
- [4] D.C. Barnett, R.G. Halburd, R.J. Korhonen, W. Morgan, Nevanlinna theory for the q-difference operator and meromorphic solutions of q-difference equations, Proc. Roy. Soc. Edinburgh Sect. A 137 (3) (2007) 457–474.
- [5] Z.X. Chen, Z.B. Huang, X.M. Zheng, On properties of difference polynomials, Acta Math. Sci. Ser. B Engl. Ed. 31 (2011) 627–633.
- [6] Y.M. Chiang, S.J. Feng, On the Nevanlinna characteristic of  $f(z+\eta)$  and difference equations in the complex plane, Ramanujan J. 16 (2008) 105–129.
- [7] R.G. Halburd, R.J. Korhonen, Difference analogue of the lemma on the logarithmic derivative with applications to difference equations, J. Math. Anal. Appl. 314 (2006) 477-487.
- [8] R.G. Halburd, R.J. Korhonen, Nevanlinna theory for the difference operator, Ann. Acad. Sci. Fenn. Math. 31 (2006) 463–478.
- [9] J. Heittokangas, R.J. Korhonen, I. Laine, J. Rieppo, Uniqueness of meromorphic functions sharing values with their shifts, Complex Var. Elliptic Equ. 56 (2011) 81–92.
- [10] W.K. Hayman, Meromorphic Functions, The Clarendon Press, Oxford (1964).
- [11] I. Lahiri, Weighted sharing and uniqueness of meromorphic functions, Nagoya Math. J. 161 (2001) 193–206.
- [12] I. Lahiri, Weighted value sharing and uniqueness of meromorphic functions, Complex Var. Theory Appl. 46 (2001) 241–253.
- [13] I. Lahiri, A. Banerjee, Weighted sharing of two sets. Kyungpook Math. J. 46 (2006) 79–87.

- [14] I. Lahiri, A. Sarkar, Uniqueness of meromorphic function and its derivative, J. Inequal. Pure Appl. Math. 5, 1, Art.20, 2004 [http://jipam.vu.edu.au/].
- [15] I. Lahiri, S. Dewan, Value distribution of the product of a meromorphic function and its derivative, Kodai Math. J. 26 (2003) 95–100.
- [16] I. Laine, C.C. Yang, Value distribution of difference polynomials, Proc. Japan Acad. Ser. A 83 (2007) 148–151.
- [17] K. Liu, Meromorphic functions sharing a set with applications to difference equations, J. Math. Anal. Appl. 359 (2009) 384–393.
- [18] K. Liu, L. Z. Yang, Value distribution of the difference operator, Arch. Math. 92 (2009) 270– 278.
- [19] K. Liu, X.L. Liu, T.B. Cao, Value distributions and uniqueness of difference polynomials, Adv. Differential Equations 2011(2011) Art. ID 234215, pp.12.
- [20] K. Liu, X.G. Qi, Meromorphic solutions of q-shift difference equations, Ann. Polon. Math. 101 (2011) 215–225.
- [21] K. Liu, X.G. Qi, Meromorphic solutions of q-shift difference equations, Ann. Pol. Math. 101 (2011) 215–225.
- [22] X.D. Luo, W.C. Lin, Value sharing results for shifts of meromorphic functions, J. Math. Anal. Appl. 377 (2011) 441–449.
- [23] A.Z. Mohon'ko, On the Nevanlinna characteristics of some meromorphic functions, Theory of Funct. Func. Anal. Appl. 14 (1971) 83–87.
- [24] X. Qi, K. Liu, L. Yang, Value sharing results of a meromorphic function f(z) and f(qz), Bull. Korean Math. Soc. 48 (2011) 1235–1243.
- [25] L.N. Wang, H.Y. Xu, T.S. Zhang, Properties of q-shift difference-differential polynomials of meromorphic functions, Adv. Difference Equ. 2014 (2014), Art. 249, doi:10.1186/1687-1847-2014-249.
- [26] H.Y. Xu, K. Liu, T.B. Cao, Uniqueness and value distribution for q-shifts of meromorphic functions, Math. Commun. 20 (2015) 97–112.
- [27] H.X. Yi, Meromorphic functions that share three sets, Kodai Math. J. 20 (1997) 22–32.
- [28] H.X. Yi, Meromorphic functions that share one or two values II, Kodai Math. J. 22 (1999) 264–272.
- [29] J. Zhang, Value distribution and shared sets of differences of meromorphic functions, J. Math. Anal. Appl. 367 (2) (2010) 401–408.
- [30] J.L. Zhang, R. Korhonen, On the Nevanlinna characteristic of f(qz) and its applications, J. Math. Anal. Appl. 369 (2010) (2) 537–544.
- [31] J.L. Zhang, Value distribution and shared sets of differences of meromorphic functions, J. Math. Anal. Appl. 367 (2010) 401–408.
- [32] J.L. Zhang, R.J. Korhonen, On the Nevanlinna characteristic of f(qz) and its applications, J. Math. Anal. Appl. 369 (2010) 537–544.