

# Measure equivalence rigidity of $\text{Out}(F_N)$

Vincent Guirardel and Camille Horbez

March 30, 2021

## Abstract

We prove that for every  $N \geq 3$ , the group  $\text{Out}(F_N)$  of outer automorphisms of a free group of rank  $N$  is superrigid from the point of view of measure equivalence: any countable group that is measure equivalent to  $\text{Out}(F_N)$ , is in fact virtually isomorphic to  $\text{Out}(F_N)$ .

We introduce three new constructions of canonical splittings associated to a subgroup of  $\text{Out}(F_N)$  of independent interest. They encode respectively the collection of invariant free splittings, invariant cyclic splittings, and maximal invariant free factor systems. Our proof also relies on the following improvement of an amenability result by Bestvina and the authors: given a free factor system  $\mathcal{F}$  of  $F_N$ , the action of  $\text{Out}(F_N, \mathcal{F})$  (the subgroup of  $\text{Out}(F_N)$  that preserves  $\mathcal{F}$ ) on the space of relatively arational trees with amenable stabilizer is a Borel amenable action.

## 1 Introduction

Measure equivalence theory is concerned with classifying countable groups up to the following equivalence relation.

**Definition 1.1** (Gromov [Gro93, Definition 0.5.E1]). *Two countable groups  $\Gamma_1$  and  $\Gamma_2$  are measure equivalent if there exists a measure-preserving action of  $\Gamma_1 \times \Gamma_2$  by Borel automorphisms on a standard measure space  $\Sigma$  (i.e. a Polish space equipped with its Borel  $\sigma$ -algebra and endowed with a nonzero  $\sigma$ -finite Borel measure) such that for every  $i \in \{1, 2\}$ , the  $\Gamma_i$ -action on  $\Sigma$  is free and has a finite measure fundamental domain, i.e. there exists a Borel subset  $X_i \subseteq \Sigma$  of finite measure such that*

$$\Sigma = \coprod_{\gamma \in \Gamma_i} \gamma X_i.$$

A typical example is that two lattices in the same locally compact second countable group are always measure equivalent (see [Gro93, Example 0.5.E2]). This notion can be viewed as a measure-theoretic analogue to the notion of quasi-isometry between finitely generated groups. Indeed, a theorem of Gromov [Gro93, 0.2.C'2] asserts that two finitely generated groups  $\Gamma_1$  and  $\Gamma_2$  are quasi-isometric if and only if they are *topologically equivalent*, i.e. there exists a continuous action of  $\Gamma_1 \times \Gamma_2$  on some locally compact space

$X$ , such that for every  $i \in \{1, 2\}$ , the  $\Gamma_i$ -action on  $X$  is properly discontinuous and cocompact. We insist however that this remains purely at the level of analogy: there is no implication in either direction between measure equivalence and quasi-isometry.

An alternative point of view on measure equivalence comes from studying measure-preserving group actions on standard probability spaces, where one forgets everything but the partition into orbits. Indeed, by works of Furman [Fur99b] and Gaboriau [Gab02b, Theorem 2.3], two countable groups are measure equivalent if and only if they admit stably orbit equivalent free actions on standard probability spaces by measure-preserving Borel automorphisms – see Definition 1.3 below.

An easy way in which two countable groups  $\Gamma_1$  and  $\Gamma_2$  can be measure equivalent is if they are *virtually isomorphic*. This means that there exist finite-index subgroups  $\Lambda_i \subseteq \Gamma_i$  and finite normal subgroups  $F_i \trianglelefteq \Lambda_i$  for all  $i \in \{1, 2\}$ , such that  $\Lambda_1/F_1$  is isomorphic to  $\Lambda_2/F_2$ . A countable group  $\Gamma$  is said to be *ME-superrigid* if every countable group which is measure equivalent to  $\Gamma$  is in fact virtually isomorphic to  $\Gamma$ .

Given  $N \in \mathbb{N}$ , we let  $F_N$  be a free group of rank  $N$ , and  $\text{Out}(F_N)$  be its outer automorphism group. The group  $\text{Out}(F_N)$  has been subject to intense study from the point of view of geometric group theory over the past decades, often in analogy to mapping class groups of surfaces. The main theorem of the present paper is the following.

**Theorem 1.2.** *For every  $N \geq 3$ , the group  $\text{Out}(F_N)$  is ME-superrigid: every countable group which is measure equivalent to  $\text{Out}(F_N)$ , is in fact virtually isomorphic to  $\text{Out}(F_N)$ .*

Notice that this does not hold when  $N = 2$ , as  $\text{Out}(F_2)$  is virtually free, and the class of groups that are measure equivalent to a free group is wide, see e.g. [Gab05, BTW07]. To our knowledge, the topological counterpart to Theorem 1.2, i.e. knowing whether  $\text{Out}(F_N)$  is quasi-isometrically rigid, is still wide open. Actually, the geometry of  $\text{Out}(F_N)$  equipped with the word metric is still very mysterious, in large part due to the lack of a satisfying analogue for  $\text{Out}(F_N)$  of the Masur–Minsky theory [MM00] yielding a distance formula for mapping class groups of surfaces.

## 1.1 Some history

### 1.1.1 Measure equivalence rigidity and flexibility

We would like to mention here some previously known results that are milestones in the (still growing) theory of measure equivalence rigidity. We refer the reader to [Sha05, Gab10, Fur11b] for general surveys on measured group theory.

A first striking flexibility result in the classification of groups up to measure equivalence (in an orthogonal direction to that of rigidity results) was the proof by Ornstein and Weiss [OW80], following previous work of Dye [Dye59, Dye63] (and further extended by Connes, Feldman and Weiss [CFW81]), that all countably infinite amenable groups are measure equivalent. More precisely, any two free ergodic measure-preserving actions of countably infinite amenable groups on standard probability spaces are orbit equivalent. In contrast, work by Hjorth, Gaboriau, Lyons, Popa, Ioana and Epstein shows that any

non-amenable countable group has uncountably many ergodic actions that are pairwise not orbit equivalent [Hjo05, GP05, GL09, Ioa11, Eps07].

Still on the flexibility side, Gaboriau showed that many groups are measure equivalent to free groups [Gab05] (see also [BTW07], showing more generally that every finitely generated group with the same elementary theory as a free group is measure equivalent to a finitely generated free group).

A groundbreaking rigidity result was the proof by Furman [Fur99a], building on earlier work of Zimmer and most notably on Zimmer’s celebrated cocycle superrigidity theorem [Zim80, Zim91], that any countable group which is measure equivalent to a lattice in a higher rank simple Lie group  $G$ , is in fact virtually isomorphic to a lattice in  $G$ .

Another remarkable result by Gaboriau shows that  $\ell^2$ -Betti numbers are (projective) invariants of measure equivalence [Gab02a]. By [GN19], the  $\ell^2$ -Betti numbers of  $\text{Out}(F_N)$  do not vanish in degree equal to the virtual cohomological dimension  $2N - 3$ , which gives a short proof that  $\text{Out}(F_N)$  is not measure equivalent to  $\text{Out}(F_{N'})$  if  $N' \neq N$ . Monod and Shalom [MS06] used bounded cohomology techniques to establish striking measure equivalence rigidity results for products of negatively curved groups.

Later, Kida established that all mapping class groups of non-exceptional finite-type orientable surfaces are ME-superrigid [Kid10]. Kida’s approach to measure equivalence rigidity of mapping class groups has had a large influence on our work: the general strategy of our proof is similar to his, though many new difficulties arose for  $\text{Out}(F_N)$ .

Kida’s theorem has then inspired further ME-superrigidity results [Kid11, CK15, HH20a]. Also, there are situations of groups that fail to be ME-superrigid, but for which one can still get interesting measure equivalence classification results within a given class, like Baumslag–Solitar groups [HR15] or right-angled Artin groups [HH20b].

In another direction, superrigidity results have been obtained for certain particular actions with few restrictions on the acting group: this started with Popa’s remarkable cocycle superrigidity theorem for Bernoulli actions of Property (T) groups [Pop07], leading to many further developments.

We finally mention that measure equivalence theory has strong connections with the theory of von Neumann algebras, see e.g. [Sin55]. In combination to the aforementioned work of Kida, the notion of *proper proximality* of a countable group, recently introduced by Boutonnet, Ioana and Peterson [BIP18], led to strong rigidity results for von Neumann algebras associated to weakly compact (e.g. profinite) free probability measure-preserving ergodic actions of mapping class groups [HHL20]. A natural question that arises from our work is to which extent an ergodic action of  $\text{Out}(F_N)$  is determined by its associated von Neumann algebra.

### 1.1.2 Rigidity in $\text{Out}(F_N)$

Rigidity of  $\text{Out}(F_N)$  also has a long history, starting with works of Khramtsov [Khr90] and of Bridson and Vogtmann [BV00] establishing that all automorphisms of  $\text{Out}(F_N)$  with  $N \geq 3$  are inner. This algebraic rigidity statement is in close relation to more geometric rigidity statements concerning the symmetries of various spaces equipped with

natural  $\text{Out}(F_N)$ -actions, starting with the spine of reduced Outer space [BV01]. Later, Farb and Handel proved [FH07] that  $\text{Out}(F_N)$  is its own abstract commensurator for all  $N \geq 4$ , i.e. every isomorphism between finite-index subgroups of  $\text{Out}(F_N)$  is given by a conjugation. A recent new proof of this result by Wade and the second named author [HW20] extends it to the  $N = 3$  case and allows to compute the abstract commensurator of various interesting subgroups of  $\text{Out}(F_N)$  such as its Torelli subgroup. Our work on measure equivalence rigidity actually recovers the Farb–Handel theorem (see Remark 1.18); the general approach of the present paper can also be viewed in a sense as a broad generalization of the strategy used in [HW20].

## 1.2 Some consequences

We now mention two applications of the techniques we develop in the present paper.

### 1.2.1 Orbit equivalence rigidity of ergodic actions of $\text{Out}(F_N)$

Using standard techniques from measured group theory originating from work of Furman [Fur99b], Theorem 1.2 can be used to deduce a strong rigidity result regarding the orbit structure of ergodic actions of  $\text{Out}(F_N)$  on standard probability spaces. Before we state it, we recall the following definition.

**Definition 1.3.** *Let  $\Gamma_1$  and  $\Gamma_2$  be two countable groups, and for every  $i \in \{1, 2\}$ , let  $\Gamma_i \curvearrowright X_i$  be an ergodic measure-preserving action by Borel automorphisms on a standard probability space  $X_i$ . The two actions  $\Gamma_1 \curvearrowright X_1$  and  $\Gamma_2 \curvearrowright X_2$  are orbit equivalent if there exists a measure space isomorphism  $f : X_1 \rightarrow X_2$  such that for a.e.  $x \in X_1$ , one has  $f(\Gamma_1 \cdot x) = \Gamma_2 \cdot f(x)$ .*

*More generally, the actions  $\Gamma_1 \curvearrowright X_1$  and  $\Gamma_2 \curvearrowright X_2$  are stably orbit equivalent if there exist Borel subsets  $Y_i \subseteq X_i$  of positive measure and a measure-scaling<sup>1</sup> isomorphism  $f : Y_1 \rightarrow Y_2$  such that for a.e.  $y \in Y_1$ , one has  $f((\Gamma_1 \cdot y) \cap Y_1) = (\Gamma_2 \cdot f(y)) \cap Y_2$ .*

An easy example of orbit equivalent actions is when the actions  $\Gamma_1 \curvearrowright X_1$  and  $\Gamma_2 \curvearrowright X_2$  are *conjugate*. This means that there exists a group isomorphism  $\rho : \Gamma_1 \rightarrow \Gamma_2$  and a measure space isomorphism  $f : X_1 \rightarrow X_2$  such that for every  $\gamma \in \Gamma_1$  and a.e.  $x \in X_1$ , one has  $f(\gamma \cdot x_1) = \rho(\gamma) \cdot f(x_1)$ .

**Theorem 1.4** (OE-superrigidity, see Theorem 4.6). *Let  $N \geq 3$ . Let  $\Gamma \subseteq \text{Out}(F_N)$  be a finite index subgroup, and  $\Lambda$  a countable group with no non-trivial finite normal subgroup. Consider two standard probability spaces  $X_1, X_2$  and two free measure-preserving actions  $\Gamma \curvearrowright X_1, \Lambda \curvearrowright X_2$ . Assume that these actions are aperiodic, i.e. that every finite index subgroup acts ergodically.*

*If the actions  $\Gamma \curvearrowright X_1$  and  $\Lambda \curvearrowright X_2$  are stably orbit equivalent, then they are conjugate.*

---

<sup>1</sup>i.e.  $f$  induces a measure space isomorphism after rescaling the measures to turn  $Y_1$  and  $Y_2$  into probability spaces

### 1.2.2 Lattice embeddings of $\text{Out}(F_N)$ into locally compact groups

As mentioned earlier, a typical example of measure equivalent groups comes from two lattices in a common locally compact second countable group. A consequence of our work is that there are no interesting lattice embeddings of  $\text{Out}(F_N)$  into any locally compact second countable group.

**Theorem 1.5.** *Let  $N \geq 3$ , and let  $\Gamma$  be a finite-index subgroup of  $\text{Out}(F_N)$ . Let  $H$  be a locally compact second countable group (equipped with its left Haar measure), and let  $\sigma : \Gamma \rightarrow H$  be an injective homomorphism, such that  $\sigma(\Gamma)$  is a lattice in  $H$  (i.e.  $\sigma(\Gamma)$  is discrete in  $H$  and has finite Haar covolume).*

*Then there exists a continuous homomorphism  $\Phi_0 : H \rightarrow \text{Out}(F_N)$  with compact kernel such that for all  $\gamma \in \Gamma$ , one has  $\Phi_0(\sigma(\gamma)) = \gamma$ .*

This implies in particular that  $H$  has infinitely many connected components and  $\sigma(\Gamma)$  is cocompact in  $H$ . By viewing  $\text{Out}(F_N)$  as a lattice in the automorphism group of its Cayley graph, we reach the following corollary, which was suggested to us by Jingyin Huang.

**Corollary 1.6.** *Let  $N \geq 3$ . For any finite generating set  $S$  of  $\text{Out}(F_N)$ , any automorphism of the Cayley graph  $\text{Cay}(\text{Out}(F_N), S)$  is at bounded distance from an automorphism induced by the left multiplication by an element of  $\text{Out}(F_N)$ .*

*If  $\Gamma$  is a torsion-free finite-index subgroup of  $\text{Out}(F_N)$ , and  $S$  is a finite generating set of  $\Gamma$ , then the group of automorphisms of the Cayley graph  $\text{Cay}(\Gamma, S)$  is isomorphic to a subgroup of  $\text{Out}(F_N)$  containing  $\Gamma$ . In particular, it is countable.*

In this statement,  $\text{Cay}(G, S)$  is the simple graph with vertices  $G$ , and such that there is an edge between two distinct elements  $g$  and  $h$  if and only if  $gh^{-1} \in S \cup S^{-1}$ . As was recently proved by Leemann and de la Salle, every finitely generated group has a Cayley graph (for some finite generating set) whose automorphism group is countable [LdlS20, Corollary 1.3]. But knowing that this holds for every Cayley graph is a much more restrictive condition, which fails for free groups, for instance. On the other hand, if a finitely generated infinite group  $G$  has a non-trivial finite order element, then it has a Cayley graph whose automorphism group is uncountable [dlST19, Lemma 6.1].

### 1.3 New constructions of canonical splittings for subgroups of $\text{Out}(F_N)$

A substantial part of the present paper (Part II) is devoted to three constructions of canonical splittings associated to a subgroup  $H$  of  $\text{Out}(F_N)$  which we believe to be of independent interest. This is reminiscent of Ivanov's *canonical reduction system* of a reducible subgroup of the mapping class group, which is a canonical decomposition of the surface along a multicurve. As a consequence of these constructions, we obtain the following result which has the same flavor.

**Theorem 1.7** (see Theorem 8.32). *Let  $N \geq 2$ , and let  $H \subseteq \text{Out}(F_N)$  be an infinite subgroup. If  $H$  has a proper free factor whose conjugacy class is periodic, so does its normalizer in  $\text{Out}(F_N)$ .*

We will not use this result directly, we will instead rely on a similar statement for a particular class of splittings (see Corollary 1.16). One has to be careful with the class of splittings considered. Indeed, there exists a subgroup  $H \subseteq \text{Out}(F_N)$  having a non-trivial periodic free splitting, but whose normalizer has none (Example 6.1 gives an instance). We actually carry out three separate constructions of canonical splittings. The first two encode the sets of  $H$ -invariant free splittings and  $H$ -invariant  $\mathcal{Z}_{\max}$ -splittings, respectively, relying on JSJ techniques. The third has a different flavor and deals with  $H$ -invariant free factor systems.

Recall that a *splitting* of  $F_N$  is an action of  $F_N$  on a simplicial tree with no proper nonempty invariant subtree. It is a *free splitting* if edge stabilizers are trivial. Thus, one-edge free splittings (i.e. those for which the quotient graph under the  $F_N$ -action has a single edge) are dual to free product decompositions  $F_N = A * B$  or HNN extensions over the trivial group  $F_N = A *$ . A  $\mathcal{Z}_{\max}$ -*splitting* is a splitting of  $F_N$  whose edge stabilizers are isomorphic to  $\mathbb{Z}$  and maximal for the inclusion.

We will actually work in the finite index subgroup  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z}) \subseteq \text{Out}(F_N)$  consisting of outer automorphisms acting trivially on homology mod 3. The point of this subgroup is that it avoids many finite order phenomena: this group is torsion-free, any free splitting which is periodic by some element or subgroup of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is in fact invariant; and so is any periodic conjugacy class of free factor, or any periodic conjugacy class of element ([HM20], see Section 2.10).

### 1.3.1 Canonical splittings from collections of free splittings

The following theorem is at the heart of Section 6 of the present paper (see Theorem 6.12 for a more complete statement). It associates to any subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  a canonical splitting from which one can read all  $H$ -invariant free splittings.

**Theorem 1.8** (see Theorem 6.12). *Let  $N \geq 2$ . To any subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , one can canonically assign a splitting  $U_H^1$  of  $F_N$  which is non-trivial as soon as  $H$  is infinite and preserves some non-trivial free splitting of  $F_N$ . The assignment  $H \mapsto U_H^1$  is  $\text{Out}(F_N)$ -equivariant.*

*The splitting  $U_H^1$  comes with a bipartition  $V(U_H^1) = V^0 \sqcup V^1$  of its vertex set, and the set of  $H$ -invariant one-edge free splittings coincides with the set of all one-edge free splittings that can be obtained by blowing up a vertex of  $V^1$ .*

Our proof of Theorem 1.8 uses techniques coming from JSJ theory to construct  $U_H^1$  from the collection of all  $H$ -invariant free splittings. It might be surprising that the result of the construction is a canonical splitting as opposed to a deformation space, as is usually the case when working with free splittings. What rigidifies the situation here is the existence of the infinite group  $H$  that preserves all these free splittings, and  $U_H^1$  is constructed using the action of  $H$ . More precisely, denoting by  $\tilde{H}$  the preimage of  $H$  in  $\text{Aut}(F_N)$ , we view  $H$ -invariant free splittings of  $F_N$  as splittings of  $\tilde{H}$ , and construct  $U_H^1$  as a tree of cylinders for a deformation space of  $\tilde{H}$ -trees made of maximal  $H$ -invariant free splittings.

The above theorem yields in particular a description of the stabilizer of a collection of free splittings of  $F_N$  (see Proposition 6.19), and leads to the following chain condition for stabilizers of collections of free splittings.

**Theorem 1.9** (see Proposition 6.17). *There is a bound, only depending on the rank  $N$  of the free group, on the length of a chain  $\mathcal{C}_1 \subseteq \cdots \subseteq \mathcal{C}_k$  of collections of free splittings of  $F_N$  such that for all  $i \in \{1, \dots, k-1\}$ , we have*

$$\bigcap_{S \in \mathcal{C}_{i+1}} \text{Stab}(S) \subsetneq \bigcap_{S \in \mathcal{C}_i} \text{Stab}(S),$$

where  $\text{Stab}(S)$  denotes the  $\text{Out}(F_N)$ -stabilizer of  $S$ .

### 1.3.2 Canonical splittings from collections of $\mathcal{Z}_{\max}$ -splittings

In the case where  $H$  does not preserve any non-trivial free splitting, the splitting  $U_H^1$  constructed above is just the trivial splitting. But in this case, one may use the collection of invariant  $\mathcal{Z}_{\max}$ -splittings, if any, to construct a more interesting canonical splitting.

**Theorem 1.10** (see Theorem 7.4). *Let  $N \geq 2$ . To any subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  that preserves a non-trivial  $\mathcal{Z}_{\max}$ -splitting but does not preserve any non-trivial free splitting, one can canonically assign a non-trivial  $\mathcal{Z}_{\max}$ -splitting  $U_H^z$ . The assignment  $H \mapsto U_H^z$  is  $\text{Out}(F_N)$ -equivariant.*

The splitting  $U_H^z$  is obtained as a JSJ splitting of all  $H$ -invariant  $\mathcal{Z}_{\max}$ -splittings of  $F_N$ : it is maximal for domination among all universally elliptic  $H$ -invariant  $\mathcal{Z}_{\max}$ -splittings, and equal to its own tree of cylinders. Although surface groups are examples of hyperbolic groups having non-trivial  $\mathcal{Z}_{\max}$ -splittings but trivial JSJ decomposition, this possibility of a trivial JSJ decomposition is ruled out here because the infinite group  $H$  has to act trivially on surfaces of the JSJ decomposition.

A collection of  $\mathcal{Z}_{\max}$ -splittings is FS-averse if its elementwise stabilizer in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  does not fix any non-trivial free splitting of  $F_N$ . As above, we obtain a chain condition as a consequence of Theorem 1.10.

**Theorem 1.11** (see Proposition 7.25). *There is a bound, only depending on the rank  $N$  of the free group, on the length of a chain  $\mathcal{C}_0 \subseteq \cdots \subseteq \mathcal{C}_k$  of FS-averse collections of  $\mathcal{Z}_{\max}$ -splittings of  $F_N$  such that for all  $i \in \{0, \dots, k-1\}$ , one has*

$$\bigcap_{S \in \mathcal{C}_{i+1}} \text{Stab}(S) \subsetneq \bigcap_{S \in \mathcal{C}_i} \text{Stab}(S).$$

### 1.3.3 The dynamical decomposition

Our third construction of a canonical splitting, carried out in Section 8, is of a different, more dynamical, flavor. It encodes the interaction between the maximal invariant free factor systems of a subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  that does not fix any non-trivial free



splitting. It is remarkable that from free factor systems, one obtains a canonical splitting, which is a much more rigid object than a free factor system.

To every subgroup  $H$  of the mapping class group of a closed, connected, orientable, hyperbolic surface  $\Sigma$ , Ivanov associated a canonical subsurface decomposition which singles out *active* parts of  $\Sigma$  (where  $H$  acts with at least one pseudo-Anosov homeomorphisms) and *inactive* parts (where, up to a finite-index subgroup,  $H$  acts trivially). By analogy, our *dynamical decomposition* for subgroups  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  extracts some *active parts* of the free group – although we cannot expect  $H$  to act trivially on the complement.

A set  $\mathcal{F}$  of conjugacy classes of non-trivial proper subgroups of  $F_N$  is a *free factor system* if there exist representatives  $P_1, \dots, P_k$  of the conjugacy classes such that  $\mathcal{F} = \{[P_1], \dots, [P_k]\}$  and  $F_N$  decomposes as a free product  $F_N = P_1 * \dots * P_k * F_r$  for some free subgroup  $F_r$  of  $F_N$  (with  $0 \leq r \leq N$ ). We order free factor systems by saying that  $\mathcal{F} \preceq \mathcal{F}'$  if every group in  $\mathcal{F}$  is conjugate in a group of  $\mathcal{F}'$ . Since chains of free factor systems have bounded length, any subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  has at least one maximal  $H$ -invariant free factor system  $\mathcal{F}$  (we allow  $\mathcal{F} = \emptyset$ ).

We now assume that  $H$  does not preserve any non-trivial free splitting. By [HM14, GH19b], for any maximal  $H$ -invariant free factor system  $\mathcal{F}$ , the group  $H$  contains a fully irreducible automorphism relative to  $\mathcal{F}$ . However, there may exist  $H$ -invariant conjugacy classes of free factors which are not  $\mathcal{F}$ -peripheral<sup>2</sup> (i.e. not conjugate into any of the subgroups  $P_1, \dots, P_k$ ); these potential extra invariant free factors are the basis of our construction.




Figure 1: The dynamical decomposition of the group  $H \subseteq \text{Out}(F_N)$  induced by homeomorphisms of the three surfaces in Example 1.12. The conjugacy class of the free factor  $Q_1$  is  $H$ -invariant, and  $\hat{\mathcal{F}}_1 = \{[Q_1], [c_1]\}$  is a maximal  $H$ -invariant almost free factor system.

<sup>2</sup>For experts, the maximality of  $\mathcal{F}$  only says that there is no proper free factor *relative to*  $\mathcal{F}$  whose conjugacy class is  $H$ -invariant.



*Example 1.12* (see Example 8.19). The following example might be helpful (see Figure 1). We identify  $F_N$  with the fundamental group of the space obtained by gluing 3 surfaces on a circle by homeomorphism along one of their boundary components, as shown in Figure 1. We take for  $H$  the group of automorphisms induced by homeomorphisms of the surfaces that are the identity on the boundary. Let  $Q_i$  be the fundamental group of the complement of  $\Sigma_i$ . Then  $\mathcal{F}_i = \{[Q_i]\}$  is a maximal  $H$ -invariant free factor system, and  $Q_2$  is a free factor whose conjugacy class is  $H$ -invariant and which is not  $\mathcal{F}_1$ -peripheral.

For technical reasons, we need to work with  $H$ -invariant *almost free factor systems*, usually denoted by  $\hat{\mathcal{F}}$ , which include extra situations involving unused boundary components of surfaces, see Definition 8.6. For example, in Example 1.12,  $\hat{\mathcal{F}}_i = \{[Q_i], [c_i]\}$  is an almost free factor system (in fact a maximal  $H$ -invariant almost free factor system). In the case where  $H$  has no invariant conjugacy class, every  $H$ -invariant almost free factor system is a free factor system, so one may ignore these subtleties at first.

If there is a unique maximal  $H$ -invariant almost free factor system, we say that  $H$  is *pure*. In this case, the dynamical decomposition of  $H$  will be trivial. In Example 1.12,  $H$  is not pure, but its restriction to each surface group is pure.

When there are several maximal  $H$ -invariant almost free factor systems, the dynamical decomposition produces a canonical splitting having some vertex groups which are free factors in restriction to which  $H$  is pure. These vertices are called *active*. In the case of Example 1.12, the dynamical decomposition is dual to the decomposition shown in Figure 1, and the active vertices correspond to the three surfaces.

**Theorem 1.13** (Dynamical decomposition, see Theorem 8.15). *Let  $N \geq 2$ , and let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup which does not preserve any non-trivial free splitting. Then there exists a canonical  $H$ -invariant splitting  $U_H^d$  coming with a bipartition of its vertex set  $U_H^d = V_p \sqcup V_a$  such that*

1. *for every  $v \in V_p$ , the stabilizer  $G_v$  is peripheral in every maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}$ , i.e.  $G_v$  is conjugate into one of the factors in  $\hat{\mathcal{F}}$ ;*
2. *for every  $v \in V_a$ , the stabilizer  $G_v$  is a free factor of  $F_N$  whose conjugacy class is  $H$ -invariant, and the restriction of  $H$  to  $G_v$  is pure.*

Vertices in  $V_a$  are called *active*. More details about this decomposition are given in Section 8.

Our proof involves considering, for each maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}$ , an element  $\alpha \in H$  which is irreducible relative to  $\hat{\mathcal{F}}$ , and an  $\alpha$ -invariant  $\mathbb{R}$ -tree  $T$ . If  $A$  is any free factor whose conjugacy class is  $H$ -invariant, and which is not  $\hat{\mathcal{F}}$ -peripheral, then one gets a splitting of  $F_N$  from a transverse covering of  $T$  by the translates of the  $A$ -minimal subtree. One then proves that there is a unique minimal such  $H$ -invariant conjugacy class of free factor  $A_{\hat{\mathcal{F}}}$  which is not  $\hat{\mathcal{F}}$ -peripheral and one can check that the associated splitting  $T_{\hat{\mathcal{F}}}$  depends only on  $\hat{\mathcal{F}}$  and  $H$ , not on the choice of  $\alpha$  and  $T$ . Then as  $\hat{\mathcal{F}}$  varies among maximal  $H$ -invariant almost free factor systems, the splittings  $T_{\hat{\mathcal{F}}}$  are compatible and yield the desired canonical dynamical decomposition.

In the end, the groups  $A_{\mathcal{F}}$  appear as the stabilizers of the active vertices of  $U_H^d$ . As a consequence, we obtain a bound on the number of maximal  $H$ -invariant free factor systems.

**Corollary 1.14** (see Corollary 8.30). *There exists  $K > 0$ , only depending on the rank of the free group, such that for every subgroup  $H \subseteq \text{Out}(F_N)$ , assuming that no free splitting is invariant under a finite-index subgroup of  $H$ , then there are at most  $K$  maximal free factor systems of  $F_N$  that are invariant under a finite-index subgroup of  $H$ .*

The assumption that  $H$  does not preserve any free splitting is crucial: one easily builds examples (including examples coming from punctured surfaces) of subgroups  $H \subseteq \text{Out}(F_N)$  with infinitely many  $H$ -invariant free splittings, giving infinitely many invariant conjugacy classes of maximal free factor systems (which are *sporadic*, i.e. of the form  $\mathcal{F} = \{[P_1]\}$  with  $F_N = P_1 * \mathbb{Z}$  or  $\mathcal{F} = \{[P_1], [P_2]\}$  with  $F_N = P_1 * P_2$ ).

### 1.3.4 Nice splittings

Theorems 1.8 and 1.10 construct from an  $H$ -invariant splitting, a canonical one, which is in particular invariant under the normalizer of  $H$ . The output is no longer a cyclic splitting in general, and neither is the dynamical decomposition. To iterate this kind of argument in a unified fashion, we use the following definition.

**Definition 1.15.** *A splitting  $S$  of  $F_N$  is nice if either*

1.  *$S$  is a non-trivial free splitting, or*
2.  *$S$  is a non-trivial  $Z_{\max}$ -splitting, or*
3.  *$S$  is bi-nonsporadic, i.e. all edge stabilizers of  $S$  are finitely generated and non-abelian, and  $S$  contains two vertices in distinct  $F_N$ -orbits such that the Grushko decomposition of their stabilizer with respect to the collection of all incident edge stabilizers is non-sporadic.*

With this definition at hand one can deduce the following result from the three constructions above (strictly speaking, the canonical splittings described above are not always nice, but they are easily turned into nice splittings in a canonical way, see Corollary 9.5).

**Corollary 1.16** (see Corollary 9.3). *Let  $N \geq 2$ . If  $H \subseteq \text{Out}(F_N)$  is an infinite subgroup preserving a nice splitting, then its normalizer preserves a nice splitting.*

As a matter of analogy, when  $\Sigma$  is a closed, connected, oriented, hyperbolic surface, and  $H$  is an infinite subgroup of  $\text{Mod}(\Sigma)$  that preserves the isotopy class of a multicurve, then  $H$  has a canonical reduction system, i.e. a multicurve whose isotopy class is also invariant by its normalizer. Thus, Corollary 1.16 and the three canonical constructions on which it is based, can be viewed as providing a (nice!) analogue of canonical reduction systems for free groups.

## 1.4 On the proof of the measure equivalence rigidity theorem

We now present the techniques we use in the proof of our main theorem.

### 1.4.1 Classical measure group theoretic reductions

We first use a standard argument in measured group theory, developed in the successive works of Furman [Fur99a, Fur99b], Monod and Shalom [MS06] and Kida [Kid10], to reduce the proof of our main theorem to a statement about measured groupoids associated to measure-preserving  $\text{Out}(F_N)$ -actions on probability spaces.

It is actually enough (see e.g. [Kid10, Theorem 6.1]) to prove a rigidity statement for self measure equivalence couplings of a finite index subgroup of  $\text{Out}(F_N)$ . As above, we will work in the finite index subgroup  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  made of automorphisms acting trivially on homology mod 3. A *self coupling* of a countable group  $\Gamma$  is a measure-preserving action  $\Gamma \times \Gamma \curvearrowright \Sigma$  by Borel automorphisms on a standard measure space  $\Sigma$  such that the action of each factor is free and has a finite measure fundamental domain. The restriction to self couplings should not surprise the reader familiar with quasi-isometry rigidity results: self couplings may be thought as analogues of self quasi-isometries of  $\Gamma$ , and quasi-isometric rigidity of groups is often deduced from rigidity properties of these self quasi-isometries.

There is a *standard* self coupling  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z}) \times \text{IA}_N(\mathbb{Z}/3\mathbb{Z}) \curvearrowright \Sigma_0 = \text{Out}(F_N)$ , where the action is by left and right multiplication. Our goal is to show that every self coupling  $\Sigma$  of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  factors through this standard coupling, i.e. that there always exists a Borel map  $\Sigma \rightarrow \text{Out}(F_N)$  which is almost everywhere equivariant with respect to the actions of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z}) \times \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . This is called *coupling rigidity* in [Kid11] and *tautness* in [BFS13].

As in [Fur99b], this is then rephrased in the language of stable orbit equivalence. The space  $\Sigma$  is equipped with an action of  $\Gamma_1 \times \Gamma_2 = \text{IA}_N(\mathbb{Z}/3\mathbb{Z}) \times \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Letting  $X_1$  be a finite measure fundamental domain for the  $\Gamma_1$ -action on  $\Sigma$ , there is a natural action of  $\Gamma_2$  on  $X_1$  (identified with  $\Sigma/\Gamma_1$ ). Likewise, letting  $X_2$  be a finite measure fundamental domain for the  $\Gamma_2$ -action on  $\Sigma$ , there is a natural action of  $\Gamma_1$  on  $X_2$ . We choose the fundamental domains  $X_1$  and  $X_2$  such that, letting  $Y = X_1 \cap X_2$ , one has  $(\Gamma_1 \times \Gamma_2)Y = \Sigma$  up to a null set. The orbit partitions coming from the  $\Gamma_1$ -action on  $X_2$  and from the  $\Gamma_2$ -action on  $X_1$  restrict to the same equivalence relation on  $Y$ .

Informally, one can think of  $Y$  as being equipped with arrows: there is an arrow from  $x$  to  $y$  whenever  $x$  and  $y$  are in the same  $\Gamma_1$ -orbit, or equivalently in the same  $\Gamma_2$ -orbit. The arrows are naturally equipped with two  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ -valued labels, indicating the element of  $\Gamma_i$  that sends  $x$  to  $y$ .

This is made formal in the language of measured groupoids (see Section 3 for precise definitions): we have a measured groupoid  $\mathcal{G}$  (the set of all arrows) over the base space  $Y$ , with two cocycles to  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  (the two labelings). In our terminology, these cocycles coming from actions of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  are important instances of *action-type* cocycles (see Definition 3.20). By careful bookkeeping of the above construction, the proof of measure equivalence rigidity of  $\text{Out}(F_N)$  reduces to the proof of the following statement,

as explained in Section 4.

**Theorem 1.17.** *Let  $N \geq 3$ . Let  $\mathcal{G}$  be a measured groupoid over a standard measure space  $Y$ . Then any two action-type cocycles  $\rho, \rho' : \mathcal{G} \rightarrow \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  are  $\mathrm{Out}(F_N)$ -cohomologous, i.e. there exist a Borel map  $\varphi : Y \rightarrow \mathrm{Out}(F_N)$  and a conull Borel subset  $Y^* \subseteq Y$  such that for every arrow  $g \in \mathcal{G}$  joining two points  $x, y \in Y^*$ , one has  $\rho'(g) = \varphi(y)\rho(g)\varphi(x)^{-1}$ .*

*Remark 1.18.* Restricting to the case where  $Y$  is reduced to a point, one recovers commensurator rigidity as follows: an isomorphism  $j$  between two finite index subgroups  $\Gamma_1, \Gamma_2$  of  $\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  yields two cocycles on the groupoid  $\Gamma_1$  given by  $j$  and the inclusion, and Theorem 1.17 shows that  $j$  is the restriction of an inner automorphism of  $\mathrm{Out}(F_N)$ .

### 1.4.2 Exploiting the rigidity of the free splitting graph

Of particular importance in the study of  $\mathrm{Out}(F_N)$  is the *free splitting graph*  $\mathbb{FS}$ , or its sibling, the *non-separating free splitting graph*  $\mathbb{FS}^{\mathrm{ns}}$ . The vertices of  $\mathbb{FS}^{\mathrm{ns}}$  are the decompositions of  $F_N$  as an HNN extension  $F_N = A*$ , up to equivariant isomorphism of their Bass–Serre trees (in  $\mathbb{FS}$ , one also allows for decompositions as a free product  $F_N = A * B$ ). Two such decompositions  $S, S'$  are joined by an edge when there is a two-edge graph of groups decomposition of  $F_N$  collapsing to  $S$  and  $S'$ . The group  $\mathrm{Out}(F_N)$  naturally acts on  $\mathbb{FS}^{\mathrm{ns}}$  and  $\mathbb{FS}$ . Building on work of Bridson and Vogtmann [BV01], Pandit proved [Pan14] that for  $N \geq 3$ ,  $\mathrm{Out}(F_N)$  is exactly the group of simplicial automorphisms of  $\mathbb{FS}^{\mathrm{ns}}$  (the same is true for  $\mathbb{FS}$ , see [AS11], but we will not use it). We will use additionally that every injective map  $\mathbb{FS}^{\mathrm{ns}} \rightarrow \mathbb{FS}$  preserving adjacency and non-adjacency takes values in the subgraph  $\mathbb{FS}^{\mathrm{ns}}$  (Proposition 15.1).

To prove Theorem 1.17, we follow a strategy initiated by Ivanov [Iva97] and pursued by Kida [Kid10] in the mapping class group setting, and associate to every  $y \in Y$  an automorphism of  $\mathbb{FS}^{\mathrm{ns}}$ . For this, ideally, we should characterize in a purely groupoid-theoretic way (i.e. with no reference to the cocycles  $\rho$  and  $\rho'$ ) those subgroupoids of  $\mathcal{G}$  that preserve a vertex of  $\mathbb{FS}^{\mathrm{ns}}$  for either  $\rho$  or  $\rho'$  (and characterize adjacency in a similar manner). In reality, we rather prove a partial characterization showing, roughly speaking, that the  $\rho$ -stabilizer of a splitting in  $\mathbb{FS}^{\mathrm{ns}}$  preserves a splitting in  $\mathbb{FS}$ . More precisely, given a vertex  $v \in \mathbb{FS}^{\mathrm{ns}}$ , one associates to the subgroupoid  $\rho^{-1}(\mathrm{Stab}(v))$  a countable Borel partition  $Y^* = \bigsqcup_i Y_i$  of a conull subset  $Y^* \subseteq Y$  such that  $\rho'(\mathcal{G}_{|Y_i})$  fixes a unique vertex  $v_i \in \mathbb{FS}$  (here  $\rho'(\mathcal{G}_{|Y_i})$  is the set of  $\rho'$ -labels of arrows with both endpoints in  $Y_i$ ). This defines for almost every  $y \in Y$  a map  $\Psi(y)$  from the vertex set of  $\mathbb{FS}^{\mathrm{ns}}$  to  $\mathbb{FS}$ , and we prove that  $\Psi$  preserves compatibility of splittings. Using the aforementioned facts, we show that in fact, for almost every  $y \in Y$ , the map  $\Psi(y)$  is an automorphism of  $\mathbb{FS}^{\mathrm{ns}}$ , i.e. an element of  $\mathrm{Out}(F_N)$ .

### 1.4.3 Towards a characterization of stabilizers of free splittings

The above discussion shows that the main theorem reduces to a groupoidal (partial) characterization of the stabilizers of non-separating free splittings of  $F_N$  (Theorem 13.10), an analogue of the algebraic version proved in [HW20].

In this introduction, we will focus on one crucial statement towards this goal, first presented in a group-theoretic fashion, before explaining the additional arguments required to improve it to a groupoid-theoretic version.

**Finding invariant nice splittings (group version).** Recall that a one-edge non-separating free splitting is the same as a decomposition as an HNN extension of the form  $F_N = A*$ . An important observation in their characterization, already exploited in [HW20], is that for  $N \geq 3$ , the stabilizer of such a splitting contains a normal subgroup (the *group of twists*) isomorphic to a direct product of two non-abelian free groups (made of all automorphisms having a representative in  $\text{Aut}(F_N)$  which acts as the identity on  $A$ , and multiply a stable letter  $t$  by elements of  $A$ , on the left and on the right). The next proposition provides a partial converse.

**Proposition 1.19.** *Let  $N \geq 3$ , and let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup. Assume that  $H$  contains a subgroup which is isomorphic to a direct product  $K_1 \times K_2$  of two normal non-amenable subgroups.*

*Then  $H$  fixes a nice splitting of  $F_N$ .*

The proof goes as follows. Since by Corollary 1.16, the property of preserving a nice splitting remains true by going to the normalizer, we are done if  $K_1$  or  $K_2$  preserves a nice splitting, hence also if some infinite cyclic subgroup  $\langle a_1 \rangle \subseteq K_1$  preserves a nice splitting. We can therefore assume that  $\langle a_1 \rangle$  is pure (as otherwise it preserves its dynamical decomposition), and to simplify the discussion, let us assume that it has a unique maximal invariant free factor system. The relative free factor graph  $\mathbb{FF}$  is a Gromov-hyperbolic graph on which  $a_1$  acts loxodromically, and its invariant endpoints in the Gromov boundary correspond to equivalence classes of  $\mathbb{R}$ -trees  $T_-, T_+$ . It follows that  $K_2$  preserves the pair of equivalence classes of  $T_\pm$ . Using arguments from [GL], the fact that  $T_\pm$  has non-amenable stabilizer allows to extract from  $T_\pm$  a canonical nice splitting that will thus be  $K_2$ -invariant.

**Finding invariant nice splittings (groupoid version): Adams' argument.** We need a groupoid-theoretic version of Proposition 1.19, given in Lemma 13.12. Its statement is similar, using a notion of amenable subgroupoid that originates in the work of Zimmer [Zim78], a notion of normal subgroupoid coined by Kida in [Kid08a] as a generalization of earlier work of Feldman, Sutherland and Zimmer [FSZ89], and a notion of pseudo-product of subgroupoids coined to replace the direct product in the statement (see Definition 13.5).

Our groupoid-theoretic proof stays very close to its group-theoretic version as long as it only involves objects taking values in a countable set, like nice splittings or free

factor systems. One reason is that a map from the base space  $Y$  to a countable set  $\mathcal{D}$  becomes constant after restriction to each part of a countable partition of  $Y$ , and all our groupoid-theoretic statements will allow to perform such partitions on the base space. In fact, the chain conditions given in the previous section also play a role here, to ensure essentially that here one never needs to perform infinitely many countable partitions at once. But there is a major difference when dealing with objects taking values in non-discrete spaces, like actions on  $\mathbb{R}$ -trees.

Thus, the proof of the groupoid-theoretic version of Proposition 1.19 requires the following key additional statement. All relevant definitions are given in Section 3.

**Theorem 1.20** (see Theorem 12.1). *Let  $N \geq 3$ . Let  $\mathcal{G}$  be a measured groupoid over a standard measure space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle. Let  $\mathcal{A}_1$  be an amenable subgroupoid of  $\mathcal{G}$  of infinite type, and let  $\mathcal{K}_2$  be a subgroupoid of  $\mathcal{H}$  that normalizes  $\mathcal{A}_1$ .*

*Then either  $\mathcal{K}_2$  is amenable, or there exists a Borel subset  $U \subseteq Y$  of positive measure such that  $(\mathcal{K}_2|_U, \rho)$  preserves a nice splitting.*

Here  $\mathcal{A}_1$  and  $\mathcal{K}_2$  are the groupoid analogues of the groups  $\langle a_1 \rangle$ ,  $K_2$  from the above argument.

The strategy of our proof of Theorem 1.20 (carried out in Section 12) relies on work of Adams [Ada94] that was already exploited by Kida in the context of mapping class groups. As above, we can assume that  $\mathcal{A}_1$  is pure, and let us assume it has a unique maximal invariant free factor system  $\mathcal{F}$ . Amenability of  $\mathcal{A}_1$  provides an equivariant map  $\mu$  that assigns to every point  $y \in Y$  a probability measure  $\mu_y$  on the compactification of Outer space (relative to  $\mathcal{F}$ ). We then take advantage of Reynolds' partition of this compactification into *arational* and *non-arational* trees [Rey12], and reduce to the case where  $\mu_y$  is supported almost everywhere on arational trees (otherwise we find invariant relative free factors, contradicting the maximality of  $\mathcal{F}$ ). Arational trees are in fact precisely those that describe points of  $\partial_\infty \mathbb{FF}$  by [BR15, Ham12, GH19b]. We then use a *barycenter map* constructed by Lécureux and the authors in [GHL20], which associates a finite set of conjugacy classes of free factors to every triple of pairwise inequivalent arational trees. This is used to ensure that  $\mu_y$  is almost everywhere supported on at most two equivalence classes of arational trees  $T_\pm(y)$ . Among the possible maps  $\mu$ , there is an essentially unique one  $\mu_{\max}$  with maximal support, so  $\mu_{\max}$  is also equivariant under the groupoid  $\mathcal{K}_2$ . If in restriction to a subset  $U \subseteq Y$  of positive measure, the pair  $\{T_-(y), T_+(y)\}$  has non-amenable stabilizer, we can build a canonical nice splitting which is invariant under  $\mathcal{A}_1$  and  $\mathcal{K}_2$  as in the group-theoretic case. Otherwise, the stabilizer of  $\{T_-(y), T_+(y)\}$  is amenable for almost every  $y \in Y$ , and we reach a contradiction to the non-amenable of  $\mathcal{K}_2$  from the following statement, a variation on work of Bestvina and the authors [BGH17] showing the amenability of the  $\text{Out}(F_N)$ -action on the space of arational trees. See Definition 3.36 for the definition of Borel amenability of a group action.

**Theorem 1.21** (see Corollary 12.16). *Let  $N \geq 2$ , and let  $\mathcal{F}$  be a non-sporadic free factor system of  $F_N$ . Let  $\mathcal{AT}^a$  be the space of all arational  $(F_N, \mathcal{F})$ -trees whose stabilizer in*



$\text{Out}(F_N, \mathcal{F})$  is amenable.

The action of  $\text{Out}(F_N, \mathcal{F})$  on  $\mathcal{AT}^a$  is Borel amenable.

Similarly, the action of  $\text{Out}(F_N, \mathcal{F})$  on the set of pairs of arational  $(F_N, \mathcal{F})$ -trees  $\{T_+, T_-\}$  whose stabilizer is amenable, is a Borel amenable action.

**Additional arguments.** Once the groupoid-theoretic version of Proposition 1.19 has been established, more work is still required to distinguish free splittings among nice splittings. Briefly,  $\mathcal{Z}_{\max}$ -splittings are distinguished from the fact that their stabilizer contains a normal non-trivial abelian subgroup (its *group of twists*). The groupoid-theoretic statement used to distinguish bi-nonsporadic splittings is quite technical; roughly, the two vertices from the definition of such a splitting enable us to find two commuting normal subgroups in its stabilizer that both contain direct products of free groups, and this does not happen in the stabilizer of a non-separating free splitting.

More work is also needed to establish a characterization of adjacency in  $\mathbb{FS}^{\text{ns}}$ . Again, the details are quite involved (see Section 14); as in [HW20], one underlying idea is that two one-edge non-separating free splittings of  $F_N$  are adjacent in  $\mathbb{FS}^{\text{ns}}$  if and only if their common stabilizer does not fix any third one-edge free splitting.

## 1.5 Structure of the paper

The paper has three parts.

The first part gives preliminary background, both on  $\text{Out}(F_N)$  (Section 2) and on measured groupoids (Section 3) and their use in the study of measure equivalence (Section 4). In particular, Section 4 explains how to reduce the main theorem of the present paper and its consequences to the rigidity statement for groupoids and their action-type cocycles (Theorem 1.17 in this introduction).

The second part does not involve any groupoids, it is devoted to our constructions of canonical splittings and our descriptions of stabilizers of collections of splittings. Section 5 collects further background regarding invariant splittings and trees of cylinders. The construction of a canonical splitting encoding all invariant free splittings is carried out in Section 6, the variant for  $\mathcal{Z}_{\max}$ -splittings is the contents of Section 7, and the dynamical decomposition is constructed in Section 8. Finally, Section 9 contains the consequence to the fact that having an invariant nice splitting passes to the normalizer.

The third part is the heart of our proof of measure equivalence rigidity of  $\text{Out}(F_N)$ . In Section 10, we translate to the groupoid setting the fact that having invariant nice splittings passes to the normalizer. We also set up the notions of pure, nice-averse and stably nice subgroupoids that provide the groupoid-theoretic framework in which the later sections are phrased. As observed in this introduction, direct products of free groups play an important role in the present paper, and Section 11 is concerned with measured groupoids with cocycles towards such direct products. In Section 12, we run our version of Adams' argument and prove Theorem 1.20. Section 13 is the heart of our proof, culminating in Theorem 13.10; it gives the necessary conditions for stabilizing a non-separating free splitting, and sufficient conditions for stabilizing a free splitting.



Section 14 deals with the characterization of adjacency in  $\mathbb{F}\mathbb{S}^{\text{ns}}$ . In Section 15, we prove that every injective map from  $\mathbb{F}\mathbb{S}^{\text{ns}}$  to  $\mathbb{F}\mathbb{S}$  which preserves adjacency and non-adjacency must take its values in  $\mathbb{F}\mathbb{S}^{\text{ns}}$  (as explained above, this compensates the slight asymmetry in the statement of Theorem 13.10). The proofs of our main theorem and its consequences are completed in Section 16.

**Acknowledgments.** We would like to warmly thank Damien Gaboriau for suggesting this problem, organizing a session at the IHP in 2011 with this problem in view within the framework of the trimester *Von Neumann algebras and ergodic theory of group actions*, and for helpful discussions and encouragements. We thank Ric Wade and Martin Bridson for pointing out a missing assumption in an earlier version of Corollary 1.6.

The first named author acknowledges support from the Centre Henri Lebesgue (Labex ANR-11-LABX-0020-01). The second named author acknowledges support from the Agence Nationale de la Recherche under Grant ANR-16-CE40-0006.

# Contents

1	Introduction	1
<b>I</b>	<b>Preliminaries</b>	<b>18</b>
2	Preliminaries on $\text{Out}(F_N)$	18
3	Preliminaries on measured groupoids	27
4	Groupoids, measure equivalence, and applications	37
<b>II</b>	<b>Stabilizers of collections of splittings or factors</b>	<b>43</b>
5	$H$ -invariant splittings and trees of cylinders	43
6	Invariant free splittings and stabilizers of collections of free splittings	44
7	Invariant $\mathcal{Z}_{\max}$ -splittings and stabilizers of collections of $\mathcal{Z}_{\max}$ -splittings	57
8	Pure subgroups of $\text{Out}(F_N)$ and dynamical decomposition	71
9	Canonical nice splittings	89
<b>III</b>	<b>Measure equivalence rigidity of <math>\text{Out}(F_N)</math></b>	<b>93</b>
10	Canonical invariant splittings for groupoids	93
11	Groupoids with cocycles to direct products of free groups	100
12	Nice-averse amenable subgroupoids: Adams' argument	103
13	Detecting when a subgroupoid stabilizes a free splitting	118
14	Characterizing compatibility	132
15	Maps from $\mathbb{F}\mathbb{S}^{\text{ns}}$ to $\mathbb{F}\mathbb{S}$	135
16	Measure equivalence rigidity of $\text{Out}(F_N)$	137

## Part I

# Preliminaries

## 2 Preliminaries on $\text{Out}(F_N)$

### 2.1 Splittings and deformation spaces

**General definitions.** Let  $G$  be a group, and let  $T$  be either a simplicial  $G$ -tree, or an  $\mathbb{R}$ -tree equipped with an isometric  $G$ -action. An element  $g \in G$  acts *elliptically* on  $T$  if  $g$  fixes a point in  $T$ , and *hyperbolically* otherwise. The  $G$ -action on  $T$  is *minimal* if  $T$  does not contain any proper nonempty  $G$ -invariant subtree. Assuming that some element of  $G$  acts hyperbolically on  $T$ , the  $G$ -action on  $T$  always contains a unique nonempty minimal  $G$ -invariant subtree [CM87, Proposition 3.1].

A *splitting* of  $G$  is a minimal, simplicial  $G$ -action on a simplicial tree with no edge inversion. Splittings are always considered up to equivariant homeomorphism.

Given a splitting  $S$  of  $G$  and a vertex  $v \in S$ , we denote by  $G_v$  the  $G$ -stabilizer of  $v$ . Likewise, given an edge  $e \subseteq S$ , we denote by  $G_e$  the  $G$ -stabilizer of  $e$ . We will also always denote by  $\text{Inc}_v$  the set of all  $G_v$ -conjugacy classes of stabilizers of edges incident on  $v$ .

Let  $S$  and  $S'$  be two splittings of  $G$ . The splitting  $S$  *dominates*  $S'$  if there exists a  $G$ -equivariant map  $S \rightarrow S'$ . This is equivalent to saying that every subgroup of  $G$  which fixes a point in  $S$ , also fixes a point in  $S'$ . Two splittings are *in the same deformation space* if they dominate each other.

The splitting  $S$  *refines*  $S'$  if  $S$  collapses onto  $S'$ , i.e.  $S'$  is obtained from  $S$  by collapsing every edge in a  $G$ -invariant collection of edges to a point (in other words, there exists a  $G$ -equivariant alignment-preserving map  $S \rightarrow S'$ ). Two splittings  $S$  and  $S'$  are *compatible* if they admit a common refinement. In this case  $S$  and  $S'$  have a minimal common refinement, onto which every other refinement collapses.

A subgroup  $A \subseteq G$  is *elliptic* in  $S$  if it fixes a point in  $S$ . Given a collection  $\mathcal{F}$  of conjugacy classes of subgroups of  $G$ , we say that a splitting  $S$  is *relative to  $\mathcal{F}$*  (or that  $S$  is a splitting of  $(G, \mathcal{F})$ ) if every subgroup in  $\mathcal{F}$  is elliptic in  $S$ .

A *one-edge* splitting of  $G$  is a splitting of  $G$  that contains a single orbit of edges.

**Free splittings and Grushko deformation spaces.** A *free splitting* of  $G$  is a splitting of  $G$  whose edge stabilizers are all trivial.

Given a collection  $\mathcal{F}$  of conjugacy classes of subgroups of  $G$ , a *Grushko splitting* of  $G$  relative to  $\mathcal{F}$  is a free splitting  $S$  of  $G$  relative to  $\mathcal{F}$  which is maximal for domination. Any two such splittings are in the same deformation space. The set of all Grushko splittings of  $G$  relative to  $\mathcal{F}$  is called the *Grushko deformation space* of  $G$  relative to  $\mathcal{F}$ .

**$\mathcal{Z}_{\max}$ -splittings.** A  $\mathcal{Z}_{\max}$ -*splitting* of  $F_N$  is a splitting of  $F_N$  in which all edge stabilizers are infinite cyclic and root-closed. The following lemma follows for instance from [BF91].

**Lemma 2.1.** *There is a bound, only depending on  $N$ , on the number of orbits of edges of a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$  with no vertex of valence 2.*

We mention that given a splitting  $S$  of  $F_N$  with non-trivial cyclic edge stabilizers, one can construct a canonical  $\mathcal{Z}_{\max}$ -splitting  $S_{\mathcal{Z}_{\max}}$  as follows.

**Definition 2.2** (see [GL17, Lemma 9.27]). *Let  $S$  be a splitting of  $F_N$  whose edge stabilizers are all isomorphic to  $\mathbb{Z}$ . One defines  $S_{\mathcal{Z}_{\max}}$  as the minimal subtree of the quotient of  $S$  by the equivalence relation defined by  $e \sim e'$  if  $e = ge'$  for some  $g \in F_N$  such that  $g^k$  fixes  $e$  for some  $k \geq 1$ .*

By [GL17, Lemma 9.27], every  $\mathcal{Z}_{\max}$ -splitting of  $F_N$  dominated by  $S$  is dominated by  $S_{\mathcal{Z}_{\max}}$ . It will be useful to describe  $S$  assuming that  $S_{\mathcal{Z}_{\max}}$  is trivial.

**Lemma 2.3** (See Figure 2). *Let  $S$  be a splitting of  $F_N$  with infinite cyclic edge stabilizers. Assume that  $S_{\mathcal{Z}_{\max}}$  is trivial (i.e. reduced to a point).*

*Then  $S/F_N$  is a tree of groups, with a unique vertex  $v$  whose stabilizer is not abelian.*

*In particular, assuming moreover that every edge of  $S$  is adjacent to at least one vertex with non-abelian stabilizer, it follows that  $S/F_N$  is a star, all vertices  $u_1, \dots, u_n$  distinct from  $v$  are joined by a unique edge  $e_i$  to  $v$ ,  $G_{u_i}$  is cyclic and  $[G_{u_i} : G_{e_i}] \geq 2$ .*




Figure 2: Splittings such that  $S_{\mathcal{Z}_{\max}}$  is trivial.

*Proof.* Denote by  $\tilde{S}_{\mathcal{Z}_{\max}}$  the quotient of  $S$  by the equivalence relation of Definition 2.2, and note that  $S/F_N$  and  $\tilde{S}_{\mathcal{Z}_{\max}}/F_N$  are isomorphic as graphs. Our assumption says that  $F_N$  fixes a point in  $\tilde{S}_{\mathcal{Z}_{\max}}$ . In particular  $\tilde{S}_{\mathcal{Z}_{\max}}/F_N$  is a tree, hence so is  $S/F_N$ .

If  $v, v' \in S$  are two vertices with non-abelian stabilizer and in distinct orbits, then  $v$  and  $v'$  have distinct images in  $\tilde{S}_{\mathcal{Z}_{\max}}/F_N$ , and since edge stabilizers of  $S_{\mathcal{Z}_{\max}}$  are cyclic, this prevents  $F_N$  from having a global fixed point in  $\tilde{S}_{\mathcal{Z}_{\max}}$ .  $\square$

## 2.2 Free factor systems

A *free factor* of  $F_N$  is a subgroup of  $F_N$  that arises as a point stabilizer in a (maybe trivial) free splitting of  $F_N$  (in particular,  $\{1\}$  and  $F_N$  are free factors). Given a collection  $\mathcal{F}$  of conjugacy classes of subgroups of  $F_N$ , a *free factor* of  $F_N$  relative to  $\mathcal{F}$  is a subgroup of  $F_N$  that arises as a point stabilizer in some free splitting of  $F_N$  relative to  $\mathcal{F}$ . A free factor  $A$  of  $F_N$  relative to  $\mathcal{F}$  is *proper* if  $A \neq F_N$ ,  $A \neq \{1\}$  and  $A$  is not conjugate in a group appearing in  $\mathcal{F}$ .

A *free factor system* of  $F_N$  is a collection of conjugacy classes of non-trivial free factors that arise as the non-trivial point stabilizers in some non-trivial (i.e. not reduced to a point) free splitting of  $F_N$ . For instance  $\emptyset$  is a free factor system, but  $\{[F_N]\}$  is not.

A free factor system  $\mathcal{F}$  of  $F_N$  is *sporadic* if either  $F_N$  splits as an HNN extension  $F_N = A*$  and  $\mathcal{F} = \{[A]\}$ , or  $F_N$  splits as a free product  $F_N = A*B$  and  $\mathcal{F} = \{[A], [B]\}$ . Otherwise  $\mathcal{F}$  is *non-sporadic*.

If  $\mathcal{F} = \{[P_1], \dots, [P_k]\}$  is a collection of conjugacy classes of subgroups of  $F_N$ , we say that a subgroup  $A \subseteq F_N$  is  $\mathcal{F}$ -*peripheral* if  $A$  is trivial or conjugate in some  $P_i$ . If  $\mathcal{F}, \mathcal{F}'$  are two collections of conjugacy class of subgroups of  $F_N$ , we write  $\mathcal{F} \preceq \mathcal{F}'$  if every group in  $\mathcal{F}$  is  $\mathcal{F}'$ -peripheral. When  $\mathcal{F}, \mathcal{F}'$  are free factor systems, this happens if and only if any Grushko  $(F_N, \mathcal{F})$ -tree dominates any Grushko  $(F_N, \mathcal{F}')$ -tree. This defines an ordering on free factor systems of  $F_N$ .

If  $\mathcal{F}$  is a free factor system of  $F_N$ , and  $A$  is a proper free factor of  $F_N$  relative to  $\mathcal{F}$ , then there is a smallest free factor system  $\mathcal{F}'$  with  $\{[A]\}, \mathcal{F} \preceq \mathcal{F}'$ , it consists of  $[A]$  together with the conjugacy classes of groups in  $\mathcal{F}$  that are not conjugate in  $A$ . We denote it by  $\mathcal{F} \vee \{[A]\}$ .

If  $\mathcal{F} = \{[P_1], \dots, [P_n]\}$  is a collection of conjugacy classes of subgroups of  $F_N$ , and  $A$  is a subgroup of  $F_N$ , we let  $\mathcal{F}|_A$  be the set of all  $A$ -conjugacy classes of non-trivial subgroups of  $A$  of the form  $P_i^g \cap A$  for  $g \in F_N$ . When  $\mathcal{F}$  is a free factor system,  $\mathcal{F}|_A$  also coincides with the set of all conjugacy classes (in  $A$ ) of non-trivial vertex stabilizers in the  $A$ -minimal subtree of any Grushko  $(F_N, \mathcal{F})$ -tree. It is a free factor system of  $A$  unless  $A$  is  $\mathcal{F}$ -peripheral.

### 2.3 Relative automorphism groups

Given a collection  $\mathcal{F}$  of conjugacy classes of subgroups of  $F_N$ , we denote by  $\text{Out}(F_N, \mathcal{F})$  the subgroup of  $\text{Out}(F_N)$  made of all outer automorphisms  $\alpha$  such that for every conjugacy class  $[A] \in \mathcal{F}$ , one has  $\alpha([A]) = [A]$ . We denote by  $\text{Out}(F_N, \mathcal{F}^{(t)})$  the subgroup made of all automorphisms  $\alpha$  such that for every representative  $\tilde{\alpha}$  of  $\alpha$  in  $\text{Aut}(F_N)$ , and every subgroup  $A \subseteq F_N$  whose conjugacy class belongs to  $\mathcal{F}$ , there exists  $g_{\tilde{\alpha}, A} \in F_N$  such that  $\tilde{\alpha}|_A$  is given by the conjugation by  $g_{\tilde{\alpha}, A}$ .

An automorphism  $\alpha \in \text{Out}(F_N, \mathcal{F})$  is *atoroidal* relative to  $\mathcal{F}$  if any element whose conjugacy class is periodic under  $\alpha$  is  $\mathcal{F}$ -peripheral. We say that  $\alpha$  is *fully irreducible* relative to  $\mathcal{F}$  if no power  $\alpha^k$  with  $k \neq 0$  preserves the conjugacy class of a proper free factor of  $F_N$  relative to  $\mathcal{F}$ .

### 2.4 Outer space and its closure

Let  $\mathcal{F}$  be a free factor system of  $F_N$ . A *Grushko  $(F_N, \mathcal{F})$ -tree* is a minimal, simplicial metric  $(F_N, \mathcal{F})$ -tree whose underlying simplicial tree is a Grushko splitting of  $F_N$  relative to  $\mathcal{F}$ . The *unprojectivized relative Outer space*  $\mathcal{O}(F_N, \mathcal{F})$  is the space of all  $F_N$ -equivariant isometry classes of Grushko  $(F_N, \mathcal{F})$ -trees (see [GL07b]). When  $\mathcal{F} = \emptyset$ , this is nothing but the unprojectivized version of Culler and Vogtmann's Outer space [CV86]. When  $\mathcal{F}$  is clear from the context, we will simply write  $\mathcal{O}$  instead of  $\mathcal{O}(F_N, \mathcal{F})$ . We denote by

$\mathbb{P}\mathcal{O}$  the projectivized outer space, where trees are considered up to homothety instead of isometry. The group  $\text{Out}(F_N, \mathcal{F})$  naturally acts on  $\mathcal{O}$  and  $\mathbb{P}\mathcal{O}$  by precomposition of the action.

The projectivized Outer space  $\mathbb{P}\mathcal{O}$  can be compactified by looking at its closure  $\overline{\mathbb{P}\mathcal{O}}$  in the space of all projective classes of  $(F_N, \mathcal{F})$ -trees (i.e.  $\mathbb{R}$ -trees equipped with an isometric action of  $F_N$  for which every subgroup in  $\mathcal{F}$  is elliptic), equipped with the equivariant Gromov–Hausdorff topology introduced by Paulin in [Pau88] (or equivalently, see [Pau89], with the length function topology studied by Culler and Morgan [CM87]). The closure is identified with the space of all projective classes of *very small*  $(F_N, \mathcal{F})$ -trees, i.e.  $(F_N, \mathcal{F})$ -trees in which all stabilizers of nondegenerate arcs are either trivial or nonperipheral, cyclic and root-closed, and all stabilizers of nondegenerate tripods are trivial ([CL95, BF94] for the case where  $\mathcal{F} = \emptyset$ , and [Hor17] in the relative setting). We also denote by  $\overline{\mathcal{O}}$  the unprojectivized version of  $\overline{\mathbb{P}\mathcal{O}}$ , where trees are considered up to isometry instead of homothety. The  $\text{Out}(F_N, \mathcal{F})$ -actions on  $\mathcal{O}$  and  $\mathbb{P}\mathcal{O}$  extend to actions on  $\overline{\mathcal{O}}$  and  $\overline{\mathbb{P}\mathcal{O}}$ . We let  $\partial\mathcal{O} = \overline{\mathcal{O}} \setminus \mathcal{O}$ .

## 2.5 Arational trees

Arational trees were first introduced by Reynolds [Rey12] as a free group analogue of arational measured foliations on surfaces. A tree  $T \in \partial\mathcal{O}(F_N, \mathcal{F})$  is *arational* if no proper  $(F_N, \mathcal{F})$ -free factor acts elliptically on  $T$ , and for every proper  $(F_N, \mathcal{F})$ -free factor  $A$ , the  $A$ -action on its minimal subtree  $T_A$  is a Grushko  $(A, \mathcal{F}|_A)$ -tree.

For every  $T \in \partial\mathcal{O}(F_N, \mathcal{F})$  and every  $\alpha \in \text{Out}(F_N, \mathcal{F})$ , it follows from the definition of the action of  $\text{Out}(F_N, \mathcal{F})$  on  $\overline{\mathbb{P}\mathcal{O}}(F_N, \mathcal{F})$  that  $\alpha$  fixes the homothety class of  $T$  if and only if for every  $\tilde{\alpha} \in \text{Aut}(F_N)$  representing  $\alpha$ , there exists a homothety  $H_{\tilde{\alpha}}$  of  $T$  (i.e. a map that multiplies all distances by the same factor  $\lambda$ ) which is  $\tilde{\alpha}$ -equivariant in the sense that

$$H_{\tilde{\alpha}}(gx) = \tilde{\alpha}(g)H_{\tilde{\alpha}}(x)$$

for every  $g \in F_N$  and every  $x \in T$ . Once the representative  $\tilde{\alpha}$  is chosen, the homothety  $H_{\tilde{\alpha}}$  is unique. The factor  $\lambda$  does not depend on the chosen representative and is called the *dilatation* of  $\alpha$  with respect to  $T$ .

**Proposition 2.4.** *Let  $\mathcal{F}$  be a non-sporadic free factor system of  $F_N$ , and let  $\alpha \in \text{Out}(F_N, \mathcal{F})$  be an automorphism which is fully irreducible relative to  $\mathcal{F}$ .*

*Then there exists an arational  $(F_N, \mathcal{F})$ -tree  $T$  whose homothety class is  $\alpha$ -invariant, and such that the dilatation of  $\alpha$  with respect to  $T$  is not equal to 1.*

*Proof.* It follows from work of Francaviglia and Martino [FM15, Theorem 8.23] that  $\alpha$  has an invariant axis in  $\mathcal{O}(F_N, \mathcal{F})$ . Let  $T$  be a limiting tree of such an axis. Then  $T$  is arational, as otherwise  $\alpha$  would have a non-trivial power that preserves one of the finitely many canonical reducing factors of  $T$  (see [Rey12] or [Hor14, Section 4.4]). That the dilatation of  $\alpha$  acting on  $T$  is not equal to 1 follows from [GH19b, Remark 6.4].  $\square$

## 2.6 Transverse families and transverse coverings

Let  $T$  be an  $F_N$ -tree. A *transverse family* in  $T$  is an  $F_N$ -invariant collection  $\mathcal{Y}$  of nondegenerate subtrees of  $T$  (i.e. nonempty and not reduced to a single point) such that any two distinct trees in  $\mathcal{Y}$  intersect in at most a point. A *transverse covering* of  $T$  is a transverse family  $\mathcal{Y}$  made of closed subtrees such that every segment in  $T$  is covered by finitely many subtrees in  $\mathcal{Y}$ .

The *skeleton* of a transverse covering  $\mathcal{Y}$  is the bipartite simplicial tree having one vertex  $v_Y$  for every subtree in  $\mathcal{Y}$ , one vertex  $v_x$  for every point  $x \in T$  that belongs to at least two trees in  $\mathcal{Y}$ , with an edge joining  $v_x$  to  $v_Y$  whenever  $x \in Y$ , see [Gui04, Definition 4.8].

An  $F_N$ -tree  $T$  is *mixing* if given any two segments  $I, J \subseteq T$ , there exists a finite collection  $\{g_1, \dots, g_k\}$  of elements of  $F_N$  such that  $J \subseteq g_1 I \cup \dots \cup g_k I$ . In a mixing tree, every transverse family made of closed subtrees is a transverse covering.

Given a non-sporadic free factor system  $\mathcal{F}$  of  $F_N$ , arational  $(F_N, \mathcal{F})$ -trees are always mixing [Hor14, Lemma 4.9].

## 2.7 QH vertex groups

The notions introduced under this heading will only be used in Sections 7 and 8.

QH vertex groups are a key notion in the theory of JSJ decompositions of groups, they are used in the description of certain canonical splittings. Unformally, this is a surface attached to the rest of the group by its boundary. QH stands for *quadratically hanging*, and refers to the fact that the attachment can be described algebraically by quadratic words.

Let  $\Sigma$  be a compact connected surface with boundary with negative Euler characteristic. A *boundary subgroup* is a subgroup of  $\pi_1(\Sigma)$  which is conjugate to a subgroup of the fundamental group of a boundary component of  $\Sigma$ . Maximal boundary subgroups are the conjugates of the fundamental groups of the boundary components of  $\Sigma$ .

**Definition 2.5.** *Let  $G$  be a group, and let  $S$  be a splitting of  $G$ . A vertex  $v$  of  $S$  is a clean QH vertex if it comes with an identification  $G_v \simeq \pi_1(\Sigma)$  where  $\Sigma$  is a (maybe non-orientable) compact hyperbolic surface with geodesic boundary, and each incident edge group is a maximal boundary subgroup of  $\pi_1(\Sigma)$ , and this induces an injection from the set of incident edges to the set of boundary components of  $\Sigma$ .*

*Boundary components not in the image of this injection are called unused.*

This definition will be sufficient in most situations, but some arguments require a more general definition.

**Definition 2.6** ([GL17, Definition 5.13]). *Let  $A$  be a group and  $\mathcal{Q}$  a collection of conjugacy classes of subgroups of  $A$ . The pair  $(A, \mathcal{Q})$  is QH with trivial fiber if there exists a connected compact hyperbolic 2-orbifold  $\Sigma$  with boundary, and an identification  $A \simeq \pi_1(\Sigma)$  such that, under this identification, each group in  $\mathcal{Q}$  is either finite or a boundary subgroup of  $\Sigma$ .*



A pair  $(A, \mathcal{Q})$  is QH with fiber  $F$  if  $F \triangleleft A$ , and  $(A/F, \bar{\mathcal{Q}})$  is QH with trivial fiber, where  $\bar{\mathcal{Q}}$  is the image of  $\mathcal{Q}$  in  $A/F$ .

Given a collection  $\mathcal{P}$  of conjugacy classes of subgroups of  $G$ , a vertex  $v$  of a splitting of  $(G, \mathcal{P})$  is a QH vertex relative to  $\mathcal{P}$  if  $(G_v, \text{Inc}_v \cup \mathcal{P}|_{G_v})$  is QH (maybe with fiber).

## 2.8 Twists of splittings

Given a group  $G$  and an element  $g \in G$ , we denote by  $\text{ad}_g$  the inner automorphism  $x \mapsto gxg^{-1}$ .

Let  $S$  be a splitting of a group  $G$ , and  $e = uv$  an edge of  $S$ . Following Levitt [Lev05], given an element  $z \in G_u$  that centralizes  $G_e$ , one defines as follows an outer automorphism  $\tau \in \text{Out}(G)$  called the *twist* by  $z$  around  $e$  near  $u$ .

First assume that the image of  $e$  in  $S/G$  is separating, and consider the corresponding amalgam  $G = A *_{G_e} B$ , where  $A$  contains  $G_u$  and  $B$  contains  $G_v$ . Then  $\tau$  is the outer automorphism of  $G$  defined as the identity on  $A$  and as  $\text{ad}_z$  in restriction to  $B$ . Similarly, if  $e$  is non-separating, and  $t$  maps the origin of  $e$  to its terminus after collapsing all edges that are not in the orbit of  $e$ , then the corresponding HNN extension gives a presentation of the form  $G = \langle A, t | t^{-1}gt = \varphi(g) \rangle$ . Then  $\tau$  is the outer automorphism of  $G$  defined as the identity on  $A$  and sending  $t$  to  $zt$ . Notice that the outer automorphism  $\tau$  defined in this way does not depend on the choice of the stable letter  $t$ , as any other choice is of the form  $t' = ta$  for some  $a \in G_u$ .

In both cases (separating and non-separating), changing  $e, u$  by a translate, and conjugating  $z$  accordingly does not change  $\tau$  as an outer automorphism, so it may be described from the data in the graph of groups. Moreover if  $\tau$  is a twist around  $e$  near  $u$  and  $\tau'$  is a twist around  $e'$  near  $u'$  and if the pair  $(e, u)$  is not in the orbit of  $(e', u')$ , then  $\tau$  and  $\tau'$  commute. The *group of twists* of  $S$  is the subgroup  $\text{Tw}$  of  $\text{Out}(G)$  generated by all twists.

An important property is that  $\tau$  preserves  $S$ : for any representative  $\tilde{\tau} \in \text{Aut}(G)$  of  $\tau$ , there is a  $\tilde{\tau}$ -equivariant isomorphism  $f : S \rightarrow S$  (i.e. such that  $f(gx) = \tilde{\tau}(g)f(x)$  for all  $x \in S$  and all  $g \in F_N$ ).

We now give three examples that will play an important role.

If  $S$  is a splitting of  $F_N$  such that all edge stabilizers are non-abelian, then the group of twists is trivial because edge stabilizers have trivial centralizer.

If  $S$  is a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ , and  $e = uv$  is an edge in  $S$ , then  $G_e$  is its own centralizer. It follows that twists around  $e$  near  $u$  agree with twists around  $e$  near  $v$ : we just say that  $\tau$  is a twist around  $e$ . Moreover, in this case, the group of twists is abelian.

If  $S$  is a splitting of  $F_N$  with trivial edge stabilizers, the group of twists can be much larger. For instance, if  $S$  is dual to the HNN extension  $A*$  (with  $A$  a corank one free factor), then assuming that  $A$  is non-abelian, the group of twists of  $S$  is isomorphic to  $A \times A$ , generated by twists around  $e$  near both sides of  $e$ . Explicitly, if  $t$  is a stable letter of the HNN extension, then  $\text{Tw}$  is the set of outer automorphisms that have a representative in  $\text{Aut}(F_N)$  acting as the identity on  $A$  and sending the stable letter  $t$  to  $atb^{-1}$  for some  $(a, b) \in A \times A$ . If  $A$  is cyclic, then there is an extra relation among the

twists and  $\text{Tw} \simeq A$  (in fact, if  $A$  is an arbitrary group, the group of twists of the HNN extension  $A*$  is isomorphic to  $(A \times A)/Z(A)$  where  $Z(A)$  is the center of  $A$ , diagonally embedded in  $A \times A$ ).

If  $S$  is dual to a splitting  $F_N = A * B$  with  $A, B$  non-abelian, then  $\text{Tw} \simeq A \times B$ . If  $A$  is abelian, then twists near  $A$  are trivial as outer automorphisms.

We note that a twist around an edge of a splitting with trivial stabilizer can be seen as a twist of a cyclic splitting. For example, if  $S$  is dual to a free product decomposition  $F_N = A * B$ , and  $z \in A$ , then the twist  $\tau$  by  $z$  around  $e$  near  $u$  can be seen as a twist of the cyclic splitting  $S'$  dual to the decomposition  $F_N = A *_{\langle z \rangle} \langle B, z \rangle$ . More generally, if  $e = uv$  is an edge of  $S$  with trivial stabilizer, and  $z \in G_u$ , then the twist  $\tau$  by  $z$  around  $e$  near  $u$  can be seen as a twist of the cyclic splitting  $S'$  obtained by folding  $e$  with its translate  $ze$ . More precisely, if  $e' = u'v'$  is the image of  $e$  in  $S'$ , then  $\tau$  is also the twist by  $z$  around  $e'$  near  $u'$ .

## 2.9 The exact sequence for automorphisms of splittings

Given a splitting  $S$  of  $F_N$ , we denote by  $\text{Stab}(S)$  the stabilizer of  $S$  in  $\text{Out}(F_N)$ . Recall that  $\alpha \in \text{Stab}(S)$  if and only if for some (equivalently every)  $\tilde{\alpha} \in \text{Aut}(F_N)$  representing  $\alpha$  there exists a (unique if  $S$  is not a line)  $\tilde{\alpha}$ -equivariant isometry  $f_{\tilde{\alpha}} : S \rightarrow S$ . The subgroup  $\text{Stab}^0(S)$  consisting of outer automorphisms  $\alpha$  such that  $f_{\tilde{\alpha}}$  acts trivially on  $S/F_N$  (this does not depend on the choice of  $\tilde{\alpha}$ ) has finite index. For each  $\alpha \in \text{Stab}^0(S)$ , the restriction of  $\alpha$  to  $G_v$  gives a well defined outer automorphism of  $G_v$ . One thus gets a restriction map  $\rho : \text{Stab}^0(S) \rightarrow \prod_{v \in S/F_N} \text{Out}(G_v)$ .

**Proposition 2.7** (Levitt [Lev05]). *Let  $S$  be a splitting of  $F_N$  with finitely generated edge stabilizers. Then one has an exact sequence*

$$1 \rightarrow \text{Tw} \rightarrow \text{Stab}^0(S) \xrightarrow{\rho} \prod_{v \in S/F_N} \text{Out}(G_v, \text{Inc}_v)$$

and the image of  $\rho$  contains  $\prod_{v \in S/F_N} \text{Out}(G_v, \text{Inc}_v^{(t)})$ .

*Proof.* The assertion on the image of  $\rho$  is [Lev05, Proposition 2.1]. By [Lev05, Proposition 2.2],  $\ker \rho$  is generated by *bitwists* (see Section 2.4 in [Lev05]). Thus, we just have to check that bitwists are twists.

Consider an edge  $e = uv$  of  $S$ , and assume that  $e$  is separating in  $S/F_N$  and induces a splitting of the form  $F_N = A *_{G_e} B$ . Given  $z \in G_u$  and  $z' \in G_v$  normalizing  $G_e$  and such that  $zz'^{-1}$  centralizes  $G_e$ , one defines a bitwist by  $\text{ad}_z$  in restriction to  $A$  and  $\text{ad}_{z'}$  in restriction to  $B$ . Because  $G_e$  is finitely generated, it has finite index in its normalizer  $N(G_e)$ , so  $N(G_e)$  fixes a point in  $S$ . It follows that  $z$  and  $z'$  fix this point so one of them fixes  $e$ . This implies that  $zz'^{-1}$  fixes one endpoint of  $e$ , and the bitwist is actually a twist by  $zz'^{-1}$  at  $e$  near one of its endpoints. The case of an HNN extension is similar and left to the reader.  $\square$

*Remark 2.8.* In many situations, one knows that a subgroup  $H \subseteq \text{Stab}^0(S)$  acts trivially on edge groups in the following sense: for every  $\tilde{\alpha} \in \text{Aut}(F_N)$  representing an element of  $H$ , and every edge group  $G_e$ , there exists  $g \in F_N$  such that  $\tilde{\alpha}|_{G_e} = (\text{ad}_g)|_{G_e}$ . If each edge stabilizer is its own normalizer, this implies that  $\rho(H)$  is contained in  $\prod_{v \in S/F_N} \text{Out}(G_v, \text{Inc}_v^{(t)})$ .

The following lemma is an immediate consequence.

**Lemma 2.9.** *Let  $S$  be a splitting of  $F_N$  whose edge stabilizers are non-abelian and finitely generated.*

*Then  $\text{Stab}^0(S)$  contains a subgroup that maps isomorphically to  $\prod_{v \in S/F_N} \text{Out}(G_v, \text{Inc}_v^{(t)})$  under the restriction map.*

*Proof.* Since edge stabilizers are non-abelian, they have trivial centralizer in  $F_N$  so  $\text{Tw}$  is trivial and  $\rho$  is one-to-one. Since  $\prod_{v \in S/F_N} \text{Out}(G_v, \text{Inc}_v^{(t)})$  is contained in the image of  $\rho$ , the lemma follows.  $\square$

## 2.10 Some properties of $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$

The group  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is the finite-index subgroup of  $\text{Out}(F_N)$  defined as the kernel of the natural homomorphism  $\text{Out}(F_N) \rightarrow \text{GL}_N(\mathbb{Z}/3\mathbb{Z})$ . In the present paper, it will often be convenient to work in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  to avoid finite-order phenomena. We now collect the various properties of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  we will use.

**Theorem 2.10** (Handel–Mosher [HM20, Theorem II.3.1]). *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and assume that there exists a free factor  $A$  of  $F_N$  whose conjugacy class is invariant under a finite index subgroup of  $H$ .*

*Then the conjugacy class of  $A$  is  $H$ -invariant.*

**Theorem 2.11** (Handel–Mosher [HM20, Theorem II.4.1]). *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and assume that there exists an element  $g$  of  $F_N$  whose conjugacy class is invariant under a finite index subgroup of  $H$ .*

*Then the conjugacy class of  $g$  is  $H$ -invariant.*

The following lemma was established for example in [HW20] as an easy consequence of the above hard theorem of Handel and Mosher.

**Lemma 2.12** (see e.g. [HW20, Lemma 2.6]). *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and let  $S$  be a free splitting of  $F_N$  which is invariant under a finite index subgroup of  $H$ . Then  $S$  is  $H$ -invariant, and  $H$  acts as the identity on  $S/F_N$ .*

We say that a vertex  $v$  in a splitting  $S$  of  $F_N$  is *Grushko* if the Grushko deformation space of its stabilizer  $G_v$  relative to the collection of all incident edge stabilizers is non-trivial.

**Lemma 2.13.** *Let  $S$  be a splitting of  $F_N$  and  $v \in S$ . Assume that  $v$  is a Grushko vertex or that there exists an edge incident on  $v$  with trivial stabilizer. Let  $\alpha \in \text{Stab}_{\text{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$ ,*

let  $\tilde{\alpha} \in \text{Aut}(F_N)$  be a representative of  $\alpha$ , and let  $I_{\tilde{\alpha}}$  be an  $\tilde{\alpha}$ -equivariant isomorphism of  $S$ .

Then  $I_{\tilde{\alpha}}(v)$  belongs to the same  $F_N$ -orbit as  $v$ .

*Proof.* First assume that there is an edge  $e$  with trivial stabilizer with origin at  $v$ . Let  $\bar{S}$  be the free splitting obtained from  $S$  by collapsing all edges with non-trivial stabilizer. Then  $\bar{S}$  is  $\alpha$ -invariant. By Lemma 2.12,  $I_{\tilde{\alpha}}$  induces the identity on  $\bar{S}/F_N$ , so  $I_{\tilde{\alpha}}$  maps  $e$  to  $ge$  for some  $g \in F_N$ . It follows that  $I_{\tilde{\alpha}}(v) = gv$ , and we are done.

We now assume that  $v$  is a Grushko vertex. Let  $\hat{S}$  be a refinement of  $S$  obtained by blowing up  $v$  into a Grushko decomposition  $S_v$  of  $G_v$  relative to the incident edge groups, and attaching back every incident edge  $e$  to the unique point of  $S_v$  fixed by  $G_e$ . We denote by  $\mathcal{F}_v$  the free factor system of  $F_N$  determined by collapsing all edges of  $\hat{S}$  with non-trivial stabilizer to a point. Notice that this free factor system only depends on  $S$  and  $v$  and not on the choice of the refinement  $\hat{S}$ : indeed,  $\mathcal{F}_v$  is equal to the smallest free factor system of  $F_N$  relative to the Grushko factors of  $(G_v, \text{Inc}_v)$  and to all other vertex groups of  $S$ .

We now observe that if  $v$  and  $v'$  are two Grushko vertices of  $S$  that belong to different  $F_N$ -orbits, then  $\mathcal{F}_v \neq \mathcal{F}_{v'}$ : this is because  $G_{v'}$  is conjugate into a subgroup of  $\mathcal{F}_v$ , while  $G_v$  is not.

The element  $\alpha$  permutes all free factor systems of the form  $\mathcal{F}_v$  where  $v$  varies among the finitely many orbits of Grushko vertices of  $S$ . As  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , we deduce using Theorem 2.10 that  $\alpha$  fixes each free factor system  $\mathcal{F}_v$ . The conclusion follows.  $\square$

As a consequence of Lemma 2.13, we get the following fact.

**Corollary 2.14.** *Let  $S$  be a splitting of  $F_N$  with all edge stabilizers non-trivial, and let  $v \in S$  be a Grushko vertex. Let  $N_v$  be the subgroup of  $\text{Stab}_{\text{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$  made of all automorphisms that act trivially on all vertex stabilizers but  $G_v$ .*

*Then  $N_v$  is normal in  $\text{Stab}_{\text{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$ .*  $\square$

**Proposition 2.15.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , and let  $S$  be a  $\mathcal{Z}_{\max}$ -splitting which is  $H$ -invariant. Then  $H$  acts trivially on  $S/F_N$ .*

*Proof.* Let  $\Gamma = S/F_N$ . Let  $K$  be the quotient of  $H_1(F_N; \mathbb{Z}/3\mathbb{Z})$  by the subgroup generated by the image of all edge groups of  $\Gamma$ . As  $S$  is a minimal  $\mathcal{Z}_{\max}$ -splitting, for each terminal vertex  $v$  of  $\Gamma$ , the group  $G_v$  is not cyclic. It follows that the image  $K_v$  of  $G_v$  in  $K$  is non-trivial. Moreover, if  $v, v'$  are distinct terminal vertices, then  $K_v \cap K_{v'}$  is trivial, so in particular  $K_v \neq K_{v'}$ . Since  $H$  acts trivially on  $K$ , it follows that  $H$  acts as the identity on the set of terminal vertices of  $\Gamma$ . This concludes the proof if  $\Gamma$  is a tree.

Otherwise, let  $\Gamma_0 \subseteq \Gamma$  be the core of  $\Gamma$ , i.e. the union of embedded loops. If  $\Gamma_0$  is not a circle, then  $H$  acts as the identity on  $\Gamma_0$  because it acts as the identity on  $H_1(\Gamma_0; \mathbb{Z}/3\mathbb{Z})$ . This implies that  $H$  acts as the identity on  $\Gamma$ . If  $\Gamma_0$  is a circle, then  $H$  acts on  $\Gamma_0$  by orientation-preserving homeomorphisms because it acts trivially on  $H_1(\Gamma_0; \mathbb{Z}/3\mathbb{Z})$ . So it suffices to prove that  $H$  fixes a point in  $\Gamma_0$ . Since  $H$  fixes all terminal points, this is clearly the case if  $\Gamma \neq \Gamma_0$  so we can assume that  $\Gamma$  itself is a circle. By [Hor17,

Proposition 6.6],  $\Gamma$  has at least one Grushko vertex, and Lemma 2.13 shows that this vertex is fixed by  $H$ , concluding the proof.  $\square$

### 3 Preliminaries on measured groupoids

We do not assume the reader to be familiar with the notion of a measured groupoid and its use in proving measure equivalence rigidity results. In this section and the next one, we therefore thoroughly review standard facts from this theory. Most of the material we need can be found in [AD07, Section 2.1] or [Kid09] and the references therein.

#### 3.1 Measured groupoids: basic definitions

##### 3.1.1 Groupoids, subgroupoids, restrictions

A *standard Borel space* is a measurable space which is isomorphic to the Borel  $\sigma$ -algebra of a separable complete metric space.

**Definition 3.1** (Discrete Borel groupoids). *A discrete Borel groupoid over a standard Borel space  $Y$  is a standard Borel space  $\mathcal{G}$  whose elements are called arrows, equipped with the following data:*

1. two Borel maps  $s, r : \mathcal{G} \rightarrow Y$  called the source map and the range map, such that for every  $y \in Y$ , the set of arrows  $g \in \mathcal{G}$  such that  $s(g) = y$  is at most countable.
2. a Borel map (called the composition law)

$$\begin{aligned} \mathcal{G}^{(2)} &\rightarrow \mathcal{G} \\ (g_1, g_2) &\mapsto g_1 g_2 \end{aligned}$$

defined on the set  $\mathcal{G}^{(2)} := \{(g_1, g_2) \in \mathcal{G} \times \mathcal{G} \mid s(g_1) = r(g_2)\}$  of composable arrows, such that for all  $(g_1, g_2), (g_2, g_3) \in \mathcal{G}^{(2)}$ , the following hold:

- **Source/Range:**  $s(g_1 g_2) = s(g_2)$  and  $r(g_1 g_2) = r(g_1)$ ,
- **Associativity:**  $(g_1 g_2) g_3 = g_1 (g_2 g_3)$ ;

3. a Borel map (called the unit map, whose image is called the space of units)

$$\begin{aligned} Y &\rightarrow \mathcal{G} \\ y &\mapsto e_y \end{aligned}$$

such that for all  $y \in Y$ , the following hold:

- **Source/Range:**  $s(e_y) = r(e_y) = y$  for all  $y \in Y$ ,
- **Neutral element:**  $g e_y = g$ ,  $e_y g' = g'$  for all arrows  $g, g' \in \mathcal{G}$  with  $s(g) = y$  and  $r(g') = y$ ;

4. a Borel map (called the inverse map)

$$\begin{aligned} \iota: \mathcal{G} &\rightarrow \mathcal{G} \\ g &\mapsto g^{-1} \end{aligned}$$

such that for all  $g \in \mathcal{G}$ , the following hold:

- **Source/Range:**  $s(g^{-1}) = r(g)$ ,  $r(g^{-1}) = s(g)$ ,
- **Inverse:**  $gg^{-1} = e_{r(g)}$  and  $g^{-1}g = e_{s(g)}$ .

The word ‘discrete’ in ‘discrete Borel groupoid’ refers to the first condition from the above definition – countability of the set of arrows having a given source. As all groupoids considered in the present paper are discrete, we will often drop the term ‘discrete’ from the terminology and simply call them Borel groupoids.

**Definition 3.2** (Subgroupoids). *Let  $\mathcal{G}$  be a Borel groupoid over a base space  $Y$ . A Borel subgroupoid of  $\mathcal{G}$  is a Borel subset  $\mathcal{H} \subseteq \mathcal{G}$  such that*

1. for every  $(g_1, g_2) \in \mathcal{G}^{(2)} \cap (\mathcal{H} \times \mathcal{H})$ , one has  $g_1g_2 \in \mathcal{H}$ ,
2. for every  $g \in \mathcal{H}$ , one has  $g^{-1} \in \mathcal{H}$ , and
3. for every  $y \in Y$ , one has  $e_y \in \mathcal{H}$ .

Every Borel subgroupoid of  $\mathcal{G}$  has a natural structure of a Borel groupoid over the same base space  $Y$ .

**Definition 3.3.** *Given a Borel groupoid  $\mathcal{G}$  and two Borel subgroupoids  $\mathcal{H}, \mathcal{H}'$  of  $\mathcal{G}$ , the subgroupoid  $\langle \mathcal{H}, \mathcal{H}' \rangle$  generated by  $\mathcal{H}$  and  $\mathcal{H}'$  is the smallest subgroupoid of  $\mathcal{G}$  that contains  $\mathcal{H}$  and  $\mathcal{H}'$ .*

More explicitly, this is the set of all arrows of the form  $h_1 \dots h_k$ , with each  $h_i$  in either  $\mathcal{H}$  or  $\mathcal{H}'$ .

The subgroupoid  $\langle \mathcal{H}, \mathcal{H}' \rangle$  is a Borel subgroupoid of  $\mathcal{G}$  as can be seen by writing  $\mathcal{H}$  and  $\mathcal{H}'$  as a countable union of bisections (see below).

**Definition 3.4** (Restrictions). *Given a Borel groupoid over a base space  $Y$ , and a Borel subset  $A \subseteq Y$ , the set*

$$\mathcal{G}|_A := \{g \in \mathcal{G} \mid s(g), r(g) \in A\}$$

has a natural structure of a Borel groupoid over the base space  $A$ , induced from the groupoid structure of  $\mathcal{G}$ . The groupoid  $\mathcal{G}|_A$  is called the restriction of  $\mathcal{G}$  to  $A$ .

*Remark 3.5.* Note that  $\mathcal{G}|_A$  is not a subgroupoid of  $\mathcal{G}$  (it is not defined on the same base space).

Every countable group has a natural structure of a Borel groupoid over a point. A more interesting – and crucial – example comes from restrictions of group actions, as explained right below (another interesting class of examples comes from equivalence relations, but we will not need those in the present paper). All countable groups will always be equipped with the discrete topology.

*Example 3.6* (Group actions). Let  $Y$  be a standard Borel space, equipped with an action of a countable group  $\Gamma$  by Borel automorphisms. Then the direct product  $\Gamma \times Y$  has a natural structure of a discrete Borel groupoid over the base space  $Y$ : one defines  $s(\gamma, y) = y$  and  $r(\gamma, y) = \gamma \cdot y$ , and the composition law is given by

$$(\gamma_1, \gamma_2 \cdot y)(\gamma_2, y) = (\gamma_1 \gamma_2, y),$$

the unit map is given by  $e_y = (1_\Gamma, y)$ , and the inverse map by  $(\gamma, y)^{-1} = (\gamma^{-1}, \gamma \cdot y)$ . This groupoid is denoted as  $\Gamma \times Y$ .

If  $A \subseteq Y$  is any Borel subset (without any invariance hypothesis), then the groupoid  $(\Gamma \times Y)|_A$  is well defined; its set of arrows is the set of pairs  $(a, \gamma) \in A \times \Gamma$  such that  $\gamma \cdot a \in A$ .

In the above example, the groupoid  $(\Gamma \times Y)|_A$  may be written as the countable union of subsets  $B_\gamma = \{\gamma\} \times (A \cap \gamma^{-1}A)$  for  $\gamma \in \Gamma$ . There is a partial isomorphism from  $s(B_\gamma) = A \cap \gamma^{-1}A$  to  $r(B_\gamma) = \gamma A \cap A$  defined by  $y \mapsto \gamma \cdot y$ . More generally, a bisection is a collection of arrows that defines a partial isomorphism in the following sense.

**Definition 3.7** (Bisections). *Let  $\mathcal{G}$  be a Borel groupoid over a base space  $Y$ . A bisection of  $\mathcal{G}$  is a Borel subset  $B \subseteq \mathcal{G}$  such that  $s|_B$  and  $r|_B$  are Borel isomorphisms to Borel subsets  $s(B), r(B) \subseteq Y$ .*

*The partial isomorphism associated to a bisection  $B$  is the Borel map  $f_B : s(B) \rightarrow r(B)$  defined by  $f_B = r \circ s|_B^{-1}$ .*

A theorem by Lusin and Novikov (see [Kec95, Theorem 18.10]) shows that any discrete Borel groupoid  $\mathcal{G}$  is always a countable union of bisections (this is obvious in the case of a restriction of a group action, but for the general case, it is crucial that Borel spaces are standard).

**Definition 3.8** (Saturation). *Let  $\mathcal{G}$  be a Borel groupoid over a base space  $Y$ , and let  $y_0 \in Y$ . The  $\mathcal{G}$ -orbit of  $y_0$  is the set of all  $y \in Y$  such that there exists  $g \in \mathcal{G}$  with  $s(g) = y_0$  and  $r(g) = y$ . The saturation  $\mathcal{G}A$  of a Borel subset  $A \subseteq Y$  is the set of points that are in the orbit of some point in  $A$ , in other words  $\mathcal{G}A = r(s^{-1}(A))$ .*

If  $A \subseteq Y$  is a Borel subset, then so is  $\mathcal{G}A$ : indeed, writing  $\mathcal{G}$  as a countable union of bisections  $B_n$ , one has  $\mathcal{G}A = \cup_n f_{B_n}(A)$ .

### 3.1.2 Measured groupoids

Let  $\mathcal{G}$  be a Borel groupoid over a base space  $Y$ . A measure  $\mu$  on  $Y$  is  $\mathcal{G}$ -invariant if for every bisection  $B \subseteq \mathcal{G}$ , one has  $\mu(s(B)) = \mu(r(B))$ . It is  $\mathcal{G}$ -quasi-invariant if for every bisection  $B \subseteq \mathcal{G}$ , one has  $\mu(s(B)) = 0$  if and only if  $\mu(r(B)) = 0$ . Note that in the case of an action of a countable group  $\Gamma$  by Borel automorphisms on  $Y$ , the group  $\Gamma$  preserves  $\mu$  if and only if the groupoid  $\Gamma \times Y$  does.

A standard measure space  $(Y, \mu)$  is a standard Borel space  $Y$  equipped with a  $\sigma$ -finite positive Borel measure  $\mu$ .



**Definition 3.9** (Measured groupoid). *Let  $(Y, \mu)$  be a standard measure space. A measured groupoid over  $(Y, \mu)$  is a discrete Borel groupoid  $\mathcal{G}$  over  $Y$  such that  $\mu$  is  $\mathcal{G}$ -quasi-invariant.*

*Remark 3.10.* If we change the measure  $\mu$  into another measure  $\mu'$  in the same measure class, then  $\mathcal{G}$  is still a measured groupoid over  $(Y, \mu')$ , and this does not change  $\mathcal{G}$  up to isomorphism (see Definition 3.11 below). Thus, we can always assume without loss of generality that  $\mu$  is a probability measure.

When  $\mathcal{G}$  is a measured groupoid over a base space  $(Y, \mu)$ , and  $\mathcal{H}$  is a Borel subgroupoid of  $\mathcal{G}$ , then  $\mu$  is  $\mathcal{H}$ -quasi-invariant. The Borel subgroupoid  $\mathcal{H}$  together with the measure  $\mu$  is called a *measured subgroupoid* of  $\mathcal{G}$ .

From a measure  $\mu$  on the base space  $Y$ , one can define a measure  $\nu$  on the set of arrows  $\mathcal{G}$  by

$$\nu(A) = \int_Y \sum_{g \in s^{-1}(y)} \mathbb{1}_A(g) d\mu(y)$$

for any Borel subset  $A \subseteq \mathcal{G}$ . Then  $\mu$  is preserved (resp. quasi-preserved) by  $\mathcal{G}$  if and only if  $\nu$  is preserved by the inversion map:  $\iota_*\nu = \nu$  (resp.  $\nu$  and  $\iota_*\nu$  have the same null sets).

### 3.1.3 Cocycles

**Definition 3.11** (Groupoid homomorphisms and isomorphisms). *Let  $\mathcal{G}_1, \mathcal{G}_2$  be measured groupoids over standard measure spaces  $(Y_1, \mu_1)$  and  $(Y_2, \mu_2)$ , respectively. A homomorphism of measured groupoids from  $\mathcal{G}_1$  to  $\mathcal{G}_2$  is a Borel map  $\varphi : \mathcal{G}_1 \rightarrow \mathcal{G}_2$  such that there exists a conull Borel subset  $Y_1^* \subseteq Y_1$  on which the following hold:*

1. for all  $(g, h) \in ((\mathcal{G}_1)_{|Y_1^*})^{(2)}$ , one has  $\varphi(gh) = \varphi(g)\varphi(h)$ , and
2. there exists a Borel map  $\varphi_Y : Y_1^* \rightarrow Y_2$  such that
  - (a) for every  $y \in Y_1^*$ , one has  $e_{\varphi_Y(y)} = \varphi(e_y)$ , and
  - (b) the measures  $(\varphi_Y)_*((\mu_1)_{|Y_1^*})$  and  $\mu_2$  have the same null sets.

*It is an isomorphism if in addition, there exist a groupoid homomorphism  $\psi : \mathcal{G}_2 \rightarrow \mathcal{G}_1$  and conull Borel subsets  $Y_1^* \subseteq Y_1$  and  $Y_2^* \subseteq Y_2$  such that  $\psi \circ \varphi$  and  $\varphi \circ \psi$  are the identity maps on  $(\mathcal{G}_1)_{|Y_1^*}$  and  $(\mathcal{G}_2)_{|Y_2^*}$ , respectively.*

**Definition 3.12** (Cocycles). *Let  $\mathcal{G}$  be a measured groupoid and  $\Gamma$  be a measurable group. A cocycle  $\rho : \mathcal{G} \rightarrow \Gamma$  is a Borel map such that  $\rho(gh) = \rho(g)\rho(h)$  for all  $(g, h) \in \mathcal{G}^{(2)}$ .*

*Remark 3.13.* Usually, in the definition of a cocycle one only asks that  $\rho(gh) = \rho(g)\rho(h)$  for almost every (and not all)  $(g, h) \in \mathcal{G}^{(2)}$ . Cocycles satisfying Definition 3.12 are usually called *strict*. When  $\Gamma$  is countable, for any non-strict cocycle, there exists a conull subset  $Y^* \subseteq Y$  such that  $\rho|_{Y^*}$  is strict. More generally, when  $\Gamma$  is second countable, then any non-strict cocycle coincides almost everywhere with a strict cocycle ([Zim84, Theorem B.9]). All the cocycles we consider in this paper take values in second countable groups (and in fact mostly in countable discrete groups). For these reasons, all the cocycles we consider are strict as in Definition 3.12.

**Definition 3.14.** *The kernel of a cocycle  $\rho$  is the set of all arrows  $g \in \mathcal{G}$  such that  $\rho(g) = 1_\Gamma$ . We say that  $\rho$  has trivial kernel if its kernel is reduced to the set of unit elements of  $\mathcal{G}$ .*

Notice that the kernel of a cocycle  $\mathcal{G} \rightarrow \Gamma$  is always a Borel subgroupoid of  $\mathcal{G}$ . More generally, if  $\rho : \mathcal{G} \rightarrow \Gamma$  is a cocycle and  $H \subseteq \Gamma$  is a subgroup, then  $\rho^{-1}(H)$  is a subgroupoid of  $\mathcal{G}$ .

*Example 3.15* (Natural cocycle associated to a group action). Let  $\mathcal{G} := \Gamma \ltimes Y$  be a groupoid associated to a quasi-measure-preserving action of a countable group  $\Gamma$  by Borel automorphisms on a standard Borel space  $Y$  equipped with a  $\sigma$ -finite Borel measure  $\mu$ . Let  $A \subseteq Y$  be a Borel subset of positive measure. The *natural cocycle* associated to the restriction to  $A$  of the  $\Gamma$ -action on  $Y$  is the cocycle

$$\begin{aligned} \rho : (\Gamma \ltimes Y)|_A &\rightarrow \Gamma \\ (\gamma, y) &\mapsto \gamma \end{aligned}$$

Notice that  $\rho$  is indeed a cocycle and has trivial kernel.

The following lemma shows that conversely, every measured groupoid which admits a cocycle with trivial kernel towards a countable group  $\Gamma$ , is a restriction of a groupoid coming from a measure-preserving action of  $\Gamma$ .

**Lemma 3.16.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $(Y, \mu)$ , let  $\Gamma$  be a countable group, and assume that there exists a cocycle  $\rho : \mathcal{G} \rightarrow \Gamma$  with trivial kernel.*

*Then there exist a standard Borel space  $S$  with a  $\sigma$ -finite Borel measure  $\nu$  and a quasi-measure-preserving  $\Gamma$ -action on  $S$  by Borel automorphisms such that  $\mathcal{G}$  is isomorphic to a restriction of  $\Gamma \ltimes S$ , and  $\rho$  is the restriction of the natural cocycle  $\Gamma \ltimes S \rightarrow \Gamma$ .*

*Remark 3.17.* If the measure  $\mu$  is  $\mathcal{G}$ -invariant, the constructed measure  $\nu$  is  $\Gamma \ltimes S$ -invariant but infinite in general even if  $\mu$  is a probability measure.

*Proof.* This is an adaptation of an argument given by Kida in his proof of [Kid08a, Proposition 4.33]. Let  $\eta$  be the counting measure on  $\Gamma$ , and let  $\tilde{\mu} := \mu \otimes \eta$  be the product measure on  $Y \times \Gamma$ . Let  $\tilde{\mathcal{G}} := \mathcal{G} \times \Gamma$ . The space  $\tilde{\mathcal{G}}$  has a natural structure of a Borel groupoid over  $Y \times \Gamma$  with  $s(g, \gamma) = (s(g), \gamma)$  and  $r(g, \gamma) = (r(g), \rho(g)\gamma)$  for all  $(g, \gamma) \in \tilde{\mathcal{G}}$ , and one checks that  $\tilde{\mathcal{G}}$  quasi-preserves the measure  $\tilde{\mu}$ .

Let  $\sim$  be the  $\tilde{\mathcal{G}}$ -orbit equivalence relation on  $Y \times \Gamma$ , i.e. two points  $(y_1, \gamma_1)$  and  $(y_2, \gamma_2)$  are equivalent whenever there exists  $h \in \tilde{\mathcal{G}}$  such that  $s(h) = (y_1, \gamma_1)$  and  $r(h) = (y_2, \gamma_2)$ . We claim that the equivalence relation  $\sim$  has a fundamental domain  $D$ . Indeed, triviality of the kernel of  $\rho$  shows that for every  $\gamma \in \Gamma$ , the equivalence relation  $\sim|_{Y \times \{\gamma\}}$  is trivial. Now choose an enumeration  $\{\gamma_i\}_{i \in \mathbb{N}}$  of  $\Gamma$ , and define inductively  $D_0 = Y \times \{\gamma_0\}$ , and  $D_{i+1} = (Y \times \{\gamma_i\}) \setminus \tilde{\mathcal{G}}D_i$  where  $\tilde{\mathcal{G}}D_i$  is the saturation of  $D_i$ . Then  $D = \cup_i D_i$  is a Borel fundamental domain for  $\sim$ .

Consider the space  $S := (Y \times \Gamma)/\sim$ . Then  $S$  is naturally isomorphic to  $D$  (in particular  $S$  is a standard Borel space), and we define  $\nu$  on  $S$  as  $\tilde{\mu}|_D$  through this identification. Given  $(y, \gamma) \in Y \times \Gamma$ , we denote by  $[y, \gamma]$  the class of  $(y, \gamma)$  in  $S$ . The

group  $\Gamma$  acts on  $S$  via  $\gamma \cdot [y', \gamma'] := [y', \gamma' \gamma^{-1}]$  in a quasi-measure-preserving way. Then  $\mathcal{G}$  is isomorphic to  $(\Gamma \ltimes S)_{|Y \times \{e\}}$  via  $g \mapsto (\rho(g), [s(g), e])$ , see [Kid08a, Lemmas 4.34 and 4.35].  $\square$

The following definition will be used at various places in the paper.

**Definition 3.18** (Nowhere trivial, stably trivial cocycle). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$  and let  $\Gamma$  be a countable group. A cocycle  $\rho : \mathcal{G} \rightarrow \Gamma$  is*

- nowhere trivial if there does not exist any Borel subset  $U \subseteq Y$  of positive measure such that  $\rho(\mathcal{G}|_U) = \{1\}$ ;
- stably trivial if there exists a Borel partition  $Y^* = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets such that for every  $i \in I$ , one has  $\rho(\mathcal{G}|_{Y_i}) = \{1\}$ .

### 3.1.4 Groupoids of infinite type and action-type cocycles

We adopt the following definition of a measured groupoid of infinite type, which is equivalent to Kida's definition [Kid10, Definition 3.1] in the case where  $\mu$  is a finite  $\mathcal{G}$ -invariant measure, see the discussion above [Kid09, Definition 2.25]. It is a notion of recurrence for a measured groupoid.

**Definition 3.19** (Groupoids of infinite type). *A measured groupoid  $\mathcal{G}$  over a base space  $(Y, \mu)$  is of infinite type if for every Borel subset  $U \subseteq Y$  of positive measure, and  $\mu$ -a.e.  $y \in U$ , the set*

$$\{g \in \mathcal{G}|_U \mid s(g) = y\}$$

*is infinite.*

Notice that if  $\mathcal{G}$  is of infinite type, then so is  $\mathcal{G}|_U$  for every Borel subset  $U \subseteq Y$  of positive measure. A crucial example comes from Poincaré recurrence: if an infinite group acts by preserving a probability measure on  $Y$ , then the associated groupoid is of infinite type (here the fact that the measure is finite and invariant is essential). The following definition reformulates this property in terms of the associated natural cocycle. It will be of central importance in the present work.

**Definition 3.20** (Action-type cocycles). *Let  $\mathcal{G}$  be a measured groupoid, let  $\Gamma$  be a countable group. A cocycle  $\rho : \mathcal{G} \rightarrow \Gamma$  is action-type if it has trivial kernel, and for every infinite subgroup  $H \subseteq \Gamma$ , the subgroupoid  $\rho^{-1}(H)$  is of infinite type.*

**Lemma 3.21** (See for instance [Kid09, Proposition 2.26]). *Let  $\Gamma$  be a countable group, let  $Y$  be a standard Borel space equipped with a  $\Gamma$ -action by Borel automorphisms which preserves a Borel probability measure  $\mu$ . Let  $\mathcal{G} = \Gamma \ltimes (Y, \mu)$ , and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be the natural cocycle.*

*Then  $\rho$  is action-type.*

We will systematically use the following obvious facts without mention.

**Lemma 3.22.** *Let  $\mathcal{G}$  be a measured groupoid over a standard measure space  $Y$ , let  $\Gamma$  be a countable group, and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be an action-type cocycle.*

1. *For every Borel subset  $U \subseteq Y$  of positive measure, the restricted cocycle  $\rho : \mathcal{G}|_U \rightarrow \Gamma$  is action-type.*
2. *For every subgroup  $\Gamma' \subseteq \Gamma$ , the induced cocycle  $\rho^{-1}(\Gamma') \rightarrow \Gamma'$  is action-type.  $\square$*

*Remark 3.23.* We will often use the following consequence of the action-type property. Let  $\rho : \mathcal{G} \rightarrow \Gamma$  be action-type,  $H \subseteq \Gamma$  be a subgroup, and  $\mathcal{H} = \rho^{-1}(H)$ . Then for every element  $h \in H$  of infinite order and every Borel subset  $U \subseteq Y$  of positive measure,  $\rho(\mathcal{H}|_U)$  contains a power of  $h$ .

### 3.1.5 Stable equality of measured groupoids

In the present paper, we will often consider measured groupoids up to stable equality, defined as follows.

**Definition 3.24** (Stable containment, stable equality). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\mathcal{H}$  and  $\mathcal{H}'$  be two measured subgroupoids.*

*The groupoid  $\mathcal{H}$  is stably equal to  $\mathcal{H}'$  (resp. stably contained in  $\mathcal{H}'$ ) if there exist a conull subset  $Y^* \subseteq Y$  and a countable Borel partition  $Y^* = \bigsqcup_{n \in \mathbb{N}} Y_n$  such that for every  $n \in \mathbb{N}$ , one has  $\mathcal{H}|_{Y_n} = \mathcal{H}'|_{Y_n}$  (resp.  $\mathcal{H}|_{Y_n} \subseteq \mathcal{H}'|_{Y_n}$ ).*

*Remark 3.25.* In the present paper, we will often look for properties  $(P)$  of measured groupoids that are preserved by restriction and stabilization:

1.  $(P)$  is preserved by restriction if, for every measured groupoid  $\mathcal{G}$  over a base space  $(Y, \mu)$ , if  $\mathcal{G}$  satisfies  $(P)$ , then so does its restriction  $\mathcal{G}|_A$  for every Borel subset  $A \subseteq Y$  of positive measure.
2.  $(P)$  is preserved by stabilization if, for every measured groupoid  $\mathcal{G}$  over a base space  $(Y, \mu)$ , and for every countable Borel partition  $Y^* = \bigsqcup_{n \in \mathbb{N}} Y_n$  of a conull Borel subset  $Y^* \subseteq Y$ , if the groupoid  $\mathcal{G}|_{Y_n}$  satisfies  $(P)$  for every  $n \in \mathbb{N}$ , then so does  $\mathcal{G}$ .

### 3.1.6 Invariant objects, equivariant maps

The following definition is a generalization to the groupoid setting of the notion of fixed points of a group action.

**Definition 3.26** ((Stably) equivariant maps). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $(Y, \mu)$ , let  $\Gamma$  be a countable group, and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be a cocycle. Let  $\Delta$  be a measurable space equipped with a measurable  $\Gamma$ -action, and let  $\varphi : Y \rightarrow \Delta$  be a measurable map.*

*An element  $x \in \Delta$  is  $(\mathcal{G}, \rho)$ -invariant if there exists a conull Borel subset  $Y^* \subseteq Y$  such that for all  $g \in \mathcal{G}|_{Y^*}$ , one has  $\rho(g)x = x$ .*

We say that  $\varphi$  is  $(\mathcal{G}, \rho)$ -equivariant if there exists a conull Borel subset  $Y^* \subseteq Y$  such that for all  $g \in \mathcal{G}|_{Y^*}$ , one has  $\varphi(r(g)) = \rho(g)\varphi(s(g))$ .

We say that  $\varphi$  is stably  $(\mathcal{G}, \rho)$ -equivariant if there exist a conull Borel subset  $Y^* \subseteq Y$  and a countable Borel partition  $Y^* = \bigsqcup_{n \in \mathbb{N}} Y_n$  such that for every  $n \in \mathbb{N}$ , the map  $\varphi|_{Y_n}$  is  $(\mathcal{G}|_{Y_n}, \rho)$ -equivariant.

*Example 3.27.* Let  $\mathcal{G} := \Gamma \ltimes Y$  be a groupoid associated to a measure-preserving action of a countable group  $\Gamma$  on some standard measure space  $Y$ , let  $\rho : \mathcal{G} \rightarrow \Gamma$  be the natural cocycle. Let  $\Delta$  be a measurable space equipped with a  $\Gamma$ -action by measurable automorphisms. Then a measurable map  $Y \rightarrow \Delta$  is  $(\mathcal{G}, \rho)$ -equivariant if and only if it is almost everywhere  $\Gamma$ -equivariant.

*Remark 3.28.* Although stably invariant maps are not always invariant, the following is true: if there exists a stably  $(\mathcal{G}, \rho)$ -invariant measurable map  $Y \rightarrow \Delta$ , then there exists another map  $Y \rightarrow \Delta$  which is  $(\mathcal{G}, \rho)$ -invariant. The key point is that any map whose restriction to a Borel subset  $A \subseteq Y$  of positive measure is  $(\mathcal{G}|_A, \rho)$ -equivariant extends uniquely to an equivariant map on the saturation  $\mathcal{G}A$  (see [Ada94, Lemma 2.7] or [Kid08a, Lemma 4.16]).

**Definition 3.29** (Stabilizers). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ . Let  $\Gamma$  be a countable group, and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be a cocycle. Let  $\Delta$  be a measurable space equipped with a measurable  $\Gamma$ -action, and let  $\varphi : Y \rightarrow \Delta$  be a measurable map.*

*The  $(\mathcal{G}, \rho)$ -stabilizer of  $\varphi$  is the subgroupoid of  $\mathcal{G}$  made of all  $g \in \mathcal{G}$  such that  $\varphi(r(g)) = \rho(g)\varphi(s(g))$ .*

### 3.2 Normal subgroupoids

A notion of normal subgroupoid of a measured groupoid was coined by Kida in [Kid08a, Section 4.6.1], building on earlier work by Feldman, Sutherland and Zimmer [FSZ89]. One can check that the definition of normality given below is equivalent to Kida's, but we will not use this fact.

Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ , and let  $B \subseteq \mathcal{G}$  be a Borel subset (typically,  $B$  may be a bisection, see Definition 3.7, but it need not be). We say that  $\mathcal{H}$  is  $B$ -invariant if there exists a conull Borel subset  $Y^* \subseteq Y$  such that for all  $g_1, g_2 \in B \cap \mathcal{G}|_{Y^*}$  and all  $h \in \mathcal{G}|_{Y^*}$  such that  $g_2, h, g_1^{-1}$  are composable, one has  $h \in \mathcal{H}$  if and only if  $g_2 h g_1^{-1} \in \mathcal{H}$ .

**Definition 3.30** ((Stable) normalization). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\mathcal{H}$  and  $\mathcal{H}'$  be measured subgroupoids of  $\mathcal{G}$ .*

*The groupoid  $\mathcal{H}'$  normalizes  $\mathcal{H}$  if there exists a countable collection of Borel subsets  $B_n \subseteq \mathcal{G}$  such that  $\mathcal{H}' = \bigcup_{n \in \mathbb{N}} B_n$ , and for every  $n \in \mathbb{N}$ , the subgroupoid  $\mathcal{H}$  is  $B_n$ -invariant.*

*The groupoid  $\mathcal{H}'$  stably normalizes  $\mathcal{H}$  if there exists a countable Borel partition  $Y^* = \bigsqcup_{n \in \mathbb{N}} Y_n$  of a conull Borel subset  $Y^* \subseteq Y$  such that for every  $n \in \mathbb{N}$ , the restriction  $\mathcal{H}'|_{Y_n}$  normalizes  $\mathcal{H}|_{Y_n}$ .*

*Remark 3.31.* Note that  $B_n$  is not assumed to be a subgroupoid. Since  $\mathcal{G}$  is a countable union of bisections, one may assume without loss of generality that every  $B_n$  is a bisection of  $\mathcal{G}$ .

When  $\mathcal{H} \subseteq \mathcal{H}'$ , we say that  $\mathcal{H}$  is *normal* in  $\mathcal{H}'$ , denoted by  $\mathcal{H} \trianglelefteq \mathcal{H}'$ . If  $\mathcal{G} = \Gamma \times Y$  is the groupoid associated to the action of a countable group  $\Gamma$  on a standard measure space  $Y$ , and if  $H \trianglelefteq \Gamma$  is a normal subgroup, then the subgroupoid  $H \times Y$  is normal in  $\mathcal{G}$ . Indeed, this is shown by letting  $B_\gamma := \{(\gamma, y) | y \in Y\}$ : then the Borel subsets  $B_\gamma$  are bisections that cover  $\mathcal{G}$ , and normality of  $H$  in  $\Gamma$  implies that  $H \times Y$  is  $B_\gamma$ -invariant for every  $\gamma \in \Gamma$ .

If  $Y^* \subseteq Y$  is a conull Borel subset, then  $\mathcal{H}'$  normalizes  $\mathcal{H}$  if and only if  $\mathcal{H}'|_{Y^*}$  normalizes  $\mathcal{H}|_{Y^*}$ . The notion of stable normalization is clearly preserved under restriction and stabilization. Notice also that the preimage of a normal subgroupoid by a groupoid homomorphism is again a normal groupoid. In particular, we record the following statement for future use.

**Lemma 3.32.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $(Y, \mu)$ , let  $\Gamma$  be a countable group, and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be a cocycle. Let  $H_1, H_2$  be two subgroups of  $\Gamma$  with  $H_1 \trianglelefteq H_2$ .*

*Then  $\rho^{-1}(H_1) \trianglelefteq \rho^{-1}(H_2)$ . □*

### 3.3 Amenability

#### 3.3.1 Amenability of a measured groupoid

We review the definition of the amenability of a measured groupoid, which is a generalization of Zimmer's definition [Zim78] of an amenable group action, see e.g. [ADR00]. Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ . Let  $B$  be a separable real Banach space. We denote by  $\text{Isom}(B)$  the group of all linear isometries of  $B$ , equipped with the strong operator topology. We denote by  $B_1^*$  the unit ball of the dual Banach space  $B^*$ , equipped with the weak-\* topology. Given a linear isometry  $T$  of  $B$ , we denote by  $T^*$  the restriction to  $B_1^*$  of the adjoint operator of  $T$ . The *adjoint cocycle*  $\rho^*$  of a cocycle  $\rho : \mathcal{G} \rightarrow \text{Isom}(B)$  is defined as  $\rho^*(g) := (\rho(g)^{-1})^*$ . We denote by  $\text{Conv}$  the set of all nonempty convex compact subsets of  $B_1^*$ . A map  $K : Y \rightarrow \text{Conv}$  (which we will refer to as a *convex field over  $Y$* ) is *measurable* if

$$\{(y, k) \in Y \times B_1^* | k \in K(y)\}$$

is a Borel subset of  $Y \times B_1^*$ . A *section* of a measurable convex field  $K$  over  $Y$  is a Borel map  $s : Y \rightarrow B_1^*$  such that for a.e.  $y \in Y$ , one has  $s(y) \in K(y)$ .

**Definition 3.33** (Amenable groupoids). *A measured groupoid  $\mathcal{G}$  over a base space  $(Y, \mu)$  is amenable if for every separable real Banach space  $B$  and every cocycle  $\rho : \mathcal{G} \rightarrow \text{Isom}(B)$ , every stably  $(\mathcal{G}, \rho^*)$ -invariant measurable convex field over  $Y$  has a stably  $(\mathcal{G}, \rho^*)$ -invariant section.*

*Remark 3.34.* The definition of an amenable groupoid is usually given in terms of invariant convex fields and invariant sections instead of stably invariant ones. However, the above definition is equivalent to the usual one in view of Remark 3.28.

Every restriction and every subgroupoid of an amenable measured groupoid is amenable, see e.g. [Kid09, Theorem 4.16] whose proof relies on work of Connes, Feldman and Weiss [CFW81]. In addition, it is clear from the definition that amenability is preserved by stabilization, i.e. if  $Y^* = \bigsqcup_{n \in \mathbb{N}} Y_n$  for some conull Borel subset  $Y^* \subseteq Y$ , and if for every  $n \in \mathbb{N}$ , the groupoid  $\mathcal{G}_{|Y_n}$  is amenable, then  $\mathcal{G}$  is amenable.

**Definition 3.35.** *A measured groupoid  $\mathcal{G}$  over a base space  $(Y, \mu)$  is everywhere non-amenable if for every Borel subset  $U \subseteq Y$  of positive measure,  $\mathcal{G}_{|U}$  is not amenable.*

### 3.3.2 Amenable groupoids from Borel amenable actions

For a groupoid associated to an action of a countable group with a quasi-invariant measure, the notion of amenability coincides with the Zimmer amenability of the group action [Zim78]. We will need the notion of Borel amenability of a Borel group action that applies without reference to a quasi-invariant measure.

We denote by  $\text{Prob}(\Gamma)$  the space of all probability measures on the countable group  $\Gamma$ , equipped with the topology of pointwise convergence.

**Definition 3.36** (Borel amenability of a group action [Ren15]). *Let  $\Gamma$  be a countable group, and let  $\Delta$  be a standard Borel space equipped with a  $\Gamma$ -action by Borel automorphisms. The  $\Gamma$ -action on  $\Delta$  is Borel amenable if there exists a sequence of Borel maps  $\nu_n : \Delta \rightarrow \text{Prob}(\Gamma)$  such that for every  $\delta \in \Delta$ , the norm  $\|\nu_n(\gamma \cdot \delta) - \gamma_* \nu_n(\delta)\|_1$  converges to 0 as  $n$  goes to  $+\infty$ .*

*Remark 3.37.* In [Ren15, Definition 2.1], Renault gives the definition in the more general context of Borel amenable groupoids. For equivalence relations, this corresponds to 1-amenability in [JKL02, Definition 2.12].

The following proposition gives a sufficient criterion to ensure the amenability of a measured groupoid.

**Proposition 3.38** (Kida [Kid08a, Proposition 4.33]). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $(Y, \mu)$ , let  $\Gamma$  be a countable group, and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be a cocycle with trivial kernel. Let  $\Delta$  be a standard Borel space equipped with a Borel amenable  $\Gamma$ -action. Assume that there exists a stably  $(\mathcal{G}, \rho)$ -invariant Borel map  $Y \rightarrow \Delta$ .*

*Then  $\mathcal{G}$  is amenable.*

*Proof.* The proof is due to Kida [Kid08a, Proposition 4.33], but as he phrased it in the specific context of mapping class groups, we include an argument for completeness of the exposition.

Since amenability is preserved by stabilization, we may assume without loss of generality that there exists a  $(\mathcal{G}, \rho)$ -invariant Borel map  $f : Y \rightarrow \Delta$ . By Lemma 3.16, there exist a standard measure space  $(\hat{Y}, \hat{\mu})$  containing  $(Y, \mu)$  and a measure-class preserving  $\Gamma$ -action on  $\hat{Y}$ , such that  $\mathcal{G}$  identifies with the restriction  $(\Gamma \times \hat{Y})_{|Y}$  in such a way that  $\rho$  is the restriction of the natural cocycle  $\hat{\rho} : (\Gamma \times \hat{Y}) \rightarrow \Gamma$ .

We can assume that  $\hat{Y}$  is the saturation of  $Y$  for the groupoid  $\Gamma \times \hat{Y}$ . Let  $\nu$  be a probability measure on  $\hat{Y}$  in the same measure class as  $\hat{\mu}$ . Using Remark 3.28, the map



$f : Y \rightarrow \Delta$  can be extended to a  $(\Gamma \times \hat{Y}, \hat{\rho})$ -equivariant Borel map  $\hat{Y} \rightarrow \Delta$ . In other words, there exists a  $\Gamma$ -invariant conull subset  $\hat{Y}^* \subseteq \hat{Y}$  and  $\Gamma$ -equivariant Borel map  $\hat{f} : \hat{Y}^* \rightarrow \Delta$ . Since the  $\Gamma$ -action on  $\Delta$  is Borel amenable, the  $\Gamma$ -action on  $(\Delta, \hat{f}_* \nu)$  is Zimmer amenable [GHL20, Proposition 2.5]. It then follows from [AEG94, Corollary C] that the  $\Gamma$ -action on  $(\hat{Y}, \hat{\mu})$  is Zimmer amenable. In other words the groupoid  $\Gamma \times \hat{Y}$  is amenable, and so is  $\mathcal{G}$ .  $\square$

As a special case of Proposition 3.38, we mention the following fact.

**Corollary 3.39.** *Let  $\mathcal{G}$  be a measured groupoid, let  $\Gamma$  be an amenable countable group. Assume that there exists a cocycle  $\mathcal{G} \rightarrow \Gamma$  with trivial kernel.*

*Then  $\mathcal{G}$  is amenable.*  $\square$

On the other hand, when the cocycle is action-type, we have the following.

**Lemma 3.40** (Kida [Kid10, Lemma 3.20]). *Let  $\mathcal{G}$  be a measured groupoid, let  $\Gamma$  be a countable group which contains a non-abelian free group. Assume that there exists an action-type cocycle  $\mathcal{G} \rightarrow \Gamma$ .*

*Then  $\mathcal{G}$  is everywhere non-amenable (see Definition 3.35).*

### 3.3.3 Amenability and invariant probability measures

One way in which amenability is often used is through the following proposition which is essentially due to Zimmer [Zim84, Proposition 4.3.9], see e.g. [Kid09, Proposition 4.14] for a proof in the language of measured groupoids. Given a compact, metrizable space  $K$ , we denote by  $\text{Prob}(K)$  the space of all probability measures on  $K$ : this is a convex closed subset (in the weak- $*$  topology) of the unit ball of the dual of  $C(K)$ , the space of continuous  $\mathbb{C}$ -valued functions on  $K$  equipped with the sup norm.

**Proposition 3.41.** (Zimmer [Zim84, Proposition 4.3.9]) *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ . Let  $\Gamma$  be a countable group, and let  $K$  be a compact metrizable space on which  $\Gamma$  acts continuously. Let  $\rho : \mathcal{G} \rightarrow \Gamma$  be a cocycle. Assume that  $\mathcal{G}$  is amenable.*

*Then there exists a  $(\mathcal{G}, \rho)$ -equivariant Borel map  $Y \rightarrow \text{Prob}(K)$ .*

## 4 Groupoids, measure equivalence, and applications

In this section, we review how proving that a countable discrete group  $\Gamma$  is rigid for measure equivalence can be reduced to a statement about measured groupoids with cocycles towards  $\Gamma$ . The criterion we present follows from the work of Kida [Kid10, Kid08b], based on earlier works of Furman [Fur99a, Fur99b] and Monod and Shalom [MS06], see also [BFS13].

Given a measured groupoid  $\mathcal{G}$  over a base space  $Y$  and a countable group  $\Gamma$ , two cocycles  $\rho, \rho' : \mathcal{G} \rightarrow \Gamma$  are  $\Gamma$ -cohomologous if there exist a conull Borel subset  $Y^* \subseteq Y$  and a Borel map  $\varphi : Y \rightarrow \Gamma$  such that for every  $g \in \mathcal{G}_{Y^*}$ , one has

$$\rho'(g) = \varphi(r(g))\rho(g)\varphi(s(g))^{-1}.$$

We consider the following rigidity property of a countable group  $\Gamma$ .

**Definition 4.1.** *A countable group  $\Gamma$  is rigid with respect to action-type cocycles if there exists a finite index subgroup  $\Gamma^0 \subseteq \Gamma$  such that for every measured groupoid  $\mathcal{G}$ , any two action-type cocycles  $\rho, \rho' : \mathcal{G} \rightarrow \Gamma^0$  are  $\Gamma$ -cohomologous.*

As noted in Remark 1.18 in the introduction, if  $\Gamma$  is rigid with respect to action-type cocycles, applying this property for groupoids over a point yields commensurator rigidity in the following sense: the natural map from  $\Gamma$  to its abstract commensurator is surjective. It is injective if and only if  $\Gamma$  is *ICC* (standing for *infinite conjugacy classes*), i.e. every nontrivial conjugacy class of  $\Gamma$  is infinite.

We will now explain why checking rigidity with respect to action-type cocycles for  $\text{Out}(F_N)$  is enough to deduce all the results announced in the introduction.

#### 4.1 Every self coupling of $\Gamma$ factors through the standard one

Given a free action of a group  $G$  by Borel automorphisms on a standard Borel space  $\Sigma$ , a *Borel fundamental domain* for the  $G$ -action of  $\Sigma$  is a Borel subset  $X \subseteq \Sigma$  such that  $\Sigma = \bigcup_{g \in G} gX$  and  $gX \cap hX = \emptyset$  for any two distinct  $g, h \in G$ . Recall from the introduction that two countable groups are *measure equivalent* if they admit a coupling in the following sense.

**Definition 4.2.** *Let  $\Gamma$  and  $\Lambda$  be two countable groups. A coupling between  $\Gamma$  and  $\Lambda$  is a standard measure space  $\Sigma$  equipped with an action of  $\Gamma \times \Lambda$  by measure-preserving Borel automorphisms, such that both the  $\Gamma$ -action and the  $\Lambda$ -action on  $\Sigma$  are free and have a Borel fundamental domain of finite measure.*

A *self coupling* of a countable group  $\Gamma$  is a coupling between  $\Gamma$  and  $\Gamma$ . If  $\Gamma^1 \subseteq \Gamma$  is a finite index subgroup, then there is a *standard self coupling* of  $\Gamma^1$  with respect to  $\Gamma$ , given by  $\Sigma = \Gamma$ , with  $\Gamma^1 \times \Gamma^1$  acting on  $\Gamma$  by

$$(g, g') \cdot h = ghg'^{-1}.$$

**Proposition 4.3** (Furman [Fur99b], Kida [Kid10]). *Let  $\Gamma$  be a countable ICC group which is rigid with respect to action-type cocycles.*

*Then for every finite-index subgroup  $\Gamma^1 \subseteq \Gamma$ , every self coupling  $\Sigma$  of  $\Gamma^1$  factors through the standard self coupling of  $\Gamma^1$  with respect to  $\Gamma$ , in the following sense: there exists a Borel map  $\Phi : \Sigma \rightarrow \Gamma$  which is almost everywhere  $(\Gamma^1 \times \Gamma^1)$ -equivariant.*

*Remark 4.4.* For an ICC group, the conclusion of Proposition 4.3 coincides with the notion of coupling rigidity in [Kid11] (for the pair  $(\Gamma, \pi)$  where  $\pi : \Gamma_1 \rightarrow \Gamma$  is the inclusion). This also agrees with the notion of tautness in [BFS13].

*Proof.* By definition of rigidity with respect to action-type cocycles (Definition 4.1), there exists a finite-index subgroup  $\Gamma^0$  of  $\Gamma$  such that for every measured groupoid  $\mathcal{G}$ , any two action-type cocycles  $\mathcal{G} \rightarrow \Gamma^0$  are  $\Gamma$ -cohomologous. Let  $\Gamma^2 := \Gamma^0 \cap \Gamma^1$ . Let  $\Gamma^3$

be a finite-index subgroup of  $\Gamma^2$  which is normal in  $\Gamma^1$ . The proof relies on the standard construction [Fur99b] of a groupoid isomorphism between two measured groupoids coming from restrictions of measure-preserving actions of  $\Gamma^3$ , from a self coupling  $\Sigma$  of  $\Gamma^1$  (which is also a self coupling of  $\Gamma^3$ ).

Let  $X$  be a Borel fundamental domain for the action of  $\Gamma^3 \times \{1\}$  on  $\Sigma$ . One can choose a Borel fundamental domain  $X'$  for the action of  $\{1\} \times \Gamma^3$  on  $\Sigma$  such that  $Y := X \cap X'$  satisfies that  $(\Gamma^3 \times \Gamma^3)Y = \Sigma$  up to a null set (see e.g. [Kid09, Lemma 2.27]).

Consider the groupoid  $\mathcal{G} = ((\Gamma^3 \times \Gamma^3) \ltimes \Sigma)|_Y$ , and recall that  $\mathcal{G}$  is the set of pairs  $((\gamma, \gamma'), y)$  such that both  $y$  and  $(\gamma, \gamma')y$  lie in  $Y$ . Let  $\rho : \mathcal{G} \rightarrow \Gamma^3$  and  $\rho' : \mathcal{G} \rightarrow \Gamma^3$  be the cocycles defined by  $\rho((\gamma, \gamma'), y) = \gamma$  and  $\rho'((\gamma, \gamma'), y) = \gamma'$ . We claim that the cocycles  $\rho$  and  $\rho'$  are action-type. Indeed,  $X$  comes equipped with a natural action of  $\{1\} \times \Gamma^3$ , by letting  $(1, \gamma') \cdot x$  be the unique element of  $X$  in the same  $(\Gamma^3 \times \{1\})$ -orbit as  $(1, \gamma')x \in \Sigma$ . There is a natural isomorphism between  $\mathcal{G}$  and  $((\{1\} \times \Gamma^3) \ltimes X)|_Y$  given by  $((\gamma, \gamma'), y) \mapsto ((1, \gamma'), y)$ , and under this identification,  $\rho'$  becomes the natural cocycle of  $((\{1\} \times \Gamma^3) \ltimes X)|_Y$ . By Lemmas 3.21 and 3.22, this shows that  $\rho'$  is action-type. By a symmetric argument, so is  $\rho$ .

By cocycle rigidity, our choice of  $\Gamma^0$  ensures that there exist a conull Borel subset  $Y^* \subseteq Y$  and a Borel map  $\varphi : Y^* \rightarrow \Gamma$  such that for all  $g \in \mathcal{G}|_{Y^*}$ , one has

$$\rho'(g) = \varphi(r(g))\rho(g)\varphi(s(g))^{-1}. \quad (1)$$

Consider the conull Borel subset  $\Sigma^* = (\Gamma^3 \times \Gamma^3)Y^*$  in  $\Sigma$ . We then claim that the assignment

$$(\gamma, \gamma')y \mapsto \gamma\varphi(y)^{-1}\gamma'^{-1}$$

for all  $(\gamma, \gamma') \in \Gamma^3 \times \Gamma^3$  and all  $y \in Y^*$  determines a  $(\Gamma^3 \times \Gamma^3)$ -equivariant Borel map  $\Phi : \Sigma^* \rightarrow \Gamma$ . To check that this is well defined, it suffices to check that if  $(\gamma, \gamma')y = y'$  with  $y, y' \in Y^*$ , then  $\gamma\varphi(y)^{-1}\gamma'^{-1} = \varphi(y')^{-1}$ . The arrow  $g = ((\gamma, \gamma'), y) \in \mathcal{G}$  has source  $s(g) = y$  and range  $r(g) = y'$ , so Equation (1) above says that  $\gamma' = \varphi(y')\gamma\varphi(y)^{-1}$  and concludes that  $\Phi$  is well defined.

The map  $\Phi$  is clearly  $(\Gamma^3 \times \Gamma^3)$ -equivariant. That  $\Phi$  is actually almost everywhere  $(\Gamma^1 \times \Gamma^1)$ -equivariant then follows from [Kid10, Lemma 5.8], using the fact that the group  $\Gamma$  is ICC.  $\square$

## 4.2 Measure equivalence rigidity

We recall from the introduction that two countable groups  $\Gamma_1$  and  $\Gamma_2$  are *virtually isomorphic* if there exist finite-index subgroups  $\Gamma_i^0 \subseteq \Gamma_i$  and finite normal subgroups  $F_i \trianglelefteq \Gamma_i^0$  such that  $\Gamma_1^0/F_1$  and  $\Gamma_2^0/F_2$  are isomorphic. A countable group  $\Gamma$  is *ME-superrigid* if for every countable group  $\Lambda$ , if  $\Lambda$  is measure equivalent to  $\Gamma$ , then  $\Gamma$  and  $\Lambda$  are virtually isomorphic.

**Theorem 4.5** (Kida [Kid10, Theorem 6.1]). *Let  $\Gamma$  be a countable ICC group which is rigid with respect to action-type cocycles.*

*Then  $\Gamma$  is ME-superrigid.*

*Proof.* This is a consequence of Proposition 4.3 and [Kid10, Theorem 6.1] (see also [BFS13, Theorem 2.6]).  $\square$

### 4.3 Orbit equivalence rigidity

Let  $\Gamma_1$  and  $\Gamma_2$  be two countable groups. Two free measure-preserving actions  $\Gamma_1 \curvearrowright X_1$  and  $\Gamma_2 \curvearrowright X_2$  on standard probability spaces are *stably orbit equivalent* if there exist Borel subsets  $Y_i \subseteq X_i$  of positive measure and a measure-scaling isomorphism  $f : Y_1 \rightarrow Y_2$  such that for a.e.  $y \in Y_1$ , one has

$$f((\Gamma_1 \cdot y) \cap Y_1) = (\Gamma_2 \cdot f(y)) \cap Y_2.$$

A particular case of stably orbit equivalent actions is when the two actions are conjugate, or virtually conjugate, in the following sense.

They are *conjugate* if there exists an isomorphism  $\varphi : \Gamma_1 \rightarrow \Gamma_2$  and a measure preserving isomorphism  $f : X_1 \rightarrow X_2$  such that  $f(\gamma x) = \varphi(\gamma)f(x)$  for all  $\gamma \in \Gamma_1$  and almost every  $x \in X_1$ .

More generally, they are *virtually conjugate* if there exist short exact sequences  $1 \rightarrow F_i \rightarrow \Gamma_i \rightarrow \bar{\Gamma}_i \rightarrow 1$  with  $F_i$  finite for every  $i \in \{1, 2\}$ , finite-index subgroups  $\bar{\Gamma}_i^0 \subseteq \bar{\Gamma}_i$ , and conjugate actions  $\bar{\Gamma}_1^0 \curvearrowright X_1'$  and  $\bar{\Gamma}_2^0 \curvearrowright X_2'$  such that for every  $i \in \{1, 2\}$ , the action  $\bar{\Gamma}_i \curvearrowright X_i/F_i$  is induced from the action  $\bar{\Gamma}_i^0 \curvearrowright X_i'$  as in [Kid08b, Definition 2.1]. Notice that this implies in particular that the groups  $\Gamma_1$  and  $\Gamma_2$  are virtually isomorphic.

In the particular case where the groups  $\Gamma_i$  have no non-trivial finite normal subgroup and the actions  $\Gamma_i \curvearrowright X_i$  are aperiodic, i.e. every finite index subgroup of  $\Gamma_i$  acts ergodically on  $X_i$ , the actions are conjugate if and only if they are virtually conjugate. Indeed, an action induced from a proper subgroup of finite index is never aperiodic. Aperiodicity holds for instance for Bernoulli actions of  $\Gamma^1$ .

An ergodic measure-preserving free action  $\alpha$  of a countable group  $\Gamma$  on a standard probability space is *OE-superrigid* if for every ergodic measure-preserving free action  $\beta$  of a countable group  $\Lambda$  on a standard probability space, if  $\alpha$  and  $\beta$  are stably orbit equivalent, then  $\alpha$  and  $\beta$  are virtually conjugate.

**Theorem 4.6** (Furman [Fur99b, Theorem A], [Kid08b, Theorem 1.1]). *Let  $\Gamma$  be a countable ICC group which is rigid with respect to action-type cocycles.*

*Then for every finite-index subgroup  $\Gamma^1 \subseteq \Gamma$ , every ergodic measure-preserving free action of  $\Gamma^1$  on a standard probability space is OE-superrigid.*

*Proof.* The proof of [Kid08b, Theorem 1.1] derives Theorem 4.6 from Theorem 4.5. The idea is to use again the dictionary between measure equivalence and stable orbit equivalence developed by Furman in [Fur99b]. See also [Fur11b, Lemma 4.18].  $\square$

See [Kid08b, Theorem 1.2] for variants of the theorem.

#### 4.4 Lattice embeddings

Let  $H$  be a locally compact second countable group with a bi-invariant Haar measure  $\lambda_H$ . Given a discrete group  $\Gamma$ , a homomorphism  $\varphi : \Gamma \rightarrow H$  is a *lattice embedding* if  $\varphi$  is one-to-one and  $\varphi(\Gamma)$  is a lattice in  $H$ , i.e.  $\varphi(\Gamma)$  is discrete in  $H$  and has finite  $\lambda_H$ -covolume.

**Theorem 4.7** (Furman [Fur11a], Kida [Kid10, Theorem 8.1]). *Let  $\Gamma$  be a countable ICC group which is rigid with respect to action-type cocycles.*

*Then for every finite-index subgroup  $\Gamma^1 \subseteq \Gamma$ , every locally compact second countable group  $H$ , and every lattice embedding  $\sigma : \Gamma^1 \rightarrow H$ , there exists a continuous homomorphism  $\Phi_0 : H \rightarrow \Gamma$  with compact kernel such that for all  $\gamma \in \Gamma^1$ , one has  $\Phi_0(\sigma(\gamma)) = \gamma$ .*

*Proof.* We give the argument for completeness, following [Kid10, Theorem 8.1]. The group  $\Gamma^1 \times \Gamma^1$  acts on  $H$  via  $(\gamma, \gamma') \cdot h := \sigma(\gamma)h\sigma(\gamma')^{-1}$ . As  $\sigma$  is a lattice embedding, this turns  $H$  (equipped with the bi-invariant Haar measure  $\lambda_H$ ) into a self coupling of  $\Gamma^1$ . Since  $\Gamma$  is ICC and rigid with respect to action-type cocycles, Proposition 4.3 yields the existence of an almost  $(\Gamma^1 \times \Gamma^1)$ -equivariant Borel map  $\Phi : H \rightarrow \Gamma$ .

One first proves that for a.e.  $(h_1, h_2) \in H$ , one has  $\Phi(h_1h_2) = \Phi(h_1)\Phi(h_2)$ . To see this, consider the map

$$\begin{aligned} F : H \times H &\rightarrow \Gamma \\ (h_1, h_2) &\mapsto \Phi(h_1^{-1})^{-1}\Phi(h_1^{-1}h_2)\Phi(h_2)^{-1}. \end{aligned}$$

An easy computation shows that  $F(h_1\gamma, h_2\gamma') = F(h_1, h_2)$  and that  $F(\gamma h_1, \gamma h_2) = \gamma F(h_1, h_2)\gamma^{-1}$  for almost every  $h_1, h_2 \in H$  and all  $\gamma, \gamma' \in \Gamma^1$ . Denote by  $X = H/\Gamma^1$  the right quotient, endowed with the left  $H$ -invariant probability measure  $\mu$ . Then  $F$  induces a map  $\bar{F} : X \times X \rightarrow \Gamma$ . The image of  $\mu \otimes \mu$  under  $\bar{F}$  is a probability measure on  $\Gamma$  which is invariant under conjugation by  $\Gamma^1$ . Since  $\Gamma$  is ICC and  $\Gamma^1$  has finite index in  $\Gamma$ , this measure must be the Dirac mass at the identity. This precisely means that for a.e.  $(h_1, h_2) \in H$ , one has  $\Phi(h_1h_2) = \Phi(h_1)\Phi(h_2)$ .

By [Zim84, Theorems B.2 and B.3],  $\Phi$  agrees almost everywhere with a continuous homomorphism  $\Phi_0 : H \rightarrow \Gamma$ . Since  $\Phi^{-1}(\{1_\Gamma\})$  is a Borel fundamental domain for the left action of  $\Gamma^1$  on  $H$ ,  $\Phi_0^{-1}(\{1_\Gamma\})$  is a closed subgroup of finite positive Haar measure, hence compact. Finally, using the equivariance of  $\Phi$  and the fact that  $\Phi$  and  $\Phi_0$  agree almost everywhere, we see that for all  $\gamma \in \Gamma^1$  and almost every  $h \in H$ , one has

$$\Phi_0(\sigma(\gamma))\Phi_0(h) = \Phi_0(\sigma(\gamma)h) = \Phi(\sigma(\gamma)h) = \gamma\Phi(h) = \gamma\Phi_0(h),$$

and therefore  $\Phi_0(\sigma(\gamma)) = \gamma$ . □

Given a finitely generated group  $\Gamma$ , and a finite generating set  $S$  of  $\Gamma$ , define the *Cayley graph*  $\text{Cay}(\Gamma, S)$  to be the simple graph with vertices  $\Gamma$ , and such that there is an edge between two distinct elements  $g$  and  $h$  if and only if  $gh^{-1} \in S \cup S^{-1}$ . The following consequence of Theorem 4.7 was suggested to us by Jingyin Huang.

**Corollary 4.8.** *Let  $\Gamma$  be a finitely generated ICC group which is rigid with respect to action-type cocycles.*

1. *For every finite generating set  $S$  of  $\Gamma$ , every automorphism of  $\text{Cay}(\Gamma, S)$  is at bounded distance from an automorphism induced by the left multiplication by an element of  $\Gamma$ .*
2. *For every torsion-free finite-index subgroup  $\Gamma^1 \subseteq \Gamma$ , and every finite generating set  $S$  of  $\Gamma^1$ , the automorphism group of  $\text{Cay}(\Gamma^1, S)$  is isomorphic to a subgroup of  $\Gamma$  containing  $\Gamma^1$ . In particular, it is countable.*

Notice that the first conclusion does not hold as such if  $\Gamma$  is replaced by an arbitrary finite-index subgroup. Indeed, if  $\Gamma^1$  is a normal finite index subgroup of  $\Gamma$  and  $S$  is a finite generating set of  $\Gamma^1$  which is invariant under conjugation by a finite order element  $\gamma \in \Gamma \setminus \Gamma^1$ , then the conjugation by  $\gamma$  is an automorphism of  $\text{Cay}(\Gamma^1, S)$ , and using the ICC condition, one can check that it is not at bounded distance from the left translation by any element of  $\Gamma^1$ .

Also, the second conclusion does not hold if  $\Gamma^1$  has torsion by [dlST19, Lemma 6.1].

*Proof.* Let  $\Gamma^1$  be any finite index subgroup of  $\Gamma$  (we will take  $\Gamma^1 = \Gamma$  or assume  $\Gamma^1$  torsion-free below). Denote by  $G$  the group of all graph isomorphisms of  $\text{Cay}(\Gamma^1, S)$ , equipped with the compact-open topology. As  $\text{Cay}(\Gamma^1, S)$  is a locally finite graph, this is a locally compact second countable group and point stabilizers are compact subgroups of  $G$ . As  $\Gamma^1$  acts transitively on the vertex set of  $\text{Cay}(\Gamma^1, S)$ , it follows that  $\Gamma^1$  is a (cocompact) lattice inside  $G$ . Theorem 4.7 thus yields the existence of a group homomorphism  $\Phi_0 : G \rightarrow \Gamma$  with compact kernel  $K$ , such that for every  $\gamma \in \Gamma^1$ , denoting by  $L_\gamma \in G$  the automorphism of  $\text{Cay}(\Gamma^1, S)$  induced by the left multiplication by  $\gamma$ , one has  $\Phi_0(L_\gamma) = \gamma$ .

As  $K$  is compact, the  $K$ -orbit of  $1_{\Gamma^1}$  is finite, and since  $K$  is normalized by  $\Gamma^1$ , all  $K$ -orbits have the same diameter so  $K$  is at bounded distance of the identity.

If  $\Gamma^1 = \Gamma$ , every isometry of  $\text{Cay}(\Gamma, S)$  can be written as a product of an element of  $K$  and of the left multiplication by an element of  $\Gamma$ , so the first conclusion follows.

We now assume that  $\Gamma^1$  is torsion-free and derive the second conclusion. Since  $K$  is normal,  $\Gamma^1$  permutes the  $K$ -orbits in  $\text{Cay}(\Gamma^1, S)$ :  $\gamma.K(\gamma') = K(\gamma\gamma')$ . It follows that  $K(1_{\Gamma^1})$  is a finite subgroup of  $\Gamma^1$ , so  $K(1_{\Gamma^1}) = \{1_{\Gamma^1}\}$ . It follows that  $K(\gamma) = \{\gamma\}$  for all  $\gamma \in \text{Cay}(\Gamma^1, S)$ , hence  $K = \{\text{id}\}$  and  $\Phi_0$  gives the desired isomorphism.  $\square$

## Part II

# Stabilizers of collections of splittings or factors

The goal of the second part of our work is to provide three constructions of canonical splittings associated to a subgroup of  $\text{Out}(F_N)$  (with  $N \geq 2$ ). The first construction relies on the existence of invariant non-trivial free splittings (see Section 6). The second relies on the existence of invariant  $\mathcal{Z}_{\max}$ -splittings assuming that there is no invariant non-trivial free splitting (see Section 7). The third one produces a splitting from the existence of several maximal invariant almost free factor systems, still assuming that there is no invariant non-trivial free splitting (see Section 8). A convenient unifying statement is given in Section 9.

## 5 $H$ -invariant splittings and trees of cylinders

This section records preliminary material that will be used in the next three sections.

### 5.1 $H$ -invariant splittings

Let  $H \subseteq \text{Out}(F_N)$  be a subgroup and let  $\tilde{H}$  be the full preimage of  $H$  in  $\text{Aut}(F_N)$ . In this subsection, we recall the relation between  $H$ -invariant splittings of  $F_N$  and actions of  $\tilde{H}$  on trees.

Given a splitting  $S$  of  $F_N$ , recall that  $S$  being  $H$ -invariant means that for every  $\tilde{\alpha} \in \tilde{H}$ , there exists an isometry  $I_{\tilde{\alpha}} : S \rightarrow S$  which is  $\tilde{\alpha}$ -equivariant in the following sense: for every  $g \in F_N$  and every  $x \in S$ , one has

$$I_{\tilde{\alpha}}(gx) = \tilde{\alpha}(g)I_{\tilde{\alpha}}(x).$$

As soon as  $S$  is not a line (which will always be the case in our setting),  $I_{\tilde{\alpha}}$  is unique, and  $\tilde{\alpha} \in \tilde{H} \mapsto I_{\tilde{\alpha}} \in \text{Aut}(S)$  defines an action of  $\tilde{H}$  on  $S$  for which the action of an inner automorphism  $\text{ad}_g$  is the same as the original action of  $g$ . We will thus consistently identify  $F_N$  with  $\text{Inn}(F_N)$ , thus viewing  $F_N$  as a normal subgroup  $F_N \triangleleft \tilde{H}$ . This way, the restriction to  $F_N$  of the action of  $\tilde{H}$  on  $S$  is the same as the original  $F_N$ -action. Conversely, given an action of  $\tilde{H}$  on a tree  $S$ , restricting the action to  $F_N$  yields an  $H$ -invariant  $F_N$ -tree.

We will denote by  $\tilde{H}_e, \tilde{H}_v$  the  $\tilde{H}$ -stabilizers of an edge  $e$  or a vertex  $v$  of  $S$ , and by  $G_e, G_v$  their  $F_N$ -stabilizers. Since  $G_e = \tilde{H}_e \cap F_N$ ,  $S$  is a free splitting (resp. a cyclic splitting) of  $F_N$  if and only if for each edge  $e \subseteq S$ ,  $\tilde{H}_e \cap F_N$  is trivial (resp. cyclic).

### 5.2 Trees of cylinders

Trees of cylinders were introduced in [GL11] as a way to produce a canonical splitting of a group  $G$  out of a deformation space. The construction is based on a notion of cylinders



in a tree, associated to an equivalence relation on edge stabilizers. This notion will be used in the next three sections, with  $G$  being either the free group  $F_N$  or a subgroup  $\tilde{H} \subseteq \text{Aut}(F_N)$  as above.

Let  $G$  be a group, and  $\mathcal{E}$  be a collection of subgroups of  $G$ , stable under conjugation (this is the collection of allowed edge groups). Following [GL11, Definition 3.1], we say that an equivalence relation  $\sim$  on  $\mathcal{E}$  is *admissible* if for every  $A, B \in \mathcal{E}$ , and every  $G$ -tree  $T$  with edge groups in  $\mathcal{E}$ , the following three conditions hold:

1. for every  $g \in G$ , if  $A \sim B$ , then  $A^g \sim B^g$ , and
2. if  $A \subseteq B$ , then  $A \sim B$ , and
3. for every pair of points  $a, b \in T$  fixed by  $A, B$  respectively, if  $A \sim B$ , then for every edge  $e \subseteq [a, b]$ , one has  $A \sim G_e \sim B$ .

More generally, given a collection  $\mathcal{P}$  of subgroups of  $G$ , one says that an equivalence relation  $\sim$  on  $\mathcal{E}$  is *admissible relative to  $\mathcal{P}$*  if the third condition is only required to hold for  $G$ -trees  $T$  relative to  $\mathcal{P}$  with edge groups in  $\mathcal{E}$ .

One then defines the *cylinder* of an edge  $e$  in  $T$  as the union of all edges  $e'$  with  $G_e \sim G_{e'}$ . Admissibility of  $\sim$  implies that this is a connected subset of  $T$ .

A simple example of admissible relation is the commutation relation (equivalently, commensurability) on the class  $\mathcal{E}$  of non-trivial cyclic subgroups of  $F_N$ . This will be used in Section 7.

One then defines the *tree of cylinders*  $T_c$  of  $T$ : this is the bipartite simplicial tree having one vertex  $v_Y$  for each cylinder  $Y \subseteq S$ , one vertex  $v_x$  for each point  $x \in S$  that belongs to at least two cylinders, and an edge between  $v_x$  and  $v_Y$  whenever  $x \in Y$ . We denote by  $V^1(T_c)$  be the set of vertices corresponding to cylinders, and let  $V^0(T_c)$  be the set of all other vertices. The tree  $T_c$  is minimal [Gui04, Lemma 4.9]. It may happen that  $T_c$  is reduced to a point even if  $T$  is non-trivial. In general, it may also happen that edge stabilizers of  $T_c$  are not in  $\mathcal{E}$ .

Recall that two  $G$ -trees *belong to the same deformation space* if they have the same elliptic subgroups. By [GL11, Theorem 1], any two trees with edge stabilizers in  $\mathcal{E}$  that lie in the same deformation space have the same tree of cylinders.

## 6 Invariant free splittings and stabilizers of collections of free splittings

Throughout the section, we fix an integer  $N \geq 2$ . Given a subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , let  $\text{FS}^H$  be the collection of all  $H$ -invariant non-trivial free splittings of  $F_N$ . The goal of this section is to construct a canonical splitting  $U_H^1$  of  $F_N$  (not a free splitting in general) encoding all the  $H$ -invariant free splittings of  $F_N$ . The splitting  $U_H^1$  will be non-trivial as soon as  $H$  is infinite and  $\text{FS}^H \neq \emptyset$ . To make things a bit more concrete, let us start with an example.




Figure 3: The canonical splitting  $U_H^1$  encoding all  $H$ -invariant free splittings.

*Example 6.1.* Let  $\Sigma$  be a connected surface with one boundary component, and let  $c \subseteq \Sigma$  be an essential simple closed curve on  $\Sigma$  which separates  $\Sigma$  into two connected components  $\Sigma_0$  and  $\Sigma_1$ , with  $\Sigma_0$  containing the boundary curve. Choose an identification  $F_N \simeq \pi_1(\Sigma)$ . Let  $H_1 \subseteq \text{Out}(F_N)$  be the subgroup induced by all homeomorphisms of  $\Sigma$  which restrict to the identity on  $\Sigma_0$ , and let  $H = H_1 \cap \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Then any properly embedded arc in  $\Sigma_0$  defines a free splitting of  $F_N$  which is  $H$ -invariant. These arcs fill the subsurface  $\Sigma_0$ , and the canonical splitting  $U_H^1$  will be the splitting dual to the amalgamated product  $F_N = A_0 *_{\langle a \rangle} A_1$  corresponding to the decomposition  $\Sigma = \Sigma_0 \cup \Sigma_1$  (see the argument right after Remark 6.10). Then any free splitting of  $A_0$  relative to  $\langle a \rangle$  induces a free splitting of  $F_N$  which is  $H$ -invariant. It will follow from our construction that conversely, any  $H$ -invariant free splitting is of this form (but it might not correspond to an arc drawn on  $\Sigma_0$ ).

Dually, given any collection  $\mathcal{C}$  of free splittings of  $F_N$ , the *elementwise stabilizer*  $\Gamma_{\mathcal{C}}$  of  $\mathcal{C}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is defined as the subgroup of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  made of all automorphisms that fix all the splittings in  $\mathcal{C}$ . One can then describe  $\Gamma_{\mathcal{C}}$  as a particular subgroup of the stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$ . In the example above, starting with the collection  $\mathcal{C}$  of all free splittings dual to properly embedded arcs in  $\Sigma_0$ , the elementwise stabilizer of  $\Gamma_{\mathcal{C}}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  coincides with the subgroup of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  made of all automorphisms that preserve the decomposition  $F_N = A_0 *_{\langle a \rangle} A_1$  and act as the identity on  $A_0$ . Additionally, one can describe the normalizer of  $\Gamma_{\mathcal{C}}$  as the stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$ . See Section 6.5.

Our construction also allows us to give a bounded chain condition for stabilizers of collections of free splittings, see Section 6.4.

Our construction of the splitting  $U_H^1$  goes in two steps: starting from any group  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  and the collection  $\text{FS}^H$  of non-trivial  $H$ -invariant free splittings, we first construct in Section 6.1 a maximal  $H$ -invariant free splitting, which we also view as an  $\tilde{H}$ -tree where  $\tilde{H}$  is the preimage of  $H$  in  $\text{Aut}(F_N)$ . Such an  $\tilde{H}$ -tree is not unique, but the collection of all of them forms a deformation space of  $\tilde{H}$ -trees. In Section 6.2, we then construct a tree of cylinders to get a canonical splitting from this deformation space.

## 6.1 The deformation space of maximal $H$ -invariant free splittings

Let  $H \subseteq \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup and  $\tilde{H}$  be the full preimage of  $H$  in  $\mathrm{Aut}(F_N)$ . Identifying  $F_N$  with  $\mathrm{Inn}(F_N)$ , we view  $F_N$  as a normal subgroup of  $\tilde{H}$ . Recall that  $H$ -invariant splittings of  $F_N$  correspond to actions of  $\tilde{H}$  that extend the action of  $F_N$  (see Section 5.1).

Recall that given two  $\tilde{H}$ -trees  $S, S'$ , one says that  $S$  dominates  $S'$  if the  $\tilde{H}$ -stabilizer of every vertex in  $S$  is elliptic in  $S'$ . Two  $\tilde{H}$ -trees *belong to the same deformation space of  $\tilde{H}$ -trees* if they dominate each other, i.e. if they have the same elliptic subgroups (in  $\tilde{H}$ ).

If  $S, S'$  are two  $H$ -invariant free splittings of  $F_N$ , and if  $S$  dominates  $S'$  as an  $\tilde{H}$ -tree, then this also holds as  $F_N$ -tree. We will see in Remark 6.4 that the converse holds.

**Proposition 6.2.** *Let  $H \subseteq \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup.*

*There exists a (maybe trivial) maximum  $H$ -invariant free splitting, i.e. that dominates all  $H$ -invariant free splittings.*

*All maximum  $H$ -invariant free splittings are in the same deformation space of  $\tilde{H}$ -trees.*




Figure 4: Example 6.1 continued. A maximum  $H$ -invariant free splitting, dual to the two arcs  $\gamma_1, \gamma_2 \subseteq \Sigma$ .

*Example 6.1 continued.* The splitting  $S$  dual to the two arcs  $\gamma_1, \gamma_2 \subseteq \Sigma_0$  depicted on Figure 4 is  $H$ -invariant. Since  $\pi_1(\Sigma_1)$  is elliptic in any  $H$ -invariant free splitting,  $S$  dominates any  $H$ -invariant free splitting and therefore satisfies the conclusion of Proposition 6.2.

In the sequel, we will denote by  $\mathcal{D}_H$  the deformation space of  $\tilde{H}$ -trees provided by Proposition 6.2. The main point in the proof of Proposition 6.2 is the following lemma.

**Lemma 6.3.** *Let  $H \subseteq \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and let  $S, S' \in \mathrm{FS}^H$ . Then the  $\tilde{H}$ -stabilizer of every edge of  $S$  is elliptic in  $S'$ .*

*Proof.* As  $H \subseteq \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , every automorphism  $\tilde{\alpha} \in \tilde{H}$  acts trivially on the quotient graph  $S/F_N$  (Lemma 2.12), and therefore induces an isometry of every collapse of  $S$ . Therefore, without loss of generality, we can assume that  $S$  has a single  $F_N$ -orbit of edges. Let  $e \subseteq S$  be an edge, and let  $\tilde{\alpha} \in \mathrm{Stab}_{\tilde{H}}(e)$ . We denote by  $I_{\tilde{\alpha}}$  the  $\tilde{\alpha}$ -equivariant

isometry of  $S$ , and by  $I'_\alpha$  the  $\tilde{\alpha}$ -equivariant isometry of  $S'$ . Our goal is to find a point in  $S'$  that is fixed by  $I'_\alpha$ , independently of  $\tilde{\alpha} \in \text{Stab}_{\tilde{H}}(e)$ .

Let  $A$  and  $B$  be the  $F_N$ -stabilizers of the two extremities  $v_A, v_B$  of  $e$ . Then  $A$  and  $B$  are proper free factors of  $F_N$ , and they are both non-trivial because  $S$  has a single orbit of edges. Since  $I_\alpha(e) = e$ , we have in particular  $I_\alpha(v_A) = v_A$ . Therefore, for every  $g \in A$ , we have  $\tilde{\alpha}(g)v_A = I_\alpha(gv_A) = I_\alpha(v_A) = v_A$ , so  $\tilde{\alpha}(g) \in A$ . This shows that  $\tilde{\alpha}(A) \subseteq A$ , and since  $\tilde{\alpha}$  is an automorphism, we deduce that  $\tilde{\alpha}(A) = A$ . Likewise  $\tilde{\alpha}(B) = B$ .

If  $A$  (or  $B$ ) fixes a point in  $S'$ , then this point is unique (because  $S'$  is a free splitting of  $F_N$ ), and it is fixed by  $I'_\alpha$ . We can therefore assume that neither  $A$  nor  $B$  fixes a point in  $S'$ .

Let  $S'_A$  (resp.  $S'_B$ ) be the minimal  $A$ -invariant (resp.  $B$ -invariant) subtree of  $S'$ . The isometry  $I'_\alpha$  sends the axis of every element  $a \in A$  acting hyperbolically on  $S'$  to the axis of  $\tilde{\alpha}(a)$ . As  $S'_A$  is equal to the union of the axes of all elements of  $A$  acting hyperbolically on  $S'$ , we deduce that  $S'_A$  is  $I'_\alpha$ -invariant. Likewise  $S'_B$  is  $I'_\alpha$ -invariant.

As  $A \cap B = \{1\}$ , the intersection  $S'_A \cap S'_B$  is compact (possibly empty). In addition this intersection is  $I'_\alpha$ -invariant. If  $S'_A \cap S'_B \neq \emptyset$ , then  $I'_\alpha$  fixes the circumcenter of  $S'_A \cap S'_B$ . If  $S'_A \cap S'_B = \emptyset$ , then  $I'_\alpha$  fixes the bridge between  $S'_A$  and  $S'_B$ . In both cases, we have found a point fixed by all  $I'_\alpha$ , as required.  $\square$

*Remark 6.4.* If  $S, S'$  are  $H$ -invariant free splittings such that  $S$  dominates  $S'$  as an  $F_N$ -tree, then  $S$  dominates  $S'$  as an  $\tilde{H}$ -tree. Indeed, if  $v \in S$  is a vertex whose  $F_N$ -stabilizer  $G_v$  is trivial, then  $\tilde{H}_v$  is commensurable to the stabilizer of an incident edge, so Lemma 6.3 shows that  $\tilde{H}_v$  fixes a point in  $S'$ . If  $G_v$  is non-trivial, then it fixes a unique point in  $S'$  and this point is  $\tilde{H}_v$  invariant since  $G_v$  is normal in  $\tilde{H}_v$ .

**Lemma 6.5.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and let  $S, S' \in \text{FS}^H$ . Assume that there does not exist any splitting in  $\text{FS}^H$  that properly refines  $S$ . Then  $S$  dominates  $S'$  as an  $\tilde{H}$ -tree.*

*Proof.* Assume not. Then there exists a point  $v \in S$  whose  $\tilde{H}$ -stabilizer  $\tilde{H}_v$  is not elliptic in  $S'$ . Let  $U$  be the minimal  $\tilde{H}_v$ -invariant subtree of  $S'$ . As  $\tilde{H}$ -edge stabilizers of  $S$  are elliptic in  $S'$  (Lemma 6.3), we can blow-up  $S$  to an  $\tilde{H}$ -tree  $\hat{S}$ , using the  $\tilde{H}_v$ -action on  $U$  (see e.g. [GL17, Proposition 2.2]).

We claim that  $\hat{S} \in \text{FS}^H$ , which will contradict our hypothesis. First,  $\hat{S}$  is  $H$ -invariant (because it is an  $\tilde{H}$ -tree). Second,  $F_N$ -stabilizers of edges of  $\hat{S}$  are trivial: indeed, every  $\tilde{H}$ -edge stabilizer of  $\hat{S}$  stabilizes an edge in either  $S$  or  $S'$ , and  $\tilde{H}$ -edge stabilizers of  $S$  and  $S'$  have trivial intersection with  $F_N$  because  $S$  and  $S'$  are free splittings of  $F_N$ . This proves our claim, and concludes the proof of the lemma.  $\square$

*Proof of Proposition 6.2.* Consider an  $H$ -invariant free splitting  $S$  of  $F_N$  without redundant vertex, whose number of orbits of edges is maximum (the trivial splitting if  $\text{FS}^H = \emptyset$ ). By Lemma 6.5, the tree  $S$  is a maximum  $H$ -invariant free splitting.

If  $S, S'$  are two maximum  $H$ -invariant free splittings, then they dominate each other as  $\tilde{H}$ -trees by definition, so they are in the same deformation space of  $\tilde{H}$ -trees.  $\square$

## 6.2 The tree of cylinders of maximal $H$ -invariant free splittings

In this section we are going to construct a tree of cylinders of the deformation space  $\mathcal{D}_H$  following [GL11] (see Section 5.2 for relevant definitions).

Let  $\mathcal{E}$  be the collection of all  $\tilde{H}$ -stabilizers of edges of  $H$ -invariant free splittings of  $F_N$ . To construct a tree of cylinders, we need an admissible equivalence relation on  $\mathcal{E}$ . We will take the trivial equivalence relation (i.e. equality), and the following lemma implies that this is an admissible equivalence relation on  $\mathcal{E}$ .

**Lemma 6.6.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and let  $S, S' \in \text{FS}^H$ . Let  $e \subseteq S$  and  $e' \subseteq S'$  be edges. If the  $\tilde{H}$ -stabilizer  $\tilde{H}_e$  of  $e$  is contained in the  $\tilde{H}$ -stabilizer  $\tilde{H}_{e'}$  of  $e'$ , then  $\tilde{H}_e = \tilde{H}_{e'}$ .*

Lemma 6.6 was proved in [BGH17, Lemma 6.9], we recall the proof for completeness.

*Proof.* Let  $\tilde{\alpha} \in \tilde{H}_{e'}$ ; we aim to prove that  $\tilde{\alpha} \in \tilde{H}_e$ . As  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  and  $S$  is a free splitting of  $F_N$ , Lemma 2.12 ensures that every automorphism in  $\tilde{H}$  acts trivially on the quotient graph  $S/F_N$ , so there exists  $g \in F_N$  such that  $\tilde{\alpha}e = ge$ . Then  $\text{ad}_g^{-1}\tilde{\alpha} \in \tilde{H}_e$ , and hence  $\text{ad}_g^{-1}\tilde{\alpha} \in \tilde{H}_{e'}$  because  $\tilde{H}_e \subseteq \tilde{H}_{e'}$ . As  $\text{ad}_g^{-1}\tilde{\alpha}$  and  $\tilde{\alpha}$  both fix  $e'$ , we deduce that  $g$  fixes  $e'$ , and therefore  $g = 1$ . Hence  $\tilde{\alpha} \in \tilde{H}_e$ , which concludes our proof.  $\square$

Recall that the cylinder of an edge  $e \subseteq S$  is the union of all edges whose  $\tilde{H}$ -stabilizers are equal to  $\tilde{H}_e$  (admissibility implies that this is a connected subset).

**Lemma 6.7.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, let  $S \in \mathcal{D}_H$ , and let  $Y \subseteq S$  be a cylinder. The subgroup  $\tilde{H}_Y$  of elements of  $\tilde{H}$  fixing  $Y$  pointwise is a lift of  $H$  in  $\text{Aut}(F_N)$ : the restriction of  $\text{Aut}(F_N) \rightarrow \text{Out}(F_N)$  to  $\tilde{H}_Y$  is an isomorphism  $\tilde{H}_Y \xrightarrow{\sim} H$ .*

*Remark 6.8.* Note that  $\tilde{H}_Y$  coincides with  $\tilde{H}_e$  for any edge  $e \subseteq Y$ . Indeed, if  $\tilde{\alpha} \in \tilde{H}_e$ , then for any edge  $e' \subseteq Y$ ,  $\tilde{H}_{e'} = \tilde{H}_e \ni \tilde{\alpha}$ .

*Proof.* Since  $F_N$ -stabilizers of edges of  $S$  are trivial,  $\tilde{H}_Y \cap F_N = \{1\}$  which proves injectivity.

To prove surjectivity, consider  $\alpha \in H$ . Let  $e \subseteq Y$  be an edge. Since  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , it acts trivially on the quotient graph  $S/F_N$ , so  $\alpha$  has a representative  $\tilde{\alpha} \in \tilde{H}$  that fixes the edge  $e$ . As noted above, we conclude that  $\tilde{\alpha} \in \tilde{H}_e = \tilde{H}_Y$ .  $\square$

Given an automorphism  $\tilde{\alpha}$  of  $F_N$ , we denote by  $\text{Fix}_{F_N}(\tilde{\alpha})$  the subgroup of  $F_N$  made of all elements  $g \in F_N$  such that  $\tilde{\alpha}(g) = g$ . More generally, if  $\mathcal{X}$  is any collection of automorphisms of  $F_N$ , we let

$$\text{Fix}_{F_N}(\mathcal{X}) := \bigcap_{\tilde{\alpha} \in \mathcal{X}} \text{Fix}_{F_N}(\tilde{\alpha}).$$

**Lemma 6.9.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, let  $S \in \mathcal{D}_H$ , and let  $Y \subseteq S$  be a cylinder. Denote by  $\tilde{H}_{(Y)}$  and  $G_{(Y)}$  the global stabilizer of  $Y$  in  $\tilde{H}$  or in  $F_N$  respectively, and by  $\tilde{H}_Y$  the pointwise stabilizer of  $Y$  in  $\tilde{H}$ .*

*Then  $G_{(Y)} = \text{Fix}_{F_N}(\tilde{H}_Y)$  and  $\tilde{H}_{(Y)} = \tilde{H}_Y \times G_{(Y)}$ .*

*Proof.* Fix an edge  $e \subseteq Y$ . To prove that  $G_{(Y)} \subseteq \text{Fix}_{F_N}(\tilde{H}_Y)$ , consider  $g \in G_{(Y)}$  and  $\tilde{\alpha} \in \tilde{H}_Y$ . As  $e$  and  $ge$  both belong to  $Y$ , we have  $ge = I_{\tilde{\alpha}}(ge) = \tilde{\alpha}(g)I_{\tilde{\alpha}}(e) = \tilde{\alpha}(g)e$ . As the  $F_N$ -stabilizer of  $e$  is trivial, this implies that  $\tilde{\alpha}(g) = g$ . This shows that  $G_{(Y)} \subseteq \text{Fix}_{F_N}(\tilde{H}_Y)$ .

Conversely, let  $g \in F_N$  be such that for every  $\tilde{\alpha} \in \tilde{H}_Y$ , one has  $\tilde{\alpha}(g) = g$ . To prove that  $g \in G_{(Y)}$  it suffices to check that  $ge \subseteq Y$ . For every  $\tilde{\alpha} \in \tilde{H}_e$ , one has  $I_{\tilde{\alpha}}(ge) = \tilde{\alpha}(g)I_{\tilde{\alpha}}(e) = ge$ , showing that  $\tilde{\alpha} \in \tilde{H}_{ge}$ . Therefore  $\tilde{H}_e \subseteq \tilde{H}_{ge}$ . By Lemma 6.6, this implies that  $\tilde{H}_e = \tilde{H}_{ge}$ . Thus  $e$  and  $ge$  belong to the same cylinder  $Y$ , and we conclude that  $G_{(Y)} = \text{Fix}_{F_N}(\tilde{H}_Y)$ .

Now the groups  $\tilde{H}_Y$  and  $G_{(Y)} = \tilde{H}_{(Y)} \cap F_N$  are normal in  $\tilde{H}_{(Y)}$ , and they intersect trivially because  $F_N$ -stabilizers of edges of  $S$  are trivial. To conclude, it suffices to check that  $\tilde{H}_{(Y)} = \tilde{H}_Y \cdot G_{(Y)}$ . Given  $\tilde{\alpha} \in \tilde{H}_{(Y)}$ , there exists  $g \in F_N$  such that  $I_{\tilde{\alpha}}(e) = ge$  (as follows from Lemma 2.12 since  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ ). Therefore  $g \in G_{(Y)}$ . Since  $\text{ad}_g^{-1}\tilde{\alpha}$  fixes  $e$ , we have  $\text{ad}_g^{-1}\tilde{\alpha} \in \tilde{H}_Y$  by Remark 6.8, which concludes the proof.  $\square$

We define the tree  $U_H^1$  as the tree of cylinders of any  $\tilde{H}$ -tree  $S \in \mathcal{D}_H$  ([GL11], see Section 5.2). We denote by  $V^1$  the set of vertices  $v_Y$  of  $U_H^1$  corresponding to cylinders  $Y$  of  $S$ , and by  $V^0$  the set of vertices  $v_x$  of  $U_H^1$  corresponding to points  $x \in S$  lying in at least two distinct cylinders. The tree  $U_H^1$  is minimal as an  $\tilde{H}$ -tree [Gui04, Lemma 4.9] (it may be reduced to a point). Since  $F_N$  is normal in  $\tilde{H}$ , the tree  $U_H^1$  is also minimal as an  $F_N$ -tree. In general,  $F_N$ -stabilizers of edges of  $U_H^1$  can be non-abelian. We will see below that it is not trivial when  $H$  is infinite and trees in  $\mathcal{D}_H$  are non-trivial (i.e.  $\text{FS}^H \neq \emptyset$ ), see Theorem 6.12.

*Remark 6.10.* Another more geometric interpretation of the tree of cylinders in terms of cross-connected components of  $H$ -invariant almost invariant sets is given in [GL10].

*Example 6.1 continued.* Recall that  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is the group of automorphisms induced by homeomorphisms of the surface  $\Sigma$  that are the identity on  $\Sigma_0$ . Let  $S$  be the splitting dual to the arcs  $\gamma_1, \gamma_2$  as in Figure 4. Let  $\tilde{\Sigma}$  be the universal cover of  $\Sigma$ , and  $\tilde{\Sigma}_0 \subseteq \tilde{\Sigma}$  a connected component of the preimage of  $\Sigma_0$ . Let  $\tilde{E}$  be the collection of arcs of  $\tilde{\Sigma}$  that are lifts of  $\gamma_1$  or  $\gamma_2$ . Thus,  $\tilde{E}$  is in bijection with the set of edges of  $S$ . Denote by  $\tilde{E}_{\tilde{\Sigma}_0}$  the collection of all arcs of  $\tilde{E}$  contained in  $\tilde{\Sigma}_0$ , and let  $Y_{\tilde{\Sigma}_0} \subseteq S$  the corresponding collection of edges. Any automorphism  $h \in H$  has a lift acting as the identity on  $\tilde{\Sigma}_0$ , thus fixing all the arcs in  $\tilde{E}_{\tilde{\Sigma}_0}$ . The collection of these lifts defines a lift  $\tilde{H} \subseteq \text{Aut}(F_N)$  of  $H$  which fixes all the edges in  $Y_{\tilde{\Sigma}_0}$ , so  $\tilde{H} = \tilde{H}_e$  for all  $e \in Y$ . In particular, all edges in  $Y$  are in the same cylinder. Using the fact that  $H$  contains a Dehn twist around  $c$ , one can check that edges outside  $Y$  are not fixed by all elements of  $\tilde{H}$ , so  $Y$  is precisely a cylinder of  $S$ , and the tree of cylinders  $U_H^1$  of  $S$  is the cyclic splitting dual to the curve  $c \subseteq \Sigma$ .

*Example 6.11.* The following three examples show that as an  $F_N$ -tree,  $U_H^1$  is not determined by the sole collection of  $H$ -invariant free splittings, but it really depends on  $H$ .

Let  $S$  be the free splitting obtained as the Bass–Serre tree of the decomposition  $F_N = A * B$ , with  $A, B$  non-abelian. Denote by  $e \subseteq S$  the edge joining the vertices fixed by  $A$  and  $B$ . Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be the stabilizer of  $S$ . The group  $\text{Aut}(A) \times \text{Aut}(B)$  naturally embeds in  $\text{Aut}(F_N)$ , and  $\tilde{H}_e \simeq H$  has finite index in  $\text{Aut}(A) \times \text{Aut}(B)$ . Then  $S$  is the only  $H$ -invariant non-trivial free splitting, so  $S \in \mathcal{D}_H$ , and we claim that every cylinder is reduced to a single edge. Indeed, for any  $a \in A \setminus \{1\}$ , the edges  $e$  and  $ae$  cannot be in the same cylinder because  $a \notin \text{Fix}_{F_N}(\tilde{H}_e)$ . Using the same argument in  $B$ , we see that the cylinder of  $e$  contains no edge adjacent to  $e$ , which proves our claim. It follows that  $U_H^1$  is isomorphic to the barycentric subdivision of  $S$ , with  $V^1$  corresponding to the midpoints of edges of  $S$ .

From now on we further assume that  $\text{rk}(A) \geq 3$ . Let  $a \in A$  be an element which is part of a free basis of  $A$ , and let  $H' \subseteq H$  be the stabilizer (inside  $H$ ) of the cyclic splitting  $S'$  dual to the cyclic decomposition  $F_N = A *_{\langle a \rangle} (\langle a \rangle * B)$ . Now  $\tilde{H}'_e \simeq H'$  has finite index in  $\text{Aut}(A, a) \times \text{Aut}(B)$ . Again,  $S$  is the only  $H'$ -invariant free splitting. But now, the cylinder of  $e$  is the orbit of  $e$  under  $\langle a \rangle$  because  $\text{Fix}_{F_N}(\tilde{H}'_e) = \langle a \rangle$ . As an  $F_N$ -tree, the tree  $U_{H'}^1$  is then dual to the two-edge splitting

$$A \xrightarrow{\langle a \rangle} \langle a \rangle \xrightarrow{\{1\}} B$$

and as an  $\tilde{H}'$ -tree, it is isomorphic to

$$(\tilde{H}'_e \times A) \xrightarrow{\tilde{H}'_e \times \langle a \rangle} \tilde{H}'_e \times \langle a \rangle \xrightarrow{\tilde{H}'_e} (\tilde{H}'_e \times B).$$

The product structure on the central vertex illustrates Lemma 6.9 and the second assertion of Theorem 6.12 below.

Finally, assume furthermore that  $\text{rk}(B) \geq 3$ , and let  $b \in B$  be an element in a free basis, and let  $H'' \subseteq H$  be the stabilizer of the cyclic splitting  $T$  given by

$$A \xrightarrow{\langle a \rangle} \langle a, b \rangle \xrightarrow{\langle b \rangle} B.$$

Now  $\tilde{H}''_e \simeq H''$  has finite index in  $\text{Aut}(A, a) \times \text{Aut}(B, b)$ . Again,  $S$  is the unique  $H''$ -invariant non-trivial free splitting. The cylinder containing  $e$  is now the orbit of  $e$  under  $\langle a, b \rangle$  (it is unbounded) and  $U_{H''}^1 = T$  as an  $F_N$ -tree (it is no more in the same deformation space as  $S$ ). As an  $\tilde{H}''$ -tree, it is isomorphic to

$$(\tilde{H}''_e \times A) \xrightarrow{\tilde{H}''_e \times \langle a \rangle} \tilde{H}''_e \times \langle a, b \rangle \xrightarrow{\tilde{H}''_e \times \langle b \rangle} (\tilde{H}''_e \times B).$$

**Theorem 6.12.** *Consider a subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Recall that the vertex set of the tree  $U_H^1$  is bipartitioned into  $V^0 \sqcup V^1$ . Then the following properties hold.*

- (1) *the  $F_N$ -stabilizers of vertices in  $V^0$  are the maximal subgroups of  $F_N$  that are elliptic in all  $H$ -invariant free splittings; in particular, they are free factors of  $F_N$ .*



(2) for every vertex  $v \in V^1$ , its  $\tilde{H}$ -stabilizer splits as  $\tilde{H}_v = G_v \times \tilde{H}_Y$  where

$$G_v = \text{Fix}_{F_N}(\tilde{H}_Y)$$

and  $\tilde{H}_Y$  maps isomorphically onto  $H$  under  $\text{Aut}(F_N) \rightarrow \text{Out}(F_N)$ ;

(3) Consider the free splittings of  $F_N$  obtained from  $U_H^1$  by blowing up some vertices in  $V^1$  (using for each  $v \in V^1$  a maybe non-minimal free splitting of  $G_v$  relative to  $\text{Inc}_v$ ) and collapsing all edges coming from  $U_H^1$ . Then any such free splitting is  $H$ -invariant, and conversely every  $H$ -invariant free splitting may be obtained in this way.

(4) for every vertex  $v \in V^1$  such that every edge incident on  $v$  has non-trivial  $F_N$ -stabilizer, the collection of all conjugacy classes of  $F_N$ -stabilizers of incident edges forms a proper free factor system of  $G_v$ .

Moreover,  $U_H^1$  is non-trivial if  $H$  is infinite and  $\text{FS}^H \neq \emptyset$ .

*Remark 6.13.* In Assertion (3), one allows to blow up  $v \in V^1$  using a non-minimal splitting of  $G_v$  to get all  $H$ -invariant free splittings; on the other hand, this is only necessary when some edge incident on  $v$  has trivial stabilizer.

*Proof.* Let  $S$  be any  $\tilde{H}$ -tree in  $\mathcal{D}_H$ . Then  $U_H^1$  is the tree of cylinders of  $S$ . For each vertex  $v_x \in V^0$  corresponding to a vertex  $x \in S$ , one has  $\tilde{H}_{v_x} = \tilde{H}_x$  and  $G_{v_x} = G_x$ . Since  $x$  lies in two distinct cylinders,  $G_x \neq \{1\}$ . By definition of  $\mathcal{D}_H$ , the subgroup  $G_x$  is elliptic in all  $H$ -invariant splittings, and is maximal for this property because  $F_N \curvearrowright S$  is an  $H$ -invariant splitting of  $F_N$ . This proves the first assertion.

The second assertion follows from Lemmas 6.7 and 6.9 noting that if  $v \in V^1$  corresponds to a cylinder  $Y \subseteq S$ , then  $\tilde{H}_v = \tilde{H}_{(Y)}$  is the global stabilizer of  $Y$ .

To prove the third assertion, given a vertex  $v \in V^1$ , consider a free splitting  $Z$  of  $G_v$  relative to  $\text{Inc}_v$  (we do not assume that it is invariant under any group of automorphisms, and we allow it to be non-minimal). Writing  $\tilde{H}_v = G_v \times \tilde{H}_Y$ , the action of  $G_v$  on  $Z$  naturally extends to an action of  $\tilde{H}_v$  where  $\tilde{H}_Y$  acts trivially. For each edge  $e$  incident on  $v$ , there exists a point  $p_e \in Z$  fixed by  $\tilde{H}_e = \tilde{H}_Y \times G_e$  (not unique if  $G_e$  is trivial). Doing this in each orbit of vertices in  $V^1$ , this allows to blow up  $U_H^1$  into an  $\tilde{H}$ -tree  $\hat{U}_H^1$ , and collapsing all edges of  $\hat{U}_H^1$  coming from  $U_H^1$  yields an  $\tilde{H}$ -tree  $T$  (which is minimal if  $Z$  is the convex hull of the orbits of the attaching points  $p_e$ ). Edges of  $T$  have trivial  $F_N$ -stabilizer, so  $F_N \curvearrowright T$  is an  $H$ -invariant free splitting.

Conversely, consider any  $H$ -invariant free splitting  $T$  of  $F_N$ , and view it as an  $\tilde{H}$ -tree. By maximality, there is an  $\tilde{H}$ -equivariant map  $S \rightarrow T$ . By [GL11, Proposition 8.1],  $U_H^1$  is compatible with  $T$ , and the proof actually shows that one can construct a common refinement  $\hat{U}$  by blowing up vertices in  $V^1$  (using the corresponding cylinder in  $T$ ), and that one obtains  $T$  from  $\hat{U}$  by collapsing all edges coming from  $U_H^1$ . This concludes the proof of Assertion (3).

We now prove Assertion (4): assuming that all groups in  $\text{Inc}_v$  are non-trivial, we will prove that  $\text{Inc}_v$  is a free factor system of  $G_v$ . We use the fact that  $G_v = G_{(Y)}$  is

the global stabilizer of  $Y$ , and incident edge stabilizers are the intersections  $G_{(Y)} \cap G_x$  where  $x \in Y$  is a point lying in at least two cylinders. Since  $G_{(Y)} \curvearrowright Y$  has trivial edge stabilizers, it follows that either  $\text{Inc}_v = \{[G_{(Y)}]\}$  or  $\text{Inc}_v$  is a free factor system of  $G_v = G_{(Y)}$ . To prove Assertion (4), it thus suffices to prove that  $G_{(Y)}$  does not fix a point in  $Y$ . So assume that  $G_{(Y)}$  fixes a point  $x_0 \in Y$ . Recall that each edge  $\varepsilon$  of  $U_H^1$  incident on  $v$  corresponds to a pair  $(x, Y)$  with  $x \in Y$  lying in at least two cylinders. Since by assumption, all groups in  $\text{Inc}_v$  are non-trivial,  $G_\varepsilon$  fixes no other point apart from  $x_0$  so  $x = x_0$ . It follows that there is a single edge in  $U_H^1$  incident on  $v$ , contradicting the minimality of  $U_H^1$ .

Finally, assume that  $U_H^1$  is reduced to a point  $v$ . If  $v \in V^0$ , then  $G_v = F_N$  is elliptic in every  $H$ -invariant free splitting, so  $\text{FS}^H = \emptyset$ . If  $v \in V^1$ , then by Assertion (2),  $\tilde{H}_v = G_v \times \tilde{H}_Y = F_N \times \tilde{H}_Y$  and  $F_N = \text{Fix}_{F_N}(\tilde{H}_Y)$  i.e.  $\tilde{H}_Y = \{\text{id}\}$ . Since  $\tilde{H}_Y \simeq H$ ,  $H$  is trivial.  $\square$

### 6.3 The canonical splitting $U_H^1$ is biflexible

**Definition 6.14.** Let  $A$  be a group, and  $\mathcal{P}$  a finite collection of conjugacy classes of subgroups of  $A$ . We say that  $(A, \mathcal{P})$  is flexible if  $\text{Out}(A, \mathcal{P}^{(t)})$  is infinite.

A splitting  $S$  of  $F_N$  is biflexible if all its edge stabilizers are finitely generated non-abelian, and  $S$  has two vertices  $v_1, v_2$  in distinct  $F_N$ -orbits such that for every  $i \in \{1, 2\}$ , the pair  $(G_{v_i}, \text{Inc}_{v_i})$  is flexible.

**Corollary 6.15.** Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup. If  $H$  is infinite and  $\text{FS}^H \neq \emptyset$  then, as an  $F_N$ -tree, either  $U_H^1$  has an edge with trivial or maximal-cyclic stabilizer, or  $U_H^1$  is biflexible.

*Proof.* We know that  $U_H^1$  is a non-trivial splitting. We view it as an  $F_N$ -tree. Note that edge stabilizers of  $U_H^1$  are root-closed because vertex stabilizers are by Assertions (1) and (2) of Theorem 6.12. Denote by  $\bar{V}^0, \bar{V}^1$  the images of  $V^0, V^1$  in  $U_H^1/F_N$ . Assume that all  $F_N$ -stabilizers of edges of  $U_H^1$  are non-cyclic (in particular non-trivial). Consider a vertex  $v \in \bar{V}^1$ . By Theorem 6.12(4), the collection  $\text{Inc}_v$  forms a proper free factor system of  $G_v$ , so  $\text{Out}(G_v, \text{Inc}_v^{(t)})$  is infinite; indeed the group of twists of any Grushko decomposition of  $G_v$  relative to  $\text{Inc}_v$  is infinite (and non-abelian) because  $\text{Inc}_v$  contains a non-abelian free group (see Section 2.8 or [Lev05, Proposition 3.1]). Thus, we are done if  $\#\bar{V}^1 \geq 2$ .

To conclude, we claim that there is at least one vertex  $v \in \bar{V}^0$  such that  $\text{Out}(G_v, \text{Inc}_v^{(t)})$  is infinite. Let  $H^0 \subseteq H$  be the finite index subgroup of  $H$  acting trivially on the quotient graph  $U_H^1/F_N$ . Since the group of twists of  $U_H^1$  is trivial, the natural homomorphism

$$\rho : H^0 \rightarrow \prod_{v \in \bar{V}} \text{Out}(G_v, \text{Inc}_v)$$

is injective (Proposition 2.7). By Theorem 6.12(2),  $H$  acts trivially on each vertex group  $G_v$  for  $v \in \bar{V}^1$ , so  $\rho$  takes its values in  $\prod_{v \in \bar{V}^0} \text{Out}(G_v, \text{Inc}_v)$ . This also implies that  $H$  acts trivially on each edge group, and that  $\rho$  takes values in  $\prod_{v \in \bar{V}^0} \text{Out}(G_v, \text{Inc}_v^{(t)})$

(see Remark 2.8; each edge stabilizer is its own normalizer as an intersection of a fixed subgroup and a free factor). Since  $H$  is infinite, there is at least one vertex  $v \in \bar{V}^0$  such that  $\text{Out}(G_v, \text{Inc}_v^{(t)})$  is infinite.  $\square$

#### 6.4 A bounded chain condition

**Proposition 6.16.** *There is a bound, depending only on the rank  $N$  of the free group, on the length of a chain of subgroups  $H^1 \subseteq \dots \subseteq H^r$  of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  such that*

$$\text{FS}^{H^1} \supsetneq \dots \supsetneq \text{FS}^{H^r}.$$

*Proof.* Choose  $S_i \in \mathcal{D}_{H^i}$  a maximal  $H^i$ -invariant free splitting of  $F_N$ . Since for  $i \leq j$ ,  $S_j$  is  $H^i$ -invariant,  $S_i$  dominates  $S_j$ . Since there is a bound on domination chains of deformation spaces of free splittings, we may assume that the trees  $S_1, \dots, S_r$  all belong to the same deformation space of  $F_N$ -trees.

Denote  $S = S_r$ . For all  $i \leq r$ ,  $S$  is a maximal  $H^i$ -invariant free splitting. Viewing  $S$  as an  $\tilde{H}^i$ -tree, let  $U_{H^i}^1$  be the tree of cylinders of  $S$ . We fix an edge  $e \subseteq S$ , and for every  $i \leq r$ , we let  $Y_i \subseteq S$  be the cylinder of  $e$  when  $S$  is viewed as an  $\tilde{H}^i$ -tree. In other words  $Y_i$  is the union of all edges  $e' \subseteq S$  such that  $\tilde{H}_{e'}^i = \tilde{H}_e^i$ . Since for  $i \leq j$ ,  $\tilde{H}_{e'}^i = \tilde{H}_{e'}^j \cap \tilde{H}^i$ , one has  $Y_1 \supseteq Y_2 \supseteq \dots \supseteq Y_r$ .

We claim that  $Y_i$  takes boundedly many values. Indeed, by Lemma 6.9, we have  $G_{(Y_i)} = \text{Fix}_{F_N}(\tilde{H}_e^i)$ , so by the bounded chain condition of auto-fixed groups [MV04], the group  $G_{(Y_i)}$  takes boundedly many values. Since  $Y_i \cap (F_N \cdot e) = G_{(Y_i)} \cdot e$ , and since the number of edges of  $S/F_N$  is finite (in fact bounded, with a bound only depending on  $N$ ), it follows that  $Y_i$  takes boundedly many values (with a bound only depending on  $N$ ).

Using again the bound on the number of  $F_N$ -orbits of edges of  $S$ , we conclude that  $U_{H^i}^1$  takes boundedly many values, as an  $F_N$ -tree. Since  $\text{FS}^{H^i}$  can be recovered from  $U_{H^i}^1$  by Theorem 6.12(3),  $\text{FS}^{H^i}$  takes boundedly many values.  $\square$

#### 6.5 The stabilizer of a collection of free splittings

We conclude this section by applying the previous results to study the elementwise stabilizer  $\Gamma_{\mathcal{C}}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  of any collection  $\mathcal{C}$  of free splittings. We denote by  $\hat{\mathcal{C}} = \text{FS}^{\Gamma_{\mathcal{C}}}$  the collection of all  $\Gamma_{\mathcal{C}}$ -invariant free splittings. Notice that  $\mathcal{C} \subseteq \hat{\mathcal{C}}$ , and  $\Gamma_{\mathcal{C}} = \Gamma_{\hat{\mathcal{C}}}$ .

Proposition 6.16 can be reformulated as follows.

**Proposition 6.17.** *There is a bounded chain condition on the set of elementwise stabilizers (in either  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  or  $\text{Out}(F_N)$ ) of collections of free splittings.*

*Proof.* We first work in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . If  $\Gamma_{\mathcal{C}_1} \subsetneq \dots \subsetneq \Gamma_{\mathcal{C}_r}$ , then  $\text{FS}^{\Gamma_{\mathcal{C}_1}} \supsetneq \dots \supsetneq \text{FS}^{\Gamma_{\mathcal{C}_r}}$ , so Proposition 6.16 concludes.

The chain condition for elementwise stabilizers in  $\text{Out}(F_N)$  follows from the chain condition for elementwise stabilizers in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , since the length of a chain of subgroups of  $\text{Out}(F_N)$  having the same given intersection with  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is bounded by the index  $[\text{Out}(F_N) : \text{IA}_N(\mathbb{Z}/3\mathbb{Z})]$ .  $\square$

Now let  $\Gamma_{(\hat{\mathcal{C}})}$  be the subgroup of all elements of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  that globally preserve the collection of free splittings  $\hat{\mathcal{C}}$ . We note that any subgroup of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  normalizing  $\Gamma_{\mathcal{C}}$  globally preserves  $\hat{\mathcal{C}}$ , and that  $\Gamma_{(\hat{\mathcal{C}})}$  normalizes  $\Gamma_{\hat{\mathcal{C}}} = \Gamma_{\mathcal{C}}$ ; this shows that  $\Gamma_{(\hat{\mathcal{C}})}$  is precisely the normalizer of  $\Gamma_{\mathcal{C}}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ .

Our next result allows to recover  $\Gamma_{\mathcal{C}}$  and  $\Gamma_{(\hat{\mathcal{C}})}$  from  $U_{\Gamma_{\mathcal{C}}}^1$ ; before we state it, we start with a definition.

**Definition 6.18.** *The  $V^1$ -rigid stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$  is the group of all automorphisms  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  such that for every  $v \in V^1(U_{\Gamma_{\mathcal{C}}}^1)$ , there exists a lift  $\tilde{\alpha}$  of  $\alpha$  to  $\text{Aut}(F_N)$  and an  $\tilde{\alpha}$ -equivariant isometry  $I_{\tilde{\alpha}} : U_{\Gamma_{\mathcal{C}}}^1 \rightarrow U_{\Gamma_{\mathcal{C}}}^1$  acting as the identity on the star of  $v$ .*

**Proposition 6.19.** *Let  $\mathcal{C}$  be any collection of free splittings, and view  $U_{\Gamma_{\mathcal{C}}}^1$  as an  $F_N$ -tree. Let  $\Gamma_{\mathcal{C}}$  be the elementwise stabilizer of  $\mathcal{C}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , and  $\Gamma_{(\hat{\mathcal{C}})}$  be the global stabilizer of  $\hat{\mathcal{C}}$ .*

*The normalizer in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  of  $\Gamma_{\mathcal{C}}$  is  $\Gamma_{(\hat{\mathcal{C}})}$ . Moreover,*

- $\Gamma_{(\hat{\mathcal{C}})}$  is the stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ ;
- $\Gamma_{\mathcal{C}}$  is the  $V^1$ -rigid stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$ .

*Proof.* We already noticed that  $\Gamma_{(\hat{\mathcal{C}})}$  is the normalizer of  $\Gamma_{\mathcal{C}}$ .

The deformation space  $\mathcal{D}_{\Gamma_{\mathcal{C}}}$  consists of all maximal  $\Gamma_{\mathcal{C}}$ -invariant free splittings, i.e. of maximal elements of  $\hat{\mathcal{C}}$ , viewed as  $\tilde{\Gamma}_{\mathcal{C}}$ -trees. Since  $\Gamma_{\mathcal{C}}$  is normalized by  $\Gamma_{(\hat{\mathcal{C}})}$ ,  $\mathcal{D}_{\Gamma_{\mathcal{C}}}$  is invariant under  $\Gamma_{(\hat{\mathcal{C}})}$ , and the fact that all trees in  $\mathcal{D}_{\Gamma_{\mathcal{C}}}$  have the same tree of cylinders implies that  $U_{\Gamma_{\mathcal{C}}}^1$  is  $\Gamma_{(\hat{\mathcal{C}})}$ -invariant as a  $\tilde{\Gamma}_{\mathcal{C}}$ -tree, hence also as an  $F_N$ -tree.

Conversely, let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be the stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$ . By Lemma 2.13,  $H$  preserves the  $F_N$ -orbit of every vertex in  $V^1$ . Since the splittings in  $\hat{\mathcal{C}}$  are precisely those that can be read from  $V^1$  as in Theorem 6.12(3),  $H$  globally preserves  $\hat{\mathcal{C}}$ , so  $H \subseteq \Gamma_{(\hat{\mathcal{C}})}$  which proves the first assertion.

To prove the second assertion, choose a  $\tilde{\Gamma}_{\mathcal{C}}$ -tree  $S \in \mathcal{D}_{\Gamma_{\mathcal{C}}}$ , then  $U_{\Gamma_{\mathcal{C}}}^1$  is the tree of cylinders of  $S$ . Fix a vertex  $v \in V^1(U_{\Gamma_{\mathcal{C}}}^1)$ , and let  $Y \subseteq S$  be the cylinder corresponding to  $v$ . By Lemma 6.7, the group  $\Gamma_{\mathcal{C}}$  has a lift  $\tilde{\Gamma}_{\mathcal{C}}^Y$  to  $\text{Aut}(F_N)$  which acts as the identity on  $Y$ . Now, every edge  $\varepsilon$  of  $U_{\Gamma_{\mathcal{C}}}^1$  incident on  $v$  corresponds to a pair  $(Y, x)$  with  $x \in Y$ . As  $\tilde{\Gamma}_{\mathcal{C}}^Y$  acts trivially on  $Y$ , it fixes  $x$ , and therefore  $\tilde{\Gamma}_{\mathcal{C}}^Y$  fixes the edge  $\varepsilon$  in  $U_{\Gamma_{\mathcal{C}}}^1$ . It follows that  $\tilde{\Gamma}_{\mathcal{C}}^Y$  acts trivially on the star of  $v$  in  $U_{\Gamma_{\mathcal{C}}}^1$ , so  $\Gamma_{\mathcal{C}}$  is contained in the  $V^1$ -rigid stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$ .

Conversely, let  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be contained in the  $V^1$ -rigid stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$ . Let  $\tilde{\alpha}$  be a lift of  $\alpha$  to  $\text{Aut}(F_N)$ . We denote by  $I_{\tilde{\alpha}}$  the  $\tilde{\alpha}$ -equivariant isometry of  $U_{\Gamma_{\mathcal{C}}}^1$ . Let  $S \in \hat{\mathcal{C}}$ . Theorem 6.12(3) can be reformulated as follows: there is an  $F_N$ -tree  $\hat{U}$  which is a common refinement of  $S$  and  $U_{\Gamma_{\mathcal{C}}}^1$ , with a collapse map  $p : \hat{U} \rightarrow U_{\Gamma_{\mathcal{C}}}^1$  such that  $p^{-1}(v)$  is reduced to a point for all  $v \in V^0(U_{\Gamma_{\mathcal{C}}}^1)$ , and such that  $S$  is obtained from  $\hat{U}$  by collapsing every edge  $e \subseteq \hat{U}$  which is not collapsed by  $p$ .

For every vertex  $v \in V^1$ , we claim that there exists a unique element  $g(\tilde{\alpha}, v) \in F_N$  such that  $I_{\tilde{\alpha}}$  coincides with the action of  $g(\tilde{\alpha}, v)$  on the star of  $v$ . Indeed, existence follows from the assumption on  $\alpha$ . Uniqueness comes from the fact that for any two distinct vertices  $w, w' \in V^0$ , the stabilizers  $G_w$  and  $G_{w'}$  are  $F_N$ -stabilizers of distinct vertices of the same free splitting of  $F_N$  (namely  $S$ ), so  $G_w \cap G_{w'} = \{1\}$ . In particular, if  $e$  and  $e'$  are two distinct edges in  $U_{\hat{\mathcal{C}}}^1$  incident on  $v$ , then  $G_e \cap G_{e'} = \{1\}$ . This is enough to deduce the uniqueness of  $g(\tilde{\alpha}, v)$ .

For every point  $w$  in the star of  $v$  and every  $h \in F_N$ , one has  $g(\tilde{\alpha}, hv)hw = I_{\tilde{\alpha}}(hw) = \tilde{\alpha}(h)I_{\tilde{\alpha}}(w) = \tilde{\alpha}(h)g(\tilde{\alpha}, v)w$ . By uniqueness, this shows that

$$\tilde{\alpha}(h)g(\tilde{\alpha}, v) = g(\tilde{\alpha}, hv)h.$$

Let  $\hat{I}_{\tilde{\alpha}} : \hat{U} \rightarrow \hat{U}$  be the map defined in the following way:

- for every vertex  $v \in V^1$  and every point  $x \in p^{-1}(v)$ , we let  $\hat{I}_{\tilde{\alpha}}(x) := g(\tilde{\alpha}, v)x$ , and
- for every point  $y \in \hat{U}$  whose  $p$ -image does not belong to  $V^1$ , we let  $\hat{I}_{\tilde{\alpha}}(y)$  be the unique preimage of  $I_{\tilde{\alpha}}(y)$  in  $\hat{U}$ .

Then  $\hat{I}_{\tilde{\alpha}}$  is an isometry, and the above relation shows that  $\hat{I}_{\tilde{\alpha}}$  is  $\tilde{\alpha}$ -equivariant. This shows that  $\hat{U}$ , whence  $S$ , is  $\alpha$ -invariant. Since this holds for all  $S \in \hat{\mathcal{C}}$ , we conclude that  $\alpha \in \Gamma_{\mathcal{C}}$  which finishes the proof.  $\square$

## 6.6 When all splittings in $\mathcal{C}$ are fixed by the stabilizer of $U_{\Gamma_{\mathcal{C}}}^1$

The following technical results will be used to characterize compatibility of free splittings in terms of their stabilizers in Section 14.

**Lemma 6.20.** *Let  $\mathcal{C}$  be a collection of non-trivial free splittings with infinite elementwise stabilizer  $\Gamma_{\mathcal{C}}$ . Denote by  $V^0 \sqcup V^1$  the bipartition of the vertex set of  $U_{\Gamma_{\mathcal{C}}}^1$ . Assume that  $\Gamma_{\mathcal{C}}$  almost coincides with the stabilizer of  $U_{\Gamma_{\mathcal{C}}}^1$  in the following sense: for every  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  preserving  $U_{\Gamma_{\mathcal{C}}}^1$ , there exists  $k \geq 1$  such that  $\alpha^k \in \Gamma_{\mathcal{C}}$ .*

*Then for every  $v \in V^1$ ,*

1. *if  $G_v$  is non-abelian, then the Grushko decomposition of  $G_v$  relative to  $\text{Inc}_v$  is sporadic with free rank 0 (i. e. dual to an amalgam  $G_v = A * B$ );*
2. *if  $G_v$  is non-abelian, then there is no edge incident on  $v$  with trivial stabilizer;*
3. *if  $G_v$  is infinite cyclic, then there is exactly one  $G_v$ -orbit of edges with trivial stabilizer incident on  $v$ .*

*Proof.* To prove the first assertion, assume that  $G_v$  is non-abelian, and that the Grushko decomposition of  $G_v$  relative to  $\text{Inc}_v$  is non-sporadic, and argue towards a contradiction. Denote by  $\mathcal{F}_v$  the peripheral factors of this decomposition. Then consider an automorphism  $\alpha_v \in \text{Out}(G_v, \mathcal{F}_v^{(t)})$  which is fully irreducible relative to  $\mathcal{F}_v$ . In particular, no power of  $\alpha_v$  preserves a non-trivial free splitting of  $G_v$  relative to  $\mathcal{F}_v$ . Since  $\alpha_v$  acts

trivially on incident edge groups, it extends to an automorphism  $\alpha \in \text{Out}(F_N)$  preserving  $U_{\Gamma_C}^1$ . By Theorem 6.12(3) every free splitting of  $F_N$  obtained by blowing up  $v$  is  $\Gamma_C$ -invariant. This prevents that  $\alpha^k \in \Gamma_C$  for some  $k \geq 1$ .

Assume now that  $G_v$  is non-abelian and that  $G_v$  has a Grushko decomposition relative to  $\text{Inc}_v$  of the form of an HNN extension  $G_v = A*$  with all incident edge groups conjugate in  $A$ . Then there we claim that exists a free splitting  $S_v$  of  $G_v$  relative to  $A$  and an automorphism  $\alpha_v \in \text{Out}(G_v, A^{(t)})$  such that no power of  $\alpha_v$  preserves  $S_v$ . The claim then leads to a contradiction as in the first case, thus concluding the proof of the first assertion. To prove the claim, let  $t$  be a stable letter of the HNN extension, so that  $G_v = A*\langle t \rangle$ , and let  $S_v$  be the free splitting dual to this amalgamated product. Choose  $a \in A \setminus \{1\}$  and consider the automorphism  $\alpha_v$  acting as the identity on  $A$  and sending  $t$  to  $ta$ . Then no power of  $\alpha_v$  preserves  $S_v$ , which proves the claim.

To prove the second assertion, assume that  $G_v$  is non-abelian, but there is an edge  $e$  with trivial stabilizer incident on  $v$ .

Let  $a \in G_v \setminus \{1\}$ , and let  $\alpha \in \text{Out}(F_N)$  be the twist by  $a$  around  $e$  near  $v$ : this has a representative  $\tilde{\alpha} \in \text{Aut}(F_N)$  such that  $\tilde{\alpha}|_{G_v} = \text{id}$  and  $I_{\tilde{\alpha}}(e) = ae$ .

Then for all  $g \in G_v$ , one has  $I_{\tilde{\alpha}}(ge) = \tilde{\alpha}(g)I_{\tilde{\alpha}}(e) = gae$  and since  $G_v$  is not abelian,  $\alpha$  is not in the  $V^1$ -rigid stabilizer of  $U_{\Gamma_C}^1$ , and neither are its non-trivial powers, a contradiction which concludes the proof of the second assertion.

To prove the third assertion, notice first that there has to be at least one  $G_v$ -orbit of edges with trivial stabilizer incident on  $v$ , as otherwise there would be no possible blowup of  $v$  into a free splitting. The uniqueness of this orbit of edges is proved as follows: if  $e, e'$  are two edges with trivial stabilizer incident on  $v$  which are not in the same orbit under  $G_v$ , one can consider a twist  $\tilde{\alpha}$  around  $e$  near  $v$  that acts as the identity on  $G_v$ , and such that  $I_{\tilde{\alpha}}$  fixes  $e$  and maps  $e'$  to  $ae'$  for some non-trivial  $a \in G_v$ . Then  $\alpha$  and its non-trivial powers are not in the  $V^1$ -rigid stabilizer of  $U_{\Gamma_C}^1$ , which proves the third assertion.  $\square$

**Corollary 6.21.** *Under the assumptions of Lemma 6.20, then any one-edge free splitting obtained (as in Theorem 6.12(3)) from  $U_{\Gamma_C}^1$  by blowing up a vertex  $v \in V^1$  with non-trivial stabilizer, is compatible with all splittings in  $\mathcal{C}$ .*

*Proof.* Let  $S$  be a one-edge free splitting as in the statement. It suffices to prove that  $S$  is compatible with any  $\Gamma_C$ -invariant one-edge free splitting  $S'$ . By Theorem 6.12(3),  $S'$  can be obtained by from  $U_{\Gamma_C}^1$  by blowing up a vertex  $v' \in V^1$ . If  $v' \neq v$ , then  $S$  and  $S'$  are clearly compatible. If  $v' = v$ , and  $G_v$  is non-abelian, then by Assertions 1 and 2 of Lemma 6.20, there is a unique possible free splitting (obtained by blowing up  $v$  using the decomposition  $G_v = A*B$ ), so  $S = S'$ . If  $G_v$  is infinite cyclic, then up to the action of  $G_v$ , there is exactly one edge  $\varepsilon$  with trivial stabilizer incident on  $v$  (Assertion 3 of Lemma 6.20). If there is an edge with non-trivial stabilizer incident on  $v$ , then there is a unique possible free splitting, namely the one dual to  $\varepsilon$ , so again  $S = S'$ .

In the remaining case,  $v$  corresponds to a terminal vertex  $\bar{v}$  of  $U_{\Gamma_C}^1/F_N$  with infinite cyclic vertex group, and the incident edge  $\bar{\varepsilon}$  carries the trivial group. Then there are exactly two possible blow ups of  $U_{\Gamma_C}^1$  at  $v$ , namely one dual to  $\bar{\varepsilon}$ , and one obtained by

replacing  $\bar{v}$  by a circle. These two splittings are compatible, so in any case,  $S$  and  $S'$  are compatible.  $\square$

## 7 Invariant $\mathcal{Z}_{\max}$ -splittings and stabilizers of collections of $\mathcal{Z}_{\max}$ -splittings

Throughout this section, we fix an integer  $N \geq 2$ .

### 7.1 Basic facts on cyclic splittings and trees of cylinders

We recall that a  $\mathcal{Z}$ -splitting of  $F_N$  is a splitting of  $F_N$  whose edge stabilizers are either trivial or isomorphic to  $\mathbb{Z}$ . A  $\mathcal{Z}_{\max}$ -splitting is a splitting whose edge stabilizers are maximally cyclic (in particular non-trivial).

**Lemma 7.1.** *Let  $S$  be a  $\mathcal{Z}$ -splitting of  $F_N$ , and let  $A \subseteq F_N$  be a subgroup isomorphic to  $\mathbb{Z}$  which is elliptic in  $S$ .*

*Then the subtree  $Y_A \subseteq S$  made of all points fixed by  $A$  is a finite subtree.*

*Proof.* Consider an edge  $e \subseteq Y_A$ . Let  $\hat{A}$  be the maximal cyclic group containing  $A$ . We claim that  $Y_A$  contains only finitely many edges in the  $F_N$ -orbit of  $e$ . Indeed, if  $e' = ge \subseteq Y_A$ , then  $G_{e'} = gG_e g^{-1}$  and  $G_e$  and  $G_{e'}$  both contain  $A$ . It follows that  $g \in \hat{A}$ , so there are at most  $\#\hat{A}/A$  possibilities for  $e'$ . This concludes the proof since there are only finitely many orbits of edges in  $S$ .  $\square$

In the case where  $S$  is a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ , we have the following more precise version.

**Lemma 7.2.** *Let  $S$  be a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ . Then two distinct edges of  $S$  with the same stabilizer belong to different  $F_N$ -orbits.*

*Proof.* Let  $g \in F_N$ , and let  $e$  be an edge with stabilizer  $G_e$ . Then the stabilizer of  $ge$  is equal to  $gG_e g^{-1}$ . Hence either  $g$  belongs to the  $\mathcal{Z}_{\max}$  subgroup  $G_e$  and  $e = ge$ , or else  $e$  and  $ge$  have distinct stabilizers.  $\square$

For a  $\mathcal{Z}_{\max}$ -splitting  $S$  of  $F_N$ , equality among edge stabilizers is an admissible equivalence relation (which coincides with commensurability and commutation). The cylinder of an edge  $e$  is thus the union of edges of  $S$  having the same stabilizer. See Section 5.2 for the definition of admissible equivalence relations and of the tree of cylinders.

**Lemma 7.3.** *Let  $S$  be a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ , and let  $S_c$  be its tree of cylinders.*

*Then  $S_c$  is a  $\mathcal{Z}_{\max}$ -splitting in the same deformation space as  $S$ , and every edge stabilizer of  $S_c$  is an edge stabilizer of  $S$ .*

*Proof.* Lemma 7.1 shows that each cylinder  $Y$  is finite, so its global stabilizer  $G_{(Y)}$  contains  $G_e$  with finite index. Since  $G_e$  is maximal cyclic, it follows that  $G_{(Y)} = G_e$ . Since  $G_{(Y)}$  is elliptic in  $S$ , this implies that  $S$  and  $S_c$  are in the same deformation space.



Additionally, if  $\varepsilon$  is an edge of  $S_c$ , it corresponds to a pair  $(x, Y)$  with  $x \in Y$ , and  $G_\varepsilon$  contains the stabilizer of any edge  $e$  of  $Y$  incident on  $x$ , so  $G_e \subseteq G_\varepsilon \subseteq G(Y) = G_e$ . This shows that  $S_c$  is a  $\mathcal{Z}_{\max}$ -splitting and concludes the proof of the lemma.  $\square$

One can check that a  $\mathcal{Z}_{\max}$ -splitting  $S$  is its own tree of cylinders if its vertex set is bipartite into  $V_{cyc} \sqcup V_{nab}$  where  $G_v$  is not abelian for  $v \in V_{nab}$ , and for all  $v \in V_{cyc}$ ,  $G_v$  is maximal cyclic and the set of fixed points of  $G_v$  is precisely the star of  $v$  in  $S$ .

## 7.2 Statement of the result

Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and assume that there is no non-trivial  $H$ -invariant free splitting of  $F_N$ . Let  $\mathcal{Z}_{\max}S^H$  be the collection of all  $H$ -invariant non-trivial  $\mathcal{Z}_{\max}$ -splittings of  $F_N$ . Out of this collection of splittings we are going to construct a JSJ decomposition that will be a canonical splitting associated to  $H$ .

If  $S, S'$  are two splittings of a group  $A$ , we say that  $S$  is *elliptic* with respect to  $S'$  if all its edge stabilizers are elliptic in  $S'$ . If  $\mathcal{T}$  is a collection of splittings of a group  $A$ , we say that a subgroup  $B \subseteq A$  is  $\mathcal{T}$ -*universally elliptic* if  $B$  is elliptic in all splittings in  $\mathcal{T}$ . We say that a splitting of  $A$  is  $\mathcal{T}$ -*universally elliptic* if all its edge stabilizers are  $\mathcal{T}$ -universally elliptic.

**Theorem 7.4** (Canonical JSJ decomposition for  $H$ -invariant  $\mathcal{Z}_{\max}$ -splittings). *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, and assume that there is no non-trivial  $H$ -invariant free splitting of  $F_N$ .*

*There exists a unique  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting  $U_H^z$  with the following properties:*

- (1)  $U_H^z$  is  $\mathcal{Z}_{\max}S^H$ -universally elliptic;
- (2)  $U_H^z$  dominates every  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting which is  $\mathcal{Z}_{\max}S^H$ -universally elliptic;
- (3)  $U_H^z$  is its own tree of cylinders.

Moreover,  $U_H^z$  is non-trivial if  $\mathcal{Z}_{\max}S^H \neq \emptyset$ .

*Remark 7.5.* The assumption that  $H$  preserves no non-trivial free splitting implies that  $H$  is non-trivial, i.e. infinite.

To prove the existence of this JSJ decomposition, one has to work consistently with  $H$ -invariant splittings, which we interpret as  $\tilde{H}$ -splittings, where  $\tilde{H}$  is the full preimage of  $H$  in  $\text{Aut}(F_N)$  (see Section 5.1). So in fact, we will construct  $U_H^z$  as a JSJ decomposition of  $\tilde{H}$  over a suitable collection of edge groups. Proving the existence and uniqueness of  $U_H^z$  is actually easy, and done in Section 7.3 below; the hard part of the work is to prove the non-triviality of  $U_H^z$  in Section 7.4.

### 7.3 Uniqueness and existence

Let us start with the proof of the uniqueness of  $U_H^z$  which is almost immediate.

*Uniqueness of  $U_H^z$ .* Let  $U_1, U_2$  be two  $H$ -invariant  $\mathcal{Z}_{\max}$ -splittings as in Theorem 7.4. We view  $U_1$  and  $U_2$  as  $F_N$ -trees. Combining Assertions (1) and (2) shows that  $U_1$  and  $U_2$  are in the same deformation space. Since they are their own trees of cylinders, this implies that  $U_1 = U_2$  ([GL11, Theorem 1]).  $\square$

**Lemma 7.6.** *Let  $S$  be an  $H$ -invariant  $\mathcal{Z}$ -splitting of  $F_N$ . Let  $\tilde{B}$  be a subgroup of  $\tilde{H}$  such that  $\tilde{B} \cap F_N$  is non-trivial and fixes a point in  $S$ .*

*Then  $\tilde{B}$  fixes a point in  $S$ .*

*Proof.* Since  $A := \tilde{B} \cap F_N$  is normal in  $\tilde{B}$ , the subtree  $Y_A \subseteq S$  consisting of all points fixed by  $A$  is  $\tilde{B}$ -invariant. If  $A$  is not cyclic, then  $Y_A$  is reduced to a point because edge stabilizers are cyclic. Otherwise,  $Y_A$  has finite diameter by Lemma 7.1, and we deduce that  $\tilde{B}$  fixes the circumcenter of  $Y_A$ .  $\square$

**Lemma 7.7.** *Let  $S, S' \in \mathcal{Z}_{\max}S^H$ . The tree  $S$  is elliptic with respect to  $S'$  as an  $F_N$ -tree if and only if  $S$  is elliptic with respect to  $S'$  as an  $\tilde{H}$ -tree.*

*Proof.* Given an edge  $e \subseteq S$ , we denote by  $G_e$  the  $F_N$ -stabilizer of  $e$ , and by  $\tilde{H}_e$  its  $\tilde{H}$ -stabilizer. Since  $G_e \subseteq \tilde{H}_e$ , if  $S$  is elliptic with respect to  $S'$  as an  $\tilde{H}$ -tree, it is also elliptic with respect to  $S'$  as an  $F_N$ -tree.

Conversely, assume that  $S$  is elliptic with respect to  $S'$  as an  $F_N$ -tree, and fix an edge  $e$  of  $S$ . Since  $G_e = \tilde{H}_e \cap F_N$  is non-trivial and elliptic in  $S'$ , Lemma 7.6 shows that  $\tilde{H}_e$  fixes a point in  $S'$ . This shows that  $S$  is elliptic with respect to  $S'$  as an  $\tilde{H}$ -tree.  $\square$

**Corollary 7.8.** *Let  $S, S' \in \mathcal{Z}_{\max}S^H$ , and assume that  $S$  is elliptic with respect to  $S'$ .*

*Then there exists an  $\tilde{H}$ -tree  $\hat{S}$  which is a blowup of  $S$  and dominates  $S'$ , and such that each edge stabilizer of  $\hat{S}$  fixes an edge in  $S$  or in  $S'$ .*

*Moreover, as an  $F_N$ -tree,  $\hat{S}$  is a ( $H$ -invariant)  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ .*

*Proof.* View  $S, S'$  as  $\tilde{H}$ -trees. Since  $S$  is elliptic with respect to  $S'$  as an  $\tilde{H}$ -tree by Lemma 7.7, one can apply [GL17, Proposition 2.2] to construct an  $\tilde{H}$ -tree  $\hat{S}$  that refines  $S$  and dominates  $S'$  and whose edge stabilizers fix an edge in  $S$  or  $S'$  (as  $\tilde{H}$ -trees, therefore also as  $F_N$ -trees). Thus,  $\hat{S}$  can be viewed as an  $H$ -invariant  $F_N$ -tree, and  $\hat{S}$  is still a  $\mathcal{Z}_{\max}$ -splitting by [GL17, § 9.5].  $\square$

We can now prove the existence of  $U_H^z$ .

*Existence of  $U_H^z$ .* Note that the trivial splitting is  $\mathcal{Z}_{\max}S^H$ -universally elliptic. By Lemma 2.1, there is a bound on the number of orbits of edges of a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$  without vertex of valence 2. Therefore, there exists an  $H$ -invariant  $\mathcal{Z}_{\max}S^H$ -universally elliptic  $\mathcal{Z}_{\max}$ -splitting  $U$  such that no proper refinement of  $U$  satisfies this property. We claim that  $U$  satisfies Assertion (2). Indeed, consider  $U' \in \mathcal{Z}_{\max}S^H$  any  $\mathcal{Z}_{\max}S^H$ -universally elliptic splitting. Apply Corollary 7.8 to construct an  $\tilde{H}$ -tree  $\hat{U}$  that refines  $U$  and dominates

$U'$ . Since edge stabilizers of  $\hat{U}$  fix an edge in  $U$  or  $U'$ ,  $\hat{U}$  is  $\mathcal{Z}_{\max}S^H$ -universally elliptic. By maximality of  $U$ , we thus have  $\hat{U} = U$ , showing that  $U$  dominates  $U'$ . This proves that  $U$  satisfies Assertions (1) and (2).

Now let  $U_H^z$  be the tree of cylinders of  $U$ , viewed as a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ . By Lemma 7.3,  $U_H^z$  is a  $\mathcal{Z}_{\max}$ -splitting in the same deformation space as  $U$ , and edge stabilizers of  $U_H^z$  are edge stabilizers of  $U$ , so  $U_H^z$  is  $\mathcal{Z}_{\max}S^H$ -universally elliptic. It follows that  $U_H^z$  satisfies (1), (2) and (3).  $\square$

#### 7.4 Description and non-triviality of $U_H^z$

The goal of the present section is to prove the non-triviality of  $U_H^z$  when  $H$  has no invariant non-trivial free splittings and  $\mathcal{Z}_{\max}S^H \neq \emptyset$  (Theorem 7.15 below). This relies on arguments coming from JSJ theory, the key tool being to understand *flexible* vertices of  $U_H^z$ , defined as follows, and show that they come from QH situations (all relevant definitions are given right below).

**Definition 7.9.** *A vertex  $v$  of  $U_H^z$  is flexible if its  $F_N$ -stabilizer  $G_v$  is not  $\mathcal{Z}_{\max}S^H$ -universally elliptic.*




Figure 5: A socket graph of groups with 3 proper sockets and 2 improper ones.

A *socket graph of groups*  $\Lambda$  is a minimal tree of groups whose underlying graph is a star over a central vertex  $v$ , where  $v$  is a clean QH vertex (see Definition 2.5) and the other vertex groups are cyclic (see [Sel97]). Equivalently, denoting by  $u_1, \dots, u_r$  the vertices of  $\Gamma \setminus \{v\}$ , and by  $e_i$  the edge joining  $u_i$  to  $v$ ,

- the group  $G_v = \pi_1(\Sigma)$  is identified to the fundamental group of a connected compact hyperbolic surface  $\Sigma$  with  $n \geq r$  boundary components, each edge group  $G_{e_i}$  being conjugate in  $\pi_1(\Sigma)$  to  $B_i$ , where  $B_i$  is the fundamental group of the  $i$ -th boundary component of  $\Sigma$ ;
- each vertex group  $G_{u_i}$  is cyclic and  $[G_{u_i} : G_{e_i}] \geq 2$ .

For  $i \leq n$ , let  $C_i$  be the maximal cyclic subgroup of  $\pi_1(\Lambda)$  containing  $B_i$ . The conjugates of the groups  $C_i$  are called the *sockets*. A socket is *improper* if  $C_i = B_i$ . A socket is *Möbius* if  $[C_i : B_i] = 2$ .

*Remark 7.10.* If  $\Gamma$  is a socket graph of groups, then the associated  $\mathcal{Z}_{\max}$ -splitting  $\Gamma_{\mathcal{Z}_{\max}}$  defined in Definition 2.2 is trivial. The following lemma gives a converse.

**Lemma 7.11.** *Let  $S$  be a splitting of  $F_N$  with infinite cyclic edge stabilizers. Assume that  $S$  has at least one QH vertex with trivial fiber, and that every edge is adjacent to at least one vertex with non-abelian stabilizer.*

*If the associated  $\mathcal{Z}_{\max}$ -splitting  $S_{\mathcal{Z}_{\max}}$  (Definition 2.2) is trivial, then  $S/F_N$  is a socket graph of groups.*

*Proof.* By Lemma 2.3,  $S/F_N$  is a tree of groups which is a star with all vertex groups cyclic except the central one. The central vertex  $v$  is QH with trivial fiber by assumption. There remains to prove that  $v$  is clean QH. First, the underlying orbifold  $\Sigma$  is a surface since  $F_N$  is torsion-free and the fiber is trivial. Second, no two edges of  $S/F_N$  are incident on the same boundary component of  $\Sigma$ , as otherwise a generator of its fundamental group would have two distinct roots in  $F_N$ , which is not possible. For the same reason, each incident edge group is a maximal boundary subgroup of  $\pi_1(\Sigma)$ .  $\square$

**Definition 7.12.** *Let  $A$  be a finitely generated group, and let  $\mathcal{P}$  be a finite collection of subgroups of  $A$ . The pair  $(A, \mathcal{P})$  is weakly QH with sockets if  $A$  splits as a socket graph of groups  $A \simeq \pi_1(\Lambda)$  and  $\mathcal{P}$  is a subset of the conjugacy classes of the socket groups.*

*It is QH with sockets if there exists a socket graph of groups as above where additionally  $\mathcal{P}$  contains all conjugacy classes of improper sockets. Such a decomposition of  $A$  is called a socket decomposition of  $(A, \mathcal{P})$ .*

*Remark 7.13.* If  $(A, \mathcal{P})$  is weakly QH with sockets but not QH with sockets, then there is a free splitting of  $A$  relative to  $\mathcal{P}$  dual to a properly embedded arc in the underlying surface, whose endpoints are on a boundary component corresponding to an improper socket not appearing in  $\mathcal{P}$ . Thus, if  $(A, \mathcal{P})$  is weakly QH with sockets and if there is no free splitting of  $A$  relative to  $\mathcal{P}$ , then  $(A, \mathcal{P})$  is in fact QH with sockets.

*Remark 7.14.* If  $(A, \mathcal{P})$  is QH with sockets, then one can choose a socket decomposition of  $(A, \mathcal{P})$  such that all the Möbius sockets appear in  $\mathcal{P}$ . Indeed, starting with any socket decomposition  $\Lambda$  of  $(A, \mathcal{P})$  with underlying surface  $\Sigma$ , one can glue a Möbius band on  $\Sigma$  for each Möbius socket not appearing in  $\mathcal{P}$  and remove the corresponding edge of  $\Lambda$ . This decomposition of  $(A, \mathcal{P})$  is its cyclic JSJ decomposition by [DG11, Proposition 4.24]. In particular, it is unique.

In the following statement, we view  $U_H^z$  as an  $F_N$ -tree. Given a vertex  $v \in U_H^z$ , we denote by  $\text{Inc}_v$  the set of all  $F_N$ -stabilizers of edges that are incident on  $v$ .

We denote by  $H|_{G_v} \subseteq \text{Out}(G_v)$  the subgroup obtained by restriction of outer automorphisms in  $H$ . More precisely, since  $U_H^z$  is  $H$ -invariant and  $H$  acts trivially on the quotient graph  $U_H^z/F_N$  (Proposition 2.15), the conjugacy class of  $G_v$  is  $H$ -invariant. Thus for every  $\tilde{\alpha} \in H$ , there is a representative  $\tilde{\alpha} \in \text{Aut}(F_N)$  such that  $\tilde{\alpha}(G_v) = G_v$ , and the image of  $\tilde{\alpha}|_{G_v}$  in  $\text{Out}(G_v)$  does not depend on the choice of  $\tilde{\alpha}$  because  $G_v$  is its own normalizer, we denote it by  $\alpha|_{G_v}$ . The map  $\alpha \mapsto \alpha|_{G_v}$  defines a morphism  $H \rightarrow \text{Out}(G_v)$  whose image we denote by  $H|_{G_v}$ . An element  $\alpha$  lies in its kernel if it has a lift  $\tilde{\alpha} \in \text{Aut}(F_N)$  such that  $\tilde{\alpha}|_{G_v} = \text{id}$ ; we then say that  $\alpha$  acts trivially on  $G_v$ .

**Theorem 7.15.** *Assume that  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  (is infinite and) preserves no non-trivial free splitting, and that  $\mathcal{Z}_{\max}S^H \neq \emptyset$ . Then  $U_H^z$  is non-trivial.*

*Moreover, if  $v$  is a flexible vertex of  $U_H^z$ , then  $(G_v, \text{Inc}_v)$  is QH with sockets, and  $H|_{G_v} = \{1\}$ .*

*Remark 7.16.* The second assertion implies the first. Indeed, if  $U_H^z$  is reduced to a single point  $v$ , then  $v$  is flexible because there exists a non-trivial  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting (we assume  $\mathcal{Z}_{\max}S^H \neq \emptyset$ ). Since  $G_v = F_N$ ,  $H|_{G_v} = H$  so  $H$  is trivial by the second assertion, a contradiction.

**Lemma 7.17.** *Let  $v$  be a flexible vertex of  $U_H^z$ , and let  $\mathcal{T}$  be the set of all non-trivial  $\mathcal{Z}_{\max}$ -splittings of  $G_v$  relative to  $\text{Inc}_v$  which are  $H|_{G_v}$ -invariant. Then*

- $G_v$  has no non-trivial  $H|_{G_v}$ -invariant free splitting relative to  $\text{Inc}_v$ ;
- $\mathcal{T} \neq \emptyset$  i. e.  $G_v$  has some non-trivial  $H|_{G_v}$ -invariant  $\mathcal{Z}_{\max}$ -splitting relative to  $\text{Inc}_v$ ;
- no splitting in  $\mathcal{T}$  is  $\mathcal{T}$ -universally elliptic.

*Proof.* We claim that any  $H|_{G_v}$ -invariant cyclic splitting  $S$  of  $G_v$  relative to  $\text{Inc}_v$  can be used to blow up  $U_H^z$  into an  $H$ -invariant cyclic splitting  $\hat{U}$  of  $F_N$ . Indeed, view  $U_H^z$  as an  $\tilde{H}$ -tree, and let  $\tilde{H}_v$  be the stabilizer of  $v$  in  $\tilde{H}$ . Since  $S$  is  $H|_{G_v}$ -invariant, it can be viewed as a  $\tilde{K}$ -tree where  $\tilde{K} \subseteq \text{Aut}(G_v)$  is the preimage of  $H|_{G_v}$  in  $\text{Aut}(G_v)$ . We note that  $\tilde{K}$  coincides with the image of the natural restriction map  $r : \tilde{H}_v \rightarrow \text{Aut}(G_v)$ . The map  $r : \tilde{H}_v \rightarrow \tilde{K}$  allows to view the  $\tilde{K}$ -tree  $S$  as an  $\tilde{H}_v$ -tree. There remains to check that the  $\tilde{H}$ -stabilizer  $\tilde{H}_e$  of each edge  $e$  incident on  $v$  fixes a point in  $S$ . Since  $S$  is relative to  $\text{Inc}_v$ , the  $F_N$ -stabilizer  $G_e$  of  $e$  fixes a point in  $S$ . By Lemma 7.6 (applied to  $\tilde{K} \curvearrowright S$  and  $\tilde{B} = r(\tilde{H}_e)$ ),  $\tilde{H}_e$  is elliptic in  $S$  which concludes the proof of the claim.

The claim implies the first assertion: any non-trivial  $H|_{G_v}$ -invariant free splitting of  $G_v$  relative to  $\text{Inc}_v$  would induce an  $H$ -invariant splitting  $\hat{U}$  of  $F_N$  having an edge with trivial  $F_N$ -stabilizer, contradicting the assumption that  $H$  preserves no non-trivial free splitting.

Since  $G_v$  is flexible, there exists an  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting  $S$  of  $F_N$  in which  $G_v$  is not elliptic. Let  $S_v \subseteq S$  be the minimal  $G_v$ -invariant subtree, with its action by  $G_v$ . Then  $S_v$  is an  $H|_{G_v}$ -invariant  $\mathcal{Z}_{\max}$ -splitting of  $G_v$ , and since  $U_H^z$  is universally elliptic, the groups in  $\text{Inc}_v$  fix a point in  $S$  hence in  $S_v$ . This proves the second assertion.

Finally, let  $S \in \mathcal{T}$ ; we aim to prove that  $S$  is not  $\mathcal{T}$ -universally elliptic. Let  $\hat{U}$  be an  $H$ -invariant blowup of  $U_H^z$  given by the claim above, using  $S$  to blow up  $U_H^z$  at  $v$ . Note that  $G_v$  is root closed in  $H$  because  $U_H^z$  is a  $\mathcal{Z}_{\max}$ -splitting. Edge stabilizers of  $\hat{U}$  are  $\mathcal{Z}_{\max}$  because  $S$  is a  $\mathcal{Z}_{\max}$ -splitting and  $G_v$  is root closed. Since  $U_H^z$  does not dominate  $\hat{U}$ ,  $\hat{U}$  is not  $\mathcal{Z}_{\max}S^H$ -universally elliptic so one can find some edge  $e \subseteq S$  whose stabilizer  $G_e \subseteq G_v$  is not elliptic in some  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting  $T$  of  $F_N$ . Let  $T_v \subseteq T$  be the minimal  $G_v$ -invariant tree. The action  $G_v \curvearrowright T_v$  is an  $H|_{G_v}$ -invariant  $\mathcal{Z}_{\max}$ -splitting of  $G_v$  relative to  $\text{Inc}_v$ . Thus this is a splitting in  $\mathcal{T}$  in which  $G_e$  is not elliptic. In particular  $S$  is not  $\mathcal{T}$ -universally elliptic.  $\square$

Theorem 7.15 is an easy consequence of the following result.

**Proposition 7.18.** *Let  $G$  be a non-cyclic finitely generated free group,  $\mathcal{P}$  be a finite family of maximal cyclic subgroups of  $G$ , and  $K \subseteq \text{Out}(G, \mathcal{P})$ . Let  $\mathcal{T}$  be the set of all non-trivial minimal  $\mathcal{Z}_{\max}$ -splittings of  $G$  relative to  $\mathcal{P}$  which are  $K$ -invariant. Assume that*

- $G$  has no non-trivial  $K$ -invariant free splitting relative to  $\mathcal{P}$ ;
- $\mathcal{T} \neq \emptyset$ ;
- no splitting in  $\mathcal{T}$  is  $\mathcal{T}$ -universally elliptic.

Then  $(G, \mathcal{P})$  is weakly QH with sockets and  $K$  is finite.

*Proof of Theorem 7.15 from Proposition 7.18.* As noted in Remark 7.16, it suffices to prove the second assertion. So consider a flexible vertex  $v$  of  $U_H^z$ . Notice that  $G_v$  is a finitely generated free group (being a point stabilizer in a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ ), and is not cyclic (because being flexible, it has a non-trivial splitting relative to incident edge groups).

We apply Proposition 7.18 with  $G = G_v$ ,  $\mathcal{P} = \text{Inc}_v$  and  $K = H|_{G_v}$ . Lemma 7.17 shows that the hypotheses of Proposition 7.18 are satisfied, so  $(G_v, \text{Inc}_v)$  is weakly QH with sockets and  $H|_{G_v}$  is finite.

We now deduce that  $H|_{G_v}$  is trivial using that  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Fix any  $\alpha \in H|_{G_v}$ , and consider  $\tilde{\alpha} \in \tilde{H}$  preserving  $G_v$  and inducing  $\alpha$  in  $\text{Out}(G_v)$ . By [KSS06, Theorem B(3)], generic elements  $u \in G_v$  satisfy that their stabilizer in  $\text{Aut}(G_v)$  is generated by  $\text{ad}_u$  (the inner automorphism associated to  $u$ ). Choose such an element  $u \in G_v$  not conjugate in an incident edge group. Since  $H|_{G_v}$  is finite, the  $H$ -orbit of the  $F_N$ -conjugacy class of  $u$  is finite. Since  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , it follows from [HM20, Theorem 4.1] that the  $F_N$ -conjugacy class of  $u$  is fixed by  $H$ . Then  $\tilde{\alpha}(u) \in G_v$  is conjugate to  $u$  in  $F_N$ , and since  $u$  fixes no edge,  $\tilde{\alpha}(u)$  is conjugate to  $u$  in  $G_v$ . Up to composing  $\tilde{\alpha}$  by the conjugation by an element in  $G_v$ , we may assume that  $\tilde{\alpha}(u) = u$ . By choice of  $u$ ,  $\tilde{\alpha}|_{G_v}$  is inner, i.e.  $\alpha$  is trivial. We thus conclude that  $H|_{G_v} = \{1\}$ .

Finally, Lemma 7.17 ensures that  $G_v$  has no non-trivial free splitting relative to  $\text{Inc}_v$ , so  $(G_v, \text{Inc}_v)$  is QH with sockets (and not only weakly QH with sockets; see Remark 7.13).  $\square$

*Proof of Proposition 7.18.* Let  $\tilde{K}$  be the full preimage of  $K$  under the quotient map  $\text{Aut}(G) \rightarrow \text{Out}(G)$ . We will see  $K$ -invariant splittings of  $G$  as splittings of  $\tilde{K}$ .

We are going to follow [GL17, §6] giving the description of flexible vertices of the JSJ decomposition of the group  $\tilde{K}$  relative to  $\mathcal{P}$  over a well chosen class of allowed edge groups. We cannot readily apply the results because we want the extra information that  $K$  is finite, and we do not know a priori that  $\tilde{K}$  is finitely presented.

If  $\mathcal{A}$  is a collection of subgroups of  $\tilde{K}$ , an  $(\mathcal{A}, \mathcal{P})$ -tree  $S$  is an action of  $\tilde{K}$  on a tree with edge stabilizers in  $\mathcal{A}$  and in which groups in  $\mathcal{P}$  are elliptic. We will work with the following collections of subgroups. We identify  $G$  with  $\text{Inn}(G)$  so that  $G \triangleleft \tilde{K}$ . Let  $\mathcal{A}_{\max}$  (respectively  $\mathcal{A}_{\mathbb{Z}}$ ,  $\mathcal{A}_{\text{cyc}}$ ) be the collection of subgroups  $E \subseteq \tilde{K}$  such that  $E \cap G$  is

maximal cyclic (respectively isomorphic to  $\mathbb{Z}$ , respectively cyclic (maybe trivial)). For instance, trees in  $\mathcal{T}$  correspond to  $(\mathcal{A}_{max}, \mathcal{P})$ -trees.

If  $E \subseteq \tilde{K}$  and  $S$  is a splitting of  $\tilde{K}$ , we say that  $E$  is *slender in  $S$*  if it preserves a line or a point of  $S$ . Let  $\mathcal{A}_{sl}$  the collection of subgroups  $E \subseteq \tilde{K}$  such that  $E \in \mathcal{A}_{cyc}$  and  $E$  is slender in any  $(\mathcal{A}_{cyc}, \mathcal{P})$ -tree. Unlike the other two classes,  $\mathcal{A}_{sl}$  and  $\mathcal{A}_{cyc}$  are stable under taking subgroups.

We have the following inclusions

$$\mathcal{A}_{max} \subseteq \mathcal{A}_{\mathbb{Z}} \subseteq \mathcal{A}_{sl} \subseteq \mathcal{A}_{cyc}.$$

We explain the second inclusion, the two others being obvious. Let  $E \in \mathcal{A}_{\mathbb{Z}}$ , and let  $C = E \cap G \simeq \mathbb{Z}$ , a normal subgroup of  $E$ . If  $T$  is an  $(\mathcal{A}_{cyc}, \mathcal{P})$ -tree in which  $C$  is not elliptic, then  $C$  has an axis  $A_C$  in  $T$ , and  $A_C$  is  $E$ -invariant. If  $C$  is elliptic in  $T$ , then its set of fixed points is bounded (Lemma 7.1) so  $E$  is also elliptic in  $T$ . This shows that  $E$  is slender in  $(\mathcal{A}_{cyc}, \mathcal{P})$ -trees, so  $\mathcal{A}_{\mathbb{Z}} \subseteq \mathcal{A}_{sl}$ .

Since there is no non-trivial  $K$ -invariant free splitting of  $G$  relative to  $\mathcal{P}$ , any  $(\mathcal{A}_{cyc}, \mathcal{P})$ -tree is in fact an  $(\mathcal{A}_{\mathbb{Z}}, \mathcal{P})$ -tree. Thus,  $(\mathcal{A}_{\mathbb{Z}}, \mathcal{P})$ -trees,  $(\mathcal{A}_{sl}, \mathcal{P})$ -trees, and  $(\mathcal{A}_{cyc}, \mathcal{P})$ -trees are the same objects: we denote by  $\mathcal{T}' \supseteq \mathcal{T}$  this set of actions of  $\tilde{K}$  on trees.

**Claim 7.19.** *The  $\tilde{K}$ -action on every non-trivial tree  $S \in \mathcal{T}'$  is faithful.*

*More generally, let  $S \in \mathcal{T}'$ ,  $A \subseteq G$  not elliptic in  $S$ , and  $\tilde{N} \subseteq \tilde{K}$  acting as the identity on an  $A$ -invariant subtree  $S' \subseteq S$ . Then for all  $\tilde{\alpha} \in \tilde{N}$  and all  $a \in A$ , one has  $\tilde{\alpha}(a) = a$ .*

*Proof.* One deduces the first assertion from the second remembering that  $\tilde{K} \subseteq \text{Aut}(G)$  by taking  $A = G$ , and  $\tilde{N}$  the kernel of the action  $\tilde{K} \curvearrowright S$ .

In general, for  $\tilde{\alpha} \in \tilde{N}$ , we denote by  $I_{\tilde{\alpha}}$  the isometry of  $S$  determined by the action of  $\tilde{\alpha}$ . Fix  $\tilde{\alpha} \in \tilde{N}$  and  $a \in A$ . Then for all  $x \in S'$ , we have  $a.x = I_{\tilde{\alpha}}(a.x) = \tilde{\alpha}(a).I_{\tilde{\alpha}}(x) = \tilde{\alpha}(a).x$ , so  $a^{-1}\tilde{\alpha}(a)$  is an element of  $G$  acting as the identity on  $S'$ . Since  $S'$  is unbounded, Lemma 7.1 implies that  $\tilde{\alpha}(a) = a$ .  $\square$

We claim that  $\mathcal{A}_{sl}$  satisfies the stability condition (SC) of [GL17, Definition 6.1] with  $\mathcal{F} = \mathcal{A}_{sl}$  as a class of fibers. Recall that this means that for every short exact sequence of groups

$$1 \rightarrow F \rightarrow B \rightarrow Q \rightarrow 1$$

with  $B \subseteq \tilde{K}$ , the following two conditions hold:

1. if  $B \in \mathcal{A}_{sl}$  and  $Q$  is isomorphic to either  $\mathbb{Z}$  or the infinite dihedral group  $D_{\infty}$ , then  $F \in \mathcal{A}_{sl}$ , and
2. if  $F \in \mathcal{A}_{sl}$  and  $Q$  is isomorphic to a quotient of  $\mathbb{Z}$  or  $D_{\infty}$ , then  $B \in \mathcal{A}_{sl}$ .

Indeed, the first condition is obvious since  $\mathcal{A}_{sl}$  is stable under taking subgroups. To check the second condition, assume that  $F \in \mathcal{A}_{sl}$ , and that  $B$  is an extension of  $F$  as above. Then  $B \cap G$  is an extension of the cyclic group  $F \cap G$  by a cyclic or dihedral group, so  $B \cap G$  is cyclic, i.e.  $B \in \mathcal{A}_{cyc}$ . To check the slenderness condition, let  $S \in \mathcal{T}'$ .



Since  $F \in \mathcal{A}_{sl}$ , either  $F$  preserves a unique line  $l$  in  $S$ , or  $F$  fixes a point. In the first case,  $l$  is  $B$ -invariant because  $F$  is normal in  $B$ , so  $B$  is slender in  $S$ . In the second case, let  $Y \subseteq S$  be the subtree consisting of all points fixed by  $F$ . Then  $Y$  is  $B$ -invariant and the action of  $B$  factors through an action of  $Q$ . Since  $Q$  is cyclic or dihedral, it preserves a line or fixes a point in  $Y$ , so  $B$  is slender in  $S$ . This concludes the proof that  $\mathcal{A}_{sl}$  satisfies the stability condition (SC).

We recall some more definitions from [GL17, Definition 6.9], which generalize a usual assumption asking that  $G$  does not split over a subgroup commensurable with a subgroup of infinite index in an allowed edge group. Given two subgroups  $A, B \in \mathcal{A}_{sl}$ , we write  $A \ll B$  if  $A \subseteq B$  and there exists a tree  $S \in \mathcal{T}'$  such that  $A$  is elliptic in  $S$  while  $B$  is not. A subgroup  $B \in \mathcal{A}_{sl}$  is *minuscule* if whenever  $A \ll B$  and  $A$  fixes an edge  $e$  in some tree in  $\mathcal{T}'$ , then  $A$  has infinite index in the  $\tilde{K}$ -stabilizer  $\tilde{K}_e$  of  $e$ . A tree  $S \in \mathcal{T}'$  is *minuscule* if all edge stabilizers of  $S$  are minuscule.

**Claim 7.20.** *Every tree  $S \in \mathcal{T}'$  is minuscule.*

*Proof.* Let  $e \subseteq S$  be an edge. We assume that there exists  $A \ll \tilde{K}_e$ , with  $A$  of finite index in the stabilizer  $\tilde{K}_{e'}$  of an edge  $e'$  of another tree  $S' \in \mathcal{T}'$ , and aim for a contradiction. As noted above,  $S'$  is an  $\mathcal{A}_{\mathbb{Z}}$ -tree so  $A \cap G \simeq \mathbb{Z}$ . Let  $C$  be the maximal cyclic subgroup of  $G$  that contains  $A \cap G$ . The group  $G_e = \tilde{K}_e \cap G$  is cyclic, of finite index in  $C$ , and therefore  $\tilde{K}_e$  normalizes  $C$ . As  $A \ll \tilde{K}_e$ , there exists a tree  $S'' \in \mathcal{T}'$  such that  $A$  is elliptic in  $S''$  while  $\tilde{K}_e$  is not. Then  $A \cap G$ , whence  $C$ , is elliptic in  $S''$ . By Lemma 7.6,  $\tilde{K}_e$  fixes a point in  $S''$ , a contradiction.  $\square$

We are now going to construct the underlying surface following the approach by Fujiwara and Papsoglu [FP06] as implemented in [GL17, §6]. We will use the general definition of QH vertices of [GL17, Definition 5.13] recalled in Definition 2.6: if  $S$  is a  $\tilde{K}$ -tree,  $v \in S$  is a QH vertex with fiber  $F$  relative to  $\mathcal{P}$  if  $F \triangleleft G_v$  and there is an identification  $G_v/F = \pi_1(\Sigma)$  with the fundamental group of a hyperbolic 2-orbifold with boundary so that incident edge stabilizers and groups in  $\mathcal{P}$  intersect  $G_v$  into subgroups that map in finite or boundary subgroups of  $\pi_1(\Sigma)$ .

Let now  $U$  be a tree in  $\mathcal{T}$  whose number of orbits of edges is maximal. Our hypothesis that no tree in  $\mathcal{T}$  is  $\mathcal{T}$ -universally elliptic applies to every one-edge collapse of  $U$ . Since all trees are minuscule, we can apply [GL17, Proposition 6.28]<sup>3</sup> saying that there exists a tree  $V \in \mathcal{T}'$  such that  $U$  and  $V$  are fully hyperbolic with respect to each other (i.e. every edge stabilizer in one of these trees is hyperbolic in the other tree). We mention that  $U$  and  $V$  are then also fully hyperbolic with respect to each other as  $G$ -trees (this follows from Lemma 7.6 as in the proof of Lemma 7.7).

We will now consider the asymmetric core of  $U$  and  $V$ , as defined by Fujiwara and Papsoglu in [FP06] (see [GL17, Definition 6.16]). Notice that the core  $\mathcal{C}_{\tilde{K}}(U, V)$  of  $U$  and  $V$  viewed as  $\tilde{K}$ -trees is the same as the core  $\mathcal{C}_G(U, V)$  as  $G$ -trees; we simply denote

<sup>3</sup>The assumption that  $G$  is totally flexible in [GL17, Proposition 6.28] is only used to ensure that each edge stabilizer of  $U$  is hyperbolic in some  $\tilde{K}$ -tree with edge groups in  $\mathcal{A}_{sl}$ .

it by  $\mathcal{C}(U, V)$ . Indeed, given an edge  $e \subseteq U$  with  $G$ -stabilizer  $G_e$ , the axis of  $G_e$  in  $V$  is  $\tilde{K}_e$ -invariant, so  $\tilde{K}_e$  and  $G_e$  have the same axis, and similarly, the minimal subtrees of  $G_v$  and  $\tilde{K}_v$  in  $V$  agree for every vertex  $v \in U$ .

As  $U$  and  $V$  are minuscule and fully hyperbolic with respect to each other, the core  $\mathcal{C}(U, V)$  is symmetric [GL17, Proposition 6.19], and actually coincides with the core defined by the first author in [Gui05], see [Gui05, Proposition 12.1]. We denote by  $V(\mathcal{C})$  the set of vertices of  $\mathcal{C}(U, V)$  and by  $\text{Cut}(\mathcal{C}) \subseteq V(\mathcal{C})$  the set of cut vertices. By [GL17, Lemma 6.18], each connected component  $Z$  of  $\mathcal{C}(U, V) \setminus V(\mathcal{C})$  is a connected surface, tessellated by squares with missing vertices.

Let  $R$  be the regular neighborhood of  $U$  and  $V$  as defined in [GL17, Definition 6.23] following [FP06]. This is the bipartite simplicial tree defined as follows. The vertex set of  $R$  is  $V_{\text{cut}} \sqcup V_{\mathcal{S}}$ , having one vertex  $v_x \in V_{\text{cut}}$  for every cut vertex  $x \in \text{Cut}(\mathcal{C})$ , one vertex  $v_Z \in V_{\mathcal{S}}$  for every connected component  $Z$  of  $\mathcal{C}(U, V) \setminus \text{Cut}(\mathcal{C})$ , with an edge joining  $v_x$  to  $v_Z$  whenever  $x \in \bar{Z}$ . By [GL17, Proposition 6.25 (RN1)],  $v_Z$  is a QH vertex of  $R$  relative to  $\mathcal{P}$ , with fiber  $F \in \mathcal{A}_{sl}$ . If we view  $U, V$  as  $G$ -trees, then their regular neighborhood as  $G$ -trees is nothing but the tree  $R$  where one restricts the action to  $G \subseteq \tilde{K}$ . Each vertex  $v_Z$  of  $R$ , viewed as a  $G$ -tree, is also a QH vertex relative to  $\mathcal{P}$ , with fiber  $F' = F \cap G$ . In fact, the fiber  $F'$  is trivial because it is at most cyclic and normal in  $G_{v_Z}$  which cannot be cyclic. Additionally, since  $G$  is torsion-free and the fiber is trivial, the underlying orbifold is a surface (we do not claim that this is the case when viewing  $R$  as a  $\tilde{K}$ -tree).

We first assume that  $R$  is trivial (as a  $\tilde{K}$ -tree or equivalently as a  $G$ -tree), i.e. that  $\text{Cut}(\mathcal{C}) = \emptyset$  and  $Z = \mathcal{C}(U, V) \setminus V(\mathcal{C})$  is connected. Then by [GL17, Proposition 6.25 (RN1)],  $(\tilde{K}, \mathcal{P})$  is QH with fiber  $F \in \mathcal{A}_{sl}$ , and  $(G, \mathcal{P})$  is QH with trivial fiber. Since  $\mathcal{P}$  consists of maximal cyclic subgroups,  $(G, \mathcal{P})$  is QH with sockets (all sockets being improper). We claim that  $K$  is finite, i.e. that  $G$  has finite index in  $\tilde{K}$ . Indeed, the index of  $G$  in  $\tilde{K}$  is bounded by the number of  $G$ -orbits of squares in  $\mathcal{C}(U, V)$ : if there exists a square  $C \in \mathcal{C}(U, V)$  such that  $kC = gC$  with  $k \in \tilde{K}$  and  $g \in G$ , then  $k^{-1}g$  fixes  $C$ , and since  $Z = \mathcal{C}(U, V) \setminus V(\mathcal{C})$  is a connected tessellated surface, this implies that  $k^{-1}g$  acts as the identity on the core. Since the natural projection from  $\mathcal{C}(U, V)$  to  $U$  is onto,  $k^{-1}g$  acts as the identity on  $U$  so  $k^{-1}g = \text{id}$  by Claim 7.19.

Now we do not assume that  $R$  is trivial any more. We are going to prove that the  $\mathcal{Z}_{\text{max}}$ -splitting  $R_{\mathcal{Z}_{\text{max}}}$  of  $G$  associated to  $R$  is trivial (see Definition 2.2 for a definition of  $R_{\mathcal{Z}_{\text{max}}}$ ). We thus view  $R$  as a  $K$ -invariant cyclic splitting of  $G$  relative to  $\mathcal{P}$ . Then  $R_{\mathcal{Z}_{\text{max}}}$  is a  $K$ -invariant  $\mathcal{Z}_{\text{max}}$ -splitting of  $G$ , i.e.  $R_{\mathcal{Z}_{\text{max}}} \in \mathcal{T}$ . By [GL17, Proposition 6.25 (RN3)], the tree  $U$  is dual to a family of curves in QH-vertices of  $R$ . Hence  $R_{\mathcal{Z}_{\text{max}}}$  is compatible with  $U$ . Let  $S$  be a common refinement of  $U$  and  $R_{\mathcal{Z}_{\text{max}}}$  in  $\mathcal{T}$ . Since  $U$  has the maximal possible number of orbits of edges among trees in  $\mathcal{T}$ , it follows that  $U = S$ . Assuming by contradiction that  $R_{\mathcal{Z}_{\text{max}}}$  is not reduced to a point, let  $e$  be an edge of  $U = S$  not collapsed in  $R_{\mathcal{Z}_{\text{max}}}$ . Since  $U$  is fully hyperbolic with respect to  $V$ , the stabilizer of  $e$  is hyperbolic in  $V$ . On the other hand, edge stabilizers of  $R$  (hence of  $R_{\mathcal{Z}_{\text{max}}}$ ) are elliptic in  $U$  and  $V$  by [GL17, Proposition 6.25 (RN2)], a contradiction.

This shows that  $R_{\mathcal{Z}_{\text{max}}}$  is trivial. Viewing it as a  $G$ -tree, this implies by Lemma 7.11 that  $R/G$  is a socket graph of groups.

We now prove that  $K$  is finite. Let  $v \in R$  be a QH vertex and  $Z$  the corresponding component of  $\mathcal{C}(U, V) \setminus \text{Cut}(\mathcal{C})$ . By Proposition 2.7, up to replacing  $K$  with a finite index subgroup, one has an extension

$$1 \rightarrow K \cap \text{Tw} \rightarrow K \xrightarrow{\rho} K|_{G_v} \rightarrow 1$$

where  $\text{Tw}$  is the group of twists, and  $K|_{G_v}$  is the image of  $K$  in  $\prod_{u \in R/F_N} \text{Out}(G_u, \text{Inc}_u) = \text{Out}(G_v, \text{Inc}_v)$  under the restriction map  $\rho$ . Since  $R/G$  is a socket graph of groups, its group of twists is trivial so it suffices to prove that  $K|_{G_v}$  is finite. The argument is similar to the one above. Since  $Z \setminus V(\mathcal{C})$  is a connected tessellated surface, the  $\tilde{K}$ -stabilizer  $\tilde{N}$  of a square  $C$  in  $Z$  acts as the identity on  $Z$ . By Claim 7.19, for every element  $\tilde{\alpha} \in \tilde{N}$ , one has  $\tilde{\alpha}|_{G_v} = \text{id}_{G_v}$ . Since  $\tilde{K}_v$  permutes the  $G_v$ -orbits of squares, for  $\tilde{\alpha}$  in finite index subgroup  $\tilde{K}_0 \subseteq \tilde{K}_v$ , there exists  $g \in G$  such that  $\tilde{\alpha}C = gC$ , so  $\text{ad}_g^{-1}\tilde{\alpha} \in \tilde{N}$ . This shows that all elements in a finite index subgroup of  $K$  act trivially on  $G_v$  and concludes the proof of Proposition 7.18.  $\square$

## 7.5 Characterization of invariant $\mathcal{Z}_{\max}$ -splittings

Recall that an outer automorphism  $\alpha \in \text{Out}(F_N)$  is said to act trivially on a subgroup  $A \subseteq F_N$  if it has a representative  $\tilde{\alpha}$  in  $\text{Aut}(F_N)$  such that  $\tilde{\alpha}|_A = \text{id}_A$ .

**Proposition 7.21.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup with no invariant non-trivial free splitting. Then  $U_H^z$  is compatible with every  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ . More precisely, any  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting of  $F_N$  may be obtained from  $U_H^z$  by blowing up each flexible vertex  $v$  into a (minimal, maybe trivial)  $\mathcal{Z}_{\max}$ -splitting  $Y_v$  relative to  $\text{Inc}_v$ , and by blowing up each vertex  $v$  with cyclic stabilizer into a finite tree on which  $G_v$  acts as the identity, and then by collapsing all the edges coming from  $U_H^z$ .*

*Conversely, let  $S$  be any  $\mathcal{Z}_{\max}$ -splitting of  $F_N$  obtained by blowing up  $U_H^z$  in this way. Then  $S$  is  $H$ -invariant. More generally,  $S$  is invariant under any automorphism  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  that preserves  $U_H^z$  and that acts trivially on each flexible vertex group of  $U_H^z$ .*

*Remark 7.22.* The fact that any  $\alpha \in H$  acts trivially on stabilizers of flexible vertices of  $U_H^z$  is a consequence of Theorem 7.15. See Proposition 7.27 below for the precise relationship between the group of all such automorphisms and the stabilizer of  $U_H^z$ .

In order to prove Proposition 7.21, we will use the following fact.

**Lemma 7.23.** *Let  $(A, \mathcal{P})$  be QH with sockets. Let  $\Lambda$  be a socket decomposition of  $(A, \mathcal{P})$  such that every Möbius socket appears in  $\mathcal{P}$ . Let  $Y_A$  be a  $\mathcal{Z}_{\max}$ -splitting of  $(A, \mathcal{P})$ .*

*Then  $Y_A$  is dual to a family of non-boundary parallel simple closed curves on the underlying surface of  $\Lambda$  and every group in  $\mathcal{P}$  fixes a unique point in  $Y_A$ .*

*Proof.* Let  $v$  be the QH vertex of  $\Lambda$  and denote by  $\tilde{\Lambda}$  the Bass–Serre tree of  $\Lambda$ .

Let  $Y_v \subseteq Y_A$  be the minimal  $G_v$ -invariant subtree of  $Y_A$ . The groups in  $\mathcal{P}$  and the edge groups of  $\Lambda$  are elliptic in  $Y_A$  because  $\Lambda$  is the cyclic JSJ decomposition of  $(A, \mathcal{P})$  by

[DG11, Proposition 4.24]. Since all improper sockets appear in  $\mathcal{P}$  by Definition 7.12, the fundamental group of each boundary component of  $\Sigma$  is elliptic in  $Y_A$ , so  $Y_v$  is dual to a collection of disjoint simple closed curves on  $\Sigma$  not parallel to the boundary. Observe in particular that for each edge  $e$  of  $\tilde{\Lambda}$  incident on  $v$ ,  $G_e$  and any finite index subgroup of  $G_e$  fixes a unique point  $p_e$  in  $Y_v$ . We also observe for future use that if  $gv \neq v$  and  $e_1, e_2$  are two edges in  $Y_v$  and  $gY_v$  respectively, then  $G_{e_1}$  and  $G_{e_2}$  do not commute: if they did, then  $G_{e_1} \cap G_{e_2}$  would fix the segment  $[v, gv]$  in  $\tilde{\Lambda}$ , hence some edge incident on  $v$ , contradicting the previous observation.

Let  $S$  be the blow up of  $\tilde{\Lambda}$  obtained by replacing  $v$  by  $Y_v$  and gluing back each incident edge  $e$  to the point  $p_e \in Y_v$ .

The inclusion  $Y_v \hookrightarrow Y_A$  extends to an equivariant map  $f : S \rightarrow Y_A$  which can be chosen linear on edges. Let  $e = uv$  be an edge of  $\tilde{\Lambda}$  incident on  $v$ , and let  $\tilde{e} = up_e$  be the corresponding edge in  $S$ . Note that  $G_u$  is a conjugate of some group in  $\mathcal{P}$ . Since edge stabilizers of  $Y_A$  are  $\mathcal{Z}_{\max}$ ,  $f$  maps the translates of  $\tilde{e}$  by  $G_u$  to the same segment  $[f(u), p_e] \subseteq Y_A$ , and one may redefine  $f$  to map  $\tilde{e}$  to the point  $p_e$ . Thus  $f$  factors through the tree  $S'$  obtained from  $S$  by collapsing all edges coming from  $\tilde{\Lambda}$ . Since  $f$  is isometric in restriction to each  $Y_v$ , and  $f$  cannot fold two edges in distinct translates of  $Y_v$  by the observation above,  $S' \simeq Y_A$ . Since no edge in  $Y_v$  has a stabilizer conjugate to some group in  $\mathcal{P}$ , this concludes the proof.  $\square$

*Proof of Proposition 7.21.* Let  $S$  be any  $H$ -invariant  $\mathcal{Z}_{\max}$ -splitting.

To each vertex  $v \in U_H^z$  we associate in an equivariant way a subtree  $Y_v \subseteq S$  as follows. If  $G_v$  is not cyclic, we define  $Y_v \subseteq S$  as the minimal  $G_v$ -invariant subtree of  $S$  (if  $G_v$  fixes a point, this point is unique because  $G_v$  is not cyclic). Since flexible vertices of  $U_H^z$  are QH with sockets (Theorem 7.15), Lemma 7.23 applies to  $G_v \curvearrowright Y_v$  for  $v$  flexible (with  $\mathcal{P} = \text{Inc}_v$ ), and in particular, the stabilizer  $G_v \cap G_e$  of every edge  $e \subseteq Y_v$  is non-trivial and fixes no edge in  $U_H^z$ .

If  $G_v$  is cyclic, then we define  $Y_v \subseteq S$  as the set of points fixed by  $G_v$ ; this is non-empty because edges of  $U_H^z$  incident on  $v$  have stabilizer  $G_v$  and are  $\mathcal{Z}_{\max}S^H$ -universally elliptic.

We note that in all cases, if  $e$  is an edge in  $Y_v$ , then the stabilizer of  $e$  for the action  $G_v \curvearrowright Y_v$  is equal to its stabilizer  $G_e$  for the  $G$ -action on  $S$ , in other words  $G_e \cap G_v = G_e$ : this is because  $G_v$  is root-closed and  $G_v \cap G_e$  is infinite cyclic.

We claim that if  $u, v \in U_H^z$  are joined by an edge, then  $Y_u \cap Y_v \neq \emptyset$ . Since  $U_H^z$  is a tree of cylinders, we may assume that  $G_u$  is cyclic and  $G_v$  is not, and since edge stabilizers of  $U_H^z$  are maximal cyclic,  $G_u \subseteq G_v$ . Since  $Y_v$  is a  $G_u$ -invariant tree and  $G_u$  is elliptic in  $S$ ,  $G_u$  fixes a point in  $Y_v$  which proves the claim.

We next claim that if  $v_1 \neq v_2$  are distinct vertices of  $U_H^z$ , then  $Y_{v_1} \cap Y_{v_2}$  contains at most one point. Indeed, assuming by contradiction that there is an edge  $e$  in  $Y_{v_1} \cap Y_{v_2}$ , then  $G_e \subseteq G_{v_1} \cap G_{v_2}$  so  $G_e$  fixes all the edges in the segment joining  $v_1$  to  $v_2$  in  $U_H^z$ . As noticed above, this prevents  $v_1$  and  $v_2$  from being flexible vertices. Since  $Y_{v_1}, Y_{v_2}$  are not reduced to a point, this implies that  $v_1$  and  $v_2$  are vertices with cyclic stabilizer. Since  $G_{v_1} \cap G_{v_2} \neq \{1\}$ , and since  $U_H^z$  is a tree of cylinders, this implies  $v_1 = v_2$ , a contradiction.

Now we construct a blowup  $\hat{U}$  of  $U_H^z$  as follows: we start from the disjoint union of the trees  $Y_v$  (where  $v$  ranges over the set of vertices of  $U_H^z$ ) and the disjoint union of the edges of  $U_H^z$ , and for each edge  $e = uv$  of  $U_H^z$ , we let  $p$  be the unique point in  $Y_u \cap Y_v$  and we glue the endpoints of  $e$  to the copy of  $p$  in  $Y_u$  and  $Y_v$  respectively. We claim that collapsing all edges of  $\hat{U}$  coming from  $U_H^z$  gives an  $F_N$ -tree  $\bar{U}$  isomorphic to  $S$ . Indeed, the inclusions  $Y_v \subseteq S$  extend to an equivariant map  $\hat{U} \rightarrow S$  which maps each edge coming from  $U_H^z$  to a point, hence factors through  $\bar{U}$ . Since  $Y_u \cap Y_v$  contains at most one point for  $u \neq v$ , the induced map  $\bar{U} \rightarrow S$  is injective, which proves the first assertion of Proposition 7.21.

We now prove the converse. Let  $\hat{U}$  be obtained by blowing up  $U_H^z$  by replacing in a  $F_N$ -equivariant way each flexible vertex  $v \in U_H^z$  by a minimal action  $G_v \curvearrowright Y_v$  relative to  $\text{Inc}_v$ , each vertex  $u$  with cyclic stabilizer by a finite tree  $Y_u$  on which  $G_u$  acts as the identity, and choosing for each edge  $\varepsilon = uv$  attaching points  $p_\varepsilon \in Y_u, q_\varepsilon \in Y_v$  fixed by  $G_\varepsilon$ . It suffices to check that  $\hat{U}$  is  $H$ -invariant, and more generally, invariant under any automorphism  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  preserving  $U_H^z$  and acting trivially on flexible vertex groups of  $U_H^z$ .

Let  $\tilde{\alpha}$  be a preimage of  $\alpha$  in  $\text{Aut}(G)$ . We have an  $\tilde{\alpha}$ -equivariant isometry  $I_{\tilde{\alpha}} : U_H^z \rightarrow U_H^z$ , and we now define a lift  $J_{\tilde{\alpha}} : \hat{U} \rightarrow \hat{U}$ . For each edge  $\varepsilon \subseteq \hat{U}$  coming from  $U_H^z$ , we define  $J_{\tilde{\alpha}}(\varepsilon) = I_{\tilde{\alpha}}(\varepsilon)$ . We now define  $J_{\tilde{\alpha}}$  on each tree  $Y_v$ . If  $v$  is rigid with non-cyclic stabilizer,  $Y_v$  is a point and we define  $J_{\tilde{\alpha}}(Y_v) = Y_{J_{\tilde{\alpha}}(v)}$ . If  $v$  is flexible, we know that there exists  $g_v \in F_N$  such that  $\text{ad}_{g_v}^{-1} \circ \tilde{\alpha}$  restricts to the identity on  $G_v$ . Since  $U_H^z$  is a tree of cylinders, there is no pair of edges incident on  $v$  with the same stabilizer. It follows that  $g_v^{-1} I_{\tilde{\alpha}}$  is the identity on the star of  $v$ . We then define  $J_{\tilde{\alpha}}(x) = g_v \cdot x$  for all  $x \in Y_v$ , and this is compatible with the attaching points. Now consider  $u \in U_H^z$  with cyclic stabilizer. Recall that as  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , Proposition 2.15 shows that  $\tilde{\alpha}$  acts as the identity on  $U_H^z/F_N$ . Therefore, there exists a (non-unique)  $g_u \in F_N$  such that  $\tilde{\alpha}$  coincides with  $\text{ad}_{g_u}$  in restriction to  $G_u$ , and we define  $J_{\tilde{\alpha}}(x) = g_u \cdot x$  for all  $x \in Y_u$ . If  $g'_u$  is another choice for  $g_u$ , then  $g_u^{-1} g'_u$  centralizes  $G_u$  so lies in  $G_u$ , so  $g_u \cdot x = g'_u \cdot x$  for all  $x \in Y_u$  since  $G_u$  acts as the identity on  $Y_u$ . To check that this is compatible with the attaching data, consider  $\varepsilon = uv$  an edge of  $U_H^z$  with origin  $u$ . Then  $I_{\tilde{\alpha}}(\varepsilon) = g_\varepsilon \varepsilon$  for some  $g_\varepsilon \in F_N$  because  $\tilde{\alpha}$  acts trivially on  $U_H^z/F_N$ , so  $g_\varepsilon u = g_u u$  and  $g_\varepsilon^{-1} g_u \in G_u$  fixes the attaching point  $p_\varepsilon \in Y_u$ .

This shows that  $J_{\tilde{\alpha}}$  induces an  $\tilde{\alpha}$ -equivariant isometry  $\hat{U} \rightarrow \hat{U}$  and concludes the proof.  $\square$

## 7.6 Bounded chain condition

**Proposition 7.24.** *There exists a bound, depending only on the rank of the free group  $F_N$ , on the length of any chain*

$$H_1 \subseteq H_2 \subseteq \cdots \subseteq H_n$$

*of subgroups of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  such that  $H_1$  (hence every  $H_i$ ) fixes no non-trivial free splitting of  $F_N$ , and such that*

$$\mathcal{Z}_{\max} S^{H_1} \supsetneq \mathcal{Z}_{\max} S^{H_2} \supsetneq \cdots \supsetneq \mathcal{Z}_{\max} S^{H_n}.$$

*Proof.* Denote  $U_i = U_{H_i}^z$ . For  $j \geq i$ ,  $U_j$  is an  $H_i$ -invariant  $\mathcal{Z}_{\max}$ -splitting, so  $U_i$  is compatible with  $U_j$  by Proposition 7.21. By [GL17, Proposition A.17], all the splittings  $U_1, \dots, U_n$  have a common refinement  $\hat{U}$ , i.e. each  $U_i$  can be obtained from  $\hat{U}$  by collapsing some set of edges. Since there is a bound on the number of edges of a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ , this gives a bound on the possible number of distinct trees  $U_i$ . If  $U_i = U_j$  and if the set of flexible vertices of  $U_i$  agrees with the set of flexible vertices of  $U_j$ , then  $\mathcal{Z}_{\max}S^{H_i} = \mathcal{Z}_{\max}S^{H_j}$  by Proposition 7.21. The proposition follows.  $\square$

## 7.7 The stabilizer of a collection of $\mathcal{Z}_{\max}$ -splittings

We conclude this section by applying the previous results to study the elementwise stabilizer  $\Gamma_{\mathcal{C}}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  of a collection  $\mathcal{C}$  of  $\mathcal{Z}_{\max}$ -splittings, assuming that this stabilizer does not preserve a non-trivial free splitting. We denote by  $\hat{\mathcal{C}} = \mathcal{Z}_{\max}S^{\Gamma_{\mathcal{C}}}$  the collection of all  $\Gamma_{\mathcal{C}}$ -invariant  $\mathcal{Z}_{\max}$ -splittings of  $F_N$ . Notice that  $\mathcal{C} \subseteq \hat{\mathcal{C}}$ , and  $\Gamma_{\mathcal{C}} = \Gamma_{\hat{\mathcal{C}}}$ .

Proposition 7.24 may be reformulated as follows:

**Proposition 7.25.** *There exists a bound, depending only on the rank of the free group  $F_N$ , on the length of any chain*

$$\mathcal{C}_1 \subseteq \dots \subseteq \mathcal{C}_n$$

*of collections of  $\mathcal{Z}_{\max}$ -splittings of  $F_N$  such that no  $\Gamma_{\mathcal{C}_i}$  fixes a non-trivial free splitting of  $F_N$ , and*

$$\Gamma_{\mathcal{C}_1} \supsetneq \dots \supsetneq \Gamma_{\mathcal{C}_n}.$$

*Proof.* We apply Proposition 7.24 with  $H_i = \Gamma_{\mathcal{C}_i}$ . Denote by  $\hat{\mathcal{C}}_i = \mathcal{Z}_{\max}S^{H_i}$ . If  $n$  is too large, then there exists  $i_1 < i_2$  such that  $\hat{\mathcal{C}}_{i_1} = \hat{\mathcal{C}}_{i_2}$ . Since  $\Gamma_{\mathcal{C}_i} = \Gamma_{\hat{\mathcal{C}}_i}$  for all  $i \leq n$ , we get  $\Gamma_{\mathcal{C}_{i_1}} = \Gamma_{\mathcal{C}_{i_2}}$ , a contradiction.  $\square$

*Remark 7.26.* As in Proposition 6.17, the same conclusion holds if one replaces  $\Gamma_{\mathcal{C}_i}$  (i.e. the elementwise stabilizer of  $\mathcal{C}_i$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ ) by the elementwise stabilizer of  $\mathcal{C}_i$  in  $\text{Out}(F_N)$ .

As in Section 6, we denote by  $\Gamma_{(\hat{\mathcal{C}})}$  the subgroup of all elements of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  that globally preserve the collection of free splittings  $\hat{\mathcal{C}}$ , and we note that  $\Gamma_{(\hat{\mathcal{C}})}$  is the normalizer of  $\Gamma_{\mathcal{C}}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ .

The following result allows to recover  $\Gamma_{\mathcal{C}}$  and  $\Gamma_{(\hat{\mathcal{C}})}$  from  $U_{\Gamma_{\mathcal{C}}}^z$ .

**Proposition 7.27.** *Let  $\mathcal{C}$  be any collection of  $\mathcal{Z}_{\max}$ -splittings, and assume that  $\Gamma_{\mathcal{C}}$  fixes no non-trivial free splitting. Then*

- $\Gamma_{(\hat{\mathcal{C}})}$  is the stabilizer of  $U_{\Gamma_{\mathcal{C}}}^z$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ ;
- $\Gamma_{\mathcal{C}}$  is the set of automorphisms  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  that stabilize  $U_{\Gamma_{\mathcal{C}}}^z$  and act trivially on each flexible vertex group.

*Remark 7.28.* Note that  $\alpha$  acts trivially on a flexible vertex stabilizer  $G_v$  of  $U_{\Gamma_{\mathcal{C}}}^z$  if and only if  $\alpha$  has a lift acting as the identity on the star of  $v$  (compare Definition 6.18). This is because two distinct edges incident on  $v$  have distinct stabilizer.



*Proof.* Since  $\Gamma_{(\hat{c})}$  preserves the collection of all  $\Gamma_C$ -invariant  $\mathcal{Z}_{\max}$ -splittings, the uniqueness statement in Theorem 7.4 ensures that  $\Gamma_{(\hat{c})}$  preserves  $U_{\Gamma_C}^z$ . Conversely, if  $\alpha \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  preserves  $U_{\Gamma_C}^z$ , then  $\alpha$  induces the identity on the quotient graph by Proposition 2.15. In particular, it preserves the collection of flexible vertices, so by Proposition 7.21, it preserves the collection of all  $\Gamma_C$ -invariant  $\mathcal{Z}_{\max}$ -splittings, i.e.  $\alpha \in \Gamma_{(\hat{c})}$ . This proves the first assertion.

Consider the group  $H' \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  consisting of outer automorphisms that preserve  $U_{\Gamma_C}^z$  and act trivially on each flexible vertex group. Theorem 7.15 shows that  $\Gamma_C \subseteq H'$ . Conversely, Proposition 7.21 says that every  $\Gamma_C$ -invariant splitting can be read from  $U_{\Gamma_C}^z$  and that every such splitting is  $H'$ -invariant. This implies that  $H' \subseteq \Gamma_C$ .  $\square$

## 8 Pure subgroups of $\text{Out}(F_N)$ and dynamical decomposition

Throughout this section, we fix an integer  $N \geq 2$ .

### 8.1 Almost free factor systems

Recall from Definition 2.5 that a vertex  $v$  in a graph of groups is a clean QH vertex if  $G_v = \pi_1(\Sigma)$  and incident edge groups correspond bijectively to the fundamental groups of a subset of the boundary components of  $\Sigma$ .

**Definition 8.1.** A QH splitting of  $F_N$  is a cyclic splitting  $S$  of  $F_N$  such that the vertices of the quotient graph of groups  $S/F_N$  are  $v, u_1, \dots, u_n$  (maybe  $n = 0$ ) where  $v$  is a clean QH vertex identified with the fundamental group of a surface  $\Sigma$  with at least one unused boundary component, and every edge joins  $v$  to some  $u_i$ .

Its factor system is  $\hat{\mathcal{F}} = \{[G_{u_1}], \dots, [G_{u_n}], [\langle b_1 \rangle], \dots, [\langle b_r \rangle]\}$  where  $[\langle b_1 \rangle], \dots, [\langle b_r \rangle]$  are the conjugacy classes of the fundamental groups of unused boundary components.

The conjugates of the groups  $\langle b_1 \rangle, \dots, \langle b_r \rangle$  are called the unused boundary subgroups.

An extracted free factor system is a collection of the form  $\hat{\mathcal{F}} \setminus \{[\langle b_i \rangle]\}$  for some  $i \leq r$ .




Figure 6: A QH splitting with 2 unused boundary components. Its factor system is  $\hat{\mathcal{F}} = \{[G_{u_1}], [G_{u_2}], [G_{u_3}], [\langle b_1 \rangle], [\langle b_2 \rangle]\}$ . The two extracted free factor systems are  $\{[G_{u_1}], [G_{u_2}], [G_{u_3}], [\langle b_1 \rangle]\}$  and  $\{[G_{u_1}], [G_{u_2}], [G_{u_3}], [\langle b_2 \rangle]\}$ .



The quotient graph  $S/F_N$  is not assumed to be a tree, there may be several edges joining  $u_i$  to  $v$ . Note that  $F_N$  is one-ended relative to  $\hat{\mathcal{F}}$  because the fundamental group of a surface is one-ended relative to its boundary components.

*Remark 8.2.* Each extracted free factor system is indeed a free factor system of  $F_N$ . This is shown by considering a maximal system of disjoint pairwise non-isotopic properly embedded arcs with both endpoints on the boundary component of  $\Sigma$  whose fundamental group is  $\langle b_i \rangle$ : such an arc system yields a free splitting of  $F_N$  in which the elliptic subgroups are precisely given by the collection  $\hat{\mathcal{F}} \setminus \{\langle b_i \rangle\}$ .

In particular, if the number  $r$  of unused boundary components is at least 2, all the factors in  $\hat{\mathcal{F}}$  are free factors of  $F_N$  (although the factor system  $\hat{\mathcal{F}}$ , as a collection, is never a free factor system of  $F_N$ , see Lemma 8.3 and Remark 8.4 below). When  $r = 1$ , it may happen that the unused boundary subgroup is not a free factor of  $F_N$ .

**Lemma 8.3.** *Let  $S$  be a QH splitting of  $F_N$ , and  $\hat{\mathcal{F}}$  be its factor system. Then  $F_N$  is one-ended relative to  $\hat{\mathcal{F}}$ , and  $S$  is a JSJ decomposition of  $F_N$  relative to  $\hat{\mathcal{F}}$ .*

*Proof.* One-endedness follows from the fact that  $\pi_1(\Sigma)$  is one-ended relative to its boundary subgroups. The splitting  $S$  is universally elliptic because all its edge groups are conjugate in  $\hat{\mathcal{F}}$ , hence elliptic in any cyclic splitting of  $F_N$  relative to  $\hat{\mathcal{F}}$ . Maximality follows from the fact that any cyclic splitting of  $\pi_1(\Sigma)$  relative to its boundary components is dual to a multicurve on  $\Sigma$ , and each non-peripheral curve on  $\Sigma$  is intersected non-trivially by another one. This shows that  $\pi_1(\Sigma)$  is elliptic in any JSJ decomposition (see also [GL17, Corollary 5.30]) so  $S$  itself is a JSJ decomposition.  $\square$

*Remark 8.4.* One-endedness implies that  $\hat{\mathcal{F}}$  cannot be a free factor system. The lemma also implies that  $\hat{\mathcal{F}}$  determines  $S$  uniquely up to deformation, but one can easily deduce that it actually determines  $S$  up to equivariant isomorphism: the tree of cylinders  $S_c$  of  $S$  is determined by  $\hat{\mathcal{F}}$  up to equivariant isomorphism and  $S$  is obtained from  $S_c$  by collapsing the edges of  $S$  not adjacent to the QH vertex.

The following corollary implies that boundary subgroups and extracted free factor systems are permuted by the stabilizer of  $\hat{\mathcal{F}}$ .

**Corollary 8.5.** *The notions of boundary subgroups, unused boundary subgroups, and extracted free factor systems depend only of  $\hat{\mathcal{F}}$ .*

*Proof.* All JSJ decompositions are in the same deformation space so have the same non-cyclic vertex groups [GL07a, Corollary 4.4]. The group  $\pi_1(\Sigma)$  is thus characterized as the unique vertex group in any JSJ decomposition that is not  $\hat{\mathcal{F}}$ -peripheral. Boundary subgroups of  $\pi_1(\Sigma)$  are characterized as the maximal cyclic subgroups of  $\pi_1(\Sigma)$  which are  $\hat{\mathcal{F}}$ -peripheral. Unused boundary subgroups are characterized as those that do not fix an edge in  $S$ , hence in any reduced JSJ decomposition of  $F_N$  relative to  $\hat{\mathcal{F}}$  [GL07a, Proposition 4.6]. Extracted free factor systems are obtained from  $\hat{\mathcal{F}}$  by removing any unused boundary subgroup.  $\square$

**Definition 8.6** (Almost free factor systems). An almost free factor system  $\hat{\mathcal{F}}$  of  $F_N$  is a collection of conjugacy classes of subgroups of  $F_N$  which is either a free factor system or the factor system of a QH splitting of  $F_N$ .

A free factor system  $\mathcal{F}$  is extracted from  $\hat{\mathcal{F}}$  if  $\hat{\mathcal{F}}$  is a free factor system and  $\mathcal{F} = \hat{\mathcal{F}}$  or if  $\hat{\mathcal{F}}$  the factor system of a QH splitting  $S$  of  $F_N$  and  $\mathcal{F}$  is an extracted free factor system.

*Remark 8.7.* Any almost free factor system  $\hat{\mathcal{F}}$  is a *malnormal family*: for each  $[P], [Q] \in \hat{\mathcal{F}}$  and every  $g \in F_N$ , if  $P^g \cap Q \neq \{1\}$  then  $Q = P^g$  and  $g \in P$ .

**Lemma 8.8.** Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup that does not preserve any non-trivial free splitting.

- (1). For any maximal  $H$ -invariant free factor system  $\mathcal{F}$ , there exists a unique maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}$  such that  $\mathcal{F} \preceq \hat{\mathcal{F}}$ . Moreover  $\mathcal{F}$  is extracted from  $\hat{\mathcal{F}}$ .
- (2). If  $\hat{\mathcal{F}}$  is any maximal  $H$ -invariant almost free factor system, then any free factor system  $\mathcal{F}$  extracted from  $\hat{\mathcal{F}}$  is a maximal  $H$ -invariant free factor system.
- (3). If  $\mathcal{F}$  and  $\hat{\mathcal{F}}$  are as in (1) or (2), then every cyclic group whose conjugacy class is  $H$ -invariant is  $\hat{\mathcal{F}}$ -peripheral, and  $H$  contains an element which is fully irreducible relative to  $\mathcal{F}$ , and atoroidal relative to  $\hat{\mathcal{F}}$ .

*Remark 8.9.* Notice that as  $H$  does not preserve any non-trivial free splitting, every  $H$ -invariant free factor system is non-sporadic.

*Proof of Lemma 8.8.* We first prove (1) and (3) assuming that every  $H$ -periodic (or equivalently,  $H$ -invariant) conjugacy class is  $\mathcal{F}$ -peripheral. We claim that every  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}$  such that  $\mathcal{F} \preceq \hat{\mathcal{F}}$  is a free factor system. This claim will imply that  $\mathcal{F} = \hat{\mathcal{F}}$ , so (1) holds and (3) follows from [GH19b, Theorem 2].

To prove the claim, assume towards a contradiction that  $\hat{\mathcal{F}}$  is not a free factor system. Consider a QH splitting defining  $\hat{\mathcal{F}}$ , and let  $\Sigma$  be the underlying surface. By Corollary 8.5, every boundary subgroup of  $\pi_1(\Sigma)$  is  $H$ -periodic, hence  $\mathcal{F}$ -peripheral by assumption. Since  $\pi_1(\Sigma)$  is freely indecomposable relative to its boundary components,  $\pi_1(\Sigma)$  is  $\mathcal{F}$ -peripheral hence  $\hat{\mathcal{F}}$ -peripheral, a contradiction.

We now consider the case where there exists a maximal cyclic subgroup  $\langle b \rangle$  which is not  $\mathcal{F}$ -peripheral and whose conjugacy class is  $H$ -periodic. We claim that such a conjugacy class is unique. Indeed, by [GH19b, Theorem 1], there exists an element  $\alpha \in H$  which is fully irreducible relative to  $\mathcal{F}$ . Proposition 2.4 ensures that there exists an  $\alpha$ -invariant tree  $T$  which is arational relative to  $\mathcal{F}$  and in which  $b$  is elliptic. It follows that  $T$  is an arational surface tree [Rey12, Hor14], and, up to conjugacy,  $\langle b \rangle$  is the unique point stabilizer of  $T$  which is not  $\mathcal{F}$ -peripheral. In particular,  $\mathcal{F} \cup \{\langle b \rangle\}$  is an almost free factor system, and we have proved that  $\alpha$  is fully irreducible relative to  $\mathcal{F}$  and atoroidal relative to  $\mathcal{F} \cup \{\langle b \rangle\}$ .

We now check that  $\hat{\mathcal{F}} = \mathcal{F} \cup \{\langle b \rangle\}$  is the unique maximal  $H$ -invariant almost free factor system relative to which  $\mathcal{F}$  is peripheral. This will conclude the proof of Assertions (1) and (3). So consider  $\hat{\mathcal{F}}'$  a maximal  $H$ -invariant almost free factor system with  $\mathcal{F} \preceq \hat{\mathcal{F}}'$  and let us prove that  $\hat{\mathcal{F}} = \hat{\mathcal{F}}'$ . Consider a QH splitting defining  $\hat{\mathcal{F}}'$ , and let  $\Sigma'$  be the underlying surface. By Corollary 8.5, the boundary subgroups of  $\pi_1(\Sigma')$  are  $H$ -periodic. As above, at least one of these boundary subgroups is not  $\mathcal{F}$ -peripheral, so it is conjugate to a subgroup of  $\langle b \rangle$ , so  $\mathcal{F} \cup \{\langle b \rangle\}$  is  $\hat{\mathcal{F}}'$ -peripheral.

If  $\langle b \rangle$  is an unused boundary subgroup of  $\hat{\mathcal{F}}'$ , then write  $\hat{\mathcal{F}}' = \mathcal{F}' \cup \{\langle b \rangle\}$  where  $\mathcal{F}'$  is the corresponding extracted free factor system. Since  $\mathcal{F} \preceq \hat{\mathcal{F}}'$  and  $b$  is not  $\mathcal{F}$ -peripheral, it follows that  $\mathcal{F} \preceq \mathcal{F}'$ . By maximality of  $\mathcal{F}$ , it follows that  $\mathcal{F} = \mathcal{F}'$ , so  $\hat{\mathcal{F}} = \hat{\mathcal{F}}'$  and we are done.

If  $\langle b \rangle$  is not an unused boundary subgroup of  $\hat{\mathcal{F}}'$ , we argue towards a contradiction. The QH splitting  $S'$  associated to  $\hat{\mathcal{F}}'$  has an edge  $e$  whose stabilizer is commensurable with  $\langle b \rangle$ . Let  $U$  be the splitting of  $F_N$  obtained from  $S'$  by collapsing every edge not in the orbit of  $e$ . Denote by  $\Sigma$  the surface underlying the QH splitting  $S$  associated to  $\hat{\mathcal{F}}$ , so that  $\pi_1(\Sigma)$  is identified with a vertex stabilizer of  $S$ . Since  $\hat{\mathcal{F}} \preceq \hat{\mathcal{F}}'$ , the boundary subgroups of  $\pi_1(\Sigma)$  are elliptic in  $S'$  hence in  $U$ . If  $\pi_1(\Sigma)$  does not fix a point in  $U$ , then the action of  $\pi_1(\Sigma)$  on its minimal subtree in  $U$  is dual to a collection of simple closed curves in  $\Sigma$  whose stabilizer is conjugate to a power of  $b$ . But since  $\langle b \rangle$  is a boundary subgroup of  $\pi_1(\Sigma)$ , there is no such curve so  $\pi_1(\Sigma)$  fixes a point in  $U$ . This implies that there is an equivariant map  $f : S \rightarrow U$ , but this map has to collapse every edge in  $S$  because no edge stabilizer of  $S$  is commensurable with  $\langle b \rangle$ . It follows that  $f$  is constant, a contradiction.

This concludes the proof of Assertions (1) and (3) in all cases.

Assertion (2) is trivial if  $\hat{\mathcal{F}}$  is a free factor system, so write  $\hat{\mathcal{F}} = \mathcal{F} \cup \{\langle b \rangle\}$  for some unused boundary subgroup  $\langle b \rangle$  of  $\hat{\mathcal{F}}$ . The free factor system  $\mathcal{F}$  is  $H$ -invariant because  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  (Theorem 2.10), and it suffices to show that  $\mathcal{F}$  is maximal. Assume that  $\mathcal{F} \preceq \mathcal{F}'$  for some maximal  $H$ -invariant free factor system  $\mathcal{F}'$ . By Assertion (1), there exists a unique maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}'$  such that  $\mathcal{F}' \preceq \hat{\mathcal{F}}'$ . Note that  $\langle b \rangle$  cannot be  $\mathcal{F}'$ -peripheral because  $F_N$  is freely indecomposable relative to  $\hat{\mathcal{F}}$ ; but as  $\langle b \rangle$  is  $H$ -invariant, Assertion (3) implies that  $\langle b \rangle$  is  $\hat{\mathcal{F}}'$ -peripheral. Therefore  $\hat{\mathcal{F}}' = \mathcal{F}' \cup \{\langle b \rangle\}$ . Hence  $\hat{\mathcal{F}} \preceq \hat{\mathcal{F}}'$ , and by maximality of  $\hat{\mathcal{F}}$  it follows that  $\hat{\mathcal{F}} = \hat{\mathcal{F}}'$ , so  $\mathcal{F} = \mathcal{F}'$ .  $\square$

**Lemma 8.10.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup that does not preserve any non-trivial free splitting, and  $\hat{\mathcal{F}}$  be a maximal  $H$ -invariant almost free factor system.*

*Then there is a maximal  $H$ -invariant almost free factor system distinct from  $\hat{\mathcal{F}}$  if and only if there exists a proper free factor  $A$  which is not  $\hat{\mathcal{F}}$ -peripheral and whose conjugacy class is  $H$ -invariant.*

*Proof.* The ‘if’ direction follows from the fact that every proper free factor whose conjugacy class is  $H$ -invariant is contained in a maximal  $H$ -invariant almost free factor system. For the ‘only if’ direction, let  $\hat{\mathcal{F}}'$  be a maximal  $H$ -invariant almost free factor system distinct from  $\hat{\mathcal{F}}$ . If  $\hat{\mathcal{F}}'$  is a free factor system, then it contains a conjugacy class

of free factor  $A$  which is not  $\hat{\mathcal{F}}$ -peripheral, and the conjugacy class of  $A$  is  $H$ -invariant (as  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ ). So we assume that  $\hat{\mathcal{F}}'$  is not a free factor system, and write  $\hat{\mathcal{F}}' = \mathcal{F}' \cup \{\langle b \rangle\}$  where  $\mathcal{F}'$  is a free factor system extracted from  $\hat{\mathcal{F}}'$ . By Lemma 8.8(2),  $\mathcal{F}'$  is a maximal  $H$ -invariant free factor system. If some free factor in  $\mathcal{F}'$  is not  $\hat{\mathcal{F}}$ -peripheral, then we are done. Otherwise, we have  $\mathcal{F}' \preceq \hat{\mathcal{F}}$ . But by Lemma 8.8(3), the conjugacy class of  $\langle b \rangle$  is also  $\hat{\mathcal{F}}$ -peripheral, so  $\hat{\mathcal{F}}' \preceq \hat{\mathcal{F}}$ , and by maximality  $\hat{\mathcal{F}}' = \hat{\mathcal{F}}$ , a contradiction.  $\square$

**Restricting an almost free factor system to a subgroup.** Recall from Section 2.2 that if  $A$  is a subgroup of  $F_N$ , and  $\mathcal{Q}$  a collection of conjugacy classes of subgroups of  $F_N$ , we define  $\mathcal{Q}|_A$  as the set of all  $A$ -conjugacy classes of non-trivial groups of the form  $Q \cap A$ , where  $Q \subseteq F_N$  is a subgroup whose conjugacy class belongs to  $\mathcal{Q}$ .

If  $\mathcal{F}$  is a free factor system of  $F_N$ , and  $A$  is any subgroup, not  $\mathcal{F}$ -peripheral, then the Kurosh lemma says that  $\mathcal{F}|_A$  is a free factor system of  $A$ . Our next lemma describes what happens for the restriction of an almost free factor system.

A *Grushko factor* of a group  $G$  relative to a collection of subgroups  $\mathcal{P}$  is a vertex stabilizer in any Grushko  $(G, \mathcal{P})$ -splitting: with our usual conventions, if  $G = G_1 * \cdots * G_p * F_r$  is a Grushko decomposition of  $(G, \mathcal{P})$ , then its Grushko factors are the groups conjugate to  $G_i$  for some  $i \leq p$ . Note that  $G_i$  is one-ended relative to  $\mathcal{P}|_{G_i}$ .

**Lemma 8.11.** *Let  $\hat{\mathcal{F}}$  be an almost free factor system of  $F_N$ , and  $A \subseteq F_N$  be a finitely generated malnormal subgroup which is not  $\hat{\mathcal{F}}$ -peripheral.*

*If  $\hat{\mathcal{F}}|_A$  is not an almost free factor system of  $A$ , then the Grushko decomposition of  $A$  relative to  $\hat{\mathcal{F}}|_A$  is non-trivial and there exists a Grushko factor  $A_0$  of  $A$  relative to  $\hat{\mathcal{F}}|_A$  which is not  $\hat{\mathcal{F}}|_A$ -peripheral.*

*Proof.* If the Grushko decomposition of  $A$  relative to  $\hat{\mathcal{F}}|_A$  is non-trivial and all its factors are  $\hat{\mathcal{F}}|_A$ -peripheral, then  $\hat{\mathcal{F}}|_A$  is a free factor system of  $A$ .

We thus assume that  $A$  is freely indecomposable relative to  $\hat{\mathcal{F}}|_A$  and prove that  $\hat{\mathcal{F}}|_A$  is an almost free factor system. We may assume that  $\hat{\mathcal{F}}$  is the factor system of a QH splitting  $S$  of  $F_N$  with QH vertex  $v \in S$ , with underlying surface  $\Sigma$ .

We first treat the case where  $A$  is contained in the surface group  $G_v = \pi_1(\Sigma)$ . If  $A$  has finite index in  $G_v$ , then being malnormal, this implies that  $A = G_v$  and  $\hat{\mathcal{F}}|_A$  consists in the conjugacy classes of fundamental groups of boundary components of  $\Sigma$ , so  $\hat{\mathcal{F}}|_A$  is an almost free factor system of  $A$ . We now assume that  $A$  has infinite index in  $G_v$ . Since  $A$  is not  $\hat{\mathcal{F}}$ -peripheral, it is not conjugate in a boundary subgroup of  $G_v$ , so  $A$  has a non-trivial free splitting relative to  $\hat{\mathcal{F}}|_A$  (see for instance [Per11, Lemma 3.11] for this well known fact).

We may now assume that  $A$  does not fix any point in  $S$ . Indeed, the case where  $A$  fixes a point in the orbit of  $v$  was treated above, and if  $A$  fixes another vertex, then  $A$  is  $\hat{\mathcal{F}}$ -peripheral.

Let  $V_p \sqcup V_s$  be the bipartition of vertices of  $S$  where  $V_s = F_N \cdot \{v\}$ . Let  $S_A \subseteq S$  be the minimal  $A$ -invariant subtree. Since  $A$  is freely indecomposable relative to  $\hat{\mathcal{F}}|_A$ , for

every edge  $e$  of  $S_A$ , the intersection  $G_e \cap A$  is non-trivial (in fact maximally cyclic in  $F_N$  by malnormality of  $A$ ). In particular, for every vertex  $w \in V_s \cap S_A$ , the intersection  $G_w \cap A$  is not cyclic since by minimality, there are at least two edges in  $S_A$  incident on  $w$ . Therefore, if  $G_w \cap A$  has infinite index in  $G_w$ , the argument above yields a non-trivial free splitting of  $G_w \cap A$  relative to the incident edge groups, hence a non-trivial free splitting of  $A$  relative to  $\hat{\mathcal{F}}_{|A}$ , a contradiction.

We can therefore assume that for every vertex  $w \in V_s \cap S_A$ , the intersection  $G_w \cap A$  has finite index in  $G_w$ . As  $A$  is malnormal, this implies that  $A \cap G_w = G_w$ . It also follows that all vertices in  $V_s \cap S_A$  are in the same  $A$ -orbit: indeed, if  $w, w' \in V_s \cap S_A$ , then  $w = gw'$  for some  $g \in F_N$ , so  $G_w \subseteq A \cap A^{g^{-1}}$  so  $g \in A$  by malnormality.

We now check that  $S_A$  is a QH splitting of  $A$ . Up to conjugating  $A$ , we may assume that  $v \in S_A$ , and we claim that  $v$  is a clean QH vertex in  $S_A$ . Indeed,  $G_v = \pi_1(\Sigma)$ , and the stabilizers of incident edges in  $S_A$  are subgroups of the boundary components of  $\Sigma$ . Moreover, since  $A \cap G_v = G_v$ , the  $A$ -stabilizer of any edge in  $S_A$  incident on  $v$  agrees with its  $F_N$ -stabilizer. It follows that  $v$  is clean QH as a vertex of  $S_A$ . Since  $\Sigma$  has an unused boundary component in  $S$ , the same is true in  $S_A$ , so  $S_A$  is a QH splitting of  $A$ .

There remains to check that  $\hat{\mathcal{F}}_{|A}$  coincides with the set of all conjugacy classes of  $A$ -stabilizers of vertices in  $V_p \cap S_A$  together with the conjugacy classes of fundamental groups of unused (in  $S_A$ ) boundary components of  $\Sigma$ . Each group  $Q$  in  $\hat{\mathcal{F}}$  is either of the form  $Q = G_p$  for some vertex  $p \in V_p$ , or  $Q$  is conjugate to the fundamental group  $G_b$  of a boundary component  $b$  of  $\Sigma$  unused in  $S$ . If  $Q = G_p$  with  $p \in S_A$ , then  $Q \cap A$  is the stabilizer of the vertex  $p$  for the action  $A \curvearrowright S$ . If  $p \notin S_A$ , then either  $G_p \cap A$  is trivial, or  $p$  is adjacent to a vertex  $w \in V_s \cap S_A$  (this is because any segment in  $S$  with non-trivial stabilizer has length at most 2 and contains at most one vertex in  $V_p$ ). In this case, up to conjugation we may assume that  $v = w$ ; then  $G_p \cap A = G_p \cap G_v$  is conjugate to the fundamental group of a boundary component of  $\Sigma$  that is unused in  $S_A$ . If  $Q$  is conjugate to the stabilizer  $G_b$  of an unused (in  $S$ ) boundary component of  $\Sigma$ , then  $Q$  fixes a unique vertex  $w \in S$  (and  $w \in V_s$ ). If  $w \in S_A$ , then  $Q \cap A = Q$  because  $G_w \subseteq A$ , so  $Q$  is conjugate in  $A$  to the stabilizer of an unused boundary component of  $\Sigma$ . Finally, if  $w \notin S_A$ , then  $Q \cap A = \{1\}$ . Otherwise,  $Q \cap A$  has finite index in the cyclic group  $Q$ . But  $Q \cap A$  fixes the arc joining  $w$  to  $S_A$ , hence so does  $Q$  since edge stabilizers are maximal cyclic, contradicting that  $Q$  fixes no edge in  $S$ . This concludes the proof.  $\square$

## 8.2 The dynamical decomposition: statement and examples

Given a subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , the goal of this section is to construct a *dynamical decomposition* of the free group  $F_N$  for  $H$ . As was informally explained in Section 1.3.3 of the introduction, this is an analogue of Ivanov's natural  $H$ -invariant decomposition of a surface into *active* and *inactive* subsurfaces, when  $H$  is a subgroup of the mapping class group.

**Definition 8.12.** *We say that a subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is pure if there is a unique maximal  $H$ -invariant almost free factor system, but no  $H$ -invariant non-trivial free splitting.*

We insist that in this definition, the unique maximal  $H$ -invariant almost free factor system is allowed to be empty, which happens if  $H$  contains an atoroidal fully irreducible automorphism.

*Remark 8.13.* One can naturally extend this definition to any subgroup of  $\text{Out}(F_N)$  as follows: define  $H \subseteq \text{Out}(F_N)$  as *pure* if  $H \cap \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is pure. Equivalently,  $H$  is pure if there is a unique maximal  $H$ -periodic almost free factor system, and no  $H$ -periodic non-trivial free splitting. The equivalence follows from the fact that any  $H$ -periodic almost free factor system (or free splitting) is invariant under  $H \cap \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ .

**Definition 8.14.** Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup. A subgroup  $P \subseteq F_N$  is *universally peripheral with respect to  $H$*  if it is  $\hat{\mathcal{F}}$ -peripheral for any maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}$ .

We denote by  $\mathcal{P}_H$  the set of all subgroups of  $F_N$  that are universally peripheral with respect to  $H$ , and by  $\mathcal{P}_H^{\max}$  the subset of maximal universally peripheral subgroups.

We note that  $\mathcal{P}_H$  is stable under taking subgroups. Since any intersection of free factors is a free factor, every group  $P \in \mathcal{P}_H^{\max}$  is either a free factor or a maximal cyclic subgroup of  $F_N$ . The case where  $H$  is fully irreducible and atoroidal (i.e. does not preserve the conjugacy class of any proper free factor or of any cyclic group) is a bit peculiar: there is no non-empty  $H$ -invariant almost free factor system (i.e.  $\hat{\mathcal{F}} = \emptyset$  is the only one) so  $\mathcal{P}_H = \mathcal{P}_H^{\max}$  is reduced to the trivial subgroup. But anyway the dynamical decomposition is clearly trivial in this case, so we may as well exclude this case from now on.

**Theorem 8.15** (Dynamical decomposition). Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup that does not preserve any non-trivial free splitting of  $F_N$ .

Then there exists a unique  $H$ -invariant splitting  $U_H^d$  of  $F_N$  relative to  $\mathcal{P}_H$  whose vertex set  $V$  has a bipartition  $V = V_p \sqcup V_a$  (for peripheral vs. active vertices) with the following properties:

- (1) for every  $v \in V_p$ , one has  $G_v \in \mathcal{P}_H^{\max}$ , and  $G_v \cap G_{v'} = \{1\}$  for distinct vertices  $v, v' \in V_p$ ;
- (2) for every  $v \in V_a$ , the group  $G_v$  is a free factor of  $F_N$  whose conjugacy class is  $H$ -invariant, the restriction of  $H$  to  $G_v$  is pure, and  $(\mathcal{P}_H^{\max})|_{G_v}$  is the unique maximal  $H$ -invariant almost free factor system in  $G_v$ ; moreover, the collection of incident edge groups is a non-sporadic free factor system of  $G_v$ ;
- (3) for every free factor  $A$  whose conjugacy class is  $H$ -invariant and which is not universally peripheral, there exists  $v \in V_a$  such that  $G_v \subseteq A$ ;
- (4) for every vertex  $v \in V_a/F_N$ , there exists a unique maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}_v$  such that  $G_v$  is not  $\hat{\mathcal{F}}_v$ -peripheral. Moreover the map  $v \mapsto \hat{\mathcal{F}}_v$  is a bijection from  $V_a/F_N$  to the set of maximal  $H$ -invariant almost free factor systems.




Figure 7: The splitting in Example 8.19.

As usual, the uniqueness of  $U_H^d$  is up to  $F_N$ -equivariant isomorphism.

*Remark 8.16.* It is *a priori* not obvious that there are only finitely many maximal  $H$ -invariant almost free factor systems (under the assumption that  $H$  preserves no non-trivial free splitting), but this is a consequence of Assertion (4). In fact, Theorem 8.15 can be used to derive a bound on their numbers, as will be done in Corollary 8.30 below. We insist that the assumption that  $H$  preserves no free splitting is crucial here, see Example 6.1 for a situation where there are infinitely many invariant splittings dual to arcs on a surface, hence infinitely many sporadic  $H$ -invariant free factor systems.

*Remark 8.17.* Since  $H$  does not preserve any non-trivial free splitting, edge stabilizers of  $U_H^d$  are non-trivial. By Assertion (1), they are universally peripheral with respect to  $H$ , and we will see that they are free factors of  $F_N$  (see Proposition 8.33).

*Remark 8.18.* If  $H$  is pure,  $U_H^d$  is the trivial splitting, reduced to a vertex in  $V_a$ . If  $H$  is not pure, then  $U_H^d$  is non-trivial, and there are at least two orbits of active vertices. One can refine each of these vertices into a non-sporadic free splitting.

*Example 8.19.* We give an example (see Figure 7) illustrating the dynamical decomposition of Theorem 8.15.

Let  $\Sigma_1, \Sigma_2, \Sigma_3$  be three connected orientable surfaces of positive genus, each of which has exactly two boundary components. Let  $F_N$  be the free group obtained by amalgamating the fundamental groups of these surfaces over the fundamental group of one of their boundary components, and let  $S$  be the splitting of  $F_N$  represented in Figure 7. Let  $H$  be the subgroup of  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  induced by the homeomorphisms of the three surfaces whose restriction to the boundary is the identity.

Let us describe the  $H$ -invariant conjugacy classes of proper free factors. Note that  $\pi_1(\Sigma_i)$  is a free factor of  $F_N$  as can be seen by drawing arcs on  $\Sigma_{i-1}$  and  $\Sigma_{i+1}$  with endpoints in their unused boundary components. Similarly, so is  $\langle c_i \rangle$ , the fundamental



group of the unused boundary curve in  $\Sigma_i$ , and  $\langle c \rangle$  the central vertex group. Finally, the subgroup  $Q_i = \pi_1(\Sigma_{i-1} \cup \Sigma_{i+1})$  is also a free factor of  $F_N$ . The conjugacy classes of all these free factors are clearly  $H$ -invariant. Conversely, one may use the argument in Lemma 8.20 below and the fact that  $H$  contains elements whose restriction to  $\pi_1(\Sigma_i)$  is pseudo-Anosov to prove that these are the only invariant free factors. It follows in particular that  $H$  does not preserve any non-trivial free splitting.

The almost free factor system  $\hat{\mathcal{F}}_i = \{[Q_i], [\langle c_i \rangle]\}$  is  $H$ -invariant and maximal. On the other hand,  $\pi_1(\Sigma_i)$  is a free factor which is not  $\hat{\mathcal{F}}_i$ -peripheral and whose conjugacy class is  $H$ -invariant, and it is minimal for this property. Minimal non-peripheral free factors will play an important role in the proof and appear as stabilizers of active vertices by Assertion (3).

One can check that the dynamical decomposition  $U_H^d$  of Theorem 8.15 is the splitting dual to the decomposition shown in Figure 7 where  $V_p/F_N$  is the central vertex, and  $V_a/F_N$  corresponds to the three surfaces.

In this example, active vertices correspond to surfaces, but this is not a general fact. For instance, one may replace any of the surfaces of the example by a free group  $F = \langle x_1, \dots, x_p \rangle$ , and attach it to the central vertex along  $x_1$ , and add to  $H$  all the automorphisms of  $F$  fixing  $x_1$ .

### 8.3 A minimal non-peripheral invariant free factor

The main step in the proof of the existence part of Theorem 8.15 is the following result.

**Lemma 8.20.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup that does not preserve any non-trivial free splitting. Let  $\hat{\mathcal{F}}$  be a maximal  $H$ -invariant almost free factor system.*

*Among all conjugacy classes of non-trivial free factors of  $F_N$  (including  $F_N$  itself) which are  $H$ -invariant and not  $\hat{\mathcal{F}}$ -peripheral, there is a unique minimal one, that we denote by  $A_{\hat{\mathcal{F}}}$ .*

*In addition,  $A_{\hat{\mathcal{F}}}$  is non-abelian and there is a minimal  $H$ -invariant splitting  $T_{\hat{\mathcal{F}}}$  of  $F_N$  relative to  $\hat{\mathcal{F}}$  coming with a bipartition of its vertex set as  $V^0 \sqcup V^1$  that satisfies the following properties.*

- (1) *For every  $v \in V^0$ , one has  $[G_v] \in \hat{\mathcal{F}}$  and  $G_v$  is not an unused boundary subgroup of  $\hat{\mathcal{F}}$ . Moreover  $G_v \cap G_{v'} = \{1\}$  for distinct vertices  $v, v' \in V^0$ ;*
- (2)  *$V^1$  consists of a single orbit of vertices, the stabilizer of every vertex in  $V^1$  is conjugate to  $A_{\hat{\mathcal{F}}}$ ;*
- (3)  *$\hat{\mathcal{F}}|_{A_{\hat{\mathcal{F}}}}$  is a non-sporadic almost free factor system of  $A_{\hat{\mathcal{F}}}$ , and the collection of incident edge groups is a non-sporadic free factor system of  $A_{\hat{\mathcal{F}}}$ .*

*Remark 8.21.* In this statement, we do not claim that  $A_{\hat{\mathcal{F}}}$  is a proper free factor; in fact  $A_{\hat{\mathcal{F}}} = F_N$  if and only if  $H$  is pure (see Lemma 8.10). More generally, the defining property of  $A_{\hat{\mathcal{F}}}$  implies that the restriction of  $H$  to  $A_{\hat{\mathcal{F}}}$  is pure, and its unique maximal  $H$ -invariant maximal almost free factor system is  $\hat{\mathcal{F}}|_{A_{\hat{\mathcal{F}}}}$ .

*Example 8.22.* Continuing with the notations from Example 8.19, let  $\hat{\mathcal{F}} := \hat{\mathcal{F}}_1 = \{[Q_1], [\langle c_1 \rangle]\}$ . Up to conjugation, among all  $H$ -invariant free factors which are not  $\hat{\mathcal{F}}$ -peripheral, the unique minimal one is  $A_{\hat{\mathcal{F}}_1} = \pi_1(\Sigma_1)$ . The splitting  $T_{\hat{\mathcal{F}}_1}$  is the Bass–Serre tree of the decomposition  $F_N = \pi_1(\Sigma_1) *_Z Q_1$ ; the vertex with stabilizer  $\pi_1(\Sigma_1)$  belongs to  $V^1$ , and the vertex with stabilizer  $Q_1$  belongs to  $V^0$ . Moreover,  $\hat{\mathcal{F}}_{|\pi_1(\Sigma_1)} = \{[\langle b_1 \rangle], [\langle c_1 \rangle]\}$ .

In our proof of Lemma 8.20, we will make use of the following fact.

**Lemma 8.23.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Let  $A$  and  $A'$  be two proper free factors of  $F_N$  whose conjugacy classes are  $H$ -invariant.*

*Then for every  $g \in F_N$ , the conjugacy class of  $A \cap (A')^g$  is  $H$ -invariant.*

*Proof.* We claim that up to conjugation, there are only finitely many subgroups of the form  $A \cap (A')^g$  with  $g \in F_N$ . To see this, let  $S_{A'}$  be a Grushko splitting of  $F_N$  relative to  $A'$ . Then the minimal  $A$ -invariant subtree of  $S_{A'}$  is a free splitting of  $A$  (in particular there is a bound on the number of  $A$ -orbits of vertices), and the collection of all its point stabilizers is exactly equal to the collection of all subgroups of the above form.

Since  $[A]$  and  $[A']$  are  $H$ -invariant, the group  $H$  permutes the conjugacy classes  $[A \cap (A')^g]$ . As  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , these are actually  $H$ -invariant.  $\square$

*Proof of Lemma 8.20.* Let  $A, A'$  be two free factors of  $F_N$  which are not  $\hat{\mathcal{F}}$ -peripheral, whose conjugacy classes are  $H$ -invariant with  $A$  of minimal rank. We aim to prove that  $A$  is conjugate into  $A'$ .

Let  $\mathcal{F}$  be a free factor system extracted from  $\hat{\mathcal{F}}$ . By Lemma 8.8,  $\mathcal{F}$  is a maximal  $H$ -invariant free factor system (non-sporadic because  $H$  does not fix any free splitting of  $F_N$ ) and  $H$  contains an automorphism  $\alpha$  which is fully irreducible relative to  $\mathcal{F}$  and atoroidal relative to  $\hat{\mathcal{F}}$ .

Let  $T$  be an  $\alpha$ -invariant arational tree (rel.  $\mathcal{F}$ ) on which  $\alpha$  acts by homothety of dilatation factor  $\lambda \neq 1$ : this exists by Proposition 2.4. Notice that a subgroup is elliptic in  $T$  if and only if it is  $\hat{\mathcal{F}}$ -peripheral.

Since  $A$  is not elliptic in  $T$ , we can consider the minimal  $A$ -invariant subtree  $T_A$  of  $T$ . Since  $[A]$  is  $\alpha$ -invariant, we deduce that  $T_A$  is  $\tilde{\alpha}$ -invariant for some preimage  $\tilde{\alpha} \in \text{Aut}(F_N)$  of  $\alpha$ . Since  $\tilde{\alpha}$  acts on  $T_A$  by homothety of dilatation  $\lambda \neq 1$ ,  $T_A$  has no simplicial edges, so  $A$  acts on  $T_A$  with dense orbits.

Using an argument of Reynolds (as formulated in [GH19a, Corollary 11.9]), we deduce that the collection  $\{gT_A\}_{g \in F_N}$  is a transverse family of  $T$ , and more precisely, for every  $g \in F_N \setminus A$ , the intersection  $gT_A \cap T_A$  contains at most one point (in particular  $\text{Stab}_{F_N}(T_A) = A$ ). As  $T$  is arational, it is mixing [Hor14, Lemma 4.9], so  $\{gT_A\}_{g \in F_N}$  is a transverse covering of  $T$ , i.e. the subtree  $T_A$  is closed and every segment of  $T$  is covered by finitely many subtrees from this family.

Likewise, the above argument also yields another transverse covering  $\{gT_{A'}\}_{g \in F_N}$  of  $T$ . We can then form the refinement of the two transverse coverings, which consists of all nondegenerate subtrees of  $T$  of the form  $gT_A \cap g'T_{A'}$  with  $g, g' \in F_N$ . Since this forms a transverse covering of  $T$ , the stabilizer of each of these trees acts with dense orbits on it (see e.g. [HW20, Lemma 5.2]) and in particular is not  $\hat{\mathcal{F}}$ -peripheral and not cyclic.

We observe that if  $T_A \cap gT_{A'}$  is nondegenerate, then its stabilizer is equal to  $A \cap (A')^g$ . Indeed, it is clear that  $A \cap (A')^g$  preserves both  $T_A$  and  $gT_{A'}$ , and therefore preserves their intersection. Conversely, if  $h \in F_N$  preserves  $T_A \cap gT_{A'}$ , then in particular  $hT_A \cap T_A$  is nondegenerate, so  $h \in A$ , and similarly,  $h \in (A')^g$ .

Therefore, there exists  $g \in F_N$  such that  $A \cap (A')^g$  is not elliptic in  $T$ , hence not  $\hat{\mathcal{F}}$ -peripheral. By Lemma 8.23, its conjugacy class is  $H$ -invariant. By minimality of the rank of  $A$ , we have  $A \subseteq (A')^g$ , as desired. This concludes our proof of the existence and the uniqueness of  $A_{\hat{\mathcal{F}}}$ , and of the fact that  $A_{\hat{\mathcal{F}}}$  is non-abelian.

We now construct the tree  $T_{\hat{\mathcal{F}}}$ . Our construction a priori depends on the choice of  $\alpha \in H$  and of the  $\mathbb{R}$ -tree  $T$ , but we will check at the end of the proof that in fact  $T_{\hat{\mathcal{F}}}$  does not depend on these choices. We keep the notations from above. Let  $T_{\hat{\mathcal{F}}}$  be the skeleton (in the sense of [Gui04, Definition 4.8]) of the transverse covering  $\{gT_A\}_{g \in F_N}$ , i.e.  $T_{\hat{\mathcal{F}}}$  is the bipartite simplicial tree having one vertex  $v_Y$  for every subtree  $Y$  of the form  $gT_A$ , one vertex  $v_x$  for every point  $x \in T$  that belongs to at least two subtrees from the family, with an edge between  $v_x$  and  $v_Y$  whenever  $x \in Y$ . Minimality of  $T_{\hat{\mathcal{F}}}$  follows from [Gui04, Lemma 4.9]. We denote by  $V^1$  the set of all vertices of the form  $v_Y$ , and by  $V^0$  the set of all vertices of the form  $v_x$ . Notice that  $V^1$  consists of a single orbit of vertices, and the stabilizer of every vertex in  $V^1$  is conjugate to  $A_{\hat{\mathcal{F}}}$ , proving Assertion (2). Edge stabilizers of  $T_{\hat{\mathcal{F}}}$  are non-trivial because otherwise, this would produce an  $(F_N, \mathcal{F})$ -free factor acting non-discretely on  $T$ , contradicting arationality. In particular, the stabilizer of any vertex  $v = v_x \in V^0$  is non-trivial, and since non-trivial point stabilizers of  $T$  are precisely groups in  $\hat{\mathcal{F}}$ , we deduce that  $[G_v] \in \hat{\mathcal{F}}$ . Since arc stabilizers of  $T$  are trivial, it follows that the stabilizers of distinct vertices in  $V^0$  have trivial intersection.

To prove Assertion (1), there remains to prove that  $G_v$  is not an unused boundary subgroup  $\langle b \rangle$  of  $\hat{\mathcal{F}}$ . So assume otherwise that  $G_v = \langle b \rangle$ . Since there are at least two edges incident on  $v$  in  $T_{\hat{\mathcal{F}}}$ , and since edge stabilizers are non-trivial, there exists  $k \geq 1$  such that  $b^k$  is contained in two distinct conjugates of  $A$ , say  $b^k \in A^g \cap A^{g'}$ . By malnormality,  $A^g = A^{g'}$ , a contradiction. This proves Assertion (1).

We now prove Assertion (3). We first apply Lemma 8.11 to prove that  $\hat{\mathcal{F}}|_A$  is an almost free factor system. If not, then there is Grushko factor  $A_0 \subsetneq A$  of  $A$  relative to  $\hat{\mathcal{F}}|_A$  which is not  $\hat{\mathcal{F}}|_A$ -peripheral. Since there are only finitely many conjugacy classes of such factors,  $A_0$  is  $H$ -periodic hence  $H$ -invariant because  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . This contradicts the choice of  $A$ . This proves that  $\hat{\mathcal{F}}|_A$  is an almost free factor system of  $A$ . If it was sporadic, then the action of  $A$  on  $T_A$  would be simplicial, a contradiction.

In order to prove the last part of Assertion (3), denote by  $\mathcal{F}_0 \subseteq \hat{\mathcal{F}}$  the collection of all conjugacy classes of stabilizers of  $V^0$ -vertices, and note that  $(\mathcal{F}_0)|_A$  coincides with the set of  $A$ -conjugacy classes of incident edge groups. By Assertion (1),  $\mathcal{F}_0$  contains no unused boundary subgroup of  $\hat{\mathcal{F}}$ , so  $\mathcal{F}_0$  is a free factor system of  $F_N$ , hence  $(\mathcal{F}_0)|_A$  is a free factor system of  $A$ . If  $(\mathcal{F}_0)|_A$  was sporadic, then as above, the action of  $A$  on  $T_A$  would be simplicial, a contradiction.

We finally give an alternative description of the tree  $T_{\hat{\mathcal{F}}}$  which shows that it does not depend on the choice of  $\alpha$  and of the  $\mathbb{R}$ -tree  $T$ , and which also implies that it is  $H$ -invariant. We note that since  $T_{\hat{\mathcal{F}}}$  is bipartite with non-trivial edge stabilizers, and

since  $G_v \cap G_{v'} = \{1\}$  for any two distinct vertices  $v, v' \in V^0$ , it follows that two vertices  $v \in V^0$  and  $v' \in V^1$  are adjacent if and only if  $G_v \cap G_{v'} \neq \{1\}$ . So let  $S$  be the bipartite graph on the vertex set  $V^0(S) \sqcup V^1(S)$ , where  $V^0(S)$  is defined as the set of subgroups of  $F_N$  whose conjugacy class is in  $\hat{\mathcal{F}}$ , and  $V^1(S)$  is the set of minimal non- $\hat{\mathcal{F}}$ -peripheral free factors whose conjugacy class is  $H$ -invariant, and where two vertices (one in  $V^0(S)$  and one in  $V^1(S)$ ) are adjacent if and only if the corresponding subgroups have non-trivial intersection. The action of  $F_N$  by conjugation on its subgroups induces an action on  $S$ . The map  $f : v \in V(T_{\hat{\mathcal{F}}}) \mapsto G_v \in V(S)$  is injective, and preserves adjacency and non-adjacency. We claim every vertex in  $S \setminus f(T_{\hat{\mathcal{F}}})$  is a terminal vertex of  $S$  (in particular,  $S$  is a tree). Indeed, let  $P$  be the subgroup associated to a vertex  $v \in S \setminus f(T_{\hat{\mathcal{F}}})$ . Then  $[P] \in \hat{\mathcal{F}}$  (in other words  $v \in V^0(S)$ ) and the point  $x \in T$  fixed by  $P$  is not in  $V^0(T_{\hat{\mathcal{F}}})$ . This means that  $x$  lies in a single translate  $gT_A$  of  $T_A$ , so  $P \subseteq A^g$ . Since arc stabilizers of  $T$  are trivial, all other conjugates of  $A$  intersect  $P$  trivially. This means that  $v$  is adjacent to a unique vertex in  $V^1(S)$ , and the bipartite structure of  $S$  implies that  $v$  is a terminal vertex. This shows that  $T_{\hat{\mathcal{F}}}$  can be described as the complement of terminal vertices in  $S$  and concludes the proof.  $\square$

The factors  $A_{\hat{\mathcal{F}}}$  have the following property.

**Lemma 8.24.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup that does not preserve any non-trivial free splitting, and let  $\hat{\mathcal{F}}$  and  $\hat{\mathcal{F}}'$  be two distinct maximal  $H$ -invariant almost free factor systems.*

*Then  $A_{\hat{\mathcal{F}}}$  is  $\hat{\mathcal{F}}'$ -peripheral, and it is not contained in an unused boundary subgroup of  $\hat{\mathcal{F}}'$ .*

*Proof.* Since  $\hat{\mathcal{F}}' \not\leq \hat{\mathcal{F}}$ , consider a factor  $F'$  in  $\hat{\mathcal{F}}'$  which is not  $\hat{\mathcal{F}}$ -peripheral. By Lemma 8.8(3),  $F'$  is not cyclic, hence is not an unused boundary subgroup of  $\hat{\mathcal{F}}'$ . In particular,  $F'$  is a free factor of  $F_N$ , and its conjugacy class is  $H$ -invariant. Therefore,  $F'$  contains a conjugate of  $A_{\hat{\mathcal{F}}}$ . This concludes that  $A_{\hat{\mathcal{F}}}$  is  $\mathcal{F}'$ -peripheral. Since  $A_{\hat{\mathcal{F}}}$  is not abelian by Theorem 8.20, it cannot be contained in an unused boundary subgroup of  $\hat{\mathcal{F}}'$ .  $\square$

We record the following consequences of Lemma 8.24.

**Corollary 8.25.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup that does not preserve any non-trivial free splitting, and let  $\hat{\mathcal{F}}$  be a maximal  $H$ -invariant almost free factor system. Then all edge stabilizers of  $T_{\hat{\mathcal{F}}}$  are universally peripheral with respect to  $H$ .*

*Proof.* Lemma 8.24 implies that edge stabilizers of  $T_{\hat{\mathcal{F}}}$  are  $\hat{\mathcal{F}}'$ -peripheral for  $\hat{\mathcal{F}}' \neq \hat{\mathcal{F}}$  because  $A_{\hat{\mathcal{F}}}$  is. They are also  $\hat{\mathcal{F}}$ -peripheral by Lemma 8.20(1).  $\square$

**Lemma 8.26.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup that does not preserve any non-trivial free splitting, and let  $\hat{\mathcal{F}}_1, \hat{\mathcal{F}}_2$  be two distinct maximal  $H$ -invariant almost free factor systems.*

*Then  $\hat{\mathcal{F}}_1 \cup \hat{\mathcal{F}}_2$  is filling in the following sense: there is no almost free factor system  $\hat{\mathcal{F}}$  such that  $\hat{\mathcal{F}}_1 \preceq \hat{\mathcal{F}}$  and  $\hat{\mathcal{F}}_2 \preceq \hat{\mathcal{F}}$ .*

*Remark 8.27.* The point in the lemma is that we do not assume that  $\hat{\mathcal{F}}$  is  $H$ -invariant.

*Proof.* Assume on the contrary that there exists an almost free factor system  $\hat{\mathcal{F}}$  such that  $\hat{\mathcal{F}}_1 \preceq \hat{\mathcal{F}}$  and  $\hat{\mathcal{F}}_2 \preceq \hat{\mathcal{F}}$ . If  $\hat{\mathcal{F}}$  is a free factor system, then the Grushko decomposition of  $F_N$  relative to  $\hat{\mathcal{F}}_1 \cup \hat{\mathcal{F}}_2$  is non-trivial, and provides a free factor system  $\mathcal{G}$  such that  $\hat{\mathcal{F}}_1 \not\preceq \mathcal{G}$  and  $\hat{\mathcal{F}}_2 \not\preceq \mathcal{G}$ , and which is  $H$ -invariant. This contradicts maximality of  $\hat{\mathcal{F}}_1$  and  $\hat{\mathcal{F}}_2$ .

So assume that  $\hat{\mathcal{F}} = \{[A_1], \dots, [A_n], [\langle b_1 \rangle], \dots, [\langle b_r \rangle]\}$  is the factor system of a QH splitting  $S$  with  $r \geq 1$  unused boundary components. Let  $\Gamma'$  be the non-minimal graph of groups obtained from  $S/F_N$  by adding  $r$  terminal edges with stabilizers  $\langle b_1 \rangle, \dots, \langle b_r \rangle$ , and let  $S'$  be the Bass-Serre tree of this splitting. The splitting  $S'$  has the property that for every  $\hat{\mathcal{F}}$ -peripheral subgroup  $A \subseteq F_N$ , there is a unique vertex  $v_A \in S'$  such that  $A$  fixes  $v_A$  and the stabilizer of  $v_A$  is in  $\hat{\mathcal{F}}$ .

We claim that all vertex stabilizers of  $T_{\hat{\mathcal{F}}_1}$  are  $\hat{\mathcal{F}}$ -peripheral. Indeed,  $A_{\hat{\mathcal{F}}_1}$  is  $\hat{\mathcal{F}}$ -peripheral because it is  $\hat{\mathcal{F}}_2$ -peripheral by Lemma 8.24, and the stabilizers of  $V^0$ -vertices of  $T_{\hat{\mathcal{F}}_1}$  are  $\hat{\mathcal{F}}_1$ -peripheral (Lemma 8.20(1)). It follows that there is an equivariant map  $f : T_{\hat{\mathcal{F}}_1} \rightarrow S'$  that sends each vertex  $u \in T_{\hat{\mathcal{F}}_1}$  to the vertex  $v_{G_u}$  defined above. Edge stabilizers of  $T_{\hat{\mathcal{F}}_1}$  are non-trivial because  $H$  preserves no non-trivial free splitting. Since  $\hat{\mathcal{F}}$  is a malnormal family, this forces  $f$  to be constant, say  $f(T_{\hat{\mathcal{F}}_1}) = \{w\}$  with  $[G_w] \in \hat{\mathcal{F}}$ . Thus  $F_N$  is  $\hat{\mathcal{F}}$ -peripheral, a contradiction.  $\square$

#### 8.4 End of the proof via a tree of cylinders construction

If  $\hat{\mathcal{F}}_1, \dots, \hat{\mathcal{F}}_n$  are maximal  $H$ -invariant free factor systems, Corollary 8.25 implies that for all  $i, j \leq n$ , the trees  $T_{\hat{\mathcal{F}}_i}, T_{\hat{\mathcal{F}}_j}$  are elliptic with respect to each other, i.e. each edge stabilizer fixes a point in the other tree. This allows to construct a refinement of  $T_{\hat{\mathcal{F}}_1}$  dominating  $T_{\hat{\mathcal{F}}_2}$ , and repeating this procedure, a splitting that dominates all the trees  $T_{\hat{\mathcal{F}}_i}$ . To make things more canonical it will be convenient to use a suitable notion of tree of cylinders (see Section 5.2). We take as a class of allowed edge groups the class  $\mathcal{E}$  of non-trivial universally peripheral subgroups (Definition 8.14), i.e.  $\mathcal{E} = \mathcal{P}_H \setminus \{\{1\}\}$ . We consider the following equivalence relation on  $\mathcal{E}$ .

**Lemma 8.28.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup. If  $P, P' \in \mathcal{P}_H$  and  $P \cap P' \neq \{1\}$ , then  $\langle P, P' \rangle \in \mathcal{P}_H$ .*

*The relation  $P \sim P'$  defined by  $\langle P, P' \rangle \in \mathcal{P}_H$  is an equivalence relation on the set  $\mathcal{E}$  of non-trivial universally peripheral subgroups.*

*Proof.* To prove the first assertion, let  $\hat{\mathcal{F}}$  be a maximal  $H$ -invariant almost free factor system. Let  $F, F'$  be the factors in  $\hat{\mathcal{F}}$  containing  $P$  and  $P'$  respectively. Since  $F \cap F'$  contains  $P \cap P'$ , malnormality of  $\hat{\mathcal{F}}$  implies that  $F = F'$ . Thus  $\langle P, P' \rangle$  is  $\hat{\mathcal{F}}$ -peripheral. Since this holds for all  $\hat{\mathcal{F}}$ , the first assertion follows.

We deduce the transitivity of the relation  $\sim$ . If  $P \sim P'$  and  $P' \sim P''$ , then  $\langle P, P' \rangle$  and  $\langle P', P'' \rangle$  are universally peripheral and have non-trivial intersection, so  $\langle P, P', P'' \rangle \in \mathcal{P}_H$  by the first assertion.  $\square$

The lemma implies that for any  $P \in \mathcal{E}$ , there exists a unique  $\hat{P} \in \mathcal{P}_H^{\max}$  such that  $P \subseteq \hat{P}$ , and  $P \sim P'$  if and only if  $\hat{P} = \hat{P}'$ .

Notice that if  $P, P' \in \mathcal{E}$  with  $P \sim P'$ , then  $\langle P, P' \rangle$  is elliptic in any  $F_N$ -tree relative to  $\mathcal{P}_H$ . It thus follows from [GL11, Lemma 3.2] that this equivalence relation is admissible relative to  $\mathcal{P}_H$ , and can therefore be used to construct the tree of cylinders of any tree  $U$  relative to  $\mathcal{P}_H$  with edge groups in  $\mathcal{E}$  (see Section 5.2): cylinders are subtrees of  $U$  defined by the partition of edges induced by the equivalence relation  $G_e \sim G_{e'}$ . One then defines the tree of cylinders  $U^c$  of  $U$  as the tree having one vertex  $v_Y$  for each cylinder  $Y \subseteq U$ , one vertex  $v_x$  for each vertex  $x \in U$  belonging to at least two cylinders, with an edge between  $v_x$  and  $v_Y$  if and only if  $x \in Y$ . We denote by  $V_p = \{v_Y\}$  and  $V_a = \{v_x\}$  the two sets of vertices of  $U^c$ .

**Lemma 8.29.** *Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a subgroup, let  $U$  be a splitting of  $F_N$  relative to  $\mathcal{P}_H$  with edge stabilizers in  $\mathcal{E}$ , and let  $U^c$  be its tree of cylinders, coming with the bipartition  $V = V_a \sqcup V_p$  of its vertex set.*

*For every  $v \in V_p$ , one has  $G_v \in \mathcal{P}_H^{\max}$ , and  $G_v \cap G_{v'} = \{1\}$  if  $v \neq v' \in V_p$ . On the other hand, if  $v \in V_a$ , then  $G_v$  is not universally peripheral.*

*Moreover,  $U$  and  $U^c$  are in the same deformation space (i.e. every vertex stabilizer of  $U^c$  fixes a vertex in  $U$ ).*

*Proof.* Consider the cylinder  $Y_P$  corresponding to all edge groups contained in a given subgroup  $P \in \mathcal{P}_H^{\max}$ . Then the stabilizer of  $Y_P$  is the normalizer of  $P$ , which coincides with  $P$ . In particular,  $G_{v_{Y_P}} = P \in \mathcal{P}_H^{\max}$ . If  $Y, Y'$  are two distinct cylinders corresponding to  $P \neq P' \in \mathcal{P}_H^{\max}$ , then  $G_{v_Y} \cap G_{v_{Y'}} = P \cap P' = \{1\}$ .

If the stabilizer  $G_x$  of a vertex  $x \in U$  is universally peripheral, then all incident edge groups are in the same equivalence class so  $x$  lies in a unique cylinder and cannot correspond to a vertex in  $V_a$ .

Since every subgroup  $P \in \mathcal{P}_H^{\max}$  is elliptic in  $U$ , all vertex stabilizers of  $U^c$  fix a point in  $U$ , and it follows that  $U$  and  $U^c$  are in the same deformation space.  $\square$

*Proof of Theorem 8.15.* We first prove the uniqueness (see Proposition 8.34 for an alternative proof). Let  $U, U'$  be as in the theorem. We first prove that  $U$  and  $U'$  are in the same deformation space, i.e. that they have the same elliptic subgroups. For any  $u \in V_p(U)$ ,  $G_u$  is universally peripheral hence elliptic in  $U'$ . For  $v \in V_a(U)$ ,  $G_v$  is not universally peripheral by Assertion (4). Since  $G_v$  is a free factor by (2), and since its conjugacy class is  $H$ -invariant, (3) implies that  $G_v$  contains the stabilizer  $G_{v'}$  of a vertex  $v' \in V_a(U')$ . Symmetrically  $G_{v'}$  contains  $G_{v''}$  for some  $v'' \in V_a(U)$ , which imposes  $G_v = G_{v''}$  hence  $G_v = G_{v'}$ . This concludes that  $U$  and  $U'$  are in the same deformation space.

To conclude that  $U = U'$ , we claim that  $U$  and  $U'$  are their own trees of cylinders. If  $e_1, e_2$  are two edges of  $U$  with  $G_{e_1} \sim G_{e_2}$ , let  $p_i$  denote the endpoint of  $e_i$  in  $V_p$ . Then  $G_{p_1} \sim G_{e_1} \sim G_{e_2} \sim G_{p_2}$  so  $\langle G_{p_1}, G_{p_2} \rangle$  is universally peripheral. Since  $G_{p_i} \in \mathcal{P}_H^{\max}$ , we get  $G_{p_1} = \langle G_{p_1}, G_{p_2} \rangle = G_{p_2}$  hence  $p_1 = p_2$  by Assertion (1). This shows that each cylinder is the star of a  $V_p$ -vertex so that  $U$  is its own tree of cylinders, and so is  $U'$  by the same argument. By [GL11, Corollary 4.10], it follows that  $U$  and  $U'$  are isomorphic.



We now construct  $U_H^d$ . Choose a numbering  $\hat{\mathcal{F}}_1, \hat{\mathcal{F}}_2, \dots$  of the set of maximal  $H$ -invariant almost free factor systems (we do not know yet that it is finite), and let  $T_i = T_{\hat{\mathcal{F}}_i}$  and  $A_i = A_{\hat{\mathcal{F}}_i}$ . By Corollary 8.25, edge stabilizers of  $T_i$  are universally peripheral. We start with  $U_1 = T_1$  and construct inductively an  $H$ -invariant tree  $U_i$  with edge groups in  $\mathcal{P}_H$  and such that a subgroup  $K \subseteq F_N$  is elliptic in  $U_i$  if and only if it is elliptic in each of the trees  $T_1, \dots, T_i$ . Since  $H$  preserves no free splitting, edge stabilizers are necessarily non-trivial. Since edge stabilizers of  $U_{i-1}$  are elliptic in  $T_i$  one can construct a blowup  $U_i$  of  $U_{i-1}$  that dominates  $T_i$ : replace each vertex  $v$  such that  $G_v$  is not elliptic in  $T_i$  by the minimal  $G_v$ -invariant subtree of  $T_i$ , and attach every edge  $e$  of  $U_{i-1}$  incident on  $v$  to a point  $p_e$  fixed by  $G_e$ . This can be done so that  $U_i$  is  $H$ -invariant by choosing  $p_e$  in a canonical way: if  $G_e$  fixes a point in  $V^0(T_i)$ , then it is unique by Lemma 8.20(1), and one can choose  $p_e$  to be this vertex; otherwise,  $G_e$  fixes no edge hence fixes a unique vertex in  $V^1(T_i)$ . This concludes the construction of  $U_i$ .

Given any  $i$  and  $j$ , the group  $A_j$  fixes a unique vertex  $v_{A_j} \in U_i$ : the existence of  $v_{A_j}$  follows from the fact that  $A_j$  is elliptic in all trees  $T_\ell$  (Lemma 8.24), and its uniqueness follows from the fact that  $A_j$  is not universally peripheral, so cannot fix an edge.

Let  $U_i^c$  be the tree of cylinders of  $U_i$ . We are going to prove that  $T_j$  is a collapse of  $U_i^c$  for all  $j \leq i$ . We construct for all  $j \leq i$  a map  $f_{ij} : U_i^c \rightarrow T_j$  as follows. For every vertex  $v \in U_i^c$ , the group  $G_v$  is elliptic in  $U_i$  by Lemma 8.29, hence also in  $T_j$ . If  $G_v$  fixes a vertex  $v' \in V^0(T_j)$ , then this vertex is unique because the stabilizers of two distinct vertices in  $V^0(T_j)$  have trivial intersection. If  $G_v$  fixes no vertex in  $V^0(T_j)$ , then it fixes a necessarily unique vertex  $v' \in V^1(T_j)$ . In both cases, we let  $f_{ij}(v) = v'$ . We extend  $f_{ij}$  linearly on edges. Notice that  $f_{ij}$  is  $F_N$ -equivariant, so its image is an  $F_N$ -invariant subtree of  $T_j$ ; the minimality of  $T_j$  thus implies that  $f_{ij}$  is surjective.

(i) We first claim that if  $v_1, v_2 \in V_a(U_i^c)$  are such that  $f_{ij}(v_1) = f_{ij}(v_2) = v'$  with  $v' \in V^1(T_j)$ , then  $v_1 = v_2$ . Indeed,  $G_{v'}$  is a conjugate of  $A_j$  and therefore fixes a vertex  $w \in U_i^c$ . Now  $G_{v_1} \subseteq G_{v'} \subseteq G_w$ , and since  $G_{v_1}$  fixes no edge of  $U_i^c$  because  $G_{v_1} \notin \mathcal{P}_H$  (Lemma 8.29), it follows that  $v_1 = w$ . Symmetrically, one gets  $v_2 = w = v_1$ , which proves the claim.

(ii) We deduce that  $f_{ij}(V_p(U_i^c)) \subseteq V^0(T_j)$ . Indeed, arguing by contradiction, consider  $u \in V_p(U_i^c)$  such that  $f_{ij}(u) \in V^1(T_j)$ . By definition of  $f_{ij}$ , this means that  $G_u$  fixes no vertex in  $V^0(T_j)$ . Let  $e$  be an edge incident on  $u$ . If  $G_e$  fixes a vertex  $w' \in V^0(T_j)$ , then  $G_u \cap G_{w'} \neq \{1\}$ , and since  $[G_{w'}] \in \hat{\mathcal{F}}_j$  and  $G_u$  is universally peripheral (Lemma 8.29), it follows that  $G_u \subseteq G_{w'}$ , a contradiction. This shows that no edge incident on  $u$  fixes a vertex in  $V^0(T_j)$ , so  $f_{ij}$  collapses the star of  $u$  to  $f_{ij}(u)$ , contradicting (i) and concluding the proof of (ii).

(iii) We now claim that the image of an edge  $e \subseteq U_i^c$  under  $f_{ij}$  is either an edge or a vertex. Write  $e = uv$  with  $u \in V_p(U_i^c)$  and  $v \in V_a(U_i^c)$ . By (ii),  $f_{ij}(u) \in V^0(T_j)$ . Since stabilizers of distinct vertices in  $V^0(T_j)$  intersect trivially and  $G_e \neq \{1\}$ , it follows that  $f_{ij}(e)$  is contained in the star of  $f_{ij}(u)$  which proves (iii).

To prove that  $f_{ij}$  is a collapse map, consider  $e_1 = u_1v_1$ ,  $e_2 = u_2v_2$  two distinct edges of  $U_i^c$  that are mapped to the same edge  $e' = u'v'$  with  $u' = f_{ij}(u_1) = f_{ij}(u_2) \in V^0(T_j)$  and  $v' = f_{ij}(v_1) = f_{ij}(v_2) \in V^1(T_j)$ . By (ii),  $v_1, v_2 \in V_a(U_i^c)$  hence  $u_1, u_2 \in V_p(U_i^c)$ . By



(i), it follows that  $v_1 = v_2$ . Since  $\langle G_{e_1}, G_{e_2} \rangle$  fixes an edge in  $T_j$ , it is universally peripheral (Corollary 8.25). It follows that  $G_{u_1} \sim G_{e_1} \sim \langle G_{e_1}, G_{e_2} \rangle \sim G_{e_2} \sim G_{u_2}$ , and  $U_i^c$  being a tree of cylinders, this implies  $u_1 = u_2$ . This shows that no pair of noncollapsed edges gets identified by  $f_{ij}$ . This concludes that  $f_{ij}$  is a collapse map.

Additionally, for any vertex  $v \in f_{ij}^{-1}(V^1(T_j))$ , the map  $f_{ij}$  induces a bijection between the star of  $v$  and the star of  $f_{ij}(v)$ . Indeed, as  $f_{ij}$  is surjective, it suffices to show that it does not collapse any edge  $uv$  incident on  $v$ , but if it did, then  $f_{ij}(u) \in V^1(T_j)$  would contradict (ii).

By Lemma 8.20 (3), this shows that the incident edge groups at  $v_{A_j}$  form a non-sporadic free factor system of  $A_j$ . Refining  $U_i^c$  at each vertex  $v_{A_j}$  into a non-sporadic free splitting gives an a priori bound on  $i$  (precisely  $i \leq \max\{1, \frac{N-1}{2}\}$ , see Corollary 8.30), and shows that there are only finitely many maximal  $H$ -invariant almost free factor systems, so the process stops.

We define  $U_H^d = U_i^c$  as the last constructed tree, and we denote by  $f_j = f_{ij} : U_H^d \rightarrow T_j$ . The tree  $U_H^d$  is relative to  $\mathcal{P}_H$  because every  $T_j$  is. Assertion (1) follows from the properties of the trees of cylinders (Lemma 8.29).

To prove (4), we check that each vertex  $v \in V_a(U_H^d)$  is in the orbit of some  $v_{A_j}$ . Indeed,  $G_v \notin \mathcal{P}_H$  so  $G_v$  is not  $\hat{\mathcal{F}}_j$ -peripheral for some  $j$ . It follows that up to changing  $v$  to a translate, the image of  $v$  in  $T_j$  under  $f_j$  is the vertex fixed by  $A_j$ , so  $G_v \subseteq A_j$ . Since  $A_j$  fixes a vertex in  $U_H^d$  and  $G_v$  fixes no edge, it follows that  $G_v = A_j$ . This proves (4).

We now prove Assertion (3). If  $A$  is a free factor whose conjugacy class is  $H$ -invariant and which is not universally peripheral, then  $A$  is not  $\hat{\mathcal{F}}_j$ -peripheral for some  $j$ , so  $A$  contains  $A_j$  by definition of  $A_j = A_{\hat{\mathcal{F}}_j}$ . Since  $A_j = G_{v_{A_j}}$ , this proves Assertion (3).

We finally prove Assertion (2). We first note that  $(\mathcal{P}_H^{\max})|_{A_j} = (\hat{\mathcal{F}}_j)|_{A_j}$  because  $A_j$  is  $\hat{\mathcal{F}}_{j'}$ -peripheral for all  $j' \neq j$  by Lemma 8.24. By Lemma 8.20(3), this is a non-sporadic almost free factor system, and we denote it by  $\hat{\mathcal{G}}$ . To prove purity, consider a proper free factor  $A \subseteq A_j$  whose conjugacy class is  $H$ -invariant. The minimality of  $A_{\hat{\mathcal{F}}_j}$  implies that  $A$  is  $\hat{\mathcal{F}}_j$ -peripheral hence  $\hat{\mathcal{G}}$ -peripheral. This implies that  $\hat{\mathcal{G}}$  is the unique maximal  $H$ -invariant almost free factor system of  $A_j$ . As already noted above, the incident edge groups form a non-sporadic free factor system of  $A_j$  which concludes the proof of Assertion (2) and of the theorem.  $\square$

## 8.5 Finiteness of maximal invariant almost free factor systems

We deduce from the previous results a bound on the number of maximal invariant almost free factor systems that will be useful in the sequel.

**Corollary 8.30.** *Let  $N \geq 2$ . For every subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  that does not preserve any free splitting of  $F_N$ , there are at most  $\max\{1, \frac{N-1}{2}\}$  maximal  $H$ -invariant almost free factor systems of  $F_N$ .*

*Consequently, under the same hypothesis on  $H$ , there are at most  $\frac{(N+1)(N-1)}{2}$  maximal  $H$ -invariant free factor systems of  $F_N$ .*

*Remark 8.31.* As already mentioned, Example 6.1 shows that Corollary 8.30 does not hold true in general if we remove the assumption that  $H$  does not preserve any free splitting of  $F_N$ .

*Proof.* Let  $U_H^d$  be the splitting of  $F_N$  associated to  $H$  given by Theorem 8.15, coming with the bipartition  $V_p \sqcup V_a$  of its vertex set. Assertion (4) from this theorem ensures that the number of maximal  $H$ -invariant almost free factor systems of  $F_N$  is equal to the cardinality  $K$  of  $V_a/F_N$ . We may assume that  $K \geq 2$ , and aim to prove that  $K \leq \frac{N-1}{2}$ .

Recall that stabilizers of edges of  $U_H^d$  are non-trivial because there is no  $H$ -invariant free splitting. By the last assertion of Theorem 8.15(2), for each vertex  $v \in V_a/F_N$  there is a free splitting of  $G_v$  relative to the incident edge groups with all vertex groups non-trivial and at least two orbits of edges (indeed, the relative Grushko decomposition is non-sporadic and has at least one non-trivial vertex group because there is at least one edge incident on  $v$ , as  $K \geq 2$ ). Blowing up every vertex in  $V_a/F_N$  using this free splitting and collapsing all edges coming from  $U_H^d/F_N$  gives a free splitting of  $F_N$  with at least  $2K$  edges and all vertex groups non-trivial. It follows that  $2K \leq N - 1$ .

For  $N = 2$ , any non-empty free factor system is sporadic, so the empty set is the unique maximal  $H$ -invariant free factor system. Otherwise, since any maximal  $H$ -invariant free factor system can be extracted from a maximal  $H$ -invariant almost free factor system by Lemma 8.8, and since an almost free factor system of  $F_N$  contains of at most  $(N + 1)$  conjugacy classes of cyclic groups, there are at most  $(N + 1) \frac{N-1}{2}$  maximal  $H$ -invariant free factor systems.  $\square$

## 8.6 Reducibility is inherited by the normalizer

The following result will not be used in the rest of the paper, but we believe that it is of independent interest.

**Theorem 8.32.** *Let  $H \subseteq \text{Out}(F_N)$  be an infinite subgroup. If  $H$  has a periodic conjugacy class of proper free factor, so does its normalizer in  $\text{Out}(F_N)$ .*

*Proof.* Up to replacing  $H$  by  $H \cap \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , we may assume that  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ .

Assume first that there is no  $H$ -invariant non-trivial free splitting. Then by Corollary 8.30, there are only finitely many maximal  $H$ -invariant free factor systems, and all of them are non-empty by assumption. The normalizer of  $H$  permutes them and the theorem follows in this case.

If there exists a non-trivial  $H$ -invariant free splitting, then by Proposition 6.2 there exists a  $H$ -invariant free splitting  $S$  which is maximum for domination among all  $H$ -invariant free splittings, and all such splittings are non-trivial and belong to the same deformation space. Let  $\mathcal{F}$  be the collection of conjugacy classes of non-trivial vertex stabilizers of  $S$ . Then  $\mathcal{F}$  is a free factor system, and  $\mathcal{F}$  is non-empty because otherwise, the stabilizer of  $S$  would be finite, contradicting that  $H$  is infinite. Since  $\mathcal{F}$  is clearly invariant under the normalizer of  $H$ , this concludes the proof.  $\square$

## 8.7 Additional properties

We conclude this section by establishing a few additional properties of the splitting  $U_H^d$ , which will not be used in the sequel of the paper.

**Proposition 8.33.** *Keeping the notations from Theorem 8.15, the following hold:*

- (a) *All edge and vertex stabilizers of  $U_H^d$  are free factors of  $F_N$ .*
- (b) *For every  $v \in V_p$ , the Grushko decomposition of  $G_v$  relative to its incident edge groups is trivial.*

*Proof.* We first claim that for every  $v \in V_p$  and every maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}$ ,  $G_v$  is not contained in an unused boundary subgroup  $\langle b \rangle$  of  $\hat{\mathcal{F}}$ . Otherwise, let  $v_{\hat{\mathcal{F}}}$  be a vertex in  $U_H^d$  whose orbit is sent to  $\hat{\mathcal{F}}$  under the bijection given in Theorem 8.15(4). Since collapsing all edges not adjacent to a vertex in the orbit of  $v_{\hat{\mathcal{F}}}$  yields the tree  $T_{\hat{\mathcal{F}}}$ , we deduce that  $\langle b \rangle$  fixes a vertex in  $V_0(T_{\hat{\mathcal{F}}})$ , contradicting Lemma 8.20(1).

To prove Assertion (a), it suffices to check that vertex stabilizers are free factors. We already know this for vertices in  $V_a$ , so consider  $v \in V_p$ . Since  $G_v \in \mathcal{P}_H^{\max}$ , for each maximal  $H$ -invariant almost free factor system  $\hat{\mathcal{F}}$ , there is a factor of  $\hat{\mathcal{F}}$  containing  $G_v$ , and the claim above says that this factor is not an unused boundary subgroup. The intersection of these free factors as  $\hat{\mathcal{F}}$  varies is a free factor  $G'_v$  containing  $G_v$  and which is universally peripheral. Since  $G_v \in \mathcal{P}_H^{\max}$ , we have  $G_v = G'_v$ , which proves Assertion (a).

To prove Assertion (b), consider  $v \in V_p$ . If the Grushko decomposition of  $G_v$  relative to its incident edge groups is non-trivial, then one can use it to blow up  $U_H^d$  at the vertex  $v$ , and thus get an  $H$ -invariant free factor system  $\mathcal{G}$  of  $F_N$ . Consider a maximal  $H$ -invariant free factor system  $\mathcal{F}$  such that  $\mathcal{G} \preceq \mathcal{F}$ . By construction, for every vertex of  $U_H^d$  not in the orbit of  $v$ , its stabilizer is  $\mathcal{G}$ -peripheral hence  $\mathcal{F}$ -peripheral. Let  $\hat{\mathcal{F}}$  be the unique maximal  $H$ -invariant almost free factor system containing  $\mathcal{F}$ . Being universally peripheral,  $G_v$  is  $\hat{\mathcal{F}}$ -peripheral and the claim above shows that  $G_v$  is in fact  $\mathcal{F}$ -peripheral. It follows that there is a domination map  $U_H^d \rightarrow R_{\mathcal{F}}$  where  $R_{\mathcal{F}}$  is a (non-trivial) free splitting whose elliptic subgroups are the elements of  $\mathcal{F}$ . As all edge stabilizers of  $U_H^d$  are non-trivial, this map has to be constant, contradicting that  $R_{\mathcal{F}}$  is non-trivial. This contradiction concludes the proof.  $\square$

We finally give a more intrinsic description of  $U_H^d$  by defining a new graph  $S$  as follows. Let  $V'_a$  be the collection of minimal free factors of  $F_N$  whose conjugacy class is  $H$ -invariant and which are not universally peripheral, i.e. the collection of subgroups that are conjugate to some  $A_{\hat{\mathcal{F}}}$  where  $\hat{\mathcal{F}}$  is a maximal  $H$ -invariant almost free factor system. Let  $V'_p$  be the set of subgroups  $P \in \mathcal{P}_H^{\max}$  such that there exist at least two distinct subgroups  $A, A' \in V'_a$  that intersect  $P$  non-trivially. We define  $S$  as the bipartite graph with vertex set  $V'_a \cup V'_p$  and with an edge between  $A \in V'_a$  and  $P \in V'_p$  if  $A \cap P \neq \{1\}$ . There is a natural action of  $F_N$  on  $S$  by conjugation.

**Proposition 8.34.** *The graph  $S$  is equivariantly isomorphic to  $U_H^d$ .*

*Proof.* For each  $v \in V_a$ , we have  $G_v \in V'_a$ , and for each  $v \in V_p$ , we have  $G_v \in V'_p$  because edge stabilizers of  $U_H^d$  are non-trivial. The assignment  $v \mapsto G_v$  then defines a map  $f$  from the vertices of  $U_H^d$  to the vertices of  $S$ .

Since edge stabilizers of  $U_H^d$  are non-trivial,  $f$  preserves adjacency. The map  $f$  is injective on  $V_p$  by Theorem 8.15(1). We claim that it is also injective on  $V_a$ . Indeed, if  $v, v' \in V_a$  are not in the same orbit, then  $G_{v'}$  is  $\hat{\mathcal{F}}_v$ -peripheral but  $G_v$  is not by Theorem 8.15(4), so  $G_v \neq G_{v'}$ . If  $v' = gv$  and  $G_{v'} = G_v$ , then  $g \in G_v$  by malnormality of the free factor  $G_v$ , so  $v' = v$ . This proves our claim that  $f$  is injective on  $V_a$ . Since groups in  $V'_p$  are universally peripheral and groups in  $V'_a$  are not,  $V'_p \cap V'_a = \emptyset$  so  $f$  is injective.

To check that  $f$  preserves non-adjacency, consider  $u \in V_p$  and  $v \in V_a$  which are not adjacent in  $U_H^d$ . Then they are at distance at least 3 and the segment  $[u, v]$  contains a vertex  $u' \in V_p$  distinct from  $u$ . Since  $G_u \cap G_{u'} = \{1\}$  by Theorem 8.15(1), it follows that  $G_u \cap G_v = \{1\}$  so  $f(u)$  is not adjacent to  $f(v)$ .

We finally check that  $f$  is onto. It is clear that  $f(V_a) = V'_a$ . So consider  $P \in V'_p$ . Since  $U_H^d$  is relative to  $\mathcal{P}_H$ , the group  $P$  fixes a vertex  $v \in U_H^d$ . If one can choose  $v \in V_p$ , then  $P = G_v$  because  $P \in \mathcal{P}_H^{\max}$  so  $P = f(v)$ . If  $P$  fixes no vertex in  $V_p$ , then  $v \in V_a$  and for any  $v' \in V_a \setminus \{v\}$ , one has  $P \cap G_{v'} = \{1\}$ . This shows that  $G_v$  is the unique group in  $V'_a$  that  $P$  intersects non-trivially, so  $P$  does not fulfill the requirements to lie in  $V'_p$ . This shows that  $f$  is surjective and defines an isomorphism  $U_H^d \rightarrow S$ .  $\square$

## 9 Canonical nice splittings

In sections 6, 7, and 8 we constructed canonical splittings using invariant free splittings, invariant  $\mathcal{Z}_{\max}$ -splittings, and maximal invariant free factor systems. The goal of this section is to unify them using the following notion of nice splittings. We let  $N \geq 2$ .

**Definition 9.1** (Nice and bi-nonsporadic splittings). *A non-trivial splitting  $S$  of  $F_N$  is nice if either*

- $S$  is a free splitting
- $S$  is a  $\mathcal{Z}_{\max}$ -splitting
- $S$  is bi-nonsporadic in the following sense: its edge stabilizers are finitely generated non-abelian, and there are at least two vertices  $v_1, v_2 \in S$  in different  $F_N$ -orbits such that for every  $i \in \{1, 2\}$ , the Grushko decomposition of  $G_{v_i}$  relative to the incident edge groups is non-sporadic.

We say that a collection  $\mathcal{C}$  of almost free factor systems is *filling* there is no almost free factor system  $\hat{\mathcal{F}}$  such that for every  $\hat{\mathcal{F}}' \in \mathcal{C}$ , one has  $\hat{\mathcal{F}}' \preceq \hat{\mathcal{F}}$ .

The main result we will prove is the following one.

**Theorem 9.2.** *There exists an  $\text{Out}(F_N)$ -equivariant map assigning a (canonical) nice splitting  $U_{\mathcal{C}}$  to  $\mathcal{C}$  when  $\mathcal{C}$  is either*

1. a non-empty collection of non-trivial free or  $\mathcal{Z}_{\max}$ -splittings whose elementwise stabilizer is infinite;
2. a filling collection of almost free factor systems whose elementwise stabilizer is infinite;
3. a non-empty collection of nice splittings whose elementwise stabilizer is infinite.

The following corollary will not be directly used in this paper, but it is good to have it in mind in preparation for the next section, where we will establish groupoid-theoretic analogues that will be crucial in the proof of our main measure equivalence rigidity theorem.

**Corollary 9.3.** *If  $H \subseteq \text{Out}(F_N)$  is an infinite group preserving a nice splitting, then its normalizer preserves a nice splitting.*

*Proof.* Let  $\mathcal{C}$  be the collection of all nice splittings preserved by  $H$ , and let  $U_{\mathcal{C}}$  be the associated nice splitting. Since  $\mathcal{C}$  is invariant under  $N(H)$ , so is  $U_{\mathcal{C}}$ .  $\square$

We prove several intermediate results before proving the theorem.

In Section 6.3, we constructed from invariant free splittings a canonical splitting which has an edge with trivial or  $\mathcal{Z}_{\max}$  stabilizer or is biflexible. Recall from Definition 6.14 that that  $(A, \mathcal{P})$  is *flexible* if  $\text{Out}(A, \mathcal{P}^{(t)})$  is infinite. A splitting  $S$  of  $F_N$  is *biflexible* if all its edge stabilizers are finitely generated non-abelian and there exist two vertices  $v_1, v_2$  in different  $F_N$ -orbits such that for every  $i \in \{1, 2\}$ ,  $(G_{v_i}, \text{Inc}_{v_i})$  is flexible.

**Lemma 9.4.** *Let  $A$  be a free group and  $\mathcal{P}$  a non-empty collection of conjugacy classes of non-abelian subgroups. Assume that  $(A, \mathcal{P})$  is flexible. Then one of the following holds:*

1. the Grushko decomposition of  $A$  relative to  $\mathcal{P}$  is non-sporadic;
2. there is a canonical non-trivial free splitting of  $A$  relative to  $\mathcal{P}$  —in particular it is invariant under  $\text{Out}(A, \mathcal{P})$ ;
3.  $A$  is one-ended relative to  $\mathcal{P}$  and the canonical  $\mathcal{Z}_{\max}$  JSJ decomposition of  $A$  relative to  $\mathcal{P}$  is non-trivial.

*Proof.* Assume that (1) does not hold. If the Grushko decomposition of  $A$  relative to  $\mathcal{P}$  is non-trivial (hence sporadic), then it contains a unique reduced free splitting (this is the splitting with a single orbit of edges), so Assertion (2) holds. Thus, we may assume that  $A$  is one-ended relative to  $\mathcal{P}$ . Since  $\text{Out}(A, \mathcal{P}^{(t)})$  is infinite, there exists a non-trivial  $\mathcal{Z}_{\max}$ -splitting of  $A$  relative to  $\mathcal{P}$  by Paulin's argument and Rips theory ([Pau91, BF95], see for instance [GL15b, Theorem 7.14] for this precise statement). If the  $\mathcal{Z}_{\max}$  JSJ decomposition of  $A$  relative to  $\mathcal{P}$  is trivial, then the pair  $(A, \mathcal{P})$  has to be QH with sockets (see [Sel97] or Proposition 7.18) and in particular, every group in  $\mathcal{P}$  is cyclic. This is a contradiction.  $\square$

**Corollary 9.5.** *To any biflexible splitting of  $F_N$ , one can associate canonically (in an  $\text{Out}(F_N)$ -equivariant way) a nice splitting.*

*Proof.* Let  $S$  be a biflexible splitting, and let  $v_1, \dots, v_k$  be the collection of flexible vertices, with  $k \geq 2$ . We construct the canonical nice splitting  $S'$  as follows.

If there are at least two vertices  $v_i$  such that the Grushko decomposition of  $(G_{v_i}, \text{Inc}_{v_i})$  is non-sporadic, then  $S$  is bi-nonsporadic and we take  $S' = S$ . Thus, we may assume that for some  $i \leq k$ , the Grushko decomposition of  $(G_{v_i}, \text{Inc}_{v_i})$  is trivial or sporadic.

If the Grushko decomposition of  $(G_{v_i}, \text{Inc}_{v_i})$  is non-trivial but sporadic for some  $i$ , then consider  $\hat{S}$  obtained from  $S$  by blowing up each such vertex  $v_i$  into the corresponding canonical free splitting of  $G_{v_i}$  relative to  $\text{Inc}_{v_i}$  (there is no choice in the attaching points of the edges of  $S$  because the groups in  $\text{Inc}_{v_i}$  are non-trivial). Collapsing all edges of  $\hat{S}$  coming from  $S$  yields a non-trivial free splitting  $S'$  that is invariant under the stabilizer of  $S$ .

If the Grushko decomposition of  $(G_{v_i}, \text{Inc}_{v_i})$  is trivial for some  $i \leq k$ , then the canonical  $\mathcal{Z}_{\max}$ -JSJ decomposition of  $G_{v_i}$  relative to  $\text{Inc}_{v_i}$  is non-trivial by Lemma 9.4. So consider  $\hat{S}$  obtained from  $S$  by blowing up each such vertex  $v_i$  using this JSJ decomposition (there is no choice in the attaching points of the edges of  $S$  because the groups in  $\text{Inc}_{v_i}$  are non-abelian). Collapsing all edges of  $\hat{S}$  coming from  $S$  yields a non-trivial  $\mathcal{Z}_{\max}$ -splitting  $S'$  that is invariant under the stabilizer of  $S$ .  $\square$

**Lemma 9.6.** *To any bi-nonsporadic splitting of  $F_N$ , one can canonically associate a filling finite set of non-empty free factor systems.*

*In particular, a subgroup  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  preserving a bi-nonsporadic splitting of  $F_N$  cannot be pure.*

*Proof.* Let  $S$  be a bi-nonsporadic splitting, and  $v_1, \dots, v_n$  with  $n \geq 2$  be the collection of vertices of  $S$  such that the Grushko decomposition of  $(G_{v_i}, \text{Inc}_{v_i})$  is non-sporadic. For each  $i \leq n$ , let  $\hat{S}_i$  be obtained from  $S$  by blowing up  $v_i$  into a Grushko decomposition of  $G_{v_i}$  relative to  $\text{Inc}_{v_i}$  and let  $S'_i$  be the free splitting obtained by collapsing all edges coming from  $S$ . Finally, let  $\mathcal{F}_i$  be the free factor system consisting of non-trivial vertex stabilizers of  $S'_i$ . Note that  $\mathcal{F}_i$  does not depend on the choice of the Grushko decomposition used to blow up  $S$ .

We claim that  $\{\mathcal{F}_1, \dots, \mathcal{F}_n\}$  is filling. Otherwise, consider a free factor system  $\mathcal{F}$  such that  $\mathcal{F}_i \preceq \mathcal{F}$  for all  $i \leq n$ . Note that for every vertex  $v \in S$  not in the orbit of  $v_i$ , the group  $G_v$  is  $\mathcal{F}_i$ -peripheral. Since  $n \geq 2$ , this implies that  $G_v$  is  $\mathcal{F}$ -peripheral for every vertex  $v$  of  $S$ . Let  $R$  be a non-trivial free splitting whose conjugacy classes of non-trivial elliptic subgroups are precisely the groups in  $\mathcal{F}$ . Since vertex stabilizers of  $S$  are elliptic in  $R$ , there is an equivariant map  $f : S \rightarrow R$ . Since edge stabilizers of  $R$  are trivial,  $f$  maps every edge of  $S$  to a point, so  $f$  is constant. Since  $R$  is non-trivial, this is a contradiction and concludes the proof.  $\square$

We now prove the main result of this section.

*Proof of Theorem 9.2.* Note that it suffices to assign to  $\mathcal{C}$  a splitting which is either bi-nonsporadic or which has an edge with trivial or  $\mathcal{Z}_{\max}$  stabilizer. Indeed, in the latter

case, collapsing all edges with non-trivial stabilizer or all edges whose stabilizer is not  $\mathcal{Z}_{\max}$  accordingly yields a nice splitting.

Assume first that  $\mathcal{C}$  is a collection of non-trivial  $\mathcal{Z}_{\max}$  or free splittings whose elementwise stabilizer  $\Gamma_{\mathcal{C}}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  is infinite. If  $\Gamma_{\mathcal{C}}$  preserves a non-trivial free splitting, then Corollary 6.15 assigns to  $\Gamma_{\mathcal{C}}$  a canonical splitting  $U_{\Gamma_{\mathcal{C}}}^1$  which has an edge with trivial or  $\mathcal{Z}_{\max}$ -stabilizer, or which is biflexible. As noted above, we may assume that  $U_{\Gamma_{\mathcal{C}}}^1$  is biflexible, and we conclude using Corollary 9.5. If  $\Gamma_{\mathcal{C}}$  preserves no non-trivial free splitting, then  $\mathcal{C}$  consists of  $\mathcal{Z}_{\max}$ -splittings. Thus Theorem 7.4 applies to  $\Gamma_{\mathcal{C}}$  so  $U_{\mathcal{C}} := U_{\Gamma_{\mathcal{C}}}^z$  is a non-trivial  $\mathcal{Z}_{\max}$  splitting of  $F_N$  canonically associated to  $\Gamma_{\mathcal{C}}$ . This proves the first assertion.

We now consider the case where  $\mathcal{C}$  is a filling collection of almost free factor systems whose elementwise stabilizer  $\Gamma_{\mathcal{C}}$  is infinite. If  $\Gamma_{\mathcal{C}}$  preserves a non-trivial free splitting of  $F_N$ , then we consider  $\mathcal{C}_1 = \text{FS}^{\Gamma_{\mathcal{C}}}$  the collection of all  $\Gamma_{\mathcal{C}}$ -invariant non-trivial free splittings and define  $U_{\mathcal{C}} := U_{\mathcal{C}_1}$  using the first assertion.

We can therefore assume that  $\Gamma_{\mathcal{C}}$  preserves no sporadic free factor system. Let  $\mathcal{C}'$  be the collection of all maximal  $\Gamma_{\mathcal{C}}$ -invariant proper almost free factor systems. Note that for any  $\hat{\mathcal{F}}$  in  $\mathcal{C}$ , there is  $\hat{\mathcal{F}}' \in \mathcal{C}'$  with  $\hat{\mathcal{F}} \preceq \hat{\mathcal{F}}'$ . Since  $\mathcal{C}$  is filling, this implies that  $\#\mathcal{C}' \geq 2$ , i.e.  $\Gamma_{\mathcal{C}}$  is not pure (Definition 8.12). Let  $U_{\Gamma_{\mathcal{C}}}^d$  be the canonical splitting provided by Theorem 8.15. By Remark 8.18, it is non-trivial and either it has an edge with  $\mathcal{Z}_{\max}$  stabilizer or else it is bi-nonsporadic.

Finally, we consider the case where  $\mathcal{C}$  is a collection of nice splittings whose elementwise stabilizer  $\Gamma_{\mathcal{C}}$  is infinite. If  $\mathcal{C}$  contains a free splitting or a  $\mathcal{Z}_{\max}$ -splitting, one can define  $U_{\mathcal{C}}$  as  $U_{\mathcal{C}_1}$  where  $\mathcal{C}_1 = \mathcal{C} \cap (\text{FS} \cup \mathcal{Z}_{\max}\text{S})$ . Thus we may assume that  $\mathcal{C}$  consists of bi-nonsporadic splittings. Using Lemma 9.6, consider for each  $S \in \mathcal{C}$  the filling collection  $\mathcal{C}_S$  of free factor systems associated to  $S$ , and let  $\mathcal{C}' = \cup_{S \in \mathcal{C}} \mathcal{C}_S$ . Then  $\mathcal{C}'$  is a filling collection of free factor systems canonically associated to  $S$  and whose stabilizer contains  $\Gamma_{\mathcal{C}}$ . One can then define  $U_{\mathcal{C}} := U_{\mathcal{C}'}$ .  $\square$



## Part III

# Measure equivalence rigidity of $\text{Out}(F_N)$

## 10 Canonical invariant splittings for groupoids

In this section, we fix a finitely generated free group  $F_k$  with  $k \geq 2$ . In the sequel of the paper, we will usually apply the results of this section to the case where  $F_k = F_N$  with  $N \geq 3$ , but we will also need to consider the case where  $F_k$  is a corank one free factor of  $F_N$  in Section 13.4.

Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , equipped with a cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . The goal of the present section is to give a criterion ensuring (roughly) that if a subgroupoid  $\mathcal{H}$  has invariant splittings (with respect to the cocycle  $\rho$ ), then so does any subgroupoid that normalizes  $\mathcal{H}$  (see Proposition 10.6 below for a precise statement). This is done by using all the results obtained in Part II and transferring them to the groupoid setting. The chain conditions proved in the previous sections are crucial for this to work.

Recall from Definition 3.26 that a splitting  $S$  of  $F_k$  is  $(\mathcal{G}, \rho)$ -invariant if there is a conull Borel subset  $Y^* \subseteq Y$  such that  $\rho(\mathcal{G}_{|Y^*}) \subseteq \Gamma_S$ , where  $\Gamma_S$  denotes the stabilizer of  $S$  in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Recall that nice splittings were defined in Definition 9.1.

**Definition 10.1** (Stably nice, nice-averse and FS-averse cocycle). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle. We say that  $(\mathcal{G}, \rho)$  is*

- stably nice if there exists a partition  $Y = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets such that for every  $i \in I$ , there is a  $(\mathcal{G}_{|Y_i}, \rho)$ -invariant nice splitting of  $F_k$ ;
- nice-averse if for every Borel subset  $U \subseteq Y$  of positive measure, there is no  $(\mathcal{G}_{|U}, \rho)$ -invariant nice splitting of  $F_k$ .
- FS-averse if for every Borel subset  $U \subseteq Y$  of positive measure, there is no  $(\mathcal{G}_{|U}, \rho)$ -invariant non-trivial free splitting of  $F_k$ .

Note that a nice-averse cocycle is in particular FS-averse.

We extend the definition of a pure subgroup of  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  (Definition 8.12) to groupoids with a cocycle towards  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ .

**Definition 10.2** (Pure cocycle). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , equipped with a cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ .*

*We say that  $(\mathcal{G}, \rho)$  is pure if  $(\mathcal{G}, \rho)$  is FS-averse and there exists a  $(\mathcal{G}, \rho)$ -invariant almost free factor system  $\hat{\mathcal{F}}$  such that for every Borel subset  $U \subseteq Y$  of positive measure, and every  $(\mathcal{G}_{|U}, \rho)$ -invariant almost free factor system  $\hat{\mathcal{F}}'$ , one has  $\hat{\mathcal{F}}' \preceq \hat{\mathcal{F}}$ .*

*We say that  $(\mathcal{G}, \rho)$  is stably pure if there is a partition  $Y = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets such that for every  $i \in I$ ,  $(\mathcal{G}_{|Y_i}, \rho)$  is pure.*

*Remark 10.3.* If  $(\mathcal{G}, \rho)$  is pure, then the corresponding almost free factor system  $\hat{\mathcal{F}}$  is necessarily unique and it is non-sporadic. We also insist that, as usual, the almost free factor system  $\hat{\mathcal{F}} = \emptyset$  is allowed in this definition.

*Remark 10.4.* Note that if  $Y^* \subseteq Y$  is a conull Borel subset such that  $(\mathcal{G}|_{Y^*}, \rho)$  is stably nice or stably pure then so is  $(\mathcal{G}, \rho)$ .

The goal of this section is to prove the following two propositions, which are groupoid analogues of Theorem 9.2 and Corollary 9.3.

**Proposition 10.5.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle.*

*There exists a Borel partition  $Y = Y_1 \sqcup Y_2$  such that  $(\mathcal{G}|_{Y_1}, \rho)$  is stably nice, and  $(\mathcal{G}|_{Y_2}, \rho)$  is stably pure and nice-averse.*

Recall that a cocycle  $\rho$  is *nowhere trivial* if there is no Borel subset of positive measure of the base space of  $\mathcal{G}$  in restriction to which  $\rho$  is trivial (Definition 3.18).

**Proposition 10.6.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , with a cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Let  $\mathcal{H}, \mathcal{H}'$  be measured subgroupoids of  $\mathcal{G}$ . Assume that  $(\mathcal{H}, \rho)$  is stably nice and that  $\rho|_{\mathcal{H}}$  is nowhere trivial.*

*If  $\mathcal{H}'$  stably normalizes  $\mathcal{H}$ , then  $(\mathcal{H}', \rho)$  is stably nice.*

*More precisely, there exists a partition  $Y = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets such that for every  $i \in I$ , there exists a nice splitting  $S_i$  which is invariant under both  $(\mathcal{H}|_{Y_i}, \rho)$  and  $(\mathcal{H}'|_{Y_i}, \rho)$ .*

## 10.1 Stably nice vs pure partition

After a partition of the base space,  $(\mathcal{G}, \rho)$  is either stably nice or nice-averse, as shown in the following lemma.

**Lemma 10.7.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle.*

*There exists a Borel partition  $Y = Y_1 \sqcup Y_2$  such that  $(\mathcal{G}|_{Y_1}, \rho)$  is stably nice, and  $(\mathcal{G}|_{Y_2}, \rho)$  is nice-averse.*

*Proof.* Without loss of generality, we assume that the measure on  $Y$  is a probability measure (see Remark 3.10). We choose for  $Y_1$  a Borel subset of maximal measure such that  $(\mathcal{G}|_{Y_1}, \rho)$  is stably nice: this exists because if  $(Y_{1,n})_{n \in \mathbb{N}}$  is a measure-maximizing sequence of such sets, then letting  $Y_{1,\infty} = \bigcup Y_{1,n}$ , we have again that  $(\mathcal{G}|_{Y_{1,\infty}}, \rho)$  is stably nice. The proof is then completed by letting  $Y_2 = Y \setminus Y_1$ : the maximality of  $Y_1$  ensures that  $(\mathcal{G}|_{Y_2}, \rho)$  is nice-averse.  $\square$

To prove Proposition 10.5, we will show that a nice-averse cocycle is stably pure (Lemma 10.12 below). A key step will be Lemma 10.11, based on the chain condition, showing the existence of a collection of maximal invariant almost free factor systems, up to partitioning the base space. We first introduce a few definitions.

**Definition 10.8** (Everywhere maximal, dominant, stable reduction collection). *Let  $\mathcal{H}$  be a measured groupoid over a base space  $Y$ , let  $\rho : \mathcal{H} \rightarrow \mathrm{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle.*

*An  $(\mathcal{H}, \rho)$ -invariant almost free factor system  $\hat{\mathcal{F}}$  is  $(\mathcal{H}, \rho)$ -everywhere maximal if for every Borel subset  $U \subseteq Y$  of positive measure, there is no almost free factor system  $\hat{\mathcal{F}}' \not\preceq \hat{\mathcal{F}}$  which is  $(\mathcal{H}|_U, \rho)$ -invariant.*

*A collection  $\mathcal{C}$  of  $(\mathcal{H}, \rho)$ -invariant almost free factor systems is  $(\mathcal{H}, \rho)$ -dominant if for every Borel subset  $U \subseteq Y$  of positive measure and every  $(\mathcal{H}|_U, \rho)$ -invariant almost free factor system  $\hat{\mathcal{F}}$ , there exists  $\hat{\mathcal{F}}' \in \mathcal{C}$  such that  $\hat{\mathcal{F}} \preceq \hat{\mathcal{F}}'$ .*

*A collection  $\mathcal{C}$  of  $(\mathcal{H}, \rho)$ -invariant almost free factor systems is a stable reduction collection for  $(\mathcal{H}, \rho)$  if it is  $(\mathcal{H}, \rho)$ -dominant and if every  $\hat{\mathcal{F}} \in \mathcal{C}$  is  $(\mathcal{H}, \rho)$ -everywhere maximal.*

*Remark 10.9.* If  $(\mathcal{H}, \rho)$  has a stable reduction collection  $\mathcal{C}$ , then it is unique. Indeed, assume that  $\mathcal{C}'$  is another stable reduction collection for  $(\mathcal{H}, \rho)$ , and let  $\hat{\mathcal{F}} \in \mathcal{C}$ . As  $\hat{\mathcal{F}}$  is  $(\mathcal{H}, \rho)$ -invariant and  $\mathcal{C}'$  is  $(\mathcal{H}, \rho)$ -dominant, we deduce that there exists  $\hat{\mathcal{F}}' \in \mathcal{C}'$  such that  $\hat{\mathcal{F}} \preceq \hat{\mathcal{F}}'$ . As  $\hat{\mathcal{F}}$  is  $(\mathcal{H}, \rho)$ -everywhere maximal, it follows that  $\hat{\mathcal{F}} = \hat{\mathcal{F}}'$ . Hence  $\mathcal{C} \subseteq \mathcal{C}'$ , and by symmetry they are actually equal.

Moreover, for every Borel subset  $U \subseteq Y$  of positive measure,  $\mathcal{C}$  is also a stable reduction collection for  $(\mathcal{H}|_U, \rho)$ .

Note that  $(\mathcal{H}, \rho)$  is pure if and only if it is FS-averse and it has a stable reduction collection  $\mathcal{C}$  consisting of a single almost free factor system (which may, as usual, be the empty almost free factor system).

**Lemma 10.10.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$  with a cocycle  $\rho : \mathcal{G} \rightarrow \mathrm{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Let  $\mathcal{H}, \mathcal{H}'$  be two measured subgroupoids. Assume that  $\mathcal{C}$  is a stable reduction collection for  $(\mathcal{H}, \rho)$  and that  $\mathcal{H}'$  normalizes  $\mathcal{H}$ .*

*Then  $\mathcal{C}$  is  $(\mathcal{H}', \rho)$ -invariant.*

*If  $\mathcal{C}$  is finite, then every element of  $\mathcal{C}$  is  $(\mathcal{H}', \rho)$ -invariant.*

We mention that finiteness of  $\mathcal{C}$  is in fact automatic when  $(\mathcal{H}, \rho)$  is FS-averse, as will be established in Lemma 10.11 below.

*Proof.* By definition of normalization of subgroupoids, one can write  $\mathcal{H}'$  as a countable union of bisections  $B_n \subseteq \mathcal{H}'$  such that for all  $h \in \mathcal{G}$  and all  $\varphi, \psi \in B_n$  such that  $\varphi h \psi^{-1}$  is defined, one has  $h \in \mathcal{H}$  if and only if  $\varphi h \psi^{-1} \in \mathcal{H}$ . Up to subdividing the bisections, we may assume additionally that the cocycle  $\rho$  maps  $B_n$  to a single value  $\gamma_n \in \mathrm{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Let  $f_{B_n} : U_n \rightarrow V_n$  be the partial isomorphism associated to  $B_n$ , with  $U_n, V_n \subseteq Y$ .

Assume that  $U_n$  and  $V_n$  have positive measure. If  $\hat{\mathcal{F}}$  is any  $(\mathcal{H}|_{U_n}, \rho)$ -invariant almost free factor system, then  $\gamma_n \hat{\mathcal{F}}$  is  $(\mathcal{H}|_{V_n}, \rho)$ -invariant. It follows that if  $\hat{\mathcal{F}}$  is  $(\mathcal{H}|_{U_n}, \rho)$ -everywhere maximal,  $\gamma_n \hat{\mathcal{F}}$  is  $(\mathcal{H}|_{V_n}, \rho)$ -everywhere maximal. Similarly, since  $\mathcal{C}$  is a stable reduction collection for  $(\mathcal{H}|_{U_n}, \rho)$ ,  $\gamma_n \mathcal{C}$  is a stable reduction collection for  $(\mathcal{H}|_{V_n}, \rho)$ . By uniqueness of the stable reduction collection,  $\gamma_n \mathcal{C} = \mathcal{C}$  if  $U_n$  has positive measure.

Discarding the bisections with zero measure, since  $\mathcal{H}'$  is the union of the bisections  $B_n$ , it follows that  $\mathcal{C}$  is  $(\mathcal{H}', \rho)$ -invariant. Under the additional assumption that  $\mathcal{C}$  is

finite, the global stabilizer of  $\mathcal{C}$  in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  coincides with its elementwise stabilizer by Theorems 2.10 and 2.11. We thus conclude that every  $\hat{\mathcal{F}} \in \mathcal{C}$  is  $(\mathcal{H}', \rho)$ -invariant.  $\square$

The following lemma shows the existence of stable reduction collections up to partitioning the base space, and shows that such collections are finite under the assumption that  $(\mathcal{H}, \rho)$  is FS-averse.

**Lemma 10.11.** *Let  $\mathcal{H}$  be a measured groupoid over a base space  $Y$ , equipped with a cocycle  $\rho: \mathcal{H} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Assume that  $(\mathcal{H}, \rho)$  is FS-averse.*

*Then there exists a finite or countable Borel partition  $Y = \bigsqcup_{i \in I} Y_i$  such that for every  $i \in I$ , there exists a stable reduction collection  $\mathcal{C}_i$  for  $(\mathcal{H}|_{Y_i}, \rho)$ , and  $\mathcal{C}_i$  is finite and nonempty.*

*Proof.* We first claim that for every subset  $Y' \subseteq Y$  of positive measure, there are at most  $\max\{1, \frac{k-1}{2}\}$  almost free factor systems which are both  $(\mathcal{H}|_{Y'}, \rho)$ -invariant and  $(\mathcal{H}|_{Y'}, \rho)$ -everywhere maximal. Indeed, assume that  $\hat{\mathcal{F}}_1, \dots, \hat{\mathcal{F}}_r$  are  $(\mathcal{H}|_{Y'}, \rho)$ -invariant and  $(\mathcal{H}|_{Y'}, \rho)$ -everywhere maximal. Let  $H = \Gamma_{\{\hat{\mathcal{F}}_1, \dots, \hat{\mathcal{F}}_r\}}$  be the elementwise stabilizer of this collection. Up to replacing  $Y'$  by a Borel subset of full measure, we have  $\rho(\mathcal{H}|_{Y'}) \subseteq H$ . Since  $(\mathcal{H}, \rho)$  is FS-averse, the group  $H$  preserves no non-trivial free splitting. Similarly, since  $\hat{\mathcal{F}}_i$  is  $(\mathcal{H}|_{Y'}, \rho)$ -everywhere maximal, this implies that  $\hat{\mathcal{F}}_i$  is a maximal  $H$ -invariant almost free factor system. Corollary 8.30 then gives the bound  $r \leq \max\{1, \frac{k-1}{2}\}$  and proves the claim.

We will use several times the following fact: if  $\hat{\mathcal{F}}$  is an almost free factor system which is  $(\mathcal{H}|_{Y'}, \rho)$ -invariant, then one can find a finite or countable Borel partition  $Y' = \bigsqcup_i Y'_i$  and an almost free factor system  $\hat{\mathcal{F}}_i \succeq \hat{\mathcal{F}}$  which is  $(\mathcal{H}|_{Y'_i}, \rho)$ -invariant and  $(\mathcal{H}|_{Y'_i}, \rho)$ -everywhere maximal. Indeed, arguing as in Lemma 10.7, one first constructs a partition  $Y' = Y'' \sqcup Z$  such that  $\hat{\mathcal{F}}$  is everywhere maximal on  $Y''$ , and  $Z$  has a countable partition  $Z = \bigsqcup_j Z_j$  where there exists  $\hat{\mathcal{F}}_j \succeq \hat{\mathcal{F}}$  that is invariant on  $Z_j$ . Since there is a bound on the size of a chain of almost free factor systems of  $F_k$ , repeating this construction finitely many times yields the desired partition.

We first use this fact (with  $\hat{\mathcal{F}}$  the empty free factor system) to find a finite or countable Borel partition  $Y = \bigsqcup_i Z_i^1$  such that for every  $i$ , there exists an  $(\mathcal{H}|_{Z_i^1}, \rho)$ -invariant almost free factor system  $\hat{\mathcal{F}}_i$  which is  $(\mathcal{H}|_{Z_i^1}, \rho)$ -everywhere maximal.

Using this fact again, one may then find a countable Borel partition  $Z_i^1 = Y_i^1 \sqcup (\bigsqcup_j Z_{i,j}^{\geq 2})$ , such that

- $(\mathcal{H}|_{Y_i^1}, \rho)$  is pure: every almost free factor system  $\hat{\mathcal{F}}'$  which is  $(\mathcal{H}|_U, \rho)$ -invariant for some Borel subset of positive measure  $U \subseteq Y_i^1$  satisfies  $\hat{\mathcal{F}}' \preceq \hat{\mathcal{F}}_i$ ;
- for each  $j$ , there are at least two everywhere maximal invariant almost free factor systems on  $Z_{i,j}^{\geq 2}$ .

Repeating this process  $r$  times, one gets a partition of  $Y$  into pieces where the lemma holds (with a stable reduction collection of cardinality at most  $r-1$ ), and pieces on which

there are at least  $r$  distinct invariant almost free factor systems which are everywhere maximal. Taking  $r > \max\{1, \frac{k-1}{2}\}$  concludes the proof.  $\square$

**Lemma 10.12.** *Let  $\mathcal{H}$  be a measured groupoid over a base space  $Y$ , equipped with a cocycle  $\rho : \mathcal{H} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ .*

*If  $(\mathcal{H}, \rho)$  is nice-averse, then it is stably pure.*

*Proof.* Being nice-averse,  $(\mathcal{H}, \rho)$  is FS-averse so one may apply Lemma 10.11 to get a partition  $Y = \bigsqcup_{i \in I} Y_i$  with associated stable reduction collections  $\mathcal{C}_i$ .

It suffices to prove that for every  $i \in I$ , the pair  $(\mathcal{H}|_{Y_i}, \rho)$  is pure, i.e. that  $\mathcal{C}_i$  consists of a single almost free factor system. Assume on the contrary that  $\#\mathcal{C}_i \geq 2$ , and let  $\Gamma_{\mathcal{C}_i}$  be the elementwise stabilizer of  $\mathcal{C}_i$ . There exists a conull Borel subset  $Y_i^* \subseteq Y_i$  such that  $\rho(\mathcal{H}|_{Y_i^*}) \subseteq \Gamma_{\mathcal{C}_i}$ . As  $(\mathcal{H}, \rho)$  is FS-averse, it follows that  $\Gamma_{\mathcal{C}_i}$  is infinite and preserves no free splitting. Since  $\mathcal{C}_i$  is a stable reduction collection, each almost free factor system  $\hat{\mathcal{F}} \in \mathcal{C}_i$  is maximal among all  $\Gamma_{\mathcal{C}_i}$ -invariant almost free factor systems. Lemma 8.26 implies that  $\mathcal{C}_i$  is filling, i.e. there does not exist any almost free factor system  $\hat{\mathcal{F}}$  such that for every  $\hat{\mathcal{F}}' \in \mathcal{C}_i$ , one has  $\hat{\mathcal{F}}' \preceq \hat{\mathcal{F}}$ . Theorem 9.2(2) then provides a nice splitting  $U_{\mathcal{C}_i}$  which is invariant under  $\Gamma_{\mathcal{C}_i}$ , hence under  $(\mathcal{H}|_{Y_i^*}, \rho)$ . This contradicts that  $(\mathcal{H}, \rho)$  is nice-averse.  $\square$

*Proof of Proposition 10.5.* Let  $Y = Y_1 \sqcup Y_2$  be a Borel partition with  $(\mathcal{H}|_{Y_1}, \rho)$  stably nice and  $(\mathcal{H}|_{Y_2}, \rho)$  nice-averse, as given by Lemma 10.7. Then by Lemma 10.12,  $(\mathcal{H}|_{Y_2}, \rho)$  is stably pure, which concludes the proof.  $\square$

## 10.2 Invariant free splittings

The proof of Proposition 10.6 occupies the next subsections, where we distinguish cases according to the nature of the nice splittings preserved. Our first lemma uses the chain condition on stabilizers of collections of free splittings.

**Lemma 10.13.** *Let  $\mathcal{H}$  be a measured groupoid over a base space  $Y$ , equipped with a cocycle  $\rho : \mathcal{H} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ .*

*Then there exist a finite or countable Borel partition  $Y = \bigsqcup_{i \in I} Y_i$  and for every  $i \in I$ , a (maybe empty) collection  $\mathcal{C}_i$  of non-trivial free splittings, such that*

1. *every splitting in  $\mathcal{C}_i$  is  $(\mathcal{H}|_{Y_i}, \rho)$ -invariant;*
2. *for every Borel subset  $U \subseteq Y_i$  of positive measure, every non-trivial free splitting which is  $(\mathcal{H}|_U, \rho)$ -invariant belongs to  $\mathcal{C}_i$ .*

Notice that for each  $i \in I$  such that  $Y_i$  has positive measure, the collection  $\mathcal{C}_i$  satisfying the two properties is unique; furthermore,  $\mathcal{C}_i$  satisfies these two properties with respect to  $\mathcal{H}|_{Y'}$  for every Borel subset  $Y' \subseteq Y_i$  of positive measure. Also note that  $(\mathcal{H}|_{Y_i}, \rho)$  is FS-averse if and only if  $\mathcal{C}_i = \emptyset$ .

*Proof.* Let  $\mathcal{C}^0$  be the (maybe empty) set of all  $(\mathcal{H}, \rho)$ -invariant non-trivial free splittings. As in Lemma 10.7, we can find a countable Borel partition  $Y = \bigsqcup_{i \in \mathbb{N}} Y_i$  such that

1. for every Borel subset  $U \subseteq Y_0$  of positive measure, every  $(\mathcal{H}|_U, \rho)$ -invariant free splitting belongs to  $\mathcal{C}^0$ ;
2. for every  $i \geq 1$ , there exists a non-trivial free splitting  $S_i \notin \mathcal{C}^0$  which is  $(\mathcal{H}|_{Y_i}, \rho)$ -invariant.

Now, for every  $i \geq 1$ , let  $\mathcal{C}_i^1$  be the set of all  $(\mathcal{H}|_{Y_i}, \rho)$ -invariant non-trivial free splittings. Notice that  $\mathcal{C}^0 \subsetneq \mathcal{C}_i^1$ . In addition, their elementwise stabilizers in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  satisfy  $\Gamma_{\mathcal{C}_i^1} \subsetneq \Gamma_{\mathcal{C}^0}$ : indeed, this inclusion is strict because  $\rho(\mathcal{H}|_{Y^*}) \subseteq \Gamma_{\mathcal{C}^0}$  for some conull Borel subset  $Y^* \subseteq Y$  whereas  $\rho(\mathcal{H}|_{Y^*}) \not\subseteq \Gamma_{\mathcal{C}_i^1}$  since  $S_i$  is not  $(\mathcal{H}, \rho)$ -invariant.

One can now repeat this construction: for every  $i \geq 1$ , we can find a countable Borel partition  $Y_i = \bigsqcup_{j \in \mathbb{N}} Y_{i,j}$  such that

1. for every Borel subset  $U \subseteq Y_{i,0}$  of positive measure, every  $(\mathcal{H}|_U, \rho)$ -invariant non-trivial free splitting belongs to  $\mathcal{C}_i^1$ ;
2. for every  $j \geq 1$ , there exists a non-trivial free splitting  $S_{i,j} \notin \mathcal{C}_i^1$  which is  $(\mathcal{H}|_{Y_{i,j}}, \rho)$ -invariant.

For every  $i, j \geq 1$ , let  $\mathcal{C}_{i,j}^2$  be the set of all  $(\mathcal{H}|_{Y_{i,j}}, \rho)$ -invariant non-trivial free splittings. As above, we have  $\Gamma_{\mathcal{C}_{i,j}^2} \subsetneq \Gamma_{\mathcal{C}_i^1}$ .

As there is a bound on the size of a chain of elementwise stabilizers of collections of free splittings (Proposition 6.17), the iteration of this process stops after finitely many steps. The output is a countable Borel partition of  $Y$  as in the lemma.  $\square$

In the following lemma, we let  $\mathcal{G}, \mathcal{H}, \mathcal{H}'$  be as in the statement of Proposition 10.6, and we apply Lemma 10.13 to  $\mathcal{H}$  to get subsets  $Y_i$  and collections  $\mathcal{C}_i$  of non-trivial free splittings.

**Lemma 10.14.** *Let  $Y_i, \mathcal{C}_i$  be as in Lemma 10.13. Let  $\mathcal{H}'$  be a measured subgroupoid of  $\mathcal{G}$  which normalizes  $\mathcal{H}$ .*

*Then for every  $i \in I$ , the collection  $\mathcal{C}_i$  is setwise  $(\mathcal{H}'|_{Y_i}, \rho)$ -invariant.*

Denoting by  $\Gamma_{(\mathcal{C}_i)}$  the setwise stabilizer in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  of the collection  $\mathcal{C}_i$ , this means that there exists a conull Borel subset  $Y_i^* \subseteq Y_i$  such that  $\rho(\mathcal{H}'|_{Y_i^*}) \subseteq \Gamma_{(\mathcal{C}_i)}$ .

*Proof.* By definition of normalization of subgroupoids, one can write  $\mathcal{H}'|_{Y_i}$  as a countable union of bisections  $B_n \subseteq \mathcal{H}'|_{Y_i}$  such that for all  $h \in \mathcal{G}|_{Y_i}$  and all  $\varphi, \psi \in B_n$  such that  $\varphi h \psi^{-1}$  is defined, one has  $h \in \mathcal{H}$  if and only if  $\varphi h \psi^{-1} \in \mathcal{H}$ . Up to subdividing the bisections, we may assume additionally that the cocycle  $\rho$  maps  $B_n$  to a single value  $\gamma_n \in \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Let  $f_{B_n} : U_n \rightarrow V_n$  be the partial isomorphism associated to the bisection  $B_n$ , with  $U_n, V_n \subseteq Y_i$ .

Any free splitting  $S \in \mathcal{C}_i$  is  $(\mathcal{H}|_{U_n}, \rho)$ -invariant, so  $\gamma_n S$  is  $(\mathcal{H}|_{V_n}, \rho)$ -invariant. By maximality of  $\mathcal{C}_i$  (Lemma 10.13(2)), it follows that  $\gamma_n S \in \mathcal{C}_i$  if  $V_n$  has positive measure. Taking for  $Y_i^*$  a subset of full measure avoiding all  $U_n$  and  $V_n$  of zero measure, we deduce that  $\rho(\mathcal{H}'|_{Y_i^*}) \subseteq \Gamma_{(\mathcal{C}_i)}$ , which concludes our proof.  $\square$

Since by Theorem 9.2, one can assign to  $\mathcal{C}_i$  a canonical nice splitting as soon as  $\mathcal{C}_i \neq \emptyset$  and  $\Gamma_{\mathcal{C}_i}$  is infinite, we get the following corollary for nowhere trivial cocycles.

**Corollary 10.15.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$  with a cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$  such that  $\rho|_{\mathcal{H}}$  is nowhere trivial. Then there exists a finite or countable Borel partition  $Y = Z \sqcup (\bigsqcup_{i \in I} Y_i)$  such that*

1.  $(\mathcal{H}|_Z, \rho)$  is FS-averse,
2. for every  $i \in I$ , there exists a nice splitting  $S_i$  which is  $(\mathcal{H}|_{Y_i}, \rho)$ -invariant, and in fact  $(\mathcal{H}'|_{Y_i}, \rho)$ -invariant for every measured subgroupoid  $\mathcal{H}'$  of  $\mathcal{G}$  such that  $\mathcal{H}'|_{Y_i}$  normalizes  $\mathcal{H}|_{Y_i}$ .

*Proof.* Consider a finite or countable Borel partition  $Y = \bigsqcup_{j \in J} Y_j$  with a collection  $\mathcal{C}_j$  of free splittings for each  $j \in J$ , as in Lemma 10.13. Denoting by  $Z$  the union of the sets  $Y_j$  such that  $\mathcal{C}_j = \emptyset$ , we see that  $(\mathcal{H}|_Z, \rho)$  is FS-averse.

Now consider  $j \in J$  such that  $\mathcal{C}_j \neq \emptyset$ . By Lemma 10.14,  $\mathcal{C}_j$  is invariant under  $(\mathcal{H}|_{Y_j}, \rho)$  and  $(\mathcal{H}'|_{Y_j}, \rho)$ . Since  $\rho|_{\mathcal{H}}$  is nowhere trivial, the elementwise stabilizer  $\Gamma_{\mathcal{C}_j}$  of  $\mathcal{C}_j$  is non-trivial, hence infinite. The nice splitting  $S_j = U_{\mathcal{C}_j}$  provided by Theorem 9.2(1) is then invariant under  $(\mathcal{H}|_{Y_j}, \rho)$  and  $(\mathcal{H}'|_{Y_j}, \rho)$ .  $\square$

### 10.3 Invariant $\mathcal{Z}_{\max}$ -splittings

**Lemma 10.16.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , equipped with a cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ , and let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ . Assume that  $(\mathcal{H}, \rho)$  is FS-averse.*

*Then there exists a finite or countable Borel partition  $Y = Z \sqcup (\bigsqcup_{i \in I} Y_i)$  such that*

1. for every Borel subset  $U \subseteq Z$  of positive measure, there is no  $(\mathcal{H}|_U, \rho)$ -invariant non-trivial  $\mathcal{Z}_{\max}$ -splitting of  $F_k$ ,
2. for every  $i \in I$ , there is a non-trivial  $\mathcal{Z}_{\max}$ -splitting  $S_i$  which is  $(\mathcal{H}|_{Y_i}, \rho)$ -invariant, and in fact  $(\mathcal{H}'|_{Y_i}, \rho)$ -invariant for every measured subgroupoid  $\mathcal{H}'$  of  $\mathcal{G}$  such that  $\mathcal{H}'|_{Y_i}$  normalizes  $\mathcal{H}|_{Y_i}$ .

*Proof.* Using the chain condition on stabilizers of  $\mathcal{Z}_{\max}$ -splittings (Proposition 7.24), one constructs as in the proof of Lemma 10.13 a finite or countable Borel partition  $Y = \bigsqcup_{j \in J} Y_j$ , and for every  $j \in J$ , a (maybe empty) set  $\mathcal{C}_j$  of non-trivial  $\mathcal{Z}_{\max}$ -splittings, such that

1. every splitting in  $\mathcal{C}_j$  is  $(\mathcal{H}|_{Y_j}, \rho)$ -invariant;
2. for every Borel subset  $U \subseteq Y_j$  of positive measure, every non-trivial  $\mathcal{Z}_{\max}$ -splitting which is  $(\mathcal{H}|_U, \rho)$ -invariant belongs to  $\mathcal{C}_j$ .



We define  $Z$  as the union of all  $Y_j$  such that  $\mathcal{C}_j$  is empty. It clearly satisfies the first assertion of the lemma.

Now fix  $j$  such that  $\mathcal{C}_j \neq \emptyset$ . As  $(\mathcal{H}, \rho)$  is FS-averse, the elementwise stabilizer  $\Gamma_{\mathcal{C}_j}$  of  $\mathcal{C}_j$  does not fix any non-trivial free splitting of  $F_k$ . As in Lemma 10.14, for every Borel subset  $U \subseteq Y_j$  of positive measure  $(\mathcal{H}', \rho)$  stabilizes  $\mathcal{C}_j$  setwise. The proposition thus follows by using Theorem 7.4, which canonically associates a non-trivial  $\mathcal{Z}_{\max}$ -splitting to the group  $\Gamma_{\mathcal{C}_j}$ .  $\square$

## 10.4 Invariant bi-nonsporadic splittings and conclusion

**Lemma 10.17.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , with a cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ . If  $(\mathcal{H}, \rho)$  is FS-averse and stabilizes a bi-nonsporadic splitting  $T$ , then for every measured subgroupoid  $\mathcal{H}'$  of  $\mathcal{G}$  that normalizes  $\mathcal{H}$ ,  $(\mathcal{H}', \rho)$  is stably nice.*

*More precisely, there is a finite or countable Borel partition  $Y = \bigsqcup_{i \in I} Y_i$  and for each  $i \in I$  a nice splitting  $S_i$  of  $F_k$  such that  $S_i$  is invariant under both  $(\mathcal{H}|_{Y_i}, \rho)$  and  $(\mathcal{H}'|_{Y_i}, \rho)$ .*

*Proof.* By Lemma 9.6, one can canonically associate to  $T$  a filling finite collection  $\mathcal{C}$  of almost free factor systems. In particular,  $\mathcal{C}$  is setwise  $(\mathcal{H}, \rho)$ -invariant, hence elementwise  $(\mathcal{H}, \rho)$ -invariant by Theorems 2.10 and 2.11. Apply Lemma 10.11 to get a partition  $Y = \bigsqcup_{i \in I} Y_i$  such that for every  $i \in I$ , there is a nonempty stable reduction collection  $\mathcal{C}_i$  of almost free factor systems. By Lemma 10.10, the set  $\mathcal{C}_i$  is also  $(\mathcal{H}'|_{Y_i}, \rho)$ -invariant.

Fix  $i \in I$ . The collection  $\mathcal{C}_i$  has the property that for every  $\hat{\mathcal{F}} \in \mathcal{C}_i$ , there exists  $\hat{\mathcal{F}}' \in \mathcal{C}_i$  such that  $\hat{\mathcal{F}} \preceq \hat{\mathcal{F}}'$ . Since  $\mathcal{C}$  is filling, so is  $\mathcal{C}_i$ . Since  $(\mathcal{H}, \rho)$  is FS-averse,  $\rho|_{\mathcal{H}}$  is nowhere trivial, so the setwise stabilizer of  $\mathcal{C}_i$  in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  is non-trivial hence infinite. It follows that the elementwise stabilizer  $\Gamma_{\mathcal{C}_i}$  of  $\mathcal{C}_i$  is infinite. Let  $S_i = U_{\mathcal{C}_i}$  be the nice splitting associated to  $\mathcal{C}_i$  by Theorem 9.2(2). Then  $S_i$  is invariant under both  $(\mathcal{H}|_{Y_i}, \rho)$  and  $(\mathcal{H}'|_{Y_i}, \rho)$ .  $\square$

*Proof of Proposition 10.6.* By Corollary 10.15, up to partitioning the base space  $Y$ , we may assume that  $(\mathcal{H}, \rho)$  is FS-averse. Since  $(\mathcal{H}, \rho)$  is stably nice, we may assume up to further partitioning that  $(\mathcal{H}, \rho)$  preserves a  $\mathcal{Z}_{\max}$  or a binonsporadic splitting  $S$ . The proposition then follows from Lemmas 10.16 and 10.17 accordingly.  $\square$

## 11 Groupoids with cocycles to direct products of free groups

In later sections, given a measured groupoid  $\mathcal{G}$  equipped with a cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , we will often want to consider subgroupoids  $\mathcal{H}$  of  $\mathcal{G}$  such that  $(\mathcal{H}, \rho)$  has an invariant one-edge non-separating free splitting  $S$ , and an interesting subgroupoid of  $\mathcal{H}$  to consider is the preimage under  $\rho$  of the group of twists of the splitting  $S$ , which is a finite-index subgroup of a direct product of free groups (see Section 2.8). This motivates the present section, which establishes a few facts about groupoids equipped with cocycles taking values in a direct product of free groups. We start by stating a first lemma which

is essentially due to Adams [Ada94], and actually follows from the next lemma which gives a more general statement.

**Lemma 11.1.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , let  $F$  be a finitely generated free group, and let  $\rho : \mathcal{G} \rightarrow F$  be a cocycle. Let  $\mathcal{A}, \mathcal{H}$  be two measured subgroupoids of  $\mathcal{G}$ . Assume that  $\rho|_{\mathcal{A}}$  is nowhere trivial and that  $\rho|_{\mathcal{H}}$  has trivial kernel (Definitions 3.18 and 3.14).*

*If  $\mathcal{A}$  is amenable and normalized by  $\mathcal{H}$ , then  $\mathcal{H}$  is amenable.*

Lemma 11.1 follows from the next statement, after replacing  $\rho : \mathcal{G} \rightarrow F$  by  $\tilde{\rho} : \mathcal{G} \rightarrow F \times F$  defined by  $\tilde{\rho}(g) = (\rho(g), \rho(g))$ , and setting  $\mathcal{A}_l = \mathcal{A}_r = \mathcal{A}$ .

**Lemma 11.2.** *Let  $\mathcal{G}$  be a measured groupoid, let  $F_l, F_r$  be two finitely generated free groups, and let  $\rho : \mathcal{G} \rightarrow F_l \times F_r$  be a cocycle. Let  $\rho_l : \mathcal{G} \rightarrow F_l$  and  $\rho_r : \mathcal{G} \rightarrow F_r$  be the two cocycles obtained by postcomposing  $\rho$  with the two projections. Let  $\mathcal{A}_l, \mathcal{A}_r, \mathcal{H}$  be measured subgroupoids of  $\mathcal{G}$ . Assume that  $(\rho_l)|_{\mathcal{A}_l}$  and  $(\rho_r)|_{\mathcal{A}_r}$  are nowhere trivial and that  $\rho|_{\mathcal{H}}$  has trivial kernel.*

*If  $\mathcal{A}_l$  and  $\mathcal{A}_r$  are both amenable and normalized by  $\mathcal{H}$ , then  $\mathcal{H}$  is amenable.*

*Proof.* The proof is based on an argument due to Adams [Ada94]. Let  $T_l$  be a Cayley tree for  $F_l$  with respect to some free basis. Then  $F_l$  acts by isometries on  $T_l$ , and this action extends continuously to an  $F_l$ -action by homeomorphisms on  $\partial_\infty T_l$ . As  $\mathcal{A}_l$  is amenable and  $\partial_\infty T_l$  is a compact metrizable space, Proposition 3.41 ensures that there exists an  $(\mathcal{A}_l, \rho_l)$ -equivariant Borel map  $\mu : Y \rightarrow \text{Prob}(\partial_\infty T_l)$ .

By [Ada94, Proposition 5.1], for a.e.  $y \in Y$ , the probability measure  $\mu(y)$  is supported on at most two points. Let us briefly recall the argument for this fact, in order to prepare for the next section where we will need a refined version. Assume towards a contradiction that there exists a Borel subset  $U \subseteq Y$  of positive measure such that for every  $y \in U$ , the probability measure  $\mu(y)$  is supported on at least three points. Then  $\mu(y) \otimes \mu(y) \otimes \mu(y)$  gives positive measure to the subspace  $(\partial_\infty T_l)^{(3)}$  made of pairwise distinct triples. By restriction and renormalization of the probability measures, we thus get an  $((\mathcal{A}_l)|_U, \rho_l)$ -equivariant Borel map  $U \rightarrow \text{Prob}((\partial_\infty T_l)^{(3)})$ . Notice that there exists an  $F_l$ -equivariant Borel *barycenter* map  $(\partial_\infty T_l)^{(3)} \rightarrow V(T_l)$ , where  $V(T_l)$  denotes the (countable) vertex set of  $T_l$ . Pushing forward the probability measures obtained above by this barycenter map, this yields an  $((\mathcal{A}_l)|_U, \rho_l)$ -equivariant Borel map  $\varphi : U \rightarrow \text{Prob}(V(T_l))$ . Denote by  $\mathcal{P}_{<\infty}(V(T_l))$  the (countable) collection of all nonempty finite subsets of  $V(T_l)$ . As  $V(T_l)$  is countable, there is also an  $F_l$ -equivariant Borel map  $\text{Prob}(V(T_l)) \rightarrow \mathcal{P}_{<\infty}(V(T_l))$ , sending a probability measure  $\nu$  to the finite set of all elements of  $V(T_l)$  with maximal  $\nu$ -measure. Combining the above two maps, we derive an  $((\mathcal{A}_l)|_U, \rho_l)$ -equivariant Borel map  $U \rightarrow \mathcal{P}_{<\infty}(V(T_l))$ . By restricting to a Borel subset  $U' \subseteq U$  of positive measure where this map is constant, we find a positive measure Borel subset of  $Y$  such that  $\rho_l((\mathcal{A}_l)|_{U'})$  is contained in the  $F_l$ -stabilizer of a finite set of vertices of  $T_l$ , whence trivial. This yields a contradiction to the fact that  $(\rho_l)|_{\mathcal{A}_l}$  is nowhere trivial.

Therefore, by post-composing  $\mu$  with the map sending every probability measure to its support, we get an  $(\mathcal{A}_l, \rho_l)$ -equivariant Borel map  $Y \rightarrow \mathcal{P}_{\leq 2}(\partial_\infty T_l)$ , where  $\mathcal{P}_{\leq 2}(\partial_\infty T_l)$

is the set of all nonempty subsets of  $\partial_\infty T_l$  of cardinality at most 2. In fact, the barycenter argument also shows that if  $\varphi_1, \varphi_2$  are two such maps, then their union  $\varphi_1 \cup \varphi_2$  must again take its values in  $\mathcal{P}_{\leq 2}(\partial_\infty T_l)$ . Therefore, by [Ada94, Lemmas 3.2 and 3.3], there exists a unique  $(\mathcal{A}_l, \rho_l)$ -equivariant Borel map  $\varphi_{\max} : Y \rightarrow \mathcal{P}_{\leq 2}(\partial_\infty T_l)$  which is everywhere maximum in the following sense: for every Borel subset  $U \subseteq Y$  of positive measure and every  $((\mathcal{A}_l)|_U, \rho_l)$ -equivariant Borel map  $\varphi : U \rightarrow \mathcal{P}_{\leq 2}(\partial_\infty T_l)$ , one has  $\varphi(y) \subseteq \varphi_{\max}(y)$  for almost every  $y \in U$ .

Using the fact that  $\mathcal{H}$  normalizes  $\mathcal{A}_l$ , we will now derive that  $\varphi_{\max}$  is also  $(\mathcal{H}, \rho_l)$ -equivariant. Indeed, one can write  $\mathcal{H}$  as a countable union of bisections  $B_n \subseteq \mathcal{H}$  such that for all  $a \in \mathcal{A}_l$  and all  $\gamma, \delta \in B_n$  such that  $\gamma a \delta^{-1}$  is defined, one has  $a \in \mathcal{A}_l$  if and only if  $\gamma a \delta^{-1} \in \mathcal{A}_l$ . Up to subdividing these bisections, we may assume that the cocycle  $\rho_l$  takes a single value  $\gamma_n$  on  $B_n$ . Let  $f_{B_n} : U_n \rightarrow V_n$  be the partial isomorphism associated to the bisection  $B_n$ , with  $U_n, V_n \subseteq Y$ . Whenever  $U_n$  has positive measure, the map

$$\begin{aligned} \varphi' : U_n &\rightarrow \mathcal{P}_{\leq 2}(\partial_\infty T_l) \\ y &\mapsto \gamma_n^{-1} \cdot \varphi_{\max}(f_{B_n}(y)) \end{aligned}$$

is an everywhere maximum  $((\mathcal{A}_l)|_{U_n}, \rho_l)$ -equivariant Borel map so there is a subset of full measure  $U'_n \subseteq U_n$  on which  $\varphi'$  and  $\varphi_{\max}$  coincide – this is also obviously true if  $U_n$  has measure zero. Since  $\mathcal{H}$  is the union of the bisections  $B_n$ , this shows that  $\varphi_{\max}$  is  $(\mathcal{H}, \rho_l)$ -equivariant.

Similarly, by working with  $\mathcal{A}_r$  instead of  $\mathcal{A}_l$ , we build an  $(\mathcal{H}, \rho_r)$ -equivariant Borel map  $Y \rightarrow \mathcal{P}_{\leq 2}(\partial_\infty T_r)$ . Combining these two maps yields an  $(\mathcal{H}, \rho)$ -equivariant Borel map  $Y \rightarrow \mathcal{P}_{\leq 2}(\partial_\infty T_l) \times \mathcal{P}_{\leq 2}(\partial_\infty T_r)$ . The action of  $F_l \times F_r$  on  $\mathcal{P}_{\leq 2}(\partial_\infty T_l) \times \mathcal{P}_{\leq 2}(\partial_\infty T_r)$  is Borel amenable (see e.g. [AD07, Example 1.4(3)] for the Borel amenability of the action of a finitely generated free group on the boundary of its Cayley tree, from which our claim easily follows). As  $\rho|_{\mathcal{H}}$  has trivial kernel, Proposition 3.38 therefore ensures that  $\mathcal{H}$  is amenable.  $\square$

As a consequence of Lemma 11.1, we deduce the following statement.

**Lemma 11.3.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , let  $F_l$  and  $F_r$  be finitely generated non-abelian free groups, and let  $\rho : \mathcal{G} \rightarrow F_l \times F_r$  be an action-type cocycle.*

*Then  $\mathcal{G}$  does not contain any normal amenable subgroupoid of infinite type.*

*Proof.* Assume towards a contradiction that  $\mathcal{G}$  contains a normal amenable subgroupoid  $\mathcal{A}$  of infinite type. Let  $\rho_l, \rho_r$  be the two cocycles obtained by postcomposing  $\rho$  with the projection to  $F_l$  and  $F_r$ , respectively.

We claim that we can then find a Borel partition  $Y = Y_l \sqcup Y_r$  such that  $(\rho_l)|_{\mathcal{A}}$  is nowhere trivial on  $Y_l$  and  $(\rho_r)|_{\mathcal{A}}$  is nowhere trivial on  $Y_r$ . Indeed, let  $Y_r$  be such that

1.  $(\rho_l)|_{\mathcal{A}}$  is stably trivial on  $Y_r$ , i.e. there exists a partition  $Y_r = \bigsqcup_{i \in I} Y_{r,i}$  into at most countably many Borel subsets such that for every  $i \in I$ , the cocycle  $(\rho_l)|_{\mathcal{A}}$  is trivial on  $Y_{r,i}$ , and
2.  $(\rho_l)|_{\mathcal{A}}$  is nowhere trivial on  $Y_l = Y \setminus Y_r$ .

We check that  $(\rho_r)_{|\mathcal{A}}$  is nowhere trivial on  $Y_r$ . Given  $U \subseteq Y_r$  of positive measure, consider  $i \in I$  such that  $U' := U \cap Y_{r,i}$  has positive measure. In particular  $\rho_l(\mathcal{A}_{|U'}) = \{1\}$ . Since  $\rho$  has trivial kernel,  $\rho_r$  has trivial kernel on  $\mathcal{A}_{|U'}$ . Since  $\mathcal{A}$  is of infinite type, this implies that  $\rho_r(\mathcal{A}_{|U'}) \neq \{1\}$  and proves our claim.

Without loss of generality, assume that  $Y_l$  has positive measure. Let  $\mathcal{H}_l = \rho^{-1}(F_l \times \{1\})$ . Since  $F_l$  is a non-abelian free group,  $\mathcal{H}_l$  is everywhere non-amenable by Lemma 3.40. As  $\rho$  has trivial kernel,  $\rho_l$  has trivial kernel on  $(\mathcal{H}_l)_{|Y_l}$ . Since  $\mathcal{A}_{|Y_l}$  is normalized by  $(\mathcal{H}_l)_{|Y_l}$ , Lemma 11.1 implies that  $(\mathcal{H}_l)_{|Y_l}$  is amenable, a contradiction.  $\square$

## 12 Nice-averse amenable subgroupoids: Adams' argument

In this section, we fix an integer  $k \geq 2$ . The main result of the present section is the following.

**Theorem 12.1.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and  $\mathcal{A}, \mathcal{H}$  two measured subgroupoids of  $\mathcal{G}$ . Let  $\rho : \mathcal{G} \rightarrow \mathrm{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle such that  $\rho_{|\mathcal{H}}$  has trivial kernel and  $\rho_{|\mathcal{A}}$  is nowhere trivial. Assume that  $\mathcal{A}$  is amenable, and that  $\mathcal{A}$  is normalized by  $\mathcal{H}$ .*

*Then either  $\mathcal{H}$  is amenable, or there exist a Borel subset  $U \subseteq Y$  of positive measure and a nice splitting which is  $(\mathcal{H}_{|U}, \rho)$ -invariant and  $(\mathcal{A}_{|U}, \rho)$ -invariant.*

**Corollary 12.2.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and  $\mathcal{A}, \mathcal{H}$  two measured subgroupoids of  $\mathcal{G}$ . Let  $\rho : \mathcal{G} \rightarrow \mathrm{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle with trivial kernel. Assume that  $\mathcal{A}$  is of infinite type and amenable, and that  $\mathcal{A}$  is stably normalized by  $\mathcal{H}$ .*

*If  $(\mathcal{H}, \rho)$ ,  $(\mathcal{A}, \rho)$  or  $(\langle \mathcal{A}, \mathcal{H} \rangle, \rho)$  is nice-averse, then  $\mathcal{H}$  is amenable.*

*Proof.* Since  $\mathcal{H}$  stably normalizes  $\mathcal{A}$ , there is a partition  $Y^* = \bigsqcup_{i \in I} Y_i$  of a conull Borel subset  $Y^* \subseteq Y$  into at most countably many Borel subsets such that  $\mathcal{H}_{|Y_i}$  normalizes  $\mathcal{A}_{|Y_i}$  for every  $i \in I$ . Since amenability of every restriction  $\mathcal{H}_{|Y_i}$  implies amenability of  $\mathcal{H}$ , we may restrict to  $Y_i$  and assume that  $\mathcal{H}$  normalizes  $\mathcal{A}$ .

Let  $\mathcal{H}' = \langle \mathcal{A}, \mathcal{H} \rangle$ , and note that  $\mathcal{H}'$  normalizes  $\mathcal{A}$ . In all cases,  $(\mathcal{H}', \rho)$  is nice-averse because if  $(\mathcal{H}, \rho)$  or  $(\mathcal{A}, \rho)$  is nice-averse, then so is  $(\mathcal{H}', \rho)$ . As  $\rho$  has trivial kernel,  $\rho_{|\mathcal{H}'}$  has trivial kernel, and as  $\mathcal{A}$  is of infinite type, we also deduce that  $\rho_{|\mathcal{A}}$  is nowhere trivial. We can therefore apply Theorem 12.1 to the subgroupoids  $\mathcal{A}$  and  $\mathcal{H}'$  and as  $(\mathcal{H}', \rho)$  is nice-averse, its conclusion ensures that  $\mathcal{H}'$  is amenable, hence so is  $\mathcal{H}$ .  $\square$

The general strategy of our proof follows an argument due to Adams [Ada94], and implemented in the mapping class group setting by Kida [Kid08a, Chapter 4]; the basic idea of this argument was in fact reviewed in our proof of Lemma 11.2. But we will also need to establish some new results about the amenability of certain actions of  $\mathrm{Out}(F_k, \mathcal{F})$ , and there are some real differences that arise between mapping class groups and  $\mathrm{Out}(F_k)$ . For instance, the analogue of Kida's statement saying that the normalizer of an irreducible amenable groupoid is amenable does not always hold here, an extra possibility being that the normalizer preserves a nice splitting.

Our proof of Theorem 12.1 will be completed in Section 12.8. It goes as follows. After a countable partition of the base space  $Y$ , we may assume that  $(\mathcal{A}, \rho)$  is pure or that  $(\mathcal{A}, \rho)$

preserves a nice splitting (Proposition 10.5), and in the latter case, Proposition 10.6 says that up to further partitioning, there is a canonical one which is both  $(\mathcal{A}, \rho)$ -invariant and  $(\mathcal{H}, \rho)$ -invariant, so we are done. We thus assume that  $(\mathcal{A}, \rho)$  is pure and consider the unique maximal  $(\mathcal{A}, \rho)$ -invariant almost free factor system  $\hat{\mathcal{F}}$ . It is also  $(\mathcal{H}, \rho)$ -invariant, and we work relative to  $\hat{\mathcal{F}}$ , and in fact relative to an extracted free factor system  $\mathcal{F} \subseteq \hat{\mathcal{F}}$  (see Definition 8.6).

At first our proof follows Kida's implementation of Adams' argument closely. We start by using the amenability of  $\mathcal{A}$  to construct an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $\mu$  from  $Y$  to the space of all probability measures on the compactification  $\mathbb{P}\bar{\mathcal{O}}$  of the relative Outer space (Section 12.1). Using a witness map assigning to any non-arational tree a finite collection of conjugacy classes of free factors, maximality of  $\mathcal{F}$  implies that for almost every  $y \in Y$ , the probability measure  $\mu(y)$  gives full measure to the set of arational trees (Section 12.2). Using a barycenter map on triples of inequivalent arational trees, one shows that the support of  $\mu(y)$  has cardinality at most two almost everywhere (Section 12.3). This argument also implies that there is an essentially unique maximal  $(\mathcal{A}, \rho)$ -equivariant Borel map from  $Y$  to the set of pairs of points in  $\mathcal{AT}/\sim$ , which is therefore  $(\mathcal{H}, \rho)$ -equivariant (Section 12.4).

One would then like to conclude that  $\mathcal{H}$  is amenable using Borel amenability of the action on  $\mathcal{AT}$ , but this property does not hold in a relative setting as some arational trees have a non-amenable stabilizer in  $\text{Out}(F_k, \mathcal{F})$ <sup>4</sup>. This is where we need new additional arguments that were not required in the mapping class group setting; we deal with this difficulty in Sections 12.5 to 12.7, which occupy most of our proof of Theorem 12.1.

What we prove is the Borel amenability of the action of  $\text{Out}(F_k, \mathcal{F})$  on the set  $\mathcal{AT}^a$  of arational trees with amenable stabilizer (and in fact, also, on the set  $\mathcal{P}_{\leq 2}^a(\mathcal{AT})$  of singletons or pairs of arational trees whose common stabilizer is amenable). We then prove that there is another equivariant witness map defined on the set of arational trees  $T$  with non-amenable stabilizer and assigning to  $T$  a nice splitting of  $F_k$  (with a variant for pairs of arational trees). As above this ensures that (almost everywhere) the support of the map  $\mu(y)$  belongs to  $\mathcal{P}_{\leq 2}^a(\mathcal{AT})$ , and we can then use the Borel amenability of the action of  $\text{Out}(F_k, \mathcal{F})$  on  $\mathcal{P}_{\leq 2}^a(\mathcal{AT})$  to conclude that  $\mathcal{H}$  is amenable. The end of the argument is summarized in Section 12.8.

**Standing assumptions:** From now on, thanks to Proposition 10.6, we will assume that  $(\mathcal{A}, \rho)$  is pure in addition to all assumptions coming from the statement of Theorem 12.1 including amenability of  $\mathcal{A}$ . We denote by  $\hat{\mathcal{F}}$  the maximal  $(\mathcal{A}, \rho)$ -invariant almost free factor system (coming from Definition 10.2 of purity). We choose a free factor system  $\mathcal{F} \subseteq \hat{\mathcal{F}}$  extracted from  $\hat{\mathcal{F}}$ . Then  $\mathcal{F}$  is  $(\mathcal{A}, \rho)$ -invariant (as  $\rho$  takes its values in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ ).

In all this section, we will work relative to  $\mathcal{F}$ , and leave  $\mathcal{F}$  implicit in the notations: we denote by  $\mathbb{P}\bar{\mathcal{O}} = \mathbb{P}\bar{\mathcal{O}}(F_k, \mathcal{F})$  the compactification of the projectivized Outer space

<sup>4</sup>There is an important subtlety here: earlier work of Bestvina and the authors, that we will use in a crucial way, shows that the action of  $\text{Out}(F_k, \mathcal{F}^{(t)})$  on  $\mathcal{AT}$  is Borel amenable [BGH17, Theorem 6.4]; but the action of the bigger group  $\text{Out}(F_k, \mathcal{F})$  is not in general; see the discussion at the beginning of Section 12.5.

of  $(F_k, \mathcal{F})$ , by  $\mathbb{P}\mathcal{AT} \subseteq \mathbb{P}\overline{\mathcal{O}}$  the set of projective arational trees, and by FF the set of conjugacy classes of proper  $(F_k, \mathcal{F})$ -free factors.

We record the following fact for future use.

**Lemma 12.3.** *The free factor system  $\mathcal{F}$  is nonsporadic, and everywhere maximal in the following sense: if  $\mathcal{F}' \succeq \mathcal{F}$  is a free factor system which is  $(\mathcal{A}|_U, \rho)$ -invariant for some Borel subset  $U \subseteq Y$  of positive measure, then  $\mathcal{F}' = \mathcal{F}$ .*

*Proof.* Non-sporadicity of  $\mathcal{F}$  follows from the fact that  $(\mathcal{A}, \rho)$  is FS-averse (this is an assumption in Definition 10.2 of purity).

For the maximality statement, the almost free factor system  $\hat{\mathcal{F}}$  is everywhere maximal by definition, so  $\mathcal{F} \preceq \mathcal{F}' \preceq \hat{\mathcal{F}}$ . Since  $\mathcal{F}'$  is a free factor system,  $\mathcal{F} = \mathcal{F}'$ .  $\square$

*Remark 12.4.* Amenability of  $\mathcal{A}$  will only be used in Section 12.1 to build an equivariant map  $Y \rightarrow \text{Prob}(\mathbb{P}\overline{\mathcal{O}})$ . In later sections, we will only use the existence of this map, and not amenability itself.

## 12.1 Equivariant probability measures on the compactified Outer space

**Lemma 12.5.** *There exists an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $\mu : Y \rightarrow \text{Prob}(\mathbb{P}\overline{\mathcal{O}})$ .*

*Proof.* This follows from the fact that  $\mathcal{A}$  is amenable and  $\mathbb{P}\overline{\mathcal{O}}$  is compact and metrizable, by applying Proposition 3.41.  $\square$

In the sequel, we will often denote by  $\mu_y$  the probability measure  $\mu(y)$ .

## 12.2 Arational trees and Reynolds' witness map

Recall that an  $\mathbb{R}$ -tree  $T$  in the boundary of Outer space is *not* arational if some proper  $(F_k, \mathcal{F})$ -free factor acts non-discretely or fixes a point. Theorem 12.6 below, established in this form in [Hor14, Lemma 5.5] by generalizing an argument of Reynolds [Rey12] to a relative setting, asserts the existence of a *witness map*, which equivariantly assigns to any non-arational tree  $T$  a finite collection of conjugacy classes of free factors, witnessing the fact that  $T$  is not arational.

We denote by  $\mathcal{AT} = \mathcal{AT}(F_k, \mathcal{F})$  the space of all arational  $(F_k, \mathcal{F})$ -trees, and by  $\mathbb{P}\mathcal{AT}$  the space of all projective classes of arational  $(F_k, \mathcal{F})$ -trees. We recall that FF denotes the countable set of all conjugacy classes of proper  $(F_k, \mathcal{F})$ -free factors (i.e. free factors relative to  $\mathcal{F}$  that are non-trivial, not  $\mathcal{F}$ -peripheral, and not equal to  $F_k$  itself, see Section 2.2). Given a set  $X$ , we denote by  $\mathcal{P}_{<\infty}(X)$  the set of all non-empty finite subsets of  $X$ .

**Theorem 12.6** (Witness map [Hor14, Lemma 5.5], [Rey12]). *There exists an  $\text{Out}(F_k, \mathcal{F})$ -equivariant Borel map*

$$\mathbb{P}\overline{\mathcal{O}} \setminus \mathbb{P}\mathcal{AT} \rightarrow \mathcal{P}_{<\infty}(\text{FF}).$$

Let now  $\mu : Y \rightarrow \mathbb{P}\overline{\mathcal{O}}$  be a map as in Lemma 12.5.

**Lemma 12.7.** *For a.e.  $y \in Y$ , the probability measure  $\mu_y$  gives full measure to  $\mathbb{P}\mathcal{AT}$ .*



*Proof.* Otherwise, let  $U \subseteq Y$  be a Borel subset of positive measure such that for every  $y \in U$ , one has  $\mu_y(\mathbb{P}\overline{\mathcal{O}} \setminus \mathbb{P}\mathcal{AT}) > 0$ . By renormalizing, we deduce an  $(\mathcal{A}|_U, \rho)$ -equivariant Borel map  $U \rightarrow \text{Prob}(\mathbb{P}\overline{\mathcal{O}} \setminus \mathbb{P}\mathcal{AT})$ . Via the witness map given by Theorem 12.6, we get for each  $y \in U$  a probability measure on  $\mathcal{P}_{<\infty}(\text{FF})$ . This yields a contradiction to the following lemma.  $\square$

**Lemma 12.8.** *For every Borel subset  $U \subseteq Y$  of positive measure, there is no  $(\mathcal{A}|_U, \rho)$ -equivariant Borel map that assigns to every  $y \in U$  a probability measure on  $\mathcal{P}_{<\infty}(\text{FF})$ .*

*Proof.* There is an  $\text{Out}(F_k, \mathcal{F})$ -equivariant Borel map  $\text{Prob}(\mathcal{P}_{<\infty}(\text{FF})) \rightarrow \mathcal{P}_{<\infty}(\text{FF})$ , assigning to every probability measure  $\nu$  the union of all finite subsets of  $\text{FF}$  with maximal  $\nu$ -measure. It is therefore enough to show that there is no  $(\mathcal{A}|_U, \rho)$ -invariant Borel map  $U \rightarrow \mathcal{P}_{<\infty}(\text{FF})$ . Assume towards a contradiction that such a map  $\psi$  exists. Since  $\psi$  takes values in the countable set  $\mathcal{P}_{<\infty}(\text{FF})$ , let  $V \subseteq U$  be a Borel subset of positive measure such that  $\psi|_V$  is constant, with value a nonempty finite collection  $\mathcal{C} \subseteq \text{FF}$ . Since  $\rho$  takes its values in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ , it follows that every free factor in  $\mathcal{C}$  is  $(\mathcal{A}|_V, \rho)$ -invariant. Choosing  $[A] \in \mathcal{C}$ , it follows that the free factor system  $\mathcal{F} \vee \{[A]\}$  (see Section 2.2) is  $(\mathcal{A}|_V, \rho)$ -invariant. This contradicts that  $\mathcal{F}$  is everywhere maximal (Lemma 12.3).  $\square$

### 12.3 A barycenter argument

Two arational  $(F_k, \mathcal{F})$ -trees  $T$  and  $T'$  are said to be *equivalent* (which we denote as  $T \sim T'$ ) if there exist  $F_k$ -equivariant alignment-preserving maps from  $T$  to  $T'$  and from  $T'$  to  $T$  (recall that a map  $T \rightarrow T'$  is *alignment-preserving* if it maps every segment in  $T$  to a segment in  $T'$ ).

Given a space  $X$ , we denote by  $X^{(3)}$  the subspace of  $X^3$  made of all pairwise distinct triples.

The following theorem was proved in a joint work with Lécureux [GHL20] – as a matter of fact  $\mathbb{P}\mathcal{AT}/\sim$  is also the Gromov boundary of the free factor graph of  $(F_k, \mathcal{F})$  [BR15, Ham12, GH19b], and the following statement can therefore be viewed as giving a barycenter map on the free factor graph.

**Theorem 12.9** ([GHL20, Theorem 8.1]). *There exists an  $\text{Out}(F_k, \mathcal{F})$ -equivariant Borel map*

$$(\mathbb{P}\mathcal{AT}/\sim)^{(3)} \rightarrow \mathcal{P}_{<\infty}(\text{FF}).$$

In fact [GHL20, Theorem 8.1] gives a stronger statement assigning to every triple in  $(\mathbb{P}\mathcal{AT}/\sim)^{(3)}$  a finite set of free splittings of  $(F_k, \mathcal{F})$ . Theorem 12.9 follows as there is a canonical way to associate a finite set of conjugacy classes of proper free factors of  $(F_k, \mathcal{F})$  to every free splitting  $S$  of  $(F_k, \mathcal{F})$ , by considering all factors that are elliptic in either  $S$  or some collapse of  $S$ .

By Lemma 12.7, we have for a.e.  $y \in Y$  a probability measure  $\mu_y$  on  $\mathbb{P}\mathcal{AT}$ . The following lemma shows that for a.e.  $y \in Y$ , its image under the quotient map  $\mathbb{P}\mathcal{AT} \rightarrow \mathbb{P}\mathcal{AT}/\sim$  is supported on a set of cardinality at most 2.



**Lemma 12.10.** *For every subset of positive measure  $U \subseteq Y$ , every  $(\mathcal{A}|_U, \rho)$ -equivariant Borel map  $\nu : U \rightarrow \text{Prob}(\mathbb{P}\mathcal{AT}/\sim)$ , and a.e.  $y \in U$ , the support of  $\nu_y$  has cardinality at most 2.*

*Proof.* Assume by contradiction that the support of  $\nu_y$  has cardinality at least 3 on a Borel subset  $U' \subseteq U$  of positive measure. Then for all  $y \in U'$ , the probability measure  $\nu_y \otimes \nu_y \otimes \nu_y$  on  $(\mathbb{P}\mathcal{AT}/\sim)^3$  gives positive measure to the subset  $(\mathbb{P}\mathcal{AT}/\sim)^{(3)}$  of pairwise distinct triples. Therefore, we get an  $(\mathcal{A}|_{U'}, \rho)$ -equivariant Borel map  $U' \rightarrow \text{Prob}((\mathbb{P}\mathcal{AT}/\sim)^{(3)})$ . Using Theorem 12.9, we deduce an  $(\mathcal{A}|_{U'}, \rho)$ -equivariant Borel map  $U' \rightarrow \text{Prob}(\mathcal{P}_{<\infty}(\text{FF}))$ . This contradicts Lemma 12.8.  $\square$

Given a set  $X$ , we denote by  $\mathcal{P}_{\leq 2}(X)$  the collection of all nonempty subsets of  $X$  of cardinality at most 2.

**Corollary 12.11.** *There exists an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $\psi_0 : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$ .*

*Proof.* Lemmas 12.5 and 12.7 show that there exists an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $\mu : Y \rightarrow \text{Prob}(\mathbb{P}\mathcal{AT})$ , and  $\mu$  then induces an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $\nu : Y \rightarrow \text{Prob}(\mathbb{P}\mathcal{AT}/\sim)$ . Lemma 12.10 ensures that for a.e.  $y \in Y$ , the support of  $\nu_y$  has cardinality at most 2. The map sending  $y$  to the support of  $\nu_y$  is then an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  (defined almost everywhere).  $\square$

## 12.4 Canonicity

**Lemma 12.12.** *There exists an essentially unique  $(\mathcal{A}, \rho)$ -equivariant Borel map  $\varphi : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  which is everywhere maximal in the following sense: for every Borel subset  $U \subseteq Y$  of positive measure, every  $(\mathcal{A}|_U, \rho)$ -equivariant Borel map  $\psi : U \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$ , and almost every  $y \in U$ ,  $\psi(y)$  is contained in  $\varphi(y)$ .*

*Proof.* The argument is exactly Adams' in [Ada94, Lemmas 3.2 and 3.3]. The uniqueness is clear.

To prove the existence, we observe that if  $\psi, \psi' : U \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  are two  $(\mathcal{A}|_U, \rho)$ -equivariant Borel maps, then for almost every  $y \in U$ ,  $\psi(y) \cup \psi'(y)$  has cardinality at most 2 by Lemma 12.10. So one would like to construct  $\varphi$  as the union of all maps  $\psi$ , but one has to be careful because of measurability issues.

For  $U \subseteq Y$  we denote by  $\bar{U}$  the  $\mathcal{A}$ -saturation of  $U$  i.e. the set of points  $y \in Y$  such that there exists an element of  $\mathcal{A}$  with source  $y$  and range in  $U$ . We observe that if  $\psi : U \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  is  $(\mathcal{A}|_U, \rho)$ -equivariant, it extends naturally to an  $(\mathcal{A}|_{\bar{U}}, \rho)$ -equivariant map  $\bar{\psi} : \bar{U} \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  by letting  $\bar{\psi}(r(g)) = \rho(g)\psi(s(g))$  for all  $g \in \mathcal{G}_{\bar{U}}$  with  $s(g) \in U$  (this does not depend on choices). It can then be extended to  $Y$  by  $\bar{\psi}(y) = \psi_0(y)$  for all  $y \notin \bar{U}$  (where  $\psi_0$  is a map provided by Corollary 12.11).

Let  $U \subseteq Y$  be a Borel subset of maximal measure such that there exists a stably  $(\mathcal{A}|_U, \rho)$ -equivariant Borel map  $\varphi'$  from  $U$  to the set  $\mathcal{P}_{=2}(\mathbb{P}\mathcal{AT}/\sim)$  of all subsets of  $\mathbb{P}\mathcal{AT}/\sim$  of cardinal exactly 2. Then there is actually an  $(\mathcal{A}|_U, \rho)$ -equivariant Borel map  $\varphi : U \rightarrow \mathcal{P}_{=2}(\mathbb{P}\mathcal{AT}/\sim)$  (this is proved by starting from a Borel subset  $V \subseteq U$  on which  $\varphi'$  is

actually equivariant, extending it to an equivariant map on  $\bar{V}$  as above, and repeating the process on the complement of  $\bar{V}$ , at most countably many times). The set  $U$  is then almost equal to its saturation  $\bar{U}$ , and extending  $\varphi$  on  $Y \setminus \bar{U}$  using  $\psi_0$ , one gets a map  $\varphi : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  which is  $(\mathcal{A}, \rho)$ -equivariant and maximal as required.  $\square$

The advantage of the maximal map constructed in Lemma 12.12 is that it is also invariant under the given subgroupoid  $\mathcal{H}$  which normalizes  $\mathcal{A}$ .

**Corollary 12.13.** *There exists a Borel map  $\varphi : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  which is both  $(\mathcal{A}, \rho)$ -equivariant and  $(\mathcal{H}, \rho)$ -equivariant.*

*Proof.* Let  $\varphi : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  be the essentially unique maximal  $(\mathcal{A}, \rho)$ -equivariant map given by the previous lemma. Since  $\mathcal{H}$  normalizes  $\mathcal{A}$ , one can write  $\mathcal{H}$  as a countable union of bisections  $B_n \subseteq \mathcal{H}$  such that for all  $a \in \mathcal{A}$  and all  $\gamma, \delta \in B_n$  such that  $\gamma a \delta^{-1}$  is defined, one has  $a \in \mathcal{A}$  if and only if  $\gamma a \delta^{-1} \in \mathcal{A}$ . Up to subdividing  $B_n$ , we may assume without loss of generality that the cocycle  $\rho$  takes a single value on  $B_n$ , say  $\rho(B_n) = \{\beta_n\}$ . Let  $f_{B_n} : U_n \rightarrow V_n$  be the partial isomorphism associated to the bisection  $B_n$ , with  $U_n, V_n \subseteq Y$ .

Assume that  $U_n$  has positive measure. The map

$$\begin{aligned} \varphi' : U_n &\rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim) \\ y &\mapsto \beta_n^{-1} \cdot \varphi(f_{B_n}(y)) \end{aligned}$$

is a maximal  $(\mathcal{A}|_{U_n}, \rho)$ -equivariant Borel map so there is a subset of full measure  $U'_n \subseteq U_n$  such that  $\varphi'$  and  $\varphi$  coincide on  $U'_n$ . This still holds if  $U_n$  has measure zero taking  $U'_n = \emptyset$ . Taking  $Y' = \cup_n U'_n$ , this shows that  $\varphi|_{Y'}$  is  $(\mathcal{H}|_{Y'}, \rho)$ -equivariant.  $\square$

At this point, we would like to make an observation, which will be useful later in Section 12.9 in order to establish a slight refinement of the main theorem of the present section. We insist (as already mentioned in Remark 12.4) that in Sections 12.2, 12.3, 12.4, we never really used the amenability of  $\mathcal{A}$ . Instead, what we used is the existence of an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $\mu : Y \rightarrow \text{Prob}(\mathbb{P}\mathcal{O})$  established in Section 12.1. We record this in the following proposition for future use.

**Proposition 12.14.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , let  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle, and let  $\mathcal{A}, \mathcal{H}$  be measured subgroupoids of  $\mathcal{G}$ . Assume that  $(\mathcal{A}, \rho)$  is pure, and let  $\hat{\mathcal{F}}$  be the maximal  $(\mathcal{A}, \rho)$ -invariant almost free factor system, and  $\mathcal{F} \subseteq \hat{\mathcal{F}}$  be an extracted free factor system.*

*Assume that  $\mathcal{H}$  normalizes  $\mathcal{A}$  and that there exists an  $(\mathcal{A}, \rho)$ -equivariant Borel map  $Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$ .*

*Then there exists a Borel map  $Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  which is both  $(\mathcal{A}, \rho)$ -equivariant and  $(\mathcal{H}, \rho)$ -equivariant.  $\square$*

As in our standing assumptions, the space  $\mathbb{P}\mathcal{AT}$  consists of projective arational trees relative to the free factor system  $\mathcal{F}$  defined in the statement. Since  $\mathcal{H}$  normalizes  $\mathcal{A}$ , the unique maximal  $(\mathcal{A}, \rho)$ -invariant almost free factor system  $\hat{\mathcal{F}}$  is  $(\mathcal{H}, \rho)$ -invariant, and as we are working in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ , so is  $\mathcal{F}$ . However,  $\mathcal{H}$  could preserve larger almost free factor systems.

## 12.5 Amenability or decomposability: setup and strategy

Remember that we aim to either prove amenability of the groupoid  $\mathcal{H}$ , or provide an  $(\mathcal{H}|_U, \rho)$ -invariant nice splitting for some Borel subset  $U \subseteq Y$  of positive measure. When  $\mathcal{F} = \emptyset$  we rely on a joint work with Bestvina [BGH17] showing that the  $\text{Out}(F_k)$ -action on  $\mathbb{P}\mathcal{AT}$  is Borel amenable [BGH17, Theorem 6.4] – see Definition 3.36 for the notion of Borel amenability of a group action.

More generally, for every free factor system  $\mathcal{F}$ , it is proved in [BGH17, Theorem 6.4] that the action of  $\text{Out}(F_k, \mathcal{F}^{(t)})$  on  $\mathbb{P}\mathcal{AT}$  is Borel amenable (recall that  $\text{Out}(F_k, \mathcal{F}^{(t)})$  denotes the subgroup of  $\text{Out}(F_k, \mathcal{F})$  made of all automorphisms acting trivially on each free factor in  $\mathcal{F}$ ).

But when  $\mathcal{F}$  contains a non-cyclic free factor (and is non-sporadic), the  $\text{Out}(F_k, \mathcal{F})$ -action on  $\mathbb{P}\mathcal{AT}(F_k, \mathcal{F})$  is not Borel amenable because there exist arational  $(F_k, \mathcal{F})$ -trees with non-amenable stabilizer. A typical example is the following: let  $\Sigma$  be a hyperbolic surface with one boundary component, let  $c$  be an essential simple closed curve on  $\Sigma$  that separates  $\Sigma$  into two connected components  $A$  and  $B$  (where  $A$  contains the boundary, say), and let  $T$  be a tree dual to an arational measured lamination on  $A$ . Then  $T$  is an arational  $(F_k, \pi_1(B))$ -tree, but its stabilizer contains all automorphisms induced by homeomorphisms of  $B$  acting as the identity on  $c$ .

We view  $\mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  as a subset of  $\mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT}/\sim)$ , and we consider the partition

$$\mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT}/\sim) = \mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim) \sqcup \mathcal{P}_{< \infty}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$$

where  $\mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim)$  (resp.  $\mathcal{P}_{< \infty}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$ ) is the set of non-empty finite collections of equivalence classes of projective arational trees, whose setwise (or equivalently elementwise) stabilizer is amenable (resp. non-amenable). These are Borel subsets of  $\mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT}/\sim)$ . Building on [BGH17], we will prove in Section 12.6 that the action of  $\text{Out}(F_k, \mathcal{F})$  on  $\mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim)$  is Borel amenable. On the other hand, using ideas from [GL], we show in Section 12.7 that there is an  $\text{Out}(F_k, \mathcal{F})$ -equivariant map assigning to any collection in  $\mathcal{P}_{< \infty}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$  a nice splitting witnessing that its stabilizer is non-amenable. This will allow us to complete the argument in Section 12.8 below.

## 12.6 Amenability of the action on $\mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim)$

The goal of this section is to prove the following theorem.

**Theorem 12.15.** *The action of  $\text{Out}(F_k, \mathcal{F})$  on  $\mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim)$  is Borel amenable.*

**Corollary 12.16.** *The action of  $\text{Out}(F_k, \mathcal{F})$  on the subset of  $\mathbb{P}\mathcal{AT}/\sim$  made of points with amenable stabilizer is Borel amenable. This also holds for the action on the set of pairs in  $\mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  whose stabilizer is amenable.  $\square$*

Given a tree  $T \in \mathcal{AT}$ , we denote by  $[T]$  and  $[[T]]$  its images in  $\mathbb{P}\mathcal{AT}$  and  $\mathbb{P}\mathcal{AT}/\sim$ , respectively. Similarly, given  $\mathcal{T} = \{T_1, \dots, T_p\} \in \mathcal{P}_{< \infty}(\mathcal{AT})$  we denote by  $[\mathcal{T}]$  and  $[[\mathcal{T}]]$  its images in  $\mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT})$  and in  $\mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT}/\sim)$ , respectively.

**Lemma 12.17.** *For  $\mathcal{T} \in \mathcal{P}_{< \infty}(\mathcal{AT})$ , the following are equivalent:*

1.  $\text{Stab}(\mathcal{T})$  is amenable;
2.  $\text{Stab}([\mathcal{T}])$  is amenable;
3.  $\text{Stab}([\![\mathcal{T}]\!])$  is amenable.

*Proof.* As  $\text{Stab}(\mathcal{T}) \subseteq \text{Stab}([\mathcal{T}]) \subseteq \text{Stab}([\![\mathcal{T}]\!])$ , the implications  $3 \Rightarrow 2 \Rightarrow 1$  are clear, and it is enough to prove that  $1 \Rightarrow 3$ . We thus assume that  $\text{Stab}(\mathcal{T})$  is amenable, and aim to show that  $\text{Stab}([\![\mathcal{T}]\!])$  is amenable. Let  $\text{Stab}^0(\mathcal{T})$  be the finite-index subgroup made of all elements that fix the isometry class of every  $T \in \mathcal{T}$  (as opposed to permuting them).

Write  $\mathcal{T} = \{T_1, \dots, T_p\}$ . For every  $i \in \{1, \dots, p\}$ , the preimage of  $[[T_i]]$  in  $\mathcal{AT}$  is a cone  $\mathcal{C}_i$  on a finite set of trees  $T_{i,1}, \dots, T_{i,\ell_i}$  (called *ergometric*), see [GH19a, Proposition 13.5] relying on [Gui00, Section 5]. A finite index subgroup  $\text{Stab}^1([\![\mathcal{T}]\!]) \subseteq \text{Stab}([\![\mathcal{T}]\!])$  acts by homothety on each of the trees  $T_{i,j}$ , and the scaling factors define a homomorphism from  $\text{Stab}^1([\![\mathcal{T}]\!])$  to the abelian group  $(\mathbb{R}_+^*)^n$  (with  $n = \ell_1 + \dots + \ell_p$ ). The kernel  $K$  of this homomorphism acts by isometry on each tree  $T_{i,j}$  and fixes each cone  $\mathcal{C}_i$  pointwise. In particular  $K \subseteq \text{Stab}^0(\mathcal{T})$ . As  $\text{Stab}^0(\mathcal{T})$  is amenable, we deduce that  $K$  is amenable, and therefore so are  $\text{Stab}^1([\![\mathcal{T}]\!])$  and  $\text{Stab}([\![\mathcal{T}]\!])$ . This concludes our proof.  $\square$

We will use a multisection of the quotient map  $\pi : \mathbb{P}\mathcal{AT} \rightarrow \mathbb{P}\mathcal{AT}/\sim$  as in the following lemma.

**Lemma 12.18.** *There exists an  $\text{Out}(F_k, \mathcal{F})$ -equivariant Borel map  $\sigma : \mathbb{P}\mathcal{AT}/\sim \rightarrow \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$  such that for all  $[[T]] \in \mathbb{P}\mathcal{AT}/\sim$ , the set  $\sigma([[T]])$  is a finite set of representatives of  $[[T]]$  in  $\mathbb{P}\mathcal{AT}$ .*

*Proof.* By [GHL20, Lemmas 7.4 and 7.5], there exists a sequence of (not equivariant) Borel maps  $f_n : \mathbb{P}\mathcal{AT} \rightarrow \mathbb{P}\mathcal{AT}$  such that for all  $[T] \in \mathbb{P}\mathcal{AT}$ , the set  $\{f_n([T]) \mid n \in \mathbb{N}\}$  is the finite set of all projective ergometric trees that are  $\sim$ -equivalent to  $T$ . Thus, the map  $\tilde{\sigma} : \mathbb{P}\mathcal{AT} \rightarrow \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$  defined by  $[T] \mapsto \{f_n([T]) \mid n \in \mathbb{N}\}$  is Borel. It descends to a Borel map  $\sigma$  on  $\mathbb{P}\mathcal{AT}/\sim$  which is equivariant because  $\sigma([[T]])$  is exactly the set of ergometric trees in  $\mathbb{P}\mathcal{AT}$  in the preimage of  $[[T]]$ .  $\square$

We denote by  $\mathcal{P}_{<\infty}^a(\mathbb{P}\mathcal{AT}) \subseteq \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$  the set of finite collections of projective trees  $\{[T_1], \dots, [T_p]\}$  whose stabilizer is amenable. Theorem 12.15 will follow from the following proposition.

**Proposition 12.19.** *The action of  $\text{Out}(F_k, \mathcal{F})$  on  $\mathcal{P}_{<\infty}^a(\mathbb{P}\mathcal{AT})$  is Borel amenable.*

*Proof of Theorem 12.15 using Proposition 12.19.* By Proposition 12.19, there exists a sequence of Borel maps

$$\nu_n : \mathcal{P}_{<\infty}^a(\mathbb{P}\mathcal{AT}) \rightarrow \text{Prob}(\text{Out}(F_k, \mathcal{F}))$$

which is asymptotically  $\text{Out}(F_k, \mathcal{F})$ -equivariant. Let  $\sigma : \mathbb{P}\mathcal{AT}/\sim \rightarrow \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$  be a Borel equivariant section, as provided by Lemma 12.18.

Note that if the stabilizer of  $\{[[T_1]], \dots, [[T_p]]\}$  is amenable, then so is the stabilizer of  $\{[S_1], \dots, [S_p]\}$  for any  $\{[S_1], \dots, [S_p]\} \in \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$  such that  $[[S_i]] = [[T_i]]$  for all  $i \leq p$ . Define  $\nu'_n : \mathcal{P}_{<\infty}^a(\mathbb{P}\mathcal{AT}/\sim) \rightarrow \text{Prob}(\text{Out}(F_k, \mathcal{F}))$  by

$$\nu'_n(\{[[T_1]], \dots, [[T_p]]\}) = \frac{1}{\#\sigma([[T_1]]) \cdots \#\sigma([[T_p]])} \sum_{[S_i] \in \sigma([[T_i]]), i \leq p} \nu_n(\{[S_1], \dots, [S_p]\}).$$

Since  $\nu_n$  is asymptotically  $\text{Out}(F_k, \mathcal{F})$ -equivariant, so is  $\nu'_n$ .  $\square$

*Proof of Proposition 12.19.* We first explain the amenability of the action on the set  $\mathbb{P}\mathcal{AT}^a$  of projective trees with amenable stabilizer. We will adapt the proof from [BGH17, Theorem 6.4], establishing that the action of  $\text{Out}(F_k, \mathcal{F}^{(t)})$  on  $\mathbb{P}\mathcal{AT}$  is Borel amenable; our proof will make use of the chain condition on stabilizers of collections of free splittings provided by Proposition 6.17.

Given a simplex  $\Delta$  of the projectivized Outer space  $\mathbb{P}\mathcal{O}$  relative to  $\mathcal{F}$ , we denote by  $\tilde{\Delta}$  its preimage in  $\mathcal{O}$ . We recall that a map  $f : S \rightarrow T$  between two  $F_k$ -trees is a *morphism* if every segment  $I \subseteq S$  can be subdivided into finitely many subsegments  $I_1, \dots, I_n$ , so that the restriction of  $f$  to each subsegment  $I_j$  is an isometry. It is *optimal* if every point  $x \in S$  is contained in an open interval  $I_x$  such that  $f|_{I_x}$  is injective.

Let  $T \in \mathcal{AT}^a$ , let  $S \in \mathcal{O}$ , and let  $f : S \rightarrow T$  be an optimal morphism. Given positive real numbers  $s < t$ , we denote by  $[s, t]_f$  the collection of all simplices  $\Delta$  of  $\mathbb{P}\mathcal{O}$  such that there exists a tree  $S' \in \tilde{\Delta}$  through which  $f$  factors, with  $e^{-t} \leq \text{vol}(S'/F_k) \leq e^{-s}$  (here  $\text{vol}(S'/F_k)$  denotes the volume of the quotient graph  $S'/F_k$ , i.e. the sum of its edge lengths). By [BGH17, Proposition 3.6 and Remark 3.9], the set  $[s, t]_f$  is always finite.

Let  $\mathcal{D}$  be the countable set of all finite collections of simplices of  $\mathbb{P}\mathcal{O}$ , and let  $\mathcal{D}^a \subseteq \mathcal{D}$  be the subset consisting of collections whose elementwise stabilizer in  $\text{Out}(F_k, \mathcal{F})$  is amenable. For every optimal morphism  $f$  whose source lies in  $\mathcal{O}$  and whose range is a tree in  $\mathcal{AT}^a$ , and every  $n \in \mathbb{N}$ , we define a probability measure  $\mu_n(f)$  on  $\mathcal{D}^a$  as follows.

The chain condition in Proposition 6.17 ensures that for each  $t \in \mathbb{R}$ , the elementwise stabilizer  $H_{t,m}$  of  $[t, t+m]_f$  in  $\text{Out}(F_k, \mathcal{F})$  stabilizes to a group  $H_{t,\infty}$  as  $m$  goes to infinity. We claim that  $H_{t,\infty}$  is amenable. Indeed, the subgroup of finite index  $H_{t,\infty}^0 = H_{t,\infty} \cap \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  acts as the identity on each simplex occurring in  $\bigcup_{m \geq 0} [t, t+m]_f$ . Now let  $\rho : [0, \infty) \rightarrow \mathcal{O}$  be a folding path from  $S$  to  $T$  such that  $f$  factors through  $\rho(t)$  for all  $t \geq 0$  (for the existence, see for instance [GL07b], and [BGH17, Lemma 2.3] for the fact that it stays in the interior of Outer space). Since  $H_{t,\infty}^0$  fixes a subray of the image of  $\rho$  in  $\mathbb{P}\mathcal{O}$ , it fixes  $[T]$ . Since the stabilizer of  $[T]$  is amenable, so are  $H_{t,\infty}^0$  and  $H_{t,\infty}$  which proves the claim.

We now assume we are given  $M_n(f) \in \mathbb{N}$  depending measurably on  $f$ , such that the elementwise stabilizer of  $[t, t+M_n(f)]_f$  in  $\text{Out}(F_k, \mathcal{F})$  is amenable for all  $t \in [n, 2n]$  (the construction of  $M_n(f)$  with these properties will be carried out in the next paragraph). We then define

$$\mu_n(f) := \frac{1}{n} \int_n^{2n} \delta_{[t, t+M_n(f)]_f} dt,$$

where  $\delta_{[t, t+M_n(f)]_f}$  is the Dirac measure on  $[t, t+M_n(f)]_f$  viewed as a point in  $\mathcal{D}^a$  – so  $\mu_n(f)$  is a probability measure on  $\mathcal{D}^a$ . The map  $f \mapsto \mu_n(f)$  is measurable. Since  $\mathcal{AT}$  has namable turning classes in the sense of [BGH17], the proof of [BGH17, Proposition 3.24] allows us to associate to every projective tree  $[T] \in \mathbb{P}\mathcal{AT}^a$  an asymptotically equivariant sequence of probability measures  $\mu_n([T])$  on  $\mathcal{D}^a$  that depend measurably on  $[T]$ <sup>5</sup>. Since the stabilizer of every point in  $\mathcal{D}^a$  is amenable, it follows from [BGH17, Proposition 2.11] that the  $\text{Out}(F_k, \mathcal{F})$ -action on  $\mathbb{P}\mathcal{AT}^a$  is Borel amenable.

We now explain how to construct  $M_n(f)$ . As already noted, for every  $t \geq 0$ ,  $H_{t,m}$  is amenable for  $m$  large enough. Let  $m_n(f)$  be the smallest integer  $m$  such that  $H_{n,m}$  is amenable, and let  $M_n(f) := 1 + \max_{n \leq k \leq 2n} m_k(f)$ . One checks as in the proof of [BGH17, Lemma 3.13] that for all  $n, m \in \mathbb{N}$ , the set  $[n, n+m]_f$  depends measurably on the optimal morphism  $f$ . Therefore, for all  $n \in \mathbb{N}$ , the integers  $m_n(f)$  and  $M_n(f)$  depend measurably on  $f$ . This concludes the proof that the action on  $\mathbb{P}\mathcal{AT}^a$  is Borel amenable.

To prove that the action on  $\mathcal{P}_{<\infty}^a(\mathbb{P}\mathcal{AT})$  is amenable, we proceed in a similar fashion. Consider a finite collection  $\mathcal{F} = \{f_1, \dots, f_p\}$  of optimal morphisms  $f_i : S_i \rightarrow T_i$  with  $\mathcal{T} = \{[T_1], \dots, [T_p]\} \in \mathcal{P}_{<\infty}^a(\mathbb{P}\mathcal{AT})$ . We define  $[t, t+M]_{\mathcal{F}} = [t, t+M]_{f_1} \cup \dots \cup [t, t+M]_{f_p}$  and construct  $M_n(\mathcal{F})$  so that for all  $t \in [n, 2n]$ , the elementwise stabilizer of  $[t, t+M]_{\mathcal{F}}$  is amenable. This is possible because by the chain condition given by Proposition 6.17, the stabiliser of  $[n, n+m]_{\mathcal{F}}$  in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  stabilizes to a group that fixes folding paths joining  $S_i$  to  $T_i$ . One then defines

$$\mu_n(\mathcal{F}) := \frac{1}{n} \int_n^{2n} \delta_{[t, t+M_n(\mathcal{F})]_{\mathcal{F}}} dt$$

and concludes as above. □

## 12.7 A witness map for trees with infinite stabilizer

We denote by  $\mathcal{P}_{\leq 2}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$  the set of pairs of (maybe not distinct) equivalence classes of trees  $[[T]], [[T']] \in \mathbb{P}\mathcal{AT}/\sim$ , such that the common stabilizer of  $[[T]]$  and  $[[T']]$  is not amenable.

**Proposition 12.20.** *Let  $\mathcal{F}$  be a non-sporadic free factor system of  $F_k$ .*

*There exists an  $\text{Out}(F_k, \mathcal{F})$ -equivariant Borel map assigning to any  $\{[[T]], [[T']]\} \in \mathcal{P}_{\leq 2}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$  a nice splitting  $S_{\{[[T]], [[T']]\}}$ .*

The proof follows the ideas introduced in [GL], but we give a self-contained proof. The proposition is based on the following result – the definition of a bifflexible splitting was given in Definition 6.14.

---

<sup>5</sup>Roughly, the idea is the following: to every arational tree  $T$ , one can associate in a measurable and  $\text{Out}(F_k, \mathcal{F})$ -equivariant way a finite set of preferred turning classes of  $T$  (a turning class should be seen as a prescription of turns in  $T$  that are taken by a morphism to  $T$ ), and then average the measures  $\mu_n(f)$  over a set of representatives  $f$  of morphisms taking these turning classes. Composing with a (not equivariant) continuous section  $\mathbb{P}\mathcal{AT} \rightarrow \mathcal{AT}$  does not destroy the asymptotic equivariance.



**Proposition 12.21.** *Let  $\mathcal{F}$  be a non-sporadic free factor system of  $F_k$ .*

*There exists an  $\text{Out}(F_k, \mathcal{F})$ -equivariant Borel map assigning to any finite set of trees  $[\mathcal{T}] = \{[T_1], \dots, [T_p]\} \in \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$  a splitting  $S_{[\mathcal{T}]}$  compatible with  $T_1, \dots, T_p$ , such that:*

- *if the common isometric stabilizer of  $T_1, \dots, T_p$  is infinite, then  $S_{[\mathcal{T}]}$  is a non-trivial splitting which is either biflexible or has an edge with  $\mathcal{Z}_{\max}$  stabilizer;*
- *if the common isometric stabilizer of  $T_1, \dots, T_p$  is finite, then  $S_{[\mathcal{T}]}$  is trivial.*

Note that if  $T, T' \in \mathcal{AT}$  represent the same projective tree  $[T] = [T'] \in \mathbb{P}\mathcal{AT}$ , then  $T$  and  $T'$  have the same isometric stabilizer, and the same compatible splittings.

*Proof of Proposition 12.20 from Proposition 12.21.* Consider an equivariant Borel multisection  $\sigma : \mathbb{P}\mathcal{AT}/\sim \rightarrow \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$ , provided by Lemma 12.18. Given  $\{[[T]], [[T']]\} \in \mathcal{P}_{\leq 2}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$ , consider  $[\mathcal{T}] = \sigma([[T]]) \cup \sigma([[T']]) = \{[T_1], \dots, [T_p]\} \in \mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$ , and the splitting  $S_{[\mathcal{T}]}$  from Proposition 12.21.

By Lemma 12.17, the common isometric stabilizer of  $T_1, \dots, T_p$  is non-amenable and in particular infinite so  $S_{[\mathcal{T}]}$  is either biflexible or has an edge with  $\mathcal{Z}_{\max}$  stabilizer. By Corollary 9.5, one can equivariantly assign a nice splitting to  $S_{[\mathcal{T}]}$ . This concludes the proof.  $\square$

Recall that we say that a subgroup  $\Gamma$  of  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  acts trivially on a subgroup  $A$  if every  $\alpha \in \Gamma$  has a lift in  $\text{Aut}(F_k)$  acting as the identity on  $A$ .

**Lemma 12.22.** *Let  $T \in \mathcal{AT}$ , and  $H \subseteq \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  a subgroup of the isometric stabilizer of  $T$ . Up to conjugacy, there exists a unique maximal subgroup  $A_H \subseteq F_k$  which is non-abelian and not  $\mathcal{F}$ -peripheral and such that  $H$  acts trivially on  $A_H$ .*

*Moreover, let  $Y \subseteq T$  be the minimal  $A_H$ -invariant subtree. Then the translates of  $Y$  form a transverse covering of  $T$ , and  $A_H$  is the global stabilizer of  $Y$ .*

*Proof.* Given  $\alpha \in H$  and a representative  $\tilde{\alpha} \in \text{Aut}(F_k)$ , we denote by  $I_{\tilde{\alpha}}$  the  $\tilde{\alpha}$ -equivariant isometry of  $T$ . Let  $\eta$  be a germ of direction at a branch point of  $T$ . Let  $H^0$  be the subgroup of  $H$  consisting of elements preserving the  $F_k$ -orbit of  $\eta$ . As there are finitely many  $F_k$ -orbits of directions at branch points in  $T$  ([GL95]), the subgroup  $H^0$  has finite index in  $H$ . We will see below that in fact  $H^0 = H$ . For each  $\alpha \in H^0$ , there exists a unique representative  $\tilde{\alpha} \in \text{Aut}(F_k)$  such that  $I_{\tilde{\alpha}}$  fixes  $\eta$ . The set of all such representatives  $\tilde{\alpha}$  defines a lift  $\tilde{H}^0$  of  $H^0$  in  $\text{Aut}(F_k)$ .

Given  $\tilde{\alpha}_1, \dots, \tilde{\alpha}_n \in \tilde{H}^0$ , let  $Y_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$  be the set of points in  $T$  fixed by  $I_{\tilde{\alpha}_i}$  for all  $i \leq n$ , and let  $A_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n} = \text{Fix}_{F_k}(\langle \tilde{\alpha}_1, \dots, \tilde{\alpha}_n \rangle)$  be the set of elements of  $F_k$  fixed by  $\tilde{\alpha}_i$  for all  $i \leq n$ . The subtree  $Y_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$  contains at least a segment representing the direction  $\eta$ .

**Claim.** *If  $g \in F_k$  is such that  $g.Y_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n} \cap Y_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$  contains a non-degenerate segment, then  $g \in A_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$ .*

*Proof of the claim.* The subtree  $g.Y_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n} \cap Y_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$  is fixed by  $I_{\tilde{\alpha}_i}$  and by  $gI_{\tilde{\alpha}_i}g^{-1}$  hence by  $gI_{\tilde{\alpha}_i}g^{-1}I_{\tilde{\alpha}_i}^{-1} = g\tilde{\alpha}_i(g^{-1})$ . Since  $F_k$ -stabilizers of arcs of  $T$  are trivial, it follows that  $\tilde{\alpha}_i(g) = g$ . We conclude that  $g \in A_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$ .  $\square$



Let now  $J \subseteq Y_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$  be a non-degenerate segment. Since  $T$  is mixing,  $J$  contains infinitely many germs of segments representing a direction in the  $F_k$ -orbit of  $\eta$ . The claim then implies that  $A_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$  contains elements of arbitrary small positive translation length, and is therefore non-abelian.

By the bounded chain condition on fixed subgroups in  $F_k$  [MV04], there exists  $\tilde{\alpha}_1, \dots, \tilde{\alpha}_n \in \tilde{H}^0$  such that  $A_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n} = A_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n, \tilde{\alpha}}$  for any  $\tilde{\alpha} \in \tilde{H}^0$ . We define  $A_H = A_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_n}$  and we note that  $H^0$  acts trivially on  $A_H$ .

We now use the fact  $H$  is contained in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  to deduce that  $H$  itself acts trivially on  $A_H$ . Since  $H^0$  preserves the conjugacy class of every element  $g \in A_H$ , its orbit under every  $\alpha \in H$  is finite. Since  $\alpha \in \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ , this implies that  $\alpha$  preserves  $[g]$  (Theorem 2.11 by Handel and Mosher). Since this holds for every  $g \in A_H$ , [GL15a, Corollary 1.6] implies that  $\alpha$  acts trivially on  $A_H$ , hence so does  $H$ .

To prove the uniqueness of  $A_H$ , assume that  $H$  acts trivially on some subgroup  $A' \subseteq F_k$  which is not abelian and not  $\mathcal{F}$ -peripheral. Since the centralizer of  $A'$  in  $F_k$  is trivial, every  $\alpha \in H$  has a unique lift  $\tilde{\alpha} \in \text{Aut}(F_k)$  acting as the identity on  $A'$ . This defines a lift  $\tilde{H}'$  of  $H$  in  $\text{Aut}(F_k)$  with  $A' \subseteq \text{Fix}_{F_k} \tilde{H}'$ . Since  $A'$  is non-abelian and not  $\mathcal{F}$ -peripheral,  $A'$  is not elliptic in  $T$  and its minimal subtree  $Y' \subseteq T$  is not a line. Since  $\tilde{H}'$  fixes  $A'$ , for every  $\tilde{\alpha} \in \tilde{H}'$ ,  $I_{\tilde{\alpha}}$  preserves the axis of every element of  $A'$  and therefore acts as the identity on  $Y'$ . Up to conjugating  $A'$  and  $\tilde{H}'$  by some element of  $F_k$ , we may assume that  $Y'$  contains a representative of  $\eta$ , so  $\tilde{H}' \subseteq \tilde{H}^0$ . Since  $\tilde{H}'$  is a lift of  $H$ , it follows that  $\tilde{H}' = \tilde{H}^0$  and  $A' \subseteq A_H$  (and it also follows that  $H^0 = H$ ). This proves the maximality and the uniqueness of  $A_H$ .

The fact that translates of the minimal  $A_H$ -invariant tree  $Y$  are transverse and that  $A_H$  is the global stabilizer of  $Y$  follows from the claim above. Since the action of  $F_k$  on  $T$  is mixing ([Rey12, Proposition 8.3], [Hor14, Lemma 4.9]), this implies that the translates of  $Y$  form a transverse covering.  $\square$

We denote by  $S_{T,H}$  the skeleton of the transverse covering of  $T$  by the translates of  $Y$ . Its vertex set is  $V_0 \sqcup V_1$  where  $V_1$  is the set of translates of  $Y$ ,  $V_0$  is the set of points  $x \in T$  lying in at least two distinct translates of  $Y$ , and there is an edge between  $x$  and  $gY$  if and only if  $x \in gY$ . Stabilizers of vertices in  $V_0$  are point stabilizers in  $T$  (as  $T$  is arational, these are either groups in  $\mathcal{F}$  or maximal cyclic), and stabilizers of vertices in  $V_1$  are conjugates of  $A_H$  (which are fixed subgroups of  $F_k$ ). In particular, vertex stabilizers of  $S_{T,H}$  are finitely generated, hence so are edge stabilizers by Howson's property. It follows that each edge stabilizer  $G_e$  is its own normalizer: being a finitely generated subgroup of  $F_k$ ,  $G_e$  has finite index in its normalizer, and since vertex stabilizers are stable under taking roots, so is  $G_e$ .

**Lemma 12.23.** *The tree  $S_{T,H}$  does not depend on  $T$ : if  $T' \in \mathcal{AT}$  is  $H$ -invariant, then  $S_{T,H} = S_{T',H}$ .*

*Remark 12.24.* Although this is not apparent in the notations, in this lemma and until the end of this section, we have fixed a free factor system  $\mathcal{F}$ , and  $S_{T,H}$  and even  $\mathcal{AT}$  do depend on  $\mathcal{F}$ .

*Proof.* We give an alternative description of  $S_{T,H}$ , independent of  $T$ . Let  $V'_1$  be the set of conjugates of  $A_H$ , and  $V'_0$  be the set of free factors whose conjugacy class belongs to  $\mathcal{F}$  and that intersect at least two distinct conjugates of  $A_H$  non-trivially. We put an edge between  $A_H^g \in V'_1$  and  $P \in V'_0$  if and only if  $A_H^g \cap P \neq \{1\}$ . Clearly, the graph  $S'$  defined in this way does not depend on  $T$ . We claim that  $S'$  is equivariantly isomorphic to  $S_{T,H}$ .

We are going to show that there is a well defined map  $f : S_{T,H} \rightarrow S'$  sending a vertex  $v$  to its stabilizer  $G_v$ . The map is well defined on  $V_1(S_{T,H})$  by sending  $v$  to  $G_v \in V'_1$ .

Every edge of  $S_{T,H}$  has non-trivial stabilizer, since otherwise, this would define a free factor of  $(F_k, \mathcal{F})$  acting non-discretely on  $T$ , contradicting arationality of  $T$ . This shows in particular that if  $v_0 \in V_0(S_{T,H})$  then  $G_{v_0}$  intersects at least two distinct conjugates of  $A_H$  non-trivially, namely the stabilizers of the neighbors of  $v_0$ . It also follows that  $G_{v_0}$  is non-trivial, and we claim that the conjugacy class of  $G_{v_0}$  lies in  $\mathcal{F}$ . If the claim does not hold, then  $T$  is not relatively free, so  $T$  has to be an arational surface tree as defined in [Hor14, Definition 4.3] (see [Hor14, Lemma 4.6]), and  $G_{v_0}$  is cyclic and non- $\mathcal{F}$ -peripheral. In particular some edge stabilizer of  $S_{T,H}$  is cyclic and non- $\mathcal{F}$ -peripheral, and as  $T$  is compatible with  $S_{T,H}$ , we get a contradiction to [Hor14, Lemma 4.8]. This proves that  $G_{v_0} \in V'_0$  and that  $f$  is well defined and sends edge to edge.

Since  $f$  is clearly injective and surjective on the set of vertices, there remains to show that  $f$  preserves non-adjacency. It suffices to prove that for  $u \in V_0$ ,  $v \in V_1$  at distance at least 3, then  $G_u \cap G_v = \{1\}$ . Consider  $u' \in [u, v] \cap V_0$  at distance 1 from  $v$ . Since  $\mathcal{F}$  is a malnormal family,  $G_u \cap G_{u'} = \{1\}$  so  $G_u \cap G_v = \{1\}$ . This concludes the proof.  $\square$

**Lemma 12.25.** *For  $\mathcal{T} = \{T_1, \dots, T_p\} \in \mathcal{P}_{<\infty}(\mathcal{AT})$ , denote by  $\Gamma_{\mathcal{T}}$  the intersection of the isometric stabilizers of  $T_1, \dots, T_p$  in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ . Let  $S_{\mathcal{T}} = S_{T_i, \Gamma_{\mathcal{T}}}$  for any  $i \leq p$ . The following are equivalent:*

1.  $\Gamma_{\mathcal{T}}$  is infinite;
2.  $S_{\mathcal{T}}$  is a non-trivial splitting of  $F_k$ ;
3.  $S_{\mathcal{T}}$  is a non-trivial splitting of  $F_k$  which is biflexible or has an edge with  $\mathcal{Z}_{\max}$  stabilizer.

*Proof.* Clearly, 3  $\Rightarrow$  2. If  $\Gamma_{\mathcal{T}}$  is finite, then it is trivial because  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  is torsion-free, so its fixed subgroup  $A_{\Gamma_{\mathcal{T}}}$  is  $F_k$ . It follows that  $S_{\mathcal{T}}$  is a point. This proves 2  $\Rightarrow$  1.

Now assume that  $\Gamma_{\mathcal{T}}$  is infinite. Then  $A_{\Gamma_{\mathcal{T}}}$  is a proper subgroup of  $F_k$ . This implies that  $F_k$  does not fix a vertex in  $S_{\mathcal{T}}$ , so  $S_{\mathcal{T}}$  is non-trivial. We know that edge stabilizers of  $S_{\mathcal{T}}$  are non-trivial (as follows from arationality of the trees  $T_i$ , see the proof of Lemma 12.23), and they are root-closed because vertex stabilizers are.

So we assume that all edge stabilizers are non-abelian, and prove that  $S_{\mathcal{T}}$  is biflexible. We first note that the group of twists of  $S_{\mathcal{T}}$  is trivial. The vertex set  $\bar{V}$  of  $S_{\mathcal{T}}/F_k$  is  $\{v_1\} \cup \bar{V}_0$  where  $G_{v_1}$  is conjugate to  $A_{\Gamma_{\mathcal{T}}}$ , and  $\bar{V}_0$  is the image of  $V_0$  in  $S_{\mathcal{T}}/F_k$ . Since  $S_{\mathcal{T}}$  is  $\Gamma_{\mathcal{T}}$ -invariant, let  $\Gamma_{\mathcal{T}}^0$  the finite index subgroup of  $\Gamma_{\mathcal{T}}$  acting as the identity on the

quotient graph  $S_{\mathcal{T}}/F_k$ . By Proposition 2.7, the natural map

$$\rho : \Gamma_{\mathcal{T}}^0 \rightarrow \prod_{v \in \bar{V}} \text{Out}(G_v, \text{Inc}_v)$$

is injective. Since  $\Gamma_{\mathcal{T}}$  acts trivially on  $A_{\Gamma_{\mathcal{T}}}$ , the image  $\rho(\Gamma_{\mathcal{T}}^0)$  is contained in  $\prod_{v \in \bar{V}_0} \text{Out}(G_v, \text{Inc}_v)$ . But since each edge stabilizer of  $S_{\mathcal{T}}$  is contained in a conjugate of  $A_{\Gamma_{\mathcal{T}}}$ ,  $\Gamma_{\mathcal{T}}$  also acts trivially on all edge groups, so  $\rho(\Gamma_{\mathcal{T}}^0)$  is in fact contained in  $\prod_{v \in \bar{V}_0} \text{Out}(G_v, \text{Inc}_v^{(t)})$  (see Remark 2.8). It follows that there exists at least one vertex  $v_0 \in \bar{V}_0$  such that  $\text{Out}(G_{v_0}; \text{Inc}_{v_0}^{(t)})$  is infinite. Since  $v_1$  has a non-sporadic Grushko decomposition relative to incident edge groups (as follows from the existence of an  $A_H$ -tree with dense orbits and trivial arc stabilizers, namely  $Y$  in Lemma 12.22), we conclude that  $S_{\mathcal{T}}$  is biflexible.  $\square$

View  $\mathcal{P}_{<\infty}(\mathcal{AT})$  as the disjoint union over  $n$  of  $\mathcal{AT}^n/\mathfrak{S}_n$ , and endow it with the corresponding Borel structure.

**Lemma 12.26.** *The map  $\mathcal{T} \in \mathcal{P}_{<\infty}(\mathcal{AT}) \mapsto S_{\mathcal{T}}$  is a Borel map.*

*Proof.* Using the description of  $S_{\mathcal{T}}$  given in the proof of Lemma 12.23, it suffices to check that the map which sends  $\mathcal{T}$  to the conjugacy class of  $A_{\Gamma_{\mathcal{T}}}$  is Borel. To this end, it is enough to check that for every finitely generated non- $\mathcal{F}$ -peripheral non-abelian subgroup  $A \subseteq F_k$ , the set of all  $\mathcal{T} \in \mathcal{P}_{<\infty}(\mathcal{AT})$  such that  $A$  is conjugate to a subgroup of  $A_{\Gamma_{\mathcal{T}}}$  is Borel.

By Lemma 12.22, the group  $A$  is conjugate into  $A_{\Gamma_{\mathcal{T}}}$  if and only if  $\Gamma_{\mathcal{T}} \subseteq \Gamma_A$ , where  $\Gamma_A$  is the subgroup of all elements of  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  acting trivially on  $A$ . This in turn is equivalent to requiring that for every  $\alpha \notin \Gamma_A$ , there exists  $T \in \mathcal{T}$  such that  $\alpha \cdot T \neq T$ , a Borel condition.  $\square$

*Proof of Proposition 12.21.* The map  $\mathcal{T} \mapsto S_{\mathcal{T}}$  is an equivariant Borel map on  $\mathcal{P}_{<\infty}(\mathcal{AT})$ . Since any two homothetic trees have the same isometric stabilizer  $\Gamma_{\mathcal{T}}$  and since  $S_{\mathcal{T}} = S_{T, \Gamma_{\mathcal{T}}}$  for any tree  $T \in \mathcal{AT}$  whose isometry class is fixed by  $\Gamma_{\mathcal{T}}$ , this map passes to the quotient into a map on  $\mathcal{P}_{<\infty}(\mathbb{P}\mathcal{AT})$ . Also, as  $S_{\mathcal{T}}$  is the skeleton of a transverse covering of any tree in  $\mathcal{T}$ , it is compatible with every tree in  $\mathcal{T}$  (see e.g. [Gui04, Lemma 4.7]). By Lemma 12.25,  $S_{\mathcal{T}}$  is biflexible or has a  $\mathcal{Z}_{\max}$  edge stabilizer if the common isometric stabilizer of the trees in  $\mathcal{T}$  is infinite.  $\square$

## 12.8 Conclusion

*Proof of Theorem 12.1.* By Proposition 10.5 applied to  $\mathcal{A}$ , there exists a Borel partition  $Y = Y_1 \sqcup Y_2$  such that  $(\mathcal{A}|_{Y_1}, \rho)$  is stably nice and  $(\mathcal{A}|_{Y_2}, \rho)$  is stably pure. As  $\rho|_{\mathcal{A}}$  is nowhere trivial, if  $Y_1$  has positive measure, Proposition 10.6 shows the existence of a subset  $U \subseteq Y_1$  of positive measure and of a nice splitting which is invariant under both  $(\mathcal{A}|_U, \rho)$  and  $(\mathcal{H}|_U, \rho)$ . We can therefore assume without loss of generality that  $(\mathcal{A}, \rho)$  is pure. Let  $\hat{\mathcal{F}}$  be the unique maximal  $(\mathcal{A}, \rho)$ -invariant almost free factor system and  $\mathcal{F}$  be an extracted free factor system. Since  $\mathcal{H}$  normalizes  $\mathcal{A}$ , we have that  $(\mathcal{H}, \rho)$  preserves  $\hat{\mathcal{F}}$ .

Since  $\rho$  takes values in  $\text{IA}_k(\mathbb{Z}/3\mathbb{Z})$ , it follows that  $(\mathcal{H}, \rho)$  also preserves  $\mathcal{F}$ . We now work relative to  $\mathcal{F}$ .

By Corollary 12.13, there is a Borel map  $\varphi : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim) \subseteq \mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT}/\sim)$  which is both  $(\mathcal{A}|_U, \rho)$  and  $(\mathcal{H}|_U, \rho)$ -equivariant. Recall the partition  $\mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT}/\sim) = \mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim) \sqcup \mathcal{P}_{< \infty}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$ .

Assume that for all  $y$  in some subset  $U \subseteq Y$  of positive measure,  $\varphi(y)$  has non-amenable stabilizer, i.e.  $\varphi(y) \in \mathcal{P}_{< \infty}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$ . Then composing with the map of Proposition 12.20, one gets a Borel map  $y \mapsto S_{\varphi(y)}$  assigning a nice splitting to every  $y \in U$ , and which is both  $(\mathcal{A}|_U, \rho)$  and  $(\mathcal{H}|_U, \rho)$ -equivariant. Since there are only countably many nice splittings, there is a subset  $V \subseteq U$  of positive measure in restriction to which  $\varphi$  is constant and defines an  $(\mathcal{H}|_V, \rho)$ -invariant nice splitting.

Thus, we may assume that  $\varphi(y) \in \mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim)$  for almost every  $y \in Y$ . Since the  $\text{Out}(F_k, \mathcal{F})$ -action on  $\mathcal{P}_{< \infty}^{\text{a}}(\mathbb{P}\mathcal{AT}/\sim)$  is Borel amenable (Theorem 12.15) and since  $\rho|_{\mathcal{H}}$  has trivial kernel, Proposition 3.38 implies that  $\mathcal{H}$  is amenable.  $\square$

## 12.9 An extension

We now establish a refined version of Theorem 12.1, that we will use in the sequel of the paper (precisely, in the proof of Lemma 13.30).

**Theorem 12.27.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\mathcal{N}_1, \dots, \mathcal{N}_p$  be measured subgroupoids of  $\mathcal{G}$ , such that for every  $i \in \{1, \dots, p-1\}$ , the subgroupoid  $\mathcal{N}_i$  is normalized by  $\mathcal{N}_{i+1}$ . Let  $\rho : \mathcal{G} \rightarrow \text{IA}_k(\mathbb{Z}/3\mathbb{Z})$  be a cocycle such that  $\rho|_{\mathcal{N}_i}$  is nowhere trivial for all  $i \in \{1, \dots, p\}$ . Assume that  $\mathcal{N}_1$  is amenable and that  $(\mathcal{N}_1, \rho)$  is pure, and let  $\hat{\mathcal{F}}$  be the unique maximal  $(\mathcal{N}_1, \rho)$ -invariant almost free factor system.*

*Then one of the following three assertions holds:*

1. *there exists a Borel subset  $U \subseteq Y$  of positive measure such that  $((\mathcal{N}_p)|_U, \rho)$  has a nice invariant splitting;*
2. *there exists a Borel subset  $U \subseteq Y$  of positive measure such that  $((\mathcal{N}_p)|_U, \rho)$  fixes an almost free factor system  $\hat{\mathcal{F}}' \not\preceq \hat{\mathcal{F}}$ ;*
3. *for every measured subgroupoid  $\mathcal{H}' \subseteq \mathcal{N}_p$ , if  $\rho|_{\mathcal{H}'}$  has trivial kernel, then  $\mathcal{H}'$  is amenable.*

*Proof.* In view of Proposition 10.6, and using the fact that the cocycles  $\rho|_{\mathcal{N}_i}$  are all nowhere trivial, the first conclusion holds as soon as there exist some  $i \in \{1, \dots, p\}$  and some Borel subset  $U \subseteq Y$  of positive measure for which there exists an  $((\mathcal{N}_i)|_U, \rho)$ -invariant nice splitting. We now assume otherwise. By Proposition 10.5, we can find a partition  $Y = \bigsqcup_{j \in J} Y_j$  into at most countably many Borel subsets such that for every  $i \in \{1, \dots, p\}$  and every  $j \in J$ , the pair  $((\mathcal{N}_i)|_{Y_j}, \rho)$  is pure, with maximum invariant almost free factor system  $\hat{\mathcal{F}}_{i,j}$ . As  $\mathcal{N}_i$  is normalized by  $\mathcal{N}_{i+1}$ , for every  $i \in \{1, \dots, p-1\}$  and every  $j \in J$ , the almost free factor system  $\hat{\mathcal{F}}_{i,j}$  is also  $((\mathcal{N}_{i+1})|_{Y_j}, \rho)$ -invariant, so  $\hat{\mathcal{F}}_{i,j} \preceq \hat{\mathcal{F}}_{i+1,j}$ . We can therefore assume that for every  $j \in J$ , one has  $\hat{\mathcal{F}}_{p,j} = \hat{\mathcal{F}}$ , as otherwise the second conclusion of the theorem holds and we are done.

Let  $\mathcal{F} \subseteq \hat{\mathcal{F}}$  be an extracted free factor system. Recall that purity implies that  $\mathcal{F}$  is non-sporadic (Remark 10.3). Moreover, as in Lemma 12.3, for every  $j \in J$  and every  $i \in \{1, \dots, p\}$ ,  $\mathcal{F}$  is an everywhere maximal  $((\mathcal{N}_i)_{|Y_j}, \rho)$ -invariant free factor system. In the sequel, we always work relative to  $\mathcal{F}$  (in particular, Outer spaces, proper free factors and arational trees are understood relative to  $\mathcal{F}$ ).

As  $\mathcal{N}_1$  is amenable, Corollary 12.13 ensures that there exists a Borel map  $\varphi : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  which is both  $(\mathcal{N}_1, \rho)$ -equivariant and  $(\mathcal{N}_2, \rho)$ -equivariant. Applying Proposition 12.14 iteratively  $p - 2$  times ensures that there exists a Borel map  $\varphi : Y \rightarrow \mathcal{P}_{\leq 2}(\mathbb{P}\mathcal{AT}/\sim)$  which is  $(\mathcal{N}_p, \rho)$ -equivariant.

We will finally use the partition  $\mathcal{P}_{< \infty}(\mathbb{P}\mathcal{AT}/\sim) = \mathcal{P}_{< \infty}^a(\mathbb{P}\mathcal{AT}/\sim) \sqcup \mathcal{P}_{< \infty}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$ . Assume first that there is a Borel subset  $U \subseteq Y$  such that  $\varphi(y) \in \mathcal{P}_{< \infty}^{\text{na}}(\mathbb{P}\mathcal{AT}/\sim)$  for all  $y \in U$ . Using Proposition 12.20, we deduce an  $((\mathcal{N}_p)_{|U}, \rho)$ -equivariant Borel map from  $U$  to the countable set of all nice splittings. We can then restrict to a Borel subset  $V \subseteq U$  where this map is constant to get an  $((\mathcal{N}_p)_{|V}, \rho)$ -invariant nice splitting, so the first conclusion of the theorem holds. We are thus left with the case where  $\varphi(y) \in \mathcal{P}_{< \infty}^a(\mathbb{P}\mathcal{AT}/\sim)$  for a.e.  $y \in Y$ . In this case, as the  $\text{Out}(F_k, \mathcal{F})$ -action on  $\mathcal{P}_{< \infty}^a(\mathbb{P}\mathcal{AT}/\sim)$  is amenable (Theorem 12.15), Proposition 3.38 ensures that a measured subgroupoid  $\mathcal{H}' \subseteq \mathcal{N}_p$  is amenable whenever  $\rho_{|\mathcal{H}'}$  has trivial kernel, so the third conclusion of the theorem holds.  $\square$

### 13 Detecting when a subgroupoid stabilizes a free splitting

The present section is of central importance in the proof of our main theorem; its goal is the following. Let  $\mathcal{G}$  be a measured groupoid coming with an action-type cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , with  $N \geq 3$ . Given a measured subgroupoid  $\mathcal{H}$  of  $\mathcal{G}$ , we want to relate the fact that  $(\mathcal{H}, \rho)$  preserves a non-separating free splitting of  $F_N$  (up to a partition of the base space of  $\mathcal{H}$  into at most countably many pieces), to some purely groupoid-theoretic properties of the pair  $(\mathcal{G}, \mathcal{H})$ , without reference to the cocycle  $\rho$ . This will be crucial in later sections to get a map at the level of (a variation of) the free splitting graph intertwining a pair of action-type cocycles  $\mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  as above.

To provide some intuition, let us mention that the analogous group-theoretic problem is to characterize subgroups of  $\text{Out}(F_N)$  that stabilize a non-separating free splitting  $S$  of  $F_N$  in a purely group-theoretic way. A crucial observation to obtain such a group-theoretic characterization is that the  $\text{Out}(F_N)$ -stabilizer of  $S$  contains a normal subgroup (namely, the group of twists about the splitting  $S$ ) which virtually splits as a direct product of two non-abelian free groups. See [HW20] for an implementation of this strategy to the computation of the abstract commensurator of  $\text{Out}(F_N)$  and some of its subgroups.

In all this section, we fix  $N \geq 3$ .

### 13.1 Schottky tuples and pseudo-products

**Definition 13.1** (Schottky tuples of subgroupoids). *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ . A Schottky tuple of subgroupoids of  $\mathcal{G}$  is a tuple  $(\mathcal{A}^1, \dots, \mathcal{A}^k)$  of amenable subgroupoids of  $\mathcal{G}$  of infinite type such that for every Borel subset  $U \subseteq Y$  of positive measure, the subgroupoid  $\langle \mathcal{A}_{|U}^1, \dots, \mathcal{A}_{|U}^k \rangle$  is non-amenable.*

*Remark 13.2.* This definition implies that  $\langle \mathcal{A}^1, \dots, \mathcal{A}^k \rangle$  is everywhere non-amenable. But this is not equivalent because in general,  $\langle \mathcal{A}_{|U}^1, \dots, \mathcal{A}_{|U}^k \rangle$  does not coincide with the restriction of  $\langle \mathcal{A}^1, \dots, \mathcal{A}^k \rangle$  to  $U$ . The point of this notion is that it is stable under restriction and stabilization. More precisely,

- If  $V \subseteq Y$  is a Borel subset of positive measure, and if  $(\mathcal{A}^1, \dots, \mathcal{A}^k)$  is a Schottky tuple of subgroupoids of  $\mathcal{G}$ , then  $(\mathcal{A}_{|V}^1, \dots, \mathcal{A}_{|V}^k)$  is a Schottky tuple of subgroupoids of  $\mathcal{G}_{|V}$ .
- If there exists a partition  $Y = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets, and if  $\mathcal{A}^1, \dots, \mathcal{A}^k$  are subgroupoids of  $\mathcal{G}$  such that for every  $i \in I$ , the tuple  $(\mathcal{A}_{|Y_i}^1, \dots, \mathcal{A}_{|Y_i}^k)$  is a Schottky tuple of subgroupoids of  $\mathcal{G}_{|Y_i}$ , then  $(\mathcal{A}^1, \dots, \mathcal{A}^k)$  is a Schottky tuple of subgroupoids of  $\mathcal{G}$ .

An important source of examples, which justifies the name *Schottky*, is given by the following lemma.

**Lemma 13.3** (Kida [Kid10, Lemma 3.20]). *Let  $\mathcal{G}$  be a measured groupoid, let  $\Gamma$  be a countable group, and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be an action-type cocycle. Let  $A_1$  and  $A_2$  be two infinite cyclic subgroups of  $\Gamma$  such that  $\langle A_1, A_2 \rangle$  is a non-abelian free group.*

*Then the groupoids  $\mathcal{A}_1 = \rho^{-1}(A_1)$  and  $\mathcal{A}_2 = \rho^{-1}(A_2)$  form a Schottky pair of subgroupoids of  $\mathcal{G}$ .*

We now introduce pseudo-products of subgroupoids. These give a counterpart for groupoids to the notion of a direct product  $G_1 \times \dots \times G_k$  of groups that all contain non-abelian free subgroups. Our definition in the groupoid setting translates the fact that each of the factors  $G_i$  of the direct product contains two infinite cyclic subgroups  $\langle a_i \rangle, \langle b_i \rangle$  such that  $\langle a_i, b_i \rangle \simeq F_2$  with  $\langle a_i \rangle$  and  $\langle b_i \rangle$  both normalized by all other factors  $G_j$ .

**Definition 13.4.** *Given a measured subgroupoid  $\mathcal{H}$  of  $\mathcal{G}$  and a Schottky tuple  $(\mathcal{A}^1, \dots, \mathcal{A}^k)$  of subgroupoids of  $\mathcal{G}$ , we say that  $(\mathcal{A}^1, \dots, \mathcal{A}^k)$  is stably normalized by  $\mathcal{H}$  if each  $\mathcal{A}^i$  is.*

**Definition 13.5** (Pseudo-products). *Let  $\mathcal{G}$  be a measured groupoid, let  $k \in \mathbb{N}$ , and let  $\mathcal{G}_1, \dots, \mathcal{G}_k$  be measured subgroupoids of  $\mathcal{G}$ .*

*The subgroupoids  $\mathcal{G}_1, \dots, \mathcal{G}_k$  form a pseudo-product if each  $\mathcal{G}_i$  contains a Schottky tuple of subgroupoids which is stably normalized by all  $\mathcal{G}_j$  with  $j \neq i$ .*

*Remark 13.6.* In view of Remark 13.2, this notion is stable under restriction and stabilization. More precisely, denoting by  $Y$  the base space of  $\mathcal{G}$ ,

- If  $U \subseteq Y$  is a Borel subset of positive measure, and if the subgroupoids  $\mathcal{G}_i$  form a pseudo-product, then so do the subgroupoids  $(\mathcal{G}_i)|_U$  of  $\mathcal{G}|_U$ .
- If the subgroupoids  $\mathcal{G}_i$  stably form a pseudo-product, then they form a pseudo-product.

A typical example, which justifies the terminology and will be of particular importance in the sequel, is the following.

**Lemma 13.7.** *Let  $\mathcal{G}$  be a measured groupoid, let  $\Gamma$  be a countable group, and let  $\rho : \mathcal{G} \rightarrow \Gamma$  be an action-type cocycle. Assume that  $\Gamma$  contains two subgroups  $G_1$  and  $G_2$  that generate their direct product in  $\Gamma$ , and such that both  $G_1$  and  $G_2$  contain a non-abelian free subgroup.*

*Then the groupoids  $\mathcal{G}_1 = \rho^{-1}(G_1)$  and  $\mathcal{G}_2 = \rho^{-1}(G_2)$  form a pseudo-product.*

*Proof.* For every  $i \in \{1, 2\}$ , let  $A_i^1$  and  $A_i^2$  be two infinite cyclic subgroups of  $G_i$  that generate a non-abelian free subgroup. By Lemma 13.3, for every  $i \in \{1, 2\}$ , the pair  $(\rho^{-1}(A_i^1), \rho^{-1}(A_i^2))$  is a Schottky pair of subgroupoids of  $\rho^{-1}(G_i)$ . In addition, since  $G_1$  and  $G_2$  form a direct product, the subgroupoids  $\rho^{-1}(A_i^1)$  and  $\rho^{-1}(A_i^2)$  are normalized by  $\rho^{-1}(G_j)$  for  $j \neq i$ .  $\square$

*Remark 13.8.* If  $\mathcal{G}_1, \dots, \mathcal{G}_k$  form a pseudo-product, then each  $\mathcal{G}_i$  is everywhere non-amenable since it contains a Schottky tuple of subgroupoids.

## 13.2 Main statement

Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ . We consider the following properties of the pair  $(\mathcal{G}, \mathcal{H})$ .

- ( $P_1$ ) The groupoid  $\mathcal{H}$  is everywhere non-amenable and stably normalizes two subgroupoids of  $\mathcal{G}$  that form a pseudo-product.
- ( $P_1^{\max}$ ) The pair  $(\mathcal{G}, \mathcal{H})$  satisfies ( $P_1$ ), and for every Borel subset  $U \subseteq Y$  of positive measure, if  $\mathcal{H}'$  is a subgroupoid of  $\mathcal{G}|_U$  such that  $(\mathcal{G}|_U, \mathcal{H}')$  satisfies ( $P_1$ ), with  $\mathcal{H}|_U$  stably contained in  $\mathcal{H}'$ , then  $\mathcal{H}|_U$  is stably equal to  $\mathcal{H}'$ .
- ( $P_2$ ) For every Borel subset  $U \subseteq Y$  of positive measure, the groupoid  $\mathcal{H}|_U$  does not normalize any amenable subgroupoid of  $\mathcal{G}|_U$  of infinite type.
- ( $P_3$ ) There do not exist a Borel subset  $U \subseteq Y$  of positive measure and two normal subgroupoids  $\mathcal{N}^-, \mathcal{N}^+$  of  $\mathcal{H}|_U$  which contain respectively three subgroupoids  $\mathcal{B}_1^-, \mathcal{B}_2^-, \mathcal{B}_3^-$  and  $\mathcal{B}_1^+, \mathcal{B}_2^+, \mathcal{B}_3^+$  such that for all  $\varepsilon \in \{\pm\}$ , the groupoids  $\mathcal{B}_1^\varepsilon, \mathcal{B}_2^\varepsilon, \mathcal{B}_3^\varepsilon, \mathcal{N}^{-\varepsilon}$  form a pseudo-product.

*Remark 13.9.* Notice that the assumptions ( $P_1$ ), ( $P_1^{\max}$ ), ( $P_2$ ), and ( $P_3$ ) are preserved under restriction and stabilization. More precisely,



1. If  $(\mathcal{G}, \mathcal{H})$  satisfies any of these properties, and  $U \subseteq Y$  is a Borel subset of positive measure, then  $(\mathcal{G}|_U, \mathcal{H}|_U)$  satisfies the property.
2. If there exists a partition  $Y = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets, such that for every  $i \in I$ , the pair  $(\mathcal{G}|_{Y_i}, \mathcal{H}|_{Y_i})$  satisfies one of these properties, then  $(\mathcal{G}, \mathcal{H})$  satisfies the corresponding property.

Our goal in the sequel of the present section is to consider an action-type cocycle  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  and to relate the above purely groupoid-theoretic properties of the pair  $(\mathcal{G}, \mathcal{H})$  to the fact that  $(\mathcal{H}, \rho)$  stabilizes a free splitting of  $F_N$ .

A *one-edge non-separating free splitting* of  $F_N$  is a free splitting of  $F_N$  equal to the Bass–Serre tree of an HNN extension of the form  $F_N = P*$ , with  $P$  a subgroup of  $F_N$  of rank  $N - 1$ . We denote by  $\text{FS}^{\text{ns}}$  the countable collection of all one-edge non-separating free splittings. We say that a free splitting of  $F_N$  is a  $(2, 2^+)$ -free splitting if it is a free splitting of  $F_N$  equal to the Bass–Serre tree of a decomposition of the form  $F_N = P * P'$  with  $\text{rank}(P) = 2$  and  $\text{rank}(P') \geq 2$ . We denote by  $\text{FS}^{2,2^+}$  the countable set of all  $(2, 2^+)$ -free splittings of  $F_N$ . Note that when  $N = 3$ ,  $\text{FS}^{2,2^+} = \emptyset$ .

The goal of the present section is to prove the following theorem.

**Theorem 13.10.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle with  $N \geq 3$ . Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ .*

1. *If  $(\mathcal{G}, \mathcal{H})$  satisfies Properties  $(P_1^{\text{max}})$ ,  $(P_2)$  and  $(P_3)$ , then there exists a conull Borel subset  $Y^* \subseteq Y$  and a partition  $Y^* = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets, such that for every  $i \in I$ , the groupoid  $\mathcal{H}|_{Y_i}$  is equal to the  $(\mathcal{G}|_{Y_i}, \rho)$ -stabilizer of a splitting  $S_i \in \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$ , i.e.  $\mathcal{H}|_{Y_i} = \rho^{-1}(\text{Stab}_{\text{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S_i))|_{Y_i}$ .*
2. *If there exist a conull Borel subset  $Y^* \subseteq Y$  and a partition  $Y^* = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets, such that for every  $i \in I$ , the groupoid  $\mathcal{H}|_{Y_i}$  is equal to the  $(\mathcal{G}|_{Y_i}, \rho)$ -stabilizer of some splitting  $S_i \in \text{FS}^{\text{ns}}$ , then  $(\mathcal{G}, \mathcal{H})$  satisfies Properties  $(P_1^{\text{max}})$ ,  $(P_2)$  and  $(P_3)$ .*

*Proof.* The first part is proved in Section 13.3 below, and the second is proved in Section 13.4 (Lemmas 13.26, 13.27 and 13.30).  $\square$

*Remark 13.11.* We do not know however whether stabilizers of splittings  $\text{FS}^{2,2^+}$  satisfy Property  $(P_3)$  or not, whence the slight asymmetry in the statement of Theorem 13.10.

### 13.3 A sufficient condition to stabilize a free splitting

Our goal in this section is to prove the first part of Theorem 13.10.

### 13.3.1 Step 1: Getting an invariant nice splitting from Property $(P_1)$

**Lemma 13.12.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be a cocycle with trivial kernel. Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ . Assume that  $(\mathcal{G}, \mathcal{H})$  satisfies Property  $(P_1)$ .*

*Then  $\mathcal{H}$  is stably nice.*

*Proof.* By assumption  $\mathcal{H}$  stably normalizes two subgroupoids  $\mathcal{N}^-, \mathcal{N}^+$  of  $\mathcal{G}$  which form a pseudo-product. In particular,  $\mathcal{H}$  stably normalizes the subgroupoid  $\mathcal{N}^-$ , which is everywhere non-amenable (Remark 13.8), and  $\mathcal{N}^-$  stably normalizes an amenable subgroupoid  $\mathcal{A}^+$  of  $\mathcal{N}^+$  of infinite type. Applying Lemma 10.7 to  $\mathcal{H}$ , consider a Borel partition  $Y = Y_1 \sqcup Y_2$  with  $(\mathcal{H}|_{Y_1}, \rho)$  stably nice and  $(\mathcal{H}|_{Y_2}, \rho)$  nice-averse. We assume that  $Y_2$  has positive measure and argue towards a contradiction. Because the property of stably preserving a nice splitting goes to the normalizer (Proposition 10.6),  $(\mathcal{A}^+_{|Y_2}, \rho)$  and  $(\mathcal{N}^-_{|Y_2}, \rho)$  are nice-averse. As  $\mathcal{A}^+_{|Y_2}$  is of infinite type and amenable, and  $\rho$  has trivial kernel, Corollary 12.2 shows that  $\mathcal{N}^-_{|Y_2}$  is amenable, a contradiction.  $\square$

### 13.3.2 Step 2: Excluding $\mathcal{Z}_{\max}$ -splittings thanks to Property $(P_2)$

**Lemma 13.13.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle. Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ .*

*If there exists an  $(\mathcal{H}, \rho)$ -invariant  $\mathcal{Z}_{\max}$ -splitting, then  $\mathcal{H}$  normalizes an amenable subgroupoid of  $\mathcal{G}$  of infinite type.*

*Proof.* Let  $S$  be a  $\mathcal{Z}_{\max}$ -splitting which is  $(\mathcal{H}, \rho)$ -invariant. Then there exists a conull Borel subset  $Y^* \subseteq Y$  such that  $\mathcal{H}|_{Y^*} \subseteq \rho^{-1}(\mathrm{Stab}_{\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S))$ . Let  $A \subseteq \mathrm{Stab}_{\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$  be the group of twists of the splitting  $S$ , an infinite amenable normal subgroup of  $\mathrm{Stab}_{\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$ , and let  $\mathcal{A} = \rho^{-1}(A)$ : this is an amenable subgroupoid of  $\mathcal{G}$ , of infinite type because  $\rho$  is action-type. In addition, as  $A$  is normal in  $\mathrm{Stab}_{\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$ , it follows that  $\mathcal{A}|_{Y^*}$  is normal in  $\rho^{-1}_{|Y^*}(\mathrm{Stab}_{\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S))$  (Lemma 3.32), and therefore  $\mathcal{A}$  is normalized by  $\mathcal{H}$ .  $\square$

Combining Lemmas 13.12 and 13.13, we reach the following statement.

**Corollary 13.14.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle. Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ . Assume that  $(\mathcal{G}, \mathcal{H})$  satisfies Properties  $(P_1)$  and  $(P_2)$ .*

*Then there exists a partition  $Y^* = \bigsqcup_{i \in I} Y_i$  of a conull Borel subset  $Y^* \subseteq Y$  into at most countably many Borel subsets such that for every  $i \in I$ , there exists a non-trivial free splitting or bi-nonsporadic splitting which is  $(\mathcal{H}, \rho)$ -invariant.  $\square$*

### 13.3.3 Step 3: Excluding bi-nonsporadic splittings via $(P_1^{\max})$ and $(P_3)$

**Lemma 13.15.** *Let  $S$  be a bi-nonsporadic splitting of  $F_N$ , and let  $H := \mathrm{Stab}_{\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$ .*

*Then  $H$  contains a direct product  $N^- \times N^+$ , with  $N^-, N^+$  normal in  $H$ , and such that for every  $\varepsilon \in \{\pm\}$ , the group  $N^\varepsilon$  contains a subgroup isomorphic to  $F_2 \times F_2 \times F_2$ .*

*Proof.* As  $S$  is bi-nonsporadic, there exist two vertices  $v^-, v^+ \in S$  in distinct  $F_N$ -orbits such that for every  $\varepsilon \in \{\pm\}$ , the Grushko decomposition of  $G_{v^\varepsilon}$  relative to the set  $\text{Inc}_{v^\varepsilon}$  of incident edge groups is non-sporadic.

We claim that the group  $\text{Out}(G_{v^\varepsilon}, \text{Inc}_{v^\varepsilon}^{(t)})$  made of all automorphisms of  $G_{v^\varepsilon}$  acting by conjugation on every incident edge group contains a copy of  $F_2 \times F_2 \times F_2$ . Indeed, let  $S_{v^\varepsilon}$  be a Grushko splitting of  $G_{v^\varepsilon}$  relative to  $\text{Inc}_{v^\varepsilon}$  with all vertex stabilizers non-trivial (hence non-abelian because incident edge groups are non-abelian). By non-sporadicity, the splitting  $S_{v^\varepsilon}$  contains at least four half-edges in distinct orbits based at vertices with non-trivial stabilizer. Therefore the group of twists associated to three of these half-edges is isomorphic to a direct product of three non-abelian free groups, which proves our claim.

Now by Lemma 2.9, the product  $\text{Out}(G_{v^+}, \text{Inc}_{v^+}^{(t)}) \times \text{Out}(G_{v^-}, \text{Inc}_{v^-}^{(t)})$  embeds in  $\text{Stab}_{\text{Out}(F_N)}(S)$ . We then denote by  $H$  the stabilizer of  $S$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , and by  $N^\varepsilon$  the intersection of  $H$  with  $\text{Out}(G_{v^\varepsilon}, \text{Inc}_{v^\varepsilon}^{(t)})$ . Corollary 2.14 ensures that  $N^+$  and  $N^-$  are normal in  $H$ , which concludes the proof.  $\square$

**Corollary 13.16.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle. Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$ . Assume that  $(\mathcal{G}, \mathcal{H})$  satisfies Properties  $(P_1^{\max})$ ,  $(P_2)$ , and  $(P_3)$ .*

*Then  $(\mathcal{H}, \rho)$  stably preserves a free splitting.*

*Proof.* We argue by contradiction, so assume otherwise. Then by Corollary 13.14, up to restricting to a subset of positive measure we may assume that  $(\mathcal{H}, \rho)$  preserves a bi-nonsporadic splitting  $S$ , and let  $\Gamma_S \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be its stabilizer. Let  $N^+, N^-$  be the normal subgroups of  $\Gamma_S$  be given by Lemma 13.15.

We claim that the groupoid  $\hat{\mathcal{H}} = \rho^{-1}(\Gamma_S)$  satisfies  $(P_1)$ . Indeed, the subgroupoids  $\mathcal{N}^+ = \rho^{-1}(N^+)$  and  $\mathcal{N}^- = \rho^{-1}(N^-)$  form a pseudo-product by Lemma 13.7 and are normalized by  $\hat{\mathcal{H}}$ , which proves the claim.

Since  $(\mathcal{G}, \mathcal{H})$  satisfies Property  $(P_1^{\max})$  it follows that  $\mathcal{H}$  is stably equal to  $\hat{\mathcal{H}}$ , so up to further restricting to a Borel subset of positive measure, we may assume that  $\mathcal{H} = \rho^{-1}(\Gamma_S)$ . For each  $\varepsilon \in \{\pm\}$ , denote by  $B_1^\varepsilon \times B_2^\varepsilon \times B_3^\varepsilon \simeq F_2 \times F_2 \times F_2$  the subgroup of  $N^\varepsilon$  given by Lemma 13.15. For each  $\varepsilon \in \{\pm\}$ , and each  $j \leq 3$ , consider  $\mathcal{B}_j^\varepsilon = \rho^{-1}(B_j^\varepsilon)$ . Then by Lemma 13.7,  $(\mathcal{N}^+, \mathcal{B}_1^-, \mathcal{B}_2^-, \mathcal{B}_3^-)$  form a pseudo-product, and so do  $(\mathcal{N}^-, \mathcal{B}_1^+, \mathcal{B}_2^+, \mathcal{B}_3^+)$ . This contradicts Property  $(P_3)$ .  $\square$

### 13.3.4 Step 4: Excluding some types of free splittings

We say that a free splitting  $S$  of  $F_N$  is *acyclic* if  $S/F_N$  does not contain any terminal vertex with vertex group isomorphic to  $\mathbb{Z}$ . When  $S$  is a one-edge splitting, this is equivalent to saying that  $S$  is not the Bass–Serre tree of a decomposition of  $F_N$  of the form  $F_N = \mathbb{Z} * F_{N-1}$ . Note that the stabilizer in  $\text{Out}(F_N)$  of the splitting  $F_N = \mathbb{Z} * F_{N-1}$  is contained in the stabilizer of the acyclic splitting  $F_N = F_{N-1} *$ . This observation leads to the following result.

**Lemma 13.17.** *Let  $S$  be a free splitting of  $F_N$ , and let  $H$  be its stabilizer in  $\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Then  $H$  stabilizes an acyclic one-edge free splitting of  $F_N$ .*

*Proof.* As  $H \subseteq \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , it stabilizes every one-edge collapse of  $S$ . In addition, if  $H$  stabilizes a splitting of the form  $F_N = A * B$  with  $B$  cyclic, then it also stabilizes the acyclic splitting  $F_N = A*$ .  $\square$

In order to be able to use Property  $(P_1^{\max})$ , we now prove that the stabilizer of a free splitting satisfies Property  $(P_1)$ . This will also be useful later for proving the second part of Theorem 13.10.

**Lemma 13.18.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle. Let  $S$  be a one-edge acyclic free splitting of  $F_N$ . Then the  $(\mathcal{G}, \rho)$ -stabilizer of  $S$  satisfies Property  $(P_1)$ .*

*Proof.* We denote by  $H$  the stabilizer of  $S$  in  $\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , and we let  $\mathcal{H} = \rho^{-1}(H)$ , so that  $\mathcal{H}$  is the  $(\mathcal{G}, \rho)$ -stabilizer of  $S$ .

We first assume that  $S$  is a separating free splitting, equal to a Bass–Serre tree of a free product decomposition  $F_N = A * B$ , with both  $A$  and  $B$  noncyclic. The group of twists of  $S$  in  $\mathrm{Out}(F_N)$  is isomorphic to the direct product  $A \times B$  (see Section 2.8). We then let  $N^- := (A \times \{1\}) \cap \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  and  $N^+ := (\{1\} \times B) \cap \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Since  $H \subseteq \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , both  $N^-$  and  $N^+$  are normal subgroups of  $H$ . For every  $\varepsilon \in \{\pm\}$ , we then let  $\mathcal{N}^\varepsilon := \rho^{-1}(N^\varepsilon)$ , a normal subgroupoid of  $\mathcal{H}$  by Lemma 3.32. In addition, as  $N^+$  and  $N^-$  generate their direct product in  $H$  and both contain a non-abelian free subgroup, Lemma 13.7 ensures that  $\mathcal{N}^-$  and  $\mathcal{N}^+$  form a pseudo-product. This proves that  $(\mathcal{H}, \rho)$  satisfies  $(P_1)$  when  $S$  is separating.

We now assume that  $S$  is a non-separating free splitting, equal to a Bass–Serre tree of a decomposition  $F_N = A*$  (where  $A$  is a corank one free factor of  $F_N$ ). The group of twists of  $S$  in  $\mathrm{Out}(F_N)$  is isomorphic to a direct product  $A^- \times A^+$  of two copies of  $A$  (see Section 2.8). We let  $N^- := (A^- \times \{1\}) \cap \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$  and  $N^+ := (\{1\} \times A^+) \cap \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Since  $H \subseteq \mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , both  $N^-$  and  $N^+$  are normal subgroups of  $H$ . For every  $\varepsilon \in \{\pm\}$ , we then let  $\mathcal{N}^\varepsilon := \rho^{-1}(N^\varepsilon)$ . As above,  $\mathcal{N}^-$  and  $\mathcal{N}^+$  are normalized by  $\mathcal{H}$  and form a pseudo-product.  $\square$

Property  $(P_3)$  will allow us to rule out one-edge acyclic free splittings not in  $\mathrm{FS}^{\mathrm{ns}} \cup \mathrm{FS}^{2,2^+}$  thanks to the following lemma.

**Lemma 13.19.** *Let  $S$  be a separating free splitting of  $F_N$  corresponding to a decomposition of the form  $F_N = F_k * F_{N-k}$  with  $k, N - k \geq 3$ , and let  $H := \mathrm{Stab}_{\mathrm{IA}_N(\mathbb{Z}/3\mathbb{Z})}(S)$ .*

*Then  $H$  contains a direct product  $N^- \times N^+$ , with  $N^-, N^+$  normal in  $H$ , and such that for every  $\varepsilon \in \{\pm\}$ , the group  $N^\varepsilon$  contains a subgroup isomorphic to  $F_2 \times F_2 \times F_2$ .*

*Proof.* The group  $\mathrm{Stab}_{\mathrm{Out}(F_N)}(S)$  has a finite-index subgroup isomorphic to  $\mathrm{Aut}(F_k) \times \mathrm{Aut}(F_{N-k})$ . The intersections of  $H$  with each of the factors  $\mathrm{Aut}(F_k)$  and  $\mathrm{Aut}(F_{N-k})$  are normal in  $H$  and have finite index in  $\mathrm{Aut}(F_k)$  and  $\mathrm{Aut}(F_{N-k})$ . The conclusion thus follows from the fact that for every  $k \geq 3$ , the group  $\mathrm{Aut}(F_k)$  contains a subgroup isomorphic to  $F_2 \times F_2 \times F_2$ , see e.g. [HW20, Theorem 6.2].  $\square$

We are now ready to complete the proof of the first part of Theorem 13.10.

*Proof of Theorem 13.10, Part 1.* Assume that  $(\mathcal{G}, \mathcal{H})$  satisfies Properties  $(P_1^{\max})$ ,  $(P_2)$  and  $(P_3)$ . By Corollary 13.16,  $\mathcal{H}$  stably preserves a free splitting: there is a countable Borel partition  $Y^* = \bigsqcup_{i \in I} Y_i$  of a conull subset  $Y^* \subseteq Y$  into subsets of positive measure such that  $\rho(\mathcal{H}_{|Y_i}) \subseteq \Gamma_{S_i}$  where  $S_i$  is a free splitting and  $\Gamma_{S_i}$  is its stabilizer in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . By Lemma 13.17, up to increasing  $\Gamma_{S_i}$ , we may assume that  $S_i$  is a one-edge acyclic free splitting. Since the subgroupoid  $\mathcal{H}' = \rho^{-1}(\Gamma_{S_i})_{|Y_i}$  contains  $\mathcal{H}_{|Y_i}$  and satisfies Property  $(P_1)$  by Lemma 13.18, Assumption  $(P_1^{\max})$  on  $\mathcal{H}$  ensures that, up to refining our partition,  $\mathcal{H}_{|Y_i} = \rho^{-1}(\Gamma_{S_i})_{|Y_i}$ .

We assume that  $S_i \notin \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$  and argue towards a contradiction. Then Lemma 13.19 applies to  $S_i$  and yields two normal subgroups  $N^+, N^- \trianglelefteq \Gamma_{S_i}$  which generate their direct product, and such that for every  $\varepsilon \in \{\pm\}$ , the group  $N^\varepsilon$  contains a direct product  $B_1^\varepsilon \times B_2^\varepsilon \times B_3^\varepsilon$  of three non-abelian free groups. The subgroupoids  $\mathcal{N}^+ = \rho^{-1}(N^+)_{|Y_i}$  and  $\mathcal{N}^- = \rho^{-1}(N^-)_{|Y_i}$  are normal subgroupoids of  $\mathcal{H}_{|Y_i}$ . For all  $\varepsilon \in \{\pm\}$  and all  $j \in \{1, 2, 3\}$ , we let  $\mathcal{B}_j^\varepsilon = \rho^{-1}(B_j^\varepsilon)_{|Y_i}$ . Then by Lemma 13.7,  $(\mathcal{N}^+, \mathcal{B}_1^-, \mathcal{B}_2^-, \mathcal{B}_3^-)$  form a pseudo-product, and so do  $(\mathcal{N}^-, \mathcal{B}_1^+, \mathcal{B}_2^+, \mathcal{B}_3^+)$ . This contradicts that  $\mathcal{H}$  satisfies Property  $(P_3)$ .  $\square$

### 13.4 A necessary condition to stabilize a non-separating free splitting

The goal of the present section is to prove the second part of Theorem 13.10.

Since Properties  $(P_1)$ ,  $(P_1^{\max})$ ,  $(P_2)$  and  $(P_3)$  are preserved under stabilization, it is enough to check that the  $(\mathcal{G}, \rho)$ -stabilizer  $\mathcal{H}$  of any non-separating free splitting satisfies Properties  $(P_1)$ ,  $(P_1^{\max})$ ,  $(P_2)$  and  $(P_3)$ . We already checked that  $\mathcal{H}$  satisfies Property  $(P_1)$  in Lemma 13.18.

In the rest of this subsection, we fix a measured groupoid  $\mathcal{G}$  over a base space  $Y$ , and  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle (with  $N \geq 3$ ). Let  $S_0$  be a one-edge non-separating free splitting of  $F_N$ , let  $H_0 \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be the stabilizer of  $S_0$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , and let  $\mathcal{H}_0 := \rho^{-1}(H_0)$ . We let  $A_0$  be a free factor of corank 1 stabilizing a vertex of  $S_0$  so that  $S_0$  is dual to the HNN extension  $F_N = A_0*$ . We denote by  $v_0$  the vertex of  $S_0$  fixed by  $A_0$ , choose an edge  $e_0$  incident on  $v_0$ , and consider  $t \in F_N$  such that  $e_0 = [v_0, tv_0]$ , so that  $t$  is a stable letter of the HNN extension. For each  $w, w' \in A_0$ , consider the automorphism  $\tilde{\tau}_{w,w'} \in \text{Aut}(F_N)$  that acts as the identity on  $A_0$  and sends  $t$  to  $wtw'$ : this is the lift to  $\text{Aut}(F_N)$  of an element of the group of twists of  $S_0$ , in particular it stabilizes  $S_0$ . The associated  $\tilde{\tau}_{w,w'}$ -equivariant isometry  $I_{\tilde{\tau}_{w,w'}}$  of  $S_0$  maps  $e_0$  to  $we_0$  and  $t^{-1}e_0$  to  $w'^{-1}t^{-1}e_0$ .

Finally, we let  $\rho_{A_0} : \mathcal{H}_0 \rightarrow \text{Out}(A_0)$  be the cocycle obtained by postcomposing  $\rho$  with the natural map  $\text{Stab}(S_0) \rightarrow \text{Out}(A_0)$ .

#### 13.4.1 Proof of Property $(P_1^{\max})$

**Lemma 13.20.** *Consider an element  $a \in F_N$  whose conjugacy class is  $H_0$ -invariant. Then  $a$  is conjugate in  $A_0$ .*

*Remark 13.21.* This uses the fact that  $N \geq 3$ . When  $N > 3$  (so that  $A_0$  has rank at least 3),  $a$  has in fact to be trivial. When  $N = 3$  (so that  $A_0$  has rank 2), one may have  $a = [x, y]^k$  for some free basis  $\{x, y\}$  of  $A_0$  and some  $k \in \mathbb{Z}$ .

*Proof.* Let  $a \in F_N$  be an element which is not conjugate in  $A_0$ ; we are going to prove that the conjugacy class of  $a$  is not  $H_0$ -invariant.

Given a non-trivial element  $w \in A_0$  which is not a proper power, consider the  $\mathcal{Z}_{\max}$ -splitting  $S_w$  of  $F_N$  dual to the HNN extension with vertex group  $\langle A_0, twt^{-1} \rangle$  and with edge group  $\langle w \rangle$  with the two obvious inclusions. The splitting  $S_w$  may be obtained from  $S_0$  as follows: recall that  $v_0 \in S_0$  is the vertex fixed by  $A_0$ , and  $e_0 = [v_0, tv_0]$ ; then  $S_w$  is obtained by folding  $e_0$  with  $we_0$ .

As  $a$  is not conjugate in  $A_0$ , it is hyperbolic in  $S_0$ . For all but finitely many choices of  $w$ , the axis of  $a$  in  $S_0$  isometrically embeds in  $S_w$  under the folding map. It follows that we may choose  $w$  so that  $a$  is hyperbolic in  $S_w$ . Let  $\tau$  be the twist associated to the splitting  $S_w$ : this is the automorphism restricting to the identity on  $A_0$  and sending  $t$  to  $tw$ , so in particular  $\tau$  has a power in  $H_0$ .

Let  $R$  be a Cayley graph of  $F_N$  with respect to some basis, equipped with the simplicial metric where every edge is assigned length 1. By [CL95], the rescaled trees  $\frac{1}{n}(\tau^n.R)$  converge to a tree homothetic to  $S_w$  as  $n \rightarrow \infty$ . It follows that the translation length  $\|\tau^n(a)\|_R$  tends to infinity, so the conjugacy class of  $a$  is not  $H_0$ -invariant.  $\square$

**Lemma 13.22.** *Let  $B$  be any proper free factor of  $F_N$  whose conjugacy class is  $H_0$ -invariant.*

*Then  $B$  is conjugate to  $A_0$ .*

*Proof.* Observe that since  $H_0$  surjects onto  $\text{IA}(A_0, \mathbb{Z}/3\mathbb{Z})$ , there is no  $H_0$ -invariant proper free factor of  $F_N$  strictly contained in  $A_0$ . We can therefore assume that  $B$  is not conjugate into  $A_0$ , and aim for a contradiction.

We first claim that  $B$  intersects every conjugate of  $A_0$  trivially, equivalently that  $A_0$  intersects every conjugate of  $B$  trivially. Indeed, for every  $g \in F_N$ , the subgroup  $A_0 \cap B^g$  is a free factor contained in  $A_0$  whose conjugacy class is  $H_0$ -invariant (Lemma 8.23). Thus,  $A_0 \cap B^g$  is either trivial or equal to  $A_0$ , but the latter case implies  $B^g = A_0$  since  $A_0$  has corank 1.

We next claim that  $H_0$  acts trivially on  $B$ . Let  $S_B \subseteq S_0$  be the minimal  $B$ -invariant subtree. As  $B$  intersects every conjugate of  $A_0$  trivially, it acts freely on  $S_B$ , so  $S_B$  defines a point in the Outer space of  $B$ , and its stabilizer in  $\text{IA}(B, \mathbb{Z}/3\mathbb{Z})$  is trivial. For all  $\alpha \in H_0$  and all  $b \in B$ , one has  $\|\alpha(b)\|_{S_B} = \|\alpha(b)\|_{S_0} = \|b\|_{S_0} = \|b\|_{S_B}$ , so  $S_B$  is invariant by the image of  $H_0$  in  $\text{IA}(B, \mathbb{Z}/3\mathbb{Z})$  (indeed every tree in the Outer space of  $B$  is determined by its length function by [CM87, Theorem 3.7]). The image of  $H_0$  in  $\text{Out}(B)$  is therefore trivial, which proves our claim.

Together with Lemma 13.20, the claim implies that every element of  $B$  is conjugate into  $A_0$ , and therefore  $B$  is conjugate into  $A_0$ , a contradiction.  $\square$

Recall that an outer automorphism  $\alpha \in \text{Out}(A_0)$  is *fully irreducible* if no non-trivial power of  $\alpha$  fixes the conjugacy class of a proper free factor of  $A_0$ .



**Lemma 13.23.** *For every fully irreducible element  $\alpha \in \text{Out}(A_0)$ , the following two assertions hold:*

1. *no non-trivial power of  $\alpha$  preserves the conjugacy class a non-cyclic finitely generated subgroup of  $A_0$  of infinite index;*
2. *no non-trivial power of  $\alpha$  preserves a non-trivial splitting of  $A_0$  with finitely generated edge or vertex stabilizers.*


*Proof.* For the first assertion, observe that the attracting  $\mathbb{R}$ -tree  $T$  of  $\alpha$  is indecomposable by [CH12, Theorem 2.1], so by [Rey11, Theorem 4.5], any finitely generated subgroup  $B \subseteq A_0$  of infinite index acts discretely on  $T$ . Thus, if the conjugacy class of  $B$  is invariant under a power of  $\alpha$ , then  $B$  has to be elliptic in  $T$ . As  $T$  is arational [Rey12], it follows that  $B$  is cyclic [Rey12, Theorem 1.1].

We now prove that  $\alpha$  satisfies the second conclusion of the lemma. Recall that a splitting with finitely generated edge stabilizers also has finitely generated vertex stabilizers. So let  $S$  be a splitting of  $A_0$  with finitely generated vertex stabilizers which is invariant under a power of  $\alpha$ . Then the vertex and edge stabilizers have to be cyclic. We claim that some edge stabilizer has to be trivial. Otherwise, any two edges incident on the same vertex have commuting stabilizers, so by commutative transitivity, all edge stabilizers of  $S$  commute, a contradiction. We may thus assume that  $S$  is a free splitting. Since there is no free factor invariant by a power of  $\alpha$ , this concludes the proof.  $\square$

**Lemma 13.24.** *Let  $S'$  be a nice splitting such that every element of  $H_0$  has a power that preserves  $S'$ .*

*Then  $S' = S_0$ .*

*Proof.* Assume first that  $S'$  is a free splitting. By assumption,  $S'$  is periodic under the action of every element of  $H_0$ . As  $H_0 \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , Lemma 2.12 implies that  $S'$  is in fact  $H_0$ -invariant, and so is every collapse of  $S'$ . By Lemma 13.22, point stabilizers of  $S'$  are either trivial or conjugate to  $A_0$ , and the same is true for collapses of  $S'$ . If the action of  $F_k$  on  $S'$  was free, then one could construct some collapse of  $S'$  with a cyclic vertex stabilizer, a contradiction. It follows that  $A_0$  fixes a point in  $S'$ , so either  $S' = S_0$  or  $S'$  is dual to a graph of groups of the form



in which case  $S'$  has a collapse with a cyclic vertex stabilizer, a contradiction.

We now assume that  $S'$  is a  $\mathcal{Z}_{\max}$ -splitting of  $F_N$ . Theorem 2.11 ensures that the the conjugacy class of any edge group of  $S'$  is  $H_0$ -invariant so if  $N > 3$ , we may use Lemma 13.20 and Remark 13.21 to exclude the existence of  $S'$ . The following argument works for all  $N \geq 3$ .

Let  $H'$  be the stabilizer of  $S'$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . The action of  $A_0$  on its minimal subtree  $S'_{A_0} \subseteq S'$  is a splitting of  $A_0$  with finitely generated edge stabilizers, which is invariant under the image  $H'_{A_0}$  of  $H'$  in  $\text{Out}(A_0)$ . As every element of  $H_0$  has a power contained in



$H'$ , it follows that every element of  $\text{Out}(A_0)$  has a power contained in  $H'_{A_0}$ , in particular  $H'_{A_0}$  contains a fully irreducible outer automorphism of  $A_0$ . By Lemma 13.23, it follows that  $A_0$  fixes a (unique) point in  $S'$ .

Let  $f : S_0 \rightarrow S'$  be the unique equivariant map that is linear on edges. Being unique,  $f$  is invariant under  $H'$ . Since  $f$  is not an isomorphism, there exists a pair of adjacent edges  $e, e'$  in  $S$  such that  $f$  identifies an initial segment of  $e$  with an initial segment of  $e'$ . Without loss of generality, the common endpoint of  $e$  and  $e'$  is the vertex  $v_0 \in S$  fixed by  $A_0$ .

Assume first that  $e' = ae$  for some  $a \in A_0 \setminus \{1\}$ . Let  $w \in A_0$  be an element that does not commute with  $a$ . We use the fact that there exists a twist  $\alpha \in \text{Out}(F_N)$  of  $S_0$  which has a lift  $\tilde{\alpha} \in \text{Aut}(F_N)$  acting as the identity on  $A_0$  and such that  $I_{\tilde{\alpha}}(e) = we$  and  $I_{\tilde{\alpha}}(ae) = awe$ . Take  $n \geq 1$  such that  $\alpha^n \in H'$ . Since  $\tilde{\alpha}$  acts as the identity on  $A_0$ ,  $I_{\tilde{\alpha}^n}(e) = w^n e$  and  $I_{\tilde{\alpha}^n}(e') = aw^n e$ . It follows that  $e$  and  $w^{-n}aw^n e$  have initial segments that are identified by  $f$ . Thus, the non-abelian group  $\langle a, w^{-n}aw^n \rangle$  fixes a segment in  $S'$ , a contradiction.

Now assume that  $e, e'$  are not in the same  $A_0$ -orbit. Choose  $a \in A_0 \setminus \{1\}$ . There exists a twist  $\alpha \in \text{Stab}(S)$  and a preimage  $\tilde{\alpha} \in \text{Aut}(F_N)$  acting as the identity on  $A_0$  and such that the isometry  $I_{\tilde{\alpha}}$  of  $S$  fixes  $e'$  and sends  $e$  to  $ae$ . In particular, if  $n \geq 1$  is such that  $\alpha^n \in H'$ , we get that  $f$  identifies some initial segments of  $e$  and  $a^n e$ . The previous case then yields a contradiction.

We finally assume that  $S'$  is a bi-nonsporadic splitting. By Lemma 9.6 there is a finite collection  $\{\mathcal{F}_1, \dots, \mathcal{F}_n\}$  of  $n \geq 2$  non-empty free factor systems of  $F_N$  such that every element of  $H_0$  has a power that preserves each  $\mathcal{F}_i$ . Since  $H_0 \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , each  $\mathcal{F}_i$  is in fact  $H_0$ -invariant. This contradicts Lemma 13.22.  $\square$

**Lemma 13.25.** *For every Borel subset  $U \subseteq Y$  of positive measure,  $S_0$  is the only nice splitting of  $F_N$  which is  $(\mathcal{H}_{0|U}, \rho)$ -invariant.*

*Moreover, there is no non-trivial splitting of  $A_0$  with finitely generated edge stabilizers which is  $(\mathcal{H}_{0|U}, \rho_{A_0})$ -invariant.*

*Proof.* Let  $S$  be an  $(\mathcal{H}_{0|U}, \rho)$ -invariant nice splitting. Then there exists a conull Borel subset  $U^* \subseteq U$  such that  $S$  is invariant by  $\rho(\mathcal{H}_{0|U^*})$ . Since  $\mathcal{H}_0 = \rho^{-1}(H_0)$ , Remark 3.23 implies that every element  $\alpha \in H_0$  has a power that preserves  $S$ . Lemmas 13.24 and 13.23 thus conclude the proof (to apply Lemma 13.23, notice indeed that  $H_0$  contains an element that restricts to a fully irreducible outer automorphism of  $A_0$ ).  $\square$

**Lemma 13.26.** *The pair  $(\mathcal{G}, \mathcal{H}_0)$  satisfies Property  $(P_1^{\max})$ .*

*Proof.* By Lemma 13.18, the pair  $(\mathcal{G}, \mathcal{H}_0)$  satisfies Property  $(P_1)$ , so we only need to check maximality. Let  $U \subseteq Y$  be a Borel subset of positive measure, and let  $\mathcal{H}'$  be a subgroupoid of  $\mathcal{G}|_U$  such that  $(\mathcal{G}|_U, \mathcal{H}')$  satisfies Property  $(P_1)$ , and such that  $\mathcal{H}_{0|U}$  is stably contained in  $\mathcal{H}'$ . By Lemma 13.12, there exists a partition  $U = \bigsqcup_{i \in I} U_i$  into at most countably many Borel subsets of positive measure, such that for every  $i \in I$ , there exists a nice splitting  $S_i$  of  $F_N$  which is  $(\mathcal{H}'_{|U_i}, \rho)$ -invariant, and up to refining the partition it is also  $(\mathcal{H}_{0|U_i}, \rho)$ -invariant.

Lemma 13.25 implies that  $S_i = S_0$  for every  $i \in I$ . Thus, there exists a conull Borel subset  $U_i^* \subseteq U_i$  such that  $\mathcal{H}'_{|U_i^*}$  is contained in  $\rho^{-1}(H_0)_{|U_i^*} = \mathcal{H}_0_{|U_i^*}$ , and hence  $\mathcal{H}_0_{|U}$  is stably equal to  $\mathcal{H}'$ .  $\square$

### 13.4.2 Proof of Property $(P_2)$

**Lemma 13.27.** *The pair  $(\mathcal{G}, \mathcal{H}_0)$  satisfies Property  $(P_2)$ .*

*Proof.* Assume towards a contradiction that there exists a Borel subset  $U \subseteq Y$  of positive measure such that  $\mathcal{H}_0_{|U}$  normalizes an amenable subgroupoid  $\mathcal{A}$  of  $\mathcal{G}_{|U}$  of infinite type.

We first claim that there exists a Borel subset  $V \subseteq U$  of positive measure such that  $S_0$  is  $(\mathcal{A}_{|V}, \rho)$ -invariant. Indeed, by Lemma 10.7, there exists a Borel partition  $Y = Y_1 \sqcup Y_2$  such that  $(\mathcal{A}_{|Y_1}, \rho)$  is nice-averse and  $(\mathcal{A}_{|Y_2}, \rho)$  is stably nice. As  $\mathcal{A}_{|Y_1}$  is amenable and normalized by  $\mathcal{H}_0_{|Y_1}$ , and as  $(\mathcal{A}_{|Y_1}, \rho)$  is nice-averse and  $\rho$  has trivial kernel, Corollary 12.2 implies that  $\mathcal{H}_0_{|Y_1}$  is amenable, so  $Y_1$  is a null set. Using Proposition 10.6, we deduce that there exists a partition  $Y = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets of positive measure such that for every  $i \in I$ , there exists a nice splitting  $S_i$  which is both  $(\mathcal{A}_{|Y_i}, \rho)$ -invariant and  $(\mathcal{H}_0_{|Y_i}, \rho)$ -invariant. Lemma 13.25 then implies that  $S_i = S_0$  for every  $i \in I$ . This completes the proof of our claim.

Up to replacing  $V$  by a conull Borel subset, we can assume that  $\mathcal{A}_{|V}$  is contained in  $\mathcal{H}_0_{|V}$ . In particular, the cocycle  $\rho_{A_0} : \mathcal{H}_0 \rightarrow \text{Out}(A_0)$  is defined on  $\mathcal{A}_{|V}$ .

We aim to prove that  $\rho_{A_0}$  is stably trivial in restriction to  $\mathcal{A}_{|V}$  (in the sense of Definition 3.18). So consider a partition  $V = V_0 \sqcup V_1 \sqcup V_2$  where  $\rho_{A_0}$  is stably trivial in restriction to  $\mathcal{A}_{|V_0}$ ,  $(\mathcal{A}_{|V_1}, \rho_{A_0})$  is nice-averse and  $(\mathcal{A}_{|V_2}, \rho_{A_0})$  is stably nice and nowhere trivial.

If  $V_2$  has positive measure, then by Proposition 10.6,  $(\mathcal{H}_0_{|V_2}, \rho_{A_0})$  is stably nice, which contradicts Lemma 13.25. So  $V_2$  is a null set.

Assume towards a contradiction that  $V_1$  has positive measure. Let  $F \subseteq H_0$  be a non-abelian free subgroup that injects in  $\text{Out}(A_0)$  under the natural restriction map  $H_0 \rightarrow \text{Out}(A_0)$ . Let  $\mathcal{H}' = \rho^{-1}(F)$ . Then  $(\rho_{A_0})_{|\mathcal{H}'}$  has trivial kernel. Since  $\mathcal{A}_{|V_1}$  is amenable and normalized by  $\mathcal{H}_0_{|V_1}$ , and  $(\mathcal{A}_{|V_1}, \rho_{A_0})$  is nowhere trivial, Theorem 12.1 implies that either  $\mathcal{H}'_{|V_1}$  is amenable or there exist a Borel subset  $W_1 \subseteq V_1$  of positive measure and a nice splitting of  $A_0$  which is both  $(\mathcal{A}_{|W_1}, \rho_{A_0})$ -invariant and  $(\mathcal{H}'_{|W_1}, \rho_{A_0})$ -invariant. The former case is excluded by Lemma 3.40 because  $F$  is a non-abelian free group, and the latter case is excluded because  $(\mathcal{A}_{|W_1}, \rho_{A_0})$  is nice-averse.

We thus conclude that  $\rho_{A_0}$  is stably trivial in restriction to  $\mathcal{A}_{|V}$ . So consider a Borel subset  $W \subseteq V$  of positive measure such that for all  $g \in \mathcal{A}_{|W}$ , one has  $\rho_{A_0}(g) = 1$ . Then  $\rho_{|\mathcal{A}_{|W}}$  takes its values in the group of twists  $\text{Tw} \simeq A_0 \times A_0$  of the splitting  $S_0$ . Let now  $\mathcal{H}'' = \rho^{-1}(\text{Tw})_{|W}$ . Then  $\mathcal{A}_{|W}$  is an amenable subgroupoid of infinite type contained in  $\mathcal{H}''$  and normalized by  $\mathcal{H}''$ . Notice that  $\rho$  restricts to an action-type cocycle  $\mathcal{H}'' \rightarrow \text{Tw}$ . We thus get a contradiction to Lemma 11.3.  $\square$

### 13.4.3 Proof of Property $(P_3)$

**Lemma 13.28.** *If  $\text{rank}(A_0) \geq 3$ , then for every Borel subset  $U \subseteq Y$  of positive measure,  $\emptyset$  is the only almost free factor system which is  $(\mathcal{H}_{0|U}, \rho_{A_0})$ -invariant.*

*If  $A_0 = \langle a, b \rangle \simeq F_2$ , then the almost free factor system  $\hat{\mathcal{F}}_{A_0} = \{[\langle aba^{-1}b^{-1} \rangle]\}$  is  $\text{Out}(A_0)$ -invariant and for every Borel subset  $U \subseteq Y$  of positive measure,  $\hat{\mathcal{F}}_{A_0}$  is the only non-empty almost free factor system which is  $(\mathcal{H}_{0|U}, \rho_{A_0})$ -invariant.*

*Proof.* If  $\text{rank}(A_0) \geq 3$ , there exists  $\alpha \in \text{IA}(A_0)$  which is atoroidal and fully irreducible: indeed, by [Sta82] and [GS91], there exists an outer automorphism which is both fully irreducible and non-geometric (i.e. not induced by a surface homeomorphism), and by [BH92, Theorem 4.1] this implies that it is atoroidal. Then for all  $k \geq 1$ ,  $\emptyset$  is the only almost free factor system which is  $\alpha^k$ -invariant. By Remark 3.23, for every Borel subset  $U \subseteq Y$  of positive measure,  $\rho(\mathcal{H}_{0|U})$  contains a non-trivial power of  $\alpha$ , which proves the first assertion.

If  $\text{rank}(A_0) = 2$ , then  $\text{Out}(A_0)$  is isomorphic to the mapping class group of a punctured torus, with  $\hat{\mathcal{F}}_{A_0}$  corresponding to the peripheral element. To prove the second assertion, it suffices to consider  $\alpha \in \text{Out}(F_2)$  induced by a pseudo-Anosov homeomorphism. Then  $[\langle aba^{-1}b^{-1} \rangle]$  is the only  $\alpha$ -periodic conjugacy class of maximal cyclic subgroup, and one concludes using Remark 3.23 as above.  $\square$

**Lemma 13.29.** *There exists a measured subgroupoid  $\mathcal{H}' \subseteq \mathcal{H}_0$  which is everywhere non-amenable and such that  $(\rho_{A_0})|_{\mathcal{H}'}$  has trivial kernel.*

*Proof.* Let  $F_0 = \langle \bar{\alpha}, \bar{\beta} \rangle \subseteq \text{IA}(A_0)$  be a non-abelian free group, and let  $F = \langle \alpha, \beta \rangle \subseteq H_0$  be a subgroup mapping isomorphically to  $F_0$  under the restriction map. The subgroupoid  $\mathcal{H}' = \rho^{-1}(F)$  is everywhere non-amenable by Lemma 3.40. As  $F$  injects into  $\text{Out}(A_0)$  and  $\rho$  has trivial kernel, it follows that  $(\rho_{A_0})|_{\mathcal{H}'}$  has trivial kernel.  $\square$

**Lemma 13.30.** *The pair  $(\mathcal{G}, \mathcal{H}_0)$  satisfies Property  $(P_3)$ .*

*Proof.* Assume by contradiction that  $(P_3)$  does not hold, and let  $U \subseteq Y$ , and subgroupoids  $\mathcal{N}^+, \mathcal{N}^-, \mathcal{B}_i^+, \mathcal{B}_i^-$  ( $i \leq 3$ ) of  $\mathcal{H}_{0|U}$  be such that  $\mathcal{N}^-, \mathcal{N}^+$  are normal in  $\mathcal{H}_{0|U}$ , and for all  $\varepsilon \in \{\pm\}$ , the four subgroupoids  $\mathcal{B}_1^\varepsilon, \mathcal{B}_2^\varepsilon, \mathcal{B}_3^\varepsilon, \mathcal{N}^{-\varepsilon}$  form a pseudo-product (Definition 13.5), with all  $\mathcal{B}_i^+$  contained in  $\mathcal{N}^+$  and all  $\mathcal{B}_i^-$  contained in  $\mathcal{N}^-$ .

Thus consider for all  $\varepsilon \in \{\pm\}$  and all  $i \leq 3$  a pair of amenable subgroupoids  $\mathcal{A}_{i,1}^\varepsilon, \mathcal{A}_{i,2}^\varepsilon \subseteq \mathcal{B}_i^\varepsilon$  stably normalized by  $\mathcal{B}_{i'}^\varepsilon$  for all  $i' \neq i$ , and by  $\mathcal{N}^{-\varepsilon}$  and such that for every Borel subset  $V \subseteq U$  of positive measure, the groupoid  $\langle (\mathcal{A}_{i,1}^\varepsilon)|_V, (\mathcal{A}_{i,2}^\varepsilon)|_V \rangle$  is non-amenable. Notice in particular that  $\mathcal{N}^+$  and  $\mathcal{N}^-$  are everywhere non-amenable. Up to restricting to a smaller subset of positive measure, we may assume that  $\mathcal{A}_{i,j}^\varepsilon$  is actually normalized (and not only stably normalized) by  $\mathcal{B}_i^\varepsilon$  and  $\mathcal{N}^{-\varepsilon}$ .

Since  $\mathcal{N}^+, \mathcal{N}^- \subseteq \mathcal{H}_{0|U}$ , the cocycle  $\rho_{A_0} : \mathcal{H}_0 \rightarrow \text{Out}(A_0)$  is well-defined on all the subgroupoids under consideration.

**Claim.** *There exists  $\varepsilon \in \{\pm\}$  and a Borel subset  $V \subseteq U$  of positive measure such that the cocycle  $\rho_{A_0}$  is trivial on the six subgroupoids  $(\mathcal{A}_{i,j}^\varepsilon)|_V$  for  $i \leq 3, j \leq 2$ .*

*Proof of the claim.* We can assume that  $(\rho_{A_0})|_{\mathcal{N}^-}$  is nowhere trivial since otherwise, the claim would hold as  $\rho_{A_0}$  would be trivial on the six subgroupoids  $(\mathcal{A}_{i,j}^-)|_V$  for some  $V \subseteq U$  of positive measure. Let  $k \leq 6$  be maximal such that there exists  $V \subseteq U$  of positive measure such that  $\rho_{A_0}$  is trivial on  $k$  of the six subgroupoids  $(\mathcal{A}_{i,j}^+)|_V$ . We assume that  $k < 6$  and argue towards a contradiction.

Up to restricting to  $V$ , the cocycle  $\rho_{A_0}$  is nowhere trivial on the restriction of each of the remaining  $6 - k$  subgroupoids to  $V$ , so up to renumbering, we may assume that  $\rho_{A_0}$  is nowhere trivial on  $\mathcal{A}_{1,1}^+$ . Using Proposition 10.5, we may additionally assume up to restricting to a Borel subset of positive measure that  $(\mathcal{A}_{1,1}^+, \rho_{A_0})$  is either stably nice, or pure and nice-averse. If stably nice, then so is  $(\mathcal{N}^-, \rho_{A_0})$  by Proposition 10.6. Since  $\rho_{A_0}$  is nowhere trivial on  $\mathcal{N}^-$ , one can apply Proposition 10.6 again and get that  $(\mathcal{H}_0, \rho_{A_0})$  is stably nice. This contradicts Lemma 13.25, so  $(\mathcal{A}_{1,1}^+, \rho_{A_0})$  is pure and nice-averse. We denote by  $\hat{\mathcal{F}}_1$  the unique maximal  $(\mathcal{A}_{1,1}^+, \rho_{A_0})$ -invariant almost free factor system of  $A_0$ .

We will now set up a contradiction to Theorem 12.27. First, by Lemma 13.29, we can find a measured subgroupoid  $\mathcal{H}'$  of  $\mathcal{H}_0$  which is everywhere non-amenable, such that  $(\rho_{A_0})|_{\mathcal{H}'}$  has trivial kernel. As  $\mathcal{A}_{1,1}^+$  is amenable and normalized by  $\mathcal{N}^-$ , and  $\mathcal{N}^-$  is normalized by  $\mathcal{H}'$ , we can therefore apply Theorem 12.27 to  $(\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_3) = (\mathcal{A}_{1,1}^+, \mathcal{N}^-, \mathcal{H}_0)$  with respect to  $\rho_{A_0}$ . Since  $(\mathcal{H}_0, \rho_{A_0})$  is nice-averse (Lemma 13.25) and  $\mathcal{H}'$  is everywhere non-amenable, the only possible conclusion from this theorem is Assertion 2 saying that  $\mathcal{H}_0$  preserves an almost free factor system  $\hat{\mathcal{F}} \not\supseteq \hat{\mathcal{F}}_1$ . In particular,  $\hat{\mathcal{F}} \neq \emptyset$  which contradicts Lemma 13.28 if  $\text{rank}(A_0) \geq 3$  (i.e.  $N \geq 4$ ). If  $\text{rank}(A_0) = 2$ , then one has  $\hat{\mathcal{F}}_{A_0} \preceq \hat{\mathcal{F}}_1 \not\supseteq \hat{\mathcal{F}}$ , again a contradiction.  $\square$

Thanks to the claim, up to swapping signs, we may find a Borel subset  $V \subseteq U$  of positive measure such that for every  $i \in \{1, 2, 3\}$  and every  $j \in \{1, 2\}$ , one has  $\rho_{A_0}((\mathcal{A}_{i,j}^+)|_V) = \{\text{id}\}$ . Since  $\rho$  takes values in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , this means that  $\rho((\mathcal{A}_{i,j}^+)|_V) \subseteq \text{Tw}$  where  $\text{Tw}$  is the group of twists of the splitting  $S_0$  in  $\text{Out}(F_N)$ . Recall that  $\text{Tw}$  is isomorphic to the direct product  $A_l \times A_r$  of two copies of  $A_0$ . We denote by  $\rho_l$  and  $\rho_r$  the corresponding cocycles (given by postcomposing  $\rho$  with the projection to one of the two factors).

We now consider the four groupoids  $\mathcal{A}_{i,j}^+$  for  $i, j \leq 2$ . Consider a Borel subset  $W \subseteq V$  of positive measure such that one of the following holds:

1.  $\rho_l((\mathcal{A}_{i,j}^+)|_W) = \{\text{id}\}$  for all  $i, j \leq 2$ ;
2.  $\rho_r((\mathcal{A}_{i,j}^+)|_W) = \{\text{id}\}$  for all  $i, j \leq 2$ ;
3. there exists  $(i_l, j_l), (i_r, j_r) \in \{1, 2\} \times \{1, 2\}$  such that  $\rho_l$  is nowhere trivial on  $(\mathcal{A}_{i_l, j_l}^+)|_W$ , and  $\rho_r$  is nowhere trivial on  $(\mathcal{A}_{i_r, j_r}^+)|_W$ .

First assume that Assertion 1 holds. Since  $\rho$  has trivial kernel, then  $\rho_r$  has trivial kernel on the groupoid  $\langle (\mathcal{A}_{1,1}^+)|_W, (\mathcal{A}_{1,2}^+)|_W \rangle$  which is everywhere non-amenable. This groupoid normalizes the amenable groupoid  $(\mathcal{A}_{2,2}^+)|_W$  on which  $\rho_r$  is nowhere trivial. This contradicts Lemma 11.1.

Similarly, Assertion 2 leads to a contradiction.

If Assertion 3 holds, we apply Lemma 11.2 to the two amenable groupoids  $(\mathcal{A}_{i_r, j_r}^+)_{|W}$ ,  $(\mathcal{A}_{i_r, j_r}^+)_{|W}$  which are normalized by the groupoid  $\langle (\mathcal{A}_{3,1}^+)_{|W}, (\mathcal{A}_{3,2}^+)_{|W} \rangle$  on which  $\rho$  has trivial kernel. We get that  $\langle (\mathcal{A}_{3,1}^+)_{|W}, (\mathcal{A}_{3,2}^+)_{|W} \rangle$  is amenable, a contradiction.  $\square$

## 14 Characterizing compatibility

Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle, with  $N \geq 3$ . Given two splittings  $S_1, S_2 \in \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$ , the goal of the present section is to characterize the compatibility of  $S_1$  and  $S_2$  in terms of a purely groupoid-theoretic statement about their  $(\mathcal{G}, \rho)$ -stabilizers.

We refer to Section 13 for Properties  $(P_1^{\text{max}})$ ,  $(P_2)$  and  $(P_3)$ . We introduce the following property of a triple  $(\mathcal{G}; \mathcal{G}_1, \mathcal{G}_2)$ , where  $\mathcal{G}$  is a measured groupoid over a base space  $Y$ , and  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are measured subgroupoids of  $\mathcal{G}$ .

- $(P_{\text{comp}}^1)$  Every measured subgroupoid of  $\mathcal{G}$  that stably normalizes  $\mathcal{G}_1 \cap \mathcal{G}_2$  is stably contained in  $\mathcal{G}_1 \cap \mathcal{G}_2$ .
- $(P_{\text{comp}}^2)$  For every Borel subset  $Z \subseteq Y$  of positive measure, and every measured subgroupoid  $\mathcal{H}$  of  $\mathcal{G}_{|Z}$  that stably contains  $(\mathcal{G}_1 \cap \mathcal{G}_2)_{|Z}$  and such that  $(\mathcal{G}_{|Z}, \mathcal{H})$  satisfies Properties  $(P_1^{\text{max}})$ ,  $(P_2)$  and  $(P_3)$ , there exists a Borel partition  $Z = Z_1 \sqcup Z_2$  such that the groupoids  $\mathcal{H}_{|Z_1}$  and  $\mathcal{H}_{|Z_2}$  are stably contained in  $(\mathcal{G}_1)_{|Z_1}$  and  $(\mathcal{G}_2)_{|Z_2}$  respectively.

Notice that Properties  $(P_{\text{comp}}^1)$  and  $(P_{\text{comp}}^2)$  are preserved by restriction to a Borel subset of positive measure, and by stabilization.

**Theorem 14.1.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle with  $N \geq 3$ . Let  $S_1, S_2 \in \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$ , and for every  $j \in \{1, 2\}$ , let  $\mathcal{G}_j$  be the  $(\mathcal{G}, \rho)$ -stabilizer of  $S_j$ . The following assertions are equivalent.*

- (i) *The splittings  $S_1$  and  $S_2$  are compatible.*
- (ii) *The triple  $(\mathcal{G}; \mathcal{G}_1, \mathcal{G}_2)$  satisfies Properties  $(P_{\text{comp}}^1)$  and  $(P_{\text{comp}}^2)$ .*

We will use the following lemma from [HW20].

**Lemma 14.2** ([HW20, Lemma 2.7]). *Let  $U$  be a one-edge  $\mathcal{Z}_{\text{max}}$ -splitting of  $F_N$ , let  $\tau_U$  be a Dehn twist about  $U$ , and let  $S$  be a free splitting of  $F_N$ .*

*If  $\tau_U^k(S) = S$  for some  $k \neq 0$ , then  $S$  is compatible with  $U$ .*

We recall from Section 13.3.4 that a free splitting  $S$  of  $F_N$  is *acyclic* if the quotient graph  $S/F_N$  does not contain any terminal vertex with vertex group isomorphic to  $\mathbb{Z}$ .

**Lemma 14.3.** *Let  $S$  be an acyclic free splitting of  $F_N$ , such that all vertex stabilizers of  $S$  are non-trivial. Let  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be the stabilizer of  $S$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ .*

*If  $S'$  is a free splitting of  $F_N$  such that every element in  $H$  has a non-trivial power that fixes  $S'$ , then  $S'$  is a collapse of  $S$ .*

*Remark 14.4.* The conclusion does not hold if  $S$  is not supposed to be acyclic. Indeed, in the case where  $S/F_N$  has a terminal vertex with edge group isomorphic to  $\mathbb{Z}$ , there is a canonical way to blow up that vertex into a loop-edge, and this blowup is  $H$ -invariant (but is not a collapse of  $S$ ). Likewise, if  $S$  contains a vertex  $v$  of valence at least 4 with trivial stabilizer, the conclusion of the lemma fails. Indeed, in this case, there are finitely possible blowups of  $S$  at  $v$ , and these are permuted by  $H$ , whence  $H$ -invariant because  $H \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ .

*Proof.* Let  $E$  be a set of representatives of the  $F_N$ -orbits of half-edges of  $S$ . For every  $e \in E$  with initial vertex  $v$ , choose an element  $g_e \in G_v$  which is not a proper power and is not contained in any proper free factor of  $G_v$ . This is possible because all vertex stabilizers are non-trivial.

Let  $U$  be the  $\mathcal{Z}_{\max}$ -splitting of  $F_N$  obtained from  $S$  by folding every half-edge  $e \in E$  with its  $g_e$ -translate, and extending this operation equivariantly to the whole tree  $S$ . The fact that  $S$  is acyclic implies that  $U$  is minimal. Notice that the splitting  $U$  is naturally bipartite over the vertex set  $V = V_0 \sqcup V_1$ , where vertices in  $V_0$  are the images in  $U$  of vertices of  $S$ , and vertices in  $V_1$  are the images in  $U$  of midpoints of edges in  $S$ . Notice that the stabilizer of every vertex in  $V_0$  is the same in  $S$  and in  $U$ , and that the stabilizer of every vertex in  $V_1$  is isomorphic to  $F_2$ : more precisely, if  $v$  is a vertex in  $V_1$ , and if  $e$  is an edge of  $S$  whose midpoint is mapped to  $v$  under the folding map from  $S$  to  $U$ , denoting by  $e_1$  and  $e_2$  the two half-edges of  $e$  (based at two distinct vertices of  $S$ , and folded with respective translates  $g_1e_1$  and  $g_2e_2$ ), then  $G_v$  is the subgroup of  $F_N$  generated by  $\langle g_1, g_2 \rangle$ .

Given a half-edge  $e \subseteq S$ , we let  $U_e$  be the collapse of  $U$  obtained by collapsing all edges of  $U$  except those in the orbit of the image of  $e$ . Let  $\tau_e$  be the twist of  $S$  by  $g_e$  around  $e$  near  $v$  (see Section 2.8), or equivalently, the Dehn twist about the cyclic splitting  $U_e$ . Since  $\tau_e$  preserves  $S$ , some power of  $\tau_e$  lies in  $H$  so there exists  $k \geq 1$  such that  $\tau_e^k(S') = S'$ .

By Lemma 14.2, this implies that  $S'$  is compatible with  $U_e$ . As this is true for every one-edge collapse  $U_e$  of  $U$ , we deduce that  $S'$  is compatible with  $U$  ([GL17, Theorem A.17]). In other words  $S'$  is obtained from  $U$  by first blowing up some vertices  $v$  of  $U$  using a free splitting of  $G_v$  relative to the incident edge stabilizers, and then collapsing every edge coming from  $U$  to a point. By choice of the elements  $g_e$ , for every vertex  $v \in V_0$ , there does not exist any free splitting of  $G_v$  relative to the incident edge stabilizers. For every vertex  $v \in V_1$ , since  $G_v$  is isomorphic to  $F_2$  and generated by two incident edge groups, there is exactly one free splitting of  $G_v$  relative to the incident edge stabilizers; blowing up  $U$  at  $v$  determines a one-edge free splitting of  $F_N$  which is a collapse of  $S$ . This shows that  $S'$  is a collapse of  $S$ .  $\square$

We are now in position to prove Theorem 14.1.

*Proof of Theorem 14.1.* We first prove that  $(P_{\text{comp}}^1)$  holds assuming that  $S_1$  and  $S_2$  are compatible. Note that  $S_1$  and  $S_2$  are acyclic. Let  $\mathcal{H}$  be a measured subgroupoid of  $\mathcal{G}$  that stably normalizes  $\mathcal{G}_1 \cap \mathcal{G}_2$ . Let  $S$  be the two-edge common refinement of  $S_1$  and  $S_2$ .



Then  $S$  is acyclic as otherwise one of its one-edge collapses (i.e.  $S_1$  or  $S_2$ ) would fail to be acyclic. We denote by  $K$  the stabilizer of  $S$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , which coincides with the intersection of the stabilizers of  $S_1$  and  $S_2$ . Notice that  $\mathcal{G}_1 \cap \mathcal{G}_2 = \rho^{-1}(K)$ .

Lemma 14.3 ensures that  $S, S_1, S_2$  are the only non-trivial free splittings of  $F_N$  that are invariant by a power of every element of  $H$ . Using the fact that  $\rho$  is action-type (in the form of Remark 3.23), we deduce that for every Borel subset  $U \subseteq Y$  of positive measure, the splittings  $S, S_1, S_2$  are the only non-trivial free splittings of  $F_N$  which are  $((\mathcal{G}_1 \cap \mathcal{G}_2)|_U, \rho)$ -invariant.

As  $\mathcal{H}$  stably normalizes  $\mathcal{G}_1 \cap \mathcal{G}_2$ , we deduce that  $S$  is stably  $(\mathcal{H}, \rho)$ -invariant. This shows that  $\mathcal{H}$  is stably contained in  $\mathcal{G}_1 \cap \mathcal{G}_2$ , so  $(P_{\text{comp}}^1)$  holds.

To prove  $(P_{\text{comp}}^2)$ , let  $Z \subseteq Y$  be a Borel subset of positive measure, and  $\mathcal{H}$  be a subgroupoid of  $\mathcal{G}|_Z$  that stably contains  $(\mathcal{G}_1 \cap \mathcal{G}_2)|_Z$ , and such that  $(\mathcal{G}|_Z, \mathcal{H})$  satisfies Properties  $(P_1^{\text{max}})$ ,  $(P_2)$  and  $(P_3)$ . By the first part of Theorem 13.10, we can find a partition  $Z = \bigsqcup_{i \in I} Z_i$  into at most countably many Borel subsets such that for every  $i \in I$ , there exists a non-trivial free splitting  $S'_i \in \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$  which is  $(\mathcal{H}|_{Z_i}, \rho)$ -invariant. Up to further partitioning the base space and replacing  $Z$  by a conull Borel subset, we can also assume that for every  $i \in I$ , one has  $(\mathcal{G}_1 \cap \mathcal{G}_2)|_{Z_i} \subseteq \mathcal{H}|_{Z_i}$ . Since  $\rho$  is action-type and  $\mathcal{G}_1 \cap \mathcal{G}_2 = \rho^{-1}(K)$ , every element of  $K$  has a power that fixes  $S'_i$  (see Remark 3.23). By Lemma 14.3, this implies that  $S'_i$  is a collapse of  $S$ , so either  $S_1$  or  $S_2$  is  $(\mathcal{H}|_{Z_i}, \rho)$ -invariant and  $(P_{\text{comp}}^2)$  follows.

We now prove that  $(ii) \Rightarrow (i)$ . Assume that  $(\mathcal{G}; \mathcal{G}_1, \mathcal{G}_2)$  satisfies Properties  $(P_{\text{comp}}^1)$  and  $(P_{\text{comp}}^2)$ . Let  $\mathcal{C} = \{S_1, S_2\}$ , and as above let  $K = \Gamma_{\mathcal{C}}$  be the elementwise stabilizer of  $\mathcal{C}$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Observe that  $K$  is infinite: otherwise  $K$  is trivial (since we are working in the torsion-free subgroup  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ ), and as  $\rho$  has trivial kernel, the groupoid  $\mathcal{G}_1 \cap \mathcal{G}_2 = \rho^{-1}(K)$  would be trivial, contradicting Property  $(P_{\text{comp}}^1)$ .

Let  $U = U_K^1$  be the splitting constructed from the group  $K$  and the collection of  $K$ -invariant free splittings in Theorem 6.12. Recall that  $U$  comes with a bipartition of its vertex set  $V = V^0 \sqcup V^1$  and that  $S_1$  and  $S_2$  may be obtained from  $U$  by blowing up some vertices  $v_1, v_2 \in V^1$  (using a possibly non-minimal free splitting relative to the incident edge groups) and collapsing all edges coming from  $U$ . If  $v_1$  and  $v_2$  are in different orbits, then one can perform the two blowups simultaneously and conclude that  $S_1$  and  $S_2$  are compatible, so up to changing  $v_2$  to another vertex in the same orbit, we may assume that  $v_1 = v_2$ , and we denote  $v = v_1 = v_2$ .

Let  $H$  be the stabilizer of  $U$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ . Then  $H$  normalizes  $K$  (see Proposition 6.19) so the subgroupoid  $\mathcal{H} = \rho^{-1}(H)$  normalizes  $\mathcal{K} = \rho^{-1}(K) = \mathcal{G}_1 \cap \mathcal{G}_2$ . By  $(P_{\text{comp}}^1)$ ,  $\mathcal{H}$  is stably contained in  $\mathcal{K}$ . As  $\rho$  is action-type, this implies in particular that every element of  $H$  has a power contained in  $K$ . We can therefore apply Corollary 6.21, and deduce that if  $v$  has non-trivial stabilizer, then  $S_1$  and  $S_2$  are compatible. So from now on, we assume that  $G_v$  is trivial.

We denote the quotient graph  $\bar{U} = U/F_N$ . For each edge  $e$  in  $\bar{U}$  with trivial stabilizer, we denote by  $U_e$  the free splitting obtained from  $U$  by collapsing all the edges whose image in  $\bar{U}$  is different from  $e$ . Since  $U$  is compatible with every splitting in  $\mathcal{C}$ , so is  $U_e$  hence we are done if  $S_1$  or  $S_2$  coincides with one of the trees  $U_e$ .



We now prove the result assuming that  $\bar{U}$  has some non-separating edge  $e$  with trivial stabilizer. Let  $H'_e$  be the stabilizer in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  of  $U_e$ , and let  $\mathcal{H}'_e := \rho^{-1}(H'_e)$ . Since  $K \subseteq H \subseteq H'_e$ , we have  $\mathcal{G}_1 \cap \mathcal{G}_2 \subseteq \mathcal{H}'_e$ . As  $e$  is non-separating, Theorem 13.10 shows that  $(\mathcal{G}, \mathcal{H}'_e)$  satisfies Properties  $(P_1^{\max})$ ,  $(P_2)$  and  $(P_3)$ . We can therefore apply Property  $(P_{\text{comp}}^2)$ , and get a Borel partition  $Y = Y_1 \sqcup Y_2$  such that for every  $i \in \{1, 2\}$ , the groupoid  $\mathcal{H}'_{|Y_i}$  is stably contained in  $(\mathcal{G}_i)_{|Y_i}$ . This implies that there exists  $j \in \{1, 2\}$  such that every element of  $H'_e$  has a power contained in the stabilizer of  $S_j$ . As  $H'_e \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , this implies that  $H'_e$  is contained in the stabilizer of  $S_j$ . By Lemma 14.3,  $U_e = S_j$ , so as noticed above, we are done in this case.

In the remaining case, every edge of  $U$  with trivial stabilizer in  $\bar{U}$  is separating. We denote by  $E_v = \{e_1, \dots, e_p\}$  the set of edges incident on  $v$ . They all have trivial stabilizer since  $G_v = \{1\}$ . This gives a decomposition of  $F_N$  as a free product  $F_N \simeq A_1 * \dots * A_p$  where for every  $k \in \{1, \dots, p\}$ ,  $A_k$  is the fundamental group (as a graph of groups) of the connected component of  $\bar{U} \setminus (\hat{e}_1 \cup \dots \cup \hat{e}_p \cup \{v\})$  containing the endpoint of  $e_k$ . Notice that  $A_k$  is non-trivial. The tree  $S_1$  is obtained from the graph of groups  $\bar{U}$  by choosing a partition  $E_v = E_a \sqcup E_b$  of the edges incident on  $v$ , and blowing-up  $v$  into a new edge that separates the two subsets  $E_a$  and  $E_b$ . It follows that  $S_1$  is a separating free splitting. Since  $S_1$  belongs to  $\text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$ , we deduce that up to exchanging the roles of  $E_a$  and  $E_b$ , the splitting  $S_1$  is the Bass–Serre tree of a decomposition of the form  $F_N = A * B$ , where  $A \simeq F_2$  is the free product of the groups  $A_k$  with  $e_k$  ranging in  $E_a$ , and  $B \simeq F_{N-2}$ . We may assume that  $E_a$  and  $E_b$  both contain at least 2 edges since otherwise  $S_1 = U_{e_k}$  for some  $k \leq p$ , in which case we are done by the above. It follows that  $E_a$  contains exactly two edges, and that the two corresponding groups  $A_k$  are infinite cyclic.

Since all edges with trivial stabilizer are separating,  $\bar{U}$  contains a terminal vertex whose stabilizer is isomorphic to  $\mathbb{Z}$ . Blowing up this vertex yields a non-separating free splitting  $U'$  of  $F_N$  which is  $H$ -invariant. As above,  $(P_{\text{comp}}^2)$  implies that  $U' = S_1$  or  $U' = S_2$ , a contradiction.  $\square$

## 15 Maps from $\text{FS}^{\text{ns}}$ to $\text{FS}$

The *free splitting graph*  $\text{FS}$  is the simplicial graph whose vertex set is the set  $\text{FS}$  of one-edge free splittings of  $F_N$ , and where two splittings are joined by an edge if they are compatible. The *non-separating free splitting graph*  $\text{FS}^{\text{ns}} \subseteq \text{FS}$  is the induced subgraph whose vertex set  $\text{FS}^{\text{ns}}$  is the set of non-separating one-edge free splittings of  $F_N$ . The goal of the present section is to prove the following proposition.

**Proposition 15.1.** *Let  $N \geq 3$ , and let  $\theta : \text{FS}^{\text{ns}} \rightarrow \text{FS}$  be an injective map that preserves adjacency and non-adjacency.*

*Then  $\theta(\text{FS}^{\text{ns}}) \subseteq \text{FS}^{\text{ns}}$ .*

*Remark 15.2.* In fact, the proposition remains true for  $N = 2$ , as in this case one can check that a vertex of  $\text{FS}$  is in  $\text{FS}^{\text{ns}}$  if and only if it has infinite valence (there are exactly two free splittings that are compatible with a given separating free splitting, given by blowing up each of the two vertex groups isomorphic to  $\mathbb{Z}$ ).

Recall that the *link* of a vertex  $x$  in a simplicial graph  $\mathbb{G}$  is the induced subgraph  $\text{lk}_{\mathbb{G}}(x) \subseteq \mathbb{G}$  whose vertex set is the set of vertices  $v \neq x$  in  $\mathbb{G}$  that are adjacent to  $x$ . Recall that a graph with vertex set  $V$  splits non-trivially as a join  $X_1 \star X_2$  if  $V = X_1 \sqcup X_2$  with  $X_1, X_2 \neq \emptyset$  and for all  $x_1 \in X_1$  and all  $x_2 \in X_2$  there is an edge joining  $x_1$  to  $x_2$ .

**Lemma 15.3.** *For any  $S \in \text{FS}^{\text{ns}}$ , the link of  $S$  in  $\mathbb{FS}^{\text{ns}}$  does not split non-trivially as a join.*

*Proof.* It is enough to show that given any two compatible non-separating free splittings  $S_1, S_2$  of  $F_N$  in the link of  $S$ , there exists a non-separating free splitting  $S'$  of  $F_N$  which is neither compatible with  $S_1$  nor with  $S_2$ . Indeed, if  $\text{lk}_{\mathbb{FS}^{\text{ns}}}(S)$  decomposes non-trivially as a join  $\text{lk}_{\mathbb{FS}^{\text{ns}}}(S) = X_1 \star X_2$ , then choosing any splittings  $S_1 \in X_1$  and  $S_2 \in X_2$ , one has that  $S_1$  and  $S_2$  are compatible, and every splitting  $S' \in \text{FS}^{\text{ns}}$  is then compatible with  $S_1$  if  $S' \in X_2$  or with  $S_2$  otherwise.

Let  $A$  be a corank one free factor of  $F_N$  such that  $S$  is the Bass–Serre tree of the HNN extension  $F_N = A*$ . Then the  $A$ -minimal invariant subtrees of  $S_1$  and  $S_2$  yield two (possibly isomorphic) non-trivial free splittings of  $A$ , which we denote by  $(S_1)_{|A}$  and  $(S_2)_{|A}$ . As  $\text{rk}(A) \geq 2$ , we can find a one-edge non-separating free splitting  $S'_A$  of  $A$  that is neither compatible with  $(S_1)_{|A}$  nor with  $(S_2)_{|A}$ . Let  $\hat{S}$  be a refinement of  $S$  such that  $\hat{S}/F_N$  is a two-petalled rose, and whose  $A$ -minimal subtree  $(\hat{S})_{|A}$  is isomorphic to  $S'_A$ : such a splitting can be obtained by blowing up  $S$  at the vertex stabilized by  $A$ , using the splitting  $S'_A$ . Let  $S' \in \text{FS}^{\text{ns}}$  be the splitting obtained from  $\hat{S}$  by collapsing every edge coming from  $S$  to a point. Then  $S'$  is a one-edge non-separating free splitting of  $F_N$  in the link of  $S$ . It is neither compatible with  $S_1$  nor with  $S_2$ , as otherwise  $S'_A$  would be compatible with either  $(S_1)_{|A}$  or  $(S_2)_{|A}$ .  $\square$

**Lemma 15.4.** *For any  $S \in \text{FS}^{\text{ns}}$ , the link of  $S$  in  $\mathbb{FS}^{\text{ns}}$  contains a complete subgraph on  $3N - 4$  vertices.*

*Proof.* The splitting  $S$  has a refinement  $\hat{S}$  such that  $\hat{S}/F_N$  has  $3N - 3$  edges and no separating edge (recall that  $3N - 3$  is the maximal number of edges of a graph in Culler–Vogtmann’s reduced Outer space [CV86]). The one-edge splittings associated to the edges of  $\hat{S}/F_N$  yield in addition to  $S$ ,  $3N - 4$  distinct non-separating one-edge free splittings in the link of  $S$  that are pairwise compatible.  $\square$

**Lemma 15.5.** *For any  $S \in \text{FS} \setminus \text{FS}^{\text{ns}}$ , the link of  $S$  in  $\mathbb{FS}$  decomposes as a non-trivial join  $\text{lk}_{\mathbb{FS}}(S) = X_1 \star X_2$  where neither  $X_1$  nor  $X_2$  contains a complete subgraph on  $3N - 4$  vertices.*

*Remark 15.6.* We allow  $X_1$  or  $X_2$  to be reduced to a point (this occurs if one of the factors of the splitting is cyclic).

*Proof.* Let  $A_1, A_2$  be proper free factors of  $F_N$  such that  $S$  is the Bass–Serre tree of the free product  $F_N = A_1 * A_2$ . Every one-edge free splitting of  $F_N$  that is compatible with  $S$  but distinct from  $S$  is obtained in the following way: choose  $i \in \{1, 2\}$ , construct a refinement  $\hat{S}$  of  $S$  by blowing up  $S$  at the vertex fixed by  $A_i$ , using a one-edge free

splitting of  $A_i$ , and then collapse every edge of  $\hat{S}$  that comes from  $S$  to a point. We say that such a splitting is a *compatible splitting of type  $i$* . Then the link  $\text{lk}_{\mathbb{F}\mathbb{S}}(S)$  decomposes as a non-trivial join  $\text{lk}_{\mathbb{F}\mathbb{S}}(S) = X_1 \star X_2$ , where  $X_i$  consists of all compatible splittings of type  $i$ .

Let  $i \in \{1, 2\}$ . Assume towards a contradiction that  $X_i$  contains a complete subgraph on  $3N - 4$  vertices. Then there exists a refinement of  $S$  with  $3N - 3$  orbits of edges where  $A_j$  is elliptic (with  $j \neq i$ ). This is a contradiction because every free splitting of  $F_N$  with  $3N - 3$  orbits of edges corresponds to a free action of  $F_N$  (this simple observation may be viewed as a consequence of the deeper fact that the boundary of Outer space has dimension  $3N - 5$  [GL95]).  $\square$

*Proof of Proposition 15.1.* Assume towards a contradiction that there exists a vertex  $S \in \text{FS}^{\text{ns}}$  such that  $\theta(S)$  is a separating splitting. By Lemma 15.5, the link  $\text{lk}_{\mathbb{F}\mathbb{S}}(\theta(S))$  decomposes as a non-trivial join  $\text{lk}_{\mathbb{F}\mathbb{S}}(\theta(S)) = X_1 \star X_2$ , where no  $X_i$  contains a complete subgraph on  $3N - 4$  vertices. By Lemma 15.3, the link  $\text{lk}_{\mathbb{F}\mathbb{S}^{\text{ns}}}(S)$  does not decompose non-trivially as a join, and as  $\theta$  preserves both adjacency and non-adjacency, neither does  $\theta(\text{lk}_{\mathbb{F}\mathbb{S}^{\text{ns}}}(S))$ . Since

$$\theta(\text{lk}_{\mathbb{F}\mathbb{S}^{\text{ns}}}(S)) = (\theta(\text{lk}_{\mathbb{F}\mathbb{S}^{\text{ns}}}(S)) \cap X_1) \star (\theta(\text{lk}_{\mathbb{F}\mathbb{S}^{\text{ns}}}(S)) \cap X_2),$$

we deduce that  $\theta(\text{lk}_{\mathbb{F}\mathbb{S}^{\text{ns}}}(S))$  is contained in either  $X_1$  or  $X_2$ . On the other hand, Lemma 15.4 ensures that  $\text{lk}_{\mathbb{F}\mathbb{S}^{\text{ns}}}(S)$  contains a complete subgraph on  $3N - 4$  vertices, a contradiction.  $\square$

## 16 Measure equivalence rigidity of $\text{Out}(F_N)$

The goal of this section is to complete the proof of the main theorem of the paper.

**Theorem 16.1.** *For every  $N \geq 3$ , the group  $\text{Out}(F_N)$  is ME-superrigid.*

The following lemma is folklore. It follows for instance from [FH07, HW20] since any element whose conjugacy class is finite is in the kernel of the map from a group to its abstract commensurator. The lemma also follows from [GG13] or [AKP17], or could be proved using an argument along the lines of [Kid10, Theorem 2.9].

**Lemma 16.2.** *For every  $N \geq 3$ , the group  $\text{Out}(F_N)$  is ICC: the conjugacy class of every non-trivial element of  $\text{Out}(F_N)$  is infinite.*  $\square$

*Proof of Theorem 16.1.* Since  $\text{Out}(F_N)$  is ICC, the theorem follows from Theorem 4.5 together with the following statement, which shows that  $\text{Out}(F_N)$  is rigid with respect to action-type cocycles (Definition 4.1).  $\square$

**Theorem 16.3.** *For every  $N \geq 3$ , the group  $\text{Out}(F_N)$  is rigid with respect to action-type cocycles.*

*More precisely, let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho, \rho' : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be two action-type cocycles.*

Then  $\rho$  and  $\rho'$  are  $\text{Out}(F_N)$ -cohomologous, i.e. there exist a Borel map  $\varphi : Y \rightarrow \text{Out}(F_N)$  and a conull Borel subset  $Y^* \subseteq Y$  such that for all  $g \in \mathcal{G}|_{Y^*}$ , one has  $\rho'(g) = \varphi(r(g))\rho(g)\varphi(s(g))^{-1}$ .

We are now left proving Theorem 16.3.

**Lemma 16.4.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be an action-type cocycle. Let  $S_1, S_2 \in \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$  be two distinct splittings and  $H_1, H_2 \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  their stabilizers.*

*Then for every Borel subset  $U \subseteq Y$  of positive measure, one has  $\rho^{-1}(H_1)|_U \neq \rho^{-1}(H_2)|_U$ .*

*Proof.* The splitting  $S_1$  is acyclic (i.e. its quotient graph does not contain any terminal vertex with vertex group isomorphic to  $\mathbb{Z}$ ), so Lemma 14.3 shows that there exists an element in  $H_1$  none of whose powers belongs to  $H_2$ . As  $\rho$  is action-type, this implies that  $\rho^{-1}(H_1)|_U$  is not contained in  $\rho^{-1}(H_2)|_U$  (see Remark 3.23), which concludes the proof.  $\square$

Recall that the  $(\mathcal{G}, \rho)$ -stabilizer of a Borel map  $\sigma : Y \rightarrow \text{FS}$  is the subgroupoid of  $\mathcal{G}$  consisting of all  $g \in \mathcal{G}$  such that  $\sigma(r(g)) = \rho(g)\sigma(s(g))$ . Note that if  $Y = \sqcup Y_i$  is a finite or countable Borel partition such that  $\sigma$  is constant on  $Y_i$  with value  $S_i$ , and if  $H_i \subseteq \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  denotes the stabilizer of  $S_i$ , then the  $(\mathcal{G}, \rho)$ -stabilizer of  $\sigma$  stably coincides with the groupoid  $\cup_i(\rho^{-1}(H_i))|_{Y_i}$ .

*Remark 16.5.* As a consequence of Lemma 16.4, if  $\sigma, \sigma' : Y \rightarrow \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$  are two Borel maps such that the  $(\mathcal{G}, \rho)$ -stabilizers of  $\sigma$  and  $\sigma'$  are stably equal, then  $\sigma$  and  $\sigma'$  coincide almost everywhere.

The following key lemma is based on the groupoidal (almost) characterization of stabilizers of non-separating free splittings of Section 13.

**Lemma 16.6.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho, \rho' : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be two action-type cocycles.*

*For every Borel map  $\sigma : Y \rightarrow \text{FS}^{\text{ns}}$ , there exists an essentially unique Borel map  $\sigma' : Y \rightarrow \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$  such that the  $(\mathcal{G}, \rho)$ -stabilizer of  $\sigma$  is stably equal to the  $(\mathcal{G}, \rho')$ -stabilizer of  $\sigma'$ .*

We say that  $\sigma'$  is the  $(\mathcal{G}, \rho, \rho')$ -image of  $\sigma$ . Equivalently, a Borel map  $\sigma' : Y \rightarrow \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$  is (up to measure 0) the  $(\mathcal{G}, \rho, \rho')$ -image of  $\sigma$  if and only if there exist a conull Borel subset  $Y^* \subseteq Y$  and a partition  $Y^* = \bigsqcup_{i \in I} Y_i$  into at most countably many Borel subsets such that for every  $i \in I$ , the maps  $\sigma$  and  $\sigma'$  are constant on  $Y_i$  with respective values  $S_i, S'_i$ , and denoting by  $H_i, H'_i$  the respective stabilizers of  $S_i, S'_i$  in  $\text{IA}_N(\mathbb{Z}/3\mathbb{Z})$ , one has  $(\rho^{-1}(H_i))|_{Y_i} = (\rho'^{-1}(H'_i))|_{Y_i}$ .

Note that *a priori*, there is a slight asymmetry between  $\sigma$  and  $\sigma'$  (coming from the asymmetry in Theorem 13.10): indeed  $\sigma$  is assumed to take its values in  $\text{FS}^{\text{ns}}$  while  $\sigma'$  is assumed to take its values in  $\text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$ . However, using the results from Section 15, we will see in the course of the proof of Theorem 16.3 that  $\sigma'$  actually takes its values in  $\text{FS}^{\text{ns}}$ .

*Proof of Lemma 16.6.* The essential uniqueness of  $\sigma'$  follows from Remark 16.5, so we now prove the existence of  $\sigma'$ .

Let  $\mathcal{H}$  be the  $(\mathcal{G}, \rho)$ -stabilizer of  $\sigma$ . By the second part of Theorem 13.10 (applied to the cocycle  $\rho$ , with the partition given by  $\sigma$ -preimages of points in  $\text{FS}^{\text{ns}}$ ), the pair  $(\mathcal{G}, \mathcal{H})$  satisfies Properties  $(P_1^{\text{max}})$ ,  $(P_2)$  and  $(P_3)$ . By the first part of Theorem 13.10 (applied to the cocycle  $\rho'$ ), there exist a conull subset  $Y^* \subseteq Y$ , a finite or countable Borel partition  $Y^* = \sqcup Y_i$ , and splittings  $S'_i \in \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$ , such that  $\mathcal{H}|_{Y_i}$  is equal to the  $(\mathcal{G}|_{Y_i}, \rho')$ -stabilizer of  $S'_i$ . Consider the map  $\sigma'$  assigning  $S'_i$  to  $y \in Y_i$  (and defined in an arbitrary way on  $Y \setminus Y^*$ ). Then  $\mathcal{H}$  is stably equal to the  $(\mathcal{G}, \rho')$ -stabilizer of  $\sigma'$ .  $\square$

*Remark 16.7.* If  $\sigma'$  is the  $(\mathcal{G}, \rho, \rho')$ -image of  $\sigma$ , and if additionally  $\sigma'$  takes its values in  $\text{FS}^{\text{ns}}$ , then by symmetry,  $\sigma$  is the  $(\mathcal{G}, \rho', \rho)$ -image of  $\sigma'$ .

*Remark 16.8.* If  $\sigma'$  is the  $(\mathcal{G}, \rho, \rho')$ -image of  $\sigma$ , then for every Borel subset  $U \subseteq Y$  of positive measure,  $\sigma'_{|U}$  is the  $(\mathcal{G}|_U, \rho, \rho')$ -image of  $\sigma|_U$ .

The following lemma is based on the groupoidal characterization of compatibility in Section 14.

**Lemma 16.9.** *Let  $\mathcal{G}$  be a measured groupoid over a base space  $Y$ , and let  $\rho, \rho' : \mathcal{G} \rightarrow \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  be two action-type cocycles. Consider  $S_1, S_2, S'_1, S'_2 \in \text{FS}^{\text{ns}} \cup \text{FS}^{2,2^+}$ , and assume that for every  $i \in \{1, 2\}$ , the  $(\mathcal{G}, \rho)$ -stabilizer of  $S_i$  is stably equal to the  $(\mathcal{G}, \rho')$ -stabilizer of  $S'_i$ .*

*Then  $S_1$  is compatible with  $S_2$  if and only if  $S'_1$  is compatible with  $S'_2$ .*

*Proof.* It suffices to assume that  $S_1$  and  $S_2$  are compatible and prove that so are  $S'_1$  and  $S'_2$ . For every  $i \in \{1, 2\}$ , let  $\mathcal{H}_i$  be the  $(\mathcal{G}, \rho)$ -stabilizer of  $S_i$ , and let  $\mathcal{H}'_i$  be the  $(\mathcal{G}, \rho')$ -stabilizer of  $S'_i$ .

Theorem 14.1 (applied to the cocycle  $\rho$ ) ensures that the triple  $(\mathcal{G}; \mathcal{H}_1, \mathcal{H}_2)$  satisfies Properties  $(P_{\text{comp}}^1)$  and  $(P_{\text{comp}}^2)$ . As  $\mathcal{H}_i$  is stably equal to  $\mathcal{H}'_i$ , we deduce that  $(\mathcal{G}; \mathcal{H}'_1, \mathcal{H}'_2)$  also satisfies Properties  $(P_{\text{comp}}^1)$  and  $(P_{\text{comp}}^2)$ . Therefore, the converse implication from Theorem 14.1 (applied to the cocycle  $\rho'$ ) ensures that  $S'_1$  and  $S'_2$  are compatible.  $\square$

*Proof of Theorem 16.3.* In this proof, given two simplicial graphs  $\mathbb{X}$  and  $\mathbb{Y}$  with vertex sets  $V_{\mathbb{X}}$  and  $V_{\mathbb{Y}}$ , we denote by  $\text{Inj}(\mathbb{X} \rightarrow \mathbb{Y})$  the set of all injective maps from  $V_{\mathbb{X}}$  to  $V_{\mathbb{Y}}$  that preserve adjacency and non-adjacency. We denote by  $\text{Aut}(\mathbb{X})$  the group of all graph automorphisms of  $\mathbb{X}$ , i.e. maps in  $\text{Inj}(\mathbb{X} \rightarrow \mathbb{X})$  which are bijective.

Before actually starting the proof, let us briefly explain the general strategy. We will use the graph  $\mathbb{FS}$  of free splittings and its subgraph  $\mathbb{FS}^{\text{ns}}$  of non-separating free splittings. We will start by constructing a map  $\Psi : Y \rightarrow \text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS})$  such that for every Borel map  $\sigma : Y \rightarrow \mathbb{FS}^{\text{ns}}$ , the  $(\mathcal{G}, \rho, \rho')$ -image  $\sigma'$  of  $\sigma$  is obtained by applying  $\Psi$  to  $\sigma$ , i.e.  $\sigma'(y) = \Psi(y)(\sigma(y))$ . There is also a symmetric map  $\Psi'$  obtained by reversing the roles of  $\rho$  and  $\rho'$ . Using the fact that  $\text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}) = \text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}^{\text{ns}})$  (Proposition 15.1), we will actually deduce that  $\Psi'(y)$  is (almost everywhere) an inverse of  $\Psi(y)$ . Therefore  $\Psi$  takes its values in  $\text{Aut}(\mathbb{FS}^{\text{ns}})$ . By a theorem of Pandit ([Pan14], which builds on work of Bridson and Vogtmann regarding the symmetries of Outer space [BV01]), we

know that  $\text{Aut}(\mathbb{FS}^{\text{ns}})$  is isomorphic to  $\text{Out}(F_N)$ . This will allow us to construct the map  $\varphi : Y \rightarrow \text{Out}(F_N)$  that conjugates  $\rho$  to  $\rho'$ .

Let us now construct the map  $\Psi$ . Given  $S \in \mathbb{FS}^{\text{ns}}$ , we let  $\sigma_S^0 : Y \rightarrow \mathbb{FS}^{\text{ns}}$  be the constant map with value  $S$ , and  $\sigma'_S : Y \rightarrow \mathbb{FS}^{\text{ns}} \cup \mathbb{FS}^{2,2^+}$  be the  $(\mathcal{G}, \rho, \rho')$ -image of  $\sigma_S^0$  (the map  $\sigma'_S$  is not necessarily constant). This allows us to define a Borel map

$$\begin{aligned} \tilde{\Psi} : Y \times \mathbb{FS}^{\text{ns}} &\rightarrow \mathbb{FS}^{\text{ns}} \cup \mathbb{FS}^{2,2^+} \\ (y, S) &\mapsto \sigma'_S(y). \end{aligned}$$

Then for each  $y \in Y$ , we define  $\Psi_y : \mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}^{\text{ns}} \cup \mathbb{FS}^{2,2^+}$  by  $\Psi_y = \tilde{\Psi}(y, \cdot)$ .

We note that for every Borel map  $\sigma : Y \rightarrow \mathbb{FS}^{\text{ns}}$ , the pointwise evaluation

$$\begin{aligned} \Psi \bullet \sigma : Y &\rightarrow \mathbb{FS}^{\text{ns}} \cup \mathbb{FS}^{2,2^+} \\ y &\mapsto \Psi_y(\sigma(y)) = \tilde{\Psi}(y, \sigma(y)) \end{aligned}$$

coincides almost everywhere with the  $(\mathcal{G}, \rho, \rho')$ -image of  $\sigma$ . Indeed, using countability of  $\mathbb{FS}$ , this follows from the definition after partitioning  $Y$  into countably many Borel subsets on which  $\sigma$  is constant.

Let us prove that for a.e.  $y \in Y$ , the map  $\Psi_y : \mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}^{\text{ns}} \cup \mathbb{FS}^{2,2^+}$  is injective. Otherwise, there exists a Borel subset  $U \subseteq Y$  of positive measure, two distinct splittings  $S_1, S_2 \in \mathbb{FS}^{\text{ns}}$  and  $S' \in \mathbb{FS}^{\text{ns}} \cup \mathbb{FS}^{2,2^+}$  such that for all  $y \in U$ , one has  $\Psi_y(S_1) = \Psi_y(S_2) = S'$ . Thus, the  $(\mathcal{G}|_U, \rho)$ -stabilizer  $\mathcal{H}_1$  of  $S_1$  and the  $(\mathcal{G}|_U, \rho)$ -stabilizer  $\mathcal{H}_2$  of  $S_2$  are both stably equal to the  $(\mathcal{G}|_U, \rho')$ -stabilizer of  $S'$ . This is a contradiction to Lemma 16.4 which shows that  $\mathcal{H}_1$  is not stably equal to  $\mathcal{H}_2$ .

Second, for a.e.  $y \in Y$ , the map  $\Psi_y$  preserves adjacency and non-adjacency. Indeed, given a pair of splittings  $S_1, S_2 \in \mathbb{FS}^{\text{ns}}$ , up to partitioning  $Y$ , one may restrict to a Borel subset of positive measure  $U \subseteq Y$  such that  $S'_1 := \Psi_y(S_1)$  and  $S'_2 := \Psi_y(S_2)$  do not depend on  $y \in U$ . Lemma 16.9 (applied to  $\mathcal{G}|_U$ ) then says that  $S_1$  and  $S_2$  are compatible if and only if  $S'_1$  and  $S'_2$  are.

Until now, we have proved that for a.e.  $y \in Y$ , the map  $\Psi_y$  belongs to  $\text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS})$ . As recalled above (Proposition 15.1),  $\text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}) = \text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}^{\text{ns}})$ , so we get a Borel map

$$\begin{aligned} \Psi : Y &\rightarrow \text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}^{\text{ns}}) \\ y &\mapsto \Psi_y. \end{aligned}$$

Note that in particular, for any Borel map  $\sigma : Y \rightarrow \mathbb{FS}^{\text{ns}}$ , the  $(\mathcal{G}, \rho, \rho')$ -image of  $\sigma$  also takes its values (essentially) in  $\mathbb{FS}^{\text{ns}}$ .

We finally claim that for a.e.  $y \in Y$ , the map  $\Psi_y$  is actually an automorphism of  $\mathbb{FS}^{\text{ns}}$ . Indeed, let  $\Psi' : Y \rightarrow \text{Inj}(\mathbb{FS}^{\text{ns}} \rightarrow \mathbb{FS}^{\text{ns}})$  be the map obtained in place of  $\Psi$  by reversing the roles of  $\rho$  and  $\rho'$ . The map  $\Psi'$  has the property that for any Borel map  $\sigma' : Y \rightarrow \mathbb{FS}^{\text{ns}}$ , the pointwise evaluation  $\Psi' \bullet \sigma'$  is the  $(\mathcal{G}, \rho', \rho)$ -image of  $\sigma'$ . Then for any  $S \in \mathbb{FS}^{\text{ns}}$ , the map  $\sigma''_S := \Psi' \bullet (\Psi \bullet \sigma_S^0)$  is the essentially unique map  $\sigma''_S : Y \rightarrow \mathbb{FS}^{\text{ns}} \cup \mathbb{FS}^{2,2^+}$  whose  $(\mathcal{G}, \rho)$ -stabilizer stably coincides with the  $(\mathcal{G}, \rho)$ -stabilizer of  $\sigma_S^0$ , so  $\sigma''_S = \sigma_S^0$  almost everywhere

(see Remark 16.7). Since this holds for every  $S \in \text{FS}^{\text{ns}}$ , this means that for almost every  $y \in Y$ , we have  $\Psi'_y \circ \Psi_y = \text{id}_{\text{FS}^{\text{ns}}}$ , and for symmetrical reasons,  $\Psi_y \circ \Psi'_y = \text{id}_{\text{FS}^{\text{ns}}}$  for almost every  $y \in Y$ , which proves our claim.

We have thus constructed a Borel map  $\Psi : Y \rightarrow \text{Aut}(\text{FS}^{\text{ns}})$ . Composing with the isomorphism  $\text{Aut}(\text{FS}^{\text{ns}}) \rightarrow \text{Out}(F_N)$  yields a map  $\varphi : Y \rightarrow \text{Out}(F_N)$  with the property that for every Borel map  $\sigma : Y \rightarrow \text{FS}^{\text{ns}}$ , its  $(\mathcal{G}, \rho, \rho')$ -image is  $\sigma' = \Psi \bullet \sigma$  i.e.  $\sigma'(y) = \varphi(y) \cdot \sigma(y)$  for almost every  $y \in Y$ .

We now check that there exists a conull Borel subset  $Y^* \subseteq Y$  such that the desired cohomology relation  $\rho'(g) = \varphi(r(g))\rho(g)\varphi(s(g))^{-1}$  holds for all  $g \in \mathcal{G}_{|Y^*}$ .

Since  $\mathcal{G}$  is a countable union of bisections, it suffices to fix a bisection  $B$  of  $\mathcal{G}$  and show that the above cohomology relation holds for almost every  $g \in B$ . Without loss of generality, we can also assume that there exist  $\gamma, \gamma' \in \text{IA}_N(\mathbb{Z}/3\mathbb{Z})$  such that  $\rho(B) = \{\gamma\}$  and  $\rho'(B) = \{\gamma'\}$ . Let  $f : U \rightarrow V$  be the partial isomorphism between Borel subsets of  $Y$  associated to  $B$ . It suffices to check that for almost every  $x \in U$ , one has

$$\gamma' = \varphi(f(x))\gamma\varphi(x)^{-1}.$$

Since the action of  $\text{Out}(F_N)$  on  $\text{FS}^{\text{ns}}$  is faithful, it suffices to show that for every Borel map  $\sigma : U \rightarrow \text{FS}^{\text{ns}}$  and almost every  $x \in U$ , one has

$$\gamma'\varphi(x) \cdot \sigma(x) = \varphi(f(x))\gamma \cdot \sigma(x)$$

(it is even enough to prove it when  $\sigma$  is constant).

Let  $\mathcal{H}_U$  be the  $(\mathcal{G}_{|U}, \rho)$ -stabilizer of  $\sigma$ . The  $(\mathcal{G}_{|U}, \rho, \rho')$ -image of  $\sigma$  is  $\sigma' = \Psi \bullet \sigma = \varphi \cdot \sigma$  (i.e.  $\sigma'(x) = \varphi(x) \cdot \sigma(x)$  for almost every  $x \in U$ ), so the  $(\mathcal{G}_{|U}, \rho')$ -stabilizer of  $\sigma'$  stably coincides with  $\mathcal{H}_U$ . Conjugating by  $B$ , we see that the  $(\mathcal{G}_{|V}, \rho)$ -stabilizer of the map  $\sigma_V := \gamma \cdot \sigma \circ f^{-1}$  is  $B\mathcal{H}_UB^{-1}$ . Similarly, the  $(\mathcal{G}_{|V}, \rho')$ -stabilizer of the map  $\sigma'_V := \gamma' \cdot \sigma' \circ f^{-1}$  stably coincides with  $B\mathcal{H}_UB^{-1}$ . It follows that  $\sigma'_V$  is the  $(\mathcal{G}_{|V}, \rho, \rho')$ -image of  $\sigma_V$ , so for almost every  $y \in V$ ,

$$\sigma'_V(y) = \varphi(y) \cdot \sigma_V(y).$$

Replacing  $\sigma'_V$  and  $\sigma_V$  by their definition, we get that for almost every  $y \in V$ ,

$$\gamma' \cdot \sigma'(f^{-1}(y)) = \varphi(y)\gamma \cdot \sigma(f^{-1}(y))$$

and applying this equality to  $y = f(x)$  for almost every  $x \in U$ , we get

$$\gamma' \cdot (\varphi(x) \cdot \sigma(x)) = \varphi(f(x))\gamma \cdot \sigma(x)$$

as desired. □

We conclude this section by proving the remaining theorems from the introduction.

*Proof of Theorems 1.4, 1.5, and Corollary 1.6.* The group  $\text{Out}(F_N)$  is ICC by Lemma 16.2. Theorem 16.3 establishes that  $\text{Out}(F_N)$  is rigid with respect action-type cocycles. The results then follow from Theorem 4.6, Theorem 4.7, and Corollary 4.8. □



## References

- [AD07] C. Anantharaman-Delaroche. Amenability and exactness for groups, group actions and operator algebras. *École thématique. Erwin Schrödinger Institute, Mars 2007. cel-00360390*, 2007.
- [Ada94] S. Adams. Indecomposability of equivalence relations generated by word hyperbolic groups. *Topol.*, 33(4):785–798, 1994.
- [ADR00] C. Anantharaman-Delaroche and J. Renault. *Amenable groupoids*, volume 36 of *Monographies de l’Enseignement Mathématique*. L’Enseignement Mathématique, Geneva, 2000.
- [AEG94] S. Adams, G.A. Elliott, and T. Giordano. Amenable actions of groups. *Trans. Amer. Math. Soc.*, 344(2):803–822, 1994.
- [AKP17] Y. Algom-Kfir and C. Pfaff. Normalizers and centralizers of cyclic subgroups generated by lone axis fully irreducible outer automorphisms. *New York J. Math.*, 23:365–381, 2017.
- [AS11] J. Aramayona and J. Souto. Automorphisms of the graph of free splittings. *Michigan Math. J.*, 60(3):483–493, 2011.
- [BF91] M. Bestvina and M. Feighn. Bounding the complexity of simplicial group actions on trees. *Invent. Math.*, 103(3):449–469, 1991.
- [BF94] M. Bestvina and M. Feighn. Outer limits. 1994. preprint, available at <http://andromeda.rutgers.edu/~feighn/papers/outer.pdf>.
- [BF95] M. Bestvina and M. Feighn. Stable actions of groups on real trees. *Invent. Math.*, 121(2):287–321, 1995.
- [BFS13] U. Bader, A. Furman, and R. Sauer. Integrable measure equivalence and rigidity of hyperbolic lattices. *Invent. Math.*, 194(2):313–379, 2013.
- [BGH17] M. Bestvina, V. Guirardel, and C. Horbez. Boundary amenability of  $\text{Out}(F_N)$ . *arXiv:1705.07017*, 2017.
- [BH92] M. Bestvina and M. Handel. Train tracks and automorphisms of free groups. *Ann. of Math. (2)*, 135(1):1–51, 1992.
- [BIP18] R. Boutonnet, A. Ioana, and J. Peterson. Properly proximal groups and their von Neumann algebras. *arXiv:1809.01881*, 2018.
- [BR15] M. Bestvina and P. Reynolds. The boundary of the complex of free factors. *Duke Math. J.*, 164(11):2213–2251, 2015.
- [BTW07] M.R. Bridson, M. Tweedale, and H. Wilton. Limit groups, positive-genus towers and measure-equivalence. *Ergodic Theory Dynam. Systems*, 27(3):703–712, 2007.
- [BV00] M. Bridson and K. Vogtmann. Automorphisms of automorphism groups of free groups. *J. Algebra*, 229(2):785–792, 2000.
- [BV01] M. Bridson and K. Vogtmann. The symmetries of outer space. *Duke Math. J.*, 106(2):391–409, 2001.
- [CFW81] A. Connes, J. Feldman, and B. Weiss. An amenable equivalence relation is generated by a single transformation. *Ergodic Theory Dynam. Systems*, 1(4):431–450, 1981.
- [CH12] T. Coulbois and A. Hilion. Botany of irreducible automorphisms of free groups. *Pacific J. Math.*, 256(2):291–307, 2012.
- [CK15] I. Chifan and Y. Kida.  $OE$  and  $W^*$  superrigidity results for actions by surface braid groups. *Proc. Lond. Math. Soc. (3)*, 111(6):1431–1470, 2015.
- [CL95] M.M. Cohen and M. Lustig. Very small group actions on  $\mathbb{R}$ -trees and Dehn twist automorphisms. *Topol.*, 34(3):575–617, 1995.
- [CM87] M. Culler and J.W. Morgan. Group actions on  $\mathbb{R}$ -trees. *Proc. London Math. Soc.*, 55(3):571–604, 1987.

- [CV86] M. Culler and K. Vogtmann. Moduli of graphs and automorphisms of free groups. *Invent. Math.*, 84(1):91–119, 1986.
- [DG11] F. Dahmani and V. Guirardel. The isomorphism problem for all hyperbolic groups. *Geom. Funct. Anal.*, 21(2):223–300, 2011.
- [dlST19] M. de la Salle and R. Tessera. Characterizing a vertex-transitive graph by a large ball. *J. Topol.*, 12(3):705–743, 2019.
- [Dye59] H.A. Dye. On groups of measure preserving transformation. I. *Amer. J. Math.*, 81:119–159, 1959.
- [Dye63] H.A. Dye. On groups of measure preserving transformations. II. *Amer. J. Math.*, 85:551–576, 1963.
- [Eps07] I. Epstein. Orbit inequivalent actions of non-amenable groups. *arXiv:0707.4215*, 2007.
- [FH07] B. Farb and M. Handel. Commensurations of  $\text{Out}(F_n)$ . *Publ. Math. Inst. Hautes Études Sci.*, 105(1):1–48, 2007.
- [FM15] S. Francaviglia and A. Martino. Stretching factors, metrics and train tracks for free products. *Illinois J. Math.*, 59(4):859–899, 2015.
- [FP06] K. Fujiwara and P. Papasoglu. JSJ-decompositions of finitely presented groups and complexes of groups. *Geom. Funct. Anal.*, 16(1):70–125, 2006.
- [FSZ89] J. Feldman, C.E. Sutherland, and R.J. Zimmer. Subrelations of ergodic equivalence relations. *Ergodic Theory Dynam. Systems*, 9(2):239–269, 1989.
- [Fur99a] A. Furman. Gromov’s measure equivalence rigidity of higher-rank lattices. *Ann. of Math. (2)*, 150:1059–1081, 1999.
- [Fur99b] A. Furman. Orbit equivalence rigidity. *Ann. of Math. (2)*, 130(3):1083–1108, 1999.
- [Fur11a] A. Furman. Mostow–Margulis rigidity with locally compact targets. *Geom. Funct. Anal.*, 11(1):30–59, 2011.
- [Fur11b] A. Furman. A survey of measured group theory. In *Geometry, rigidity, and group actions*, Chicago Lectures in Math., pages 296–374, Chicago, IL, 2011. Univ. Chicago Press.
- [Gab02a] D. Gaboriau. Invariants  $l^2$  de relations d’équivalence et de groupes. *Publ. Math. Inst. Hautes Études Sci.*, (95):93–150, 2002.
- [Gab02b] D. Gaboriau. On orbit equivalence of measure preserving actions. In *Rigidity in dynamics and geometry (Cambridge, 2000)*, pages 167–186, Berlin, 2002. Springer.
- [Gab05] D. Gaboriau. Examples of groups that are measure equivalent to the free group. *Ergodic Theory Dynam. Systems*, 25(6):1809–1827, 2005.
- [Gab10] D. Gaboriau. Orbit equivalence and measured group theory. In *Proceedings of the International Congress of Mathematicians*, volume III, pages 1501–1527, New Dehli, 2010. Hindustan Book Agency.
- [GG13] S. Garion and Y. Glasner. Highly transitive actions of  $\text{Out}(F_n)$ . *Groups Geom. Dyn.*, 7(2):357–376, 2013.
- [GH19a] V. Guirardel and C. Horbez. Algebraic laminations for free products and arational trees. *Algebr. Geom. Topol.*, 19(5):2283–2400, 2019.
- [GH19b] V. Guirardel and C. Horbez. Boundaries of relative factor graphs and subgroup classification for automorphisms of free products. *arXiv:1901.05046*, 2019.
- [GHL20] V. Guirardel, C. Horbez, and J. Lécureux. Cocycle superrigidity from higher rank lattices to  $\text{Out}(F_N)$ . *arXiv:2005.07477*, 2020.
- [GL] V. Guirardel and G. Levitt. *in preparation*.
- [GL95] D. Gaboriau and G. Levitt. The rank of actions on  $\mathbb{R}$ -trees. *Ann. Sci. École Norm. Sup. (4)*, 28(5):549–570, 1995.

- [GL07a] V. Guirardel and G. Levitt. Deformation spaces of trees. *Groups Geom. Dyn.*, 1(2):135–181, 2007.
- [GL07b] V. Guirardel and G. Levitt. The outer space of a free product. *Proc. London Math. Soc.*, 94(3):695–714, 2007.
- [GL09] D. Gaboriau and R. Lyons. A measurable-group-theoretic solution to von Neumann’s problem. *Invent. Math.*, 177(3):533–540, 2009.
- [GL10] V. Guirardel and G. Levitt. Scott and Swarup’s regular neighbourhood as a tree of cylinders. *Pacific J. Math.*, 245(1):79–98, 2010.
- [GL11] V. Guirardel and G. Levitt. Trees of cylinders and canonical splittings. *Geom. Topol.*, 15(2):977–1012, 2011.
- [GL15a] V. Guirardel and G. Levitt. McCool groups of toral relatively hyperbolic groups. *Algebr. Geom. Topol.*, 15(6):3485–3534, 2015.
- [GL15b] V. Guirardel and G. Levitt. Splittings and automorphisms of relatively hyperbolic groups. *Groups Geom. Dyn.*, 9(2):599–663, 2015.
- [GL17] V. Guirardel and G. Levitt. JSJ decompositions of groups. *Astérisque*, (395):vii+165, 2017.
- [GN19] D. Gaboriau and C. Noûs. On the top-dimensional  $\ell^2$ -Betti numbers. *arXiv:1909.01633*, 2019.
- [GP05] D. Gaboriau and S. Popa. An uncountable family of nonorbit equivalent actions of  $\mathbb{F}_n$ . *J. Amer. Math. Soc.*, 18(3):547–559, 2005.
- [Gro93] M. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory*, volume 2 of *London Math. Soc. Lecture Note Ser.*, pages 1–295, Cambridge Univ. Press, Cambridge, 1993.
- [GS91] S.M. Gersten and J.R. Stallings. Irreducible outer automorphisms of a free group. *Proc. Amer. Math. Soc.*, 111(2):309–314, 1991.
- [Gui00] V. Guirardel. Dynamics of  $\text{Out}(F_n)$  on the boundary of outer space. *Ann. Sci. École Norm. Sup. (4)*, 33(4):433–465, 2000.
- [Gui04] V. Guirardel. Limit groups and groups acting freely on  $\mathbb{R}^n$ -trees. *Geom. Topol.*, 8:1427–1470, 2004.
- [Gui05] V. Guirardel. Cœur et nombre d’intersection pour les actions de groupes sur les arbres. *Ann. Sci. École Norm. Sup. (4)*, 38(6):847–888, 2005.
- [Ham12] U. Hamenstädt. The boundary of the free splitting graph and the free factor graph. *arXiv:1211.1630*, 2012.
- [HH20a] C. Horbez and J. Huang. Boundary amenability and measure equivalence rigidity among two-dimensional Artin groups of hyperbolic type. *arXiv:2004.09325*, 2020.
- [HH20b] C. Horbez and J. Huang. Measure equivalence classification of transvection-free right-angled Artin groups. *arXiv:2010.03613*, 2020.
- [HHL20] C. Horbez, J. Huang, and J. Lécureux. Proper proximality in non-positive curvature. *arXiv:2005.08756*, 2020.
- [Hjo05] G. Hjorth. A converse to Dye’s theorem. *Trans. Amer. Math. Soc.*, 357(8):3083–3103, 2005.
- [HM14] M. Handel and L. Mosher. Relative free splitting and free factor complexes I: Hyperbolicity. *arXiv:1407.3508*, 2014.
- [HM20] M. Handel and L. Mosher. Subgroup decomposition in  $\text{Out}(F_n)$ . *Mem. Amer. Math. Soc.*, 2020.
- [Hor14] C. Horbez. The Tits alternative for the automorphism group of a free product. *arXiv:1408.0546v2*, 2014.
- [Hor17] C. Horbez. The boundary of the outer space of a free product. *Israel J. Math.*, 221(1):179–234, 2017.

- [HR15] C. Houdayer and S. Raum. Baumslag-solitar groups, relative profinite completions and measure equivalence rigidity. *J. Topol.*, 8(1):295–313, 2015.
- [HW20] C. Horbez and R.D. Wade. Commensurations of subgroups of  $\text{Out}(F_N)$ . *Trans. Amer. Math. Soc.*, 373(4):2699–2742, 2020.
- [Ioa11] A. Ioana. Orbit inequivalent actions for groups containing a copy of  $\mathbb{F}_2$ . *Invent. Math.*, 185(1):55–73, 2011.
- [Iva97] N.V. Ivanov. Automorphism of complexes of curves and of Teichmüller spaces. *Int. Math. Res. Not.*, 14:651–666, 1997.
- [JKL02] S. Jackson, A.S. Kechris, and A. Louveau. Countable Borel equivalence relations. *J. Math. Logic*, 2(1):1–80, 2002.
- [Kec95] A.S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [Khr90] D.G. Khramtsov. Completeness of groups of outer automorphisms of free groups. In *Group-theoretic investigations (Russian)*, pages 128–143, Sverdlovsk, 1990. Akad. Nauk SSSR Ural. Otdel.
- [Kid08a] Y. Kida. The mapping class group from the viewpoint of measure equivalence theory. *Mem. Amer. Math. Soc.*, 196(916), 2008.
- [Kid08b] Y. Kida. Orbit equivalence rigidity for ergodic actions of the mapping class group. *Geom. Dedic.*, 131(1):99–109, 2008.
- [Kid09] Y. Kida. Introduction to measurable rigidity of mapping class groups. In *Handbook of Teichmüller theory, volume II*, volume 13 of *IRMA Lect. Math. Theor. Phys.*, pages 297–367, Zürich, 2009. Eur. Math. Soc.
- [Kid10] Y. Kida. Measure equivalence rigidity of the mapping class group. *Ann. of Math. (2)*, 171(3):1851–1901, 2010.
- [Kid11] Y. Kida. Rigidity of amalgamated free products in measure equivalence. *J. Topol.*, 4(3):687–735, 2011.
- [KSS06] I. Kapovich, P. Schupp, and V. Shpilrain. Generic properties of Whitehead’s algorithm and isomorphism rigidity of random one-relator groups. *Pacific J. Math.*, 223(1):113–140, 2006.
- [LdlS20] P.-H. Leemann and M. de la Salle. Cayley graphs with few automorphisms: the case of infinite groups. *arXiv:2010.06020*, 2020.
- [Lev05] G. Levitt. Automorphisms of Hyperbolic Groups and Graphs of Groups. *Geom. Dedic.*, 114(1):49–70, 2005.
- [MM00] H.A. Masur and Y.N. Minsky. Geometry of the complex of curves II. Hierarchical structure. *Geom. Funct. Anal.*, 10(4):902–974, 2000.
- [MS06] N. Monod and Y. Shalom. Orbit equivalence rigidity and bounded cohomology. *Ann. of Math. (2)*, 164:825–878, 2006.
- [MV04] A. Martino and E. Ventura. Fixed subgroups are compressed in free groups. *Comm. Algebra*, 32(10):3921–3935, 2004.
- [OW80] D.S. Ornstein and B. Weiss. Ergodic theory of amenable group actions. I. The Rohlin lemma. *Bull. Amer. Math. Soc. (N.S.)*, 2(1):161–164, 1980.
- [Pan14] S. Pandit. The complex of non-separating embedded spheres. *Rocky Mountain J. Math.*, 44(6):2029–2054, 2014.
- [Pau88] F. Paulin. Topologie de Gromov équivariante, structures hyperboliques et arbres réels. *Invent. Math.*, 94(1):53–80, 1988.
- [Pau89] F. Paulin. The Gromov topology on  $\mathbb{R}$ -trees. *Topology Appl.*, 32(3):197–221, 1989.
- [Pau91] F. Paulin. Outer automorphisms of hyperbolic groups and small actions on  $\mathbb{R}$ -trees. In *Arboreal group theory (Berkeley, CA, 1988)*, pages 331–343. Springer, 1991.

- [Per11] C. Perin. Elementary embeddings in torsion-free hyperbolic groups. *Ann. Sci. Éc. Norm. Supér. (4)*, 44(4):631–681, 2011.
- [Pop07] S. Popa. Cocycle and orbit equivalence superrigidity for malleable actions of  $w$ -rigid groups. *Invent. Math.*, 170(2):243–295, 2007.
- [Ren15] J. Renault. Topological Amenability Is a Borel Property. *Math. Scand.*, 117(1):5–30, 2015.
- [Rey11] P. Reynolds. On indecomposable trees in the boundary of outer space. *Geom. Dedicata*, 153:59–71, 2011.
- [Rey12] P. Reynolds. Reducing systems for very small trees. *arXiv:1211.3378*, 2012.
- [Sel97] Z. Sela. Structure and rigidity in (Gromov) hyperbolic groups and discrete groups in rank 1 Lie groups. II. *Geom. Funct. Anal.*, 7(3):561–593, 1997.
- [Sha05] Y. Shalom. Measurable group theory. In *European Congress of Mathematics*, pages 391–423, Eur. Math. Soc., Zürich, 2005.
- [Sin55] I.M. Singer. Automorphisms of finite factors. *Amer. J. Math.*, 77:117–133, 1955.
- [Sta82] J.R. Stallings. Topologically unrealizable automorphisms of free groups. *Proc. Amer. Math. Soc.*, 84(1):21–24, 1982.
- [Zim78] R.J. Zimmer. Amenable ergodic group actions and an application to Poisson boundaries of random walks. *J. Funct. Anal.*, 27(3):350–372, 1978.
- [Zim80] R.J. Zimmer. Strong rigidity for ergodic actions of semisimple groups. *Ann. of Math. (2)*, 112(3):511–529, 1980.
- [Zim84] R.J. Zimmer. *Ergodic Theory and Semisimple Groups*, volume 81 of *Monographs in Mathematics*. Birkhäuser, Basel; Boston; Stuttgart, 1984.
- [Zim91] R.J. Zimmer. Groups generating transversals to semisimple Lie group actions. *Israel J. Math.*, 73(2):151–159, 1991.

Vincent Guirardel  
 Univ Rennes, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France  
*e-mail*: [vincent.guirardel@univ-rennes1.fr](mailto:vincent.guirardel@univ-rennes1.fr)

Camille Horbez  
 Université Paris-Saclay, CNRS, Laboratoire de mathématiques d'Orsay, 91405, Orsay, France  
*e-mail*: [camille.horbez@universite-paris-saclay.fr](mailto:camille.horbez@universite-paris-saclay.fr)